

Table 2.70C Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	797	783	883	828	832	803
GEOGRAPHIC DIVISION						
Northeast	374	328	372	301	345	304
New England	183	138	149	169	147	165
Middle Atlantic	327	298	341	249	312	254
Midwest	311	298	361	350	347	322
East North Central	252	234	300	295	283	266
West North Central	183	186	202	188	199	180
South	434	485	573	545	529	558
South Atlantic	340	365	454	382	397	397
East South Central	157	190	250	253	252	256
West South Central	220	256	245	303	246	302
West	458	428	418	410	408	369
Mountain	147	175	177	171	182	149
Pacific	434	391	374	362	362	328
COUNTY TYPE¹						
Large Metro	641	621	677	600	635	571
Small Metro	389	399	451	483	414	479
250K – 1 Mil. Pop.	304	322	365	402	334	402
< 250K Pop.	230	239	270	269	251	247
Nonmetro	323	264	345	322	328	304
Urbanized	188	181	211	227	196	222
Less Urbanized	228	183	248	210	235	197
Completely Rural	117	86	128	121	129	105

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.70D Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.34	0.33	0.38	0.35	0.35	0.34
GEOGRAPHIC DIVISION						
Northeast	0.83	0.72	0.83	0.66	0.77	0.67
New England	1.55	1.16	1.26	1.42	1.24	1.39
Middle Atlantic	0.98	0.89	1.03	0.75	0.94	0.76
Midwest	0.58	0.55	0.68	0.65	0.65	0.60
East North Central	0.67	0.62	0.80	0.78	0.76	0.71
West North Central	1.14	1.15	1.26	1.17	1.24	1.12
South	0.52	0.57	0.68	0.64	0.63	0.66
South Atlantic	0.78	0.82	1.04	0.86	0.91	0.89
East South Central	1.11	1.33	1.76	1.78	1.78	1.80
West South Central	0.85	0.98	0.95	1.16	0.95	1.15
West	0.87	0.80	0.79	0.76	0.77	0.69
Mountain	0.96	1.12	1.16	1.09	1.19	0.96
Pacific	1.15	1.02	1.00	0.95	0.96	0.86
COUNTY TYPE¹						
Large Metro	0.51	0.49	0.54	0.48	0.50	0.45
Small Metro	0.56	0.56	0.65	0.68	0.60	0.67
250K – 1 Mil. Pop.	0.67	0.69	0.80	0.86	0.73	0.86
< 250K Pop.	0.97	0.98	1.13	1.10	1.06	1.01
Nonmetro	0.81	0.66	0.86	0.80	0.82	0.76
Urbanized	1.17	1.01	1.32	1.27	1.22	1.24
Less Urbanized	1.16	1.01	1.26	1.15	1.20	1.08
Completely Rural	2.64	2.10	2.88	2.96	2.90	2.57

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.70P Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.4499		0.0635		0.1893	
GEOGRAPHIC DIVISION						
Northeast	0.2119		0.0969		0.4805	
New England	0.8446		0.7498		0.6268	
Middle Atlantic	0.1118		0.0897		0.5848	
Midwest	0.9944		0.9230		0.8104	
East North Central	0.7884		0.7240		0.3601	
West North Central	0.7348		0.7443		0.3789	
South	0.3570		0.7562		0.9479	
South Atlantic	0.8681		0.1248		0.1791	
East South Central	0.1143		0.4192		0.7003	
West South Central	0.3641		0.3183		0.1120	
West	0.6236		0.0691		0.0956	
Mountain	0.8798		0.9848		0.9121	
Pacific	0.5727		0.0428		0.0683	
COUNTY TYPE¹						
Large Metro	0.1683		0.0059		0.0259	
Small Metro	0.6294		0.6596		0.5160	
250K – 1 Mil. Pop.	0.2684		0.2414		0.2146	
< 250K Pop.	0.0116		0.3037		0.4666	
Nonmetro	0.1277		0.7808		0.8273	
Urbanized	0.0918		0.5786		0.4111	
Less Urbanized	0.4142		0.4626		0.4259	
Completely Rural	0.7563		0.7091		0.5076	

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.71C Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	108	113	92	93	81	80
GEOGRAPHIC DIVISION						
Northeast	41	44	36	39	33	32
New England	18	23	18	18	15	16
Middle Atlantic	37	37	31	35	30	28
Midwest	41	43	40	40	35	37
East North Central	32	34	30	30	26	30
West North Central	25	27	27	26	24	22
South	75	70	61	58	53	51
South Atlantic	51	46	41	39	39	36
East South Central	38	30	34	27	28	23
West South Central	38	44	28	35	24	28
West	52	64	43	47	35	38
Mountain	23	28	22	24	21	19
Pacific	46	57	38	40	29	32
COUNTY TYPE¹						
Large Metro	83	84	69	69	58	60
Small Metro	55	58	48	46	42	40
250K – 1 Mil. Pop.	46	49	39	40	35	35
< 250K Pop.	32	33	29	28	25	24
Nonmetro	39	44	38	39	32	31
Urbanized	27	28	24	25	22	20
Less Urbanized	28	29	27	27	22	22
Completely Rural	14	14	15	12	13	10

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.71D Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.44	0.45	0.37	0.37	0.33	0.32
GEOGRAPHIC DIVISION						
Northeast	0.93	0.97	0.80	0.87	0.75	0.71
New England	1.57	1.96	1.52	1.50	1.27	1.38
Middle Atlantic	1.12	1.11	0.94	1.06	0.90	0.83
Midwest	0.72	0.76	0.71	0.71	0.62	0.66
East North Central	0.81	0.86	0.75	0.76	0.65	0.75
West North Central	1.45	1.57	1.59	1.55	1.39	1.30
South	0.86	0.78	0.69	0.65	0.61	0.57
South Atlantic	1.17	1.03	0.94	0.87	0.88	0.81
East South Central	2.66	2.07	2.36	1.85	1.94	1.58
West South Central	1.27	1.48	0.95	1.17	0.80	0.93
West	0.89	1.09	0.74	0.80	0.60	0.64
Mountain	1.37	1.63	1.29	1.37	1.20	1.08
Pacific	1.13	1.38	0.91	0.97	0.69	0.78
COUNTY TYPE¹						
Large Metro	0.63	0.63	0.52	0.52	0.44	0.45
Small Metro	0.76	0.78	0.67	0.62	0.59	0.53
250K – 1 Mil. Pop.	0.94	0.99	0.80	0.80	0.72	0.71
< 250K Pop.	1.36	1.36	1.23	1.15	1.05	0.99
Nonmetro	0.92	1.03	0.87	0.90	0.74	0.73
Urbanized	1.57	1.53	1.38	1.35	1.25	1.09
Less Urbanized	1.33	1.46	1.28	1.35	1.05	1.13
Completely Rural	2.88	3.23	3.12	2.79	2.68	2.29

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.71P Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.0002		0.0438		0.0797	
GEOGRAPHIC DIVISION						
Northeast	0.0921		0.2614		0.4065	
New England	0.3664		0.7460		0.4195	
Middle Atlantic	0.1539		0.2789		0.5536	
Midwest	0.0014		0.1323		0.1169	
East North Central	0.0189		0.2556		0.1312	
West North Central	0.0270		0.3203		0.4958	
South	0.0085		0.0312		0.0446	
South Atlantic	0.0003		0.0003		0.0032	
East South Central	0.2217		0.7917		0.7507	
West South Central	0.4608		0.2749		0.4860	
West	0.7092		0.3578		0.2016	
Mountain	0.9012		0.7004		0.7732	
Pacific	0.7195		0.4024		0.1771	
COUNTY TYPE¹						
Large Metro	0.0000		0.0055		0.0330	
Small Metro	0.7055		0.8946		0.8418	
250K – 1 Mil. Pop.	0.6291		0.5258		0.5455	
< 250K Pop.	0.9707		0.5261		0.5916	
Nonmetro	0.3450		0.8614		0.7304	
Urbanized	0.7889		0.4956		0.6990	
Less Urbanized	0.7889		0.5318		0.9772	
Completely Rural	0.0058		0.1050		0.1833	

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.72C Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	132	142	154	152	150	151
GEOGRAPHIC DIVISION						
Northeast	53	52	63	58	60	60
New England	33	23	40	26	26	20
Middle Atlantic	41	46	49	52	54	56
Midwest	54	56	53	69	62	68
East North Central	47	45	44	59	51	59
West North Central	27	33	30	35	35	33
South	79	93	90	92	89	92
South Atlantic	62	74	66	66	63	68
East South Central	32	33	42	39	46	40
West South Central	38	46	45	51	45	48
West	70	77	86	79	80	76
Mountain	29	29	36	41	35	35
Pacific	64	71	77	68	71	67
COUNTY TYPE¹						
Large Metro	96	108	117	120	108	117
Small Metro	75	79	88	78	89	77
250K – 1 Mil. Pop.	56	67	71	66	71	63
< 250K Pop.	50	43	53	44	55	44
Nonmetro	49	45	52	50	59	53
Urbanized	30	28	32	33	37	36
Less Urbanized	36	32	38	36	41	38
Completely Rural	15	10	18	14	17	14

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.72D Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.43	0.45	0.50	0.48	0.48	0.48
GEOGRAPHIC DIVISION						
Northeast	0.97	0.92	1.14	1.03	1.09	1.06
New England	2.32	1.59	2.79	1.78	1.79	1.39
Middle Atlantic	1.01	1.12	1.20	1.24	1.33	1.35
Midwest	0.76	0.76	0.74	0.94	0.86	0.93
East North Central	0.95	0.89	0.89	1.17	1.03	1.17
West North Central	1.21	1.45	1.33	1.55	1.59	1.47
South	0.72	0.82	0.81	0.81	0.80	0.81
South Atlantic	1.15	1.32	1.23	1.18	1.16	1.21
East South Central	1.68	1.70	2.20	2.00	2.40	2.05
West South Central	1.02	1.20	1.21	1.35	1.22	1.25
West	0.96	1.03	1.18	1.07	1.09	1.02
Mountain	1.31	1.29	1.66	1.82	1.61	1.54
Pacific	1.24	1.37	1.50	1.32	1.39	1.30
COUNTY TYPE¹						
Large Metro	0.60	0.65	0.74	0.72	0.68	0.70
Small Metro	0.74	0.77	0.87	0.76	0.88	0.76
250K – 1 Mil. Pop.	0.87	1.01	1.10	0.99	1.11	0.94
< 250K Pop.	1.33	1.20	1.43	1.25	1.47	1.25
Nonmetro	0.99	0.92	1.05	1.01	1.18	1.08
Urbanized	1.32	1.18	1.42	1.40	1.67	1.52
Less Urbanized	1.58	1.46	1.67	1.65	1.81	1.75
Completely Rural	3.24	2.82	3.83	3.78	3.70	3.91

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.72P Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.2277		0.0977		0.4260	
GEOGRAPHIC DIVISION						
Northeast	0.2555		0.2508		0.1427	
New England	0.7773		0.8315		0.3832	
Middle Atlantic	0.2271		0.1951		0.2292	
Midwest	0.8324		0.1417		0.1732	
East North Central	0.5926		0.1116		0.1313	
West North Central	0.6105		0.8277		0.9256	
South	0.0443		0.3053		0.7281	
South Atlantic	0.1808		0.4379		0.7188	
East South Central	0.0183		0.2335		0.8288	
West South Central	0.8322		0.8935		0.1324	
West	0.3744		0.9906		0.7325	
Mountain	0.8471		0.3220		0.3888	
Pacific	0.2980		0.6002		0.4479	
COUNTY TYPE¹						
Large Metro	0.1345		0.0604		0.1760	
Small Metro	0.7680		0.6789		0.2536	
250K – 1 Mil. Pop.	0.9107		0.3592		0.3096	
< 250K Pop.	0.7491		0.5889		0.5753	
Nonmetro	0.6404		0.3219		0.4039	
Urbanized	0.6582		0.9460		0.7565	
Less Urbanized	0.9164		0.1362		0.2637	
Completely Rural	0.2950		0.6122		0.4244	

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.73C Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	763	729	848	774	797	748
GEOGRAPHIC DIVISION						
Northeast	355	325	366	278	338	284
New England	173	136	159	156	150	158
Middle Atlantic	312	296	330	229	303	234
Midwest	283	267	336	327	324	297
East North Central	221	212	280	270	267	241
West North Central	177	163	185	183	183	174
South	416	449	549	501	500	512
South Atlantic	317	330	433	362	374	369
East South Central	147	181	230	216	228	219
West South Central	223	241	247	275	243	281
West	450	392	406	396	400	354
Mountain	138	155	179	168	180	148
Pacific	427	359	360	348	352	312
COUNTY TYPE¹						
Large Metro	620	581	658	576	618	540
Small Metro	384	363	426	447	387	449
250K – 1 Mil. Pop.	298	297	338	375	303	381
< 250K Pop.	229	215	251	251	239	232
Nonmetro	307	253	335	305	314	288
Urbanized	185	169	207	214	193	212
Less Urbanized	216	179	237	194	221	182
Completely Rural	118	88	118	119	122	103

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.73D Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.43	0.40	0.47	0.43	0.44	0.41
GEOGRAPHIC DIVISION						
Northeast	1.01	0.92	1.04	0.79	0.96	0.81
New England	1.87	1.47	1.72	1.69	1.62	1.70
Middle Atlantic	1.20	1.14	1.28	0.88	1.17	0.90
Midwest	0.70	0.66	0.83	0.80	0.80	0.73
East North Central	0.78	0.74	0.98	0.95	0.94	0.84
West North Central	1.46	1.34	1.53	1.51	1.52	1.43
South	0.65	0.70	0.86	0.78	0.78	0.79
South Atlantic	0.93	0.96	1.28	1.05	1.10	1.07
East South Central	1.36	1.67	2.13	1.99	2.11	2.02
West South Central	1.16	1.25	1.29	1.42	1.26	1.45
West	1.13	0.97	1.02	0.98	1.01	0.87
Mountain	1.21	1.33	1.56	1.44	1.58	1.27
Pacific	1.51	1.24	1.27	1.21	1.24	1.08
COUNTY TYPE¹						
Large Metro	0.64	0.60	0.68	0.60	0.64	0.56
Small Metro	0.74	0.68	0.82	0.83	0.75	0.83
250K – 1 Mil. Pop.	0.87	0.84	0.99	1.06	0.89	1.08
< 250K Pop.	1.29	1.16	1.42	1.36	1.35	1.26
Nonmetro	1.00	0.82	1.09	0.98	1.02	0.93
Urbanized	1.54	1.24	1.72	1.57	1.60	1.55
Less Urbanized	1.41	1.27	1.55	1.38	1.45	1.29
Completely Rural	3.37	2.66	3.39	3.62	3.49	3.13

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.73P Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.9643		0.1433		0.2724	
GEOGRAPHIC DIVISION						
Northeast	0.3555		0.1526		0.6788	
New England	0.7178		0.7908		0.7652	
Middle Atlantic	0.1873		0.1406		0.7535	
Midwest	0.6123		0.7745		0.8562	
East North Central	0.4676		1.0000		0.6205	
West North Central	0.9212		0.6297		0.3297	
South	0.9261		0.9706		0.9578	
South Atlantic	0.2838		0.2658		0.2876	
East South Central	0.3125		0.2814		0.6277	
West South Central	0.2967		0.3952		0.1846	
West	0.6982		0.0524		0.0590	
Mountain	0.9061		0.8672		1.0000	
Pacific	0.6545		0.0250		0.0324	
COUNTY TYPE¹						
Large Metro	0.5516		0.0223		0.0533	
Small Metro	0.6526		0.6301		0.5921	
250K – 1 Mil. Pop.	0.2291		0.2604		0.2504	
< 250K Pop.	0.0095		0.3679		0.4175	
Nonmetro	0.1013		0.8971		0.9830	
Urbanized	0.1024		0.5516		0.3643	
Less Urbanized	0.3895		0.5017		0.4948	
Completely Rural	0.9504		0.8395		0.5481	

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.74C Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	780	755	876	810	829	784
GEOGRAPHIC DIVISION						
Northeast	360	334	373	296	345	296
New England	173	142	150	165	147	162
Middle Atlantic	317	303	341	245	312	247
Midwest	298	281	355	348	338	316
East North Central	237	222	295	292	277	261
West North Central	181	173	199	189	194	179
South	426	455	566	524	524	537
South Atlantic	324	337	445	372	391	384
East South Central	155	180	247	231	250	237
West South Central	228	244	249	291	247	293
West	455	415	416	403	413	365
Mountain	143	163	181	168	182	146
Pacific	431	381	370	355	367	324
COUNTY TYPE¹						
Large Metro	627	601	668	586	630	554
Small Metro	391	381	451	472	417	471
250K – 1 Mil. Pop.	308	308	361	395	331	397
< 250K Pop.	228	227	270	263	255	243
Nonmetro	320	258	342	319	325	302
Urbanized	189	174	210	224	196	221
Less Urbanized	225	180	245	207	233	195
Completely Rural	120	89	125	120	127	104

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.74D Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.37	0.35	0.42	0.38	0.39	0.37
GEOGRAPHIC DIVISION						
Northeast	0.89	0.82	0.92	0.72	0.85	0.73
New England	1.62	1.33	1.41	1.54	1.38	1.51
Middle Atlantic	1.06	1.01	1.14	0.81	1.04	0.82
Midwest	0.63	0.59	0.74	0.72	0.71	0.66
East North Central	0.71	0.66	0.88	0.87	0.83	0.77
West North Central	1.27	1.20	1.39	1.31	1.35	1.24
South	0.57	0.60	0.75	0.69	0.70	0.71
South Atlantic	0.82	0.84	1.13	0.93	0.99	0.96
East South Central	1.22	1.41	1.94	1.81	1.97	1.86
West South Central	1.00	1.06	1.09	1.26	1.08	1.27
West	0.97	0.87	0.89	0.84	0.88	0.76
Mountain	1.05	1.17	1.33	1.21	1.34	1.05
Pacific	1.29	1.12	1.11	1.04	1.10	0.95
COUNTY TYPE¹						
Large Metro	0.56	0.53	0.59	0.52	0.56	0.49
Small Metro	0.63	0.60	0.73	0.74	0.67	0.74
250K – 1 Mil. Pop.	0.76	0.73	0.89	0.94	0.82	0.95
< 250K Pop.	1.06	1.03	1.26	1.19	1.19	1.10
Nonmetro	0.89	0.72	0.96	0.89	0.91	0.84
Urbanized	1.32	1.09	1.47	1.40	1.37	1.38
Less Urbanized	1.29	1.11	1.40	1.27	1.33	1.20
Completely Rural	3.03	2.43	3.15	3.27	3.21	2.84

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.74P Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.7895		0.0973		0.2451	
GEOGRAPHIC DIVISION						
Northeast	0.2729		0.1198		0.5192	
New England	0.7622		0.7811		0.6757	
Middle Atlantic	0.1334		0.1074		0.6102	
Midwest	0.6557		0.9389		0.9433	
East North Central	0.5581		0.8108		0.4391	
West North Central	0.9833		0.6447		0.3339	
South	0.6208		0.9393		0.8882	
South Atlantic	0.4477		0.2429		0.2897	
East South Central	0.1626		0.4010		0.6713	
West South Central	0.2971		0.3760		0.1193	
West	0.6001		0.0500		0.0721	
Mountain	0.8759		0.9691		0.8813	
Pacific	0.5470		0.0305		0.0505	
COUNTY TYPE¹						
Large Metro	0.3958		0.0132		0.0410	
Small Metro	0.6311		0.6545		0.5113	
250K – 1 Mil. Pop.	0.2633		0.2156		0.1970	
< 250K Pop.	0.0090		0.2680		0.4388	
Nonmetro	0.0950		0.7585		0.8435	
Urbanized	0.0975		0.6409		0.4521	
Less Urbanized	0.4011		0.4162		0.4274	
Completely Rural	0.9512		0.8301		0.5865	

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.