

**Table 6.22C Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003**

Age Category	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	571	562	123	150	445	401	528	529
18	29	30	9	13	24	23	22	26
19	27	29	9	9	22	22	22	25
20	27	28	7	9	22	21	23	24
21	29	30	10	12	25	24	24	28
22	29	31	9	9	26	24	24	27
23	31	32	11	8	26	27	27	28
24	26	29	5	8	22	26	23	26
25	28	32	9	6	24	25	24	26
26-29	127	106	40	38	108	92	105	95
30-34	146	147	23	45	120	122	121	127
35-39	154	167	40	34	132	126	141	152
40-44	178	161	52	45	141	130	164	149
45-49	169	186	47	56	136	145	156	168
50-54	235	236	23	59	201	173	218	217
55-59	202	195	33	50	153	128	197	190
60-64	139	155	37	25	98	99	125	152
65 or Older	266	236	50	80	169	146	252	220

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.22D Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003**

Age Category	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	0.27	0.26	0.06	0.07	0.21	0.19	0.25	0.25
18	0.66	0.66	0.20	0.29	0.55	0.49	0.51	0.57
19	0.67	0.74	0.22	0.23	0.55	0.58	0.53	0.64
20	0.67	0.71	0.19	0.23	0.56	0.52	0.57	0.60
21	0.73	0.77	0.25	0.32	0.61	0.60	0.60	0.70
22	0.74	0.77	0.22	0.21	0.67	0.61	0.61	0.67
23	0.82	0.79	0.30	0.21	0.67	0.67	0.72	0.69
24	0.75	0.80	0.15	0.22	0.61	0.72	0.64	0.70
25	0.84	0.87	0.26	0.18	0.73	0.69	0.73	0.73
26-29	0.83	0.72	0.26	0.26	0.71	0.63	0.69	0.64
30-34	0.74	0.73	0.12	0.22	0.61	0.61	0.61	0.63
35-39	0.70	0.80	0.18	0.16	0.60	0.60	0.64	0.73
40-44	0.82	0.71	0.24	0.20	0.64	0.57	0.75	0.66
45-49	0.80	0.87	0.22	0.26	0.64	0.68	0.74	0.79
50-54	1.25	1.18	0.12	0.30	1.07	0.87	1.16	1.09
55-59	1.38	1.31	0.23	0.34	1.04	0.86	1.35	1.28
60-64	1.20	1.32	0.32	0.21	0.85	0.85	1.09	1.29
65 or Older	0.79	0.69	0.15	0.23	0.50	0.43	0.74	0.64

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.22P Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Age Category	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	0.5909		0.1282		0.3304		0.1790	
18	0.3969		0.2102		0.6638		0.1587	
19	0.7222		0.3584		0.5100		0.4711	
20	0.5694		0.3114		0.3825		0.3440	
21	0.8667		0.7987		0.4874		0.1829	
22	0.4068		0.6341		0.3313		0.3271	
23	0.6839		0.7817		0.6166		0.5991	
24	0.5902		0.3519		0.1158		0.3932	
25	0.4853		0.3026		0.8780		0.8243	
26-29	0.6779		0.9576		0.6545		0.9451	
30-34	0.9667		0.2875		0.8452		0.7350	
35-39	0.6143		0.5544		0.9799		0.2264	
40-44	0.0790		0.5145		0.3516		0.1088	
45-49	0.1431		0.1817		0.7997		0.0287	
50-54	0.6693		0.2092		0.9863		0.5641	
55-59	0.6105		0.2975		0.3095		0.6740	
60-64	0.1628		0.5944		0.6207		0.0702	
65 or Older	0.7953		0.3533		0.3485		0.6165	

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.23C Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older with Past Year Serious Mental Illness, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003**

Age Category	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	294	336	77	126	252	280	281	312
18	18	20	5	9	14	15	14	18
19	19	22	5	8	15	19	15	19
20	16	20	5	6	14	14	15	18
21	20	20	4	8	18	17	18	18
22	20	19	8	8	18	14	18	18
23	23	21	9	6	19	19	21	20
24	19	24	3	4	16	20	16	20
25	22	22	7	6	19	18	18	19
26-29	68	81	13	37	57	73	62	73
30-34	99	90	19	35	79	76	88	81
35-39	107	97	18	31	93	80	102	92
40-44	114	102	42	30	100	86	109	95
45-49	103	105	40	49	81	90	95	99
50-54	*	149	16	59	*	121	*	139
55-59	*	*	*	*	*	*	*	*
60-64	*	*	*	*	*	*	*	*
65 or Older	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.23D Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older with Past Year Serious Mental Illness, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003**

Age Category	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	1.27	1.13	0.45	0.62	1.22	1.12	1.29	1.12
18	2.95	2.68	0.99	1.31	2.36	2.20	2.51	2.48
19	2.95	3.14	0.89	1.20	2.42	2.74	2.53	2.85
20	2.84	2.76	1.03	0.99	2.55	2.29	2.70	2.59
21	2.78	3.10	0.67	1.45	2.66	2.81	2.69	2.94
22	3.19	3.06	1.43	1.45	2.96	2.52	2.97	2.92
23	3.25	3.38	1.65	1.20	3.03	3.17	3.15	3.23
24	3.50	3.41	0.77	0.76	3.20	3.08	3.21	3.18
25	3.45	3.29	1.50	1.15	3.40	3.10	3.18	3.13
26-29	3.63	3.68	0.89	2.05	3.32	3.57	3.43	3.49
30-34	3.27	3.31	0.87	1.74	3.02	3.25	3.19	3.20
35-39	3.52	3.17	0.92	1.33	3.28	2.95	3.75	3.21
40-44	3.34	3.25	1.81	1.30	3.38	3.09	3.40	3.27
45-49	3.28	3.39	2.21	2.08	3.76	3.12	3.43	3.32
50-54	*	5.16	1.17	3.23	*	5.33	*	5.33
55-59	*	*	*	*	*	*	*	*
60-64	*	*	*	*	*	*	*	*
65 or Older	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.23P Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older with Past Year Serious Mental Illness, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Age Category	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	0.7214		0.0193		0.2959		0.8898	
18	0.6772		0.4479		0.9418		0.2646	
19	0.7311		0.9637		0.5348		0.8629	
20	0.4141		0.8772		0.8508		0.4473	
21	0.7290		0.0751		0.8318		0.8154	
22	0.2223		0.4818		0.5753		0.2552	
23	0.7291		0.8866		0.3071		0.5868	
24	0.4743		0.9923		0.2128		0.7112	
25	0.5419		0.3976		0.5434		0.7434	
26-29	0.6316		0.0672		0.4095		0.6869	
30-34	0.8230		0.8408		0.9164		0.6326	
35-39	0.3456		0.4219		0.3107		0.7245	
40-44	0.1054		0.5764		0.2574		0.1656	
45-49	0.0078		0.3593		0.5426		0.0744	
50-54	*		0.0737		*		*	
55-59	*		*		*		*	
60-64	*		*		*		*	
65 or Older	*		*		*		*	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.24C Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Demographic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	571	562	123	150	445	401	528	529
<b>AGE</b>								
18-25	78	90	25	28	66	66	68	78
26-49	361	356	93	95	295	283	317	322
50 or Older	424	425	76	113	309	281	405	409
<b>GENDER</b>								
Male	318	319	83	112	242	226	279	285
Female	476	448	91	102	400	323	432	433
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	555	528	112	131	437	384	513	495
White	518	504	88	101	395	362	483	473
Black or African American	138	151	60	82	127	110	124	133
American Indian or Alaska Native	40	33	15	19	30	19	37	26
Native Hawaiian or Other Pacific Islander	11	*	1	0	3	3	10	*
Asian	118	89	23	11	120	62	134	81
Two or More Races	56	59	10	9	40	48	49	49
Hispanic or Latino	153	165	54	69	113	119	136	159
<b>EDUCATION</b>								
< High School	233	200	81	80	152	141	210	194
High School Graduate	323	309	68	100	231	218	302	294
Some College	282	279	37	67	241	192	261	271
College Graduate	328	286	53	48	270	231	291	251
<b>CURRENT EMPLOYMENT</b>								
Full-Time	382	401	62	56	289	287	338	364
Part-Time	190	200	43	38	154	160	179	183
Unemployed	103	87	33	32	91	66	94	73
Other <sup>3</sup>	362	338	93	131	270	231	346	320

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.24D Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Demographic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.27	0.26	0.06	0.07	0.21	0.19	0.25	0.25
<b>AGE</b>								
18-25	0.25	0.28	0.08	0.09	0.21	0.21	0.22	0.25
26-49	0.36	0.36	0.09	0.10	0.29	0.28	0.32	0.32
50 or Older	0.54	0.53	0.10	0.14	0.39	0.35	0.51	0.51
<b>GENDER</b>								
Male	0.32	0.31	0.08	0.11	0.24	0.22	0.28	0.28
Female	0.44	0.41	0.08	0.09	0.37	0.29	0.40	0.39
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.30	0.28	0.06	0.07	0.24	0.21	0.28	0.27
White	0.35	0.34	0.06	0.07	0.26	0.24	0.32	0.31
Black or African American	0.60	0.64	0.26	0.34	0.55	0.47	0.54	0.56
American Indian or Alaska Native	3.12	3.02	1.15	1.72	2.35	1.76	2.91	2.41
Native Hawaiian or Other Pacific Islander	1.49	*	0.08	0.02	0.48	0.82	1.43	*
Asian	1.48	1.02	0.29	0.13	1.49	0.71	1.67	0.93
Two or More Races	2.65	2.74	0.46	0.44	1.91	2.24	2.32	2.29
Hispanic or Latino	0.61	0.64	0.22	0.27	0.45	0.46	0.54	0.61
<b>EDUCATION</b>								
< High School	0.63	0.53	0.22	0.21	0.41	0.37	0.57	0.51
High School Graduate	0.48	0.46	0.10	0.15	0.34	0.33	0.45	0.44
Some College	0.54	0.53	0.07	0.13	0.46	0.37	0.50	0.52
College Graduate	0.62	0.52	0.10	0.09	0.51	0.42	0.55	0.46
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.33	0.34	0.05	0.05	0.25	0.25	0.29	0.31
Part-Time	0.69	0.70	0.16	0.13	0.56	0.56	0.65	0.64
Unemployed	1.36	1.08	0.44	0.39	1.21	0.82	1.24	0.91
Other <sup>3</sup>	0.62	0.58	0.16	0.22	0.46	0.39	0.59	0.54

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 6.24P Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Demographic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.5909		0.1282		0.3304		0.1790	
<b>AGE</b>								
18-25	0.1252		0.5458		0.8159		0.0175	
26-49	0.9440		0.5744		0.6133		0.2910	
50 or Older	0.6813		0.1332		0.4691		0.6141	
<b>GENDER</b>								
Male	0.6772		0.4833		0.8383		0.8279	
Female	0.3384		0.1491		0.2190		0.0742	
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.5091		0.2526		0.3904		0.1668	
White	0.2308		0.3275		0.9213		0.0983	
Black or African American	0.9908		0.4748		0.1617		0.4574	
American Indian or Alaska Native	0.1943		0.7059		0.3396		0.1295	
Native Hawaiian or Other Pacific Islander	*		0.3031		0.5083		*	
Asian	0.0461		0.5855		0.0363		0.2144	
Two or More Races	0.8159		0.9191		0.7128		0.7685	
Hispanic or Latino	0.8098		0.2532		0.7107		0.8774	
<b>EDUCATION</b>								
< High School	0.1614		0.6921		0.5497		0.4831	
High School Graduate	0.2010		0.0633		0.8823		0.3063	
Some College	0.4812		0.0177		0.3476		0.0688	
College Graduate	0.9721		0.4804		0.4864		0.8248	
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.2098		0.8342		0.6944		0.0357	
Part-Time	0.7593		0.4358		0.7786		0.6840	
Unemployed	0.4846		0.6303		0.2017		0.5388	
Other <sup>3</sup>	0.5972		0.0354		0.1264		0.8628	

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.25C Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older with Past Year Serious Mental Illness, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Demographic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	294	336	77	126	252	280	281	312
<b>AGE</b>								
18-25	60	63	17	20	51	51	52	57
26-49	220	215	65	79	183	186	203	200
50 or Older	192	236	36	95	166	182	187	220
<b>GENDER</b>								
Male	181	188	54	94	150	151	172	171
Female	226	272	54	85	199	222	217	257
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	278	319	75	112	241	266	266	297
White	248	307	59	85	211	247	236	288
Black or African American	111	97	40	70	105	87	100	73
American Indian or Alaska Native	*	*	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	27	*	11	*	*	*	*
Two or More Races	*	*	*	*	*	*	*	*
Hispanic or Latino	87	90	20	57	74	74	81	88
<b>EDUCATION</b>								
< High School	129	136	49	65	101	113	121	129
High School Graduate	176	187	35	80	152	156	170	176
Some College	160	159	29	58	146	125	150	154
College Graduate	130	154	39	48	107	131	123	139
<b>CURRENT EMPLOYMENT</b>								
Full-Time	190	208	31	42	151	171	175	188
Part-Time	119	124	34	25	85	114	114	118
Unemployed	76	55	12	23	70	48	73	50
Other <sup>3</sup>	187	205	63	113	173	162	182	197

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.25D Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older with Past Year Serious Mental Illness, by Demographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Demographic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	1.27	1.13	0.45	0.62	1.22	1.12	1.29	1.12
<b>AGE</b>								
18-25	1.17	1.10	0.42	0.44	1.07	0.97	1.08	1.03
26-49	1.54	1.49	0.68	0.75	1.53	1.44	1.56	1.47
50 or Older	3.91	3.35	0.92	1.91	3.65	3.08	3.87	3.35
<b>GENDER</b>								
Male	2.24	2.07	0.87	1.30	2.09	1.92	2.21	2.00
Female	1.54	1.36	0.47	0.65	1.49	1.34	1.55	1.34
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	1.32	1.18	0.48	0.63	1.29	1.20	1.34	1.19
White	1.50	1.32	0.46	0.59	1.41	1.32	1.51	1.33
Black or African American	3.88	3.63	1.91	3.21	3.92	3.58	3.85	3.20
American Indian or Alaska Native	*	*	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	4.54	*	1.89	*	*	*	*
Two or More Races	*	*	*	*	*	*	*	*
Hispanic or Latino	3.91	3.20	1.11	2.29	3.67	2.74	3.82	3.11
<b>EDUCATION</b>								
< High School	3.09	2.59	1.34	1.47	2.57	2.33	2.96	2.56
High School Graduate	2.23	2.03	0.59	1.20	2.11	2.00	2.19	2.10
Some College	2.09	1.91	0.58	1.04	2.22	1.94	2.07	1.88
College Graduate	2.81	3.02	1.32	1.32	2.58	2.90	2.93	2.76
<b>CURRENT EMPLOYMENT</b>								
Full-Time	1.57	1.63	0.37	0.44	1.45	1.49	1.56	1.53
Part-Time	2.85	2.82	1.25	0.85	2.59	3.02	2.98	2.92
Unemployed	5.13	3.41	1.13	1.83	5.06	3.23	5.12	3.24
Other <sup>3</sup>	2.76	2.31	1.17	1.74	2.71	2.24	2.75	2.49

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.25P Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older with Past Year Serious Mental Illness, by Demographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Demographic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.7214		0.0193		0.2959		0.8898	
<b>AGE</b>								
18-25	0.5307		0.4014		0.9783		0.2975	
26-49	0.0202		0.1766		0.1301		0.1477	
50 or Older	0.1293		0.0595		0.9465		0.2294	
<b>GENDER</b>								
Male	0.7451		0.2913		0.8961		0.7680	
Female	0.8996		0.0219		0.2417		0.6304	
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.6823		0.0892		0.5640		0.4676	
White	0.6786		0.1088		0.8157		0.4032	
Black or African American	0.9516		0.3926		0.4591		0.9351	
American Indian or Alaska Native	*		*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	*		*		*		*	
Two or More Races	*		*		*		*	
Hispanic or Latino	0.0897		0.0362		0.2353		0.2028	
<b>EDUCATION</b>								
< High School	0.9439		0.7061		0.8751		0.9141	
High School Graduate	0.7909		0.0573		0.7098		0.6821	
Some College	0.7751		0.0453		0.0850		0.4505	
College Graduate	0.9973		0.7924		0.9908		0.9370	
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.2414		0.2471		0.2304		0.8448	
Part-Time	0.4767		0.7082		0.6788		0.6800	
Unemployed	0.3383		0.2505		0.5648		0.2596	
Other <sup>3</sup>	0.1652		0.0251		0.6517		0.3004	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.26C Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic/Socioeconomic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	610	659	123	151	472	447	552	599
<b>GEOGRAPHIC REGION</b>								
Northeast	281	264	57	83	236	200	257	244
Midwest	263	255	42	46	164	191	242	237
South	335	412	82	93	239	261	310	381
West	334	361	60	72	288	234	290	313
<b>COUNTY TYPE<sup>3</sup></b>								
Large Metro	468	480	93	119	374	336	426	435
Small Metro	332	405	65	82	243	254	292	359
250K - 1 Mil. Pop.	274	321	58	70	191	216	233	287
< 250K Pop.	192	248	29	43	153	143	173	221
Nonmetro	231	247	46	51	162	166	214	230
Urbanized	171	184	23	46	118	113	156	176
Less Urbanized	154	181	40	19	108	120	144	166
Completely Rural	59	85	4	9	32	51	51	85
<b>FAMILY INCOME</b>								
Less Than \$20,000	304	301	87	113	218	209	289	272
\$20,000 - \$49,999	360	362	73	88	273	244	322	330
\$50,000 - \$74,999	269	263	35	48	186	185	250	246
\$75,000 or More	305	323	33	24	255	232	259	279
<b>GOVERNMENT ASSISTANCE<sup>4</sup></b>								
Yes	245	257	81	104	188	175	226	248
No	569	610	91	113	440	408	507	541

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

<sup>4</sup> One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.26D Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic/Socioeconomic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.27	0.26	0.06	0.07	0.21	0.19	0.25	0.25
<b>GEOGRAPHIC REGION</b>								
Northeast	0.61	0.62	0.14	0.20	0.51	0.47	0.57	0.60
Midwest	0.51	0.45	0.09	0.09	0.33	0.36	0.49	0.43
South	0.47	0.45	0.11	0.12	0.33	0.30	0.44	0.43
West	0.59	0.62	0.13	0.15	0.54	0.44	0.53	0.56
<b>COUNTY TYPE<sup>3</sup></b>								
Large Metro	0.38	0.37	0.08	0.10	0.31	0.27	0.36	0.35
Small Metro	0.47	0.51	0.10	0.13	0.36	0.34	0.42	0.46
250K - 1 Mil. Pop.	0.60	0.60	0.14	0.17	0.44	0.43	0.51	0.56
< 250K Pop.	0.78	0.93	0.14	0.19	0.66	0.56	0.72	0.83
Nonmetro	0.59	0.57	0.13	0.14	0.43	0.41	0.54	0.54
Urbanized	0.96	0.88	0.16	0.28	0.73	0.64	0.90	0.89
Less Urbanized	0.79	0.86	0.22	0.12	0.57	0.64	0.75	0.83
Completely Rural	1.41	2.02	0.09	0.26	0.81	1.28	1.19	2.03
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.68	0.60	0.21	0.27	0.51	0.45	0.65	0.57
\$20,000 - \$49,999	0.41	0.41	0.09	0.11	0.32	0.29	0.37	0.38
\$50,000 - \$74,999	0.65	0.61	0.09	0.12	0.47	0.45	0.61	0.57
\$75,000 or More	0.56	0.55	0.07	0.05	0.47	0.41	0.49	0.50
<b>GOVERNMENT ASSISTANCE<sup>4</sup></b>								
Yes	0.83	0.77	0.30	0.37	0.66	0.58	0.78	0.77
No	0.29	0.29	0.05	0.06	0.23	0.20	0.26	0.26

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

<sup>4</sup> One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.26P Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Geographic or Socioeconomic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic/Socioeconomic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.5909		0.1282		0.3304		0.1790	
<b>GEOGRAPHIC REGION</b>								
Northeast	0.8966		0.4574		0.6302		0.8115	
Midwest	0.0447		0.7370		0.1680		0.0064	
South	0.2679		0.4062		0.8644		0.1054	
West	0.0778		0.2908		0.0100		0.1457	
<b>COUNTY TYPE<sup>3</sup></b>								
Large Metro	0.9265		0.1147		0.2505		0.5137	
Small Metro	0.9847		0.5566		0.6137		0.6495	
250K - 1 Mil. Pop.	0.9007		0.6731		0.7676		0.7830	
< 250K Pop.	0.8455		0.6472		0.6558		0.6846	
Nonmetro	0.2236		0.9866		0.6659		0.0935	
Urbanized	0.6318		0.2039		0.1750		0.9750	
Less Urbanized	0.1916		0.0694		0.1950		0.1810	
Completely Rural	0.1285		0.1189		0.2911		0.0405	
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.9921		0.3467		0.7783		0.9975	
\$20,000 - \$49,999	0.9482		0.2819		0.3952		0.3219	
\$50,000 - \$74,999	0.6866		0.3130		0.9475		0.4391	
\$75,000 or More	0.4290		0.8370		0.5134		0.3821	
<b>GOVERNMENT ASSISTANCE<sup>4</sup></b>								
Yes	0.6707		0.4006		0.7324		0.3322	
No	0.7499		0.2295		0.3370		0.3335	

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

<sup>4</sup> One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.27C Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older with Past Year Serious Mental Illness, by Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic/Socioeconomic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	294	336	77	126	252	280	281	312
<b>GEOGRAPHIC REGION</b>								
Northeast	155	146	49	75	128	120	148	126
Midwest	119	135	21	27	97	112	107	132
South	185	200	50	77	161	161	179	201
West	119	182	23	61	109	160	117	153
<b>COUNTY TYPE<sup>3</sup></b>								
Large Metro	210	234	59	98	178	208	199	213
Small Metro	173	198	41	68	155	147	166	178
250K - 1 Mil. Pop.	126	154	39	58	110	110	119	146
< 250K Pop.	118	114	12	36	110	93	114	96
Nonmetro	119	140	26	45	95	114	115	141
Urbanized	92	114	14	40	78	85	87	111
Less Urbanized	71	91	22	17	47	69	69	85
Completely Rural	*	*	3	*	*	*	*	*
<b>FAMILY INCOME</b>								
Less Than \$20,000	167	168	57	94	147	137	157	156
\$20,000 - \$49,999	192	210	42	71	157	176	183	197
\$50,000 - \$74,999	118	123	27	46	96	102	112	121
\$75,000 or More	122	137	10	19	109	105	116	131
<b>GOVERNMENT ASSISTANCE<sup>4</sup></b>								
Yes	159	150	54	87	134	125	149	152
No	258	294	56	94	220	240	245	262

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

<sup>4</sup> One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 6.27D Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older with Past Year Serious Mental Illness, by Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic/Socioeconomic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	1.27	1.13	0.45	0.62	1.22	1.12	1.29	1.12
<b>GEOGRAPHIC REGION</b>								
Northeast	2.84	2.64	1.44	2.06	2.99	2.65	2.95	2.67
Midwest	2.39	1.92	0.54	0.63	2.13	1.99	2.34	1.96
South	2.17	1.88	0.81	1.02	2.14	1.78	2.20	2.00
West	2.85	2.71	0.64	1.27	2.67	2.74	2.87	2.33
<b>COUNTY TYPE<sup>3</sup></b>								
Large Metro	1.79	1.73	0.68	0.95	1.71	1.75	1.84	1.67
Small Metro	1.97	1.99	0.76	1.06	2.22	1.86	2.02	1.92
250K - 1 Mil. Pop.	2.44	2.20	1.11	1.35	2.66	2.09	2.32	2.26
< 250K Pop.	3.29	3.71	0.70	1.65	3.82	3.61	3.48	3.30
Nonmetro	3.00	2.43	0.81	1.19	2.58	2.43	2.91	2.61
Urbanized	4.14	3.90	1.09	2.29	4.44	3.71	4.21	4.07
Less Urbanized	4.25	3.74	1.30	1.02	3.08	3.36	4.09	3.65
Completely Rural	*	*	0.96	*	*	*	*	*
<b>FAMILY INCOME</b>								
Less Than \$20,000	2.38	2.07	1.04	1.50	2.28	1.97	2.33	2.15
\$20,000 - \$49,999	1.95	1.85	0.61	0.84	1.86	1.77	1.99	1.88
\$50,000 - \$74,999	2.95	3.06	1.04	1.56	2.87	3.00	2.99	3.13
\$75,000 or More	3.23	2.92	0.36	0.65	3.13	2.83	3.22	3.06
<b>GOVERNMENT ASSISTANCE<sup>4</sup></b>								
Yes	2.54	2.36	1.34	1.76	2.67	2.24	2.59	2.47
No	1.45	1.34	0.41	0.62	1.36	1.28	1.47	1.27

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

<sup>4</sup> One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.27P Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older with Past Year Serious Mental Illness, by Geographic or Socioeconomic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic/Socioeconomic Characteristic	Received Mental Health Treatment/Counseling <sup>1</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>					
			Inpatient		Outpatient		Prescription Medication	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.7214		0.0193		0.2959		0.8898	
<b>GEOGRAPHIC REGION</b>								
Northeast	0.6855		0.3791		0.7571		0.5440	
Midwest	0.1012		0.6674		0.1273		0.0096	
South	0.9689		0.1308		0.2261		0.8671	
West	0.1631		0.0495		0.2029		0.2579	
<b>COUNTY TYPE<sup>3</sup></b>								
Large Metro	0.5221		0.0545		0.4084		0.8786	
Small Metro	0.4893		0.3128		0.1025		0.6882	
250K - 1 Mil. Pop.	0.7162		0.3481		0.5003		0.3161	
< 250K Pop.	0.1225		0.7016		0.0710		0.0435	
Nonmetro	0.2917		0.3381		0.2415		0.2881	
Urbanized	0.6334		0.0865		0.5943		0.8627	
Less Urbanized	0.4276		0.3137		0.2238		0.4902	
Completely Rural	*		*		*		*	
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.8831		0.1215		0.5575		0.9452	
\$20,000 - \$49,999	0.1361		0.2660		0.4479		0.2488	
\$50,000 - \$74,999	0.8128		0.1946		0.8562		0.4331	
\$75,000 or More	0.2557		0.1801		0.7073		0.1512	
<b>GOVERNMENT ASSISTANCE<sup>4</sup></b>								
Yes	0.1018		0.2705		0.1776		0.3438	
No	0.7231		0.0555		0.6073		0.5786	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>2</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>3</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

<sup>4</sup> One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.