

Table 6.28C Perceived a Great Deal or a Lot of Help in the Past Year from Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Past Year Mental Health Treatment/Counseling, by Past Year Serious Mental Illness and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Demographic Characteristic	Total		SERIOUS MENTAL ILLNESS ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	480	494	218	251	423	410
AGE						
18-25	58	60	38	41	42	42
26-49	303	317	160	162	246	248
50 or Older	365	362	139	184	340	308
GENDER						
Male	250	238	126	130	217	201
Female	407	410	173	213	362	344
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	460	458	202	239	408	382
White	436	438	186	225	391	367
Black or African American	104	132	*	*	87	*
American Indian or Alaska Native	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	*	*	*	*	*	*
Two or More Races	*	*	*	*	*	*
Hispanic or Latino	119	135	*	*	93	121
EDUCATION						
< High School	170	147	76	94	153	114
High School Graduate	256	260	138	149	209	213
Some College	245	245	115	113	211	211
College Graduate	280	253	92	119	256	212
CURRENT EMPLOYMENT						
Full-Time	325	335	146	144	278	290
Part-Time	168	173	88	108	145	136
Unemployed	89	57	*	*	*	47
Other ²	305	282	132	153	262	232

*Low precision; no estimate reported.

NOTE: Respondents were asked how much the counseling or medicine they received in the past year improved their ability to manage daily activities. Respondents with missing information are excluded.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

¹ Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

² Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.28D Perceived a Great Deal or a Lot of Help in the Past Year from Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Past Year Mental Health Treatment/Counseling, by Past Year Serious Mental Illness and Demographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Demographic Characteristic	Total		SERIOUS MENTAL ILLNESS ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	1.08	1.03	1.85	1.71	1.29	1.31
AGE						
18-25	1.28	1.21	1.82	1.89	1.70	1.56
26-49	1.36	1.27	2.26	2.06	1.63	1.47
50 or Older	2.27	2.24	5.41	4.71	2.49	2.55
GENDER						
Male	1.98	1.87	3.88	3.57	2.26	2.26
Female	1.27	1.22	2.04	1.94	1.56	1.54
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1.10	1.06	1.90	1.76	1.32	1.35
White	1.18	1.10	1.90	1.93	1.42	1.34
Black or African American	4.41	4.72	*	*	4.65	*
American Indian or Alaska Native	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	*	*	*	*	*	*
Two or More Races	*	*	*	*	*	*
Hispanic or Latino	4.16	4.30	*	*	5.08	5.45
EDUCATION						
< High School	2.95	2.84	4.43	4.38	3.67	3.76
High School Graduate	1.81	1.82	3.14	3.08	2.35	2.39
Some College	1.98	2.00	3.27	2.88	2.41	2.48
College Graduate	1.94	1.93	3.46	3.74	2.28	2.27
CURRENT EMPLOYMENT						
Full-Time	1.50	1.39	2.67	2.49	1.70	1.64
Part-Time	2.52	2.45	4.27	4.54	3.06	3.17
Unemployed	4.90	4.17	*	*	*	5.74
Other ²	2.23	1.98	3.71	3.03	2.54	2.71

*Low precision; no estimate reported.

NOTE: Respondents were asked how much the counseling or medicine they received in the past year improved their ability to manage daily activities. Respondents with missing information are excluded.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

¹ Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

² Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.28P Perceived a Great Deal or a Lot of Help in the Past Year from Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Past Year Mental Health Treatment/Counseling, by Past Year Serious Mental Illness and Demographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Demographic Characteristic	Total		SERIOUS MENTAL ILLNESS ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	0.4249		0.8209		0.5308	
AGE						
18-25	0.5685		0.7513		0.6300	
26-49	0.5150		0.5950		0.6445	
50 or Older	0.1298		0.3582		0.3364	
GENDER						
Male	0.6931		0.9169		0.7947	
Female	0.4048		0.8473		0.4682	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4610		0.8155		0.4177	
White	0.3885		0.6534		0.6201	
Black or African American	0.8537		*		*	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	*		*		*	
Two or More Races	*		*		*	
Hispanic or Latino	0.7508		*		0.5741	
EDUCATION						
< High School	0.2387		0.9961		0.2440	
High School Graduate	0.9151		0.9569		0.9138	
Some College	0.2400		0.2142		0.5399	
College Graduate	0.8073		0.2694		0.8373	
CURRENT EMPLOYMENT						
Full-Time	0.7174		0.9936		0.6852	
Part-Time	0.5982		0.3255		0.1693	
Unemployed	0.0519		*		*	
Other ²	0.8346		0.7284		0.7696	

*Low precision; no estimate reported.

NOTE: Respondents were asked how much the counseling or medicine they received in the past year improved their ability to manage daily activities. Respondents with missing information are excluded.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

¹ Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

² Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.29C Perceived a Great Deal or a Lot of Help in the Past Year from Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Past Year Mental Health Treatment/Counseling, by Past Year Serious Mental Illness and Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic/Socioeconomic Characteristic	Total		SERIOUS MENTAL ILLNESS ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	480	494	218	251	423	410
GEOGRAPHIC REGION						
Northeast	226	183	113	107	221	142
Midwest	211	186	93	96	172	159
South	246	291	127	145	213	254
West	273	302	100	145	235	242
COUNTY TYPE²						
Large Metro	382	376	149	174	348	314
Small Metro	248	287	128	140	211	231
250K - 1 Mil. Pop.	209	222	103	111	176	184
< 250K Pop.	133	184	76	84	116	144
Nonmetro	161	182	94	109	133	157
Urbanized	117	133	*	87	92	100
Less Urbanized	109	142	59	74	94	118
Completely Rural	*	*	*	*	*	*
FAMILY INCOME						
Less Than \$20,000	219	200	110	124	184	157
\$20,000 - \$49,999	265	285	138	163	239	234
\$50,000 - \$74,999	226	217	93	93	203	188
\$75,000 or More	246	242	97	97	221	216
GOVERNMENT ASSISTANCE³						
Yes	190	189	119	116	150	153
No	445	471	184	224	403	389

*Low precision; no estimate reported.

NOTE: Respondents were asked how much the counseling or medicine they received in the past year improved their ability to manage daily activities. Respondents with missing information are excluded.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

¹ Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.29D Perceived a Great Deal or a Lot of Help in the Past Year from Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Past Year Mental Health Treatment/Counseling, by Past Year Serious Mental Illness and Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic/Socioeconomic Characteristic	Total		SERIOUS MENTAL ILLNESS ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	1.08	1.03	1.85	1.71	1.29	1.31
GEOGRAPHIC REGION						
Northeast	2.42	2.07	4.67	4.25	2.67	2.69
Midwest	1.89	1.80	3.05	2.82	2.36	2.26
South	1.92	1.77	3.14	2.71	2.28	2.36
West	2.39	2.63	4.11	4.40	2.84	3.06
COUNTY TYPE²						
Large Metro	1.64	1.58	2.72	2.69	1.95	1.86
Small Metro	1.76	1.75	3.31	2.80	2.08	2.30
250K - 1 Mil. Pop.	2.13	2.07	4.32	3.58	2.39	2.84
< 250K Pop.	3.14	3.09	5.18	4.72	4.01	3.69
Nonmetro	2.27	2.18	3.59	3.51	2.77	3.02
Urbanized	3.34	3.46	*	5.18	3.71	4.45
Less Urbanized	3.45	3.52	4.55	5.05	4.60	4.83
Completely Rural	*	*	*	*	*	*
FAMILY INCOME						
Less Than \$20,000	2.24	2.12	3.38	3.16	2.96	3.02
\$20,000 - \$49,999	1.70	1.79	2.96	2.79	2.17	2.19
\$50,000 - \$74,999	2.50	2.40	4.16	4.18	2.99	2.74
\$75,000 or More	2.29	2.16	4.66	4.37	2.50	2.47
GOVERNMENT ASSISTANCE³						
Yes	2.39	2.34	3.81	3.43	2.95	3.51
No	1.23	1.15	2.08	1.99	1.48	1.39

*Low precision; no estimate reported.

NOTE: Respondents were asked how much the counseling or medicine they received in the past year improved their ability to manage daily activities. Respondents with missing information are excluded.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

¹ Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.29P Perceived a Great Deal or a Lot of Help in the Past Year from Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Past Year Mental Health Treatment/Counseling, by Past Year Serious Mental Illness and Geographic or Socioeconomic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic/Socioeconomic Characteristic	Total		SERIOUS MENTAL ILLNESS ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	0.4249		0.8209		0.5308	
GEOGRAPHIC REGION						
Northeast	0.0005		0.4375		0.0003	
Midwest	0.1335		0.1362		0.4570	
South	0.0999		0.5909		0.0778	
West	0.5221		0.3620		0.6142	
COUNTY TYPE²						
Large Metro	0.3333		0.5321		0.5110	
Small Metro	0.2583		0.7982		0.2841	
250K - 1 Mil. Pop.	0.0994		0.2172		0.4002	
< 250K Pop.	0.7658		0.2009		0.5728	
Nonmetro	0.1459		0.3760		0.2448	
Urbanized	0.7540		*		0.3854	
Less Urbanized	0.1282		0.8430		0.0714	
Completely Rural	*		*		*	
FAMILY INCOME						
Less Than \$20,000	0.3636		0.9798		0.3145	
\$20,000 - \$49,999	0.4418		0.6555		0.4262	
\$50,000 - \$74,999	0.4605		0.1904		0.9456	
\$75,000 or More	0.2467		0.9799		0.2000	
GOVERNMENT ASSISTANCE³						
Yes	0.0370		0.5512		0.0297	
No	0.9667		0.9356		0.8632	

*Low precision; no estimate reported.

NOTE: Respondents were asked how much the counseling or medicine they received in the past year improved their ability to manage daily activities. Respondents with missing information are excluded.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

¹ Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.