

Table 6.46C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Demographic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
TOTAL	84	85	47	49	53	50	46	50
GENDER								
Male	57	56	36	35	32	32	30	32
Female	60	62	30	33	38	37	34	38
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	75	78	43	44	46	44	41	44
White	66	66	36	37	39	36	36	37
Black or African American	32	34	21	20	18	21	15	19
American Indian or Alaska Native	*	*	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	18	18	11	10	11	11	9	10
Two or More Races	9	10	6	7	5	5	5	*
Hispanic or Latino	37	36	20	21	24	22	19	23

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.46D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Demographic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
TOTAL	0.34	0.34	0.57	0.59	0.63	0.60	0.58	0.60
GENDER								
Male	0.45	0.44	0.82	0.81	0.76	0.77	0.74	0.76
Female	0.49	0.51	0.75	0.81	0.91	0.93	0.85	0.92
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.36	0.37	0.62	0.63	0.66	0.63	0.61	0.64
White	0.42	0.42	0.71	0.71	0.73	0.70	0.69	0.72
Black or African American	0.90	0.90	1.58	1.56	1.53	1.66	1.33	1.58
American Indian or Alaska Native	*	*	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.78	1.85	3.19	3.30	3.29	3.41	2.86	2.81
Two or More Races	2.41	2.56	3.47	4.15	4.50	3.64	4.95	*
Hispanic or Latino	0.95	0.90	1.50	1.49	1.75	1.71	1.58	1.74

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.46P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Demographic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
TOTAL	0.0052		0.3528		0.0459		0.0569	
GENDER								
Male	0.1063		0.6157		0.0486		0.1257	
Female	0.0156		0.0370		0.2878		0.2327	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.0147		0.4723		0.0463		0.1014	
White	0.1610		0.8547		0.2074		0.2728	
Black or African American	0.1610		0.8228		0.2012		0.1131	
American Indian or Alaska Native	*		*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.1400		0.2478		0.3311		0.5115	
Two or More Races	0.2264		0.1222		0.8380		*	
Hispanic or Latino	0.1222		0.4336		0.5992		0.2642	

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.47C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic/Socioeconomic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
TOTAL	101	107	57	58	58	57	50	58
GEOGRAPHIC REGION								
Northeast	40	45	23	19	28	25	21	24
Midwest	41	43	21	22	22	25	22	23
South	69	65	41	36	34	32	33	35
West	46	59	25	35	30	30	24	33
COUNTY TYPE¹								
Large Metro	79	87	46	45	45	44	40	48
Small Metro	56	57	29	29	33	32	27	26
250K - 1 Mil. Pop.	45	49	24	26	26	28	23	21
< 250K Pop.	33	30	16	14	19	16	15	15
Nonmetro	34	39	21	21	17	19	16	18
Urbanized	22	27	12	15	10	13	11	12
Less Urbanized	27	26	17	15	13	13	11	13
Completely Rural	13	18	7	6	7	7	5	8
FAMILY INCOME								
Less Than \$20,000	45	46	26	26	24	27	23	26
\$20,000 - \$49,999	59	61	34	34	36	34	29	35
\$50,000 - \$74,999	40	40	23	20	24	25	22	24
\$75,000 or More	56	56	30	29	29	28	28	31
GOVERNMENT ASSISTANCE²								
Yes	48	52	28	30	25	30	21	24
No	92	96	50	50	53	50	46	52

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

² One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.47D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic/Socioeconomic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
TOTAL	0.34	0.34	0.57	0.59	0.63	0.60	0.58	0.60
GEOGRAPHIC REGION								
Northeast	0.80	0.81	1.33	1.16	1.61	1.45	1.27	1.45
Midwest	0.64	0.67	0.97	1.00	1.07	1.22	1.04	1.06
South	0.60	0.55	1.02	1.02	1.07	0.92	1.08	0.97
West	0.73	0.75	1.27	1.43	1.38	1.39	1.18	1.46
COUNTY TYPE¹								
Large Metro	0.48	0.47	0.85	0.80	0.92	0.85	0.85	0.90
Small Metro	0.62	0.66	1.02	1.05	1.21	1.15	1.00	0.95
250K - 1 Mil. Pop.	0.69	0.81	1.25	1.31	1.42	1.50	1.27	1.12
< 250K Pop.	1.23	1.02	1.85	1.59	2.12	1.66	1.67	1.57
Nonmetro	0.72	0.72	1.34	1.26	1.07	1.16	1.13	1.13
Urbanized	1.09	1.24	1.81	1.99	1.70	1.84	1.84	1.84
Less Urbanized	1.09	0.98	2.20	2.04	1.62	1.74	1.60	1.62
Completely Rural	1.97	2.25	3.33	3.15	2.96	2.97	2.62	3.28
FAMILY INCOME								
Less Than \$20,000	0.83	0.89	1.42	1.44	1.40	1.57	1.45	1.62
\$20,000 - \$49,999	0.57	0.59	0.99	1.01	1.03	1.05	0.96	1.03
\$50,000 - \$74,999	0.73	0.76	1.24	1.25	1.31	1.33	1.24	1.35
\$75,000 or More	0.64	0.64	1.10	1.04	1.15	1.10	1.08	1.16
GOVERNMENT ASSISTANCE²								
Yes	0.94	0.91	1.56	1.50	1.60	1.74	1.61	1.57
No	0.36	0.37	0.61	0.64	0.67	0.64	0.63	0.64

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

² One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.47P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic/Socioeconomic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
TOTAL	0.0052		0.3528		0.0459		0.0569	
GEOGRAPHIC REGION								
Northeast	0.5861		0.0498		0.2882		0.1449	
Midwest	0.1172		0.4950		0.0231		0.2214	
South	0.7325		0.7114		0.6308		0.4496	
West	0.0011		0.0038		0.1007		0.4835	
COUNTY TYPE¹								
Large Metro	0.0473		0.8760		0.1148		0.0749	
Small Metro	0.1577		0.0864		0.6671		0.6100	
250K - 1 Mil. Pop.	0.0975		0.0323		0.3668		0.9600	
< 250K Pop.	0.8340		0.9744		0.6475		0.3139	
Nonmetro	0.1298		0.6212		0.0954		0.3888	
Urbanized	0.4863		0.7074		0.2958		0.9648	
Less Urbanized	0.2945		0.6076		0.2226		0.6813	
Completely Rural	0.3219		0.7781		0.7144		0.0301	
FAMILY INCOME								
Less Than \$20,000	0.0156		0.4390		0.0330		0.1431	
\$20,000 - \$49,999	0.0598		0.5151		0.1366		0.2001	
\$50,000 - \$74,999	0.2155		0.4437		0.4968		0.4917	
\$75,000 or More	0.8719		0.7368		0.8269		0.7977	
GOVERNMENT ASSISTANCE²								
Yes	0.0156		0.8320		0.0335		0.0363	
No	0.1031		0.4314		0.3239		0.2884	

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

² One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.48C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Any Illicit Drug Use and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Demographic Characteristic	Total		ANY ILLICIT DRUG USE ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	84	85	45	42	69	71
AGE						
12-13	47	49	16	16	46	45
14-15	53	50	26	26	43	42
16-17	46	50	32	32	32	37
GENDER						
Male	57	56	27	28	50	50
Female	60	62	32	32	49	52
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	75	78	39	39	62	65
White	66	66	36	34	53	55
Black or African American	32	34	14	15	27	30
American Indian or Alaska Native	*	*	*	3	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	18	18	*	*	17	16
Two or More Races	9	10	5	*	8	9
Hispanic or Latino	37	36	20	20	32	30

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.48D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Any Illicit Drug Use and Demographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Demographic Characteristic	Total		ANY ILLICIT DRUG USE ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	0.34	0.34	0.82	0.78	0.36	0.36
AGE						
12-13	0.57	0.59	2.18	2.16	0.61	0.59
14-15	0.63	0.60	1.36	1.42	0.66	0.66
16-17	0.58	0.60	1.13	1.12	0.62	0.68
GENDER						
Male	0.45	0.44	0.96	1.02	0.51	0.50
Female	0.49	0.51	1.19	1.19	0.52	0.54
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.36	0.37	0.85	0.85	0.39	0.40
White	0.42	0.42	0.98	0.96	0.45	0.46
Black or African American	0.90	0.90	2.14	2.15	0.93	0.99
American Indian or Alaska Native	*	*	*	4.98	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.78	1.85	*	*	1.93	1.93
Two or More Races	2.41	2.56	5.04	*	2.73	2.85
Hispanic or Latino	0.95	0.90	2.44	2.35	1.03	0.96

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.48P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Any Illicit Drug Use and Demographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Demographic Characteristic	Total		ANY ILLICIT DRUG USE ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	0.0052		0.2251		0.0072	
AGE						
12-13	0.3528		0.9844		0.3111	
14-15	0.0459		0.0849		0.1522	
16-17	0.0569		0.7780		0.0149	
GENDER						
Male	0.1063		0.3102		0.1745	
Female	0.0156		0.4842		0.0150	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0147		0.4292		0.0088	
White	0.1610		0.6932		0.0848	
Black or African American	0.1610		0.3961		0.2534	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1400		*		0.2320	
Two or More Races	0.2264		*		0.3080	
Hispanic or Latino	0.1222		0.2276		0.4236	

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.49C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Any Illicit Drug Use and Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic/Socioeconomic Characteristic	Total		ANY ILLICIT DRUG USE ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	84	85	45	42	69	71
GEOGRAPHIC REGION						
Northeast	36	36	16	17	29	32
Midwest	36	38	19	19	30	34
South	52	49	28	24	43	40
West	42	44	24	24	34	36
COUNTY TYPE²						
Large Metro	63	62	34	33	54	53
Small Metro	45	49	23	22	36	39
250K - 1 Mil. Pop.	34	40	18	18	28	33
< 250K Pop.	29	25	15	13	21	19
Nonmetro	31	31	15	17	25	25
Urbanized	19	23	9	13	15	17
Less Urbanized	23	19	11	10	20	17
Completely Rural	9	10	5	*	6	7
FAMILY INCOME						
Less Than \$20,000	36	40	19	19	31	35
\$20,000 - \$49,999	51	52	26	26	42	42
\$50,000 - \$74,999	35	36	19	17	29	31
\$75,000 or More	42	44	24	24	37	36
GOVERNMENT ASSISTANCE³						
Yes	38	40	19	20	32	33
No	75	76	39	38	62	65

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.49D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Any Illicit Drug Use and Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic/Socioeconomic Characteristic	Total		ANY ILLICIT DRUG USE ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	0.34	0.34	0.82	0.78	0.36	0.36
GEOGRAPHIC REGION						
Northeast	0.80	0.81	1.59	1.85	0.85	0.92
Midwest	0.64	0.67	1.50	1.55	0.68	0.75
South	0.60	0.55	1.47	1.27	0.63	0.57
West	0.73	0.75	1.89	1.74	0.76	0.80
COUNTY TYPE²						
Large Metro	0.48	0.47	1.15	1.20	0.52	0.51
Small Metro	0.62	0.66	1.44	1.26	0.64	0.68
250K - 1 Mil. Pop.	0.69	0.81	1.62	1.63	0.75	0.85
< 250K Pop.	1.23	1.02	2.96	2.13	1.14	1.03
Nonmetro	0.72	0.72	1.60	1.77	0.74	0.76
Urbanized	1.09	1.24	2.27	2.85	1.12	1.24
Less Urbanized	1.09	0.98	2.30	2.32	1.20	1.08
Completely Rural	1.97	2.25	4.36	*	1.76	2.04
FAMILY INCOME						
Less Than \$20,000	0.83	0.89	1.99	1.92	0.91	0.99
\$20,000 - \$49,999	0.57	0.59	1.31	1.31	0.61	0.61
\$50,000 - \$74,999	0.73	0.76	1.82	1.67	0.76	0.83
\$75,000 or More	0.64	0.64	1.60	1.68	0.72	0.66
GOVERNMENT ASSISTANCE³						
Yes	0.94	0.91	1.94	1.94	1.05	0.98
No	0.36	0.37	0.88	0.87	0.38	0.40

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.49P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Any Illicit Drug Use and Geographic or Socioeconomic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic/Socioeconomic Characteristic	Total		ANY ILLICIT DRUG USE ¹			
			Yes		No	
	2002	2003	2002	2003	2002	2003
TOTAL	0.0052		0.2251		0.0072	
GEOGRAPHIC REGION						
Northeast	0.5861		0.3726		0.7948	
Midwest	0.1172		0.9535		0.0538	
South	0.7325		0.5524		0.9007	
West	0.0011		0.3517		0.0011	
COUNTY TYPE²						
Large Metro	0.0473		0.0682		0.1838	
Small Metro	0.1577		0.2443		0.0145	
250K - 1 Mil. Pop.	0.0975		0.4659		0.0111	
< 250K Pop.	0.8340		0.3551		0.5101	
Nonmetro	0.1298		0.1908		0.2697	
Urbanized	0.4863		0.5867		0.6874	
Less Urbanized	0.2945		0.3351		0.4443	
Completely Rural	0.3219		*		0.2454	
FAMILY INCOME						
Less Than \$20,000	0.0156		0.8758		0.0064	
\$20,000 - \$49,999	0.0598		0.5479		0.0506	
\$50,000 - \$74,999	0.2155		0.9323		0.1346	
\$75,000 or More	0.8719		0.1428		0.3864	
GOVERNMENT ASSISTANCE³						
Yes	0.0156		0.5998		0.0080	
No	0.1031		0.2828		0.1854	

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ One or more household family members received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.50C Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

Source of Treatment/Counseling	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES¹	101	107	57	58	58	57	50	58
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	67	70	34	36	39	39	36	40
School Counselors, School Psychologists, or Having Regular Meetings with Teachers ²	61	70	37	44	38	38	30	36
Mental Health Clinic or Center	35	31	15	19	20	17	22	19
In-home Therapist, Counselor, or Family Preservation Worker	33	33	20	19	22	18	17	21
Pediatrician or Other Family Doctor	34	38	20	20	21	21	20	21
Overnight or Longer Stay in Any Type of Hospital	29	31	15	15	16	19	18	19
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School ²	40	42	27	22	21	23	19	26
Partial Day Hospital or Day Treatment Program	28	26	15	16	17	14	16	15
Overnight or Longer Stay in a Residential Treatment Center	20	23	7	10	10	11	16	15
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	16	19	9	11	8	11	9	11

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

¹ Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.50D Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2002 and 2003

Source of Treatment/Counseling	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES¹	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	1.02	0.94	1.64	1.64	1.62	1.56	1.77	1.64
School Counselors, School Psychologists, or Having Regular Meetings with Teachers ²	0.96	0.92	1.69	1.75	1.59	1.51	1.69	1.58
Mental Health Clinic or Center	0.69	0.60	0.86	1.04	1.10	0.94	1.39	1.13
In-home Therapist, Counselor, or Family Preservation Worker	0.65	0.61	1.10	1.07	1.21	0.98	1.15	1.19
Pediatrician or Other Family Doctor	0.64	0.68	1.10	1.11	1.10	1.11	1.25	1.17
Overnight or Longer Stay in Any Type of Hospital	0.56	0.58	0.86	0.82	0.89	1.01	1.14	1.12
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School ²	0.74	0.75	1.43	1.20	1.13	1.19	1.22	1.48
Partial Day Hospital or Day Treatment Program	0.55	0.48	0.85	0.87	0.95	0.74	1.05	0.88
Overnight or Longer Stay in a Residential Treatment Center	0.41	0.43	0.45	0.56	0.58	0.59	1.08	0.91
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	0.32	0.37	0.53	0.63	0.48	0.59	0.65	0.66

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

¹ Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.50P Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Source of Treatment/Counseling	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES¹	N/A		N/A		N/A		N/A	
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	0.2889		0.4278		0.0148		0.7372	
School Counselors, School Psychologists, or Having Regular Meetings with Teachers ²	0.0140		0.4506		0.3374		0.0048	
Mental Health Clinic or Center	0.6480		0.0237		0.4302		0.0561	
In-home Therapist, Counselor, or Family Preservation Worker	0.0588		0.1920		0.0163		0.6482	
Pediatrician or Other Family Doctor	0.5864		0.3121		0.4032		0.3288	
Overnight or Longer Stay in Any Type of Hospital	0.7872		0.7010		0.5225		0.5873	
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School ²	0.2943		0.1778		0.3166		0.0218	
Partial Day Hospital or Day Treatment Program	0.1681		0.6890		0.0364		0.5147	
Overnight or Longer Stay in a Residential Treatment Center	0.7489		0.4419		0.8886		0.3073	
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	0.6919		0.5145		0.6938		0.7245	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

¹ Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.51C Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

Reason for Treatment/Counseling ¹	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES	95	103	53	56	56	55	49	56
Thought About Killing Self or Tried to Kill Self	41	38	17	18	24	22	25	24
Felt Depressed ²	63	67	31	33	36	34	36	39
Felt Very Afraid or Tense	35	42	20	22	24	23	20	26
Breaking Rules or "Acting Out" ³	45	44	26	24	26	28	23	23
Eating Problems	25	27	11	11	17	18	15	18
Family/Home Problems ⁴	31	35	16	20	19	19	18	20
Social/Friend Problems ⁴	23	28	11	15	13	17	13	14
School-related Issues ⁴	25	28	14	15	17	17	13	16
Other Diagnosed Mental/Neurological Disorder ⁴	13	14	8	7	7	8	7	8
Other Reason/Reasons	40	47	22	27	22	23	24	27

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they received treatment/counseling from one specific source (i.e., "special education services while in a regular classroom or in a special classroom, a special program, or in a special school") were not asked for the reasons they received these services and were therefore excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive.

² Includes other specify responses of "feeling inadequate, hopeless, lonely, or having low self-esteem."

³ Includes other specify responses of "difficulty controlling anger or fighting."

⁴ Respondents were permitted to specify other reasons for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.51D Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2002 and 2003

Reason for Treatment/Counseling ¹	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Thought About Killing Self or Tried to Kill Self	0.91	0.80	1.19	1.22	1.45	1.34	1.75	1.52
Felt Depressed ²	1.05	1.08	1.71	1.90	1.75	1.70	1.89	1.82
Felt Very Afraid or Tense	0.80	0.88	1.32	1.45	1.46	1.40	1.43	1.57
Breaking Rules or "Acting Out" ³	0.94	0.89	1.66	1.51	1.58	1.62	1.63	1.47
Eating Problems	0.60	0.62	0.84	0.80	1.06	1.12	1.18	1.23
Family/Home Problems ⁴	0.73	0.73	1.20	1.33	1.23	1.15	1.31	1.29
Social/Friend Problems ⁴	0.57	0.61	0.87	1.10	0.92	1.05	1.01	0.92
School-related Issues ⁴	0.61	0.62	1.00	1.09	1.08	1.13	1.00	1.09
Other Diagnosed Mental/Neurological Disorder ⁴	0.31	0.31	0.63	0.54	0.49	0.53	0.53	0.54
Other Reason/Reasons	0.87	0.94	1.45	1.72	1.35	1.40	1.66	1.59

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they received treatment/counseling from one specific source (i.e., "special education services while in a regular classroom or in a special classroom, a special program, or in a special school") were not asked for the reasons they received these services and were therefore excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive.

² Includes other specify responses of "feeling inadequate, hopeless, lonely, or having low self-esteem."

³ Includes other specify responses of "difficulty controlling anger or fighting."

⁴ Respondents were permitted to specify other reasons for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.51P Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Reason for Treatment/Counseling ¹	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2002	2003	2002	2003	2002	2003	2002	2003
RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES	N/A		N/A		N/A		N/A	
Thought About Killing Self or Tried to Kill Self	0.5848		0.2715		0.4333		0.3037	
Felt Depressed ²	0.6481		0.5059		0.5736		0.5209	
Felt Very Afraid or Tense	0.1139		0.2275		0.9801		0.0968	
Breaking Rules or "Acting Out" ³	0.4402		0.7515		0.8644		0.4161	
Eating Problems	0.7289		0.6226		0.9621		0.9170	
Family/Home Problems ⁴	0.5984		0.1570		0.3647		0.5938	
Social/Friend Problems ⁴	0.3025		0.1900		0.4601		0.8969	
School-related Issues ⁴	0.4234		0.9184		0.5764		0.3661	
Other Diagnosed Mental/Neurological Disorder ⁴	0.6446		0.5897		0.5456		0.4567	
Other Reason/Reasons	0.7597		0.3965		0.3109		0.6777	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they received treatment/counseling from one specific source (i.e., "special education services while in a regular classroom or in a special classroom, a special program, or in a special school") were not asked for the reasons they received these services and were therefore excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive.

² Includes other specify responses of "feeling inadequate, hopeless, lonely, or having low self-esteem."

³ Includes other specify responses of "difficulty controlling anger or fighting."

⁴ Respondents were permitted to specify other reasons for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.52C Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG³	111	110	50	51	99	92
Marijuana and Hashish	90	88	41	42	80	73
Cocaine	32	28	21	16	26	22
Crack	14	13	8	8	11	10
Heroin	11	8	4	6	10	5
Hallucinogens	41	38	25	20	34	31
LSD	25	15	16	11	21	12
PCP	13	12	9	7	9	9
Ecstasy	33	23	19	12	27	19
Inhalants	45	50	23	27	39	42
Nonmedical Use of Any Psychotherapeutic ⁴	68	70	35	36	56	58
Pain Relievers	62	64	32	33	51	53
Tranquilizers	35	34	20	20	27	28
Stimulants	36	31	22	18	26	25
Methamphetamine	21	17	15	11	15	13
Sedatives	17	16	10	9	13	13
ANY ILLICIT DRUG OTHER THAN MARIJUANA³	82	89	40	43	71	72

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.52D Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2002 and 2003

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG³	0.38	0.36	0.86	0.82	0.43	0.39
Marijuana and Hashish	0.32	0.31	0.76	0.71	0.35	0.33
Cocaine	0.13	0.11	0.43	0.30	0.13	0.11
Crack	0.06	0.05	0.17	0.15	0.06	0.05
Heroin	0.04	0.03	0.09	0.11	0.05	0.03
Hallucinogens	0.16	0.15	0.50	0.37	0.17	0.15
LSD	0.10	0.06	0.32	0.21	0.10	0.06
PCP	0.05	0.05	0.18	0.13	0.05	0.05
Ecstasy	0.13	0.09	0.40	0.24	0.13	0.10
Inhalants	0.18	0.19	0.47	0.51	0.19	0.20
Nonmedical Use of Any Psychotherapeutic ⁴	0.26	0.26	0.66	0.65	0.27	0.27
Pain Relievers	0.24	0.24	0.63	0.61	0.24	0.25
Tranquilizers	0.14	0.13	0.41	0.38	0.13	0.14
Stimulants	0.14	0.12	0.45	0.34	0.13	0.12
Methamphetamine	0.09	0.07	0.31	0.21	0.08	0.07
Sedatives	0.07	0.06	0.20	0.18	0.06	0.06
ANY ILLICIT DRUG OTHER THAN MARIJUANA³	0.31	0.32	0.75	0.75	0.33	0.33

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.52P Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG³	0.4872		0.3607		0.4668	
Marijuana and Hashish	0.0780		0.1188		0.1471	
Cocaine	0.0730		0.0815		0.2853	
Crack	0.7667		0.5186		0.8896	
Heroin	0.2709		0.7065		0.1132	
Hallucinogens	0.0011		0.0301		0.0065	
LSD	0.0000		0.0102		0.0000	
PCP	0.2074		0.0541		0.8470	
Ecstasy	0.0000		0.0010		0.0000	
Inhalants	0.6422		0.5899		0.9966	
Nonmedical Use of Any Psychotherapeutic ⁴	0.8584		0.5382		0.8601	
Pain Relievers	0.7254		0.8469		0.8041	
Tranquilizers	0.7938		0.4471		0.6514	
Stimulants	0.0562		0.0295		0.3211	
Methamphetamine	0.0490		0.0232		0.4030	
Sedatives	0.7149		0.7879		0.4221	
ANY ILLICIT DRUG OTHER THAN MARIJUANA³	0.8608		0.9179		0.6025	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.53C Tobacco and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
PAST YEAR						
Any Tobacco ³	113	111	50	50	100	94
Cigarettes	105	101	48	46	94	84
Smokeless Tobacco	42	42	19	18	37	36
Cigars	69	71	34	32	59	61
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	141	137	58	61	122	116
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Any Tobacco ³	87	87	43	40	76	71
Cigarettes	80	78	40	38	71	64
Smokeless Tobacco	27	30	13	12	24	26
Cigars	47	44	28	20	38	39
Pipe Tobacco	14	18	9	9	11	15
Alcohol	90	95	45	44	79	79
Binge Alcohol Use ⁵	72	73	35	37	60	59
Heavy Alcohol Use ⁵	31	34	15	18	28	29

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

³ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.53D Tobacco and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2002 and 2003

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
PAST YEAR						
Any Tobacco ³	0.37	0.37	0.86	0.82	0.42	0.40
Cigarettes	0.35	0.36	0.84	0.79	0.40	0.38
Smokeless Tobacco	0.17	0.16	0.40	0.35	0.18	0.18
Cigars	0.26	0.26	0.66	0.59	0.28	0.28
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	0.42	0.42	0.92	0.92	0.47	0.46
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Any Tobacco ³	0.33	0.32	0.80	0.73	0.35	0.33
Cigarettes	0.30	0.29	0.75	0.71	0.33	0.30
Smokeless Tobacco	0.11	0.12	0.28	0.23	0.12	0.13
Cigars	0.19	0.17	0.57	0.40	0.19	0.19
Pipe Tobacco	0.06	0.07	0.20	0.17	0.05	0.08
Alcohol	0.32	0.33	0.82	0.73	0.35	0.35
Binge Alcohol Use ⁵	0.27	0.27	0.67	0.66	0.28	0.28
Heavy Alcohol Use ⁵	0.12	0.13	0.31	0.34	0.14	0.15

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

³ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.53P Tobacco and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
PAST YEAR						
Any Tobacco ³	0.0438		0.0281		0.1900	
Cigarettes	0.0051		0.0263		0.0300	
Smokeless Tobacco	0.5002		0.3652		0.7417	
Cigars	0.6166		0.3418		0.8247	
Pipe Tobacco ⁴	--		--		--	
Alcohol	0.5577		0.5626		0.5934	
Binge Alcohol Use ⁵	--		--		--	
Heavy Alcohol Use ⁵	--		--		--	
PAST MONTH						
Any Tobacco ³	0.0797		0.0127		0.4491	
Cigarettes	0.0511		0.1019		0.1576	
Smokeless Tobacco	0.6430		0.6385		0.4654	
Cigars	0.8391		0.0079		0.1924	
Pipe Tobacco	0.3391		0.7076		0.2005	
Alcohol	0.8725		0.6223		0.8061	
Binge Alcohol Use ⁵	0.9274		0.3159		0.3890	
Heavy Alcohol Use ⁵	0.7209		0.6566		0.5697	

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

³ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.54C Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2002 and 2003

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
DEPENDENCE OR ABUSE³						
Any Illicit Drug or Alcohol ⁴	65	65	38	37	53	52
Any Illicit Drug ⁴	53	47	31	27	41	38
Alcohol	54	52	32	29	44	42
DEPENDENCE³						
Any Illicit Drug or Alcohol ⁴	46	45	30	26	35	34
Any Illicit Drug ⁴	39	35	24	21	30	28
Alcohol	31	32	23	20	23	23

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.54D Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2002 and 2003

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
DEPENDENCE OR ABUSE³						
Any Illicit Drug or Alcohol ⁴	0.24	0.25	0.73	0.66	0.25	0.25
Any Illicit Drug ⁴	0.20	0.18	0.61	0.50	0.20	0.19
Alcohol	0.20	0.20	0.62	0.54	0.21	0.20
DEPENDENCE³						
Any Illicit Drug or Alcohol ⁴	0.18	0.17	0.59	0.50	0.17	0.17
Any Illicit Drug ⁴	0.16	0.14	0.49	0.41	0.15	0.14
Alcohol	0.12	0.12	0.46	0.39	0.11	0.11

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.54P Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
DEPENDENCE OR ABUSE³						
Any Illicit Drug or Alcohol ⁴	0.8360		0.2334		0.8248	
Any Illicit Drug ⁴	0.1231		0.0290		0.4812	
Alcohol	0.9592		0.2687		0.5474	
DEPENDENCE³						
Any Illicit Drug or Alcohol ⁴	0.2386		0.0452		0.8111	
Any Illicit Drug ⁴	0.0624		0.0801		0.2290	
Alcohol	0.6372		0.1144		0.6388	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.50 for a list of the 10 specific sources of treatment.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.