

Table 7.113A Illicit Drug Use in the Past Month, by Levels of Past Month Alcohol Use and Past Month Cigarette Use and Age Group: Numbers in Thousands, 2002 and 2003

| Level of Alcohol Use/ Cigarette Use | Total | | AGE GROUP (Years) | | | | | |
|--|--------|--------|-------------------|-------|-------|-------|-------------|--------|
| | | | 12-17 | | 18-25 | | 26 or Older | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 19,522 | 19,470 | 2,878 | 2,811 | 6,264 | 6,435 | 10,380 | 10,224 |
| Heavy Alcohol Use ¹ | 5,178 | 5,247 | 422 | 421 | 2,277 | 2,336 | 2,479 | 2,490 |
| Binge Alcohol Use But Not Heavy Use ¹ | 6,307 | 6,216 | 887 | 913 | 2,255 | 2,307 | 3,165 | 2,997 |
| Alcohol Use But Not Binge Use ¹ | 3,853 | 4,136 | 435 | 428 | 906 | 1,013 | 2,512 | 2,694 |
| No Alcohol Use | 4,185 | 3,871 | 1,134 | 1,049 | 827 | 778 | 2,224 | 2,043 |
| PAST MONTH CIGARETTE USE | 11,916 | 11,974 | 1,543 | 1,473 | 4,500 | 4,490 | 5,873 | 6,011 |
| Heavy Alcohol Use ¹ | 3,911 | 3,951 | 334 | 342 | 1,820 | 1,801 | 1,757 | 1,808 |
| Binge Alcohol Use But Not Heavy Use ¹ | 4,150 | 4,102 | 583 | 579 | 1,628 | 1,633 | 1,939 | 1,890 |
| Alcohol Use But Not Binge Use ¹ | 1,996 | 2,086 | 236 | 218 | 593 | 630 | 1,167 | 1,237 |
| No Alcohol Use | 1,858 | 1,836 | 390 | 335 | 459 | 426 | 1,009 | 1,075 |
| NO PAST MONTH CIGARETTE USE | 7,606 | 7,496 | 1,335 | 1,338 | 1,764 | 1,945 | 4,507 | 4,213 |
| Heavy Alcohol Use ¹ | 1,267 | 1,296 | 88 | 79 | 457 | 535 | 722 | 682 |
| Binge Alcohol Use But Not Heavy Use ¹ | 2,157 | 2,114 | 304 | 334 | 627 | 674 | 1,226 | 1,106 |
| Alcohol Use But Not Binge Use ¹ | 1,856 | 2,050 | 199 | 210 | 313 | 383 | 1,345 | 1,457 |
| No Alcohol Use | 2,326 | 2,035 | 744 | 715 | 368 | 353 | 1,214 | 968 |

*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.113B Illicit Drug Use in the Past Month, by Levels of Past Month Alcohol Use and Past Month Cigarette Use and Age Group: Percentages, 2002 and 2003

| Level of Alcohol Use/ Cigarette Use | Total | | AGE GROUP (Years) | | | | | |
|--|-------|------|-------------------|------|-------|------|-------------|------|
| | | | 12-17 | | 18-25 | | 26 or Older | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 8.3 | 8.2 | 11.6 | 11.2 | 20.2 | 20.3 | 5.8 | 5.6 |
| Heavy Alcohol Use ¹ | 32.6 | 32.5 | 67.0 | 64.5 | 49.2 | 48.8 | 23.4 | 23.3 |
| Binge Alcohol Use But Not Heavy Use ¹ | 16.6 | 16.5 | 44.1 | 45.5 | 27.9 | 27.5 | 11.4 | 11.0 |
| Alcohol Use But Not Binge Use ¹ | 5.8 | 6.3 | 25.2 | 24.2 | 14.9 | 16.1 | 4.3 | 4.7 |
| No Alcohol Use | 3.6 | 3.3 | 5.6 | 5.1 | 6.7 | 6.3 | 2.7 | 2.4 |
| PAST MONTH CIGARETTE USE | 19.5 | 19.8 | 48.1 | 48.4 | 35.6 | 35.2 | 13.0 | 13.5 |
| Heavy Alcohol Use ¹ | 40.2 | 39.7 | 74.5 | 72.4 | 55.3 | 53.2 | 29.4 | 29.6 |
| Binge Alcohol Use But Not Heavy Use ¹ | 25.0 | 25.2 | 59.1 | 59.9 | 36.3 | 35.7 | 17.4 | 17.6 |
| Alcohol Use But Not Binge Use ¹ | 13.9 | 15.5 | 51.5 | 50.8 | 28.0 | 30.4 | 9.9 | 11.3 |
| No Alcohol Use | 9.1 | 8.9 | 29.6 | 28.5 | 16.7 | 15.6 | 6.2 | 6.4 |
| NO PAST MONTH CIGARETTE USE | 4.4 | 4.2 | 6.2 | 6.1 | 9.6 | 10.3 | 3.4 | 3.1 |
| Heavy Alcohol Use ¹ | 20.6 | 20.9 | 48.4 | 43.8 | 34.3 | 38.1 | 15.6 | 14.8 |
| Binge Alcohol Use But Not Heavy Use ¹ | 10.1 | 9.9 | 29.7 | 32.1 | 17.4 | 17.6 | 7.3 | 6.7 |
| Alcohol Use But Not Binge Use ¹ | 3.6 | 4.0 | 15.7 | 15.7 | 7.9 | 9.1 | 2.9 | 3.2 |
| No Alcohol Use | 2.5 | 2.1 | 3.9 | 3.7 | 3.9 | 3.7 | 1.8 | 1.4 |

*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.