

Table 7.25A Tobacco Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY TOBACCO²	11,259	11,444	19,345	19,298	17,309	16,245	23,585	23,771
Cigarettes	9,722	9,953	16,601	16,305	14,365	13,489	20,449	20,687
Smokeless Tobacco	1,667	1,582	2,054	2,115	1,314	1,508	2,751	2,520
Cigars	3,039	2,975	4,319	4,436	2,937	2,823	2,456	2,603
Pipe Tobacco	277	325	453	358	520	335	566	601

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.25B Tobacco Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY TOBACCO²	71.0	70.9	51.0	51.3	26.2	24.9	20.5	20.0
Cigarettes	61.3	61.7	43.8	43.3	21.8	20.7	17.7	17.4
Smokeless Tobacco	10.5	9.8	5.4	5.6	2.0	2.3	2.4	2.1
Cigars	19.2	18.4	11.4	11.8	4.4	4.3	2.1	2.2
Pipe Tobacco	1.7	2.0	1.2	1.0	0.8	0.5	0.5	0.5

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.26A Tobacco Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY TOBACCO²	511	525	1,124	1,113	547	495	1,585	1,477
Cigarettes	449	472	987	966	458	430	1,316 ^a	1,176
Smokeless Tobacco	101	114	155	153	44	46	188	197
Cigars	214	213	407	384	143	147	354	371
Pipe Tobacco	26	30	38	61	16	9	58	60

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.26B Tobacco Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY TOBACCO²	81.1	80.4	55.9	55.5	31.7	28.0	7.8	7.2
Cigarettes	71.2	72.4	49.1	48.1	26.6	24.3	6.5 ^a	5.7
Smokeless Tobacco	16.1	17.5	7.7	7.6	2.5	2.6	0.9	1.0
Cigars	33.9	32.7	20.2	19.1	8.3	8.3	1.7	1.8
Pipe Tobacco	4.1	4.6	1.9	3.0	0.9	0.5	0.3	0.3

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.27A Tobacco Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY TOBACCO²	3,654	3,758	4,979	5,141	2,375	2,324	3,042	2,978
Cigarettes	3,293	3,383	4,483	4,577	2,120	2,075	2,754	2,728
Smokeless Tobacco	659	638	475	505	132	159	237	184
Cigars	1,133	1,140	1,258 ^a	1,423	473	502	554	557
Pipe Tobacco	88	96	142 ^a	99	45	40	58	58

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.27B Tobacco Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY TOBACCO²	79.0	78.5	61.7	61.2	39.2	37.0	24.8	24.3
Cigarettes	71.2	70.7	55.5	54.5	35.0	33.1	22.5	22.3
Smokeless Tobacco	14.2	13.3	5.9	6.0	2.2	2.5	1.9	1.5
Cigars	24.5	23.8	15.6	16.9	7.8	8.0	4.5	4.5
Pipe Tobacco	1.9	2.0	1.8 ^a	1.2	0.7	0.6	0.5	0.5

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.28A Tobacco Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY TOBACCO²	7,095	7,161	13,242	13,044	14,388	13,426	18,959	19,315
Cigarettes	5,981	6,098	11,131	10,762	11,786	10,984	16,379	16,782
Smokeless Tobacco	907	830	1,425	1,457	1,138	1,302	2,327	2,139
Cigars	1,693	1,622	2,654	2,628	2,322	2,174	1,547	1,675
Pipe Tobacco	163	199	274	198	460	286	451	483

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.28B Tobacco Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY TOBACCO²	66.9	66.9	47.6	47.9	24.7	23.5	22.9	22.5
Cigarettes	56.4	57.0	40.0	39.5	20.2	19.2	19.8	19.5
Smokeless Tobacco	8.6	7.8	5.1	5.4	2.0	2.3	2.8	2.5
Cigars	16.0	15.2	9.5	9.7	4.0	3.8	1.9	1.9
Pipe Tobacco	1.5	1.9	1.0	0.7	0.8	0.5	0.5	0.6

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.