

Table 7.113C Illicit Drug Use in the Past Month, by Levels of Past Month Alcohol Use and Past Month Cigarette Use and Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

Level of Alcohol Use/ Cigarette Use	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
TOTAL	401	430	74	73	146	155	360	380
Heavy Alcohol Use ¹	178	178	23	27	86	83	154	146
Binge Alcohol Use But Not Heavy Use ¹	210	202	40	44	79	81	192	174
Alcohol Use But Not Binge Use ¹	202	223	27	28	45	51	198	215
No Alcohol Use	191	181	50	45	44	45	174	165
PAST MONTH CIGARETTE USE	285	310	55	52	120	116	259	270
Heavy Alcohol Use ¹	157	152	22	25	77	68	132	128
Binge Alcohol Use But Not Heavy Use ¹	158	166	34	33	69	66	142	149
Alcohol Use But Not Binge Use ¹	123	136	21	21	35	39	118	126
No Alcohol Use	111	121	28	24	34	33	101	113
NO PAST MONTH CIGARETTE USE	269	269	50	50	68	81	249	253
Heavy Alcohol Use ¹	92	88	13	10	32	43	85	79
Binge Alcohol Use But Not Heavy Use ¹	134	119	22	27	38	41	127	104
Alcohol Use But Not Binge Use ¹	162	161	16	18	27	33	156	157
No Alcohol Use	150	133	40	38	31	30	140	119

*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.113D Illicit Drug Use in the Past Month, by Levels of Past Month Alcohol Use and Past Month Cigarette Use and Age Group: Standard Errors of Percentages, 2002 and 2003

Level of Alcohol Use/ Cigarette Use	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
TOTAL	0.17	0.17	0.29	0.27	0.37	0.40	0.20	0.20
Heavy Alcohol Use ¹	1.02	0.93	2.22	2.50	1.11	1.08	1.34	1.23
Binge Alcohol Use But Not Heavy Use ¹	0.52	0.51	1.39	1.51	0.78	0.76	0.65	0.61
Alcohol Use But Not Binge Use ¹	0.31	0.33	1.40	1.42	0.68	0.74	0.34	0.37
No Alcohol Use	0.17	0.15	0.24	0.21	0.36	0.36	0.21	0.19
PAST MONTH CIGARETTE USE	0.45	0.45	1.21	1.17	0.66	0.70	0.55	0.55
Heavy Alcohol Use ¹	1.42	1.29	2.47	2.71	1.27	1.32	2.00	1.85
Binge Alcohol Use But Not Heavy Use ¹	0.87	0.90	2.11	2.03	1.17	1.10	1.18	1.24
Alcohol Use But Not Binge Use ¹	0.81	0.94	3.00	3.17	1.41	1.59	0.96	1.08
No Alcohol Use	0.56	0.54	1.73	1.69	1.12	1.13	0.62	0.62
NO PAST MONTH CIGARETTE USE	0.15	0.15	0.23	0.21	0.34	0.39	0.18	0.18
Heavy Alcohol Use ¹	1.43	1.37	4.65	4.51	1.91	2.12	1.72	1.66
Binge Alcohol Use But Not Heavy Use ¹	0.60	0.54	1.84	2.06	0.93	0.92	0.72	0.62
Alcohol Use But Not Binge Use ¹	0.31	0.31	1.24	1.29	0.65	0.76	0.34	0.34
No Alcohol Use	0.16	0.13	0.21	0.19	0.33	0.31	0.21	0.17

*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.113P Illicit Drug Use in the Past Month, by Levels of Past Month Alcohol Use and Past Month Cigarette Use and Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Level of Alcohol Use/ Cigarette Use	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
TOTAL	0.6300		0.3311		0.8639		0.6286	
Heavy Alcohol Use ¹	0.9168		0.4541		0.7689		0.9522	
Binge Alcohol Use But Not Heavy Use ¹	0.8866		0.5030		0.6717		0.7008	
Alcohol Use But Not Binge Use ¹	0.2621		0.6285		0.2315		0.4334	
No Alcohol Use	0.1000		0.1469		0.4348		0.2763	
PAST MONTH CIGARETTE USE	0.6156		0.8496		0.6767		0.5208	
Heavy Alcohol Use ¹	0.7812		0.5792		0.2545		0.9239	
Binge Alcohol Use But Not Heavy Use ¹	0.9008		0.7888		0.6991		0.9311	
Alcohol Use But Not Binge Use ¹	0.2048		0.8757		0.2474		0.3433	
No Alcohol Use	0.7832		0.6297		0.5048		0.7781	
NO PAST MONTH CIGARETTE USE	0.5089		0.7426		0.2022		0.3078	
Heavy Alcohol Use ¹	0.8765		0.4667		0.1993		0.7443	
Binge Alcohol Use But Not Heavy Use ¹	0.8118		0.3890		0.8960		0.5251	
Alcohol Use But Not Binge Use ¹	0.3972		0.9842		0.2496		0.5902	
No Alcohol Use	0.0713		0.4469		0.7165		0.1130	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.