

HIW Indicators

Version 1.6.1

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Abstinence discussion with parent before age 18: females (percent)	Percent of adolescents females who talked to a parent or guardian about abstinence before they were 18 years old	Yes				810
Abstinence discussion with parent before age 18: males (percent)	Percent of adolescents males who talked to a parent or guardian about abstinence before they were 18 years old	Yes				811
Abstinence instruction receipt before age 18: females (percent)	Percent of adolescents females who received formal instruction on abstinence before they were 18 years old	Yes				802
Abstinence instruction receipt before age 18: males (percent)	Percent of adolescents males who received formal instruction on abstinence before they were 18 years old	Yes				803
Activity limitation due to arthritis: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who experience a limitation in activity due to arthritis or joint symptoms	Yes				419
Activity limitation due to chronic back conditions (per 1,000)	Activity limitation due to chronic back conditions	Yes				428
Activity limitations: adults 45+ years with COPD (percent)	Percent of adults aged 45 years and over with chronic obstructive pulmonary disease (COPD) who have activity limitations	Yes				1405
Activity limitations: persons with asthma (percent)	Percent of persons with current asthma who have activity limitations	Yes				1404
Acute chemical poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: acute chemical poisoning	Yes				718
Acute hospital readmissions (count)	Total inpatient readmissions within 30 days of an acute hospital stay	Yes	Yes		Yes	278
Acute hospital readmissions (percent)	Inpatient readmissions within 30 days of an acute hospital stay	Yes	Yes		Yes	279

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Acute inpatient prospective payment system Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who had covered acute inpatient prospective payment system services	Yes	Yes		Yes	167
Acute inpatient prospective payment system Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered acute inpatient prospective payment system services	Yes	Yes		Yes	168
Added sugar consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake from added sugars for persons aged 2 years and over	Yes				1237
Adolescents having an adult to discuss problems with (percent)	Percent of adolescents who have an adult in their lives with whom they can talk about serious problems	Yes				409
Adolescents' parents attend activities/events (percent)	Percent of parents who attend events and activities in which their adolescents participate	Yes				410
Aerobic physical activity meeting guidelines: adults (percent)	Percent of adults who engage in aerobic physical activity of at least moderate intensity for at least 150 minutes/week, or 75 minutes/week of vigorous intensity, or an equivalent combination	Yes				1319
Aerobic physical activity meeting guidelines: teens (percent)	Percent of adolescents who meet current Federal physical activity guidelines for Aerobic physical activity	Yes	Yes			1323
Aerobic physical activity meeting high guidelines: adults (percent)	Percent of adults who engage in aerobic physical activity of at least moderate intensity for more than 300 minutes/week, or more than 150 minutes/week of vigorous intensity, or an equivalent combination	Yes				1320
Aerobic physical and muscle-strengthening activity: adults (percent)	Percent of adults who meet the objectives for aerobic physical activity and for muscle-strengthening activity	Yes				1322
African American Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months who are African American	Yes	Yes		Yes	312
AIDS, perinatally acquired (number)	Number of new cases of perinatally acquired AIDS	Yes				939
AIDS: heterosexual persons 13+ years (number)	Number of new AIDS cases among adolescent and adult heterosexuals	Yes				943

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
AIDS: injection drug users (number)	Number of new AIDS cases among adolescents and adults who inject drugs	Yes				931
AIDS: men who have sex with men (number)	Number of new AIDS cases among adolescent and adult men who have sex with men	Yes				930
AIDS: persons 13+ years (per 100,000)	Rate of new AIDS cases among adolescents and adults	Yes				929
Air quality management program: schools (percent)	Percent of the Nation's elementary, middle, and high schools that engage in practices that promote a healthy and safe physical school environment: Have an indoor air quality management program	Yes				683
Airborne toxins from area sources (number)	Air toxic emissions to decrease the risk of adverse health effects caused by airborne toxics - area sources	Yes				671
Airborne toxins from major sources (number)	Air toxic emissions to decrease the risk of adverse health effects caused by airborne toxics - major sources	Yes				672
Airborne toxins from mobile sources (number)	Air toxic emissions to decrease the risk of adverse health effects caused by airborne toxics - mobile sources	Yes				670
Alcohol abstinence, preconception (percent)	Percent of women delivering a live birth who did not drink alcohol prior to pregnancy	Yes				1181
Alcohol abstinence, prenatal (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Alcohol	Yes				1142
Alcohol and/or illicit drug treatment receipt (percent)	Percent of persons who need alcohol and/or illicit drug treatment and received specialty treatment for abuse or dependence in the past year-Alcohol and/or illicit drug treatment	Yes				1455
Alcohol Screening and Brief Intervention: primary care & trauma centers (number)	Number of Level I and Level II trauma centers and primary care settings that implement evidence-based alcohol Screening and Brief Intervention (SBI)	Yes				1460
Alcohol sold (number)	Average annual alcohol consumption rate	Yes				1439

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Alcohol treatment receipt (percent)	Percent of persons who need alcohol and/or illicit drug treatment and received specialty treatment for abuse or dependence in the past year-Alcohol abuse or dependence	Yes				1456
Alcohol, never use: at risk teens (percent)	Percent of at risk adolescents aged 12 to 17 years who, in the past year, refrained from using alcohol for the first time	Yes				1428
Alcohol, never use: high school seniors (percent)	Percent of high school seniors never using substances-Alcoholic beverages	Yes				1430
Alcohol-attributable deaths (number)	Number of deaths attributable to alcohol	Yes				1467
Alcohol-impaired driving deaths (per 100 vehicle miles)	Alcohol-impaired driving (08+ blood alcohol content [BAC]) fatality rate	Yes				1466
Allergic reactions to food: adults (percent)	Percent of adults with a physician-diagnosed food allergy with a severe allergic reactions to food	Yes				820
Ambulatory surgery center Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used ambulatory surgery center services	Yes	Yes		Yes	233
Ambulatory surgery center Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used ambulatory surgery center services	Yes	Yes		Yes	234
Ambulatory surgical center Medicare service events (per 1,000 beneficiaries)	Rate of ambulatory surgery center service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2020
Ambulatory Surgical Centers (number)	Number of Medicare participating Ambulatory Surgical Centers		Yes		Yes	10033
American Indian or Alaska Native Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months who are American Indian or Alaska Native	Yes	Yes		Yes	315
Amputation among diabetics rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Rate of lower extremity amputation among patients with diabetes among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	2046
Anencephaly (per 100,000 live births)	Rate of anencephaly	Yes				1170

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Antibiotic use: persons with the common colds (per 100,000)	Rate of courses of antibiotics prescribed for the sole diagnosis of the common cold	Yes				975
Antibiotics for ear infections: children <5 years (percent)	Rate of physician visits by children less than 5 years of age diagnosed with an ear infection that have an antibiotic course ordered, supplied, administered, or continued	Yes				974
Antimicrobial resistance to 3+ classes of antimicrobials (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to three or more classes of antimicrobial agents	Yes				833
Antimicrobial resistance to Ampicillin (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to ampicillin	Yes				832
Antimicrobial resistance to Ceftriaxone (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to Ceftriaxone (third-generation cephalosporin)	Yes				830
Antimicrobial resistance to Erythromycin (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Campylobacter jejuni isolates from humans that are resistant to Erythromycin	Yes				834
Antimicrobial resistance to Gentamicin (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to gentamicin	Yes				831
Antimicrobial resistance to Nalidixic acid (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to nalidixic acid (quinolone)	Yes				829

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Appropriate treatment use: adults with balance/dizziness problems (percent)	Percent of adults who have tried recommended methods for treating their balance or dizziness problem	Yes				765
Appropriate treatment use: adults with tinnitus (percent)	Percent of adults for whom tinnitus is a moderate to severe problem who have tried appropriate treatments	Yes				761
Arsenic level (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: arsenic	Yes				695
Arsenic poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: arsenic poisoning	Yes				716
Arteriovenous fistula use: hemodialysis patients (percent)	Percent of adult hemodialysis patients who use arteriovenous fistulas or have a maturing fistula as the primary mode of vascular access at the start of renal replacement therapy	Yes				471
Arthritis education: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who have had effective, evidence-based arthritis education as an integral part of the management of their condition	Yes				426
Artificial UV tanning: adults (percent)	Percent of adults aged 18 and older who report using artificial sources of ultraviolet light for tanning	Yes				502
Artificial UV tanning: teens (percent)	Percent of adolescents in grades 9 through 12 who report using artificial sources of ultraviolet light for tanning	Yes				501
Asian American or Pacific Islander Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months who are Asian American or Pacific Islander	Yes	Yes		Yes	314
Assaults, work-related (per 10,000)	Rate of work-related assaults	Yes				1311
Assistive listening device use: older adults (per 1,000)	Rate of adults aged 70 years and older with hearing loss who use assistive listening devices	Yes				747
Assistive/adaptive devices use: adults with visual impairment (percent)	Percent of persons aged 18 years and over with visual impairment who use assistive and adaptive devices	Yes				1575

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Associate degrees in public health: 2-year colleges (percent)	Percent of 2-year colleges that offer public health or related Associate degrees	Yes				1388
Asthma deaths: adults 35-64 years (per 1,000,000)	Asthma death rate: Adults aged 35 to 64 years old	Yes				1408
Asthma deaths: older adults (per 1,000,000)	Asthma death rate: Adults aged 65 years and over	Yes				1409
Asthma deaths: persons <35 years (per 1,000,000)	Asthma death rate: Children and adults under age 35 years	Yes				1407
Asthma ED visits: children <5 years (per 10,000)	Rate of hospital emergency department visits for asthma: Children under age 5 years	Yes				1401
Asthma ED visits: older adults (per 10,000)	Rate of hospital emergency department visits for asthma: Adults aged 65 years and over	Yes				1403
Asthma ED visits: persons 5-64 years (per 10,000)	Rate of hospital emergency department visits for asthma: Children and adults aged 5 to 64 years	Yes				1402
Asthma exposure risks, advise to reduce: persons with asthma (percent)	Percent of persons with current asthma who have been advised by a health professional to change things in their home, school, and work environments to reduce exposure to irritants or allergens to which they are sensitive	Yes				1417
Asthma hospitalizations: children <5 years (per 10,000)	Rate of hospitalizations for asthma: Children under age 5 years	Yes				1398
Asthma hospitalizations: older adults (per 10,000)	Rate of hospitalizations for asthma: Adults aged 65 years and over	Yes				1400
Asthma hospitalizations: persons 5-64 years (per 10,000)	Rate of hospitalizations for asthma: Children and adults aged 5 to 64 years	Yes				1399
Asthma inhalers, receipt of instructions (percent)	Percent of persons with current asthma with prescribed inhalers who receive instruction on their use	Yes				1414
Asthma management plans receipt from health care provider (percent)	Percent of persons with current asthma who receive written asthma management plans from their health care provider	Yes				1413

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Asthma Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have asthma	Yes	Yes		Yes	50007
Asthma Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have asthma	Yes	Yes		Yes	50008
Asthma medication regimen, appropriate (percent)	Percent of persons with current asthma who do not use more than one canister of short-acting inhaled beta agonist per month	Yes				1416
Asthma patient education receipt (percent)	Percent of persons with current asthma who receive formal patient education	Yes				1412
Asthma signs/symptoms and responses, education receipt (percent)	Percent of persons with current asthma who receive education about appropriate response to an asthma episode, including recognizing early signs and symptoms or monitoring peak flow results	Yes				1415
Asthma surveillance systems: states (number)	Numbers of states, territories, and the District of Columbia with a comprehensive asthma surveillance system for tracking asthma cases, illness, and disability at the State level	Yes				1421
Atrial fibrillation Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have atrial fibrillation	Yes	Yes		Yes	285
Atrial fibrillation Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have atrial fibrillation	Yes	Yes		Yes	286
Autism spectrum disorder screening (percent)	Percent of young children who are screened for an autism spectrum disorder (ASD) and other developmental delays by 24 months of age	Yes				1188
Average age of Medicare beneficiaries 65+ (mean)	Average age of Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	308
Average total hierarchical condition category score for Medicare beneficiaries 65+ (mean)	Average hierarchical condition category score of Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	318

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Bacterial pneumonia admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Bacterial pneumonia admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	2043
BDE 47 level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: BDE 47, (2,2',4,4'-tetrabromodiphenyl ether)	Yes				712
Beaches open and safe (percent)	Percent of days that beaches are open and safe for swimming	Yes				674
Beta-HCH level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: beta-hexachlorocyclohexane or beta-HCH	Yes				702
Bicycle helmet laws: states (number)	Number of states and the District of Columbia with laws requiring bicycle helmets for bicycle riders	Yes				1049
Bicycling for transportation (percent)	Percent of trips to work using of alternative modes of transportation - bicycling	Yes				666
Binge drinking abstinence, prenatal (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Binge drinking	Yes				1143
Binge drinking: adults (percent)	Percent of persons engaging in binge drinking during the past month-Adults aged 18 years and older	Yes				1437
Binge drinking: adults (percent)	Percent of adults 18 years and over that report binge drinking		Yes	Yes		17
Binge drinking: college students (percent)	Percent of students engaging in binge drinking during the past 2 weeks-College students	Yes				1436
Binge drinking: high school seniors (percent)	Percent of students engaging in binge drinking during the past 2 weeks-High school seniors	Yes				1435
Binge drinking: teens (percent)	Percent of persons engaging in binge drinking during the past month-Adolescents aged 12 to 17 years	Yes				1438

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Birth control discussion with parent before age 18: females (percent)	Percent of adolescents females who talked to a parent or guardian about birth control methods before they were 18 years old	Yes				812
Birth control discussion with parent before age 18: males (percent)	Percent of adolescents males who talked to a parent or guardian about birth control methods before they were 18 years old	Yes				813
Birth control instruction receipt before age 18: females (percent)	Percent of adolescents females who received formal instruction on birth control methods before they were 18 years old	Yes				804
Birth control instruction receipt before age 18: males (percent)	Percent of adolescents males who received formal instruction on birth control methods before they were 18 years old	Yes				805
Births, moderately preterm (32-36 wk.) (percent)	Percent of live births that are moderately preterm (32-36 weeks gestation)	Yes	Yes	Yes		10008
Births: unmarried women 18-54 years (percent)	Percent of all live births to women ages 18-54 years who report not being married	Yes	Yes	Yes		99
Births: women 40-54 years (percent)	Percent of all live births to women 40-54 years of age	Yes	Yes	Yes		98
Births: women under 18 years (percent)	Percent of all live births to mothers less than 18 years of age	Yes	Yes	Yes		97
Births:females 15-19 years (per 1,000)	Rate per 1,000 of births to females 15-19 years of age	Yes	Yes	Yes		22
Bisphenol A level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Bisphenol A	Yes				709
Blood donations (percent)	Percent of persons who donate blood	Yes				436
Blood foliate concentration: reproductive-aged women (percent)	Percent of women of childbearing potential who have low red blood-cell foliate concentrations	Yes				1172
Blood pressure control: adults with hypertension (percent)	Percent of adults with hypertension whose blood pressure is under control	Yes				882

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Blood pressure screening: adults (percent)	Percent of adults who have had their blood pressure measured within the preceding 2 years and can state whether their blood pressure was normal or high	Yes				883
Blood pressure under control: adults with diabetes (percent)	Percent of adults with diagnosed diabetes whose blood pressure is under control	Yes				560
Body mass index assessment of adults by physicians (percent)	Percent of primary care physicians who regularly assess body mass index (BMI) in their adult patients	Yes				1234
Body mass index assessment of children by physicians (percent)	Percent of primary care physicians who regularly assess body mass index (BMI) for age and sex in their child or adolescent patients	Yes				1235
Breakfast program: schools (percent)	Percent of schools with a school breakfast program	Yes				417
Breast cancer deaths (per 100,000)	Breast cancer death (female) rate	Yes	Yes	Yes		488
Breast cancer Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have breast cancer	Yes	Yes		Yes	299
Breast cancer Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have breast cancer	Yes	Yes		Yes	300
Breastfeeding, at 1 year (percent)	Percent of infants who are breastfed-At 1 year	Yes				1149
Breastfeeding, at 6 months (percent)	Percent of infants who are breastfed-At 6 months	Yes				1148
Breastfeeding, ever (percent)	Percent of infants who are breastfed-Ever	Yes				1147
Breastfeeding, exclusively through 3 months (percent)	Percent of infants who are breastfed-Exclusively through 3 months	Yes				1150
Breastfeeding, exclusively through 6 months (percent)	Percent of infants who are breastfed-Exclusively through 6 months	Yes				1151
Broadband access to internet (percent)	Percent of persons with broadband access to the Internet	Yes				858
Bullying: teens (percent)	Percent of bullying among adolescents	Yes	Yes			1102
c. Listeria monocytogenes infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Reduction in infections caused by Listeria monocytogenes	Yes				824

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Cadmium level (µg/L)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: cadmium	Yes				696
Cadmium poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: cadmium poisoning	Yes				717
Calcium consumption (milligrams)	Mean total daily calcium intake (in mg) by persons aged 2 years and over	Yes				1223
Campylobacter species infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Campylobacter species	Yes				822
Cancer survival (percent)	Percent of cancer survivors who are living 5 years or longer after diagnosis	Yes				498
Carbon monoxide poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: carbon monoxide poisoning	Yes				719
Cardiac surgery patients with controlled 6 AM postoperative blood glucose (percent)	Hospital Compare: Cardiac surgery patients with controlled 6 AM postoperative blood glucose	Yes	Yes		Yes	339
Cardiovascular death: hemodialysis patients (per 1,000)	Cardiovascular death rate for persons on hemodialysis	Yes				465
Cardiovascular death: kidney transplant patients (per 1,000)	Cardiovascular death rate for persons with a functioning kidney transplant	Yes				467
Catheter use for vascular access: hemodialysis patients (percent)	Percent of adult hemodialysis patients who use catheters as the only mode of vascular access	Yes				470
Central line-associated bloodstream infections (Standardized Infection Ratio (SIR))	Central line-associated bloodstream infections	Yes				851

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Certificate programs in public health: 2-year colleges (percent)	Percent of 2-year colleges that offer public health or related Certificate programs	Yes				1389
Certificates of birth, use of latest standard form: states (number)	Number of States that record vital events using the latest U.S. standard certificates and report-States using the standard certificate of birth	Yes				1382
Certificates of death, use of latest standard form: states (number)	Number of States that record vital events using the latest U.S. standard certificates and report - States using the standard certificate of death	Yes				1383
Cervical cancer screening: women 21-65 years (percent)	Percent of women who receive a cervical cancer screening based on the most recent guidelines	Yes				505
Cesarean delivery, first (percent)	Percent of live births that are cesarean deliveries among low-risk (full-term, singleton, vertex presentation) women who have never previously had a cesarean birth	Yes				1133
Cesarean delivery, repeat (percent)	Percent of live births that are cesarean among low-risk (full-term, singleton, vertex presentation) women who have had a prior cesarean birth	Yes				1134
Child developmental concerns addressed by doctors: parents (percent)	Percent of parents who receive information from their doctors or other Health Care professionals when they have a concern about their children's learning, development, or behavior	Yes				741
Child maltreatment deaths (per 100,000)	Child maltreatment death rate	Yes				1085
Child maltreatment, nonfatal (per 1,000)	Nonfatal child maltreatment	Yes				1084
Childrens' Hospitals (number)	Number of Children's Hospitals participating in the Medicare program		Yes		Yes	10034
Chlamydia (per 100,000)	Chlamydia rate		Yes	Yes		20
Chlamydia cases (number)	Number of reported cases of chlamydia		Yes	Yes		10064

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Chlamydia screening: females 16-20 years with Medicaid (percent)	Percent of sexually active females aged 16 to 20 years enrolled in Medicaid plans who are screened for genital Chlamydia infections during the measurement year	Yes				1486
Chlamydia screening: females 16-20 years with private health insurance (percent)	Percent of sexually active females aged 16 to 20 years enrolled in commercial health insurance plans who are screened for genital Chlamydia infections during the measurement year	Yes				1488
Chlamydia screening: females 21-24 years with Medicaid (percent)	Percent of sexually active females aged 21 to 24 years enrolled in Medicaid plans who are screened for genital Chlamydia infections during the measurement year	Yes				1487
Chlamydia screening: females 21-24 years with private health insurance (percent)	Percent of sexually active females aged 21 to 24 years enrolled in commercial health insurance plans who are screened for genital Chlamydia infections during the measurement year	Yes				1489
Chlamydia: females <24 years in a national job training program (percent)	Percent of adolescents and young adults with Chlamydia trachomatis infections-Among females aged 24 years and under enrolled in a National Job Training Program	Yes				1476
Chlamydia: females 15-24 years attending family planning clinics (percent)	Percent of adolescents and young adults with Chlamydia trachomatis infections-Among females aged 15 to 24 years attending family planning clinics	Yes				1475
Chlamydia: males <24 years in a national job training program (percent)	Percent of adolescents and young adults with Chlamydia trachomatis infections-Among males aged 24 years and under enrolled in a National Job Training Program	Yes				1477
Chlordane level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Chlordane (Oxychlordane)	Yes				700

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Chlorpyrifos level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: 3,4,6-trichloro-2-pyridinol (chlorpyrifos)	Yes				704
Cholesterol levels: adults (mean)	Mean total blood cholesterol levels among adults	Yes				884
Cholesterol screening: adults (percent)	Percent of adults who have had their blood cholesterol checked within the preceding 5 years	Yes				886
Chronic diseases prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in chronic disease programs	Yes				602
Chronic hepatitis B: children 1-24 months (number)	Number of chronic hepatitis B virus infections in infants and young children (perinatal infections)	Yes				962
Chronic kidney disease (percent)	Percent of the adults with chronic kidney disease	Yes				483
Chronic kidney disease deaths (per 100)	Death rate among adults 18 years or older with chronic kidney disease	Yes				484
Chronic kidney disease Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have chronic kidney disease	Yes	Yes		Yes	287
Chronic kidney disease Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have chronic kidney disease	Yes	Yes		Yes	288
Chronic obstructive pulmonary disease (COPD) Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have chronic obstructive pulmonary disease	Yes	Yes		Yes	289
Chronic obstructive pulmonary disease (COPD) Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have chronic obstructive pulmonary disease	Yes	Yes		Yes	290

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Chronic obstructive pulmonary disease admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Chronic obstructive pulmonary disease admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	2039
Cigar smoking: adults (percent)	Percent of adults aged 18 years and over smoking cigars	Yes				1500
Cigar use in past month: teens (percent)	Percent of adolescent reporting use of cigars in the past month - Students (grades 9 through 12)	Yes	Yes			1504
Cigar use initiation: adults 18-25 years (percent)	Percent of young adults aged 18 to 25 years initiating cigar use in the past 12 months	Yes				1512
Cigar use initiation: teens (percent)	Percent of children and adolescents aged 12 to 17 years initiating cigar use in past 12 months	Yes				1508
Cigarette abstinence, prenatal (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Cigarette smoking	Yes				1144
Cigarette smoking: adults (percent)	Percent of adults aged 18 years and over smoking cigarettes	Yes				1498
Cigarette use in past month: teens (percent)	Percent of adolescent reporting use of cigarettes in the past month - Students (grades 9 through 12)	Yes	Yes			1502
Cigarette use initiation: adults 18-25 years (percent)	Percent of young adults aged 18 to 25 years initiating cigarette use in the past 12 months	Yes				1510
Cigarette use initiation: teens (percent)	Percent of children and adolescents aged 12 to 17 years initiating cigarette use in past 12 months	Yes				1506
Cirrhosis deaths (per 100,000)	Cirrhosis death rate	Yes	Yes			1425
Clean hands, food safety practice (percent)	Percent of consumers who follow key food safety practices-- Clean: Wash hands and surfaces often	Yes				835
Clinical Laboratory Improvement Act (CLIA) Providers (number)	Number of Clinical Laboratory Improvement Act (CLIA) providers participating in the Medicare program		Yes		Yes	10035
Clinical preventive measure use: females 65+ years (percent)	Percent of older adults who are up to date on a core set of clinical preventive measures-Females	Yes				1262

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Clinical preventive measure use: males 65+ years (percent)	Percent of older adults who are up to date on a core set of clinical preventive measures-Males	Yes				1261
Cochlear implant surgeries: deaf persons (per 10,000)	Rate of persons who are deaf or very hard of hearing and who have new cochlear implants	Yes				745
College degree: adults 25+ (percent)	Percent of population age 25+ with 4-year college degree or higher		Yes	Yes		34
Colonoscopy/sigmoidoscopy: adults 50+ (percent)	Percent of adults age 50+ that report ever having had a sigmoidoscopy or colonoscopy		Yes	Yes		117
Colorectal cancer deaths (per 100,000)	Colorectal cancer death rate per 100,000	Yes	Yes	Yes		86
Colorectal cancer deaths, including unspecified sites (per 100,000)	Colorectal cancer death rate, including unspecified sites, per 100,000	Yes				490
Colorectal cancer Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have colorectal cancer	Yes	Yes		Yes	301
Colorectal cancer Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have colorectal cancer	Yes	Yes		Yes	302
Colorectal cancer screening: persons 50-75 years (percent)	Percent of adults who receive a colorectal cancer screening based on the most recent guidelines	Yes				506
Community Mental Health Centers (number)	Number of Community Mental Health Centers participating in the Medicare program		Yes		Yes	10036
Competencies for public health professionals: Local agencies (percent)	Percent of Federal, Tribal, State, and local public health agencies that incorporate core competencies for public health professionals into job descriptions and performance evaluations-Local public health agencies	Yes				1372
Complete vaccination: children 19-35 months (percent)	Percent of children aged 19 to 35 months who receive the recommended doses of DTaP, polio, MMR, Hib, hepatitis B, varicella and PCV	Yes				1008
Comprehensive care systems: children with special needs (percent)	Percent of children with special health care needs who receive their care in family-centered, comprehensive, coordinated systems-Children aged 0 to 11 years	Yes				1154

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Comprehensive care systems: teens with special needs (percent)	Percent of children with special health care needs who receive their care in family-centered, comprehensive, coordinated systems-Children aged 12 to 17 years	Yes				1155
Comprehensive Outpatient Rehabilitation Facilities (number)	Number of Comprehensive Outpatient Rehabilitation (CORF) Facilities participating in the Medicare program		Yes		Yes	10037
Condition not improving: adults with balance/dizziness problems (percent)	Percent of adults with balance and dizziness problems in the past 12 months who reported their condition got worse or did not improve	Yes				766
Condom & hormonal/intrauterine contraception use at first intercourse: females 15-19 years (percent)	Percent of sexually active females aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at first intercourse	Yes				798
Condom & hormonal/intrauterine contraception use at first intercourse: males 15-19 years (percent)	Percent of sexually active males aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at first intercourse	Yes				799
Condom & hormonal/intrauterine contraception use at last intercourse: females 15-19 years (percent)	Percent of sexually active females aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at last intercourse	Yes				800
Condom & hormonal/intrauterine contraception use at last intercourse: males 15-19 years (percent)	Percent of sexually active males aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at last intercourse	Yes				801
Condom use at first intercourse: females 15-19 years (percent)	Percent of sexually active females aged 15 to 19 years who use a condom at first intercourse	Yes				794
Condom use at first intercourse: males 15-19 years (percent)	Percent of sexually active males aged 15 to 19 years who use a condom at first intercourse	Yes				795
Condom use at last intercourse: females 15-19 years (percent)	Percent of sexually active females aged 15 to 19 years who use a condom at last intercourse	Yes				796
Condom use at last intercourse: males 15-19 years (percent)	Percent of sexually active males aged 15 to 19 years who use a condom at last intercourse	Yes				797

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Condom use: unmarried females 15-44 years (percent)	Percent of sexually active persons who use condoms- Unmarried females aged 15 to 44 years	Yes				940
Condom use: unmarried males 15-44 years (percent)	Percent of sexually active persons who use condoms- Unmarried males aged 15 to 44 years	Yes				941
Congenital Rubella Syndrome (number)	Number of Congenital Rubella Syndrome (CRS) among children aged 1 year and under (U.S.-acquired cases)	Yes				978
Congestive heart failure admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Congestive heart failure admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	2041
Congregate care: adults with disabilities (number)	Number of adults with disabilities (aged 22 years and older) living in congregate care facilities with 16 beds or more	Yes				524
Congregate care: persons <22 years with disabilities (number)	Number of children and youth with disabilities (aged 21 years and under) living in congregate care facilities with 16 beds or more	Yes				525
Contraceptive services/supplies need and receipt through public sector (percent)	Percent of females in need of publicly supported contraceptive services and supplies who receive those services and supplies	Yes				819
Contraceptive use at most recent sexual intercourse (percent)	Percent of females or their partners at risk of unintended pregnancy who used contraception at most recent sexual intercourse	Yes				785
Cook to proper temperature, food safety practice (percent)	Percent of consumers who follow key food safety practices-- Cook: Cook to proper temperatures	Yes				837
COPD deaths: adults 45+ years (per 100,000)	Chronic obstructive pulmonary disease (COPD) death rate among adults aged 45 years and over	Yes				1406
COPD ED visits: adults 45+ years (per 10,000)	Rate of hospital emergency department visits for chronic obstructive pulmonary disease (COPD) among adults aged 45 years and over	Yes				1423
COPD hospitalizations: adults 45+ years (per 10,000)	Hospitalization rate for chronic obstructive pulmonary disease (COPD) among adults aged 45 years and over	Yes				1422

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Core Competencies in Public Health in curricula (percent)	Percent of Council on Education for Public Health (CEPH) accredited schools of public health, CEPH accredited academic programs, and schools of nursing (with a public health or community health component) that integrate Core Competencies for Public Health Professionals into curricula	Yes				1373
Coronary heart disease deaths (per 100,000)	Coronary heart disease death rate	Yes	Yes	Yes		877
Critical Access Hospital Medicare admissions (per 1,000 beneficiaries)	Rate of critical access hospital (CAH) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2003
Critical access hospital Medicare days (per 1,000 beneficiaries)	Average number of critical access hospital (CAH) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2011
Critical access hospital services Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who had covered critical access hospital services	Yes	Yes		Yes	174
Critical access hospital services Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered critical access hospital services	Yes	Yes		Yes	175
Critical Access Hospitals (number)	Number of Critical Access Hospitals participating in the Medicare program		Yes		Yes	10038
D.O. training, counseling for health promotion (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-DO-granting medical schools	Yes				612
D.O. training, cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-DO-granting medical schools	Yes				613
D.O. training, environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-DO-granting medical schools	Yes				615

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
D.O. training, evaluation of health sciences literature (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-DO-granting medical schools	Yes				614
D.O. training, global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-DO-granting medical schools	Yes				617
D.O. training, public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-DO-granting medical schools	Yes				616
DDT level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: DDT (DDE)	Yes				701
Death: kidney transplant patients (per 1,000)	Total death rate for persons with a functioning kidney transplant	Yes				466
Deaths, all causes (per 100,000)	Death rate per 100,000 population from all causes of death	Yes	Yes	Yes		71
Deaths: adults 20-24 years (per 100,000)	Rates of adolescent and young adult deaths-Young adults aged 20 to 24 years	Yes	Yes			1128
Deaths: children 1-4 years (per 100,000)	Rate of child deaths-Children aged 1 to 4 years	Yes	Yes			1124
Deaths: children 10-14 years (per 100,000)	Rates of adolescent and young adult deaths-Adolescents aged 10 to 14 years	Yes	Yes			1126
Deaths: children 5-9 years (per 100,000)	Rate of child deaths-Children aged 5 to 9 years	Yes	Yes			1125
Deaths: infants with Down syndrome (per 1,000)	One-year mortality rate for infants with Down syndrome	Yes				1187
Deaths: teens 15-19 years (per 100,000)	Rates of adolescent and young adult deaths-Adolescents aged 15 to 19 years	Yes	Yes			1127

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Dehydration admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Dehydration admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	2042
Dental care programs: school-based health centers (percent)	Percent of school-based health centers with an oral health component that includes dental care	Yes				1285
Dental caries experience: children 3-5 years (percent)	Percent of young children aged 3 to 5 years with dental caries experience in their primary teeth	Yes				1269
Dental caries experience: children 6-9 years (percent)	Percent of children aged 6 to 9 years with dental caries experience in their primary and permanent teeth	Yes				1270
Dental caries experience: teens 13-15 years (percent)	Percent of adolescents aged 13 to 15 years with dental caries experience in their permanent teeth	Yes				1271
Dental decay, untreated: adults 35-44 years (percent)	Percent of adults aged 35 to 44 years with untreated dental decay	Yes				1275
Dental decay, untreated: adults 65-74 years (percent)	Percent of older adults aged 65 to 74 years with untreated coronal caries	Yes				1276
Dental decay, untreated: adults 75+ years (percent)	Percent of older adults aged 75 years and older with untreated root surface caries	Yes				1277
Dental decay, untreated: children 3-5 years (percent)	Percent of young children aged 3 to 5 years with untreated dental decay in their primary teeth	Yes				1272
Dental decay, untreated: children 6-9 years (percent)	Percent of children aged 6 to 9 years with untreated dental decay in their primary and permanent teeth	Yes				1273
Dental decay, untreated: teens 13-15 years (percent)	Percent of adolescents aged 13 to 15 years with untreated dental decay in their permanent teeth	Yes				1274
Dental examinations, annual: adults with diabetes (percent)	Percent of persons with diagnosed diabetes who have at least an annual dental examination	Yes				553
Dental professional directing dental program: IHS programs (number)	Number of Indian Health Service Areas and Tribal health programs that serve jurisdictions of 30,000 or more persons with a dental public health program directed by a dental professional with public health training	Yes				1292

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Dental professional directing dental program: state & local agencies (percent)	Percent of States (including the District of Columbia) and local health agencies that serve jurisdictions of 250,000 or more persons with a dental public health program directed by a dental professional with public health training	Yes				1291
Dental sealant programs: school-based health centers (percent)	Percent of school-based health centers with an oral health component that includes dental sealants	Yes				1284
Dental sealants: children 3-5 years (percent)	Percent of children aged 3 to 5 years who have received dental sealants on one or more of their primary molar teeth	Yes				1281
Dental sealants: children 6-9 years (percent)	Percent of children aged 6 to 9 years who have received dental sealants on one or more of their permanent first molar teeth	Yes				1282
Dental sealants: teens 13-15 years (percent)	Percent of adolescents aged 13 to 15 years who have received dental sealants on one or more of their permanent molar teeth	Yes				1283
Dental service use: low-income children (percent)	Percent of low-income children and adolescents who received any preventive dental service during the past year	Yes				1267
Dentist rate	Dentists per capita		Yes	Yes		130
Depression Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have depression	Yes	Yes		Yes	291
Depression Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have depression	Yes	Yes		Yes	292
Depression screening by primary care providers: adults (percent)	Percent of primary care physician office visits that screen adults aged 19 years and older for depression	Yes				1120
Depression screening by primary care providers: teens (percent)	Percent of primary care physician office visits that screen youth aged 12 to 18 years for depression	Yes				1121
Diabetes education (percent)	Percent of adults with diagnosed diabetes who receive formal diabetes education	Yes	Yes			545

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Diabetes long-term complications admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Diabetes long-term complications admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	2038
Diabetes management benefit use: older adults with diabetes (percent)	Percent of older adults who receive Diabetes Self-Management Benefits	Yes				1263
Diabetes Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have diabetes	Yes	Yes		Yes	293
Diabetes Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have diabetes	Yes	Yes		Yes	294
Diabetes, new cases (per 1,000)	Rate of newly diagnosed diabetes among adults aged 18-84	Yes				555
Diabetes-related deaths (per 100,000)	Diabetes death rate	Yes				547
Diabetes: adults (percent)	Percent of adults that report having been diagnosed as having diabetes		Yes	Yes		125
Diagnosed diabetes: adults with diabetes (percent)	Percent of adults aged 20 years and older with diabetes whose condition has been diagnosed	Yes				546
Dialysis Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used dialysis services	Yes	Yes		Yes	50005
Dialysis Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used dialysis services	Yes	Yes		Yes	50006
Dialysis Medicare visits (per 1,000 beneficiaries)	Rate of dialysis visits among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	50004
Dilated eye examinations: adults (percent)	Percent of adults who have a comprehensive eye examination, including dilation, within the past 2 years	Yes				1566

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Dioxin level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: 1,2,3,6,7,8-hexachlorodibenzo-p-dioxin, representative of the dioxin class	Yes				708
Disapproval of heavy drinking: 10th graders (percent)	Percent of adolescents who disapprove of having one or two alcoholic drinks nearly every day-10th graders	Yes				1445
Disapproval of heavy drinking: 8th graders (percent)	Percent of adolescents who disapprove of having one or two alcoholic drinks nearly every day-8th graders	Yes				1444
Disapproval of heavy drinking: high school seniors (percent)	Percent of adolescents who disapprove of having one or two alcoholic drinks nearly every day-12th graders	Yes				1446
Disapproval of trying marijuana: 10th graders (percent)	Percent of adolescents who disapprove of trying marijuana or hashish once or twice-10th graders	Yes				1448
Disapproval of trying marijuana: 8th graders (percent)	Percent of adolescents who disapprove of trying marijuana or hashish once or twice-8th graders	Yes				1447
Disapproval of trying marijuana: high school seniors (percent)	Percent of adolescents who disapprove of trying marijuana or hashish once or twice-12th graders	Yes				1449
Disorders diagnosed through newborn blood spot screening (percent)	Percent of children diagnosed with a disorder through newborn blood spot screening who experience developmental delay requiring special education services	Yes				1167
Domestic water use (gallons)	Per capita domestic water withdrawals with respect to use and conservation	Yes				673
Drownings (per 100,000)	Drowning death rate	Yes	Yes	Yes		1081
Drug poisoning deaths (per 100,000)	Rate per 100,000 of deaths with underlying causes of drug related poisonings	Yes	Yes	Yes		10016
Drug-induced deaths (per 100,000)	Drug-induced death rate	Yes	Yes			1426
DTaP vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-4 doses diphtheria-tetanus-acellular pertussis (DTaP) vaccine by age 19 to 35 months	Yes				993

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
DTaP vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-DTaP vaccine (4 or more doses)	Yes				1003
Durable medical equipment Medicare service events (per 1,000 beneficiaries)	Rate of durable medical equipment service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2026
Durable medical equipment Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used durable medical equipment services	Yes	Yes		Yes	257
Durable medical equipment Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used durable medical equipment services	Yes	Yes		Yes	258
E-code collection at hospital discharge: states (percent)	Percent of states and the District of Columbia with statewide hospital discharge data systems that routinely collect external-cause-of injury codes for 90 percent or more of injury-related discharges	Yes				1094
E-code collection in emergency department: states (percent)	Percent of states and the District of Columbia with statewide emergency department data systems that routinely collect external-cause-of injury codes for 90 percent or more of injury-related visits	Yes				1093
Early death: hemodialysis patients (per 1,000)	Death rate in hemodialysis patients within the first 3 months of initiation of renal replacement therapy	Yes				464
Early infant diagnostic hearing evaluation (percent)	Percent of infants who did not pass the hearing screening that received audiologic evaluation at no later than age 3 months	Yes				757
Early intervention services: children with disabilities (percent)	Percent of children with disabilities, birth through age 2 years, who receive early intervention services in home or community-based settings	Yes				539
Early intervention services: infants with hearing loss (percent)	Percent of infants with confirmed hearing loss enrolled for intervention services no later than age 6 months	Yes				758
Eating disorders: teens (percent)	Percent of adolescents who engage in disordered eating behaviors in an attempt to control their weight	Yes	Yes			1108

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Elder abuse, neglect, and exploitation information: States (number)	Number of States-and the District of Columbia-that collect and make publicly available information on the characteristics of victims, perpetrators, and cases of elder abuse, neglect, and exploitation	Yes				1256
Electronic health records in medical practices (percent)	Percent of medical practices that use electronic health records	Yes				872
Emergency department visit rate (per 1,000 beneficiaries)	Rate among beneficiaries of inpatient or hospital outpatient emergency department visits	Yes	Yes		Yes	282
Emergency department visits (count)	Total inpatient or hospital outpatient emergency department visits	Yes	Yes		Yes	281
Emergency personnel response time (minutes)	The time necessary to activate designated personnel in response to public health emergency in minutes	Yes				1394
Employment at indoor workplaces that prohibit smoking (percent)	Percent of persons covered by indoor worksite policies that prohibit smoking	Yes				1492
Employment: persons 16-64 years with disabilities (percent)	Employment rates among people with disabilities	Yes				516
Employment: persons with serious mental illness (percent)	Percent of persons with serious mental illness (SMI) who are employed	Yes				1116
End Stage Renal Disease Facilities (number)	Number of End Stage Renal Disease (ESRD) facilities participating in the Medicare program		Yes		Yes	10039
End-stage kidney failure due to diabetes (per 1,000,000)	Incidence rate of kidney failure due to diabetes	Yes				475
End-stage kidney failure: persons with diabetes (per 1,000,000)	Incidence rate of kidney failure due to diabetes among persons with diabetes	Yes				476
End-stage renal disease (per 1,000,000)	Incidence rate of end-stage renal disease (ESRD)	Yes				462
End-stage renal disease deaths: hemodialysis patients (per 1,000)	Total death rate for persons on hemodialysis	Yes				463

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Enforcement of illegal tobacco sales to minors laws: States (number)	Illegal tobacco sales rate of <=5% to minors through enforcement of laws prohibiting the sale of tobacco products to minors-number of States and the District of Columbia	Yes				1493
Enforcement of illegal tobacco sales to minors laws: Territories (number)	Illegal tobacco sales rate of <=5% to minors through enforcement of laws prohibiting the sale of tobacco products to minors-number of Territories	Yes				1494
Epidemiology services: Local public health agencies (percent)	Percent of Tribal, State, and local public health agencies that provide or assure comprehensive epidemiology services to support essential public health services-Local public health agencies	Yes				1363
Epidemiology services: State public health agencies (percent)	Percent of Tribal, State, and local public health agencies that provide or assure comprehensive epidemiology services to support essential public health services-State public health agencies	Yes				1362
Escherichia coli infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Reduction of Shiga toxin-producing Escherichia coli (STEC) O157	Yes				823
Estimated Poverty (percent)	Small area estimates of population living below the poverty line	Yes	Yes	Yes		10004
Excessive drinking: adults (percent)	Percent of adults who drank excessively in the previous 30 days	Yes				1458
Excessive drinking: adults (percent)	Percent of adults that report excessive drinking, either chronic high alcohol consumption or binge drinking		Yes	Yes		150
Extracurricular activities: teens aged 12-17 (percent)	Percent of adolescents who participate in extracurricular and out-of-school activities	Yes				404
Eye examinations, annual: adults with diabetes (percent)	Percent of adults with diabetes who have an annual dilated eye examination	Yes				551
Fair or poor health: adults (percent)	Percent of adults that report fair or poor health		Yes	Yes		5

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Fall deaths (per 100,000)	Fall-related death rate: All persons	Yes	Yes	Yes		1079
Fall deaths: older adults (per 100,000)	Fall-related death rate: Adults aged 65 years and over	Yes				1080
Fall ED visits: older adults (per 100,000)	Rate of emergency department visits due to falls among older adults	Yes				1259
Falling: adults with balance/dizziness problems (percent)	Percent of adults with balance and dizziness problems who have fallen in the past 5 years while experiencing symptoms of dizziness, vertigo, or imbalance	Yes				769
Family planning clinics offering all contraception methods (percent)	Percent of publicly funded family planning clinics that offer the full range of FDA-approved methods of contraception onsite	Yes				781
Family planning clinics offering emergency contraception (percent)	Percent of publicly funded family planning clinics that offer emergency contraception onsite	Yes				782
Family planning services income eligibility: states (number)	Number of States that set the income eligibility level for Medicaid-covered family planning services to at least the same level used to determine eligibility for Medicaid-covered, pregnancy-related care	Yes				818
Federally Qualified Health Center and Rural Health Clinic Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used Federally Qualified Health Center and Rural Health Clinic services	Yes	Yes		Yes	50002
Federally Qualified Health Center and Rural Health Clinic Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used Federally Qualified Health Center and Rural Health Clinic services	Yes	Yes		Yes	50003
Federally Qualified Health Center and Rural Health Clinic visits (per 1,000 beneficiaries)	Rate of Federally Qualified Health Center and Rural Health Clinic visits among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	50001
Federally Qualified Health Centers (number)	Number of Federally Qualified Health Centers (FQHC) participating in the Medicare program		Yes		Yes	10040
Female Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months who are female	Yes	Yes		Yes	309

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Fetal alcohol syndrome (per 10,000 live births)	Rate of fetal alcohol syndrome (FAS)	Yes				1146
Fetal deaths (per 1,000 live births plus fetal deaths)	Rate of fetal deaths at 20 or more weeks of gestation	Yes				1156
Few fruits/vegetables: adults (percent)	Adults that report fewer than 5 servings of fruits/vegetables per day		Yes	Yes		121
Fibrinolytic therapy timeliness: heart attack patients (percent)	Percent of eligible patients with heart attacks who receive fibrinolytic therapy within 30 minutes of hospital arrival	Yes				878
Firearm-related deaths (per 100,000)	Firearm-related death rate	Yes	Yes	Yes		1042
Flu vaccination: adults 18-64 years (percent)	Percent of noninstitutionalized adults aged 18 to 64 years vaccinated annually against seasonal influenza	Yes				1033
Flu vaccination: adults 65+ (percent)	Percent of adults age 65+ that report having Influenza vaccine in past 12 months		Yes	Yes		119
Flu vaccination: children 2-4 years (percent)	Percent of children aged 2 to 4 years vaccinated annually against seasonal influenza	Yes				1030
Flu vaccination: children 5-12 years (percent)	Percent of children aged 5 to 12 years vaccinated annually against seasonal influenza	Yes				1031
Flu vaccination: children 6-23 months (percent)	Percent of children aged 6 to 23 months vaccinated annually against seasonal influenza- (1 or 2 doses, depending on age-appropriateness and previous doses received)	Yes				1029
Flu vaccination: health care workers (percent)	Percent of health care personnel vaccinated annually against seasonal influenza	Yes				1037
Flu vaccination: high-risk adults (percent)	Percent of noninstitutionalized high risk adults aged 18 to 64 years vaccinated annually against seasonal influenza	Yes				1034
Flu vaccination: nursing home residents (percent)	Percent of institutionalized adults aged 18 years and older in long-term or nursing homes vaccinated annually against seasonal influenza	Yes				1036

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Flu vaccination: older adults (percent)	Percent of noninstitutionalized adults aged 65 years and older vaccinated annually against seasonal influenza	Yes				1035
Flu vaccination: pregnant women (percent)	Percent of pregnant women vaccinated annually against seasonal influenza	Yes				1038
Flu vaccination: teens 13-17 years (percent)	Percent of children aged 13 to 17 years vaccinated annually against seasonal influenza	Yes				1032
Fluoridation of community water (percent)	Percent of the U.S. population served by community water systems with optimally fluoridated water	Yes	Yes			1265
Fluoride programs: school-based health centers (percent)	Percent of school-based health centers with an oral health component that includes topical fluoride	Yes				1286
Folic acid intake: reproductive-aged women (percent)	Percent of women of childbearing potential with intake of at least 400 µg of folic acid from fortified foods or dietary supplements	Yes				1171
Follow-up to newborn screening (percent)	Percent of screen-positive children who receive follow-up testing within the recommended time period	Yes				1174
Food insecurity: households (percent)	Percent of households with low food security	Yes				1243
Food insecurity: households with children (percent)	Percent of household with children with very low food security	Yes				1228
Foot examinations, annual: adults with diabetes (percent)	Percent of adults with diabetes who have at least an annual foot examination	Yes	Yes			552
Formula supplementation: breastfed newborns (percent)	Percent of breastfed newborns who receive formula supplementation within the first 2 days of life	Yes				1185
Free school lunch eligibility (percent)	Percent of total students eligible to participate in the National School Lunch Program under the Free Lunch Act		Yes	Yes		10005
Fruit consumption (cup equivalents per 1,000 calories)	Mean daily intake of fruit (in cup equivalents per 1,000 calories) by persons aged 2 years and over	Yes				1217
Fruit/vegetable availability required: school districts (percent)	Percent of school districts that require schools to make fruits or vegetables available whenever other food is offered or sold.	Yes				10066

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Functional limitations: older adults (percent)	Percent of older adults 65 years and over who have moderate to severe functional limitations	Yes				1245
Genetic counseling: women with family history of breast/ovarian cancer (percent)	Percent of women with a family history of breast and/or ovarian cancer who receive genetic counseling	Yes				850
Genital herpes simplex type 2: adults 20-29 years (percent)	Percent of young adults with genital herpes infection due to herpes simplex type 2	Yes				1482
Geriatric dentists (percent)	Percent of the health care workforce with geriatric certification-Dentists	Yes				1253
Geriatric dieticians (percent)	Percent of the health care workforce with geriatric certification-Registered dieticians	Yes				1255
Geriatric nurses (percent)	Percent of the health care workforce with geriatric certification-Registered nurses	Yes				1252
Geriatric physical therapists (percent)	Percent of the health care workforce with geriatric certification-Physical therapists	Yes				1254
Geriatric physicians (percent)	Percent of the health care workforce with geriatric certification-Physicians	Yes				1250
Geriatric psychiatrists (percent)	Percent of the health care workforce with geriatric certification-Geriatric psychiatrists	Yes				1251
Giardiasis (number)	Number of reported cases of giardiasis	Yes	Yes	Yes		10021
Gini coefficient (ratio)	Gini coefficient of household income inequality		Yes	Yes		37
Global Disease Detection increased testing capacity (number)	Diagnostic testing capacity in host countries and regionally through the Global Disease Detection (GDD) Regional Centers	Yes				848
Global Disease Detection Regional Centers (number)	Number of Global Disease Detection (GDD) Regional Centers worldwide to detect and contain emerging health threats	Yes				846

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Global Disease Detection training of public health professionals (number)	Number of public health professionals trained by Global Disease Detection: Global Disease Detection (GDD) programs worldwide	Yes				847
Glucose A1c < 7%: adults with diabetes (percent)	Percent of adults with diagnosed diabetes with an HbA1c value less than 7 percent	Yes				559
Glucose A1c > 9%: adults with diabetes (percent)	Percent of adults with diagnosed diabetes with an HbA1c value greater than 9 percent	Yes				558
Glucose A1c test, biannual: adults with diabetes (percent)	Percent of adults with diabetes who have a glycosylated hemoglobin measurement at least twice a year	Yes	Yes			550
Glucose monitoring, daily: adults with diabetes (percent)	Percent of adults with diabetes who perform self-blood glucose-monitoring at least once daily	Yes	Yes			554
Gonorrhea rate (per 100,000)	Rate of gonorrhea cases per 100,000 population			Yes		10062
Gonorrhea: females 15-44 years (per 100,000)	Gonorrhea rate among females aged 15 to 44 years	Yes				1478
Gonorrhea: males 15-44 years (per 100,000)	Gonorrhea rate among males aged 15 to 44 years	Yes				1479
Graduated driver licensing laws: states (number)	Number of states and the District of Columbia with "good" graduated driver licensing (GDL) laws	Yes				1058
Group B streptococcal disease: newborns (per 1,000)	Early onset group B streptococcal disease rate	Yes				973
Hazardous material plan: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Have a plan for how to use, label, store, and dispose of hazardous materials	Yes				685
Hazardous sites posing risk (number)	Number of hazardous sites that pose risks to human health and the environment	Yes				677
HbA1c test: diabetic Medicare beneficiaries 65-75 years (percent)	Percent of diabetic Medicare enrollees ages 65-75 receiving a hemoglobin A1c test in the past year		Yes	Yes		29

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Health care evaluation: adults with balance/dizziness problems (percent)	Percent of adults with balance or dizziness problems in the past 12 months who have ever seen a Health Care provider about their balance or dizziness problems	Yes				762
Health care evaluation: adults with tinnitus (percent)	Percent of adults bothered by tinnitus in the past 12 months who have seen a doctor	Yes				759
Health care organizations monitoring adverse events (percent)	Percent of health care organizations that are monitoring and analyzing adverse events associated with medical therapies within their systems	Yes				1196
Health care provider explaining things (percent)	Percent of persons who report that their Health Care provider always explained things so they could understand them	Yes				854
Health care provider listening skills (percent)	Percent of persons who report that their Health Care provider always listened carefully to them	Yes				853
Health care provider referrals to hearing specialists (per 1,000)	Rate of persons who are referred by their primary care physician or other health care provider for hearing evaluation and treatment	Yes				751
Health care provider showing respect (percent)	Percent of persons who report that their Health Care provider always showed respect for what they had to say	Yes				855
Health care provider spending enough time with patient (percent)	Percent of persons who report that their health care provider always spent enough time with them	Yes				856
Health ed. certification requirements: elementary school (percent)	Percent of schools that require newly hired staff who teach required health instruction to be certified, licensed, or endorsed by the State in health education-Elementary schools	Yes				731
Health ed. certification requirements: high school (percent)	Percent of schools that require newly hired staff who teach required health instruction to be certified, licensed, or endorsed by the State in health education-High schools	Yes				733
Health ed. certification requirements: middle school (percent)	Percent of schools that require newly hired staff who teach required health instruction to be certified, licensed, or endorsed by the State in health education-Middle schools	Yes				732

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Health ed. goals, advocacy: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Advocating for personal, family, and community health (skills)	Yes				638
Health ed. goals, communication skills: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Using interpersonal communication skills to enhance health (skills)	Yes				642
Health ed. goals, concept comprehension: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Comprehending concepts related to health promotion and disease prevention (knowledge)	Yes				636
Health ed. goals, decision-making skills: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Using goal-setting and decision-making skills to enhance health (skills)	Yes				641
Health ed. goals, external influences: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Analyzing the influence of culture, media, technology, and other factors on health (skills)	Yes				639

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Health ed. goals, healthy behaviors: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)-Practicing health-enhancing behaviors and reducing health risks (skills)	Yes				640
Health ed. goals, information access: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)-Accessing valid information and health promoting products and services (skills)	Yes				637
Health ed. teachers trained in behavior change (percent)	Percent of required health education classes or courses with a teacher who has had professional development related to teaching personal and social skills for behavior change within the past 2 years	Yes				737
Health ed. training requirements: elementary schools (percent)	Percent of schools that require newly hired staff who teach required health education to have undergraduate or graduate training in health education-Elementary schools	Yes				728
Health ed. training requirements: high schools (percent)	Percent of schools that require newly hired staff who teach required health education to have undergraduate or graduate training in health education-High schools	Yes				730
Health ed. training requirements: middle schools (percent)	Percent of schools that require newly hired staff who teach required health education to have undergraduate or graduate training in health education-Middle schools	Yes				729
Health education, alcohol/drug use: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in alcohol and other drug use	Yes				571
Health education, all priority areas: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in All priority areas	Yes				566

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Health education, benefits of sleep: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Benefits of rest and sleep	Yes				656
Health education, growth & development: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Growth and development	Yes				654
Health education, hand washing info: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Hand washing or hand hygiene	Yes				652
Health education, health screening: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Importance of health screenings and checkups	Yes				658
Health education, injury prevention: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in unintentional injury	Yes				567
Health education, nutrition: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in unhealthy dietary patterns	Yes				573
Health education, oral health: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Dental and oral health	Yes				653
Health education, physical activity: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in inadequate physical activity	Yes				574
Health education, pregnancy, HIV/AIDS, STD: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in unintended pregnancy, HIV/AIDS, and STD infection	Yes				572

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Health education, suicide prevention: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in suicide	Yes				569
Health education, tobacco use: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in tobacco use and addiction	Yes				570
Health education, violence prevention: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in violence	Yes				568
Health education, vision & hearing loss prevention: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Ways to prevent vision and hearing loss	Yes				657
Health educations, sun safety: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Sun safety or skin cancer prevention	Yes				655
Health information, alcohol/drug: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on alcohol and other drug use	Yes				580
Health information, HIV & STD: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on HIV, AIDS and STD infection	Yes				582
Health information, injury: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on unintentional injury	Yes				576
Health information, nutrition: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on unhealthy dietary patterns	Yes				583
Health information, physical activity: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on inadequate physical activity	Yes				584

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Health information, suicide: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on suicide	Yes				578
Health information, tobacco use: post-secondary schools (percent)	percent of college and university students who receive information from their institution on tobacco use and addiction	Yes				579
Health information, unintended pregnancy: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on unintended pregnancy	Yes				581
Health information, violence: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on violence	Yes				577
Health instruction meets time standards: elementary schools (percent)	Percent of schools that require cumulative instruction in health education that meet the US National Health Education Standards for elementary, middle, and senior high schools-Elementary schools	Yes				734
Health instruction meets time standards: high schools (percent)	Percent of schools that require cumulative instruction in health education that meet the US National Health Education Standards for elementary, middle, and senior high schools-High schools	Yes				736
Health instruction meets time standards: middle schools (percent)	Percent of schools that require cumulative instruction in health education that meet the US National Health Education Standards for elementary, middle, and senior high schools-Middle schools	Yes				735
Health insurance: <65 years (percent)	Percent of persons under age 65 years with health (medical) insurance	Yes				368
Health Professions Shortage Area (HPSA) Indicator (yes/no)	Indicator for single county designated Health Professional Shortage Area			Yes		132
Health promotion programs for caregivers: States (number)	States and the District of Columbia with health promotion programs for caregivers	Yes				529
Health promotion programs for people with disabilities: States (number)	States and the District of Columbia with health promotion programs for people with disabilities	Yes				527

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Health provider seen: adults with joint-related symptoms (percent)	Percent of adults with chronic joint symptoms who have seen a health care provider for their symptoms	Yes				425
Health websites meeting 3+ criteria to assess reliability (percent)	Percent of health-related Websites that meet three or more evaluation criteria disclosing information that can be used to assess information reliability	Yes				863
Healthy food outlets (percent)	Percent of zip codes in county with a healthy food outlet		Yes	Yes		48
Healthy weight prior to pregnancy (percent)	Percent of women delivering a live birth who had a healthy weight prior to pregnancy	Yes				1182
Healthy weight: adults (percent)	Percent of adults who are at a healthy weight	Yes				1207
Hearing aid use: older adults (per 1,000)	Rate of adults aged 70 years and older with hearing loss who have ever used a hearing aid	Yes				746
Hearing loss, noise-induced, work-related (per 10,000 full-time workers)	Rate of work-related, noise-induced hearing loss	Yes				1302
Hearing loss, noise-induced: adults (per 1,000)	Rate of adults who have elevated hearing thresholds, or audiometric notches, in high frequencies (3, 4, or 6 kHz) in both ears, signifying noise-induced hearing loss	Yes				755
Hearing loss, noise-induced: teens (per 1,000)	Rate of adolescents who have elevated hearing thresholds, or audiometric notches, in high frequencies (3, 4, or 6 kHz) in both ears, signifying noise-induced hearing loss	Yes				754
Heart attack awareness: importance of calling 9-1-1 (percent)	Percent of adults aged 20 years and over who are aware of the importance of accessing rapid emergency care by calling 9-1-1 or another emergency number	Yes				896
Heart attack awareness: symptoms (percent)	Percent of adults aged 20 years and over who are aware of the early warning symptoms and signs of a heart attack	Yes				895
Heart attack awareness: symptoms and importance of calling 9-1-1 (percent)	Percent of adults aged 20 years and over who are aware of the early warning symptoms and signs of a heart attack and the importance of accessing rapid emergency care by calling 9-1-1 or another emergency number	Yes				894

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Heart attack Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have had a heart attack	Yes	Yes		Yes	283
Heart attack Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have had a heart attack	Yes	Yes		Yes	284
Heart attack patients given aspirin at hospital arrival (percent)	Hospital Compare: Heart attack patients given aspirin at hospital arrival	Yes	Yes		Yes	319
Heart attack patients prescribed ACE inhibitor or angiotensin receptor blocker at discharge (percent)	Hospital Compare: Heart attack patients prescribed angiotensin converting enzyme inhibitor or angiotensin receptor blocker at hospital discharge	Yes	Yes		Yes	321
Heart attack patients with aspirin prescribed at hospital discharge (percent)	Hospital Compare: Heart attack patients with aspirin prescribed at hospital discharge	Yes	Yes		Yes	320
Heart attack patients with beta blocker prescribed at hospital discharge (percent)	Hospital Compare: Heart attack patients with beta blocker prescribed at hospital discharge	Yes	Yes		Yes	323
Heart attack patients with fibrinolytic received within 30 minutes of hospital arrival (percent)	Hospital Compare: Heart attack patients with fibrinolytic received within 30 minutes of hospital arrival	Yes	Yes		Yes	324
Heart attack patients with percutaneous coronary intervention within 90 minutes of hospital arrival (percent)	Hospital Compare: Heart attack patients with percutaneous coronary intervention within 90 minutes of hospital arrival	Yes	Yes		Yes	325
Heart attack patients with smoking cessation counseling during hospital stay (percent)	Hospital Compare: Heart attack patients with smoking cessation counseling during hospital stay	Yes	Yes		Yes	322
Heart disease deaths (per 100,000)	Heart disease death rate	Yes	Yes	Yes		83
Heart failure hospitalizations: Medicare beneficiaries 65-74 years (per 1,000)	Rate of hospitalizations of older adults with heart failure as the principal diagnosis-Adults aged 65 to 74 years	Yes				891

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Heart failure hospitalizations: Medicare beneficiaries 75-84 years (per 1,000)	Rate of hospitalizations of older adults with heart failure as the principal diagnosis-Adults aged 75 to 84 years	Yes				892
Heart failure hospitalizations: Medicare beneficiaries 85+ years (per 1,000)	Rate of hospitalizations of older adults with heart failure as the principal diagnosis-Adults aged 85 years and over	Yes				893
Heart failure Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have heart failure	Yes	Yes		Yes	295
Heart failure Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have heart failure	Yes	Yes		Yes	296
Heart failure patients prescribed ACE inhibitor or angiotensin receptor blocker at discharge (percent)	Hospital Compare: Heart failure patients prescribed angiotensin converting enzyme inhibitor or angiotensin receptor blocker at hospital discharge	Yes	Yes		Yes	328
Heart failure patients with discharge instructions (percent)	Hospital Compare: Heart failure patients with discharge instructions	Yes	Yes		Yes	326
Heart failure patients with evaluation of left ventricular systolic function (percent)	Hospital Compare: Heart failure patients with evaluation of left ventricular systolic function	Yes	Yes		Yes	327
Heart failure patients with smoking cessation counseling (percent)	Hospital Compare: Heart failure patients with smoking cessation counseling	Yes	Yes		Yes	329
Hearing aid use: adults 20-69 years (per 1,000)	Rate of adults aged 20 to 69 years with hearing loss who have ever used a hearing aid	Yes				744
Hep B vaccination, birth dose (percent)	Percent of young children who have received universally recommended vaccines-A birth dose of hepatitis B vaccine (0 to 3 days between birth date and date of vaccination, reported by annual birth cohort)	Yes				1001
Hep B vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-3 doses hepatitis B (hep B) vaccine by age 19 to 35 months	Yes				995
Hepatitis A (per 100,000)	Hepatitis A rate	Yes				967

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Hepatitis A vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-2 doses hepatitis A vaccine by age 19 to 35 months	Yes				1000
Hepatitis A, acute (number)	Number of reported cases of acute Hepatitis A	Yes	Yes	Yes		10022
Hepatitis B vaccination: health care workers (percent)	Percent of Hepatitis B vaccine coverage among high-risk populations-Health care personnel	Yes				1016
Hepatitis B vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-Hepatitis B vaccine (3 or more doses)	Yes				1006
Hepatitis B, acute (number)	Number of reported cases of acute Hepatitis B	Yes	Yes	Yes		10023
Hepatitis B: adults (per 100,000)	Rate of new hepatitis B infections in adults aged 19 and older	Yes				988
Hepatitis B: children (per 100,000)	Rate of new hepatitis B cases among persons aged 2 to 18 years	Yes				980
Hepatitis B: injection drug users (number)	Number of new hepatitis B infections among high-risk populations: Injection drug users	Yes				989
Hepatitis B: men who have sex with men (number)	Number of new hepatitis B infections among high-risk populations: Men who have sex with men	Yes				991
Hepatitis C (per 100,000)	Rate of new hepatitis C infections	Yes				992
Hepatitis C infection status awareness (percent)	Percent of persons aware they have a hepatitis C infection	Yes				1022
Hib invasive disease: children <5 years (per 100,000)	Rate of serotype b cases of Haemophilus influenzae (Hib) invasive disease (children aged 5 years and under)	Yes				979
Hib vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-3 doses or more Haemophilus influenzae type b (Hib) vaccine by age 19 to 35 months	Yes				994
High blood cholesterol: adults with chronic kidney disease (percent)	Percent of adults with chronic kidney disease who have elevated lipid levels	Yes				481

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
High blood pressure: adults 18+ (pounds)	Percent of adults that report having been diagnosed with high blood pressure		Yes	Yes		123
High cholesterol: adults (percent)	Percent of adults with high total blood cholesterol levels	Yes				885
High school completion: adults 18-24 years (percent)	Percent of persons aged 18-24 that completes high school education	Yes				565
High school diploma receipt by students with disabilities (percent)	Percent of students who are served under the Individuals with Disabilities Education Act who graduate high school with a diploma	Yes				395
High school graduates, on-time with regular diploma (percent)	Percent of public school students who graduate with a regular diploma 4 years after starting 9th grade	Yes				394
High school on-time graduation (percent)	Percent of public high school students who graduate with a regular diploma 4 years after starting 9th grade		Yes	Yes		33
Hip fractures: female older adults (per 100,000)	Hip fracture hospitalizaion rate among females aged 65 years and over	Yes				429
Hip fractures: male older adults (per 100,000)	Hip fracture hospitalizaion rate among males aged 65 years and over	Yes				430
Hispanic Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months who are Hispanic	Yes	Yes		Yes	313
HIV deaths (per 100,000)	HIV infection mortality rate	Yes	Yes	Yes		935
HIV infection status awareness (percent)	Percent of HIV-infected persons who are aware of their infection	Yes				942
HIV prevalence (per 100,000)	Rate of adolescents and adults aged 13 years and over living with HIV		Yes	Yes		10018
HIV testing: persons 15-24 years (percent)	Percent of adolescents and young adults who have been tested for HIV in the past 12 months	Yes				947
HIV testing: persons 15-44 years (percent)	Percent of adolescents and adults who have been tested for HIV in the past 12 months	Yes				944

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
HIV testing: pregnant women (percent)	Percent of pregnant women who have been tested for HIV during their pregnancy	Yes				946
HIV testing: TB patients (percent)	Percent of adults with tuberculosis (TB) who have been tested for HIV	Yes				934
HIV transmission rate: persons living with HIV (per 100)	Rate of HIV transmission per 100 persons with HIV among adolescents and adults	Yes				959
HIV/AIDS education: substance abuse treatment programs (percent)	Percent of substance abuse treatment facilities that offer HIV/AIDS education, counseling, and support	Yes				933
HIV/AIDS prevention discussion with parent before age 18: females (percent)	Percent of adolescents females who talked to a parent or guardian about HIV/AIDS prevention before they were 18 years old	Yes				814
HIV/AIDS prevention discussion with parent before age 18: males (percent)	Percent of adolescents males who talked to a parent or guardian about HIV/AIDS prevention before they were 18 years old	Yes				815
HIV/AIDS prevention instruction receipt before age 18: females (percent)	Percent of adolescents females who received formal instruction on HIV/AIDS prevention before they were 18 years old	Yes				806
HIV/AIDS prevention instruction receipt before age 18: males (percent)	Percent of adolescents males who received formal instruction on HIV/AIDS prevention before they were 18 years old	Yes				807
Home Health Agencies (number)	Number of Home Health Agencies participating in the Medicare program		Yes		Yes	10041
Home health Medicare episodes (per 1,000 beneficiaries)	Average number of home health episodes among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2018
Home health Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used home health services	Yes	Yes		Yes	221
Home health Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used home health services	Yes	Yes		Yes	222

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Home health Medicare visits (per 1,000 beneficiaries)	Average number of home health visits among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2017
Homes with dust-lead hazards (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US homes that have dust-lead hazards	Yes				723
Homes with lead paint hazards (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US homes that have paint-lead hazards	Yes				722
Homes with lead-based paint (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US homes that are found to have lead-based paint	Yes				721
Homes with soil-lead hazards (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US homes that have soil-lead hazards	Yes				724
Homicide deaths (per 100,000)	Homicide mortality rate	Yes	Yes	Yes		1041
Homicides, work-related (per 10,000)	Number of deaths due to work-related homicides	Yes				1299
Hospice Facilities (number)	Number of Hospice Facilities participating in the Medicare program		Yes		Yes	10042
Hospice Medicare admissions (per 1,000 beneficiaries)	Rate of hospice admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2007
Hospice Medicare days (per 1,000 beneficiaries)	Average number of hospice days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2015
Hospice Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who had covered hospice services	Yes	Yes		Yes	227
Hospice Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered hospice services	Yes	Yes		Yes	228
Hospital 30-day death (mortality) for heart attack patients (percent)	Hospital Compare: Hospital 30-day death (mortality) rates for heart attack patients	Yes	Yes		Yes	343

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Hospital 30-day death (mortality) for heart failure patients (percent)	Hospital Compare: Hospital 30-day death (mortality) rates for heart failure patients	Yes	Yes		Yes	344
Hospital 30-day death (mortality) for pneumonia patients (percent)	Hospital Compare: Hospital 30-day death (mortality) rates for pneumonia patients	Yes	Yes		Yes	345
Hospital 30-day readmission for heart attack patients (percent)	Hospital Compare: Hospital 30-day readmission rates for heart attack patients	Yes	Yes		Yes	346
Hospital 30-day readmission for heart failure patients (percent)	Hospital Compare: Hospital 30-day readmission rates for heart failure patients	Yes	Yes		Yes	347
Hospital 30-day readmission for pneumonia patients (percent)	Hospital Compare: Hospital 30-day readmission rates for pneumonia patients	Yes	Yes		Yes	348
Hospital inpatient Medicare admissions (per 1,000 beneficiaries)	Rate of hospital inpatient admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2001
Hospital inpatient Medicare days (per 1,000 beneficiaries)	Average number of hospital inpatient (IP) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2009
Hospital inpatient Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who had covered hospital inpatient services	Yes	Yes		Yes	160
Hospital inpatient Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered hospital inpatient services	Yes	Yes		Yes	161
Hospital outpatient Medicare visits (per 1,000 beneficiaries)	Rate of hospital outpatient (OP) visits among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2019
Housing units with physical problems (percent)	Percent of occupied housing units that have moderate or severe physical problems	Yes				663
HPV vaccination: girls 13-15 years (percent)	Percent of female adolescents who have 3 doses Human papillomavirus vaccine (HPV) by age 13 to 15 years	Yes				1013
Hypertension admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Hypertension admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	2040
Hypertension Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have hypertension	Yes	Yes		Yes	50009

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Hypertension Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have hypertension	Yes	Yes		Yes	50010
Hypertension medication compliance (percent)	Percent of adults with hypertension who are taking the prescribed medications to lower their blood pressure	Yes				912
Hypertension: adults (percent)	Percent of adults with hypertension	Yes				897
Hypertension: adults with chronic kidney disease (percent)	Percent of adults with chronic kidney disease who have elevated blood pressure	Yes				480
Hypertension: children 8-17 years (percent)	Percent of children and adolescents with hypertension	Yes				898
Identification of "people with disabilities" in data systems (number)	Percent of major applicable Healthy People 2020 data systems with a standardized set of questions that identify "people with disabilities"	Yes				513
Illegal drugs on school property (percent)	Percent of adolescents who have been offered, sold, or given an illegal drug on school property	Yes	Yes			411
Illicit drug abstinence, prenatal (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Illicit drugs	Yes				1145
Illicit drug treatment receipt (percent)	Percent of persons who need illicit drug treatment and received specialty treatment for abuse or dependence in the past year-Illicit drug treatment	Yes				1454
Illicit drugs, never use: high school seniors (percent)	Percent of high school seniors never using substances-Illicit drugs	Yes				1431
Illicit drugs, recent use: adults (percent)	Percent of adults reporting use of any illicit drug during the past 30 days	Yes				1434
Illicit drugs, recent use: teens (percent)	Percent of adolescents reporting use of alcohol or any illicit drugs during the past 30 days	Yes				1432
Imaging Medicare service events (per 1,000 beneficiaries)	Rate of imaging service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2023
Imaging Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used imaging services	Yes	Yes		Yes	251

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Imaging Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used imaging services	Yes	Yes		Yes	252
Immunization information system participation: States (number)	Number of States that have 80 percent of adolescents with two or more age-appropriate immunizations recorded in immunization information systems (adolescents aged 11 to 18 years)	Yes				1025
Impaired fecundity: women 18-44 years (percent)	Percent of women aged 18 to 44 years who have impaired fecundity	Yes				1193
Improvement plan development: health emergencies/exercises (number)	Time reported for State public health agencies to establish improvement plans following responses to public health emergencies and exercises in minutes	Yes				1397
Inability to obtain or delay in obtaining dental care (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary dental care	Yes				381
Inability to obtain or delay in obtaining medical care (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary medical care	Yes				380
Inability to obtain or delay in obtaining prescription meds (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary prescription medicines	Yes				382
Inability to obtain or delayed access to medical/dental care or prescriptions (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary medical care, dental care, or prescription medicines	Yes				379
Incentive policies for food retail: states (number)	Number of states and the District of Columbia that have state-level policies that incentivize food retail outlets to provide foods that are encouraged by the Dietary Guidelines	Yes				1239
Inclusion in regular education: students with disabilities (percent)	Percent of children and youth with disabilities in classroom settings who spend at least 80 percent of their time in regular education programs	Yes				517
Increasing physical activity: adults with pre-diabetes (percent)	Percent of persons at high risk for diabetes (with pre-diabetes) who report increasing their levels of physical activity	Yes				562

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Indoor allergen levels, cockroach (units of allergen/grams of settled dust)	Indoor allergen levels: cockroach	Yes				679
Indoor allergen levels, mouse (micrograms of allergen/grams of settled dust)	Indoor allergen levels: mouse	Yes				680
Infant deaths between 28 days-1 year (per 1,000 live births)	Rate of postneonatal deaths (between 28 days and 1 year)	Yes	Yes	Yes		1160
Infant deaths due to birth defects (per 1,000 live births)	Rate of infant deaths related to birth defects (all birth defects)	Yes				1161
Infant deaths due to congenital heart defects (per 1,000 live births)	Rate of infant deaths related to birth defects (congenital heart defects)	Yes				1162
Infant deaths from sudden infant death syndrome (per 1,000 live births)	Rate of infant deaths from sudden infant death syndrome (SIDS)	Yes				1163
Infant deaths from sudden unexpected/unexplained causes (per 1,000 live births)	Rate of infant deaths from sudden unexpected/unexplained causes (includes SIDS, Unknown Cause, Accidental Suffocation, and Strangulation in Bed)	Yes				1164
Infant deaths within first 28 days of life (per 1,000 live births)	Rate of neonatal deaths (within the first 28 days of life)	Yes	Yes	Yes		1159
Infant deaths, all (per 1,000 live births)	Rate of all infant deaths (within 1 year)	Yes	Yes	Yes		1158
Infants put to sleep on their backs (percent)	Percent of infants who are put down to sleep on their backs	Yes				1141
Influenza-virus resistance to antivirals monitoring: states (number)	Number of public health laboratories monitoring influenza-virus resistance to antiviral agents	Yes				1028
Information systems used for environmental health: states (number)	Quality, utility, awareness, and use of existing information systems for environmental health among states	Yes				664
Inhalant use: teens (percent)	Percent of adolescents who use inhalants	Yes				1443
Injection drug use admissions (number)	Number of admissions to substance abuse treatment for injection drug use	Yes				1453

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Injuries due to falls: adults with balance/dizziness problems (percent)	Percent of adults with balance and dizziness problems who have been injured as a result of a fall for any reason in the past 12 months	Yes				770
Injuries, non-fatal, work-related (per 100)	Rate of injuries in private sector industries resulting in medical treatment, lost time from work, or restricted work activity, as reported by employers	Yes				1308
Injuries, nonfatal unintentional (per 100,000)	Nonfatal unintentional injury rate	Yes				1075
Injuries, sports and recreation (per 1,000)	Sports and recreation injury rate	Yes				1098
Injuries, treated in ED, non-fatal, work-related (per 100)	Rate of injuries treated in emergency departments from work, or restricted work activity, as reported by employers	Yes				1309
Injuries, treated in ED, non-fatal, work-related: teens (per 100)	Rate of nonfatal work-related injuries among dolescent workers aged 15 to 19 years	Yes				1310
Injury deaths (per 100,000)	Injury death rate	Yes	Yes	Yes		1071
Injury deaths, unintentional (per 100,000)	Unintentional injury death rate	Yes	Yes	Yes		1074
Injury deaths, work-related, all industries (per 100,000)	Rate of work-related injury deaths - All industry	Yes				1303
Injury deaths, work-related, construction (per 100,000)	Rate of work-related injury deaths - Construction	Yes				1305
Injury deaths, work-related, farming/forestry/fishing/hunting (per 100,000)	Rate of work-related injury deaths - Agriculture, forestry, fishing, and hunting	Yes				1307
Injury deaths, work-related, mining (per 100,000)	Rate of work-related injury deaths - Mining	Yes				1304
Injury deaths, work-related, transportation (per 100,000)	Rate of work-related injury deaths - Transportation	Yes				1306
Injury ED visits (per 100,000)	Rate of emergency department visits for nonfatal injuries	Yes				1073

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Injury hospitalizations (per 100,000)	Hospitalization rate for nonfatal injuries	Yes				1072
Injury prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in injury	Yes				596
Inpatient prospective payment system Medicare admissions (per 1,000 beneficiaries)	Rate of acute inpatient prospective payment system (PPS) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2002
Inpatient prospective payment system Medicare days (per 1,000 beneficiaries)	Average number of acute inpatient prospective payment system (PPS) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2010
Inpatient rehab Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who had covered inpatient rehabilitation services	Yes	Yes		Yes	207
Inpatient rehab Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered inpatient rehabilitation facility services	Yes	Yes		Yes	208
Inpatient rehabilitation facility Medicare admissions (per 1,000 beneficiaries)	Rate of inpatient rehabilitation facility (IRF) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2005
Inpatient rehabilitation facility Medicare days (per 1,000 beneficiaries)	Average number of inpatient rehabilitation facility (IRF) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2013
Inspection of drinking water for bacteria: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Among schools with community water systems, test drinking water for bacteria	Yes				690

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Inspection of drinking water for coliforms: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Among schools with community water systems, test drinking water for coliforms	Yes				691
Inspection of drinking water for lead: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Test drinking water outlets for lead	Yes				689
Intended pregnancies (percent)	Percent of pregnancies that are intended	Yes				779
Internet access (percent)	Percent of persons with access to the Internet	Yes				857
Internet use to communicate with providers (percent)	Percent of persons who use the Internet to communicate with their provider	Yes				868
Internet use to track personal health information (percent)	Percent of persons who use the Internet to keep track of personal health information, such as care received, test results, or upcoming medical appointments	Yes				867
Invasive colorectal cancer (per 100,000)	Invasive colorectal cancer incidence	Yes				509
Invasive pneumococcal infections: children <5 years (per 100,000)	Rate of new invasive pneumococcal infections among children under age 5 years	Yes				963
Invasive pneumococcal infections: older adults (per 100,000)	Rate of new invasive pneumococcal infections among adults aged 65 years and older	Yes				964
Invasive uterine cervical cancer: females (per 100,000)	Invasive uterine cervical cancer incidence	Yes				510
Involvement in health care decision making (percent)	Percent of persons who report that their health care providers always involved them in decisions about their health care as much as they wanted	Yes				866
Iron deficiency: children 1-2 years (percent)	Percent of young children aged 1 to 2 years who are iron deficient	Yes				1209

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Iron deficiency: children 3-4 years (percent)	Percent of young children aged 3 to 4 years who are iron deficient	Yes				1210
Iron deficiency: females of childbearing age (percent)	Percent of females of childbearing age, aged 12 to 49 years, who are iron deficient	Yes				1211
Iron deficiency: pregnant females (percent)	Percent of pregnant females who are iron deficient	Yes				1212
Ischemic heart disease Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have ischemic heart disease	Yes	Yes		Yes	297
Ischemic heart disease Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have ischemic heart disease	Yes	Yes		Yes	298
Joint difficulty grasping: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-related activities- Use fingers to grasp or handle small objects	Yes				434
Joint difficulty stooping/bending/ kneeling: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-related activities- Stoop, bend, or kneel	Yes				433
Joint difficulty walking 3 blocks: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-related activities- Walk a quarter of a mile-about 3 city blocks	Yes				431
Joint difficulty walking up 10 steps: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-related activities- Walk up 10 steps without resting	Yes				432
Joint mobility: persons with hemophilia (percent)	Percent of persons with hemophilia who develop reduced joint mobility due to bleeding into joints	Yes				444
Joint pain: adults with arthritis (mean)	Mean level of joint pain among adults with doctor-diagnosed arthritis	Yes				418
Kidney transplantation within 3 years: hemodialysis patients (percent)	Percent of patients receiving a kidney transplant within 3 years of end-stage renal disease (ESRD)	Yes				473

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Kidney waitlisting/transplantation: hemodialysis patients (percent)	percent of hemodialysis patients wait-listed and/or receiving a deceased donor kidney transplant within 1 year of end-stage renal disease (ESRD) start (among patients under 70 years of age)	Yes				472
Kindergarten vaccination record keeping: states (number)	Number of States collecting kindergarten vaccination coverage data according to CDC minimum standards	Yes				1026
Laboratories support data management: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Integrated data management	Yes				1350
Laboratories support disease prevention/control: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Disease prevention, control, and surveillance	Yes				1349
Laboratories support emergency response: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Emergency response	Yes				1356
Laboratories support environmental health: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Environmental health and protection	Yes				1352
Laboratories support food safety: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Food safety	Yes				1353
Laboratories support health research: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Public health-related research	Yes				1357

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Laboratories support lab improvement: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Laboratory improvement and regulation	Yes				1354
Laboratories support partnerships: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Partnerships and communication	Yes				1359
Laboratories support policy development: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Policy development	Yes				1355
Laboratories support reference & specialized testing: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Reference and specialized testing	Yes				1351
Laboratories support training/education: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Training and education	Yes				1358
Laboratory proficiency, chemical threat agents (percent)	Percent of LRN chemical laboratories that meet proficiency standards for chemical threat agents	Yes				1396
Laboratory test Medicare service events (per 1,000 beneficiaries)	Rate of laboratory test service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2024
Laboratory test Medicare users (count)	Count of Medicare fee-for-service beneficiaries 65+ years of age who used laboratory test services	Yes	Yes		Yes	2027
Laboratory test Medicare utilization (percent)	Percent of Medicare fee-for-service beneficiaries 65+ years of age who used laboratory test services	Yes	Yes		Yes	2028
Lack of leisure-time physical activity: adults (percent)	Percent of adults who engage in no leisure-time physical activity	Yes				1313

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Lactation care in birthing facilities (percent)	Percent of live births that occur in facilities that provide recommended care for lactating mothers and their babies	Yes				1186
Late-stage breast cancer: females (per 100,000)	Late-stage female breast cancer incidence	Yes				511
Lead level (µg/dL)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: lead	Yes				697
Lead level elevated: children 1-5 years (percent)	Percent of children with elevated blood lead levels	Yes				675
Lead levels: children 1-5 years (mean)	Mean blood lead levels in children	Yes				676
Lead poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: Lead poisoning	Yes				713
Lead, elevated level, work-related (per 100,000)	Rate of elevated blood lead concentrations from work exposures among adults	Yes				1300
Leisure-time physical activity: older adults (percent)	Percent of older adults with reduced physical or cognitive function who engage in light, moderate, or vigorous leisure-time physical activities	Yes				1258
Limitation of activity: adults with balance/dizziness problems (percent)	Percent of adults with balance and dizziness problems in the past 12 months who were prevented from doing regular activities within the home or outside	Yes				767
Lipids under control: adults with diabetes (percent)	Lipid control among persons with diagnosed diabetes	Yes				561
Liquor stores	Liquor store density rate		Yes	Yes		51
Live births (number)	Number of live births	Yes	Yes	Yes		10009
Long term care hospital Medicare admissions (per 1,000 beneficiaries)	Rate of long term care (LTC) hospital admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2006

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Long term care hospital Medicare days (per 1,000 beneficiaries)	Average number of long term care (LTC) hospital days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2014
Long term care hospital Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who had covered long-term care hospital services	Yes	Yes		Yes	200
Long term care hospital Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered long-term care hospital services	Yes	Yes		Yes	201
Long-Term Care Hospitals (number)	Number of long-term care hospitals participating in the Medicare program		Yes		Yes	10043
Low birth weight infant (percent)	Percent of live births that are low birth weight (LBW)	Yes	Yes	Yes		1135
Low birth weight: children with cerebral palsy (percent)	Percent of children with cerebral palsy born as low birth weight infants (less than 2,500 grams)	Yes				1168
Lower extremity amputations: persons with diabetes (per 1,000)	Rate of lower extremity amputations in persons with diagnosed diabetes	Yes				548
Lung cancer deaths (per 100,000)	Lung cancer death rate	Yes				487
Lung cancer Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have lung cancer	Yes	Yes		Yes	303
Lung cancer Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have lung cancer	Yes	Yes		Yes	304
Lung, trachea, and bronchus cancer deaths (per 100,000)	Lung, trachea, and bronchus cancer death rate per 100,000	Yes	Yes	Yes		88
Lyme disease (number)	Number of reported cases of Lyme disease	Yes	Yes	Yes		10024
M.D. training, counseling for health promotion (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-MD-granting medical schools	Yes				606
M.D. training, cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-MD-granting medical schools	Yes				607

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
M.D. training, environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-MD-granting medical schools	Yes				609
M.D. training, evaluation of health sciences literature. (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-MD-granting medical schools	Yes				608
M.D. training, global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-MD-granting medical schools	Yes				611
M.D. training, public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-MD-granting medical schools	Yes				610
Major depressive episode: adults (percent)	Percent of adults aged 18 years and older with Major Depressive Episode	Yes				1122
Major depressive episode: teens (percent)	Percent of adolescents aged 12 to 17 years with Major Depressive Episode	Yes				1123
Malaria (number)	Number of cases of malaria reported in the United States	Yes				844
Male Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months who are male	Yes	Yes		Yes	310
Mammogram: women 50+ (percent)	Percent of women age 50+ that report a mammogram in the past 2 years		Yes	Yes		116
Mammography counseling: women 50-74 years (percent)	Percent of women who were counseled by their providers about mammograms	Yes				494
Mammography: women 50-74 years (percent)	Percent of women who receive a breast cancer screening based on the most recent guidelines	Yes				507
Mandatory ignition interlock laws: States (number)	Number of States with mandatory ignition interlock laws for first and repeat impaired driving offenders in the United States	Yes				1468

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Marijuana, never use: at risk teens (percent)	Percent of at risk adolescents aged 12 to 17 years who, in the past year, refrained from using marijuana for the first time	Yes				1429
Marijuana, recent use: teens (percent)	Percent of adolescents reporting use of marijuana during the past 30 days	Yes				1433
Mass transit for transportation (percent)	Percent of trips to work using of alternative modes of transportation - mass transit	Yes				668
Maternal deaths (per 100,000 live births)	Rate of maternal death	Yes				1129
Maternal illness and complications due to pregnancy (percent)	Maternal illness and complications due to pregnancy (complications during hospitalized labor and delivery)	Yes				1130
Measles, total (number)	Number of reported cases of measles	Yes	Yes	Yes		10032
Measles, U.S.-acquired (number)	Number of measles cases (U.S.-acquired cases)	Yes				981
Median Household Income (dollars)	Small area estimates of median household Income	Yes	Yes	Yes		10003
Medicaid beneficiaries (number in thousands)	Number of Medicaid beneficiaries	Yes	Yes			10014
Medicaid coverage for nicotine treatment: States (number)	Comprehensive Medicaid insurance coverage of evidence-based treatment for nicotine dependency in States and the District of Columbia	Yes				1516
Medicaid managed care beneficiaries (percent)	Percent of Medicaid beneficiaries in managed care plans	Yes	Yes			10015
Medicaid payments per beneficiary (dollars)	Average payment per Medicaid beneficiary	Yes	Yes			10017
Medical evaluation: older adults with chronic kidney disease & diabetes (percent)	Percent of persons 65 years of age and over with type 1 or type 2 diabetes and chronic kidney disease who receive medical evaluation with serum creatinine, microalbuminuria, HbA1c, lipids, and eye examinations	Yes				478

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Medical evaluation: older adults with chronic kidney disease (percent)	Percent of persons 65 years of age and over with chronic kidney disease who receive medical evaluation with serum creatinine, lipids, and microalbuminuria	Yes				477
Medical home: children (percent)	Percent of children who have access to a medical home	Yes				1152
Medical home: children with special needs (percent)	Percent of children with special health care needs who have access to a medical home	Yes				1153
Medical treatment: older adults with chronic kidney disease & diabetes (percent)	Percent of persons 65 years of age and over with Part D insurance with diabetes and chronic kidney disease who receive recommended medical treatment with angiotensin-converting enzyme (ACE) inhibitors or angiotensin II receptor blockers (ARBS)	Yes				479
Medicare beneficiaries 65+ (count)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	307
Medicare beneficiaries 65+ who are also eligible for Medicaid (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months who are also eligible for Medicaid	Yes	Yes		Yes	317
Medicare managed care beneficiaries (percent)	Percent of Medicare beneficiaries enrolled in Medicare Advantage managed care plans	Yes	Yes			10020
Melanoma cancer deaths (per 100,000)	Melanoma cancer death rate	Yes				493
Meningococcal disease (per 100,000)	Meningococcal disease rate	Yes				968
Meningococcal vaccination: teens 13-15 years (percent)	Percent of adolescents who have 1 dose Meningococcal vaccine (MCV) by age 13 to 15 years	Yes				1012
Mental health screening: juvenile justice facilities (percent)	Percent of juvenile residential facilities that screen admissions for mental health problems	Yes				1111
Mental health services receipt: homeless adults (percent)	Percent of homeless adults with mental health problems who receive mental health services	Yes				1107
Mental health treatment provision: primary care facilities (percent)	Percent of primary care facilities that provide mental health treatment onsite or by paid referral	Yes				1109

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Mental illness prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in mental illness	Yes				598
Mentally unhealthy days: adults (per person)	Average number of reported mentally unhealthy days per month among adults 18 years and over		Yes	Yes		9
Mercury level: children 1-5 years (µg/L)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Mercury, children aged 1 to 5 years	Yes				698
Mercury level: females 16-49 years (µg/L)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Mercury, females aged 16 to 49 years	Yes				699
Mercury poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: mercury poisoning	Yes				715
Methicillin-resistant Staphylococcus aureus infections (per 100,000)	Invasive healthcare-associated methicillin-resistant Staphylococcus aureus (MRSA) infections	Yes				852
Missed school days: children 5-17 with asthma (percent)	Percent of children aged 5 to 17 years with asthma who miss school days	Yes				1410
Missed work days: adults 18-64 with asthma (percent)	Percent of adults aged 18 to 64 years with asthma who miss work days	Yes				1411
MMR vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-1 dose measles-mumps-rubella (MMR) vaccine by age 19 to 35 months	Yes				996
MMR vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-MMR vaccines (2 or more doses)	Yes				1004

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Mold problem policy: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Have a plan for how to address mold problems	Yes				684
Motor vehicle crash deaths (per 100,000,000 vehicle miles traveled)	Motor vehicle crash-related deaths per 100 million vehicle miles traveled	Yes				1077
Motor vehicle crashes involving drowsy driving (per 100,000,000)	Motor vehicle crashes per 100 million miles traveled involving drowsy driving	Yes				1470
Motor vehicle deaths (per 100,000)	Unintentional motor vehicle related death rate	Yes	Yes	Yes		18
Motor vehicle injuries, nonfatal (per 100,000)	Nonfatal motor vehicle crash-related injuries	Yes				1078
Motor vehicle traffic deaths (per 100,000)	Motor vehicle traffic-related death rate per 100,000	Yes	Yes	Yes		1076
Motorcycle helmet use (percent)	Percent of motorcycle operators and passengers using helmets	Yes				1048
Multivitamins/folic acid use, preconception (percent)	Percent of women delivering a live birth who took multivitamins/folic acid prior to pregnancy	Yes				1179
Mumps (number)	Number of reported cases of mumps	Yes	Yes	Yes		10031
Mumps, U.S.-acquired (number)	Number of mumps cases (U.S.-acquired cases)	Yes				982
Muscle-strengthening activity meeting guidelines: adults (percent)	Percent of adults who perform muscle-strengthening activities on 2 or more days of the week	Yes				1321
Neonatal hearing screening (percent)	Percent of infants that receive screening for hearing loss no later than age 1 month	Yes				756
Nephrologist seen at least 12 months before hemodialysis initiation (percent)	Percent of chronic kidney disease patients receiving care from a nephrologist at least 12 months before the start of renal replacement therapy	Yes				468
Never had sexual intercourse: females <16 years (percent)	Percent of female adolescents aged 15 years and under who have never had sexual intercourse	Yes				792

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Never had sexual intercourse: females 15-17 years (percent)	Percent of female adolescents aged 15 to 17 years who have never had sexual intercourse	Yes				790
Never had sexual intercourse: males <16 years (percent)	Percent of male adolescents aged 15 years and under who have never had sexual intercourse	Yes				793
Never had sexual intercourse: males 15-17 years (percent)	Percent of male adolescents aged 15 to 17 years who have never had sexual intercourse	Yes				791
Newborn screening verification: states (number)	Number of States and the District of Columbia that verify through linkage with vital records that all newborns are screened shortly after birth for conditions mandated by their State-sponsored screening program	Yes				1173
Nitrophenol level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Paranitrophenol (methyl parathion and parathions)	Yes				703
No exercise: adults (percent)	Adults that report no leisure-time exercise in past month		Yes	Yes		120
Non-FDA-approved pain medications (number)	Number of non-FDA-approved pain medications	Yes				1198
Non-Hispanic White Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months who are Non-Hispanic white	Yes	Yes		Yes	311
Nonfatal firearm-related injuries (per 100,000)	Nonfatal firearm-related injury rate	Yes				1043
Number of days the Air Quality Index (AQI) exceeds 100 (number)	Reduce the number of days the Air Quality Index (AQI) exceeds 100, weighted by population and AQI value	Yes				665
Nurse practitioner training, cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-Nurse practitioner	Yes				625

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Nurse practitioner training, environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-Nurse practitioner	Yes				627
Nurse practitioner training, evaluation of health literature (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-Nurse practitioner	Yes				626
Nurse practitioner training, global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-Nurse practitioner	Yes				629
Nurse practitioner training, health promotion counseling (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-Nurse practitioner	Yes				624
Nurse practitioner training, public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-Nurse practitioner	Yes				628
Nursing home (Title 18/19) facilities (number)	Number of Medicare or Medicaid-certified Title 18/19 (nursing home) facilities		Yes		Yes	10053
Nursing training, counseling for health promotion (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-Undergraduate nursing	Yes				618
Nursing training, cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-Undergraduate nursing	Yes				619
Nursing training, environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-Undergraduate nursing	Yes				621

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Nursing training, evaluation of health sciences literature (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-Undergraduate nursing	Yes				620
Nursing training, global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-Undergraduate nursing	Yes				623
Nursing training, public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-Undergraduate nursing	Yes				622
Nutrition counseling at physician visits (percent)	Percent of physician visits made by all child or adult patients that include counseling about nutrition or diet	Yes				1227
Nutrition counseling: patients with CVD, diabetes, or hyperlipidemia (percent)	Percent of physician office visits made by patients with a diagnosis of cardiovascular disease, diabetes, or hyperlipidemia that include counseling or education related to diet and nutrition	Yes				1225
Nutrition services: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in nutrition	Yes				603
Nutrition standards for child care: states (number)	Number of states and the District of Columbia with nutrition standards for foods and beverages provided to preschool-aged children in child care	Yes				1240
Obesity: adult (percent)	Percent of adults 18 years and over that report BMI \geq 30		Yes	Yes		15
Obesity: adults (percent)	Percent of adults who are obese	Yes				1208
Obesity: children 2-19 years (percent)	Percent of children and adolescents aged 2 to 19 years who are considered obese	Yes				1216
Obesity: children 2-5 years (percent)	Percent of children and adolescents aged 2 to 5 years who are considered obese	Yes				1213

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Obesity: children 6-11 years (percent)	Percent of children and adolescents aged 6 to 11 years who are considered obese	Yes				1214
Obesity: teens 12-19 years (percent)	Percent of children and adolescents aged 12 to 19 years who are considered obese	Yes				1215
Obstructive sleep apnea evaluation: adults (percent)	Percent of adults with symptoms of obstructive sleep apnea who seek medical evaluation	Yes				1469
Occupational eye injuries resulting in lost work days (per 10,000)	Rate of occupational eye injuries resulting in lost work days	Yes				1564
Occupational eye injuries treated in emergency departments (per 10,000)	Rate of occupational eye injuries treated in emergency departments	Yes				1565
Online health information access (percent)	Percent of online health information seekers who report easily accessing health information	Yes				865
Oral and craniofacial health surveillance: states (number)	Number of States and the District of Columbia that have an oral and craniofacial health surveillance system	Yes				1268
Oral and pharyngeal cancer early detection (percent)	Percent of oral and pharyngeal cancers detected at the earliest stage	Yes				1264
Oral health care system use (percent)	Percent of children, adolescents, and adults who used the oral health care system in the past year	Yes				1266
Oral health program: Federally qualified health centers (percent)	Percent of Federally Qualified Health Centers that have an oral health care program	Yes				1287
Oral health program: Local health departments (percent)	Percent of local health departments that have oral health prevention or care programs	Yes				1288
Oral health services receipt at health centers (percent)	Percent of patients who receive oral health services at Federally Qualified Health Centers each year	Yes				1296
Organ Procurement Organizations (number)	Number of Organ Procurement Organizations participating in the Medicare program		Yes		Yes	10044
Oropharyngeal cancer deaths (per 100,000)	Oropharyngeal cancer death rate	Yes				491

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Osteoporosis: adults 50+ years (percent)	Percent of adults aged 50 years and over with osteoporosis	Yes				427
Other hospital inpatient Medicare admissions (per 1,000 beneficiaries)	Rate of other hospital inpatient (IP) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2004
Other hospital inpatient Medicare days (per 1,000 beneficiaries)	Average number of other hospital inpatient (IP) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2012
Other hospital inpatient Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who had covered other hospital inpatient services	Yes	Yes		Yes	181
Other hospital inpatient Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered other hospital inpatient services	Yes	Yes		Yes	182
Other or Unknown race Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months who are Other or Unknown race	Yes	Yes		Yes	316
Other Part B Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used other Part B	Yes	Yes		Yes	275
Other Part B Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used other Part B	Yes	Yes		Yes	276
Other tests (Non-laboratory) Medicare service events (per 1,000 beneficiaries)	Rate of non-laboratory test service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2025
Other tests (non-laboratory) Medicare users (count)	Count of Medicare fee-for-service beneficiaries 65+ years of age who used non-laboratory test services	Yes	Yes		Yes	2029
Other tests (non-laboratory) Medicare utilization (percent)	Percent of Medicare fee-for-service beneficiaries 65+ years of age who used non-laboratory test services	Yes	Yes		Yes	2030
Otitis media visits in ambulatory care settings: children (per 1,000)	Rate of otitis media in children and adolescents	Yes				743

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Outbreak-associated infections associated with beef (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: beef	Yes				839
Outbreak-associated infections associated with chicken (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: poultry	Yes				843
Outbreak-associated infections associated with dairy (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: dairy	Yes				840
Outbreak-associated infections associated with fruits/nuts (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: fruits and nuts	Yes				841
Outbreak-associated infections associated with vegetables (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: leafy vegetables	Yes				842
Outpatient Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used outpatient services	Yes	Yes		Yes	187
Outpatient Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used outpatient services	Yes	Yes		Yes	188
Outpatient physical therapy/speech pathology facilities (number)	Number of Outpatient physical therapy/speech pathology facilities participating in the Medicare program		Yes		Yes	10045
Overall cancer deaths (per 100,000)	Overall cancer death rate	Yes	Yes	Yes		486
Overdose ED visits for injectable antidiabetic agents (per 10,000)	Rate of emergency department (ED) visits for overdoses from injectable antidiabetic agents	Yes				1204

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Overdose ED visits for narrow-therapeutic-index medications, (per 10,000)	Rate of emergency department (ED) visits for overdoses of narrow-therapeutic-index medications	Yes				1205
Overdose ED visits for oral anticoagulants (per 10,000)	Rate of emergency department (ED) visits for overdoses from oral anticoagulants	Yes				1203
Overdose ED visits: children <5 years (per 10,000)	Rate of emergency department (ED) visits for medication overdoses among children less than 5 years of age	Yes				1206
Overexertion or repetitive motion injuries/illness from work (per 10,000)	Rate of injury and illness cases involving days away from work due to overexertion or repetitive motion	Yes				1297
Ozone days (number)	Annual number of days with daily 8-hour maximum ozone concentration over the National Ambient Air Quality Standard			Yes		45
Pap test counseling: women 21-65 years (percent)	Percent of women who were counseled by their providers about Pap tests	Yes				495
Pap test: women 18+ (percent)	Percent of women age 18+ that report Pap test in the last 3 years		Yes	Yes		115
Part B drug Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used Part B drugs	Yes	Yes		Yes	269
Part B drugs Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used Part B drugs	Yes	Yes		Yes	270
Particulate matter days (percent)	Percent of days with maximum 24-hour average PM 2.5 concentration over the National Ambient Air Quality Standard			Yes		44
PCB level, dioxin-like: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: PCB 126, representative of dioxin-like PCBs	Yes				707

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
PCB level, nondioxin-like: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: PCB 153, representative of nondioxin-like PCBs	Yes				706
PCV vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-4 doses pneumococcal conjugate vaccine (PCV) by age 19 to 35 months	Yes				999
Pedal-cyclist deaths (per 100,000)	Death rate of pedal-cyclists on public roads	Yes				1103
Pedestrian deaths (per 100,000)	Rate of pedestrian deaths on public roads	Yes				1045
Pedestrian injuries, nonfatal (per 100,000)	Rate of nonfatal pedestrian injuries on public roads	Yes				1046
Pelvic inflammatory disease (percent)	Percent of females aged 15 to 44 years who have ever required treatment for pelvic inflammatory disease (PID)	Yes				1473
Penicillin-resistant pneumococcal infections: children <5 years (per 100,000)	Rate of invasive penicillin-resistant pneumococcal infections among children under age 5 years	Yes				965
Penicillin-resistant pneumococcal infections: older adults (per 100,000)	Rate of invasive penicillin-resistant pneumococcal infections among adults aged 65 years and older	Yes				966
Perchlorate level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: perchlorate	Yes				710
Percutaneous intervention timeliness: heart attack patients (percent)	Percent of eligible patients with heart attacks who receive Percutaneous intervention (PCI) within 90 minutes of hospital arrival	Yes				879
Perinatal deaths (per 1,000 live births plus fetal deaths)	Rate of fetal and infant deaths during perinatal period (28 weeks of gestation to 7 days after birth)	Yes				1157
Periodontal disease: adults 45-74 years (percent)	Percent of adults aged 45 to 74 years with periodontitis	Yes				1280

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Personal care limitations: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who have difficulty in performing two or more personal care activities, thereby preserving independence	Yes				420
Pertussis (number)	Number of reported cases of pertussis	Yes	Yes	Yes		10029
Pertussis: infants (number)	Number of pertussis cases among children under age 1 year	Yes				983
Pertussis: teens (number)	Number of pertussis cases among adolescents aged 11 to 18 years	Yes				984
Pesticide area marking policy: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Exposure to pesticides by marking areas treated with pesticides	Yes				687
Pesticide exposures causing health care facility visits (number)	Number of pesticide exposures that result in visits to a health care facility	Yes				661
Pesticide information policy: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Exposure to pesticides by notifying staff, students, and families prior to application of the pesticide	Yes				688
Pesticide limited use policy: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Exposure to pesticides by using spot treatments and baiting rather than widespread application of pesticide	Yes				686
Pesticide poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: pesticide poisoning	Yes				714

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Phthalate level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Mono-n-butyl phthalate	Yes				711
Physical activity counseling: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who receive health care provider counseling-For physical activity or exercise	Yes				422
Physical activity duration policies for child care settings: States (number)	Number of States with licensing regulations for physical activity provided in child care-Require number of minutes of physical activity per day or by length of time in care	Yes				1345
Physical activity intensity policies for child care settings: States (number)	Number of States with licensing regulations for physical activity provided in child care-Require children to engage in vigorous or moderate physical activity	Yes				1344
Physical activity policies for child care settings: States (number)	Number of States with licensing regulations for physical activity provided in child care-Require activity programs providing large muscle or gross motor activity, development, and/or equipment	Yes				1343
Physical activity: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in physical activity	Yes				604
Physical assault injuries, nonfatal (per 100,000)	Nonfatal physical assault injury rate	Yes				1099
Physical assaults: persons 12+ years (per 1,000)	Rate of physical assaults among persons 12 years of age and over	Yes				1051
Physical education required daily: elementary schools (percent)	Percent of the Nation's public and private schools that require daily physical education for all students-Elementary schools	Yes				1314

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Physical education required daily: high schools (percent)	Percent of the Nation's public and private schools that require daily physical education for all students-Senior high schools	Yes				1316
Physical education required daily: middle/junior high schools (percent)	Percent of the Nation's public and private schools that require daily physical education for all students-Middle and junior high schools	Yes				1315
Physical education, daily at school: teens (percent)	Percent of adolescents who participate in daily school physical education	Yes	Yes			1317
Physical fighting: teens (percent)	Percent of physical fighting among adolescents	Yes	Yes			1052
Physically or mentally unhealthy days: adults (days)	Average number of reported physically or mentally unhealthy days per month among persons 18 years and over		Yes	Yes		75
Physically unhealthy days: adults (per person)	Average number of reported physically unhealthy days per month among adults 18 years of age and over		Yes	Yes		7
Physican physical activity counseling: adults with chronic conditions (percent)	Percent of office visits made by patients with a diagnosis of cardiovascular disease, diabetes, or hyperlipidemia that include counseling or education related to exercise	Yes				1341
Physician asst. training in cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-Physician assistant	Yes				631
Physician asst. training, counseling for health promotion (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-Physician assistant	Yes				630
Physician asst. training, environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-Physician assistant	Yes				633
Physician asst. training, evaluation of health literature (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-Physician assistant	Yes				632

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Physician asst. training, global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-Physician assistant	Yes				635
Physician asst. training, public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-Physician assistant	Yes				634
Physician evaluation and management Medicare service events (per 1,000 beneficiaries)	Rate of physician evaluation and management service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2021
Physician evaluation and management Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used physician evaluation and management services	Yes	Yes		Yes	239
Physician evaluation and management Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used physician evaluation and management services	Yes	Yes		Yes	240
Physician physical activity counseling (percent)	Percent of physician visits made by all child and adult patients that include counseling about exercise	Yes				1342
Physician procedures Medicare service events (per 1,000 beneficiaries)	Rate of physician procedure service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2022
Physician procedures Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used physician procedures services	Yes	Yes		Yes	245
Physician procedures Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used physician procedures services	Yes	Yes		Yes	246
Physician use delayed due to cost (percent)	Percent of adults 18+ who needed to see a doctor but could not because of cost		Yes	Yes		10019
Pneumococcal vaccination: adults 65+ (percent)	Percent of adults age 65+ that report ever having a pneumococcal vaccination		Yes	Yes		118
Pneumococcal vaccination: high-risk adults (percent)	Percent of adults who are vaccinated against pneumococcal disease-Noninstitutionalized high-risk adults aged 18 to 64 years	Yes				1019

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Pneumococcal vaccination: nursing home residents (percent)	Percent of adults who are vaccinated against pneumococcal disease-Institutionalized adults (persons aged 18 years and older in long-term or nursing homes)	Yes				1020
Pneumococcal vaccination: older adults (percent)	Percent of adults who are vaccinated against pneumococcal disease-Noninstitutionalized adults aged 65 years and older	Yes				1018
Pneumoconiosis deaths (number)	Annual number of pneumoconiosis deaths	Yes				1298
Pneumonia patient with blood cultures in emergency department before antibiotic administered (percent)	Hospital Compare: Pneumonia patient with blood cultures in emergency department before antibiotic administered	Yes	Yes		Yes	332
Pneumonia patients with appropriate initial antibiotic selection for community-acquired pneumonia (percent)	Hospital Compare: Pneumonia patients with appropriate initial antibiotic selection for community-acquired pneumonia in immunocompetent patients	Yes	Yes		Yes	331
Pneumonia patients with influenza vaccination (percent)	Hospital Compare: Pneumonia patients with influenza vaccination	Yes	Yes		Yes	333
Pneumonia patients with initial antibiotic received within 6 hours of hospital arrival (percent)	Hospital Compare: Pneumonia patients with initial antibiotic received within 6 hours of hospital arrival	Yes	Yes		Yes	335
Pneumonia patients with pneumococcal vaccination (percent)	Hospital Compare: Pneumonia patients with pneumococcal vaccination	Yes	Yes		Yes	330
Pneumonia patients with smoking cessation counseling (percent)	Hospital Compare: Pneumonia patients with smoking cessation counseling	Yes	Yes		Yes	334
Poisoning deaths (per 100,000)	Poisoning death rate: All persons	Yes	Yes	Yes		1064
Poisoning deaths, unintentional/undetermined (per 100,000)	Poisoning death rate of unintentional or undetermined intent among all persons	Yes				1066
Poisoning deaths, unintentional/undetermined: adults 35-54 years (per 100,000)	Poisoning death rate of unintentional or undetermined intent among persons aged 35 to 54 years	Yes				1067

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Poisoning deaths: adults 35-54 years (per 100,000)	Poisoning death rate: Persons aged 35 to 54 years	Yes				1065
Poisonings, nonfatal (per 100,000)	Rate of nonfatal poisonings	Yes				1040
Polio vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-3 doses polio vaccine by age 19 to 35 months	Yes				997
Polio vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-Polio vaccine (3 or more doses)	Yes				1005
Polio, U.S.-acquired (number)	Number of acute paralytic poliomyelitis cases (U.S.-acquired cases)	Yes				985
Population density (per square mile)	Population density of United States geographic area per square mile	Yes	Yes	Yes		10001
Population distributions (percent)	Bridged-race population distributions according to 1977 OMB standard	Yes	Yes	Yes		10002
Population estimates (number)	Bridged-race population estimates according to 1977 OMB standard	Yes	Yes	Yes		10000
Portable X-Ray providers (number)	Number of portable X-Ray providers participating in the Medicare Program		Yes		Yes	10046
Positive communication with children: parents (percent)	Percent of parents who use positive communication with their child	Yes				739
Post acute care Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used post-acute care services	Yes	Yes		Yes	194
Post-acute care Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who used post-acute care services	Yes	Yes		Yes	193
Postdiarrheal hemolytic-uremic syndrome infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Postdiarrheal hemolytic uremic syndrome (HUS) in children less than 5 years of age	Yes				826
Poverty: children < 18 years (percent)	Percent of children under 18 years of age living in poverty	Yes	Yes	Yes		36

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Preemptive smoke-free air laws: States (number)	Number of States with State laws that preempt stronger local tobacco control laws-Preemption on smoke-free indoor air	Yes				1545
Preemptive tobacco advertising laws: States (number)	Number of States with State laws that preempt stronger local tobacco control laws-Preemption in advertising	Yes				1546
Preemptive tobacco control laws, youth access: States (number)	Number of States with State laws that preempt stronger local tobacco control laws-Preemption on youth access	Yes				1547
Preemptive transplantation in end stage kidney failure (percent)	Percent of patients who receive a preemptive transplant at the start of ESRD	Yes				474
Pregnancies conceived within 18 months of a previous birth (percent)	Percent of pregnancies conceived within 18 months of a previous birth	Yes				784
Pregnancies: females 15-17 years (per 1,000)	Pregnancy rate among adolescent females aged 15 to 17 years	Yes				788
Pregnancies: females 18-19 years (per 1,000)	Pregnancy rate among adolescent females aged 18 to 19 years	Yes				789
Pregnancy despite use of reversible contraceptive method (percent)	Percent of females experiencing pregnancy despite use of a reversible contraceptive method	Yes				780
Prenatal care, early and adequate (percent)	Percent of pregnant women who receive early and adequate prenatal care	Yes				1132
Prenatal care, first trimester (percent)	Percent of pregnant women who receive prenatal care beginning in the first trimester of pregnancy	Yes				1131
Prescription pain reliever abuse (percent)	Percent of person 12 years and older with nonmedical use of prescription pain relievers in the past year	Yes				1461
Prescription psychotherapeutic drug abuse (percent)	Percent of persons 12 years and older with nonmedical use of any prescription psychotherapeutic drug in the past year	Yes				1465
Prescription sedatives abuse (percent)	Percent of persons 12 years and older with nonmedical use of prescription sedatives in the past year	Yes				1464
Prescription stimulants abuse (percent)	Percent of persons 12 years and older with nonmedical use of prescription stimulants in the past year	Yes				1463

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Prescription tranquilizers abuse (percent)	Percent of persons 12 years and older with nonmedical use of prescription tranquilizers in the past year	Yes				1462
Pressure ulcer-related hospitalizations: older adults (per 100,000)	Rate of pressure ulcer-related hospitalizations among older adults	Yes				1249
Preterm births, <32 weeks of gestation (percent)	Percent of live births that are very preterm (less than 32 weeks of gestation)	Yes	Yes	Yes		1140
Preterm births, 32-33 weeks of gestation (percent)	Percent of live births at 32 to 33 weeks of gestation	Yes	Yes	Yes		1139
Preterm births, 34-36 weeks of gestation (percent)	Percent of live births that are late preterm (34 to 36 weeks of gestation)	Yes	Yes	Yes		1138
Preterm births, total (percent)	Percent of live births that are preterm (< 37 weeks of gestation)	Yes	Yes	Yes		1137
Preventable hospitalizations: Medicare beneficiaries (per 1,000)	Discharge rate per 1,000 Medicare fee-for-service enrollees for ambulatory care sensitive conditions		Yes	Yes		27
Proficiency in math: 12th grade (percent)	Percent of 12th graders whose mathematics skills are at or above the proficient achievement level for 12th grade	Yes				401
Proficiency in math: 4th grade (percent)	Percent of 4th graders whose mathematics skills are at or above the proficient achievement level for 4th grade	Yes				399
Proficiency in math: 8th grade (percent)	Percent of 8th graders whose mathematics skills are at or above the proficient achievement level for 8th grade	Yes				400
Proficiency in reading: 12th grade (percent)	Percent of 12th graders whose reading skills are at or above the proficient achievement level for 12th grade	Yes				398
Proficiency in reading: 4th grade (percent)	Percent of 4th graders whose reading skills are at or above the proficient achievement level for 4th grade	Yes				396
Proficiency in reading: 8th grade (percent)	Percent of 8th graders whose reading skills are at or above the proficient achievement level for 8th grade	Yes				397
Prostate cancer deaths (per 100,000)	Prostate cancer death rate	Yes				492

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Prostate cancer Medicare beneficiaries (count)	Medicare beneficiaries age 65+ years with full FFS Medicare who have prostate cancer	Yes	Yes		Yes	305
Prostate cancer Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have prostate cancer	Yes	Yes		Yes	306
Protective eyewear use at home: adults (percent)	Percent of adults aged 18 years and over who use protective eyewear in recreational activities and hazardous situations around the home	Yes				1573
Protective eyewear use at home: children (percent)	Percent of children and adolescents aged 6 to 17 years who use personal protective eyewear in recreational activities and hazardous situations around the home	Yes				1572
Protective gear use in intramural activities: schools (percent)	Percent of public and private schools that require students to wear appropriate protective gear when engaged in school-sponsored intramural activities or physical activity clubs	Yes				1083
Protective gear use in physical education: schools (percent)	Percent of public and private schools that require students to wear appropriate protective gear when engaged in school-sponsored physical education	Yes				1082
Psychiatric hospitals (number)	Number of Psychiatric hospitals participating in the Medicare program		Yes		Yes	10047
Public health majors: 4-year colleges/universities (percent)	Percent of 4-year colleges and universities that offer public health or related majors	Yes				1385
Public health minors: 4-year colleges/universities (percent)	Percent of 4-year colleges and universities that offer public health or related minors	Yes				1386
Public health system assessment: Local public health systems (percent)	Percent of State and local public health jurisdictions that conduct a public health system assessment using national performance standards-Local public health systems	Yes				1376
Public health system assessment: State public health systems (percent)	Percent of State and local public health jurisdictions that conduct a public health system assessment using national performance standards-State public health systems	Yes				1375

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Pyrethroid level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: 3-phenoxybenzoic acid	Yes				705
Rabies surveillance, electronic: states (number)	Number of States that use electronic data from rabies animal surveillance to inform public health prevention programs	Yes				1024
Radon mitigation systems: homes at risk for radon exposure (percent)	Percent of homes with an operating radon mitigation system for persons living in homes at risk for radon exposure	Yes				681
Radon-reducing construction: new homes (percent)	Percent of new single family homes (SFH) constructed with radon-reducing features, especially in high-radon-potential areas	Yes				682
Read to children: parents (percent)	Percent of parents who read to their young child	Yes				740
Recent hearing exams: adults 20-69 years (percent)	Percent of adults aged 20 to 69 years who have had a hearing examination in the past 5 years	Yes				748
Recent hearing exams: older adults (percent)	Percent of adults aged 70 years and older who have had a hearing examination in the past 5 years	Yes				749
Recent hearing exams: teens (percent)	Percent of adolescents aged 12 to 19 years who have had a hearing examination in the past 5 years	Yes				750
Recess for 20+ min. in elementary school: school districts (percent)	Percent of school districts that require or recommend elementary school recess for an appropriate period of time	Yes				1340
Recess, required in elementary school: school districts (percent)	Percent of school districts that require regularly scheduled elementary school recess	Yes				1339
Recess, required in elementary school: States (number)	Number of States that require regularly scheduled elementary school recess	Yes				1338
Recycling of municipal waste (percent)	Percent of municipal solid waste that is recycled or composted	Yes				662

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Reducing fat/calories in diet: adults with pre-diabetes (percent)	Percent of persons at high risk for diabetes (with pre-diabetes) who report reducing the amount of fat or calories in their diet	Yes				564
Refrigerate promptly, food safety practice (percent)	Percent of consumers who follow key food safety practices-- Chill: Refrigerate promptly	Yes				838
Rehabilitation Hospitals (number)	Number of Rehabilitation hospitals participating in the Medicare program		Yes		Yes	10048
Religious Nonmedical Health Care Institutions (number)	Number of religious nonmedical health care institutions participating in the Medicare program		Yes		Yes	10049
Renal evaluation after kidney injury: Medicare beneficiaries (percent)	Percent of hospital patients who incurred acute kidney injury and have followup renal evaluation in 6 months post discharge	Yes				482
Renal function knowledge: adults aged 45+ with chronic kidney disease (percent)	Percent of adults 45 years and older with chronic kidney disease (CKD) who know they have impaired renal function	Yes				485
Reports of fetal death, use of latest standard form: states (number)	Number of States that record vital events using the latest U.S. standard certificates and report - States using the standard report of fetal death	Yes				1384
Reproductive health services receipt: sexually active females (percent)	Percent of sexually active females aged 15 to 44 years who received reproductive health services	Yes				786
Reproductive health services receipt: sexually active males (percent)	Percent of sexually active males aged 15 to 44 years who received reproductive health services	Yes				787
Residential fire deaths (per 100,000)	Residential fire death rate	Yes				1050
Riding with a driver who has been drinking: teens (percent)	Percent of adolescents who report that they rode, during the previous 30 days, with a driver who had been drinking alcohol	Yes	Yes			1427

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Risk perception, binge drinking: at risk teens (percent)	Percent of adolescents aged 12 to 17 years perceiving great risk associated with substance abuse-Consuming five or more alcoholic drinks at a single occasion once or twice a week	Yes				1450
Risk perception, cocaine: at risk teens (percent)	Percent of adolescents aged 12 to 17 years perceiving great risk associated with substance abuse-Using cocaine once per month	Yes				1452
Risk perception, marijuana: at risk teens (percent)	Percent of adolescents aged 12 to 17 years perceiving great risk associated with substance abuse-Smoking marijuana once per month	Yes				1451
Rotavirus vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-2 or more doses rotavirus vaccine (aged 19 to 35 months)	Yes				1002
Rubella, U.S.-acquired (number)	Number of rubella cases (U.S.-acquired cases)	Yes				986
Rural Health Clinics (number)	Number of rural health clinics participating in the Medicare program		Yes		Yes	10050
Safe drinking water: persons served by community water (percent)	Percent of persons served by community water systems who receive a supply of drinking water that meets the regulations of the Safe Drinking Water Act	Yes				659
Safety at school: perception of teens' parents (percent)	Percent of adolescents whose parents consider them to be safe at school	Yes				407
Safety belt use (percent)	Percent of persons observed using safety belts	Yes				1047
Salmonella species infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Reduction in infections caused by Salmonella species	Yes				825
Salmonellosis (number)	Number of reported cases of salmonellosis	Yes	Yes	Yes		10026
Saturated fat consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake from saturated fat for persons aged 2 years and over	Yes				1221

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
School absenteeism from illness/injury: teens aged 12-17 (percent)	Percent of adolescents missing school 11 or more days due to illness or injury	Yes				403
School athletic facilities, public access (percent)	Percent of the Nation's public and private schools that provide access to their physical activity spaces and facilities for all persons outside of normal school hours (that is, before and after the school day, on weekends, and during summer and other vacations)	Yes				1318
School nurse-to-student ratio: elementary schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered school nurse-to-student ratio of at least 1:750-Elementary schools	Yes				588
School nurse-to-student ratio: high schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered school nurse-to-student ratio of at least 1:750-Senior high schools	Yes				586
School nurse-to-student ratio: middle schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered school nurse-to-student ratio of at least 1:750-Middle schools	Yes				587
School nurse-to-student ratio: schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered school nurse-to-student ratio of at least 1:750-All elementary, middle, and senior high schools	Yes				585
School work meaningful: teens aged 12-17 (percent)	Percent of adolescents who consider their school work to be meaningful and important	Yes				402
Schools located near highways (percent)	Number of new schools sited within 500 feet of an interstate or Federal or State highway	Yes				725
Screen time, computer: children 2-5 years (percent)	Percent of children aged 2 to 5 years who use a computer or play computer games outside of school (for nonschool work) for no more than 2 hours a day	Yes				1330
Screen time, computer: children 6-14 years (percent)	Percent of children aged 6 to 14 years who use a computer or play computer games outside of school (for nonschool work) for no more than 2 hours a day	Yes				1331

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Screen time, computer: teens (percent)	Percent of adolescents in grades 9 through 12 who use a computer or play computer games outside of school (for nonschool work) for no more than 2 hours a day	Yes	Yes			1332
Screen time: children 0-2 years (percent)	Percent of children aged 0 to 2 years who view no television or videos on an average weekday	Yes				1326
Screen time: children 2-5 years (percent)	Percent of children aged 2 to 5 years who view television, videos, or play video games for no more than 2 hours a day	Yes				1327
Screen time: children 6-14 years (percent)	Percent of children aged 6 to 14 years who view television, videos, or play video games for no more than 2 hours a day	Yes				1328
Screen time: teens (percent)	Percent of adolescents in grades 9 through 12 who view television, videos, or play video games for no more than 2 hours a day	Yes	Yes			1329
Secondhand smoke exposure: adult nonsmokers (percent)	Percent of nonsmokers exposed to secondhand smoke- Adults aged 18 years and older	Yes				1519
Secondhand smoke exposure: nonsmokers 12-17 years (percent)	Percent of nonsmokers exposed to secondhand smoke- Adolescents aged 12 to 17 years	Yes				1518
Secondhand smoke exposure: nonsmokers 3-11 years (percent)	Percent of nonsmokers exposed to secondhand smoke- Children aged 3 to 11 years	Yes				1517
Self-harm injuries, nonfatal (per 100,000)	Nonfatal intentional self-harm injury rate	Yes				1100
Separate food, food safety practice (percent)	Percent of consumers who follow key food safety practices-- Separate: Don't cross-contaminate	Yes				836
Serious property crimes with perpetrators 10-24 years (per 100,000)	Rate of minor and young adult perpetration of serious property crimes	Yes				414
Serious psychological distress: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who report serious psychological distress	Yes				435
Serious violent incidents in public schools (percent)	Percent of public schools with a serious violent incident	Yes				408
Shigellosis (number)	Number of reported cases of shigellosis	Yes	Yes	Yes		10027

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Shingles vaccination: older adults (percent)	Percent of adults who are vaccinated against zoster (shingles)	Yes				1027
Short stay hospitals (number)	Number of short stay hospitals participating in the Medicare program		Yes		Yes	10051
Sick leave use: adults with balance/dizziness problems (percent)	Percent of adults who have missed days of work or school in the past 12 months because of balance and dizziness problems	Yes				768
Single-parent households (percent)	Percent of households that are single-parent households		Yes	Yes		40
Skilled nursing facility Medicare admissions (per 1,000 beneficiaries)	Rate of skilled nursing facility (SNF) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2008
Skilled nursing facility Medicare days (per 1,000 beneficiaries)	Average number of skilled nursing facility (SNF) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes		Yes	2016
Skilled nursing facility Medicare users (count)	Medicare fee-for-service beneficiaries 65+ years of age who had covered skilled nursing facility services	Yes	Yes		Yes	214
Skilled nursing facility Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered skilled nursing facility services	Yes	Yes		Yes	215
Skilled Nursing Home Title 18 Facilities (number)	Number of skilled nursing home Title 18 (Medicare only) facilities		Yes		Yes	10052
Skin cancer protective measures: adults (percent)	Percent of adults aged 18 years and older who follow protective measures that may reduce the risk of skin cancer	Yes				504
Skin cancer protective measures: teens (percent)	Percent of adolescents in grades 9 through 12 who follow protective measures that may reduce the risk of skin cancer	Yes				503
Skin diseases or disorders, work-related (per 10,000)	Rate of occupational skin diseases or disorders among full-time workers	Yes				1301
Smoke-free air laws covering bars: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in bars	Yes				1527

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Smoke-free air laws covering commercial daycare centers: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in commercial daycare centers	Yes				1529
Smoke-free air laws covering home based daycare centers: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in home-based daycare centers	Yes				1530
Smoke-free air laws covering hotels and motels: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in hotels and motels	Yes				1532
Smoke-free air laws covering multiunit housing: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in multiunit housing	Yes				1533
Smoke-free air laws covering Prisons: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in prisons and correctional facilities	Yes				1535
Smoke-free air laws covering private workplaces: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in private workplaces	Yes				1524
Smoke-free air laws covering public transportation: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in public transportation	Yes				1531
Smoke-free air laws covering public workplaces: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in public workplaces	Yes				1525
Smoke-free air laws covering restaurants: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in restaurants	Yes				1526
Smoke-free air laws covering vehicles with children: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in vehicles with children	Yes				1534

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Smoke-free homes: adults (percent)	Percent of adults living in smoke-free homes	Yes				1561
Smokeless tobacco use in past month: teens (percent)	Percent of adolescent reporting use of smokeless tobacco products in the past month - Students (grades 9 through 12)	Yes	Yes			1503
Smokeless tobacco use initiation: adults 18-25 years (percent)	Percent of young adults aged 18 to 25 years initiating smokeless tobacco use in the past 12 months	Yes				1511
Smokeless tobacco use initiation: teens (percent)	Percent of children and adolescents aged 12 to 17 years initiating smokeless tobacco use in past 12 months	Yes				1507
Smokeless tobacco use: adults (percent)	Percent of adults aged 18 years and over using smokeless tobacco products	Yes				1499
Smoking abstinence, preconception (percent)	Percent of women delivering a live birth who did not smoke prior to pregnancy	Yes				1180
Smoking cessation attempts: adult smokers (percent)	Percent of adult smokers aged 18 years and over who attempted smoking cessation in the past 12 months	Yes				1513
Smoking cessation attempts: teen smokers (percent)	Percent of adolescent smokers in grades 9 through 12 who attempted smoking cessation in the past 12 months	Yes	Yes			1515
Smoking cessation during pregnancy (percent)	Percent of females aged 18 to 49 years who ceased smoking during the first trimester of pregnancy	Yes				1491
Smoking cessation success (percent)	Percent of adults 18 years and older with smoking cessation success--6 months to 1 year ago	Yes				1555
Smoking: adults (percent)	Percent of adults 18 years and over that report currently smoking cigarettes		Yes	Yes		13
Social and emotional support: adults with disabilities (percent)	Percent of adults with disabilities who report sufficient social and emotional support	Yes	Yes			515
Social support for health concerns (percent)	Percent of adults who report having friends or family members whom they talk with about their health	Yes				870
Social-emotional support: Adults (percent)	Percent of adults 18 years and over that report receiving sufficient social-emotional support		Yes	Yes		39

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Sodium consumption (milligrams)	Mean total daily sodium intake (in mg) by persons aged 2 years and over	Yes				1222
Solid fat and added sugar consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake from solid fats and added sugars for persons aged 2 years and over	Yes				1238
Solid fat consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake from solid fats for persons aged 2 years and over	Yes				1236
Source of ongoing care: <18 years (percent)	Percent of persons who have a specific source of ongoing care-children and youth aged 17 years and under	Yes				376
Source of ongoing care: 18 to 64 years (percent)	Percent of persons who have a specific source of ongoing care-adults aged 18 to 64 years	Yes				377
Source of ongoing care: 65+ years (percent)	Percent of persons who have a specific source of care - adults aged 65 years and older	Yes				378
Source of ongoing care: all ages (percent)	Percent of persons who have a specific source of ongoing care-all ages	Yes				375
Special services: children with Autism spectrum disorders (percent)	Percent of children with an ASD enrolled in special services by 48 months of age	Yes				1190
Specialty health care use: adults with balance/dizziness problems (percent)	Percent of adults with moderate to severe balance or dizziness problems who have seen or been referred to a Health Care specialist for evaluation or treatment	Yes				763
Specialty health care use: adults with tinnitus (percent)	Percent of adults bothered by the onset of tinnitus in the past 5 years, for whom it is a moderate, big, or very big problem, who have seen or been referred to an audiologist or otolaryngologist (ENT physician)	Yes				760
Spina bifida (per 100,000 live births)	Rate of spina bifida	Yes				1169
Spinal cord injury deaths (per 100,000)	Spinal cord injury death rate	Yes	Yes			1062
Spinal cord injury hospitalizations (per 100,000)	Hospitalization rate for nonfatal traumatic spinal cord injuries	Yes				1063
Spotted Fever Rickettsiosis (number)	Number of reported cases of spotted fever rickettsiosis	Yes	Yes	Yes		10028

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
State epidemiologists with formal training in epidemiology (percent)	Percent of Tribal, State, and local public health agencies that provide or assure comprehensive epidemiology services to support essential public health services-State epidemiologists with formal training in epidemiology	Yes				1360
Statewide cancer registries (number)	Number of central, population-based registries from the 50 States and the District of Columbia that capture case information on at least 95 percent of the expected number of reportable cancers	Yes				497
STD discussion with parent before age 18: females (percent)	Percent of adolescents females who talked to a parent or guardian about STDs before they were 18 years old	Yes				816
STD discussion with parent before age 18: males (percent)	Percent of adolescents males who talked to a parent or guardian about STDs before they were 18 years old	Yes				817
STD instruction receipt before age 18: females (percent)	Percent of adolescents females who received formal instruction on STDs before they were 18 years old	Yes				808
STD instruction receipt before age 18: males (percent)	Percent of adolescents males who received formal instruction on STDs before they were 18 years old	Yes				809
Steroid use: 10th graders (percent)	Percent of 10th graders using steroids	Yes				1441
Steroid use: 8th graders (percent)	Percent of 8th graders using steroids	Yes				1440
Steroid use: high school seniors (percent)	Percent of 12th graders using steroids	Yes				1442
Streptococcus pneumoniae, drug-resistant (number)	Number of reported cases of drug-resistant streptococcus pneumoniae	Yes	Yes	Yes		10025
Stroke awareness: importance of calling 9-1-1 (percent)	Percent of adults aged 20 years and over who are aware of the importance of accessing rapid emergency care by calling 9-1-1 or another emergency number	Yes				889
Stroke awareness: symptoms (percent)	Percent of adults aged 20 years and over who are aware of the early warning symptoms and signs of a stroke	Yes				888

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Stroke awareness: symptoms and importance of calling 9-1-1 (percent)	Percent of adults who are aware of the early warning symptoms and signs of a stroke and the importance of accessing rapid emergency care by calling 9-1-1 or another emergency number	Yes				887
Stroke deaths (per 100,000)	Stroke death rate	Yes	Yes	Yes		881
Substance abuse prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in substance abuse	Yes				600
Sufficient sleep: adults (percent)	Percent of adults who get sufficient sleep	Yes				1472
Sufficient sleep: teens (percent)	Percent of students in grades 9 through 12 who get sufficient sleep	Yes				1471
Suffocation deaths, unintentional (per 100,000)	Unintentional suffocation death rate: All persons	Yes				1068
Suffocation deaths, unintentional: infants (per 100,000)	Unintentional suffocation death rate: Infants 0 to 12 months	Yes				1069
Suffocation deaths, unintentional: older adults (per 100,000)	Unintentional suffocation death rate: Adults aged 65 years and over	Yes				1070
Suicide attempts requiring medical attention: teens (per 100)	Rate of suicide attempts by adolescents that require medical treatment	Yes	Yes			1106
Suicide deaths (per 100,000)	Suicide death rate	Yes	Yes	Yes		1105
Surgery patients who received appropriate venous thromboembolism prophylaxis (percent)	Hospital Compare: Surgery patients who received appropriate venous thromboembolism prophylaxis	Yes	Yes		Yes	342
Surgery patients with appropriate hair removal (percent)	Hospital Compare: Surgery patients with appropriate hair removal	Yes	Yes		Yes	340

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Surgery patients with appropriate prophylactic antibiotic selection (percent)	Hospital Compare: Surgery patients with appropriate prophylactic antibiotic selection	Yes	Yes		Yes	337
Surgery patients with prophylactic antibiotic received within one hour prior to surgery incision (percent)	Hospital Compare: Surgery patients with prophylactic antibiotic received within one hour prior to surgery incision	Yes	Yes		Yes	336
Surgery patients with prophylactic antibiotics discontinued within 24 hours after surgery end time (percent)	Hospital Compare: Surgery patients with prophylactic antibiotics discontinued within 24 hours after surgery end time	Yes	Yes		Yes	338
Surgery patients with recommended venous thromboembolism prophylaxis ordered (percent)	Hospital Compare: Surgery patients with recommended venous thromboembolism prophylaxis ordered	Yes	Yes		Yes	341
Surveillance programs for caregivers: States (number)	States and the District of Columbia with surveillance programs for caregivers	Yes				528
Survival after AIDS diagnosis (percent)	Percent of persons surviving more than 3 years after a diagnosis with AIDS	Yes				937
Sweetened beverages not offered: schools (percent)	Percent of schools that do not sell or offer calorically sweetened beverages to students.	Yes				10065
Syphilis, congenital (per 100,000)	Congenital syphilis rate	Yes				1474
Syphilis, primary and secondary: females (per 100,000)	Sustained domestic transmission of primary and secondary syphilis among females	Yes				1480
Syphilis, primary and secondary: males (per 100,000)	Sustained domestic transmission of primary and secondary syphilis among males	Yes				1481
TB curative therapy completion (percent)	Percent of all tuberculosis patients who complete curative therapy within 12 months	Yes				970
TB test confirmation timeliness (percent)	Average time for a laboratory to confirm and report tuberculosis cases	Yes				972
TB treatment: persons with latent TB (percent)	Percent of persons with latent tuberculosis infection who complete a course of treatment	Yes				971

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Tdap vaccination: teens 13-15 years (percent)	Percent of adolescents who have 1 dose of tetanus-diphtheria-acellular pertussis (Tdap) booster vaccine by 13 to 15 years	Yes				1010
Telecommuting (percent)	Percent of trips to work using of alternative modes of transportation - telecommute	Yes				669
Timely evaluation: children with Autism spectrum disorder (percent)	Percent of children with an ASD with a first evaluation by 36 months of age	Yes				1189
Tobacco advertising exposure, internet: teens (percent)	Percent of adolescents and young adults grades 6 through 12 who are exposed to tobacco advertising and promotion-Internet advertising and promotion	Yes				1541
Tobacco advertising exposure, magazine/newspaper: teens (percent)	Percent of adolescents and young adults grades 6 through 12 who are exposed to tobacco advertising and promotion-Magazine and newspaper advertising and promotion	Yes				1542
Tobacco cessation counseling in hospital outpatient dept. visits: adults (percent)	Percent of tobacco users ages 18 and older who receive tobacco cessation counseling at hospital ambulatory care visits	Yes				1552
Tobacco cessation counseling in physician office visits: adults (percent)	Percent of tobacco users ages 18 and older who receive tobacco cessation counseling at office based ambulatory care visits	Yes				1551
Tobacco free environments: high schools (percent)	Percent of senior high schools with tobacco-free environments	Yes				1522
Tobacco free environments: junior high schools (percent)	Percent of Junior high schools with tobacco-free environments	Yes				1520
Tobacco free environments: middle schools (percent)	Percent of middle schools with tobacco-free environments	Yes				1521
Tobacco screening in hospital outpatient dept. visits: adults (percent)	Percent of persons aged 18 years and older who received tobacco screening in hospital ambulatory care settings	Yes				1558
Tobacco screening in physician office visits: adults (percent)	Percent of persons aged 18 years and older who received tobacco screening in office-based ambulatory care settings	Yes				1557

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Tobacco tax increases on cigarettes: States/Federal (number)	Number of States, DC, and Federal government that have increased tax on cigarettes by \$1.50	Yes				1548
Tobacco tax increases on smokeless tobacco: States/Federal (number)	Number of States, DC, and Federal government that have increased tax on smokeless tobacco products by \$1.50	Yes				1549
Tobacco use in past month: teens (percent)	Percent of adolescent reporting use of tobacco products in the past month - Students (grades 9 through 12)	Yes	Yes			1501
Tobacco use initiation: adults 18-25 years (percent)	Percent of young adults aged 18 to 25 years initiating tobacco products use in the past 12 months	Yes				1509
Tobacco use initiation: teens (percent)	Percent of children and adolescents aged 12 to 17 years initiating tobacco use in past 12 months	Yes				1505
Tobacco use prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in tobacco	Yes				599
Tooth loss, all permanent: adults 65-74 years (percent)	Percent of older adults aged 65 to 74 years who have lost all of their natural teeth	Yes				1279
Tooth loss, any permanent: adults 45-64 years (percent)	Percent of adults aged 45 to 64 years who have ever had a permanent tooth extracted because of dental caries or periodontitis	Yes				1278
Toxic chemicals (pounds)	Toxic chemicals released annually	Yes	Yes	Yes		105
Toxic pollutants (tons)	Amount of toxic pollutants released into the environment	Yes				678
Transition planning: teens with special health care needs (percent)	Percent of youth with special health care needs whose health care provider has discussed transition planning from pediatric to adult Health Care	Yes				536
Trauma care access, by land mass (percent)	Percent of the land mass of the continental United States with access to trauma care	Yes				1097
Trauma care access, by population (percent)	Percent of the population residing within the continental United States with access to trauma care	Yes				1096

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Traumatic brain injury deaths (per 100,000)	Traumatic brain injury death rate	Yes	Yes			1059
Traumatic brain injury ED visits (per 100,000)	Rate of emergency department visits for nonfatal traumatic brain injuries	Yes				1061
Traumatic brain injury hospitalizations (per 100,000)	Hospitalization rate for nonfatal traumatic brain injuries	Yes				1060
Treatment: adults with both substance abuse and mental disorders (percent)	Percent of persons with co-occurring substance abuse and mental disorders who receive treatment for both disorders	Yes				1119
Treatment: adults with major depressive episode (percent)	Percent of adults aged 18 years and older with major depressive episode	Yes				1118
Treatment: adults with serious mental illness (percent)	Percent of adults aged 18 years and older with serious mental illness (SMI)	Yes				1117
Treatment: children with mental health problems (percent)	Percent of children with mental health problems who receive treatment	Yes				1110
Trying to lose weight: adults with pre-diabetes (percent)	Percent of persons at high risk for diabetes (with pre-diabetes) who report trying to lose weight	Yes				563
Tuberculosis (per 100,000)	Tuberculosis (TB) rate	Yes				969
Tuberculosis: foreign born persons (per 100,000)	Tuberculosis (TB) case rate for foreign-born persons living in the United States	Yes				845
Unemployed: persons 16+ (percent)	Percent of population age 16+ unemployed and looking for work		Yes	Yes		35
Unemployment: adults 18-64 years with arthritis (percent)	Unemployment rate among adults aged 18-64 years with doctor-diagnosed arthritis	Yes				423
Unemployment: persons 16-64 years with disabilities (percent)	Unemployment rates among adults aged 16-64 with disabilities	Yes				541
Uninsured: persons less than 65 years (percent)	Percent of persons under 65 years of age without health insurance		Yes	Yes		23

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Unintended pregnancy prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in unintended pregnancy	Yes				601
Unintentional Injury deaths: No MVA (per 100,000)	Unintentional injury death rate (excluding motor vehicle injuries)	Yes	Yes	Yes		79
Urinary microalbumin measurement: diabetic Medicare beneficiaries (percent)	Percent of persons with diagnosed diabetes who obtain an annual urinary microalbumin measurement	Yes				549
Urinary tract infection admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Urinary tract infection admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes		Yes	2044
Use of hearing protection: adults (per 1,000)	Rate of adults aged 20 to 69 years who have ever used hearing protection devices (earplugs, earmuffs) when exposed to loud sounds or noise	Yes				752
Use of hearing protection: teens (per 1,000)	Rate of adolescents aged 12 to 19 years who have ever used hearing protection devices (earplugs, earmuffs) when exposed to loud sounds or noise	Yes				753
Use of mobile devices for internet access (percent)	Percent of persons who use mobile devices for internet use	Yes				859
Usual primary care provider (percent)	Percent of persons with a usual primary care provider	Yes				372
Uterine cervix cancer deaths (per 100,000)	Uterine cervix cancer death rate	Yes				489
Vaccination coverage assessment: private health providers (percent)	Percent of private providers who have had vaccination coverage levels among children in their practice population measured within the past year	Yes				977
Vaccination coverage assessment: public health providers (percent)	Percent of public health providers who have had vaccination coverage levels among children in their practice population measured within the past year	Yes				976

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Vaccination record keeping (percent)	Percent of children under age 6 years of age whose immunization records are in fully operational, population-based immunization information systems	Yes				1009
Vaccination, zero doses: children 18-35 months (percent)	Percent of children in the United States who receive zero doses of recommended vaccines by age 19 to 35 months	Yes				1039
Varicella vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-1 dose varicella vaccine by age 19 to 35 months	Yes				998
Varicella vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-Varicella vaccine (2 or more doses)	Yes				1007
Varicella vaccination: teens 13-15 years (percent)	Percent of adolescents who have 2 doses of varicella vaccine by age 13 to 15 years (excluding children who have had varicella)	Yes				1011
Varicella: children (number)	Rate of varicella (chicken pox) cases among persons aged 17 years of age and under	Yes				987
Vascular access for hemodialysis (percent)	Percent of adult hemodialysis patients who use arteriovenous fistulas as the primary mode of vascular access	Yes				469
Vegetable consumption (cup equivalents per 1,000 calories)	Mean daily intake of vegetables (in cup equivalents per 1,000 calories) by persons aged 2 years and over	Yes				1218
Vegetable consumption-dark green, orange, and legumes (cup equivalents per 1,000 calories)	Mean daily intake of dark green vegetables, orange vegetables, and legumes (in cup equivalents per 1,000 calories) by persons aged 2 years and over	Yes				1219
Vehicle restraint proper use: children 1-3 years (percent)	Percent of age-appropriate vehicle restraint system use in children aged 1 to 3 years	Yes				1055
Vehicle restraint proper use: children 4-7 years (percent)	Percent of age-appropriate vehicle restraint system use in children aged 4 to 7 years	Yes				1056
Vehicle restraint proper use: children 8-12 years (percent)	Percent of age-appropriate vehicle restraint system use in children aged 8 to 12 years	Yes				1057

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Vehicle restraint proper use: infants 0-12 months (percent)	Percent of age-appropriate vehicle restraint system use in infants 0 to 12 months	Yes				1054
Venous thromboembolism physician visits (per 10,000)	Rate of visits for venous thromboembolism (VTE)	Yes				441
Very low birth weight infant (percent)	Percent of live births that are very low birth weight (VLBW)	Yes	Yes	Yes		1136
Very low birth weight infants born at level III hospitals (percent)	Percent of very low birth weight (VLBW) infants born at level III hospitals or subspecialty perinatal centers	Yes				1195
Vibrio species infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Vibrio species	Yes				827
Violence exposure: children (percent)	Percent of children who are exposed to violence	Yes				1104
Violence prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in violence	Yes				597
Violent crime (per 100,000)	Violent crime rate		Yes	Yes		42
Violent crimes: perpetrators 10-24 years (per 100,000)	Rate of minor and young adult perpetration of violent crimes	Yes				413
Violent death surveillance: states (number)	Number of states that link data on violent deaths from death certificates, law enforcement, and coroner and medical examiner reports to inform prevention efforts at the state and local levels	Yes				1101
Vision screening: children <5 years (percent)	Percent of preschool children aged 5 years and under who receive vision screening	Yes				1562
Visitable features: homes and residential buildings (percent)	Percent of newly constructed or retrofitted U.S. homes and residential buildings that have visitable features	Yes				542
Visual impairment due to cataracts: older adults (per 1,000)	Rate of visual impairment due to cataract	Yes				1570

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Visual impairment due to diabetic retinopathy: adults with diabetes (per 1,000)	Rate of visual impairment due to diabetic retinopathy	Yes				1568
Visual impairment due to macular degeneration: adults 45+ years (per 1,000)	Rate of visual impairment due to age-related macular degeneration (AMD)	Yes				1571
Visual impairment due to refractive errors: persons 12+ years (per 1,000)	Rate of visual impairment due to uncorrected refractive error	Yes				1567
Visual impairment: children <18 years (per 1,000)	Percent of children and adolescents aged 17 years and under with blindness and visual impairment	Yes				1563
Visual impairments due to glaucoma: adults 45+ years (per 1,000)	Rate of visual impairment due to glaucoma	Yes				1569
Visual rehabilitation services use: adults with visual impairments (per 1,000)	Rate of use of vision rehabilitation services by persons with visual impairment	Yes				1574
von Willebrand Disease diagnosis: women (percent)	Percent of women with von Willebrand disease (vWD) who are timely and accurately diagnosed	Yes				443
Walking for transportation (percent)	Percent of trips to work using of alternative modes of transportation - walking	Yes				667
Water quality/sanitation/hygiene deaths: global (number)	Global burden of disease resulting in death due to poor water quality, sanitation, and insufficient hygiene	Yes				720
Waterborne disease outbreaks (number)	Number of waterborne disease outbreaks arising from water intended for drinking among persons served by community water systems	Yes				660
Weapon carrying on school property: teens (percent)	Percent of weapon carrying by adolescents on school property	Yes	Yes			1053
Weight loss counseling: overweight adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who receive health care provider counseling-For weight reduction among overweight and obese persons	Yes				421

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Weight loss, nutrition or physical activity counseling: obese patients (percent)	Percent of physician office visits made by adult patients who are obese that include counseling or education related to weight reduction, nutrition, or physical activity	Yes				1226
Welcome to Medicare benefit use (percent)	Percent of older adults who use the Welcome to Medicare benefit	Yes				1260
Wellness check-up: teens aged 12-17 (percent)	Percent of adolescents who have had a wellness checkup in the past 12 months	Yes				405
Whole grain consumption (ounce equivalents per 1,000 calories)	Mean daily intake of whole grains (in ounce equivalents per 1,000 calories) by persons aged 2 years and over	Yes				1220
Work limitations: adults 18-64 years with arthritis (percent)	Percent of adults aged 18-64 years with doctor-diagnosed arthritis who are limited in their ability to work for pay due to arthritis	Yes				424
Worksite lactation support programs (percent)	Percent of employers that have worksite lactation support programs	Yes				1184
Years of potential life lost before age 75 (per 100,000)	Rate per 100,000 of years of potential life lost before age 75	Yes	Yes	Yes		3
Yersinia species infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Yersinia species	Yes				828