
**INCIDENTAL HARASSMENT AUTHORIZATION APPLICATION
FOR THE NAVY'S EHW-1 PILE REPLACEMENT PROJECT
CONDUCTED AT
NAVAL BASE KITSAP BANGOR**



Submitted to:
**Office of Protected Resources,
National Marine Fisheries Service,
National Oceanographic and Atmospheric Administration**

Prepared by:
Naval Facilities Engineering Command
For:
Naval Base Kitsap Bangor and Naval Strategic Systems Program

December 2010

This Page Intentionally Left Blank

TABLE OF CONTENTS

ACRONYMS AND ABBREVIATIONS.....	vii
EXECUTIVE SUMMARY.....	ix
1 DESCRIPTION OF ACTIVITIES	1
1.1 Introduction	1
1.2 Proposed Action	1
1.3 Description of Pile Installation/Removal and Other Construction Activities	1
2 DATES, DURATION, AND LOCATION OF ACTIVITIES	9
2.1 Dates of Construction	9
2.2 Duration of Activities	9
2.3 Study Area Description	10
2.3.1 Bathymetric Setting	10
2.3.2 Tides	10
2.3.3 Circulation and Currents	13
2.3.4 Sea State	14
2.3.5 Water Temperature	14
2.3.6 Dissolved Oxygen	14
2.3.7 Stratification and Salinity	16
2.3.8 Sediments	16
2.3.9 Ambient Underwater Soundscape	16
3 MARINE MAMMAL SPECIES AND NUMBERS	19
3.1 ESA-Listed Marine Mammals	20
3.1.1 Steller Sea Lion	20
3.2 Non-ESA Listed Marine Mammals	20
3.2.1 California Sea Lion	20
3.2.2 Harbor Seal	21
3.2.3 Killer Whale	21
3.2.4 Dall's Porpoise	22
3.2.5 Harbor Porpoise	23
3.3 Marine Mammal Modeling Parameters	23
3.3.1 Spatial Distribution	23
3.3.2 Submergence	24
4 STATUS AND DISTRIBUTION OF MARINE MAMMAL SPECIES OR STOCKS THAT COULD POTENTIALLY BE AFFECTED	25
4.1 ESA-Listed Marine Mammals	25
4.1.1 Steller Sea Lion (<i>Eumetopias jubatus</i>), Eastern U.S. Stock	25

4.2	Non-ESA Listed Marine Mammals	26
4.2.1	California Sea Lion (<i>Zalophus californianus</i>), U.S. Stock	26
4.2.2	Harbor Seal (<i>Phoca vitulina</i>), WA Inland Waters Stock	28
4.2.3	Killer whale (<i>Orcinus orca</i>), West Coast Transient Stock	30
4.2.4	Dall's Porpoise (<i>Phocoenoides dalli</i>), CA/OR/WA Stock	31
4.2.5	Harbor Porpoise (<i>Phocoena phocoena</i>), WA Inland Waters Stock	32
5	HARASSMENT AUTHORIZATION REQUESTED	35
5.1	Take Authorization Request	35
5.2	Method of Incidental Taking	36
6	NUMBERS AND SPECIES EXPOSED	37
6.1	Introduction	37
6.2	Fundamentals of Sound	37
6.3	Effects of Pile Installation and Removal Activities	39
6.3.1	Description of Noise Sources	39
6.3.2	Sound Exposure Criteria and Thresholds	40
6.3.3	Limitations of Existing Noise Criteria	40
6.3.4	Ambient Noise	41
6.3.5	Distance to Sound Thresholds	42
6.3.6	Auditory Masking	61
6.3.7	Basis for Estimating Take by Harassment	61
6.3.8	Description of Take Calculation	63
6.3.9	Summary of Impacts for Pile Installation and Removal	71
6.4	Effects of Other Construction Activities	72
7	IMPACTS TO MARINE MAMMAL SPECIES OR STOCKS	74
7.1	Potential Effects of Pile Driving on Marine Mammals	75
7.1.1	Underwater Noise Effects	75
7.1.2	Airborne Noise Effects	77
7.2	Conclusions Regarding Impacts to Species or Stocks	77
8	IMPACT ON SUBSISTENCE USE	79
8.1	Subsistence Harvests by Northwest Treaty Indian Tribes	79
8.1.1	Harbor Seals	79
8.1.2	California Sea Lions	80
8.1.3	Gray Whales	80
8.2	Summary	81
9	IMPACTS TO THE MARINE MAMMAL HABITAT AND THE LIKELIHOOD OF RESTORATION	83

9.1	Pile Driving Effects on Potential Prey (Fish)	83
9.2	Pile Driving Effects on Potential Foraging Habitat	83
9.3	Summary of Impacts to Marine Mammal Habitat	84
10	IMPACTS TO MARINE MAMMALS FROM LOSS OR MODIFICATION OF HABITAT	85
11	MEANS OF EFFECTING THE LEAST PRACTICABLE ADVERSE IMPACTS – MITIGATION MEASURES	87
11.1	Mitigation for Pile Driving Activities	87
11.2	Mitigation Effectiveness	89
12	MINIMIZATION OF ADVERSE EFFECTS ON SUBSISTENCE USE	91
13	MONITORING AND REPORTING MEASURES	93
13.1	Monitoring Plan	93
13.1.1	Acoustic Measurements	93
13.1.2	Visual Marine Mammal Observations	95
13.1.3	Methods of Monitoring	95
13.1.4	Data Collection	96
13.2	Reporting	96
14	RESEARCH	99
15	LIST OF PREPARERS	100
16	REFERENCES	101

LIST OF FIGURES

Figure 1-1 Map of the Surrounding Vicinity	2
Figure 1-2 Proposed Project Area	3
Figure 1-3 Locations of the Proposed Construction Activities	5
Figure 1-4 Example Repairs to the EHW-1 Facility – Section A	6
Figure 2-1 NBK Bangor Restricted Areas.....	11
Figure 2-2 EHW Maximum Fetch Diagram.....	12
Figure 2-3 Sediment Sampling Locations.....	17
Figure 4-1 Pinniped Haulouts within the	29
Figure 6-1 Distance(s) (m) to Underwater Sound Thresholds for Cetaceans from Impact & Vibratory Pile Driving During Pile Installation.....	48
Figure 6-2 Distance(s) (m) to Underwater Sound Thresholds for Pinnipeds from Impact & Vibratory Pile Driving During Pile Installation.....	49
Figure 6-3 Distance(s) (m) to Underwater Sound Thresholds for all Marine Mammals from Vibratory Pile Driving During Steel Pile Removal	50
Figure 6-4 Distance(s) (m) to Underwater Sound Thresholds for all Marine Mammals from a Chipping Hammer During Concrete Pile Removal.....	51
Figure 6-5 Distance(s) (m) to Airborne Sound Thresholds for Pinnipeds (except harbor seals) from Impact & Vibratory Pile Driving During Pile Installation.....	55
Figure 6-6 Distance(s) (m) to Airborne Sound Thresholds for Harbor Seals from Impact & Vibratory Pile Driving During Pile Installation.....	56
Figure 6-7 Distance(s) (m) to Airborne Sound Thresholds for Pinnipeds (except Harbor Seals) from Vibratory Pile Driving During Steel Pile Removal	57
Figure 6-8 Distance(s) (m) to Airborne Sound Thresholds for Harbor Seals from Vibratory Pile Driving During Steel Pile Removal	58
Figure 6-9 Distance(s) (m) to Airborne Sound Thresholds for Pinnipeds (except Harbor Seals) from a Chipping Hammer During Concrete Pile Removal.....	59
Figure 6-10 Distance(s) (m) to Airborne Sound Thresholds for Harbor Seals from a Chipping Hammer During Concrete Pile Removal.....	60

LIST OF TABLES

Table 3-1 Marine Mammals Present in the Hood Canal in the Vicinity of NBK Bangor	19
Table 6-1 Definitions of Acoustical Terms	38
Table 6-2 Representative Noise Levels of Anthropogenic Sources	39
Table 6-3 Injury and Disturbance Thresholds for Underwater and Airborne Sounds	41
Table 6-4 Underwater Sound Pressure Levels Expected During Impact Installation Based on Similar In-situ Monitored Construction Activities.....	43
Table 6-5 Underwater Sound Pressure Levels Expected During Vibratory Installation Based on Similar In-situ Monitored Construction Activities.....	43
Table 6-6 Underwater Sound Pressure Levels Expected During Steel Pile Removal Based on Similar In-situ Monitored Construction Activities.....	44
Table 6-7 Underwater Sound Pressure Levels Expected During Concrete Pile Removal Based on Similar In-situ Monitored Construction Activities.....	44
Table 6-8 Calculated Distance(s) to and the Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Pile Installation	45
Table 6-9 Calculated Distance(s) to and the Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Steel Pile Removal	45
Table 6-10 Calculated Distance(s) to and the Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Concrete Pile Removal	46
Table 6-11 Actual Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Pile Installation	46
Table 6-12 Actual Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Steel Pile Removal	47
Table 6-13 Actual Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Concrete Pile Removal	47
Table 6-14 Airborne Sound Pressure Levels Expected During Impact Installation from Similar In-situ Monitored Construction Activities	53
Table 6-15 Airborne Sound Pressure Levels Expected During Vibratory Installation from Similar In-situ Monitored Construction Activities.....	53
Table 6-16 Airborne Sound Pressure Levels Expected During Steel Pile Removal from Similar In-situ Monitored Construction Activities.....	53
Table 6-17 Airborne Sound Pressure Levels Expected During Concrete Pile Removal from Similar In-situ Monitored Construction Activities.....	53
Table 6-18 Calculated Distances (m) to and the Area(s) Encompassed by the Marine Mammal Noise Thresholds In-Air from Pile Installation.....	54
Table 6-19 Calculated Distances (m) to and the Area(s) Encompassed by the Marine Mammal Noise Thresholds In-Air from Steel Pile Removal.....	54
Table 6-20 Calculated Distances (m) to and the Area(s) Encompassed by the Marine Mammal Noise Thresholds In-Air from Concrete Pile Removal.....	54

Table 6-21 Number of Potential Exposures of California Sea Lions within Various Acoustic Threshold Zones66

Table 6-22 Number of Potential Exposures of Harbor Seals within Various Acoustic Threshold Zones.....68

Table 6-23 Number Potential Exposures of Killer Whales within Various Acoustic Threshold Zones 69

Table 6-24 Number of Potential Exposures of Dall's Porpoise within Various Acoustic Threshold Zones.....70

Table 6-25 Number of Potential Exposures of Harbor Porpoise within Various Acoustic Threshold Zones.....71

Table 6-26 Summary of Potential Exposures for All Species During the EHW-1 Pile Replacement Project's Timeframe (July - Oct)72

ACRONYMS AND ABBREVIATIONS

BSS	Beaufort Sea State
°C	Celsius
CA	California
CISS	Cast in Steel Shells
CV	Coefficient of Variation
dB	Decibel
dBA	Decibel with A-weighting filter
DoN	Department of the Navy
EEZ	Exclusive Economic Zone
EHW	Explosive Handling Wharf
EIS	Environmental Impact Statement
ESA	Endangered Species Act
°F	Fahrenheit
FEIS	Final Environmental Impact Statement
FERC	Federal Energy Regulatory Commission
Ft.	Feet
Hz	Hertz
IHA	Incidental Harassment Authorization
kHz	Kilohertz
kg	Kilogram
km	Kilometer
lbs	Pounds
m	Meter
mi	Mile
MMO	Marine Mammal Observer
MMPA	Marine Mammal Protection Act
NAVBASE	Naval Base
NBK	Naval Base Kitsap
NMFS	National Marine Fisheries Service
OR	Oregon
Pa	Pascal
PDA	Pile Dynamic Analyzer
PSU	Practical Salinity Units
PTS	Permanent Threshold Shift
RMS	Root Mean Square
SEL	Sound Exposure Level
SFOBB	San Francisco-Oakland Bay Bridge
SPL	Sound Pressure Level
TL	Transmission Loss
TS	Threshold Shift
TTS	Temporary Threshold Shift
TRIDENT	Trident Fleet Ballistic Missile
U.S.	United States
USACE	United States Army Corp of Engineers
° W	West
WA	Washington
WDFW	Washington Department of Fish and Wildlife
WSDOT	Washington State Department of Transportation
ZOI	Zone of Influence

This Page Intentionally Left Blank

EXECUTIVE SUMMARY

In accordance with the Marine Mammal Protection Act (MMPA) of 1972, as amended, the U.S. Navy (Navy) is applying for an Incidental Harassment Authorization (IHA) to initiate the EHW-1 Pile Replacement Project at Naval Base Kitsap (NBK) Bangor to complete necessary repairs and maintenance to the EHW-1 facility. Six species of marine mammals are present within the waters surrounding NBK Bangor: the Steller sea lion (*Eumetopias jubatus*), the California sea lion (*Zalophus californianus*), the Harbor seal (*Phoca vitulina*), the Killer whale (*Orcinus orca*), the Dall's porpoise (*Phocoenoides dalli*), and the Harbor porpoise (*Phocoena phocoena*). These species may occur year-round in the Hood Canal, except the Steller sea lion which is only present from fall to late spring (Nov – June), outside of the project's timeline (16 July – 31 October). Additionally, while the Southern Resident killer whale (SRKW) is resident to the inland waters of Washington State and British Columbia it has not been observed in the Hood Canal in decades and was therefore excluded from further analysis. Only the five species which will be present during the project's timeline could be exposed to sound pressure levels associated with pneumatic chipping devices, and vibratory and impulsive pile driving and removal.

The proposed action includes the removal of the fragmentation barrier, walkway, and 138 steel and concrete piles. Of the piles requiring removal, ninety six are 24-inch diameter hollow pre-cast concrete piles which will be excised down to the mudline using a pneumatic chipping hammer. An additional three 24-inch steel fender piles, and thirty nine 12-inch steel fender piles will be extracted using a vibratory hammer. Also included in the repair work, is the installation of twenty eight new 30-inch diameter steel pipe piles, the construction of new cast-in-place pile caps (concrete formwork may be located below Mean Higher High Water [MHHW]), the installation of the pre-stressed superstructure, the installation of five sled mounted cathodic protection systems, and the installation/re-installation of related appurtenances. The proposed action will occur over two years.

All of the steel piles will be installed and/or removed with a vibratory hammer. Concrete piles will be removed using a pneumatic chipping hammer or other similar concrete demolition tool. During pile installation, depending on local site conditions, some piles may be driven (proofed¹) for the final few feet with an impact hammer. Impact pile driving will not occur on more than 5 days for the duration of any pile driving window during the implementation of the project and no more than one pile will be proofed in a given day. Furthermore, impact driving or proofing would be limited to 15 minutes per pile (up to 5 piles total per pile driving window). All piles driven with an impact hammer will be surrounded by a bubble curtain or other sound attenuation device over the full water column to minimize in-water noise. Vibratory pile driving is restricted to the time period between July 16 and October 31 while impact driving will only be performed between July 16th and September 30th during each construction window of this two year project. Non-pile driving, in-water work can be performed between July 16th and February 15th of each year.

For pile installation and removal activities, the Navy used NMFS promulgated thresholds for assessing pile driving impacts (NMFS 2005b, NMFS 2009), outlined in Section 6. The Navy used recommended spreading loss formulas (the practical spreading loss equation for underwater sounds,

¹ "Proofing" is driving the pile the last few feet into the substrate to determine the capacity of the pile. The capacity during proofing is established by measuring the resistance of the pile to a hammer that has a piston with a known weight and stroke (distance the hammer rises and falls) so that the energy on top of the pile can be calculated. The blow count in "blows per inch" is measured to verify resistance, and pile compression capacities are calculated using a known formula.

and the spherical spreading loss equation for airborne sounds) and empirically measured source levels from other similar pile driving events to estimate potential marine mammal exposures. Predicted exposures are outlined in Section 6. The modeling predicted no Level A harassments would occur associated with pile driving activities (including pneumatic chipping). The modeling predicts that 2,488 Level B harassments may occur during the Pile Replacement Project from underwater sound. No incidents of harassment were predicted from airborne sounds associated with pile driving (including pneumatic chipping). No incidents of harassment are anticipated from non-pile driving construction activities. Conservative assumptions (including marine mammal densities and other assumptions) used to estimate the exposures are likely to overestimate the potential number of exposures and their severity.

Pursuant to the Marine Mammal Protection Act (MMPA) Section 101(a)(5)(D)², the Navy submits this application to the National Marine Fisheries Service (NMFS) for an Incidental Harassment Authorization (IHA) for the incidental, but not intentional, taking of five marine mammal species during pile driving activities as part of the EHW-1 Pile Replacement Project between July 2011 and July 2012. The taking would be in the form of non-lethal, temporary harassment and is expected to have a negligible impact on these species. In addition, the taking would not have an unmitigable adverse impact on the availability of these species for subsistence use.

Regulations governing the issuance of incidental take under certain circumstances are codified at 50 Code of Federal Regulations (CFR) Part 216, Subpart I (Sections 216.101 – 216.108). Section 216.104 sets out 14 specific items that must be addressed in requests for take pursuant to Section 101 (a)(5)(D) of the MMPA. These 14 items are addressed in Sections 1 through 14 of this IHA application.

² 16 U.S.C. § 1371(a)(5); 50 C.F.R. Part 216, Subpart I.

1 DESCRIPTION OF ACTIVITIES

A detailed description of the specific activity or class of activities that can be expected to result in incidental taking of marine mammals.

1.1 Introduction


Naval Base Kitsap (NBK) Bangor, Washington is located on the Hood Canal approximately 20 miles west of Seattle, WA (Figure 1-1 and 1-2). NBK Bangor provides berthing and support services to United States (U.S.) Navy submarines and other fleet assets. The entirety of NBK Bangor, including the land areas and adjacent water areas in the Hood Canal are restricted from general public access. The Navy proposes to complete necessary repairs and maintenance at the Explosive Handling Wharf #1 (EHW-1) facility at NBK Bangor as part of a Pile Replacement Project to maintain the structure integrity of the wharf and ensure its continued functionality to support the operational requirements of the TRIDENT program. Sections 1.2 and 1.3 describe the proposed activities to be conducted in detail. The proposed actions with the potential to affect marine mammals within the waterways adjacent to NBK Bangor that could result in harassment under the Marine Mammal Protection Act (MMPA) of 1972, as amended in 1994, are pile installation and removal activities associated with the Pile Replacement Project.

1.2 Proposed Action

As part of the U.S. Navy's sea-based strategic deterrence mission, the Navy Strategic Systems Programs (SSP) directs research, development, manufacturing, test, evaluation, and operational support of the TRIDENT Fleet Ballistic Missile (TRIDENT) program. The proposed action (also called the Pile Replacement Project) for this harassment authorization request includes the removal of the fragmentation barrier, walkway, and 138 steel and concrete piles at EHW-1. Of the piles requiring removal, ninety six are 24-inch diameter hollow pre-cast concrete piles which will be excised down to the mudline. An additional three 24-inch steel fender piles, and thirty nine 12-inch steel fender piles will be extracted using a vibratory hammer. Also included in the repair work, is the installation of twenty eight new 30-inch diameter steel pipe piles. Additional construction activities will occur once the new steel piles are installed. This includes the installation of new cast-in-place pile caps, five sled mounted cathodic protection systems, and a new pre-stressed superstructure and related appurtenances. Sound propagation data will be collected through hydroacoustic monitoring during pile installation and removal to support environmental analyses for future repair work that may be necessary to maintain the EHW-1 facility. The presence of marine mammals will also be monitored during pile installation and removal.

1.3 Description of Pile Installation/Removal and Other Construction Activities

The EHW-1 Pile Replacement Project has been designed to restore the structural integrity of the EHW-1 facility which has been compromised due to the deterioration of the wharf's existing piling sub-structure. Under the proposed action, ninety six 24-inch diameter concrete piles, thirty nine 12-inch steel fender piles, and three 24-inch diameter steel fender piles will be removed. In addition, a total of twenty eight 30-inch diameter steel pipe piles will be installed and filled with concrete on the southwest corner of EHW-1. The proposed action will occur over a two year construction period scheduled to begin in July 2011.



(Source: Navy 2002; ESRI 2000)

Figure 1-1 Map of the Surrounding Vicinity




Figure 1-2 Proposed Project Area

The removal and installation of piles at EHW-1 is broken up into three components described in detail below and depicted in Figure 1-3.

The first component of this project would entail (Section A on Figure 1-3):

- The removal of one 24-inch diameter steel fender pile and its associated fender system components at the outboard support. A fender pile is set beside slips, wharves, etc., to guide approaching vessels and driven so as to yield slightly when struck in order to lessen the shock of contact. The fender system components are the components that attach the fender piles to the structure. These components are above the water line.
- The installation of sixteen 30-inch diameter hollow steel pipe piles (approximately 130 feet [40 meters] long). The piles would be installed to the tip elevation approximately 110 feet (34 meters [m]) (Mean Lower Low Water [MLLW]). This means that 100 feet of the pile will be below the MLLW mark.
- The construction of two cast-in-place concrete pile caps. The pile caps will be situated on the tops of the steel piles which are located directly beneath the structure (Figure 1-4) and function as a load transfer mechanism between the superstructure and the piles. Concrete formwork may be located below Mean Higher High Water [MHHW].
- The installation of three sled mounted passive cathodic protection systems would follow. The passive cathodic protection system is a metallic rod or anode that is attached to a metal object to protect it from corrosion. The anode is composed of more active metal which is more easily oxidized, corroding first and acting as a barrier against corrosion for the object to which it's attached. This system will be banded to the steel piles to prevent metallic surfaces of the wharf from corroding due to the saline conditions in Hood Canal.
- The piles will be removed/installed between July 16 and October 31 during each year of construction. The installation of the concrete pile caps and sled mounted passive cathodic protection systems will occur out of the water and will be installed on the tops of the piles themselves or attached the wharf's superstructure. While sound transmission from these activities could occur along the piles length and enter the water, this is expected to be minimal. However, to be cautious, these activities will occur between the window of July 16 to February 15 during each year of construction to minimize impacts to listed species, particularly fish.
- Figure 2-2 provides a diagram of Section A.

The second component of this project would require (Section B on Figure 1-3):

- The removal of two 24-inch diameter steel fender piles at the main wharf and associated fender system components.
- The installation of twelve 30-inch diameter hollow steel pipe piles (approximately 74-122 feet (23-37 m) long). The embedment depth of the piles would range from 30-50 feet (9-15 m).
- The construction of four concrete pile caps.
- The installation of a pre-stressed concrete superstructure. The superstructure is the pre-stressed concrete deck of the wharf found above, or supported by, the caps or sills, including the deck, girders, and stringers.
- The installation of two sled mounted passive cathodic protection systems.




Figure 1-3 Locations of the Proposed Construction Activities




Figure 1-4 Example Repairs to the EHW-1 Facility – Section A

- The installation/re-installation of related appurtenances would follow. Appurtenances are the associated parts of the superstructure that connects the superstructure to the piles. These pieces include all of the components such as bolts, welded metal hangers and fittings, brackets, etc.
- The piles will be removed/installed between July 16 and October 31 during each year of construction. The installation of the concrete pile caps, the concrete superstructure, sled mounted passive cathodic protection systems, will all occur out of the water and will be installed on the tops of the piles or attached to the wharf's superstructure. While sound transmission from these activities could occur along the piles length and enter the water, this is expected to be minimal. However, to be cautious, these activities will occur between the window of July 16 to February 15 during each year of construction to minimize impacts to listed species, particularly fish.

The last component of this project would be (Section C on Figure 1-3):

- The removal of the concrete fragmentation barrier and walkway. The walkway is used to get from the Wharf Apron to the Outboard Support. These structures will likely be removed by cutting the concrete into sections (potentially 3 or 4 total) using a saw, or other equipment, and removed using a crane. The crane will lift the sections from the existing piles and will be placed on a barge.
- The removal of the piles supporting the fragmentation barrier including:
 - Thirty nine 12-inch diameter steel fender piles,
 - Ninety six 24-inch diameter hollow pre-cast concrete piles cut to the mudline (includes 72 at fragmentation barrier, 4 at walkway, 4 at Bent 8 outboard support, and 8 at Bents 9 and 10).
- Concrete piles would be removed with a pneumatic chipping hammer or another tool capable of cutting through concrete. A pneumatic chipping hammer is similar to an electric power tool, such as a jackhammer, but uses the energy of compressed air instead of electricity. The pneumatic chipping hammer basically consists of a steel piston that is reciprocated (moved backward and forward alternately) in a steel barrel by compressed air. On its forward stroke the piston strikes the end of the chisel. The reciprocating motion of the piston occurs as such a rate that the chisel edge vibrates against the concrete with enough force to fragment or splinter the pile. The concrete debris would be captured using debris curtains/sheeting and removed from the project area.
- The piles will be removed between July 16 and October 31 during each year of construction. The removal of the fragmentation barrier and walkway will occur above the water. While sound transmission from these activities could occur along the piles length and enter the water, this is expected to be minimal. However, to be cautious, these activities will occur between the window of July 16 to February 15 during each year of construction to minimize impacts to listed species, particularly fish.

All concrete piles will be removed with a pneumatic chipping hammer or other similar device. All of the steel piles will be installed/removed with a vibratory hammer. During pile installation,

depending on local site conditions, some piles may be driven (proofed³) for the final few feet with an impact hammer. Per consultation with USFWS under the ESA, impact pile driving will not occur on more than 5 days for the duration of any pile driving window during the implementation of the project and no more than one pile will be proofed in a given day. Furthermore, impact driving or proofing would be limited to 15 minutes per pile (up to 5 piles total). Based on the Navy's experiences replacing piles during previous repair cycles at the EHW-1 facility, the Navy felt that this measure could be complied with because an impact pile driver has yet to be required to accomplish installation. Additionally, during typical construction projects impact proofing is only required on every 4-5 piles. All piles driven with an impact hammer will be surrounded by a bubble curtain or other sound attenuation device over the full water column to minimize in-water noise. Sound attenuation measures (*e.g.* bubble curtain) will be used during all impact hammer operations. The Navy will monitor hydroacoustic levels, as well as the presence and behavior of marine mammals during pile installation and removal. Section 11 provides the details proposed to reduce or mitigate the impacts from the proposed action.

³ "Proofing" is driving the pile the last few feet into the substrate to determine the capacity of the pile. The capacity during proofing is established by measuring the resistance of the pile to a hammer that has a piston with a known weight and stroke (distance the hammer rises and falls) so that the energy on top of the pile can be calculated. The blow count in "blows per inch" is measured to verify resistance, and pile compression capacities are calculated using a known formula.

2 DATES, DURATION, AND LOCATION OF ACTIVITIES

The dates and duration of such activity and the specific geographical region where it will occur.

2.1 Dates of Construction

The proposed action will occur over a two year construction period scheduled to begin in July 2011. This incidental harassment authorization (IHA) will cover only those activities which will occur between July 16, 2011 and July 15, 2012. A subsequent IHA will be required for the remaining work which will occur in the second construction window from July 16, 2012 through July 15, 2013. Vibratory pile driving is restricted to the time period between July 16th and October 31st while impact driving will only be performed between July 16th and September 30th of each construction window. Non-pile driving, in-water work can be performed between July 16th and February 15th of each year. These in-water timeframe restrictions were determined in consultation with NMFS NW region and USWFS under the ESA. All in-water construction activities within the Hood Canal are only permitted during July 16th – February 15th, and restricted at any other time of the year in order to protect spawning fish populations.

2.2 Duration of Activities

No work will begin on the proposed action until all required permits and approvals are in place. Under the proposed action, twenty eight 30-inch piles will be installed and 138 steel and concrete piles will be removed. The work will occur over a two year construction window scheduled to begin in July 2011. In-water construction will be limited to July 16 – Oct 31 of each year, therefore the potential duration of activities is 108 days per year or 216 days over the two year period.

The contractor estimates that steel pile installation and removal will occur at an average rate of two piles per day. For each pile installed, the driving time is expected to be no more than one hour for the vibratory portion of the project. The impact driving portion of the project is anticipated to take approximately 15 minutes per pile, with a maximum of 5 piles per construction window permitted to be impact driven. Impact pile driving will not occur on more than 5 days for the duration of any pile driving window and no more than one pile will be proofed in a given day. Steel piles will be extracted using a vibratory hammer. Extraction is anticipated to take approximately 30 minutes per pile. Concrete piles will be removed using a pneumatic chipping hammer or other similar concrete demolition tool. It is estimated that concrete pile removal could occur at a rate of five piles per day maximum, but removal will more likely occur at a rate of 3 piles per day. It is expected to take a couple of hours to remove each concrete pile with a pneumatic chipping hammer.

For steel piles, this results in a maximum of two hours of pile driving per pile or potentially four hours per day. For concrete piles, this results in a maximum of two hours of pneumatic chipping per pile, or potentially 6 hours per day. Therefore, while 216 days of in-water work time is proposed (108 days per construction period), only a fraction of the total work time per day will actually be spent pile driving. An average work day (two hours post-sunrise to two hours prior to sunset) is approximately 8-9 hours, depending on the month. While it is anticipated that only 4 hours of pile driving would be needed per day for steel piles, or 6 hours of pneumatic chipping would be needed for concrete piles, to take into account deviations from the estimated times for

pile installation and removal, the Navy modeled potential impact as if the entire day could be spent pile driving.

Based on the proposed action, the total time from vibratory pile driving during steel pile installation would be approximately 14 days (28 piles at an average of 2 per day). The total time from impact pile driving during steel pile installation would be 5 days (5 piles at 1 per day). The total time from vibratory pile driving during steel pile removal would be 21 days (42 piles at an average of 2 per day). The total time from using a pneumatic chipping hammer during concrete pile removal would be 32 days (96 piles at an average of 3 per day).

2.3 Study Area Description

EHW-1 is located along the eastern shoreline of Hood Canal in Kitsap County, WA. The Hood Canal is a long, narrow fjord-like basin of the western Puget Sound. Throughout its 67-mile length, the width of the canal varies from 1-2 miles and exhibits strong depth/elevation gradients and irregular seafloor topography in many areas. Although no official boundaries exist along the waterway, the northeastern section of the canal extending from the mouth of the canal at Admiralty Inlet to the southern tip of Toandos Peninsula is referred to as the northern Hood Canal. The proposed Project Area is located within this region. The wharf is a U-shaped concrete structure built in 1977 for ordnance handling operations in support of the Trident Submarine squadron home ported at NBK Bangor. EHW-1 consists of two 30 meter (100 ft) access trestles and a main pier deck which measures approximately 215 meters (700 ft) in length. The wharf is supported by both 16-inch and 14-inch hollow octagonal pre-cast concrete piles. Additionally, there are steel and timber fender piles on the outboard and inboard edges of the wharf.

Two restricted areas are associated with NBK Bangor, Naval Restricted Areas 1 and 2 (33 CFR 334.1220), which are depicted in Figure 2-1 relative to the Project Area. The regulations associated with Naval Restricted Area 1 indicated that no persons or vessels shall enter this area without permission from the Commander, Naval Submarine Base Bangor, or his/her authorized representative. The regulations associated with Naval Restricted Area 2 indicate that Navigation will be permitted within that portion of the circular area not lying within Area 1 at all times except when magnetic silencing operations are in progress.

2.3.1 Bathymetric Setting

In the northern Hood Canal, water depths in the center of the waterway near Admiralty Inlet vary between 300 to 420 feet. As the canal extends southwestward toward the Olympic Mountain Range and Thorndyke Bay, water depths shoal to approximately 160 feet over a moraine deposit. This deposit forms a sill across the short axis of the canal in the vicinity of Thorndyke Bay, which has an important impact on deep circulation and seawater exchange. The NBK Bangor waterfront occupies approximately 5 miles of the shoreline within northern Hood Canal (1.7 percent of the entire Hood Canal coastline) and lies just south of the sill feature. Depths of the in-water project site are provided in Figure 2-2. The width of the canal is approximately 1.5 miles at the site, 2.2 miles at the northern end of NBK Bangor, and constricts to approximately 1.1 miles near the southern end near Brown Point. The furthest direct line of site from the project site is 8.4 miles to the north and 4.2 miles to the south (see Figure 2-2).

2.3.2 Tides

The tides in Hood Canal are mixed, diurnal-semidiurnal with a range directly dependent upon the phase and alignment of the lunar and solar gravitational influences on the regional tides (URS





Figure 2-1 NBK Bangor Restricted Areas



1994; Morris *et al.* 2008). The astronomic influences (tides) on water level within Puget Sound and Hood Canal result in one flood and one ebb tidal event with a small to moderate range (1 to 6 feet) and a second flood and second ebb with a larger range (8 to 16 feet) during a 24-hour and 50-minute tidal day. As a result, higher high, lower high, higher low, and lower low water levels are recorded within each tide day. Since the tides within Hood Canal are mixed diurnal to semi-diurnal, this body of water is subject to one major flushing event per tide day when approximately 1.1326×10^9 cubic yards (or 3 percent of the total canal volume) is exchanged over a 6-hour period. Due to the wide range of tidal heights that can occur in this body of water, the actual seawater exchange volume for Hood Canal ranges from 1 percent during a minor tide to 4 percent during a major tide.

Despite considerable tidally driven seawater influx within the basin, some studies have estimated water residence time in the southern and middle portions of Hood Canal can be up to one year due to the natural limitation on seawater exchange (i.e., bathymetry; Warner *et al.* 2001; Warner 2007). However, at the project site, the majority of the daily volume of seawater exchange flows directly across the NBK Bangor waterfront area. As a result, the degree of flushing that occurs at the Project Area is relatively high and the characteristics of this seawater more closely track the physical, chemical, and biological conditions of Puget Sound than southern Hood Canal.

2.3.3 Circulation and Currents

Tidal currents and resulting circulation patterns within Hood Canal are complex due to the configuration of the basin, as well as the mixed diurnal-semidiurnal tidal regime. Current measurements obtained from the reaches of northern Hood Canal in the summer of 2007 indicate that tidal phase and range have a significant impact on the velocity of currents associated with the flood and ebb tides (Morris *et al.* 2008). The larger tidal ranges promote higher velocity currents and increased flushing of the basin, while small to moderate tidal ranges yield a diminished tidal current regime and limit the volume of seawater exchange between Hood Canal and Puget Sound. Seawater that enters the canal from Puget Sound during an incoming flood tide tends to be cooler, more saline, and well-oxygenated relative to the Hood Canal waters. As a result, the incoming Puget Sound water has a tendency to sink to the bottom of the canal as it flows over the sill and move south during each flood tide, while the lower density Hood Canal water tends to remain in the upper water column.

Current flow (speed and direction) at the Project Area is primarily a function of tidal action based on the phase and range of each tide within the mixed diurnal-semidiurnal regime, and current velocities in the shallower water areas (less than 50 feet) around the Project Area are variable and complex. The magnitude or instantaneous velocity of these fluctuating water column currents range from 0 to 0.88 ft/sec within the 30- to 65-foot water depth interval. However, current flow in any one direction is short-lived and inconsistent in magnitude, with relatively few periods of time when sufficient energy (0.7 ft/sec) exists to exceed the threshold for re-suspending deposits of unconsolidated material on the seafloor (Boggs 1995). Statistical summaries show that time-averaged net flow is within the 0.07 to 0.10 ft/sec range in the upper water column and less than 0.03 ft/sec in proximity to the seafloor.

The nearshore current observations at the Project Area and other NBK Bangor piers and wharves in the summer of 2006 suggest that tidal currents were inconsistent with water level (tide) measurements. Rather than the typical relationship where maximum current corresponds to mid-flood or mid-ebb in the water level record, maximum flow velocities at the Project Area align

with water levels at the high and low tide. Furthermore, the direction of nearshore flow often ran counter to expectations in a normal system, with flood tide coinciding with northeastward currents and ebb tide resulting in southwesterly currents (Morris et al. 2008).

2.3.4 Sea State

Apart from larger impacts associated with large-scale changes in weather and ocean circulation in the Pacific Basin, seasonal variability in Hood Canal circulation can occur in the winter, when strong meteorological events (e.g., storms, high winds) are more prevalent. Regardless of direction, winds with velocities in excess of 25 knots occur relatively infrequently in the Puget Sound region (Morris et al. 2008). The typically light winds afforded by the surrounding highlands (Olympic and Cascade Mountain Ranges) coupled with the fetch-limited environment of Hood Canal result in relatively calm wind conditions throughout most of the year. However, the northern and middle sections of Hood Canal are oriented in the southwest to northeast direction. Therefore, organized coastal storm events that reach land in the late autumn and winter months, as well as fair weather systems in the spring and summer exhibiting wind speeds in excess of 20 knots, have the capability to generate substantial wind waves due to increased fetch and/or alter normal tidal flow within the basin.

However, the Project Area is afforded some protection by the coastline of both Kitsap and Toandos Peninsulas (see Figure 2-2). Using a maximum fetch of 8.4 miles between the Project Area and the north shore of Thorndyke Bay to the north-northeast, estimates indicate that a 20-knot sustained wind has the capability to generate average wave heights of 1.9 feet (Beaufort Sea State [BSS] of 2) and a 30-knot wind event could produce wave heights of 3.1 feet (BSS = 3) (CERC 1984). The maximum fetch to the southwest is one-half that to the northeast (4.2 miles), and could yield average waves of 1.3 feet in height (BSS = 2) in a 20-knot wind, and 1.9 feet (BSS = 2) in a 30 knot wind. Maximum wave heights that would be expected in these weather conditions would actually be 67 percent higher than average estimates reported above. Thus, a weather event capable of generating waves with an average height of 3.1 feet (BSS = 3) could also yield waves with maximum heights of 5.1 feet (BSS = 4) (CERC 1984).

2.3.5 Water Temperature

Water temperatures in the Strait of Juan de Fuca and Puget Sound typically range from 44 to 46 degrees Fahrenheit (°F) throughout the winter months (mid-December through mid-March). Surface waters slowly warm throughout the spring and summer due to increased solar heating, reaching temperatures of 50°F in mid-May or early June to a maximum temperature of 54°F during the month of August. Beginning in September, water temperatures begin to decrease over time, falling 6 to 8°F over the next 3 months due to decreasing levels of solar radiation. Occasionally, anomalies in this pattern of heating and cooling are detected in the data record, but are often short in duration (1 to 2 weeks). Monthly mean water temperatures along the NBK Bangor waterfront are summarized in Table 2-1. Nearshore areas (water depths range from 1-60 m) are susceptible to greater temperature variations due to seasonal fluxes in solar radiation input.

2.3.6 Dissolved Oxygen

Concentrations of dissolved oxygen (DO) in extraordinary quality marine surface waters should exceed 7.0 mg/L of DO, allowing for only 0.2 mg/L reductions in the natural condition by human-caused activities (WAC 173-201A). According to the WDOE Marine Water Quality

Table 2-1. Monthly Mean Surface Water Temperatures (°C/°F)

SAMPLING MONTH (2005, 2006) ¹	NEARSHORE TEMPERATURE	OFFSHORE TEMPERATURE
July 2005	14.3°C (57.8°F)	11.6°C (52.9°F)
August 2005	13.8°C (56.8°F)	13.5°C (56.3°F)
September 2005	14.9°C (58.8°F)	11.6°C (52.9°F)
January 2006	8.2°C (46.8°F)	---
February 2006	8.1°C (46.6°F)	---
March 2006	8.5°C (47.3°F)	8.3°C (46.9°F)
April 2006	9.6°C (49.3°F)	9.3°C (48.7°F)
May 2006	10.9°C (51.6°F)	11.0°C (51.8°F)
June 2006	13.2°C (55.8°F)	---

Source: Phillips et al. 2009.

Data are from 13 nearshore and 4 offshore stations along the NAVBASE Kitsap Bangor waterfront. Those stations near the EHW-2 project site are shown in Figure 3-4.

--- No data were collected at this depth during this sampling month

Report for 1998 to 2000, fish are negatively affected by DO concentrations of less than 4.5 mg/L (Newton et al. 2002). Data from WDOE's Marine Water Quality Monitoring Program for 1998 to 2000 and the Hood Canal Dissolved Oxygen Program (HCDOP) for 2002 to 2004 show that Hood Canal is particularly susceptible to low DO levels (Newton et al. 2002; HCDOP 2005). The 2008 Clean Water Act Section 303(d) list, the most recent list approved by the U.S. Environmental Protection Agency (USEPA), includes five segments within northern Hood Canal impaired by low DO levels (WDOE 2009a). Two of these segments are located along the NBK Bangor waterfront. The low DO for both of those segments is believed to be due to or influenced by human actions (WDOE 2009a). However, these stations are offshore in deep water and would not necessarily be representative of nearshore conditions at the NBK Bangor waterfront.

Although some waters along the NBK Bangor waterfront are on the 303(d) list, mean DO measurements during July 2005 through June 2006 indicate that nearshore stations at the NBK Bangor waterfront consistently met extraordinary quality standards for DO. From July 2005 through June 2006 and January 2007 through April 2008, DO levels met the extraordinary standard for surface waters (0 to 20 feet in depth) year round and for deep water (66 to 197 feet in depth) most of the year (deeper waters can drop to only a fair standard for DO in late summer). In late summer-early fall, DO levels in the action area drop from typical ranges of approximately 6 to 10 mg/L to a range of 4.7 to 9.1 mg/L (Phillips et al. 2009). The variation in mean DO measurements for deeper waters (66 to 197 feet in depth) near the project site was consistent with DO patterns within the rest of Hood Canal. During the late summer and early fall period (July through September 2005), mean DO measurements met fair to excellent quality standards. At 66 to 197 feet in depth, these measurements are on the upper range of low DO conditions measured historically throughout Hood Canal during the late summer and fall periods (Warner 2007). Mean DO measurements at 66 to 197 feet in depth from March through May 2006 met Extraordinary Quality standards.

2.3.7 Stratification and Salinity

The waters of Hood Canal surrounding the Project Area are stratified, with less saline, warmer water overlying colder, more saline bottom waters. The salinity of the upper water layer is sensitive to the amount of freshwater input and may become more diluted during heavy precipitation (URS 1994). Variances due to seasonal changes (such as freshwater input, wind-induced mixing, and solar heating) are common (URS 1994).

Freshwater input into Hood Canal comes from creeks, rivers, groundwater (including artesian wells [deep underground aquifer]), and stormwater outfalls. The freshwater inputs affect the salinity in Hood Canal. Artesian wells also contribute to freshwater inputs, with estimated flows of 2,000 to 2,500 gallons per minute (WDOE 1981). Overland flow from much of the western portion of NBK Bangor is routed to Hood Canal through a series of stormwater outfalls. Saltwater and freshwater mixing zones exist at the mouths of each of these streams and outfalls (URS 1994).

Between June 2005 and July 2006, surface water salinity levels along the NBK Bangor waterfront ranged from 26 to 35 practical salinity units (PSU) (Phillips et al. 2009). Salinity measurements with depth reflected a stratified water column, with less saline surface water overlying cooler saline water at depth. The transition between the lower salinity surface waters and higher salinity subsurface waters occurred at a depth of about 33 feet (Phillips et al. 2009). The lowest surface water salinity (26.7 PSU) was measured in January 2006 when input from fresh water may have been high due to winter storms and runoff. The range of salinity along the NBK Bangor waterfront is typical for marine waters in Puget Sound (Newton et al. 1998, 2002).

2.3.8 Sediments

Existing sediment information is based on results from sampling near the Project Area during 2007 (Hammermeister and Hafner 2009); sampling locations are shown in Figure 2-3. Sediment quality at the project site is generally good; levels of contaminants meet applicable state standards. Marine sediments are composed of gravelly sands with some cobbles in the intertidal zone, transitioning to silty sands in the subtidal zone (Hammermeister and Hafner 2009). Subsurface coring studies conducted in 1994 found the presence of glacial till approximately 6 feet below mud line in the intertidal zone, increasing to over 10 feet in the subtidal zone (URS 1994). The composition of sediment samples from the Project Area ranged from 65 to 100 percent for sand, less than 1 to 7 percent for gravel, 2 to 32 percent silt, and 2 to 11 percent clay.

2.3.9 Ambient Underwater Soundscape

Underwater ambient noise at the Project Area is widely variable over time due to a number of natural and anthropogenic sources. A number of sources of underwater sound exist in the vicinity of the Project Area. Sources of naturally caused underwater noise include wind, waves, precipitation, and biological sources (such as shrimp, fish, and cetaceans). Noise derived from biological organisms can be absent or dominant over narrow and broad frequency ranges. Precipitation can contribute up to 35 dB to the existing sound level, and increases in wind speed of 5 to 10 knots can cause a 5 dB increase in ambient ocean noise across most frequencies (Urick 1983). The highest noise levels occur in nearshore areas where the sound of surf can increase underwater noise levels by 20 dB or more within 200 yards from the surf zone in the 200 Hz to 2 kHz regime (Wilson et al. 1985). In addition, wakes from boat traffic causes breaking waves in the surf zone.



Figure 2-3 Sediment Sampling Locations



Projection: Lambert Conformal Conic
Datum: NAD 83

0 75 150 300 450 600 Feet

There is also human-generated noise from ship or boat traffic and other mechanical sources (Urick 1983). Small powerboats generate peak narrow band sound pressure levels of 150 to 165 dB re 1 μ Pa at 3 feet in the 350 to 1,200 Hz region, with mean sound pressure levels of 148 dB re 1 μ Pa at 3 feet (Barlett and Wilson 2002). Fishing vessels can generate peak spectral densities of 140 dB re 1 μ Pa at 3 feet in the 250 to 1,000 Hz regime (Hildebrand 2007). Underwater sound from human activities includes ship traffic noise, use of sonar and echo sounders in commercial fishing to locate fish schools, industrial ship noise, and recreational boat use. Ship and small boat noise comes from propellers and other on-board rotating equipment. Other sources of underwater noise at industrial waterfronts could come from cranes, generators, and other types of mechanized equipment on wharves or the adjacent shoreline.

Slater (2009) measured ambient noise levels along the NBK waterfront from several locations over an approximately one month period from July 10, 2007 – Aug 14, 2007. The location closest to the Project Area, designated as Mid-Point in the report, recorded data from two hydrophones deployed southwest of the existing EHW-1 facility. Recordings were made 5 minutes per hour throughout the entire study period. Average broadband ambient noise levels measured in the vicinity of the Project Area were 114 dB re 1 μ Pa between 100 Hz and 20 kHz (Slater 2009). Peak spectral noise from industrial activity was noted below the 300 Hz frequency, with maximum levels of 110 dB re 1 μ Pa noted in the 125 Hz band. In the 300 Hz to 5 kHz range, average levels ranged between 83 and 99 dB re 1 μ Pa. Wind-driven wave noise dominated the background noise environment at approximately 5 kHz and above, and ambient noise levels flattened above 10 kHz. The primary source of noise was due to industrial activity along the waterfront (such as at EHW-1, Marginal Wharf, and Delta Pier), small boat traffic, and wind-driven wave noise. No substantial precipitation was noted during the study period, although this noise would be undoubtedly present during seasonal periods.

Carlson et al. (2005) measured the underwater baseline noise at Hood Canal Bridge and found that underwater noise levels ranged from 115 to 135 dB re 1 μ Pa. The Washington State Department of Transportation (WSDOT) summarized underwater noise at ferry terminals with no construction activity as ranging from 80 to 90 dB at the Everett Home Port, 145 dB at Mukilteo ferry terminal, and 131 to 136 dB (peak levels) at Friday Harbor (WSDOT 2007), which demonstrates the range over which localized anthropogenic noise can vary by specific locations and time periods. Average underwater broadband noise levels measured at the Project Area, inclusive of existing human activities but in the absence of construction activities, fell within the minimum and maximum range of measurements taken at similar environments within Puget Sound. For the purposes of further noise analyses, the average background underwater noise levels at the Project Area were considered to be 114 decibels (dB) re 1 μ Pa between 100 hertz (Hz) and 20 kilohertz (kHz).

3 MARINE MAMMAL SPECIES AND NUMBERS

The species and numbers of marine mammals likely to be found within the activity area.

There are six marine mammal species, three cetaceans and three pinnipeds, which may inhabit or transit through the waters nearby NBK Bangor in the Hood Canal. These include the transient killer whale, harbor porpoise, Dall's porpoise, Steller sea lion, California sea lion, and the harbor seal. While the Southern Resident killer whale (SRKW) is resident to the inland waters of Washington State and British Columbia, it has not been observed in the Hood Canal in decades, and therefore was excluded from further analysis. The Steller sea lion is the only marine mammal that occurs within the Hood Canal which is listed under the Endangered Species Act (ESA), though it is unlikely to be present during the project duration (July – Oct); The U.S. Eastern stock/Distinct Population Segment (DPS) is listed as threatened. All marine mammal species are protected under the Marine Mammal Protection Act (MMPA). Section 3 summarizes the population status and abundance of these species, while Section 4 contains detailed life history information. Table 3-1 lists the marine mammal species that occur in the vicinity of NBK Bangor and their estimated densities within the Project Area.

Table 3-1 Marine Mammals Present in the Hood Canal in the Vicinity of NBK Bangor

SPECIES	STOCK(S) ABUNDANCE ¹	RELATIVE OCCURRENCE IN HOOD CANAL, WASHINGTON	SEASON(S) OF OCCURRENCE	DENSITY IN THE WORK WINDOW (INDIVIDUALS PER KM ²) ^a
Steller sea lion <i>Eumetopias jubatus</i> Eastern U.S. stock/DPS	45,095 – 55,832 ²	Rare to occasional use	Fall to late spring (Nov – mid April)	0.00
California sea lion <i>Zalophus californianus</i> U.S. Stock	238,000 ⁴	Common	Fall to late spring (Aug – May)	0.410 ^b
Harbor seal <i>Phoca vitulina</i> WA inland waters stock	14,612 ³ (CV = 0.15)	Common	Year-round; resident species in Hood Canal	1.31 ^c
Killer whale <i>Orcinus orca</i> West Coast transient stock	314 ⁵	Rare to occasional use	Year-round	0.038 ^d
Dall's porpoise <i>Phocoenoides dalli</i> CA/OR/WA stock	48,376 ³ (CV = 0.24)	Rare to occasional use	Year-round	0.043 ^e
Harbor porpoise <i>Phocoena phocoena</i> WA inland waters stock	10,682 ³ (CV=0.38)	Rare to occasional use	Year-round	0.011 ^e

Sources: ¹ NMFS marine mammal stock assessment reports at: <http://www.nmfs.noaa.gov/pr/sars/species.htm> ² Angliss and Outlaw, 2009; ³ Carretta *et al.*, 2008; ⁶ Carretta *et al.*, 2007; ⁷ Allen and Angliss, 2010; ^a Work Window refers to the period from July - Oct; ^b DoN, 2010a; ^c Jeffries *et al.*, 2003 and Huber *et al.*, 2001; ^d London, 2006; ^e Agness and Tannenbaum 2009a.

3.1 ESA-Listed Marine Mammals

3.1.1 Steller Sea Lion

Species Description

Steller sea lions are the largest members of the Otariid (eared seal) family. Steller sea lions show marked sexual dimorphism, in which adult males are noticeably larger and have distinct coloration patterns from females. Males average approximately 1,500 lbs and 10 feet in length; females average about 700 lbs and 8 feet in length. Adult females have a tawny to silver-colored pelt. Males are characterized by dark, dense fur around their necks that appears like a mane and light tawny coloring over the rest of their body (NMFS 2008).

There are two distinct populations of Steller sea lions based on genetics and population trends, separated at 144°W longitude (Loughlin 1997; Angliss and Outlaw 2005). The Eastern U.S. stock, which is the population which may occur within the Study Area, includes the animals east of Cape Suckling, Alaska (144°W) (NMFS 1997; Loughlin 2002; Angliss and Outlaw 2005). Steller sea lions west of 144°W longitude residing in the central and western Gulf of Alaska, Aleutian islands, as well as those that inhabit coastal waters and breed in Asia (e.g. Japan and Russia) are part of the Western U.S. Stock. The Eastern U.S. stock breeds on rookeries (places where they give birth and mate) located in southeast Alaska, British Columbia, Oregon, and California; there are no rookeries located in Washington.

Population Abundance

The U.S. Eastern stock was estimated to number between 46,000 and 58,000 animals in 2002, and has been increasing approximately 3 percent per year since the late 1970s (NMFS 2008; Pitcher *et al.* 2007). The most recent population estimate for the Eastern North Pacific stock of the Steller sea lion, which occurs along the WA coast and Puget Sound, is 48,519 individuals (Angliss and Outlaw 2008). The U.S. Eastern stock is stable or increasing throughout the northern portion of its range (Southeast Alaska and British Columbia) and stable or increasing slowly in the central portion of its range (Oregon through central California). Since the mid-1970s the annual rate of increase has been approximately 3 percent (Angliss and Outlaw 2008). Although the stock size has increased, the status of this stock relative to its optimum sustainable population (OSP) is unknown (Angliss and Outlaw 2008).

3.2 Non-ESA Listed Marine Mammals

3.2.1 California Sea Lion

Species Description

California sea lions are also members of the Otarrid family. The species *Zalophus californianus* includes three subspecies: *Z. c. wollebaeki* (on the Galapagos Islands), *Z. c. japonicus* (in Japan, but now thought to be extinct), and *Z. c. californianus* (found from southern Mexico to southwestern Canada; referred to here as the California sea lion) (Carretta *et al.* 2007).

The California sea lion is sexually dimorphic. Males may reach 1,000 pounds and 8 feet in length; females grow to 300 pounds and 6 feet in length. Their color ranges from chocolate brown in males to a lighter, golden brown in females. At around 5 years of age, males develop a bony bump on top of the skull called a sagittal crest. The crest is visible in the “dog-like” profile of male sea lion heads, and hair around the crest gets lighter with age.

Population Abundance

The U.S. stock of California sea lions may occur in the marine waters nearby NBK Bangor. The estimated stock is 238,000 and the minimum population size of this stock is 141,842 individuals (Carretta *et al.* 2007). These numbers are from counts during the 2001 breeding season of animals that were ashore at the four major rookeries in southern California and at haulout sites north to the Oregon/California border. Sea lions that were at-sea or hauled out at other locations were not counted (Carretta *et al.* 2007). An estimated 3,000 to 5,000 California sea lions migrate to Washington and British Columbia waters during the non-breeding season from September to May (Jeffries *et al.* 2000). Peak numbers of up to 1,000 sea lions occur in Puget Sound (including Hood Canal) during this time period (Jeffries *et al.* 2000).

3.2.2 Harbor Seal

Species Description

Harbor seals, which are members of the family Phocidae (“true seals”), inhabit coastal and estuarine waters and shoreline areas from Baja California to western Alaska. For management purposes, differences in mean pupping date (i.e., birthing) (Temte 1986), movement patterns (Jeffries 1985; Brown 1988), pollutant loads (Calambokidis *et al.* 1985) and fishery interactions have led to the recognition of three separate harbor seal stocks along the west coast of the continental U.S. (Boveng 1988). The three distinct stocks are: 1) inland waters of Washington State (including Hood Canal, Puget Sound, and the Strait of Juan de Fuca out to Cape Flattery), 2) outer coast of Oregon and Washington, and 3) California (Carretta *et al.* 2007). The inland waters of Washington state stock is the only stock that is expected to occur within the Study Area.

The average weight for adult seals is about 180 pounds and males are slightly larger than females. Male harbor seals weight up to 245 lbs and measure approximately 5 feet in length. The basic color of harbor seals’ coat is gray and mottled but highly variable, from dark with light color rings or spots to light with dark markings (NMFS 2008c).

Population Abundance

Estimated population numbers for the inland waters of Washington, including the Hood Canal, Puget Sound, and the Strait of Juan de Fuca out to Cape Flattery, are 14,612 (CV = 0.15) individuals (Carretta *et al.* 2007). The Harbor seal is the only species of marine mammals that is consistently abundance and considered resident in the Hood Canal (Jeffries *et al.* 2003). The population of harbor seals in Hood Canal is a closed population, meaning they do not have much movement outside of Hood Canal (London 2006). The abundance of harbor seals in Hood canal has stabilized, and the population may have reached its carrying capacity in the mid-1990s with an approximate abundance of 1,000 harbor seals (Jeffries *et al.* 2003).

3.2.3 Killer Whale

Species Description

Killer whales are members of the Delphinid family and are the most widely distributed cetacean (*e.g.* whales, dolphins, and porpoises) species in the world. Killer whales have a distinctive color pattern, with black dorsal (top) and white ventral (bottom) portions. They also have a conspicuous white patch above and behind the eye and a highly variable gray or white saddle area behind the dorsal fin. The species shows considerable sexual dimorphism. Adult males

develop larger pectoral flippers, dorsal fins, tail flukes, and girths than females. Male adult killer whales can reach up to 32 ft in length and weight nearly 22,000 lbs (10,000 kg); females reach 28 ft in length and weigh up to 16,500 lbs (7,500 kg).

Based on appearance, feeding habits, vocalizations, social structure, and distribution and movement patterns there are three types of populations of killer whales (Wiles 2004; NMFS 2005). The three distinct forms or types of killer whales recognized in the North Pacific Ocean are: 1) Residents, 2) Transients, and 3) Offshores. The resident and transient populations have been divided further into different subpopulations based mainly on genetic analyses and distribution; not enough is known about the offshore whales to divide them into subpopulations (Wiles, 2004).

Within the transient ecotype, association data (Ford et al. 1994, Ford and Ellis 1999, Matkin et al. 1999), acoustic data (Saulitis 1993, Ford and Ellis 1999) and genetic data (Hoelzel et al. 1998, 2002; Barrett-Lennard 2000) confirms that three communities of transient whales exist and represent three discrete populations: 1) Gulf of Alaska, Aleutian Islands, and Bering Sea transients, 2) AT1 transients, and 3) West Coast transients. Among the genetically distinct assemblages of transient killer whales in the northeastern Pacific, only the West Coast transient stock, which occurs from southern California to southeastern Alaska, may occur in the Project Area.

Population Abundance

The West Coast Transient stock is a trans-boundary stock, with minimum counts for the population of “transient” killer whales coming from various photographic datasets. Combining these counts of cataloged “transient” whales gives a minimum number of 314 individuals for the West Coast Transient stock (Allen and Angliss 2010). However, the number in Washington waters at any one time is probably fewer than 20 individuals (Wiles 2004).

3.2.4 Dall's Porpoise

Species Description

Dall's porpoises are members of the Phocoenid (porpoise) family and are common in the North Pacific Ocean. They can reach a maximum length of just under 8 ft and weight up to 480 lbs. Males are slightly larger and thicker than females, which reach lengths of just under 7 ft long. The body of Dall's porpoises is a very dark gray or black in coloration with variable contrasting white “throracic” panels and white “frosting” on the dorsal fin and tail that distinguish them from other cetacean species. These markings and colorations vary with geographic region and lifestage, with adults having more distinct patterns.

Based on NMFS stock assessment reports, Dall's porpoises within the Pacific U.S. Exclusive Economic Zone (EEZ) are divided into two discrete, noncontiguous areas: 1) waters off California, Oregon, and Washington, and 2) those in Alaskan waters (Carretta *et al.* 2008). Only individuals from the CA/OR/WA stock may occur within the Project Area.

Population Abundance

The NMFS population estimate, recently updated in 2008 for the California/Oregon/Washington stock, is 48,376 (CV – 0.24) which is based on vessel line transect surveys by Barlow and Forney (2007) and Forney (2007) (Carretta *et al.* 2008). Additional numbers of Dall's porpoise occur in the inland waters of WA state, but the most recent estimate obtained in 1996 (900

animals; CV = 0.40) is over 10 years old (Calambokidis et al. 1997) and is not included in the overall estimate of abundance for this stock due to the need for more up-to-date information.

3.2.5 Harbor Porpoise

Species Description

Harbor porpoises belong to the Phocoenid (porpoise) family and are found extensively along the Pacific U.S. coast. Harbor porpoises are small with males reaching average lengths of approximately 5 ft; Females are slightly larger with average length of 5.5 ft. The average adult harbor porpoise weights between 135 and 170 lbs. Harbor porpoises have a dark grey coloration on their backs, which their belly and throats are white. They have a dark grey chin patch and intermediate shades of grey along their sides.

Recent preliminary genetic analyses of samples ranging from Monterey, CA to Vancouver Island, BC indicate that there is small-scale subdivision within the U.S. portion of this range (Chivers *et al.* 2002). Although geographic structure exists along an almost continuous distribution of harbor porpoise from California to Alaska, stock boundaries are difficult to draw because any rigid line is (to a greater or lesser extent) arbitrary from a biological perspective. Nevertheless, based on genetic data and density discontinuities identified from aerial surveys, NMFS identifies 8 stocks in the Northeast Pacific Ocean. Pacific coast harbor porpoise stocks include: 1) a Monterey Bay stock, 2) a San Francisco-Russian River stock, 3) a northern California/southern Oregon stock, 4) an Oregon/Washington coast stock, 5) an Inland Washington stock, 6) a Southeast Alaska stock, 7) a Gulf of Alaska stock, and 8) a Bering Sea stock. Only individuals from the Inland waters of Washington stock may occur in the Project Area.

Population Abundance

Aerial surveys of the inside waters of Washington and southern British Columbia were conducted during August of 2002 and 2003 (J. Laake, unpubl. data). These aerial surveys included the Strait of Juan de Fuca, San Juan Islands, Gulf Islands, and Strait of Georgia, which includes waters inhabited by the Washington Inland Waters stock of harbor porpoise as well as harbor porpoise from British Columbia. An average of the 2002 and 2003 estimates of abundance in U.S. waters resulted in an uncorrected abundance of 3,123 (CV= 0.10) harbor porpoises in Washington inland waters (J. Laake, unpubl. data). When corrected for availability and perception bias, using a correction factor of 3.42 ($1/g(0)$; $g(0)=0.292$, CV=0.366) (Laake et al. 1997), the estimated abundance for the Washington Inland Waters stock of harbor porpoise is 10,682 (CV=0.38) animals (Carretta *et al.* 2008).

3.3 Marine Mammal Modeling Parameters

3.3.1 Spatial Distribution

Density assumes that marine mammals are uniformly distributed within a given area, although this is rarely the case. Marine mammals are usually clumped in areas of greater importance, for example, areas of high productivity, lower predation, safe calving, foraging, etc. Density can occasionally be calculated for smaller areas that are used regularly by marine mammals, but more often than not there are insufficient scientific data to represent the spatial distribution of animals for small regions such as the construction area encompassed by the Project Area.

Therefore, given the lack of availability of NBK Bangor specific marine mammal data, this IHA application assumes that marine mammals are uniformly distributed in the study area.

3.3.2 Submergence

Cetaceans spend their entire lives in the water and spend most of their time (>90% for most species) entirely submerged below the surface. When at the surface, cetacean bodies are almost entirely below the water's surface, with only the blowhole exposed to allow breathing. This makes cetaceans difficult to locate visually and also exposes them to underwater noise, both natural and anthropogenic, essentially 100% of the time because their ears are nearly always below the water's surface.

Seals and sea lions (pinnipeds) spend significant amounts of time out of the water during breeding, molting, and hauling out periods. A few seals and sea lions have been sighted hauling out in the vicinity of NBK Bangor. In the water, pinnipeds (seals and sea lions) spend varying amounts of time underwater. California sea lions are known to rest at the surface in large groups for long amounts of time. When not actively diving, pinnipeds at the surface often orient their bodies vertically in the water column and often hold their heads above the water surface. Consequently, pinnipeds may not be exposed to underwater sounds to the same extent as cetaceans.

For the purpose of assessing impacts from underwater sound at NBK Bangor, the Navy assumed that all three cetacean species and two pinniped species that may be found in the vicinity of NBK Bangor (Steller sea lion, California sea lion, killer whale, Dall's porpoise, and harbor porpoise) spend 100% of the time underwater. This approach could be considered conservative because sea lions spend a portion of their time hauled out and therefore are expected to be exposed to less sound than is estimated by this approach. The harbor seal was the only species for which detailed information regarding the percentage of time spent underwater, in-water but at the surface, and hauled out was available (Jeffries et al 2003 and Huber et al 2001).

4 STATUS AND DISTRIBUTION OF MARINE MAMMAL SPECIES OR STOCKS THAT COULD POTENTIALLY BE AFFECTED

A description of the status, distribution, and seasonal distribution (when applicable) of the affected species or stocks of marine mammals likely to be affected by such activities.

There are six marine mammals species within the marine waters adjacent to NBK Bangor with confirmed or historic occurrence in the Project Area. Only one of these species, the Steller sea lion, is listed as threatened or endangered under the Endangered Species Act (ESA).

4.1 ESA-Listed Marine Mammals

4.1.1 Steller Sea Lion (*Eumetopias jubatus*), Eastern U.S. Stock

Status and Management

The Steller sea lion was originally listed as threatened under the ESA in 1990. In 1997, the NMFS reclassified Steller sea lions as two subpopulations, listing the Western Stock as endangered under the ESA, and maintaining threatened status for the Eastern stock (NMFS 1997). There is a final revised species recovery plan that addresses both stocks (NMFS 2008).

Critical habitat has been designated for the Steller sea lion (NMFS 1993). Critical habitat includes so-called "aquatic zones" that extend 3,000 ft (1 km) seaward in state and federally managed waters from the baseline or basepoint of each major rookery in Oregon and California (NMFS 2008). Three major rookery sites in Oregon (Rogue Reef, Pyramid Rock; and Long Brown Rock and Seal Rock on Orford Reef at Cape Blanco) and three rookery sites in California (Ano Nuevo I; Southeast Farallon I; and Sugarloaf Island and Cape Mendocino) are designated critical habitat (NMFS 1993). There is no designated critical habitat for the species in Washington.

Distribution

Steller sea lions are found along the coasts of Washington, Oregon, and northern California where they occur at breeding rookeries and numerous haulout locations along the coastline (Jeffries *et al.* 2000; Scordino 2006). From breeding rookeries in northern California (St. George Reef) and southern Oregon (Rogue Reef), male Steller sea lions often disperse widely outside of the breeding season (Scordino 2006). Based on mark recapture sighting studies, males migrate back into these Oregon and California locations from winter feeding areas in Washington, British Columbia, and Alaska (Scordino 2006).

In Washington, Steller sea lions use haulout sites primarily along the outer coast from the Columbia River to Cape Flattery, as well as along the Vancouver Island side of the Strait of Juan de Fuca (Jeffries *et al.* 2000). Numbers vary seasonally in Washington with peak numbers present during the fall and winter months (Jeffries *et al.* 2000). Steller Sea lions are occasionally present in the Puget Sound at the Toliva Shauls haul-out site in south Puget Sound (Jeffries *et al.* 2000). At NBK Bangor, Steller sea lions were observed hauled out on submarines at Delta Pier on several occasions from 2008 through 2010 during winter and spring months (Bhuthimethee 2008, personal communication; Walters 2010, personal communication). Steller sea lions likely occupy habitats in Hood Canal similar to those of the California sea lion and harbor seal, which include marine water habitats for foraging and manmade structures for haul out.

Behavior and Ecology

Steller sea lions are opportunistic predators, feeding primarily on fish and cephalopods, and their diet varies geographically and seasonally (Merrick et al. 1997). Foraging habitat is primarily shallow, nearshore and continental shelf waters; some Steller sea lions feed in freshwater rivers (Reeves et al. 1992, Robson 2002). They also are known to feed in deep waters past the continental shelf break (Jefferson 2005). Steller sea lions are gregarious animals that often travel or haul out in large groups of up to 45 individuals (Keple 2002). At sea, groups usually consist of female and subadult males; adult males are usually solitary while at sea (Loughlin 2002). Haulout and rookery sites are located on isolated islands, rocky shorelines, and jetties. Steller sea lions also haul out on buoys, rafts, floats, and Navy submarines in the Puget Sound (Jeffries et al. 2000, DoN 2001a). Females reach sexual maturity at 4 to 5 years of age (Pitcher and Calkins 1981). In the Pacific Northwest, breeding rookeries are located in British Columbia, Oregon, and northern California. There are no rookeries in Washington (NMFS 1992b, Angliss and Outlaw 2005).

Acoustics

Like all pinnipeds, the Steller sea lion is amphibious; while all foraging activity takes place in the water, breeding behavior is carried out on land in coastal rookeries (Mulsow and Reichmuth 2008, in prep). On land, territorial male Steller sea lions regularly use loud, relatively low-frequency calls/roars to establish breeding territories (Schusterman et al. 1970; Loughlin et al 1987). The calls of females range from 0.03 to 3 kHz, with peak frequencies from 0.15 to 1 kHz; typical duration is 1.0 to 1.5 sec (Campbell et al 2002). Mulsow and Reichmuth (2008) measured the unmasked aerial hearing sensitivity of one male Steller sea lion. The range of best hearing sensitivity was between 5 and 14.1 kHz (Mulsow and Reichmuth 2008). Maximum sensitivity was found at 10 kHz, where the subject had a mean threshold of 7 dB re 20 μ Pa.

The underwater hearing of two Steller sea lions were tested, the hearing threshold of the male was significantly different from that of the female. The range of best hearing for the male was from 1 to 16 kHz, with maximum sensitivity (77 dB re 1 μ Pa-m) at 1 kHz. The range of best hearing for the female was from 16 to above 25 kHz, with maximum sensitivity (73 dB re 1 μ Pa-m) at 25 kHz. However, because of the small number of animals tested, the findings could not be attributed to individual differences in sensitivity or sexual dimorphism (Kastelein et al. 2005).

4.2 Non-ESA Listed Marine Mammals

4.2.1 California Sea Lion (*Zalophus californianus*), U.S. Stock

Distribution

The geographic distribution of California sea lions includes a breeding range from Baja California to southern California. During the summer, California sea lions breed on islands from the Gulf of California to the Channel Islands and seldom travel more than about 31 miles (50 km) from the islands (Bonnell *et al.* 1983). The primary rookeries are located on the California Channel Islands of San Miguel, San Nicolas, Santa Barbara, and San Clemente (Le Boeuf and Bonnell 1980; Bonnell and Dailey 1993). Their distribution shifts to the northwest in fall and to the southeast during winter and spring, probably in response to changes in prey availability (Bonnell and Ford 1987).

The non-breeding distribution extends from Baja California north to Alaska for males, and encompasses the waters of California and Baja California for females (Reeves et al. 2008;

Maniscalco et al. 2004). In the non-breeding season, an estimated 3,000 to 5,000 adult and sub-adult males migrate northward along the coast to central and northern California, Oregon, Washington, and Vancouver Island from September to May (Jeffries et al. 2000) and return south the following spring (Mate 1975; Bonnell et al. 1983). Along their migration, they are occasionally sighted hundreds of miles (kilometers) offshore (Jefferson et al. 1993). Females and juveniles tend to stay closer to the rookeries (Bonnell et al 1983).

Peak abundance in the Puget Sound is September to May. Although there are no regular California sea lion haulouts within the Hood Canal (Jeffries et al. 2000), they often haul out at several opportune areas. They are known to utilize man-made structures such as piers, jetties, offshore buoys, and oil platforms (Riedman 1990). California sea lions in the Puget Sound even haul out on log booms and U.S. Navy submarines, and are often seen rafted off river mouths (Jeffries et al. 2000; DoN 2001). As many as 40 California sea lions have been observed hauled out at NBK Bangor on manmade structures – submarines, the floating security fence, and barges (Agness and Tannenbaum 2009a; Tannenbaum et al 2009a; Walters 2009, personal communication). California sea lions have also been observed swimming in the Hood Canal in the vicinity of the Project Area on several occasions and likely forage in both nearshore marine and inland marine deeper waters (Navy 2001).

Behavior and Ecology

California sea lions feed on a wide variety of prey, including many species of fish and squid (Everitt et al. 1981; Roffe and Mate 1984; Antonelis et al. 1990; Lowry et al. 1991). In the Puget Sound region, they feed primarily on fish such as hake, walleye pollock, herring, and spiny dogfish (Calambokidis and Baird 1994). In some locations where sea lions and salmon runs exist, California sea lions also feed on returning adult and out-migrating juvenile salmonids (London 2006). Sexual maturity occurs at around four to five years of age for California sea lions (Heath 2002). California sea lions are gregarious during the breeding season and social on land during other times.

Acoustics

On land, California sea lions make incessant, raucous barking sounds; these have most of their energy at less than 2 kHz (Schusterman et al. 1967). Males vary both the number and rhythm of their barks depending on the social context; the barks appear to control the movements and other behavior patterns of nearby conspecifics (Schusterman 1977). Females produce barks, squeals, belches, and growls in the frequency range of 0.25 to 5 kHz, while pups make bleating sounds at 0.25 to 6 kHz. California sea lions produce two types of underwater sounds: clicks (or short-duration sound pulses) and barks (Schusterman et al. 1966, 1967, Schusterman and Baillet 1969). All underwater sounds have most of their energy below 4 kHz (Schusterman et al. 1967).

The range of maximal hearing sensitivity underwater is between 1 and 28 kHz (Schusterman et al. 1972). Functional underwater high frequency hearing limits are between 35 and 40 kHz, with peak sensitivities from 15 to 30 kHz (Schusterman et al. 1972). The California sea lion shows relatively poor hearing at frequencies below 1 kHz (Kastak and Schusterman 1998). Peak hearing sensitivities in air are shifted to lower frequencies; the effective upper hearing limit is approximately 36 kHz (Schusterman 1974). The best range of sound detection is from 2 to 16 kHz (Schusterman 1974). Kastak and Schusterman (2002) determined that hearing sensitivity generally worsens with depth—hearing thresholds were lower in shallow water, except at the highest frequency tested (35 kHz), where this trend was reversed. Octave band noise levels of 65

to 70 dB above the animal's threshold produced an average TTS of 4.9 dB in the California sea lion (Kastak et al. 1999). Center frequencies were 1,000 hertz (Hz) for corresponding threshold testing at 1000 Hz and 2,000 Hz for threshold testing at 2,000 Hz; the duration of exposure was 20 minutes.

4.2.2 Harbor Seal (*Phoca vitulina*), WA Inland Waters Stock

Distribution

Harbor seals are coastal species, rarely found more than 12 miles (20 km) from shore, and frequently occupy bays, estuaries, and inlets (Baird 2001). Individual seals have been observed several miles upstream in coastal rivers. Ideal harbor seal habitat includes haulout sites, shelter during the breeding periods, and sufficient food (Bjorge 2002). Haulout areas can include intertidal and subtidal rock outcrops, sandbars, sandy beaches, peat banks in salt marshes, and manmade structures such as log booms, docks, and recreational floats (Wilson 1978; Prescott 1982; Schneider and Payne 1983; Gilber and Guldager 1998; Jeffries *et al.* 2000). Human disturbance can affect haul-out choice (Harris *et al.* 2003).


Harbor seals occur throughout Hood Canal and are seen relatively commonly in the area. They are year-round, non-migratory residents, and pup (give birth) in Hood Canal. Surveys in the Hood Canal from the mid-1970s to 2000 show a fairly stable population between 600-1,200 seals (Jeffries *et al.* 2003). Harbor seals have been observed swimming in the waters along NBK Bangor in every month of surveys conducted from 2007 to 2010 (Agness and Tannenbaum 2009b; Tannenbaum *et al.* 2009b). On the NBK Bangor waterfront, harbor seals have not been observed hauling out in the intertidal zone, but have been observed hauled out on manmade structures such as the floating security fence, buoys, barges, marine vessels, and logs (Agness and Tannenbaum 2009a; Tannenbaum *et al.* 2009a). The main haul-out locations for harbor seals in Hood Canal are located on river delta and tidal exposed areas at Quilcene, Dosewallips, Duckabush, Hamma Hamma, and Skokomish River mouths (see Figure 4-1), with the closest haul-out area to the Project Area being 10 miles southwest of NBK Bangor at Dosewallips River Mouth (London 2006).

Behavior and Ecology

Harbor seals are typically seen in small groups resting on tidal reefs, boulders, mudflats, man-made structures, and sandbars. Harbor seals are opportunistic feeders that adjust their patterns to take advantage of locally and seasonally abundant prey ((Payne and Selzer 1989, Baird 2001, Bjørge 2002). Diet consists of fish and invertebrates (Bigg 1981, Roffe and Mate 1984, Orr *et al.* 2004). Although harbor seals in the Pacific Northwest are common in inshore and estuarine waters, they primarily feed at sea (Orr *et al.* 2004) during high tide. Researchers have found that they complete both shallow and deep dives during hunting depending on the availability of prey (Tollit *et al.* 1997). Their diet in Puget Sound consists of many of the prey resources that are present in the nearshore and deeper waters of NBK Bangor, including Pacific hake and Pacific herring and adult and out-migrating juvenile salmonids. Harbor seals in Hood Canal are known to feed on returning adult salmon, including threatened summer-run chum. Over a five year study of harbor seal predation in the Hood Canal, the average percent escapement of summer-run chum consumed was 8 percent (London 2006).

Harbor seals mate at sea and females live birth during the spring and summer; although the "pupping season" varies by latitude. In coastal and inland regions of Washington, pups are born

Incidental Harassment Authorization Application for the Navy's EHW-1 Pile Replacement Project
 Conducted at Naval Base Kitsap Bangor, WA



from April through January. Pups are generally born earlier in the coastal areas and later in the Puget Sound/Hood Canal region (Calambokidis and Jeffries 1991; Jeffries et al. 2000). Suckling harbor seal pups spend as much as 40 percent of their time in the water (Bowen et al. 1999).

Acoustics

In air, harbor seal males produce a variety of low-frequency (<4 kHz) vocalizations, including snorts, grunts, and growls. Male harbor seals produce communication sounds in the frequency range of 100 to 1,000 Hz (Richardson et al. 1995). Pups make individually unique calls for mother recognition that contain multiple harmonics with main energy below 0.35 kHz (Bigg 1981, Thomson and Richardson 1995). Harbor seals hear nearly as well in air as underwater and had lower thresholds than California sea lions (Kastak and Schusterman 1998). Kastak and Schusterman (1998) reported airborne low frequency (100 Hz) sound detection thresholds at 65.4 dB re 20 μ Pa for harbor seals. In air, they hear frequencies from 0.25 kHz - 30 kHz and are most sensitive from 6 to 16 kHz (Richardson 1995, Terhune & Turnbull 1995, Wolski et al. 2003).

Adult males also produce underwater sounds during the breeding season that typically range from 0.025 to 4 kHz (duration range: 0.1 s to multiple seconds; Hanggi and Schusterman 1994). Hanggi and Schusterman (1994) found that there is individual variation in the dominant frequency range of sounds between different males, and Van Parijs et al. (2003) reported oceanic, regional, population, and site-specific variation that could be vocal dialects. In water, they hear frequencies from 1 to 75 kHz (Southall 2007) and can detect sound levels as weak as 60 to 85 dB re 1 μ Pa within that band. They are most sensitive at frequencies below 50 kHz; above 60 kHz sensitivity rapidly decreases.

4.2.3 Killer whale (*Orcinus orca*), West Coast Transient Stock

Distribution

The geographical range of transient killer whales includes the northeast Pacific, with preference for coastal waters of southern Alaska and British Columbia (Krahn et al. 2002). Transient killer whales in the eastern North Pacific spend most of their time along the outer coast, but visit Hood Canal and the Puget Sound in search of harbor seals, sea lions, and other prey. Transient occurrence in inland waters appears to peak during August and September (Morton 1990; Baird and Dill 1995; Ford and Ellis 1999) which is the peak time for harbor seal pupping, weaning, and post-weaning (Baird and Dill 1995). In 2003 and 2005, small groups of transient killer whales (11 and 6 individuals, respectively) visited Hood Canal to feed on harbor seals and remained in the area for significant periods of time (59 and 172 days, respectively) between the months of January and July.

Behavior and Ecology

Transient killer whales show greater variability in habitat use, with some groups spending most of their time foraging in shallow waters close to shore while others hunt almost entirely in open water (Felleman et al. 1991, Baird and Dill 1995, Matkin and Saulitis 1997). Transient killer whales feed on marine mammals and some seabirds, but apparently no fish (Morton 1990, Baird and Dill 1996, Ford et al. 1998, Ford and Ellis 1999, Ford et al. 2005). While present in Hood Canal in 2003 and 2005, transient killer whales preyed on harbor seals in the subtidal zone of the nearshore marine and inland marine deeper water habitats (London 2006). Other observations of foraging transient killer whales indicate they prefer to forage on pinnipeds in shallow, protected waters (Heimlich-Boran 1988; Saulitis et al. 2000). Transient killer whales travel in small,

matrilineal groups, but they typically contain fewer than 10 animals and their social organization generally is more flexible than the resident killer whale (Morton 1990, Ford and Ellis 1999). These differences in social organization probably relate to differences in foraging (Baird and Whitehead 2000). There is no information on the reproductive behavior of killer whales in this area.

Acoustics

Killer whales produce a wide variety of clicks and whistles, but most of their sounds are pulsed with frequencies ranging from 0.5 to 25 kHz (dominant frequency range: 1 to 6 kHz) (Thomson and Richardson 1995; Richardson et al. 1995). Source levels of echolocation signals range between 195 and 224 dB re 1 μ Pa-m peak-to-peak, dominant frequencies ranging from 20 to 60 kHz, and durations of about 0.1 sec (Au et al. 2004). Source levels associated with social sounds have been calculated to range between 131 to 168 dB re 1 μ Pa-m and vary with vocalization type (Veirs 2004).

Both behavioral and auditory brainstem response technique indicate killer whales can hear in a frequency range of 1 to 100 kHz and are most sensitive at 20 kHz. This is one of the lowest maximum-sensitivity frequencies known among toothed whales (Szymanski et al. 1999).

4.2.4 Dall's Porpoise (*Phocoenoides dalli*), CA/OR/WA Stock

Distribution

The Dall's porpoise is found from northern Baja California, Mexico, north to the northern Bering Sea and south to southern Japan (Jefferson et al. 1993). The species is only common between 32°N and 62°N in the eastern North Pacific (Morejohn 1979; Houck and Jefferson 1999). North-south movements in California, Oregon, and Washington have been suggested. Dall's porpoises shift their distribution southward during cooler-water periods (Forney and Barlow 1998). Norris and Prescott (1961) reported finding Dall's porpoise in southern California waters only in the winter, generally when the water temperature was less than 15°C. Seasonal movements have also been noted off Oregon and Washington, where higher densities of Dall's porpoises were sighted offshore in winter and spring and inshore in summer and fall (Green et al. 1992).

In Washington, they are most abundant in offshore waters. They are year-round residents in Washington (Green et al. 1992), but their distribution is highly variable between years likely due to changes in oceanographic conditions (Forney and Barlow 1998). Dall's porpoises are observed throughout the year in the Puget Sound north of Seattle (Osborne et al. 1998) and are seen occasionally in southern Puget Sound. Dall's porpoises may also occasionally occur in Hood Canal (Jeffries 2006, personal communication). Nearshore habitats used by Dall's porpoise could include the marine habitats found in the inland marine waters of the Hood Canal. A Dall's porpoise was observed in the deeper water at NBK Bangor in summer 2008 (Tannenbaum et al. 2009a).

Behavior and Ecology

Dall's porpoises can be opportunistic feeders but primarily consume schooling forage fish. They are known to eat squid, crustaceans, and fishes such as eelpout, herring, Pollock, whiting, and sand lance (Walker et al. 1998). Groups of Dall's porpoises generally include fewer than 10 individuals and are fluid, probably aggregating for feeding (Jefferson 1990 and 1991, Houck and Jefferson 1999). Dall's porpoises become sexually mature at 3.5 to 8 years of age (Houck and Jefferson 1999) and give birth to a single calf after 10-12 months. Breeding and calving typically

occurs in the spring and summer (Angell and Balcomb 1982). In the North Pacific, there is a strong summer calving peak from early June through August (Ferrero and Walker 1999), and a smaller peak in March (Jefferson 1989). Resident Dall's porpoise breed in Puget Sound from August to September.

Acoustics

Only short duration pulsed sounds have been recorded for Dall's porpoise (Houck and Jefferson 1999); this species apparently does not whistle often (Richardson et al. 1995). Dall's porpoises produce short duration (50 to 1,500 μ s), high-frequency, narrow band clicks, with peak energies between 120 and 160 kHz (Jefferson 1988). There is no published data on the hearing abilities of this species.

4.2.5 Harbor Porpoise (*Phocoena phocoena*), WA Inland Waters Stock

Distribution

Harbor porpoise are generally found in cool temperature to subarctic waters over the continental shelf in both the North Atlantic and North Pacific (Read 1999). This species is seldom found in waters warmer than 17°C (Read 1999) or south of Point Conception (Hubbs 1960; Barlow and Hanan 1995). Harbor porpoises can be found year-round primarily in the coastal shallow waters of harbors, bays, and river mouths (Green et al. 1992). Along the Pacific coast, harbor porpoises occur from Monterey Bay, California to the Aleutian Islands and west to Japan (Reeves et al. 2002). Harbor porpoises are known to occur in Puget Sound year round (Osmek et al. 1996, 1998; Carretta et al. 2007), and may occasionally occur in Hood Canal (Jeffries 2006, personal communication). Harbor porpoise observations in northern Hood Canal have increased in recent years (Calambokidis 2010, personal communication). A harbor porpoise was seen in deeper water at NBK Bangor during 2010 field observations (SAIC staff observations 2010).

Behavior and Ecology

Harbor porpoises are non-social animals usually seen in small groups of 2 to 5 animals. Little is known about their social behavior. Harbor porpoises can be opportunistic foragers but primarily consume schooling forage fish (Osmek et al. 1996; Bowen and Siniff 1999; Reeves et al. 2002). Along the coast of Washington, harbor porpoise primarily feed on Pacific herring (*Clupea pallasii*), market squid and smelts (Gearin et al. 1994). Females reach sexual maturity at 3-4 years and may give birth every year for several years in a row. Calves are born in late spring (Read 1990; Read and Hohn 1995). Dall's and harbor porpoises appear to hybridize relatively frequently in the Puget Sound area (Willis et al. 2004).

Acoustics

Harbor porpoise vocalizations include clicks and pulses (Ketten 1998), as well as whistle-like signals (Verboom and Kastelein 1995). The dominant frequency range is 110 to 150 kHz, with source levels of 135 to 177 dB re 1 μ Pa-m (Ketten 1998). Echolocation signals include one or two low-frequency components in the 1.4 to 2.5 kHz range (Verboom and Kastelein 1995).

A behavioral audiogram of a harbor porpoise indicated the range of best sensitivity is 8 to 32 kHz at levels between 45 and 50 dB re 1 μ Pa-m (Andersen 1970); however, auditory-evoked potential studies showed a much higher frequency of approximately 125 to 130 kHz (Bibikov 1992). The auditory-evoked potential method suggests that the harbor porpoise actually has two frequency ranges of best sensitivity. More recent psycho-acoustic studies found the range of best

hearing to be 16 to 140 kHz, with a reduced sensitivity around 64 kHz (Kastelein et al. 2002). Maximum sensitivity occurs between 100 and 140 kHz (Kastelein et al. 2002).

This Page Intentionally Left Blank

5 HARASSMENT AUTHORIZATION REQUESTED

The type of incidental taking authorization that is being requested (i.e., takes by harassment only, takes by harassment, injury and/or death), and the method of incidental taking.

Under Section 101 (a)(5)(D) of the MMPA, the Navy requests an Incidental Harassment Authorization (IHA) for the take of small numbers of marine mammals, by Level B behavioral harassment only, incidental to conduction pile driving operations associated with the EHW-1 Pile Replacement Project at NBK Bangor, Washington. The Navy requests an IHA for incidental take of marine mammals described within this application for one year commencing in July 2011 (or the issuance date, whichever is later). It is anticipated that the Navy would request an annual renewal of the IHA, since the proposed activities will occur over two years. The Navy is not requesting a multi-year Letter of Authorization (LOA) at this time because the activities described herein are not expected to rise to the level of injury or death, which would require a LOA.

Except with respect to certain activities not pertinent here, the MMPA defines “harassment” as: *any act of pursuit, torment, or annoyance which (i) has the potential to injure a marine mammal or marine mammal stock in the wild [Level A harassment]; or (ii) has the potential to disturb a marine mammal or marine mammal stock in the wild by causing disruption of behavioral patterns, including, but not limited to, migration, breathing, nursing, breeding, feeding, or sheltering [Level B harassment] (50 C.F.R, Part 216, Subpart A, Section 216.3-Definitions).*

Level A is the more severe form of harassment because it may result in injury, whereas Level B only results in disturbance without the potential for injury (Norberg pers. comm. 2007a).

5.1 Take Authorization Request

Under Section 101 (a)(5)(D) of the MMPA, the Navy requests an IHA from NMFS for: Level B take (behavioral harassment) of small numbers of marine mammals described within this application as a result of in-water pile driving activities. The Navy requests the IHA to begin coverage on July 16, 2011.

The exposure assessment methodology taken in this IHA application attempts to quantify potential exposures to marine mammals resulting from pile driving. Section 6 presents a detailed description of the acoustic exposure assessment methodology. Results from this approach tend to provide an overestimation of exposures because all animals are assumed to be available to be exposed 100% of the time, and the formulas used to estimate transmission loss used idealized parameters, which are unrealistic in nature. Modeling was conducted for the work window from 16 July – 31 October.

The analysis for the Pile Replacement Project predicts 2,488 potential exposures (see Section 6 for estimates of exposures by species and installation type) from pile installation and removal activities over the course of the project that could be classified as Level B harassment as defined under MMPA. The Navy’s mitigation procedures, presented in Section 11, include monitoring of mitigation zones prior to the initiation of pile driving, the use of sound attenuation devices (*e.g.* bubble curtain/wall) on all impact driven piles, and instantaneous in-situ hydroacoustic recordings. These mitigation measures, decrease the likelihood that marine mammals will be

exposed to sound pressure levels that would cause Level B harassment, though the amount of that decrease cannot be quantified.

The Navy does not anticipate that 2,488 actual harassment incidents will result from the Pile Replacement Project. However, to allow for scientific uncertainty regarding the exact mechanisms of the physical and behavioral effects, and as a conservative approach, the Navy is requesting authorization for take (Level B harassment) of 2,488 marine mammals over the course of one year in this IHA application.

5.2 Method of Incidental Taking

Construction activities associated with the Pile Replacement Project as outlined in Sections 1 and 2 have the potential to disturb or displace small numbers of marine mammals. Specifically, only underwater sounds generated from pile installation and removal activities (impact/vibratory pile driving and pneumatic chipping) may result in “take” in the form of Level B harassment (behavioral disturbance). Level B harassment is not anticipated from airborne sounds generated during pile installation or removal, or during other non-pile driving construction activities. Level A harassment is not anticipated to result from any of the construction activities, and measures designed to minimize the possibility of injury to marine mammals. Specifically, vibratory hammers will be the primary method of installation, which are not expected to cause injury to marine mammals due to the relatively low source levels (<190 dB). Also, no impact pile driving will occur without the use of a noise attenuation system (*e.g.* bubble curtain/wall), and pile driving will either not start or be halted if marine mammals approach the shutdown zone. See Section 11 for more details on the impact reduction and mitigation measures proposed. Furthermore, the pile driving activities analyzed are similar to other nearby construction activities within the Hood Canal, for instance, test piles driven in 2005 for the Hood Canal Bridge (SR-104) constructed by WSDOT, which have taken place with no reported injuries or mortality to marine mammals.

6 NUMBERS AND SPECIES EXPOSED

By age, sex, and reproductive condition (if possible), the number of marine mammals (by species) that may be taken by each type of taking identified in [Section 5], and the number of times such takings by each type of taking are likely to occur.

6.1 Introduction

The National Marine Fisheries Service (NMFS) application for Incidental Harassment Authorizations (IHAs) requires applicants to determine the number of marine mammals that are expected to be incidentally harassed by an action and the nature of the harassment (Level A or Level B). Section 5 defines MMPA Level A and Level B and Section 6 below presents how these definitions were relied on to develop the quantitative acoustic analysis methodologies used to assess the potential for the proposed action to affect marine mammals.

The project construction and operation as outlined in Sections 1 and 2 have the potential to take marine mammals by harassment only, primarily through construction activities involving in-water pile driving. Other activities are not expected to result in take as defined under the MMPA.

In-water pile driving would temporarily increase the local underwater and airborne noise environment in the vicinity of the Project Area. Research suggests that increased noise may impact marine mammals in several ways and depends on many factors. This will be discussed in more detail in Section 7. The following text provides a background on underwater sound, description of noise sources in the Project area, applicable noise criteria, and the basis for the calculation of take by Level B harassment. Level A harassment of cetaceans and pinnipeds for this project is not expected to occur; therefore, Level A harassment is not discussed in this application.

6.2 Fundamentals of Sound

Sound is a physical phenomenon consisting of minute vibrations that travel through a medium, such as air or water. Sound is generally characterized by several factors, including frequency and intensity. Frequency describes the sound's pitch and is measured in hertz (Hz), while intensity describes the sound's loudness. Due to the wide range of pressure and intensity encountered during measurements of sound, a logarithmic scale is used. In acoustics, the word "level" denotes a sound measurement in decibels. A decibel (dB) expresses the logarithmic strength of a signal relative to a reference. Because the decibel is a logarithmic measure, each increase of 20 dB reflects a ten-fold increase in signal amplitude (whether expressed in terms of pressure or particle motion), i.e., 20 dB means ten times the amplitude, 40 dB means one hundred times the amplitude, 60 dB means one thousand times the amplitude, and so on. Because the decibel is a relative measure, any value expressed in decibels is meaningless without an accompanying reference. In describing underwater sound pressure, the reference amplitude is usually 1 microPascal (μPa , or 10^{-6} Pascals), and is expressed as "dB re 1 μPa ." For in-air sound pressure, the reference amplitude is usually 20 μPa and is expressed as "dB re 20 μPa ."

The method commonly used to quantify airborne sounds consists of evaluating all frequencies of a sound according to a weighting system that reflects that human hearing is less sensitive at low frequencies and extremely high frequencies than at the mid-range frequencies. This is called A-weighting, and the decibel level measured is called the A-weighted sound level (dBA). A filtering method that reflects hearing of marine mammals has not yet been developed. Therefore,

underwater sound levels are not weighted and measure the entire frequency range of interest. In the case of marine construction work, the frequency range of interest is 10 to 10,000 Hz.

Table 6-1 summarizes commonly used terms to describe underwater sounds. Two common descriptors are the instantaneous peak sound pressure level (SPL) and the root mean square (rms) SPL (dB rms) during the pulse or over a defined averaging period. The peak pressure is the instantaneous maximum or minimum overpressure observed during each pulse or sound event and is presented in Pascals (Pa) or dB referenced to a pressure of one microPascal (dB re 1 μ Pa). The rms level is the square root of the energy divided by a defined time period. All underwater sound levels throughout the remainder of this application are presented in dB re 1 μ Pa unless otherwise noted.

Table 6-1 Definitions of Acoustical Terms

Term	Definition
Decibel, dB	A unit describing the amplitude of sound, equal to 20 times the logarithm to the base 10 of the ratio of the pressure of the sound measured to the reference pressure. The reference pressure for water is 1 microPascal (μ Pa) and for air is 20 μ Pa (approximate threshold of human audibility).
Sound Pressure Level, SPL	Sound pressure is the force per unit area, usually expressed in microPascals (or 20 micro Newtons per square meter), where 1 Pascal is the pressure resulting from a force of 1 Newton exerted over an area of 1 square meter. The sound pressure level is expressed in decibels as 20 times the logarithm to the base 10 of the ratio between the pressure exerted by the sound to a reference sound pressure. Sound pressure level is the quantity that is directly measured by a sound level meter.
Frequency, Hz	Frequency is expressed in terms of oscillations, or cycles, per second. Cycles per second are commonly referred to as hertz (Hz). Typical human hearing ranges from 20 Hz to 20,000 Hz.
Peak Sound Pressure (unweighted), dB re 1 μ Pa	Peak sound pressure level is based on the largest absolute value of the instantaneous sound pressure over the frequency range from 20 Hz to 20,000 Hz. This pressure is expressed in this application as dB re 1 μ Pa.
Root-Mean-Square (rms), dB re 1 μ Pa	The rms level is the square root of the energy divided by a defined time period. For pulses, the rms has been defined as the average of the squared pressures over the time that comprise that portion of waveform containing 90 percent of the sound energy for one impact pile driving impulse. ⁴
Sound Exposure Level (SEL), dB re 1 μ Pa ² sec	Sound exposure level is a measure of energy. Specifically, it is the dB level of the time integral of the squared-instantaneous sound pressure, normalized to a 1-second period. It can be an extremely useful metric for assessing cumulative exposure because it enables sounds of differing duration, to be compared in terms of total energy.
Waveforms, μ Pa over time	A graphical plot illustrating the time history of positive and negative sound pressure of individual pile strikes shown as a plot of μ Pa over time (i.e., seconds).
Frequency Spectra, dB over frequency range	A graphical plot illustrating the 6 to 12 Hz band-center frequency sound pressure over a frequency range (e.g., 10 to 5,000 Hz in this application).
A-Weighting Sound Level, dBA	The sound pressure level in decibels as measured on a sound level meter using the A- or C-weighting filter network. The A-weighting filter de-emphasizes the low and high frequency components of the sound in a manner similar to the frequency response of the human ear and correlates well with subjective human reactions to noise.
Ambient Noise Level	The background sound level, which is a composite of noise from all sources near and far. The normal or existing level of environmental noise at a given location.

⁴ Underwater sound measurement results obtained by Illingworth & Rodkin (2001) for the Pile Installation Demonstration Project in San Francisco Bay indicated that most impact pile driving impulses occurred over a 50 to 100 millisecond (ms) period. Most of the energy was contained in the first 30 to 50 ms. Analyses of that underwater acoustic data for various pile strikes at various distances demonstrated that the acoustic signal measured using the standard "impulse exponential time-weighting" on the sound level meter (35-ms rise time) correlated to the rms level measured over the duration of the pulse.

6.3 Effects of Pile Installation and Removal Activities

6.3.1 Description of Noise Sources

Underwater sound levels are comprised of multiple sources, including physical noise, biological noise, and anthropogenic noise. Physical noise includes waves at the surface, earthquakes, ice, and atmospheric noise. Biological noise includes sounds produced by marine mammals, fish, and invertebrates. Anthropogenic noise consists of vessels (small and large), dredging, aircraft overflights, and construction noise. Known noise levels and frequency ranges associated with anthropogenic sources similar to those that would be used for this project are summarized in Table 6-2. Details of each of the sources are described in the following text.

Table 6-2 Representative Noise Levels of Anthropogenic Sources

Noise Source	Frequency Range (Hz)	Underwater Noise Level (dB re 1 μ Pa)	Reference
Small vessels	250 – 1,000	151 dB rms at 1 meter (m)	Richardson et al. 1995
Tug docking gravel barge	200 – 1,000	149 dB rms at 100 m	Blackwell and Greene 2002
Vibratory driving of 30-inch Steel Pipe pile	10 – 1,500	~168 dB rms at 10m	WSDOT 2010a, 2010b
Impact driving of 30-inch Steel Pipe pile	10 – 1,500	~193 dB rms at 10m	WSDOT 2005, 2008; Caltrans 2007; Reyff 2005

In-water construction activities associated with the Project would include the use of impact and vibratory pile driving, as well as pneumatic chipping tools. The sounds produced by these activities fall into one of two sound types: pulsed and non-pulsed (defined below). Impact pile driving produces pulsed sounds, while vibratory pile driving and pneumatic chippers produce non-pulsed (or continuous) sounds. The distinction between these two general sound types is important because they have differing potential to cause physical effects, particularly with regard to hearing (e.g. Ward 1997 as cited in Southall et al. 2007).

Pulsed sounds (e.g. explosions, gunshots, sonic booms, seismic airgun pulses, and impact pile driving) are brief, broadband, atonal transients (ANSI 1986; Harris 1998) and occur either as isolated events or repeated in some succession (Southall et al. 2007). Pulsed sounds are all characterized by a relatively rapid rise from ambient pressure to a maximal pressure value followed by a decay period that may include a period of diminishing, oscillating maximal and minimal pressures (Southall et al. 2007). Pulsed sounds generally have an increased capacity to induce physical injury as compared with sounds that lack these features (Southall et al. 2007).

Non-pulse (intermittent or continuous sounds) can be tonal, broadband, or both (Southall et al. 2007). Some of these non-pulse sounds can be transient signals of short duration but without the essential properties of pulses (e.g. rapid rise time) (Southall et al. 2007). Examples of non-pulse sounds include vessels, aircraft, machinery operations such as drilling or dredging, vibratory pile driving, pneumatic chipping, and active sonar systems (Southall et al. 2007). The duration of such sounds, as received at a distance, can be greatly extended in highly reverberant environments (Southall et al. 2007).

6.3.2 Sound Exposure Criteria and Thresholds

Under the MMPA, NMFS has defined levels of harassment for marine mammals. Level A harassment is defined as “Any act of pursuit, torment, or annoyance which has the potential to injure a marine mammal or marine mammal stock in the wild.” Level B harassment is defined as “Any act of pursuit, torment, or annoyance which has the potential to disturb a marine mammal or marine mammal stock in the wild by causing disruption of behavioral patterns, including but not limited to migration, breathing, nursing, breeding, feeding or sheltering.”

Since 1997, NMFS has used generic sound exposure thresholds to determine when an activity in the ocean that produces sound might result in impacts to a marine mammal such that a take by harassment might occur (NMFS 2005b). To date, no studies have been conducted that examine impacts to marine mammal from pile driving sounds from which empirical noise thresholds have been established. Current NMFS practice regarding exposure of marine mammals to high level sounds is that cetaceans and pinnipeds exposed to impulsive sounds of 180 and 190 dB rms or above, respectively, are considered to have been taken by Level A (*i.e.*, injurious) harassment. Behavioral harassment (Level B) is considered to have occurred when marine mammals are exposed to sounds at or above 160dB rms for impulse sounds (*e.g.*, impact pile driving) and 120dB rms for continuous noise (*e.g.*, vibratory pile driving, pneumatic chipping), but below injurious thresholds. The application of the 120 dB rms threshold can sometimes be problematic because this threshold level can be either at or below the ambient noise level of certain locations. In fact, there is no evidence that pinnipeds will react to continuous sounds at this level and more research is needed (Hollingshead pers. comm. 2008). As a result, these levels are considered precautionary (NMFS 2009 74 FR 41684). NMFS is developing new science-based thresholds to improve and replace the current generic exposure level thresholds, but the criteria have not been finalized (Southall et al. 2007). The current Level A (injury) and Level B (disturbance) thresholds are provided in Table 6-3.

6.3.3 Limitations of Existing Noise Criteria

The 120 dB rms threshold level for continuous noise originated from research conducted by Malme et al. (1984) for California gray whale response to industrial sounds (Hollingshead pers. comm. 2008). This 120 dB *continuous* sound threshold should not be confused with the 120 dB *pulsed* sound criterion established for migrating bowhead whales in the Arctic as a result of research in the Beaufort Sea by Miller et al. 1999.

To date, there is no research or data supporting a response by pinnipeds or odontocetes to continuous sounds from vibratory pile driving as low as the 120 dB threshold. Southall et al. 2007 reviewed studies conducted to document behavioral responses of harbor seals and northern elephant seals to continuous sounds under various conditions, and concluded that those limited studies suggest that exposures between 90 and 140 dB re 1 μ Pa rms generally do not appear to induce strong behavioral responses. In addition, Moulton et al. 2005 concluded that ringed seal densities were not significantly reduced by intense construction activities at the study site (Northstar). Ringed seal hearing in water (Terhune and Ronald 1975) and presumably in air is probably similar to that of other phocinid seals (*e.g.* harbor seals) (Richardson et al. 1995).

Table 6-3 Injury and Disturbance Thresholds for Underwater and Airborne Sounds

Marine Mammals	Airborne Marine Construction Criteria (Impact & Vibratory Pile Driving) (re 20 µPa)	Underwater Vibratory Driving & Chipping Hammer ¹ Criteria (e.g. non-pulsed/continuous sounds) (re 1 µPa)		Underwater Impact Pile Driving Criteria (e.g. pulsed sounds) (re 1 µPa)	
	Disturbance Guideline Threshold (Haulout) ²	Level A Injury Threshold	Level B Disturbance Threshold	Level A Injury Threshold	Level B Disturbance Threshold
Cetaceans (whales, dolphins, porpoises)	N/A	180 dB rms	120 dB rms	180 dB rms	160 dB rms
Pinnipeds (seals, sea lions, walrus; except harbor seal)	100 dB rms (unweighted)	190 dB rms	120 dB rms	190 dB rms	160 dB rms
Harbor seal	90 dB rms (unweighted)	190 dB rms	120 dB rms	190 dB rms	160 dB rms

¹ Specific criteria for pneumatic chipping hammers do not exist. These tools produce continuous sounds similar to vibratory pile driving and therefore use the same criteria for the analysis of effects. ² Sound level at which pinniped haulout disturbance has been documented. Not an official threshold, but used as a guideline. dB = decibel; N/A = not applicable; rms = root mean square

6.3.4 Ambient Noise

Ambient noise by definition is background noise and it has not single source or point (Richardson et al. 1995). Ambient noise varies with location, season, time of day, and frequency. Ambient noise is continuous, but with much variability on time scales ranging from less than one second to one year (Richardson et al. 1995). Ambient underwater noise at the Project Area is widely variable over time due to a number of natural and anthropogenic sources. Sources of naturally occurring underwater noise include wind, waves, precipitation, and biological noise (such as shrimp, fish, and cetaceans). There is also human generated noise from ship or boat traffic and other mechanical means (Urick 1983). Other sources of underwater noise at industrial waterfronts could come from cranes, generators, and other types of mechanized equipment on wharves or the adjacent shoreline.

In the vicinity of the Project Area, the average broadband ambient underwater noise levels were measured at 114 dB re 1µPa between 100 Hz and 20 kHz (Slater 2009). Peak spectral noise from industrial activity was noted below the 300 Hz frequency, with maximum levels of 110 dB re 1µPa noted in the 125 Hz band. In the 300 Hz to 5 kHz range, average levels ranged between 83 and 99 dB re 1µPa. Wind-driven wave noise dominated the background noise environment at approximately 5 kHz and above, and ambient noise levels flattened above 10 kHz.

Airborne noise levels at NBK Bangor vary based on location but are estimated to average around 65 dBA (A-weighted decibels) in the residential and office park areas, with traffic noise ranging from 60 to 80 dBA during daytime hours (Cavanaugh and Tocci 1998). The highest levels of airborne noise are produced along the waterfront and at the ordnance handling areas where

estimated noise levels range from 70 to 90 dBA and may peak at 99 dBA for short durations. These higher noise levels are produced by a combination of sound sources including heavy trucks, fork lifts, cranes, marine vessels, mechanized tools and equipment, and other sound-generating industrial/military activities. All references to noise relate to noise in the air as opposed to underwater noise, and noise measurements are not corrected for distance unless specifically indicated.

6.3.5 Distance to Sound Thresholds

6.3.5.1 Underwater Sound Propagation Formula

Pile driving would generate underwater noise that potentially could result in disturbance to marine mammals swimming by the Project Area. Transmission loss (TL) underwater is the decrease in acoustic intensity as an acoustic pressure wave propagates out from a source. TL parameters vary with frequency, temperature, sea conditions, current, source and receiver depth, water depth, water chemistry, and bottom composition and topography. The formula for transmission loss is:

$$TL = B * \log_{10}(R) + C * R, \text{ where}$$

B = logarithmic (predominantly spreading) loss
C = linear (scattering and absorption) loss
R = range from source in meters

For all underwater calculations in this assessment, linear loss (C) was not used (i.e. C=0) and transmission loss was calculated using only logarithmic spreading. Therefore, using practical spreading (B=15), the revised formula for transmission loss is $TL = 15 \log_{10}(R)$.

6.3.5.2 Underwater Noise from Pile Driving

The intensity of pile driving sounds, is greatly influenced by factors such as the type of piles, hammers, and the physical environment in which the activity takes place. A large quantity of literature regarding sound pressure levels recorded from pile driving projects is available for consideration. In order to determine reasonable sound pressure levels and their associated affects on marine mammals that are likely to result from pile driving at NBK Bangor, studies with similar properties to the proposed action were evaluated. Sound levels associated with vibratory pile removal are the same as those during vibratory installation (Caltrans 2007) and have been taken into consideration in the modeling analysis. A lack of empirical data exists regarding the acoustic output of chipping hammers. As a result, acoustic information for similar types of concrete breaking instruments, such as jackhammers, concrete saws, etc. was also consulted. Additionally, NMFS' recent opinion in the Port of Anchorage LOA (NMFS 2009, 74 FR 35136) provided guidance with our acoustic assessment. For instance, NMFS noted that "chipping hammers operate at 19 percent of the energy that is required for a vibratory pile driving hammer". Overall, studies which met the following parameters were considered:

1. Pile materials: Installation - steel pipe piles (30" diameter); Removal – steel pipe piles (12 – 24" diameter); Removal – concrete piles (24" diameter)
2. Hammer machinery: Installation (steel)- vibratory and impact hammer, Removal (steel) – vibratory hammer; Removal (concrete)- pneumatic chipping and/or jackhammer
3. Physical environment - shallow depth (<100 feet [30 m]).

The tables below detail representative pile driving sound pressure levels that have been recorded from similar construction activities in recent years. Due to the similarity of these actions and the Navy's proposed action, they represent reasonable sound pressure levels which could be anticipated and these values were used in the acoustic modeling and analysis. Table 6-4 represents SPLs that may be expected during the installation of the 30-inch steel pipe piles using an impact hammer. Table 6-5 represents SPLs that may be expected during the installation of the 30-inch steel piles using a vibratory hammer. Table 6-6 represents SPLs that may be expected during the removal of the 12 to 24-inch steel pipe piles. Table 6-7 represents SPLs that may be expected during the removal of the 24-inch concrete pilings.

Table 6-4 Underwater Sound Pressure Levels Expected During Impact Installation Based on Similar In-situ Monitored Construction Activities

Project & Location	Pile Size & Type	Installation Method	Water Depth	Measured Sound Pressure Levels
Richmond San Rafael Bridge, CA ¹	30-inch Steel Pipe Pile	Impact	4-5 m	190 dB re 1 μPa (rms) at 10 m
Eagle Harbor Maintenance Facility, WA ²	30-inch Steel Pipe Pile	Impact	10 m (33 feet)	193 dB re 1 μPa (rms) at 10 m
Friday Harbor Ferry Terminal, WA ³	30-inch Steel Pipe Pile	Impact	10 m (33 feet)	196 dB re 1 μPa (rms) at 10 m
Various Projects ⁴	30-inch Steel CISS Pile	Impact	?	192 dB re 1 μPa (rms) at 10 m
			Average	~ 193 dB re 1 μPa (rms) at 10 m

Sources: ¹Caltrans, 2007; ²WSDOT, 2008; ³WSDOT, 2005; ⁴Reyff, 2005

Table 6-5 Underwater Sound Pressure Levels Expected During Vibratory Installation Based on Similar In-situ Monitored Construction Activities

Project & Location	Pile Size & Type	Installation Method	Water Depth	Measured Sound Pressure Levels
Keystone Ferry Terminal, WA ¹	30-inch Steel Pipe Pile	Vibratory	~5 m (15 ft)	166 dB re 1 μPa (rms) at 10 m
Keystone Ferry Terminal, WA ¹	30-inch Steel Pipe Pile	Vibratory	~8 m (28 ft)	171 dB re 1 μPa (rms) at 10 m
Vashon Ferry Terminal, WA ²	30-inch Steel Pipe Pile	Vibratory	10-12 m (36-40 ft)	165 dB re 1 μPa (rms) at 10 m
			Average	~ 168 dB re 1 μPa (rms) at 10 m

Sources: ¹WSDOT, 2010a; ²WSDOT, 2010b;

Table 6-6 Underwater Sound Pressure Levels Expected During Steel Pile Removal Based on Similar In-situ Monitored Construction Activities

Project & Location	Pile Size & Type	Installation Method	Water Depth	Measured Sound Pressure Levels
Unknown, CA ¹	24-inch Steel Pipe Pile	Vibratory	~15 m	165 dB re 1 µPa (rms) at 10 m

Sources: ¹Caltrans, 2007;

Table 6-7 Underwater Sound Pressure Levels Expected During Concrete Pile Removal Based on Similar In-situ Monitored Construction Activities

Project & Location	Pile Size & Type	Installation Method	Water Depth	Measured Sound Pressure Levels
United Kingdom ¹	Unknown size ² , Concrete	Jackhammer	?	161 dB re 1 µPa (rms) at 1 m

Sources: ¹Nedwell & Howell, 2004

² This is the only underwater reading available for the use of a jackhammer/pneumatic chipping tool. The size of the pile was not recorded. Since these tools operate to chip portions of concrete from the pile, its sound output may not be tied to the size of the pile itself as impact and vibratory pile drivers are. Therefore, this data was found to be representative for this project.

Several noise reduction measures can be employed during pile driving to reduce the high source pressures associated with impact pile driving. Among these is the use of bubble curtains, cofferdams, pile caps, or the use of vibratory installation. The efficacy of bubble curtains is dependent upon a variety of site-specific factors, including environmental conditions such as water current, sediment type, and bathymetry; the type and size of the pile; and the type and energy of the hammer. Thorson and Reyff (2004) determined that a properly designed bubble curtain could provide a reduction of 5 to 20 dB. Under certain conditions, bubble curtains may not provide the appropriate level of attenuation. In the event that the underwater monitoring demonstrates the proper attenuation is not being achieved, other mitigation techniques may be used (see Section 11). The use of a dewatered cofferdam (i.e., there is no water inside the cofferdam) represents the most effective way of reducing sound because the pile driving is completely decoupled from the surrounding water column. Reyff (2003) conducted measurements in the shallow water surrounding Pier 16E for the San Francisco-Oakland Bay Bridge (SFOBB) project during the driving of 2.4 meter cast-in-steel-shell (CISS) piles in a dewatered cofferdam. That study determined that the dewatered cofferdam reduced underwater sound pressures by 30 dB. However, this study also noted that low frequency levels were higher in one direction most likely based on the complex propagation path.

In addition to these techniques, cap materials have been used on the pile to reduce pile driving noise. Laughlin (2006) measured sound levels of a 12-inch diameter standard steel pile with bubble curtains and different pile cap materials. He found that using wood as a pile cap may provide 11 to 26 dB reductions; however, wood compressed easily or caught on fire and therefore does not warrant regular use. Conbest provided a 7 to 8 dB reduction, nylon provided 4 to 5 dB reduction, and Micarta provided 1 to 5 dB reduction. The use of vibratory installation versus impact driving can be considered a mitigation technique as well. Noise levels associated with vibratory pile driving are typically 15 to 20 dB less than impact pile driving, (Hastings and Popper 2005; WSDOT 2008).

For the EHW-1 Pile Replacement Project, the Navy intends to employ noise reduction techniques during impact pile driving, including the use of sound attenuation systems, such as a bubble curtain/wall. See Section 11 for more details on the impact reduction and mitigation measures proposed. Additionally, vibratory pile driving will be the primary installation method, which has lower source levels than impact pile driving. The calculations of the distances to the marine mammal noise thresholds were calculated for impact installation with and without consideration for mitigation measures. Distances calculated with consideration for mitigation assumed a 10 dB reduction in source levels from the use of sound attenuation devices. The Navy will be using the mitigated distances for impact pile driving for all further analysis in this IHA. All calculated distances to and the total area encompassed by the marine mammal noise thresholds are provided in Tables 6-8 through 6-10, respectively.

Table 6-8 Calculated Distance(s) to and the Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Pile Installation

Species	Threshold	Without Mitigation (m) ¹	-10 dB Mitigation (m) ¹	Distance in (km)	Area in (km ²)
Pinnipeds	Impact Driving Injury (190 dB rms)	16	4	0.004	0.00005
Cetaceans	Impact Driving Injury (180 dB rms)	74	16	0.016	0.0008
All Marine Mammals	Impact Driving Disturbance (160 dB rms)	1,585	342	0.342	0.367
Pinnipeds	Vibratory Driving Injury (190 dB rms)	0	0	0.000	0.000
Cetaceans	Vibratory Driving Injury (180 dB rms)	2	2	0.002	0.00001
All Marine Mammals	Vibratory Driving Disturbance (120 dB rms)	15,849	15,849 ²	15.849 ²	789.139 ²

All sound levels expressed in dB re 1 µPa rms. dB = decibel; rms = root-mean-square; µPa = microPascal

Practical spreading loss (15 log, or 4.5 dB per doubling of distance) used for calculations.

¹Sound pressure levels used for calculations were: 193 dB re 1 µPa @ 10m for impact and 168 dB re 1 µPa @ 10m for vibratory

²Range calculated is greater than what would be realistic. Hood Canal average width at site is 2.4 km, and is fetch limited from N to S at 20.3 km.

Table 6-9 Calculated Distance(s) to and the Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Steel Pile Removal

Species	Threshold	Distance in (m) ¹	Distance in (km)	Area in (km ²)
Pinnipeds	Vibratory Driving Injury (190 dB rms)	0	0.000	0.000
Cetaceans	Vibratory Driving Injury (180 dB rms)	1	0.001	0.000003
All Marine Mammals	Vibratory Driving Disturbance (120 dB rms)	10,000 ²	10.0 ²	314.159 ²

All sound levels expressed in dB re 1 μ Pa rms.

dB = decibel; rms = root-mean-square; μ Pa = microPascal

Practical spreading loss (15 log, or 4.5 dB per doubling of distanced) used for calculations.

¹Sound pressure levels used for calculations were: 165 dB re1 μ Pa @ 10m for vibratory

²Range calculated is greater than what would be realistic. Hood Canal average width at site is 2.4 km, and is fetch limited from N to S at 20.3 km.

Table 6-10 Calculated Distance(s) to and the Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Concrete Pile Removal

Species	Threshold ¹	Distance in (m) ²	Distance in (km)	Area in (km ²)
Pinnipeds	Chipping Hammer Injury (190 dB rms)	0	0.000	0.000
Cetaceans	Chipping Hammer Injury (180 dB rms)	0	0.000	0.000
All Marine Mammals	Chipping Hammer Disturbance (120 dB rms)	542 ³	0.542 ³	0.929 ³

All sound levels expressed in dB re 1 μ Pa rms.

dB = decibel; rms = root-mean-square; μ Pa = microPascal

Practical spreading loss (15 log, or 4.5 dB per doubling of distanced) used for calculations.

¹Specific criteria for pneumatic chipping hammers does not exist. These tools produce continuous sound similar to vibratory pile driving and therefore use the same criteria for the analysis of effects.

²Sound pressure levels used for calculations were: 161 dB re1 μ Pa @ 1m for jackhammer

³Range calculated is greater than what would be realistic. Hood Canal average width at site is 2.4 km, and is fetch limited from N to S at 20.3 km.

The calculations presented in Tables 6-8 through 6-10 assumed a field free of obstruction, which is unrealistic, however, because Hood Canal does not represent open water conditions (free field) and therefore, sounds would attenuate as they encountered land masses or bends in the canal. As a result, some of the distances and areas of impact calculated cannot actually be attained at the project area. The actual distances to the behavioral disturbance thresholds for impact and vibratory pile driving and pneumatic chipping may be shorter than those calculated due to the irregular contour of the waterfront, the narrowness of the canal, and the maximum fetch (furthest distance sound waves travel without obstruction [i.e. line of site]) at the project area. Tables 6-11 through 6-13 depict the actual areas encompassed by the marine mammal thresholds during each stage of the EHW-1 Pile Replacement Project. Figures 6-1 through 6-4 depict the areas of each underwater sound threshold that are predicted to occur at the project area due to pile driving for marine mammals (cetaceans and pinnipeds) during each stage of the Pile Replacement project.

Table 6-11 Actual Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Pile Installation

Species	Threshold	-10 dB Mitigation (m)	Distance in (km)	Predicted Area in (km ²)	Actual Area in (km ²)
Pinnipeds	Impact Driving Injury (190 dB rms)	4	0.004	0.00005	0.000
Cetaceans	Impact Driving Injury (180 dB rms)	16	0.016	0.0008	0.001

*Incidental Harassment Authorization Application for the Navy's EHW-1 Pile Replacement Project
Conducted at Naval Base Kitsap Bangor, WA*

Species	Threshold	-10 dB Mitigation (m)	Distance in (km)	Predicted Area in (km ²)	Actual Area in (km ²)
All Marine Mammals	Impact Driving Disturbance (160 dB rms)	342	0.342	0.367	0.287
Pinnipeds	Vibratory Driving Injury (190 dB rms)	0	0.000	0.000	0.000
Cetaceans	Vibratory Driving Injury (180 dB rms)	2	0.002	0.00001	0.000
All Marine Mammals	Vibratory Driving Disturbance (120 dB rms)	15,849	15.849	789.139	40.273

Table 6-12 Actual Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Steel Pile Removal

Species	Threshold	Distance in (m)	Distance in (km)	Predicted Area in (km ²)	Actual Area in (km ²)
Pinnipeds	Vibratory Driving Injury (190 dB rms)	0	0.000	0.000	0.000
Cetaceans	Vibratory Driving Injury (180 dB rms)	1	0.001	0.000003	0.000
All Marine Mammals	Vibratory Driving Disturbance (120 dB rms)	10,000	10.0	314.159	35.870

Table 6-13 Actual Area(s) Encompassed by the Underwater Marine Mammal Noise Thresholds During Concrete Pile Removal

Species	Threshold	Distance in (m)	Distance in (km)	Predicted Area in (km ²)	Actual Area in (km ²)
Pinnipeds	Chipping Hammer Injury (190 dB rms)	0	0.000	0.000	0.000
Cetaceans	Chipping Hammer Injury (180 dB rms)	0	0.000	0.000	0.000
All Marine Mammals	Chipping Hammer Disturbance (120 dB rms)	542	0.542	0.929	0.608




Figure 6-1 Distance(s) (m) to Underwater Sound Thresholds for Cetaceans from Impact & Vibratory Pile Driving During Pile Installation




Figure 6-2 Distance(s) (m) to Underwater Sound Thresholds for Pinnipeds from Impact & Vibratory Pile Driving During Pile Installation




Figure 6-3 Distance(s) (m) to Underwater Sound Thresholds for all Marine Mammals from Vibratory Pile Driving During Steel Pile Removal



Figure 6-4 Distance(s) (m) to Underwater Sound Thresholds for all Marine Mammals from a Chipping Hammer During Concrete Pile Removal

6.3.5.3 Airborne Sound Propagation Formula

Pile driving can generate airborne noise that could potentially result in disturbance to marine mammals (pinnipeds) which are hauled out or at the water's surface. As a result, the Navy analyzed the potential for pinnipeds hauled out or swimming at the surface near the NBK Bangor to be exposed to airborne sound pressure levels that could result in Level B behavioral harassment. The appropriate airborne noise thresholds for behavioral disturbance for all pinnipeds, except harbor seals is 100 dB re 20 μ Pa rms (unweighted) and for harbor seals is 90 dB re 20 μ Pa rms (unweighted) (see Table 6-3). A spherical spreading loss model, assuming average atmospheric conditions, was used to estimate the distance to the 100 dB and 90 dB re 20 μ Pa rms (unweighted) airborne thresholds. The formula for calculating spherical spreading loss is:

$$TL = 20\log r$$

where:

TL = Transmission loss

r = Distance from source to receiver

*Spherical spreading results in a 6 dB decrease in sound pressure level per doubling of distance.

6.3.5.4 Airborne Sound from Pile Driving

The intensity of pile driving sounds is greatly influenced by factors such as the type of piles, hammers, and the physical environment in which the activity takes place. A large quantity of literature regarding sound pressure levels recorded from pile driving projects is available for consideration. In order to determine reasonable airborne sound pressure levels and their associated affects on marine mammals that are likely to result from pile driving at NBK Bangor, studies with similar properties to the proposed action were evaluated. Studies which met the following parameters were considered:

1. Pile materials: Installation - steel pipe piles (24-42" diameter); Removal – steel pipe piles (12 – 30" diameter); Removal – concrete piles (24" diameter)
2. Hammer machinery: Installation (steel)- vibratory and impact hammer, Removal (steel) – vibratory hammer; Removal (concrete)- pneumatic chipping and/or jackhammer
3. Physical environment - shallow depth (<100 feet [30 m]).

The tables below detail representative airborne pile driving sound pressure levels that have been recorded from similar construction activities in recent years. Due to the similarity of these actions and the Navy's proposed action, they represent reasonable sound pressure levels which could be anticipated and these values were used in the acoustic modeling and analysis. Table 6-14 represents SPLs that may be expected during the installation of the 30-inch steel pipe piles using an impact hammer. Table 6-15 represents SPLs that may be expected during the installation of the 30-inch steel piles using a vibratory hammer. Table 6-16 represents SPLs that may be expected during the removal of the 12 to 24-inch steel pipe piles. Table 6-17 represents SPLs that may be expected during the removal of the 24-inch concrete pilings.

Table 6-14 Airborne Sound Pressure Levels Expected During Impact Installation from Similar In-situ Monitored Construction Activities

Project & Location	Pile Size & Type	Installation Method	Water Depth	Measured Sound Pressure Levels
Northstar Island, AK ¹	42- inch Steel Pipe Pile	Impact	~12 m (40 feet)	97 dB re 20 µPa (rms) at 525 feet
Friday Harbor Ferry Terminal, WA ²	24-inch Steel Pipe	Impact	~10 m (33 feet)	112 dB re 20 µPa (rms) at 160 feet
			Average	120 dB re 20 µPa (rms) at 50 feet

Sources: ¹Blackwell et al., 2004; ²WSDOT, 2005

Table 6-15 Airborne Sound Pressure Levels Expected During Vibratory Installation from Similar In-situ Monitored Construction Activities

Project & Location	Pile Size & Type	Installation Method	Water Depth	Measured Sound Pressure Levels
Keystone Ferry Terminal, WA ¹	30- inch Steel Pipe Pile	Vibratory	~9 m (30 feet)	98 dB re 20 µPa (rms) at 36 feet

Sources: ¹WSDOT, 2010c

Table 6-16 Airborne Sound Pressure Levels Expected During Steel Pile Removal from Similar In-situ Monitored Construction Activities

Project & Location	Pile Size & Type	Installation Method	Water Depth	Measured Sound Pressure Levels
Wahkiakum Ferry Terminal ¹	18- inch Steel Pipe Pile	Vibratory	~3-4 m (10-12 feet)	87.5 dB re 20 µPa (rms) at 50 feet
Keystone Ferry Terminal, WA ¹	30- inch Steel Pipe Pile	Vibratory	~9 m (30 feet)	98 dB re 20 µPa (rms) at 36 feet
			Average	92 dB re 20 µPa (rms) at 50 feet

Sources: ¹WSDOT, 2010c

Table 6-17 Airborne Sound Pressure Levels Expected During Concrete Pile Removal from Similar In-situ Monitored Construction Activities

Project & Location	Pile Size & Type	Installation Method	Water Depth	Measured Sound Pressure Levels
Unknown ¹	Unknown ² , Concrete	Chipping Hammer	?	92 dB re 20 µPa (rms) at 10 m

Sources: ¹Cheremisinoff, 1996;

² This is the only underwater reading available for the use of a jackhammer/pneumatic chipping tool. The size of the pile was not recorded. Since these tools operate to chip portions of concrete from the pile, its sound output may not be tied to the size of the pile itself as impact and vibratory pile drivers are. Therefore, this data was found to be representative for this project.

The distances to the airborne thresholds were calculated with the airborne transmission loss formula presented in section 6.5.3. All calculated distances to and the total area encompassed by the marine mammal noise thresholds are provided in Tables 6-18 through 6-20. All, airborne distances are less than those calculated for underwater sound thresholds, with the exception of the behavioral disturbance distances from impact pile driving for harbor seals. Therefore, the

monitoring buffer zone for behavioral disturbance will be expanded to encompass this distance. All construction noise associated with the Project would not extend beyond the buffer zone that would be established to protect seals and sea lions. Figures 6-5 through 6-10 depict the actual distances for each airborne sound threshold that are predicted to occur at the project area due to pile driving for pinnipeds.

Table 6-18 Calculated Distances (m) to and the Area(s) Encompassed by the Marine Mammal Noise Thresholds In-Air from Pile Installation

Species	Threshold	Airborne Behavioral Disturbance		
		Distance (m)	Distance (km)	Area (km ²)
Pinnipeds (except harbor seal)	100dB rms (impact disturbance)	159 m (522 feet)	0.159	0.079
Pinnipeds (except harbor seal)	100dB rms (vibratory disturbance)	9 m (30 feet)	0.009	0.00025
Harbor seal	90dB rms (impact disturbance)	501 m (1643 feet)	0.501	0.789
Harbor seal	90dB rms (vibratory disturbance)	29 m (95 feet)	0.029	0.0026

All sound pressure levels are reported re 20 µPa rms (unweighted)

Table 6-19 Calculated Distances (m) to and the Area(s) Encompassed by the Marine Mammal Noise Thresholds In-Air from Steel Pile Removal

Species	Threshold	Airborne Behavioral Disturbance		
		Distance (m)	Distance (km)	Area (km ²)
Pinnipeds (except harbor seal)	100dB rms (vibratory disturbance)	7 m (23 feet)	0.007	0.00015
Harbor seal	90dB rms (vibratory disturbance)	20 m (66 feet)	0.020	0.00126

All sound pressure levels are reported re 20 µPa rms (unweighted)

Table 6-20 Calculated Distances (m) to and the Area(s) Encompassed by the Marine Mammal Noise Thresholds In-Air from Concrete Pile Removal

Species	Threshold	Airborne Behavioral Disturbance		
		Distance (m)	Distance (km)	Area (km ²)
Pinnipeds (except harbor seal)	100dB rms (vibratory disturbance)	4 m (13 feet)	0.004	0.00005
Harbor seal	90dB rms (vibratory disturbance)	13 m (43 feet)	0.013	0.0005

All sound pressure levels are reported re 20 µPa rms (unweighted)




Figure 6-5 Distance(s) (m) to Airborne Sound Thresholds for Pinnipeds (except harbor seals) from Impact & Vibratory Pile Driving During Pile Installation




Figure 6-6 Distance(s) (m) to Airborne Sound Thresholds for Harbor Seals from Impact & Vibratory Pile Driving During Pile Installation



Figure 6-7 Distance(s) (m) to Airborne Sound Thresholds for Pinnipeds (except Harbor Seals) from Vibratory Pile Driving During Steel Pile Removal



Figure 6-8 Distance(s) (m) to Airborne Sound Thresholds for Harbor Seals from Vibratory Pile Driving During Steel Pile Removal



Figure 6-9 Distance(s) (m) to Airborne Sound Thresholds for Pinnipeds (except Harbor Seals) from a Chipping Hammer During Concrete Pile Removal



Figure 6-10 Distance(s) (m) to Airborne Sound Thresholds for Harbor Seals from a Chipping Hammer During Concrete Pile Removal

6.3.6 Auditory Masking

Natural and artificial sounds can disrupt behavior by masking, or interfering with a marine mammal's ability to hear other sounds. Masking occurs when the receipt of a sound is interfered with by another coincident sound at similar frequencies and at similar or higher levels. If the second sound were man-made, it could be potentially harassing (according to the MMPA) if it disrupted hearing-related behavior such as communications or echolocation. It is important to distinguish temporary threshold shift (TTS) and permanent threshold shift (PTS), which persist after the sound exposure, from masking, which occurs during the sound exposure. Because masking (without a resulting in a threshold shift [TS]) is not associated with abnormal physiological function, it is not considered a physiological effect in this IHA application, but rather a potential behavioral effect.

The most intense underwater sounds in the proposed action are those produced by impact pile driving. Given that the energy distribution of pile driving covers a broad frequency spectrum, sound from these sources would likely be within the audible range of Steller sea lions, California sea lions, harbor seals, transient killer whales, harbor porpoise, and Dall's porpoise. Impact pile driving activity is relatively short-term, with rapid pulses occurring for approximately 15 minutes per pile. The probability for impact pile driving resulting from this proposed action masking acoustic signals important to the behavior and survival of marine mammal species is therefore negligible. Vibratory pile driving is also relatively short-term, with rapid oscillations occurring for a few hours per pile. It is possible that vibratory pile driving and to a lesser degree pneumatic chipping, resulting from this proposed action may mask acoustic signals important to the behavior and survival of marine mammal species, but the short-term duration and limited affected area would result in a negligible impact from masking. Any masking event that could possibly rise to Level B harassment under the MMPA would occur concurrently within the zones of behavioral harassment already estimated for pneumatic chipping and vibratory and impact pile driving, and which have already been taken into account in the exposure analysis.

6.3.7 Basis for Estimating Take by Harassment

The U.S. Navy analyzed the potential for taking of small numbers of Steller sea lions, California sea lions, harbor seals, transient killer whales, Dall's porpoises, and harbor porpoises in the Hood Canal as a result of pile installation and removal during construction activities associated with the Pile Replacement Project. Based on densities available for the Steller sea lion, this species is not likely to be present during the short duration of the project, therefore no takes were estimated. The takes requested for the other species are expected to have no more than a minor effect on individual animals and no effect on the populations of these species. Any effects experienced by individual marine mammals are anticipated to be limited to short-term disturbance of normal behavior or temporary displacement of animals near source of the noise.

6.3.7.1 Steller Sea Lion

Steller sea lions are present in the Hood Canal, but are only expected as far as the project area during November through mid-April (cold season). Because this action will occur between July 16 – October 31, when Steller sea lions are not likely to be present in the project area, no acoustic impacts from pile driving operations are expected for this species.

6.3.7.2 California Sea Lion

California sea lions are present in the Hood Canal almost year-round with the exception of mid-June through August. The Navy conducted year round waterfront surveys for marine mammals at NBK Bangor in 2008 and 2009 (DoN 2010a). During these surveys, the daily maximum number of California sea lions hauled out for the months July – October (the timeframe of the Pile Replacement Project), were 0, 0, 12, and 47 in 2008 and 0, 1, 32, and 44 in 2009, respectively. The monthly average of the maximum number of California sea lions observed per day was 17 individuals. Females are rarely observed north of the California-Oregon border (NMFS 2008c), therefore only adult and sub-adult males are expected in the Hood Canal. Breeding rookeries are in California; therefore pups are not expected to be present in the Hood Canal.

Potential takes would likely involve sea lions that are moving through the area en route to submarine haulout or during the return trip to the ocean when pile driving would occur. California sea lions that are taken could exhibit behavioral changes such as increased swimming speeds, increased surfacing time, or decreased foraging. Most likely, California sea lions may move away from the sound source and be temporarily displaced from the areas of pile driving. With the absence of any major rookeries and only a few isolated haul-out areas near or adjacent to the project site, potential takes by disturbance will have a negligible short-term effect on individual California sea lions and would not result in population-level impacts.

6.3.7.3 Harbor Seal

Harbor seals are present in the Hood Canal year-round and would be expected at the project site. Harbor seal numbers increase from January through April and then decrease from May through August as the harbor seals move to adjacent bays on the outer coast of Washington for the pupping season. Harbor seals are the most abundant marine mammal in the Hood Canal. Jeffries et al. (2003) did a stock assessment of harbor seals in the Hood Canal in 1999 and counted 711 harbor seals hauled out. This abundance was adjusted using a correction factor of 1.53 to account for seals in the water and not counted to provide a population estimate of 1,088 harbor seals in the Hood Canal. The Navy conducted boat surveys of the waterfront area in 2008 from July to September (Agness and Tannenbaum 2009a). Harbor seals were sited during every survey and were found in all marine habitats including near and hauled out on man-made objects such as piers and buoys. During most of the year, all age and sex classes (except newborn pups) could occur in the Project area throughout the period of construction activity. From April through mid-July, female harbor seals haul out on the outer coast of Washington at pupping sites to give birth. Pups may be encountered near these areas during this time. Since there are no known pupping sites in the vicinity of the Project, harbor seal pups are not expected to be present during pile driving.

Potential takes would likely involve seals that are moving through the area on foraging trips when pile driving would occur. Harbor seals that are taken could exhibit behavioral changes such as increased swimming speeds, increased surfacing time, or decreased foraging. Most likely, harbor seals may move away from the sound source and be temporarily displaced from the areas of pile driving. With the absence of any major rookeries and only a few potential haul-out areas near the project site, potential takes by disturbance will have a negligible short-term effect on individual harbor seals and would not result in population-level impacts.

6.3.7.4 Transient Killer Whales

Transient killer whales are uncommon visitors to Hood Canal. Transients may be present in the Hood Canal anytime during the year and traverse as far as the project site. Resident killer whales have not been observed in Hood Canal, but transient pods were observed in Hood Canal for lengthy periods of time in 2003 (January-March) and 2005 (February-June), feeding on harbor seals (London 2006).

Potential takes would likely involve transient killer whales that are moving through the area on foraging trips when pile driving would occur. Killer whales that are taken could exhibit behavioral changes such as increased swimming speeds, increased surfacing time, or decreased foraging. Most likely, killer whales may move away from the sound source and be temporarily displaced from the areas of pile driving. With the absence of any regular occurrence adjacent to the project site, potential takes by disturbance will have a negligible short-term effect on individual killer whales and would not result in population-level impacts.

6.3.7.5 Dall's Porpoise

Dall's porpoise may be present in the Hood Canal year-round and could occur as far as the project site. Their use of inland Washington waters, however, is mostly limited to the Strait of Juan de Fuca. The Navy conducted boat surveys of the waterfront area in 2008 from July to September (Agness and Tannenbaum 2009a). During one of the surveys a Dall's porpoise was sighted in August in the deeper waters off Carlson Spit.

Potential takes would likely involve Dall's porpoise that are moving through the area on foraging trips when pile driving would occur. Dall's porpoise that are taken could exhibit behavioral changes such as increased swimming speeds, increased surfacing time, or decreased foraging. Most likely, Dall's porpoise may move away from the sound source and be temporarily displaced from the areas of pile driving. With the absence of any regular occurrence adjacent to the project site, Potential takes by disturbance will have a negligible short-term effect on individual Dall's porpoise and would not result in population-level impacts.

6.3.7.6 Harbor Porpoise

Harbor porpoises may be present in the Hood Canal year-round, however their presence is rare. During waterfront survey of NBK Bangor over the past two years (2008 – present) only one harbor porpoise has been seen in 24 surveys.

Potential takes could occur if harbor porpoises move through the area on foraging trips when pile driving would occur. Harbor porpoises that are taken could exhibit behavioral changes such as increased swimming speeds, increased surfacing time, or decreased foraging. Most likely, harbor porpoises may move away from the sound source and be temporarily displaced from the areas of pile driving. With the absence of any regular occurrence adjacent to the project site, Potential takes by disturbance will have a negligible short-term effect on individual harbor porpoises and would not result in population-level impacts.

6.3.8 Description of Take Calculation

The take calculations presented here relied on the best data currently available for marine mammal populations in the Hood Canal. The population data used is discussed in within each species take calculation subsection 6.7.1 – 6.7.5. The formula was developed for calculating

take due to impact pile driving and applied to each group specific noise impact threshold. The formula is founded on the following assumptions:

- Each species population is at least as large as any previously documented highest population estimate.
- All pilings to be installed would have a noise disturbance distance equal to the piling that causes the greatest noise disturbance (i.e. the piling furthest from shore).
- Pile driving could potentially occur every day of the in-water work window; However, it is estimated no more than a few hours of pile driving will occur per day. An average of 2 steel piles will be installed and removed per day or an average of 3 concrete piles will be removed per day.
- Some degree of mitigation (i.e. sound attenuation system, etc.) will be utilized during all impact pile driving, as discussed previously.
- That an individual can only be taken once per method of installation/removal during a 24 hour period.

The calculation for marine mammal takes is estimated by:

$$\text{Take estimate} = (n * \text{ZOI}) * X \text{ days of total activity}$$

where:

n = density estimate used for each species/season

X = number of days of pile driving, estimated based on the total number of piles and the average number of piles that the contractor can install per day.

ZOI⁵ = noise threshold zone of influence (ZOI) impact area

n * ZOI produces an estimate of the abundance of animals that could be present in the area for exposure, this must be a whole number, therefore, this value was rounded (down if <0.5, up if >0.5).

The ZOI impact area is the estimated range of impact to the noise criteria. The formula for determining the area of a circle ($\pi * \text{radius}^2$) was used to calculate the ZOI around each pile, for each threshold. The distances specified in Tables 6-11 through 6-13 and Tables 6-18 through 6-20, were used for the radius in the equation. All impact pile driving take calculations were based on the estimated threshold ranges using a bubble curtain with 10 dB attenuation as a mitigation measure. The ZOI impact area took into consideration the possible effected area of the Hood Canal from the furthest from shore pile driving site with attenuation due to land shadowing from bends in the canal. As described in Section 6.5.2 with regard to the distances, because of the close proximity of some of the piles to the shore, the narrowness of the canal at the project area, and the maximum fetch, the ZOIs for each threshold aren't necessarily spherical and may be truncated.

While pile driving can occur any day throughout the in-water work window, only a "fraction" of that time is actually spent pile driving. Some days there will be only 30 minutes of pile driving, other days several hours. The contractor estimates that pile installation could occur at a

⁵ Zone of Influence (ZOI) is the area encompassed by all locations where the sound pressure levels equal or exceed the threshold being evaluated.

maximum rate of four piles per day, however, it's more likely that an average of two piles will be installed and removed per day. The contractor estimates that a maximum of five concrete piles can be removed per day, with an average of three being removed per day. For each pile installed, vibratory pile driving is expected to be no more than one hour. The impact driving portion of the project is anticipated to take approximately 15 minutes per pile, per day with a maximum of 5 piles requiring proofing. All steel piles will be extracted using a vibratory hammer. Extraction is anticipated to take approximately 30 minutes per pile. Concrete piles will be removed using a pneumatic chipping hammer or other similar concrete demolition tool. It is expected to take a couple of hours to remove each concrete pile with a pneumatic chipping hammer. For steel piles, this results in a maximum of two hours of pile driving per pile or potentially four hours per day. For concrete piles, this results in a maximum of two hours of pneumatic chipping per pile, or potentially 6 hours per day.

Therefore, while 216 days of in-water work time is proposed (108 days per construction period), only a fraction of the total work time per day will actually be spent pile driving. An average work day (two hours post-sunrise to two hours prior to sunset) is approximately 8-9 hours, depending on the month. While it is anticipated that only 4 hours of pile driving would be needed per day for steel piles, or 6 hours of pneumatic chipping would be needed for concrete piles, to take into account deviations from the estimated times for pile installation and removal, the Navy modeled potential impacts as if the entire day could be spent pile driving.

Based on the proposed action, the total pile driving time from vibratory pile driving during installation would be approximately 14 days (28 piles at an average of 2 per day). The total pile driving time from vibratory pile driving during steel pile removal would be 21 days (42 piles at an average of 2 per day). The total pile driving time for utilizing a pneumatic chipping hammer during concrete pile removal would be 32 days (96 piles at an average of 3 per day). Therefore, impacts for installation, steel pile removal, and concrete pile removal were modeled as if these actions were to occur throughout the duration of 14, 21, and 32 days, respectively. During installation, there is the potential for the contractor to need to utilize an impact hammer to proof a select number of piles, although past repairs on the EHW-1 pier have never required the use of an impact pile driver. However, if the use of an impact hammer is required, impact pile driving will occur on no more than five piles, with only one pile being impact driven per day. Therefore, impact pile driving during installation was modeled as occurring for 5 days.

The exposure assessment methodology is an estimate of the numbers of individuals exposed to the effects of pile driving activities exceeding NMFS established thresholds. Of significant note in these exposure estimates, additional mitigation methods (i.e. visual monitoring and the use of shutdown zones) were not quantified within the assessment and successful implementation of this mitigation is not reflected in exposure estimates. Results from acoustic impact exposure assessments should be regarded as conservative estimates that are strongly influenced by limited biological data. While the numbers generated from the pile driving exposure calculations provide conservative overestimates of marine mammal exposures for consultation with NMFS, the short duration and limited geographic extent of Pile Replacement Project would further limit actual exposures.

6.3.8.1 Steller Sea Lion

Although Steller sea lions have been documented in Hood Canal, the numbers (at least at present) are still fairly low, and their presence is only expected in the project area during

November through mid-April. Because this action will occur between July 16 – Oct 31, when Steller sea lions are not likely to be present in the project area no acoustic impacts from pile driving operations (including the use of a pneumatic chipping hammer) are expected for this species.

6.3.8.2 California Sea Lion

California sea lions are present in the Hood Canal almost year-round with the exception of mid-June through August. California sea lions are likely present in the Hood Canal as far as the project site from January through mid-June, although this is changing every year. The Navy conducted year round waterfront surveys for marine mammals at NBK Bangor in 2008 and 2009 (DoN 2010a). During these surveys, the daily maximum number of California sea lions hauled out for the months July – October (the timeframe of the Pile Replacement Project), were 0, 0, 12, and 47 in 2008 and 0, 1, 32, and 44 in 2009, respectively. Because the proportion of pile driving that could occur in a given month is dependent on several factors (i.e. availability of materials, weather, etc.) the Navy assumed that pile driving operations could occur at any time in the construction window. Therefore, an average of the maximum number of California sea lions observed per day across the months of July – October was used in the modeling analysis. The monthly average of the maximum number of California sea lions observed per day was 17 individuals. Exposures were calculated using a density derived from this value (17 individuals), divided by the area encompassed by the maximum fetch at the project site (41.5 km²) and the formula in Section 6.7. Table 6-21 depicts the number of acoustic harassments that are estimated from vibratory and impact pile driving and pneumatic chipping both underwater and in-air.

Table 6-21 Number of Potential Exposures of California Sea Lions within Various Acoustic Threshold Zones

Density at Project Area (July- Oct)	Stage of EHW-1 Action	Underwater			Airborne	
		Impact Injury Threshold (190 dB)	Impact Disturbance Threshold (160dB)	Vibratory ¹ Disturbance Threshold (120 dB)	Impact Disturbance Threshold ² (100dB)	Vibratory Disturbance Threshold ² (100dB)
0.410	Steel Pile Installation	0	5*	238	0	0
	Steel Pile Removal	N/A	N/A	315	N/A	0
	Concrete Pile Removal	N/A	N/A	0	N/A	0
Total Action		0	5*	553	0	0

Note: The take estimates include those from impact & vibratory pile driving and pneumatic chipping.

¹ Pneumatic chipping hammers are assessed under the same criteria as vibratory pile driving.

² The airborne exposure calculations assumed that 100% of the in-water densities were available at the surface to be exposed to airborne sound.

*The modeling indicated that zero California sea lions were likely to be exposed to sounds that would qualify as behavioral harassment during impact pile driving (160 dB zone). However, the Navy feels based on the abundance of this species in the waters along NBK, including their presence at nearby haulouts, that it's likely that an individual could pass through this zone in transit to or from a haulout, Therefore, the Navy is requesting a behavioral take of California sea lion by impact pile driving each day of pile driving, for a total of 5 takes.

California sea lions that are taken could exhibit behavioral reactions. Disturbance from underwater noise impacts is not expected to be significant because it is estimated that only a small number of California sea lions may be affected by acoustic harassment. Additionally, marine mammal observers will be monitoring the shutdown and buffer zones (see Section 11 for a detailed discussion of mitigation measures) for the presence of marine mammals, and will alert work crews when to begin or stop work due to presence of sea lions in or near the shutdown and buffer zones, reducing the potential for acoustic harassment. Based on the exposure analysis, no California sea lions are anticipated to experience airborne sound pressure levels that would qualify as harassment.

6.3.8.3 Harbor Seal

Harbor seals are the most abundant marine mammal in the Hood Canal. The Navy conducted boat surveys of the waterfront area in 2008 from July to September (Agness and Tannenbaum 2009a). Harbor seals were sighted during every survey and were found in all marine habitats including near and hauled out on man-made objects such as piers and buoys. Jeffries et al. (2003) completed a more comprehensive stock assessment of the Hood Canal in 1999 and counted 711 harbor seals hauled out. This abundance was adjusted using a correction factor of 1.53 to account for seals in the water and not counted to provide a population estimate of 1,088 harbor seals in the Hood Canal (Jeffries et al 2003). Research by Huber et al. (2001) indicates that approximately 35% of harbor seals are in the water at any one time. Exposures were calculated using a density derived from the number of harbor seals that are present in the water at any one time (35% of 1,088 or ~381 individuals), divided by the area of the Hood Canal (291 km²) and the formula presented in Section 6.7.

While Huber et al.'s (2001) data suggests that harbor seals typically spend 65% of their time hauled out; the Navy's waterfront surveys found that it is extremely rare for harbor seals to haul out in the vicinity of the Project Area. Therefore, the only population of harbor seals that could potentially be exposed to airborne sounds are those that are in-water but at the surface. Based on the diving cycle of tagged harbor seals near the San Juan Islands we can estimate that seals are on the surface approximately 16.4 percent of the of their total in-water duration (Suryan and Harvey 1998). Therefore, by multiplying the percentage of time spent at the surface (16.4%) by the total in-water population of harbor seals at any one time (~381 individuals), the population of harbor seals with the potential to experience airborne impacts (~63 individuals) can be obtained. Airborne exposures were calculated using a density derived from the maximum number of harbor seals available at the surface (~63 individuals), divided by the area of the Hood Canal (291 km²) and the formula presented in Section 6.7. Table 6-22 depicts the number of acoustic harassments that are estimated from vibratory and impact pile driving and pneumatic chipping both underwater and in-air.

Harbor seals that are taken could exhibit behavioral reactions. Disturbance from underwater noise impacts is not expected to be significant because it is estimated that only a small number of harbor seals may be affected by acoustic harassment. Additionally, marine mammal observers will be monitoring the shutdown and buffer zones (see Section 11 for a detailed discussion of mitigation measures) for the presence of marine mammals, and will alert work crews when to begin or stop work due to presence of seals in or near the shutdown and buffer zones, reducing the potential for acoustic harassment. Based on the exposure analysis, no harbor seals are anticipated to experience airborne sound pressure levels that would qualify as harassment.

Table 6-22 Number of Potential Exposures of Harbor Seals within Various Acoustic Threshold Zones

Density at Project Area (July - Oct)	Stage of EHW-1 Action	Underwater			Airborne	
		Impact Injury Threshold (190 dB)	Impact Disturbance Threshold (160dB)	Vibratory ¹ Disturbance Threshold (120 dB)	Impact Disturbance Threshold ² (90dB)	Vibratory Disturbance Threshold ² (90dB)
1.31	Steel Pile Installation	0	5*	742	0	0
	Steel Pile Removal	N/A	N/A	987	N/A	0
	Concrete Pile Removal	N/A	N/A	32	N/A	0
	Total Action	0	5*	1761	0	0

Note: The take estimates include those from impact & vibratory pile driving and pneumatic chipping.

¹ Pneumatic chipping hammers are assessed under the same criteria as vibratory pile driving.

² Airborne densities were based on the percentage (16.4%) of in-water density available on surface to be exposed (Suryan and Harvey, 1998).

* The modeling indicated that zero harbor seals were likely to be exposed to sounds that would qualify as behavioral harassment during impact pile driving (160 dB zone). However, the Navy feels based on the abundance of this species in the waters along NBK, including their presence at nearby haulouts, that it's likely that an individual could pass through this zone in transit to or from a haulout. Therefore, the Navy is requesting a behavioral take of harbor seals by impact pile driving each day of pile driving, for a total of 5 takes.

6.3.8.4 Killer Whale

Transients are uncommon visitors to Hood Canal. In 2003 and 2005, small groups of transient killer whales (6 – 11 individuals per event) visited Hood Canal to feed on harbor seals and remained in the area for significant periods of time (59 – 172 days) between the months of January and July (London 2006). These whales used the entire expanse of Hood Canal for feeding. Subsequent aerial surveys suggest that there has not been a sharp decline in the local seal population from these sustained feeding events (London 2006). Based on this data, the density for Transient killer whales in the Hood Canal for January to June is 0.038/km² (11 individuals divided by the area of the Hood Canal [291 km²]). Since this timeframe overlaps the period in which the Pile Replacement Project will occur (July – Oct), this density was used for all exposure calculations. Exposures were calculated using the formula presented in Section 6.7. Table 6-23 depicts the number of acoustic harassments that are estimated from vibratory and impact pile driving and pneumatic chipping.

Killer whales that are taken could exhibit behavioral changes. Disturbance from underwater noise impacts is not expected to be significant because it is estimated that only a small number of killer whales may be affected by acoustic harassment. Additionally, marine mammal observers will be monitoring the shutdown and buffer zones (see Section 11 for a detailed discussion of mitigation measures) for the presence of marine mammals, and will alert work crews when to begin or stop work due to presence of seals in or near the shutdown and buffer zones, reducing the potential for acoustic harassment.

Table 6-23 Number Potential Exposures of Killer Whales within Various Acoustic Threshold Zones

Density at Project Area (July- Oct)	Stage of EHW-1 Action	Underwater		
		Impact Injury Threshold (180 dB)	Impact Disturbance Threshold (160dB)	Vibratory ¹ Disturbance Threshold (120 dB)
0.038	Steel Pile Installation	0	9*	28
	Steel Pile Removal	N/A	N/A	21
	Concrete Pile Removal	N/A	N/A	0
	Total Action	0	9*	49

Note: The take estimates include those from impact & vibratory pile driving and pneumatic chipping.

¹ Pneumatic chipping hammers are assessed under the same criteria as vibratory pile driving.

*The modeling indicated that zero killer whales were likely to be exposed to sounds that would qualify as behavioral harassment during impact pile driving (160 dB zone). However, while Transient killer whales are rare in the Hood Canal, when these animals are present they occur in pods, so their density in the project area is unlikely to be uniform, as was modeled. If they are present during impact pile driving it's possible that one or more individuals within a pod could travel through the behavioral harassment zone. Therefore, the Navy is requesting nine behavioral takes of Transient killer whales – based on the average size of pods seen previously in the Hood Canal - by impact pile driving.

6.3.8.5 Dall's Porpoise

Dall's porpoise may be present in the Hood Canal year-round and may be expected as the project site. Their use of inland Washington waters, however, is mostly limited to the Strait of Juan de Fuca. The Navy conducted boat surveys of the waterfront area in 2008 from July to September (Agness and Tannenbaum 2009a). During one of the surveys a single Dall's porpoise was sighted in August in the deeper waters off Carlson Spit. In the absence of an abundance estimate for the entire Hood Canal, a seasonal density (warm season only) was derived from the waterfront survey by the number of individuals seen divided by total number of kilometers of survey effort (6 surveys with approximately 3.9km² of effort), assuming strip transect surveys. In absence of any other survey data for the Hood Canal, this density is assumed to be throughout the Project Area. Exposures were calculated using the formula presented in Section 6.7. Table 6-24 depicts the number of acoustic harassments that are estimated from vibratory and impact pile driving and pneumatic chipping underwater.

Dall's porpoise that are taken could exhibit behavioral reactions. Disturbance from underwater noise impacts is not expected to be significant because it is estimated that only a small number of killer whales may be affected by acoustic harassment. Additionally, marine mammal observers will be monitoring the shutdown and buffer zones (see Section 11 for a detailed discussion of mitigation measures) for the presence of marine mammals, and will alert work crews when to begin or stop work due to presence of seals in or near the shutdown and buffer zones, reducing the potential for acoustic harassment.

Table 6-24 Number of Potential Exposures of Dall's Porpoise within Various Acoustic Threshold Zones

Density at Project Area (July- Oct)	Stage of EHW-1 Action	Underwater		
		Impact Injury Threshold (180 dB)	Impact Disturbance Threshold (160dB)	Vibratory ¹ Disturbance Threshold (120 dB)
0.043	Steel Pile Installation	0	1*	28
	Steel Pile Removal	N/A	N/A	42
	Concrete Pile Removal	N/A	N/A	0
	Total Action	0	1*	70

Note: The take estimates include those from impact & vibratory pile driving and pneumatic chipping.

¹ Pneumatic chipping hammers are assessed under the same criteria as vibratory pile driving.

* The modeling indicated that zero Dall's porpoise were likely to be exposed to sounds that would qualify as behavioral harassment during impact pile driving (160 dB zone). Dall's porpoises are rare in the Hood Canal; only one animal, seen located in deep waters offshore the base has been seen in the project area in the past few years. However, it's possible that additional animals exist or that this single individual could pass through the behavioral harassment zone (160 dB) while transiting along the waterfront. Therefore, the Navy is requesting a single behavioral take of Dall's porpoise by impact pile driving.

6.3.8.6 Harbor Porpoise

Harbor porpoises may be present in the Hood Canal year-round, however their presence is rare. The Navy conducted boat surveys of the waterfront area from July to September over the past few years (2008 – present) (Agness and Tannenbaum 2009a). During one of the surveys a single Dall's porpoise was sighted in the deeper waters offshore the waterfront. In the absence of an abundance estimate for the entire Hood Canal, a seasonal density (warm season only) was derived from the waterfront survey by the number of individuals seen divided by total number of kilometers of survey effort (24 surveys with approximately 3.9 km² of effort each), assuming strip transect surveys. In the absence of any other survey data for the Hood Canal, this density is assumed to be throughout the Project Area. Exposures were calculated using the formula presented in Section 6.7. Table 6-25 depicts the number of acoustic harassments that are estimated from vibratory and impact pile driving and pneumatic chipping underwater.

Disturbance from underwater noise impacts is not expected to be significant. Additionally, marine mammal observers will be monitoring the shutdown and buffer zones (see Chapter 4 for a detailed discussion of mitigation measures) for the presence of marine mammals, and will alert work crews when to begin or stop work due to presence of marine mammals in or near the shutdown zones, reducing the potential for acoustic harassment. Potential takes by disturbance would have a negligible short-term effect on individual harbor porpoises and would not result in population-level impacts.

Table 6-25 Number of Potential Exposures of Harbor Porpoise within Various Acoustic Threshold Zones

Density at Project Area (May- Oct)	Stage of EHW-1 Action	Underwater		
		Impact Injury Threshold (180 dB)	Impact Disturbance Threshold (160dB)	Vibratory ¹ Disturbance Threshold (120 dB)
0.011	Steel Pile Installation	0	0	14*
	Steel Pile Removal	N/A	N/A	21*
	Concrete Pile Removal	N/A	N/A	0
	Total Action	0	0	35*

Note: The take estimates include those from impact & vibratory pile driving and pneumatic chipping.

¹ Pneumatic chipping hammers are assessed under the same criteria as vibratory pile driving.

* The modeling indicated that zero harbor porpoise were likely to be exposed to sounds that would qualify as behavioral harassment during vibratory pile driving (120 dB zone). However, while harbor porpoises are rare, one has been sighted in surveys over the last few years in the deep waters offshore the base. It's possible this offshore region is encapsulated within the vibratory disturbance zone during vibratory steel pile installation and removal due to its size (40.273 and 35.87 sq. km, respectively). Therefore the Navy feels based on the possibility of this animal to be present in the offshore waters during every day of construction, the Navy is requesting a single behavioral take of harbor porpoise by vibratory pile driving each day of pile driving, for a total of 35 takes (14 during installation and 21 during removal). The area of disturbance during pneumatic chipping is relatively small 0.608 sq. km) therefore the Navy does not feel harbor porpoises are likely to occur in this area and no additional takes are requested.

6.3.9 Summary of Impacts for Pile Installation and Removal

Based on the modeling results presented above, the total number of takes that the Navy is requesting for the five marine mammals species that may occur within the Project Area during the duration of the EHW-1 Pile Replacement project are presented below in Table 6-26. During the first construction period of the project from July 16, 2011 – October 31, 2011, there is the potential for 20 Level B disturbance takes (160 dB) of various species from impulsive pile driving operations, and an additional 2,468 Level B disturbance takes (120 dB) of various species from vibratory pile installation/removal and pneumatic chipping due to underwater sound. The following species and numbers of Level B disturbance takes could occur due to underwater sound as a result of impact pile driving operations: 5 California sea lions, 5 harbor seals, 9 Transient killer whales, and 1 Dall's porpoise. The following species and numbers of Level B disturbance takes could occur due to underwater sounds as a result of vibratory pile driving and pneumatic chipping operations: 553 California sea lions, 1,761 harbor seals, 49 Transient killer whales, 70 Dall's porpoises, and 35 harbor porpoises. In total, the Navy is requesting 2,488 Level B disturbance takes due to underwater noise from all pile driving operations (including pneumatic chipping) for the EHW-1 Pile Replacement Project. Due to their lack of presence within the project area during the timeframe of the EHW-1 Pile Replacement Project (July 16 – October 31), no ESA-listed Steller sea lions would be acoustically harassed. Lastly, no species of pinnipeds are expected to be exposed to airborne sound pressure levels that would cause harassment.

Table 6-26 Summary of Potential Exposures for All Species During the EHW-1 Pile Replacement Project's Timeframe (July - Oct)

Species	Underwater				Airborne			
	Impact Injury Threshold (190 dB)	Impact Injury Threshold (180dB)	Impact Disturbance Threshold (160dB)	Vibratory Disturbance Threshold (120dB)	Impact Disturbance Threshold (100dB)*	Vibratory Disturbance Threshold (100dB)*	Vibratory Disturbance Threshold (90dB)*	Impact Disturbance Threshold (90dB)*
California sea lion	0	N/A	5*	553	0	0	N/A	N/A
Harbor seal	0	N/A	5*	1761	N/A	N/A	0	0
Transient killer whale	N/A	0	9*	49	N/A	N/A	N/A	N/A
Dall's porpoise	N/A	0	1*	70	N/A	N/A	N/A	N/A
Harbor porpoise	N/A	0	0	35*	N/A	N/A	N/A	N/A
Total	0	0	20*	2468	0	0	0	0

6.4 Effects of Other Construction Activities

Several non-pile installation/removal construction activities will also occur at the project area as part of the proposed action. Among them are the removal of the fragmentation barrier and walkway and the installation of cast-in-place concrete pile caps, passive cathodic protection systems, and the new pre-stressed wharf superstructure and related appurtenances. All of these activities will occur above the water and are likely to have similar impacts to all marine mammal species.

The fragmentation barrier and walkway will be removed from the existing piling supports by cutting the concrete into sections (potentially 3 or 4) using a concrete cutting saw. Each section will be lifted from wharf using a crane and transported to a barge. Pre-cast concrete pile caps will be situated on the tops of steel pipe piles which are located directly beneath the structure (see Figure 1-4) and function as a load transfer mechanism between the superstructure and the piles. The passive cathodic protection system is a metallic rod or anode that is attached to a metal object to protect it from corrosion. The anode is composed of more active metal which is more easily oxidized, corroding first and acting as a barrier against corrosion for the object to which it's attached. At the EHW-1 facility, the passive cathodic protection systems will be banded to the steel piles to prevent the metallic surfaces of the wharf from corroding due to the saline conditions in Hood Canal. The superstructure is the pre-stressed concrete deck for the new wharf section. It will be installed using a crane to situate the concrete slab above the piles. It is supported by the caps or sills. Appurtenances are the associated parts of the superstructure that connect the superstructure to the piles. These pieces include all of the components such as bolts, welded metal hangers and fittings, brackets, etc.

All of these construction activities will occur out of the water/above the water's surface and will be installed on the tops of the piles or attached to the wharf's superstructure. Each of these activities could involve the generation of low levels of noise from the operation of associated installation machinery (i.e. concrete cutting saw, bolt gun, welder, etc.). While no empirical data exists for these construction activities they are expected to be significantly lower than those

estimated for pile installation and removal using an impact/vibratory pile driver or pneumatic chipping hammer. As a result, airborne disturbance is not anticipated for any marine mammal species. It's possible that sound could be transmitted from these activities along the piles' length and enter the water. However, since these activities will be occurring at the top of the pile or on the superstructure, tens of feet above the water, sounds transmitted into the water would be significantly reduced. As a result, underwater acoustic impacts from these construction operations are expected to be minimal and are unlikely to result in harassment of any marine mammal species. Therefore, the Navy is not requesting any additional takes from non-pile installation/removal construction activities.

This Page Intentionally Left Blank

7 IMPACTS TO MARINE MAMMAL SPECIES OR STOCKS

The anticipated impact of the activity upon the species or stock of marine mammals

7.1 Potential Effects of Pile Driving on Marine Mammals

7.1.1 Underwater Noise Effects

The effects of pile driving on marine mammals are dependent on several factors, including the size, type, and depth of both the animal; the depth, intensity, and duration of the pile driving sound; the depth of the water column; the substrate of the habitat; the standoff distance between the pile and the animal; and the sound propagation properties of the environment. Impacts to marine mammals from pile driving activities are expected to result primarily from acoustic pathways. As such, the degree of effect is intrinsically related to the received level and duration of the sound exposure, which are in turn influenced by the distance between the animal and the source. The further away from the source, the less intense the exposure should be. The substrate and depth of the habitat affect the sound propagation properties of the environment. Shallow environments are typically more structurally complex which leads to rapid sound attenuation. In addition, substrates which are soft (i.e. sand) will absorb or attenuate the sound more readily than hard substrates (rock) which may reflect the acoustic wave. Soft porous substrates would also likely require less time to drive the pile, and possibly less forceful equipment, which would ultimately decrease the intensity of the acoustic source.

Impacts to marine species are expected to be the result of physiological responses to both the type and strength of the acoustic signature (Viada et al., 2008). Behavioral impacts are also expected, though the type and severity of these effects are more difficult to define due to limited studies addressing the behavioral effects of impulsive sounds on marine mammals. Potential effects from impulsive sound sources can range from brief acoustic effects such as behavioral disturbance, tactile perception, physical discomfort, slight injury of the internal organs and the auditory system, to death of the animal (Yelverton *et al.*, 1973; O'Keefe and Young, 1984; DoN, 2001b).

Physiological Responses

Direct tissue responses to impact/impulsive sound stimulation may range from mechanical vibration or compression with no resulting injury, to tissue trauma (injury). Because the ears are the most sensitive organ to pressure, they are the organs most sensitive to injury (Ketten, 2000). Sound related trauma can be lethal or sub-lethal. Lethal impacts are those that result in immediate death or serious debilitation in or near an intense source (Ketten, 1995). Sub-lethal impacts include hearing loss, which is caused by exposure to perceptible sounds. Severe damage, from a pressure wave, to the ear can include rupture of the tympanum, fracture of the ossicles, damage to the cochlea, hemorrhage, and cerebrospinal fluid leakage into the middle ear (NMFS, 2008b). Moderate injury implies partial hearing loss. Permanent hearing loss can occur when the hair cells are damaged by one very loud event, as well as prolonged exposure to noise. Instances of temporary threshold shifts (TTS) and/or auditory fatigue are well documented in marine mammal literature as being one of the primary avenues of acoustic impact. Temporary loss of hearing sensitivity (TTS) has been documented in controlled settings using captive marine mammals exposed to strong sound exposure levels at various frequencies (Ridgway et al. 1997; Kastak et al. 1999; Finneran et al. 2005), but it has not been documented in wild marine

mammals exposed to pile driving. While injuries to other sensitive organs are possible, they are less likely since pile driving impacts are almost entirely acoustically mediated, versus explosive sounds which also include a shock wave which can result in damage.

No physiological responses are expected from pile driving operations (including the use of a pneumatic chipping hammer) occurring during the EHW-1 Pile Replacement Project within the Project Area for several reasons. Firstly, vibratory pile driving which is being utilized as the primary installation method, does not generate high enough peak sound pressure levels that are commonly associated with physiological damage. Any use of impulsive pile driving will only occur from a short period of time (~15 min per pile) and only to proof a maximum of five piles. Additionally, the mitigation measures which the Navy will be employing (see Section 11) will greatly reduce the chance that a marine mammal may be exposed to sound pressure levels that could cause physical harm. During impact pile driving the Navy will employ a sound attenuation system (*e.g.* bubble curtain/wall) to attenuate initial sound pressure levels (-10 dB reduction assumed). Furthermore, the Navy will have trained biologists monitoring a shutdown zone equivalent to the Level A Harassment zone (inclusive of the 180 dB re 1 μ Pa (cetaceans) and 190 dB re 1 μ Pa (pinnipeds) isopleths) to ensure no marine mammals are injured.

Behavioral Responses

Behavioral responses to sound are highly variable and context specific. For each potential behavioral change, the magnitude of the change ultimately determines the severity of the response. A number of factors may influence an animal's response to noise, including its previous experience, its auditory sensitivity, its biological and social status (including age and sex), and its behavioral state and activity at the time of exposure.

Habituation can occur when an animal's response to a stimulus wanes with repeated exposure, usually in the absence of unpleasant associated events (Wartzok et al. 2003/04). Animals are most likely to habituate to sounds that are predictable and unvarying. The opposite process is sensitization, when an unpleasant experience leads to subsequent responses, often in the form of avoidance, at a lower level of exposure. Behavioral state may affect the type of response as well. For example, animals that are resting may show greater behavioral change in response to disturbing noise levels than animals that are highly motivated to remain in an area for feeding (Richardson et al. 1995; NRC 2003; Wartzok et al. 2003/04).

Controlled experiments with captive marine mammals showed pronounced behavioral reactions, including avoidance of loud sound sources (Ridgway et al. 1997; Finneran et al. 2003). Observed responses of wild marine mammals to loud pulsed sound sources (typically seismic guns or acoustic harassment devices, and also including pile driving) have been varied but often consist of avoidance behavior or other behavioral changes suggesting discomfort (Morton and Symonds 2002; CALTRANS 2001, 2006; also see reviews in Gordon et al. 2004; Wartzok et al. 2003/04; and Nowacek et al. 2007). Responses to continuous noise, such as vibratory pile installation, have not been documented as well as responses to pulsed sounds.

With regard to pile driving (and the use of a pneumatic chipping hammer), it is likely that the onset of pile driving could result in temporary, short term changes in the animal's typical behavior and/or avoidance of the affected area. A marine mammal may show signs that it is startled by the noise and/or may swim further away from the sound source and avoid the area. Other potential behavioral changes could include increased swimming speed, increased surfacing time, and decreased foraging in the affected area. Since pile driving will likely occur for a few

hours a day, over a short period of time, it is unlikely to result in permanent displacement. Any potential impacts from pile driving activities could be experienced by individual marine mammals, but would not cause population level impacts, or affect the long-term fitness of the species.

7.1.2 Airborne Noise Effects

Marine mammals that occur in the project area could be exposed to airborne sounds associated with pile driving that have the potential to cause harassment, depending on their distance from pile driving activities. Airborne pile driving noise would have less impact on cetaceans than pinnipeds because noise from atmospheric sources does not transmit well underwater (Richardson et al. 1995); thus airborne noise would only be an issue for hauled-out pinnipeds in the Project Area. Most likely, airborne sound would cause behavioral responses similar to those discussed above in relation to underwater noise. For instance, anthropogenic sound could cause hauled out pinnipeds to exhibit changes in their normal behavior, such as reduction in vocalizations, or cause them to temporarily abandon their habitat and move further from the source. Studies by Blackwell et al. (2004) and Moulton et al. (2005) indicate a tolerance or lack of response to unweighted airborne sounds as high as 112 dB peak and 96 dB rms. Based on these observations marine mammals could exhibit temporary behavioral reactions to airborne noise, however, exposure is not likely to result in population level impacts. Despite taking into consideration all known and incidental haulout locations nearby the Project Area the exposure modeling indicated that no pinniped species would be exposed to airborne noise levels at sound pressure levels that would constitute Level B behavioral harassment during either impact or vibratory pile driving (see Section 6 for modeling results). Injury or Level A harassment is not expected to occur from airborne noise. In conclusion, this is a negligible impact.

7.2 Conclusions Regarding Impacts to Species or Stocks

Individual marine mammals may be exposed to sound pressure levels during pile driving operations at NBK Bangor which may result in Level B Behavioral harassment. Any marine mammals which are taken (harassed), may change their normal behavior patterns (i.e. swimming speed, foraging habits, etc.) or be temporarily displaced from the area of construction. Any takes would likely have only a minor effect on individuals and no effect on the population. The sound generated from vibratory pile driving is non-pulsed (e.g., continuous) which is not known to cause injury to marine mammals. Mitigation is likely to avoid most potential adverse underwater impacts to marine mammals from impact pile driving. Nevertheless, some level of impact is unavoidable. The expected level of unavoidable impact (defined as an acoustic or harassment "take") is described in Sections 6 and 7. This level of effect is not anticipated to have any detectable adverse impact on population recruitment, survival or recovery (i.e., no more than a negligible adverse effect).

This Page Intentionally Left Blank

8 IMPACT ON SUBSISTENCE USE

The anticipated impact of the activity on the availability of the species or stock of marine mammals for subsistence uses.

8.1 Subsistence Harvests by Northwest Treaty Indian Tribes

Historically, Pacific Northwest treaty Indian tribes were known to utilize (hunt) several species of marine mammals including, but not limited to: harbor seals, Steller sea lions, northern fur seals, gray whales, and humpback whales (Norberg pers comm. 2007). Recently, several Pacific Northwest treaty Indian tribes have promulgated⁶ tribal regulations allowing tribal members to exercise treaty rights for subsistence harvest of California sea lions and harbor seals (Caretta et al. 2007). The Makah Indian Tribe (Makah) has specifically passed hunting regulations for gray whales (Norberg pers comm. 2007b). However, the directed take of marine mammals (not just gray whales) for ceremonial and/or subsistence purposes was enjoined⁷ by the Ninth Circuit Court of Appeals in a ruling against the Makah in 2002, 2003, and 2004 (Norberg pers comm. 2007b; NMFS 2007). The issues surrounding the Makah gray whale hunt, (in addition to the hunt for marine mammals in general) is currently in litigation or not yet clarified in recent court decisions (Wright pers. comm. 2007). These issues also require National Environmental Policy Act (NEPA) and MMPA compliance, which has not yet been completed. Presently, there are no known active ceremonial and/or subsistence hunts for marine mammals in Puget Sound or the San Juan Islands.

- Tribes along the coast are most likely to still have regulations in place allowing a small number of directed takes for subsistence purposes. It is unlikely that those regulations have been exercised in recent years, but they are still likely on the books (Wright pers. comm. 2007).
- Many tribes in Puget Sound and on the coast do have additional regulations that allow their fishermen to protect their life, gear, and catch from seals and California sea lions by lethal means. These rare takes, which are not for subsistence or ceremonial needs, are reported annually to NMFS by each tribe (T. Wright pers. comm. 2007).

8.1.1 Harbor Seals

There have been only a few reported takes of harbor seals from directed tribal subsistence hunts (Caretta et al. 2007). It is possible that very few seals have been taken in directed hunts because tribal fishers use seals caught incidental to fishing operations in the northern Washington marine set gillnet and Washington Puget Sound Region treaty salmon gillnet fisheries, for their subsistence needs before undertaking a ceremonial or subsistence hunt (Caretta et al. 2007). From communications with the tribes, the NMFS Northwest Regional Office believes that zero to five harbor seals from this stock (the Washington Inland Waters Stock) may be taken annually in directed subsistence harvests (Caretta et al. 2007).

⁶ To make known by open declaration; publish; proclaim formally or put into operation (a law, decree of a court, etc.).

⁷ To prohibit or forbid

No impacts to the availability of the species or stock to the Pacific Northwest treaty tribes are expected as a result of the proposed activities.

8.1.2 California Sea Lions

Current estimates of annual subsistence take are zero to two animals per year (Caretta et al. 2007).

No impacts to the availability of the species or stock to the Pacific Northwest treaty tribes are expected as a result of the proposed activities.

8.1.3 Gray Whales

The Makah ceased whaling in the 1920's after commercial whaling decimated the eastern North Pacific (ENP) gray whale population (NMFS 2007). On June 16, 1994, gray whales were removed from the endangered species list after a determination that the population has "recovered to near its estimated original population size and is neither in danger of extinction throughout all or a significant portion of its range, nor likely to again become endangered within the foreseeable future throughout all or a significant portion of its range." (59 FR 31094). On May 5, 1995 the Makah formally notified the U.S. Government of their interest in resuming treaty ceremonial and subsistence harvest of ENP gray whales, asking the Department of Commerce to represent them in seeking approval from the International Whaling Commission (IWC) for an annual quota (NMFS 2007). On October 18, 1997 the IWC approved an aboriginal subsistence quota of 620 ENP gray whales (with an annual cap of 140) for the Russian Chukotka people and the Makah (Angliss and Outlaw 2005; NMFS 2007). On May 17, 1999 the Makah hunt, strike and land one ENP gray whale (sic) (NMFS 2005).

On December 20, 2002 the Ninth Circuit Court of Appeals ruled that an Environmental Impact Statement (EIS) (rather than an Environmental Assessment [EA]) should have been prepared under the NEPA and that the Makah must comply with the process prescribed in the MMPA for authorizing take of marine mammals otherwise prohibited by a moratorium (NMFS 2007). This was further upheld at rulings in 2003 and 2004 (NMFS 2007).

At the most recent meetings of the IWC (59th Annual Meeting in Anchorage, Alaska from May 28 - 31, 2007), an aboriginal subsistence quota for gray whales was again approved for natives in Russia and 20 whales or 4 per year for 5 years for the Makah (Norberg pers comm. 2007), but under the Ninth Circuit Court ruling the Makah must first obtain a waiver of the MMPA take moratorium before harvesting under their IWC quota (Norberg pers comm. 2007b). NMFS is currently preparing an EIS to examine the alternatives for a decision to approve or deny such a waiver (Norberg pers comm. 2007b).

Gray whales migrate north and south along the coast of Washington and there is a regular group of gray whales that enter the Puget Sound waters (specifically Saratoga passage on the eastern side of Whidbey Island) to feed during early spring and summer (March through May/June) (Calambokidis pers comm. 2007). These whales movements do not overlap with the proposed activities.

Should the Makah tribe resume hunting gray whales, this hunt would occur along the outer coast of Washington, but not likely within the region of activity located in the inland waters of Washington. In addition, all in-water work for Hood Canal is restricted to only take place between July 16 through February 15 of any year (or a more compressed schedule within those dates), thus the proposed activities would not occur when the majority of the gray whales are

present in Puget Sound (from March through May/June). Furthermore, underwater noise disturbance would not displace the whales because of the distance between Hood Canal and Saratoga Passage and the presence of landmasses between the two locations. Therefore, the proposed activities would not directly or indirectly interfere with or affect the hunt, should it occur during the proposed project activities.

8.2 Summary

Potential impacts resulting from the Proposed Action will be limited to individuals of marine mammal species located in the marine waters near NBK Bangor and will be limited to Level B harassment. Therefore, no impacts on the availability of species or stocks for subsistence use were found.

This Page Intentionally Left Blank

9 IMPACTS TO THE MARINE MAMMAL HABITAT AND THE LIKELIHOOD OF RESTORATION

The anticipated impact of the activity upon the habitat of the marine mammal populations, and the likelihood of restoration of the affected habitat.

The proposed activities at NBK Bangor would have potential short-term impacts to food sources such as forage fish, but would not result in any permanent impacts on habitats used by marine mammals. There are no rookeries or major haul-out sites within 10km, foraging hotspots, or other ocean bottom structure of significant biological importance to marine mammals that may be present in the marine waters in the vicinity of the Project Area. Therefore, the main impact issue associated with the proposed activity will be temporarily elevated noise levels and the associated direct effects on marine mammals, as discussed in Sections 6 and 7. The most likely impact to marine mammal habitat occurs from pile driving effects on likely marine mammal prey (i.e., fish) nearby NBK Bangor and minor impacts to the immediate substrate during installation and removal of piles during the EHW-1 Pile Replacement Project.

9.1 Pile Driving Effects on Potential Prey (Fish)

Construction activities will produce both pulsed (i.e. impact pile driving) and continuous sounds (i.e. vibratory pile driving, pneumatic chipping). Fish react to sounds which are especially strong and/or intermittent low-frequency sounds. Short duration, sharp sounds can cause overt or subtle changes in fish behavior and local distribution. Hastings and Popper (2005, 2009) identified several studies that suggest fish may relocate to avoid certain areas of noise energy. Additional studies have documented effects of pile driving (or other types of continuous sounds) on fish, although several are based on studies in support of large, multiyear bridge construction projects (Scholik and Yan 2001, 2002, Govoni et al. 2003, Hawkins 2005, Hastings 1990, 2007, Popper et al. 2006, Popper and Hastings 2009). Sound pulses at received levels of 160 dB re 1 μ Pa may cause subtle changes in fish behavior. SPLs of 180 dB may cause noticeable changes in behavior (Chapman and Hawkins 1969; Pearson et al. 1992; Skalski et al. 1992). SPLs of sufficient strength have been known to cause injury to fish and fish mortality (CalTrans 2001; Longmuir and Lively 2001). Fish that occur in the immediate project area would be exposed to underwater noise that could injure or disturb fish during pile driving activity. Because vibratory pile driving and pneumatic chipping are the primary installation and removal methodologies, the most likely impact to fish from pile driving activities at the project area would be temporary behavioral disturbance or avoidance of the area. The duration of fish avoidance of this area after pile driving stops is unknown, but a rapid return to normal recruitment, distribution and behavior is anticipated. In general, impacts to marine mammal prey species are expected to be minor and temporary due to the short-time frame for the Pile Replacement Project. However, moderate impacts may occur to a few species of rockfish (bocaccio, yellowweye, and canary rockfish) and salmon (chinook and summer run chum) which may still be present in the project area despite operating in a reduced work-window in an attempt to avoid important fish spawning time periods. Impacts to these species could result from potential impacts to their eggs and larvae.

9.2 Pile Driving Effects on Potential Foraging Habitat

In addition, the area likely impacted by the EHW-1 Pile Replacement Project is relatively small compared to the available habitat in the Hood Canal. Potentially a maximum of 0.46 m² (based

on a 30-inch diameter pile) of marine mammal foraging habitat may have decreased foraging value as each pile is driven. Avoidance by potential prey (i.e. fish) of the immediate area due to the temporary loss of this foraging habitat is also possible. The duration of fish avoidance of this area after pile driving stops is unknown, but a rapid return to normal recruitment, distribution and behavior is anticipated. Any behavioral avoidance by fish of the disturbed area would still leave significantly large areas of fish and marine mammal foraging habitat in the Hood Canal and nearby vicinity.

9.3 Summary of Impacts to Marine Mammal Habitat

Given the short daily duration of noise associated with individual pile driving\removal, the short duration of the entire Pile Replacement Project (<110 days per year), and the relatively small areas being affected, pile driving activities associated with the proposed action are not likely to have a permanent, adverse effect on any EFH, or population of fish species. Therefore, pile driving\removal is not likely to have a permanent, adverse effect on marine mammal foraging habitat at the Project Area.

10 IMPACTS TO MARINE MAMMALS FROM LOSS OR MODIFICATION OF HABITAT

The anticipated impact of the loss or modification of the habitat on the marine mammal populations involved.

The proposed activities at NBK Bangor are not expected to have any habitat-related effects that could cause significant or long-term consequences for individual marine mammals or their populations, since pile driving will be temporary and all pile are being installed within the existing footprint of the current EHW-1 facility. Based on the discussions in Section 9, there will be no impacts to marine mammals resulting from loss or modification of marine mammal habitat.

This Page Intentionally Left Blank

11 MEANS OF EFFECTING THE LEAST PRACTICABLE ADVERSE IMPACTS – MITIGATION MEASURES

The availability and feasibility (economic and technological) of equipment, methods, and manner of conducting such activity or other means of effecting the least practicable adverse impact upon the affected species or stocks, their habitat, and on their availability for subsistence uses, paying particular attention to rookeries, mating grounds, and areas of similar significance.

The exposures outlined in Section 6 represent the maximum expected number of marine mammals that could be exposed to acoustic sources reaching Level B harassment levels. Navy proposes to employ a number of mitigation measures, discussed below, in an effort to minimize the number of marine mammals potentially affected.

11.1 Mitigation for Pile Driving Activities

The modeling results for zones of influences (ZOIs) discussed in Section 6 were used to develop mitigation measures for pile installation/removal activities at NBK Bangor. The ZOIs effectively represent the mitigation zone that would be established around each pile to prevent Level A harassment to marine mammals. While the ZOIs vary between the different diameter piles and types of installation and removal methodologies, the Navy is proposing to establish mitigation zones for the maximum zone of influence for all pile installation and removal activities conducted to support of the EHW-1 Pile Replacement Project.

1. Shutdown and Buffer Zone -

- The shutdown zone shall include all areas where the underwater sound pressure levels (SPLs) are anticipated to equal or exceed the Level A (injury) Harassment criteria for marine mammals (180 dB isopleth for cetaceans; 190 dB isopleth for pinnipeds).
- The buffer zone shall include all areas where the underwater or airborne sound pressure levels are anticipated to equal or exceed the Level B (disturbance) Harassment criteria for marine mammals (160 dB re 1 μ Pa or 90 dB re 20 μ Pa isopleths). The distance encompassing these zones will be adjusted to accommodate any difference between predicted and measured sound levels.
- The shutdown and buffer zones will be monitored throughout the time required to drive a pile. If a marine mammal is observed entering the buffer zone, a “take” would be recorded and behaviors documented. However, that pile segment would be completed without cessation, unless the animal approaches/enters the shutdown zone, at which point all pile driving activities will be halted.
- All buffer and shutdown zones will initially be based on the distances from the source which were predicted for each threshold level. However, in-situ acoustic monitoring will be utilized to determine the actual distances to these threshold zones, and the size of the shutdown and buffer zones will be adjusted accordingly (increased or decreased) based on received sound pressure levels.

2. Visual Monitoring -

- a. Impact Installation: Monitoring will be conducted for a 50 meter* shutdown zone (Level A harassment) and a 501 meter buffer zone (Level B harassment) surrounding each pile for the presence of marine mammals before, during, and after pile driving activities. The buffer zone was set at 501 meters, since this is the largest Level B behavioral disturbance zone calculated for impact pile driving. It is based on the calculations for airborne noise for harbor seals. Monitoring will take place from 30 minutes prior to initiation through 30 minutes post-completion of pile installation/removal activities.

Vibratory Installation: Monitoring will be conducted for a 50 meter* shutdown zone. The 120 dB disturbance criterion predicts an affected area of 40.3 sq. km. Due to the difficulty of effectively monitoring such a large area, the Navy intends to monitor a buffer zone equivalent to the size of the Level B disturbance zone for impact pile driving (501 m) surrounding each pile for the presence of marine mammals before, during, and after pile driving activities. Sightings occurring outside this area will still be recorded and noted as a take, but detailed observations outside this zone will not be possible. Monitoring will take place from 30 minutes prior to initiation through 30 minutes post-completion of pile driving activities.
 - b. Monitoring will be conducted by qualified observers. A trained observer will be placed from the best vantage point(s) practicable (*e.g.* from a small boat, the pile driving barge, on shore, or any other suitable location) to monitor for marine mammals and implement shut-down/delay procedures when applicable by calling for the shut-down to the hammer operator.
 - c. Prior to the start of pile driving activity, the shutdown and safety zones will be monitored for 30 minutes to ensure that it is clear of marine mammals. Pile driving will only commence once observers have declared the shutdown zone clear of marine mammals; Animals will be allowed to remain in the buffer zone and their behavior will be monitored and documented.
 - d. If a marine mammals approaches/enters the shutdown zone during the course of pile driving operations, pile driving will be halted and delayed until either the animal has voluntarily left and been visually confirmed beyond the shutdown zone or 30 minutes have passed without re-detection of the animal.
2. Sound Attenuation Devices – Sound attenuation devices (*e.g.* bubble curtain/wall, etc.) will be utilized during all impact pile driving operations. Impact pile driving will only be used on a maximum of five piles (one pile per day) for 15 minutes each. Historically, impact pile

* Based on coordination with NMFS HQ, a minimum shutdown zone of 50 meters was recommended to standardize monitoring for future activities, even though this zone is slightly larger than the modeled Level A harassment zone. This mitigation only applies to marine mammals. This measure will be carried out for impact and vibratory pile driving and pneumatic chipping. However, Navy may seek future revision of this mitigation as it applies to pneumatic chipping hammers, in subsequent IHA/LOA requests, as additional empirical data becomes available regarding the sound pressure levels produced by this machinery.

driving has not be required at the EHW-1 location in the past, and is only expected to be required to “proof” or drive the last 10-15 ft of each pile.

3. Acoustic Measurements – Acoustic measurements will be used to empirically verify the proposed shutdown and buffer zones. For further detail regarding our acoustic monitoring plan see Section 13.
4. Timing Restrictions - The Navy, in consultation with NMFS NW region and USFWS under ESA, has set timing restrictions for pile installation and removal activities to avoid in-water work when ESA-listed fish populations are most likely to be present. Therefore, all pile driving would occur only during the work window from July 16 through October 31st (impact pile driving only between July 16 to September 30th) of any year, to minimize the number of fish exposed to underwater noise and other disturbance. These months (July – Oct) also overlap with times when Steller sea lions and most California sea lions are not expected to be present within the study area.
5. Soft Start - The use of a soft-start procedure is believed to provide additional protection to marine mammals by providing a warning and/or giving marine mammals a chance to leave the area prior to the hammer operating at full capacity. The Pile Replacement Project will use existing soft-start (ramp-up/dry fire) techniques recommended by NMFS for impact pile driving. These measures are as follows:

“The soft-start requires contractors to initiate noise from vibratory hammers for 15 seconds at reduced energy followed by a 1-minute waiting period. This procedure should be repeated two additional times. If an impact hammer is used, contractors are required to provide an initial set of three strikes from the impact hammer at 40 percent energy, followed by a 1-minute waiting period, then two subsequent 3-strike sets.”

6. Daylight Construction – Pile driving will only be conducted between two hours post-sunrise through two hours prior to sunset.

11.2 Mitigation Effectiveness

It should be recognized that although marine mammals will be protected from Level A harassment by the utilization of a bubble curtain/wall and marine mammal observers (MMOs) monitoring the near-field injury zones, mitigation may not be one hundred percent effective at all times in locating marine mammals in the buffer zone. The efficacy of visual detection depends on several factors including the observer’s ability to detect the animal, the environmental conditions (visibility and sea state), and monitoring platforms.

All observers utilized for mitigation activities will be experienced biologists with training in marine mammal detection and behavior. Due to their specialized training the Navy expects that visual mitigation will be highly effective. Trained observers have specific knowledge of marine mammal physiology, behavior, and life-history which may improve their ability to detect individuals or help determine if observed animals are exhibiting behavioral reactions to construction activities.

The Puget Sound region, including the Hood Canal, only infrequently experience winds with velocities in excess of 25 knots (Morris et al. 2008). The typically light winds afforded by the surrounding highlands coupled with the fetch limited environment of the Hood Canal result in

relatively calm wind and sea conditions throughout most of the year. The EHW-1 Pile Replacement project site has a maximum fetch of 8.4 miles to the north, and 4.2 miles to the south, resulting in maximum wave heights of from 2.85-5.1 feet (BSS between 2-4), even in extreme conditions (30 knot winds) (CERC 1984). Visual detection conditions are considered optimal in BSS conditions of 3 or less, which align with the conditions that should be expected for the Pile Replacement Project at NBK Bangor.

Observers will be positioned in locations which provide the best vantage point(s) for monitoring. This will probably be an elevated position as they provide a better range of viewing angles. Also, the shutdown and buffer zone has a relatively small radius to monitor which should improve detectability.

12 MINIMIZATION OF ADVERSE EFFECTS ON SUBSISTENCE USE

Where the proposed activity would take place in or near a traditional Arctic subsistence hunting area and/or may affect the availability of a species or stock of marine mammal for Arctic subsistence uses, the applicant must submit either a plan of cooperation or information that identifies what measures have been taken and/or will be taken to minimize any adverse effects on the availability of marine mammals for subsistence uses. A plan must include the following:

- (i) A statement that the applicant has notified and provided the affected subsistence community with a draft plan of cooperation;*
- (ii) A schedule for meeting with the affected subsistence communities to discuss proposed activities and to resolve potential conflicts regarding any aspects of either the operation or the plan of cooperation;*
- (iii) A description of what measures the applicant has taken an/or will take to ensure that proposed activities will not interfere with subsistence whaling or sealing; and*
- (iv) What plans the applicant has to continue to meet with the affected communities, both prior to and while conducting activity, to resolve conflicts and to notify the communities of any changes in the operation.*

Subsistence use is the traditional exploitation of marine mammals by native peoples for their own consumption. Based on the discussions in Section 8, there are no adverse effects on the availability of species or stocks for subsistence use.

This Page Intentionally Left Blank

13 MONITORING AND REPORTING MEASURES

The suggested means of accomplishing the necessary monitoring and reporting that will result in increased knowledge of the species, the level of taking or impacts on populations of marine mammals that are expected to be present while conducting activities and suggested means of minimizing burdens by coordinating such reporting requirements with other schemes already applicable to persons conducting such activity. Monitoring plans should include a description of the survey techniques that would be used to determine the movement and activity of marine mammals near the activity site(s) including migration and other habitat uses, such as feeding.

13.1 Monitoring Plan

The following monitoring measures would be implemented along with the mitigation measures (Section 11) in order to reduce impacts to marine mammals to the lowest extent practicable. The monitoring plan includes the following components: acoustic measurements and visual observations.

13.1.1 Acoustic Measurements

The Navy will conduct acoustic monitoring for impact driving of steel piles in order to determine the actual distances to the 190 dB re 1 μ Pa rms/180 dB re 1 μ Pa rms and the 160 dB re 1 μ Pa rms isopleths and to determine the relative effectiveness of the bubble curtain/wall system at attenuating noise underwater. The Navy will also conduct acoustic monitoring for vibratory pile driving in order to determine the actual distance to the 120 dB re 1 μ Pa rms isopleth for behavioral harassment relative to background levels. Acoustic monitoring will occur for each type of pile installation and removal methodology, including impact and vibratory pile driving and pneumatic chipping. The monitoring plan addresses both underwater and airborne sounds from the EHW-1 Pile Replacement Project.

At a minimum, the methodology includes:

- For underwater recordings, a stationary hydrophone placed at mid-water depth and 10 meters from the source pile to measure the effectiveness of the bubble curtain system; A weighted tape measure will be used to determine the depth of the water. The hydrophone will be attached to a nylon cord or steel chain if current is swift enough, to maintain a constant distance from the pile. The nylon cord or chain will be attached to a float or tied to a static line at the surface 10 meters from the piles.
- For each monitored location, a two-hydrophone set-up will be used, with the first hydrophone at mid-depth and the second hydrophone at ~1 meter from the bottom in order to evaluate site specific attenuation and propagation characteristics that by be present throughout the water column.
- For underwater measurements, in addition to determining the area encompassed by the 190, 180, 160, and 120 dB RMS isopleths for marine mammals, hydrophones would also be placed at other distances and or depths as appropriate to accurately capture the spreading loss which occurs at the EHW-1 project area, or to determine the distance to the thresholds for fish and birds (these include peak, rms, and sound exposure levels [SEL]);

- For airborne recordings, a stationary hydrophone will be placed at 50 feet (15.24 meters) from the source for initial reference recordings.
- For airborne measurements, in addition to determining the area encompassed by the 100 and 90 dB RMS isopleths for pinnipeds and harbor seals, hydrophones will be placed at other distances as appropriate to accurately capture spreading loss which occurs at the EHW-1 project area, or to determine the distance to thresholds for birds.
- All hydrophones will be calibrated at the start of the action and will be checked at the beginning of each day of monitoring activity.
- Ambient conditions, both airborne and underwater, would be measured at the project site in the absence of construction activities to determine background sound levels. Ambient levels are intended to be recorded over the frequency range from 10 Hz to 20 kHz.
- Sound levels associated with soft-start techniques will also be measured
- Underwater sound pressure levels would be continuously monitored during the entire duration of each pile being driven. Sound pressure levels will be monitored in real time. Sound levels will be measured in Pascals which are easily converted to decibel (dB) units.
- Airborne levels would be recorded as unweighted, as well as in dBA and the distance to marine mammal and/or avian thresholds (respectively) would be measured;
- The effectiveness of the bubble curtain/wall system will be tested in-situ. The following on/off regime is anticipated to be utilized during impact pile driving:

Pile Driving Timeframe	Sound Attenuation Device Condition
Initial 30 seconds	Off
Next minute (minimum)	On
Middle of pile driving segment 30 seconds	Off
Next minute (minimum)	On
Final 30 seconds	Off

- Environmental data would be collected including but not limited to: wind speed and direction, air temperature, humidity, surface water temperature, water depth, wave height, weather conditions and other factors that could contribute to influencing the airborne and underwater sound levels (e.g. aircraft, boats, etc.);
- The chief inspector would supply the acoustics specialist with the substrate composition, hammer model and size, hammer energy settings and any changes to those settings during the piles being monitored, depth of the pile being driven, and blows per foot for the piles monitored.
- Post-analysis of the sound level signals will include determination of absolute peak overpressure and under pressure levels recorded for each pile, Root Mean Square (RMS) value for each absolute peak pile strike, rise time, average duration of each pile strike, number of strikes per pile, Sound Exposure Level (SEL) of the absolute peak pile strike, mean SEL, and cumulative SEL (Accumulated SEL = single strike SEL + 10*log (# hammer strikes) and a frequency spectrum both with and without mitigation, between 10 and 20,000 Hz for up to eight successive strikes with similar sound levels.

13.1.2 Visual Marine Mammal Observations

The Navy will collect sighting data and behavioral responses to construction for marine mammal species observed in the region of activity during the period of construction. All observers will be trained in marine mammal identification and behaviors. NMFS requires that the observers have no other construction related tasks while conducting monitoring.

13.1.3 Methods of Monitoring

The Navy will monitor the shut down zone and safety zone before, during, and after pile driving. Based on NMFS requirements, the Marine Mammal Monitoring Plan would include the following procedures for impact pile driving:

- MMOs would be located at the best vantage point(s) in order to properly see the entire shut down zone and safety zone. This may require the use of a small boat to monitor certain areas while also monitoring from one or more land based vantage points;
- During all observation periods, observers would use binoculars and the naked eye to search continuously for marine mammals;
- To verify the required monitoring distances, the zones would be clearly marked with buoys or other suitable aquatic markers;
- If the shut down or safety zones are obscured by fog or poor lighting conditions, pile driving would not be initiated until all zones are visible;
- The shut down and safety zones around the pile will be monitored for the presence of marine mammals before, during, and after any pile driving activity;
- Pre-Activity Monitoring:
 - The shut down and buffer zones will be monitored for 30 minutes prior to initiating the soft start for pile driving. If marine mammal(s) are present within the shut down prior to pile driving or during the soft start, the start of pile driving would be delayed until the animal(s) leave the shut down zone. Pile driving would resume only after the MMO has determined, through sighting or by waiting approximately 30 minutes that the animal(s) has moved outside the shut down zone.
- During Activity Monitoring:
 - The shutdown and buffer zones will also be monitored throughout the time required to drive a pile. If a marine mammal is observed entering the buffer zone, a “take” would be recorded and behaviors documented. However, that pile segment would be completed without cessation, unless the animal enters or approaches the shutdown zone, at which point all pile driving activities will be halted. Pile driving can only resume once the animal has left the shutdown zone of its own volition or has not been re-sighted for a period of 30 minutes.
- Post-Activity Monitoring: Monitoring of the shutdown and buffer zones would continue for 30 minutes following the completion of pile driving.

13.1.4 Data Collection

NMFS requires that the MMOs use NMFS-approved sighting forms. NMFS requires that a minimum, the following information be collected on the sighting forms:

- Date and time that pile driving begins or ends;
- Construction activities occurring during each observation period;
- Weather parameters identified in the acoustic monitoring (e.g. wind, humidity, temperature);
- Tide state and water currents;
- Visibility;
- Species, numbers, and if possible sex and age class of marine mammals;
- Marine mammal behavior patterns observed, including bearing and direction of travel, and if possible, the correlation to sound pressure levels;
- Distance from pile driving activities to marine mammals and distance from the marine mammal to the observation point;
- Locations of all marine mammal observations;
- Other human activity in the area.

Additionally, based on recent discussions with NMFS HQ, they request that the Navy record behavioral observations such that, if possible, the Navy can attempt to determine whether animals can be (or are) “taken” by more than one sound source in a day’s operations. For instance, the Navy has agreed to: “Note in behavioral observations, to the extent practicable, if an animal has remained in the area during construction activities. Therefore, it may be possible to identify if the same animal or a different individuals are being taken.”

13.2 Reporting

A draft report would be submitted to NMFS within 45 days of the completion of acoustic measurements and marine mammal monitoring. The results would be summarized in graphical form and include summary statistics and time histories of sound values for each pile. Acoustic measurements will be reported for each type of pile installation and removal methodology, including impact and vibratory pile driving and pneumatic chipping. A final report would be prepared and submitted to the NMFS within 30 days following receipt of comments on the draft report from the NMFS. At a minimum, the report shall include:

- Size and type of piles.
- A detailed description of the bubble curtain/wall, including design specifications.
- The impact or vibratory hammer force used to drive/extract the piles.
- A description of the monitoring equipment.
- The distance between hydrophone(s) and pile.
- The depth of the hydrophone(s).

- The depth of water in which the pile was driven.
- The depth into the substrate that the pile was driven.
- The physical characteristics of the bottom substrate into which the piles were driven.
- The ranges and means for peak, RMS, and SEL's for each pile.
- The results of the acoustic measurements, including the frequency spectrum, peak and RMS SPL's, and single-strike and cumulative SEL with and without the attenuation system.
- The results of the airborne noise measurements including dBA and unweighted levels.
- A description of any observable marine mammal, fish, or bird behavior in the immediate area and, if possible, the correlation to underwater sound levels occurring at that time.
- Results: Including the detectability of marine mammals, species and numbers observed, sighting rates and distances, behavioral reactions within and outside of safety zones.
- A refined take estimate based on the number of marine mammals observed in the safety and buffer zones. This may be reported as one or both of the following: a rate of take (number of marine mammals per hour), or take based on density (number of individuals within the area).

This Page Intentionally Left Blank

14 RESEARCH

Suggested means of learning of, encouraging, and coordinating research opportunities, plans, and activities relating to reducing such incidental taking and evaluating its effects.

To minimize the likelihood that impacts will occur to the species, stocks and subsistence use of marine mammals, all construction activities will be conducted in accordance with all federal, state and local regulations and minimization measures proposed by the Navy will be implemented to protect marine mammals. The Navy will coordinate all activities with the relevant federal and state agencies. These include, but are not limited to: the NMFS, U.S. Fish and Wildlife Service, U.S. Coast Guard, Federal Energy Regulatory Commission (FERC), U.S. Army Corps of Engineers (USACE), and the Washington Department of Fish and Wildlife (WDFW). The Navy will share field data and behavioral observations on all marine mammals that occur in the project area. Results of each monitoring effort will be provided to NMFS in one summary report within 45 days of the conclusion of monitoring. This information could be made available to regional, state and federal resource agencies, scientists, professors, and other interested private parties upon written request to NMFS.

Additionally the Navy provides a significant amount of funding and support for marine research. The Navy provided \$26 million in Fiscal Year 2008 and \$22 million in Fiscal Year 2009 to universities, research institutions, federal laboratories, private companies, and independent researchers around the world to study marine mammals. Over the past five years the Navy has funded over \$100 million in marine mammal research, with several projects ongoing in Washington.

The Navy sponsors 70% of all U.S. research concerning the effects of human-generated sound on marine mammals and 50% of such research conducted worldwide. Major topics of Navy-supported research include the following:

- Gaining a better understanding of marine species distribution and important habitat areas,
- Developing methods to detect and monitor marine species before and during training,
- Understanding the effects of sound on marine mammals, and
- Developing tools to model and estimate potential effects of sound.

The Navy has sponsored several workshops to evaluate the current state of knowledge and potential for future acoustic monitoring of marine mammals. The workshops brought together acoustic experts and marine biologists from the Navy and other research organizations to present data and information on current acoustic monitoring research efforts and to evaluate the potential for incorporating similar technology and methods in Navy activities. The Navy supports research efforts on acoustic monitoring and will continue to investigate the feasibility of passive acoustics as a potential monitoring tool. Overall, the Navy will continue to research and contribute to university/external research to improve the state of the science regarding marine species biology and acoustic effects. These efforts include monitoring programs; data sharing with National Marine Fisheries Service (NMFS) from research and development efforts; and future research as described previously.

15 LIST OF PREPARERS

Danielle Buonantony, Marine Resource Specialist, NAVFAC Atlantic
M.E.M. Coastal Environmental Management, Duke University
B.S. Zoology, University of Maryland – College Park
Years of Experience: 4

16 REFERENCES

- Agness, A., and B.R. Tannenbaum. 2009a. Naval Base Kitsap at Bangor marine mammal resource report. Prepared by Science Applications International Corporation, Bothell, WA. Prepared for BAE Systems Applied Technologies, Inc., Rockville, MD.
- Agness, A., and B.R. Tannenbaum. 2009b. Naval Base Kitsap at Bangor marine bird resource report. Prepared by Science Applications International Corporation, Bothell, WA. Prepared for BAE Systems Applied Technologies, Inc., Rockville, MD.
- Allen, B. M., and R. P. Angliss. 2010. Alaska Marine Mammal Stock Assessments, 2009. U.S. Dep. Commer., NOAA Technical Memorandum NMFS-AFSC-206, 276 p.
- Andersen, S. 1970. Auditory sensitivity of then harbour porpoise *Phocoena phocoena*. Invest Cetacea. 2: 255-259.
- Angell, T. and K.C. Balcomb III. 1982. *Marine birds and mammals of Puget Sound*. University of Washington Press: Seattle, 145 pp.
- Angliss, R.P. and R.B. Outlaw. 2008. Alaska Marine Mammal Stock Assessments, 2007. NOAA Technical Memorandum NMFS-AFSC-180.
- Angliss, R.P. and R.B. Outlaw. 2005. Alaska Marine Mammal Stock Assessment, 2005. NOAA Technical Memorandum NMFS-AFSC-161.
- ANSI. 1986. Methods for measurement of impulse noise (ANSI S12.7-1986). New York: Acoustical Society of America.
- Antonelis, G.A., Jr., B.S. Stewart, and W.F. Perryman. 1990. Foraging characteristics of female northern fur seals (*Callorhinus ursinus*) and California sea lions (*Zalophus californianus*). Canadian Journal of Zoology 68:150-158.
- Au, W.W.L., J.K.B. Ford, J.K. Horne, and K.A. Newman Allman. 2004. Echolocation signals of free ranging killer whales (*Orcinus orca*) and modeling of foraging for chinook salmon (*Oncorhynchus tshawytscha*). Journal of the Acoustical Society of America 115(2):901-909.
- Baird, R.W. 2001. Status of harbour seals, *Phoca vitulina*, in Canada. Canadian Field-Naturalist 115(4):663-675.
- Baird, R.W. and H. Whitehead. 2000. Social organization of mammal-eating killer whales: Group stability and dispersal patterns. Canadian Journal of Zoology 78:2096-2105.
- Baird, R.W. and L.M. Dill. 1995. Occurrence and behaviour of transient killer whales: Seasonal and pod-specific variability, foraging behaviour, and prey handling. Canadian Journal of Zoology 73:1300- 1311.
- Baird, R.W. and L.M. Dill. 1996. Ecological and social determinants of group size in transient killer whales. Behavioral Ecology 7(4):408-416.
- Barlow, J. and K.A. Forney. 2007. Abundance and population density of cetaceans in the California Current ecosystem. Fishery Bulletin 105:509-526.
- Barlow, J. and D. Hanan. 1995. An assessment of the status of harbor porpoise in central California. Rept. Int. Whal., Special Issue 16:123-140.
- Barrett-Lennard, L. G. 2000. Population structure and mating patterns of killer whales (*Orcinus orca*) as revealed by DNA analysis. Ph.D. Thesis, University of British Columbia, Vancouver, BC, Canada, 97 pp.

- Barlett, M.L., and G.R. Wilson. 2002. Characteristics of small boat signatures. *The Journal of the Acoustical Society of America*. 112(5): 2221.
- Bhuthimethee, M. 2008. Mary Bhuthimethee, Marine Scientist, Science Applications International Corporation, Bothell, WA. November 25, 2008. Personal communication with Bernice Tannenbaum, Wildlife Biologist, Science Applications International Corporation, Bothell, WA, re: Steller sea lions at NAVBASE Kitsap Bangor.
- Bibikov, N.G. 1992. Auditory brainstem responses in the harbor porpoise (*Phocoena phocoena*). In *Marine Mammal Sensory Systems* (ed. J. A. Thomas, R. A. Kastelein and A. Y. Supin), pp. 197- 211. New York: Plenum Press.
- Bigg, M.A. 1981. Harbour seal *Phoca vitulina* Linnaeus, 1758 and *Phoca largha* Pallas, 1811. Pages 1-27 IN: S.H. Ridgway and R. Harrison, eds. *Handbook of marine mammals, Volume 2: Seals*. San Diego: Academic Press.
- BjØrge, A. 2002. How persistent are marine mammal habitats in an ocean of variability? Pages 63-91 IN: P.G.H. Evans, and J.A. Riga, eds. *Marine Mammals: Biology and Conservation*. Kluwer Academic/Plenum Publishers, New York.
- Blackwell, S.B. and C.R. Greene Jr. 2002. Acoustic measurements in Cook Inlet, Alaska during August 2001. Greeneridge Report 271-2. Report from Greeneridge Sciences, Inc., Santa Barbara for National Marine Fisheries Service, Anchorage, AK. 43 p.
- Blackwell, S.B., J.W. Lawson, and M.T. Williams. 2004. Tolerance by ringed seals (*Phoca hispida*) to impact pipe-driving and construction sounds at an oil production island. *Journal of the Acoustical Society of America*. 115(5): 2346-2357
- Boggs, S., Jr. 1995. *Principles in Sedimentology and Stratigraphy, Second Edition*. Prentice-Hall, Inc., Upper Saddle River, NJ.
- Bonnell, M.L. and M.D. Dailey. 1993. Marine mammals. Pages 604-681 in Dailey, M.D., D.J. Reish, and J.W. Anderson, eds. *Ecology of the Southern California Bight: A synthesis and interpretation*. Berkeley, California: University of California Press.
- Bonnell, M.L. and R.G. Ford. 1987. California sea lion distribution: A statistical analysis of aerial transect data. *Journal of Wildlife Management* 51(1):13-20.
- Bonnell, M.L., M.O. Pierson, and G.D. Farrens. 1983. Pinnipeds and sea otters of central and northern California, 1980 - 1983: Status, abundance, and distribution. Volume III, Book 1. OCS Study MMS 84-0044. Los Angeles, California: Minerals Management Service.
- Boveng, P. 1988. Status of the Pacific harbor seal population on the U.S. west coast. Admin. Rep. LJ-88- 06. Southwest Fisheries Science Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA 92038. 43 pp.
- Bowen, W.D., and D.B. Siniff. 1999. Distribution, population biology, and feeding ecology of marine mammals. In *Biology of marine mammals*, ed. Reynolds, J.E. and S.A. Rommel. Washington: Smithsonian Institution Press. 423-484.
- Bowen, W.D., D.J. Boness, and S.J. Iverson. 1999. Diving behaviour of lactating harbour seals and their pups during maternal foraging trips. *Canadian Journal of Zoology* 77:978-988.

- Brown, R. F. 1988. Assessment of pinniped populations in Oregon. Processed Report 88-05, National Marine Fisheries Service, Northwest and Alaska Fisheries Center, Seattle, Washington.
- Calambokidis, J. 2010. John Calambokidis, senior marine mammal biologist and co-founder of Cascadia Research, Olympia, WA. September 15, 2001. Personal communication with Chris Hunt, Marine Scientist, Science Applications International Corporation, Bothell, WA, re: the rare occurrence of large whales (e.g., gray/humpback whales) occurring south of the Hood Canal Bridge since its construction.
- Calambokidis, J., and R.W. Baird. 1994. Status of marine mammals in the Strait of Georgia, Puget Sound, and the Juan de Fuca Strait, and potential human impacts. *Canadian Technical Report of Fisheries and Aquatic Sciences*. 1948: 282-300.
- Calambokidis, J. and S.J. Jeffries. 1991. Censuses and disturbance of harbor seals at Woodard Bay and recommendations for protection. Final report. Prepared for Washington Department of Natural Resources, Olympia, Washington by Cascadia Research Collective, Olympia, Washington and Washington Department of Wildlife.
- Calambokidis et al. 1985. Biology of Puget Sound marine mammals and marine birds: Population health and evidence of pollution effects. NOAA Tech. Memo. NOS OMA 18, National Technical Information Service, Springfield, Virginia 159 p.
- Calambokidis, J., S. Osmek, and J. L. Laake. 1997. Aerial surveys for marine mammals in Washington and British Columbia inside waters. Final Contract Report for Contract 52ABNF-6-00092, available from Cascadia Research Collective, Waterstreet Building 218 ½ West Fourth Avenue, Olympia, Washington 98501.
- CALTRANS. 2001. Marine Mammal Impact Assessment for the San Francisco-Oakland Bay Bridge Pile Installation Demonstration Project. PIDP EA 012081
- CALTRANS. 2007. Compendium of Pile Driving Sound Data. Report. Published Sept. 27, 2007.
- Campbell, G.S., R.C. Gisiner, D.A. Helweg, and L.L. Milette. 2002. Acoustic identification of female Steller sea lions (*Eumetopias jubatus*). *Journal of the Acoustical Society of America* 111 (6):2920- 2928.
- Carretta, J.V., K.A. Forney, M.S. Lowry, J. Barlow, J. Baker, B. Hanson, and M.M. Muto. 2007. U.S. Pacific Marine Mammal Stock Assessments: 2007. NOAA Technical Memorandum NMFS-SWFSC-414.
- Caretta, J.V., K.A. Forney, M.S. Lowry, J. Barlow, J. Baker, D. Johnston, B. Hanson, M.M. Muto, D. Lynch, and L. Carswell. 2008. U.S. Pacific Marine Mammal Stock Assessments: 2008. NOAA Technical Memorandum NMFS-SWFSC-434.
- Carlson, T.J., D.A. Woodruff, G.E. Johnson, N.P. Kohn, G.R. Plosky, M.A. Weiland, J.A. Southard, and S.L. Southard. 2005. Hydroacoustic measurements during pile driving at the Hood Canal Bridge, September through November 2004. Battelle Marine Sciences Laboratory Sequim, WA.
- Cavanaugh, W.J., and G.C. Tocci. 1998. Environmental noise: The invisible pollutant. Environmental Excellence in South Carolina (E2SC). USC Institute of Public Affairs, Los Angeles, CA. Vol. 1, No. 1.
- CERC (Coastal Engineering Research Center). 1984. *Shore Protection Manual*, Fourth ed., U.S. Army Corps of Engineers, Washington, D.C.

- Chapman, C.J. and A.D. Hawkins. 1969. The importance of fish behaviour in relation to capture by trawls. *FAO Fisheries Report* 62(3): 717-729.
- Cheremisinoff, N.P. 1996. Noise Control in industry: a practical guide. Noyes Publications. Westwood: NJ.
- Chivers, S. J., A. E. Dizon, P. J. Gearin, and K. M. Robertson. 2002. Small-scale population structure of eastern North Pacific harbour porpoises (*Phocoena phocoena*) indicated by molecular genetic analyses. *J. Cetacean Res. Manage.* 4(2):111-122.
- DoN. 2001a. Integrated Natural Resource Management Plan Naval Submarine Base Bangor, Silverdale, Washington.
- DoN. 2001b. Shock trial of the WINSTON S. CHURCHILL (DDG 81): final environmental impact statement.
- DoN 2010a. Marine mammal surveys at Naval Base Kitsap Bangor – sighting reports.
- Everitt, R.D., P.J. Gearin, J.S. Skidmore, and R.L. DeLong. 1981. Prey items of harbor seals and California sea lions in Puget Sound, Washington. *Murrelet* 62(3):83-86.
- Felleman, F.L., J.R. Heimlich-Boran, and R.W. Osborne. 1991. The feeding ecology of killer whales (*Orcinus orca*) in the Pacific Northwest. Pages 113-147 in Pryor, K. and K.S. Norris, eds. *Dolphin societies: Discoveries and puzzles*. Berkeley: University of California Press.
- Ferrero, R. C., and W. A. Walker. 1999. Age, growth, and reproductive patterns of Dall's porpoise (*Phocoenoides dalli*) in the central North Pacific Ocean. *Marine Mammal Science* 15:273-313.
- Finneran, J. J., D. A. Carder, C. E. Schlundt, and S. H. Ridgway, 2005. Temporary threshold shift in bottlenose dolphins (*Tursiops truncatus*) exposed to mid-frequency tones. *Journal of the Acoustical Society of America*, Vol 118, pp 2696–2705.
- Ford, J.K.B. 2002. Dialects. Pages 322-323 in Perrin, W.F., B. Würsig, and J.G.M. Theewissen, eds. *Encyclopedia of marine mammals*. San Diego, California: Academic Press.
- Ford, J. K. B., and G. M. Ellis. 1999. *Transients: Mammal-Hunting Killer Whales of British Columbia, Washington, and Southeastern Alaska*. University of British Columbia Press, Vancouver, BC. 96 pp.
- Ford, J. K. B., G. M. Ellis, and K. C. Balcomb. 1994. *Killer Whales: The Natural History and Genealogy of *Orcinus orca* in British Columbia and Washington State*. University of British Columbia Press, Vancouver, BC, and University of Washington Press, Seattle. 102 pp.
- Ford, J.K.B., G.M. Ellis, L.G. Barrett-Lennard, A.B. Morton, R.S. Palm, and K.C. Balcomb III. 1998. Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. *Canadian Journal of Zoology* 76:1456-1471.
- Ford, J.K.B., G.M. Ellis, and P.F. Olesiuk. 2005. Linking prey and population dynamics: Did food limitation cause recent declines of 'resident' killer whales (*Orcinus orca*) in British Columbia? Canadian Science Advisory Secretariat Research document 2005/042. Department of Fisheries and Oceans.

- Forney, K.A. 2007. Preliminary estimates of cetacean abundance along the U.S. west coast and within four National Marine Sanctuaries during 2005. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SWFSC-406. 27 p.
- Forney, K.A. and J. Barlow. 1998. Seasonal patterns in the abundance and distribution of California cetaceans, 1991-1992. *Marine Mammal Science* 14(3):460-489.
- Gilbert, J.R. and N. Guldager. 1998. Status of harbor and gray seal populations in northern New England. Woods Hole, Massachusetts: National Marine Fisheries Service.
- Govoni, J.J., L.R. Settle, and M.A. West. 2003. Trauma to juvenile pinfish and spot inflicted by submarine detonations. *Journal of Aquatic Animal Health* 15:111-119.
- Green, G.A., J.J. Brueggeman, R.A. Grotefendt, C.E. Bowlby, M.L. Bonnell, and K.C. Balcomb III. 1992. Cetacean distribution and abundance off Oregon and Washington, 1989-1990. Pages 1-1 to 1-100 in Brueggeman, J.J., ed. Oregon and Washington marine mammal and seabird surveys. OCS Study MMS 91-0093. Los Angeles, California: Minerals Management Service.
- Hammermeister, T., and W. Hafner. 2009. Naval Base Kitsap sediment quality investigation: data report. Prepared by Science Applications International Corporation, Bothell, WA. Prepared for BAE Systems Applied Technologies, Inc., Rockville, MD.
- Hanggi, E.B. and R.J. Schusterman. 1994. Underwater acoustic displays and individual variation in male harbour seals, *Phoca vitulina*. *Animal Behaviour* 48:1275-1283
- Harris. C.M. 1998. Handbook of acoustical measurements and noise control (3rd Edition). Huntington, NY: Acoustical Society of America.
- Harris, D.E., B. Lelli, and S. Gupta. 2003. Long-term observations of a harbor seal haul-out site in a protected cove in Casco Bay, Gulf of Maine. *Northeastern Naturalist* 10(2):141-148.
- Hastings, M. C. 1990. Effects of Underwater Sound on Fish. Document No. 46254-900206-01IM. Project No. 401775-1600, AT&T Bell Laboratories.
- Hastings, M. 2007. Calculation of SEL for Govoni et al. (2003, 2007) and Popper et al. (2007) Studies. Report for Amendment to Project 15218, J&S Working Group, 14 December 2007. 7 pp.
- Hastings, M.C., and A.N. Popper. 2005. Effects of Sound on Fish. Report prepared by Jones & Stokes for California Department of Transportation, Contract No. 43A0139, Task Order 1.
- Hawkins, A. 2005. Assessing the impact of pile driving upon fish. UC Davis: Road Ecology Center. Retrieved from: <http://www.escholarship.org/uc/item/28n858z1>
- Heath, C. B. 2002. California, Galapagos, and Japanese sea lions— *Zalophus californianus*, *Z. wollebaeki*, and *Z. japonicus*. Pages 180 to 186 in: Perrin, W. F., B. Würsig, and J. G. M. Thewissen, editors. 2002. Encyclopedia of Marine Mammals. Academic Press.
- Heimlich-Boran, J.R. 1988. Behavioral ecology of killer whales (*Orcinus orca*) in the Pacific Northwest. *Canadian Journal of Zoology* 66:565-578.
- Hildebrand, J. 2007. Sources of anthropogenic sound in the marine environment. Marine Mammal Commission.
<http://www.mmc.gov/sound/internationalwrkshp/pdf/hildebrand.pdf>

- Hoelzel, A. R., M. E. Dahlheim, and S. J. Stern. 1998. Low genetic variation among killer whales (*Orcinus orca*) in the Eastern North Pacific, and genetic differentiation between foraging specialists. *J. Heredity* 89:121-128.
- Hollingshead, K. 2008. Personal communication via email between Ken Hollingshead (Fishery Biologist in Resource Management at NMFS headquarters) and Andrea Balla-Holden (URS Corporation Fisheries and Marine Mammal Biologist) in March 2008 regarding the origin of the 120 dB re 1 μ Pa rms threshold and application to pinnipeds.
- Houck W.J. and T.A. Jefferson. 1999. Dall's porpoise - *Phocoenoides dalli* (True, 1885). In: *Handbook of Marine Mammals* (Ridgway SH, Harrison SR Eds.) Vol. 6: The second book of dolphins and porpoises. pp. 443-472
- Huber, H. R., S. J. Jeffries, R. F. Brown, R. L. DeLong, and G. VanBlaricom. 2001. Correcting aerial survey counts of harbor seals (*Phoca vitulina richardsi*) in Washington and Oregon. *Mar. Mammal Sci.* 17(2):276-293.
- Hubbs, C. 1960. The marine invertebrates of the outer coast. In: *The biogeography of Baja California and adjacent seas, part 2. Systematic Zoology* 9:134-147.
- Jefferson, T.A. 1988. *Phocoenoides dalli*. *Mammalian Species* 319:1-7.
- Jefferson, T.A. 1989. Status of Dall's porpoise, *Phocoenoides dalli*, in Canada. *Canadian Field-Naturalist* 104:112-116.
- Jefferson, T.A. 1990. Sexual dimorphism and development of external features in Dall's porpoise *Phocoenoides dalli*. *Fishery Bulletin* 88:119-132.
- Jefferson, T.A. 1991. Observations on the distribution and behaviour of Dall's porpoise (*Phocoenoides dalli*) in Monterey Bay, California. *Aquatic Mammals* 17(1):12-19.
- Jefferson, T.A. 2005. NMFS-SWFSC, Personal Communication., 14-18 March 2005.
- Jefferson, T.A., S. Leatherwood, and M.A. Webber. 1993. *FAO species identification guide. Marine mammals of the world.* Rome, Italy: Food and Agriculture Organization of the United Nations.
- Jeffries, S. J. 1985. Occurrence and distribution patterns of marine mammals in the Columbia River and adjacent coastal waters of northern Oregon and Washington. In: *Marine Mammals and Adjacent Waters, 1980-1982. Processed Report 85-04,* National Marine Fisheries Service, Northwest and Alaska Fisheries Center, Seattle, Washington.
- Jeffries, S. 2006. Steve Jeffries, Marine Mammal Specialist, Washington Department of Fish and Wildlife. December 14, 2006. Personal communication with Alison Agness, Science Applications International Corporation, re: occurrence of marine mammals in Hood Canal.
- Jeffries, S.J., P.J. Gearin, H.R. Huber, D.L. Saul, and D.A. Pruett. 2000. Atlas of seal and sea lion haul-out sites in Washington. Washington State Department of Fish and Wildlife, Wildlife Science Division, Olympia, WA. 150 pp.
http://wdfw.wa.gov/wlm/research/papers/seal_haulout/
- Jeffries, S., H. Huber, J. Calambokidis, and J. Laake. 2003. Trends and status of harbor seals in Washington State:1978-1999. *J. Wildl. Manage.* 67(1):208-219.
- Kastak, D. and R.J. Schusterman. 1998. Low-frequency amphibious hearing in pinnipeds: methods, measurements, noise, and ecology. *J. Acoust. Soc. Am.* 103(4):2216-2228.

- Kastak, D. and R.J. Schusterman. 2002. Changes in auditory sensitivity with depth in a free-diving California sea lion (*Zalophus californianus*). *Journal of the Acoustical Society of America* 112(1):329-333.
- Kastak, D., R.J. Schusterman, B.L. Southall and C.J. Reichmuth. 1999. Underwater temporary threshold shift induced by octave-band noise in three species of pinniped. *Journal of the Acoustic Society of America*. *Journal of the Acoustic Society of America*. 106(2):1142-1148.
- Kastelein, R. A., P. Bunskoek, M. Hagedoorn, W. W. L. Au, and D. de Haan. 2002. Audiogram of a harbor porpoise (*Phocoena phocoena*) measured with narrow-band frequency-modulated signals. *Journal of the Acoustical Society of America* 112(1):334-344.
- Kastelein, R. A. R. van Schie, W. C. Verboom, and D. de Haan. 2005. Underwater hearing sensitivity of a male and a female Steller sea lion (*Eumetopias jubatus*). *J. Acoust. Soc. Am.* 118 (3):1820-1829.
- Keple, A.R. 2002. Seasonal abundance and distribution of marine mammals in the southern Strait of Georgia, British Columbia. Master's thesis, University of British Columbia.
- Ketten, D.R. 1995. Estimates of blast injury and acoustic trauma zones for marine mammals from underwater explosions. Pp. 391-407. In: R.A. Kastelein, J.A. Thomas, and P.E. Nachtigall (eds.). *Sensory Systems of Aquatic Mammals*. Woerden, The Netherlands: De Spil Publishers.
- Ketten, D.R. 1998. Marine mammal auditory systems: A summary of audiometric and anatomical data and its implications for underwater acoustic impacts. NOAA Technical Memorandum NMFS-SWFSC-256:1-74.
- Ketten, D.R. 2000. Cetacean ears. Pp. 43-108. In: W.W.L. Au, A.N. Popper, and R.R. Fay (eds.). *Hearing by Whales and Dolphins*. New York: Springer-Verlag.
- Krahn, M.M., M.J. Ford, W.F. Perrin, P.R. Wade, R.P. Angliss, M.B. Hanson, B.L. Taylor, G.M. Ylitalo, M.E. Dahlheim, J.E. Stein, and R.S. Waples. 2002. Status review of Southern Resident killer whales (*Orcinus orca*) under the Endangered Species Act. U.S. Dept. Commerce., NOAA Tech. Memo. NMFS-NWFSC-54.
- Laake, J. L. National Marine Mammal Laboratory, AFSC, NMFS, 7600 Sand Point Way NE, Seattle, WA 98115.
- Le Boeuf, B.J. 2002. Status of pinnipeds on Santa Catalina Island. *Proceedings of the California Academy of Sciences* 53(2):11-21.
- Le Boeuf, B. J., and M. L. Bonnell. 1980. Pinnipeds of the California islands: abundance and distribution. Pages 475-493 in D. Power, ed. *The California islands*. Santa Barbara Mus. Nat. Hist. 787 pp.
- London, J.M. 2006. "Harbor Seals in Hood Canal: Predators and Prey." Ph.D. dissertation, University of Washington. Available from: <http://www.marinemammal.org>.
- Longmuir, C. and T. Lively. 2001. Bubble curtain systems for use during marine pile driving. Report by Fraser River Pile and Dredge Ltd., New Westminster, BC.
- Laughlin, J. 2006. Underwater sound levels associated with pile driving at the Cape Disappointment Boat Launch Facility, Wave Barrier Project. Prepared by Washington State Department of Transportation, Office of Air Quality and Noise, Seattle, WA.

- Loughlin, T. R. 1997. Using the phylogeographic method to identify Steller sea lion stocks. Pp. 329-341 In A. Dizon, S. J. Chivers, and W. Perrin (eds.), Molecular genetics of marine mammals, incorporating the proceedings of a workshop on the analysis of genetic data to address problems of stock identity as related to management of marine mammals. Soc. Mar. Mammal., Spec. Rep. No. 3.
- Loughlin, T.R. 2002. Steller's sea lion, *Eumetopias jubatus*. Pages 1181-1185 in Perrin, W.F., B. Würsig, and J.G.M. Thewissen, eds. Encyclopedia of marine mammals. San Diego, California: Academic Press.
- Loughlin, T.R., M.A. Perez, and R.L. Merrick. 1987. *Eumetopias jubatus*. Mammalian Species 283:1-7.
- Lowry, M.S., B.S. Stewart, C.B. Heath, P.K. Yochem, and J.M. Francis. 1991. Seasonal and annual variability in the diet of California sea lions *Zalophus californianus* at San Nicolas Island, California, 1981-86. Fishery Bulletin 89:331-336.
- Maniscalco, J.M., K. Wynne, K.W. Pitcher, M.B. Hanson, S.R. Melin, and S. Atkinson. 2004. The occurrence of California sea lions (*Zalophus californianus*) in Alaska. *Aquatic Mammals*. 30(3): 427-433.
- Mate, B.R. 1975. Annual migrations of the sea lions *Eumetopias jubatus* and *Zalophus californianus* along the Oregon coast. Rapports et Proces-Verbaux des Reunions Commission Internationale pour l'Exploration Scientifique de la Mer Mediterranee Monaco 169:455-461.
- Matkin, C. and E. Saulitis. 1997. Killer whale *Orcinus orca*. Restoration Notebook (Publication of the Exxon Valdez Oil Spill Trustee Council) November:1-12.
- Matkin, C., G. Ellis, E. Saulitis, L. Barrett-Lennard, and D. Matkin. 1999. Killer Whales of Southern Alaska. North Gulf Oceanic Society. 96 pp.
- Merrick, R. L., M. K. Chumbley, and G. V. Byrd. 1997. Diet diversity of Steller sea lions (*Eumetopias jubatus*) and their population decline in Alaska: a potential relationship. *Can J. Fish Aquat. Sci.* 54:1342-1348.
- Miller, G.W., R.E. Elliott, W.R. Koski, V.D. Moulton, and W.J. Richardson. 1999. Whales. In: Marine Mammal and Acoustical Monitoring of Western Geophysical's Open-Water Seismic Program in the Alaskan Beaufort Sea, 1998, LGL and Greeneridge, eds. LGL Report TA 2230-3. King City, Ont., Canada: LGL Ecological Research Associates, Inc., 109 pp.
- Morejohn, G.V. 1979. The natural history of Dall's porpoise in the North Pacific Ocean. Pages 45-83 in Behavior of Marine Animals, Vol. 3, Cetaceans. H.E. Winn and B.L. Olla (Eds). Plenum Press, New York.
- Morris, J.T., V.I. Osychny, and P.J. Luey. 2008. Naval Base Kitsap Bangor – Supplemental Current Measurement Survey: August 2007 field data report. Final. Prepared by Science Applications International Corporation, Newport, RI. Prepared for BAE Systems Applied Technologies, Inc., Rockville, MD.
- Morton, A. B. 1990. A quantitative comparison of the behaviour of resident and transient forms of the killer whale off the central British Columbia coast. Reports of the International Whaling Commission (Special Issue 12):245-248.
- Moulton, V. D., Richardson, W. J., Elliott, R. E., McDonald, T. L., Nations, C., & Williams, M. T. 2005. Effects of an offshore oil development on local abundance and distribution of ringed seals (*Phoca hispida*) of the Alaskan Beaufort Sea. *Marine Mammal Science*, 21, 217-242.

- Mulsow, J. and C. Reichmuth. 2008 in prep. Aerial Hearing Sensitivity in a Steller Sea Lion. Extended abstract presented at the Acoustic Communication by Animals, Second International Conference. Corvallis, Oregon. August 12 – 15, 2008. Citations were used with permission from the authors.
- Navy. 2005a. Environmental Assessment. Installation and Operation of Underwater Surveillance System (USS) at Sub-base Bangor, Silverdale, WA. January 2005.
- Nedwell, J. and D. Howell. A review of offshore windfarm related underwater noise sources. Subacoustech Report. October 2004.
- Newton, J.A., S.L. Albertson, K. Nakata, and C. Clishe. 1998. Washington State marine water quality in 1996 and 1997. Washington State Department of Ecology, Environmental Assessment Program, Publication No. 98-338.
<http://www.ecy.wa.gov/pubs/98338.pdf>
- Newton, J.A., S.L. Albertson, K. Van Voorhis, C. Maloy, and E. Siegel. 2002. Washington State marine water quality, 1998 through 2000. Washington State Department of Ecology Environmental Assessment Program, Publication No. 02-03-056. <http://www.ecy.wa.gov/pubs/0203056.pdf>
- National Marine Fisheries Service (NMFS). 1992. Final recovery plan for Steller sea lions *Eumetopias jubatus*. NMFS Office of Protected Resources, Silver Spring, MD. 92pp.
- NMFS. 1993. Final Rule: Designated Critical Habitat; Steller Sea Lion. 58 FR 45269.
- NMFS. 2005. Final Rule; Endangered Status for Southern Resident Killer Whales. 70 FR 69903.
- NMFS. 2005b. Endangered Fish and Wildlife; Notice of intent to prepare an environmental impact statement. 70 FR 1871.
- NMFS. 2007. Final Rule: Threatened Fish and Wildlife; Change in Listing Status of Steller Sea Lions Under the Endangered Species Act. 62 FR 24345.
- NMFS. 2008a. Recovery Plan for the Steller Sea Lion (*Eumetopias jubatus*). Revision. National Marine Fisheries Service, Silver Spring, MD. 325 pages.
- NMFS. 2008b. Taking of marine mammals incidental to specified activities; construction of the east span of the San Francisco-Oakland Bay Bridge. 73 FR 38180, July 3, 2008.
- NMFS. 2008c. Draft Environmental Assessment: Reducing the impact on at-risk salmon and steelhead by California sea lions in the area downstream of Bonneville Dam on the Columbia River, Oregon and Washington. NOAA National Marine Fisheries Service, Northwest Region, Seattle, Washington. pp. 127.
- NMFS. 2009. Taking of marine mammals incidental to specified activities; construction of the East Span of the San Francisco-Oakland Bay Bridge. 74 FR 41684.
- Norberg, B. 2007a. Personal email communication between Brent Norberg (National Marine Mammal Laboratory Biologist) and Andrea Balla-Holden (URS Corporation Fisheries and Marine Mammal Biologist) on Monday April 30, 2007.
- Norberg, B. 2007b. Personal email communication between Brent Norberg (National Marine Mammal Laboratory Biologist) and Andrea Balla-Holden (URS Corporation Fisheries and Marine Mammal Biologist) on Wednesday June 13, 2007.

- Norris, K.S., and J.H. Prescott. 1961. Observations on Pacific cetaceans of Californian and Mexican waters. University of California Publications in Zoology 63:291-402.
- O'Keefe, D.J. and G.A. Young. 1984. Handbook on the environmental effects of underwater explosions. Naval Surface Weapons Center, Dahlgren and Silver Spring, NSWC TR 83-240.
- Orr, A.J., A.S. Banks, S. Mellman, H.R. Huber, R.L. DeLong, and R.F. Brown. 2004. Examination of the foraging habits of Pacific harbor seal (*Phoca vitulina richardsi*) to describe their use of the Umpqua River, Oregon, and their predation on salmonids. Fishery Bulletin 102:108-117.
- Osborne, R., J. Calambokidis, and E.M. Dorsey. 1988. *A guide to marine mammals of Greater Puget Sound*. Anacortes, WA: Island Publishers.
- Osmek, S.D., J. Calambokidis, J. Laake, P. Gearin, R. Delong, J. Scordino, S. Jeffries, and R. Brown. 1996. Assessment of the status of harbor porpoise (*Phocoena phocoena*) in Oregon and Washington Waters. December 1996. NOAA Technical Memorandum NMFS-AFSC-76.
- Osmek, S.D., J. Calambokidis, and J.L. Laake. 1998. Abundance and distribution of porpoise and other marine mammals of the inside waters of Washington and British Columbia. In Proceedings of the Fourth Puget Sound Research Conference, Strickland, R., ed. *Puget Sound Water Quality Action Team, Olympia, WA*. 868-880 pp; March 12-13, 1998, Seattle, WA.
- Payne, P.M. and L.A. Selzer. 1989. The distribution, abundance and selected prey of the harbor seal, *Phoca vitulina concolor*, in southern New England. Marine Mammal Science 5(2):173-192.
- Pearson, W. H., J.R. Skalski, and C.I. Malme. 1992. Effects of sounds from a geophysical survey device on behavior of captive rockfish (*Sebastes* spp.). *Canadian Journal of Fisheries and Aquatic Sciences* 49: 1343-1356.
- Phillips, C., B. Dolan, and W. Hafner. 2009. Naval Base Kitsap at Bangor water quality 2005 and 2006 field survey report. Prepared by Science Applications International Corporation, Bothell, WA. Prepared for BAE Systems Applied Technologies, Inc., Rockville, MD.
- Pitcher, K. W., and D. G. Calkins. 1981. Reproductive biology of Steller sea lions in the Gulf of Alaska. *J. Mamm.* 62:599-605.
- Pitcher, K. W., P. F. Olesiuk, R. F. Brown, M. S. Lowry, S. J. Jeffries, J. L. Sease, W. L. Perryman, C. E. Stinchcomb, and L. F. Lowry. 2007. Status and trends in a bundance and distribution of the eastern Steller sea lion (*Eumetopias jubatus*) population. *Fish. Bull.* 107(1):102-115.
- Popper, A.N. and M. Hastings. 2009. The effects of human-generated sound on fish. *Integrative Zoology* 4: 43-52.
- Popper, A.N., T.J. Carlson, B.L. Southall, and R.L. Gentry. 2006. Interim Criteria for Injury of Fish Exposed to Pile Driving Operations: A White Paper.
- Popper, A. N., M.B. Halvorsen, A. Kane, D.L. Miller, M.E. Smith, J. Song, P. Stein, and L.E. Wysocki. 2007. The effects of high-intensity, low-frequency active sonar on rainbow trout. *Journal of the Acoustic Society of America* 122: 623-635.
- Prescott, R. 1982. Harbor seals: Mysterious lords of the winter beach. *Cape Cod Life* 3(4):24-29.

- Read, A.J. 1990. Reproductive seasonality in harbour porpoises, *Phocoena phocoena*, from the Bay of Fundy. *Canadian Journal of Zoology* 68:284-288.
- Read, A.J., 1999. Harbour porpoise *Phocoena phocoena* (Linnaeus, 1758). Pages 323-355 in Ridgway, S.H. and R. Harrison, eds. *Handbook of marine mammals*. Volume 6: The second book of dolphins and the porpoises. San Diego, California: Academic Press.
- Read, A.J. and A.A. Hohn. 1995. Life in the fast lane: The life history of harbor porpoises from the Gulf of Maine. *Marine Mammal Science* 11(4):423-440.
- Reeves, R.R., B.S. Stewart, and S. Leatherwood. 1992. *The Sierra Club handbook of seals and sirenians*. San Francisco, California: Sierra Club Books.
- Reeves, R.R., P.A. Folkens, and National Audubon Society. 2002. *Guide to marine mammals of the world*. New York: Alfred A. Knopf.
- Reeves RR, Dalebout ML, Jefferson TA, Karczmarski L, Laidre K, O'Corry-Crowe G, Rojas-Bracho L, Secchi ER, Slooten E, Smith BD, Wang JY, Zhou K (2008) IUCN 2009. IUCN Red List of Threatened Species. Version 2009.2. <www.iucnredlist.org>.
- Reyff, J. 2003. Memo to Caltrans District 4 regarding SFOBB East Span Construction Pier 16E. Dated July 24, 2003.
- Richardson, W.J., G.R. Greene, Jr., C.I. Malme, and D.H. Thomson. 1995. *Marine mammals and noise*. San Diego, CA: Academic Press. 576 pp.
- Richardson, W.J. 1995. Marine mammal hearing. Pages 205-240 in Richardson, W.J., C.R. Greene, Jr., C.I. Malme, and D.H. Thomson, eds. *Marine mammals and noise*. San Diego, California: Academic Press.
- Ridgway, S. H., D. A. Carder, R. R. Smith, T. Kamolnick, C. E. Schlundt, and W. R. Elsberry, 1997. Behavioral responses and temporary shift in masked hearing threshold of bottlenose dolphins, *Tursiops truncatus*, to 1-second tones of 141 to 201 dB re 1 μ Pa. Technical Report 1751, Revision 1. San Diego, California: Naval Sea Systems Command.
- Riedman, M. 1990. *The pinnipeds: Seals, sea lions, and walruses*. Berkeley, California: University of California Press.
- Riedman, M.L. and J.A. Estes. 1990. *The sea otter (Enhydra lutris): Behavior, ecology, and natural history*. U.S. Fish and Wildlife Service Biological Report 90(14). Washington, D.C.: U.S. Fish and Wildlife Service.
- Robson, B.W., ed. 2002. *Fur seal investigations, 2000-2001*. NOAA Technical Memorandum NMFS-AFSC-1 34:1-80.
- Roffe, T. and B. Mate. 1984. Abundance and feeding habits of pinnipeds in the Rogue River, OR. *Journal of Wildlife Management*. Volume 48, pages 1,262 to 1,277.
- Saulitis, E. L. 1993. The behavior and vocalizations of the "AT" group of killer whales (*Orcinus orca*) in Prince William Sound, Alaska. M.S. Thesis, University of Alaska Fairbanks, Fairbanks, AK, 193 pp.
- Saulitis, E., C.O. Matkin, L.G. Barrett-Lennard, K. Heise, and G.M. Ellis. 2000. Foraging strategies of sympatric killer whale (*Orcinus orca*) populations in Prince William Sound, Alaska. *Marine Mammal Science*. 16: 94-109.
- Schneider, D.C. and P.M. Payne, 1983. Factors affecting haul-out of harbor seals at a site in southeastern Massachusetts. *Journal of Mammalogy* 64(3):518-520.

- Scholik, A. R., and H.Y. Yan. 2001. Effects of underwater noise on auditory sensitivity of a cyprinid fish. *Hearing Research* 152:17-24.
- Scholik, A. R., and H.Y. Yan. 2002. The effects of noise on the auditory sensitivity of the bluegill sunfish, *Lepomis macrochirus*. *Comp. Biochemical Physiology A* 133:43-52.
- Schusterman, R.J. 1974. Auditory sensitivity of a California sea lion to airborne sound. *Journal of the Acoustical Society of America*, 56:1248-1251.
- Schusterman, R.J. 1977. Temporal patterning in sea lion barking (*Zalophus californianus*). *Behavioral Biology*, 20:404-408.
- Schusterman, R.J. and R.F. Balliet. 1969. Underwater barking by male sea lions (*Zalophus californianus*). *Nature* 222(5199):1179-1181.
- Schusterman, R.J., R. Gentry, and J. Schmook. 1966. Underwater vocalization by sea lions: Social and mirror stimuli. *Science* 154(3748):540-542.
- Schusterman, R.J., Gentry, R., and Schmook, J. 1967. Underwater sound production by captive California sea lions. *Zoologica*, 52:21-24.
- Schusterman, R.J., R.F. Balliet, and S. St. John. 1970. Vocal displays under water by the gray seal, the harbor seal, and the stellar [sic] sea lion. *Psychonomic Science* 18(5):303-305.
- Schusterman, R.J., Balliet, R.F., and Nixon, J. 1972. Underwater audiogram of the California sea lion by the conditioned vocalization technique. *Journal of the Experimental Analysis of Behavior*, 17:339-350.
- Scordino, J. 2006. Steller sea lions (*Eumetopias jubatus*) of Oregon and Northern California: Seasonal haulout abundance patterns, movements of marked juveniles, and effects of hot-iron branding on apparent survival of pups at Rogue Reef. Master of Science thesis, Oregon State University, Corvallis, OR. 92 pages.
- Skalski, J.R., W.H. Pearson, and C.I. Malme. 1992. Effects of sounds from a geophysical survey device on catch-per-unit-effort in a hook-and-line fishery for rockfish (*Sebastes* spp.). *Canadian Journal of Fisheries and Aquatic Sciences* 49: 1357-1365.
- Slater, M.C. 2009. Naval Base Kitsap, Bangor baseline underwater noise survey report. Prepared by Science Applications International Corporation, Bremerton, WA. Prepared for BAE Systems Applied Technologies, Inc., Rockville, MD.
- Southall, B.L., Bowles, A.E., Ellison, W.T., Finneran, J.J., Gentry, R.L., Greene, C.R. Jr., Kastak, D., Ketten, D.K., Miller, J.H., Nachtigall, P.E., Richardson, W.J., Thomas, J.A. and Tyack, P.L. 2007. Marine mammal noise exposure criteria: initial scientific recommendations. *Special Issue of Aquatic Mammals*. 33(4): 412-522.
- Suryan, R.M. and J.T. Harvey. 1998. Tracking harbor seals (*Phoca vitulina richardsi*) to determine dive behavior, foraging activity, and haul-out site use. *Marine Mammal Science*. 14(2): 361-372.
- Szymanski, M.D., D.E. Bain, K. Kiehl, S. Pennington, S. Wong, and K.R. Henry. 1999. Killer whale (*Orcinus orca*) hearing: auditory brainstem response and behavioral audiograms. *The Journal of the Acoustical Society of America*. 106(2):1134-1141.
- Tannenbaum, B.R., M. Bhuthimethee, L. Delwiche, G. Vadera, and J.M. Wallin. 2009a. Naval Base Kitsap at Bangor 2008 Marine Mammal Survey Report. Prepared by Science Applications International Corporation, Bothell, WA. Prepared for BAE Systems Applied Technologies, Inc., Rockville, MD.

- Tannenbaum, B.R. M. Bhuthimethee, L. Delwiche, G. Veder, and J.M. Wallin. 2009b. Naval Base Kitsap at Bangor 2008 Marine Bird Survey Report. Prepared by Science Applications International Corporation, Bothell, WA. Prepared for BAE Systems Applied Technologies, Inc., Rockville, MD.
- Temte, J. L. 1986. Photoperiod and the timing of pupping in the Pacific harbor seal (*Phoca vitulina richardsi*) with notes on reproduction in northern fur seals and Dall porpoises. Thesis, Oregon State University, Corvallis, USA.
- Terhune, J. M., and Ronald, K. 1975. Underwater hearing sensitivity of two ringed seals (*Pusa hispida*) Can. J. Zool. 53(3), 227–231.
- Terhune, J. and S. Turnbull. 1995. Variation in the psychometric functions and hearing thresholds of a harbour seal. Pages 81-93. IN: R.A. Kastelein, J.A. Thomas, and P.E. Nachtigall, eds. Sensory Systems of Aquatic Mammals. De Spil Publishers, Woerden, Netherlands.
- Thomson, D. H. and W. J. Richardson. 1995. Marine mammal sounds. Pages 159-204 in Richardson, W. J., C. R. Greene, Jr., C. I. Malme, and D. H. Thomson, eds. Marine mammals and noise. San Diego: Academic Press.
- Thorson, P. and J.A. Reyff. 2004. Marine mammal and acoustic monitoring for the eastbound structure. San Francisco-Oakland Bay Bridge East Span Seismic Safety Project. Report submitted for Incidental Harassment Authorization issued November 14, 2003 to Caltrans.
- Tollit, D. J., Greenstreet, S. P. R. & Thompson, P. M. 1997. Prey selection by harbour seals (*Phoca vitulina*) in relation to variations in prey abundance. *Canadian Journal of Zoology*, 75, 1508–1518.
- Urick, Robert J. 1983. *Principles of underwater sound*. 3rd ed. New York: McGraw-Hill.
- URS Consultants, Inc. 1994. Final remedial investigation report for the Comprehensive Long-Term Environmental Action Navy (CLEAN) Program, Northwest Area. Remedial investigation for Operable Unit 7, CTO-0058, SUBASE Bangor, Bremerton, WA. Prepared by URS Consultants, Inc., Seattle, WA. Prepared for Engineering Field Activity, Northwest, Western Division, Naval Facilities Engineering Command, Silverdale, WA. June 13, 1994.
- Van Parijs, S.M., P.J. Corkeron, J. Harvey, S.A. Hayes, D.K. Mellinger, P.A. Rouget, P.M. Thompson, M. Wahlberg, and K.M. Kovacs. 2003. Patterns in the vocalizations of male harbor seals. *Journal of the Acoustical Society of America* 113(6):3403-3410.
- Veirs, V. 2004. Source levels of free-ranging killer whale (*Orcinus orca*) social vocalizations. *Journal of the Acoustical Society of America* 116(No. 4, Pt. 2):2615.
- Verboom, W. C. and R.A. Kastelein. 1995. Acoustic signals by harbour porpoises (*Phocoena phocoena*). In Harbour Porpoises – Laboratory Studies to Reduce Bycatch (ed. P. E. Nachtigall, J. Lien, W. W. L. Au and A. J. Read), pp. 1-39. Woerden, The Netherlands: De Spil Publishers.
- Viada, S.T., R.M. Hammer, R. Racca, D. Hannay, M.J. Thompson, B.B. Balcom, and N.W. Phillips. 2008. Review of potential impacts to sea turtles from underwater explosive removal of offshore structures. *Environmental Impact Assessment*. 28: 267-285.

- Walker, W.A., M.B. Hanson, R.W. Baird and T.J. Guenther. 1998. Food habits of the harbor porpoise, *Phocoena phocoena*, and Dall's porpoise, *Phocoenoides dalli*, in the inland waters of British Columbia and Washington. Pages 63-75 in Marine Mammal Protection Act and Endangered Species Act Implementation Program 1997. AFSC Processed Report 98-10.
- Walters, A. 2009. Allison Walters, Naval Base Kitsap Environmental, Bangor, WA. January 23, 2009. Personal communication, email, with Bernice Tannenbaum, Science Applications International Corporation, Bothell, WA. re: occurrence of Steller sea lions, California sea lions, and harbor seals at Naval Base Kitsap Bangor.
- Ward, W.D. 1997. Effects of high intensity sound. In M.J. Crocker (Ed.) *Encyclopedia of acoustics, Volume III*. (pp 1497-1507). New York: John Wiley & Sons.
- Warner, M.J. 2007. Historical comparison of average dissolved oxygen in Hood Canal. Hood Canal Dissolved Oxygen Program. February 2007.
<http://www.hoodcanal.washington.edu/observations/historicalcomparison.jsp>
- Warner, M.J., M. Kawase, and J.A. Newton. 2001. Recent studies of the overturning circulation in Hood Canal. In Proceedings of the 2001 Puget Sound Research Conference, Puget Sound Action Team, Olympia, WA. 9 pp.
<http://www.hoodcanal.washington.edu/documents/document.jsp?id=1561>
- WDOE. 1991. Net shore-drift in Washington State Volume 4: Hood Canal Region. Ecology Report 00-06-03. Shorelands and Environmental Assistance Program. Washington Department of Ecology, Olympia, WA.
<http://www.ecy.wa.gov/pubs/93520.pdf>
- Wiles, G. J. 2004. Washington State status report for the killer whale. Washington Department Fish and Wildlife, Olympia. 106 pp.
http://wdfw.wa.gov/science/articles/orca/final_orca_status.pdf
- Willis, P.M., B.J. Crespi, L.M. Dill, R.W. Baird, and M.B. Hanson. 2004. Natural hybridization between Dall's porpoises (*Phocoenoides dalli*) and harbour porpoises (*Phocoena phocoena*). *Canadian Journal of Zoology* 82:828-834.
- Wilson, S.C. 1978. Social organization and behavior of harbor seals, *Phoca vitulina concolor*, in Maine. Final report to the U.S. Marine Mammal Commission. Washington, D.C.: Smithsonian Institution Press.
- Wilson, O.B.J., S.N. Wolf, and F. Ingenito. 1985. Measurements of acoustic ambient noise in shallow water due to breaking surf. *The Journal of the Acoustical Society of America*. 78(1): 190-195.
- Wolski, L.F., R.C. Anderson, A.E. Bowles, and P.K. Yochem. 2003. Measuring hearing in the harbor seal (*Phoca vitulina*): Comparison of behavioral and auditory brainstem response techniques. *Journal of the Acoustical Society of America* 113(1):629-637.
- Wright, Bryan. 2008. Personal communication via email between Bryan Wright (Biometrician for the Oregon Department of Fish and Game) and Andrea Balla- Holden (URS Corporation Fisheries and Marine Mammal Biologist) in March 2008 regarding California and Steller sea lion presence and numbers counted at Willamette Falls on the Willamette River.
- WSDOT, 2005. Friday Harbor Ferry Terminal Restoration Project: Underwater sound levels associated with restoration of the Friday harbor ferry terminal. May 2005.

- WSDOT. 2005b. Hydroacoustic Measurements during Pile Driving at the Hood Canal Bridge, September through November 2004.
- WSDOT. 2006. Underwater sound levels associated with pile driving at the Cape Disappointment Boat Launch Facility, Wave Barrier Project. March 2006.
- WSDOT. 2007. Underwater sound levels associated with driving steel and concrete piles near the Mukilteo Ferry Terminal. March 2007.
- WSDOT, 2008. Eagle Harbor Hydroacoustic Pressure Monitoring Technical Memorandum. May 2008.
- WSDOT. 2008b. Advanced Training Manual, Biological Assessment Preparations for Transportation Projects. April 2008.
- WSDOT. 2010. Keystone Ferry Terminal – vibratory pile monitoring technical memorandum. May 2010.
- WSDOT, 2010b. Vashon Ferry Terminal – vibratory pile monitoring technical memorandum. May 2010.
- WSDOT, 2010c. Airborne Noise Measurements (A-weighted and unweighted) during Vibratory Pile Installation – Technical Memorandum. June 2010
- Yelverton, J.T., D.R. Richmond, E.R. Fletcher, and R.K. Jones. 1973. Safe distances from underwater explosions for mammals and birds. Lovelace Foundation, Albuquerque, DNA 3114T.
<http://stinet.dtic.mil/cgi-bin/GetTRDoc?AD=AD766952&Location=U2&doc=GetTRDoc.pdf>.
- Yurk, H., L. Barrett-Lennard, J.K.B. Ford, and C.O. Matkin. 2002. Cultural transmission within maternal lineages: Vocal clans in resident killer whales in southern Alaska. *Animal Behaviour* 63:1103- 1119.