


# Diabetic Eye Disease Rates Increase Among U.S. Population

*1.2 million Hispanics at risk of losing their sight as a result of complications from diabetes*



Diabetes affects nearly 26 million people in the United States. In addition, another 79 million people are estimated to have pre-diabetes, a condition that puts people at increased risk for diabetes. Many Hispanics/Latinos are included in these statistics. All people with diabetes, both type 1 and type 2, are at risk for diabetic eye disease, a leading cause of vision loss and blindness.

Diabetic eye disease refers to a group of eye problems that people with diabetes may face as a complication of the disease and includes cataract, diabetic retinopathy, and glaucoma. Diabetic retinopathy, the most common diabetic eye disease, is the leading cause of blindness in adults 20–74 years of age. According to the National Eye Institute (NEI), 1.2 million Hispanics/Latinos have diabetic retinopathy and this number is projected to increase to approximately 3 million people by 2030.



This graph shows that 1.2 million Hispanics have diabetic retinopathy and this number is projected to increase to approximately 3 million by 2030.

Source: *Vision Problems in the U.S., 2012*

“The longer a person has diabetes the greater is his or her risk of developing diabetic eye disease,” said Paul A. Sieving, M.D., Ph.D., director of the NEI. “If you have diabetes, be sure to have a comprehensive dilated eye exam at least once a year. Diabetic eye disease often has no early warning signs, but can be detected early and treated before vision loss occurs. Don’t wait until you notice an eye problem to have an exam because vision that is lost often cannot be restored.”

While all people with diabetes can develop diabetic eye disease, Hispanics/Latinos, African Americans, and American Indians/Alaska Natives with diabetes are at higher risk of losing vision or going blind from it. All people with diabetes

should have a dilated eye exam at least once a year to detect vision problems early. “In fact, with early detection, timely laser surgery, and appropriate follow-up care, people with advanced diabetic retinopathy can reduce their risk of blindness by 90 percent,” adds Suber Huang, M.D., M.B.A., chair of the Diabetic Eye Disease Subcommittee for the NEI’s National Eye Health Education Program.

Research has shown that when people with diabetes maintain good control of blood sugar, blood pressure, and cholesterol they can slow the development and progression of diabetic eye disease. In addition to having annual comprehensive dilated eye exams, people with diabetes should do the following to keep their health on **TRACK**:

- Take your medications.
- Reach and maintain a healthy weight.
- Add physical activity to your daily routine.
- Control your blood sugar, blood pressure, and cholesterol.
- Kick the smoking habit.

NEI supports more than \$40 million in diabetes research and outreach, including the Diabetic Retinopathy Clinical Research Network (DRCR.net), a collaboration of more than 300 physicians at more than 100 clinical sites across the United States.

For more information on diabetic eye disease and tips on finding an eye care professional or financial assistance for eye care, visit [www.nei.nih.gov/diabetes](http://www.nei.nih.gov/diabetes) or call NEI at 301-496-5248.

*The National Eye Institute (NEI), part of the National Institutes of Health, leads the federal government’s research on the visual system and eye diseases. NEI supports basic and clinical science programs that result in the development of sight-saving treatments. For more information, visit [www.nei.nih.gov](http://www.nei.nih.gov).*

*About the National Institutes of Health (NIH): NIH, the Nation’s medical research agency, includes 27 Institutes and Centers and is a component of the U.S. Department of Health and Human Services. NIH is the primary federal agency conducting and supporting basic, clinical, and translational medical research, and is investigating the causes, treatments, and cures for both common and rare diseases. For more information about NIH and its programs, visit [www.nih.gov](http://www.nih.gov).*

- People with diabetes should have a comprehensive dilated eye exam at least once a year to detect diabetic eye disease in its early stages.
- Early detection and timely treatment can reduce the risk of vision loss.

###