

Table I-23
AGE-ADJUSTED SEER INCIDENCE RATES AND TRENDS FOR THE TOP 15 CANCER SITES^a BY RACE/ETHNICITY

Both Sexes

	All Races		White		Black	
	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005
All Sites	467.4	-0.8*	475.3	-0.7*	500.6	-1.1*
Prostate ^f	72.0	-0.6	70.4	-1.6*	102.9	-1.9*
Breast	68.5	-1.5*	69.8	-0.5	76.0	-1.7*
Lung and Bronchus	63.9	-1.4*	65.1	-1.3*	67.4	-0.8*
Colon and Rectum	50.6	-2.0*	50.1	-2.2*	61.2	-0.9
Urinary Bladder	21.2	0.0	23.0	0.1	15.2	-0.7
Non-Hodgkin Lymphoma	19.5	0.2	22.6	2.2*	14.9	0.4
Melanoma of the Skin	19.4	1.6*	20.3	0.3	14.8	1.9*
Kidney and Renal Pelvis	13.2	2.2*	13.7	2.3*	12.8	0.6
Corpus and Uterus, NOS ^f	12.6	-1.0*	13.1	-1.2*	12.4	-3.4*
Leukemia	12.3	-0.8*	12.9	-0.8*	11.7	1.5
Pancreas	11.5	0.3	11.4	0.5*	11.6	-0.5
Oral Cavity and Pharynx	10.4	-1.6*	10.6	-1.3*	10.8	-3.0*
Thyroid	9.1	5.8*	9.5	6.1*	10.1	-0.4
Stomach	8.0	-1.7*	7.6	-1.6*	8.0	3.9*
Ovary ^{fh}	7.3	-1.4*	7.1	-0.3	6.0	-4.9*
	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005
All Sites	313.5	-0.7*	312.7	-0.4	357.6	-0.7*
Breast	49.0	-0.4	46.0	0.8	60.0	-0.7
Colon and Rectum	40.9	-1.6*	43.5	-0.4	48.4	-0.8*
Prostate ^f	40.5	-0.3	40.5	0.0	39.2	-1.2*
Lung and Bronchus	39.1	-0.7*	32.3	-2.6	33.3	-2.0*
Liver & IBD ^g	14.3	0.0	15.8	0.9	16.7	-0.3
Stomach	13.9	-2.9*	11.6	-1.3	13.0	2.8*
Non-Hodgkin Lymphoma	13.2	0.1	11.0	1.5	12.1	-1.8*
Urinary Bladder	9.3	0.4	9.9	4.6	11.7	0.2
Pancreas	9.0	0.0	9.3	-0.4	10.6	-0.1
Corpus and Uterus, NOS ^f	9.0	0.8	8.2	-1.2	10.0	1.2
Thyroid	8.9	3.4*	7.5	-0.3	9.6	-0.8
Oral Cavity and Pharynx	7.8	-2.1*	7.1	-	9.6	0.9
Leukemia	7.3	-1.3	7.1	-2.8	8.0	4.5*
Kidney and Renal Pelvis	6.6	1.8*	6.2	-3.6	6.8	-3.8*
Ovary ^{fh}	5.3	-0.1	5.9	2.2	6.4	-0.5

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Top 15 cancer sites selected based on 2001-2005 age-adjusted rates for the race/ethnic group.

^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval. Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^f Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky.

^g The rates for sex-specific cancer sites are calculated using the population for both sexes combined.

^h IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

* The APC is significantly different from zero (p<.05).

Table I-24
AGE-ADJUSTED SEER INCIDENCE RATES AND TRENDS FOR THE TOP 15 CANCER SITES^a BY RACE/ETHNICITY

			Males					
All Races			White			Black		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2001-2005	1996-2005		2001-2005	1996-2005		2001-2005	1996-2005
All Sites	549.3	-1.0*	All Sites	551.4	-0.9*	All Sites	651.5	-1.9*
Prostate	163.0	-1.1	Prostate	156.7	-1.1	Prostate	248.5	-2.2*
Lung and Bronchus	79.4	-2.2*	Lung and Bronchus	79.3	-2.1*	Lung and Bronchus	107.6	-3.2*
Colon and Rectum	59.2	-2.4*	Colon and Rectum	58.9	-2.6*	Colon and Rectum	71.2	-0.8
Urinary Bladder	37.3	-0.1	Urinary Bladder	40.6	-0.1	Kidney and Renal Pelvis	21.3	2.8*
Melanoma of the Skin	24.6	1.7*	Melanoma of the Skin	28.5	2.2*	Urinary Bladder	20.4	0.5
Non-Hodgkin Lymphoma	23.5	-0.1	Non-Hodgkin Lymphoma	24.3	0.0	Non-Hodgkin Lymphoma	18.4	-0.1
Kidney and Renal Pelvis	18.3	1.9*	Kidney and Renal Pelvis	18.8	2.0*	Stomach	17.4	-3.6*
Leukemia	16.0	-1.1*	Leukemia	16.7	-1.0*	Oral Cavity and Pharynx	17.2	-3.9*
Oral Cavity and Pharynx	15.5	-1.7*	Oral Cavity and Pharynx	15.7	-1.3*	Pancreas	16.2	-1.3
Pancreas	13.0	0.2	Pancreas	13.0	0.3	Myeloma	14.4	0.2
Stomach	11.3	-2.2*	Stomach	10.0	-2.3*	Liver & IBD ^f	13.2	4.3*
Liver & IBD ^f	9.9	2.2*	Brain and ONS ^f	8.4	-0.6	Leukemia	13.0	-1.2
Esophagus	7.8	-0.1	Liver & IBD ^f	8.2	2.2*	Larynx	10.9	-3.4*
Brain and ONS ^f	7.6	-0.8*	Esophagus	8.0	0.9	Esophagus	9.9	-5.3*
Myeloma	7.0	-0.6	Myeloma	6.6	-0.7	Brain and ONS ^f	4.7	-0.4
Asian/Pacific Islander			American Indian/Alaska Native ^d			Hispanic ^e		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2001-2005	1996-2005		2001-2005	1996-2005		2001-2005	1996-2005
All Sites	354.0	-0.9*	All Sites	336.6	-1.3	All Sites	419.4	-1.1*
Prostate	93.8	-0.2	Prostate	73.3	-2.8	Prostate	138.0	-1.2
Lung and Bronchus	53.9	-1.2*	Lung and Bronchus	54.3	-2.2	Colon and Rectum	47.3	-1.8*
Colon and Rectum	48.0	-2.2*	Colon and Rectum	46.0	-0.7	Lung and Bronchus	44.2	-2.3*
Liver & IBD ^f	21.7	-0.2	Kidney and Renal Pelvis	19.5	-0.7	Urinary Bladder	20.2	0.1
Stomach	18.6	-2.7*	Stomach	16.8	-	Non-Hodgkin Lymphoma	19.2	-0.6
Urinary Bladder	16.3	0.9	Liver & IBD ^f	14.4	-	Kidney and Renal Pelvis	17.4	3.2*
Non-Hodgkin Lymphoma	16.0	-0.1	Non-Hodgkin Lymphoma	13.3	-	Stomach	15.5	-2.9*
Oral Cavity and Pharynx	10.5	-2.8*	Urinary Bladder	12.4	-	Liver & IBD ^f	15.0	1.4
Pancreas	10.1	0.2	Pancreas	10.9	-	Leukemia	11.7	-0.9
Leukemia	9.1	-1.8	Oral Cavity and Pharynx	9.7	-	Pancreas	10.9	-0.2
Kidney and Renal Pelvis	9.1	1.1	Leukemia	7.9	-	Oral Cavity and Pharynx	9.4	-2.1*
Thyroid	4.1	1.7	Esophagus	7.1	-	Myeloma	6.8	-0.6
Esophagus	4.0	-3.2*	Myeloma	4.6	-	Brain and ONS ^f	6.2	2.4
Brain and ONS ^f	4.0	-1.2	Testis	4.2	-	Esophagus	5.5	-1.0
Myeloma	3.8	-2.4*	Brain and ONS ^f	4.0	-	Larynx	5.1	-3.3*

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Top 15 cancer sites selected based on 2001-2005 age-adjusted rates for the race/ethnic group.

^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval. Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero (p<.05).

Table I-25
AGE-ADJUSTED SEER INCIDENCE RATES AND TRENDS FOR THE TOP 15 CANCER SITES^a BY RACE/ETHNICITY

	Females					
	All Races		White		Black	
	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005
All Sites	411.0	-0.7*	423.6	-0.6*	398.9	-0.4*
Breast	126.1	-1.4*	130.6	-1.4*	117.5	-0.7*
Lung and Bronchus	52.6	-0.5*	54.9	-0.6*	54.6	0.4
Colon and Rectum	43.8	-1.7*	43.2	-1.9*	54.5	-0.9*
Corpus and Uterus, NOS	23.4	-0.7*	24.3	-0.9*	20.3	1.6*
Non-Hodgkin Lymphoma	16.3	0.7	18.5	2.3*	14.3	-0.2
Melanoma of the Skin	15.6	1.6*	17.1	0.8*	12.1	1.0
Thyroid	13.4	6.1*	14.1	6.4*	10.8	-4.8*
Ovary ^g	13.3	-1.2*	14.1	-1.4*	10.1	0.5
Pancreas	10.3	0.4	10.0	0.6	10.1	0.4
Leukemia	9.5	-0.8	10.0	-0.2	9.8	-1.2
Urinary Bladder	9.4	-0.3	9.9	-0.9	8.9	-3.1*
Kidney and Renal Pelvis	9.2	2.3*	9.5	2.5*	8.0	0.1
Cervix Uteri	8.4	-3.6*	8.2	-3.0*	8.0	6.5*
Oral Cavity and Pharynx	6.1	-1.6*	6.1	-1.8*	7.7	0.4
Stomach	5.5	-1.2*	6.0	0.0	5.9	-1.1
	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005
All Sites	287.8	-0.3	296.4	0.4	317.8	-0.4
Breast	89.6	-0.5	75.0	0.1	90.1	-0.6
Colon and Rectum	35.4	-0.9	41.2	0.2	32.8	-0.8*
Lung and Bronchus	28.0	0.4	39.7	5.1*	25.4	-1.7*
Corpus and Uterus, NOS	16.5	0.8	15.3	-1.1	17.8	1.2
Thyroid	13.3	3.9*	12.7	-	14.6	0.2
Non-Hodgkin Lymphoma	11.1	0.7	11.3	-3.5	13.2	-3.5*
Stomach	10.5	-2.9*	9.3	-	12.5	5.1*
Ovary ^g	9.8	-0.2	8.8	0.9	11.7	-0.2
Liver & IBD ^f	8.3	0.7	8.2	-	10.3	-0.2
Pancreas	8.2	-0.3	7.7	-	9.6	1.8*
Cervix Uteri	8.0	-5.7*	6.9	-	9.5	-0.7
Leukemia	5.9	-0.6	6.9	-3.6	8.0	-0.8
Oral Cavity and Pharynx	5.4	-0.9	6.3	-	5.8	0.1
Kidney and Renal Pelvis	4.6	3.2*	5.1	-	5.5	0.0
Urinary Bladder	4.1	-0.1	4.9	-	4.9	-0.1

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Top 15 cancer sites selected based on 2001-2005 age-adjusted rates for the race/ethnic group.

^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval. Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

^g Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

* The APC is significantly different from zero (p<.05).

Table I-26
AGE-ADJUSTED U.S. DEATH RATES AND TRENDS FOR THE TOP 15 CANCER SITES^a BY RACE/ETHNICITY

Both Sexes

	All Races		White		Black	
	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005
All Sites	189.8	-1.3*	188.0	-1.1*	234.0	-1.9*
Lung and Bronchus	54.1	-1.0*	54.4	-0.9*	60.9	-1.7*
Colon and Rectum	18.8	-2.4*	18.3	-2.5*	26.1	-1.7*
Breast	14.1	-2.2*	13.7	-2.4*	20.9	-3.8*
Pancreas	10.6	0.2*	10.4	0.4*	19.9	-1.5*
Prostate ^f	10.1	-3.5*	9.3	-3.4*	13.7	-0.8*
Leukemia	7.4	-0.9*	7.6	-2.9*	7.8	-3.5*
Non-Hodgkin Lymphoma	7.3	-2.9*	7.6	-0.7*	6.9	-1.8*
Liver & IBD ^g	5.0	1.9*	5.1	-0.3	6.7	1.9*
Ovary ^f	5.0	-0.4	4.7	-1.0*	6.5	-1.5*
Esophagus	4.4	0.3*	4.6	1.9*	5.7	-4.4*
Brain and ONS ^g	4.4	-1.1*	4.5	0.0	5.1	-3.1*
Urinary Bladder	4.3	-0.2	4.4	1.2*	4.5	-0.2
Kidney and Renal Pelvis	4.2	-0.3	4.3	-0.2	4.3	0.1
Stomach	4.1	-3.2*	3.6	-3.3*	4.1	-0.5
Myeloma	3.7	-1.0*	3.4	-0.8*	3.7	-3.4*

	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005
All Sites	113.8	-1.7*	161.7	-0.4	127.0	-1.4*
Lung and Bronchus	26.6	-1.3*	40.9	0.0	23.2	-1.6*
Colon and Rectum	12.0	-2.1*	16.9	-1.6	13.3	-1.3*
Liver & IBD ^g	10.4	-0.7	9.7	1.0	8.7	-2.7*
Stomach	7.6	-4.2*	8.5	2.4	8.3	0.6*
Pancreas	7.4	0.1	8.2	-1.7	8.1	-3.0*
Breast	7.0	0.1	7.8	0.8	7.8	1.7*
Non-Hodgkin Lymphoma	4.6	-2.7*	7.1	-1.6	6.5	-2.3*
Prostate ^f	4.5	-4.6*	6.5	-0.1	5.5	-3.5*
Leukemia	3.9	-1.4*	5.4	-0.6	5.0	-1.2*
Ovary ^f	2.7	0.8	4.9	0.8	3.7	0.7*
Oral Cavity and Pharynx	2.2	-2.2*	4.2	0.4	3.4	0.1
Brain and ONS ^g	2.0	1.2	4.0	4.3*	3.0	-1.8*
Esophagus	1.8	-3.2*	3.9	-1.1	2.8	-0.2
Urinary Bladder	1.8	-0.3	2.4	-1.4	2.4	-0.4
Kidney and Renal Pelvis	1.7	-3.0*	2.1	-0.3	2.3	-2.2*

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Top 15 cancer sites selected based on 2001-2005 age-adjusted rates for the race/ethnic group.

^b Mortality data used in calculating the rates are analyzed from US mortality files provided by the National Center for Health Statistics, CDC. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). The rates shown for sex-specific cancer sites are calculated using the population for both sexes combined.

^c The APC is the Annual Percent Change over the time interval. Mortality data used in calculating the trends are analyzed from US mortality files provided by the National Center for Health Statistics, CDC. Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. The 2001-2005 Hispanic death rates do not include deaths from Minnesota, New Hampshire and North Dakota. The 1996-2005 Hispanic mortality trends do not include deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.

^f The rates for sex-specific cancer sites are calculated using the population for both sexes combined.

^g IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero ($p < .05$).

Table I-27
AGE-ADJUSTED U.S. DEATH RATES AND TRENDS FOR THE TOP 15 CANCER SITES^a BY RACE/ETHNICITY

			Males					
All Races			White			Black		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2001-2005	1996-2005		2001-2005	1996-2005		2001-2005	1996-2005
All Sites	234.4	-1.7*	All Sites	230.7	-1.5*	All Sites	313.0	-2.5*
Lung and Bronchus	72.0	-2.0*	Lung and Bronchus	71.3	-1.8*	Lung and Bronchus	93.1	-2.8*
Prostate	26.7	-4.1*	Prostate	24.6	-4.0*	Prostate	59.4	-4.0*
Colon and Rectum	22.7	-2.6*	Colon and Rectum	22.1	-2.7*	Colon and Rectum	31.8	-1.6*
Pancreas	12.2	0.1	Pancreas	12.1	0.2*	Pancreas	15.4	-0.9*
Leukemia	9.9	-0.8*	Leukemia	10.2	-0.7*	Stomach	11.5	-3.8*
Non-Hodgkin Lymphoma	9.3	-2.7*	Non-Hodgkin Lymphoma	9.7	-2.6*	Liver & IBD ^f	10.3	2.4*
Esophagus	7.8	0.5*	Urinary Bladder	7.9	-0.2	Esophagus	9.8	-4.6*
Urinary Bladder	7.5	-0.3*	Esophagus	7.8	1.3*	Leukemia	8.5	-1.8*
Liver & IBD ^f	7.3	2.1*	Liver & IBD ^f	6.7	2.1*	Myeloma	8.3	-2.0*
Kidney and Renal Pelvis	6.0	-0.4	Kidney and Renal Pelvis	6.2	-0.3	Oral Cavity and Pharynx	6.7	-3.0*
Stomach	5.7	-3.7*	Brain and ONS ^f	5.7	-1.0*	Non-Hodgkin Lymphoma	6.4	-3.7*
Brain and ONS ^f	5.4	-1.0*	Stomach	5.0	-3.8*	Kidney and Renal Pelvis	6.1	-0.5
Myeloma	4.6	-1.1*	Melanoma of the Skin	4.4	-0.1	Urinary Bladder	5.4	-1.0
Oral Cavity and Pharynx	4.0	-1.9*	Myeloma	4.3	-0.9*	Larynx	4.8	-3.3*
Melanoma of the Skin	3.9	-0.2	Oral Cavity and Pharynx	3.8	-1.7*	Brain and ONS ^f	3.2	-0.4
Asian/Pacific Islander			American Indian/Alaska Native ^d			Hispanic ^e		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2001-2005	1996-2005		2001-2005	1996-2005		2001-2005	1996-2005
All Sites	138.8	-2.0*	All Sites	190.0	-0.7	All Sites	159.0	-1.9*
Lung and Bronchus	37.5	-1.6*	Lung and Bronchus	50.2	-2.3*	Lung and Bronchus	35.1	-2.5*
Liver & IBD ^f	15.2	-1.3*	Prostate	21.1	-1.9	Prostate	20.6	-3.5*
Colon and Rectum	14.4	-2.1*	Colon and Rectum	20.5	-1.0	Colon and Rectum	16.5	-2.0*
Prostate	11.0	-4.1*	Liver & IBD ^f	10.6	1.4	Liver & IBD ^f	11.1	1.4*
Stomach	10.1	-3.9*	Stomach	9.9	0.1	Pancreas	9.1	0.3
Pancreas	8.0	-0.4	Kidney and Renal Pelvis	9.3	0.2	Stomach	8.7	-2.6*
Non-Hodgkin Lymphoma	5.6	-3.7*	Pancreas	8.6	3.3	Non-Hodgkin Lymphoma	6.7	-4.3*
Leukemia	5.1	-0.7	Esophagus	7.1	5.3*	Leukemia	6.3	-1.1
Oral Cavity and Pharynx	3.3	-3.0*	Non-Hodgkin Lymphoma	5.8	-2.1	Kidney and Renal Pelvis	5.3	0.5
Esophagus	3.1	-3.5	Leukemia	5.7	1.0	Esophagus	4.2	-2.1*
Urinary Bladder	2.9	0.0	Myeloma	4.6	1.8	Urinary Bladder	4.0	-1.0
Brain and ONS ^f	2.6	2.2	Oral Cavity and Pharynx	3.8	-	Myeloma	3.6	-1.8
Kidney and Renal Pelvis	2.4	-4.0*	Urinary Bladder	2.9	-	Brain and ONS ^f	3.4	-0.5
Myeloma	1.9	-2.9*	Brain and ONS ^f	2.8	3.1	Oral Cavity and Pharynx	2.6	-4.0*
Soft Tissue including Heart	0.9	-2.5	Larynx	2.1	-	Larynx	1.9	-4.1*

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Top 15 cancer sites selected based on 2001-2005 age-adjusted rates for the race/ethnic group.

^b Mortality data used in calculating the rates are analyzed from US mortality files provided by the National Center for Health Statistics, CDC. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval. Mortality data used in calculating the trends are analyzed from US mortality files provided by the National Center for Health Statistics, CDC. Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. The 2001-2005 Hispanic death rates do not include deaths from Minnesota, New Hampshire and North Dakota. The 1996-2005 Hispanic mortality trends do not include deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero ($p < .05$).

Table I-28
AGE-ADJUSTED U.S. DEATH RATES AND TRENDS FOR THE TOP 15 CANCER SITES^a BY RACE/ETHNICITY

	Females					
	All Races		White		Black	
	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005
All Sites	159.9	-1.0*	159.2	-0.9*	186.7	-1.2*
Lung and Bronchus	41.0	0.1	42.0	0.2	39.9	0.1
Breast	25.0	-2.1*	24.4	-2.2*	33.5	-1.5*
Colon and Rectum	15.9	-2.5*	15.3	-2.6*	22.4	-1.9*
Pancreas	9.3	0.2*	9.2	-0.1	12.4	-0.8*
Ovary	8.8	-0.1	9.0	0.4*	7.5	-0.1
Non-Hodgkin Lymphoma	5.9	-3.2*	6.2	-3.3*	7.1	0.3
Leukemia	5.6	-1.1*	5.7	-1.0*	6.0	-1.7*
Corpus and Uterus, NOS	4.1	0.1	3.9	0.1	5.5	-3.3*
Brain and ONS ^f	3.6	-1.3*	3.9	-1.1*	5.2	-1.3*
Liver & IBD ^f	3.1	1.1*	2.9	1.0*	4.7	-4.3*
Myeloma	3.0	-1.2*	2.8	-0.4	4.2	-2.3*
Stomach	2.9	-2.7*	2.8	-1.1*	3.9	0.6
Kidney and Renal Pelvis	2.7	-0.5*	2.5	-2.8*	2.8	-3.9*
Cervix Uteri	2.5	-3.4*	2.3	-3.2*	2.8	-1.7*
Urinary Bladder	2.3	-0.8*	2.2	-0.6*	2.7	-0.6
	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005	Rate ^b 2001-2005	APC ^c 1996-2005
All Sites	95.9	-1.1*	142.0	-0.2	105.2	-1.1*
Lung and Bronchus	18.5	-0.7	33.8	2.9*	15.8	-2.5*
Breast	12.6	0.0	17.1	1.0	14.6	-0.5
Colon and Rectum	10.2	-2.1*	14.2	-2.2	10.8	-0.9
Pancreas	6.9	0.5	7.5	0.6	7.6	0.7
Liver & IBD ^f	6.6	0.5	7.2	-1.3	6.0	0.4
Stomach	5.9	-4.4*	6.6	3.2	5.1	1.6*
Ovary	4.9	0.7	5.2	-3.5	4.9	-2.0*
Non-Hodgkin Lymphoma	3.8	-1.4	4.9	0.5	4.7	-2.8*
Leukemia	3.0	-2.1*	4.3	-0.8	4.0	-1.5*
Corpus and Uterus, NOS	2.4	0.4	4.1	-	3.2	-3.1*
Cervix Uteri	2.2	-5.2*	3.7	-2.8	3.2	0.0
Brain and ONS ^f	1.5	-0.1	3.6	-3.7	2.6	-1.7
Myeloma	1.4	-1.9	3.1	-	2.4	0.8
Oral Cavity and Pharynx	1.4	-0.5	2.2	-	2.4	0.0
Kidney and Renal Pelvis	1.2	-1.2	1.6	-	1.3	0.4

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Top 15 cancer sites selected based on 2001-2005 age-adjusted rates for the race/ethnic group.

^b Mortality data used in calculating the rates are analyzed from US mortality files provided by the National Center for Health Statistics, CDC. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval. Mortality data used in calculating the trends are analyzed from US mortality files provided by the National Center for Health Statistics, CDC. Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. The 2001-2005 Hispanic death rates do not include deaths from Minnesota, New Hampshire and North Dakota. The 1996-2005 Hispanic mortality trends do not include deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero ($p < .05$).