
Charmonium from Lattice QCD

Jo Dudek

Jefferson Lab / Old Dominion University

Charm '07 - Cornell

'parameters' of a lattice computation

- discretised charm & light quark actions
 - various varieties available (*clover, Fermilab scheme, domain-wall, overlap, staggered, HISQ, ...*)
- lattice spacing a (might be differing a_s, a_t) - $a \rightarrow 0$ desired
 - certain actions designed to speed this approach, e.g. domain-wall fermions: $X(a) = X(0) + \mathcal{O}(a^2)$
 - current lowest $\sim 0.06\text{fm}$ - perform extrapolations
- mass of light & strange quarks 'in the sea', $m_{q,s}$ - $m_q \rightarrow m_q^{\text{phys}} \sim 0$ desired
 - dynamical lattices, current lowest $m_\pi \sim 200\text{ MeV}$
 - quenched lattices neglect these quarks altogether
- volume of spatial lattice box, L^3 - $L \rightarrow \infty$ desired
 - sensitivity to this depends upon the states under study
- inclusion of disconnected diagrams (OZI)
 - usually just connected diagrams - effects probably small


which quantities can be computed?

- “easy” & approaching maturity:
 - spectrum of states below open charm decay threshold
 - leptonic decay constants
- relatively “easy” but only recently begun:
 - radiative transitions between charmonium states
 - two-photon decays of charmonium states
- “hard”:
 - hadronic decays of charmonia
 - accurate determination of state masses above threshold when they can decay

lightest five states

- have been studied the most
- relatively easy to extract
- use simple interpolators $\mathcal{O} = \bar{\psi}\Gamma\psi$
- can get masses and leptonic decay constants
- relative ease of computation means can devote effort to dealing with lattice systematics
- many groups have worked on this - no time to summarise them all
- a single recent example: *Phys.Rev.D75:054502,2007 (HPQCD & UKQCD)*
 - highly improved action (small effect in extrapolating $a \rightarrow 0$)
 - fine dynamical lattice $a \sim 0.09 \text{ fm}$, $m_\pi \sim 250 \text{ MeV}$
 - decay constant analysis is underway (C.Davies private communication)

$$\langle \mathcal{O}_f \mathcal{O}_i \rangle = \sum_N \frac{\langle 0 | \mathcal{O}_f | N \rangle \langle N | \mathcal{O}_i | 0 \rangle}{2m_N} e^{-m_N t}$$



higher spectrum


- higher spectrum results not at the same level of ‘minimised’ lattice systematics
- need larger set of interpolating fields (to get spin ≥ 2 and exotics)
- e.g. derivative based operators

$$\begin{aligned} & \bar{\psi} \Gamma \psi \\ & \bar{\psi} \Gamma \overleftrightarrow{D}_k \psi \\ & \bar{\psi} \Gamma \overleftrightarrow{D}_j \overleftrightarrow{D}_k \psi \end{aligned}$$

- recent study in **quenched** lattice QCD
 - somewhat improved Clover action
 - anisotropic lattice action $a_s = 3a_t$
 - establish if sophisticated analysis method can extract multiple excited states from lattice correlators

excited states

- an example of the difficulty in analysis
- the charmonium vector channel below and close to threshold:



- need a reliable excited state extraction procedure
- variational method utilises the **orthogonality** of states

excited states

● first results are promising



arXiv:0707.4162v1 [hep-lat] 27 Jul 2007

JLAB-THY-07-689

Charmonium excited state spectrum in lattice QCD

Joze J. Dudek
Jefferson Laboratory MS 12H2, 12000 Jefferson Avenue, Newport News, VA 23606, USA and
Department of Physics, Old Dominion University, Norfolk, VA 23529, USA*

Robert G. Edwards, Nilmani Mathur, and David G. Richards
Jefferson Laboratory MS 12H2, 12000 Jefferson Avenue, Newport News, VA 23606, USA



$a \rightarrow 0$



$$\langle 0 | (\bar{\psi} \Gamma \overleftrightarrow{D}_k \psi)_{T_2} | J \rangle = Z_J \cdot K_{T_2}^J$$

$$\langle 0 | (\bar{\psi} \Gamma \overleftrightarrow{D}_k \psi)_E | J \rangle = Z_J \cdot K_E^J$$

to my knowledge, the first time this has been seen



PC = --

[quenched & $a \neq 0$]





PC = ++

[quenched & $a \neq 0$]



PC = +-


[quenched & $a \neq 0$]




$$PC = -+$$

[quenched & $a \neq 0$]

naive analysis puts
states in lowest spin



exotic




other lattice studies claim a 1^+ near 4300 MeV


$$PC = -+$$

[quenched & $a \neq 0$]

equally plausible to assign the lightest state to be **non-exotic** 4^-



exotic



then the exotic 1^- is heavier (>4600 MeV)

radiative transitions

- real photon transitions ($Q^2=0$)
- lattice method will yield transition form-factors (at multiple Q^2)
- will extrapolate back to $Q^2=0$


PHYSICAL REVIEW D 73, 074507 (2006)

Radiative transitions in charmonium from lattice QCD

Jozef J. Dudek,* Robert G. Edwards, and David G. Richards

Jefferson Laboratory Mail-Stop 12H2, 12000 Jefferson Avenue, Newport News, Virginia 23606, USA

(Received 17 January 2006; published 20 April 2006)



radiative transitions

- extract from three-point functions involving the vector current

$$\begin{aligned}\Gamma(t_f, t; \vec{p}, \vec{q}) &= \sum_{\vec{x}, \vec{y}} e^{-i\vec{p}\cdot\vec{x}} e^{i\vec{q}\cdot\vec{y}} \langle \varphi_f(\vec{x}, t_f) j^\mu(\vec{y}, t) \varphi(\vec{0}, 0) \rangle \\ &\sim \sum_{n, m} e^{-E_{f_n}(t_f - t)} \langle 0 | \varphi_f(0) | f_n(\vec{p}) \rangle \\ &\quad \times \langle f_n(\vec{p}) | j^\mu(0) | i_m(\vec{p} + \vec{q}) \rangle \\ &\quad \times \langle i_m(\vec{p} + \vec{q}) | \varphi_i(0) | 0 \rangle e^{-E_{i_m} t}\end{aligned}$$

- overlaps and energies come from the spectrum analysis (two-point functions)

- matrix element related to the decay width


- e.g. $J/\psi \rightarrow \eta_c \gamma$

$$\langle \eta_c(\vec{p}') | j^\mu(0) | \psi(\vec{p}, r) \rangle = \frac{2V(Q^2)}{m_{\eta_c} + m_\psi} \epsilon^{\mu\alpha\beta\gamma} p'_\alpha p_\beta \epsilon_\gamma(\vec{p}, r)$$

$$\Gamma(\psi \rightarrow \eta_c \gamma) = \alpha_{\text{em}} \frac{|\vec{q}|^3}{(m_{\eta_c} + m_\psi)^2} \frac{64}{27} |\hat{V}(0)|^2$$


$J/\psi \rightarrow \eta_c \gamma$ transition

- statistically most precise channel, but very sensitive to the hyperfine splitting which is not correct on this quenched lattice ($\delta m_{\text{lat.}} \approx 80$ MeV, $\delta m_{\text{expt.}} \approx 117$ MeV)



- the Crystal Ball experimental value needs confirmation
- all eyes turn to Matt Shepherd & Ryan Mitchell at CLEO




$\chi_{c0} \rightarrow J/\psi \gamma$ $E1$ transition



$1P \rightarrow 1S$ transitions

- fit form inspired by potential models with spin-dependent corrections

$$E_1(Q^2) = E_1(0) \left(1 + \frac{Q^2}{\rho^2} \right) e^{-\frac{Q^2}{16\beta^2}}$$



$$\chi_{c0} \rightarrow J/\psi \gamma E1$$

$$\beta = 542(35) \text{ MeV}$$

$$\rho = 1.08(13) \text{ GeV}$$

$$\chi_{c1} \rightarrow J/\psi \gamma E1$$

$$\beta = 555(113) \text{ MeV}$$

$$\rho = 1.65(59) \text{ GeV}$$

$$h_c \rightarrow \eta_c \gamma E1$$

$$\beta = 689(133) \text{ MeV}$$

$$\rho \rightarrow \infty$$

simplest quark model has all β equal and $\rho(\chi_{c0}) = 2\beta$, $\rho(\chi_{c1}) = \sqrt{2} \cdot \rho(\chi_{c0})$, $\rho(h_c) \rightarrow \infty$

two-photon decays





- this is non-trivial in Euclidean lattice QCD
- the photon is not an eigenstate of QCD
- how do we 'make' one on the lattice
 - solution is to realise that it is a suitable sum of QCD eigenstates
 - like a 'vector dominance' picture
 - *exactly* expressed in the LSZ reduction of field theory
- explained carefully in *Ji & Jung PRL86, 208* & *Dudek & Edwards PRL97, 172001*

$$\begin{aligned} \langle \gamma(q_1, \lambda_1) \gamma(q_2, \lambda_2) | M(p) \rangle = \\ - \lim_{\substack{q'_1 \rightarrow q_1 \\ q'_2 \rightarrow q_2}} \epsilon_\mu^*(q_1, \lambda_1) \epsilon_\nu^*(q_2, \lambda_2) q_1'^2 q_2'^2 \int d^4x d^4y e^{iq'_1 \cdot y + iq'_2 \cdot x} \langle 0 | T \{ A^\mu(y) A^\nu(x) \} | M(p) \rangle \\ \rightarrow e^2 \epsilon_\mu^{(1)*} \epsilon_\nu^{(2)*} \int d^4y e^{-iq_1 \cdot y} \langle 0 | T \{ j^\mu(0) j^\nu(y) \} | M(p) \rangle \end{aligned}$$

- the 'extra' integral becomes a sum of a correlator over timeslices on the lattice

two-photon decays

- integrand is peaked and can be summed
- result is the form-factor for $\eta_c \rightarrow \gamma\gamma^*$



PRL 97, 172001 (2006)

PHYSICAL REVIEW LETTERS

week ending
27 OCTOBER 2006

Two-Photon Decays of Charmonia from Lattice QCD

Jozef J. Dudek* and Robert G. Edwards

Jefferson Laboratory MS 12H2, 12000 Jefferson Avenue, Newport News, Virginia 23606, USA
(Received 12 July 2006; published 23 October 2006)

Charmonium from Lattice QCD


What can lattice do for the onia?

- a lot, in principle
- but it is a long way behind established techniques like potential models
- many lattice groups calculate charmonium spectra
 - but not all are primarily interested in charmonium physics
 - use ‘precise’ comparison with the lower part of the spectrum to set the charm quark mass for D-meson flavor physics
- smaller number of groups trying to compute quantities beyond the spectrum
 - new techniques take time to get working
 - will initially not use “the best lattice systematics”
- US lattice groups have to beg for computing time every year
 - decided by a committee of lattice QCD theorists
 - explicit support from experimentalists is always helpful
 - if you think we’re computing the right quantities and want better calculations, please cite us

important physics
also important physics


Manke & Liao

- earlier study with similar operators
- less sophisticated analysis
- somewhat heavier I^{-+} reported



lattice technique

- fairly straightforward application of three-point correlators
- similar to pion, proton form-factor, $N \leftrightarrow \Delta$... calculations
- compute three-point functions with sequential-source technology
 - completely specify the sink (operator & momentum)
 - can insert any momentum



- obtain correlators at various values of photon Q^2

radiative transitions

- usually expressed in terms of multipoles
- covariant expressions can be derived
- e.g. $\chi_{c0} \rightarrow J/\psi \gamma$

$$\langle \chi_{c0}(\vec{p}_\chi) | V^\mu(0) | \psi(\vec{p}_\psi, r) \rangle = \Omega^{-1}(Q^2) \left(E_1(Q^2) \left[\Omega(Q^2) \epsilon^\mu(\vec{p}_\psi, r) - \epsilon(\vec{p}_\psi, r) \cdot p_\chi \left(p_\chi \cdot p_\psi p_\psi^\mu - m_\psi^2 p_\chi^\mu \right) \right] \right. \\ \left. + \frac{C_1(Q^2)}{\sqrt{Q^2}} m_\psi \epsilon(\vec{p}_\psi, r) \cdot p_\chi \left[p_\chi \cdot p_\psi (p_\chi + p_\psi)^\mu - m_\chi^2 p_\psi^\mu - m_\psi^2 p_\chi^\mu \right] \right)$$

- the multipole form-factors can be obtained from the three-point functions as an overconstrained linear problem
- need the E 's and Z 's from two point function fits
- deals with all the data at a given Q^2 simultaneously - in principle can simultaneously extract excited state transitions

$$\Gamma(p_f, p_i; t) = \sum_n P(p_f, p_i; t) \cdot K_n(p_f, p_i) \cdot f_n(Q^2)$$

$$P = \frac{Z_f Z_i}{4E_f E_i} e^{-E_f t_f} e^{-(E_i - E_f)t}$$

$$\begin{bmatrix} \Gamma(a; t) \\ \Gamma(b; t) \\ \Gamma(c; t) \\ \vdots \end{bmatrix} = \begin{bmatrix} P(a; t)K_1(a) & P(a; t)K_2(a) & \cdots \\ P(b; t)K_1(b) & P(b; t)K_2(b) & \cdots \\ P(c; t)K_1(c) & P(c; t)K_2(c) & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} f_1(Q^2)[t] \\ f_2(Q^2)[t] \\ \vdots \end{bmatrix}$$

first results

- quenched, anisotropic lattice
- $a_s = 0.1$ fm, $\xi = 3.0$, $12^3 \times 48$
- domain wall fermions ($L_5 = 16$)
 - charm quark mass tuning is not perfect (5% low)

- ground state to ground state transitions only


PHYSICAL REVIEW D **73**, 074507 (2006)

Radiative transitions in charmonium from lattice QCD

Jozef J. Dudek,* Robert G. Edwards, and David G. Richards

Jefferson Laboratory Mail-Stop 12H2, 12000 Jefferson Avenue, Newport News, Virginia 23606, USA

(Received 17 January 2006; published 20 April 2006)



$\chi_{c1} \rightarrow J/\psi \gamma$ transition

- derived the covariant multipole decomposition

$$\langle A(\vec{p}_A, r_A) | j^\mu(0) | V(\vec{p}_V, r_V) \rangle = \frac{i}{4\sqrt{2}\Omega(Q^2)} \epsilon^{\mu\nu\rho\sigma} (p_A - p_V)_\sigma \times$$

$$\times \left[E_1(Q^2) (p_A + p_V)_\rho \left(2m_A [\epsilon^*(\vec{p}_A, r_A) \cdot p_V] \epsilon_\nu(\vec{p}_V, r_V) + 2m_V [\epsilon(\vec{p}_V, r_V) \cdot p_A] \epsilon_\nu^*(\vec{p}_A, r_A) \right) \right.$$

$$+ M_2(Q^2) (p_A + p_V)_\rho \left(2m_A [\epsilon^*(\vec{p}_A, r_A) \cdot p_V] \epsilon_\nu(\vec{p}_V, r_V) - 2m_V [\epsilon(\vec{p}_V, r_V) \cdot p_A] \epsilon_\nu^*(\vec{p}_A, r_A) \right)$$

$$+ \frac{C_1(Q^2)}{\sqrt{q^2}} \left(-4\Omega(Q^2) \epsilon_\nu^*(\vec{p}_A, r_A) \epsilon_\rho(\vec{p}_V, r_V) \right.$$

$$\left. \left. + (p_A + p_V)_\rho \left[(m_A^2 - m_V^2 + q^2) [\epsilon^*(\vec{p}_A, r_A) \cdot p_V] \epsilon_\nu(\vec{p}_V, r_V) + (m_A^2 - m_V^2 - q^2) [\epsilon(\vec{p}_V, r_V) \cdot p_A] \epsilon_\nu^*(\vec{p}_A, r_A) \right] \right) \right].$$

- $E_1(Q^2)$ - electric dipole - experimentally measured at $Q^2 = 0$
- $M_2(Q^2)$ - magnetic quadrupole - experimentally measured (via photon angular dependence) at $Q^2 = 0$
- $C_1(Q^2)$ - longitudinal - goes to zero at $Q^2 = 0$
- this lattice $\delta m(\chi_{c1} - J/\psi)$ close to experiment, so small phase-space ambiguity

$\chi_{c1} \rightarrow J/\psi \gamma$ transition

no $Q^2 < 0$ points owing to kinematical structure of matrix element

