



Strong interaction physics with an Electron–Ion Collider

C. Weiss (JLab), Temple U. Physics Colloquium, Philadelphia, 12–Mar–12 weiss@jlab.org



Theoretical methods, simulations

- Internal structure of proton/neutron

Quantum Chromodynamics

Many-body system: relativistic, quantum-mechanical, strongly coupled
Cf. condensed matter, atomic physics

- High-energy electron scattering

Fixed-target [JLab 12 GeV](#)

Colliding beams [Electron–Ion Collider EIC](#)

- QCD with an Electron–Ion Collider

Brief summary: <http://arxiv.org/abs/arXiv:1110.1031>

Sea quark and gluon polarization

Spatial imaging of quarks/gluons

Gluons in nuclei and coherent effects

High gluon densities, saturation

- Road ahead

Why Quantum Chromodynamics

- Fundamental structure of matter

Origin of mass: $> 99\%$ from energy in strong fields

Phases of matter at high density/temperature: Early universe

Conversion of energy/radiation into matter: Cosmic ray physics

- Nuclei and nuclear reactions from “first principles”

Origin of nucleon–nucleon interaction

Nuclear energy, stellar structure in astrophysics

Fundamental symmetries: Neutrino interactions with nuclei

- Concepts and methods


Quantum field theory: Perturbative methods, renormalization group, topology, symmetry breaking

Geometric methods: Gauge \leftrightarrow string correspondence

Numerical simulations: Lattice gauge theory

Nucleon structure: Short distances

3



- Pointlike objects: Quarks

Practically massless $m_{u,d} < 0.01 m_p$

Fermions with spin $1/2$

Electromagnetic and weak charge:
Coupling to external probes!

- Quantum Chromodynamics

Gauge theory with $SU(3)$ group charge:
“Color,” cf. Electrodynamics


Effective coupling decreases with distance:
“Asymptotic freedom” Gross, Politzer, Wilczek 73

- Larger distances $r \gtrsim 0.3 \text{ fm}$

Strong non-perturbative fields create
condensate of quark–antiquark pairs


Dynamical mass generation

Dynamics changes with resolution scale!



Nucleon structure: Fields vs. particles

4



- Understand/describe nucleon structure in terms of QCD degrees of freedom!

Uniquely challenging problem:
relativistic + QM + strongly coupled

- Nucleon at rest: Interacting fields

Imaginary time $t \rightarrow i\tau$:
Statistical mechanics, lattice simulations


No concept of particle content:
Cannot separate “constituents”
from vacuum fluctuations!

- Nucleon fast: Particle content

Closed system: Wave function description
Gribov, Feynman


Components with different particle number:
 $|N\rangle = |qqq\rangle + |qqqq\bar{q}\rangle + |qqqg\rangle + \dots$

High-energy scattering process:
Snapshot with resolution $1/Q$



Nucleon structure: Many-body system

5



- Different components of wave function


Few particles with large $x \equiv$ fractional momentum
Many particles with small x

- Measurable properties

Particle number densities, incl. spin/flavor dependence
Transverse spatial distributions
Orbital motion: Transverse momenta, polarization
Particle-particle correlations

} change with
resolution scale
 $1/Q!$

Electron scattering: Probing short distances



- Electron–nucleon scattering

Also: positron, muon

EM interaction well-known

Momentum transfer \rightarrow resolution $1/Q$
 Energy transfer \rightarrow configuration x

Variety of final states: Inclusive, semi-inclusive, exclusive


- Accessible region


Want spatial resolution $1/Q \ll 0.2$ fm,
 or $Q \gg 1$ GeV

$Q^2 < x E_{CM}^2$ from kinematics.
 High Q^2 or small x require high energies!

- Alt: Nucleon–nucleon scattering with high-momentum transfer processes

Sensitive to antiquarks, gluons. LHC, Tevatron, RHIC





- Beam on fixed target

High rates from density of particles in target

Center-of-mass energy grows as $E_{\text{CM}}^2 = 2E_e M_p$

- Colliding beams



Higher energies: $E_{\text{CM}}^2 = 4E_e E_p$ Product of beam energies!


Energy-efficient: Beams collide multiple times

Clean: No scattering from atomic electrons



Detection: Recoil proton/nucleus, variable angles

Demands much higher beam quality:
Focusing, cooling, time structure

Integration of detectors and accelerator
elements at interaction point



Experience with storage rings: e^+e^- (LEP, PEP-II, KEK, DAΦNE),
 $pp/p\bar{p}$ (RHIC, Tevatron, LHC), AA (RHIC, LHC), ep (HERA)



- Luminosity

$$\text{Rate} = \text{Luminosity} \times \text{Cross section}$$

High luminosity required for rare processes
 exclusive channels, high p_T
 multidimensional binning spatial imaging
 precision measurements Q^2 dependence

Limiting factor in most nucleon structure experiments!

- JLab 12 GeV

Energy \times luminosity frontier in fixed-target scattering

- Electron-Ion Collider EIC

A high-luminosity, polarized ep/eA collider for QCD and nuclear physics!

Electron scattering: JLab 12 GeV

9

- “Race track” accelerator with linacs + arcs, extensible to 24 GeV

Uses unique superconducting RF technology and energy recovery

- Experimental halls

A, C Magnetic spectrometers
B Large acceptance CLAS

- 12 GeV Upgrade

Double beam energy 6 → 12 GeV

Add Hall D (γ beam, GlueX detector)


Upgrade existing halls

DOE project (CD0 2004, CD3 2008)

Construction on-going, beam exp. 2013

Total cost ~ 300M\$


<http://www.jlab.org/12GeV/>



CW beam $\sim 100 \mu A$

Present beam energy 6 GeV

Operating since 1994




- JLab ring–ring design MEIC/ELIC

11 GeV CEBAF as injector *continued fixed-target op*
 Medium–energy: 1 km ring, 3–11 on 60/96 GeV
 High–energy: 2.5 km ring, 3–11 on 250 GeV
 Luminosity $\sim 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ over wide energy range
 Figure–8 for polarization transport, up to four IP's

- BNL linac–ring design eRHIC

RHIC proton/ion beam up to 325 GeV
 5–20 (30) GeV electrons from linac in tunnel *staged*
 Luminosity $\sim 10^{34} (10^{33})$ over wide range
 Re-use RHIC detectors? *ePHENIX*



- Related proposals


CERN LHeC: 20–150 GeV on 7 TeV ep
 Ring–ring and linac–ring discussed, $L \sim 10^{33}$
 Mainly particle physics after LHC, but also high–energy QCD

GSI ENC: 3.3 GeV on 15 GeV $\vec{e}\vec{p}$
 Ring–ring using FAIR HESR, $L \sim 10^{32}$ *PANDA detector*

Convergence in parameters, “staging”
 Differences in technological challenges, cost (?)

Nucleon structure: Many-body system


11



- Measurable properties

- Particle densities, including spin/ flavor dependence
- Transverse spatial distributions
- Orbital motion: Transverse momenta, polarization
- Particle-particle correlations

} change with resolution scale $1/Q!$




- How are valence quarks in nucleon polarized at $x \rightarrow 1$?

Basic $3q$ component of nucleon wave fn

Non-perturbative QCD interactions?

Orbital angular momentum $L = 1$?



- d quark polarization from inclusive scattering on neutron


d in proton = u in neutron isospin symmetry

Poorly constrained by present data
SLAC, HERMES

- JLab12: Map d quark polarization precisely up to $x \sim 0.8$

Combination of energy and luminosity!


Many more applications: Spatial imaging, orbital motion, nuclei, . . .



- How are sea quarks polarized in nucleon?

Non-perturbative QCD interactions connecting valence ↔ sea quarks?

Flavor asymmetry related to mesonic degrees of freedom? “Pion cloud”




- Semi-inclusive scattering: Identify particles produced from struck quark

“Tag” charge and flavor of struck quark


Flavor asymmetries poorly determined from present data HERMES

- EIC: Map sea quark distributions and their spin dependence

High energy ensures independent fragmentation of struck quark



- How do gluons respond to nucleon spin?
 - Origin of non-perturbative gluon fields?
 - Gluon contribution to nucleon spin?
 - “Spin puzzle”
 - Orbital angular momentum in nucleon wave function?




- $\Delta G(x)$ presently poorly constrained

Q^2 dependence of polarized nucleon structure function $g_1(x, Q^2)$
 EMC/SMC, SLAC, HERMES, COMPASS, JLab 6/12 GeV

Hard processes in $\vec{p}\vec{p}$ RHIC Spin

- EIC: Fully quantitative determination of gluon polarization

Wide kinematic coverage enables study of Q^2 evolution



- How are quarks/gluons distributed in transverse space?

Fundamental size and “shape” of nucleon in QCD

Distributions change with x :
Diffusion, chiral dynamics

Input for modeling pp collisions at LHC


- Exclusive processes $\gamma^* + N \rightarrow J/\psi + N$


Gluonic form factor of nucleon:
Generalized parton distribution

Other channels γ, ρ^0, π, K
sensitive to quarks

- EIC: “Gluon imaging” of nucleon

Luminosity for low rates,
differential measurements






- How are the fundamental quark/gluon densities affected by nuclear binding?

$x > 0.1$ Modification of free nucleon structure:
Quark–gluon basis of NN interaction?

$x \ll 0.1$ QM interference of gluon fields
of different nucleons: “Shadowing”





- Nuclear gluon density poorly constrained by present data

Q^2 dependence of nuclear structure function $F_{2A}(x, Q^2)$

- EIC: Accurate determination of quark/gluon densities of nuclei!

Wide coverage in x, Q^2



- New dynamical scale in wave function at small x : $Q_s(x)$
 - Gluon density grows through QCD radiation
 - Theory: Non-linear QCD evolution, Classical fields “Color Glass Condensate”
 McLerran, Venugopalan; Balitsky, Kovchegov, JIMWLK
- New phenomena
 - Breakdown of Bjorken scaling in F_L, F_2
 - High p_T in forward particle production
 - Multiple hard processes, correlations
- Expected to be enhanced in nuclei
 - $Q_s(x) \sim A^{1/3}$ without shadowing, depends on nuclear gluon density
- EIC: Study saturation through inclusive/diffractive/exclusive processes

- Orbital motion of quarks and gluons
 - Transverse momenta and polarization effects in semi-inclusive hadron production
 - Quark/gluon orbital angular momentum, QCD spin-orbit interactions
- Color transparency: Interaction of small-size $q\bar{q}$ configurations with color fields
 - High- Q^2 meson production on nuclei
- Hadronization: Conversion of color charge to hadrons
 - Quark fragmentation, target break-up, correlations
 - Hadronization in the nuclear medium
- Electroweak physics
 - Neutral/charged current nucleon structure functions
 - Standard model parameters

EIC: Project status and next steps

19

- Informal recommendation in 2007 DOE/NSF NSAC Long-Range Plan
<http://science.energy.gov/np/nsac>
- EIC accelerator and physics R&D at Brookhaven and Jefferson Lab
<http://www.jlab.org/meic/>
International EIC Advisory Committee, three reviews
of physics and accelerator designs Feb-09, Nov-09, Apr-11

Increasingly supported by lab users [JLab User Workshops 2010](#)
- EIC Collaboration <http://web.mit.edu/eicc/>
Formed 2007, over 100 physicists from > 20 institutions, advancing EIC physics
and accelerator R&D. Semi-annual collaboration meetings/workshops
- EIC science discussed at 2011 Institute of Nuclear Theory INT Program
Very strong participation. Talks on-line at <http://www.int.washington.edu/PROGRAMS/10-3/>
Summary report available at <http://arxiv.org/abs/arXiv:1108.1713>
- Working toward full recommendation in 2013 NSAC LRP
[Further timeline tentative. Site selection? CD0? Budget realities](#)

Needs support of the nuclear physics and broader scientific community!

- Quantum Chromodynamics remains a uniquely challenging problem
 - Mature field with 40+ years of experience
 - Full of surprises! Revolutions sure to come
- Nucleon as a many-body system — a unifying perspective
 - Relativity + quantum mechanics + strong interactions
 - Connections with nuclear/condensed matter physics
- High-luminosity Electron-Ion Collider as the next-generation machine for QCD and nucleon structure
 - Complements/extends JLab 12 GeV, RHIC Spin, HERA small- x , and LHC/RHIC Heavy-Ion programs