

health concern in industrialized and resource-poor settings. Few reports are available from Africa, although hospital-associated ESBL producers have been described in Cameroon and the Central African Republic (6,7). ESBL-producing bacteria have been recovered from different sources in the community, including food and companion animals (8,9), and 1 recent study from India reported that a substantial number of tap water samples were contaminated with carbapenemase *bla*_{NDM-1} producing organisms (10).

Kinshasa is the second-largest city in sub-Saharan Africa. In 2008, of its estimated 8.7 million inhabitants, only 46% had access to safe drinking water, and 23% had access to improved sanitation facilities according to the World Bank. Opportunistic pathogens in drinking water and poor sanitary conditions may increase the risk of developing infectious enterocolitis for consumers, especially for those who are immunocompromised. It can eventually lead to chronic intestinal carriage of multidrug-resistant organisms. The presence of ESBL producers in the intestinal flora could also lead to horizontal transfer of drug resistance genes from commensal flora to enteric pathogens. This emergence of ESBL-producing bacteria and further community-associated infections poses a public threat, especially in low-resource countries where surveillance is suboptimal and empiric treatment of invasive infections often includes third-generation cephalosporins.

**Hilde De Boeck,
Berthe Miwanda,
Octavie Lunguya-Metila,
Jean-Jacques Muyembe-Tamfum,
Ellen Stobberingh,
Youri Glupczynski,
and Jan Jacobs**

Author affiliations: Institute of Tropical Medicine, Antwerp, Belgium (H. De Boeck, J. Jacobs); Institut National de Recherche Biomédicale, Kinshasa, Democratic Republic of the Congo (B. Miwanda, O. Lunguya-Metila, J.J. Muyembe-Tamfum); Maastricht University Medical Center, Maastricht, the Netherlands (E. Stobberingh); and Cliniques Universitaires Université Catholique de Louvain de Mont-Godinne, Yvoir, Belgium (Y. Glupczynski)

DOI: <http://dx.doi.org/10.3201/eid1806.111214>

References

- Okeke IN, Laxminarayan R, Bhutta ZA, Duse AG, Jenkins P, O'Brien TF, et al. Antimicrobial resistance in developing countries. Part I: recent trends and current status. *Lancet Infect Dis*. 2005;5:481–93. [http://dx.doi.org/10.1016/S1473-3099\(05\)70189-4](http://dx.doi.org/10.1016/S1473-3099(05)70189-4)
- Vlieghe E, Phoba MF, Tamfun JJ, Jacobs J. Antibiotic resistance among bacterial pathogens in central Africa: a review of the published literature between 1955 and 2008. *Int J Antimicrob Agents*. 2009;34:295–303. <http://dx.doi.org/10.1016/j.ijantimicag.2009.04.015>
- Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing: 21st informational supplement. CLSI document M100-S21. Wayne (PA): The Institute; 2011.
- Endimiani A, Hujer AM, Hujer KM, Gatta JA, Schriver AC, Jacobs MR, et al. Evaluation of a commercial microarray system for detection of SHV-, TEM-, CTX-M-, and KPC-type β -lactamase genes in gram-negative isolates. *J Clin Microbiol*. 2010;48:2618–22. <http://dx.doi.org/10.1128/JCM.00568-10>
- Bradford PA. Extended-spectrum β -lactamases in the 21st century: characterization, epidemiology, and detection of this important resistance threat. *Clin Microbiol Rev*. 2001;14:933–51. <http://dx.doi.org/10.1128/CMR.14.4.933-951.2001>
- Gangoué-Piéboji J, Bedenic B, Koulla-Shiro S, Randegger C, Adiogo D, Ngassam P, et al. Extended-spectrum β -lactamase-producing *Enterobacteriaceae* in Yaounde, Cameroon. *J Clin Microbiol*. 2005;43:3273–7. <http://dx.doi.org/10.1128/JCM.43.7.3273-3277.2005>
- Frank T, Arlet G, Gautier V, Talarmin A, Bercion R. Extended-spectrum β -lactamase-producing *Enterobacteriaceae*, Central African Republic. *Emerg Infect Dis*. 2006;12:863. <http://dx.doi.org/10.3201/eid1205.050951>
- Ewers C, Grobbel M, Stamm I, Kopp PA, Diehl I, Semmler T, et al. Emergence of human pandemic O25:H4-ST131 CTX-M-15 extended-spectrum β -lactamase-producing *Escherichia coli* among companion animals. *J Antimicrob Chemother*. 2010;65:651–60. <http://dx.doi.org/10.1093/jac/dkq004>
- Warren RE, Ensor VM, O'Neill P, Butler V, Taylor J, Nye K, et al. Imported chicken meat as a potential source of quinolone-resistant *Escherichia coli* producing extended-spectrum β -lactamases in the UK. *J Antimicrob Chemother*. 2008;61:504–8. <http://dx.doi.org/10.1093/jac/dkm517>
- Walsh TR, Weeks J, Livermore DM, Toleman MA. Dissemination of NDM-1 positive bacteria in the New Delhi environment and its implications for human health: an environmental point prevalence study. *Lancet Infect Dis*. 2011;11:355–62. [http://dx.doi.org/10.1016/S1473-3099\(11\)70059-7](http://dx.doi.org/10.1016/S1473-3099(11)70059-7)

Address for correspondence: Hilde De Boeck, Institute of Tropical Medicine, Nationalestraat 155, Antwerp, Belgium; email: hdeboeck@itg.be

Novel *Chlamydiaceae* Disease in Captive Salamanders

To the Editor: Although 2 major diseases of amphibians, chytridiomycosis and ranavirosis, have been relatively well studied, enigmatic amphibian disease and death not attributable to any of the known amphibian diseases frequently occur (1). We describe an apparently new disease in salamanders that is associated with a novel genus within the family *Chlamydiaceae*.

The salamanders seen in our clinic belonged to 1 of the following species: *Salamandra corsica*, the Corsican fire salamander (5 animals from 1 collection); *Neurergus crocatus*, the yellow spotted newt (11 animals from 3 collections); or *N. strauchii*, Strauch's

spotted newt (6 animals from 2 collections). All salamanders were captive bred; housed in breeding colonies in private collections in Elsloo and Eindhoven, the Netherlands, Munich, Germany, and Brugge, Belgium; and 1–3 years of age.

Disease was characterized by anorexia, lethargy, edema, and markedly abnormal gait. Mortality rate was 100%. Animals in these collections had no histories of disease. All animals were in good nutritional condition. Necropsy did not yield any macroscopic lesions. All animals had mild intestinal nematode or protozoan infections. Results of real-time PCRs for iridoviruses in liver and skin (2) or *Batrachochytrium dendrobatidis* fungus of skin (3) were negative for all animals.

We placed liver suspensions from the dead salamanders on Columbia agar with 5% sheep blood and tryptic soy agar and then incubated the samples up to 14 days at 20°C. No consistent bacterial growth was observed. Histologic examination of 2 Corsican fire salamanders and 1 yellow spotted newt revealed hepatitis in 1 of the Corsican fire salamanders and the yellow spotted newt. Hepatitis was characterized by high numbers of melanomacrophages and a marked infiltration of granulocytic leukocytes. Immunohistochemical staining for chlamydia (IMAGEN Chlamydia; Oxoid, Basingstone, UK) showed cell-associated fluorescently stained aggregates in liver tissue, suggestive of Chlamydiales bacteria. Transmission electron microscopic examination of the liver of a yellow spotted newt revealed intracellular inclusions containing particles matching the morphology of reticulate or elementary bodies of *Chlamydiaceae* (online Technical Appendix, wwwnc.cdc.gov/EID/pdfs/11-1137-Techapp.pdf).

A PCR (4) to detect the 16S rRNA of all Chlamydiales bacteria, performed on liver tissue samples from all animals, yielded positive results

in all 5 Corsican fire salamanders; in 4/7, 1/3, and 1/1 yellow spotted newts; and in 4/5 and 1/1 Strauch's spotted newts. For taxon identification, the 16S rRNA gene of the Chlamydiales bacteria was amplified and sequenced from the livers from 2 yellow spotted newts (1 from the collection in Elsloo, the Netherlands and 1 from the collection in Munich, Germany), 1 Strauch's spotted newt, and 5 Corsican fire salamanders.

The sequences shared >90% nt identity with the 16S rRNA gene of *C. abortus* B577 (GenBank accession no. D85709) and therefore can be identified as a member of the family *Chlamydiaceae* (5). The closest 16S rRNA similarity (92%) was observed with *C. psittaci* strain CPX0308 (AB285329). The sequence obtained from all spotted newt species specimens was identical (GenBank accession no. JN392920) but differed slightly (1%) from that obtained from the fire sala-

mander species specimens (GenBank accession no. JN392919). These sequence differences point to the existence of multiple strains with possible host adaptation.

We determined the phylogenetic position of the novel taxon, named *Candidatus Amphibiichlamydia salamandrae* (online Technical Appendix), identified by using neighbor-joining analysis with Kodon software (Applied Maths, Sint-Martens-Latem, Belgium). The novel Chlamydiales forms a distinct branch in the well-supported monophyletic clade with the genera *Chlamydia* and *Candidatus Clavochlamydia salmonicola* (family *Chlamydiaceae*) (Figure). Maximum parsimony and unweighted pair group with arithmetic mean analyses yielded cladograms with the same topology (results not shown). Previous reports of members of the family *Chlamydiaceae* in amphibians concerned species occurring in other vertebrate taxa as well:




Figure. Topology of the novel amphibian *Chlamydiaceae* (*Candidatus Amphibiichlamydia salamandrae*) within the phylogenetic tree obtained by neighbor-joining and based on 16S rRNA gene data from representative species. Numbers show the percentage of times each branch was found in 1,000 bootstrap replicates. The tree has been rooted with *Verrucomicrobium spinosum* as outgroup. Scale bar indicates nucleotide substitutions per site.

C. psittaci, *C. pneumoniae*, *C. abortus*, and *C. suis* (6–10). To our knowledge, this member of the family *Chlamydiaceae* has been seen in amphibians, but not in other vertebrate hosts. The 16S rRNA analysis showed this taxon to belong to a clade with *Candidatus Clavochlamydia salmonicola*, a taxon found in fish. The phylogenetic position of the novel taxon in the family *Chlamydiaceae* thus roughly reflects the phylogenetic relation between the host species, providing evidence for host–bacterium co-evolution in the family *Chlamydiaceae*.

Although the results obtained are not conclusive with regard to the pathogenic potential of this novel genus and species of Chlamydiales, we were not able to attribute the clinical signs to any known disease. We therefore suggest that we discovered a novel bacterial taxon with possible considerable impact on amphibian health.

Acknowledgments

We are grateful to Jean Euzéby for support with the nomenclature. We thank Rob Keulers for his cooperation.

**An Martel, Connie Adriaensen,
Sergé Bogaerts,
Richard Ducatelle,
Herman Favoreel,
Sandra Crameri, Alex D. Hyatt,
Freddy Haesebrouck,
and Frank Pasmans**

Author affiliations: Ghent University, Merelbeke, Belgium (A. Martel, C. Adriaensen, R. Ducatelle, H. Favoreel, F. Haesebrouck, F. Pasmans); Waalre, the Netherlands (S. Bogaerts); Commonwealth Scientific and Industrial Research Organisation, Geelong, Victoria, Australia (S. Crameri, A.D. Hyatt)

DOI: <http://dx.doi.org/10.3201/eid1806.111137>

References

1. Daszak P, Berger L, Cunningham AA, Hyatt AD, Green DE, Speare R. Emerging infectious diseases and amphibian population declines. *Emerg Infect Dis.* 1999;5:735–48. <http://dx.doi.org/10.3201/eid0506.990601>
2. Mao J, Hedrick RP, Chichar VB. Molecular characterization, sequence analysis, and taxonomic position of newly isolated fish iridoviruses. *Virology.* 1997;229:212–20. <http://dx.doi.org/10.1006/viro.1996.8435>
3. Boyle DG, Boyle DB, Olsen V, Morgan JAT, Hyatt AD. Rapid quantitative detection of Chytridiomycosis (*Batrachochytrium dendrobatidis*) in amphibian samples using real-time Taqman PCR assay. *Dis Aquat Organ.* 2004;60:141–8. <http://dx.doi.org/10.3354/dao060141>
4. Everett KDE. *Chlamydia* and Chlamydiales: more than meets the eye. *Vet Microbiol.* 2000;75:109–26. [http://dx.doi.org/10.1016/S0378-1135\(00\)00213-3](http://dx.doi.org/10.1016/S0378-1135(00)00213-3)
5. Everett KDE, Bush RM, Andersen AA. Emended description of the order Chlamydiales, proposal of *Parachlamydiaceae* fam. nov. and *Simkaniaceae* fam. nov., each containing one monotypic genus, revised taxonomy of the family *Chlamydiaceae*, including a new genus and five new species, and standards for the identification of organisms. *Int J Syst Bacteriol.* 1999;49:415–40. <http://dx.doi.org/10.1099/00207713-49-2-415>
6. Berger L, Volp K, Mathews S, Speare R, Timms P. *Chlamydia pneumoniae* in a free-ranging giant barred frog (*Mixophyes iterates*) from Australia. *J Clin Microbiol.* 1999;37:2378–80.
7. Blumer C, Zimmermann DR, Weilenmann R, Vaughan L, Pospischil A. *Chlamydiae* in free-ranging and captive frogs in Switzerland. *Vet Pathol.* 2007;44:144–50. <http://dx.doi.org/10.1354/vp.44-2-144>
8. Howerth EW. Pathology of naturally-occurring chlamydiosis in African clawed frogs (*Xenopus laevis*). *Vet Pathol.* 1984;21:28–32.
9. Newcomer CE, Anver MR, Simmons JL, Wilcke RW, Nace GW. Spontaneous and experimental infections of *Xenopus laevis* with *Chlamydia psittaci*. *Lab Anim Sci.* 1982;32:680–6.
10. Reed KD, Ruth GR, Meyer JA, Shukla SK. *Chlamydia pneumoniae* infection in a breeding colony of African clawed frogs (*Xenopus tropicalis*). *Emerg Infect Dis.* 2000;6:196–9. <http://dx.doi.org/10.3201/eid0602.000216>

Address for correspondence: An Martel, Department of Pathology, Bacteriology and Avian Diseases, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, B-9820 Belgium; email: an.martel@ugent.be

All material published in *Emerging Infectious Diseases* is in the public domain and may be used and reprinted without special permission; proper citation, however, is required.

Novel Variant of Beilong Paramyxovirus in Rats, China

To the Editor: In 2003, two cDNA strands were identified in a human mesangial cell line during experimental screening for genes upregulated by angiotensin II (1). Sequence analysis showed that the strands were homologous to the matrix, fusion, and phosphoprotein genes of paramyxoviruses, suggesting the possibility of a novel paramyxovirus (2,3). Subsequent research found that these sequences, believed to originate from human kidney mesangial cell lines, were not amplifiable from such cell lines or human kidney samples but were amplifiable from a rat kidney mesangial cell line (4). Isolation and complete genome sequencing of the virus confirmed that it was a novel paramyxovirus of the subfamily *Paramyxovirinae*, named Beilong virus (BeV).

BeV is most closely related to J virus, discovered in auticulture of kidney tissue from a moribund house mouse, and Tailam virus from Sikkim rats (5,6). Because J virus and Tailam virus were found to originate in rodents and BeV was amplifiable from a rat kidney mesangial cell line, we hypothesized that BeV was a novel paramyxovirus originating in rats. To test this hypothesis, we conducted a territorywide molecular epidemiologic study of rats and other mammals to evaluate this novel paramyxovirus.

We tested 4,130 samples from 1,398 animals collected from various locations in Hong Kong, People's Republic of China, during September 2008–August 2009 (Table). These included 480 kidney, spleen, respiratory swab, and anal swab samples from 120 asymptomatic rats (105 brown rats [*Rattus norvegicus*] and 15 black rats [*R. rattus*]). To