

National Cancer Institute
CARCINOGENESIS
Technical Report Series
No. 200
NTP No. 80-20
1980

**BIOASSAY OF
2, 6-TOLUENEDIAMINE DIHYDROCHLORIDE
FOR POSSIBLE CARCINOGENICITY**

CAS No. 15481-70-6

NCI-CG-TR-200

NTP-80-20

**U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
National Institutes of Health**



BIOASSAY OF
2,6-TOLUENEDIAMINE DIHYDROCHLORIDE
FOR POSSIBLE CARCINOGENICITY

Carcinogenesis Testing Program
National Cancer Institute
National Institutes of Health
Bethesda, Maryland 20205
and
National Toxicology Program
Research Triangle Park
Box 12233
North Carolina 27709

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
National Institutes of Health

NIH Publication No. 80-1756
August 1980

BIOASSAY OF
2,6-TOLUENEDIAMINE DIHYDROCHLORIDE
FOR POSSIBLE CARCINOGENICITY

Carcinogenesis Testing Program
National Cancer Institute/National Toxicology Program

FOREWORD

This report presents the results of the bioassay of 2,6-toluenediamine dihydrochloride conducted for the Carcinogenesis Testing Program, National Cancer Institute (NCI)/National Toxicology Program (NTP). This is one of a series of experiments designed to determine whether selected chemicals have the capacity to produce cancer in animals. A negative result, in which the test animals do not have a greater incidence of cancer than control animals, does not necessarily mean that a test chemical is not a carcinogen, inasmuch as the experiments are conducted under a limited set of circumstances. A positive result demonstrates that a test chemical is carcinogenic for animals under the conditions of the test and indicates that exposure to the chemical is a potential risk to man. The actual determination of the risk to man from chemicals found to be carcinogenic in animals requires a wider analysis.

CONTRIBUTORS

The bioassay of 2,6-toluenediamine dihydrochloride was conducted at EG&G Mason Research Institute, Worcester, Massachusetts, under a subcontract to Tracor Jitco, Inc., the prime contractor for the NCI Carcinogenesis Testing Program.

The bioassay was conducted under the supervision of Drs. A. Handler (1,2), H. Lilja (1), E. Massaro (1,3), and E. Smith (1,4), principal investigators, and Mr. G. Wade (1). The program manager was Ms. R. Monson (1). Ms. A. Good (1) supervised the technicians in charge of animal care, and Ms. E. Zepp (1) supervised the preparation of the feed mixtures and collected samples of the diets for analysis. Ms. D. Bouthot (1) kept all daily records of the test. Dr. A. S. Krishna Murthy (1), pathologist, directed the necropsies and performed the histopathologic examinations. The pathology report and selected slides were evaluated by the NCI Pathology Working Group as described in Ward et al. (1978).

Animal pathology tables and survival tables were compiled at EG&G Mason Research Institute, Rockville, Maryland (5). The statistical analyses were performed by Dr. J. R. Joiner (6) and Ms. S. Vatsan (6), using methods selected for the bioassay program by Dr. J. J. Gart (7).

Chemicals used in this bioassay were analyzed at Midwest Research Institute (8), and dosed feed mixtures were analyzed by Dr. M. Hagopian (1).

This report was prepared at Tracor Jitco (6) under the direction of NCI. Those responsible for the report at Tracor Jitco were Dr. C. R. Angel, Director of the Bioassay Program; Dr. S. S. Olin, Deputy Director for Science; Dr. R. L. Schueler, pathologist; Dr. A. C. Jacobs, bioscience writer; and Dr. W. D. Theriault and Ms. M. W. Glasser, technical editors.

The following scientists at NCI (9) were responsible for evaluating the bioassay experiment, interpreting the results, and reporting the findings: Dr. Kenneth C. Chu, Dr. Michael P. Dieter, Dr. J. Fielding Douglas, Dr. Richard A. Griesemer, Dr. Charles K. Grieshaber, Dr. Thomas E. Hamm, Dr. William V. Hartwell, Dr. Y. Jack Lee, Dr. Harry Mahar, Dr. Harry A. Milman, Dr. Thomas W. Orme, Dr. Marcelina B. Powers, Dr. Sherman F. Stinson, Dr. Jerrold M. Ward, and Dr. Carrie E. Whitmire.

-
- (1) EG&G Mason Research Institute, 57 Union Street, Worcester, Massachusetts.
 - (2) Now with Findley Research Co., P.O. Box 375, Assonet, Massachusetts.
 - (3) Now with Pennsylvania State University, 226 Fenske Laboratory, University Park, Pennsylvania.
 - (4) Now with the University of Massachusetts Medical Center, 55 Lake Avenue, Worcester, Massachusetts.
 - (5) EG&G Mason Research Institute, 1530 East Jefferson Street, Rockville, Maryland.
 - (6) Tracor Jitco, Inc., 1776 East Jefferson Street, Rockville, Maryland.
 - (7) Mathematical Statistics and Applied Mathematics Section, Biometry Branch, Field Studies and Statistics, Division of Cancer Cause and Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.
 - (8) Midwest Research Institute, 425 Volker Boulevard, Kansas City, Missouri.
 - (9) Carcinogenesis Testing Program, National Cancer Institute, National Institutes of Health, Bethesda, Maryland, National Toxicology Program, Research Triangle Park, Box 12233, North Carolina.

SUMMARY

2,6-Toluenediamine is used as an intermediate in the production of dyes for furs and textiles, and of flexible polyurethane foams and elastomers. A bioassay of 2,6-toluenediamine dihydrochloride for possible carcinogenicity was conducted by feeding diets containing the test chemical to F344 rats and B6C3F1 mice.

Groups of 50 rats of each sex were fed the test chemical at two doses, 250 or 500 ppm, for 103 weeks and observed for 1 additional week. Groups of 50 mice of each sex were fed the test chemical at two doses, 50 or 100 ppm, for 103 weeks and then observed for 1 additional week. Groups of 50 untreated rats and 50 untreated mice of each sex were used as matched controls. All surviving animals were killed and necropsied at 104 weeks.

Weight gain depression was less than 10% for dosed groups of male rats and male and female mice, when compared with controls. Mean body weight gain was depressed 17% in low-dose female rats and 27% in high-dose female rats. Mortality was not increased in rats or mice of either sex by the test chemical. No clinical evidence indicated that mice of either sex received a maximum tolerated dose of the compound.

In male rats, islet-cell adenomas of the pancreas and neoplastic nodules or carcinomas of the liver occurred in dose-related trends that were significant using the Cochran-Armitage test ($P=0.025$ and $P=0.037$, respectively). The results of the Fisher exact test were not significant for either lesion. The occurrences of tumors in dosed female rats were not significantly different from those in control rats.

Significant results in the negative direction were observed in the incidences of C-cell tumors of the thyroid in male rats and of fibroadenomas of the mammary gland in female rats.

In male mice, in the low-dose group, lymphomas occurred at an incidence significantly higher ($P=0.046$) than that of the corresponding control group; however, the incidence was not significant when the Bonferroni criterion for multiple comparison was used.

The occurrence of hepatocellular carcinomas in female mice was dose related, but the result of the Fisher exact test comparing the incidence in the high-dose group with that in the controls was not significant.

It was concluded that, under conditions of the bioassay, 2,6-toluenediamine dihydrochloride was not carcinogenic for male and female F344 rats or for male and female B6C3F1 mice.

TABLE OF CONTENTS

	<u>Page</u>
I. Introduction	1
II. Materials and Methods	3
A. Chemical	3
B. Dietary Preparation	3
C. Animals	4
D. Animal Maintenance	4
E. Range-Finding and 14-Day Repeated Dose Studies	5
F. Subchronic Studies	7
G. Chronic Studies	10
H. Clinical Examinations and Pathology	10
I. Data Recording and Statistical Analyses	12
III. Results - Rats	15
A. Body Weights and Clinical Signs (Rats)	15
B. Survival (Rats)	15
C. Pathology (Rats)	15
D. Statistical Analyses of Results (Rats)	18
IV. Results - Mice	27
A. Body Weights and Clinical Signs (Mice)	27
B. Survival (Mice)	27
C. Pathology (Mice)	27
D. Statistical Analyses of Results (Mice)	30
V. Discussion	39
VI. Conclusions.....	41
VII. Bibliography	43

APPENDIXES

Appendix A	Summary of the Incidence of Neoplasms in Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet	45
Table A1	Summary of the Incidence of Neoplasms in Male Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet	47

	<u>Page</u>	
Table A2	Summary of the Incidence of Neoplasms in Female Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet.....	51
Appendix B	Summary of the Incidence of Neoplasms in Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet.....	55
Table B1	Summary of the Incidence of Neoplasms in Male Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet.....	57
Table B2	Summary of the Incidence of Neoplasms in Female Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet.....	61
Appendix C	Summary of the Incidence of Nonneoplastic Lesions in Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet	65
Table C1	Summary of the Incidence of Nonneoplastic Lesions in Male Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet	67
Table C2	Summary of the Incidence of Nonneoplastic Lesions in Female Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet	71
Appendix D	Summary of the Incidence of Nonneoplastic Lesions in Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet	75
Table D1	Summary of the Incidence of Nonneoplastic Lesions in Male Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet	77
Table D2	Summary of the Incidence of Nonneoplastic Lesions in Female Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet	81
Appendix E	Analysis of 2,6-Toluenediamine - Midwest Research Institute	85


	<u>Page</u>
Appendix F Analysis of 2,6-Toluenediamine dihydrochloride - Midwest Research Institute	93
Appendix G Stability Analysis of 2,6-Toluenediamine dihydrochloride - Midwest Research Institute	101
Appendix H Analytical Procedure for Analysis of 2,6-Toluenediamine dihydrochloride in Feed	105

TABLES

Table 1 Dosage and Survival of Rats and Mice Administered a Single Dose of 2,6-Toluenediamine dihydrochloride by Gavage	6
Table 2 Dosage and Survival of Rats and Mice Administered Repeated Doses of 2,6-Toluenediamine dihydrochloride in the Feed for 14 days	6
Table 3 Dosage, Survival, and Mean Body Weights of Rats Administered 2,6-Toluenediamine dihydrochloride in Diets for the 13-Week Subchronic Study	8
Table 4 Dosage, Survival, and Mean Body Weights of Mice Administered 2,6-Toluenediamine dihydrochloride in Diets for the 13-Week Subchronic Study	9
Table 5 Experimental Design of Chronic Feeding Studies with 2,6-Toluenediamine dihydrochloride in Rats and Mice	11
Table 6 Analyses of the Incidence of Primary Tumors in Male Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet	21
Table 7 Analyses of the Incidence of Primary Tumors in Female Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet	25
Table 8 Analyses of the Incidence of Primary Tumors in Male Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet	32
Table 9 Analyses of the Incidence of Primary Tumors in Female Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet	35

	<u>FIGURES</u>	<u>Page</u>
Figure 1	Growth Curves for Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet	16
Figure 2	Survival Curves for Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet	17
Figure 3	Growth Curves for Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet	28
Figure 4	Survival Curves for Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet	29
Figure 5	Infrared Absorption Spectrum of 2,6-Toluenediamine	89
Figure 6	Nuclear Magnetic Resonance Spectrum of 2,6-Toluenediamine	91
Figure 7	Infrared Absorption Spectrum of 2,6-Toluenediamine dihydrochloride.....	98
Figure 8	Nuclear Magnetic Resonance Spectrum of 2,6-Toluenediamine dihydrochloride	99

I. INTRODUCTION



2,6-TOLUENEDIAMINE DIHYDROCHLORIDE

2,6-Toluenediamine (CAS 823-40-5) is a by-product of the synthesis of 2,4-toluenediamine (CAS 95-80-7).

Mixtures of 2,6-toluenediamine and 2,4-toluenediamine are used in the synthesis of toluene diisocyanate, which is the predominant isocyanate used in the production of flexible polyurethane foams and elastomers (Milligan, 1968; Layer, 1964). Commercial mixtures of 2,4- and 2,6-toluenediamine normally contain 80% of the 2,4 isomer and 20% of the 2,6 isomer. A commercial mixture containing 65% of the 2,4 isomer and 35% of the 2,6 is also available (Buist, 1970). Annual production of the 80/20 mixture of 2,4- and 2,6-toluene diisocyanate in 1977 was 583 million pounds (United States International Trade Commission, 1978). Based on this figure, the annual production of 2,6-toluenediamine in 1977 is estimated to have been greater than 116 million pounds. 2,6-Toluenediamine (or its dihydrochloride salt) is also an intermediate used in the synthesis of several dyes used for textiles and furs (Society of Dyers and Colourists, 1971).

During production of urethane foam, significant quantities of volatile toluene diisocyanate are carried into the air above the foam and may reach concentrations as high as 10 ppm (Buist, 1970). These isocyanates are unstable in an aqueous environment and are readily converted to toluenediamine (Lowe, 1970).

2,6-Toluenediamine, activated by rat or mouse liver microsomal fractions, is weakly mutagenic in Salmonella typhimurium TA 1538, with a reversion rate similar to 2,4-toluenediamine (Dybing and Thorgeirsson, 1977).

2,6-Toluenediamine was selected for testing by the Carcinogenesis Testing Program because of its close structural relationship to 2,4-toluenediamine, a reported carcinogen (Ito et al., 1969). (The carcinogenic effects of 2,4-toluenediamine were recently confirmed -- NCI, 1979.) The dosed feed route was chosen for its convenience and comparability with results from several previous studies using structural analogs which had been administered by this route. The dihydrochloride of 2,6-toluenediamine (CAS 15481-70-6) was found to be more stable than the free amine in feed and was used in the bioassay.

II. MATERIALS AND METHODS

A. Chemical

2,6-Toluenediamine (Lot No. 012157) containing a trace impurity (unidentified) was obtained from Aldrich Chemical Company, Metuchen, New Jersey, and converted to the dihydrochloride salt (Lot No. 090857-MRI) at Midwest Research Institute. Analyses of both 2,6-toluenediamine and 2,6-toluenediamine dihydrochloride at Midwest Research Institute (elemental analysis; melting point, thin-layer and vapor-phase chromatography; and spectral analyses including infrared, ultraviolet, and nuclear magnetic resonance) were consistent with the respective structures (Appendixes E and F). Vapor-phase chromatography revealed a single homogeneous peak at two alternate settings. Thin-layer chromatography showed only a trace impurity (less than 1%) at the origin. 2,6-Toluenediamine dihydrochloride was stored at 4°C in its original container and transferred to amber stock bottles as needed.

B. Dietary Preparation

Test diets were prepared by first mixing the chemical and an aliquot of powdered Wayne® Lab Blox animal feed (Allied Mills, Chicago, Illinois) with a mortar and pestle, placing this mixture in a Patterson-Kelly® twin-shell blender with the remainder of the feed, and mixing for 20 minutes. Test diets were sealed in labelled plastic bags and stored at 4°C for no longer than 14 days.

The stability of 2,6-toluenediamine dihydrochloride in feed was determined at Midwest Research Institute by assaying sample diet mixtures containing 10% 2,6-toluenediamine dihydrochloride that had been stored at -20°, 5°, 25°, or 45°C for 2 weeks. Amounts of the test chemical present were determined by gas chromatography as described in Appendix G. The compound was stable in feed for 2 weeks at temperatures as high as 45°C.

The amounts of 2,6-toluenediamine dihydrochloride in selected batches of feed were measured at 2- to 3-month intervals. In preliminary tests, 2-gram samples were extracted with 50 ml of 95% ethanol and analyzed by gas chromatography on a Tenax column at 235°C. Subsequent samples were extracted and analyzed as described in Appendix H.

The mean concentration of 10 samples, measured in duplicate and containing a theoretical level of 500 ppm, was 450±140 ppm. The mean concentration of 11 samples, measured in duplicate and containing a theoretical level of 250, was 240±50 ppm. Variation in the concentration of the compound among samples from the same mixture indicated some lack of homogeneity in the dosed-feed preparations.

C. Animals

Three-week old F344 rats and 4- to 5-week old B6C3F1 mice were obtained from the NCI Frederick Cancer Research Center, Frederick, Maryland. The animals were acclimated for 2 weeks and then assigned to control or dosed groups in such a manner that average cage weights were approximately equal for all animals of the same sex and species.

D. Animal Maintenance

Rats were housed four per cage in suspended polycarbonate cages (Lab Products, Inc., Garfield, N.J.) equipped with disposable nonwoven fiber filter sheets (Webrex). Mice were housed five per cage in polycarbonate shoe-box type cages covered with spumbonded Filtek filter bonnets (Lab Products). Aspen-bed® hardwood chips (American Excelsior, Summerville, Mass.) were used as bedding. Clean bedding and cages were provided twice weekly. Cage racks were changed every 2 weeks and disposable filters and filter bonnets were replaced at the same time. Filter bonnets were cleaned for reuse.

Water was available ad libitum for both species. Clean glass water bottles were provided twice weekly. Powdered Wayne® Lab Blox diet was available ad libitum in stainless-steel, gang-style hoppers (Scientific Cages, Inc., Bryan, Texas) which were changed once per week.

*

The temperature in the animal rooms was 19°-28°C and the relative humidity was 8%-74%. Incoming air was filtered through Tri-Dek 15/40 denier Dacron filters, with 10 to 12 air changes per hour. Fluorescent lighting was provided 12 hours per day.

Rats and mice were housed by species in rooms in which chronic tests were also being conducted on 4,4'-oxydianiline (CAS 101-80-4).

E. Range-Finding and 14-Day Repeated Dose Studies

Range-finding and 14-day repeated dose feed studies were conducted using F344 rats and B6C3F1 mice to determine the concentrations of 2,6-toluenediamine dihydrochloride to be used in the subchronic studies.

In the range-finding study, groups of two males and two females of each species were tested at each of five doses by administering a single dose of the test substance in corn oil by gavage (as shown in Table 1). All surviving animals were killed after 14 days. At the three highest doses, 2,6-toluenediamine dihydrochloride induced hemorrhage of the fundic portion of the stomach and the intestinal tract of rats. 2,6-Toluenediamine dihydrochloride was lethal for male rats at doses greater than 1 g/kg and at 0.3 g/kg for females. Deaths occurred in mice at all doses.

In the repeated dose study, groups of five males and five females of each species were tested for 2 weeks with five dose levels of the test substance in the feed. Groups of five males and five females of each species were maintained as untreated controls. All surviving animals were killed after 2 weeks. Survival of dosed groups at 2 weeks is shown in Table 2. No deaths occurred in rats at any of the doses tested. Weight gain was depressed 10% or more in both male and female rats at doses of 1,000 ppm or higher. Weight gain depression was 12.5% for male rats fed diets containing 1,000 ppm and 47% for male rats at 3,000 ppm. Weight gain depression was 52% for female rats fed diets containing 1,000 ppm and 88% at 3,000 ppm. All of the male mice and 3/5 female mice that received 3,000 ppm 2,6-toluenediamine dihydrochloride in their feed died. Hemorrhage of the stomach, digestive tract, and renal medullae were detected in the mice that died.

Table 1. Dosage and Survival of Rats and Mice Administered A Single Dose of 2,6-Toluenediamine Dihydrochloride by Gavage

Dose (g/kg)	Survival			
	Rats		Mice	
	Male	Female	Male	Female
0.1	2/2	2/2	1/2	1/2
0.3	2/2	2/2	0/2	0/2
1.0	2/2	0/2	0/2	1/2
3.0	0/2	0/2	0/2	0/2
10.0	0/2	0/2	0/2	0/2

Table 2. Dosage and Survival of Rats and Mice Administered Repeated Doses of 2,6-Toluenediamine Dihydrochloride in the Feed for 14 Days

Dose (ppm)	Survival			
	Rats		Mice	
	Male	Female	Male	Female
0	5/5	5/5	5/5	5/5
100	5/5	5/5	5/5	5/5
300	5/5	5/5	5/5	5/5
1,000	5/5	5/5	5/5	5/5
3,000	5/5	5/5	0/5	2/5

F. Subchronic Studies

Subchronic studies were conducted to determine the two concentrations (referred to in this report as "low" and "high" doses) to be used in the chronic studies. Diets containing 0, 100, 300, 1,000, 3,000, or 10,000 ppm 2,6-toluenediamine dihydrochloride were fed for 13 weeks to groups of 10 male and 10 female rats (Table 3), and groups of 10 male and 10 female mice received diets with 0, 10, 30, 100, 300, or 1,000 ppm (Table 4).

Clinical observations were made twice daily and animals were weighed weekly. At the end of the 91-day study, survivors were killed. Necropsies were performed on all animals and tissues were taken for histopathologic analysis.

Two out of 10 male rats and 7/10 female rats that received 10,000 ppm died before the end of the study. One of the males and four of the females died during week 1 of the study. No deaths occurred at any of the other dose levels.

Weight gains were depressed in male rats at all doses and in female rats at the three highest doses (1,000, 3,000, and 10,000 ppm).

Slight to moderate thyroid enlargement and darkening of the spleen, numerous lymph nodes, liver, kidney, adrenals, and nasal turbinates were observed in the eight surviving male rats that received the 10,000-ppm dose. Darkening of the same organs as in the males was observed in the three surviving female rats receiving the 10,000-ppm dose, but only one of these females had an enlarged thyroid. Among the rats examined, diffuse bilateral adenomatous hyperplasia of the thyroid with or without macroscopic enlargement was seen in 7/10 male rats fed 3,000 and 8/8 fed 10,000 ppm, and in 3/7 female rats fed 10,000 ppm. The only thyroid lesion observed in controls was seen in one male with papillary cyst hyperplasia. Bone marrow hyperplasia was seen in 8/8 male and 7/7 female rats at 10,000 ppm. Nephrosis was seen in 5/8 males and in 1/7 female rats fed the 10,000-ppm dose. Darkening of the nasal turbinates occurred in both male and female rats that received 2,000 or 3,000 ppm. No significant gross abnormalities were observed in the rats at the 300- and 1,000-ppm dose levels or in the controls.

Table 3. Dosage, Survival, and Mean Body Weights of Rats Fed 2,6-Toluenediamine Dihydrochloride in Diets for the 13-Week Subchronic Study

Dose (ppm)	Survival(a)	Mean Body Weights (grams)			Weight Change Relative to Controls(c) (Percent)
		Initial(b)	Final	Gain	
MALE					
0	10/10	110	298	188	
100	10/10	110	271	161	-14
300	10/10	110	269	159	-15
1,000	10/10	110	258	148	-19
3,000	10/10	110	229	119	-36
10,000	8/10	110	158	48	-74
FEMALE					
0	10/10	98	180	82	
100	10/10	98	177	79	-3
300	10/10	98	176	78	-5
1,000	10/10	98	156	58	-29
3,000	10/10	98	123	25	-70
10,000	3/10	98	103	5	-91

(a) Number surviving/number per group.

(b) The initial weights were obtained from graphs submitted by the test laboratory.

(c) Weight Change Relative to Controls =

$$\frac{\text{Weight Gain (Dosed Group)} - \text{Weight Gain (Control Group)}}{\text{Weight Gain (Control Group)}} \times 100$$

Table 4. Dosage, Survival, and Mean Body Weights of Mice Fed 2,6-Toluenediamine Dihydrochloride in Diets for the 13-Week Subchronic Study

Dose (ppm)	Survival(a)	Mean Body Weights(grams)			Weight Change Relative to Controls(c) (Percent)
		Initial(b)	Final	Gain	
<u>MALE</u>					
0	10/10	20.9	35.0	14.1	
100	10/10	20.9	34.9	14.0	0.7
300	10/10	20.9	33.0	12.1	-14
1,000	10/10	20.9	32.9	12.0	-15
3,000	10/10	20.9	31.5	10.6	-25
10,000	10/10	20.9	30.0	9.1	-35
<u>FEMALE</u>					
0	10/10	17.2	24.3	7.1	
100	10/10	17.2	24.2	7.0	-1.4
300	10/10	17.2	25.3	8.1	14
1,000	10/10	17.2	23.6	6.4	-10
3,000	10/10	17.2	23.6	6.4	-10
10,000	10/10	17.2	21.3	4.1	-42

(a) Number surviving/number per group.

(b) The initial weights were obtained from graphs submitted by the test laboratory.

(c) Weight Change Relative to Controls =

$$\frac{\text{Weight Gain (Dosed Group)} - \text{Weight Gain (Control Group)}}{\text{Weight Gain (Control Group)}} \times 100$$

No deaths occurred among the mice. Male mice receiving 300 ppm and females receiving a 1,000-ppm or higher dose had weight gain depressions of 10% or more.

Among the mice fed 1,000 ppm, squamous papilloma of the forestomach was detected in 1/10 males, and renal hyperpigmentation similar to that seen in the rats occurred in a second male and 2/10 females.

The low and high doses for the chronic studies were established at 250 and 500 ppm for the rats and at 50 and 100 ppm for the mice.

G. Chronic Studies

The test groups, doses administered, and durations of the chronic studies are shown in Table 5.

H. Clinical Examinations and Pathology

Animals were inspected twice daily. Body weights were recorded every 4 weeks. Animals that were moribund and those that survived to the termination of the study were killed and necropsied following anesthetization using CO₂ inhalation.

Gross and microscopic examinations were performed on major tissues, major organs, and all gross lesions from killed animals and from animals found dead. Tissues were preserved in 10% neutral buffered formalin, embedded in paraffin, sectioned, and stained with hematoxylin and eosin. The following tissues were examined microscopically: skin, lungs and bronchi, trachea, bone and bone marrow, spleen, lymph nodes, heart, salivary gland, liver, pancreas, stomach, small intestine, large intestine, kidneys, urinary bladder, pituitary, adrenal, thyroid, parathyroid, mammary gland, prostate and seminal vesicles or uterus, testis or ovary, brain, thymus, larynx, and esophagus.

Necropsies were performed on all animals found dead unless precluded in whole or in part by autolysis or cannibalization. Thus, the number of animals from which particular organs or tissues were examined microscopically varies and does not necessarily represent the number of animals that were placed on study in each group.

Table 5. Experimental Design of Chronic Feeding Studies with 2,6-Toluenediamine Dihydrochloride in Rats and Mice

Sex, Species and Test Group	Initial No. of Animals	2,6-Toluene- diamine 2(HCl) in Diet (a) (ppm)	Time on Study	
			Dosed (weeks)	Observed (weeks)
<u>MALE RATS</u>				
Matched-Control	50	0	0	104
Low-Dose	50	250	103	1
High-Dose	50	500	103	1
<u>FEMALE RATS</u>				
Matched-Control	50	0	0	104
Low-Dose	50	250	103	1
High-Dose	50	500	103	1
<u>MALE MICE</u>				
Matched-Control	50	0	0	104
Low-Dose	50	50	103	1
High-Dose	50	100	103	1
<u>FEMALE MICE</u>				
Matched-Control	50	0	0	104
Low-Dose	50	50	103	1
High-Dose	50	100	103	1

(a) Diets were available ad libitum.

I. Data Recording and Statistical Analyses

Data on this experiment were recorded in the Carcinogenesis Bioassay Data System (Linhart et al., 1974). The data elements include descriptive information on the chemicals, animals, experimental design, clinical observations, survival, body weight, and individual pathologic results, as recommended by the International Union Against Cancer (Berenblum, 1969).

Probabilities of survival were estimated by the product-limit procedure of Kaplan and Meier (1958) and are presented in this report in the form of graphs. Animals were statistically censored as of the time that they died of other than natural causes or were found to be missing; animals dying from natural causes were not statistically censored. Statistical analyses for a possible dose-related effect on survival used the method of Cox (1972) for testing two groups for equality and Tarone's (1975) extensions of Cox's methods for testing for a dose-related trend. One-tailed P values have been reported for all tests except the departure from linearity test, which is reported only when its two-tailed P value is less than 0.05.

The incidence of neoplastic or nonneoplastic lesions has been given as the ratio of the number of animals bearing such lesions at a specific anatomic site (numerator) to the number of animals in which that site is examined (denominator). In most instances, the denominators included only those animals for which that site was examined histologically. However, when macroscopic examination was required to detect lesions prior to histologic sampling (e.g., skin or mammary tumors), or when lesions could have appeared at multiple sites (e.g., lymphomas), the denominators consist of the numbers of animals necropsied.

The purpose of the statistical analyses of tumor incidence is to determine whether animals receiving the test chemical developed a significantly higher proportion of tumors than did the control animals. As a part of these analyses, the one-tailed Fisher exact test (Cox, 1970) was used to compare the tumor incidence of a control group with that of a group of dosed animals at each dose level. When results for two dosed groups are compared simultaneously with those for a control group, a correction to ensure an overall significance level of 0.05 is made. The Bonferroni inequality (Miller, 1966) requires that the P value for any comparison be less than or

equal to 0.025. When this correction was used, it is discussed in the narrative section. It is not, however, presented in the tables, where the Fisher exact P values are shown.

The Cochran-Armitage test for linear trend in proportions, with continuity correction (Armitage, 1971), was also used. Under the assumption of a linear trend, this test determines if the slope of the dose-response curve is different from zero at the one-tailed 0.05 level of significance. Unless otherwise noted, the direction of the significant trend is a positive dose relationship. This method also provides a two-tailed test of departure from linear trend.

The approximate 95% confidence interval for the relative risk of each dosed group compared with its control was calculated from the exact interval on the odds ratio (Gart, 1971). The lower and upper limits of the confidence interval of the relative risk have been included in the tables of statistical analyses. The interpretation of the limits is that, in approximately 95% of a large number of identical experiments, the true ratio of the risk in a dosed group of animals to that in a control group would be within the interval calculated from the experiment. When the lower limit of the confidence interval is greater than one, it can be inferred that a statistically significant result has occurred (P less than 0.025 one-tailed test when the control incidence is not zero, P less than 0.050 when the control incidence is zero). When the lower limit is less than unity but the upper limit is greater than unity, the lower limit indicates the absence of a significant result while the upper limit indicates that there is a theoretical possibility of the induction of tumors by the test chemical, which could not be detected under the conditions of this test.

III. RESULTS - RATS

A. Body Weights and Clinical Signs (Rats)

Mean body weight gain was depressed 17% in low-dose female rats and 27% in high-dose female rats when compared with controls. Weight gain depression was less than 10% for dosed male rats (Figure 1). No other clinical signs were observed that could be related to administration of the test chemical.

B. Survival (Rats)

Estimates of the probabilities of survival for male and female rats administered 2,6-toluenediamine dihydrochloride in the diet at the doses of this bioassay, together with those of the matched controls, are shown by the Kaplan and Meier curves in Figure 2. The result of the Tarone test for dose-related trend in the proportions surviving is not significant in either sex.

In male rats, 33/50 (66%) of the high-dose group, 34/50 (68%) of the low-dose group, and 25/50 (50%) of the matched-control group lived to the end of the bioassay. In females, 39/50 (78%) of the high-dose group, 35/50 (70%) of the low-dose group, and 35/50 (70%) of the matched-control group lived to the end of the study.

A sufficient number of rats of each sex were at risk for the development of late-appearing tumors.

C. Pathology (Rats)

Histopathologic findings on neoplasms in rats are summarized in Appendix A, Tables A1 and A2; findings on nonneoplastic lesions are summarized in Appendix C, Tables C1 and C2.

The various types of neoplasms occurring in both control and dosed rats did not appear to be compound related and were the types commonly found in aging F344 rats.





Figure 1. Growth Curves for Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet





Figure 2. Survival Curves for Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet

A variety of nonneoplastic lesions were observed in both control and dosed rats, none of which appeared to be compound related.

According to the histopathologic examination, 2,6-toluenediamine dihydrochloride was not carcinogenic to F344 rats under the conditions of this bioassay.

D. Statistical Analyses of Results (Rats)

Tables 6 and 7 contain the statistical analyses of the incidences of those primary tumors that occurred in at least two animals in one group and at an incidence of at least 5% in one or more groups.

In male rats, a significant positive dose-related trend is observed in the incidences of animals with neoplastic nodules or hepatocellular carcinomas ($P=0.037$). The historical incidence at this laboratory of untreated male rats with neoplastic nodules or hepatocellular adenomas or carcinomas is 2/334 (0.6%), which is lower than the 4/50 (8%) observed in the high dose on this study. The historical incidence across all laboratories of these tumors in male rats is 36/2,230 (1.6%). The 4/50 (8%) incidence in the high-dose when compared with 0/50 in the control groups by the Fisher exact test has a significance level of 0.059 which is above the 0.025 level required for statistical significance when the Bonferroni inequality is applied; therefore, the evidence associating the administration of the chemical and the development of these liver tumors is not statistically conclusive.

In male rats, a significant positive dose-related trend is observed in the incidences of animals with islet-cell adenomas of the pancreas ($P=0.025$). The historical incidence at this laboratory for this tumor in control male rats is 0/334. One group of vehicle control male rats had a 2/35 (5.7%) incidence of this tumor. The Fisher exact test comparing the high-dose incidence with that of the controls has a significance level of 0.058 which is above the $P=0.025$ level required for statistical significance in this study; therefore, the evidence associating the administration of the chemical and the development of these pancreatic tumors is not statistically conclusive.

Significant trends in the negative direction are observed in the incidences of C-cell tumors of the thyroid in male rats and of the fibroadenomas of the mammary gland in female rats.

In each of the 95% confidence intervals for relative risk shown in the tables, the value of one or less than one is included: this indicates the absence of significant positive results. It should also be noted that each of the intervals, except for incidences of fibroadenomas of the mammary gland in high-dose female rats and C-cell adenomas of the thyroid in dosed groups of male rats, has an upper limit greater than one indicating that 2,6-toluenediamine dihydrochloride might induce tumors that could not be detected under the conditions of this test.

Table 6. Analyses of the Incidence of Primary Tumors in Male Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)

Topography: Morphology	Matched Dose	Low Dose	High Dose
Lung: Alveolar/Bronchiolar Adenoma (b)	3/49 (6)	1/50 (2)	2/50 (4)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		0.327	0.653
Lower Limit		0.006	0.057
Upper Limit		3.903	5.457
Weeks to First Observed Tumor	98	105	94
Hematopoietic System: All Lymphomas (b)	2/50 (4)	2/50 (4)	4/50 (8)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.000	2.000
Lower Limit		0.075	0.301
Upper Limit		13.326	21.316
Weeks to First Observed Tumor	85	95	100
Hematopoietic System: All Leukemias (b)	9/50 (18)	10/50 (20)	7/50 (14)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.111	0.778
Lower Limit		0.445	0.267
Upper Limit		2.823	2.159
Weeks to First Observed Tumor	84	83	97
Hematopoietic System: Lymphoma or Leukemia (b)	11/50 (22)	12/50 (24)	11/50 (22)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.091	1.000
Lower Limit		0.488	0.434
Upper Limit		2.465	2.303
Weeks to First Observed Tumor	84	83	97

Table 6. Analyses of the Incidence of Primary Tumors in Male Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet (a) (continued)

Topography: Morphology	Matched Control	Low Dose	High Dose
Hematopoietic System:			
Neoplasm, NOS (b)	2/50 (4)	6/50 (12)	1/50 (2)
P Values (c,d)	N.S.	N.S.	N.S.
Departure from Linear Trend (f)	P=0.029		
Relative Risk (e)		3.000	0.500
Lower Limit		0.569	0.009
Upper Limit		29.254	9.290
Weeks to First Observed Tumor	88	76	99
Liver: Neoplastic Nodule (b)			
	0/50 (0)	0/50 (0)	3/50 (6)
P Values (c,d)	P=0.037	N.S.	N.S.
Relative Risk (e)		--	Infinite
Lower Limit		--	0.601
Upper Limit		--	Infinite
Weeks to First Observed Tumor	--	--	104
Liver: Neoplastic Nodule or Hepatocellular Carcinoma (b)			
	0/50 (0)	2/50 (4)	4/50 (8)
P Values (c,d)	P=0.037	N.S.	N.S.
Relative Risk (e)		Infinite	Infinite
Lower Limit		0.296	0.927
Upper Limit		Infinite	Infinite
Weeks to First Observed Tumor	--	102	97
Pituitary: Adenoma, NOS (b)			
	10/45 (22)	11/49 (22)	15/44 (34)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.010	1.534
Lower Limit		0.433	0.727
Upper Limit		2.397	3.378
Weeks to First Observed Tumor	84	100	88

Table 6. Analyses of the Incidence of Primary Tumors in Male Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)
(continued)

Topography: Morphology	Matched Control	Low Dose	High Dose
Adrenal: Pheochromocytoma (b)	10/50 (20)	6/50 (12)	8/48 (17)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		0.600	0.833
Lower Limit		0.194	0.312
Upper Limit		1.676	2.140
Weeks to First Observed Tumor	96	97	75
Thyroid: C-cell Adenoma (b)	5/44 (11)	0/48 (0)	0/47 (0)
P Values (c,d)	P=0.005(N)	P=0.022(N)	P=0.023(N)
Relative Risk (e)		0.000	0.000
Lower Limit		0.000	0.000
Upper Limit		0.725	0.740
Weeks to First Observed Tumor	98	--	--
Thyroid: C-cell Carcinoma (b)	2/44 (5)	2/48 (4)	2/47 (4)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		0.917	0.936
Lower Limit		0.069	0.071
Upper Limit		12.189	12.441
Weeks to First Observed Tumor	101	105	105
Thyroid: C-cell Adenoma or Carcinoma(b)	7/44 (16)	2/48 (4)	2/47 (4)
P Values (c,d)	P=0.033 (N)	N.S.	N.S.
Relative Risk (e)		0.262	0.267
Lower Limit		0.028	0.028
Upper Limit		1.288	1.314
Weeks to First Observed Tumor	98	105	105

Table 6. Analyses of the Incidence of Primary Tumors in Male Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)
(continued)

Topography: Morphology	Matched Control	Low Dose	High Dose
Pancreatic Islets: Islet-cell Adenoma (b)	0/45 (0)	1/46 (2)	4/45 (9)
P Values (c,d)	P=0.025	N.S.	N.S.
Relative Risk (e)		Infinite	Infinite
Lower Limit		0.053	0.930
Upper Limit		Infinite	Infinite
Weeks to First Observed Tumor	—	105	94
Preputial Gland: Adenoma, NOS (b)	2/50 (4)	3/50 (6)	4/50 (8)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.500	2.000
Lower Limit		0.180	0.301
Upper Limit		17.329	21.316
Weeks to First Observed Tumor	107	105	88
Testis: Interstitial-cell Tumor (b)	42/49 (86)	47/49 (96)	44/49 (90)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.119	1.048
Lower Limit		0.966	0.890
Upper Limit		1.210	1.209
Weeks to First Observed Tumor	63	76	78

Table 6. Analyses of the Incidence of Primary Tumors in Male Rats
Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)
(continued)

- (a) Dosed groups received doses of 250 or 500 ppm.
- (b) Number of tumor-bearing animals/number of animals examined at site (percent).
- (c) Beneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when P is less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P is less than 0.05; otherwise, not significant (N.S.) is indicated.
- (d) A negative trend (N) indicates a lower incidence in a dose group than in a control group.
- (e) The 95% confidence interval of the relative risk between each dosed group and the control group.
- (f) The probability level for departure from linear trend is given when P is less than 0.05 for any comparison.

Table 7. Analyses of the Incidence of Primary Tumors in Female Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)

Topography: Morphology	Matched Control	Low Dose	High Dose
Hematopoietic System: All Leukemias (b)	4/50 (8)	4/50 (8)	5/50 (10)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.000	1.250
Lower Limit		0.197	0.286
Upper Limit		5.083	5.954
Weeks to First Observed Tumor	26	97	104
Hematopoietic System: Leukemia or Lymphoma (b)	5/50 (10)	5/50 (10)	5/50 (10)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.000	1.000
Lower Limit		0.245	0.245
Upper Limit		4.082	4.082
Weeks to First Observed Tumor	26	97	104
Pituitary: Adenoma, NOS (b)	19/48 (40)	23/43 (53)	25/48 (52)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.351	1.316
Lower Limit		0.827	0.814
Upper Limit		2.195	2.139
Weeks to First Observed Tumor	95	74	70
Thyroid: C-cell Adenoma or Carcinoma (b)	3/49 (6)	3/49 (6)	1/48 (2)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.000	0.340
Lower Limit		0.140	0.007
Upper Limit		7.126	4.060
Weeks to First Observed Tumor	62	106	105

Table 7. Analyses of the Incidence of Primary Tumors in Female Rats Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)
(continued)

Topography: Morphology	Matched Control	Low Dose	High Dose
Mammary Gland: Fibroadenoma (b)	11/50 (22)	6/50 (12)	2/50 (4)
P Values (c,d)	P=0.006 (N)	N.S.	P = 0.007 (N)
Relative Risk (e)		0.545	0.182
Lower Limit		0.179	0.020
Upper Limit		1.477	0.777
Weeks to First Observed Tumor	96	105	98
Uterus: Endometrial Stromal Polyp (b)	9/48 (19)	10/49 (20)	11/50 (22)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.088	1.173
Lower Limit		0.437	0.487
Upper Limit		2.758	2.916
Weeks to First Observed Tumor	62	80	92

(a) Dosed groups received doses of 250 or 500 ppm.

(b) Number of tumor-bearing animals/number of animals examined at site (percent).

(c) Beneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when P is less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P is less than 0.05; otherwise, not significant (N.S.) is indicated.

(d) A negative trend (N) indicates a lower incidence in a dose group than in a control group.

(e) The 95% confidence interval of the relative risk between each dosed group and the control group.

IV. RESULTS - MICE

A. Body Weights and Clinical Signs (Mice)

Weight gain depression when compared with controls was less than 10% for dosed male and female mice (Figure 3). No other clinical signs were observed that could be related to the administration of the test chemical.

B. Survival (Mice)

Estimates of the probabilities of survival for male and female mice administered 2,6-toluenediamine dihydrochloride in the diet at the doses of this bioassay, together with those of the matched controls, are shown by the Kaplan and Meier curves in Figure 4. The result of the Tarone test for dose-related trend in the proportions surviving is not significant in either sex.

In male mice, 32/50 (64%) of the high-dose group, 31/50 (62%) of the low-dose group, and 31/50 (62%) of the matched-control group lived to the end of the bioassay. In females, 39/50 (78%) of the high-dose group, 40/50 (80%) of the low-dose group, and 39/50 (78%) of the matched-control group lived to the end of the study.

A sufficient number of mice of each sex were at risk for the development of late-appearing tumors.

C. Pathology (Mice)

Histopathologic findings on neoplasms in mice are summarized in Appendix B, Tables B1 and B2; findings on nonneoplastic lesions are summarized in Appendix D, Tables D1 and D2.

Neoplasms occurred in both matched-control and dosed mice. A slight increase over that in controls occurred in the incidence of vascular neoplasms (hemangioma/hemangiosarcoma) of the spleen and liver in dosed male mice. Vascular neoplasms were noted in 1/50 control male mice, 5/50 (10%) male mice of the low-dose group, and 3/49 (6%) of the high-dose group.




Figure 3. Growth Curves for Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet




Figure 4. Survival Curves for Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet

Hemangiomas were small lesions with distended capillaries or sinusoids and anastomosing vascular cysts lined by transformed endothelial cells which were fusiform. Cytoplasm of the cells was eosinophilic and nuclei were hyperchromatic. A marked variation in the size of cavernous vascular spaces was characteristic of hemangiosarcoma. The cells were pleomorphic and crowded in areas. Nuclei were large and vesicular. Mitotic figures were numerous. In the smaller tumors in the liver, some of the hepatocytes encircled by the transformed endothelial cells were hypertrophic and hyperplastic. Hepatocytes were obliterated in the large tumors. In two male mice of the low-dose group, hemangiosarcoma was found in association with hepatocellular carcinoma.

A variety of nonneoplastic lesions were seen in matched-control and dosed mice. None of these lesions appeared to be related to chemical administration.

The histopathologic examination provided no evidence that 2,6-toluenediamine dihydrochloride was carcinogenic to B6C3F1 mice, under the conditions of this bioassay.

D. Statistical Analyses of Results (Mice)

Tables 8 and 9 contain the statistical analyses of the incidences of those primary tumors that occurred in at least two animals in one group and at an incidence of at least 5% in one or more groups.

In male mice, the Cochran-Armitage test indicates a departure from linear trend in the incidence of lymphomas in animals due to the higher incidence in the low-dose group (8/50, 16%) when compared with the high-dose group (2/50, 4%). The result of the Fisher exact test indicates an increased incidence in the low-dose group, but the $P=0.046$ observed in this group is above the level required for significance when the Bonferroni inequality criterion is used in the comparison of two dosed groups with a single control group ($P=0.025$). In females, a significant ($P=0.036$) positive linear trend was observed in the incidence of animals with hepatocellular carcinomas of the liver, but the results of the Fisher exact test are not significant.

In each of the 95% confidence intervals for relative risk shown in the tables, one is included: this indicates the absence of significant positive results. It should also be noted that each of the intervals has an upper limit greater than one indicating the theoretical possibility of tumor induction by 2,6-toluenediamine dihydrochloride, which could not be detected under the conditions of this test.

Table 8. Analyses of the Incidence of Primary Tumors in Male Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)

Topography: Morphology	Matched Control	Low Dose	High Dose
Lung: Alveolar/Bronchiolar Carcinoma (b)	4/50 (8)	5/50 (10)	1/50 (2)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.250	0.250
Lower Limit		0.286	0.005
Upper Limit		5.954	2.411
Weeks to First Observed Tumor	75	105	104
Lung: Alveolar/Bronchiolar Adenoma (b)	7/50 (14)	8/50 (16)	6/50 (12)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.143	0.857
Lower Limit		0.392	0.256
Upper Limit		3.423	2.766
Weeks to First Observed Tumor	101	69	104
Lung: Alveolar/Bronchiolar Carcinoma or Adenoma (b)	11/50 (22)	13/50 (26)	7/50 (14)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.182	0.636
Lower Limit		0.542	0.228
Upper Limit		2.626	1.645
Weeks to First Observed Tumor	75	69	104
Hematopoietic System: All Lymphomas (b)	2/50 (4)	8/50 (16)	2/50 (4)
P Values (c,d)	N.S.	P=0.046	N.S.
Departure from Linear Trend (f)	P=0.011		
Relative Risk (e)		4.000	1.000
Lower Limit		0.851	0.075
Upper Limit		37.147	13.326
Weeks to First Observed Tumor	94	90	91

Table 8. Analyses of the Incidence of Primary Tumors in Male Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)
(continued)

Topography: Morphology	Matched Control	Low Dose	High Dose
Circulatory System: Hemangioma, Hemangiosarcoma, or Angiosarcoma (b)	1/50 (2)	5/50 (10)	3/50 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		4.000	3.000
Lower Limit		0.415	0.251
Upper Limit		192.805	154.270
Weeks to First Observed Tumor	83	96	56
Liver: Hepatocellular Carcinoma (b)	14/50 (28)	12/50 (24)	13/49 (27)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		0.857	0.948
Lower Limit		0.404	0.459
Upper Limit		1.790	1.940
Weeks to First Observed Tumor	67	64	90
Liver: Hepatocellular Adenoma (b)	7/50 (14)	5/50 (10)	5/49 (10)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		0.714	0.729
Lower Limit		0.191	0.195
Upper Limit		2.434	2.481
Weeks to First Observed Tumor	63	105	104
Liver: Hepatocellular Carcinoma or Adenoma (b)	21/50 (42)	17/50 (34)	18/49 (37)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		0.810	0.875
Lower Limit		0.462	0.507
Upper Limit		1.405	1.496
Weeks to First Observed Tumor	63	64	90

Table 8. Analyses of the Incidence of Primary Tumors in Male Mice
Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)
(continued)

- (a) Dosed groups received doses of 50 or 100 ppm.
- (b) Number of tumor-bearing animals/number of animals examined at site (percent).
- (c) Beneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when P is less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P is less than 0.05; otherwise, not significant (N.S.) is indicated.
- (d) A negative trend (N) indicates a lower incidence in a dose group than in a control group.
- (e) The 95% confidence interval of the relative risk between each dosed group and the control group.
- (f) The probability level for departure from linear trend is given when P is less than 0.05 for any comparison.

Table 9. Analyses of the Incidence of Primary Tumors in Female Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)

Topography: Morphology	Matched Control	Low Dose	High Dose
Lung: Alveolar/Bronchiolar Adenoma (b)	4/50 (8)	7/50 (14)	3/49 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.750	0.765
Lower Limit		0.476	0.118
Upper Limit		7.682	4.288
Weeks to First Observed Tumor	92	105	105
Lung: Alveolar/Bronchiolar Adenoma or Carcinoma (b)	4/50 (8)	8/50 (16)	3/49 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		2.000	0.765
Lower Limit		0.576	0.118
Upper Limit		8.539	4.288
Weeks to First Observed Tumor	92	105	105
Hematopoietic System: All Lymphomas (b)	4/50 (8)	10/50 (20)	6/50 (12)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		2.500	1.500
Lower Limit		0.779	0.380
Upper Limit		10.246	6.820
Weeks to First Observed Tumor	91	89	88
Hematopoietic System: Neoplasm, NOS (b)	4/50 (8)	1/50 (2)	1/50 (2)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		0.250	0.250
Lower Limit		0.005	0.005
Upper Limit		2.411	2.411
Weeks to First Observed Tumor	84	90	85

Table 9. Analyses of the Incidence of Primary Tumors in Female Mice Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)
(continued)

Topography: Morphology	Matched Control	Low Dose	High Dose
Liver: Hepatocellular Carcinoma (b)	0/50 (0)	0/49 (0)	3/49 (6)
P Values (c,d)	P=0.036	--	N.S.
Relative Risk (e)		--	Infinite
Lower Limit		--	0.614
Upper Limit		--	Infinite
Weeks to First Observed Tumor	--	--	93
Liver: Hepatocellular Adenoma (b)	4/50 (8)	3/49 (6)	4/49 (8)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		0.765	1.020
Lower Limit		0.118	0.201
Upper Limit		4.288	5.183
Weeks to First Observed Tumor	106	105	105
Liver: Hepatocellular Carcinoma or Adenoma (b)	4/50 (8)	3/49 (6)	7/49 (14)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		0.765	1.786
Lower Limit		0.118	0.486
Upper Limit		4.288	7.830
Weeks to First Observed Tumor	106	105	93
Pituitary: Adenoma, NOS (b)	3/37 (8)	5/38 (13)	6/38 (16)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (e)		1.623	1.947
Lower Limit		0.342	0.452
Upper Limit		9.783	11.243
Weeks to First Observed Tumor	106	105	105

Table 9. Analyses of the Incidence of Primary Tumors in Female Mice
Administered 2,6-Toluenediamine dihydrochloride in the Diet (a)
(continued)

- (a) Dosed groups received doses of 50 or 100 ppm.
- (b) Number of tumor-bearing animals/number of animals examined at site (percent).
- (c) Beneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when P is less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P is less than 0.05; otherwise, not significant (N.S.) is indicated.
- (d) A negative trend (N) indicates a lower incidence in a dose group than in a control group.
- (e) The 95% confidence interval of the relative risk between each dosed group and the control group.

V. DISCUSSION

Mean body weight gain was depressed 17% in low-dose female rats and 27% in high-dose female rats. Weight gain depression was less than 10% for dosed groups of male rats and of male and female mice when compared with the controls.

The slight depression in weight gain among male and female mice receiving the test chemical as well as the lack of other observable clinical signs suggest that the maximum tolerated dose may not have been given to the mice.

In dosed male and female rats, no tumors occurred in statistically significant numbers when compared with those in control rats.

Diffuse bilateral adenomatous hyperplasia and enlargement of the thyroid occurred in the subchronic studies in male rats administered doses 6- to 20-fold higher than those used in the chronic studies. In contrast, in the chronic studies significant results in the negative direction were observed in the incidence of C-cell tumors of the thyroid in male rats.

In male rats, neoplastic nodules or hepatocellular carcinomas occurred with a dose-related trend that was significant ($P=0.037$), but the result of the Fisher exact test was not significant. Islet-cell adenomas of the pancreas occurred with a dose-related trend that was significant ($P=0.025$), but the result of the Fisher exact test was not significant.

In female rats, significant results in the negative direction were observed in the incidences of fibroadenoma of the mammary gland.

In male mice, lymphomas occurred in the low-dose group at an incidence that was significantly higher ($P=0.046$) than that of the corresponding matched-control group; however, this level of significance is below the level required when the Bonferroni criterion is used for multiple comparison.

In female mice, hepatocellular carcinomas occurred with a dose-related trend that was significant ($P=0.036$), but the result of the Fisher exact test comparing the incidence in the high-dose group with that in the matched controls was not significant.

In a previous study conducted independently but under the protocols of the Carcinogenesis Testing Program (NCI, 1979), the structurally related 2,4-isomer of toluenediamine was found to be carcinogenic for F344 rats, inducing

hepatocellular or neoplastic nodules in both males and females and carcinomas or adenomas of the mammary gland in females. The 2,4-isomer was also carcinogenic for B6C3F1 mice, inducing hepatocellular carcinomas. In those studies, rats received approximately 33% of the dose that animals received daily in the present studies, and the period of compound administration was 40 to 60 weeks shorter. The mice in the 2,4-toluenediamine study received a dose 2-fold higher than mice in the present study.

Primary occupational exposure is to the 80/20 or 65/35 mixtures of 2,4-toluenediamine and 2,6-toluenediamine, rather than to the 2,6-toluenediamine alone (Buist, 1970). The 2,4-isomer has been shown to be carcinogenic (NCI, 1979; Ito et al., 1969), but in the present study the 2,6-isomer has not been demonstrated to be carcinogenic.

Some aromatic amines have previously been found to be carcinogenic while others have been found to be without carcinogenic effects; for example, m-phenylenediamine has not yet been found to be carcinogenic in rats, but the addition of an ortho methyl group as in 2,4-toluenediamine enhances carcinogenic effects; 2,4,6-trimethylaniline is clearly carcinogenic for rats, but 2,4,5-trimethylaniline has an equivocal effect in rats; o-toluidine is carcinogenic, yet addition of a second amine substituent ortho to the methyl group as in 2,6-toluenediamine eliminates carcinogenic effects (Weisburger et al., 1978). Thus, it appears that a monocyclic aromatic compound that has two methyl groups ortho to an amine substituent is carcinogenic, whereas a monocyclic aromatic compound that has two amine groups ortho to a methyl group is not. While the role of steric effects in this group of compounds is not yet clear, the carcinogenicity of toluediamines appears to be determined by the exact ring positions of the methyl and amino groups.

VI. CONCLUSIONS

Under conditions of this bioassay, 2,6-toluenediamine dihydrochloride was not carcinogenic for either F344 rats or for B6C3F1 mice.

VII. BIBLIOGRAPHY

- Armitage, P., Statistical Methods in Medical Research, John Wiley & Sons, Inc., New York, 1971, pp. 362-365.
- Berenblum, I., ed., Carcinogenicity Testing: A Report of the Panel of Carcinogenicity of the Cancer Research Commission of UICC, Vol. 2, International Union Against Cancer, Geneva, 1969.
- Buist, J.M., Isocyanates in industry. Proc. R. Soc. Med. 63:365-366, 1970.
- Cox, D. R., Analysis of Binary Data, Methuen & Co., Ltd., London, 1970, pp. 48-52.
- Cox, D. R., Regression models and life tables. J. R. Statist. Soc. B34:187-220, 1972.
- Dybing, E. and Thorgeirsson, S.S., Metabolic activation of 2,4-diaminoanisole, a hair-dye component--I, Biochem. Pharmacol. 26:729-734, 1977.
- Finkleshtein, A. I. and Boitsov, E. N, Zavod. Lab 26:959, 1960.
- Gart, J. J., The comparison of proportions: a review of significance tests, confidence limits and adjustments for stratification. Rev. Int. Stat. Inst. 39:148-169, 1971.
- Ito, N., Hiasa, Y., Konishi, Y., and Marugami, M., The development of carcinoma in liver of rats treated with m-toluylenediamine and the synergistic and antagonistic effects with other chemicals. Cancer Res. 29:1137-1145, 1969.
- Kaplan, E. L. and Meier, P., Nonparametric estimation from incomplete observations. J. Amer. Statist. Assoc. 53:457-481, 1958.
- Layer, R. W., Amines, aromatic (phenylenediamines). In: Kirk-Othmer Encyclopedia of Chemical Technology, Vol. 2, Interscience Publisher, New York, 1964, pp. 348-354.
- Linhart, M. S., Cooper, J. A., Martin, R. L., Page, N. P., and Peters, J. A., Carcinogenesis bioassay data system. Comput. and Biomed. Res. 7:230-248, 1978.
- Lowe, A., The chemistry of isocyanates. Proc. R. Soc. Med. 63:367- 368, 1970.
- Miller, R. G., Jr., Simultaneous Statistical Inference, McGraw-Hill Book Co., New York, 1966, pp. 6-10.

Milligan, B., Amines, aromatic (diaminotoluenes). In: Kirk-Othmer Encyclopedia of Chemical Technology, Vol. 2., Interscience Publisher, New York, 1968, pp. 321-329.

NCI, National Cancer Institute, Bioassay of 2,4-Diaminotoluene for Possible Carcinogenicity, TR 162, U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, Bethesda, Md., 1979.

Sadtler Standard Spectra, Sadtler Research Laboratories, Philadelphia, Pa., NMR No. 12852.

Society of Dyers and Colourists, Colour Index, Vol. 4, The Society of Dyers and Colourists, Bradford, England, 1971, p. 4850.

Tarone, R. E., Tests for trend in life table analysis. Biometrika 62:679-682, 1975.

United States International Trade Commission, Synthetic Organic Chemicals - United States Production and Sales, 1977, USITC Publication 920, U. S. Government Printing Office, Washington, D.C., 1978.

Ward, J. M., Goodman, D. G., Griesemer, R. A., Hardisty, J. F., Schueler, R. L., Squire, R. A., and Strandberg, J. D., Quality assurance for pathology in rodent carcinogenesis tests. J. Environ. Pathol. Toxicol. 2:371-378, 1978.

Weisburger, E. K., Russfield, A. B., Weisburger, J. H., Boger, E., Van Dongen, C. G., and Chu, K. C., Testing of twenty-one environmental aromatic amines or derivatives for long-term toxicity or carcinogenicity. J. Environ. Pathol. Toxicol. 2:325-356, 1978.

APPENDIX A

Summary of the Incidence of Neoplasms in Rats
Administered 2,6-Toluenediamine dihydrochloride
in the Diet

|

|

TABLE A1.

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE RATS
ADMINISTERED 2,6-TOLUENEDIAMINE DIHYDROCHLORIDE IN THE DIET

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	50	50	50
INTEGUMENTARY SYSTEM			
*SKIN	(50)	(50)	(50)
BASAL-CELL CARCINOMA			1 (2%)
*SUBCUT TISSUE	(50)	(50)	(50)
CARCINOMA, NOS	1 (2%)		
SQUAMOUS CELL CARCINOMA	1 (2%)		
SARCOMA, NOS		2 (4%)	
FIBROMA		2 (4%)	
NEURILEMOMA, MALIGNANT			1 (2%)
RESPIRATORY SYSTEM			
#LUNG	(49)	(50)	(50)
ALVEOLAR/BRONCHIOLAR ADENOMA	3 (6%)	1 (2%)	2 (4%)
SARCOMA, NOS, METASTATIC		1 (2%)	
NEURILEMOMA, METASTATIC			1 (2%)
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(50)	(50)	(50)
MALIGNANT LYMPHOMA, NOS	2 (4%)	2 (4%)	4 (8%)
LEUKEMIA, NOS	9 (18%)	10 (20%)	7 (14%)
*HEMATOPOIETIC SYSTEM	(50)	(50)	(50)
NEOPLASM, NOS	2 (4%)	6 (12%)	1 (2%)
#SPLEEN	(50)	(50)	(49)
SARCOMA, NOS			1 (2%)
CIRCULATORY SYSTEM			
NONE			

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE A1. MALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
DIGESTIVE SYSTEM			
*LIP SQUAMOUS CELL CARCINOMA	(50)	(50) 1 (2%)	(50)
#LIVER NEOPLASTIC NODULE HEPATOCELLULAR CARCINOMA	(50)	(50) 2 (4%)	(50) 3 (6%) 1 (2%)
URINARY SYSTEM			
#KIDNEY TUBULAR-CELL ADENOCARCINOMA LEIOMYOSARCOMA	(50) 1 (2%)	(50) 1 (2%)	(50)
#KIDNEY/PELVIS TRANSITIONAL-CELL PAPILLOMA	(50)	(50)	(50) 1 (2%)
ENDOCRINE SYSTEM			
#PITUITARY NEOPLASM, NOS ADENOMA, NOS	(45) 1 (2%) 10 (22%)	(49) 11 (22%)	(44) 15 (34%)
#ADRENAL CORTICAL ADENOMA PHEOCHROMOCYTOMA	(50) 10 (20%)	(50) 6 (12%)	(48) 1 (2%) 8 (17%)
#THYROID FOLLICULAR-CELL CARCINOMA C-CELL ADENOMA C-CELL CARCINOMA	(44) 5 (11%) 2 (5%)	(48) 2 (4%)	(47) 1 (2%) 2 (4%)
#PANCREATIC ISLETS ISLET-CELL ADENOMA	(45)	(46) 1 (2%)	(45) 4 (9%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND ADENOCARCINOMA, NOS	(50) 1 (2%)	(50)	(50)
*PREPUTIAL GLAND ADENOMA, NOS	(50) 2 (4%)	(50) 3 (6%)	(50) 4 (8%)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE A1. MALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
#TESTIS INTERSTITIAL-CELL TUMOR	(49) 42 (86%)	(49) 47 (96%)	(49) 44 (90%)
NERVOUS SYSTEM			
#BRAIN ASTROCYTOMA	(50)	(50) 1 (2%)	(50)
SPECIAL SENSE ORGANS			
*ZYMBA'S GLAND CARCINOMA, NOS	(50)	(50) 1 (2%)	(50) 1 (2%)
MUSCULOSKELETAL SYSTEM			
*SKULL OSTEOSARCOMA	(50)	(50)	(50) 1 (2%)
BODY CAVITIES			
*TUNICA VAGINALIS MESOTHELIOMA, NOS	(50) 2 (4%)	(50) 1 (2%)	(50)
ALL OTHER SYSTEMS			
SITE UNKNOWN CARCINOMA, NOS	1		
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	50	50	50
NATURAL DEATH ^a	17	12	11
MORIBUND SACRIFICE	8	4	6
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	25	34	33
ANIMAL MISSING			

^a INCLUDES AUTOLYZED ANIMALS

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE A1. MALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	47	49	49
TOTAL PRIMARY TUMORS	95	100	104
TOTAL ANIMALS WITH BENIGN TUMORS	46	48	49
TOTAL BENIGN TUMORS	72	71	80
TOTAL ANIMALS WITH MALIGNANT TUMORS	16	20	16
TOTAL MALIGNANT TUMORS	18	22	20
TOTAL ANIMALS WITH SECONDARY TUMORS#		1	1
TOTAL SECONDARY TUMORS		1	1
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT	5	7	4
TOTAL UNCERTAIN TUMORS	5	7	4
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE A2.

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE RATS
ADMINISTERED 2,6-TOLUENEDIAMINE DIHYDROCHLORIDE IN THE DIET

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	50	50	50
INTEGUMENTARY SYSTEM			
*SKIN	(50)	(50)	(50)
BASAL-CELL CARCINOMA		1 (2%)	
*SUBCUT TISSUE	(50)	(50)	(50)
NEOPLASM, NOS			1 (2%)
SQUAMOUS CELL CARCINOMA	1 (2%)		1 (2%)
SARCOMA, NOS		2 (4%)	
FIBROMA	1 (2%)		
FIBROUS HISTIOCYTOMA			1 (2%)
LEIOMYOSARCOMA			1 (2%)
RESPIRATORY SYSTEM			
#LUNG	(50)	(50)	(50)
NEOPLASM, NOS, METASTATIC		1 (2%)	
UNDIFFERENTIATED CARCINOMA			1 (2%)
SQUAMOUS CELL CARCINOMA, METASTA	1 (2%)		
ALVEOLAR/BRONCHIOLAR ADENOMA	1 (2%)	2 (4%)	1 (2%)
C-CELL CARCINOMA, METASTATIC	1 (2%)	1 (2%)	
SARCOMA, NOS, METASTATIC		1 (2%)	
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(50)	(50)	(50)
MALIGNANT LYMPHOMA, NOS		1 (2%)	
LEUKEMIA, NOS	4 (8%)	4 (8%)	5 (10%)
*HEMATOPOIETIC SYSTEM	(50)	(50)	(50)
NEOPLASM, NOS	1 (2%)	2 (4%)	
#SPLEEN	(49)	(50)	(50)
FIBROSARCOMA, INVASIVE	1 (2%)		

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE A2. FEMALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
MALIG. LYMPHOMA, HISTIOCYTIC TYPE	1 (2%)		
#LYMPH NODE UNDIFFERENTIATED CARCINOMA METAS C-CELL CARCINOMA, METASTATIC	(47) 1 (2%)	(44)	(47) 1 (2%)
#THYMUS THYMOMA	(35)	(30)	(32) 1 (3%)
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
#LIVER NEOPLASTIC NODULE HEPATOCELLULAR CARCINOMA ENDOMETRIAL STROMAL SARCOMA, MET	(50)	(50)	(50) 1 (2%) 1 (2%)
#STOMACH FIBROSARCOMA	(50) 1 (2%)	(48)	(50)
#JEJUNUM SARCOMA, NOS	(49)	(48) 1 (2%)	(48)
URINARY SYSTEM			
#KIDNEY UNDIFFERENTIATED CARCINOMA METAS	(50)	(50)	(50) 1 (2%)
ENDOCRINE SYSTEM			
#PITUITARY ADENOMA, NOS	(48) 19 (40%)	(43) 23 (53%)	(48) 25 (52%)
#ADRENAL PHEOCHROMOCYTOMA	(50) 2 (4%)	(50) 2 (4%)	(50) 1 (2%)
#THYROID C-CELL ADENOMA	(49) 2 (4%)	(49) 1 (2%)	(48) 1 (2%)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE A2. FEMALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
C-CELL CARCINOMA	1 (2%)	2 (4%)	
#PANCREATIC ISLETS	(48)	(47)	(46)
ISLET-CELL ADENOMA	1 (2%)		
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(50)	(50)	(50)
ADENOMA, NOS	1 (2%)	1 (2%)	2 (4%)
FIBROADENOMA	11 (22%)	6 (12%)	2 (4%)
*CLITORAL GLAND	(50)	(50)	(50)
ADENOMA, NOS	1 (2%)		1 (2%)
#UTERUS	(48)	(49)	(50)
ENDOMETRIAL STROMAL POLYP	9 (19%)	10 (20%)	11 (22%)
ENDOMETRIAL STROMAL SARCOMA	1 (2%)	1 (2%)	
#OVARY	(47)	(47)	(50)
UNDIFFERENTIATED CARCINOMA METAS			1 (2%)
NERVOUS SYSTEM			
#BRAIN	(49)	(49)	(50)
NEOPLASM, NOS		1 (2%)	
ASTROCYTOMA	1 (2%)	1 (2%)	
SPECIAL SENSE ORGANS			
*ZYMBAL'S GLAND	(50)	(50)	(50)
CARCINOMA, NOS	1 (2%)		
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*ABDOMINAL CAVITY	(50)	(50)	(50)
MESOTHELIOOMA, NOS		1 (2%)	
ALL OTHER SYSTEMS			
NONE			

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
 * NUMBER OF ANIMALS NECROPSIED

TABLE A2. FEMALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	50	50	50
NATURAL DEATHS	9	10	7
MORIBUND SACRIFICE	6	5	4
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	35	35	39
ANIMAL MISSING			
a INCLUDES AUTOLYZED ANIMALS			
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	42	37	39
TOTAL PRIMARY TUMORS	60	62	57
TOTAL ANIMALS WITH BENIGN TUMORS	34	31	33
TOTAL BENIGN TUMORS	48	45	46
TOTAL ANIMALS WITH MALIGNANT TUMORS	11	12	9
TOTAL MALIGNANT TUMORS	11	13	9
TOTAL ANIMALS WITH SECONDARY TUMORS#	3	4	1
TOTAL SECONDARY TUMORS	4	4	3
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT	1	3	2
TOTAL UNCERTAIN TUMORS	1	4	2
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

APPENDIX B

**Summary of the Incidence of Neoplasms in Mice
Administered 2,6-Toluenediamine dihydrochloride
in the Diet**

TABLE B1.

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MICE
ADMINISTERED 2,6-TOLUENEDIAMINE DIHYDROCHLORIDE IN THE DIET

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	50	50	50
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(50)	(50)	(50)
FIBROMA		1 (2%)	
FIBROSARCOMA	1 (2%)	1 (2%)	
RESPIRATORY SYSTEM			
#LUNG	(50)	(50)	(50)
HEPATOCELLULAR CARCINOMA, METAST	4 (8%)	2 (4%)	4 (8%)
ALVEOLAR/BRONCHIOLAR ADENOMA	7 (14%)	8 (16%)	6 (12%)
ALVEOLAR/BRONCHIOLAR CARCINOMA	4 (8%)	5 (10%)	1 (2%)
FIBROSARCOMA, METASTATIC		1 (2%)	
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(50)	(50)	(50)
MALIGNANT LYMPHOMA, NOS	2 (4%)	5 (10%)	1 (2%)
*HEMATOPOIETIC SYSTEM	(50)	(50)	(50)
NEOPLASM, NOS			2 (4%)
#SPLEEN	(49)	(50)	(50)
MALIGNANT LYMPHOMA, NOS		1 (2%)	
#LYMPH NODE	(44)	(38)	(41)
HEPATOCELLULAR CARCINOMA, METAST		1 (3%)	
MALIGNANT LYMPHOMA, NOS		1 (3%)	1 (2%)
#PEYER'S PATCH	(47)	(48)	(45)
MALIGNANT LYMPHOMA, NOS		1 (2%)	
CIRCULATORY SYSTEM			
#SPLEEN	(49)	(50)	(50)
HEMANGIOMA		1 (2%)	1 (2%)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
HEMANGIOSARCOMA		1 (2%)	
#LIVER	(50)	(50)	(49)
HEMANGIOMA	1 (2%)	1 (2%)	1 (2%)
HEMANGIOSARCOMA		2 (4%)	1 (2%)
ANGIOSARCOMA		1 (2%)	
DIGESTIVE SYSTEM			
#LIVER	(50)	(50)	(49)
NEOPLASM, NOS	3 (6%)		
HEPATOCELLULAR ADENOMA	7 (14%)	5 (10%)	5 (10%)
HEPATOCELLULAR CARCINOMA	14 (28%)	12 (24%)	13 (27%)
FIBROSARCOMA, METASTATIC		1 (2%)	
#STOMACH	(48)	(49)	(47)
HEPATOCELLULAR CARCINOMA, METAST		1 (2%)	
URINARY SYSTEM			
NONE			
ENDOCRINE SYSTEM			
#PITUITARY	(42)	(37)	(42)
ADENOMA, NOS	1 (2%)	1 (3%)	
#ADRENAL	(47)	(43)	(47)
CORTICAL ADENOMA		1 (2%)	
#ADRENAL/CAPSULE	(47)	(43)	(47)
ADENOMA, NOS			1 (2%)
REPRODUCTIVE SYSTEM			
NONE			
NERVOUS SYSTEM			
NONE			

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
 * NUMBER OF ANIMALS NECROPSIED

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
SPECIAL SENSE ORGANS			
*HARDERIAN GLAND ADENOMA, NOS	(50)	(50) 2 (4%)	(50) 1 (2%)
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
SITE UNKNOWN CARCINOMA, NOS		1	
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	50	50	50
NATURAL DEATH ^a	17	17	15
MORIBUND SACRIFICE	2	2	3
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	31	31	32
ANIMAL MISSING			
^a INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	31	36	26
TOTAL PRIMARY TUMORS	40	51	34
TOTAL ANIMALS WITH BENIGN TUMORS	15	17	14
TOTAL BENIGN TUMORS	16	20	15
TOTAL ANIMALS WITH MALIGNANT TUMORS	19	26	16
TOTAL MALIGNANT TUMORS	21	31	17
TOTAL ANIMALS WITH SECONDARY TUMORS#	4	3	4
TOTAL SECONDARY TUMORS	4	6	4
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT	3		2
TOTAL UNCERTAIN TUMORS	3		2
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE B2.

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MICE
ADMINISTERED 2,6-TOLUENEDIAMINE DIHYDROCHLORIDE IN THE DIET

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	50	50	50
INTEGUMENTARY SYSTEM			
*SKIN	(50)	(50)	(50)
SQUAMOUS CELL PAPILLOMA		1 (2%)	
SQUAMOUS CELL CARCINOMA		1 (2%)	
*SUBCUT TISSUE	(50)	(50)	(50)
SQUAMOUS CELL CARCINOMA	1 (2%)		
FIBROSARCOMA			2 (4%)
RESPIRATORY SYSTEM			
#LUNG	(50)	(50)	(49)
ALVEOLAR/BRONCHIOLAR ADENOMA	4 (8%)	7 (14%)	3 (6%)
ALVEOLAR/BRONCHIOLAR CARCINOMA		1 (2%)	
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(50)	(50)	(50)
MALIGNANT LYMPHOMA, NOS	3 (6%)	9 (18%)	5 (10%)
MALIG. LYMPHOMA, HISTIOCYTIC TYPE	1 (2%)		
*HEMATOPOIETIC SYSTEM NEOPLASM, NOS	(50) 4 (8%)	(50) 1 (2%)	(50) 1 (2%)
#SPLEEN	(49)	(49)	(49)
MALIGNANT LYMPHOMA, NOS			1 (2%)
#LYMPH NODE	(40)	(44)	(39)
ENDOMETRIAL STROMAL SARCOMA, MET			1 (3%)
#PEYER'S PATCH	(47)	(47)	(44)
MALIGNANT LYMPHOMA, NOS		1 (2%)	

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
CIRCULATORY SYSTEM			
*SUBCUT TISSUE	(50)	(50)	(50)
HEMANGIOMA	1 (2%)		
HEMANGIOSARCOMA		1 (2%)	
#SPLEEN	(49)	(49)	(49)
HEMANGIOSARCOMA		1 (2%)	
#LIVER	(50)	(49)	(49)
HEMANGIOMA	1 (2%)		1 (2%)
DIGESTIVE SYSTEM			
#LIVER	(50)	(49)	(49)
HEPATOCELLULAR ADENOMA	4 (8%)	3 (6%)	4 (8%)
HEPATOCELLULAR CARCINOMA			3 (6%)
ENDOMETRIAL STROMAL SARCOMA, MET			1 (2%)
#DUODENUM	(47)	(47)	(44)
ADENOMATOUS POLYP, NOS		1 (2%)	
URINARY SYSTEM			
NONE			
ENDOCRINE SYSTEM			
#PITUITARY	(37)	(38)	(38)
ADENOMA, NOS	3 (8%)	5 (13%)	6 (16%)
#ADRENAL	(45)	(47)	(42)
CORTICAL ADENOMA			1 (2%)
PHEOCHROMOCYTOMA		1 (2%)	
#ADRENAL/CAPSULE	(45)	(47)	(42)
ADENOMA, NOS		1 (2%)	1 (2%)
#THYROID	(38)	(49)	(46)
FOLLICULAR-CELL ADENOMA		1 (2%)	

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
 * NUMBER OF ANIMALS NECROPSIED

TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
REPRODUCTIVE SYSTEM			
#UTERUS ENDOMETRIAL STROMAL SARCOMA	(48)	(48)	(50) 1 (2%)
#OVARY TUBULAR ADENOMA	(35)	(47) 1 (2%)	(39) 1 (3%)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
*HARDERIAN GLAND ADENOMA, NOS	(50)	(50) 1 (2%)	(50)
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS SARCOMA, NOS	(50)	(50) 1 (2%)	(50)
LOWER LEG OSTEOSARCOMA			1
OMENTUM SARCOMA, NOS		1	

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
 * NUMBER OF ANIMALS NECROPSIED

TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	50	50	50
NATURAL DEATH ^a	10	9	10
MORIBUND SACRIFICE			
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED	1	1	1
TERMINAL SACRIFICE	39	40	39
ANIMAL MISSING			
^a INCLUDES AUTOLYZED ANIMALS			
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	21	30	24
TOTAL PRIMARY TUMORS	22	39	31
TOTAL ANIMALS WITH BENIGN TUMORS	13	16	16
TOTAL BENIGN TUMORS	13	22	17
TOTAL ANIMALS WITH MALIGNANT TUMORS	5	15	12
TOTAL MALIGNANT TUMORS	5	16	13
TOTAL ANIMALS WITH SECONDARY TUMORS#			1
TOTAL SECONDARY TUMORS			2
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT	4	1	1
TOTAL UNCERTAIN TUMORS	4	1	1
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

APPENDIX C

**Summary of the Incidence of Nonneoplastic Lesions in Rats
Administered 2,6-Toluenediamine dihydrochloride
in the Diet**

TABLE C1.

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE RATS
ADMINISTERED 2,6-TOLUENEDIAMINE DIHYDROCHLORIDE IN THE DIET

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	50	50	50
INTEGUMENTARY SYSTEM			
*SKIN EPIDERMAL INCLUSION CYST	(50)	(50) 1 (2%)	(50)
*SUBCUT TISSUE NECROSIS, NOS	(50) 1 (2%)	(50)	(50)
RESPIRATORY SYSTEM			
#LUNG FIBROSIS	(49) 1 (2%)	(50)	(50)
HEMATOPOIETIC SYSTEM			
#SPLEEN FIBROSIS INFARCT, NOS HEMATOPOIESIS	(50) 2 (4%)	(50) 1 (2%) 1 (2%) 2 (4%)	(49) 3 (6%)
#SPLENIC FOLLICLES ATROPHY, NOS	(50)	(50) 2 (4%)	(49)
CIRCULATORY SYSTEM			
#HEART MINERALIZATION THROMBOSIS, NOS CALCIFICATION, NOS	(48) 1 (2%)	(50) 1 (2%)	(50) 1 (2%)
#AURICULAR APPENDAGE THROMBOSIS, NOS	(48) 1 (2%)	(50)	(50) 3 (6%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C1. MALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
#MYOCARDIUM	(48)	(50)	(50)
INFLAMMATION, NOS		1 (2%)	
DEGENERATION, NOS	2 (4%)	4 (8%)	7 (14%)
*MESENTERY	(50)	(50)	(50)
PERIVASCULITIS	2 (4%)		
#URINARY BLADDER	(45)	(45)	(44)
PERIVASCULITIS	1 (2%)		
DIGESTIVE SYSTEM			
#LIVER	(50)	(50)	(50)
NECROSIS, FOCAL	5 (10%)	1 (2%)	3 (6%)
METAMORPHOSIS FATTY	8 (16%)	5 (10%)	1 (2%)
BASOPHILIC CYTO CHANGE	3 (6%)	5 (10%)	10 (20%)
FOCAL CELLULAR CHANGE	1 (2%)	1 (2%)	5 (10%)
CLEAR-CELL CHANGE			1 (2%)
#BILE DUCT	(50)	(50)	(50)
HYPERPLASIA, NOS	1 (2%)	2 (4%)	
#STOMACH	(49)	(49)	(48)
MINERALIZATION	2 (4%)		
CALCIFICATION, NOS		1 (2%)	
HYPERPLASIA, BASAL CELL	1 (2%)	1 (2%)	
#PEYER'S PATCH	(47)	(48)	(48)
HYPERPLASIA, NOS	1 (2%)		
URINARY SYSTEM			
#KIDNEY	(50)	(50)	(50)
MINERALIZATION			1 (2%)
HYDRONEPHROSIS	1 (2%)		
NEPHROPATHY	32 (64%)	39 (78%)	41 (82%)
NECROSIS, NOS		1 (2%)	
CALCIFICATION, NOS		1 (2%)	
HYPERPLASIA, EPITHELIAL	1 (2%)		
#KIDNEY/PELVIS	(50)	(50)	(50)
HYPERPLASIA, EPITHELIAL		1 (2%)	1 (2%)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
 * NUMBER OF ANIMALS NECROPSIED

TABLE C1. MALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ENDOCRINE SYSTEM			
#PITUITARY ANGIECTASIS	(45)	(49) 1 (2%)	(44)
#ADRENAL MEDULLA HYPERPLASIA, NODULAR	(50) 1 (2%)	(50)	(48)
#THYROID CALCIFICATION, FOCAL HYPERPLASIA, C-CELL	(44)	(48) 1 (2%) 2 (4%)	(47) 2 (4%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND GALACTOCELE	(50) 1 (2%)	(50)	(50) 1 (2%)
*PREPUTIAL GLAND INFLAMMATION, NOS	(50) 1 (2%)	(50) 1 (2%)	(50) 1 (2%)
#PROSTATE INFLAMMATION, NOS INFLAMMATION, ACUTE	(42) 1 (2%)	(44) 1 (2%)	(42)
*SEMINAL VESICLE CALCULUS, NOS	(50)	(50) 1 (2%)	(50)
#TESTIS MINERALIZATION ATROPHY, NOS HYPERPLASIA, INTERSTITIAL CELL	(49) 1 (2%) 1 (2%) 2 (4%)	(49) 2 (4%) 4 (8%)	(49) 1 (2%)
NERVOUS SYSTEM			
#BRAIN HEMORRHAGE NECROSIS, NOS NECROSIS, HEMORRHAGIC	(50) 1 (2%)	(50) 1 (2%) 1 (2%)	(50)
SPECIAL SENSE ORGANS			
NONE			

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE C1. MALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*MEDIASTINUM	(50)	(50)	(50)
MINERALIZATION	1 (2%)		
NECROSIS, NOS	1 (2%)		
*ABDOMINAL CAVITY	(50)	(50)	(50)
HEMORRHAGE			1 (2%)
*MESENTERY	(50)	(50)	(50)
NECROSIS, FAT	1 (2%)		
ALL OTHER SYSTEMS			
OMENTUM			
NECROSIS, FAT	2	6	7
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED			1
AUTO/NECROPSY/HISTO PERF		1	
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2.

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE RATS
ADMINISTERED 2,6-TOLUENEDIAMINE DIHYDROCHLORIDE IN THE DIET

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	50	50	50
INTEGUMENTARY SYSTEM			
NONE			
RESPIRATORY SYSTEM			
#LUNG	(50)	(50)	(50)
INFLAMMATION, NOS	1 (2%)	1 (2%)	
INFLAMMATION, INTERSTITIAL GRANULOMA, FOREIGN BODY		1 (2%) 3 (6%)	
HEMATOPOIETIC SYSTEM			
#SPLEEN	(49)	(50)	(50)
CONGESTION, NOS	1 (2%)		
HEMATOPOIESIS	7 (14%)	10 (20%)	11 (22%)
CIRCULATORY SYSTEM			
#HEART	(50)	(49)	(50)
MINERALIZATION	1 (2%)		
#AURICULAR APPENDAGE	(50)	(49)	(50)
THROMBOSIS, NOS			1 (2%)
#MYOCARDIUM	(50)	(49)	(50)
DEGENERATION, NOS	1 (2%)		2 (4%)
DIGESTIVE SYSTEM			
#LIVER	(50)	(50)	(50)
NECROSIS, FOCAL	1 (2%)	3 (6%)	
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2. FEMALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
NECROSIS, HEMORRHAGIC	1 (2%)		
METAMORPHOSIS FATTY	7 (14%)	4 (8%)	
BASOPHILIC CYTO CHANGE	21 (42%)	29 (58%)	41 (82%)
FOCAL CELLULAR CHANGE		2 (4%)	
CLEAR-CELL CHANGE			2 (4%)
#BILE DUCT	(50)	(50)	(50)
HYPERPLASIA, NOS		1 (2%)	
#STOMACH	(50)	(48)	(50)
MINERALIZATION	1 (2%)		
HYPERPLASIA, BASAL CELL		3 (6%)	
ACANTHOSIS		1 (2%)	
#FORESTOMACH	(50)	(48)	(50)
HYPERPLASIA, BASAL CELL	1 (2%)		
#JEJUNAL SUBMUCOSA	(49)	(48)	(48)
INFLAMMATION, NOS			1 (2%)
URINARY SYSTEM			
#KIDNEY	(50)	(50)	(50)
MINERALIZATION	3 (6%)	3 (6%)	4 (8%)
NEPHROPATHY	14 (28%)	10 (20%)	7 (14%)
#KIDNEY/PELVIS	(50)	(50)	(50)
HYPERPLASIA, EPITHELIAL		1 (2%)	
ENDOCRINE SYSTEM			
#PITUITARY	(48)	(43)	(48)
HEMORRHAGIC CYST	1 (2%)		
#THYROID	(49)	(49)	(48)
HYPERPLASIA, C-CELL	1 (2%)		1 (2%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(50)	(50)	(50)
GALACTOCELE	4 (8%)	4 (8%)	6 (12%)
*VAGINA	(50)	(50)	(50)
POLYP			1 (2%)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE C2. FEMALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
#UTERUS HYDROMETRA	(48) 1 (2%)	(49)	(50)
#UTERUS/ENDOMETRIUM HYPERPLASIA, NOS	(48)	(49)	(50) 1 (2%)
NERVOUS SYSTEM			
#BRAIN HEMORRHAGE	(49)	(49) 1 (2%)	(50)
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
FOOT INFLAMMATION, NOS	1		
NECROSIS, NOS	1		
OMENTUM NECROSIS, FAT	2	3	1
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	2	1	2
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

APPENDIX D

Summary of the Incidence of Nonneoplastic Lesions in Mice
Administered 2,6-Toluenediamine dihydrochloride
in the Diet

.

TABLE D1.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MICE
ADMINISTERED 2,6-TOLUENEDIAMINE DIHYDROCHLORIDE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	50	50	50
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(50)	(50)	(50)
INFLAMMATION, ACUTE/CHRONIC		1 (2%)	
NECROSIS, NOS	3 (6%)		4 (8%)
RESPIRATORY SYSTEM			
#LUNG	(50)	(50)	(50)
INFLAMMATION, NOS			1 (2%)
GRANULOMA, FOREIGN BODY		1 (2%)	
HEMATOPOIETIC SYSTEM			
#BONE MARROW	(47)	(43)	(49)
HYPERPLASIA, HEMATOPOIETIC	2 (4%)		
#SPLEEN	(49)	(50)	(50)
HYPERPLASIA, NOS		1 (2%)	1 (2%)
HYPERPLASIA, HEMATOPOIETIC	1 (2%)		
HEMATOPOIESIS	6 (12%)	9 (18%)	3 (6%)
#LYMPH NODE	(44)	(38)	(41)
HYPERPLASIA, NOS			1 (2%)
HEMATOPOIESIS	4 (9%)	2 (5%)	6 (15%)
#MESENTERIC L. NODE	(44)	(38)	(41)
HYPERPLASIA, NOS	1 (2%)		
CIRCULATORY SYSTEM			
*MULTIPLE ORGANS	(50)	(50)	(50)
PERIARTERITIS		1 (2%)	

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
PERIVASCULITIS		1 (2%)	
#HEART	(50)	(49)	(50)
MINERALIZATION			1 (2%)
FIBROSIS	2 (4%)	1 (2%)	
CALCIFICATION, FOCAL			1 (2%)
#MYOCARDIUM	(50)	(49)	(50)
INFLAMMATION, NOS	2 (4%)	3 (6%)	8 (16%)
DEGENERATION, NOS	1 (2%)		
DIGESTIVE SYSTEM			
#LIVER	(50)	(50)	(49)
MINERALIZATION			1 (2%)
HEMORRHAGE		1 (2%)	
FIBROSIS		1 (2%)	
NECROSIS, FOCAL	2 (4%)		4 (8%)
METAMORPHOSIS FATTY	1 (2%)	4 (8%)	7 (14%)
CLEAR-CELL CHANGE			1 (2%)
HEPATOCYTOMEGALY		1 (2%)	
*GALLBLADDER	(50)	(50)	(50)
HYPERPLASIA, PAPILLARY		1 (2%)	
#PANCREAS	(43)	(48)	(43)
NECROSIS, FOCAL	1 (2%)		
#PANCREATIC ACINUS	(43)	(48)	(43)
ATROPHY, NOS	1 (2%)		
ATROPHY, FOCAL			1 (2%)
#STOMACH	(48)	(49)	(47)
MINERALIZATION	1 (2%)		
INFLAMMATION, NOS	1 (2%)	2 (4%)	1 (2%)
HYPERKERATOSIS	1 (2%)		
ACANTHOSIS	1 (2%)		1 (2%)
#SMALL INTESTINE	(47)	(48)	(45)
INFLAMMATION, NOS			1 (2%)
URINARY SYSTEM			
#KIDNEY	(49)	(50)	(49)
MINERALIZATION		10 (20%)	6 (12%)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
HYDRONEPHROSIS INFLAMMATION, INTERSTITIAL DEGENERATION, CYSTIC NEPHROSIS, NOS NECROSIS, NOS	1 (2%) 1 (2%)	1 (2%) 1 (2%)	1 (2%)
#KIDNEY/TUBULE DEGENERATION, NOS NECROSIS, FOCAL REGENERATION, NOS	(49) 1 (2%) 1 (2%)	(50)	(49) 1 (2%)
ENDOCRINE SYSTEM			
#PITUITARY CYST, NOS	(42)	(37)	(42) 1 (2%)
#ADRENAL/CAPSULE HYPERPLASIA, NOS	(47) 1 (2%)	(43)	(47)
#THYROID FOLLICULAR CYST, NOS HYPERPLASIA, FOLLICULAR-CELL	(45)	(47) 1 (2%)	(48) 3 (6%)
REPRODUCTIVE SYSTEM			
*PREPUTIAL GLAND ABSCCESS, NOS NECROSIS, NOS	(50) 1 (2%)	(50) 4 (8%)	(50) 1 (2%)
#PROSTATE ABSCCESS, NOS	(41)	(41) 1 (2%)	(49)
#TESTIS MINERALIZATION	(50) 1 (2%)	(49)	(50)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
*HARDERIAN GLAND MINERALIZATION	(50)	(50) 1 (2%)	(50)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
INFLAMMATION, NOS		1 (2%)	
HYPERPLASIA, NOS		1 (2%)	
*EAR	(50)	(50)	(50)
GRANULOMA, FOREIGN BODY	1 (2%)		
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
OMENTUM			
NECROSIS, FAT	7	4	3
CALCIFICATION, NOS	1		
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	6	5	3
‡ NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D2.

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MICE
ADMINISTERED 2,6-TOLUENEDIAMINE DIHYDROCHLORIDE IN THE DIET

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	50	50	50
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE NECROSIS, NOS	(50)	(50) 1 (2%)	(50)
RESPIRATORY SYSTEM			
#LUNG CONGESTION, NOS HEMORRHAGE	(50)	(50) 1 (2%)	(49) 1 (2%) 2 (4%)
HEMATOPOIETIC SYSTEM			
#SPLEEN HYPERPLASIA, NOS HEMATOPOIESIS	(49) 5 (10%) 7 (14%)	(49) 1 (2%) 6 (12%)	(49) 1 (2%) 7 (14%)
#LYMPH NODE HEMATOPOIESIS	(40) 1 (3%)	(44) 1 (2%)	(39) 1 (3%)
#MESENTERIC L. NODE CONGESTION, NOS	(40) 1 (3%)	(44)	(39)
#LIVER HEMATOPOIESIS	(50) 2 (4%)	(49) 2 (4%)	(49)
CIRCULATORY SYSTEM			
#HEART MINERALIZATION	(49)	(50)	(49) 1 (2%)
DIGESTIVE SYSTEM			
#LIVER NECROSIS, FOCAL	(50) 3 (6%)	(49) 1 (2%)	(49)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
METAMORPHOSIS FATTY	1 (2%)		
#PANCREATIC ACINUS DEGENERATION, NOS	(45) 1 (2%)	(47)	(44)
#STOMACH INFLAMMATION, NOS ACANTHOSIS	(48)	(47) 1 (2%)	(48) 2 (4%) 1 (2%)
#DUODENUM CONGESTION, NOS	(47) 1 (2%)	(47)	(44)
#JEJUNUM CONGESTION, NOS	(47) 1 (2%)	(47)	(44)
#ILEAL MUCOUS MEMBRAN HEMORRHAGE	(47) 1 (2%)	(47)	(44)
URINARY SYSTEM			
#KIDNEY MINERALIZATION CONGESTION, NOS INFLAMMATION, INTERSTITIAL GLOMERULOSCLEROSIS, NOS	(50)	(50) 1 (2%) 1 (2%)	(49) 1 (2%) 1 (2%)
#URINARY BLADDER HYPERPLASIA, EPITHELIAL	(48)	(48) 1 (2%)	(46)
ENDOCRINE SYSTEM			
#ADRENAL/CAPSULE HYPERPLASIA, NOS	(45) 3 (7%)	(47) 1 (2%)	(42) 1 (2%)
#THYROID HYPERPLASIA, FOLLICULAR-CELL	(38) 1 (3%)	(49)	(46)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND HYPERPLASIA, NOS	(50) 1 (2%)	(50)	(50)
*VAGINA NECROSIS, NOS	(50) 1 (2%)	(50)	(50)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
#UTERUS	(48)	(48)	(50)
HYDROMETRA	12 (25%)	4 (8%)	6 (12%)
INFLAMMATION, NOS	4 (8%)	2 (4%)	
PYOMETRA	1 (2%)	1 (2%)	2 (4%)
METAPLASIA, SQUAMOUS	1 (2%)		
#UTERUS/ENDOMETRIUM	(48)	(48)	(50)
HYPERPLASIA, NOS	1 (2%)	3 (6%)	2 (4%)
HYPERPLASIA, CYSTIC	8 (17%)	23 (48%)	25 (50%)
#OVARY	(35)	(47)	(39)
MINERALIZATION			1 (3%)
CYST, NOS	2 (6%)	1 (2%)	4 (10%)
FOLLICULAR CYST, NOS		1 (2%)	
CONGESTION, NOS		1 (2%)	
INFLAMMATION, NOS			1 (3%)
DEGENERATION, CYSTIC		2 (4%)	3 (8%)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*ABDOMINAL CAVITY	(50)	(50)	(50)
ABSCESS, NOS		1 (2%)	
ALL OTHER SYSTEMS			
OMENTUM			
MINERALIZATION			1

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
 * NUMBER OF ANIMALS NECROPSIED

TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
NECROSIS, FAT	4	5	1
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	6	2	3
AUTO/NECROPSY/HISTO PERF	1		1

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
* NUMBER OF ANIMALS NECROPSIED

APPENDIX E

**Analysis of 2,6-Toluenediamine
Midwest Research Institute**

Appendix E
Analysis of 2,6-Toluenediamine (Lot 012157)
Midwest Research Institute

A. ELEMENTAL ANALYSIS

Element	C	H	N
Theory	68.81	8.25	22.94
Determined	68.75	8.11	22.73
	68.94	8.09	22.81

B. MELTING POINT

<u>Determined</u>	<u>Literature Values</u>
105°C (Dupont 900 DTA)	105°C (Sadtlter Standard Spectra)
104°-105.5°C (visual, capillary)	

C. THIN-LAYER CHROMATOGRAPHY

Plates: Silica gel, F-254
Amount Spotted: 100 and
300 μ g
System 1: Ethyl acetate,
100%
 R_f : 0.37, origin (trace
visualized at 254 nm only)
 R_{st} : 0.58, origin

Ref. Standard: Aniline
Visualization: Ultraviolet
254 and 366nm, and furfural
System 2: Tetrahydrofuran,
100%
 R_f : 0.61, origin (trace
visualized at 254 nm only)
 R_{st} : 0.87, origin

D. VAPOR-PHASE CHROMATOGRAPHY

Instrument: Bendix 2500

Column: 3% OV-1 on 80/100 Supelcoport, 1.8 m x 4 mm I.D.

Inlet temperature: 280°C

Detector: Flame Ionization

Oven Temperature Program: 5 min at 50°C, then
50°-250°C at
10°C/min

Results: One homogeneous peak, retention time 11.8 min

E. SPECTRAL DATA

a. Infrared

Instrument: Beckman IR-12

Cell: 0.8% KBr pellet

Results: See Figure 5

Consistent with literature
spectrum (Finkleshtein
and Boitsov, 1960).

b. Ultraviolet/Visible

Instrument: Cary 118

No literature reference
found.

$$\epsilon_{\max} 289.5 = (1.48 \pm 0.01 (\delta)) \\ \times 10^3$$

Another maximum near the solvent
cut-off of 210 nm.

No absorption in the visible region,
350 - 800 nm at 1 mg/ml

Solvent: Methanol




Figure 5. Infrared Absorption Spectrum of 2,6-Toluediamine

c. Nuclear Magnetic Resonance

Instrument: Varian HA-100	Consistent with
Solvent: CDCl_3 with internal TMS	literature spectrum (Sadtler, Standard Spectra)
Assignments: (Refer to figure 6)	
(a) 1.87 δ (b) 3.42 δ (c) 6.08 δ	
(d) 6.74 δ , $J_{\text{cd}} = 8$ cps	
Integration Ratios: (a)3.06 (b) 3.83 (c) 2.02 (d) 0.93	

CONCLUSIONS

Elemental analysis agrees with theory. Thin-layer chromatography indicates only a trace impurity at the origin. Vapor-phase chromatography gives one homogeneous peak. The infrared and nuclear magnetic resonance spectra are consistent with the structure.




Figure 6. Infrared Absorption Spectrum of 2,6-Toluenediamine dihydrochloride

APPENDIX F

**Analysis of 2,6-Toluenediamine dihydrochloride
Midwest Research Institute**

Appendix F

Analysis of 2,6-Toluenediamine dihydrochloride

Midwest Research Institute

A. SYNTHESIS OF 2,6-TOLUENEDIAMINE DIHYDROCHLORIDE

2,6-Toluenediamine (1,921 g, 15.72 eq.) was dissolved in hydrochloric acid (2,880 ml of a 1:1 mixture of concentrated HCl:water). The solution was filtered and an additional 100 ml of concentrated hydrochloric acid was added to the filtrate. The filtrate was cooled in an ice-salt bath. The colorless crystals were collected and dried in a vacuum oven overnight. The yield was 1,863 g. The filtrate was concentrated and a second crop of crystals collected and dried (993 g).

B. ELEMENTAL ANALYSIS

Element	C	H	N	Cl
Theory	43.09	6.20	14.36	36.35
Theory with 5.8% H ₂ O	42.86	6.23	14.28	36.14
Determined	43.13	6.13	14.33	35.8
	43.21	6.17	14.40	36.0

C. MELTING POINT

Determined

174° to 250°C dec.
(visual; sealed,
evacuated capillary)
186.5° to 232°C
(Dupont 900DTA)

Literature Values

No literature found.

D. THIN-LAYER CHROMATOGRAPHY

Plates: Silica gel G, F 254
Amount Spotted: 100 and 300 μ g

Ref. Standard: Aniline
Visualization: Ultraviolet
(254 nm) and furfuraldehyde

System 1: n-Butanol:conc.
aqueous ammonium hydroxide
(99:1)

R_f: 0.54

R_{st}: 0.65

System 2: Tetrahydrofuran:
conc. aqueous ammonium
hydroxide (99:1)

R_f: 0.75

R_{st}: 0.91

E. VAPOR-PHASE CHROMATOGRAPHY

(a) System 1

Instrument: Tracor MT-220

Detection: Flame ionization

Column: 3% OV-225 on Chromosorb W (HP) 80/100,
1.8 M x 4 mm I.D.

Inlet temperature: 280°C

Detector temperature: 280°C

Oven temperature program: 5 min at 100°C, then
100°-200°C at
10°C/min

Results: One homogeneous peak, retention time 11.8 min

(b) System 2

Instrument: Tracor MT-220

Detection: Flame ionization

Column: 3% OV-1 on Supelcoport, 80/100, 1.8 m x 4 mm I.D.

Inlet temperature: 270°C

Detector temperature: 280°C

Oven temperature program: 5 min at 75°C, then 75°-200°C
at 10°C/min

Results: One homogeneous peak, retention time 10.8 min

F. SPECTRAL DATA

(a) Infrared

Instrument: Beckman IR-12

Cell: 1.0% Potassium
bromide pellet.

Results: See Figure 6.

Consistent with the spectrum
of amine (Finkleshtein
and Boitsov, 1960).

(b) Ultraviolet/Visible

Literature Values

Instrument: Cary 118

No literature reference found.

$\lambda_{\max}(\text{nm})$	$\epsilon \times 10^{-3}$
282	1.928+0.003 (δ)
232.5	8.66+0.03 (δ)

No absorbance between 350 and 800 nm (visible range) at a concentration of 0.3 mg/ml.

Solvent: water

(c) Nuclear Magnetic Resonance

Consistent with literature spectrum of amine (Sadtler, Standard Spectra)

Instrument: Varian HA-100

Solvent: Dimethylsulfoxide-d6 with internal tetramethylsilane

Assignments: (refer to Figure 7)

(a) s, δ 2.39 ppm

(b) m, δ 7.32-7.74 ppm

(c) s, δ 9.67 ppm

Integration Ratios:

(a) 2.76

(b) 3.24

(c) NH₂ and HCl

G. NUCLEAR MAGNETIC RESONANCE: (undried sample)

Instrument: Varian HA100

Solvent: DMSO-d₆ with internal TMS

Assignments: (refer to Figure 8)

(a) 2.33 δ

(b) 7.17-7.60 δ

(c) 10.10 δ

Integration Ratios:

(a) 2.89

(b) 3.11

(c) 6.33

Consistent with literature spectrum of amine (Sadtler Standard Spectra).

CONCLUSIONS

The elemental analysis for chloride is slightly low; carbon, hydrogen, and nitrogen analyses agree with the theoretical values. Thin-layer chromatography shows a trace impurity at the origin. Vapor-phase chromatography indicates one homogeneous peak. The infrared, ultraviolet, visible, and nuclear magnetic resonance spectra conform to the structure.




Figure 7. Nuclear Magnetic Resonance Spectrum of 2,6-Toluenediamine

66




Figure 8. Nuclear Magnetic Resonance Spectrum of 2,6-Toluenediamine dihydrochloride

APPENDIX G

**Stability Analysis of 2,6-Toluenediamine dihydrochloride
Midwest Research Institute**

APPENDIX G

Stability Analysis of 2,6-Toluenediamine dihydrochloride

Midwest Research Institute

STABILITY TESTING

A. BULK CHEMICAL

1. METHOD: Samples of bulk chemical were stored at -20°, 5°, 25°, and 60°C for 2 weeks and then analyzed by vapor-phase chromatography using the following conditions:

Instrument: Bendix 2500

Detection: Flame Ionization

Column: 3% OV-225 on chromosorb W (HP) 80/10 mesh; glass,
1.8 M x 2 mm

Oven Temperature: 150°C isothermal

2. RESULTS:

<u>Temperature (°C)</u>	<u>Response (nm/μg)</u>
-20	16.7+0.2
5	16.6+0.2
25	16.4+0.3
60	16.7+0.2

3. CONCLUSION: 2,6-Toluenediamine dihydrochloride is stable in bulk for 2 weeks at 60°C.

B. CHEMICAL/FEED MIXTURE

1. METHOD: A 10.03% 2,6-toluenediamine hydrochloride rat-feed mixture was prepared and stored at -20°, 5°, 25°, and 45°C for 2 weeks. One-gram samples of this mixture were blended with 50 ml of 95% ethanol for 1 minute on a Brinkman Polytron mixer. These blended samples were centrifuged for 10 minutes and the supernatants decanted into 100-ml volumetric flasks. The centrifugates were blended with another 50 ml of 95% ethanol for 1 minute on a Polytron mixer. These blended samples were centrifuged and the supernatants combined with those in the 100-ml volumetric flasks from the previous centrifugations. The resulting solutions were diluted to volume with 95% ethanol and injected on a gas chromatograph under the following conditions:

Instrument: Bendix 2500
Column: 3% OV-225 on chromosorb W (HP); 1.8 m x 2 mm, glass
Temperature: 145°C, isothermal
Detection: Flame ionization
R_f of Test Compound: 2.0 minutes

2. RESULTS:

<u>Temperature (°C)</u>	<u>Percent 2,6-Toluenediamine dihydrochloride in Feed</u>
-20	11.36 <u>±</u> 1.80
5	10.37 <u>±</u> 1.69
25	11.59 <u>±</u> 1.90
45	10.25 <u>±</u> 1.66

3. CONCLUSION: 2,6-Toluenediamine dihydrochloride is stable in feed for 2 weeks at temperatures up to 45°C.

APPENDIX H

**Analytical Procedure for Analysis of
2,6-Toluenediamine dihydrochloride
in Feed**

APPENDIX H

Analytical Procedure for Analysis of 2,6-Toluenediamine dihydrochloride in Feed

Samples of 5 g were extracted twice with 25 ml of 0.6N hydrochloric acid in 100-ml ground glass stoppered graduated cylinders. The supernatant solutions were adjusted to $\text{pH} \geq 12$ by addition of approximately 10 ml of 10% sodium hydroxide solution. After addition of 5 g of sodium chloride, the aqueous phases were extracted with 50 ml of chloroform in separatory funnels. The emulsified organic phases were centrifuged for 12 minutes and 10-ml aliquots were treated with anhydrous sodium sulfate and then reduced in volume to 1 ml by evaporation under nitrogen and analyzed by vapor-phase chromatography at 150° oven temperature on a 3% OV-17, 100/200 Supelcoport glass column. The retention time of the 2,6-toluenediamine dihydrochloride was 6.2 minutes.

Review of the Bioassay of 2,6-Toluenediamine Dihydrochloride (2,6-TDA)*
for Carcinogenicity by the Data Evaluation/Risk Assessment Subgroup
of the Clearinghouse on Environmental Carcinogens

February 15, 1980

The Clearinghouse on Environmental Carcinogens was established in May, 1976, in compliance with DHEW Committee Regulations and the Provisions of the Federal Advisory Committee Act. The purpose of the Clearinghouse is to advise the Director of the National Cancer Institute (NCI) on its bioassay program to identify and to evaluate chemical carcinogens in the environment to which humans may be exposed. The members of the Clearinghouse have been drawn from academia, industry, organized labor, public interest groups, State health officials, and quasi-public health and research organizations. Members have been selected on the basis of their experience in carcinogenesis or related fields and, collectively, provide expertise in chemistry, biochemistry, biostatistics, toxicology, pathology, and epidemiology. Representatives of various Governmental agencies participate as ad hoc members. The Data Evaluation/Risk Assessment Subgroup of the Clearinghouse is charged with the responsibility of providing a peer review of reports prepared on NCI-sponsored bioassays of chemicals studied for carcinogenicity. It is in this context that the below critique is given on the bioassay of 2,6-Toluenediamine Dihydrochloride (2,6-TDA) for carcinogenicity.

The primary reviewer for the report on the bioassay of 2,6-TDA said that the selection and testing of 2,6-TDA was justified by its broad human exposure; its structural relationship to 2,4-TDA, a known carcinogen; and its mutagenicity in bacteria. After a brief review of the results of the bioassay, he indicated that a dose-related incidence of liver tumors occurred in treated male rats and female mice. He opined that the maximum tolerated dose was not reached. The reviewer pointed out that the mice receiving 2,4-TDA were given a dose about two times the one used in this study. Based on these findings and certain inconsistencies in the language of the write-up, he recommended that the report on the bioassay of 2,6-TDA not be accepted. He said that acceptance of the report would indicate that the study was adequate and the chemical need not be further tested.

The secondary reviewer agreed with some of the reservations expressed by the primary reviewer. Despite the reservations, he said the conclusion that the chemical was not carcinogenic, under the conditions of test, was valid.

A Clearinghouse member suggested that a statement be added to the report indicating the need for further testing to resolve questions raised by deficiencies in the study. Another member agreed that such a statement should be included, especially since there is a wide variety of species differences to aromatic amines. He suggested that the hamster might be an appropriate test species.

The secondary reviewer moved that the part of the report dealing with the study in rats be approved. He further moved that the part dealing with the mice was inadequate because of the use of too low dosages and that additional animal studies should be conducted. The motion was seconded and approved unanimously.

Members present were:

Arnold L. Brown (Chairman), University of Wisconsin Medical School
David B. Clayson, Eppley Institute for Research in Cancer
Joseph Highland, Environmental Defense Fund
William Lijinsky, Federick Cancer Research Center
Henry C. Pitot, University of Wisconsin Medical Center
Verne A. Ray, Pfizer Medical Research Laboratory
Louise Strong, University of Texas Health Sciences Center

-
- * Subsequent to this review, changes may have been made in the bioassay report either as a result of the review or other reasons. Thus, certain comments and criticisms reflected in the review may no longer be appropriate.

NIH Publication No. 80-1756
August 1980