

ADVERTISEMENT

Keep up with critical fields...

CHEMISTRY
MEDICINE
NURSING
BIOLOGY
PSYCHOLOGY
ENGINEERING
PHYSICS
BUSINESS
EDUCATION

▶ Get E-mail alerts from

WILEY
InterScience

My Cart My Profile Log In Athens Log In

WILEY
InterScience®
DISCOVER SOMETHING GREAT

Home / [Medical, Veterinary and Health Sciences](#) / [Geriatric Medicine](#)

PSYCHOGERIATRICS

The Official Journal of the Japanese Psychogeriatric Society



Psychogeriatrics

Volume 9 Issue 4, Pages 173 - 179

Published Online: 8 Mar 2010

Journal compilation © 2010 The Japanese Psychogeriatric Society

- [Get Sample Copy](#)
- [Recommend to Your Librarian](#)
- [Save journal to My Profile](#)
- [Set E-Mail Alert](#)
- ✉ [Email this page](#)
- 🖨 [Print this page](#)
- 📡 [RSS web feed \(What is RSS?\)](#)

The Official Journal of the Japanese Psychogeriatric Society

Society Information

[Go to Society Site](#)

- [Save Article to My Profile](#)
- [Download Citation](#)
- [Request Permissions](#)

< [Previous Abstract](#) | [Next Abstract](#) >

Abstract | [References](#) | Full Text: [HTML](#), [PDF](#) (Size: 180K) | [Related Articles](#) | [Citation Tracking](#)

ORIGINAL ARTICLE

Effect of aromatherapy on patients with Alzheimer's disease

Daiki JIMBO, ¹ Yuki KIMURA, ¹ Miyako TANIGUCHI, ¹ Masashi INOUE ² and Katsuya URAKAMI ¹

¹ Section of Environment and Health Science, Department of Biological Regulation, School of Health Science, Faculty of Medicine and ² Information Media Center, Tottori University, Yonago, Japan

Correspondence to Dr Katsuya Urakami PhD MD, Section of Environment and Health Science, Department of Biological Regulation, School of Health Science, Faculty of Medicine, Tottori University, Nishimachi 86, Yonago 683-8503, Japan. Email: kurakami@med.tottori-u.ac.jp

KEYWORDS

Alzheimer's disease • aromatherapy • cognitive function • dementia • elderly people • non-pharmacological therapy

ABSTRACT

Objective: Recently, the importance of non-pharmacological therapies for dementia has come to the fore. In the present study, we examined the curative effects of aromatherapy in dementia in 28 elderly people, 17 of whom had Alzheimer's disease (AD).

Methods: After a control period of 28 days, aromatherapy was performed over the following 28 days, with a wash out period of another 28 days. Aromatherapy consisted of the use of rosemary and lemon essential oils in the morning, and lavender and orange in the evening. To determine the effects of aromatherapy, patients were evaluated using the Japanese version of the Gottfries, Brane, Steen scale (GBSS-J), Functional Assessment Staging of Alzheimer's disease (FAST), a revised version of Hasegawa's Dementia Scale (HDS-R), and the Touch Panel-type Dementia Assessment Scale (TDAS) four times: before the control period, after the control period, after aromatherapy, and after the washout period.

Results: All patients showed significant improvement in personal orientation related to cognitive function on both the GBSS-J and TDAS after therapy. In particular, patients with AD showed significant improvement in total TDAS scores. Result of routine laboratory tests showed no significant changes, suggesting that there were no side-effects associated with the use of aromatherapy. Results from Zarit's score showed no significant changes, suggesting that caregivers had no effect on the improved patient scores seen in the other tests.

Conclusions: In conclusion, we found aromatherapy an efficacious non-pharmacological therapy for dementia. Aromatherapy may have some potential for improving cognitive function, especially in AD patients.