


Worldwide numbers represent medical diagnoses of TBI that occurred anywhere U.S. forces are located including the continental United States since 2000.

Concussion/Mild TBI is characterized by the following: Confused or disoriented state which lasts less than 24 hours; or loss of consciousness for up to 30 minutes; or memory loss lasting less than 24 hours. Excludes penetrating TBI. A CT scan is not indicated for most patients with a Mild TBI. If obtained, it is normal.

Moderate TBI is characterized by the following: Confused or disoriented state which lasts more than 24 hours; or loss of consciousness for more than 30 minutes, but less than 24 hours; or memory loss lasting greater than 24 hours but less than seven days; or meets criteria for Mild TBI except an abnormal CT scan is present. Excludes penetrating TBI. A structural brain imaging study may be normal or abnormal.

Severe TBI is characterized by the following: Confused or disoriented state which lasts more than 24 hours; or loss of consciousness for more than 24 hours; or memory loss for more than seven days. Excludes penetrating TBI. A structural brain imaging study may be normal but usually is abnormal.

Penetrating TBI, or open head injury, is characterized by the following: A head injury in which the scalp, skull and dura mater (the outer layer of the meninges) are penetrated. Penetrating injuries can be caused by high-velocity projectiles or objects of lower velocity such as knives, or bone fragments from a skull fracture that are driven into the brain.



Source: Defense Medical Surveillance System (DMSS), Theater Medical Data Store (TMDS) provided by the Armed Forces Health Surveillance Center (AFHSC)


Prepared by the Defense and Veterans Brain Injury Center (DVBIC)

2001, as of May 15, 2015

Service	Component	Severity	Diagnosed
Army	Active	Penetrating	97
		Severe	74
		Moderate	535
		Mild	3,113
		Not Classifiable	442
	Guard	Penetrating	11
		Severe	9
		Moderate	49
		Mild	367
		Not Classifiable	40
	Reserve	Penetrating	5
		Severe	4
		Moderate	37
		Mild	194
		Not Classifiable	18

2001 as of May 15, 2015

Source: Defense Medical Surveillance System (DMSS), Theater Medical Data Store (TMDS) provided by the Armed Forces Health Surveillance Center (AFHSC)




Prepared by the Defense and Veterans Brain Injury Center (DVBIC)

2001, as of May 15, 2015

Service	Component	Severity	Diagnosed
Navy	Active	Penetrating	72
		Severe	38
		Moderate	350
		Mild	1527
		Not Classifiable	344
	Reserve	Penetrating	2
		Severe	2
		Moderate	19
		Mild	91
		Not Classifiable	16

2001 as of May 15, 2015

Source: Defense Medical Surveillance System (DMSS), Theater Medical Data Store (TMDS) provided by the Armed Forces Health Surveillance Center (AFHSC)



Theater Medical Data Store (TMDS) provided by the Armed Forces Health Surveillance Center (AFHSC)


Prepared by the Defense and Veterans Brain Injury Center (DVBIC)

2001, as of May 15, 2015

Service	Component	Severity	Diagnosed
Air Force	Active	Penetrating	48
		Severe	25
		Moderate	289
		Mild	1,566
		Not Classifiable	138
	Guard	Penetrating	3
		Severe	1
		Moderate	20
		Mild	87
		Not Classifiable	11
	Reserve	Penetrating	1
		Severe	-
		Moderate	12
		Mild	41
		Not Classifiable	6

2001 as of May 15, 2015

Source: Defense Medical Surveillance System (DMSS), Theater Medical Data Store (TMDS) provided by the Armed Forces Health Surveillance Center (AFHSC)



Prepared by the Defense and Veterans Brain Injury Center (DVBIC)

2001, as of May 15, 2015

Service	Component	Severity	Diagnosed
Marines	Active	Penetrating	47
		Severe	37
		Moderate	365
		Mild	1,242
		Not Classifiable	138
	Reserve	Penetrating	2
		Severe	-
		Moderate	10
		Mild	70
		Not Classifiable	4

2001 as of May 15, 2015

Source: Defense Medical Surveillance System (DMSS), Theater Medical Data Store (TMDS) provided by the Armed Forces Health Surveillance Center (AFHSC)