

714 **MEDICATION GUIDE FOR NONSTEROIDAL ANTI-INFLAMMATORY DRUGS**
715 **(NSAIDS)**

716 (See the end of this Medication Guide
717 for a list of prescription NSAID medicines.)

718 **What is the most important information I should know about medicines called**
719 **Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)?**

720 **NSAID medicines may increase the chance of a heart attack or stroke that can lead**
721 **to death.** This chance increases:

- 722 • with longer use of NSAID medicines
723 • in people who have heart disease
724

725 **NSAID medicines should never be used right before or after a heart surgery called a**
726 **“coronary artery bypass graft (CABG).”**

727 **NSAID medicines can cause ulcers and bleeding in the stomach and intestines at any**
728 **time during treatment. Ulcers and bleeding:**

- 729 • can happen without warning symptoms
730 • may cause death
731

732 **The chance of a person getting an ulcer or bleeding increases with:**

- 733 • taking medicines called “corticosteroids” and “anticoagulants”
734 • longer use
735 • smoking
736 • drinking alcohol
737 • older age
738 • having poor health
739

740 **NSAID medicines should only be used:**

- 741 • exactly as prescribed
742 • at the lowest dose possible for your treatment
743 • for the shortest time needed
744

745 **What are Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)?**

746 NSAID medicines are used to treat pain and redness, swelling, and heat (inflammation)
747 from medical conditions such as:

- 748 • different types of arthritis
749 • menstrual cramps and other types of short-term pain
750

751 **Who should not take a Nonsteroidal Anti-Inflammatory Drug (NSAID)?**

752 **Do not take an NSAID medicine:**

- 753 • if you had an asthma attack, hives, or other allergic reaction with aspirin or any other
 754 NSAID medicine
- 755 • for pain right before or after heart bypass surgery
- 756

757 **Tell your healthcare provider:**

- 758 • about all of your medical conditions.
- 759 • about all of the medicines you take. NSAIDs and some other medicines can interact
 760 with each other and cause serious side effects. **Keep a list of your medicines to show**
 761 **to your healthcare provider and pharmacist.**
- 762 • if you are pregnant. **NSAID medicines should not be used by pregnant women late**
 763 **in their pregnancy.**
- 764 • if you are breastfeeding. Talk to your doctor.
- 765

766 **What are the possible side effects of Nonsteroidal Anti-Inflammatory Drugs**
 767 **(NSAIDs)?**

Serious side effects include:	Other side effects include:
<ul style="list-style-type: none"> • heart attack • stroke • high blood pressure • heart failure from body swelling (fluid retention) • kidney problems including kidney failure • bleeding and ulcers in the stomach and intestine • low red blood cells (anemia) • life-threatening skin reactions • life-threatening allergic reactions • liver problems including liver failure • asthma attacks in people who have asthma 	<ul style="list-style-type: none"> • stomach pain • constipation • diarrhea • gas • heartburn • nausea • vomiting • dizziness

768

769 **Get emergency help right away if you have any of the following symptoms:**

- 770 • shortness of breath or trouble breathing
- 771 • chest pain
- 772 • weakness in one part or side of your body
- 773 • slurred speech
- 774 • swelling of the face or throat
- 775

776 **Stop your NSAID medicine and call your healthcare provider right away if you have**
 777 **any of the following symptoms:**

- 778 • nausea
- 779 • more tired or weaker than usual
- 780 • itching
- 781 • your skin or eyes look yellow

- 782 • stomach pain
- 783 • flu-like symptoms
- 784 • vomit blood
- 785 • there is blood in your bowel movement or it is black and sticky like tar
- 786 • unusual weight gain
- 787 • skin rash or blisters with fever
- 788 • swelling of the arms and legs, hands and feet

789
790 These are not all the side effects with NSAID medicines. Talk to your healthcare provider
791 or pharmacist for more information about NSAID medicines.

792 **Other information about Nonsteroidal Anti-Inflammatory Drugs (NSAIDs):**

- 793 • Aspirin is an NSAID medicine but it does not increase the chance of a heart attack.
794 Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also
795 cause ulcers in the stomach and intestines.
- 796 • Some of these NSAID medicines are sold in lower doses without a prescription (over-
797 the-counter). Talk to your healthcare provider before using over-the-counter NSAIDs
798 for more than 10 days.

799
800 **NSAID medicines that need a prescription:**

Generic Name	Tradename
Celecoxib	Celebrex
Diclofenac	Cataflam, Voltaren, Arthrotec (combined with misoprostol)
Diflunisal	Dolobid
Etodolac	Lodine, Lodine XL
Fenoprofen	Nalfon, Nalfon 200
Flurbiprofen	Ansaid
Ibuprofen	Motrin, Tab-Profen, Vicoprofen* (combined with hydrocodone), Combunox (combined with oxycodone)
Indomethacin	Indocin, Indocin SR, Indo-Lemmon, Indomethagan
Ketoprofen	Oruvail
Ketorolac	Toradol
Mefenamic Acid	Ponstel
Meloxicam	Mobic

Nabumetone	Relafen
Naproxen	Naprosyn, Anaprox, Anaprox DS, EC-Naproxyn, Naprelan, Naprapac (copackaged with lansoprazole)
Oxaprozin	Daypro
Piroxicam	Feldene
Sulindac	Clinoril
Tolmetin	Tolectin, Tolectin DS, Tolectin 600

801 *Vicoprofen contains the same dose of ibuprofen as over-the-counter (OTC) NSAIDs, and is usually used
802 for less than 10 days to treat pain. The OTC NSAID label warns that long term continuous use may
803 increase the risk of heart attack or stroke.
804

805 This Medication Guide has been approved by the U.S. Food and Drug Administration.

806 Date created: June 15, 2005

807

808 Celebrex is a registered trademark of G.D. Searle LLC.

809 Cataflam, Voltaren are registered trademarks of Novartis Corporation.

810 Arthrotec (combined with misoprostol) is a registered trademark of G.D. Searle LLC.

811 Dolobid is a registered trademark of Merck & Co. Inc.

812 Lodine, Lodine XL are registered trademarks of Wyeth.

813 Nalfon, Nalfon 200 are registered trademarks of Pedinol Pharmacal Inc.

814 Ansaid is a registered trademark of Pharmacia & Upjohn Company LLC.

815 Motrin is a registered trademark of Johnson & Johnson.

816 Tab-Profen is a registered trademark of L. Perrigo Company.

817 Vicoprofen (combined with hydrocodone) is a registered trademark of BASF K & F
818 Corporation.

819 Combunox (combined with oxycodone) is a registered trademark of Forest Laboratories,
820 Inc.

821 Indocin, Indocin SR are registered trademarks of Merck & Co. Inc.

822 Oruvail is a registered trademark of Imperial Bank, As Agent (formerly registered to
823 Aventis Pharma S.A.).

824 Toradol is a registered trademark of Hoffmann-La Roche Inc.

825 Ponstel is a registered trademark of Lasalle National Bank Association.

826 Mobic is a registered trademark of Boehringer Ingelheim Pharma GMBG & Co. Kg.

827 Relafen is a registered trademark of SmithKline Beecham Corporation.

828 Naprosyn, EC-Naprosyn, Anaprox, Anaprox DS are registered trademarks of Syntex
829 Pharmaceuticals International Ltd.

830 Naprelan is a registered trademark of Elan Corporation PLC.

831 Naprapac (copackaged with lansoprazole) is a registered trademark of Syntex
832 Pharmaceuticals International Ltd.

833 Daypro is a registered trademark of G.D. Searle LLC.

834 Feldene is a registered trademark of Pfizer.

835 Clinoril is a registered trademark of Merck & Co. Inc.

836 Tolectin, Tolectin DS, Tolectin 600 are registered trademarks of Johnson & Johnson
837 Corporation.

838

839 Distributed by:



Pharmaceuticals

Roche Laboratories Inc.
340 Kingsland Street
Nutley, New Jersey 07110-1199

840

841

842 Month Year

843 Copyright © 1997-200x by Roche Laboratories Inc. All rights reserved