

GRAY SEAL (*Halichoerus grypus grypus*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

The gray seal is found on both sides of the North Atlantic, with three major populations: eastern Canada, northwestern Europe and the Baltic Sea (Katona *et al.* 1993). The western North Atlantic stock is equivalent to the eastern Canada population, and ranges from New Jersey to Labrador (Davies 1957; Mansfield 1966; Katona *et al.* 1993; Lesage and Hammill 2001; DiGiovanni, pers. comm., Riverhead Foundation for Marine Research and Preservation). This stock is separated by geography, differences in the breeding season, and mitochondrial DNA variation from the northeastern Atlantic stock (Bonner 1981; Boskovic *et al.* 1996; Lesage and Hammill 2001). There are three breeding concentrations in eastern Canada: Sable Island, Gulf of St. Lawrence, and along the coast of Nova Scotia (Lavigne and Hammill 1993). Tagging studies indicate that there is little intermixing between the two breeding groups (Zwanenberg and Bowen 1990) and, for management purposes, they are treated by the Canadian Department of Fisheries and Oceans (DFO) as separate stocks (Mohn and Bowen 1996). Outside the breeding period, there is overlap in the distribution of animals from the three colonies (Lavigne and Hammill 1993; Harvey *et al.* 2008; Breed *et al.* 2006, 2009, Hammill, pers. comm. DFO, Mont-Joli, Quebec, Canada). In the mid- 1980s, small numbers of animals and pupping were observed on several isolated islands along the Maine coast and in Nantucket-Vineyard Sound, Massachusetts (Katona *et al.* 1993; Rough 1995; Gilbert *et al.* 2005). In the late 1990s, a year-round breeding population of approximately 400+ animals was documented on outer Cape Cod and Muskeget Island (D. Murley, pers. comm., Mass. Audubon Society, Wellfleet, MA). In December 2001, NMFS initiated aerial surveys to monitor gray seal pup production on Muskeget Island and adjacent sites in Nantucket Sound, and Green and Seal Islands off the coast of Maine (Wood *et al.* 2007). To assess the stock structure of gray seals in the northwest Atlantic, tissue samples were collected from Canadian and US populations for genetic analyses (Wood *et al.* 2011). Based on examination of nine highly variable microsatellite loci, all individuals were placed into one population. This provides additional confirmation that recolonization by Canadian gray seals is the source of the U.S. population.




Figure 1. Approximate coastal range of gray seals. Isobaths are the 100-m, 1000-m, and 4000-m depth contours.

POPULATION SIZE

Current estimates of the total western Atlantic gray seal population are not available; although estimates of portions of the stock are available for select time periods. The size of the total Canadian population from 1969-2012 has been estimated using updated age-specific reproductive rate data, and accounting for higher pup mortality in the Gulf of St. Lawrence breeding colony due to years with poor ice condition (DFO, 2011d; Hammill *et al.* 2012). For Sable Island the 2012 pup production estimate is 67,000 (95% CI=56,000 to 85,000), the total population size estimate of 262,000 (95% CI 219,000-332,000). Model estimates for coastal Nova Scotia were 2,300 (95% CI

=1,100-3800) pups and a total population of 20,000 (95% CI= 17,000-23,000) in 2012. For the Gulf in 2012, pup production was estimated to be 7,000 (95% CI=2,900-15,200), and a total population of 49,000 (95% CI=27,000-102,000). The combined 2012 pup production is estimated to be 76,300 (95% CI=60,000-105,000), with a total population of 331,000 (95% CI=262,000-458,000). Difference between the total 2012 and 2010 (Thomas *et al.* 2011) estimates are due solely to differences in modeling approaches (DFO 2011d; Hammill *et al.* 2012). The new model estimates replace the 2010 pup production and total population estimates reported in Thomas *et al.* (2011). Average annual rates of total population increase were estimated to be 6% in the 1980s, 9% in the 1990s, and 6% in the 2000s. The authors note that these estimates should be treated with caution due to modeling and data concerns. In comparison to the pooled estimates, Bowen *et al.* (2003) reported that the Sable Island had been increasing by approximately 13% for nearly 40 years, but subsequently declined to 7% based on the 2004 pup production survey (Trzcinski *et al.* 2005; Bowen *et al.* 2007). The 2012 estimates suggest that the Sable Island population continued to increase at a rate of about 2.8% since 2010 (Hammill *et al.* 2012). Whereas, the coastal Nova Scotia and Gulf of St. Lawrence stocks do not appear to have shown much change in abundance since 2010 (DFO 2012, in review).

In U.S. waters, gray seals currently pup at five established colonies: Muskeget Island, Massachusetts, and Green Island, e, Seal Island, Matinicus Rock and Mount Desert Rock in Maine. Although white coated pups have stranded on eastern Long Island beaches, no pupping colonies have been detected in that region. Gray seals have been observed using the historic pupping site on Muskeget Island in Massachusetts since 1990. Pupping has taken place on Seal and Green Islands in Maine since at least the mid 1990s. Aerial survey data from these sites indicate that pup production is increasing. A minimum of 2,620 pups (Muskeget= 2,095, Green= 59, Seal= 466) were born in the U.S. in 2008 (Wood LaFond 2009). Table 2 summarizes single-day pup counts from the three U.S. pupping colonies from 2001/2002 to 2007/2008 pupping periods. The decrease in pup counts in some years is an artifact of survey timing and not indicative of true declines in those years. In recent years NMFS monitoring surveys have detected an occasional mother/pup (white coats) pair on both Monomoy Island and Nomans Land in Massachusetts. Some of the local breeders have been observed with brands and tags indicating they had been born on Sable Island, Canada (Rough 1995; L. Sette, pers. comm.). The increase in the number of gray seals observed in the U.S. is probably due to both natural increase and immigration.

Gray seals are also observed in New England outside of the pupping season. In April-May 1994 a maximum count of 2,010 was obtained for Muskeget Island and Monomoy combined (Rough 1995). Maine coast-wide surveys conducted during summer revealed 597 and 1,731 gray seals in 1993 and 2001, respectively (Gilbert *et al.* 2005). In March 1999 a maximum count of 5,611 was obtained in the region south of Maine (between Isles of Shoals, Maine and Woods Hole, Massachusetts) (Barlas 1999). In March 2011 a maximum count of 15,756 was obtained in southeastern Massachusetts coastal waters (NMFS unpubl. data). No gray seals were recorded at haul-out sites between Newport, Rhode Island and Montauk Pt., New York (Barlas 1999), currently several hundred gray seals have been recorded in surveys conducted off eastern Long Island (R. DiGiovanni, pers. comm).

Table 1. Summary of abundance estimates for the western North Atlantic gray seal: year, and area covered during each abundance survey, resulting total abundance estimate and 95% confidence interval.			
Month/Year	Area	N _{best}	CI
2012 ^a	Gulf of St Lawrence + Nova Scotia Eastern Shore + Sable Island	331,000	95% CI 263,000-458,000
^a These are model based estimates derived from pup surveys.			

Table 2. The number of pups observed on Muskeget, Seal, and Green Islands 2002-2008. Data are from aerial surveys ¹ . These are single-day counts, not estimates of total pup production (Wood LaFond 2009).			
Pupping Season	Muskeget Island	Seal Island	Green Island
2001-2	883	No data	34
2002-3	509	147	No data
2003-4	824	150	26
2004-5	992	365	33
2005-6	868	239	43
2006-7	1704	364	57
2007-8	2095	466	59
¹ Survey data for the 2008-9 through 2012-3 seasons have not been counted, including Matinicus Rock and Mount Desert Rock			

Minimum Population Estimate

Based on modeling, the total Canadian gray seal population was estimated to be 331,000 (95% CI 263,000-458,000) (Hammill *et al.* 2011d). Present data are insufficient to calculate the minimum population estimate for U.S. waters.

Current Population Trend

Gray seal abundance is likely increasing in the U.S. Atlantic Exclusive Economic Zone (EEZ), but the rate of increase is unknown. The population in eastern Canada was greatly reduced by hunting and bounty programs, and in the 1950s the gray seal was considered rare (Lesage and Hammill 2001). The Sable Island, Nova Scotia, population was less affected and has been increasing for several decades. Pup production on Sable Island increased exponentially at a rate of 12.8% per year between the 1970s and 1997 (Stobo and Zwanenburg 1990; Mohn and Bowen 1996; Bowen *et al.* 2003; Trzcinski *et al.* 2005; Bowen *et al.* 2007; DFO 2011a), but has declined to about 4% per year between 2007 and 2010, and 2.8% from 2010 to 2012 (DFO 2011a, 2012). The non-Sable Island population increased from approximately 25,000 in the mid-1980s to a peak of 71,500 in 2010 (Thomas *et al.* 2011). Modeling estimates of pup production increased from approximately 6,000 in 1985 to 17,400 in 2010 (Thomas *et al.* 2011). Approximately 70% of the western North Atlantic population is from the Sable Island stock. In the early 1990s pupping was established on Hay Island, off the Cape Breton coast (Lesage and Hammill 2001; Hammill *et al.* 2007, 2010).

Surveys of winter breeding colonies in Maine and on Muskeget Island may provide some measure of gray seal population trends and expansion in distribution. Sightings in New England increased during the 1980s as the gray seal population and range expanded in eastern Canada. Five pups were born at Muskeget in 1988. The number of pups increased to 12 in 1992, 30 in 1993, and 59 in 1994 (Rough 1995). In January 2002, 883 pups were counted on Muskeget Island and surrounding shoals (Wood Lafond 2009). In recent years NMFS monitoring surveys have detected an occasional mother/pup (white coats) pair on both Monomoy Island and Nomans Land. These observations continue the increasing trend in pup production reported by Rough (1995). The change in gray seal counts from southeastern Massachusetts (i.e., Monomoy, Muskeget and adjacent tidal bars) from 5,611 in spring 1999 to 15,756 in spring 2011 represents an annual increase of 8.6%, however, it has not been determined what proportion of the increase represents growth or immigration. For example, a few gray seals branded as pups on Sable Island in the 1970s and 2000s (Stobo and Zwanenburg 1990; C. den Heyer, pers. comm. DFO, Halifax) and satellite-tagged adults have been sighted in the Cape Cod region. Further, a branded female and pup were photographed on Seal Island, Maine in early January 2014 (unpubl. NMFS data).

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. Recent studies estimated the current annual rate of increase at 2.8% between 2010 and 2012 on Sable Island (DFO 2011d), continuing a decline in the rate of increase (Trzcinski *et al.* 2005; Bowen *et al.* 2007; Thomas *et al.* 2011). Overall, population growth in the three Canadian breeding herds appears to be leveling off (DFO 2011d). For purposes of this assessment, the maximum net productivity rate was assumed to be 0.12. This value is based on theoretical modeling showing that pinniped populations may not grow at rates much greater than 12% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is unknown. The maximum productivity rate is 0.12, the default value for pinnipeds. The recovery factor (F_R) for this stock is 1.0, the value for stocks of unknown status, but which are known to be increasing. PBR for the western North Atlantic gray seals in U.S. waters is unknown.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

For the period 2008-2012, the total estimated human caused mortality and serious injury to gray seals was 4,533 per year. The average was derived from five components: 1) 1,086 (CV=0.11) (Table 3) from the 2008-2012 U.S. observed fishery; 2) 9 from average 2008-2012 non-fishery related, human interaction stranding mortalities (NMFS unpublished data); 3) 403 from average 2008-2012 kill in the Canadian hunt (DFO in 2011d); 4) 90 from DFO scientific collections (DFO in 2011d.); and 5) 2,945 removals of nuisance animals in Canada (DFO 2011d). Analysis of bycatch rates from fisheries observer program records likely greatly under-represents sub-lethal fishery interactions. Photographic analysis of gray seals at haulout sites on Cape Cod, Massachusetts revealed 5-8% of seals

exhibited signs of entanglement (Sette *et al.* 2009).

New Serious Injury Guidelines

NMFS updated its serious injury designation and reporting process, which uses guidance from previous serious injury workshops, expert opinion, and analysis of historic injury cases to develop new criteria for distinguishing serious from non-serious injury (Angliss and DeMaster 1998; Andersen *et al.* 2008; NOAA 2012). NMFS defines serious injury as an “*injury that is more likely than not to result in mortality*”. Injury determinations for stock assessments revised in 2013 or later incorporate the new serious injury guidelines, based on the most recent 5-year period for which data are available.

Fishery Information

Detailed fishery information is given in Appendix III.

U.S.

Northeast Sink Gillnet

Annual estimates of gray seal bycatch in the Northeast sink gillnet fishery reflect seasonal distribution of the species and of fishing effort. Estimated annual mortalities (CV in parentheses) from this fishery were 618 (0.23) in 2008, 1,063(0.26) in 2009, 1,155(0.28) in 2010, 1,491 (0.22) in 2011, and 542 (0.19) in 2012 (Table 3; Orphanides 2013; Hatch and Orphanides 2014). There were 6, 8, 7, 9, and 1 unidentified seals observed during 2008-2012, respectively. Since 1997 unidentified seals have not been prorated to a species. This is consistent with the treatment of other unidentified mammals that do not get prorated to a specific species. Average annual estimated fishery-related mortality and serious injury to this stock attributable to this fishery during 2008-2012 was 974 gray seals (CV=0.12) (Table 3). The stratification design used is the same as that for harbor porpoise (Bravington and Bisack 1996).

Mid-Atlantic Gillnet

Gray seal interactions were first observed in this fishery in 2010, when nine gray seal and 2 unidentified seal mortalities were observed. In 2011, 1 unidentified seal and 2 gray seal mortalities were observed in this fishery. In 2012 one gray seal mortality was observed. Annual estimated fishery-related mortality and serious injury (CV in parentheses) to this stock attributable to this fishery was 267 (0.75) in 2010, 19 (0.60) in 2011, and 14 (0.98) in 2012 (Table 3; Orphanides 2013; Hatch and Orphanides 2014). Average annual estimated fishery-related mortality and serious injury to this stock attributable to this fishery during 2008-2012 was 60 gray seals (CV=0.67) (Table 3).

Northeast Mid-Water Trawl

One gray seal mortality was observed in 2012 in this fishery. An expanded bycatch estimate has not been generated. Until this bycatch estimate can be developed, the average annual fishery-related mortality and serious injury for 2008-2012 is calculated as 0.2 animals (1 animal /5 years).

Mid-Atlantic Mid-Water Trawl

One gray seal mortality was observed in 2010 in this fishery. An expanded bycatch estimate has not been generated. Until this bycatch estimate can be developed, the average annual fishery-related mortality and serious injury for 2008-2012 is calculated as 0.2 animals (1 animal /5 years).

Gulf of Maine Atlantic Herring Purse Seine Fishery

The Gulf of Maine Atlantic Herring Purse Seine Fishery is a Category III fishery. This fishery was not observed until 2003, and was not observed in 2006. No mortalities have been observed, but 15 gray seals were captured and released alive in 2004, 19 in 2005, 0 in 2007, 6 in 2008, 0 in 2009, 4 in 2010, 34 in 2011, and 33 in 2012. In addition, 5 seals of unknown species were captured and released alive in 2004, 2 in 2005, 1 in 2007, none in 2008-2010, 8 in 2011, and 0 in 2012.

Northeast Bottom Trawl

Vessels in the North Atlantic bottom trawl fishery, a Category III fishery under MMPA, were observed in order to meet fishery management, rather than marine mammal management needs. No mortalities were observed prior to 2005, when four mortalities were attributed to this fishery. No mortalities were observed in 2006. The estimated annual fishery-related mortality and serious injury attributable to this fishery was 0 between 2001 and 2004, and for 2006. Nine gray seal mortalities were attributed to this fishery in 2007, 4 in 2008, 5 in 2009, 9 in 2010, 19 in 2011,

and 8 in 2012. The average annual fishery-related mortality and serious injury for 2008-2012 is calculated as 33 (0.18) animals.

Mid-Atlantic Bottom Trawl

One gray seal mortality was observed in this fishery in 2009, 2 in 2011 and 1 in 2012 (Table 2). The average annual fishery-related mortality and serious injury for 2008-2012 is calculated as 19 (0.48) animals.

CANADA

Historically, an unknown number of gray seals have been taken in Newfoundland and Labrador, Gulf of St. Lawrence, and Bay of Fundy groundfish gillnets, Atlantic Canada and Greenland salmon gillnets, Atlantic Canada cod traps, and in Bay of Fundy herring weirs (Read 1994). In addition to incidental catches, some mortalities (e.g., seals trapped in herring weirs) were the result of direct shooting, and there were culls of about 1,700 animals annually during the 1970s and early 1980s on Sable Island (Anonymous 1986).

In 1996, observers recorded 3 gray seals (1 released alive) in Spanish deep-water trawl fishing on the southern edge of the Grand Banks (NAFO Areas 3) (Lens 1997). Seal bycatch occurred year-round, but interactions were highest during April-June. Many of the seals that died during fishing activities were unidentified. The proportion of sets with mortality (all seals) was 2.7 per 1,000 hauls (0.003).

Fishery	Years	Data Type ^a	Observer Coverage ^b	Observed Serious Injury ^c	Observed Mortality	Estimated Serious Injury	Estimated Mortality	Estimated Combined Mortality	Estimated CVs	Mean Annual Combined Mortality
Northeast Sink Gillnet ^c	08-12	Obs. Data, Weighout, Trip Logbook	.05, .04, .17, .19, .15	0, 0, 0, 0, 0	31, 52, 107, 222, 91	0, 0, 0, 0, 0	618, 1063, 1155, 1491, 542	618, 1063, 1155, 1491, 542	.23, .26, .28, .22, .19	974(0.12)
Mid-Atlantic Gillnet	08-12	Obs. Data, Trip Logbook, Allocated Dealer Data	.03, .03, .04, .02, .02	0, 0, 0, 0, 0	0, 0, 9, 2, 1	0, 0, 0, 0, 0	0, 0, 267, 19, 14	0, 0, 267, 19, 14	0, 0, .75, .60, .98	60 (0.67)
Northeast Bottom Trawl	08-12	Obs. Data, Trip Logbook	.08, .09, .16, .26, .17	0, 0, 0, 0, 0	4, 5, 9, 19, 8	0, 0, 0, 0, 0	16, 22, 30, 58, 37	16, 22, 30, 58, 37	.52, .46, .34, .25, .49	33 (0.18)
Mid-Atlantic Bottom Trawl	08-12	Obs. Data, Trip Logbook	.03, .05, .06, .08, .05	0, 0, 0, 0, 0	0, 3, 0, 3, 1	0, 0, 0, 0, 0	0, 38, 0, 25, 30	0, 38, 0, 25, 30	0, .70, 0, .57, 1.1	19 (0.48) ^d
Northeast Mid-water Trawl - Including Pair Trawl	08-12	Obs. Data, Trip Logbook	.199, .42, .53, .41, .45	0, 0, 0, 0, 0	0, 0, 0, 0, 1	0, 0, 0, 0, 0	0, 0, 0, 0, na	0, 0, 0, 0, na	0, 0, 0, 0, na	0.2 (na)
Mid-Atlantic Mid-water Trawl - Including Pair Trawl	08-12	Obs. Data, Trip Logbook	.13, .13, .25, .41, .21	0, 0, 0, 0, 0	0, 0, 1, 0, 0	0, 0, 0, 0, 0	0, 0, na, 0, 0	0, 0, na, 0, 0	0, 0, na, 0, 0	0.2 (na)
TOTAL										1086 (0.11)
<p>a. Observer data (Obs. Data) are used to measure bycatch rates, and the data are collected within the Northeast Fisheries Observer Program. The Northeast Fisheries Observer Program collects landings data (Weighout), and total landings are used as a measure of total effort for the sink gillnet fishery. Mandatory logbook (Logbook) data are used to determine the spatial distribution of fishing effort in the Northeast multispecies sink gillnet fishery.</p>										

b. The observer coverages for the Northeast sink gillnet fishery and the mid-Atlantic gillnet fisheries are ratios based on tons of fish landed. . North Atlantic bottom trawl mid-Atlantic bottom trawl, and mid-Atlantic mid-water trawl fishery coverages are ratios based on trips. Total observer coverage reported for bottom trawl gear and gillnet gear in the years 2010 -2012 includes traditional fisheries observers in addition to fishery monitors through the Northeast Fisheries Observer Program (NEFOP).
c. Since 1998, takes from pingered and non-pingered nets within a marine mammal time/area closure that required pingers, and takes from pingered and non-pingered nets not within a marine mammal time/area closure were pooled. The pooled bycatch rate was weighted by the total number of samples taken from the stratum and used to estimate the mortality. In 2008- 2012, respectively, 4, 13, 17, 125 and 54 takes were observed in nets with pingers. In 2008 -2012, respectively, 72, 27, 39, 90, 97, and 10 takes were observed in nets without pingers.
^d Fishery related bycatch rates for years 2008-2012 were estimated using an annual stratified ratio-estimator. These estimates replace the 2008-2011 annual estimates reported in the 2013 stock assessment report that were generated using a different method.
e. Serious injuries were evaluated for the 2008-2011 period using new guidelines (Waring <i>et al.</i> 2014, 2015)

Other Mortality

Canada: In Canada, gray seals were hunted for several centuries by indigenous people and European settlers in the Gulf of St. Lawrence and along the Nova Scotia eastern shore, and were locally extirpated (Laviguier and Hammill 1993). Between 1999 and 2012 the annual kill of gray seals by hunters in Canada was: 1999 (98), 2000 (342), 2001 (76), 2002 (126), 2003 (6), 2004 (0), 2005 (1073), 2006 (1,857) 2007 (1747), 2008 (1,471), 2009 (263), 2010 (58), and 2011 (215). (DFO 2003; 2008; 2009; 2011b; 2011d). DFO reports less than 10 gray seals were taken in commercial hunts in 2012 (<http://www.dfo-mpo.gc.ca/fm-gp/seal-phoque/faq-eng.htm>; accessed 11 January 2014). The traditional hunt of a few hundred animals is expected to continue off the Magdalen Islands and in other areas, except Sable Island where commercial hunting is not permitted (DFO 2003). DFO established a total allowable catch (TAC) of 12,000 gray seals for 2007 and 2008: 2,000 in the Gulf and 10,000 on the Scotian Shelf. The TAC for 2009 and 2010 was 50,000 seals, and for 2011 and 2012 it was set at 60,000. Since 2007, a small commercial hunt has taken place on Hay Island in Nova Scotia (<http://www.dfo-mpo.gc.ca/fm-gp/seal-phoque/faq-eng.htm>). The Hay Island TAC for 2010 was 2,220 (DFO 2011c), and for 2011 and 2012 it was set at 1,900 (<http://www.dfo-mpo.gc.ca/decisions/fm-2012-gp/atl-002-eng.htm>, accessed 27 February 2013)). The hunting of gray seals will continue to be prohibited on Sable Island (DFO 2011b).

Canada also issues personal hunting licenses which allow the holder to take six gray seals annually (Lesage and Hammill 2001; DFO 2011b). Hunting is not permitted during the breeding season and some additional seasonal/spatial restrictions are in effect (Lesage and Hammill 2001). Further, between 2005 and 2012 the lethal removal of nuisance seals was: 2005 (3105), 2006 (3437), 2007 (3373), 2008 (3334), 2009 (3381), 2010 (2933), 2011 (2076), and 2012 (3000) (DFO 2011b).

For scientific collections, DFO took 87, 320, and 90 animals, respectively in 2007, 2011, and 2012 (DFO 2012,).

U.S.: Gray seals, like harbor seals, were hunted for bounty in New England waters until the late 1960s (Katona, *et al.* 1993; Lelli, *et al.* 2009). This hunt may have severely depleted this stock in U.S. waters (Rough 1995; Lelli, *et al.* 2009). Other sources of mortality include human interactions, storms, abandonment by the mother, disease, and predation. Mortalities caused by human interactions include boat strikes, fishing gear interactions, power plant entrainment, oil spill/exposure, harassment, and shooting. Seals entangled in netting have been reported at several major haul-out sites in the Gulf of Maine.

From 2008 to 2012 468 gray seal stranding mortalities were recorded, extending from Maine to North Carolina (Table 4; NMFS unpublished data). Most stranding mortalities were in Massachusetts, which is the center of gray seal abundance in U.S. waters. Seventy-one (15.2%) of the total stranding mortalities showed signs of human interaction (21 in 2008, 14 in 2009, 12 in 2010, 20 in 2011 and 4 in 2012), 27 of which had some indication of fishery interaction (7 in 2008, 9 in 2009, 4 in 2010, 5 in 2011 and 2 in 2012). Ten gray seals are recorded in the NE stranding database during the 2008 to 2012 period as having been shot – one in Maine in 2009 and one in Maine and two in Massachusetts in 2010, 6 in Massachusetts in 2011 and none in 2012.

State	2008	2009	2010	2011	2012	Total
Maine	6 (1)	3	8 (4)	4 (2)	10(2)	31
New Hampshire	0	1 (1)	0	8 (1)	1 (1)	10

Massachusetts	53 (4)	52 (7)	43 (5)	89 (14)	38 (21)	274
Rhode Island	7	10 (2)	8 (3)	14 (2)	13 (5)	52
Connecticut	0	1(1)	0	2	0	3
New York	2 (2)	16 (7)	10 (7)	22 (6)	5 (3)	55
New Jersey	3	4	4 (1)	10	4	25
Delaware	1 (1)	0	0	0	0	1
Maryland	1	1	1	4 (2)	0	7
Virginia	1	2	1	1	0	5
North Carolina	1 (1)	1 (1)	1	2 (2)	0	5
Total	75 (9)	91 (19)	76 (20)	156 (29)	71 (32)	468
Unspecified seals (all states)	51	34	22	11	27	145
a. Mortalities include those which stranded dead, died at site, were euthanized, died during transport, or died soon after transfer to rehab.						

STATUS OF STOCK

Gray seals are not listed as threatened or endangered under the Endangered Species Act, and the western North Atlantic stock is not considered strategic under the Marine Mammal Protection Act. The level of human-caused mortality and serious injury in the U.S. Atlantic EEZ is unknown, but believed to be very low relative to the total stock size. The status of the gray seal population relative to OSP in U.S. Atlantic EEZ waters is unknown, but the stock's abundance appears to be increasing in Canadian and U.S. waters. The total U.S. fishery-related mortality and serious injury for this stock is low relative to the stock size in Canadian and U.S. waters and can be considered insignificant and approaching zero mortality and serious injury rate.

REFERENCES CITED

- Andersen, M. S., K. A. Forney, T. V. N. Cole, T. Eagle, R. Angliss, K. Long, L. Barre, L. Van Atta, D. Borggaard, T. Rowles, B. Norberg, J. Whaley and L. Engleby. 2008. Differentiating serious and non-serious injury of marine mammals: Report of the serious injury technical workshop. NOAA Tech. Memo. NMFS-OPR-39 94 pp.
- Angliss, R.P. and D.P. DeMaster. 1998. Differentiating serious and non-serious injury of marine mammals taken incidental to commercial fishing operations: Report of the serious injury workshop, 1-2 April 1997, Silver Spring, MD. NOAA Tech. Memo. NMFS-OPR-13. 48 pp.
- Anonymous. 1986. Seals and sealing in Canada. Rep. of the Royal Commission on Seals and Sealing, Vol. 1. Canadian Government Publishing Centre, Ottawa, Ontario, Canada, 65 pp.
- Barlas, M.E. 1999. The distribution and abundance of harbor seals (*Phoca vitulina concolor*) and gray seals (*Halichoerus grypus*) in southern New England, winter 1998-summer 1999. M.A. thesis. Graduate School of Arts and Sciences Boston University, Boston, MA. 52 pp.
- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Bonner, W.N. 1981. Grey seal *Halichoerus grypus* Fabricus, 1791. Pages 111-144 in: S.H. Ridgway and R.J. Harrison, (eds.) Handbook of marine mammals, Vol. 2: Seals. Academic Press, London.
- Boskovic, R., K.M. Kovacs, M.O. Hammill and B.N. White. 1996. Geographic distribution of mitochondrial DNA haplotypes in grey seals (*Halichoerus grypus*). Can. J. Zool. 74: 1787-1796.
- Bowen, W.D., J. McMillan and R. Mohn. 2003. Sustained exponential population growth of grey seals at Sable Island, Nova Scotia. ICES J. Mar. Sci. 60: 1265-1274.
- Bowen, W.D., J.I. McMillan and W. Blanchard. 2007. Reduced population growth of gray seals at Sable Island: Evidence from pup production and age of primiparity. Mar. Mamm. Sci. 23: 48-64.
- Bravington, M.V. and K.D. Bisack. 1996. Estimates of harbour porpoise bycatch in the Gulf of Maine sink gillnet fishery, 1990-1993. Rep. Int. Whal. Comm. 46: 567-574.

- Breed, G.A., W.D. Bowen, J.I. McMillan and M.L. Leonard. 2006. Sexual segregation of seasonal foraging habitats in a non-migratory marine mammal. *Proc. Royal Soc. B* 273:2319–2326.
- Breed, G.A., I.D. Jonsen, R.A. Myers, W.D. Bowen, and M.L. Leonard. 2009. Sex-specific, seasonal foraging tactics of adult grey seals (*Halichoerus grypus*) revealed by state-space analysis. *Ecology* 90(11):3209–3221.
- Davies, J.L. 1957. The geography of the gray seal. *J. Mamm.* 38: 297-310.
- deHart, P.A.P. 2002. The distribution and abundance of harbor seals (*Phoca vitulina concolor*) in the Woods Hole region. M.A. thesis. Graduate School of Arts and Sciences, Boston University, Boston, MA. 88 pp.
- DFO. 2003. Atlantic Seal Hunt: 2003-2005 management plan. Department of Fisheries and Oceans. Fisheries Resource Management – Atlantic, Ottawa, Ontario, Canada. 34 pp.
- DFO. 2008. Atlantic Seal Hunt: 2006-2010 management plan. Department of Fisheries and Oceans. Fisheries Resource Management – Atlantic, Ottawa, Ontario, Canada. 34 pp. Available at: http://www.dfo-mpo.gc.ca/seal-phoque/faq_e.htm#2.
- DFO, 2009. Frequently asked questions about Canada’s seal hunt. Department of Fisheries and Oceans. Available at: <http://www.dfo-mpo.gc.ca/fm-gp/seal-phoque/faq-eng.htm>
- DFO 2011a. Stock assessment of Northwest Atlantic grey seals (*Halichoerus grypus*), Canadian Science Advisory Secretariat (CSAS) Res. Doc. 2010/091 12 pp.
- DFO 2011b. 2011-2015 Integrated Fisheries Management Plan for Atlantic Seals, available at <http://www.dfo-mpo.gc.ca/fm-gp/seal-phoque/reports-rapports/mgtplan-planges20112015/mgtplan-planges20112015-eng.htm#c3.7.1>.
- DFO, 2011c. Frequently asked questions about Canada’s seal hunt. Department of Fisheries and Oceans. Available at: http://www.dfo-mpo.gc.ca/fm-gp/seal-phoque/faq-eng.htm#faq_4
- DFO, 2011d. Stock assessment of Northwest Atlantic grey seals (*Halichoerus Grypus*). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2010/091
- Gilbert, J.R., G.T. Waring, K.M. Wynne and N. Guldager. 2005. Changes in abundance and distribution of harbor seals in Maine, 1981-2001. *Mar. Mamm. Sci.* 21: 519-535.
- Hatch, J.M. and C.D. Orphanides 2014. Estimates of cetacean and pinniped bycatch in the 2012 New England sink and mid-Atlantic gillnet fisheries. Northeast Fish Sci Cent Ref Doc.14-02. 20 pp.
- Hammill, M.O. 2005. Abundance of Northwest Atlantic grey seals in the Gulf of St. Lawrence and along the Nova Scotia Eastern Shore. DFO Research Document 2005/036. Canadian Department of Fisheries and Oceans. Ottawa, Ontario. 11 pp. http://www.dfo-mpo.gc.ca/csas/Csas/Publications/ResDocs-DocRech/2005/2005_036_e.
- Hammill, M.O. and J.F. Gosselin. 2005. Pup production of non-Sable Island grey seals in 2004. DFO Research Document 2005/036. Canadian Department of Fisheries and Oceans. Ottawa, Ontario. 20 pp. http://www.dfo-mpo.gc.ca/csas/Csas/Publications/ResDocs-DocRech/2005/2005_033_e.htm
- Hammill, M.O., W.D.Bowen, and C. den Heyer. 2012. Northwest Atlantic grey seal population trends, 1960-2012. DFO Can. Sci. Advis. Sec. Res. Doc. 2012/169. iv + xx p. available at <http://www.dfo-mpo.gc.ca/csas>.
- Harvey, V., S.D. Côté, and M.O. Hammill 2008. The ecology of 3-D space use in a sexually dimorphic mammal. *Ecography* 31(3):371–380.
- Katona, S.K., V. Rough and D.T. Richardson. 1993. A field guide to whales, porpoises, and seals from Cape Cod to Newfoundland. Smithsonian Institution Press, Washington, DC. 316 pp.
- Laviguer, L. and M.O. Hammill. 1993. Distribution and seasonal movements of grey seals, *Halichoerus grypus*, born in the Gulf of St. Lawrence and eastern Nova Scotia shore. *Can. Field-Nat.* 107: 329-340.
- Lens, S. 1997. Interactions between marine mammals and deep water trawlers in the NAFO regulatory area. ICES [Int. Counc. Explor. Sea] C.M. 1997/Q:08:10.
- Lesage, V. and M.O. Hammill. 2001. The status of the grey seal, *Halichoerus grypus*, in the Northwest Atlantic. *Can. Field-Nat.* 115(4): 653-662.
- Lelli, B., D.E. Harris, and A-M Aboueissa. 2009. Seal bounties in Maine and Massachusetts, 1888 to 1962. *Northeast. Nat.* 16(2): 239-254.
- Mansfield, A.W. 1966. The grey seal in eastern Canadian waters. *Can. Audubon Mag.* 28: 161-166.
- Mohn, R. and W.D. Bowen. 1996. Grey seal predation on the eastern Scotian Shelf: Modeling the impact on Atlantic cod. *Can. J. Fish. Aquat. Sci* 53: 2722-2738.
- NOAA. 2012. Federal Register 77:3233. National Policy for Distinguishing Serious From Non-Serious Injuries of Marine Mammals. Available from: <http://www.nmfs.noaa.gov/op/pds/documents/02/238/02-238-01.pdf>.

- Orphanides, C. D. 2013. Estimates of cetacean and pinniped bycatch during 2010 and 2011 in the New England Sink Gillnet fishery, Mid-Atlantic Gillnet fishery, and two NMFS gillnet experiments. Northeast Fish Sci Cent Ref Doc. 13-13 38 pp. Available at: <http://nefsc.noaa.gov/publications/crd/crd1313/>.
- Read, A.J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. Rep. Int. Whal. Comm. (Special Issue) 15: 133-147
- Rough, V. 1995. Gray seals in Nantucket Sound, Massachusetts, winter and spring, 1994. Final report to Marine Mammal Commission. Contract T10155615 28 pp.
- Sette, L. A., G. T. Waring, K. Touhey, S. Sharp, B. Sharp, and S. Landry. 2009. Photographic surveys of entanglement occurrence at grey seal (*Halichoerus grypus*) and harbor seal (*Phoca vitulina*) haul-outs on Cape Cod, Massachusetts. Poster presentation, 18th Biennial Conference on the Biology of Marine Mammals
- Stobo, W.T. and K.C.T. Zwanenburg. 1990. Grey seal (*Halichoerus grypus*) pup production on Sable Island and estimates of recent production in the northwest Atlantic. Pages 171-184 in: W. D. Bowen, (ed.) Population biology of sealworm (*Pseudoterranova decipiens*) in relation to its intermediate and seal hosts. Can. Bull. Fish. Aquat. Sci. 222.
- Thomas, L., M.O. Hammill, and W.D. Bowen. 2011. Estimated size of the northwest Atlantic grey seal population, Canadian Science Advisory Secretariat (CSAS) Res. Doc. 2011/017 23 pp.
- Trzcinski, M.K., R. Mohn and W.D. Bowen. 2005. Estimation of grey seal population size and trends at Sable Island. DFO Research Document 2005/067. Canadian Department of Fisheries and Oceans. Ottawa, Ontario. 10pp. http://www.dfo-mpo.gc.ca/csas/Csas/Publications/ResDocs-DocRech/2005/2005_067_e.htm
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Waring, G.T., F.W. Wenzel M.C., E. Josephson, and M. Lyssikatos 2014. Serious injury determinations for small cetaceans and pinnipeds caught in commercial fisheries off the northeast U.S. coast, 2007–2011. Northeast Fish. Sci. Cent. Ref. Doc. 14-13. 26 p.
- Waring, G.T., M.C. Rossman and F.W. Wenzel. 2015 Serious injury determinations for small cetaceans and seals caught in commercial fisheries off the northeast U.S. coast, 2012. Northeast Fish. Sci. Cent. Ref. Doc. 15-12. 19 p.
- Wood LaFond, S. 2009. Dynamics of recolonization: a study of the gray seal (*Halichoerus grypus*) in the northeast U.S. Ph.D. Dissertation. University of Massachusetts, Boston. 83 p.
- Wood, S.A., S. Brault and J.R. Gilbert. 2007. 2002 aerial survey of grey seals in the northeastern United States. Pages 117-121 in: T. Haug, M. Hammill and D. Ólafsdóttir, (eds.) Grey seals in the North Atlantic and Baltic. NAMMCO Sci. Pub. 6, Tromsø, Norway.
- Wood, S.A., T.R. Frasier, B.A. McLeod, J.R. Gilbert, B.N. White, W.D. Bowen, M.O. Hammill, G.T. Waring, and S. Brault. 2011. The genetics of recolonization: an analysis of the stock structure of grey seals (*Halichoerus grypus*) in the northwest Atlantic. Can. J. Zool. 89:490-497.
- Zwanenberg, K.C.T. and W.D. Bowen. 1990. Population trends of the grey seal (*Halichoerus grypus*) in eastern Canada. Pages 185-197 in: W. D. Bowen, (ed.) Population biology of sealworm (*Pseudoterranova decipiens*) in relation to its intermediate and seal hosts. Can. Bull. Fish. Aquat. Sci. 222.