

PROPOSED MODIFIED BASE FLOOD ELEVATIONS—Continued

State	City/town/county	Source of flooding	Location	#Depth in feet above ground *Elevation in feet (NGVD)	
				Existing	Modified
Maps available for inspection at the City Hall, 1st and Washington, McAlester, Oklahoma. Send comments to The Honorable Randy Green, Manager of the City of McAlester, Pittsburg County, P.O. Box 578, McAlester, Oklahoma 74501.					
Washington.....	Benton (Unincorporated Areas).	Zintel Canyon.....	Approximately 400 feet northeast of the intersection of East 19th Avenue and Washington Street.	*378	*376
			Approximately 1,200 feet west of the intersection of East 23rd Street and South Gum Street.	*368	*368
			At the intersection of East 27th Avenue and South Myrtle Street.	*368	*366
			At the intersection of East 19th Avenue and South Oak Street.	*368	*366

Maps available for review at the Benton County Planning Department, 620 Market Street, Prosser, Washington.
Send comments to The Honorable Robert J. Drake, Sr., Chairman, Benton County Board of Commissioners, P.O. Box 190, Prosser, Washington 99350.

Issued: July 14, 1989.

Harold T. Duryee,

Administrator, Federal Insurance Administration.

[FR Doc. 17565 Filed 7-26-89; 8:45 am]

BILLING CODE 6718-03-M

DEPARTMENT OF THE INTERIOR

Fish and Wildlife Service

50 CFR Part 17

RIN 1018-AB31

Proposed Endangered or Threatened Status for Five Plants From the Southern San Joaquin Valley

AGENCY: Fish and Wildlife Service, Interior.

ACTION: Proposed rule.

SUMMARY: The U.S. Fish and Wildlife Service (Service) proposes endangered status pursuant to the Endangered Species Act of 1973 (Act), as amended, for four plants: *Caulanthus californicus* (California jewelflower), *Eremalche kernensis* (Kern mallow), *Lembertia congdonii* (San Joaquin wooly-threads), and *Opuntia treleasei* (Bakersfield cactus). The Service also proposes threatened status for one plant, *Eriastrum hooveri* (Hoover's wooly-star).

These species are restricted to grassland and adjacent plant communities (valley sink scrub, valley saltbush scrub, and juniper woodland) in the southern San Joaquin Valley, California, and neighboring foothills and valleys. The five plants have been variously affected and are threatened by one or more of the following: urbanization, conversion of native habitat for agriculture (ag-land conversion) and related water

development, oil and gas development and exploration, livestock grazing, competition from alien plants, utilization of habitat for ground-water recharge basins or for disposal of agricultural effluent or runoff, flood control projects, off-road vehicle use, mining, alteration of the natural fire regime, poor air quality, and stochastic extinction by virtue of the small isolated nature of the remaining populations. This proposal, if made final, would implement the Federal protection and recovery provisions afforded by the Act for these plants. The Service seeks data and comments from the public on this proposal.

DATES: Comments from all interested parties must be received by September 25, 1989. Public hearing requests must be received by September 11, 1989.

ADDRESS: Comments and materials concerning this proposal should be sent to the Field Supervisor, U.S. Fish and Wildlife Service, Endangered Species Office, 2800 Cottage Way, Room E-1823, Sacramento, California 95825. Comments and materials received will be available for public inspection, by appointment, during normal business hours at the above address.

FOR FURTHER INFORMATION CONTACT: Mr. Gail C. Kobetich, Field Supervisor, at the above address (916/978-4866 or FTS 460-4866).

SUPPLEMENTARY INFORMATION:

Background

Caulanthus californicus, *Eremalche kernensis*, *Eriastrum hooveri*, *Lembertia congdonii*, and *Opuntia treleasei* are endemic to grassland and adjacent plant communities [valley sink scrub, valley saltbush scrub, and juniper woodland (cf. Holland 1986)] of the southern San Joaquin Valley and neighboring foothills and valleys of California. This portion of

the San Joaquin Valley, often referred to as the Tulare Lake Basin, contains roughly 2.5 million acres of nearly flat, valley floor. If the neighboring valleys (i.e., Carrizo Plain, Cuyama Valley) and foothills are included with the Tulare Lake Basin, prehistoric grassland and adjacent plant communities likely totaled over 6 million acres. However, 96 percent of the native habitats of the valley floor has been lost principally to urbanization and ag-land conversion (Richard Anderson, California Energy Commission, pers. comm, July 21, 1987). The remaining non-urbanized or non-converted lands have been subject to livestock grazing, water development, oil and gas development and exploration, off-road vehicle use, mining, and/or other anthropogenic actions.

The prehistoric composition of the native grasslands and adjoining plant communities likely will remain a mystery (Brown 1982), although numerous authors have speculated as to the composition of the "pristine" flora of the Central Valley, inclusive of the San Joaquin Valley and Tulare Lake Basin (Clements 1934, Munz and Keck 1950, Biswell 1956, Twisselmann 1956, White 1967, McNaughton 1968, Bakker 1971, Ornduff 1974, Heady 1977, Bartolome and Gemmill 1981, and Wester 1981). Alien, annual grasses and forbs invaded the low-elevation, plant communities of California during the days of the Franciscan missionaries. Today, these grasses, which account for 50 to 90 percent of the vegetative cover (Heady 1956) and can stand up to a meter in height (Holland 1986), dominate most grasslands in California. Alien grasses have outcompeted the native flora throughout much of California because these exotics germinate in late fall prior to the germination of the native forbs,

including the four herbaceous species proposed herein (*Caulanthus californicus*, *Eremalche kernensis*, *Eriastrum hooveri*, and *Lembertia congdonii*). Consequently, these four herbs generally occupy sites with reduced grass cover. Although the stem succulent proposed herein (*Opuntia treleasei*) persists in areas largely dominated by alien plants (mostly annual grasses), the cactus does not necessarily prefer such "grassy" sites. The invasion of grasses has been quite thorough throughout much of the lower elevation portions of California. These exotics likely compete for nutrients and water, and may further threaten *Opuntia treleasei* by providing abundant fine (thin, slender) fuels, which probably increase the frequency and intensity of wildfires affecting the species' habitat.

The five plant taxa largely persist today in three native plant communities adjoining the non-native annual grasslands; valley sink scrub, valley saltbush scrub, or juniper woodland. However, these plant communities too have been affected somewhat by the presence of alien grasses. Valley sink scrub is an open to dense shrubland dominated by alkali-tolerant plants of the goosefoot family (Chenopodiaceae, so-called "chenopods"), like iodinebush (*Allenrolfea occidentalis*) and sea-blite (*Suaeda* spp.). This plant community, which generally lacks or produces a sparse understory of herbs, occurs about the margins of playas and on the heavy clays of the valley floor. Valley sink scrub essentially has been lost due to ag-land conversion, flood control projects, and ground water pumping (Holland 1986). Valley saltbush scrub, a scrubland of chenopods over a low understory of annual herbs, typically occurs on the gentle, rolling hills surrounding the Tulare Lake Basin on sandy to loamy soils. Similar activities, including oil and gas exploration and development, have adversely affected and threaten this plant community (Holland 1986). Juniper woodland, a compact woodland of California juniper (*Juniperus californicus*), often adjoins grassland sites immediately above the valley floor on gentle sloping terraces. Livestock grazing is the predominant activity influencing this community.

Discussion of the Five Species Proposed Herein for Listing Follows

Caulanthus californicus (California jewelflower) evidently was first collected by Mrs. A.E. Bush near Tulare, although the date and repository of this specimen are known (Taylor and Davilla 1986). Serano Watson, citing the Bush collection as the type, described the plant as *Stanfordia californica* in

1880. Although E.L. Greene (1891) had placed most species of *Caulanthus* within the genus *Streptanthus*, Edwin Payson (1923) transferred the species to the former genus. Dean Taylor and William Davilla (1986) discussed in detail the appropriate generic assignment for the jewelflower and concurred with I.A. Al-Shehbaz (1973) that the monotypic genus *Stanfordia* should be submerged within *Caulanthus*. *C. californicus*, a rosette-forming annual herb of the mustard family (Brassicaceae), grows to about 1 foot in height and produces several flowering branches. The leaves of the species have dry, wavy margins while its non-rosette leaves clasp the stem. The flowers are translucent white with purple to green tips. Its sword-shaped siliques (narrow, many-seeded pods) attain a length of 1 inch and width of about ¼ inch. The shape and size of siliques, together with an absence of hairs and an inflated stem, separate *C. californicus* from its closest relatives: *C. coulteri* var. *coulteri*, *C. coulteri* var. *lemmonii*, and *C. inflatus*. The species historically was distributed within the general area bounded by the present-day cities or communities of Coalinga and Fresno in Fresno County, New Cuyama in Santa Barbara County, and Bakersfield in Kern County (Taylor and Davilla 1986). Previously known from 40 sites, the plant now exists as one introduced population in Kern County and two natural populations (one in Santa Barbara County and another in San Luis Obispo County). The species was not known to occur naturally at the latter site. A recent status survey (Taylor and Davilla 1986) reported that intensive livestock grazing, ag-land conversion, and other anthropogenic activities likely extirpated *Caulanthus californicus* from Fresno, Kings, and Tulare Counties.

Eremalche kernensis (Kern mallow) was first collected by Carl Wolf in the Temblor Valley about 7 miles northwest of McKittrick along the Lost Hills Road in Kern County in 1937. Using his collection as the type, Wolf described *E. kernensis* in 1938. Although Phillip Munz (1959) at first placed all *Eremalche* in *Malvastrum* in his flora of California, he later concurred with the use of *Eremalche* for the California taxa in his supplement (Munz 1968). The species, a small annual herb of the mallow family (Malvaceae), typically develops an erect (rarely decumbent to prostrate) stem about 2 to 4 inches in height. The plant produces white to rose-pink or lavender, hollyhock-like flowers (Taylor and Davilla 1986). Although other characters (i.e., flower color, shape of the calyx

lobes, flower size) have been employed in the past (Wiggins 1951, Munz 1959, Leonelli 1986), differences in leaf shape, pubescence (hair type and density), color-spotting on the petal, and number of carpels (seed-bearing organs) per flower separate *E. kernensis* from other members of the genus. Contrary to Thomas Kearney (1959) and Robert Hoover (1970), Taylor and Davilla (1986) concluded that the species was valid and that morphologically similar plants often confused with *E. kernensis* were actually male-sterile *E. parryi*. Restricted to the eastern base of the Temblor Range, the species ranges from the vicinity of McKittrick to near Buttonwillow within valley saltbush scrub in Kern County (Taylor and Davilla 1986). Oil and gas development likely extirpated the species at the type locality while ag-land conversion probably eliminated one other population of *E. kernensis*. Because the remaining four populations exist near active oil and gas fields or in the vicinity of transmission corridors (Taylor and Davilla 1986), further oil and gas development in the area or transmission line maintenance or expansion likely would threaten these sites. The species, to a lesser degree, may be affected by ag-land conversion, livestock overgrazing, exotic plant competition, and off-road vehicle use.

Eriastrum hooveri (Hoover's woolly-star) was evidently first collected in 1935 by Gregory Lyons near Little Panoche Creek in Fresno County. However, Willis Jepson (1943), in describing the plant as *Huegelia hooveri*, cited a 1937 collection by Robert Hoover (the namesake for the specific epithet) as the type. Later Herbert Mason (1945) transferred the species along with the rest of the woolly-stars to *Eriastrum*. *E. hooveri*, an annual herb of the phlox family (Polemoniaceae), produces many wire-like branches and small (about ¼ inch across), white flowers. Standing about 2-3 inches tall, the species has grayish, fuzzy stems and is often branched (Taylor and Davilla 1986). Primarily flower size, and the ratio of corolla tube to the length of petal lobes separate the species from other *Eriastrum*, although stamen characteristics play a secondary role (Taylor and Davilla 1986). *E. hooveri* was historically distributed in the Temblor Range (Kern and San Luis Obispo Counties), Cuyama Valley (San Luis Obispo and Santa Barbara Counties), and in a discontinuous fashion within valley saltbush scrub and valley sink scrub from Fresno County south in the San Joaquin Valley (Taylor and Davilla 1986). Reportedly the

species never grew around the borders of the historic Tulare Lake (Kings County). Today, however, 24 percent of the historical and extant populations of the species, including the type locality (7 miles south of Shafter in Kern County), has been extirpated by various habitat modifications (Taylor and Davilla 1986). Ag-land conversion, urbanization, conversion of habitat "for ground-water recharge basins or disposal of nutrient-agricultural effluent", and oil and gas development threaten the remaining populations of the species (Taylor and Davilla 1986).

Lembertia congdonii (San Joaquin woolly-threads) was first collected by J.W. Congdon near Deer Creek in Tulare County. Using the Deer Creek collection as the type, Asa Gray described the species in 1883. Greene placed the plant in his newly-created, monotypic genus *Lembertia* in 1897. Although subsequent floras (i.e., Munz 1959, Abrams and Ferris 1960) included this species in the genus *Eatonella*, Taylor (1987) maintains that the species is sufficiently different from *Eatonella* and other relatives to warrant placement within a monotypic genus. This annual herb, a member of the sunflower family (Asteraceae), produces several, frequently-branching stems arising from the base. These white-wooly stems grow to about 10 inches in length and often trail on the ground. Aside from differences in growth habit, disk and ray flowers, and other minor characters, the presence of dimorphic achenes (one-seeded, indehiscent fruit) separate *L. congdonii* from its closest relative, *Eatonella nivea* from the Great Basin (Taylor 1987). Associated with valley saltbush scrub, only 12 populations of *L. congdonii* remain in the San Joaquin Valley and adjoining foothills from the vicinity of Panoche Pass (San Benito County) southeasterly to Caliente Creek east of Bakersfield (Kern County) (Taylor 1987). Another seven populations occur to the southwest in the Cuyama Valley (San Luis Obispo and Santa Barbara Counties) and Carrizo Plain (San Luis Obispo County). Primarily as a result of ag-land conversion, 33 populations or 63 percent of the 52 historical and extant populations of the species have been lost (Taylor 1987). Ag-land conversion, urbanization, gravel and sand extraction, oil and gas development, continued overgrazing, and off-road vehicle use threaten the remaining stands of *L. congdonii*.

Opuntia treleasei (Bakersfield cactus) evidently was first collected east of the community of Caliente in Kern County by William Trelease in 1892. After cultivating this collection in the Missouri

Botanical Garden, John Coulter (1896) described the species using this garden material as the type. James Toumey in Bailey's *Cyclopaedia of Horticulture* combined the species as a variety of the widespread *O. basilaris* in 1901. David Griffiths and Raleigh Hare described the long-spiny form of the species from along the Kern River bluffs as *O. treleasei* var. *kernii* in 1906. Although Munz (1959) and Lyman Benson (1969 and 1982) continued to treat the Bakersfield cactus as *O. basilaris* var. *treleasei*, recent work by Charlotte Chamberlain (U.S. Corps of Engineers 1986) concluded that the *O. treleasei* is morphologically distinct from *O. basilaris*. *O. treleasei*, a low-growing cactus (Cactaceae) that typically spreads to form extensive thickets, generally develops beavertail-like pads (flattened stems) 3 to 4 inches wide by 5 to 7 inches long. The areoles (eye-spots) are never depressed but flush with the pad surface or somewhat raised. All areoles have spines, although they vary in number and length. Unlike *O. basilaris*, the surface of the pads, which are nearly cylindrical at the base, is not papillate (covered with numerous small protuberances). Although the large magenta flowers of *O. treleasei* appear identical to *O. basilaris*, the characters cited above clearly separate these two taxa as species. Found chiefly within annual grassland on sandy to sandy-loam soils, the species historically grew atop the low hills northeast of Oildale southeasterly along the valley floor to the low foothills of the Tehachapi Mountains southeast and south-southwest of Arvin in Kern County. Charles Preuss, John C. Fremont's cartographer, wrote of this area in 1844, that "[t]he hilly country is bleak, without any vegetation except a beautiful species of cactus whose magnificent red blossoms grace this sad, sandy desert in a strange manner." Ernest Twisselmann (1969) claimed the species "once grew in dense almost impenetrable colonies on the mesas east of Bakersfield." A photograph in Britton and Rose (1920) attests to the species' former abundance. As late as 1937, biologists noted that the species produced a "thick growth" along Caliente Creek (Piemeisel and Lawson 1937). However, ag-land conversion (primarily for the production of potatoes and cotton), oil development, sand mining, urbanization, and perhaps wildfire have reduced this formerly widespread species to numerous, small isolated colonies. These colonies can be divided into five general population areas: the oilfields northeast of Oildale, Kern River Bluffs northeast of Bakersfield, the bluffs and rolling hills

west and north of Caliente Creek east of Bakersfield, Comanche Point on the Tejon Ranch southeast of Arvin, and northwest of the community of Wheeler Ridge. Off-road vehicle use, proposed flood control basins, and the activities cited above continue to threaten the remaining sites.

Federal government actions on these five plants began as a result of section 12 of the Endangered Species Act of 1973, which directed the Secretary of the Smithsonian Institution to prepare a report on those plants considered to be endangered, threatened, or extinct. This report designated as House Document No. 94-51, was presented to Congress on January 9, 1975. In the report, *Opuntia basilaris* var. *treleasei* was listed as an endangered species. On July 1, 1975 (40 FR 27823), the Service published a notice in the *Federal Register* of its acceptance of the report as a petition within the context of section 4(c)(2) [now section 4(b)(3)] of the Act, and of the Service's intention thereby to review the status of the plant taxa named within. *Opuntia basilaris* var. *treleasei* was included in that notice. On June 16, 1976, the Service published a proposed rule in the *Federal Register* (41 FR 24523) to determine approximately 1,700 vascular plant species to be endangered species pursuant to section 4 of the Act. The list of 1,700 plant taxa was assembled on the basis of comments and data received by the Smithsonian Institution and the Service in response to House Document No. 94-51 and the July 1, 1975, *Federal Register* publication. *Opuntia basilaris* var. *treleasei* was included in the proposed rule. General comments received in relation to the 1976 proposal were summarized in an April 26, 1978, *Federal Register* publication, which also determined 13 plant species to be endangered or threatened (43 FR 178909). On December 10, 1979, the Service published a notice of withdrawal of that portion of the June 16, 1976, proposal that had expired due to a procedural requirement of the 1978 Amendments. On December 15, 1980, the Service published a revised notice of review of native plants in the *Federal Register* (45 FR 82480); *Opuntia basilaris* var. *treleasei* was included as a category 1 candidate species (species for which data in the Service's possession indicate listing is warranted). On November 28, 1983, the Service published in the *Federal Register* (48 FR 53640) a supplement to the 1980 notice of review. This supplement added *Caulanthus californicus* as a category 2 candidate species (species for which data in the Service's possession indicate listing is probably appropriate, but for

which additional biological information is needed to support a proposed rule). Along with *Opuntia basilaris* var. *treleasei* in category 1, *Eremalche kernensis* and *Eriastrum hooveri* were included with *Caulanthus californicus* in category 2 in the September 27, 1985, revised notice of review for plants (50 FR 39526). This proposal to list *Caulanthus californicus*, *Eremalche kernensis*, *Lembertia congdonii*, and *Opuntia treleasei* as endangered and *Eriastrum hooveri* as threatened largely is based on status surveys conducted by Taylor and Davilla (1986) and Taylor (1987), field work carried out by Chamberlain (U.S. Army Corps of Engineers 1986) and Mike Foster (botanist, California Energy Commission, pers. comm., November 24, 1987, January 22, 1988), and pertinent literature (see "References Cited" below).

Section 4(b)(3)(B) of the Endangered Species Act, as amended in 1982, requires the Secretary to make findings on certain pending petitions within 12 months of their receipt. Section 2(b)(1) of the 1982 amendments further requires that all petitions pending on October 13, 1982, be treated as having been newly submitted on that date. This was the case for one of the southern San Joaquin Valley plants, *Opuntia treleasei*, because the 1975 Smithsonian report was accepted as a petition. In October 1983, 1984, 1985, 1986, 1987, and 1988, the Service found that the petitioned listing of *Opuntia treleasei* was warranted, but that the listing of this species was precluded due to other higher priority listing actions. Publication of the present proposal constitutes the next 1-year finding required by October 1988.

Summary of Factors Affecting the Species

Section 4(a)(1) of the Endangered Species Act (16 U.S.C. 1531 *et seq.*) and regulations promulgated to implement the listing provisions of the Act (50 CFR Part 424) set forth the procedures for adding species to the Federal lists. A species may be determined to be an endangered or threatened species due to one or more of the five factors described in section 4(a)(1). These factors and their application to *Caulanthus californicus* (Watson) Payson (California jewelflower); *Eremalche kernensis* C.B. Wolf (Kern mallow); *Eriastrum hooveri* (Jepson) H.L. Mason (Hoover's wooly-star); *Lembertia congdonii* (Gray) Greene (= *Eatonella congdonii* Gray) (San Joaquin wooly-threads); and *Opuntia treleasei* Coulter [= *Opuntia basilaris* Engelm. & Bigelow var. *treleasei* (Coulter) Toumey] (Bakersfield cactus) are as follows:

A. The present or threatened destruction, modification, or curtailment of its habitat or range. All five species proposed herein (*Caulanthus californicus*, *Eremalche kernensis*, *Eriastrum hooveri*, *Lembertia congdonii*, and *Opuntia treleasei*) are restricted to grassland and adjacent plant communities (valley sink scrub, valley saltbush scrub, and juniper woodland) in the southern San Joaquin Valley and neighboring foothills and valleys in California (see "Background" section for specific distributions). The primary threat facing these five species is the ongoing and threatened destruction and adverse modification of habitat. As discussed in the "Background" section, primarily ag-land conversion and urbanization have claimed 96 percent of the native habitats of the valley floor. The remaining non-urbanized or non-converted lands, which largely occur in the neighboring foothills and valleys (i.e., Carrizo Plain, Cuyama Valley), have been subject to livestock grazing, water development, oil and gas development and exploration, off-road vehicle use, mining, and/or other activities. These anthropogenic actions continue to threaten the native plant communities and habitats of these five species.

Caulanthus californicus was known from 40 sites in six counties (Fresno, Kern, Kings, San Luis Obispo, Santa Barbara, and Tulare), according to herbarium and field records detailed in the status survey by Taylor and Davilla (1986). Although once described as "abundant on the plains of the San Joaquin from Tulare southward (Greene 1891)," the species is known today from three sites: Two natural populations on private land (one near the mouth of Santa Barbara Canyon in Santa Barbara County and another in the southern portion of Carrizo Plain in San Luis Obispo County) and one introduced colony in Paul Paine Preserve (owned by The Nature Conservancy) in Kern County. Although the landowners of the Santa Barbara Canyon population have entered into voluntary agreements with The Nature Conservancy to protect the site (California Nature Conservancy 1987), no such agreement or protection exists for the recently-discovered population in San Luis Obispo County. Though no plants were observed at Santa Barbara Canyon in 1987 (Taylor, pers. comm., February 21, 1987), several thousand plants were counted in the spring of 1988. The Carrizo Plain population harbored a couple thousand individuals in 1988 (Mike Foster, pers. comm., March 14, 1988). Taylor noted only 24 plants at the introduced site, of

which only four plants flowered. Rainfall patterns probably account for the variation in population size for these colonies of *C. californicus*. Ag-land conversion likely claimed most of the valley floor sites due to the species' preference for sandy soils, which are prized for viticulture (Taylor and Davilla 1986). As suggested from herbarium records and the species' palatability, livestock grazing probably claimed the remaining extirpated sites within the last few decades (see Factor "D" for discussion). In addition, Taylor and Davilla (1986) speculate that poor air quality may have contributed to the demise of the species by promoting the growth of competing, pollution-tolerant plants (i.e., *Bromus rubens*).

Eremalche kernensis was known from six sites in western Kern County, according to herbarium and field records detailed in the status survey by Taylor and Davilla (1986). Oil and gas development likely extirpated the type locality of the species in the Temblor Valley. Another site of *E. kernensis*, 5 miles north of Lost Hills, was probably eliminated by ag-land conversion. In addition, construction of the California Aqueduct may have eliminated some unknown populations of the species. Three of the remaining four known occurrences exist on private land less than 5 miles from the South Belridge and Cymric Oil Fields and in the vicinity of transmission corridors (Taylor and Davilla 1986). Aside from maintenance or expansion of these corridors, future oil and gas development and exploration may threaten these remaining sites. One population north of McKittrick occurs on public land managed by the Bureau of Land Management (Bureau). Though the agency has not undertaken any special management of the site, the Bureau gives limited management consideration to candidate species. Nonetheless, this site still may be used for a variety of public uses (e.g., mineral extraction, oil and gas development, livestock grazing). All populations occur in areas grazed by sheep in the winter and spring. Consequently, Taylor and Davilla (1986) concluded, "[u]ncontrolled and heavy sheep grazing would be detrimental to *E. kernensis*."

Lembertia congdonii was known from 52 sites in seven counties (Fresno, Kern, Kings, San Benito, San Luis Obispo, Santa Barbara, and Tulare), according to herbarium and field records, and a recent status survey (Taylor 1987; Foster, pers. comm., March 14, 1988). Habitat alteration, principally due to ag-land conversion, eliminated 33 of these sites, including the type locality and

only known population in Tulare County. Of the remaining 19 sites, Taylor (1987) observed the species growing at six of these localities in either 1986 or 1987, while Foster (pers. comm., March 14, 1988) found an additional three populations in 1988. Population size ranged from 20 to 300 plants, the largest stand scattered over approximately 100 acres. Although no plants were located at the other 10 localities, Taylor (1987) reported that these sites still have suitable habitat. Although three of the 19 sites presumably harboring *L. congdonii* are on public land managed by the Bureau of Land Management, the agency has not undertaken any special management of these localities. Although the Bureau gives limited management consideration to candidate species, these sites still may be used for a variety of public uses (e.g., mineral extraction, oil and gas development, livestock grazing). Another population presumably still persists at Sand Ridge east of Bakersfield. Although The Nature Conservancy owns a 120-acre parcel on Sand Ridge, the northern portion of this area remains in private ownership. Off-road vehicle use, sand mining, and a flood control project proposed by the U.S. Army Corps of Engineers pose a potential threat to resources within this area. Portions of two populations were acquired by The Nature Conservancy as part of their Carrizo Plain Natural Heritage Preserve in early 1988. On August 30, 1988, the California Department of Water Resources purchased lands within the largely abandoned Strand and Canal Oil Fields, as part of the Kern Water Bank Project, that harbor the three populations found by Foster. The remaining portions of three sites owned in part by The Nature Conservancy and the other ten populations are privately owned and adjacent to lands that have been or continue to be urbanized, converted to agriculture, developed for oil and gas extraction and conveyance, or affected by off-road vehicles and grazing livestock. Similar activities are likely to continue in the near future.

Opuntia treleasei "once grew in dense almost impenetrable colonies on the mesas east of Bakersfield," according to Twisselman (1969). However, ag-land conversion (primarily for the production of potatoes and cotton), oil development, sand mining, urbanization, and perhaps wildfire have reduced this formerly widespread species to numerous, small isolated colonies. As discussed in the "Background" section, these colonies can be divided into five general population areas. Primarily oil

and gas development threaten the colonies northeast of Oildale, the northernmost population. Though this activity, to some degree, affects the population along the Kern River Bluffs northeast and east of Bakersfield, this area is rapidly being converted to housing for the ever-expanding population of Bakersfield. The construction of a small hydroelectric project and its associated accidental wildfire affected a few plants within the Kern River floodplain northeast of Bakersfield and east of Lake Ming. Off-road vehicle use, sand mining, and perhaps livestock overgrazing threaten the colonies on the bluffs and rolling hills west and north of Caliente Creek, the population located within the center of the species' range. Because the cactus provides no forage for livestock and competes with the alien grasses, ranchers may undertake eradication programs which may adversely affect the species. As discussed under *Lembertia congdonii*, The Nature Conservancy owns a portion of the Sand Ridge colony along the bluffs of Caliente Creek. However, a proposed flood control project likely will eliminate some individuals in the Sand Ridge area, including many plants on property owned by The Nature Conservancy. The Tejon Ranch, which is aware of the solitary clump of *O. Treleasei* on the ranch, has not expressed any plans to eliminate the cactus at Comanche Point. This population, however, is less than 4 miles from the Comanche Point Oil Field, which suggests the site may be subject to future oil and gas exploration. Ag-land conversion, aqueduct and transmission line maintenance, off-road vehicle use, urbanization, road widening, and illegal dumping threaten the remaining isolated colonies northwest of the community of Wheeler Ridge (Foster, pers. comm., January, 22, 1988), although one population grows on land owned by the State of California and administered by the California Department of Water Resources. In addition, the North Tejon Oil Field affects much of the Wheeler Ridge area.

Eriastrum hooveri was known from 49 sites in four counties (Fresno, Kern, San Luis Obispo, and Santa Barbara), according to individual reports (Taylor, pers. comm., November 30, 1987; Foster, pers. comm., November 24, 1987) and a recent status survey (Taylor and Davilla 1986). Primarily ag-land conversion and urbanization eliminated eleven of these sites. Of the remaining 38 sites, one population occurs on the Paul Paine Preserve, which The Nature Conservancy owns and protects. Another five sites presumably still exist

either in the Temblor Range or Alcalde Hills, either or which may harbor additional populations. Overgrazing poses the only imminent threat to these foothill populations. Although recent field survey work conducted by the California Department of Water Resources greatly expanded the area known by Taylor and Davilla (1986) to harbor the species along Warthan Creek in Fresno County (Arthur Gooch, pers. comm., July 22, 1988), this large population is threatened by the proposed Arroyo Pasajero Project. Cited as two populations in Taylor and Davilla (1986), the only populations known to occur on public land are from the Elk Hills on Naval Petroleum Reserve No. 1 (NPR-1) managed by the Department of Energy and on an adjacent parcel managed by the Bureau of Land Management. Inasmuch as Taylor and Davilla (1986) were unable to thoroughly survey NPR-1, the Department of Energy's consultant conducted a survey in 1988. Preliminary results reported finding 28 "populations" primarily along the northern and southern boundaries of NPR-1 (Thomas Kato, EG&G Energy Measurements, pers. comm., August 2, 1988). Though neither the Department of Energy nor the Bureau has undertaken any special management of these localities, the latter agency gives limited management consideration to candidate species. However, this policy does not necessarily prevent this population site from being used for a variety of public uses. Consequently, mineral extraction, oil and gas development, and livestock grazing may threaten these federally owned populations. One population, occurring on the Alkali Sink Ecological Preserve, grows on land owned by the State of California and managed by the California Department of Fish and Game. According to Taylor and Davilla (1986), the 29 remaining sites (including a portion of another population that occurs on both public and private land) occur on private property and typically on small, irregularly shaped parcels surrounded by ag-land and/or urban areas, which are often adjacent to roads. Seven of these sites harbor substantial populations (5,000 to 40,000 plants), while the remaining 23 sites consist of less than 1,000 individuals. All 29 populations occur on sites ranging from approximately an acre to less than 400 acres in size. Though many of these privately owned sites are perhaps too small to farm economically, parcels such as these continue to be converted to ag-land. Moreover, urbanization, conversion of habitat for ground-water recharge basins or disposal of nutrient-

laden agricultural effluent, off-road vehicle use, and oil and gas development continue to threaten these 29 populations (Taylor and Davilla 1986).

B. *Overutilization for commercial, recreational, scientific, or educational purposes.* Although not necessarily applicable to these species, many cacti are collected and cultivated by plant collectors, or offered for sale or trade by cactus growers. Though no data exist demonstrating such commerce in *Opuntia treleasei*, the species may still be collected and cultivated.

C. *Disease or predation.* As briefly mentioned above under Factor "A", livestock grazing probably extirpated colonies of *Caulanthus californicus* growing in the foothills and valleys adjoining the southern San Joaquin Valley. Moreover, trampling by livestock may have contributed to the endangerment of this species and *Eremalche kernensis*. Overgrazing may also threaten the other three species proposed herein.

D. *The inadequacy of existing regulatory mechanisms.* Under the Native Plant Protection Act (Chapter 1.5 section 1900 *et seq.* of the Fish and Game Code) and California Endangered Species Act (Chapter 1.5 section 2050 *et seq.*), the California Fish and Game Commission has listed one of these five species (*Caulanthus californicus*) as endangered (14 California Code of Regulations 670.2), while another species (*Opuntia treleasei*) is a State candidate (Stephen Nicola, pers. comm., September 23, 1988). Though both statutes prohibit the "take" of State-listed plants (Chapter 1.5 sections 1908 and 2080), Stat law appears to exempt the taking of such plants via habitat modification or land use change by the landowner. After the California Department of Fish and Game notifies a landowner that a State-listed plant grows on his or her property, State law evidently requires only that the landowner notify the agency "at least 10 days in advance of changing the land use to allow salvage of such plant." (Chapter 1.5 section 1913)

Opuntia treleasei, like all Cactaceae from the Americas not listed separately under Appendix I, is included under Appendix II of the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES). Although CITES regulates the international trade of listed species, the possibility of trade is not currently a threat to *Opuntia treleasei*.

E. *Other natural or manmade factors affecting its continued existence.* The invasion of alien, annual grasses has adversely affected all of the remaining

"natural" areas since the days of the Franciscan missionaries. These alien grasses, which account for 50 to 90 percent of the vegetative cover (Heady 1956) and can stand up to a meter in height (Holland 1986), largely dominate grasslands of California. As discussed in the "Background" section, the exotic annuals may alter the natural fire regime and these plants have either outcompeted or continue to compete with the native flora.

The Service has carefully assessed the best scientific and commercial information available regarding the past, present, and future threats faced by these species in determining to propose this rule. Based on this evaluation, the preferred action is to list *Caulanthus californicus*, *Eremalche kernensis*, *Lembertia congdonii*, and *Opuntia treleasei* as endangered, and to list *Eriastrum hooveri* as threatened.

Caulanthus californicus, *Eremalche kernensis*, *Lembertia congdonii*, and *Opuntia treleasei* have been extirpated from all but a small fraction of their historical ranges. Today these species generally persist as small, isolated populations or colonies surrounded by ag-land, urban areas, oil fields, and/or roads. Competition from alien grasses probably has and continues to adversely affect these species, especially the three annual herbs (*Caulanthus californicus*, *Eremalche kernensis*, and *Lembertia congdonii*). Although The Nature Conservancy owns most of the three remaining populations of *Caulanthus californicus* (California Nature Conservancy 1987), stochastic events affecting such extremely small populations still may result in the extinction of this species. All four remaining populations of *Eremalche kernensis* occur within a solitary township north of McKittrick, which may be adversely affected by livestock trampling, transmission corridor maintenance or expansion, and oil and gas development or exploration. The remaining 19 sites of *Lembertia congdonii* are variously threatened by ag-land conversion, urbanization, conversion of habitat for ground-water recharge basins, or disposal of agricultural effluent, livestock overgrazing, off-road vehicle use, and/or oil and gas development and exploration. Four populations of two of these species (*Eremalche kernensis* and *Lembertia congdonii*) occur on public land managed by the Bureau of Land Management, which accords limited management consideration to candidate species. However, this policy does not prevent the use of these sites for a variety of public uses (e.g., mineral extraction, oil and gas development,

livestock grazing). The relictual colonies of *Opuntia treleasei* are imminently threatened by ag-land conversion, oil development, sand mining, urbanization, off-road vehicle use, construction of flood control basins, aqueduct and transmission line maintenance, road widening, illegal dumping, and/or potentially alterations in the natural fire regime. Because these four plants are in danger of extinction throughout all or a significant portion of their ranges, they fit the definition of endangered as defined in the Act.

Eriastrum hooveri has been extirpated, principally as a result of ag-land conversion and urbanization, from 11 of its 49 known sites. Of the remaining 38 sites, one population is in preserve status and five sites presumably still exist either in the Temblor Range or the Alcalde Hills. Overgrazing poses the only imminent threat to these foothill populations. Two populations, for the most part, occur on public land, though the extent of the species of NPR-1 is under study. Regardless, these sites remain vulnerable to a variety of public uses (e.g., mineral extraction, oil and gas development, and livestock grazing). The 30 remaining parcels, including a portion of another population that occurs on both public and private land, are threatened by ag-land conversion, urbanization, conversion of habitat for ground-water recharge basins or disposal of agricultural effluent, off-road vehicle use, and oil and gas development and exploration (Taylor and Davilla 1986). Although the number of extant populations (38), including those located on NPR-1, provides greater flexibility in recovery and reduces the likelihood that the species will go extinct in the immediate future, the threats facing the 31 sites of *E. hooveri* on private property, at least in part, suggest that the species is likely to become an endangered species within the foreseeable future. Because of the limited threats facing the foothill populations of *E. hooveri* and the likelihood additional occurrences may be found in these upland areas, this species is not now in immediate danger of extinction throughout all or a significant portion of its range. However, if appropriate management actions are not taken, the species is likely to become in danger of extinction in the near future. As a result, *E. hooveri* fits the definition of threatened species as defined in the Act.

Critical Habitat

Section 4(a)(3) of the Act, as amended, requires that to the maximum extent

prudent and determinable, the Secretary designate critical habitat at the time a species is determined to be endangered or threatened. The Service finds that determination of critical habitat is not prudent for these species at this time. Because the five species face numerous anthropogenic threats (see Factor A in "Summary of Factors Affecting the Species") and occur predominantly on private land, the publication of precise maps and descriptions of critical habitat in the Federal Register would make these plants more vulnerable to incidents of vandalism and, therefore, could contribute to the decline of these species. The listing of these species as either endangered or threatened also publicizes the rarity of these plants and, thus, can make these plants attractive to researchers or collectors of rare plants. No Federal protection exists pursuant to section 9 of the Act for such take of listed plants on non-Federal lands. The proper agencies have been notified of the locations and management needs of these plants. Landowners will be notified of the location and importance of protecting habitat of these species. Protection of these species' habitats will be addressed through the recovery process and through the section 7 jeopardy standard. The Service believes that Federal involvement in the areas where these plants occur can be identified without the designation of critical habitat. Therefore, the Service finds that designation of critical habitat for these plants is not prudent at this time, because such designation likely would increase the degree of threat from vandalism, collecting, or other human activities.

Available Conservation Measures

Conservation measures provided to species listed as endangered or threatened under the Endangered Species Act include recognition, recovery actions, requirements for Federal protection, and prohibitions against certain practices. Recognition through listing encourages and results in conservation actions by Federal, State, and private agencies, groups, and individuals. The Endangered Species Act provides for possible land acquisition and cooperation with the States and requires that recovery actions be carried out for all listed species. Such actions are initiated by the Service following listing. The protection required of Federal agencies and the prohibitions against taking are discussed, in part, below.

Section 7(a) of the Act, as amended, requires Federal agencies to evaluate their actions with respect to any species that is proposed or listed as endangered

or threatened and with respect to its critical habitat, if any is being designated. Regulations implementing this interagency cooperation provision of the Act are codified at 50 CFR Part 402. Section 7(a)(4) requires Federal agencies to confer informally with the Service on any action that is likely to jeopardize the continued existence of a proposed species or result in destruction or adverse modification of proposed critical habitat. If a species is subsequently listed, section 7(a)(2) requires Federal agencies to ensure that activities they authorize, fund, or carry out are not likely to jeopardize the continued existence of such a species or to destroy or adversely modify its critical habitat. If a Federal action may affect a listed species or its critical habitat, the responsible Federal agency must enter into formal consultation with the Service. Seven populations of three of the proposed plant species occur on Federal (public) land; one population of *Eremalche kernensis* is managed by the Bureau of Land Management, two populations of *Eriastrum hooveri* are administered respectively by the Bureau of Land Management and Department of Energy, and four populations of *Lembertia congdonii* are managed by the Bureau of Land Management. Though some other stands occur near Federal land, all of the remaining known sites are on private land with no known Federal involvement with the following exceptions. The U.S. Army Corps of Engineers and the Bureau of Reclamation may fund or develop, at least in part, proposed flood control or water projects. Because of potential impacts to two federally listed animals, the San Joaquin kit fox (*Vulpes macrotis mutica*) and the bluntnosed leopard lizard (*Gambelia silus*), the Corps has consulted formally on a proposed flood control project for Caliente Creek. This project would potentially eliminate numerous individual plants of *Opuntia treleasei* from the Sand Ridge colony, which grows on the bluffs adjoining the creek (U.S. Army Corps of Engineers 1986). Aside from the direct effects of this and other similar proposed projects, other potential Federal actions include new allocations of water from existing Federal projects, which could increase ag-land conversion and possibly affect one or more of these five plant species. Activities involving Federal mortgage programs, including those of the U.S. Department of Agriculture (Farmers Home Administration), Veterans Administration, and U.S. Department of Housing and Urban Development (Federal Home Administration loans), may be subject to section 7 review.

The Act and its implementing regulations found at 50 CFR 17.61, 17.62, and 17.63 for endangered species and 17.71 and 17.72 for threatened species set forth a series of general trade prohibitions and exceptions that apply to all endangered and threatened plant species. With respect to the five plants from the southern San Joaquin Valley, all trade prohibitions of section 9(a)(2) of the Act, implemented by 50 CFR 17.61 and 17.71, would apply. These prohibitions, in part, make it illegal for any person subject to the jurisdiction of the United States to import or export, transport in interstate or foreign commerce in the course of a commercial activity, sell or offer for sale these species in interstate or foreign commerce, or to remove and reduce to possession these species from areas under Federal jurisdiction. Seeds from cultivated specimens of threatened plant species are exempt from these prohibitions provided that a statement of "cultivated origin" appears on their containers. Certain exceptions can apply to agents of the Service and State conservation agencies. The Act and 50 CFR 17.62, 17.63, and 17.72 also provide for the issuance of permits to carry out otherwise prohibited activities involving endangered and threatened plant species under certain circumstances. The Service anticipates few trade permits would ever be sought or issued for the five species, with the possible exception of *Opuntia treleasei* which, like other cacti, may be in cultivation. Requests for copies of the regulations on plants and inquiries regarding them may be addressed to the Office of Management Authority, U.S. Fish and Wildlife Service, Washington, DC 20240 (703/343-4955).

As a species of Cactaceae (Cactus family), *Opuntia treleasei* is included in Appendix II of the CITES Convention (See 50 CFR 23.23). The effect of this protection under the CITES Convention is that an export permit must be issued by the country of origin, or a re-export certificate must be issued by the country of re-export prior to the importation of *Opuntia treleasei*. Such CITES Convention restrictions are intended to prevent international trade from being detrimental to the survival of the species.

Public Comments Solicited

The Service intends that any final action resulting from this proposal will be as accurate and as effective as possible. Therefore, any comments or suggestions from the public, other concerned governmental agencies, the scientific community, industry, or any

other interested party concerning any aspect of this proposal are hereby solicited. Comments particularly are sought concerning:

(1) Biological, commercial trade, or other relevant data concerning any threat (or lack thereof) to *Caulanthus californicus*, *Eremalche kernensis*, *Eriastrum hooveri*, *Lembertia congdonii*, or *Opuntia treleasei*;

(2) The location of any additional populations of these species and the reasons why any habitat should or should not be determined to be critical habitat as provided by Section 4 of the Act;

(3) Additional information concerning the range and distribution of these species; and

(4) Current or planned activities in the ranges and habitats of these species and their possible impacts on these species.

Final promulgation of the regulations on these species will take into consideration the comments and any additional information received by the Service, and such communications may lead to adoption of final regulations that differ from this proposal.

The Endangered Species Act provides for a public hearing on this proposal, if requested. Requests must be filed within 45 days of the date of the proposal. Such requests must be made in writing and addressed to Field Supervisor (see ADDRESS section).

National Environmental Policy Act

The Fish and Wildlife Service has determined that an Environmental Assessment, as defined by the National Environmental Policy Act of 1969, need not be prepared in connection with regulations adopted pursuant to Section 4(a) of the Endangered Species Act of 1973, as amended. A notice outlining the Service's reasons for this determination was published in the *Federal Register* on October 25, 1983 (48 FR 49244).

References Cited

- Abrams, L., and R.S. Ferris. 1960. Illustrated flora of the Pacific states. Vol. IV. Stanford Univ. Press.
- Al-Shehbaz, I.A. 1973. The biosystematics of the genus *Thelypodium* (Cruciferae). *Contr. Gray Herb.* 204:1-148.
- Bakker, E.S. 1971. An island called California. Univ. Calif. Press, Berkeley.
- Bartolome, J.W., and B. Gemmill. 1981. The ecological status of *Stipa pulchra* (Poaceae) in California. *Madroño* 28:172-184.
- Benson, L. 1969. The native cacti of California. Stanford Univ. Press.
- Benson, L. 1982. The cacti of the United States and Canada. Stanford Univ. Press.
- Biswell, H.H. 1956. Ecology of California grasslands. *J. Range Manage.* 9:19-24.

- Britton, N.L., and J.N. Rose. 1920. The Cactaceae, 4 vol., Carnegie Inst. Washington Publ. 248. Washington, D.C.
- Brown, D.E. 1982. 143.2 Californian valley grassland. *Desert Plants* 4:132-135.
- California Nature Conservancy. 1987. Landowners protect rare wildflower. *Autumn Newsletter*:3.
- Clements, F.E. 1934. The relict method in dynamic ecology. *J. Ecology* 22:39-68.
- Coulter, J.M. 1896. Preliminary revision of the North American species of *Echinocactus*, *Cereus*, and *Opuntia*. *Contr. U.S. Nat. Herb.* 3:434.
- Gray, A. 1883. Contributions to North American botany. *Proc. Amer. Acad. Arts Sci.* 19:1-96.
- Greene, E.L. 1891. fl. Fran. 256.
- Greene, E.L. 1897. Fl. Fran. 441.
- Griffiths, D., and R.F. Hare. 1906. Prickly pear and other cacti. *N. Mex. Agr. Exp. Sta. Bull.* 60:81.
- Heady, H.F. 1956. Evaluation and measurement of the California annual type. *J. Range Manage.* 9:25-27.
- Heady, H.F. 1977. Valley grassland. In: Barbour, M.G., and J. Major (eds.), *Terrestrial vegetation of California*. John Wiley and Sons, New York.
- Holland, R.F. 1986. Preliminary descriptions of the terrestrial natural communities of California. Unpublished report, California Department of Fish and Game, 156 pp.
- Hoover, R.F. 1970. The vascular plants of San Luis Obispo County, California. Univ. Calif. Press, Berkeley.
- Jepson, W.L. 1943. A flora of California. Vol. III. Jepson Herbarium and Library, Univ. Calif., Berkeley.
- Kearney, T.H. 1956. Notes on Malvaceae. VIII. *Eremalche*. *Madroño* 13:241-243.
- Leonelli, S. 1988. An investigation of the taxonomic status of *Eremalche kernensis* C.B. Wolf (Malvaceae). Unpublished M.S. thesis, Calif. State Univ., Long Beach.
- Mason, H.L. 1945. The genus *Eriastrum* and the influence of Benth and Hooker upon the problem of generic confusion in the Polemoniaceae. *Madroño* 8:65-91.
- McNaughton, S.J. 1968. Structure and function in California grasslands. *Ecology* 49:962-972.
- Munz, P.A. 1959. A California flora. Univ. Calif. Press, Berkeley.
- Munz, P.A., and D.D. Keck. 1950. California plant communities-supplement. *Aliso* 2:199-202.
- Munz, P.A., and D.D. Keck. 1950. California plant communities-supplement. *Aliso* 2:199-202.
- Ornduff, R. 1974. An introduction to California plant life. Univ. Calif. Press, Berkeley.
- Payson, E.B. 1923. A monograph of *Thelypodium* and its immediate allies. *Ann. Missouri Bot. Gard.* 9:233-324.
- Piemeisel, R.L., and F.R. Lawson. 1937. Types of vegetation in the San Joaquin Valley of California and their relation to the beet leafhopper. U.S.D.A. Tech. Bull. No. 557, Washington, D.C.
- Preuss, C. 1844. Exploring with Fremont. Taylor, D.W. 1987. Status survey of San Joaquin woolly-threads (*Lembertia congdonii*). U.S. Fish and Wildlife Service, Sacramento, Calif.
- Taylor, D.W., and W.B. Davilla. 1986. Status survey of three plants endemic to the San Joaquin Valley. U.S. Fish and Wildlife Service, Sacramento, Calif.
- Toumey, J.W. 1901. *Opuntia*. In: Bailey, Cyclop. Hort. 4:1147.
- Twisselmann, E.C. 1956. Flora of Temblor Range and the neighboring part of the San Joaquin Valley. *Wasmann J. Biol.* 14:161-300.
- Twisselmann, E.C. 1969. Status of the rare plants of Kern County. Calif. Native Plant Soc. Newsletter 5(3):1-7.
- U.S. Corps of Engineers. 1986. Caliente Creek stream group investigation. Unpublished report, U.S. Corps of Engineers, Sacramento, Calif.
- Wester, L. 1981. Composition of native grasslands in the San Joaquin Valley, California. *Madroño* 28:231-241.
- White, K.L. 1967. Native bunch grass (*Stipa pulchra*) on Hastings Reservation, California. *Ecology* 48:949-955.
- Wiggins, I.L. 1951. Malvaceae. In: Abrams, L., illus. *Flora Pacific States*. III. Stanford Univ. Press.
- Wolf, C.B. 1938. California plant notes II. Occasional Papers Rancho Santa Ana Bot. Gard. Ser. I, No. 2:64-68.

Author

The primary author of this proposed rule is Jim A. Bartel, Endangered Species Office, U.S. Fish and Wildlife Service, 2800 Cottage Way, Room E-1823, Sacramento, California 95825 (916/978-4866, FTS 460-4866).

List of Subjects in 50 CFR Part 17

Endangered and threatened wildlife, Fish, Marine mammals, Plants (agriculture).

Proposed Regulations Promulgation

Accordingly, it is hereby proposed to amend Part 17, Subchapter B of Chapter I, Title 50 of the Code of Federal Regulations, as set forth below:

PART 17—[AMENDED]

1. The authority citation for Part 17 continues to read as follows:

Authority: Pub. L. 93-205, 87 Stat. 884; Pub. L. 94-359, 90 Stat. 911; Pub. L. 95-632, 92 Stat. 3751; Pub. L. 96-159, 93 Stat. 1225; Pub. L. 97-304, 96 Stat. 1411; Pub. L. 100-478, 102 Stat. 2306; Pub. L. 100-653, 102 Stat. 3825 (16 U.S.C. 1531 *et seq.*); Pub. L. 99-625, 100 Stat. 3500 (1986), unless otherwise noted.

2. It is proposed to amend § 17.12(h) by adding the following, in alphabetical order under the families indicated, to the List of Endangered and Threatened Plants:

§ 17.12 Endangered and threatened plants.

* * * * *

(h) * * *

Species		Historic range	Status	When listed	Critical habitat	Special rules
Scientific name	Common name					
Brassicaceae—Mustard Family:						
<i>Caulanthus californicus</i>	California jewelflower	U.S.A. (CA)	E		NA	NA
Malvaceae—Mallow Family:						
<i>Eremalche kernensis</i>	Kern mallow	U.S.A. (CA)	E		NA	NA
Polemoniaceae—Phlox family:						
<i>Eriastrum hooveri</i>	Hoover's wooly-star	U.S.A. (CA)	T		NA	NA
Asteraceae—Aster family:						
<i>Lembertia congodonii</i>	San Joaquin wooly-threads	U.S.A. (CA)	Ed		NA	NA
Cactaceae—Cactus family:						
<i>Opuntia treleasei</i>	Bakersfield cactus	U.S.A. (CA)	E		NA	NA

Dated: June 12, 1989.
 Susan Recca Lamson,
 Acting Assistant Secretary for Fish and
 Wildlife and Parks.
 [FR Doc. 89-17595 Filed 7-26-89; 8:45 am]
 BILLING CODE 4310-55-M

50 CFR Part 17

RIN 1018-AB31

Endangered and Threatened Wildlife and Plants; Proposal to List the Purple Cat's Paw Pearly Mussel as an Endangered Species

AGENCY: Fish and Wildlife Service, Interior.

ACTION: Proposed rule.

SUMMARY: The Service proposes to list a freshwater mussel, the purple cat's paw pearly mussel (*Epioblasma (=Dysnomia) obliquata obliquata (=E. sulcata sulcata)*), as an endangered species under the Endangered Species Act of 1973, as amended (Act). This freshwater mussel historically occurred in the Ohio River and its large tributaries in Ohio, Indiana, Illinois, Kentucky, Tennessee, and Alabama. Presently the purple cat's paw pearly mussel is known from only two relic, apparently nonreproducing populations—one in a reach of the Cumberland River in Tennessee and one in a reach of the Green River in Kentucky. The distribution and reproductive capacity of this species have been seriously impacted by the construction of impoundments on the large rivers it once inhabited. Unless reproducing populations are found or methods developed to maintain existing

populations, this species will likely become extinct in the foreseeable future. Comments and information are sought from the public concerning this proposal. DATES: Comments from all interested parties must be received by September 25, 1989. Public hearing requests must be received by September 11, 1989.

ADDRESS: Comments and materials, and requests for public hearing concerning this proposal should be sent to the Field Supervisor, U.S. Fish and Wildlife Service, Asheville Field Office, 100 Otis Street, Room 224, Asheville, North Carolina 28801. Comments and materials received will be available for public inspection, by appointment, during normal business hours at the above address.

FOR FURTHER INFORMATION CONTACT: Mr. Richard G. Biggins at the above address (704/259-0321 or FTS 672-0321).

SUPPLEMENTARY INFORMATION:

Background

The purple cat's paw pearly mussel (*Epioblasma (=Dysnomia) Obliquata obliquata (=E. sulcata sulcata)*), was described by Rafinesque (1820). The white cat's paw (*Epioblasma (=Dysnomia) sulcata delicata*), the northern subspecies of the cat's paw pearly mussel known from the Lake Erie system of the St. Lawrence drainage, was listed as endangered on June 14, 1976 (41 FR 24064). The purple cat's paw, which is characterized as a large river species (Bates and Dennis 1985), has a medium-size shell that is subquadrate in outline (Bogan and Parmalee 1983). The shell has fine, faint, wavy green rays with a smooth and shiny surface. The inside of the shell is purplish to deep

purple (the inside shell of the white cat's paw is white). Like other freshwater mussels, the purple cat's paw feeds by filtering food particles from the water. It has a complex reproductive cycle in which the mussel's larvae parasitize fish. The mussel's life span, fish species its larvae parasitize, and other aspects of its life history are unknown.

The purple cat's paw pearly mussel was historically distributed in the Ohio, Cumberland, and Tennessee River systems in Ohio, Illinois, Indiana, Kentucky, Tennessee, and Alabama (Bogan and Parmalee 1983, Isom *et al.* 1979, Kentucky Nature Preserves Commission 1980, Parmalee *et al.* 1980, Watters 1986, Stansbery 1970). Based on personal communication with knowledgeable experts (Steven Ahlstedt and John Jenkinson, Tennessee Valley Authority, 1987; Mark Gordon and Robert Anderson, Tennessee Technological University, 1988; Arthur Bogan, Philadelphia Academy of Sciences, 1988; Ronald Cicerello, Kentucky Nature Preserves Commission, 1988; David Stansbery, Ohio State University, 1987) and a review of current literature, the species is known to survive in only two river reaches, but apparently as nonreproducing populations. These are located in the Cumberland River, Smith County, Tennessee, and the Green River, Warren and Butler Counties, Kentucky.

The continued existence of these two populations is questionable. Unless reproducing populations can be found or methods can be developed to maintain these or create new populations, the species will become extinct in the foreseeable future. Any individuals that do still survive in these two river