

# Recovery Plan for Two Plants from Rota



*Osmoxylon  
mariannense*



*Nesogenes  
rotensis*

**Recovery Plan**  
**for Two Plants from Rota**  
**(*Nesogenes rotensis* and *Osmoxylon mariannense*)**


**Region 1**

**U.S. Fish and Wildlife Service**

**Portland, Oregon**

Approved: \_\_\_\_\_

**Acting**



**David J. Wesley**

Regional Director, Region 1, U.S. Fish and Wildlife Service

Date: \_\_\_\_\_

**FEB 14 2007**

## DISCLAIMER

Recovery plans delineate reasonable actions that are believed to be required to recover and protect listed species. We, the U.S. Fish and Wildlife Service, prepare and publish recovery plans, sometimes with the assistance of recovery teams, contractors, State agencies, Tribal agencies, and other affected and interested parties. The objectives of recovery plans will be attained and any necessary funds made available subject to budgetary and other constraints affecting the parties involved, as well as the need to address other priorities. Recovery plans do not obligate other parties to undertake specific actions and may not represent the views nor the official positions or approval of any individuals or agencies involved in recovery plan formulation, other than our own. They represent our official position *only* after they have been signed by the Regional Director or Director as *approved*. Recovery plans are reviewed by the public and submitted to peer review before we adopt them as approved final documents. Approved recovery plans are subject to modification as dictated by new findings, changes in species status, and the completion of recovery actions.

### **Literature citation of this document should read as follows:**

U.S. Fish and Wildlife Service. 2007. Recovery Plan for Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*). Portland, Oregon. 86 pp.

### **An electronic copy of this plan is available at:**

<<http://pacific.fws.gov/ecoservices/endangered/recovery/default.htm>> and also at <<http://endangered.fws.gov/recovery/index.html>>.

## ACKNOWLEDGMENTS

This recovery plan was prepared by the U.S. Fish and Wildlife Service's Pacific Islands Fish and Wildlife Office in cooperation with the Commonwealth of the Northern Mariana Islands Department of Lands and Natural Resources. Its completion would not have been possible without their assistance.

Many individuals and organizations have contributed to our knowledge of *Nesogenes rotensis* and *Osmoxylon mariannense* and work cooperatively with us to recover these species. We would especially like to thank the following individuals for their support: James Manglona, Lynn Raulerson, Estanislao Taisacan, and Laura Williams.

## EXECUTIVE SUMMARY

**Current Status:** The plants *Nesogenes rotensis* and *Osmoxylon mariannense* (no common names) are found only on the island of Rota in the Commonwealth of the Northern Mariana Islands (CNMI). *Nesogenes rotensis* is a low-growing herbaceous plant in the verbena family (*Verbenaceae*) and *Osmoxylon mariannense* is a spindly, soft-wooded tree in the ginseng family (*Araliaceae*). Both species were federally listed as endangered in 2004 (U.S. Fish and Wildlife Service [USFWS] 2004). *Osmoxylon mariannense* is also protected by the government of the Commonwealth of the Northern Mariana Islands. Currently, there are 2 known populations of *N. rotensis* of 15 to 20 plants each. The 10 known individuals of *O. mariannense* are in scattered locations on the Sabana (the cloudswept plateau that dominates the western half of Rota).

**Habitat Requirements:** *Nesogenes rotensis* is found on exposed, raised limestone flats in non-forested coastal strand habitat. These flats are subject to salt spray during severe storms. *Osmoxylon mariannense* is found in limestone forests on the Sabana, a raised plateau unique in the Mariana archipelago, that are often shrouded in clouds and mist.

**Limiting Factors:** Human activities that alter native vegetation and habitat are believed to be the primary factors leading to the small population sizes and limited distribution of *Nesogenes rotensis* and *Osmoxylon mariannense*. These activities include agriculture, ranching, nonnative plant and animal introductions, resort and beach park development in the coastal habitat of *N. rotensis*, and road construction and maintenance in the Sabana habitat of *O. mariannense*. In the last decade, several major typhoons have made landfall on Rota, severely impacting individuals of both species. Another factor that may affect the recovery of these two species is their vulnerability to extinction from reduced reproductive vigor due to their small population sizes.

**Recovery Strategy:** Recovery of *Nesogenes rotensis* and *Osmoxylon mariannense* focuses on the following actions: (1) protecting and restoring the native coastal strand and forest habitat of *N. rotensis* and *O. mariannense*, respectively; (2) establishing new populations and augmenting existing populations of these species through methods including controlled propagation and outplanting; (3) assessing the impacts of feral ungulates (deer and pigs), rats, mice, insects, diseases, and introduced plants, and determining methods for their appropriate control or eradication; (4) building public support for conservation; and (5) reassessing and refining recovery actions as appropriate.

**Recovery Priority Numbers:** *Nesogenes rotensis* has a recovery priority number of 2 on a scale of 1 (highest) to 18 (lowest), reflecting a high degree of threat, high recovery potential, and its taxonomic status as a full species. *Osmoxylon mariannense* has a recovery priority number of 5, reflecting a high degree of threat, low recovery potential, and its taxonomic status as a full species.

**Recovery Goals:** The interim goal is to control threats and increase population sizes and geographic distribution sufficient to reclassify or downlist these two endangered species to threatened status. The long-term goal is to conserve and recover *Nesogenes rotensis* and *Osmoxylon mariannense* in order to remove them from the Federal List of Endangered and Threatened Wildlife and Plants (delisting).

**Recovery Objectives:** Restore and maintain multiple naturally reproducing populations of *Nesogenes rotensis* and *Osmoxylon mariannense* on the island of Rota such that the protections of the Endangered Species Act are no longer necessary.

**Recovery Criteria:** *Nesogenes rotensis* and *Osmoxylon mariannense* may be considered for downlisting to threatened status when all of the following criteria are achieved and maintained for a minimum of 10 consecutive years:

- 1) A total of two populations of *Nesogenes rotensis* are naturally reproducing and stable, or increasing in numbers. Each population of *N. rotensis* must consist of at least 300 mature, reproducing individuals. A total of two populations of *Osmoxylon mariannense* are naturally reproducing and stable, or increasing in numbers. Each population of *O. mariannense* must consist of at least 100 mature, reproducing individuals. A stable or increasing population is defined as having a finite rate of increase (known as  $\lambda$  or lambda) greater than or equal to 1 over the requisite 10-year time period.
- 2) Sufficient habitat is protected and managed to achieve criterion 1 above.
- 3) Management and control of nonnative species by local, regional, Commonwealth, and Federal authorities are demonstrated to be successful and sufficient to achieve criterion 1 above.

*Nesogenes rotensis* and *Osmoxylon mariannense* may be considered for removal (delisting) from the Federal List of Endangered and Threatened Wildlife and Plants when all of the following criteria are achieved and maintained for a minimum of 10 consecutive years:

- 1) A total of four populations of *Nesogenes rotensis* are naturally reproducing and stable, or increasing in numbers. Each population of *N. rotensis* must consist of at least 300 mature, reproducing individuals. A total of four populations of *Osmoxylon mariannense* are naturally reproducing and stable, or increasing in numbers. Each population of *O. mariannense* must consist of at least 100 mature, reproducing individuals. A stable or increasing population is defined as having a finite rate of increase (known as  $\lambda$  or lambda) greater than or equal to 1 over the requisite 10-year time period.
- 2) Sufficient habitat is protected and managed to achieve criterion 1 above.

3) Management and control of nonnative species by local, regional, Commonwealth, and Federal authorities are demonstrated to be successful and sufficient to achieve criterion 1 above.

**Recovery Actions Needed:** 1) Coordinate and monitor recovery efforts; 2) Address factors affecting viability of the wild populations; 3) Monitor the extant *Nesogenes rotensis* and *Osmoxylon mariannense* populations, establish new populations, and augment existing populations; and 4) Provide educational and informational opportunities to build public support for conservation.

**Date of Recovery:** Due to the present uncertainties regarding the roles of introduced species, habitat loss and degradation, and the susceptibility of the wild populations to random catastrophic events, we cannot realistically estimate when delisting of *Nesogenes rotensis* and *Osmoxylon mariannense* might occur. We expect that it will likely take several decades to fully recover these species, depending on the status of controlling threats to the species and the status of their populations. As we implement the recovery actions in this plan we anticipate that we will be able to more accurately estimate when delisting might occur.

**Total Estimated Cost of Recovery:** Due to the great number of unknown variables affecting the management and potential recovery of these two species, the total cost to recover *Nesogenes rotensis* and *Osmoxylon mariannense* cannot be realistically determined at this time. Future recovery actions will be initiated, implemented, and modified as informed by research and monitoring of ongoing management actions (see Recovery Actions 1 through 3). We therefore present our best estimate of recovery costs for the next 15 years, a total of \$6,836,000. Approximately \$2,946,000, or close to 44 percent of the estimated total, will be needed during the first 5 years of recovery implementation. A detailed cost breakdown with expected annual costs for the first 5 years of recovery implementation is provided in the Implementation Schedule. We anticipate



updating and revising this recovery plan on a 5-year time schedule, as needed, and recovery cost projections will be updated accordingly at that time.

The recovery actions described in this plan are also expected to benefit the endangered Rota bridled white-eye (*Zosterops rotensis*), the endangered *Serianthes nelsonii* (tronkon guafi or hayun lagu), and the threatened Mariana fruit bat or fanihi (*Pteropus mariannus mariannus*) through habitat protection and restoration in the Sabana region of Rota.

The 15-year and first 5-year costs referenced above are broken down by recovery action priority number as follows:

**Priority 1 Recovery Actions** – Those actions that must be taken to prevent extinction or prevent the species from declining irreversibly in the foreseeable future.

**Priority 2 Recovery Actions** – Those actions that must be taken to prevent a significant decline in population or habitat quality, or some other significant negative impact short of extinction.

**Priority 3 Recovery Actions** – All other actions necessary to meet recovery objectives.

**Table 1. Recovery Plan Time and Costs (\$)**

<b>Years</b>	<b>Priority 1 Actions</b>	<b>Priority 2 Actions</b>	<b>Priority 3 Actions</b>	<b>Totals</b>
<b>2007-2012</b>	500,000	2,146,000	248,000	2,894,000
<b>2013-2022</b>	718,000	2,946,000	278,000	3,942,000
<b>Totals</b>	1,218,000	5,092,000	526,000	6,836,000

# TABLE OF CONTENTS

Executive Summary .....	iii
List of Tables and Figures.....	x
I. Background and Overview .....	1
A. Introduction .....	1
B. Status of the Species.....	3
C. Species Description and Taxonomy .....	4
1. <i>Nesogenes rotensis</i> .....	4
2. <i>Osmoxylon mariannense</i> .....	6
D. Population Trends and Distribution .....	6
1. <i>Nesogenes rotensis</i> .....	6
2. <i>Osmoxylon mariannense</i> .....	8
E. Life History and Ecology .....	10
1. <i>Nesogenes rotensis</i> .....	10
2. <i>Osmoxylon mariannense</i> .....	12
F. Habitat Descriptions .....	12
1. <i>Nesogenes rotensis</i> .....	12
2. <i>Osmoxylon mariannense</i> .....	13
G. Reasons for listing and Current Threats .....	13
1. <i>Nesogenes rotensis</i> .....	15
2. <i>Osmoxylon mariannense</i> .....	17
H. Conservation Efforts to Date.....	20
1. <i>Nesogenes rotensis</i> .....	20
2. <i>Osmoxylon mariannense</i> .....	20

II. Recovery Strategy and Goals .....	23
A. Recovery strategy .....	23
B. Recovery Goals .....	24
C. Recovery Objectives.....	24
D. Recovery Criteria .....	25
III. Recovery Program .....	29
A. Stepdown Outline of Recovery Actions .....	29
B. Narrative Outline of Recovery Actions.....	33
IV. Implementation Schedule .....	49
V. References.....	65
A. Literature Cited.....	65
B. Personal Communications.....	68
C. <i>In Litt.</i> References .....	69
VI. Appendices .....	70
Appendix A. Agency and Peer Reviewers.....	70
Appendix B. Recovery Priority System.....	82
Appendix C. Recovery actions identified to address factors currently limiting the recovery of <i>Nesogenes rotensis</i> and <i>Osmoxylon  mariannense</i> and achieve the recovery criteria.....	83
Appendix D. Summary of Comments.....	85

## LIST OF TABLES AND FIGURES

<b>Table 1.</b>	Recovery Plan Time and Costs.....	vii
<b>Figure 1.</b>	Map of the Mariana archipelago, showing the location of Rota. ....	2
<b>Figure 2.</b>	<i>Nesogenes rotensis</i> . A. Detail of flowering and fruiting stem. B. Habit. C. Flower. D. Seed. E. Fruit.....	5
<b>Figure 3.</b>	<i>Osmoxylon mariannense</i> . A. Habit. B. Immature leaf. C. Mature leaves at branch tip. D. Cross section of fruit, showing seeds. E. Flower and flower bud cluster. ....	7
<b>Figure 4.</b>	Distribution of <i>Nesogenes rotensis</i> .....	9
<b>Figure 5.</b>	Distribution of <i>Osmoxylon mariannense</i> .....	11
<b>Figure 6.</b>	Short-stature coastal strand habitat for <i>Nesogenes rotensis</i> . ....	14
<b>Figure 7.</b>	Limestone forest habitat for <i>Osmoxylon mariannense</i> .....	14
<b>Figure 8.</b>	Conservation areas on Rota. ....	21


# I. Background and Overview

## A. INTRODUCTION

The plants *Nesogenes rotensis* and *Osmoxylon mariannense* (no common names) are historically known only from Rota, an island located approximately 134 kilometers (80 miles) northeast of the territory of Guam. Rota is the fourth largest island in the Mariana archipelago (Figure 1) and is the southernmost island in the Commonwealth of the Northern Mariana Islands (CNMI). Rota is approximately 20 kilometers (12 miles) long and 6 kilometers (4 miles) wide with a land area of approximately 86 square kilometers (33 square miles). The Sabana region, a 12-square kilometer (5-square mile) plateau 450 meters (1,476 feet) in elevation, dominates the western half of the island (see Figure 6). Cliffs border the Sabana region on all sides but the northeastern side, where the plateau slopes down to 150 meters (492 feet) elevation and the Sinapalu plateau, which dominates the eastern half of the island. The airport and village of Sinapalu are situated on the Sinapalu plateau. The village of Songsong and the commercial port for the island are situated on the Taipingot Peninsula, a narrow peninsula jutting out to the southwest on the western coast of the island. Fringing reefs surround most of the island (Resources Northwest 1997).

Rota is a municipality in the Commonwealth of the Northern Mariana Islands. The human population was 3,283 in 2000, a 43 percent increase from the 1990 census estimate (U.S. Census Bureau 2003). Rota's climate is tropical marine with high humidity and uniform temperatures throughout the year. Average daytime temperatures are approximately 27 degrees Celsius (80 degrees Fahrenheit) with approximately 200 centimeters (80 inches) of rain annually and about 80 percent humidity. Rainfall averages 27 centimeters (11 inches) per month during the wet season (June to December) and 10 centimeters (4 inches) per month during the dry season (January to May) (Resources Northwest 1997).

The vegetation on Rota has been described in detail by Fosberg (1960), Falanruw *et al.* (1989), and Mueller-Dombois and Fosberg (1998). The vegetation includes primary and secondary limestone forest, atoll forest, agricultural forest, coconut plantations, *Acacia confusa* (Formosan koa) forest, secondary vegetation, open fields, grassland, and urban vegetation. Approximately 60 percent of the island is forested (Falanruw *et al.* 1989); however, much of this is of medium stature and has been degraded by



**Figure 1.** Map of the Mariana archipelago, showing the location of Rota.

development activities, introduced plants and animals, logging, and the effects of activities from World War II (e.g., roads, defense bunkers, artillery sites, etc.). Historically, native limestone forest varied from semi-dry forest to more or less dry-season deciduous forests on the lower terraces to wet cloud forest on the highest terraces (Fosberg 1960; Mueller, Dombois and Fosberg 1998).

The intent of this recovery plan is to guide the recovery and delisting of *Nesogenes rotensis* and *Osmoxylon mariannense*. The Background section of the plan outlines the basic biology, ecology, and status of the species and their habitats, and describes threats to the species and conservation actions to date. The Recovery section provides the actions needed to recover these species and specific criteria for measuring when recovery has occurred. The success of this plan depends upon the collaboration of many people and organizations to secure the future existence of these species.

## **B. STATUS OF THE SPECIES**

Federal action on these plants began with the publication on February 28, 1996, of the Notice of Review of Plant and Animal Taxa (USFWS 1996). In this document, *Nesogenes rotensis* and *Osmoxylon mariannense* were considered candidate species. These two species were again recognized as candidate species in the September 19, 1997, Candidate Notice of Review (USFWS 1997), and we proposed to list *N. rotensis* and *O. mariannense* as endangered on June 1, 2000 (USFWS 2000). The final listing decision for both species was deferred due to lack of resources because our Pacific Islands Fish and Wildlife Office (where the proposed listing was initiated) was under court order to designate critical habitat for 255 Hawaiian plants and 4 Hawaiian invertebrates. Pursuant to a settlement agreement approved by the U.S. District Court for the District of Hawaii on August 21, 2002, we had to make a final decision on whether to list these species and submit our determination to the *Federal Register* by April 1, 2004 (Center for Biological Diversity v. Norton, Civil No. 99-00603 [D. Haw.]). We listed *N. rotensis* and *O. mariannense* as endangered on April 8, 2004 (USFWS 2004).

*Nesogenes rotensis* is assigned a recovery priority number of 2 based on its taxonomic status as a full species, a high degree of threat, and its high potential for recovery (USFWS 1983 a, b). *Osmoxylon mariannense* has a recovery priority number of 5, reflecting its taxonomic status as a full species, high degree of threat, and a currently low potential for recovery (e.g., no observed

reproduction). We regularly review the status of each listed species with regard to their threats and potential for recovery and update the species' recovery priority number as appropriate.


## C. SPECIES DESCRIPTION AND TAXONOMY

### 1. *Nesogenes rotensis*

*Nesogenes rotensis* is a low-growing herbaceous (non-woody) plant with small, opposite, broadly lanceolate (lance-shaped), coarsely toothed leaves. Flowers are axillary (located on a stem in the area between the stem and the petiole) and tubular in shape, with five white petals and male and female components (Figure 2). Often a flowering branch grows upright, which might aid in pollination or seed dispersal (Raulerson and Rinehart 1997). Plants typically branch near the base at about five to seven nodes, and stature may range from not quite flat-growing (subprostrate) to upward-growing (ascending), scrambling over flattened (appressed) shrubs, with whole plants up to almost 1 meter (3 feet) in diameter (Fosberg and Herbst 1983).

The type specimen of *Nesogenes rotensis* was collected on April 23, 1982, by Derral Herbst and Marjorie Falanruw from Haaniya Point (Poña Point Fishing Cliff) in the Palie area on the island of Rota growing on exposed, dry raised limestone at 100 meters (328 feet) elevation (Fosberg and Herbst 1983). It was growing in association with *Scaevola taccada* (nanaso), *Terminalia samoensis* (talisai ganu), *Hedyotis strigulosa* (paodedo), *Pogonatherum paniceum* (no common name), and *Bikkia tetrandra* (gausali). Fosberg and Herbst (1983) formally described the species and published the name *Nesogenes rotensis* and placed it in the family Chloanthaceae, a largely Australian family. This placement was a change from the historic placement of the genus *Nesogenes* in the family Verbenaceae (verbena family) and its subsequent placement in its own family, Nesogenaceae (Nesogenes family). At present, Mabberly (1990) recognizes *Nesogenes* as a genus of Verbenaceae.





**Figure 2.** *Nesogenes rotensis*. **A.** Detail of flowering and fruiting stem. **B.** Habit. **C.** Flower. **D.** Seed. **E.** Fruit. Drawing © Gregory A. Koob 2005, used with permission.

## **2. *Osmoxylon mariannense***


*Osmoxylon mariannense* is a spindly, soft-wooded tree in the ginseng family (Araliaceae), which can reach 10 meters (33 feet) in height (Figure 3). It has several upward-growing (ascending), gray-barked branches that bear conspicuous leaf scars. Leaves vary in size; mature leaves are palmately lobed (hand-shaped) and about 30 centimeters (1 foot) long and 50 centimeters (1.7 feet) wide. The seven to nine lobes are coarsely toothed, and each lobe has a conspicuous, depressed mid-vein. The leaves are alternate or whorled and grow only at the branch tips. The petioles are 35 to 40 centimeters (1 to 1.5 feet) long and based in distinctive, conspicuous green multiple “sockets” (Raulerson and Rinehart 1991). The flowers are yellow and have both male and female components. They are borne in many-branched, compact terminal cymes or umbels. The fruits are round and maroon in color when ripe (Raulerson and Rinehart 1991).

*Osmoxylon mariannense* was first collected on Rota by French naturalist Alfred Marche, an active botanical explorer in the Mariana Islands from 1887 to 1889 (Stone 1970). It was not until 1933, however, when a study of Marche’s collection was made, that Kanehira first described the species as *Boerlagiodendron mariannense* based on a collection he made in 1932 in dense primary forest at about 400 meters (1,320 feet) elevation (Kanehira 1933). In 1980, Fosberg and Sachet (1980) published the currently accepted recombination, *Osmoxylon mariannense*, which has been upheld by Raulerson and Rinehart (1991).

## **D. POPULATION TRENDS AND DISTRIBUTION**

### **1. *Nesogenes rotensis***

One population of fewer than 100 plants was reported in 1982 by Derral Herbst at the Poña Point Fishing Cliff public park land, owned by and under the jurisdiction of the Commonwealth of the Northern Mariana Islands Department of Lands and Natural Resources (L. Mehrhoff, National Park Service, pers. comm. 1993). In 1994, Raulerson and Rinehart (1997) recorded a population of about 20 plants, occupying 200 square meters (240 square yards) of habitat at the Poña Point Fishing Cliff. This is believed to be the same population reported by Fosberg and Herbst in their 1983 publication; Herbst was uncertain of the original



**Figure 3.** *Osmoxylon mariannense*. **A.** Habit. **B.** Immature leaf. **C.** Mature leaves at branch tip. **D.** Cross section of fruit, showing seeds. **E.** Flower and flower bud cluster. Drawing © Gregory A. Koob, used with permission.


location when he made the herbarium sheet (Fosberg and Herbst 1983; L. Mehrhoff, pers. comm. 1993).

Biannual surveys for this species have been conducted since 2001 at Poña Point Fishing Cliff. A direct count was made on June 27, 2000. At that time there were 80 individuals within an approximate area of 800 square meters (960 square yards). In May and November 2001, direct counts made by staff from the Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife identified 458 and 579 adult plants, respectively. No individuals of *Nesogenes rotensis* were observed in May or November of 2003 following supertyphoon Pongsona; however, 34 adults were observed in December 2003, and about 20 individuals (both seedlings and adults) were observed in March 2005 (USFWS 2004; G. Koob, USFWS, *in litt.* 2005).

In March 2005, a new population of *Nesogenes rotensis* was discovered at Puntan Fina Atkos on public land in the I Chenchon Park conservation area (Figure 4). This area had been suggested as a potential site for surveys, since it had habitat conditions similar to that at Poña Point in Raulerson and Rinehart (1997). There are 15 to 20 individuals (both seedlings and adults) in this new population, which expands the known range of the species (G. Koob, *in litt.* 2005). To date, this species has not been propagated and does not exist in cultivation.

## **2. *Osmoxylon mariannense***

*Osmoxylon mariannense* was first collected more than 100 years ago by Marche and was not collected again until 1932 when Kanehira made at least two collections from dense primary forest at about 400 meters (1,320 feet) elevation (Kanehira 1933). However, there are no written records of the distribution and population size of *O. mariannense* until 1980. Reports from 1980 to 1995 indicate that approximately 20 individuals from 1 scattered population were in the same vicinity as reported by Kanehira (L. Raulerson, University of Guam, pers. comm. 1998; D. Grout, USFWS, and L. Mehrhoff, pers. comms. 1997). One of the larger subpopulations had approximately nine individuals in 1994, but typhoons appeared to have damaged many of the trees and only 2 were visible in 1997 (Raulerson and Rinehart 1997). *Osmoxylon mariannense* occurs as an understory species in *Pisonia umbellifera* (no common name) and *Hernandia labyrinthica* (ocshal) forests, and is often hard to see until some trunks are tall



**Figure 4.** Distribution of *Nesogenes rotensis*.

enough to mingle with the trunks of the other two species (Raulerson and Rinehart 1997).

In January 1998, shortly after typhoon Paka, a total of eight trees, known from five different locations, were observed along the Sabana road (USFWS 2000). The trees were completely defoliated and damaged by the high typhoon winds. Many of the locations had several trees present 15 years earlier but by 1998, only single trees remained in each of the areas, and none were reproducing naturally (G. Hughes, USFWS, pers. comm. 1998).


In 2000, a survey conducted by biologists with the Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife identified six living and five dead individuals of *Osmoxylon mariannense* on Rota. A survey conducted in 2002 by E. Taisacan confirmed eight occurrences in the same vicinity, again with only one living mature tree in each location. *Osmoxylon mariannense* was found on both private (two individuals) and publicly owned (Commonwealth of the Northern Mariana Islands) land (six individuals). *Osmoxylon mariannense* individuals were again defoliated in 2003 during supertyphoon Pongsona; however, in 2003, E. Taisacan reported that some individuals were leafing out and appeared to be recovering (USFWS 2004). Currently, all eight known wild individuals of this species occur along a simple system of unimproved roads crossing the top of the Sabana (Figure 5). This distribution is possibly an artifact of limited access for surveys, as large areas of the Sabana away from the roads are difficult or dangerous to survey due to natural topography and large, often hidden holes left from abandoned mining operations.

Propagation of this species has been limited in its success thus far (see Conservation Efforts to Date). An unknown number of trees currently exist in cultivation, and 2 trees that were outplanted in 2002 adjacent to wild individuals of *Osmoxylon mariannense* continue to survive, bringing the total number of currently known individuals in the wild to 10.

## **E. LIFE HISTORY AND ECOLOGY**

### **1. *Nesogenes rotensis***

Little is known of the life history or ecology of *Nesogenes rotensis*. Based on information from collections and observations, *N. rotensis* flowers in March,



**Figure 5.** Distribution of *Osmoxylon mariannense*.

April, May, and November (Raulerson and Rinehart 1997; G. Koob, *in litt.* 2005; L. Williams, CNMI Division of Fish and Wildlife, pers. comm. 2005). It has been observed in fruit in January, March, and November (Raulerson and Rinehart 1997; G. Hughes, *in litt.* 1998; G. Koob, *in litt.* 2005). All of the available information and recent observations suggest that these plants are perennials, but their above-ground parts die back annually.

## **2. *Osmoxylon mariannense***

Little is known of the life history or ecology of *Osmoxylon mariannense*. It occurs as an understory species in *Pisonia umbellifera* and *Hernandia labyrinthica* forests, and is often hard to see until some trunks are tall enough to mingle with the trunks of the other two species (Raulerson and Rinehart 1997). There are conflicting reports about the habitat requirements of *O. mariannense*. While some authorities consider *O. mariannense* to be an edge species (D. Grout, pers. comm. 1997; Raulerson and Rinehart 1997; L. Raulerson, pers. comm. 2005), others believe it requires shade and protection from the effects of wind by other canopy trees (E. Taisacan, pers. comm. 1997; L. Williams, pers. comm. 2005). Trees have been observed flowering in February, March, and October and fruiting in November, December, January, February, and March (Raulerson and Rinehart 1997; G. Hughes, *in litt.* 1998; G. Koob, *in litt.* 2005). The fruit may provide food for birds and bats, which may also be the dispersal agents, though this is not confirmed (Raulerson and Rinehart 1991; Resources Northwest 1997; J. Manglona, pers. comm. 2005). The seeds of *O. mariannense* are difficult to germinate and this may be due to production of “false seeds” (structures that appear to be seeds but aren’t) or low viability rates (J. Manglona, CNMI, pers. comm. 2005; L. Raulerson, pers. comm. 2005).

## **F. HABITAT DESCRIPTIONS**

### **1. *Nesogenes rotensis***

This species has been found in two locations, Poña Point on Rota’s southern coast and Puntan Fina Atkos on Rota’s eastern coast. At both locations a small population exists on an exposed, raised limestone flat above a 7.6- to 30.5-meter (25- to 100-foot) seaside cliff. Although these flats are up to 30.5 meters (100 feet) above the sea, they are subject to scouring winds during severe storms. *Nesogenes rotensis* grows in association with *Scaevola taccada*, *Terminalia*



*samoensis*, *Hedyotis strigulosa*, *Pogonatherum paniceum*, and *Bikkia tetrandra* (Resources Northwest 1997). (Figure 6).

## **2. *Osmoxylon mariannense***

This tree is found in limestone forests that are often shrouded in clouds and mist on the Sabana and its escarpments. These forests occur in patches in the formerly mined Sabana and are dominated by *Hernandia labyrinthica* and *Elaeocarpus joga* (yoga) interspersed with *Pandanus* (kafu) thickets. Mixed with the *Elaeocarpus* and *Hernandia* are a few *Ficus* spp., *Artocarpus* spp., and *Hibiscus tiliaceus* (pago). Understory species include *Macaranga thompsonii* (pengua) and *Pipturus argenteus* (amahadyan). Epiphytes are abundant and include *Freycinetia reineckeii* (no common name), *Asplenium nidus* (galak), *Davalia solida* (pugua-machena), and other ferns; *Coelogyne guamensis* (no common name) and other orchids; and mosses (Falunruw *et al.* 1989) (Figure 7).

## **G. REASONS FOR LISTING AND CURRENT THREATS**

The threats to the two Mariana plants are each classified according to the five factors identified under section 4(a)(1) of the Endangered Species Act in consideration of listing, delisting, and reclassification decisions. These five factors are as follows:

- (A) The present or threatened destruction, modification, or curtailment of habitat or range;
- (B) Overutilization for commercial, recreational, scientific, or educational purposes;
- (C) Disease or predation;
- (D) Inadequacy of existing regulatory mechanisms; and
- (E) Other natural or man-made factors affecting the continued existence of the species.



**Figure 6.** Short stature coastal strand habitat of *Nesogenes rotensis*. Photo by Gregory A. Koob, USFWS.



**Figure 7.** Limestone forest habitat of *Osmoxylon mariannense*. Photo by Gregory A. Koob, USFWS.

## 1. *Nesogenes rotensis*

**(a) Threats to Habitat (Factor A):** Native vegetation, including open coastal scrubland habitat for *Nesogenes rotensis* on Rota, has undergone extreme alteration due to past and present land use practices, including ranching, deliberate and unintentional nonnative animal and plant introductions, agriculture, and military activities during World War II (Falanruw *et al.* 1989).

Coastal habitat for *Nesogenes rotensis* is under increased pressure of development and is threatened by fragmentation and degradation associated with human use, development, and recreational activities. The species occurs in an area adjacent to a trail that is increasingly subject to bonfires, collecting, and trampling by fishermen and tourists, all human-caused threats to its habitat.

**(b) Overutilization (Factor B):** At this time, overutilization of *Nesogenes rotensis* is not known to be a limiting factor. Unrestricted scientific or horticultural collecting by interested individuals would significantly affect this species due to its extremely low numbers (USFWS 2004).

**(c) Disease or Predation (Factor C):** Disease is not known to impact *Nesogenes rotensis*.

**(d) Inadequacy of Regulatory Mechanisms (Factor D):** *Nesogenes rotensis* is not yet protected under Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife (DFW) regulations but the agency intends to include it in the next amendment to their regulations (L. Williams, *in litt.* 2006). The species was federally listed as endangered in 2004 (USFWS 2004). The Puntan Fina Atkos population, however, is on public land in the I Chenchon Park conservation area where regulations are in place that limit human use and prohibit removal of any plant life in the area (Rota Local Law No. 9-1, 1994). These regulations are inadequate to protect the species from threats identified under section 4(a)(1) of the Endangered Species Act (Factors A, E).

**(e) Other Factors (Factor E):** Rota has a long history of disturbances by tropical typhoons (Weir 1991). While native species are adapted to these events, these typhoons, in combination with human-caused disturbances and the relatively new presence of invasive species, threaten the continued existence of *Nesogenes rotensis*. Within the past decade, frequent typhoons have made

landfall on Rota, severely affecting the island. Most recently, in 2003 supertyphoon Pongsona affected the Mariana Islands, particularly Guam and Rota, with winds of up to 296 kilometers (184 miles) per hour. *Nesogenes rotensis* is particularly susceptible to extirpation or extinction from a natural disaster because of its limited distribution and small numbers of individuals.

Typhoons are a common occurrence in the Mariana Islands. The nearby island of Guam, for example, has been affected by typhoons in 37 of the last 50 years (based on records compiled by the U.S. Navy, Joint Typhoon Warning Center, <http://www.npmoc.navy.mil/jtwc.html>). Supertyphoons (a category of severe storm, defined as having gusts exceeding 240 kilometers [150 miles] per hour) occur about once every 5 to 10 years. There is some evidence that the frequency of severe storms (estimated gusts exceeding 160 kilometers [100 miles] per hour) is increasing in the Mariana Islands. With reference to Guam, the historical record shows increasing numbers of mild (estimated gusts in the range of 80 to 160 kilometers [50 to 100 miles] per hour) and severe storms over the last three centuries, as well as in just the last decade. While some underreporting of storms may have occurred in prior centuries, even mild storms were noticed in the colonial era because they destroyed the relatively flimsy structures used for early housing. Furthermore, these data are consistent with trends expected on the basis of increasing sea surface temperatures that have been documented in recent years (*e.g.*, Strong *et al.* 1998; U.S. Department of State 1999). The two populations of *N. rotensis* are especially vulnerable to the extreme impact of typhoons, storm surge, and high surf because their open scrubland habitats are located in coastal areas.

*Nesogenes rotensis* is also threatened by *Casuarina equisetifolia* (gagu or ironwood), which is becoming established in the coastal scrubland habitat at Poña Point Fishing Cliff. This tree is known to spread by root suckers and, as observed in other areas with similar habitat, may change the coastal scrubland into a monotypic gagu forest. *Casuarina equisetifolia* forest habitat is characterized by an absence of understory vegetation due to the restriction of available sunlight and soil nutrients, and possibly the release of a chemical agent that prevents other plants from growing beneath the trees (Smith 1985; L. Williams, pers. comm. 2004). Additionally, the native parasitic vine *Cassytha filiformis* (agasi) has been reported to parasitize individuals of *N. rotensis*, but deleterious effects have not been documented (L. Williams, pers. comm. 2005).

One population of *Nesogenes rotensis* is located in a public park (Poña Point). Human activities not only threaten the habitat for the species at this location, but the plants may also experience direct mortality and lower reproductive success due to trampling from foot traffic and bonfires set by tourists and fishermen. The small population size and extremely limited distribution of this species makes it particularly vulnerable to extinction from reduced reproductive vigor. With only 2 populations of 15 to 20 individuals each, a decline of successful reproduction in *N. rotensis* could lead to the extinction of the species (USFWS 2004).

## **2. *Osmoxylon mariannense***

**(a) Threats to Habitat (Factor A):** Native vegetation, including Sabana limestone forest habitat for *Osmoxylon mariannense*, has undergone extreme alteration due to past and present land use practices, including ranching, deliberate and unintentional nonnative animal and plant introductions, agriculture, and military activities during World War II (Falanruw *et al.* 1989).

Rota was subject to extensive agricultural development (particularly the cultivation of sugar cane in the lowland areas) by the Japanese prior to World War II. Currently, Rota retains less than 60 percent of its historical native forest (Falanruw *et al.* 1989). Continued loss of native forest is attributable to application of the Agricultural Homestead Act of 1990, which allows for the distribution of 1-hectare (2.5-acre) parcels of public land to eligible participants. Land use plans have proposed that approximately 25 to 45 percent of Rota be designated private agricultural homestead land or land likely to be converted to agricultural homesteads (Resources Northwest 1997). In 2001, the Agricultural Homestead Act of 1990 was amended to allow agricultural homestead permitting on any public lands not required for government use or reserved for other purposes by any other provision of the law. Thus, individuals awaiting permits may choose many areas of Rota's public lands for agricultural homesteads, rather than areas planned and reserved specifically for those purposes (Public Law 12–53). Therefore, the potential for agricultural development continues to threaten the remaining limestone forests on Rota, which includes habitat for *Osmoxylon mariannense* within the Sabana Protected Area. Remaining forest habitat is threatened by fragmentation and degradation associated with agricultural activities, and road maintenance and construction (D. Grout, USFWS and L. Mehrhoff, National Park Service, pers. comms. 1997). Individuals of

*O. mariannense* on Rota were almost lost during road widening activities that occurred in the late 1990s (D. Grout and L. Mehrhoff, pers. comms. 1997).

Throughout the Mariana Islands, introduced goats (*Capra hircus*), pigs (*Sus scrofa*), cattle (*Bos taurus*), and Philippine deer (*Cervus mariannus*) have severely damaged forest vegetation by browsing on plants, causing habitat degradation and erosion (Kessler 1997; Wiles *et. al.* 1999, 2005; Worthington *et. al.* 2001) that then retards forest growth and regeneration (Lemke 1992). Of these nonnative ungulates, deer and feral pigs are degrading the forests of the Sabana (L. Williams, pers. comm. 2005, G. Wiles *in litt.* 2006). These animals may also directly browse on young individuals of *Osmoxylon mariannense* (see Factor C, below).

**(b) Overutilization (Factor B):** At this time, overutilization of *Osmoxylon mariannense* is not known to be a limiting factor, although unrestricted scientific or horticultural collecting would significantly affect this species due to its extremely low numbers (USFWS 2004).

**(c) Disease or Predation (Factor C):** To date, no specific diseases have been identified for *Osmoxylon mariannense*, although individuals have suffered defoliation by an unknown agent (E. Taisacan, pers. comm. 1997). Defoliation (due to the poor health of the leaves), the lack of seedlings and juveniles of *O. mariannense*, and the death of several previously mapped older individual plants are suspected to be caused by unidentified invertebrate pests, mice (*Mus musculus*), rats (*Rattus spp.*), or disease (D. Grout, pers. comm. 1997).

Deer are reported to browse on seedlings of *Osmoxylon mariannense* (USFWS 2004; L. Williams, pers. comm. 2004). Cooperative efforts between the U.S. Fish and Wildlife Service and the Rota Division of Fish and Wildlife have resulted in the construction of fenced exclosures around two individuals in the wild. The remaining eight individuals are not currently fenced (USFWS 2004).

**(d) Inadequacy of Regulatory Mechanisms (Factor D):**

*Osmoxylon mariannense* was federally listed as endangered in 2004 (USFWS 2004) and is included in the list of species protected by the government of the Commonwealth of the Northern Mariana Islands (DFW Non-commercial fishing and hunting regulations, April 2000). These regulations prohibit the

collection or possession of protected plant species but provide no requirements for the analysis of potential adverse effects associated with new projects proposed in the Commonwealth of the Northern Mariana Islands.

**(e) Other Factors (Factor E):** The combination of frequent storm disturbances and resultant competition from invasive, nonnative plant species adversely affects the habitat occupied by *Osmoxylon mariannense* (L. Williams, pers. comm. 2004). While *O. mariannense* is expected to have adapted to high winds and typhoons, its distribution and numbers have been reduced significantly due to human activities, making the few remaining individuals particularly susceptible to extirpation or extinction from a natural disaster. Destruction of the Sabana forest canopy by typhoons has adversely affected *O. mariannense*, altering sub-canopy vegetation conditions over the long term by opening up and drying out older, closed forest habitat (E. Taisacan, pers. comm. 1998; L. Williams, pers. comm. 2004). As discussed above for *Nesogenes rotensis*, all evidence points to the increasing frequency and intensity of the threat from typhoons in this region.

*Osmoxylon mariannense* is threatened by competition from invasive, nonnative plant species including *Momordica charantia* (bitter melon), *Mikania scandens* (climbing hempvine), and *Passiflora suberosa* (corky-stem passionflower). In opened forest areas, these opportunistic, weedy vines cover the ground (Fosberg 1960; G. Hughes, pers. comm. 1998) and may alter the conditions necessary for seed germination and seedling growth provided in closed-canopy, high-stature forests covered with mosses and various epiphytic species.

Small population size and limited distribution make this species particularly vulnerable to extinction from reduced reproductive vigor or random environmental events. There are currently only 10 known individuals of *Osmoxylon mariannense* remaining in the wild, 2 of which were outplanted from past controlled propagation efforts. A single adverse environmental event or a decline of successful reproduction in *O. mariannense* could lead to the extirpation of this species.

## **H. CONSERVATION EFFORTS TO DATE**

### **1. *Nesogenes rotensis***


Attempts to propagate *Nesogenes rotensis* from seeds and cuttings have not been successful (J. Manglona, pers. comm. 2005; L. Mehrhoff, pers. comm. 2005). Surveys of the Poña Point population have been conducted twice a year every year since 2001 (L. Williams, pers. comm. 2005).

In 1994, the I Chenchon Park area was designated as a protected area through Rota Local Law No. 9-1 (Figure 8). The purpose of this protected area, according to the law, is to establish a Wildlife Conservation Area to “prohibit persons from engaging in certain activities within the ... I Chenchon Park area that may have an adverse impact on the wildlife and vegetation” (CNMI Rota Senate Local Law No. 9-1, 1994). In addition, “all persons are ... prohibited from taking or in any way harassing or disturbing ... all plant life, including any fungi, forest vegetation or grasses, with the exception of those plants that possess medicinal properties and/or those that have been used in traditional healing practices ...; and any soil, sand, or rock, within or from the area of Rota known and referred to as ... I Chenchon Park.” The definition of “take” in the law includes harvesting or gathering by any method the entire plant or any part of the plant. “Harassing or disturbing” are also partly defined as “... excavation of surface land for the removal of any type of soil or plant life ... or the destruction of plant life or soil/rock/coral compositions.”

### **2. *Osmoxylon mariannense***

Staff of the Rota Forestry Service are currently collecting seeds from wild individuals and propagating seedlings, in cooperation with E. Taisacan, Commonwealth of the Northern Mariana Islands, Division of Fish and Wildlife (retired) (J. Manglona, pers. comm. 2005). Earlier propagation efforts have had limited success. In 2001, seeds were collected from wild individuals and planted in October and March 2002. Approximately 150 individuals from the October planting had germinated by November. In March 2003, 11 individuals were surviving in a nursery, however they have since died. The seeds planted in March 2002 produced approximately 100 seedlings. Thirty-five of these individuals





**Figure 8.** Conservation areas on Rota.

initially survived but currently only 2 are living and have been outplanted adjacent to wild individuals (USFWS 2004; G. Koob, *in litt.* 2005). Fenced enclosures were constructed around these two individuals through cooperative efforts with the Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife on Rota. The remaining eight individuals are not currently fenced (USFWS 2004). In 1994, part of the Sabana region was designated as a protected area through Rota Local Law No. 9-1 (see Figure 8). The purpose of this protected area, according to the law, is to establish a Wildlife Conservation Area to, “prohibit persons from engaging in certain activities within the Sabana area that may have an adverse impact on the wildlife and vegetation.” The law also calls for groundwater protection, and the continuation of the traditional use of medicinal plants and agricultural practices (CNMI Rota Senate Local Law No. 9-1, 1994). In 1996, a management plan for the Sabana Conservation Area was developed which defined the purpose of the protected area, identified zones for each use (tourism, agriculture, wildlife conservation, firing range, and communication facilities), and suggested rules for each zone (Sabana Protected Area Management Committee 1996). As of 2005, this management plan had not been implemented and the rules, regulations, and prohibitions had not been promulgated to manage the Sabana Conservation Area as required under Rota Local Law No. 9-1 (L. Williams, DFW, pers. comm. 2005). At present, the Rota Division of Land and Natural Resources (DLNR) has been unable to implement the plan due to insufficient funding for programs, facilities, and staff (H. Apatang, *in litt.* 2006).

## II. Recovery Strategy and Goals

### A. RECOVERY STRATEGY

Due to their small population sizes and limited distributions, both *Nesogenes rotensis* and *Osmoxylon mariannense* are highly vulnerable to extirpation or even extinction from random catastrophic events such as typhoons and the accidental introduction of new predators. The first step in the recovery strategy is therefore to protect the known populations and individuals from current threats. In order to achieve full recovery, the threats to the species must be eliminated or controlled sufficiently to be confident in the long-term persistence of the species and the ecosystems upon which they depend. Unfortunately, beyond the inherent risk posed by the small number of individuals and restricted distribution of these species, the nature and relative magnitude of any specific threats to these species is not clear and therefore cannot be addressed in an efficient manner without further research. Currently available information indicates that habitat loss and degradation and predation by introduced animals are negatively impacting *O. mariannense*, but the impact of each of these threats is unclear and needs to be assessed. It is unclear why *N. rotensis* is found in such low numbers. Identifying and assessing the impacts of key threats to the species and determining methods for their control and eradication is a fundamental component of the long-term recovery strategy.

The next step in the recovery of these two species is to establish additional populations and to increase the size of extant populations. The development of a successful propagation and outplanting protocol will be necessary for both species, and will serve in both population augmentation and establishment efforts. Population monitoring and collection of demographic data will reveal whether successful reproduction is occurring, and assist in evaluating progress toward population recovery goals. The specific habitat requirements of both *Nesogenes rotensis* and *Osmoxylon mariannense* must be determined, and the protection and restoration of forests in the Sabana region and non-forested coastal shrubland on the south and east coasts must be a high priority. The potential impacts of feral ungulates (deer and pigs), rats, mice, insects, diseases, and introduced plants must be assessed and controlled as appropriate. For *O. mariannense*, the fencing of some individual trees is a stop-gap defense against deer and pigs, but we propose the effective control or elimination of feral ungulates from the forest as a more effective long-term approach to ecosystem restoration. A similar holistic

approach is proposed for any other nonnative species that are found to pose a threat to the viability of the ecosystem, as practicable. Community support and involvement will be integral to the successful recovery of these species and their habitats, and we suggest an outreach program to involve the people of Rota in the conservation of *N. rotensis*, *O. mariannense*, and other native species of the island. As we learn more about the specific needs of these species through the recovery process, we will reassess and refine the proposed recovery actions to ensure that these activities lead to their successful recovery, downlisting, and delisting.

## **B. RECOVERY GOALS**

The ultimate goal of this plan is to conserve and recover *Nesogenes rotensis* and *Osmoxylon mariannense* to the point that they may be removed from the Federal List of Endangered and Threatened Wildlife and Plants (delisted). Attaining this goal requires that the threats that led to their listing have been successfully addressed, and the increased population sizes and geographic distribution have been maintained and monitored for a period of time sufficient to ensure confidence in the long-term security of the species. The interim goal is to identify and address threats and increase population sizes and geographic distribution sufficient to reclassify or downlist these two endangered species to threatened status. This recovery plan outlines actions necessary to conserve these species and the ecosystems upon which they depend. The conservation of *N. rotensis* and *O. mariannense* in the wild will depend upon the conservation of non-forested coastal shrubland and Sabana limestone forest, respectively, in appropriate quantity and quality to support multiple self-sustaining populations of these species and their habitats.

## **C. RECOVERY OBJECTIVES**

To reach the recovery goals for *Nesogenes rotensis* and *Osmoxylon mariannense* the target recovery objectives are:

- (1) Restore and maintain multiple self-sustaining populations of *Nesogenes rotensis* at Poña Point and Puntan Fina Atkos, and of *Osmoxylon mariannense* in the Sabana region of Rota.
- (2) Secure protection of the non-forested coastal regions on the southern and northeastern coasts and of the Sabana forests of Rota for long-term

maintenance of the ecosystems upon which *Nesogenes rotensis* and *Osmoxylon mariannense* rely.

- (3) Reduce human-induced alterations of the ecosystems in which *Nesogenes rotensis* and *Osmoxylon mariannense* occur and restore ecosystem function to maintain the ecological conditions required by these and other native species.

#### **D. RECOVERY CRITERIA**

Downlisting or delisting is warranted when a listed species no longer meets the definition of threatened or endangered under the Endangered Species Act (Box 1). We set recovery criteria to serve as objective, measurable guidelines to assist us in determining when a species has recovered to the point that the protections afforded by the Endangered Species Act are no longer necessary. However, the actual change in listing status is not solely dependent upon achieving the recovery criteria set forth in a recovery plan; it requires a formal rulemaking process based upon an analysis of the same five factors considered in the listing of a species. The recovery criteria presented in this recovery plan thus represent our best assessment of the conditions that would most likely result in a determination that downlisting or delisting of *Nesogenes rotensis* and *Osmoxylon mariannense* is warranted as the outcome of a formal five-factor analysis in a subsequent regulatory rulemaking.

**Box 1.** Definitions according to section 3 of the Endangered Species Act.

**Endangered Species :** (*In part*) Any species which is in danger of extinction throughout all or a significant portion of its range.

**Threatened Species:** Any species which is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range.

In this plan, initial criteria for downlisting and delisting are provided to guide recovery efforts. These criteria are based on reaching and maintaining population goals to ensure long-term viability, protecting and preserving habitat, and removing or reducing and controlling threats to *Nesogenes rotensis* and *Osmoxylon mariannense*. However, additional threats to these species may be identified as recovery efforts continue. These criteria will be revised as additional information about the species and their threats is accumulated as we implement the recovery actions outlined in this plan.

Mature individuals are defined as those either known or believed to be capable of reproduction. In general, long-lived perennials are those taxa either known or believed to have life spans greater than 10 years and short-lived perennials are those known or believed to have life spans greater than 1 year but less than 10 years. *Nesogenes rotensis* is a short-lived perennial, while *Osmoxylon mariannense* is a long-lived perennial.

Because we have only limited knowledge of the life history of each of these species with respect to specific requirements for both their short-term and long-term survival, this plan establishes only tentative criteria for downlisting and delisting. These criteria were formulated based on recommendations by the Hawaii and Pacific Plants Recovery Coordinating Committee (HPPRCC 1994) and discussions with various biologists and knowledgeable individuals. The Hawaii and Pacific Plants Recovery Coordinating Committee was appointed by the Regional Director for the Fish and Wildlife Service in April 1993 to guide all aspects of recovery for the listed, proposed, and candidate plants of the Hawaiian and other Pacific Islands.

For the purposes of this recovery plan, a population is defined as a discrete unit with sufficient distance between it and neighboring populations that they are not affected by the same small-scale events (such as a landslide). In general, the downlisting criteria for each population of short-lived perennials are naturally reproducing, stable or increasing in numbers, with a minimum of 300 mature, reproducing individuals, and a minimum of 100 mature, reproducing individuals for each population of long-lived perennials. Additional information is needed about both of these plants so that recovery criteria can be quantified and made more meaningful. Recovery actions have been developed to address these uncertainties.

## **Downlisting Criteria**

*Nesogenes rotensis* and *Osmoxylon mariannense* may be considered for downlisting to threatened status when all of the following criteria are achieved and maintained for a minimum of 10 consecutive years:

- 1) A total of two populations of *Nesogenes rotensis* are naturally reproducing and stable, or increasing in numbers. Each population of *Nesogenes rotensis* must consist of at least 300 mature, reproducing individuals. A total of two populations of *Osmoxylon mariannense* are naturally reproducing and stable, or increasing in numbers. Each population of *Osmoxylon mariannense* must consist of at least 100 mature, reproducing individuals. A stable or increasing population is defined as having a finite rate of increase (known as lambda or  $\lambda$ ) greater than or equal to 1 over the requisite 10-year time period.
- 2) Sufficient habitat is protected and managed to achieve criterion 1 above.
- 3) Management and control of nonnative species by local, Commonwealth, and Federal authorities are demonstrated to be successful and sufficient to achieve criterion 1 above.

## **Delisting Criteria**

*Nesogenes rotensis* and *Osmoxylon mariannense* may be considered for removal from the Federal list of endangered and threatened species when all of the following criteria are achieved and maintained for a minimum of 10 consecutive years:

- 1) A total of four populations of *Nesogenes rotensis* are naturally reproducing and stable, or increasing in numbers. Each population of *Nesogenes rotensis* must consist of at least 300 mature, reproducing individuals. A total of four populations of *Osmoxylon mariannense* are naturally reproducing and stable, or increasing in numbers. Each population of *Osmoxylon mariannense* must consist of at least 100 mature, reproducing individuals. A stable or increasing

population is defined as having a finite rate of increase (known as lambda or  $\lambda$ ) greater than or equal to 1 over the requisite 10-year time period.

- 2) Sufficient habitat is protected and managed to achieve criterion 1 above.
- 3) Management and control of nonnative species by local, Commonwealth, and Federal authorities are demonstrated to be successful and sufficient to achieve criterion 1 above.



### III. Recovery Program

#### A. STEPDOWN OUTLINE OF RECOVERY ACTIONS

##### 1. Coordinate and monitor recovery efforts

- 1.1. Coordinate recovery actions with other recovery and ecosystem management efforts
- 1.2. Update or revise recovery plan as needed
- 1.3. Monitor recovery efforts and develop a post-delisting monitoring plan

##### 2. Address factors affecting viability of the wild populations

- 2.1. Protect and restore *Nesogenes rotensis* and *Osmoxylon mariannense* habitat
  - 2.1.1. Conduct research on *Nesogenes rotensis* and *Osmoxylon mariannense* habitat requirements
  - 2.1.2. Protect, manage, and restore strand habitat of *Nesogenes rotensis* by working with the Mariana Public Lands Authority (MPLA) and the Division of Land and Natural Resources (DLNR) on lands they administer
  - 2.1.3. Protect, manage, and restore limestone forest habitat of *Osmoxylon mariannense* by working with interested private landowners and the Mariana Public Lands Authority (MPLA) and the Division of Land and Natural Resources (DLNR) on their lands or lands they administer
    - 2.1.3.1. Develop techniques for restoring native forest within the historical range of *Osmoxylon mariannense*
    - 2.1.3.2. Develop and implement a reforestation plan for the Sabana
  - 2.1.4. Evaluate the impact of feral ungulates (deer and pigs) on native forest regeneration

- 2.1.4.1. Conduct research to determine the impact of feral ungulates on the native forest on the Sabana
- 2.1.4.2. Develop and implement a control program for feral ungulates
- 2.2. Conduct surveys for *Nesogenes rotensis* and *Osmoxylon mariannense* in potentially suitable habitat. Evaluate the use of remote sensing technology as a tool to identify suitable habitat for targeting surveys
- 2.3. Assess and address the impact of introduced predators on *Nesogenes rotensis* and *Osmoxylon mariannense* populations
  - 2.3.1 Evaluate impact of invertebrates on *Nesogenes rotensis* and *Osmoxylon mariannense* populations and control, if necessary
    - 2.3.1.1. Conduct research to determine the impact of invertebrates on *Nesogenes rotensis* and *Osmoxylon mariannense* populations
    - 2.3.1.2. Conduct research on methods to control invertebrates, if necessary
    - 2.3.1.3. Develop and implement an invertebrate control and/or eradication program, if necessary
  - 2.3.2. Evaluate the impact of introduced rats and mice on *Osmoxylon mariannense* populations and control or eradicate, if necessary
    - 2.3.2.1. Conduct research on *Osmoxylon mariannense* fruit predators
    - 2.3.2.2. Develop and implement a rat and mouse control or eradication program, if necessary
  - 2.3.3. Evaluate the impact of feral ungulates (deer and pigs) on *Osmoxylon mariannense* populations and control, if necessary
    - 2.3.3.1. Conduct research to determine the impact of feral ungulates on *Osmoxylon mariannense* populations
    - 2.3.3.2. Conduct research on methods to control feral ungulates
    - 2.3.3.3. Develop and implement a control program for feral ungulates (see recovery action 2.1.4.2.), if necessary

- 2.4. Evaluate the impact of disease on *Nesogenes rotensis* and *Osmoxylon mariannense* populations and control, if necessary
  - 2.4.1. Conduct research to determine the impact of disease on *Nesogenes rotensis* and *Osmoxylon mariannense* populations
  - 2.4.2. Conduct research on methods to control disease, if necessary
  - 2.4.3. Develop and implement a disease control or eradication program, if necessary
- 2.5. Evaluate the impact of the parasitic vine *Cassytha filiformis* on *Nesogenes rotensis* populations and control, if necessary
  - 2.5.1. Conduct research to determine the impact of *Cassytha filiformis* on *Nesogenes rotensis* populations
  - 2.5.2. Conduct research on methods to control *Cassytha filiformis*
  - 2.5.3. Develop and implement a control program for *Cassytha filiformis*, if necessary
- 2.6. Evaluate the impact of *Casuarina equisetifolia* on *Nesogenes rotensis* populations and control, if necessary
  - 2.6.1. Conduct research to determine the impact of *Casuarina equisetifolia* on *Nesogenes rotensis* populations
  - 2.6.2. Conduct research on methods to control *Casuarina equisetifolia*
  - 2.6.3. Develop and implement a control program for *Casuarina equisetifolia*, if necessary
- 2.7. Evaluate the impact of nonnative plants on *Nesogenes rotensis* and *Osmoxylon mariannense* populations and control, if necessary
  - 2.7.1. Conduct research to determine the impact of nonnative plants on *Nesogenes rotensis* and *Osmoxylon mariannense* populations
  - 2.7.2. Conduct research on methods to control nonnative plants

- 2.7.3 Develop and implement a control or eradication program for nonnative plants, if necessary

**3. Monitor *Nesogenes rotensis* and *Osmoxylon mariannense* populations, establish new populations, and augment existing populations**

- 3.1. Develop and implement long-term population monitoring programs for *Nesogenes rotensis* and *Osmoxylon mariannense*
- 3.2. Develop and implement a plan(s) for establishing additional populations and augmenting existing populations
  - 3.2.1. Develop collection and propagation protocols and maintain genetic stock of *Nesogenes rotensis* and *Osmoxylon mariannense*
  - 3.2.2. Determine locations for augmenting existing populations and establishing new populations of *Nesogenes rotensis* and *Osmoxylon mariannense*
  - 3.2.3. Develop outplanting protocols for *Nesogenes rotensis* and *Osmoxylon mariannense*

**4. Provide educational information opportunities to build public support for conservation**

- 4.1. Develop educational programs that can be incorporated into school curricula on Rota
  - 4.1.1. Develop educator's packets for elementary to high school students
  - 4.1.2. Develop workshops for teachers
- 4.2. Develop a public awareness campaign that targets citizens, community groups, and lawmakers
  - 4.2.1. Develop and broadcast Public Service Announcements
  - 4.2.2. Promote a poster and essay contest
  - 4.2.3. Encourage media coverage of environmental issues

- 4.3. Develop and promote “hands-on” community outreach activities that protect and conserve native species and their habitat
  - 4.3.1. Establish a series of small community-based native plant nurseries and botanical gardens, or upgrade existing facilities
  - 4.3.2. Conduct a minimum of one community “Native Seed Collection Day” per year
  - 4.3.3. Conduct a minimum of one community outplanting day per year
  - 4.3.4. Promote interpretation and educational programs on the identification and importance of native plants
  - 4.3.5. Coordinate community involvement in at least one environmental protection activity each year
  - 4.3.6. Promote the development of nature trails in the Sabana or other areas rich in native species
  - 4.3.7. Promote the use of native plants in environmentally appropriate landscaping projects

## **B. NARRATIVE OUTLINE OF RECOVERY ACTIONS**

### **1. Coordinate and monitor recovery efforts**

Due to the complexity of issues associated with *Nesogenes rotensis* and *Osmoxylon mariannense* recovery, a recovery effort that is coordinated with the major stakeholders and interested individuals is needed. In addition, the effectiveness of this effort will depend on monitoring recovery actions and goals, and refinement or modification of management plans and programs that reflect the knowledge gained to maximize the success of the recovery program for these two plants.

#### **1.1. Coordinate recovery actions with other recovery and ecosystem management efforts**

Due to the similarities in recovery issues for listed species in the Mariana Islands, we recommend that recovery groups, Federal and Commonwealth agencies, and interested parties coordinate their conservation efforts.

Holding joint meetings of agencies and other interested parties associated with recovery efforts in the Mariana Islands and maintaining open lines of communication will enable these groups and individuals to assess the progress being made in recovering listed species.

### **1.2. Update or revise recovery plan as needed**

The recovery plan should be reviewed, updated, and revised, as necessary, as we gain further information and knowledge of the life history and ecology of *Nesogenes rotensis* and *Osmoxylon mariannense*, their habitats, and the specific nature of the threats to their persistence through research and management efforts.

### **1.3. Monitor recovery efforts and develop a post-delisting monitoring plan**

A successful recovery program requires frequent and regular monitoring and reporting of recovery efforts. Each recovery action includes a monitoring component that will allow for review to determine its effectiveness. Prior to delisting, a post-delisting monitoring plan must be ready for implementation to ensure the ongoing recovery of the species and continuing effectiveness of monitoring actions. Monitoring under this plan must continue for a minimum of 5 years after delisting occurs.

## **2. Address factors affecting viability of the wild populations**

Habitat loss and degradation, and predation by introduced animals are believed to be impacting the *Nesogenes rotensis* and *Osmoxylon mariannense* populations. In addition, the accidental or intentional introduction of new predators, diseases, and invasive, nonnative plants also threaten the wild populations. These factors need to be assessed and addressed.

### **2.1. Protect and restore *Nesogenes rotensis* and *Osmoxylon mariannense* habitat**

*Nesogenes rotensis* is restricted in distribution to native coastal shrubland while *Osmoxylon mariannense* is restricted to the Sabana native limestone forest. We surmise that the most intact native ecosystems will most likely provide suitable habitat for augmenting existing wild populations and establishing new populations of *Nesogenes rotensis* and *Osmoxylon*

*mariannense*. Therefore, these ecosystems need to be protected from human and other disturbances and restored as necessary. The remaining populations of both species are very small, consisting of 2 small localized populations of *Nesogenes rotensis*, and 1 small scattered population of 10 individual *Osmoxylon mariannense* in the Sabana region. For this reason, all remaining populations and individuals making up these populations require immediate protection. Protection may involve creating protected areas, fencing, and changing public access routes/trails.

**2.1.1. Conduct research on *Nesogenes rotensis* and *Osmoxylon mariannense* habitat requirements**

Research on the specific habitat requirements of *Nesogenes rotensis* and *Osmoxylon mariannense* is needed to determine the best areas for reintroduction.

**2.1.2. Protect, manage, and restore strand habitat of *Nesogenes rotensis* by working with the Mariana Public Lands Authority and the Department of Lands and Natural Resources on lands they administer**

Currently, *Nesogenes rotensis* is found in coastal shrubland on public lands. In order to promote the recovery of this species, efforts should be undertaken to protect and manage habitat on these public lands.

**2.1.3. Protect, manage, and restore limestone forest habitat of *Osmoxylon mariannense* by working with interested private landowners, the Mariana Public Lands Authority and the Department of Lands and Natural Resources, on their lands or lands they administer**

*Osmoxylon mariannense* habitat is found on and adjacent to some private lands on the Sabana. Assistance should be provided to interested landowners to protect, restore, and manage this habitat through programs such as the U.S. Fish and Wildlife Service Landowner Incentive and Partners for Fish and Wildlife programs, and U.S. Department of Agriculture Natural Resources Conservation Service Wildlife Habitat Improvement Program. Assistance to protect, restore, and manage public lands may come from programs

administered by the Office of Insular Affairs, the U.S. Fish and Wildlife Service, or the Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife.

**2.1.3.1. Develop techniques for restoring native forest within the historical range of *Osmoxylon mariannense***

Techniques for restoring unforested and degraded areas need to be developed and evaluated to determine the most effective measures for restoring mature forest within the species' historical range (any suitable habitat within the Sabana region). These techniques might include determining how to propagate suitable native plant species, determining how to increase the regeneration of canopy and understory species in degraded areas, and determining which native plant species are best for restoring cleared areas.

**2.1.3.2. Develop and implement a reforestation plan for the Sabana**

Once effective techniques have been determined for restoration of native forest, an implementation plan should be developed for the Sabana. This plan should include the locations of proposed reforestation actions, an implementation schedule, monitoring protocols and schedule, and estimated costs to implement the plan. The plan could be used to obtain funding for implementing specific tasks.

**2.1.4. Evaluate impact of feral ungulates (deer and pigs) on regeneration of native forest on the Sabana**

Feral ungulates (deer and pigs) negatively impact the natural regeneration of native forest in the Sabana region. Ungulate exclosure plots should be established within degraded forest and open fields on the Sabana and monitored to determine the impacts of ungulate exclusion on forest regeneration.



**2.1.4.1. Conduct research to determine the impact of feral ungulates on the native forest on the Sabana**

Research on effective ungulate control methods is needed if deer and pigs are found to significantly impact the forests of the Sabana. It is not known if a reduced number of ungulates would be acceptable or if complete eradication of ungulates is needed.

**2.1.4.2. Develop and implement a feral ungulate control program**

A feral ungulate control or eradication plan should be developed and implemented to benefit the health of the forests on the Sabana.

**2.2. Conduct surveys for *Nesogenes rotensis* and *Osmoxylon mariannense* in potentially suitable habitat**

The recent discovery of a new population of *Nesogenes rotensis* indicates that potentially suitable habitat may harbor more individuals and populations of this species than are currently known (G. Koob *in litt.* 2005). There are also several areas of the Sabana limestone forests that may provide suitable habitat for *Osmoxylon mariannense* that have not been surveyed. Surveys should be conducted in areas of suitable habitat and remote sensing should be evaluated as a tool for identifying additional areas.

**2.3. Assess and address the impact of introduced predators on *Nesogenes rotensis* and *Osmoxylon mariannense* populations**

Introduced predators are probably impacting *Nesogenes rotensis* and *Osmoxylon mariannense* populations. The predators need to be identified and the extent of these impacts needs to be assessed and addressed, as necessary.

**2.3.1. Evaluate the impact of invertebrates on *Nesogenes rotensis* and *Osmoxylon mariannense* populations and control, if necessary**

It is not known if *Nesogenes rotensis* is threatened by introduced invertebrates. Slugs and various insects have been observed on

individuals of *Osmoxylon mariannense*, and insects are suspected in killing seedlings of this species (J. Manglona, pers. comm. 2005; L. Williams, pers. comm. 2005). Research identifying the species and assessing the impact of invertebrates on *N. rotensis* and *O. mariannense* needs to be conducted, and invertebrate control techniques should be developed, if needed.

**2.3.1.1. Conduct research to determine the impact of invertebrates on the *Nesogenes rotensis* and *Osmoxylon mariannense* populations**

Currently, only anecdotal evidence is available on the impact of invertebrates on *Osmoxylon mariannense* and there is no information regarding their impact on *Nesogenes rotensis*. Research to identify the species and determine the impact of invertebrates on these listed species will enable managers to determine if control of invertebrate pests is necessary for the recovery of one or both listed species.

**2.3.1.2. Conduct research on methods to control invertebrates, if necessary**

Research on effective control methods may be needed if invertebrates are found to be negatively impacting either *Nesogenes rotensis* or *Osmoxylon mariannense* populations.

**2.3.1.3. Develop and implement an invertebrate control or eradication program, if necessary**

An invertebrate control or eradication plan should be developed and implemented if adequate control techniques are available and if it is determined that invertebrate control would be beneficial to *Nesogenes rotensis* and/or *Osmoxylon mariannense* populations.

**2.3.2. Evaluate the impact of introduced rats and mice on *Osmoxylon mariannense* populations and control or eradicate, if necessary**

Rat and mouse predation on *Osmoxylon mariannense* fruit appears to affect its regeneration (D. Grout, pers. comm. 1997; E. Taisacan,

CNMI Division of Fish and Wildlife, retired, pers. comm. 2005). However, the extent of this impact is unknown at this time and should be evaluated to determine if control of these rodents is necessary for the conservation of *O. mariannense*.

**2.3.2.1. Conduct research on *Osmoxylon mariannense* fruit predators**

Currently, it is only suspected that rats or mice are affecting fruit availability of *Osmoxylon mariannense*. Rat-damaged fruits have been observed but it is unknown to what extent rodents are affecting the reproductive capabilities of *O. mariannense*.

**2.3.2.2. Develop and implement a rat and mouse control or eradication program, if necessary**

A plan for control or eradication of rats and mice should be developed and implemented if it is determined that rodent control would be beneficial to the reproductive success of *Osmoxylon mariannense*. This plan should identify the actions to be conducted, additional information needed, and permit requirements. The plan could be used to obtain funding for implementing specific rodent control tasks.

**2.3.3. Evaluate the impact of feral ungulates on *Osmoxylon mariannense* populations and control, if necessary**

The extent of negative impacts by feral ungulates (deer and pigs) on *Osmoxylon mariannense* populations should be evaluated.

**2.3.3.1. Conduct research to determine the impact of feral ungulates on *Osmoxylon mariannense* populations**

Research on the impacts of feral ungulates on individuals of *Osmoxylon mariannense* will enable managers to determine if control is needed.

#### **2.3.3.2. Conduct research on methods to control feral ungulates**

Research on effective control methods is needed if feral ungulates are found to significantly impact individuals of *Osmoxylon mariannense*. It is not known if a reduced number of ungulates (deer and pigs) would be acceptable or if complete eradication of feral ungulates in *O. mariannense* habitat is needed.

#### **2.3.3.3. Develop and implement a feral ungulate control program, if necessary**

A feral ungulate control or eradication plan should be developed and implemented if it is determined that their control would be beneficial to individuals of *Osmoxylon mariannense*.

### **2.4. Evaluate the impact of disease on *Nesogenes rotensis* and *Osmoxylon mariannense* populations and control, if necessary**

It is not known if diseases are affecting *Nesogenes rotensis*. It is possible disease is responsible for the defoliation and dieback that has been observed on *Osmoxylon mariannense*. Research assessing the impact of disease on *N. rotensis* and *O. mariannense* needs to be conducted, and pathogen control techniques need to be researched, developed, and implemented, if necessary.

#### **2.4.1. Conduct research to determine the impact of disease on *Nesogenes rotensis* and *Osmoxylon mariannense* populations**

Currently, it is only suspected that disease affects *Osmoxylon mariannense*, and there is no information regarding the impact of diseases on *Nesogenes rotensis*. Research on the impact of disease on these species will enable managers to determine if disease control is necessary for the recovery of one or both listed plants.

#### **2.4.2. Conduct research on methods to control disease, if necessary**

Research on effective disease control methods will be needed if disease(s) is found to be damaging to individuals of *Nesogenes rotensis* or *Osmoxylon mariannense*.

**2.4.3. Develop and implement a disease control or eradication program, if necessary**

If it is determined that disease control will be beneficial to the recovery of either *Nesogenes rotensis* or *Osmoxylon mariannense*, and effective control techniques are available, a disease control or eradication plan should be developed and implemented.

**2.5. Evaluate the impact of the parasitic vine *Cassytha filiformis* on *Nesogenes rotensis* populations and control, if necessary**

*Cassytha filiformis* is a native parasitic vine known to grow on individuals of *Nesogenes rotensis*. Its effect on populations of this listed species needs to be evaluated.

**2.5.1. Conduct research to determine the impact of *Cassytha filiformis* on *Nesogenes rotensis* populations**

Currently *Cassytha filiformis* is known to parasitize some individuals of *Nesogenes rotensis*. Research on the impact of this parasite on *N. rotensis* will enable managers to determine if control of *C. filiformis* is necessary for recovery.

**2.5.2. Conduct research on methods to control *Cassytha filiformis***

Research on effective control methods is needed if *Cassytha filiformis* is found to significantly impact individuals of *Nesogenes rotensis*.

**2.5.3. Develop and implement a control program for *Cassytha filiformis*, if necessary**

A control or eradication plan should be developed and implemented if effective control techniques are available, and if control or removal of this parasitic vine would be beneficial to individuals of *Nesogenes rotensis*.

**2.6. Evaluate the impact of *Casuarina equisetifolia* on *Nesogenes rotensis* populations and control, if necessary**

Research assessing the impact of *Casuarina equisetifolia* on *Nesogenes rotensis* needs to be conducted, control techniques need to be researched and developed, and implemented if necessary.

**2.6.1. Conduct research to determine the impact of *Casuarina equisetifolia* on *Nesogenes rotensis* populations**

*Casuarina* species are known to spread extensively by root suckers, and shade out and chemically hinder understory vegetation. Research on the impacts of *C. equisetifolia* on *Nesogenes rotensis* will enable managers to determine if their control is necessary for the recovery of the listed plant species.

**2.6.2. Conduct research on methods to control *Casuarina equisetifolia***

Research on effective control methods may be needed if *Casuarina equisetifolia* are found to significantly impact recovery of *Nesogenes rotensis*.

**2.6.3. Develop and implement a control program for *Casuarina equisetifolia*, if necessary**

A control plan for *Casuarina equisetifolia* should be developed and implemented if effective control techniques are available and it is determined that control would be beneficial to recover *Nesogenes rotensis*.

**2.7. Evaluate the impact of nonnative plants on *Nesogenes rotensis* and *Osmoxylon mariannense* populations and control, if necessary**

Nonnative, invasive plant species can alter the function of ecosystems and compete with native species for light, water, and space. Research assessing the impact of nonnative plant species on *Nesogenes rotensis* and *Osmoxylon mariannense* needs to be conducted, and control techniques need to be researched, developed, and implemented if necessary.

**2.7.1. Conduct research to determine the impact of nonnative plants on *Nesogenes rotensis* and *Osmoxylon mariannense* populations**

Currently nonnative plants are suspected of negatively impacting the *Nesogenes rotensis* population at Poña Point. Nonnative plants have invaded the Sabana forest and are changing the native limestone forest habitat of *Osmoxylon mariannense*. Research on the impacts of nonnative plant species on *N. rotensis* and *O. mariannense* will enable managers to determine if their control is necessary for the recovery of the listed plant species.

**2.7.2. Conduct research on methods to control nonnative plants**

Research on effective control methods may be needed if nonnative plants are found to significantly impact recovery of *Nesogenes rotensis* or *Osmoxylon mariannense*.

**2.7.3. Develop and implement a control or eradication program for nonnative plants, if necessary**

A control or eradication plan for nonnative plants should be developed and implemented if effective control techniques are available and it is determined that nonnative plant control would be beneficial to recover *Nesogenes rotensis* or *Osmoxylon mariannense* populations.

**3. Monitor *Nesogenes rotensis* and *Osmoxylon mariannense* populations, establish new populations, and augment existing populations**

Both *Nesogenes rotensis* and *Osmoxylon mariannense* exist in very low numbers. Monitoring various parameters (such as seedling establishment, population structure, etc.) is necessary to determine the effectiveness of recovery actions, evaluate recovery of the species, and indicate when timely action should be taken to prevent complete extinction or loss of individuals in the wild.

**3.1. Develop and implement long-term population monitoring programs for *Nesogenes rotensis* and *Osmoxylon mariannense***

Long-term population monitoring programs for *Nesogenes rotensis* and *Osmoxylon mariannense* will enable managers implementing recovery actions for these species to assess the current status of individuals and/or populations, evaluate population trends, determine effectiveness of recovery actions, and modify recovery actions to ensure their effectiveness.

Currently the population of *Nesogenes rotensis* at Poña Point is monitored twice a year by staff of the Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife (L. Williams, pers. comm. 2005). The population at Punta Fina Atkos was only recently discovered and a monitoring program is not yet in place for it. The 10 known individuals of *Osmoxylon mariannense*, including 8 wild and 2 outplanted trees, are occasionally monitored by Forestry Services Section staff on Rota and contactors of the Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife (J. Manglona, pers. comm. 2005; E. Taisacan, pers. comm. 2005).

Monitoring of *Nesogenes rotensis* should occur during the mid-wet season in order to address the species perennial die-back in the dry season (L. Williams, *in litt.*, 2006). Consideration will also be given to the timing of population surveys following typhoons.

**3.2. Develop and implement a plan(s) for establishing new populations and augmenting existing populations**

A plan for establishing a propagation program should be developed and implemented. The plan should be developed cooperatively by the Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife, the Forestry Services Section, and the U.S. Fish and Wildlife Service, and periodically reviewed and revised as new populations are established and extant wild populations are augmented. Specific tasks should be identified and a schedule for completing the tasks should be provided.



**3.2.1. Develop collection and propagation protocols and maintain genetic stock of *Nesogenes rotensis* and *Osmoxylon mariannense***

To prevent extinction of both species, effective *ex situ* propagation techniques must be developed and implemented. Currently little is known regarding effective methods to collect propagules and successfully propagate either species. Development of effective collection, propagation, and seed storage protocols will enable managers to effectively produce the plants needed to recover these species.

**3.2.2. Determine locations for augmenting existing populations and establishing new populations of *Nesogenes rotensis* and *Osmoxylon mariannense***

The locations for establishing new populations and augmenting existing populations should be selected once the habitat requirements of *Nesogenes rotensis* and *Osmoxylon mariannense* are determined (see Recovery Action 2.1.1.).

**3.2.3. Develop outplanting protocols for *Nesogenes rotensis* and *Osmoxylon mariannense***

Currently little is known regarding the most effective methods to outplant individuals of *Nesogenes rotensis* or *Osmoxylon mariannense*. Research on the most effective outplanting protocols will enable managers to effectively establish new populations and augment existing populations of each of the listed species.

**4. Provide educational and informational opportunities to build public support for conservation**

Public understanding and support of recovery efforts for *Nesogenes rotensis* and *Osmoxylon mariannense* and other rare species is essential to their recovery. Programs that inform teachers and educate students, lawmakers, community groups, local governmental agencies, and interested individuals should be supported, funded, and promoted.

#### **4.1. Develop educational programs that can be incorporated into school curricula on Rota**

##### **4.1.1. Develop educator’s packets for elementary through high school students**

Contract with a highly skilled individual with knowledge of Rota’s peoples and culture and expertise in developing outreach products that comply with local curriculum standards. This individual should then help develop an educational packet focusing on elementary grades through high school that incorporates the basic skills (reading, writing, arithmetic, and science), while creating a positive environmental ethic toward *Nesogenes rotensis* and *Osmoxylon mariannense* (as well as other rare species of Rota) and their habitats. Initially, the educator’s packets should be utilized for 2 years before a thorough evaluation of their effectiveness is conducted. The results of this evaluation should be used to revise the packets to ensure accuracy and effectiveness. The packets should be revised periodically, as needed.

##### **4.1.2. Develop workshops for teachers**

Teacher workshops should be developed and subsequently offered to provide orientation and guidance for the implementation of the educator’s packets.

#### **4.2. Develop a public awareness campaign that targets citizens, community groups, and lawmakers**

Develop a media campaign that promotes the conservation of native species and their habitat. Ensure that residents of Rota and other islands of the Commonwealth of the Northern Mariana Islands have direct involvement in the development and implementation of the campaign. Whenever applicable, the Chamorro language and culture should be incorporated into the materials.

##### **4.2.1. Develop and broadcast Public Service Announcements**

Develop and broadcast a minimum of one Public Service Announcement per year that promotes conservation of *Nesogenes*

*rotensis* and *Osmoxylon mariannense* (and other native species) and their habitats. Public Service Announcements should utilize radio, television, and print media.

#### **4.2.2 Promote a poster and essay contest**

Promote a poster and essay contest among local school children and adults with prizes awarded from local businesses. Contest themes should focus on native species and habitat protection. The winning selections should be highlighted in a calendar and distributed on Rota.

#### **4.2.3. Encourage media coverage of environmental issues**

Encourage media coverage of environmental issues that highlight efforts by the local community to conserve and protect native species and their habitat.

### **4.3. Develop and promote “hands-on” community outreach activities that protect and conserve native species and their habitat**

Develop a community outreach program that unites the people of Rota to actively participate in conservation activities, while learning to appreciate and conserve *Nesogenes rotensis* and *Osmoxylon mariannense* (and other native species) and their habitats. Whenever applicable, the Chamorro language and culture should be incorporated into the materials.

#### **4.3.1. Establish a series of small community-based native plant nurseries and botanical gardens, or upgrade existing facilities**

These nurseries and botanical gardens should be used to promote the propagation and outplanting of native plants. Locate at least one nursery in each local elementary and high school.

#### **4.3.2. Conduct at least one community “Native Seed Collection Day” per year**

Collecting native seeds several times a year for the nurseries will ensure genetic integrity and diversity, and allow participants to better understand and appreciate Rota’s native plants and their habitats.

**4.3.3. Conduct at least one community outplanting day per year**

Outplanting sites will focus on areas of high ecological value, watersheds, and high profile, ecologically appropriate, public places.

**4.3.4. Promote interpretation and educational programs on the identification and importance of native plants**

Utilize the school nurseries to educate school children on the identification and importance of native plant species, while beautifying school grounds and providing potential habitat for native animal species. The students could produce informational signs identifying each native species planted.

**4.3.5. Coordinate community involvement in a minimum of one environmental protection activity a year**

These activities could include beach clean-ups, streamside clean-ups, soil conservation, Earth Day, and local festivals. These activities should provide participants the opportunity to actively participate in conservation, promote *Nesogenes rotensis* and *Osmoxylon mariannense* related outreach products, raise awareness of these and other rare species on Rota, and create community pride.

**4.3.6. Promote the development of nature trails in the Sabana or other areas rich in native species**

Trails can be used as part of educational programs including school field trips, for ecotourism, and for educating local residents about their heritage and environment. Native plant species can be labeled and interpretative signs or brochures be developed.

**4.3.7. Promote the use of native plants in environmentally appropriate landscaping projects**

Work with landscape managers to incorporate native plants into landscape plantings around local hotels and resorts. Educational or interpretative information can be incorporated.

## IV. Implementation Schedule

The Implementation Schedule that follows outlines recovery actions and estimated costs for two plants from Rota, as set forth in this recovery plan. It is a *guide* for accomplishing the objectives and actions discussed in Parts II and III of this plan. This schedule indicates recovery action priority numbers (defined below), action numbers from the recovery action outline in Part III-A, action descriptions, anticipated duration of actions, the responsible parties, and lastly, estimated costs. In addition, the schedule indicates which of the five listing factors the action is intended to address and ameliorate. The initiation and completion of these recovery actions is subject to the availability of funds, as well as other constraints affecting the parties involved.

We have the statutory responsibility for implementing this recovery plan, and only Federal agencies are mandated to take part in recovery efforts for threatened and endangered species. However, recovery of the two plants from Rota will require the involvement of the full range of Federal, Commonwealth, private, and local interests. The expertise and contributions of additional agencies and interested parties is needed to implement certain recovery actions and to accomplish education and outreach objectives. For each recovery action described in the Implementation Schedule, the column titled “Responsible Parties” lists the primary agencies having the authority or responsibility for implementing recovery actions and other groups, such as Commonwealth, private, and non-profit organizations, that also may wish to be involved in recovery implementation. The listing of a party in the implementation schedule does not require, nor imply a requirement, that the identified party has agreed to implement the action(s) or to secure funding for implementing the action(s). When more than one party is listed, the most appropriate lead agency (based on authorities, mandates, and capabilities), has been identified in bold type.

## **Definition of Recovery Action Priorities:**

*Priority 1* — An action that must be taken to prevent extinction or prevent the species from declining irreversibly in the foreseeable future.

*Priority 2* — An action that must be taken to prevent a significant decline in species population or habitat quality, or some other significant negative impact short of extinction.

*Priority 3* — All other actions necessary to meet the recovery objectives.

## **Definition of Recovery Action Durations:**

*Continual (C)* — An action that will be implemented on a routine basis once begun.

*To Be Determined (TBD)* — The action duration is not known at this time or implementation of the action is dependent on the outcome of other recovery actions.

## **Definitions and Acronyms**

*Key to Acronyms used in the Implementation Schedule:*

BRD	U.S. Geological Survey, Biological Resources Discipline
DFW	Commonwealth of the Northern Mariana Islands, Division of Fish and Wildlife
DLNR	Department of Lands and Natural Resources
DOE	Commonwealth of the Northern Mariana Islands, Department of Education
OIA	Office of Insular Affairs
FWS-PIFWO	U.S. Fish and Wildlife Service, Pacific Islands Fish and Wildlife Office
MPLA	Mariana Public Lands Authority
NRCS	U.S. Department of Agriculture, Natural Resources Conservation Service
RFS	Rota Forestry Service

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
1	2.1.2.	A	Protect, manage, and restore strand habitat of <i>Nesogenes rotensis</i>	C	FWS-PIFWO DFW MPLA DLNR NRCS	120	2	2	52	2	2
1	2.1.3.	A	Protect, manage, and restore limestone forest habitat of <i>Osmoxylon mariannense</i>	C	FWS-PIFWO DFW MPLA DLNR NRCS Private	540	10	10	180	10	30
1	3.2.1.	E	Develop collection and propagation protocols and maintain genetic stock of <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i>	5	BRD DFW RFS	558	40	40	40	40	40

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
2	2.1.1.	A	Conduct research on <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i> habitat requirements	5	FWS-PIFWO BRD DFW RFS Research Institutions	90	18	18	18	18	18
2	2.2.	A, E	Conduct surveys for <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i> in potentially suitable habitat	2	FWS-PIFWO DFW	90	45	45	-	-	-
2	2.3.1.1.	C	Conduct research to determine the impact of invertebrates on <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i> populations	5	FWS-PIFWO DFW OIA BRD Research Institutions	100	20	20	20	20	20



## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
2	2.3.2.1.	C	Conduct research on <i>Osmoxylon mariannense</i> fruit predators	5	FWS-PIFWO OIA <b>BRD</b> Research Institutions	100	20	20	20	20	20
2	2.3.3.1.	C	Conduct research to determine the impact of feral ungulates (deer and pigs) on <i>Osmoxylon mariannense</i> populations	5	<b>BRD</b> FWS-PIFWO DFW Research Institutions	250	50	25	25	25	25
2	2.4.1.	C	Conduct research to determine the impact of disease on <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i> populations	3	FWS-PIFWO DFW <b>BRD</b> Research Institutions	75	25	25	25	-	-

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
2	2.6.1.	E	Conduct research to determine the impact of nonnative plants on <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i> populations	5	FWS-PIFWO DFW <b>BRD</b> Research Institutions	100	20	20	20	20	20
2	3.1.	E	Develop and implement long-term population monitoring programs for <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i>	C	<b>FWS-PIFWO</b> DFW RFS	98	8	8	8	8	8

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
2	3.2.2.	E	Determine locations for augmenting existing populations and establishing new populations of <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i>	3	FWS-PIFWO DFW MPLA RFS	60	-	-	20	20	20
2	3.2.3.	E	Develop outplanting protocols for <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i>	5	FWS-PIFWO DFW BRD	150	-	-	-	-	-
2	2.1.3.1.	A	Develop techniques for restoring native forest within the historical range of <i>Osmoxylon mariannense</i>	5	DFW RFS BRD	125	-	-	25	25	25

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
2	2.1.3.2.	A	Develop and implement a reforestation plan for the Sabana	C	<b>FWS-PIFWO</b> DFW MPLA RFS private	750	50	50	50	50	50
2	2.1.4.1.	C	Evaluate impact of feral ungulates (deer and pigs) on regeneration of native forest on the Sabana	C	<b>BRD</b> <b>DFW</b> RFS	104	-	-	-	8	8
2	2.1.4.2. and 2.3.3.3.	C	Develop and implement a feral ungulate control program, if necessary	TBD	FWS-PIFWO <b>DFW</b> RFS OIA NRCS	TBD	-	-	-	-	-

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
2	2.3.1.2.	C	Conduct research on methods to control invertebrates, if necessary	5	DFW <b>BRD</b> Research Institutions	TBD	-	-	-	-	-
2	2.3.1.3.	C	Develop and implement an invertebrate control or eradication program, if necessary	TBD	DFW FWS-PIFWO	TBD	-	-	-	-	-
2	2.3.2.2.	C	Develop and implement a rat and mouse control or eradication program, if necessary	C	FWS-PIFWO DFW OIA RFS	TBD	-	-	-	-	-
2	2.3.3.2.	C	Conduct research on methods to control feral ungulates	3	FWS-PIFWO DFW <b>BRD</b> Research Institutions	TBD	-	-	-	-	-

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
2	2.4.2.	C	Conduct research on methods to control disease	TBD	<b>BRD</b> Research Institutions	TBD	-	-	-	-	-
2	2.4.3.	C	Develop and implement a disease control or eradication program, if necessary	TBD	FWS-PIFWO <b>DFW</b> RFS	TBD	-	-	-	-	-
2	2.6.2.	E	Conduct research on methods to control <i>Casuarina equisetifolia</i>	3	FWS-PIFWO DFW <b>BRD</b>	3,000	200	200	200	200	200
2	2.6.3	E	Develop and implement a control program for <i>Casuarina equisetifolia</i> if necessary	TBD	FWS-PIFWO <b>DFW</b> RFS	TBD	-	-	-	-	-

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
2	2.7.2.	E	Conduct research on methods to control nonnative plants	TBD	FWS-PIFWO DFW <b>BRD</b> RFS Research Institutions	TBD	-	-	-	-	-
2	2.7.3.	E	Develop and implement a control or eradication program for nonnative plants, if necessary	C	FWS-PIFWO DFW <b>RFS</b>	TBD	-	-	-	-	-
3	2.5.1.	E	Conduct research to determine the impact of <i>Cassytha filiformis</i> on the <i>Nesogenes rotensis</i> populations	3	FWS-PIFWO DFW <b>BRD</b> Research Institutions	60	-	-	-	-	-

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
3	1.1.	A, C, E	Coordinate recovery actions with other recovery and ecosystem management efforts	C	<b>FWS-PIFWO</b> DFW MPLA Private Landowners RFS	6	0.4	0.4	0.4	0.4	0.4
3	1.2.	A, C, E	Update or revise recovery plan as needed	C	<b>FWS-PIFWO</b>	36	-	-	-	-	12
3	1.3.	A, C, E	Monitor recovery efforts	C	<b>FWS-PIFWO</b> DFW	30	2	2	2	2	2
3	2.5.2.	E	Conduct research on methods to control <i>Cassytha filiformis</i>	3	FWS-PIFWO DFW <b>BRD</b>	15	-	-	-	-	5



## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
3	2.5.3.	E	Develop and implement a control program for <i>Cassytha filiformis</i> , if necessary	TBD	FWS-PIFWO DFW RFS	TBD	-	-	-	-	-
3	4.1.1.	A, C, E	Develop educator's packets for elementary to high school students	3	FWS-PIFWO DFW DOE	26	4	2	20	-	-
3	4.1.2.	A, C, E	Develop workshops for teachers	3	FWS-PIFWO DFW DOE	14	2	2	10	-	-
3	4.2.1.	A, C, E	Develop and broadcast Public Service Announcements	6	FWS-PIFWO DFW DOE	30	5	5	5	-	5

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
3	4.2.2.	A, C, E	Promote a poster and essay contest	1	<b>FWS-PIFWO</b> DFW DOE	10	-	-	10	-	-
3	4.2.3.	A, C, E	Encourage media coverage of environmental issues	C	<b>FWS-PIFWO</b> DFW RFS	15	1	1	1	1	1
3	4.3.1.	A, C, E	Establish a series of small community-based native plant nurseries and botanical gardens, or upgrade existing facilities	5	<b>FWS-PIFWO</b> <b>DFW</b> RFS	50	10	10	10	10	10
3	4.3.2.	A, C, E	Conduct a minimum of one community "Native Seed Collection Day" a year.	C	<b>FWS-PIFWO</b> <b>DFW</b> <b>RFS</b>	30	2	2	2	2	2

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
3	4.3.3.	A, C, E	Conduct a minimum of one community outplanting day a year	C	FWS-PIFWO DFW RFS	30	2	2	2	2	2
3	4.3.4.	A, C, E	Promote interpretation and educational programs on the identification and importance of native plants	C	FWS-PIFWO DFW RFS	15	1	1	1	1	1
3	4.3.5.	A, C, E	Coordinate community involvement in a minimum of one environmental protection activity a year	C	FWS-PIFWO DFW RFS	30	2	2	2	2	2

## Implementation Schedule for the Recovery Plan

### Two Plants from Rota (*Nesogenes rotensis* and *Osmoxylon mariannense*)

Recovery Action Priority	Recovery Action Number	Listing Factor	Recovery Action Description	Recovery Action Duration (Years)	Responsible Parties	Total Cost (15 years)	Cost estimates by fiscal year (in \$1,000 units)				
							2007	2008	2009	2010	2011
3	4.3.6	A, C, E	Promote the development of nature trails in the Sabana area or other areas rich in native species	C	FWS-PIFWO DFW RFS	95	20	10	5	5	5
3	4.3.7	A, C, E	Promote the use of native plants in environmentally appropriate landscaping projects	C	FWS-PIFWO DFW RFS	34	4	4	2	2	2
			<b>TOTAL</b>			6,836	563.4	526.4	775.4	493.4	535.4

## V. References

### A. LITERATURE CITED

- Falanruw, M.C., T.G. Cole, and A.H. Ambacher. 1989. Vegetation survey of Rota, Tinian, and Saipan, CNMI. Resource Bulletin PSW-27. Berkeley, California: Pacific Southwest Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture: 11 pages.
- Fosberg, F.R. 1960. The vegetation of Micronesia. Bulletin of the American Museum of Natural History. 119:1-76 and plates 1-40.
- Fosberg, F.R., M.H. Sachet and R.L. Oliver. 1979. A geographical checklist of the Micronesian dicotyledonae. *Micronesica* 15:1-295.
- Fosberg, F.R. and D. Herbst. 1983. A *Nesogenes* (Chloanthaceae) from Micronesia. *Micronesica* 19(1-2):11-15.
- Fosberg, F.R. and M.H. Sachet. 1980. Systematic studies of Micronesian plants. *Smithsonian Contributions to Botany* 45:1-40.
- Hawaii and Pacific Plants Recovery Coordinating Committee. 1994. Minutes of the July 7-8, 1994 meeting. Prepared for U.S. Fish and Wildlife Service Pacific Islands Ecoregion, Honolulu, Hawaii.
- Kanehira, R. 1933. New or noteworthy trees from Micronesia 1-15. *Botanical Magazine of Tokyo* 47:676.
- Kessler, C. 1997. Feral animal monitoring and management. *In*: Annual progress report, fiscal year 1996. Pittman-Robertson Federal Aid in Wildlife Restoration Program. Division of Fish and Wildlife, CNMI, pages 8-14.
- Lemke, T.O. 1992. Status of the Mariana fruit bat (*Pteropus mariannus*) in the Northern Mariana Islands north of Saipan. Pages 68-73 *in* D.E. Wilson and G.L. Graham (eds.), *Pacific Island Flying Foxes: Proceedings of an international conservation conference*. U.S. Fish and Wildlife Service Biological Report 90(23):68-73.

- Mabberley, D.J. 1990. *The Plant Book*. Cambridge University Press. New York, New York. 707 pages.
- Mueller-Dombois, D. and Fosberg, R.F. 1998. *Vegetation of the tropical Pacific Islands*. Springer-Verlag New York, Inc., New York. 733 pages.
- Raulerson, L. and A. Rinehart. 1991. Page 15 *in* *Trees and shrubs of the Northern Mariana Islands*. Coastal Resources Management, Commonwealth of the Northern Mariana Islands.
- Raulerson, L. and A. Rinehart. 1997. Three species from Rota. University of Guam Herbarium (GUAM) Contribution No. 34, University of Guam, Territory of Guam. 18 pages.
- Resources Northwest, Inc. 1997. Draft island of Rota natural resources conservation plan, June 5, 1997. Prepared for the Commonwealth of the Northern Mariana Islands by Resources Northwest, Inc., Tigard, Oregon. 80 pages.
- Sabana Protected Area Management Committee. 1996. Management plan for the Sabana Protected Area, Rota/Luta, Commonwealth of the Northern Mariana Islands. 31 pages.
- Smith, C.W. 1985. Impact of alien plants on Hawaii's native biota. Pages 180-250 *in* C.P. Stone and J.M. Scott (eds.), *Hawaii's terrestrial ecosystems preservation and management*. Cooperative National Park Resources Studies Unit, University of Hawaii, Honolulu, Hawaii.
- Stone, B.B. 1970. The flora of Guam. *Micronesia* 6:485.
- Strong, A.E., T.J. Goreau and R.L. Hayes. 1998. Ocean hot spots and coral reef bleaching: January-July 1998. *Reef Encounter* 24:20-22.
- U.S. Census Bureau. 2003. Population and housing profile 2000: 2000 census population and housing, Commonwealth of the Northern Mariana Islands, issued May 2003. Available online at <http://www.census.gov/prod/cen2000/island/CNMIprofile.pdf>. Accessed May 12, 2005.

U.S. Department of State. 1999. Coral bleaching, coral mortality, and global climate change. Report presented by Rafe Pomerance, Deputy Assistant Secretary of State for the Environment and Development to the U.S. Coral Reef Task Force, 5 March 1999, Maui, Hawaii. Released by the Bureau of Oceans and International Environmental and Scientific Affairs, U.S. Department of State, March 5, 1999. Available online at <[http://www.state.gov/www/global/global\\_issues/coral\\_reefs/990305\\_coralreef\\_rpt.html](http://www.state.gov/www/global/global_issues/coral_reefs/990305_coralreef_rpt.html)>. Accessed September 27, 2005.

[USFWS] U.S. Fish and Wildlife Service. 1983a. Endangered and threatened species listing and recovery priority guidelines. Federal Register 48(184):43098-43105.

[USFWS] U.S. Fish and Wildlife Service. 1983b. Endangered and threatened species listing and recovery priority guidelines correction 48(221):51985.

[USFWS] U.S. Fish and Wildlife Service. 1996. Endangered and threatened wildlife and plants; review of plant and animal taxa that are candidates for listing as endangered or threatened species. Federal Register 61(40):7595-7613.

[USFWS] U.S. Fish and Wildlife Service. 1997. Endangered and threatened wildlife and plants; review of plant and animal taxa that are candidates or proposed for listing as endangered or threatened, annual notice of findings on recycled petitions, and annual description of progress on listing actions. Federal Register 62(182):49397-49411.

[USFWS] U.S. Fish and Wildlife Service. 2000. Endangered and threatened wildlife and plants; proposed endangered status for three plants from the Mariana Islands and Guam. Federal Register 65(106):35025-35033.

[USFWS] U.S. Fish and Wildlife Service. 2004. Endangered and threatened wildlife and plants; determination of endangered status and prudency determination for designation of critical habitat for two plant species from the CNMI. Federal Register 69(68):18499-18507.

- Weir, R.C. 1991. Tropical cyclones affecting Guam (1671-1991). F.P.O. San Francisco: U.S. Naval Oceanography Command Center, Joint Typhoon Warning Center, 1991. Technical Note 91-2.
- Wiles, G.J., D.W. Buden, and D.J. Worthington. 1999. History of introduction, population status, and management of Philippine deer (*Cervus mariannus*) on Micronesian islands. *Mammalia* 63:193-215.
- Wiles G.J. 2005. Decline of a population of wild seeded breadfruit (*Artocarpus mariannensis*) on Guam, Mariana Islands. *Pacific Science* 59:509-522.
- Worthington, D.J. A.P. Marshall, G.J. Wiles, and C.C. Kessler. 2001. Abundance and management of Mariana fruit bats and feral ungulates on Anatahan, Mariana Islands. *Pacific Conservation Biology* 7:134-142.

## **B. PERSONAL COMMUNICATIONS**

- Grout, Dan. 1997. Wildlife Biologist, U.S. Fish and Wildlife Service, Rota, Commonwealth of the Northern Mariana Islands.
- Hughes, Guy. 1998. Wildlife Biologist, U.S. Fish and Wildlife Service, Honolulu, Hawaii.
- Manglona, James. 2005. Forester III, Rota Forestry Services Section, Rota, Commonwealth of the Northern Mariana Islands.
- Mehrhoff, Loyal. 1993, 1997, 2005. Chief, Biological Resource Management Division, National Park Service, Fort Collins, Colorado.
- Pangelinan, Arlene. 2005. Conservation Planning and Permits Coordinator, U.S. Fish and Wildlife Service, Honolulu, Hawaii.
- Raulerson, Lynn. 1998, 2005. Professor of Biology, Biology Department, University of Guam, UOG Station Mangilao, Guam.
- Taisacan, Estanislao M. 1997, 1998, and 2005. Retired, Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife, Rota, Commonwealth of the Northern Mariana Islands.



Williams, Laura. 2004, 2005, 2006. Wildlife Supervisor. Commonwealth of the Northern Mariana Islands Division of Fish and Wildlife, Saipan, Commonwealth of the Northern Mariana Islands.

### **C. *IN LITT.* REFERENCES**

Apatang, H. 2006. Resident Director, Department of Lands and Natural Resources, Commonwealth of the Northern Mariana Islands. E-mail to Donna Ball, U.S. Fish and Wildlife Service, Pacific Islands Fish and Wildlife Office, dated September 12, 2006. Subject: Sabana Management Plan.

Hughes, Guy. 1998. Wildlife Biologist. Field notes on three plants from Rota provided to U.S. Fish and Wildlife Service. U.S. Fish and Wildlife Service, Honolulu, Hawaii.

Koob, Gregory. 2005. Mariana trip report – Guam, Rota, and Saipan; March 4 to 17, 2005 Guam time provided to U.S. Fish and Wildlife Service. U.S. Fish and Wildlife Service, Honolulu, Hawaii.

Wiles, G. 2006. Wildlife Biologist, Washington Department of Fish and Wildlife. Letter addressed to Patrick Leonard, U.S. Fish and Wildlife Service, Pacific Islands Office, dated June 11, 2006. Subject: *comment letter on the Draft Recovery Plan for Two Plants from Rota.*

Williams, L. 2006. Wildlife Supervisor, Division of Fish and Wildlife, Department of Lands and Natural Resources, Commonwealth of the Northern Mariana Islands. Letter addressed to Patrick Leonard, U.S. Fish and Wildlife Service, Pacific Islands Office, dated June 5, 2006. Subject: *comment letter on the Draft Recovery Plan for Two Plants from Rota.*

## VI. Appendices

### APPENDIX A. Agency and Peer Reviewers

#### U.S. Fish and Wildlife Service - Washington D.C.

Chief, U.S. Fish and Wildlife Service  
Division of Endangered Species  
Arlington Square Building  
4401 N. Fairfax Dr., Room 452  
Arlington, VA 22203

Chief, Office of Public Affairs  
U.S. Fish and Wildlife Service, DOI  
Main Interior Building  
1849 C. St., NW, Room 3447  
Washington, DC 20240

Chief, Office of Internal Affairs  
U.S. Fish and Wildlife Service  
Washington, DC 20240

Chief, U.S. Fish and Wildlife Service  
Division of Refuges and Wildlife  
U.S. Fish and Wildlife Service  
4401 N. Fairfax Dr., Room 670  
Arlington, VA 22203

#### Others – Washington D.C.

Environmental Protection Agency  
Hazard Evaluation Division – EEB  
(TS769C)  
401 M Street, SW  
Washington, DC 20460

Biological Resources Discipline  
U.S. Geological Survey  
Office of Research Support  
4401 N. Fairfax Dr., Room 725  
Arlington, VA 22203

Ms. Peggy Olwell  
Bureau of Land Management  
Room 204, Building LJB  
Washington, DC 20240-0001

Dr. Loyal Mehrhoff \*  
Chief, Biological Resource Management Division  
National Park Service  
102 Oak Ridge Drive, Suite 200  
Fort Collins, CO 80525

Dr. Warren L. Wagner  
Curator of Pacific Botany  
Botany Department, NHB #166  
Smithsonian Institution  
Washington, DC 20560

**U.S. Fish and Wildlife Service - Pacific Islands Offices**

Dr. Earl Campbell  
Invasive Species Coordinator  
U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50888  
Honolulu, HI 96850

Mr. Eugene Hester  
U.S. Fish and Wildlife Service  
Division of Law Enforcement  
3375 Koapaka St. B296  
Honolulu, HI 96819

Mr. Ernest Kosaka  
Federal Assistance Coordinator  
U.S. Fish and Wildlife Service  
300 Ala Moana Blvd., Room 5-207  
Box 50167  
Honolulu, HI 96850

Mr. Patrick Leonard  
Field Supervisor – Ecological Services  
U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50888  
Honolulu, HI 96850

Dr. Stephen Miller  
Ecosystem Management Coordinator  
U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50888  
Honolulu, HI 96850

Mr. Michael Molina  
Environmental Review Coordinator  
U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50888  
Honolulu, HI 96850

Mr. Jeff Newman  
Assistant Field Supervisor, Habitat Conservation  
U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50888  
Honolulu, HI 96850

Mr. Craig Rowland  
Conservation Partnerships Coordinator  
U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50888  
Honolulu, HI 96850

Ms. Christa Russell  
Prelisting and Listing Program Leader, U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50888  
Honolulu, HI 96850

Ms. Gina Shultz  
Assistant Field Supervisor, Endangered Species  
U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50888  
Honolulu, HI 96850

Mr. Barry Stieglitz  
Refuge Complex Manager  
Hawaiian and Pacific Islands National Wildlife Refuge Complex  
300 Ala Moana Blvd., Room 5-231  
P.O. Box 50167  
Honolulu, HI 96850

Ms. Marilet Zablan  
Endangered Species Recovery Program Leader, U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50888  
Honolulu, HI 96850

**Hawaii and Pacific Plants Recovery Coordinating Committee**

Ms. Marie Bruegmann\*  
U.S. Fish and Wildlife Service  
Pacific Islands Fish and Wildlife Office  
300 Ala Moana Blvd., Room 3-122  
Box 50088  
Honolulu, HI 96850

Ms. Vickie Caraway  
Hawaii Division of Forestry and Wildlife  
1151 Punchbowl Street  
Honolulu, HI 96822

Dr. Sam Gon  
The Nature Conservancy of Hawaii  
923 Nuuanu Avenue  
Honolulu, HI 96817

Mr. Robert Hobdy  
Haiku, HI

Dr. James D. Jacobi  
U.S. Geological Survey –Biological Resources Discipline  
Pacific Islands Ecosystem Research Center  
P.O. Box 44  
Hawaii Volcanoes National Park, HI 96718

Ms. Kapua Kawelo  
U.S. Army Garrison Hawaii  
Directorate of Public Works  
Attn: APVG-GWV  
Schofield Barracks, HI 96857-5000

Mr. Joel Lau  
Hawaii Natural Heritage Program  
677 Ala Moana Boulevard, Suite 705  
Honolulu, HI 96813

Dr. Lloyd Loope  
U.S. Geological Survey, Biological Resources Discipline  
Haleakala Station  
P.O. Box 369  
Makawao, HI 96768

Dr. Mike Maunder  
Fairchild Tropical Garden  
10901 Old Cutler Road  
Miami, FL 33156-4296

Dr. Cliff Morden  
Dept. of Botany  
University of Hawaii at Manoa  
3190 Maile Way  
Honolulu, HI 96822

Ms. Linda Pratt  
U.S. Geological Survey, Biological Resource Division  
Pacific Islands Ecosystem Research Center  
P.O. Box 44  
Hawaii Volcanoes National Park, HI 96718

Dr. Robert Robichaux  
University of Arizona  
Department of Ecology and Evolutionary Biology  
Tucson, AZ 85721

### **Other Federal Offices**

Ms. Jacqueline Bamba  
Biologist  
U.S. Department of Agriculture – Natural Resources Conservation Service  
P. O. Box 5082 CHRB  
Saipan, MP 96950-5082

Ms. Lauren Bjorkman  
Resource Conservationist  
U.S. Department of Agriculture – Natural Resources Conservation Service  
P.O. Box 5004  
Honolulu, HI 96850

Ms. Mary Blevins  
Environmental Protection Agency  
75 Hawthorne Street, H-32  
San Francisco, CA 94105

Mr. Jonathan B. Jarvis  
Director, Pacific Area Office  
National Park Service  
P.O. Box 50165  
Honolulu, HI 96850

Dr. Boone Kaufman, Director  
Institute of Pacific Islands Forestry  
U.S. Forest Service  
23 East Kawili St.  
Hilo, HI 96720

Mr. Ken Kramer  
Soil Conservationist  
U.S. Department of Agriculture – Natural Resources Conservation Service  
P.O. Box 5082 CHRB  
Saipan, MP 96950-5080

Mr. Bill Moran  
District Conservationist  
U.S. Department of Agriculture – Natural Resources Conservation Service  
P. O. Box 5082 CHRB  
Saipan, MP 96950-5082

Mr. Jeffrey Schorr  
Field Representative, CNMI  
Department of the Interior

Office of Insular Affairs  
Third floor, Horiguchi Bldg.  
P. O. Box 502622  
Saipan, MP 96950

Ms. Patricia Young  
Program Manager CNMI and Republic of Palau  
U. S. Environmental Protection Agency  
Pacific Insular Area Programs  
75 Hawthorne St. (CMD-5)  
San Francisco, CA 94105

**Commonwealth of the Northern Mariana Islands**

Governor Juan N. Babauta  
Office of the Governor  
Commonwealth of the Northern Mariana Islands  
Capitol Hill, Caller Box 10007  
Saipan, MP 96950

Mr. Thorne Abbott  
Natural Resource Planner  
Division of Fish and Wildlife  
P.O. Box 10007, Lower Base  
Saipan, MP 96950

Mr. Tomas Aldan, Chair  
CNMI Board of Public Lands  
c/o Division of Public Lands  
P.O. Box 380  
Saipan, MP 96950

Mr. Ben Atalig, Resident Director  
CNMI Department of Lands and Natural Resources  
Box 924  
Songsong Village  
Rota, MP 96951

Mr. Felipe Atalig, Director  
Coastal Resources Management Office  
CNMI Department of Lands and Natural Resources  
AAA 2852, Box 10001  
San Jose  
Saipan, MP 96950



Mr. Ignacio V. Cabrera, Acting Director  
CNMI Division of Environmental Quality  
P.O. Box 1034 CK  
Saipan, MP 96950

Mr. Isaac Calvo  
CNMI Department of Lands and Natural Resources  
Box 924  
Songsong Village, Rota, MP 96951

Mr. Vince Castro  
Office of Public Lands  
P.O. Box 100380  
Saipan, MP 96950

Mr. Pete A. Duenas, Jr.  
Department of Community and Cultural Affairs  
CNMI Division of Historic Preservation  
P.O. Box 827  
Rota, MP 96951

Mr. Frank M. Eliptico  
Chief, Real Estate Division  
Office of Public Lands  
P.O. Box 500380  
Saipan, MP 96950

Rep. Melvin L.O. Faisao  
Chairman, Committee on Federal and Foreign Relations  
House of Representatives  
Eleventh Northern Marianas  
Commonwealth Legislature  
P.O. Box 586 CK  
Saipan, MP 96950

Ms. Bertha C. Leon Guerrero  
Director, Office of Public Lands  
P.O. Box 500380  
Saipan, MP 96950

Mr. Joe Guerrero  
CNMI Division of Historic Preservation  
Department of Community and Cultural Affairs  
Office of the Governor  
Airport Road  
Saipan, MP 96950

Ms. Patricia Guerrero  
Federal Program Coordinator of Policy  
Technical Services Division  
CNMI Department of Public Works  
Lower Base  
Saipan, MP 96950

Mr. Henry Hofschneider, Commissioner  
Commonwealth of the Northern Mariana Islands  
Marianas Public Lands Authority  
P.O. Box 500380, CK  
Saipan, MP 96950

Mr. James C. Manglona \*  
Forester III  
Forestry Services Section  
Department of Lands and Natural Resources  
P.O. Box 924  
Rota, MP 96951

Mr. Richard Seman, Acting Secretary  
Department of Lands and Natural Resources  
P.O. Box 10007, Lower Base  
CNMI  
Saipan, MP 96950

Mr. Juan Reyes, Secretary  
CNMI Department of Public Works  
P.O. Box 10007  
Saipan, MP 96950

Mr. Joseph Ruak  
Public Information and Education Officer  
Division of Fish and Wildlife  
P.O. Box 10007, Lower Base  
Saipan, MP 96950

Mr. Paul Hamilton, Acting Director  
CNMI Division of Fish and Wildlife  
P.O. Box 10007, Lower Base  
Saipan, MP 96950

Ms. Laura L. Williams \*  
Wildlife Supervisor  
CNMI Division of Fish and Wildlife

P.O. Box 10007, Lower Base  
Saipan, MP 96950

**Counties**

Honorable Benjamin T. Manglona  
Mayor, Municipality of Rota  
Commonwealth of the Northern Mariana Islands  
P.O. Box 537  
Songsong Village, Rota, MP 96951

Mr. Thomas Bruce  
Legal Counsel, Office of the Mayor  
P.O. Box 537  
Songsong Village, Rota, MP 96951

Mr. William (Bill) Pendergrass  
Rota Coastal Resources Management  
P.O. Box 1025  
Songsong Village, Rota, MP 96951

**Other Interested Parties**

Mr. David Henkin  
223 South King Street  
Austin Building, Suite 400  
Honolulu, HI 96813

Dr. Derral Herbst \*  
Bishop Museum, Botany Department  
1525 Bernice Street  
P.O. Box 19000A  
Honolulu, HI 96817-2704

Mr. Clyde Imada  
Bishop Museum, Botany Department  
1525 Bernice Street  
P.O. Box 19000A  
Honolulu, HI 96817-2704

Dr. John Marzluff  
College of Forest Resources  
Ecosystem Science and Conservation Division  
University of Washington  
Box 352100  
Seattle, WA 98195-2100

Mr. Frank M. Norita  
Student Activities Coordinator  
Northern Mariana College  
Upward Bound Program  
P.O. Box 879  
Rota, MP 96951

Mr. David Orr  
Waimea Valley Audubon Center  
59-864 Kamehameha Highway  
Haleiwa, HI 96712-9406

Dr. Lynn Raulerson\*  
Biology Department, University of Guam  
UOG Station Mangilao, Guam 96923

Ms. Agnes Rinehart  
P.O. Box 428  
Agana, Guam 96910

Mr. Gregory Schroer  
President, Resources Northwest  
P.O. Box 675  
Kirkland, WA 98083-0675

Mr. Estanislao Taisacan\*  
Rota, MP

Dr. Gregory Witteman  
Biology Department, University of Guam  
UOG Station Mangilao, GU 96923

Mr. Gary Wiles \*  
521 Rogers St. SW  
Olympia, WA 98502

### **Libraries**

Commonwealth Judiciary Law Library  
P.O. Box 502165  
Saipan, MP 96950

Commonwealth Superior Court Law Library  
Civic Centre, Susupe  
P.O. Box 309  
Saipan, MP 96950

Department of Land and Natural Resources, Resource Center  
Caller Box 10007  
Saipan, MP 96950

CNMI Division of Fish and Wildlife Reference Library  
Department of Natural Resources  
Caller Box 10007  
Saipan, MP 96950

Joeten-Kiyu Public Library  
P.O. Box 501092  
Saipan, MP 96950

Rota High School Library  
P.O. Box 1105  
Rota, MP 96950

School Library Program  
Attn: Commissioner of Education  
P.O. Box 501370  
Saipan, MP 96950

U.S. District Court Law Library  
P.O. Box 500687  
Saipan, MP 96950

Thomas Hale Hamilton Library  
University of Hawaii at Manoa  
2550 The Mall  
Honolulu, HI 96822

Hawaii State Library  
478 S. King  
Honolulu, HI 96813

(\* ) Persons and agencies that provided information necessary for the development of the Plan.

## **APPENDIX B. Recovery Priority System**

The Recovery Priority System uses the criteria of degree of threat, recovery potential, and taxonomy (level of genetic distinctiveness). By applying these criteria, all listed species are assigned a species recovery priority number of 1 through 18. A fourth factor, conflict, is a supplementary element in determining what actions are to be implemented for recovery of a species. In addition, the fourth factor gives priority, within each category, to those species that are or may be in conflict with construction or development projects. Thus, the species retains its numerical rank and acquires the letter designation of “C,” indicating conflict (1C-18C). The conflict designation elevates the priority ranking, thus the highest recovery priority ranking is 1C (i.e., a ranking of 1C is higher than a ranking of 1).

A detailed discussion of the Recovery Priority System can be found in pages 43098 through 43105 of the Federal Register volume 48, number 184 of the issue from Wednesday, September 21, 1983 (with corrected table issued on page 51985 of volume 48, number 221, issued Tuesday, November 15, 1983). This table is reproduced on page 83.

**Recovery Priority Table in *Federal Register***

**Vol. 48, No. 221, page 51985**

Degree of Threat	Recovery Potential	Taxonomy	Priority
High	High	Monotypic genus	1
	High	Species	2
	High	Subspecies	3
	Low	Monotypic genus	4
	Low	Species	5
	Low	Subspecies	6
Moderate	High	Monotypic genus	7
	High	Species	8
	High	Subspecies	9
	Low	Monotypic genus	10
	Low	Species	11
	Low	Subspecies	12
Low	High	Monotypic genus	13
	High	Species	14
	High	Subspecies	15
	Low	Monotypic genus	16
	Low	Species	17
	Low	Subspecies	18

**APPENDIX C. Recovery actions identified to address factors currently limiting the recovery of *Nesogenes rotensis* and *Osmoxylon mariannense* and achieve the recovery criteria.**

<b>Listing Factor</b>	<b>Threats</b>	<b>Recovery Criteria</b>	<b>Recovery Actions (including all sub-actions)</b>
<b>A, C, E</b>	All	1, 2, 3	1. Coordinate and monitor recovery efforts  4. Provide educational opportunities to build public support for conservation
<b>Factor A</b>	Habitat alteration and loss	2	2.1. Protect and restore habitat  2.2. Conduct surveys
<b>Factor B</b>	None identified	Not applicable	Not applicable
<b>Factor C</b>	Predators and disease	2	2.3. Evaluate impact of introduced predators on <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i> populations
<b>Factor D</b>	None identified	Not applicable	Not applicable
<b>Factor E</b>	Native and Nonnative invasive plants	3	2.5. Evaluate impact of parasitic vine <i>Cassytha filiformis</i> on <i>Nesogenes rotensis</i> and control  2.6. Evaluate impact of <i>Casuarina equisetifolia</i> and control  2.7. Evaluate impact of nonnative plants and control
	Small population size and limited distribution	1 and 3	3. Monitor <i>Nesogenes rotensis</i> and <i>Osmoxylon mariannense</i> populations, establish new populations, and augment existing populations



## **APPENDIX D. Summary of Comments**

In April 2006, we released the Draft Recovery Plan for Two Plants from Rota for review and comment by Federal agencies, State and local governments, and members of the public. The public comment period was announced in the *Federal Register* (71 FR 23942 ) on April 25, 2006 and closed on June 26, 2006. A total of 89 copies of the draft plan were sent to interested parties for review during the comment period, and it was also made available online. We received five comment letters during the comment period, and some additional comments, information, and updates after the comment period ended. We carefully considered all comments received in finalizing this recovery plan. We thank all the commenters and peer reviewers for their time and interest in the recovery plan, and we feel the final Recovery Plan for Two Plants from Rota has been improved by the comments we received. Many of the editorial and organizational comments provided have been incorporated in the text of this document. Additional comments are addressed specifically below.

Comment 1: The draft downlisting and delisting criteria for *Nesogenes rotensis* and *Osmoxylon mariannense* should be more flexible as there is not enough information about their reproduction, habitat requirements, and historical abundance to provide for one set of criteria. In addition, a minimum of 10 years is needed to safely establish increasing population trends for both species because of the occasional occurrence of major typhoons striking Rota.

Response: We agree that we have only limited knowledge of the life history of each of these species with respect to specific requirements for both their short-term and long-term survival. The criteria for downlisting and delisting each of these species in this final recovery plan are based on recommendations by the Hawaii and Pacific Plants Recovery Coordinating Committee (HPPRCC 1994), discussions with various biologists and knowledgeable individuals, and comments received during the public review period for the draft recovery plan. These revisions are incorporated in this final recovery plan under section II.D. Recovery Criteria (“Downlisting Criteria” and “Delisting Criteria”). The criteria

for downlisting and delisting these species can be considered tentative in a sense, in that we will review and revise, as appropriate, these criteria for *Nesogenes rotensis* and *Osmoxylon mariannense* as numbers of plants increase and as new information emerges on the life histories of these two species.

Comment 2: The boundaries of the population should be defined.

Response: For the purposes of this recovery plan, a population is defined as a discrete unit with sufficient distance between it and neighboring populations that they are not affected by the same small-scale events (such as a landslide).

Comment 3: Fosberg states in his *Micronesian Checklist* that *Casuarina* is native to the Mariana Islands.

Response: Fosberg *et al* (1979) and Raulerson and Rinehart (1991) both agree that *Casuarina equisetifolia* is indigenous (not introduced) to the Mariana Islands. Therefore, we have addressed its impacts to *Nesogenes rotensis* separate from impacts of nonnative plants.

Comment 4: Although feral pigs may not currently occur in the areas occupied by *Osmoxylon* and *Nesogenes*, it would nonetheless be appropriate to mention their occurrence on the island. Reference to pig control should also be included as a possibility later in the plan.

Response: We agree that while feral pigs may not currently inhabit areas occupied by *Nesogenes* and *Osmoxylon*, they may impact the two plant species in the future if pig populations are left uncontrolled. Also, because there are other species of introduced ungulates on the island, the plan has been revised to cover any introduced ungulate which could currently impact, or which may impact *Nesogenes* or *Osmoxylon* populations in the future.

**Region 1  
U.S. Fish and Wildlife Service  
Ecological Services  
911 NE. 11th Avenue  
Portland, Oregon 97232-4181**



**February 2007**