

FEDERAL RADIATION COUNCIL
RADIATION PROTECTION GUIDANCE
FOR FEDERAL AGENCIES
 Memorandum for the President

SEPTEMBER 13, 1961.

Pursuant to Executive Order 10831 and Public Law 86-373, the Federal Radiation Council herewith transmits its second report to you concerning findings and recommendations for guidance for Federal agencies in the conduct of their radiation protection activities.

Background. On May 13, 1960, the first recommendations of the Council were approved by the President and the memorandum containing these recommendations was published in the FEDERAL REGISTER on May 18, 1960. There was also released at the same time, Staff Report No. 1 of the Federal Radiation Council, entitled, "Background Material for the Development of Radiation Protection Standards," dated May 13, 1960.

The first report of the Council provided a general philosophy of radiation protection to be used by Federal agencies in the conduct of their specific programs and responsibilities. It introduced and defined the term "Radiation Protection Guide" (RPG). It provided numerical values for Radiation Protection Guides for the whole body and certain organs of radiation workers and for the whole body of individuals in the general population, as well as an average population gonadal dose. It introduced as an operational technique, where individual whole body doses are not known, the use of a "suitable sample" of the exposed population in which the guide for the average exposure of the sample should be one-third the RPG for the individual members of the group. It emphasized that this operational technique should be modified to meet special situations. In selecting a suitable sample particular care should be taken to assure that a disproportionate fraction of the average dose is not received by the most sensitive population elements. The observations, assumptions, and comments set out in the memorandum published in the FEDERAL REGISTER, May 18, 1960, are equally applicable to this memorandum.

This memorandum contains recommendations for the guidance of Federal agencies in activities designed to limit exposure of members of population groups to radiation from radioactive materials deposited in the body as a result of their occurrence in the environment. These recommendations include: (1) Radiation Protection Guides for certain organs of individuals in the general population, as well as averages over suitable samples of exposed groups; (2) guidance on general principles of control applicable to all radionuclides occurring in the environment; and (3) specific guidance in connection with exposure

of population groups to radium-226, iodine-131, strontium-90, and strontium-89. It is the intention of the Council to release the background material leading to these recommendations as Staff Report No. 2 when the recommendations contained herein are approved.

Specific attention was directed to problems associated with radium-226, iodine-131, strontium-90, and strontium-89. Radium-226 is an important naturally occurring radioactive material. The other three were present in fallout from nuclear weapons testing. They could, under certain circumstances, also be major constituents of radioactive materials released to the environment from large scale atomic energy installations used for peaceful purposes. Available data suggest that effective control of these nuclides, in cases of mixed fission product contamination of the environment, would provide reasonable assurance of at least comparable limitation of hazard from other fission products in the body.

Establishment of the Federal Radiation Council followed a period of public concern incident to discussions of fallout. While strontium-90 received the greatest popular attention, exposures to cesium-137, iodine-131, strontium-89 and, in still lesser degrees to other radionuclides, are involved in the evaluation of over-all effects. The characteristics of cesium-137 lead to direct comparison with whole body exposures for which recommendations by the Council have already been made.

Studies by the staff of the Council indicate that observed concentrations of radioactive strontium in food and water do not result in concentrations in the skeleton (and consequently in radiation doses) as large as have been assumed in the past. However, concentrations of iodine-131 in the diets of small children, particularly in milk, equal to those permitted under current standards would lead to radiation doses to the child's thyroid which, in comparison with the general structure of current radiation protection standards, would be too high. This is because current concentration guides for exposure of population groups to radioactive materials in air, food, and water have been derived by application of a single fraction to corresponding occupational guides. In the case of iodine-131 in milk, consumption of milk and retention of iodine by the child may be at least as great as by the adult, while the relatively small size of the thyroid makes the radiation dose to the thyroid much larger than in the case of the adult. In addition, there is evidence that irradiation of the thyroid involves greater risk to children than to adults.

Recommendations as to Radiation Protection Guides. The Federal Radiation Council has previously emphasized that establishment of radiation protection standards involves a balancing of the benefits to be derived from the controlled use of radiation and atomic energy against the risk of radiation exposure.

In the development of the Radiation Protection Guides contained herein, the Council has considered both sides of this balance. The Council has reviewed available knowledge, consulted with scientists within and outside the Government, and solicited views of interested individuals and groups from the general public. In particular, the Council has not only drawn heavily upon reports published by the International Commission on Radiological Protection (ICRP), the National Committee on Radiation Protection and Measurements (NCRP), and the National Academy of Sciences (NAS), but has had during the development of the report the benefit of consultation with, and comments and suggestions by, individuals from NCRP and NAS and of their subcommittees. The Radiation Protection Guides recommended below are considered by the Council to represent an appropriate balance between the requirements of health protection and of the beneficial uses of radiation and atomic energy.

It is recommended that:

1. The following Radiation Protection Guides be adopted for normal peacetime operations.

TABLE I—RADIATION PROTECTION GUIDES FOR CERTAIN BODY ORGANS IN RELATION TO EXPOSURE OF POPULATION GROUPS

Organ	RPG for individuals	RPG for average of suitable sample of exposed population group
Thyroid.....	1.5 rem per year....	0.5 rem per year.
Bone marrow.....	0.5 rem per year....	0.17 rem per year.
Bone.....	1.5 rem per year....	0.5 rem per year.
Bone (alter-nate guide).	0.003 micrograms of Ra-226 in the adult skeleton or the biological equivalent of this amount of Ra-226.	0.001 micrograms of Ra-226 in the adult skeleton or the biological equivalent of this amount of Ra-226.

It will be noted that the preceding table provides Radiation Protection Guides to be applied to the average of a suitable sample of an exposed population group which are one-third of those applying to individuals. This is in accordance with the recommendations in the first report of the Council concerning operational techniques for controlling population exposure. Since in the case of exposure of a population group to radionuclides the radiation doses to individuals are not usually known, the organ dose to be used as a guide for the average of suitable samples of an exposed population group is also given as an RPG.

Recommendations as to general principles. Control of population exposure from radionuclides occurring in the environment is accomplished in general either by restriction on the entry of such materials into the environment or through measures designed to limit the intake by members of the population of radionuclides already in the environment. Both approaches involve the consideration of actual or potential concentrations of radioactive material in air, water, or food. Controls should be based upon an evaluation of population

exposure with respect to the RPG. For this purpose, the total daily intake of such materials, averaged over periods of the order of a year, constitutes an appropriate criterion.

The control of the intake by members of the general population of radioactive materials from the environment can appropriately involve many different kinds of actions. The character and import of these actions may vary widely, from those which entail little interference with usual activities, such as monitoring and surveillance, to those which involve a major disruption, such as condemnation of food supplies. Some control actions may require prolonged lead times before becoming effective, e.g., major changes in processing facilities or water supplies. The magnitude of control measures should be related to the degree of likelihood that the RPG may be exceeded. The use of a single numerical intake value, which in part has been the practice until now, does not in many instances provide adequate guidance for taking actions appropriate to the risk involved. For planning purposes, it is desirable that insofar as possible control actions to meet contingencies be known in advance.

It is recommended that:

3. The radiological health activities of Federal agencies in connection with environmental contamination with radioactive materials be based, within the limits of the agency's statutory responsibilities, on a graded series of appropriate actions related to ranges of intake of radioactive materials by exposed population groups.

In order to provide guidance to the agencies in adapting the graded approach to their own programs, the recommendations pertaining to the specific radionuclides in this memorandum consider three transient daily rates of intake by suitable samples of exposed population groups. For the other radionuclides, the agencies can use the same general approach, the details of which are considered in Staff Report No. 2. The general types of action appropriate when these transient rates of intake fall into the different ranges are also discussed in Staff Report No. 2. The purpose of these actions is to provide reasonable assurance that average rates of intake by a suitable sample of an exposed population group, averaged over a sample and averaged over periods of the order of one year, do not exceed the upper value of Range II. The general character of these actions is suggested in the following table.

TABLE II—GRADED SCALES OF ACTION

Ranges of transient rates of daily intake	Graded scale of action
Range I.....	Periodic confirmatory surveillance as necessary.
Range II.....	Quantitative surveillance and routine control.
Range III.....	Evaluation and application of additional control measures as necessary.

Recommendations on Ra-226, I-131, Sr-90, and Sr-89. The Council has given specific consideration to the effects on man of rates of intake of radium-226, iodine-131, strontium-90 and strontium-89 resulting in radiation doses equal to those specified in the appropriate RPG's. The Council has also reviewed past and current activities resulting in the release of these radionuclides to the environment and has given consideration to future developments. For each of the nuclides three ranges of transient daily intake are given which correspond to the guidance contained in Recommendation 2, above. Routine control of useful applications of radiation and atomic energy should be such that expected average exposures of suitable samples of an exposed population group will not exceed the upper value of Range II. For iodine-131 and radium-226, this value corresponds to the RPG for the average of a suitable sample of an exposed population group. In the cases of strontium-90 and strontium-89, the Council's study indicated that there is currently no known operational requirement for an intake value as high as the one corresponding to the RPG. Hence, a value estimated to correspond to doses to the critical organ not greater than one-third of the RPG has been used.

The guidance recommended below is given in terms of transient rates of (radioactivity) intake in micromicrocuries per day. The upper limit of Range II is based on an annual RPG (or lower, in case of radioactive strontium) considered as an acceptable risk for a lifetime. However, it is necessary to use averages over periods much shorter than a lifetime for both radiation dose rates and rates of intake for administrative and regulatory purposes. It is recommended that such periods should be of the order of one year. It is to be noted that values listed in the tables are much smaller than any single intake from which an individual might be expected to sustain injury.

It is recommended that:

3. (a) The following guidance on daily intake be adopted for normal peacetime operations to be applied to the average of suitable samples of an exposed population group:

TABLE III—RANGES OF TRANSIENT RATES OF INTAKE (MICROMICROCURIES PER DAY) FOR USE IN GRADED SCALE OF ACTIONS SUMMARIZED IN TABLE II.

Radionuclides	Range I	Range II	Range III
Radium-226.....	0-2	2-20	20-200
Iodine-131.....	0-10	10-100	100-1,000
Strontium-90.....	0-20	20-200	200-2,000
Strontium-89.....	0-200	200-2,000	2,000-20,000

¹ In the case of iodine-131, the suitable sample would include only small children. For adults, the RPG for the thyroid would not be exceeded by rates of intake higher by a factor of 10 than those applicable to small children.

(b) Federal agencies determine concentrations of these radionuclides in air, water, or items of food applicable to their particular programs which are consistent with the guidance contained herein on average daily intake for the radionuclides radium-226, iodine-131, strontium-90, and strontium-89. Some of the general considerations involved in the derivation of concentration values from intake values are given in Staff Report No. 2.

It is recommended that:

4. For radionuclides not considered in this report, agencies use concentration values in air, water, or items of food which are consistent with recommended Radiation Protection Guides and the general guidance on intake.

In the future, the Council will direct attention to the development of appropriate radiation protection guidance for those radionuclides for which such consideration appears appropriate or necessary. In particular, the Council will study any radionuclides for which useful applications of radiation or atomic energy require release to the environment of significant amounts of these nuclides. Federal agencies are urged to inform the Council of such situations.

ABRAHAM RIBICOFF,
Chairman,
Federal Radiation Council.

The recommendations numbered "1" through "4" contained in the above memorandum are approved for the guidance of Federal agencies, and the memorandum shall be published in the FEDERAL REGISTER.

JOHN F. KENNEDY.

SEPTEMBER 20, 1961.