


# **STATUS OF THE PACIFIC COAST GROUND FISH FISHERY**



## **Stock Assessment and Fishery Evaluation**

### **DESCRIPTION OF THE FISHERY**

**PREPARED BY  
THE PACIFIC FISHERY MANAGEMENT COUNCIL  
7700 NE AMBASSADOR PLACE, SUITE 101  
PORTLAND, OR 97220  
503-820-2280  
[WWW.PCOUNCIL.ORG](http://WWW.PCOUNCIL.ORG)**

**December 2014**

# Table of Contents

<b>Description of the Fishery</b> .....	<b>1</b>
<b>Table of Contents</b> .....	<b>1</b>
<b>List of Tables</b> 3	
<b>List of Figures</b> .....	<b>6</b>
<b>CHAPTER 1 Groundfish Stocks, Their Status, and Description of the Management System</b> .....	<b>8</b>
1.1 Description and Status of Groundfish Stocks.....	8
1.1.1 Productivity and Susceptibility Assessment of Stocks to Overfishing .....	15
1.1.2 Stock Assessments and Rebuilding Analyses Used to Estimate Stock Status and Inform Management Decisions .....	21
1.1.2.1 Types of Assessments Used in Managing Groundfish Stocks.....	22
1.1.2.2 Rebuilding Analyses .....	26
1.1.3 Overfished Groundfish Stocks.....	34
1.1.3.1 Bocaccio .....	34
1.1.3.2 Canary Rockfish .....	39
1.1.3.3 Cowcod.....	44
1.1.3.4 Darkblotched Rockfish.....	49
1.1.3.5 Pacific Ocean Perch .....	54
1.1.3.6 Petrale Sole.....	59
1.1.3.7 Yelloweye Rockfish.....	65
1.1.4 Non-Overfished Groundfish Stocks.....	71
1.1.4.1 Arrowtooth Flounder.....	71
1.1.4.2 Black Rockfish off California and Oregon.....	73
1.1.4.3 Black Rockfish off Washington.....	75
1.1.4.4 Cabezon off California .....	76
1.1.4.5 Cabezon off Oregon .....	77
1.1.4.6 California Scorpionfish .....	78
1.1.4.7 Chilipepper Rockfish South of 40°10' N Lat.....	79
1.1.4.8 Dover Sole .....	81
1.1.4.9 English Sole.....	83
1.1.4.10 Lingcod North and South of 40°10' N Lat.....	86
1.1.4.11 Longnose Skate .....	88
1.1.4.12 Longspine Thornyhead.....	91
1.1.4.13 Pacific Cod.....	94
1.1.4.14 Pacific Whiting .....	95
1.1.4.15 Sablefish .....	98
1.1.4.16 Shortbelly Rockfish.....	101
1.1.4.17 Shortspine Thornyhead.....	103
1.1.4.18 Spiny Dogfish .....	106
1.1.4.19 Splitnose Rockfish South of 40°10' N Lat. ....	109
1.1.4.20 Starry Flounder .....	112
1.1.4.21 Widow Rockfish .....	114
1.1.4.22 Yellowtail Rockfish North of 40°10' N lat. ....	117
1.1.5 Groundfish Stock Complexes.....	118
1.1.5.1 Nearshore Rockfish North and South of 40°10' N Lat. ....	119

1.1.5.2	Shelf Rockfish North and South of 40°10' N Lat. ....	126
1.1.5.3	Slope Rockfish North and South of 40°10' N Lat. ....	133
1.1.5.4	Other Flatfish.....	150
1.1.5.5	Other Fish.....	153
1.1.6	Ecosystem Component Species.....	153
1.2	Discard Mortality Rates Used to Manage West Coast Groundfish Stocks .....	154
1.3	The Groundfish Harvest Specification Framework and Harvest Specifications for Fisheries in 2015 and Beyond.....	158
1.3.1	Overfishing Limits.....	162
1.3.2	Acceptable Biological Catches.....	168
1.3.2.1	Considerations for Deciding the Overfishing Probability (P*) When Specifying an Acceptable Biological Catch.....	171
1.3.2.2	Preferred 2015 and 2016 Acceptable Biological Catches .....	172
1.3.3	Annual Catch Limits .....	186
<b>CHAPTER 2</b>	<b>Socioeconomics of the Groundfish Fishery .....</b>	<b>192</b>
2.1	Socioeconomic Summary.....	192
2.2	PacFIN Groundfish Sector Criteria.....	231
<b>CHAPTER 3</b>	<b>Essential Fish Habitat .....</b>	<b>237</b>
<b>CHAPTER 4</b>	<b>Safety At Sea.....</b>	<b>238</b>
<b>Literature Cited</b>	<b>.....</b>	<b>241</b>

## List of Tables

Table 1-1. Latitudinal and depth distributions of groundfish species (adults) managed under the Pacific Coast Groundfish Fishery Management Plan. <sup>a/</sup> .....	9
Table 1-2. Overall scores and results of the Productivity and Susceptibility Assessment (PSA) ranked from most to least vulnerable to overfishing relative to the current west coast fishery based on the GMT's scoring.....	16
Table 1-3. Retrospective Productivity and Susceptibility Assessment (PSA) vulnerability scores of currently overfished rockfish species ranked from most to least vulnerable to overfishing relative to stock status and the fishery circa 1998, based on the GMT's scoring. ....	18
Table 1-4. Management quantities estimated from the most recent stock assessments informing management in 2015 and beyond. ....	28
Table 1-5. Parameters estimated and/or assumed in base models in the most recent west coast groundfish stock assessments, excluding those done using XDB-SRA. ....	31
Table 1-6. Population parameters estimated and/or derived in base models in 2013 west coast groundfish stock assessments using XDB-SRA.....	33
Table 1-7. Rebuilding parameters estimated in the most recent rebuilding analyses and specified in rebuilding plans for overfished groundfish stocks at the start of the 2015-2016 management cycle. ....	34
Table 1-8. Groundfish species proposed for designation as Ecosystem Component Species under FMP Amendment 24. ....	153
Table 1-9. Discard mortality rates by commercial gear type used to manage west coast groundfish...	155
Table 1-10. Discard mortality rates by depth of groundfish species released at the surface in west coast recreational fisheries using hook-and-line gear. ....	156
Table 1-11. Proportion of recreational and non-recreational gears used in 2004-2006 Oregon nearshore commercial fisheries based on logbook data.....	157
Table 1-12. Discard mortality rates by depth of rockfish species released at the surface in west coast nearshore commercial fisheries. ....	157
Table 1-13. Discard mortality rates by depth of canary rockfish, cowcod, and yelloweye rockfish released in west coast recreational fisheries using descending devices.....	158
Table 1-14. Status quo 2014 harvest specifications and final preferred overfishing limits (OFLs in mt), acceptable biological catches (ABCs in mt), annual catch limits (ACLs in mt), and ecosystem component species designations for west coast groundfish stocks and stock complexes in 2015 and 2016 (stocks with new assessments in bold). ....	159
Table 1-15. Specified 2014 OFLs (mt) and preferred 2015 and 2016 OFLs (mt) for west coast groundfish stocks stock complexes (overfished stocks in CAPS, stocks with new assessments in bold, substock contributions to a stock OFL in <i>italics</i> ).....	165
Table 1-16. Criteria used by the SSC to categorize stocks based on the quantity and quality of data informing the estimate of OFL. Stock categories are used in deciding 2015 and 2016 ABCs that accommodate the uncertainty in estimating OFLs.....	170
Table 1-17. Relationship between P* and the percent reduction of the OFL for deciding the 2015 and 2016 ABCs for category 1, aurora rockfish, widow rockfish, category 2, and category 3 stocks based on $\sigma$ values of 0.36, 0.39, 0.41, 0.72, and 1.44, respectively. ....	173
Table 1-18. 2014 ABCs (mt) and a range of alternative 2015 ABCs (mt) varied by the probability of overfishing (P*) for west coast groundfish stocks (overfished stocks in CAPS; stocks with new assessments in bold; component stocks in stock complexes in <i>italics</i> ).....	174
Table 1-19. 2014 ABCs (mt) and a range of alternative 2016 ABCs (mt) varied by the probability of overfishing (P*) for west coast groundfish stocks (overfished stocks in CAPS; stocks with new assessments in bold; component stocks in stock complexes in <i>italics</i> ).....	178

Table 1-20. Preferred 2015 and 2016 ABCs (mt) for west coast groundfish stocks and stock complexes and the associated ABC harvest control rule (P* = overfishing probability) (overfished stocks in CAPS; stocks with new assessments in bold; component stocks in stock complexes in <i>italics</i> ). .....	182
Table 1-21. The status quo 2014 and preferred 2015 and 2016 annual catch limits (ACLs in mt) for west coast groundfish stocks and stock complexes (stocks with new assessments in bold). .....	189
Table 2-1. Landings (shoreside commercial and tribal) by species group (mt), 1981-2012. ....	192
Table 2-2. Ex-vessel revenue (shoreside commercial and tribal) by species group in current (2012) dollars, \$1,000s, 1981-2012. ....	194
Table 2-3. Groundfish landings (shoreside commercial and tribal) by species or species group (mt), 1981-2012. ....	195
Table 2-4. Groundfish ex-vessel revenue (shoreside commercial and tribal) by species or species group in current (2012) dollars, \$1,000s, 1981-2012. ....	195
Table 2-5. Groundfish landings (shoreside commercial and tribal) by gear type (mt), 1991-2012. ....	197
Table 2-6. Groundfish ex-vessel revenue (shoreside commercial and tribal) by gear type in current (2012) dollars, \$1,000s, 1981-2012. ....	198
Table 2-7. Shoreside IFQ - Trawl (whiting and non-whiting) by groundfish species or species group (mt), 2003-2012. ....	199
Table 2-8. Shoreside IFQ - Trawl (whiting and non-whiting) ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012. ....	199
Table 2-9. Shoreside IFQ – Non-trawl by groundfish species or species group (mt), 2011-2012. ....	200
Table 2-10. Shoreside IFQ - Non-trawl ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2011-2012. ....	200
Table 2-11. Limited entry fixed gear landings by groundfish species or species group (mt), 2003-2012. ....	201
Table 2-12. Limited entry fixed gear ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012. ....	201
Table 2-13. Open access landings (other than by fixed gear) by groundfish species or species group (mt), 2003-2012. ....	202
Table 2-14. Open access ex-vessel revenue (other than by fixed gear) by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012. ....	202
Table 2-15. Non-nearshore fixed gear landings by groundfish species or species group (mt), 2003-2012. ....	203
Table 2-16. Non-nearshore fixed gear ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012. ....	203
Table 2-17. Nearshore fixed gear landings by groundfish species or species group (mt), 2003-2012. ....	204
Table 2-18. Nearshore fixed gear ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012. ....	204
Table 2-19. Incidental open access landings by groundfish species or species group (mt), 2003-2012. ....	205
Table 2-20. Incidental open access ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012. ....	205
Table 2-21. Non-nearshore non-sablefish landings by groundfish species or species group (mt), 2003-2012. ....	206
Table 2-22. Non-nearshore non-sablefish ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012. ....	206
Table 2-23. Groundfish landings by shoreside commercial fishery sectors (mt), 2003-2012. ....	207
Table 2-24. Groundfish ex-vessel revenue in current (2012) dollars, \$1,000s, by shoreside commercial fishery sectors, 2003-2012. ....	208
Table 2-25. Treaty non-whiting groundfish sector landings (groundfish only) by gear group (mt), 2003-2012. ....	209

Table 2-26. Treaty non-whiting groundfish sector ex-vessel revenue (groundfish only), current (2012) dollars, \$1,000s, 2003-2012. ....	209
Table 2-27. Landings (mt) by whiting sectors, 2003-2012. ....	210
Table 2-28. Ex-vessel revenue, current (2012) dollars, \$1,000s, by whiting sectors, 2003-2012. ....	211
Table 2-29. Average monthly landings (mt) by commercial fishery sectors, 5 years, 2008-2012 (except as noted). ....	212
Table 2-30. Average monthly ex-vessel revenue in nominal dollars, \$1,000s, by commercial fishery sectors, 5 years, 2008-2012 (except as noted). ....	213
Table 2-31. Average monthly landings (mt) by Treaty fishery sectors, 5 years, 2008-2012. ....	214
Table 2-32. Average monthly ex-vessel revenue in nominal dollars, \$1,000s, by Treaty fishery sectors, 5 years, 2008-2012. ....	214
Table 2-33. Average monthly groundfish landings (mt) by species and species groups, 5 years, 2008-2012. ....	215
Table 2-34. Average monthly groundfish ex-vessel revenue in nominal dollars, \$1,000s, by species and species groups, 5 years, 2008-2012. ....	215
Table 2-35. Groundfish landings (mt) by "IOPAC port groups", 2003-2012. ....	216
Table 2-36. Groundfish ex-vessel revenue in current (2012) dollars, \$1,000s, by "IOPAC port groups", 2003-2012. ....	217
Table 2-37. Landings (mt) ex-vessel revenue in current (2012) dollars, \$1,000s, by "IOPAC port groups" and species and species groups, 2010-2012. ....	218
Table 2-38. Landings (mt) ex-vessel revenue in current (2012) dollars, \$1,000s, by "IOPAC port groups" and commercial fishery sectors, 2010-2012 (Washington and Oregon). ....	221
Table 2-39. Landings (mt) ex-vessel revenue in current (2012) dollars, \$1,000s, by "IOPAC port groups" and commercial fishery sectors, 2010-2012 (California). ....	223
Table 2-40. Number of vessels making at least one groundfish landing by "IOPAC port groups" and commercial fishery sectors, 5 years 2008-2012. ....	226
Table 2-41. Number of vessels making at least one groundfish landing by commercial fishery sector and length category (feet), 5 years 2008-2012. ....	229
Table 2-42. Engagement (groundfish ex-vessel revenue in port as percent of ex-vessel coastwide revenue) and dependence (groundfish ex-vessel revenue in port as percent of total ex-vessel revenue in port), using current (2012) dollars, 2003-2012. ....	229
Table 2-43. Groundfish landings (mt) on the west coast (Washington-Oregon-California) from inside and outside the Pacific Council management area. ....	230
Table 2-44. Sector codes. ....	235
Table 2-45. Nearshore species list. (Source: M. Bellman, Heery, E., Jannot, J., and Majewski, J. 2010. <i>Explicit Retrieval and Processing of PacFIN Data Used in Total Mortality Estimation</i> . FRAM, NWFSC, NMFS.) ....	236

# List of Figures

Figure 1-1. Fishery management lines on the U.S. west coast. .... 13

Figure 1-2. Productivity and Susceptibility Analysis (PSA) plot for species in the west coast groundfish FMP. Contours delineate areas of relative vulnerability (V, i.e. distance from the origin), with the highest vulnerability stocks above the solid red line ( $V = 2.2$ ), high vulnerability above the orange broken line ( $V=2$ ), medium vulnerability above the green dotted line ( $V=1.8$ ) and the lowest vulnerability below the green dotted line. The maximum vulnerability ( $V=2.8$ ) is indicated with the solid black line. Solid circles are based on current PSA scores. Open circles are based on PSA scores circa 1998. Numbers refer to the Stock ID in Table 1-2 and Table 1-3..... 19

Figure 1-3. Data quality plots for the productivity and susceptibility scores in the PSA for each species (represented numerically in Table 1-2 and Table 1-3) in the west coast groundfish FMP. Higher scores indicate less data quality. Vertical and horizontal lines provide a general guide to relative data quality with values above 3 on either axis considered data-poor..... 20

Figure 1-4. Relative depletion of bocaccio south of 40°10' N lat. from 1980 to 2013 based on the 2013 stock assessment update..... 37

Figure 1-5. Estimated bocaccio recruitments, 1980-2012 (from Field 2013)..... 38

Figure 1-6. Relative depletion of canary rockfish from 1980 to 2011 based on the 2011 stock assessment update..... 42

Figure 1-7. Estimated canary rockfish recruitments, 1980-2010 (from Wallace and Cope 2011). .... 43

Figure 1-8. Estimated spawning potential ratio (SPR) of canary rockfish relative to the current  $F_{MSY}$  and rebuilding harvest rates, 1995-2010. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis..... 44

Figure 1-9. Relative depletion of cowcod south of 40°10' N lat. from 1970 to 2013 based on the 2013 stock assessment. .... 47

Figure 1-10. Time series of median harvest rates (total catch divided by age-11 and older biomass) from the base model in the 2013 cowcod assessment. The gray line is the estimated median harvest rate producing MSY. .... 49

Figure 1-11. Relative depletion of darkblotched rockfish from 1980 to 2013 based on the 2013 stock assessment. .... 52

Figure 1-12. Estimated recruitments of darkblotched rockfish, 1970-2012. .... 53

Figure 1-13. Time series of estimated SPR harvest rates of darkblotched rockfish, 1960-2012. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis..... 54

Figure 1-14. Relative depletion of Pacific ocean perch from 1980 to 2011 based on the 2011 stock assessment. .... 57

Figure 1-15. Time series of estimated (age-0) POP recruitments..... 58

Figure 1-16. Time series of POP exploitation rates (catch/summary biomass), 1940-2010. .... 59

Figure 1-17. Relative depletion trend from 1980 to 2013 for petrale sole based on the 2013 stock assessment. .... 63

Figure 1-18. Time series of estimated (age 0) petrale sole recruitments, 1934-2012 (from Haltuch, *et al.* 2013). .... 64

Figure 1-19. Estimated spawning potential ratio (SPR) of petrale sole, 2004-2012. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis. The management target is plotted as a dashed horizontal line and values above this reflect harvests in excess of the overfishing proxy based on the  $F_{MSY}$  harvest rate ( $SPR = 30\%$ )..... 65

Figure 1-20. Relative depletion of yelloweye rockfish from 1980 to 2011 based on the 2011 stock assessment update. .... 68

Figure 1-21. Time series of estimated yelloweye rockfish spawning output and recruitments for the base-case model in the 2011 assessment (Taylor and Wetzel 2011). .... 69

Figure 1-22. Time series of estimated relative exploitation rates (catch/biomass of age 8+ fish) of yelloweye rockfish, 1916-2010 (Taylor and Wetzel 2011). ..... 70

Figure 1-23. Trend in median fishing intensity (relative to the SPR management target) of Pacific whiting through 2013 with 95% posterior credibility intervals. The management target defined in the Agreement is shown as a horizontal line at 1.0. .... 97

Figure 1-24. Trend in median exploitation fraction through 2013 with 95% posterior credibility intervals. .... 98

Figure 1-25. Time series of estimated recruitments from the base model in the 2011 widow rockfish assessment. .... 116

Figure 1-26. Estimated total catch of China rockfish north of 40°10' N lat. in 2004-2012 relative to the preferred 2015 OFL contribution and ACL contribution. .... 124

Figure 1-27. Time series of estimated spawning biomass and depletion of aurora rockfish, 1916-2013. .... 135

Figure 1-28. Time series of estimated age-0 recruits of aurora rockfish on the U.S. west coast, 1970-2013. .... 136

Figure 1-29. Time series of estimated relative spawning potential ratio (1-SPR/1-SPR(Target=0.50)) for aurora rockfish, 1916-2012. Values of relative SPR above 1.0 reflect harvests in excess of the current overfishing proxy. .... 137

Figure 1-30. Estimated coastwide total annual catch of aurora rockfish in 2002-2012 relative to the proposed 2015 OFLs and 2015 ABCs (summed north and south of 40°10' N lat. to compare to coastwide catches). .... 137

Figure 1-31. Time series of estimated spawning output and depletion of blackgill rockfish south of 40°10' N lat., 1950-2011. .... 143

Figure 1-32. Time series of estimated summary harvest rate for the blackgill rockfish south of 40°10' N lat., 1950-2011. The dashed line is the harvest rate at the overfishing  $F_{MSY}$  proxy. .... 144

Figure 1-33. Estimated coastwide total annual catch of blackgill rockfish in 2002-2012 relative to the proposed 2015 OFLs, 2015 ABCs, and 2015 ACLs (summed north and south of 40°10' N lat. to compare to coastwide catches). .... 144

Figure 1-34. Time series of estimated spawning biomass and depletion of rougheye/blackspotted rockfish, 1916-2013 (from Hicks et al. 2013). .... 146

Figure 1-35. Time series of estimated age-0 recruits of rougheye/blackspotted rockfish on the U.S. west coast, 1980-2013(from Hicks et al. 2013). .... 147

Figure 1-36. Time-series of estimated summary harvest rate for the west coast stocks of rougheye and blackspotted rockfish, 1916-2012. The dotted line is the harvest rate at the overfishing  $F_{MSY}$  proxy. . 148

Figure 1-37. Estimated coastwide total annual catch of rougheye and blackspotted rockfish in 2002-2012 relative to the proposed 2015 OFLs and 2015 ABCs (summed north and south of 40°10' N lat. to compare to coastwide catches). .... 149

Figure 1-38. Relationship between SPR and instantaneous fishing mortality rate (F) for a hypothetical rockfish. .... 162

Figure 1-39. Conceptual diagram of the 25-5 and 40-10 ACL harvest control rules used to manage assessed west coast flatfish and other groundfish species, respectively, that are in the precautionary zone. .... 187



# CHAPTER 1      **GROUNDFISH STOCKS, THEIR STATUS, AND DESCRIPTION OF THE MANAGEMENT SYSTEM**

---

## **1.1      Description and Status of Groundfish Stocks**

There are over 90 stocks managed under the Pacific Coast Groundfish Fishery Management Plan (FMP). The actual number of FMP stocks is equivocal since all endemic species of the genus *Sebastes* are included and new species of this diverse genus are periodically described in the literature providing results of genetic/taxonomic research. These species include over 64 species of rockfish in the family *Scorpaenidae*, 7 roundfish species, 12 flatfish species, assorted chondrichthyans (sharks, skates, and ratfishes), and a few miscellaneous bottom-dwelling marine fish species. Table 1-1 depicts the latitudinal and depth distributions of groundfish species managed under the groundfish FMP, and Figure 1-1 depicts management area divisions.

The following sections contain information on the life histories of a subset of the groundfish managed under the groundfish FMP. While reading these sections, it is important to keep in mind how certain life history traits of the species have important implications on how the stocks are sustainably managed.

In contrast to the highly variable, and often volatile, population cycles of many coastal pelagic and invertebrate populations in the California Current, many of the resident groundfish in the California Current have evolved entirely different life history approaches to coping with environmental variability. Sablefish, Dover sole, spiny dogfish and a large number of rockfish (*Sebastes* and *Sebastolobus*) species have life spans that typically span decades, and in some extreme examples may reach ages of 100 or greater (Beamish, *et al.* 2006; Love, *et al.* 2002). Although large initial catches of many rockfish had given the impression that these stocks were also highly productive, a growing body of scientific evidence soon made it clear that many of these species were incapable of sustaining high intensity fishing pressure using modern fishing methods (Francis 1986; Gunderson 1977; Gunderson 1984; Leaman and Beamish 1984).

Among the concerns raised in some of the early research and analyses were that the large standing stocks of older individuals were simply maintaining themselves within the dynamic bounds of their ecosystem, and that the failure to consider the role of such longevity in Northeast Pacific groundfish could lead to management challenges. Factors such as extreme longevity, low natural mortality, increasing fecundity with age, and infrequent reproductive success (recruitment) were explicitly considered when initial harvest rate strategies were developed for the Council (Clark 1991). However, the paucity of data and magnitude of some of these factors as related to the low productivity of many species were not fully appreciated in many early studies, and are now known to be important considerations in developing harvest rate guidelines and management policies (Clark 2002; Dorn 2002a). Consequently, harvest rates for many species have been reduced repeatedly in recent years to account for the improved knowledge regarding the overall productivity of these stocks. As new information continues to emerge regarding the significance of diverse age structures and other factors in sustaining groundfish resources (Berkeley 2004; Berkeley, *et al.* 2004; Bobko and Berkeley 2004), such information continues to be evaluated and incorporated into the stock assessment and assessment review processes that provide the scientific basis upon which management decisions are made.

Management of these groundfish species is based on principles outlined in the Magnuson-Stevens Fishery Management and Conservation Act (MSA), groundfish FMP, and National Standard Guidelines, which

provide guidance on the 10 national standards in the MSA. Stock assessments are based on resource surveys, catch trends in west coast fisheries, and other data sources.

**Table 1-1. Latitudinal and depth distributions of groundfish species (adults) managed under the Pacific Coast Groundfish Fishery Management Plan.** <sup>a/</sup>

Common name	Scientific name	Latitudinal Distribution		Depth Distribution (fm)	
		Overall	Highest Density	Overall	Highest Density
<b>Flatfish Species</b>					
Arrowtooth flounder	<i>Atheresthes stomias</i>	N 34° N lat.	N 40° N lat.	10-400	27-270
Butter sole	<i>Isopsetta isolepis</i>	N 34° N lat.	N 34° N lat.	0-200	0-100
Curlfin sole	<i>Pleuronichthys decurrens</i>	Coastwide	Coastwide	4-291	4-50
Dover sole	<i>Microstomus pacificus</i>	Coastwide	Coastwide	10-500	110-270
English sole	<i>Parophrys vetulus</i>	Coastwide	Coastwide	0-300	40-200
Flathead sole	<i>Hippoglossoides elassodon</i>	N 38° N lat.	N 40° N lat.	3-300	100-200
Pacific sanddab	<i>Citharichthys sordidus</i>	Coastwide	Coastwide	0-300	0-82
Petrale sole	<i>Eopsetta jordani</i>	Coastwide	Coastwide	10-250	160-250
Rex sole	<i>Glyptocephalus zachirus</i>	Coastwide	Coastwide	10-350	27-250
Rock sole	<i>Lepidopsetta bilineata</i>	Coastwide	N 32°30' N lat.	0-200	summer 10-44 winter 70-150
Sand sole	<i>Psettichthys melanostictus</i>	Coastwide	N 33°50' N lat.	0-100	0-44
Starry flounder	<i>Platichthys stellatus</i>	Coastwide	N 34°20' N lat.	0-150	0-82
<b>Rockfish Species <sup>b/</sup></b>					
Aurora rockfish	<i>Sebastes aurora</i>	Coastwide	Coastwide	45-420	160-270
Bank rockfish	<i>Sebastes rufus</i>	S. 39°30' N lat.	S. 39°30' N lat.	17-135	115-140
Black rockfish	<i>Sebastes melanops</i>	N 34° N lat.	N 34° N lat.	0-200	0-30
Black-and-yellow rockfish	<i>Sebastes chrysomelas</i>	S. 40° N lat.	S. 40° N lat.	0-20	0-10
Blackgill rockfish	<i>Sebastes melanostomus</i>	Coastwide	S. 40° N lat.	48-420	125-300
Blackspotted rockfish	<i>Sebastes melanostictus</i>	Coastwide	N 40° N lat.	27-400	27-250
Blue rockfish	<i>Sebastes mystinus</i>	Coastwide	Coastwide	0-300	13-50
Bocaccio <sup>c/</sup>	<i>Sebastes paucispinis</i>	Coastwide	S. 40° N lat., N 48° N lat.	15-180	54-82
Bronzespotted rockfish	<i>Sebastes gilli</i>	S. 37° N lat.	S. 37° N lat.	41-205	110-160
Brown rockfish	<i>Sebastes auriculatus</i>	Coastwide	S. 40° N lat.	0-70	0-50
Calico rockfish	<i>Sebastes dallii</i>	S. 38° N lat.	S. 33° N lat.	10-140	33-50
California scorpionfish	<i>Scorpaena gutatta</i>	S. 37° N lat.	S. 34°27' N lat.	0-100	0-100
Canary rockfish	<i>Sebastes pinniger</i>	Coastwide	Coastwide	27-460	50-100
Chameleon rockfish	<i>Sebastes phillipsi</i>	37°-33° N lat.	37°-33° N lat.	95-150	95-150

Common name	Scientific name	Latitudinal Distribution		Depth Distribution (fm)	
		Overall	Highest Density	Overall	Highest Density
Chilipepper rockfish	<i>Sebastes goodei</i>	Coastwide	34°-40° N lat.	27-190	27-190
China rockfish	<i>Sebastes nebulosus</i>	N 34° N lat.	N 35° N lat.	0-70	2-50
Copper rockfish	<i>Sebastes caurinus</i>	Coastwide	S. 40° N lat.	0-100	0-100
Cowcod	<i>Sebastes levis</i>	S. 40° N lat.	S. 34°27' N lat.	22-270	100-130
Darkblotched rockfish	<i>Sebastes crameri</i>	N 33° N lat.	N 38° N lat.	16-300	96-220
Dusky rockfish <sup>d/</sup>	<i>Sebastes ciliatus</i>	N 55° N lat.	N 55° N lat.	0-150	0-150
Dwarf-Red rockfish	<i>Sebastes rufinanus</i>	33° N lat.	33° N lat.	>100	>100
Flag rockfish	<i>Sebastes rubrivinctus</i>	S. 38° N lat.	S. 37° N lat.	17-100	shallow
Freckled rockfish	<i>Sebastes lentiginosus</i>	S. 33° N lat.	S. 33° N lat.	22-92	22-92
Gopher rockfish	<i>Sebastes carnatus</i>	S. 40° N lat.	S. 40° N lat.	0-45	5-20
Grass rockfish	<i>Sebastes rastrelliger</i>	S. 44°40' N lat.	S. 40° N lat.	0-25	0-8
Greenblotched rockfish	<i>Sebastes rosenblatti</i>	S. 38° N lat.	S. 38° N lat.	33-217	115-130
Greenspotted rockfish	<i>Sebastes chlorostictus</i>	S. 47° N lat.	S. 40° N lat.	27-110	50-100
Greenstriped rockfish	<i>Sebastes elongatus</i>	Coastwide	Coastwide	33-220	27-136
Halfbanded rockfish	<i>Sebastes semicinctus</i>	S. 36°40' N lat.	S. 36°40' N lat.	32-220	32-220
Harlequin rockfish <sup>e/</sup>	<i>Sebastes variegatus</i>	N 40° N lat.	N 51° N lat.	38-167	38-167
Honeycomb rockfish	<i>Sebastes umbrosus</i>	S. 36°40' N lat.	S. 34°27' N lat.	16-65	16-38
Kelp rockfish	<i>Sebastes atrovirens</i>	S. 39° N lat.	S. 37° N lat.	0-25	3-4
Longspine thornyhead	<i>Sebastes altivelis</i>	Coastwide	Coastwide	167->833	320-550
Mexican rockfish	<i>Sebastes macdonaldi</i>	S. 36°20' N lat.	S. 36°20' N lat.	50-140	50-140
Olive rockfish	<i>Sebastes serranoides</i>	S. 41°20' N lat.	S. 40° N lat.	0-80	0-16
Pacific ocean perch	<i>Sebastes alutus</i>	Coastwide	N 42° N lat.	50-450	110-250
Pink rockfish	<i>Sebastes eos</i>	S. 37° N lat.	S. 35° N lat.	40-200	40-200
Pinkrose rockfish	<i>Sebastes simulator</i>	S. 34° N lat.	S. 34° N lat.	54-160	108
Puget Sound rockfish	<i>Sebastes emphaeus</i>	N 40° N lat.	N 40° N lat.	6-200	6-200
Pygmy rockfish	<i>Sebastes wilsoni</i>	N 32°30' N lat.	N 32°30' N lat.	17-150	17-150
Quillback rockfish	<i>Sebastes maliger</i>	N 36°20' N lat.	N 40° N lat.	0-150	22-33
Redbanded rockfish	<i>Sebastes babcocki</i>	Coastwide	N 37° N lat.	50-260	82-245
Redstripe rockfish	<i>Sebastes proriger</i>	N 37° N lat.	N 37° N lat.	7-190	55-190
Rosethorn rockfish	<i>Sebastes helvomaculatus</i>	Coastwide	N 38° N lat.	65-300	55-190
Rosy rockfish	<i>Sebastes rosaceus</i>	S. 42° N lat.	S. 40° N lat.	8-70	30-58
Rougheye rockfish	<i>Sebastes aleutianus</i>	Coastwide	N 40° N lat.	27-400	27-250

Common name	Scientific name	Latitudinal Distribution		Depth Distribution (fm)	
		Overall	Highest Density	Overall	Highest Density
Semaphore rockfish	<i>Sebastes melanosema</i>	S. 34°27' N lat.	S. 34°27' N lat.	75-100	75-100
Sharpchin rockfish	<i>Sebastes zacentrus</i>	Coastwide	Coastwide	50-175	50-175
Shortbelly rockfish	<i>Sebastes jordani</i>	Coastwide	S. 46° N lat.	50-175	50-155
Shortraker rockfish	<i>Sebastes borealis</i>	N 39°30' N lat.	N 44° N lat.	110-220	110-220
Shortspine thornyhead	<i>Sebastes alascanus</i>	Coastwide	Coastwide	14->833	55-550
Silvergray rockfish	<i>Sebastes brevispinis</i>	Coastwide	N 40° N lat.	17-200	55-160
Speckled rockfish	<i>Sebastes ovalis</i>	S. 38° N lat.	S. 37° N lat.	17-200	41-83
Splitnose rockfish	<i>Sebastes diploproa</i>	Coastwide	Coastwide	50-317	55-250
Squarespot rockfish	<i>Sebastes hopkinsi</i>	S. 38° N lat.	S. 36° N lat.	10-100	10-100
Starry rockfish	<i>Sebastes constellatus</i>	S. 38° N lat.	S. 37° N lat.	13-150	13-150
Stripetail rockfish	<i>Sebastes saxicola</i>	Coastwide	Coastwide	5-230	5-190
Swordspine rockfish	<i>Sebastes ensifer</i>	S. 38° N lat.	S. 38° N lat.	38-237	38-237
Tiger rockfish	<i>Sebastes nigrocinctus</i>	N 35° N lat.	N 35° N lat.	30-170	35-170
Treefish	<i>Sebastes serriceps</i>	S. 38° N lat.	S. 34°27' N lat.	0-25	3-16
Vermilion rockfish	<i>Sebastes miniatus</i>	Coastwide	Coastwide	0-150	4-130
Widow rockfish	<i>Sebastes entomelas</i>	Coastwide	N 37° N lat.	13-200	55-160
Yelloweye rockfish	<i>Sebastes ruberrimus</i>	Coastwide	N 36° N lat.	25-300	27-220
Yellowmouth rockfish	<i>Sebastes reedi</i>	N 40° N lat.	N 40° N lat.	77-200	150-200
Yellowtail rockfish	<i>Sebastes flavidus</i>	Coastwide	N 37° N lat.	27-300	27-160
<b>Roundfish Species</b>					
Cabezon	<i>Scorpaenichthys marmoratus</i>	Coastwide	Coastwide	0-60	0-27
Kelp greenling	<i>Hexagrammos decagrammus</i>	Coastwide	N 40° N lat.	0-25	0-10
Lingcod	<i>Ophiodon elongatus</i>	Coastwide	Coastwide	0-233	0-40
Pacific cod	<i>Gadus macrocephalus</i>	N 34° N lat.	N 40° N lat.	7-300	27-160
Pacific whiting	<i>Merluccius productus</i>	Coastwide	Coastwide	20-500	27-270
Sablefish	<i>Anoplopoma fimbria</i>	Coastwide	Coastwide	27->1,000	110-550
<b>Shark and Skate Species</b>					
Big skate	<i>Raja binoculata</i>	Coastwide	S. 46° N lat.	2-110	27-110
California skate	<i>Raja inornata</i>	Coastwide	S. 39° N lat.	0-367	0-10
Leopard shark	<i>Triakis semifasciata</i>	S. 46° N lat.	S. 46° N lat.	0-50	0-2
Longnose skate	<i>Raja rhina</i>	Coastwide	N 46° N lat.	30-410	30-340
Southern shark	<i>Galeorhinus zyopterus</i>	Coastwide	Coastwide	0-225	0-225

Common name	Scientific name	Latitudinal Distribution		Depth Distribution (fm)	
		Overall	Highest Density	Overall	Highest Density
Spiny dogfish	<i>Squalus suckleyi</i>	Coastwide	Coastwide	0->640	0-190
<b>Other Species</b>					
Finescale codling	<i>Antimora microlepis</i>	Coastwide	N 38° N lat.	190-1,588	190-470
Pacific rattail	<i>Coryphaenoides acrolepis</i>	Coastwide	N 38° N lat.	85-1,350	500-1,350
Ratfish	<i>Hydrolagus colliciei</i>	Coastwide	Coastwide	0-499	55-82
<p>a/ Data from (Casillas, <i>et al.</i> 1998), (Eschmeyer, <i>et al.</i> 1983), (Hart 1988), (Miller and Lea 1972), (Love, <i>et al.</i> 2002), and NMFS survey data. Depth distributions refer to offshore distributions, not vertical distributions in the water column.</p> <p>b/ The category “rockfish” includes all genera and species of the family Scorpaenidae, even if not listed, that occur in the Washington, Oregon, and California area.</p> <p>c/ Only the southern stock of bocaccio south of 40° 10' N lat. is listed as overfished.</p> <p>d/ Only two occurrences of harlequin rockfish south of 51° N lat. (off Newport, OR and La Push, WA; (Casillas, <i>et al.</i> 1998)).</p>					



Figure 1-1. Fishery management lines on the U.S. west coast.

The passage of the Sustainable Fisheries Act in 1996 and the reauthorization of the MSA in 2006<sup>1</sup> incorporated the current conservation and rebuilding mandates into the MSA. These mandates—including abundance-based standard reference points for declaring the status of a stock (overfished; in a “precautionary” status; or at levels that can support maximum sustainable yield (MSY) (healthy or “rebuilt”))—were subsequently incorporated in the groundfish FMP with adoption of Amendments 11, 12, and 23. These reference points are determined relative to an estimate of “virgin” or unexploited spawning biomass of the stock, denoted as  $B_0$ , which is defined as the average equilibrium abundance of a stock’s spawning biomass before it is affected by fishing-related mortality.<sup>2</sup>  $B_0$  is then used to estimate MSY, as identified in the MSA and National Standard Guidelines. MSY represents a theoretical maximum surplus production from a population of constant size; National Standard Guidelines define it as “the largest long-term average catch or yield that can be taken from a stock or stock complex under prevailing ecological and environmental conditions.” For a given population and set of ecological conditions, there is a biomass that produces MSY (denoted as  $B_{MSY}$ ), which is less than the equilibrium size in the absence of fishing ( $B_0$ ). (Generally, population sizes above  $B_{MSY}$  are assumed to be less productive because of competition for resources or other density dependent factors.) The harvest rate used to achieve or sustain  $B_{MSY}$  is referred to as the Maximum Fishing Mortality Threshold (MFMT, denoted as  $F_{MSY}$ ). Three harvest specification reference points, defined in the groundfish FMP, provide guidance in setting the harvest rate: an overfishing limit (OFL), an acceptable biological catch (ABC), and an annual catch limit (ACL) (see section 1.2 for more information on harvest specifications). The Council identifies the ACL as the management target for each species or species complex. When the stock biomass is determined to be lower than  $B_{MSY}$ , the ACL is set to an adequately low level to rebuild the stock to a healthy level in a timely fashion.

The biomass level that produces MSY (i.e.,  $B_{MSY}$ ) is generally unknown and assumed to be variable over time due to long-term fluctuations in ocean conditions, so that no single value is appropriate. Furthermore,  $F_{MSY}$  is tightly linked to an assumed level of density dependence in recruitment, and there is insufficient information to determine that level for many west coast groundfish stocks. Therefore, the use of approximations or proxies is necessary; absent a more accurate determination of  $F_{MSY}$ , the Council applies default MSY proxies (see section 1.1.1 for more details). The Council adopts management actions aimed to maintain abundance of each stock at or above the specified  $B_{MSY}$  target. The threshold for declaring a stock overfished is when the stock’s spawning biomass declines to less than the specified Minimum Stock Size Threshold or MSST (i.e., 12.5% of  $B_0$  or  $B_{12.5\%}$  for assessed flatfish stocks and  $B_{25\%}$  for all other groundfish stocks). A rebuilding plan that specifies how total fishing-related mortality is constrained to achieve an MSY abundance level within the legally allowed time is required by the MSA and groundfish FMP when a stock is declared overfished.

Of the more than 90 species managed under the groundfish FMP, only a portion are individually managed. Thus, the remaining species are managed and accounted for in groupings or stock complexes (see section 1.1.5) because individually they comprise a small part of the landed catch and, in general, insufficient information exists to develop the stock assessments necessary to set harvest specifications based on yield estimates. The Council has also decided to continue to manage some assessed stocks in complexes to avoid management complications such as disruption to the trawl rationalization program. Catch-based methods described in section 1.1.1 are used to set OFLs for unassessed stocks. Additionally, there is a category of stocks that are incidentally caught in groundfish fisheries for which no harvest limits are

---

<sup>1</sup> The Magnuson-Stevens Act is again up for reauthorization in 2014.

<sup>2</sup> The current abundance of a stock relative to its unfished level is commonly written as a percentage or a proportion; this value represents the stock’s depletion level. In addition to using a comparison between current spawning biomass and unfished spawning biomass to determine this reference point, some stock assessment authors compare current and unfished levels of spawning output or of total stock biomass, depending on the information that is available.

specified. This category of stocks, termed Ecosystem Component (EC) species, are not considered to be in the fishery and are neither targeted nor generally retained for sale or personal use. EC species are determined not to likely become subject to overfishing or to be overfished in the absence of conservation and management measures. There is a monitoring requirement for species designated as EC to the extent that any new pertinent scientific information becomes available (e.g., catch trends, vulnerability, etc.) to determine changes in their status or their vulnerability to the fishery. The Council is proposing an EC designation for some species currently managed in the FMP, as well as other non-FMP species (see section 1.1.6).

When the total fishing mortality (i.e., landed catch plus dead discards) of a west coast groundfish stock or stock complex exceeds the specified OFL for that stock or complex, the stock is considered to be subject to overfishing. Total mortality is estimated by the NMFS Northwest Fisheries Science Center and reported for all managed west coast stocks and complexes in total mortality reports found at [http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data\\_products/species\\_management.cfm](http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data_products/species_management.cfm). There are currently no west coast groundfish stocks or complexes that are subject to overfishing. Summaries of the status of west coast groundfish stocks and complexes (and the other federally managed stocks and complexes nationally), are provided by the NMFS Fish Stock Sustainability Index (FSSI) at [http://www.nmfs.noaa.gov/sfa/fisheries\\_eco/status\\_of\\_fisheries/status\\_updates.html](http://www.nmfs.noaa.gov/sfa/fisheries_eco/status_of_fisheries/status_updates.html).

### **1.1.1 Productivity and Susceptibility Assessment of Stocks to Overfishing**

The vulnerability to potential overfishing of a stock to the fishery for each groundfish stock in the FMP was defined as a first step in assisting with two specific tasks set forth in the FMP: 1) to define species as either “in the fishery” or as an “ecosystem component,” and 2) identify stock complexes. In addition, the vulnerability scores were considered when prioritizing stock assessments and determining data collection needs.

The Productivity-Susceptibility Assessment (PSA) approach of Patrick et al. (2009) was used to characterize vulnerability and has two components: 1) productivity as defined by life histories traits, and 2) susceptibility to current fishing practices. Each vulnerability component is comprised of several attributes (10 productivity and 12 susceptibility attributes) and the weighted mean score of all attributes defines the overall productivity and susceptibility score. Table 1-2 includes the vulnerability scores for all species in the FMP relative to the current fishery. Table 1-2 shows the vulnerability scores for currently overfished rockfish species relative to the fishery circa 1998. Scores are presented in two-dimensions, with productivity on the x-axis and susceptibility on the y-axis (Figure 1-2). Cope et al. (2011) established vulnerability reference points of assessed and unassessed west coast groundfish stocks to determine vulnerability groups as follows:

- $V \geq 2.2$  indicate species of major concern.
- $2.0 \leq V < 2.2$  indicate species of high concern.
- $1.8 \leq V < 2.0$  indicate species of medium concern.
- $V < 1.8$  indicate species of low concern.

Rockfish and elasmobranchs showed the highest vulnerabilities ( $>2.0$ ), with the deepest-residing members of those groups often the most vulnerable, though there were several species of nearshore rockfish (China, quillback, and copper rockfish) with some of the highest scored vulnerabilities. Flatfishes in general showed the lowest vulnerabilities.

In addition to scoring each productivity and susceptibility attribute, the quality of the data used for each score was also recorded (Table 1-2, Table 1-3, and Figure 1-3). Data quality is scored for each



productivity and susceptibility attribute, with the overall data quality score calculated as the weighted mean of all attributes. A scoring scale of 1-5 was used, with the best data score being 5.

Recording the data quality can highlight vulnerability scores that can be improved with additional data or that should be interpreted with caution because of questionable data contribution. Data quality scores can also be used to justify future data collection on particular attributes.

In general, susceptibility was harder to score (lower data quality) than productivity. Flatfishes as a group had the least informed species, but elasmobranches and several rockfish species also showed low-quality data informing vulnerability scores (Table 1-2).

PSA analyses are anticipated to be re-done every biennial specifications cycle. Productivity scores are not expected to vary much over time since they are based on life history traits. However, susceptibility scores may vary based on changes in fishing practices and/or management, and an updated understanding of the stock's interaction with the fishery. As susceptibility scores change, so do the vulnerability scores.

**Table 1-2. Overall scores and results of the Productivity and Susceptibility Assessment (PSA) ranked from most to least vulnerable to overfishing relative to the current west coast fishery based on the GMT's scoring.**


Stock ID	Stock Name	Productivity	Susceptibility	Vulnerability
21	Copper rockfish	1.95	1.60	2.27
67	Rougeye rockfish	1.17	2.33	2.27
72	Shortraker rockfish	1.22	2.38	2.25
20	China rockfish	1.33	2.29	2.23
58	Quillback rockfish	1.31	2.43	2.22
61	Redstripe rockfish	1.31	2.33	2.16
22	Cowcod	1.25	2.00	2.13
77	Spiny dogfish	1.11	1.98	2.13
10	Bronzespotted rockfish	1.37	2.14	2.12
16	California skate	1.33	2.00	2.12
35	Greenblotched rockfish	1.28	2.24	2.12
2	Aurora rockfish	1.89	2.29	2.10
76	Speckled rockfish	1.33	2.29	2.10
65	Rosethorn rockfish	1.19	2.05	2.09
81	Starry rockfish	1.25	2.14	2.09
7	Blackgill rockfish	1.22	2.08	2.08
84	Tiger rockfish	1.25	2.10	2.06
70	Sharpchin rockfish	1.36	2.24	2.05
86	Vermilion rockfish	1.22	2.02	2.05
87	Widow rockfish	1.31	2.16	2.05
18	Chameleon rockfish	1.39	2.20	2.03
3	Bank rockfish	1.28	1.88	2.02
55	Pink rockfish	1.33	2.14	2.02
60	Redbanded rockfish	1.28	2.05	2.02
74	Silvergray rockfish	1.22	1.95	2.02
75	Soupfin shark	1.11	1.71	2.02
8	Blue rockfish	1.22	2.16	2.01

<b>Stock ID</b>	<b>Stock Name</b>	<b>Productivity</b>	<b>Susceptibility</b>	<b>Vulnerability</b>
17	Canary rockfish	1.61	2.43	2.01
43	Leopard shark	1.26	2.00	2.00
88	Yelloweye rockfish	1.22	1.92	2.00
4	Big skate	2.45	2.05	1.99
11	Brown rockfish	1.72	2.08	1.99
26	Dusky rockfish	1.75	1.76	1.99
36	Greenspotted rockfish	1.39	2.14	1.98
30	Flag rockfish	1.83	1.80	1.97
40	Honeycomb rockfish	1.36	2.10	1.97
89	Yellowmouth rockfish	1.61	2.38	1.96
5	Black rockfish	1.21	2.14	1.94
39	Harlequin rockfish	1.31	1.95	1.94
54	Petrale sole	1.70	2.44	1.94
83	Swordspine rockfish	1.33	2.00	1.94
9	Bocaccio	1.28	2.04	1.93
24	Darkblotched rockfish	1.39	2.24	1.92
34	Grass rockfish	1.61	2.29	1.89
66	Rosy rockfish	1.61	2.29	1.89
37	Greenstriped rockfish	1.28	1.76	1.88
90	Yellowtail rockfish	1.33	1.88	1.88
48	Olive rockfish	1.69	2.33	1.87
79	Squarespot rockfish	1.61	2.24	1.86
51	Pacific grenadier	1.44	1.95	1.82
56	Pinkrose rockfish	1.31	1.67	1.82
78	Splitnose rockfish	1.28	1.60	1.82
47	Mexican rockfish	1.50	2.00	1.80
73	Shortspine thornyhead	1.33	1.68	1.80
82	Stripetail rockfish	1.39	1.81	1.80
63	Rock greenling	1.78	2.29	1.77
33	Gopher rockfish	1.56	2.00	1.76
85	Treefish	1.67	2.10	1.73
59	Ratfish	1.63	2.05	1.72
6	Black-and-yellow rockfish	1.83	1.68	1.70
50	Pacific ocean perch	1.44	1.67	1.69
53	Pacific whiting	2.00	2.36	1.69
13	Cabazon	1.33	2.48	1.68
45	Longnose skate	1.53	1.80	1.68
68	Sablefish	1.61	1.88	1.64
42	Kelp rockfish	1.83	2.12	1.62
41	Kelp greenling	1.83	2.04	1.56
44	Lingcod	1.75	1.92	1.55
25	Dover sole	1.36	2.57	1.54
27	Dwarf-red rockfish	1.06	1.88	1.54
46	Longspine thornyhead	1.47	1.16	1.54


Stock ID	Stock Name	Productivity	Susceptibility	Vulnerability
29	Finescale codling	2.45	2.10	1.48
14	Calico rockfish	1.39	2.04	1.46
32	Freckled rockfish	1.80	1.96	1.44
57	Pygmy rockfish	1.78	1.71	1.42
64	Rock sole	1.95	1.95	1.42
15	California scorpionfish	1.28	0.00	1.41
19	Chilipepper	1.83	0.00	1.35
49	Pacific cod	2.11	2.00	1.34
62	Rex sole	2.05	1.86	1.28
31	Flathead sole	2.25	1.92	1.26
38	Halfbanded rockfish	2.00	1.76	1.26
52	Pacific sanddab	2.40	2.10	1.25
23	Curlfin sole	1.72	1.75	1.23
69	Sand sole	2.35	2.05	1.23
1	Arrowtooth flounder	1.33	2.05	1.21
28	English sole	2.30	2.05	1.19
12	Butter sole	1.78	1.76	1.18
71	Shortbelly rockfish	1.94	1.40	1.13
80	Starry flounder	2.15	1.60	1.04

**Table 1-3. Retrospective Productivity and Susceptibility Assessment (PSA) vulnerability scores of currently overfished rockfish species ranked from most to least vulnerable to overfishing relative to stock status and the fishery circa 1998, based on the GMT's scoring.**

Stock Name	Stock ID	Susceptibility	Vulnerability
Bocaccio	25_H	2.72	2.43
Canary	23_H	2.84	2.52
Cowcod	10_H	2.68	2.57
Darkblotched	51_H	2.76	2.39
POP	92_H	2.32	2.08
Yelloweye	18_H	2.80	2.53



**Figure 1-2. Productivity and Susceptibility Analysis (PSA) plot for species in the west coast groundfish FMP. Contours delineate areas of relative vulnerability (V, i.e. distance from the origin), with the highest vulnerability stocks above the solid red line (V = 2.2), high vulnerability above the orange broken line (V=2), medium vulnerability above the green dotted line (V=1.8) and the lowest vulnerability below the green dotted line. The maximum vulnerability (V=2.8) is indicated with the solid black line. Solid circles are based on current PSA scores. Open circles are based on PSA scores circa 1998. Numbers refer to the Stock ID in Table 1-2 and Table 1-3.**



**Figure 1-3. Data quality plots for the productivity and susceptibility scores in the PSA for each species (represented numerically in Table 1-2 and Table 1-3) in the west coast groundfish FMP. Higher scores indicate less data quality. Vertical and horizontal lines provide a general guide to relative data quality with values above 3 on either axis considered data-poor.**

### **1.1.2 Stock Assessments and Rebuilding Analyses Used to Estimate Stock Status and Inform Management Decisions**

Stock assessments are used for setting harvest specifications by providing estimates of MSY, OFL, the MFMT, the MSST, and ABC. Stock assessments are also used to determine the status of a fish population or subpopulation (stock) terms of estimating population size, reproductive status (e.g., spawning biomass, fecundity, etc.), fishing mortality, and whether current catches are sustainable. In the terms of the Groundfish FMP, stock assessments provide: 1) an estimate of the current biomass and reproductive potential (generally expressed as spawning biomass), 2) an estimate of  $F_{MSY}$  (the harvest rate estimated to produce MSY) or proxy thereof translated into exploitation rate or spawning potential ratio (SPR; see section 1.3.1 for a description of SPR), 3) the estimated biomass corresponding to MSY ( $B_{MSY}$ ), or a proxy thereof, 4) estimated unfished biomass ( $B_0$ ), and 5) the estimated variance (or a confidence interval) for the estimate of current biomass. With the exception of Pacific whiting, which is assessed annually as specified in the Agreement with Canada on Pacific Hake/Whiting, groundfish stock assessments are conducted on a two-year cycle. Given the large number of groundfish species and limited state and Federal resources, a subset of all groundfish stocks are assessed in each stock assessment cycle. Overfished species' stock assessments are typically conducted every two years, although a catch report can be substituted for an assessment to monitor compliance with adopted rebuilding plans. The process for setting groundfish specifications involves the adoption of new and updated stock assessments. During the biennial specification process, the SSC reviews stock assessments and rebuilding analyses for overfished species and makes recommendations to the Council relative to the standards of the best available science and the soundness of the scientific information relative to management decisions. The Council then approves all or a portion of the stock assessments, or recommends further analysis.

The perception of stock status and productivity may change substantially between stock assessments. Such changes can result from technical changes in the assessment model, including how a given assessment model is structured, the assumptions used to fix or estimate key parameters (i.e., whether parameters such as natural mortality and steepness are fixed, estimated freely, or estimated with an informative prior), and the evolution of methods for developing data time series and estimates of uncertainty from different sources of raw data. The population dynamics of target species themselves are responsive to a mix of complex (and often poorly-understood) biological, oceanographic, and interspecies interactions. New data sources (e.g., new data types, extensions of existing data sets, incorporation of environmental factors into assessments) can result in changes in parameter estimates and model outputs.

All stock assessments are subject to a peer review process, consistent with the MSA (§302(g)(1)(E)). The process considers components of the assessments starting with data collection and continuing through to scientific recommendations and information presented to the Council and its advisors. The terms of reference for the groundfish stock assessment process defines the expectations and responsibilities for various participants in the groundfish stock assessment review (STAR) process, and outlines the guidelines and procedures for a peer review process. The STAR process is a key element in an overall process designed to review the technical merits of stock assessments and other scientific information used by the SSC. This process allows the Council to make timely use of new fishery and survey data, to analyze and understand these data as completely as possible, to provide opportunity for public comment, and to assure that the results are as accurate and error-free as possible.

Harvest specifications and the science used as the basis for management decision-making are derived from the most recent assessments and/or rebuilding analyses prepared for those stocks, informed by an assessment. The newest assessments were those prepared and adopted in 2013 and the oldest assessments informing management decisions for fisheries in 2015 and beyond were prepared and adopted in 2005. Table 1-4 presents a summary of the management quantities estimated by the base models of the most recent assessments informing management in 2015 and beyond. Table 1-5 lists life history parameters

from the stocks assessed since 2005, excluding those conducted using XDB-SRA; steepness of the spawner-recruitment curve ( $h$ ), recruitment variability ( $\sigma_r$ ), the von Bertalanffy Equation growth constant ( $k$ ), and natural mortality ( $M$ ) are each important contributors to the understanding of the productivity and resiliency of these stocks. Table 1-6 lists life history parameters from the stocks assessed in 2013 using XDB-SRA;  $B_{MSY}$ ,  $F_{MSY}$ ,  $M$ ,  $B_{MSY}/B_0$ , and  $F_{MSY}/M$  inform the relative productivity and resiliency of these stocks.

All stock assessments, STAR panel reports, and rebuilding analyses used to inform management decisions on west coast groundfish stocks and fisheries can be found on the Council's web site at <http://www.pcouncil.org/groundfish/stock-assessments/>.

#### 1.1.2.1 Types of Assessments Used in Managing Groundfish Stocks

The Council uses various types of assessments that range from data-rich full assessments (also known as benchmark assessments) to data-poor catch-based methods used to only estimate an OFL. The Council decides which groundfish stocks will be assessed and, based on SSC recommendations, what type of assessment will be used (i.e., full, update, data-moderate) each cycle. These stock assessment priorities are decided in even years and assessments are conducted, reviewed, and adopted in odd years. Results from these assessments are used to inform management decisions for the following biennial cycle, which begins in the next odd year. The SSC reviews all assessments and recommends to the Council if they represent the best available science for the stock, and whether and how they can be used to inform Council decisions.

The SSC categorizes stocks based on the type of assessment and the quality of data informing that assessment. The FMP harvest specification framework calls for increasing uncertainty buffers translated into lower ABCs (and ACLs) for stocks informed by less certain assessments (see section 1.3.2). Stock categories range from category 1, characterized by stocks informed by full assessments with reasonably good estimates of year class strength, to unassessed category 3 stocks where there is only a data-poor estimate of the OFL. A more detailed description of the assessment models used in current groundfish management follows.

#### **Data-Poor Assessments**

Data-poor assessments employ catch-based statistics to estimate an OFL for a stock. Stock status cannot be determined using these types of assessment since there are no survey or other abundance indices used in a data-poor assessment. The most rudimentary data-poor assessment is simply setting the OFL to a proportion of the average historical catch. However, there is great uncertainty whether that is a "true" OFL since the historical catch used to compute the average could have been unsustainably high. Therefore, the SSC categorizes stocks informed by a data-poor OFL as category 3 stocks, thus mandating a higher buffer to determine the ABC. While this category of data-poor methods are being characterized as "assessments" here, stocks with OFLs informed with data-poor methods are considered unassessed since there is no estimate of relative depletion or status. Other approved data-poor methods (DCAC and DB-SRA) more sophisticated than average catch are described below.

#### **Depletion-Corrected Average Catch**

The Depletion-Corrected Average Catch (DCAC) method provides an estimate of sustainable yield (taken as the OFL) for data-poor stocks of uncertain status (MacCall 2009). DCAC adjusts historical average catch to account for one-time "windfall" catches that are the result of stock depletion, producing an estimate of yield that was likely to be sustainable over the same time period. Advantages of the DCAC

approach for determining sustainable yield for data-poor stocks include: 1) relatively minimal data requirements (i.e., an historical catch time series), 2) biologically-based adjustment to catch-based yield proxies with transparent assumptions about relative changes in abundance (e.g., a production function with compensation exists for the stock), and 3) simplicity in computing.

### **Depletion-Based Stock Reduction Analysis**

The Depletion-Based Stock Reduction Analysis (DB-SRA) method extends the DCAC method by 1) restoring the temporal link between production and biomass, and 2) evaluating and integrating alternative hypotheses regarding changes in abundance during the historical catch period (Dick and MacCall 2011). This method combines DCAC's distributional assumptions regarding life history characteristics and stock status with the dynamic models and simulation approach of stochastic stock reduction analysis.

### **Simple Stock Synthesis**

A similar approach to DB\_SRA can also be conducted in Stock Synthesis (Cope 2013).

### **Data-Moderate Assessments**

Data-moderate assessments are less complicated than full assessments and can therefore be reviewed more expeditiously. Unlike a full assessment, which is reviewed by a STAR panel and the SSC, only the SSC reviews a data-moderate assessment<sup>3</sup>.

Data-moderate assessments combine catch-based methods with a time series of relative abundance estimates from one or more surveys or other types of abundance indices (e.g., CPUE time series). This type of assessment represents the minimal structure of an assessment used to determine stock status according to the NMFS National Stock Assessment Improvement plan (Mace, *et al.* 2001). These assessments exclude compositional age and length data, which are used to determine survey and/or fishery selectivities and to estimate other parameters in a full assessment model. The addition of compositional data complicates an assessment, requiring more review time to understand what data are driving model results. Data-moderate assessments were therefore developed to increase the number of groundfish stocks assessed given the resources available to conduct and review assessments each cycle. There are two data-moderate assessment models in current use that have been reviewed and recommended by the SSC: Extended Simple Stock Synthesis (exSSS) and Extended Depletion-based Stock Reduction Analysis (XDB-SRA). These are described in more detail below.

Since data-moderate assessments are less informative than full assessments, the SSC categorizes stocks informed with such assessment as category 2 stocks.

### **Extended Simple Stock Synthesis**

Extended Simple Stock Synthesis (exSSS) is based on sampling parameters (steepness, natural mortality and depletion) from prior distributions and using Stock Synthesis to solve for virgin recruitment (R0)

---

<sup>3</sup> While this is technically true, the SSC and Council elected to do a more rigorous review of data-moderate assessments in a STAR panel in 2013, the first year data-moderate assessments were conducted on the west coast.



given inputs for selectivity, growth, and fecundity. ExSSS extends Simple Stock Synthesis, originally a data-poor method reviewed by the SSC, by allowing index data (and potentially length and age data) to be used for parameter estimation using the Stock Synthesis platform. Parameter estimation for exSSS is based on the Adaptive Importance Sampling (AIS) methods (Cope, *et al.* In Press; Wetzel and Punt In Press). ExSSS assumes that recruitment is related deterministically to the stock-recruitment relationship. The outputs from exSSS include biomass trajectories, as well as estimates of (and measures of uncertainty for) the OFL. The prior for depletion is based on the results of a regression of depletion on the PSA vulnerability score (see section 1.1.1 and (Cope, *et al.* In Press).

### **Extended Depletion-Based Stock Reduction Analysis**

Extended Depletion-Based Stock Reduction Analysis (XDB-SRA), an extension of DB-SRA, is an assessment method approved by the SSC for use in data-moderate assessments. XDB-SRA can be implemented within a Bayesian framework, with the priors for the parameters updated based on index data. The additional parameters in XDB-SRA compared with DB-SRA include the catchability coefficient ( $q$ ) for each index of abundance, and the extent of observation variance additional to that inferred from sampling error ( $a$ ). The priors for these parameters are a weakly informative log-normal and a uniform distribution, respectively.

XDB-SRA was used in the 2013 full assessment of cowcod in the Southern California Bight (see section 1.1.3.3). While XDB-SRA is an approved data-moderate assessment model, it can also be parameterized to incorporate compositional data<sup>4</sup>.

### **Full Stock Assessments**

Full, or benchmark, stock assessments are those where Stock Assessment Teams (STATs) can propose new models and explore new data to determine the status and dynamics of a fish stock. The Council has a rigorous process for first determining those stocks that will be assessed and, once determined, how they will be reviewed (the process is codified in the Stock Assessment and Review Terms of Reference, which is updated every other year; available at <http://www.pccouncil.org/groundfish/stock-assessments/terms-of-reference/>). Full assessments are more vigorously reviewed than other types of assessments since they are inherently more complicated. A week-long Stock Assessment Review (STAR) panel meeting occurs with STATs presenting assessment models to a panel of experts (typically comprised of one SSC Groundfish Subcommittee member who chairs the meeting, one west coast groundfish assessment expert, and two independent reviewers from the Center of Independent Experts. Additionally, one Groundfish Management Team representative, one Groundfish Advisory Subpanel representative, and a member of the Council staff attends STAR panel meetings as advisors. The STAR panel prepares a report recommending whether the assessment is robust enough to be used in management, along with other detailed recommendations on how to interpret assessment results and how to improve the assessment next time it is conducted. STAR panel reports also detail the model and data explorations that occurred during the review. The draft assessment and STAR panel report are then reviewed by the SSC. The assessment is only adopted for use in management decision-making if recommended by the SSC.

Stocks assessed with SSC-endorsed assessments are categorized either as category 1 or category 2 depending on the quality of data informing the assessment, relative uncertainty of model estimates, and/or whether individual year class strength (i.e., recruitment) is estimated.

---

<sup>4</sup> Note that the 2013 cowcod assessment excluded compositional data within the model. However, the model was subject of the two-step (i.e., STAR panel and SSC) review process defined for full assessments.

## Stock Synthesis

Most of the groundfish assessments on the U.S. west coast currently used to inform management decisions have been conducted using Stock Synthesis (SS). Stock Synthesis provides a statistical framework for calibration of a population dynamics model using a diversity of fishery and survey data. It is designed to accommodate both age and size structure in the population and with multiple stock sub-areas. Selectivity can be cast as age specific only, size-specific in the observations only, or size-specific with the ability to capture the major effect of size-specific survivorship. The overall model contains subcomponents which simulate the population dynamics of the stock and fisheries, derive the expected values for the various observed data, and quantify the magnitude of difference between observed and expected data. Some SS features include ageing error, growth estimation, spawner-recruitment relationship, movement between areas. SS is most flexible in its ability to utilize a wide diversity of age, size, and aggregate data from fisheries and surveys. The ADMB C++ software in which SS is written searches for the set of parameter values that maximize the goodness-of-fit, then calculates the variance of these parameters using inverse Hessian and MCMC methods. A management layer is also included in the model allowing uncertainty in estimated parameters to be propagated to the management quantities, thus facilitating a description of the risk of various possible management scenarios, including forecasts of possible annual catch limits. The structure of Stock Synthesis allows for building of simple to complex models depending upon the data available. The latest version of SS used in most of the assessments done in 2013 is version 3.24f (download available at <http://nft.nefsc.noaa.gov/SS3.html>).

## Update Assessments

An update assessment uses the model structure of the stock's last full, SSC-endorsed assessment, but is generally restricted to the addition of new data that have become available since the last full assessment. It must carry forward the fundamental structure of the last full assessment reviewed and endorsed by a STAR panel, the SSC, and the Council. Assessment structure here refers to the population dynamics model, data sources used as inputs to the model, the statistical platform used to fit model to the data, and how the management quantities used to set harvest specifications are calculated. Particularly, when an update assessment is developed, no substantial changes should be made to 1) the sources of data used, 2) the software used in programming the assessment, 3) the assumptions and structure of the population dynamics model underlying the stock assessment, 4) the statistical framework for fitting the model to the data and determining goodness of fit, and 5) the analytical treatment of model outputs in determining management reference points.

Major changes to the assessment should be postponed until the next full assessment. Minor alterations to the input data and the assessment can be considered as long as the update assessment clearly documents and justifies the need for such changes. A step-by-step transition (via sensitivity analysis) from the last full assessment to an update assessment under review should be provided. Minor alterations can be considered under only two circumstances: first, when the addition of new data reveals an unanticipated sensitivity of the model, and second, when there are clear and straightforward improvements in the input data and how it is processed and analyzed for use in the model. Examples of minor alterations include: 1) changes in how compositional data are pooled across sampling strata, 2) the weighting of the various data components (including the use of methods for tuning the variances of the data components), 3) changes in the time periods for the selectivity blocks, 4) correcting data entry errors, and 5) bug fixes in software programming. This list is not meant to be exhaustive, and other alterations can be considered if warranted. Ideally, improved data or methods used to process and analyze data would be reviewed by the SSC prior to being used in assessments.

The SSC reviews all update assessments; a STAR panel review is not needed since the assessment only updates the last full, STAR panel-reviewed assessment.

### 1.1.2.2 Rebuilding Analyses

Rebuilding analyses use the results of stock assessments and project stock rebuilding periods under alternative harvest control rules in a stochastic fashion. In other words, a rebuilding analysis involves projecting the status of the overfished resource into the future under a variety of alternative harvest strategies to determine the probability of recovery to  $B_{MSY}$  (or its proxy) within a pre-specified time-frame. Rebuilding analyses are used to develop new rebuilding plans or in consideration for modifying existing rebuilding plans; rebuilding plans dictate the target year to rebuild a stock, the harvest control rules for rebuilding the stock, and any other special management measures designed to foster rebuilding. Rebuilding analyses also are used to determine the OFLs and ACLs for overfished stocks.

The steps when conducting a rebuilding analysis are 1) estimation of  $B_0$  (and hence  $B_{MSY}$  or its proxy), 2) selection of a method to generate future recruitment, 3) specification of the mean generation time (defined as the predicted time it would take for a mature female in the population to replace herself), 4) calculation of the minimum and maximum times to recovery, and 5) identification and analysis of alternative harvest strategies and rebuilding times. Most rebuilding analyses are done using software developed by Dr. André Punt from the University of Washington (informally termed the Puntalyzer; available at <http://puntlab.washington.edu/software/>).

The Puntalyzer uses “Monte Carlo simulation” to derive a probability estimate for a given rebuilding strategy. This method projects population growth many times in separate simulations. It accounts for possible variability by randomly choosing the value of a key variable, generally the deviation in recruitment about the stock-recruitment relationship but some rebuilding also allowed for uncertainty in the estimated parameters of the stock assessment. Because of this variability in a key input value, each simulation will show a different pattern of population growth. As a result, a modeled population may reach the target biomass that defines a rebuilt stock ( $B_{MSY}$ ) in a different year in each of the simulations.

This technique is first used to calculate minimal time to rebuild a stock given its level of depletion and productivity from the time of implementing the first rebuilding plan ( $T_{MIN}$ ) in probabilistic terms, which is defined as the time needed to reach the target biomass in the absence of fishing with a 50 percent probability. In other words, in half the simulations the target biomass was reached in some year up to and including the computed  $T_{MIN}$ . Given  $T_{MIN}$ , the maximum legal time to rebuild ( $T_{MAX}$ ) is computed as 10 years or by adding the value of one mean generation time to  $T_{MIN}$ , if  $T_{MIN}$  is greater than or equal to 10 years. In cases where there is consideration for modifying an existing rebuilding plan, the shortest time to rebuild is calculated as the biological limit for the stock to rebuild in the absence of fishing beginning in the year the modified rebuilding plan is implemented; this limit is denoted, “ $T_{F=0}$ ”.

A target rebuilding year,  $T_{TARGET}$ , is set as a year at  $T_{MIN}$  (or  $T_{F=0}$ ) or greater, which does not exceed  $T_{MAX}$ , and which is as short as possible, taking into account the status and biology of the stock, the needs of fishing communities, and the interaction of the stock of fish within the marine ecosystem. Prior to Amendment 16-4, the Council set  $T_{TARGET}$  in part by considering the probability of rebuilding the stock by  $T_{MAX}$ . The Council may continue to review the probability of rebuilding the stock by  $T_{MAX}$  given differing harvest control rules, a reference parameter known as “ $P_{MAX}$ .” The Magnuson-Stevens Act, however, simply requires that rebuilding periods be as short as possible, taking into account:

- the status and biology of any overfished stocks of fish;
- the needs of fishing communities;
- recommendations by international organizations in which the United States participates; and
- the interaction of the overfished stock of fish within the marine ecosystem (§304(e)(4)(A)(i)).

It is important to recognize that some of the terms introduced and described above represent policy decisions at the national level and the Council **does not have a choice** in setting their values. The dates for  $T_{MIN}$  and  $T_{MAX}$  are determined based on guidelines established at the national level. Mean generation time is a biological characteristic that cannot be chosen by policymakers. Thus, the Council cannot choose these values and then use them as a basis for management. Defined in national guidelines,  $T_{MIN}$  is a consequence of the productivity of the fish stock and is calculated by fishery biologists based on information they estimate for a particular stock. Similarly,  $T_{MAX}$ , which is calculated from  $T_{MIN}$ , does not represent a Council choice.

Policy flexibility comes into play in determining  $T_{TARGET}$ , or the time by which the stock is projected to rebuild. As explained earlier, the time to rebuild must be as short as possible, taking into account the status and biology of the stock, the needs of fishing communities, and the interaction of the stock of fish within the marine ecosystem. When developing a management strategy the Council can choose a fishing mortality rate and corresponding annual level of fishing. However, when rebuilding overfished species, the choice of the harvest control rule is based on the value of  $T_{TARGET}$ , keeping in mind that these values cannot be chosen independently of one another. In other words, the Council may choose one value and derive the other from it, but they cannot choose these values independently of the other.

The current groundfish rebuilding plan parameters are depicted in Table 1-7.

**Table 1-4. Management quantities estimated from the most recent stock assessments informing management in 2015 and beyond.**

Stock	Year of Most Recent Assessment	Est. Depletion a/	Initial Spawning Biomass (B <sub>0</sub> )	Current Spawning Biomass a/	Current Total Biomass a/	Spawning Biomass at MSY	Harvest Rate at MSY	MSY	MSY Basis
Arrowtooth flounder	2007	0.79	80,313 mt	63,302 mt	85,175 mt	22,524 mt	0.162	5,833 mt	F <sub>30%</sub>
Aurora rockfish	2013	0.64	2,626 mt	1,673 mt	4,366 mt	1,213 mt	0.025	67 mt	F <sub>50%</sub>
Black rockfish (S of Cape Falcon)	2007	0.71	4,578.5 M larvae	3,227 M larvae	23,232 M larvae	1,831.4 M larvae	0.072	1,035.4 mt	F <sub>50%</sub>
Black rockfish (N of Cape Falcon)	2007	0.53	2,321 mt	1,239 mt	7,558 mt	928 mt	0.110	408 mt	F <sub>50%</sub>
Blackgill rockfish	2011	0.30	1,188 M larvae	359 M larvae	6,585 mt	543.0 M larvae	0.022	177 mt	F <sub>50%</sub>
Blue rockfish	2007	0.30	2,077 M larvae	622 M larvae	5,447 mt	831 mt	0.040	275 mt	F <sub>50%</sub>
Bocaccio	2013	0.31	8,117,510 M eggs	2,551,060 M eggs	19,077 mt	3,307,000 M eggs	0.067	1,341 mt	F <sub>50%</sub>
Brown rockfish	2013	0.42	1,794 mt	727 mt	1,454 mt	718 mt	0.102	149 mt	B <sub>40%</sub>
Cabezon (CA)	2009	0.48	1,298 mt	627 mt	1,342 mt	515 mt	0.118	149 mt	F <sub>45%</sub>
Cabezon (OR)	2009	0.52	409 mt	214 mt	455 mt	157 mt	0.120	49 mt	F <sub>45%</sub>
California scorpionfish	2005	0.80	1,024 mt	816 mt	1,866 mt	259 mt	0.161	127 mt	est.
Canary rockfish	2011	0.23	27,846 mt	6,458 mt	16,124 mt	9,545 mt	0.033	799 mt	F <sub>50%</sub>
Chilipepper rockfish	2007	0.71	33,390 mt	23,827 mt	32,401 mt	15,482 mt	0.088	2,099 mt	F <sub>50%</sub>
China rockfish (N of 40°10' N. lat.)	2013	0.37	243 mt	84 mt	168 mt	97 mt	0.045	9 mt	B <sub>40%</sub>
China rockfish (S of 40°10' N. lat.)	2013	0.66	405 mt	264 mt	527 mt	162 mt	0.100	32 mt	B <sub>40%</sub>
Copper rockfish (N of 34°27' N lat.)	2013	0.48	1,704 mt	795 mt	1,590 mt	681 mt	0.083	114 mt	B <sub>40%</sub>
Copper rockfish (S of 34°27' N lat.)	2013	0.76	942 mt	699 mt	1,397 mt	377 mt	0.109	84 mt	B <sub>40%</sub>
Cowcod	2013	0.34	1,549 mt	524 mt	1,049 mt	620 mt	0.050	62 mt	B <sub>40%</sub>
Darkblotched rockfish	2013	0.36	3,358 M eggs	1,214 M eggs	16,613 mt	1,343 M eggs	0.040	675 mt	B <sub>40%</sub>
Dover sole	2011	0.84	469,866 mt	393,507 mt	684,685 mt	119,033 mt	0.128	34,743 mt	F <sub>30%</sub>

Stock	Year of Most Recent Assessment	Est. Depletion a/	Initial Spawning Biomass (B <sub>0</sub> )	Current Spawning Biomass a/	Current Total Biomass a/	Spawning Biomass at MSY	Harvest Rate at MSY	MSY	MSY Basis
English sole	2013	0.88	29,238 mt	25,719 mt	46,968 mt	7,833 mt	0.404	3,875 mt	F <sub>30%</sub>
Gopher rockfish	2005	0.97	1,995 mt	1,931 mt	2,440 mt	798 mt	0.103	101 mt	F <sub>50%</sub>
Greenspotted rockfish	2011	0.35	1,357.8 B eggs	449.9 B eggs	3,110 mt	621 B eggs	.034 N; .024 S	95.6 mt	F <sub>50%</sub>
Greenstriped rockfish	2009	0.81	7,090 M eggs	5,736 M eggs	29,391 mt	3,101 M eggs	0.044	738 mt	F <sub>50%</sub>
Kelp greenling (OR)	2005	0.49	b/	b/	b/	b/	b/	b/	b/
Lingcod (WA & OR)	2009	0.62	33,075 mt	20,484 mt	32,222 mt	13,671 mt	0.082	1,710 mt	F <sub>45%</sub>
Lingcod (CA)	2009	0.74	25,311 mt	18,656 mt	31,266 mt	10,462 mt	0.084	1,492 mt	F <sub>45%</sub>
Longnose skate	2007	0.66	7,034 mt	4,634 mt	71,971 mt	844 mt	0.043	787 mt	F <sub>45%</sub>
Longspine thornyhead	2013	0.75	39,134 mt	29,436 mt	68,131 mt	15,654 mt	0.060	2,487 mt	F <sub>50%</sub>
Pacific ocean perch	2011	0.19	6,556 B eggs	1,079.4 B eggs	106,847 mt	1,311.2 B eggs	0.032	863 mt	F <sub>50%</sub>
Pacific sanddabs	2013	0.96	b/	b/	b/	b/	b/	b/	b/
Pacific whiting	2013	0.72	2.081 M mt	1.504 M mt	NA	.556 M mt	0.184	.243 M mt	F <sub>40%</sub>
Petrale sole	2013	0.22	32,426 mt	7,233 mt	15,015 mt	8,107 mt	0.170	2,750 mt	B <sub>25%</sub>
Rex sole	2013	0.80	3,808 mt	2,966 mt	18,497 mt	1,026 mt	0.464	1,646 mt	F <sub>30%</sub>
Rougeye/blackspotted rockfish	2013	0.47	5,394 mt	2,552 mt	8,176 mt	2,491 mt	0.027	194 mt	F <sub>50%</sub>
Sablefish	2011	0.33	182,136 mt	60,957 mt	205,662 mt	61,926 mt	0.050	10,021 mt	F <sub>45%</sub>
Sharpchin rockfish	2013	0.68	7,887 mt	4,947 mt	12,767 mt	3,482 mt	0.050	270 mt	F <sub>50%</sub>
Shortbelly rockfish	2007	0.73	c/	35,000 mt	NA	NA	NA	NA	NA
Shortspine thornyhead	2013	0.74	189,765 mt	140,753 mt	244,400 mt	75,906 mt	0.015	2,034 mt	F <sub>50%</sub>
Spiny dogfish	2011	0.63	70,724 K fish	44,660 K fish	215,988 K fish	28,290 K fish	0.006	831 mt	B <sub>40%</sub>

<b>Stock</b>	<b>Year of Most Recent Assessment</b>	<b>Est. Depletion a/</b>	<b>Initial Spawning Biomass (B<sub>0</sub>)</b>	<b>Current Spawning Biomass a/</b>	<b>Current Total Biomass a/</b>	<b>Spawning Biomass at MSY</b>	<b>Harvest Rate at MSY</b>	<b>MSY</b>	<b>MSY Basis</b>
Splitnose rockfish	2009	0.66	12,853 M eggs	8,426 M eggs	74,772 mt	5,006 M eggs	0.033	1,244 mt	F <sub>50%</sub>
Starry flounder	2005	0.50	7,158 mt	3,566 mt	7,638 mt	1,830 mt	0.229	1,848 mt	F <sub>30%</sub>
Stripetail rockfish	2013	>0.775	b/	b/	b/	b/	b/	b/	b/
Widow rockfish	2011	0.51	71,126 mt	36,342 mt	68,238 mt	32,315 mt	0.067	4,758 mt	F <sub>50%</sub>
Yelloweye rockfish	2011	0.21	1,028 M eggs	219 M eggs	2,188 mt	411 M eggs	0.022	58 mt	F <sub>50%</sub>
Yellowtail rockfish (N of 40°10' N. lat.)	2013	0.67	82,974 mt	50,043 mt	143,384 mt	37,036 mt	0.074	4,806 mt	F <sub>50%</sub>

a/ Estimates pertain to the most recent assessment year.

b/ The assessment results were only used for informing status since the scale of the population could not be adequately determined.

c/ A dynamic B<sub>0</sub> was modeled with an initial biomass estimate of 187,000 mt in 1950 and a mean unfished biomass of 48,000 mt.

**Table 1-5. Parameters estimated and/or assumed in base models in the most recent west coast groundfish stock assessments, excluding those done using XDB-SRA.**

Stock	ln(R0)	Steepness (h)		Sigma-r	von-Bertalanffy Growth Coefficient (K)		Natural Mortality (M)		
		value	est.?		females	males	females	males	est.?
Arrowtooth flounder	10.26	0.90	N	0.8	0.17	0.39	0.166	0.274	N
Aurora rockfish	6.64	0.78	N	0.5	0.09	0.09	0.035	0.037	a/
Black rockfish (S of Cape Falcon)	8.97	0.60	N	0.5	0.17	0.26	0.160 < 10 yrs 0.240 > 15 yrs	0.160	N
Black rockfish (N of Cape Falcon)	8.04	0.60	N	0.35	0.164	0.194	0.200	0.160	Y
Blackgill rockfish (S of 40°10' N lat.)	7.73	0.65	N	0.5	0.028	0.047	0.063	0.065	N
Blue rockfish (CA N of Pt. Con.)	8.08	0.58	N	0.5	0.147	0.295	0.100	0.120	N
Bocaccio	8.55	0.61	Y	1.0	0.22	0.27	0.150	0.150	N
Cabazon (CA N of Pt. Con.)	6.78	0.70	N	0.5	0.149	0.269	0.250	0.300	N
Cabazon (CA S of Pt. Con.)	5.33	0.70	N	0.7	0.130	0.230	0.250	0.300	N
Cabazon (OR)	5.27	0.70	N	0.5	0.190	0.178	0.250	0.300	N
California scorpionfish	7.63	0.70	N	1.0	0.13	0.12	0.250	0.250	N
Canary rockfish	8.12	0.51	N	0.5	0.125	0.162	0.060 < 6 yrs 0.092 ≥ 6 yrs	0.060	Y
Chilipepper rockfish	19.45	0.57	N	1.0	0.2 - 0.32 b/	0.2 - 0.32 b/	0.160	0.200	N
Darkblotched rockfish	7.84	0.78	N	0.75	0.2	0.26	0.050	0.067	a/
Dover sole	12.85	0.80	N	0.35	0.150	0.171	0.117	0.142	Y
English sole	11.45 c/	0.87 c/	Y	d/	0.36	0.48	0.24 c/	0.27 c/	Y
Gopher rockfish	7.92	0.65	N	0.5	0.186	0.186	0.200	0.200	N
Greenspotted rockfish (CA N of Pt. Con.)	6.15	0.76	N	0.7	0.057	0.057	0.065	0.065	N
Greenspotted rockfish (CA S of Pt. Con.)	6.65	0.76	N	0.7	0.042	0.042	0.065	0.065	N
Greenstriped rockfish	9.62	0.69	N	0.84	0.11	0.15	0.080	0.080	N
Kelp greenling (OR)	7.02	0.70	N	1.0	0.3	0.4	0.260	0.260	N
Lingcod (WA & OR)	8.06	0.80	N	0.5	0.13	0.22	0.180	0.320	N
Lingcod (CA)	8.17	0.80	N	0.5	0.11	0.23	0.180	0.320	N
Longnose skate	9.65	0.40	N	d/	0.064	0.064	0.200	0.200	N
Longspine thornyhead	11.82	0.60	N	0.6	0.109	0.109	0.111	0.111	N



Stock	ln(R0)	Steepness (h)		Sigma-r	von-Bertalanffy Growth Coefficient (K)		Natural Mortality (M)		
		value	est.?		females	males	females	males	est.?
Pacific ocean perch	9.14	0.40	N	0.7	0.159	0.195	0.050	0.051	N
Pacific whiting	21.71	0.82	Y	1.4	e/	e/	0.224	0.224	Y
Petrale sole	9.72	0.86	Y	0.4	0.13	0.21	0.150	0.169	Y
Rex sole	9.51 c/	0.89 c/	Y	d/	0.39	0.39	0.23 c/	0.12 c/	Y
Rougeye/blackspotted rockfish	6.19	0.78	N	0.4	0.081	0.081	0.042	0.042	Y
Sablefish	10.01	0.60	N	0.6	0.335	0.419	0.080	0.065	Y
Sharpchin rockfish	8.23 c/	0.77 c/	Y	d/	0.17	0.20	0.07 c/	0.07 c/	Y
Shortbelly rockfish	12.64	0.65	N	1.0	0.198	0.200	0.260	0.260	N
Shortspine thornyhead	10.32	0.60	N	0.5	0.018	0.018	0.051	0.051	N
Spiny dogfish	10.07	0.28	f/	d/	0.026	0.052	0.064	0.064	N
Splitnose rockfish	9.54	0.58	N	1.0	0.156	0.165	0.048	0.048	N
Starry flounder (OR & WA)	7.96	0.80	N	1.0	0.251	0.426	0.510	0.760	N
Starry flounder (CA)	7.23	0.80	N	1.0	0.251	0.426	0.510	0.760	N
Widow rockfish	10.06	0.76	N	0.65	0.209	0.233	0.120	0.129	Y
Yelloweye rockfish	5.43	0.44	Y	d/	0.047	0.047	0.046	0.045	Y
Yellowtail rockfish (N of 40°10' N lat.)	9.84 c/	0.78 c/	Y	d/	0.170	0.190	0.10 c/	0.11 c/	Y

a/ Female M was fixed and male M was estimated as an offset to female M.

b/ The base case model allowed growth for each sex to differ between blocks of time, based on freely estimating the K parameter.

c/ This value is the median of the posterior distribution of estimates for this parameter. These estimates were not summarized in tabular form in the 2013 data-moderate assessments' document (Cope, *et al.* 2014) but were provided by the Stock Assessment Team after the assessment was published.

d/ Recruitment variability (sigma-r) not estimated.

e/ The 2013 Pacific whiting assessment uses weight-at-age, thus there is no estimate of growth. Weight-at-age varies between years; therefore, growth is time-varying.

f/ Steepness was a derived quantity from the 2011 assessment, not an estimated parameter from an alternative stock-recruitment relationship modeled in the assessment.

**Table 1-6. Population parameters estimated and/or derived in base models in 2013 west coast groundfish stock assessments using XDB-SRA.**

Stock	$B_0$	Estimated Parameters				$B_{MSY}$	$F_{MSY}$
		M	$F_{MSY}/M$	$B_{MSY}/B_0$	$\Delta_{2000}$		
Brown rockfish	3588	0.133	0.971	0.399	0.698	1,383.4	0.130
China rockfish (N of 40°10' N lat.)	486	0.058	0.916	0.381	0.461	178.7	0.054
China rockfish (S of 40°10' N lat.)	811	0.070	1.436	0.446	0.539	361.0	0.104
Copper rockfish (N of Pt. Con.)	3407	0.089	1.090	0.402	0.717	1,334.4	0.099
Copper rockfish (S of Pt. Con.)	942	0.097	1.165	0.458	0.501	860.1	0.115
Cowcod	3099	0.054	1.051	0.422	0.801	1,239.5	0.050

### 1.1.3 Overfished Groundfish Stocks

There are six overfished west coast rockfish stocks (i.e., bocaccio south of 40°10' N lat., canary rockfish, cowcod south of 40°10' N lat., darkblotched rockfish, Pacific ocean perch, and yelloweye rockfish) and one overfished flatfish stock (i.e., petrale sole) at the start of 2013. All seven of these stocks are rebuilding and three (i.e., bocaccio south of 40°10' N lat., darkblotched rockfish, and petrale sole) are predicted to be successfully rebuilt by the start of 2015. Descriptions of these overfished groundfish stocks follows.

Stock rebuilding parameters estimated from the most recent rebuilding analyses and current rebuilding parameters specified at the start of 2015 are provided in Table 1-7.

**Table 1-7. Rebuilding parameters estimated in the most recent rebuilding analyses and specified in rebuilding plans for overfished groundfish stocks at the start of the 2015-2016 management cycle.**

Stock	T <sub>MIN</sub>	T <sub>F=0</sub>	T <sub>MAX</sub>	T <sub>TARGET</sub>	Harvest Control Rule Specification
Bocaccio	2018	2018	2031	2022	SPR 77.7%
Canary	2027	2028	2050	2030	SPR 88.7%
Cowcod	2019	2019	2057	2020	SPR 82.7% (E = 0.007)
Darkblotched	2012	2016	2037	2025	SPR 64.9%
POP	2040	2043	2071	2051	SPR 86.4%
Petrale sole	2014	2014	2021	2016	25-5 Rule
Yelloweye	2044	2047	2083	2074	SPR 76%

#### 1.1.3.1 Bocaccio

##### Distribution and Life History

Bocaccio (*Sebastes paucispinis*) is a rockfish species that ranges from Stepovak Bay on the Alaskan Peninsula (as well as Kodiak Island, Alaska) to Punta Blanca, Baja California, Mexico (Hart 1988; Miller and Lea 1972). Love, et al. (2002) and Thomas and MacCall (2001) describe bocaccio distribution and life history. Bocaccio are historically most abundant in waters off central and southern California. The southern bocaccio stock is most prevalent in the 54-82 fm depth zone (Casillas, *et al.* 1998).

Bocaccio are found in a wide variety of habitats, often on or near bottom features, but sometimes over muddy bottoms. They are found both nearshore and offshore (Sakuma and Ralston 1995). Larvae and small juveniles are pelagic (Garrison and Miller 1982) and are commonly found in the upper 100 m of the water column, often far from shore (MBC 1987). Large juveniles and adults are semi-demersal and are most often found in shallow coastal waters over rocky bottoms associated with algae (Sakuma and Ralston 1995). Adults are commonly found in eelgrass beds, or congregated around floating kelp beds love (Love, *et al.* 1990; Sakuma and Ralston 1995). Young and adult bocaccio also occur around artificial structures, such as piers and oil platforms (MBC 1987). Although juveniles and adults are usually found around vertical relief, adult aggregations also occur over firm sand-mud bottoms (MBC 1987). Bocaccio move into shallow waters during their first year of life hart (Hart 1988), then move into deeper water with increased size and age (Garrison and Miller 1982).

Bocaccio are ovoviviparous (live young are produced from eggs that hatch within the female's body) (Garrison and Miller 1982; Hart 1988). Love et al. (1990) reported the spawning season to last nearly an

entire year (>10 months). Parturition occurs during January to April off Washington, November to March off Northern and Central California, and October to March off Southern California (MBC 1987). Fecundity ranges from 20,000 to 2,300,000 eggs. In California, two or more broods may be born per year (Love, *et al.* 1990). The spawning season is not well known in northern waters. Males mature at three to seven years, with about half maturing in four to five years. Females mature at three to eight years, with about half maturing in four to six years (MBC 1987).

Maximum age of bocaccio was radiometrically determined to be at least 40 years, and perhaps more than 50 years. Bocaccio are difficult to age, and stock assessments used length measurement data and growth curves to estimate the age composition of the stock (Ralston and Ianelli 1998). Although recent assessments have described the true natural mortality rate as a key unknown for estimating stock status, recent assessments have used a value of 0.15 (which is associated with an 86 percent adult annual survival rate in the absence of fishing mortality).

Larval bocaccio eat diatoms, dinoflagellates, tintinnids, and cladocerans (Sumida and Moser 1984). Copepods and euphausiids of all life stages (adults, nauplii and egg masses) are common prey for juveniles (Sumida and Moser 1984). Both Phillips (1964) and Love *et al.* (2002) described bocaccio rockfish as almost exclusively piscivorous, and include other rockfish, Pacific whiting, sablefish, anchovy, mesopelagic fishes and squid as the key prey for large juvenile and adult bocaccio. Bocaccio are eaten by sharks, salmon, other rockfishes, lingcod, albacore, sea lions, porpoises, and whales (MBC 1987). Adult bocaccio are often caught with chilipepper rockfish and have been observed schooling with speckled, vermilion, widow, and yellowtail rockfish (Love, *et al.* 2002). As pelagic juveniles, they may compete with chilipepper, widow, yellowtail, shortbelly, and other pelagic juvenile rockfishes for both food and habitat (Reilly, *et al.* 1992).

### **Stock Status and Management History**

Bocaccio are managed as two separate west coast populations. The southern stock exists south of Cape Mendocino and the northern stock north of Cape Mendocino (the northern stock density is limited south of 48° N lat. with increasing abundance off Cape Flattery, Washington and points north). It is unclear whether this stock separation implies stock structure. The distribution of the two populations and evidence of lack of genetic intermixing suggests stock structure, although MacCall (2002) reported some evidence for limited genetic mixing of the two populations. Nonetheless, assessment scientists and managers have treated the two populations as independent stocks north and south of Cape Mendocino.

Bocaccio have long been an important component of California rockfish fisheries. Catches increased to high levels in the 1970s and early 1980s as relatively strong year-classes recruited to the stock. The Council began to recommend increasingly restrictive regulations after an assessment of the southern stock in 1990 (Bence and Hightower 1990) indicated that fishing rates were too high. The southern stock suffered poor recruitment during the warm water conditions that prevailed off Southern California beginning in the late 1980s. The 1996 assessment (Ralston, *et al.* 1996) indicated the stock was in severe decline. NMFS formally declared the stock overfished in March 1999 after the groundfish FMP was amended to incorporate the tenets of the Sustainable Fisheries Act. MacCall *et al.* (1999) confirmed the overfished status of bocaccio and estimated spawning output of the southern stock to be 2.1 percent of its unfished biomass.

In the 2002 assessment (MacCall 2002) relative abundance increased slightly from the previous assessment (4.8 percent of unfished biomass), potential productivity (as evidenced from the steepness of the spawner/recruit relationship, which reflects the level of compensatory production at low stock sizes) appeared lower than previously thought, making for a more pessimistic outlook. Furthermore, the 2002 assessment revealed that although the 1999 year class was the strongest in several years, it was weak relative to the range of possibilities considered in the 1999 assessment. The 2002 rebuilding analysis (MacCall and

He 2002) predicted the stock would not rebuild within maximum time legally possible ( $T_{MAX}$ ) even with no fishing-related mortality. Total mortality in 2003 fisheries was restricted to less than 20 mt as a means of conserving the stock while minimizing adverse socioeconomic impacts to communities.

The 2003 bocaccio assessment (MacCall 2003b) estimated a higher stock biomass (7.4% depletion) relative to the 2002 assessment. The instantaneous rate of natural mortality was changed from 0.2 to 0.15. Additional CalCOFI data indicated an increasing abundance trend due to recruitment of the 1999 year class. This was corroborated by a dramatic increase in recreational CPUE, which was at a record high level in central California north of Pt. Conception. The 2003 rebuilding analysis suggested the stock could rebuild to  $B_{MSY}$  within 25 years while sustaining an OY of approximately 300 mt in 2004 (MacCall 2003a).

The 2003 assessment was updated in 2005 and 2007 (MacCall 2006b; MacCall 2008b) using the original 2003 base model (i.e., STATc) in SS1. These assessments were used to establish annual specifications and management measures consistent with a strategy of a higher OY than the impacts anticipated under the suite of management measures adopted. This strategy was designed to buffer the effects of a large recruitment event like that observed for the 1999 year class. Such effects include disruption to fisheries as experienced in previous years when fisheries closed early to avoid young bocaccio. This buffer strategy, which addressed the large, episodic recruitment pattern inherent in the stock's dynamics, became a tenet of the bocaccio rebuilding plan.

A bocaccio rebuilding plan was adopted by the Council in 2004 under Amendment 16-3 (PFMC 2004). The rebuilding plan established a target rebuilding year of 2023 and a harvest control rule of  $F = 0.0498$ . (It was later clarified in the 2005 rebuilding analysis (MacCall 2006a) that the target rebuilding year had been incorrectly stated in the rebuilding plan to be 2023 since the 2003 rebuilding analysis indicated that a 50 percent probability rebuilding would require 23 years, and that this assumed a beginning date of 2004 (the first simulated year). Therefore, the Council amended the rebuilding plan's target year to 2026.

A new rebuilding analysis was conducted in 2007 (MacCall 2008a) based on the results of the 2007 stock assessment (MacCall 2008b). The 2007 bocaccio rebuilding analysis showed a similar rebuilding trajectory to that adopted in Amendment 16-4 and the rebuilding plan was maintained for the 2009-2010 management cycle.

A new bocaccio assessment (Field, *et al.* 2009) and rebuilding analysis (Field and He 2009) were prepared in 2009. Field *et al.* (2009) extended the assessment north of Cape Mendocino to Cape Blanco, Oregon; the U.S. west coast stock north of this point has not been assessed. Indications of strong 2009 and 2010 year classes were projected to result in increased abundance. Depletion in 2011 was estimated at 26 percent (18.7 -33.1 percent), with the stock projected to be rebuilt by 2019. Based on these analyses, the Council changed the target year for rebuilding bocaccio from 2026 to 2022; the amended rebuilding plan was implemented in 2011.

A bocaccio stock assessment update (Field 2011b) and rebuilding analysis (Field 2011a) were prepared in 2011. The 2011 bocaccio assessment was originally scheduled to be an update of the 2009 full assessment; however, the STAT some limited changes in the 2009 model structure since a strict update estimated that the 2010 year class was extraordinarily and unrealistically strong, based on length frequency data collected in the 2010 NMFS trawl survey. The modified update was ultimately reviewed, endorsed by the SSC, and adopted for use in management decision-making. The 2011 bocaccio rebuilding analysis indicated rebuilding progress was well ahead of schedule with a predicted median year to rebuild of 2021 or one year earlier than the target rebuilding year (Field 2011a). The Council elected to maintain the revised rebuilding plan implemented in 2011.

An update of the 2011 bocaccio assessment model was prepared in 2013, which confirmed the 2009 and 2010 year classes were indeed strong (Field 2013). The assessment estimated a depletion of 31.4 percent at the start of 2013 (Figure 1-4) and predicted the stock would rebuild by 2015. The SSC recommended maintaining the current rebuilding plan for the 2015-2016 management cycle and a full assessment be done in 2015 to confirm this prediction. The SSC further recommended against preparing a rebuilding analysis in 2013. A full assessment of bocaccio is planned for 2015.




Figure 1-4. Relative depletion of bocaccio south of 40°10' N lat. from 1980 to 2013 based on the 2013 stock assessment update.

### Stock Productivity Relative to Rebuilding Success

The 2013 bocaccio assessment produced a (very slightly) more optimistic estimation of steepness (from 0.595 to 0.614) relative to the 2011 model (the 2009 model had a point estimate of 0.573). Despite these modest changes, the overall trajectory of spawning output, relative spawning output, total biomass and recruitment are barely distinguishable as changed from the 2011 model, with the most important change being the relative strength of the 2010 year class. The strength of the 2010 year class is estimated with less uncertainty in the 2013 assessment.

Recruitment for bocaccio is highly variable, with a small number of year classes tending to dominate the catch in any given fishery or region. Adult abundance is highly variable even in the absence of fishing (MacCall 2002). Recruitment appears to have been at very low levels throughout most of the 1990s, but the 1999 year class was the highest since 1988, and led to a substantive increase in abundance during the early 2000s. Several year classes of moderate strength (2003, 2005) occurred in the mid-2000s, and two

recent very strong year classes (2009 and 2010) are now estimated to be comparable to (2009) and roughly double (2010) the size of the 1999 year class (Figure 1-5). These year classes were strongly evident in recreational length frequency data, in the NWFSC hook and line survey data (and length comps), and in the power plant impingement dataset modeled by Field (2013), as well as in an index (not included in the 2013 assessment update) of recreational CPUE. These strong year classes are already estimated to have resulted in an increase in abundance and spawning output, and should propel the stock spawning output to target levels by approximately 2015 as the 2010 year class continues to grow and mature. Preliminary estimates from the juvenile rockfish survey also indicate very strong abundance of young-of-the-year rockfish of many species (including bocaccio) in 2013, suggesting anecdotally that 2013 will also be a strong recruitment year for bocaccio, as well as for other species. However, these data are not incorporated in the 2013 update, which only includes data through 2012. Although poorly understood, the stock assessment suggests that recovery may be taking place more rapidly in the south, and recovery in the central/northern California region may be dependent on an influx of fish from the southern area.




Figure 1-5. Estimated bocaccio recruitments, 1980-2012 (from Field 2013).

### Fishing Mortality

The presence of a banner 2010 year class in the bocaccio stock is not entirely unexpected. Bocaccio stock production is characterized by high episodic recruitment and relatively rapid juvenile growth rates Field, et al. 2009. Juvenile bocaccio also recruit to shallow waters and are consequently caught in nearshore recreational fisheries as evidenced by dramatic spikes in both catch rates and the percentage of the total southern California rockfish catch that is bocaccio following strong recruitment events. Unlike most rockfish species where recruitment to fisheries usually takes several years due to low growth rates, juvenile bocaccio can recruit to nearshore fisheries in California within a year or two of parturition. Recruitment of the strong 1999 year class complicated management of California fisheries in 2001-2003, as this unpredictable event could not be reacted to in time given the lag in reconciling recreational catch estimates. Most species' rebuilding analyses are able to project recruitment into affected fisheries in time to decide and implement responsive management measures that will not compromise rebuilding plans. However, the fast growth and unpredictable recruitment of bocaccio poses the unique problem of having to react to a large recruitment event in real time. This experience has led the Council to a strategy of adopting higher bocaccio OYs/ACLs and more conservative management measures that are predicted to result in catches much lower than these harvest limits. The rebuilding strategy has been formalized by deciding OYs/ACLs

to determine rebuilding objectives and more stringent HGs for California. The buffer between the ACL (formerly the OY) and the HG accommodates the management uncertainty of an unforeseen recruitment event disrupting fisheries. Unlike an ACL, fisheries do not need to close upon attainment of an HG. The difference between the projected catch and the HG or ACL provides managers time to react to a strong recruitment to minimize mortality on bocaccio while minimizing disruptions to ongoing fisheries. This strategy has worked well to enhance bocaccio rebuilding while minimizing harm to California fishing communities.

Catch monitoring uncertainty is relatively high given the fact that a substantial amount of the total fishing mortality of bocaccio now occurs in the California recreational fishery, the sector with the largest bocaccio take in recent years. Recent recreational catch is estimated using the new California Recreational Fishing Survey (CRFS) program, which has been in existence since 2004. Prior to 2004, all recreational catch was estimated using the Marine Recreational Fisheries Statistical Survey (MRFSS) program, a survey methodology designed to understand long-term national trends in marine recreational catch and participation. The higher uncertainty in monitoring California recreational catches also translates into higher uncertainty in projecting recreational total mortalities. The fact that a substantial portion of the current take of bocaccio is in the California recreational fishery is another consideration for a relatively larger buffer between the predicted mortalities of bocaccio and the preferred ACLs.

### **Rebuilding Duration and Probabilities**

There is a high probability of successful bocaccio rebuilding by 2015 given the strength of recent year classes currently recruiting into the spawning population. The SSC has recommended a full assessment in 2015 to confirm this prediction.

#### **1.1.3.2 Canary Rockfish**

### **Distribution and Life History**

Canary rockfish (*Sebastes pinniger*) are distributed in the northeastern Pacific Ocean from the western Gulf of Alaska to northern Baja California; however, the species is most abundant from British Columbia to central California (Hart 1988; Love, *et al.* 2002; Miller and Lea 1972). Adults are primarily found along the continental shelf shallower than 300 m, although they are occasionally observed in deeper waters. Juvenile canary rockfish are found in shallow and intertidal areas (Love, *et al.* 2002).

Canary rockfish spawn in the winter, producing pelagic larvae and juveniles that remain in the upper water column for 3-4 months (Love, *et al.* 2002). These juveniles settle in shallow water around nearshore rocky reefs, where they may congregate for up to three years (Boehlert 1980; Sampson 1996) before moving into deeper water. The mean size of individuals captured in the trawl survey shows a characteristic ontogenetic shift to deeper water with increasing body size. The degree to which this ontogenetic shift may be accompanied by a component of latitudinal dispersal from shallow rocky reefs is unknown. Canary rockfish are a medium to large-bodied rockfish; achieving a maximum size of around 70 cm. Female canary rockfish reach slightly larger sizes than males.

Adult canary rockfish primarily inhabit areas in and around rocky habitat. They form very dense schools, leading to an extremely patchy population distribution that is reflected in both fishery and survey encounter rates.

Canary rockfish are relatively long-lived, with a maximum observed age of 95 years, however only males are commonly observed above the age of 50, while females tend to be rare above age 30. The degree to which this pattern reflects behavioral differences translating to reduced availability to fishery and survey



fishing gear, or an increase in relative mortality for older females has been the focus of much discussion and remains unclear. A similar pattern has been observed for yellowtail rockfish (*Sebastes flavidus*), a closely related, but more pelagic species with a similar distribution (Wallace and Lai 2006).

Canary rockfish off the west coast exhibit a protracted spawning period from September through March, probably peaking in December and January off Washington and Oregon (Hart 1988; Johnson, *et al.* 1982). Female canary rockfish reach sexual maturity at roughly eight years of age. Like many members of *Sebastes*, canary rockfish are ovoviviparous, whereby eggs are internally fertilized within females, and hatched eggs are released as live young (Bond 1979; Golden and Demory 1984; Kendall and Lenarz 1986). Canary rockfish are a relatively fecund species, with egg production being correlated with size (e.g., a 49-cm female can produce roughly 0.8 million eggs, and a female that has realized maximum length (approximately 60 cm) produces approximately 1.5 million eggs (Gunderson 1971).

Very little is known about the early life history strategies of canary rockfish. The limited research that has been conducted indicates that larvae are strictly pelagic (near the ocean surface) for a short period of time and begin to migrate to demersal waters during the summer of their first year of life. Larvae develop into juveniles around nearshore rocky reefs, where they may congregate for up to three years (Boehlert 1980; Sampson 1996). Evaluations of length distributions by depth demonstrate an increasing trend in mean size of fish with depth (Methot and Stewart 2006). Since 1990, stock assessments have assumed a base natural mortality rate of 0.06 (94 percent adult annual survival when there is no fishing mortality). Due to the rarity of old females in both survey and catch data, female canary rockfish have long been assumed to have increasing natural mortality rates with age (Golden and Wood 1990).

Little is known about ecological relationships between canary rockfish and other organisms. Adult canary rockfish are often caught with bocaccio, sharpchin, yelloweye, and yellowtail rockfishes, and lingcod. Researchers have also observed canary rockfish associated with silvergray and widow rockfish. Young-of-the-year feed on copepods, amphipods, and young stages of euphausiids. Adult canary rockfish feed primarily on euphausiids, as well as pelagic shrimp, cephalopods, mesopelagic fishes and other prey (Brodeur and Percy 1984; Lee 2002; Phillips 1964). Small canary rockfish are consumed by seabirds, Chinook salmon, lingcod, and marine mammals.

### **Stock Status and Management History**

Canary rockfish have long been an important component of rockfish fisheries. The Council began to recommend increasingly restrictive regulations after an assessment in 1994 (Sampson and Stewart 1994) indicated that fishing rates were too high. Prior to passage of the Sustainable Fisheries Act of 1996, there was no requirement for stock assessments to estimate biomass status; and until 1997 the Council's default target rate for fishing mortality corresponded to an SPR of 35%. Wallace and Cope (2011) estimated that the abundance of the canary rockfish stock dropped below  $B_{MSY}$  ( $B_{40\%}$ ) in 1983 and below the MSST in 1990, at which time the annual catch was more than double the current estimate of the MSY level. Harvest rates in excess of the current fishing mortality target for rockfish (SPR = 50%) is estimated to have begun in the late 1970s and persisted through 1999. Recent management actions appear to have curtailed the rate of removal such that overfishing has not occurred since 1999, and recent SPR values are in excess of 90 percent.

A 1999 stock assessment showed the stock had declined to 6.6 percent of unfished biomass in the northern area (Columbia and U.S. Vancouver management areas) (Crone, *et al.* 1999) and in the southern area (Conception, Monterey, and Eureka areas) (Williams, *et al.* 1999). The stock was declared overfished in January 2000. The first rebuilding analysis (Methot 2000) used results from the northern area assessment to project rates of potential stock recovery. The stock was found to have extremely low productivity, defined as production of recruits in excess of the level necessary to maintain the stock at its current, low

level. Rates of recovery were highly dependent upon the level of recent recruitment, which could not be estimated with high certainty. The initial rebuilding OY for 2001 and 2002 was set at 93 mt based upon a 50 percent probability of rebuilding by the year 2057, a medium level for these recent recruitments, and maintaining a constant annual catch of 93 mt through 2002.

A coastwide 2002 canary rockfish assessment estimated stock depletion to be 7.9 percent at the start of 2002 (Methot and Piner 2002b). A canary rockfish rebuilding plan was adopted in 2003 under Amendment 16-2 based on the results of the 2002 rebuilding analysis (Methot and Piner 2002a). The rebuilding plan established a target rebuilding year of 2074 and the harvest control rule of  $F = 0.022$  (with a  $P_{MAX}$  of 60 percent).

A full canary rockfish assessment was done in 2005 indicating a stock depletion of 9.0 percent at the start of 2005 (Methot and Stewart 2006). The assessment was based on two equally plausible models; one with differential male and female gear selectivities and one without gender-specific selectivities. A critical uncertainty in canary rockfish assessments was the lack of older, mature females in surveys and other assessment indices. There were two competing explanations for this observation. Older females could have a higher natural mortality rate, resulting in their disproportionate disappearance from the population. Alternatively, survey and fishing gears may be less effective at catching them, perhaps because older females are associated with habitat inaccessible to most trawl gear. If this is the case, then these fish (which, because of their higher spawning output, may make an important contribution to future recruitment) are part of the population, but remain poorly sampled. Methot and Stewart (2006) assumed a linear increase in female natural mortality from 0.06 at age 6 to approximately 0.09 at age 14. In the base model (differential male-female selectivity)  $B_0$  was estimated to be 34,798 mt, resulting in a depletion level of 5.7 percent. In the alternate model (no difference in selectivity)  $B_0$  was estimated to be 33,872 mt, with a depletion level of 11.3 percent. The steepness of the spawner-recruitment relationship, which largely determines the rate of increase in recruitment as the stock rebuilds, was estimated to be 0.33 in the base model, and 0.45 in the alternate model. The approved canary rockfish rebuilding analysis (Methot 2006) blended the two models by alternately re-sampling between the two input parameter sets.

The 2005 canary rebuilding analysis (Methot 2006) was used to inform the revised canary rebuilding plan adopted under Amendment 16-4, which specified a target rebuilding year of 2063 and a constant harvest strategy (SPR = 88.7%). Amendment 16-4 rebuilding plans were implemented in 2007.

The 2007 canary assessment estimated relative depletion level was 32.4 percent at the start of 2007 (Stewart 2008b). This was a significant departure from the previous assessment and largely driven by a higher assumed steepness ( $h = 0.51$ ) relative to past assessments. The 2007 assessment was unable to estimate steepness as had been done in the 2005 assessment, largely because the 2007 assessment treated the triennial bottom trawl survey as two separate indices due to changes between the 1992 and 1995 surveys in the seasonal timing. The 2007 canary rebuilding analysis (Stewart 2008a) predicted the SPR harvest rate in the rebuilding plan (88.7%) would rebuild 42 years earlier (2021) than the originally estimated rebuilding schedule (2063). A modification of the Amendment 16-4 canary rockfish rebuilding plan specifying a target rebuilding year of 2021 while maintaining the SPR harvest rate of 88.7% was implemented in 2009.

The 2009 canary assessment (Stewart 2009c), an update of the 2007 assessment, estimated stock depletion at 23.7% at the start of 2009. This change in stock status was due to a lower estimate of initial, unfished biomass ( $B_0$ ) largely attributable to the inclusion of revised historical California catches from a formal reconstruction of 1916-1980 California catch data (Ralston, *et al.* 2010). The 2009 canary rebuilding analysis (Stewart 2009a) predicted the stock would not rebuild to the target year of 2021 with at least a 50% probability even in the absence of fishing-related mortality starting in 2011 ( $T_{F=0}$ ). The rebuilding plan was revised by changing the target to rebuild the stock to 2027 while maintaining the 88.7% SPR harvest rate; the revised rebuilding plan was implemented in 2011.

Another update assessment was prepared in 2011 (Wallace and Cope 2011), which estimated stock depletion was 23.2 percent at the start of 2011 (Figure 1-6). This change in stock status was due to a lower estimate of initial, unfished biomass ( $B_0$ ) largely attributable to the inclusion of revised historical Oregon catches from a formal reconstruction of Oregon catch data. For the period 2000-2011, the spawning biomass was estimated to have increased from 11.2 percent to 23.2 percent of the unfished biomass level.

The 2011 canary rebuilding analysis (Wallace 2011) predicted the stock would not rebuild to the target year of 2027 with at least a 50% probability. The rebuilding plan was revised slightly by changing the target to rebuild the stock to 2030 while maintaining the 88.7% SPR harvest rate; the revised rebuilding plan was implemented in 2013.

A canary catch report was provided in 2013 ([Agenda Item F.5.a, Attachment 9, June 2013](#)), which indicated 2010-2012 total catches were below specified ACLs/OYs. A full assessment of canary rockfish is planned for 2015.




Figure 1-6. Relative depletion of canary rockfish from 1980 to 2011 based on the 2011 stock assessment update.

### Stock Productivity Relative to Rebuilding Success

Steepness is assumed to be 0.511 in the latest full assessment (Stewart 2008b) and the subsequent updates to that assessment (Stewart 2009c; Wallace and Cope 2011). This is a moderate to relatively low value of steepness for rockfish, as compared to the prior mean steepness (0.779) derived from meta-analysis of west coast rockfish stocks used in 2013 assessments (e.g., darkblotched rockfish, see section 1.1.3.4). Lower steepness implies a greater dependence on the size of the spawning population. The projected increase in

the canary rockfish biomass from the 2011 assessment is very sensitive to the value assumed for steepness and was projected to slow as below average recruitments begin to contribute to the spawning biomass.

Steepness is a difficult parameter to estimate and canary rockfish assessments are especially uninformative of steepness. The assumed canary steepness of 0.511 used in the last three assessments was based on the Dorn (2002a; Dorn 2002b) meta-analysis of west coast rockfish stocks. The value used in many 2013 assessments was based on an update of the Dorn (2002a) analysis (J. Thorson, pers. comm.).

As in previous assessments of canary rockfish, the recruitment deviations estimated by Wallace and Cope (2011) are primarily informed by the available age composition data. After a period of above average recruitments, recent year-class strengths (1997-2010) have generally been low, with only 2 of the 10 years (2001 and 2007) producing large estimated recruitments (Figure 1-7). The strength of the 2007 year class is subject to greater uncertainty than other strong recruitment events in the last 30 years because of the limited number of years in which it has been observed. As the larger recruitments from the late 1980s and early 1990s move through the population in future projections, the effects of recent poor recruitment may tend to slow the rate of recovery.





Figure 1-7. Estimated canary rockfish recruitments, 1980-2010 (from Wallace and Cope 2011).

### Fishing Mortality

Fishing mortality rates for canary rockfish in excess of the current proxy  $F_{MSY}$  harvest rates for rockfish (SPR = 50%) are estimated to have begun in the late 1970s and persisted through 1999. Figure 1-8 depicts estimated annual harvest rates relative to the overfishing limit ( $F_{MSY}$ ) and the current SPR harvest rate limit specified in the rebuilding plan. Recent management actions appear to have curtailed the rate of removal such that overfishing has not occurred since before 1999 and have maintained harvest rates below the current rebuilding SPR since 2005. Relative exploitation rates (catch/biomass of age-5 and older fish) are estimated to have been less than 1% since 2001.



**Figure 1-8. Estimated spawning potential ratio (SPR) of canary rockfish relative to the current  $F_{MSY}$  and rebuilding harvest rates, 1995-2010. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis.**

### Rebuilding Duration and Probabilities

Wallace 2013 estimated the canary rebuilding probability ( $P_{MAX}$ ) under the current SPR harvest rate of 88.7% is 75 percent. There is a 50 percent probability of recovering by the current target year of 2030.

#### 1.1.3.3 Cowcod

### Distribution and Life History

Cowcod (*Sebastes levis*) is a species of large rockfish with a distribution from Newport, Oregon, to central Baja California, Mexico (Love et al., 2002). They are most common from Cape Mendocino (California) to northern Baja California, in depths from 50-300 m. Hess et al. (2014) recently used genetic and otolith microchemistry tools to study cowcod population structure from California to Oregon. Specifically, they tested the hypothesis that a phylogeographic boundary exists at Point Conception. Their results supported a hypothesis of two primary lineages with a geographic boundary falling south of, rather than at Point Conception. Both lineages co-occur in the Southern California Bight (SCB), with no clear pattern of depth stratification or spatial structure within the Bight. Within lineages, there is evidence for considerable gene flow across the Point Conception boundary. Cowcod found north of Point Conception consist primarily of a single lineage, also found in northern areas of the SCB.

Cowcod are easily identified at all life stages, including larvae. Adults are piscivorous, with a diet consisting mainly of fishes, squids, and octopi. Cowcod are considered to be parademersal (transitional between a midwater pelagic and benthic species). Larvae develop into a pelagic juvenile stage, settling to benthic habitats after about 3 months. Juvenile cowcod were once thought to associate primarily with soft sediments, but Love and Yoklavich (2008), using visual surveys, found juveniles mainly associate with low-relief, hard substrate. Young-of-the-year were observed over a wide depth range (52-277 m), with juveniles slightly deeper, and adults mainly deeper than 150 m. Larger juveniles increasingly associate

with high-relief, complex rocky substrate, the primary habitat for adult cowcod. Adult cowcod are generally solitary, but occasionally aggregate (Love, *et al.* 1990). Although cowcod are generally not migratory, they may move, to some extent, to follow food (Love 1996).

Cowcod are a long-lived, slow-growing species that require a decade or more to reach sexual maturity. Fertilization is internal, with females giving birth to planktonic larvae mainly during winter months. Spawning peaks in January in the Southern California Bight (MacGregor 1986) and large females may produce up to three broods per season (Love, *et al.* 1990). Larvae emerge at about 5.0 mm (MacGregor 1986).

Cowcod are a highly fecund species, with large females producing 2 million eggs (fecundity is dependent on size and ranges from 181,000 to 1,925,000 eggs) (Love, *et al.* 2002). Dick *et al.* (2009) found no evidence of increasing weight-specific fecundity (i.e., spawning output is roughly proportional to spawning biomass).

Maximum observed age for cowcod is 55 years (Love, *et al.* 2002). Dick *et al.* (2007) estimated the natural mortality rate (M) using three methods, reporting a range of values from 0.027 to 0.064 based on Beverton's (1992) method, a range of total mortality (Z) estimates from 0.038 to 0.072 based on catch curve analysis, and Hoenig's geometric mean regression. Females reach 90 percent of their maximum expected size by 42 years.

Little is known about ecological relationships between cowcod and other organisms. Small cowcod feed on planktonic organisms such as copepods. Juveniles eat shrimp and crabs, and adults eat fish, octopus, and squid (Allen 1982). Adults consume a wide range of prey items, but are primarily piscivorous (Love, *et al.* 2002).

### **Stock Status and Management History**

While cowcod are not a major component of the groundfish fishery, they are highly desired by both recreational and commercial fishers because of their bright color and large size. The cowcod stock in the Conception area was first assessed in 1998 (Butler, *et al.* 1999b). Abundance indices decreased approximately tenfold between the 1960s and the 1990s, based on commercial passenger fishing vessel (CPFV) logs (Butler, *et al.* 1999b). Recreational and commercial catch also declined substantially from peaks in the 1970s and 1980s, respectively.

NMFS declared cowcod in the Conception and Monterey management areas overfished in January 2000, after Butler *et al.* (1999b) estimated the 1998 spawning biomass to be at 7 percent of  $B_0$ , well below the 25 percent overfishing threshold. Because cowcod is a fairly sedentary species, closed areas were established in 2002 to reduce cowcod mortality. Two Cowcod Conservation Areas (CCAs), in the Southern California Bight, were selected due to their high density of cowcod. The larger of the two areas (CCA West) is a 4200 square mile area west of Santa Catalina and San Clemente Islands. A smaller area (CCA East) is about 40 miles offshore of San Diego, and covers about 100 square miles. Bottom fishing is prohibited deeper than 20 fm within the CCAs.

A cowcod rebuilding analysis was completed in 2003 which validated the assumption that non-retention regulations and area closures had been effective in constraining cowcod fishing mortality (Butler, *et al.* 2003). These encouraging results were based on cowcod fishery-related landings in recreational and commercial fisheries, although the assessment included discard information only with respect to CPFV observations (which indicated negligible discards in that sector). This rebuilding review pointed out a common problem among the analyses of overfished species: reliance on landings (fishery-dependent) data for providing relative abundance values becomes increasingly difficult as the allowable catch is decreased

and fishery observer data remains low. Monitoring stock status and recovery thus becomes increasingly difficult in the absence of fishery-independent surveys.

As in the 1999 assessment, the 2005 cowcod assessment (Piner, *et al.* 2006) considered only the cowcod population in Southern California Bight (from the US-Mexico border north to Point Conception) population, as this is the area in which cowcod are most abundant, adult habitat is most common, and catches are highest. The 2005 assessment used only two data sources, the CPFV time series and the visual survey estimate data (Yoklavich, *et al.* 2007). The model was developed in Stock Synthesis 2, and although the base model estimated only three parameters (two of which were “nuisance parameters,” the other was equilibrium recruitment), the STAR Panel determined that this simplicity was appropriate given the paucity of data. The assessment provided a set of results corresponding to three different values for assumed steepness ( $h$ ), the key parameter in the stock-recruitment relationship ( $h=0.4, 0.5, \text{ and } 0.6$ ) and one the key uncertainties in the assessment. The assessment estimated that the 2005 spawning biomass was 18 percent of unfished levels and within a range of 14 to 21 percent depending on the value assumed for steepness, a considerably more optimistic result than the 1999 assessment. The corresponding 2005 cowcod rebuilding analysis (Piner 2006) was used to develop the cowcod rebuilding plan adopted in the groundfish FMP under Amendment 16-4. The rebuilding plan established a target rebuilding year of 2039 and an SPR of 90%.

A full cowcod assessment was conducted in 2007, which estimated spawning biomass to be 3.8 percent of its unfished level at the start of 2007 (Dick, *et al.* 2007). The 2007 cowcod assessment was an age-structured production model assuming a Beverton-Holt stock-recruitment function with deterministic recruitment, fit to the aggregated CPFV logbook index and the 2002 visual survey biomass estimate (Yoklavich, *et al.* 2007). Productivity parameters were fixed (steepness = 0.6, natural mortality = 0.055), leaving only virgin recruitment ( $R_0$ ) to be estimated. Spawning biomass in 2007 was estimated to be between 3.4 percent and 16.3 percent of the unfished level. The poor precision of this estimate was due to 1) a lack of data to inform estimates of stock productivity, and 2) conflicting information from fishery-dependent and fishery-independent data. However, even the most optimistic model, which assumed a high-productivity stock and ignored declines in CPFV catch rates, suggested that spawning biomass was below 25 percent since 1980. Since retention of cowcod was prohibited and bycatch was thought to be minimal, it was considered unlikely that overfishing was an issue. It is likely that the 2007 base model underestimated the uncertainty about stock status given steepness and the natural mortality rate were treated as fixed and known in the model.

The 2007 assessment was originally prepared as an “update” stock assessment; however, while preparing the update, an error was discovered in the previous assessment’s specification of the selectivity curve. Several revisions were proposed, including new estimates of historical landings, a corrected growth curve, and a two-fishery model. The 2007 assessment used Stock Synthesis 2, revised estimates of historical commercial catch, contained corrections to gear selectivity curves, utilized a revised growth curve, and separated the catch into commercial (all gears) and recreational fisheries rather than a single fishery. Recreational catches in the 2007 assessment were identical to those in the previous assessment, but estimates of commercial catches had been updated to reflect three additional data sources: 1) recovered port samples from Southern California (1983-1985), 2) regional summaries of total rockfish landings (1928-1968) provided by the NMFS SWFSC Environmental Research Division, and 3) California rockfish landings by region (1916-1927), published in CDF&G Fish Bulletin No. 105 (1958).

The 2007 rebuilding analysis (Dick and Ralston 2007) estimated a new  $T_{MAX}$  of 2098, 24 years later than the date estimated by Piner (2006), due in part to the corrections described above, but only 1 year earlier than the 2099 date estimated previously (Butler, *et al.* 2003). It was noted in the rebuilding analysis that rebuilding scenarios were extremely uncertain for this data-poor species, particularly with respect to steepness. Moreover, there was widespread concern about the ability to monitor the stock, and consequently to evaluate progress towards rebuilding in the future. The 2007 rebuilding analysis


projections indicated that it would not be possible to rebuild the cowcod stock by 2039, even if all the catches are eliminated, and the estimated time to rebuild under the current harvest rate (SPR = 90%) was 26 years greater than the target year of 2039 adopted under Amendment 16-4. Therefore, a modification of the Amendment 16-4 cowcod rebuilding plan was implemented in 2007 which prescribed a target year of 2072 and an SPR harvest rate of 82.1%.

The 2007 cowcod assessment was updated in 2009, with stock depletion estimated to be 4.5 percent of its unfished level at the start of 2009 (Dick, *et al.* 2009). Estimates of female spawning stock biomass in 2009 were highly uncertain. Spawning biomass had declined from an unfished biomass of 2,101-2,461 mt to 93-441 mt in 2009. The 2009 cowcod rebuilding analysis (Dick, *et al.* 2009) was used to reconsider the cowcod rebuilding plan adopted under Amendment 16-4 as mandated in a legal challenge (*NRDC v. Locke*). The revised rebuilding plan, implemented in 2011, prescribed a target year of 2068 and an SPR harvest rate 82.7%.

A new cowcod assessment of the stock in the Southern California Bight was conducted in 2013 (Dick and MacCall 2013), which estimated stock depletion to be 33.9 percent of unfished spawning biomass at the start of 2013 (Figure 1-9). The 2013 assessment suggested that cowcod in the Southern California Bight constitute a smaller, but more productive stock than was estimated from previous assessments. Median unfished and 2013 spawning biomasses were estimated to be 1,549 mt and 524 mt, respectively (Table 1-4).

The 2013 assessment used the XDB-SRA modeling platform to estimate stock status, scale, and productivity. Dick et al. (2013) fit five fishery-independent data sources: four time series of relative abundance (CalCOFI larval abundance survey, Sanitation District trawl surveys, NWFSC trawl survey, and NWFSC hook-and-line survey), and the 2002 Yoklavich et al. (2007) visual survey estimate of absolute abundance.

A cowcod catch report is planned for 2015.



**Figure 1-9. Relative depletion of cowcod south of 40°10' N lat. from 1970 to 2013 based on the 2013 stock assessment.**



## **Stock Productivity Relative to Rebuilding Success**


As in the previous assessment, production in the 2013 assessment is assumed to be a deterministic function of spawning biomass. Recruitment pulses may be evident in the abundance indices, but insufficient information is available to reliably estimate the relative strength of individual year classes.

## **Fishing Mortality**

Estimated harvest rates for cowcod were highest during the mid-1980s (Figure 1-10). Retention of cowcod was prohibited from January 2001 to present. Dick and MacCall 2013 estimated that removals of cowcod have been less than 0.2% of vulnerable biomass since 2003. The estimated harvest rate that produces long term MSY (5.5%) is nearly twice the proxy (SPR = 50%) harvest rate from the last assessment (2.7%). Unlike previous assessments, the recent increasing trends in fishery-independent surveys allow the model to estimate the rate of increase in stock size. However, the 95% posterior interval for the MSY harvest rate (2.2% - 12.6%) reflects uncertainty in the data regarding overall productivity of the stock.

Median harvest rates around 1930 were near the MSY rate, then declined due to shifts in fishing effort and WWII (Figure 1-10). Following the war, catch rates slowly increased until about 1970, then rose quickly to a maximum of approximately 54% of vulnerable biomass in the mid-1980s. The MSY harvest rate estimated in the 2013 assessment is 5.5%, similar to the proxy ( $B_{40\%}$ ) harvest rate of 5%, but higher than the SPR harvest rate in the 2009 assessment (2.7%). Median harvest rates were roughly 8-10 times the median MSY harvest rate in the mid-1980s, then declined to near zero after 2000, followed by steady increases in stock biomass.

Under the current SPR harvest rate specified in the rebuilding plan (82.7%), the median time to rebuild is 2020 (Dick and MacCall 2014). This SPR harvest rate is equivalent to an exploitation rate (catch over age 11+ biomass) of 0.007 based on the 2009 assessment.



**Figure 1-10. Time series of median harvest rates (total catch divided by age-11 and older biomass) from the base model in the 2013 cowcod assessment. The gray line is the estimated median harvest rate producing MSY.**

### Rebuilding Duration and Probabilities

The 2013 rebuilding analysis (Dick and MacCall 2014) was unique in that the Punt rebuilding program (Punt 2005) was not used given its incompatibility with XDB-SRA. In each rebuilding model run, 15,000 simulated trajectories were generated using draws from the joint posterior distribution. Since the XDB-SRA platform is not compatible with spawning potential ratios, harvest control rules were translated into exploitation rates ( $E$ ) calculated as catch/estimated age 11+ biomass. Similar to the previous cowcod rebuilding analysis, variability in future recruitment was expressed as a weighted set of different states of nature (parameter values), rather than random deviations from an average stock-recruitment relationship. While the previous rebuilding analysis accounted only for uncertainty in the Beverton-Holt steepness parameter, the current analysis accounts for uncertainty in all estimated model parameters. Estimates of total cowcod mortality have not exceeded the ACL (or OY) in any year since 2003. The estimate of median time to rebuild under the current harvest rate (2020) is 48 years earlier than the current target year of 2068.

#### 1.1.3.4 Darkblotched Rockfish

##### Distribution and Life History

Darkblotched rockfish (*Sebastes crameri*) are found from Santa Catalina Island off Southern California to the Bering Sea (Miller and Lea 1972; Richardson and Laroche 1979). They are most abundant from Oregon to British Columbia. Darkblotched primarily occur on the outer shelf and upper slope off Oregon, Washington, and British Columbia (Richardson and Laroche 1979). Based upon genetic information and the absence of large scale gaps in catches, there are no clear stock delineations for darkblotched rockfish in U.S. waters. This does not mean there are not more fine scale groupings to be found, and in fact,

darkblotched catches are characterized by infrequent large tows of larger fish. Distinct population groups have been found off the Oregon coast between 44°30' N lat. and 45°20' N lat. (Richardson and Laroche 1979). This species co-occurs with an assemblage of slope rockfish, including Pacific ocean perch (*Sebastes alutus*), splitnose rockfish (*Sebastes diploproa*), yellowmouth rockfish (*Sebastes reedi*), and sharpchin rockfish (*Sebastes zacentrus*).

Darkblotched rockfish mate from August to December, eggs are fertilized from October through March, and larvae are released from November through April (Love, *et al.* 2002). Older larvae and pelagic juvenile darkblotched rockfish are found closer to the surface than many other rockfish species. Pelagic juveniles settle at 4 to 6 cm in length in about 55 to 200 m (Love, *et al.* 2002). As with many other *Sebastes*, this species exhibits ontogenetic movement, with fish migrating to deeper waters as they mature and increase in size and age (Lenarz 1993; Nichol 1990).

Darkblotched rockfish are among the longer living rockfish; the data used in the most recent assessment (Gertseva and Thorson 2013) includes individuals that have been aged to be 98 years old. The maximum reported age of darkblotched rockfish is 105 years (Love, *et al.* 2002). As with many other *Sebastes* species, darkblotched rockfish exhibit sexually dimorphic growth; females reach larger sizes than males, while males attain maximum length earlier than females (Love, *et al.* 2002; Nichol 1990; Rogers, *et al.* 2000).

Darkblotched rockfish are ovoviviparous (Nichol and Pikitch 1994). Insemination of female darkblotched rockfish occurs from August to December, and fertilization and parturition occur from December to March off Oregon and California, and primarily in February off Oregon and Washington (Hart 1988; Nichol and Pikitch 1994; Richardson and Laroche 1979). Fecundity is dependent on size and ranges from 20,000 to 610,000 eggs.

Little is known about ecological relationships between darkblotched rockfish and other organisms. Pelagic juveniles feed on planktonic organisms such as copepods. Adults are often caught with other fish such as Pacific ocean perch and splitnose rockfish. Midwater animals such as euphausiids and amphipods dominate the diet of adult fish. Albacore and Chinook salmon consume pelagic juveniles (Hart 1988). Little is known about predation of adults.

### **Stock Status and Management History**

Darkblotched rockfish are caught primarily with commercial trawl gear, as part of a complex of slope rockfish, which includes Pacific ocean perch (*Sebastes alutus*), splitnose rockfish (*Sebastes diploproa*), yellowmouth rockfish (*Sebastes reedi*), and sharpchin rockfish (*Sebastes zacentrus*). Catches of darkblotched rockfish first became significant in the mid-to-late 1940s due to increased demand for fish protein during World War II. During the mid-1960s to mid-1970s darkblotched rockfish were caught by both domestic and foreign fleets (Rogers 2003b). Domestic landings rose from late 1970s until the late 1980s, although limits on rockfish catch were first instituted in 1983, when darkblotched rockfish was managed as part of a group of around 50 species (designated as the *Sebastes* complex) (Rogers, *et al.* 2000). During the 2000s, progressive steps have been taken to reduce the catch of darkblotched rockfish, following the declaration of its overfished status in 2001.

The first full assessment of the darkblotched rockfish stock was conducted in 2000, which estimated stock depletion at 14–31 percent of its unfished level, depending on assumptions regarding the historic catch of darkblotched rockfish in the foreign fishery from 1965-1978 (Rogers, *et al.* 2000). The base model assumed 10 percent of foreign catch was comprised of darkblotched, leading to the conclusion that the spawning stock biomass was at 22 percent of its unfished level. NMFS declared darkblotched rockfish to be overfished in 2001 based on these results.

The 2001 rebuilding analysis for the stock (Methot and Rogers 2001) incorporated results of the 2000 Alaska Fishery Science Center triennial slope trawl survey and modeled a more recent time series of recruitments. Incorporating these data resulted in a downward revision of the estimated recruitment and abundance throughout the time series compared to what had been used in the Rogers et al. (2000) assessment. This led to a revised estimate of spawning stock biomass at the beginning of 2002 of 14 percent of its unfished level and a longer projected rebuilding period.

A 2003 assessment and rebuilding update for darkblotched rockfish (Rogers 2003a) estimated a lower depletion ( $B_{11\%}$ ), but provided evidence of strong recent recruitment not yet recruited to the spawning population. This analysis was used to inform the darkblotched rockfish rebuilding plan adopted under Amendment 16-2, which established a target rebuilding year of 2030 and a fishing mortality rate of  $F = 0.027$ . A revised darkblotched rebuilding plan was implemented in 2004 that specified a higher harvest rate ( $F = 0.032$ ) to avoid negative socioeconomic impacts.

The 2005 full darkblotched assessment estimated a spawning stock depletion of 16 percent of unfished biomass at the start of 2005 (Rogers 2005a). The assessment estimated strong recruitment of the 1999 and 2000 year classes. The 2005 rebuilding analysis (Rogers 2005b) was used to inform a revised rebuilding plan adopted under Amendment 16-4 and implemented in 2007. The revised rebuilding plan specified a target year of 2011 and a constant harvest rate strategy ( $SPR = 60.7\%$ ).

The 2007 darkblotched rockfish assessment estimated a stock depletion of 22.7 percent at the start of 2007 (Hamel 2008c). The 2007 darkblotched rebuilding analysis (Hamel 2008a) predicted the median time to rebuild would be 19 years later than the target year of 2011 under the  $SPR$  harvest rate adopted under Amendment 16-4. The Council revised the Amendment 16-4 rebuilding plan by specifying a target year to rebuild the stock of 2028 and decreasing the harvest rate ( $SPR = 62.1\%$ ).

The 2007 darkblotched assessment was updated in 2009 and 2011. The 2009 stock assessment update estimated a stock depletion of 27.5 percent at the start of 2009 (Wallace and Hamel 2009). The 2009 darkblotched rebuilding analysis (Wallace 2009) was used to inform a revised rebuilding plan, which was implemented in 2011. The revised rebuilding plan specified a target year to rebuild the stock of 2025 and decreased the harvest rate to  $SPR = 64.9\%$ . The 2011 stock assessment update estimated a stock depletion of 30.2 percent at the start of 2009 (Stephens, *et al.* 2011). No revisions to the rebuilding plan were made based on the 2011 assessment update and accompanying rebuilding plan (Stephens 2011).

A full darkblotched stock assessment in 2013 (Gertseva and Thorson 2013) estimated a stock depletion of 36 percent at the start of 2013 (Figure 1-11). The assessment also predicts the stock will be rebuilt by the start of 2015. The improved stock status and rebuilding outlook were largely attributed to 1) reduced fishing mortality under the rebuilding program; 2) inferences that follow from more favorable perceptions of steepness, fecundity, and age at maturity of the stock; and 3) length and age data indicating relatively large recruitments in 1999, 2000, and 2008. The SSC recommended maintaining the current rebuilding plan for the 2015-2016 management cycle and a full assessment be done in 2015 to confirm this prediction. The SSC further recommended against preparing a rebuilding analysis in 2013. A full assessment of darkblotched rockfish is planned for 2015.




Figure 1-11. Relative depletion of darkblotched rockfish from 1980 to 2013 based on the 2013 stock assessment.

### Stock Productivity Relative to Rebuilding Success

Gertseva and Thorson (2013) fixed steepness at its prior mean of 0.779. This prior was estimated using a likelihood profile approximation to a maximum marginal likelihood mixed-effect model for steepness from ten category 1 rockfish species off the U.S. west coast (Pacific ocean perch, bocaccio, canary, chilipepper, black, darkblotched, gopher, splitnose, widow, and yellowtail rockfish). Both northern and southern assessments of black rockfish were used in the meta-analysis, although the log-likelihood for each was given a 0.5 weighting, to ensure that the together these two assessments had an equal weighting to the other species. This likelihood profile model is intended to synthesize observation-level data from assessed species, while avoiding the use of model output and thus improving upon previous meta-analyses (Dorn 2002a; Forrest, *et al.* 2010). This methodology has been simulation tested, and has been recommended by the SSC for use in stock assessments.




Figure 1-12. Estimated recruitments of darkblotched rockfish, 1970-2012.


### Fishing Mortality

Historically, the spawning output of darkblotched rockfish dropped below the  $B_{MSY}$  target for the first time in 1987, as a result of intense fishing by foreign and domestic fleets. It continued to decline and reached the level of 13 percent of its unfished output in 1999. Since 2000, when the stock was declared overfished, the spawning output slowly increased primarily due to management regulations implemented for the stock.

Overfishing for darkblotched has not occurred in the last 10 years (Gertseva and Thorson 2013). Historically, the darkblotched rockfish has experienced overfishing in the 1980s and 1990s, during the peak years of the Pacific ocean perch fishery, as well as in the mid-1960s when foreign trawl fleets were targeting groundfish off the west coast. Exploitation rates were effectively decreased after the stock was declared overfished in 2000 and rebuilding measures were implemented.

### Rebuilding Duration and Probabilities

The 2013 darkblotched assessment predicts the stock will be rebuilt by 2015. Therefore, rebuilding probabilities (both  $P_{MAX}$  and  $P_{TARGET}$ ) are high for darkblotched under the harvest control rule in the rebuilding plan. The SSC is recommending a new assessment be done in 2015 to confirm that prediction.



**Figure 1-13. Time series of estimated SPR harvest rates of darkblotched rockfish, 1960-2012. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis.**

#### 1.1.3.5 Pacific Ocean Perch

##### **Distribution and Life History**

Pacific ocean perch (POP, *Sebastes alutus*) are most abundant in the Gulf of Alaska, and have been observed off of Japan, in the Bering Sea, and south to Baja California, although they are sparse south of Oregon and rare in southern California. (Eschmeyer, *et al.* 1983; Gunderson 1971; Miller and Lea 1972). They primarily inhabit waters of the upper continental slope (Dark and Wilkins 1994) and are found along the edge of the continental shelf (Archibald, *et al.* 1983). Pacific ocean perch occur as deep as 825 m, but usually are at 200 m to 450 m and along submarine canyons and depressions (Love, *et al.* 2002). Throughout their range, POP are generally associated with gravel, rocky, or boulder type substrate (Ito, *et al.* 1986). Larvae and juveniles are pelagic; subadults and adults are benthopelagic (living and feeding on the bottom and in the water column). Adults form large schools 30 m wide, to 80 m deep, and as much as 1,300 m long (NOAA 1990). They also form spawning schools (Gunderson 1971). Juvenile POP form ball-shaped schools near the surface or hide in rocks (NOAA 1990).

Pacific ocean perch winter and spawn in deeper water (>275 m). In the summer (June through August) they move to feeding grounds in shallower water (180 m to 220 m) to allow gonads to ripen (Archibald, *et al.* 1983; Gunderson 1971; NOAA 1990). They are slow-growing and long-lived; the maximum age has been estimated at about 98 years (Heifetz, *et al.* 2000). They can grow up to about 54 cm and 2 kg (Archibald, *et al.* 1983; Beamish 1979; Gunderson 1971; Ito, *et al.* 1986; Mulligan and Leaman 1992; NOAA 1990). POP are carnivorous. Larvae eat small zooplankton. Small juveniles eat copepods, and larger juveniles feed on euphausiids (krill). Adults eat euphausiids, shrimps, squids, and small fish. Immature fish feed throughout the year, but adults feed only seasonally, mostly April through August (NOAA 1990). POP predators include sablefish and Pacific halibut.

## Stock Status and Management History

POP were harvested exclusively by U.S. and Canadian vessels in the Columbia and Vancouver INPFC areas prior to 1966. Large Soviet and Japanese factory trawlers began fishing for POP in 1965 in the Vancouver area and in the Columbia area a year later. Intense fishing pressure by these foreign fleets occurred from 1966 to 1975. The mandates of the MSA, passed by Congress in 1976, eventually ended foreign fishing within 200 miles of the United States coast.

The POP resource off the west coast was and is currently estimated to have been overfished before implementation of the groundfish FMP in 1982, and Council actions to conserve the resource likewise predate the FMP. Large removals of POP in the foreign trawl fishery, followed by significant declines in catch and abundance, led the Council to limit harvest beginning in 1979. A 20-year rebuilding plan for POP was adopted in 1981. Rebuilding under this original plan was largely influenced by a cohort analysis of 1966-1976 catch and age composition data (Gunderson 1979), updated with 1977-1980 data (Gunderson 1981), and an evaluation of trip limits as a management tool (Tagart, *et al.* 1980). This was the first time trip limits were used by the Council to discourage targeting and overharvest of an overfished stock, and it remains a management strategy in use today in the west coast groundfish fishery. In addition to trip limits, the Council significantly lowered the OY for POP. After twenty years of rebuilding under the original plan, the stock stabilized at a lower equilibrium than estimated in the pre-fishing condition. While continuing stock decline was abated, rebuilding was not achieved as the stock failed to increase in abundance to  $B_{MSY}$ .

Ianelli and Zimmerman (1998) estimated POP female spawning biomass in 1997 to be at 13 percent of its unfished level, thereby confirming that the stock was overfished. NMFS formally declared POP overfished in March 1999 after the groundfish FMP was amended to incorporate the tenets of the Sustainable Fisheries Act. The Council adopted and NMFS enacted more conservative management measures in 1999 as part of a redoubled rebuilding effort.

A 2000 POP assessment suggested the stock was more productive than originally thought (Ianelli, *et al.* 2000). A revised POP rebuilding analysis was completed and adopted by the Council in 2001 (Punt and Ianelli 2001). This analysis estimated a  $T_{MIN}$  of 12 years and a  $T_{MAX}$  of 42 years. It was noted in the rebuilding analysis that the ongoing retrospective analysis of historic foreign fleet catches was likely to change projections of POP rebuilding.

The 2003 POP assessment (Hamel, *et al.* 2003) incorporating updated survey and fishery data including the retrospective of foreign fleet catches (Rogers 2003b). The assessment covered areas from southern Oregon to the U.S. border with Canada, the southern extent of POP distribution. The overall conclusion was that the stock was relatively stable at approximately 28 percent of its unfished biomass ( $B_{28\%}$ ). Of all the changes and additions to the data, the historical catch estimates had the greatest effect, resulting in lower estimates of both equilibrium unfished biomass ( $B_0$ ) and  $MSY$ .

A POP rebuilding plan was adopted in 2003 under Amendment 16-2. The rebuilding plan was informed by a revised rebuilding analysis based on the 2000 assessment and conducted in 2001 (Punt and Ianelli 2001). The rebuilding plan established a target rebuilding year of 2027 and a harvest control rule of  $F = 0.0082$  (with a  $P_{MAX}$  of 70 percent).

The 2003 assessment estimated a stock depletion of 28 percent at the start of 2003 (Hamel, *et al.* 2003). The 2003 rebuilding analysis (Punt, *et al.* 2003) was used to amend the harvest control rule and set annual POP OYs for the 2004-2006 period. The amended harvest control rule was  $F = 0.0257$ .

The 2003 POP assessment was updated in 2005, 2007, and 2009. The 2005 update assessment estimated a stock depletion of 23.4 percent of its unfished level at the start of 2005 (Hamel 2006b). The 2005 POP



rebuilding analysis (Hamel 2006a) was used to inform revisions to the POP rebuilding plan. The revised rebuilding plan, which was adopted under Amendment 16-4, specified a target rebuilding year of 2017 and a constant harvest rate strategy (SPR = 86.4%).

The 2007 POP assessment update estimated a stock depletion of 27.5 percent at the start of 2007 (Hamel 2008d). The 2007 rebuilding analysis indicated rebuilding was progressing ahead of schedule (Hamel 2008b). No modifications to the rebuilding plan were made.

The 2009 POP assessment estimated a stock depletion of 28.6 percent at the start of 2009 (Hamel 2009b). The 2009 POP rebuilding analysis (Hamel 2009a) predicted rebuilding would not occur by the target year of 2017 with at least a 50% probability even in the absence of fishing-related mortality beginning in 2011 (i.e.,  $T_{F=0}$ ). Therefore the rebuilding plan was revised by changing the target rebuilding year to 2020 while maintaining the constant SPR harvest rate of 86.4%.

A full assessment in 2011 estimated a stock depletion of 19.1 percent at the start of 2011 (Hamel and Ono 2011). The significant decrease in the estimated depletion of the stock was largely due to a much higher estimate of initial, unfished biomass ( $B_0$ ). Previous assessments assumed a large recruitment in the late 1950s provided the higher biomass to support the estimated removals by the foreign fleets without any data to support that assumption. The assumption in the 2011 assessment is that the large foreign fleet catch fished the biomass down to critical levels, thus resulting in a substantially larger  $B_0$  estimate. The 2011 assessment also estimated a longer sequence of higher recruitment based on fitting to the data available for early years of the assessment period. The 2011 rebuilding analysis (Hamel 2011) predicted rebuilding would not occur by the target year of 2020 with at least a 50% probability even in the absence of fishing-related mortality beginning in 2013 (i.e.,  $T_{F=0}$ ). Therefore the rebuilding plan was revised by changing the target rebuilding year to 2051 while maintaining the constant SPR harvest rate of 86.4%.

A POP catch report was provided in 2013 ([Agenda Item F.5.a, Attachment 10, June 2013](#)), which indicated 2010-2012 total catches were below specified ACLs/OYs.

A POP catch report is planned for 2015.




Figure 1-14. Relative depletion of Pacific ocean perch from 1980 to 2011 based on the 2011 stock assessment.

### Stock Productivity Relative to Rebuilding Success

Stock-recruitment steepness was estimated external to the 2011 POP stock assessment base model at 0.4 (and then fixed in the model), which is low compared to steepness estimates from POP assessments conducted off Canada and Alaska. The 2011 assessment assumes no connectivity with the other assessed POP stocks in Canada and Alaska. POP off the U.S. west coast (mostly Washington and Oregon) are at the southern end of the range where there are enough POP to be commercially important, and the numbers seen are likely related to movement across the Canadian border, as well as reproductive success (recruitment), stock status, and fishing mortality north of the border. Given there is no evidence of stock structure in the meta-population of POP in the northeast Pacific and larval distribution of slope rockfish tends to be fairly geographically widespread, it is plausible that steepness is higher than determined in the 2011 assessment, which would tend to estimate a less depleted and more productive stock. The major axis of uncertainty in the assessment is steepness, with states of nature ranging from a low steepness of 0.35 to a higher value of 0.55. If steepness was as high as 0.55, the POP stock would be on the verge of being rebuilt at the start of 2011 (depletion = 39.9 percent) and projected to be rebuilt at the start of 2012. Under the base case model with a steepness of 0.4 and continuing to manage POP using the 86.4 percent SPR harvest rate in the current rebuilding plan, the stock is projected to be rebuilt by 2051.

Recruitment trends estimated in the 2011 POP assessment indicate that, like most assessed rockfish, recruitment has been relatively lower in the last few decades compared to the 1950s and 1960s. However,

the 1999 and 2000 year classes are estimated to be above average and the 2008 year class recruitment, while uncertain, appears to be the largest in at least the past 50 years (Figure 1-15).

Fishing practices are unlikely to have any effect on stock productivity, given the low fishing mortality implemented under the rebuilding plan limits. There is no indication that fishing operations are likely to substantially interfere with or disturb reproductive behavior or juvenile survival.




Figure 1-15. Time series of estimated (age-0) POP recruitments.

### Fishing Mortality

POP are caught almost exclusively by groundfish trawl gear and predominantly bottom trawls operating on the outer continental shelf and slope north of 43° N lat. POP are distributed from 30-350 fm, with the core distribution between 110-220 fm.

According to the base model in the 2011 assessment, the fishing level has been below the proxy  $F_{50\% F_{MSY}}$  harvest rate for the past 12 years (Figure 1-16), during which period the stock has begun to rebuild (Figure 1-14). The point estimates of summary (age 3+) biomass also show an upward trend over the past decade, increasing approximately 50 percent in that time.




Figure 1-16. Time series of POP exploitation rates (catch/summary biomass), 1940-2010.

### Rebuilding Duration and Probabilities

Hamel (2011) estimated a probability of rebuilding in the maximum time allowable ( $P_{MAX}$ ) under the SPR harvest rate specified in the rebuilding plan of 72.3%. There is a 50% probability of rebuilding by the target year of 2051.

#### 1.1.3.6 Petrale Sole

### Distribution and Life History

Petrale sole (*Eopsetta jordani*) is a right-eyed flounder in the family Pleuronectidae ranging from the western Gulf of Alaska to the Coronado Islands, northern Baja California, (Hart 1988; Kramer and O'Connell 1995; Love, *et al.* 2002) with a preference for soft substrates at depths ranging from 0-550 m (Love, *et al.* 2002). In northern and central California petrale sole are mostly found on the middle and outer continental shelf (Allen, *et al.* 2006).

There is little information regarding the stock structure of petrale sole off the U.S. Pacific coast. Tagging studies show adult petrale sole can move up to 350 - 390 miles, having the ability to be highly migratory with the possibility for homing ability (Alverson and Chatwin 1957; MBC 1987). Juveniles show little coastwide or bathymetric movement while studies suggest that adults generally move inshore and northward onto the continental shelf during the spring and summer to feeding grounds and offshore and southward during the fall and winter to deep water spawning grounds (Hart 1988; Love 1996; MBC 1987). Adult petrale sole can tolerate a wide range of bottom temperatures (Perry, *et al.* 1994).

Mixing of fish from multiple deep water spawning grounds likely occurs during the spring and summer when petrale sole are feeding on the continental shelf. Fish that were captured, tagged, and released off the northwest coast of Washington during May and September were subsequently recaptured during winter from spawning grounds off Vancouver Island (British Columbia, 1 fish), Heceta Bank (central Oregon, 2 fish), Eureka (northern California, 2 fish), and Halfmoon Bay (central California, 2 fish) (Pederson 1975). Fish tagged south of Fort Bragg (central California) during July 1964 were later recaptured off Oregon (11

fish), Washington (6 fish), and Swiftsure Bank (southwestern tip of Vancouver Island, 1 fish) (D. Thomas, California Department of Fish and Game, Menlo Park, CA, cited by Sampson and Lee (1999)).

The highest densities of spawning adults off of British Columbia, as well as of eggs, larvae and juveniles, are found in the waters around Vancouver Island. Adults may utilize nearshore areas as summer feeding grounds and non-migrating adults may stay there during winter (Starr and Fargo 2004).

Petrale sole spawn during the winter at several discrete deep water sites (270-460 m) off the U.S. west coast, from November to April, with peak spawning taking place from December to February (Best 1960; Casillas, *et al.* 1998; Castillo 1995; Castillo, *et al.* 1993; Garrison and Miller 1982; Gregory and Jow 1976; Harry 1959; Love 1996; Moser 1996; Reilly, *et al.* 1994). Females spawn once each year and fecundity varies with fish size, with one large female laying as many as 1.5 million eggs (Porter 1964). Petrale sole eggs are planktonic, ranging in size from 1.2 to 1.3 mm, and are found in deep water habitats at water temperatures of 4–10 degrees C and salinities of 25–30 ppt (Alderdice and Forrester 1971; Best 1960; Gregory and Jow 1976; Ketchen and Forrester 1966). The duration of the egg stage can range from approximately 6 to 14 days (Alderdice and Forrester 1971; Casillas, *et al.* 1998; Hart 1988; Love 1996).

Petrale sole larvae are planktonic, ranging in size from approximately 3 to 20 mm, and are found up to 150 km offshore foraging upon copepod eggs and nauplii (Casillas, *et al.* 1998; Hart 1988; MBC 1987; Moser 1996). The larval duration, including the egg stage, spans approximately 6 months with larvae settling at about 2.2 cm in length on the inner continental shelf (Pearcy, *et al.* 1977). Juveniles are benthic and found on sandy or sand-mud bottoms (Eschmeyer, *et al.* 1983; MBC 1987) and range in size from approximately 2.2 cm to the size at maturity, 50% of the population is mature at approximately 38 cm and 41 cm for males and females, respectively (Casillas, *et al.* 1998). No specific areas have been identified as nursery grounds for juvenile petrale sole. In the waters off British Columbia, Canada larvae are usually found in the upper 50 m far offshore, juveniles at 19–82 m and large juveniles at 25–125 m (Starr and Fargo 2004).

Adult petrale sole achieve a maximum size of around 50 cm and 63 cm for males and females, respectively (Best 1963; Pedersen 1975). The maximum length reported for petrale sole is 70 cm (Eschmeyer, *et al.* 1983; Hart 1988; Love, *et al.* 2002) while the maximum observed break and burn age is 31 years (Haltuch, *et al.* 2013).

Petrale sole juveniles are carnivorous, foraging on annelid worms, clams, brittle star, mysids, sculpin, amphipods, and other juvenile flatfish (Casillas, *et al.* 1998; Ford 1965; Pearsall and Fargo 2007). Predators on juvenile petrale sole include adult petrale sole as well as other larger fish (Casillas, *et al.* 1998; Ford 1965) while adults are preyed upon by marine mammals, sharks, and larger fishes (Casillas, *et al.* 1998; Love 1996; Trumble 1995).

One of the ambushing flatfishes, adult petrale sole have diverse diets that become more piscivorous at larger sizes (Allen, *et al.* 2006). Adult petrale sole are found on sandy and sand-mud bottoms (Eschmeyer, *et al.* 1983) foraging for a variety of invertebrates including, crab, octopi, squid, euphausiids, and shrimp, as well as anchovies, hake, herring, sand lance, and other smaller rockfish and flatfish (Birtwell, *et al.* 1984; Casillas, *et al.* 1998; Ford 1965; Kravitz, *et al.* 1977; Love 1996; Pearsall and Fargo 2007; Reilly, *et al.* 1994). On the continental shelf petrale sole generally co-occur with English sole, rex sole, Pacific sanddab, and rock sole (Kravitz, *et al.* 1977).

Castillo (1992) and Castillo *et al.* (1995) suggest that density-independent survival of early life stages is low and show that offshore Ekman transportation of eggs and larvae may be an important source of variation in year class strength in the Columbia INPFC area. The effects of the Pacific Decadal Oscillation (PDO) on California current temperature and productivity (Mantua, *et al.* 1997) may also contribute to non-stationary recruitment dynamics for petrale sole. The prevalence of a strong late 1990s year class for many

west coast groundfish species suggests that environmentally driven recruitment variation may be correlated among species with relatively diverse life history strategies.

### Stock Status and Management History

Petrale sole were lightly exploited during the early 1900s. By the 1950s the petrale sole fishery was well-developed and showing clear signs of depletion and declines in catches and biomass. Haltuch et al. (2013) estimated petrale sole biomass on the U.S. west coast dropped below the  $B_{25\%}$  management target during the 1960s and generally stayed there through 2013. The stock declined below the  $B_{12.5\%}$  overfished threshold from the early 1980s until the early 2000s (Figure 7). Since 2000 the stock has increased, reaching a peak of 14.2% of unfished biomass in 2005, followed by a decreasing trend through 2010. The petrale sole biomass currently shows an increasing trend with recent above-average year classes recruiting into the spawning biomass. The estimated relative depletion level in 2013 is 22.3 percent, which is above the MSST of  $B_{12.5\%}$  and below the  $B_{MSY}$  target of  $B_{25\%}$  (i.e., in the precautionary zone).

Early stock assessments only assessed petrale sole in the combined U.S.-Vancouver and Columbia INPFC areas (i.e., petrale in these areas were treated as a unit stock, using time series of data that began during the 1970s) (Demory 1984; Turnock, *et al.* 1993). The first assessment used stock reduction analysis and the second assessment used the length-based Stock Synthesis model. The third petrale sole assessment utilized the hybrid length-and-age-based Stock Synthesis 1 model, using data from 1977–1998 (Sampson and Lee 1999), and structured the data into separate seasonal fisheries - one for the winter spawning ground fishery that harvests larger fish and another for the rest of the year. Sampson and Lee (1999) estimated petrale sole stock depletion at 42 percent of unfished biomass at the start of 1999.

The 2005 petrale sole assessment (Lai, *et al.* 2006) was conducted assuming two separate stocks: the northern stock encompassing the U.S. Vancouver and Columbia INPFC areas and the southern stock including the Eureka, Monterey and Conception INPFC areas. Petrale sole in the north was estimated to be at 34 percent of unfished spawning stock biomass in 2005. In the south, the stock was estimated to be at 29 percent of unfished spawning stock biomass. Biomass trends were qualitatively similar in both areas, and also showed consistency with petrale sole trends in Canadian waters. Both stocks were estimated to have been below the Council's MSST of  $B_{25\%}$ <sup>5</sup> from the mid-1970s until very recently. Estimated harvest rates were in excess of the target fishing mortality rate of  $F_{40\%}$ <sup>6</sup> during this period as well. Petrale sole in both areas showed large recent increases in stock size, which was consistent with the strong upward trend in the shelf survey biomass index. In 2005, the STAR panel noted that the petrale sole stock trends were similar in both northern and southern areas in spite of the different modeling choices made for each area, and that a single coastwide assessment should be considered (Dorn, *et al.* 2006).

The 2009 petrale assessment estimated a stock depletion of 11.6 percent of its unfished biomass at the start of 2009 (Haltuch and Hicks 2009b). That result compelled NMFS to declare the stock overfished in 2010. The 2009 assessment treated petrale sole as a single coastwide stock, with the fleets and landings structured by state (WA, OR, CA) area of catch. The data series for historical catches was extended back to 1876, the first year of estimated exploitation for the stock.

New proxy management reference points used to manage FMP flatfish stocks, such as petrale sole, were implemented in 2011 under FMP Amendment 16-5 (also referred to as Secretarial Amendment 1) in 2011 (PFMC and NMFS 2011). The proxy  $F_{MSY}$  harvest rate or MFMT of  $F_{40\%}$ , which is applied to the estimated

---

<sup>5</sup>  $B_{25\%}$  was the MSST or overfished threshold for all groundfish stocks from the implementation of Amendment 12 in 1998 through 2010.

<sup>6</sup>  $F_{40\%}$  was the  $F_{MSY}$  proxy harvest rate for all flatfish stocks from 1997-2011. Prior to 1997, the proxy  $F_{MSY}$  harvest rate was  $F_{35\%}$ .

exploitable biomass to determine the OFL, was changed to  $F_{30\%}$ ; the  $B_{MSY}$  target of  $B_{40\%}$  was changed to  $B_{25\%}$ ; and the MSST of  $B_{25\%}$ , was changed to  $B_{12.5\%}$ . The SSC recommended these new proxy reference points to manage flatfish stocks based on a meta-analysis of the relative productivity of assessed west coast flatfish species and other assessed Pleuronectid species internationally. The precautionary ACL harvest control rule, referred to as the 25-5 rule and analogous to the 40-10 rule for other groundfish stocks (see Figure 1-39 and section 1.3.3 for more detail on these ACL harvest control rules), was also adopted for flatfish stocks under Amendment 16-5.

The 2009 rebuilding analysis (Haltuch and Hicks 2009a) was used to consider a petrale sole rebuilding plan for petrale sole, which was implemented under FMP Amendment 16-5. The rebuilding plan specified a target year of 2016 and the strategy of using the 25-5 harvest control rule after 2011 to set harvest levels (the 2011 ACL was set equal to the ABC to avoid unnecessary negative socioeconomic impacts). An emergency rule was implemented to reduce the 2010 petrale OY to 1,200 mt.

The 2011 petrale assessment estimated a stock depletion of 18 percent of its unfished biomass at the start of 2011 (Haltuch, *et al.* 2011). The assessment indicated an increasing spawning biomass trend with above average year classes recruiting into the spawning biomass. The 2011 rebuilding analysis (Haltuch 2011) indicated rebuilding was ahead of schedule and predicted spawning biomass would likely attain the  $B_{MSY}$  target of  $B_{25\%}$  by the start of 2013. No modifications were made to the rebuilding plan based on this result.

The 2013 petrale assessment (Haltuch, *et al.* 2013) estimated a stock depletion of 22.3 percent of its unfished biomass at the start of 2013 and short of the prediction from the 2011 rebuilding analysis; spawning biomass is predicted to reach the  $B_{MSY}$  target by the start of 2014. The 2013 stock assessment continued with the coastwide stock assessment, but was restructured to summarize petrale sole landings by the port of landing and combined Washington and Oregon into a single fleet, but structured seasonally based on winter (November to February) and summer (March to October) fishing seasons. The down-weighting of the trawl CPUE index used in the 2011 assessment was largely responsible for the more pessimistic result and the one year lag in rebuilding relative to the previous assessment. However, the estimation of recent recruitments indicated two very strong year classes (2007 and 2008; Figure 1-18) recruiting into the spawning population, which increases the likelihood of imminent success in rebuilding this stock. An update of the 2013 full petrale sole assessment is planned for 2015.




Figure 1-17. Relative depletion trend from 1980 to 2013 for petrale sole based on the 2013 stock assessment.

### Stock Productivity Relative to Rebuilding Success

Petrale have high stock productivity with an estimated stock-recruitment steepness of 0.86 (Haltuch, *et al.* 2013); the prior for this estimate was based on a meta-analysis of flatfish species in the family *Pleuronectidae* (Myers, *et al.* 1999). The time series of estimated recruitments shows a relationship with the decline in spawning biomass, punctuated by larger recruitments. The five weakest recruitments since 1934 are estimated to be from 1986, 1987, 1992, 1995, and 2001, while the five strongest recruitments since 1934 are estimated to be from 1939, 1966, 1998, 2007, and 2008. The 2007 and 2008 recruitments were the third and second largest estimated, respectively, behind only the record 1966 recruitment event (Figure 1-18). Until 2007, the most recent large recruitment event is estimated to be in 2006, which was smaller than the 1998 recruitment event.

The high stock productivity and the large recent recruitments contribute to a predicted quick recovery of the petrale sole stock. The 2013 petrale assessment predicts the stock will be successfully rebuilt by the start of 2014, with an estimated depletion of 26 percent.



## Age-0 Recruits




Figure 1-18. Time series of estimated (age 0) petrale sole recruitments, 1934-2012 (from Haltuch, *et al.* 2013).


### Fishing Mortality

Most of the petrale sole catch is made by deep-water demersal trawls at depths of 164-252 fm. Since discovery of petrale spawning grounds during the 1950s and 1960s, petrale sole catch statistics have exhibited marked seasonal variation, with substantial portions of the annual harvest taken from the spawning grounds in December and January. From the inception of the fishery in 1876 through the mid-1940s, the vast majority of catches occurred between March and October (the summer fishery), when the stock is dispersed over the continental shelf. The post-World War II period witnessed a steady decline in the amount and proportion of annual catches occurring during the summer months (March-October). Conversely, petrale catch during the winter season (November-February), when the fishery targets spawning aggregations, has exhibited a steadily increasing trend since the 1950s. Since the mid-1980s, catches during the winter months have been roughly equivalent to or exceeded catches throughout the remainder of the year. In 2009, catches of petrale sole began to be restricted due to declining stock size.

Petrale sole exhibit distinct seasonal depth migrations with higher abundance on the shelf during summer months and higher abundance in distinct spawning areas during winter months. Hence, RCA structures for this species could vary seasonally if RCA management is needed to control fishing mortality. The general pattern for petrale sole is a shallower depth distribution during the summer months (periods 3 and 4) and a deeper depth distribution during the winter months (periods 1 and 6). Petrale sole are typically in transition as they migrate between shallow and deeper depths during periods 2 and 5.

Petrale sole are caught almost exclusively by bottom trawl gears. Therefore, the uncertainty in catch monitoring and accounting is low, given the mandatory 100 percent observer coverage and near real-time reporting of total catches in the rationalized groundfish trawl fisheries.

Prior to 2010, when interim rebuilding measures were implemented, harvest rates were in excess of what is now considered the  $F_{MSY}$  limit of  $F_{30\%}$  (i.e.,  $SPR = 30\%$ ). Management measures implemented since 2010 have resulted in harvest rates below the  $F_{MSY}$  limit.



**Figure 1-19. Estimated spawning potential ratio (SPR) of petrale sole, 2004-2012. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis. The management target is plotted as a dashed horizontal line and values above this reflect harvests in excess of the overfishing proxy based on the  $F_{MSY}$  harvest rate (SPR = 30%).**

### Rebuilding Duration and Probabilities

The 2013 petrale assessment predicts the stock will be rebuilt by 2014. Therefore, rebuilding probabilities (both  $P_{MAX}$  and  $P_{TARGET}$ ) are high for petrale sole under the harvest control rule in the rebuilding plan. The SSC is recommending a new assessment be done in 2015 to confirm that prediction.

#### 1.1.3.7 Yelloweye Rockfish

##### Distribution and Life History

Yelloweye rockfish (*Sebastes ruberrimus*) range from the Aleutian Islands, Alaska, to northern Baja California, Mexico, and are common from Central California northward to the Gulf of Alaska (Eschmeyer, *et al.* 1983; Hart 1988; Love, *et al.* 2002; Miller and Lea 1972; O'Connell and Funk 1986). Yelloweye rockfish occur in water 25 m to 550 m deep with 95 percent of survey catches occurring from 50 m to 400 m (Allen and Smith 1988). Yelloweye rockfish are bottom dwelling, generally solitary, rocky reef fish, found either on or just over reefs (Eschmeyer, *et al.* 1983; Hart 1988; Love, *et al.* 2002; Miller and Lea 1972; O'Connell and Funk 1986). Boulder areas in deep water (>180 m) are the most densely populated habitat type, and juveniles prefer shallow-zone broken-rock habitat (O'Connell and Carlile 1993). They also reportedly occur around steep cliffs and offshore pinnacles (Rosenthal, *et al.* 1982). The presence of refuge spaces is an important factor affecting their occurrence (O'Connell and Carlile 1993).

Yelloweye rockfish are ovoviviparous and give birth to live young in June off Washington (Hart 1988). The age of first maturity is estimated at six years and all are estimated to be mature by eight years (Wyllie Echeverria 1987). They can grow to 91 cm (Eschmeyer, *et al.* 1983; Hart 1988) and males and females

probably grow at the same rates (Love 1996; O'Connell and Funk 1986). The growth rate levels off at approximately 30 years of age (O'Connell and Funk 1986) but they can live to be 118 years old (Love, *et al.* 2002). Yelloweye rockfish are a large predatory reef fish that usually feeds close to the bottom (Rosenthal, *et al.* 1982). They have a widely varied diet, including fish, crabs, shrimps and snails, rockfish, cods, sand lances, and herring (Love, *et al.* 2002). Yelloweye rockfish have been observed underwater capturing smaller rockfish with rapid bursts of speed and agility. Off Oregon the major food items of the yelloweye rockfish include cancrivora crabs, cottids, righteye flounders, adult rockfishes, and pandalid shrimps (Steiner 1978). Quillback and yelloweye rockfish have many trophic features in common (Rosenthal, *et al.* 1982).

### **Stock Status and Management History**

The first yelloweye rockfish stock assessment on the U.S. west coast was conducted in 2001 (Wallace 2002). This assessment incorporated two area assessments: one from Northern California using CPUE indices constructed from Marine Recreational Fisheries Statistical Survey (MRFSS) sample data and CDFG data collected on board commercial passenger fishing vessels, and the other from Oregon using Oregon Department of Fish and Wildlife (ODFW) sampling data. The assessment concluded yelloweye rockfish stock biomass in 2001 was at about 7 percent of unexploited biomass in Northern California and 13 percent of unexploited biomass in Oregon. The assessment revealed a thirty-year declining biomass trend in both areas with the last above average recruitment occurring in the late 1980s. The assessment's conclusion that yelloweye rockfish biomass was well below the 25 percent of unexploited biomass threshold for overfished stocks led to this stock being declared overfished in 2002. Until 2002, yelloweye rockfish were listed in the "remaining rockfish" complex on the shelf in the Vancouver, Columbia, and Eureka INPFC areas and the "other rockfish" complex on the shelf in the Monterey and Conception areas. As with the other overfished stocks, yelloweye rockfish harvest is now tracked separately and managed against a species-specific ACL.

In June 2002 the SSC recommended that managers should conduct a new assessment incorporating Washington catch and age data. This recommendation was based on evidence that the biomass distribution of yelloweye rockfish on the west coast was centered in waters off Washington and that useable data from Washington were available. Based on that testimony, the Council recommended completing a new assessment in the summer of 2002, before a final decision was made on 2003 management measures. Methot *et al.* (Methot, *et al.* 2003) did the assessment, which confirmed the overfished status (24 percent of unfished biomass) and provided evidence of higher stock productivity than originally assumed. The assessment also treated the stock as a coastwide assemblage. The 2002 rebuilding analysis (Methot and Piner 2002a) informed the yelloweye rockfish rebuilding plan adopted under FMP Amendment 16-3 in 2004. The rebuilding plan established a target rebuilding year of 2058 and a harvest control rule of  $F = 0.0153$ .

A coastwide 2006 yelloweye rockfish assessment estimated a stock depletion of 17.7 percent of the unfished level at the start of 2006 (Wallace, *et al.* 2006). New data sources in the assessment included WDFW 2002 submersible survey and the International Pacific Halibut Commission annual longline survey. Further revisions in the assessment included reducing natural mortality from 0.045 to 0.036 and increasing steepness from 0.437 to 0.45.

The 2006 rebuilding analysis (Tsou and Wallace 2006) was used to inform a revision of the yelloweye rebuilding plan under FMP Amendment 16-4. Given the significant negative socioeconomic impacts associated with the projected OYs under the constant harvest rate modeled in the rebuilding analysis, the Council elected to gradually ramp down the harvest rate beginning in 2007 before resuming a constant harvest rate rebuilding strategy in 2011. The harvest rate ramp-down strategy, which projected annual OYs of 23 mt, 20 mt, 17 mt, and 14 mt, respectively in 2007-2011, was projected to extend rebuilding by less

than one year relative to the more conservative constant harvest rate strategy analyzed. The ramp-down strategy afforded more time to consider new Yelloweye Rockfish Conservation Areas and other management measures designed to reduce the harvest rate to prescribed levels. Therefore, the Amendment 16-4 rebuilding plan incorporated the ramp-down strategy before resuming a constant harvest rate (SPR = 71.9%) in 2011. The rebuilding plan also specified a target rebuilding year of 2084.

The 2007 updated stock assessment for yelloweye rockfish estimated a stock depletion of 16.4 percent of initial, unfished biomass (Wallace 2008a). The long-term biomass trajectory in the 2007 updated assessment was very similar to that in the 2006 assessment. The 2007 rebuilding analysis (Wallace 2008b) indicated rebuilding progress was on track under the ramp-down strategy; therefore, no revisions were made to the rebuilding plan.

The benchmark 2009 yelloweye assessment estimated a stock depletion of 20.3 percent of initial, unfished biomass at the start of 2009 (Stewart, *et al.* 2009). The resource was modeled as a single stock, but with three explicit spatial areas: Washington, Oregon and California. Each area was modeled simultaneously with its own unique catch history and fishing fleets (recreational and commercial), with the stocks linked via a common stock-recruit relationship with negligible adult movement among areas. The assumed level of historical removals and estimated steepness were identified as the main axes of uncertainty.

The 2009 yelloweye rebuilding analysis (Stewart 2009b) was used to inform a revised rebuilding plan that was implemented under FMP Amendment 16-5. The revised rebuilding plan implemented in 2011 specified a constant harvest rate (SPR = 76%) strategy (the ramp-down strategy was abandoned) and a target year to rebuild the stock of 2074.

The 2011 yelloweye assessment (Taylor and Wetzel 2011), an update of the 2009 assessment, estimated stock depletion at 21.4 percent of initial, unfished biomass at the start of 2011 (Figure 1-20). The update assessment results were very similar to those in the previous assessment. The 2011 yelloweye rebuilding analysis (Taylor 2011) indicated rebuilding progress was on schedule and no revisions were made to the rebuilding plan.

A yelloweye catch report was provided in 2013 ([Agenda Item F.5.a, Attachment 11, June 2013](#)), which indicated 2010-2012 total catches were below specified ACLs/OYs.

A yelloweye rockfish catch report is planned for 2015.




Figure 1-20. Relative depletion of yelloweye rockfish from 1980 to 2011 based on the 2011 stock assessment update.

### Stock Productivity Relative to Rebuilding Success

Yelloweye year class strength is modeled as a deterministic process in the 2011 assessment with no estimation of the size of individual year classes. Therefore, the decline in estimated recruitment tracks closely to that of the spawning output (Figure 1-21). The decline is especially pronounced given the low (and likely imprecise) estimate for steepness of the stock-recruit relationship in the base-case model (0.441). The low estimated steepness in the assessment results in a prediction of very little surplus production and consequently estimates of low yields at  $B_{MSY}$  (MSY is estimated to be 58 mt under the  $F_{MSY}$  proxy SPR harvest rate of 50 percent). This relatively low stock productivity also predicts a long mean generation time of 46 years and a slow recovery rate under the very low harvest rate specified in the yelloweye rebuilding plan.




Figure 1-21. Time series of estimated yelloweye rockfish spawning output and recruitments for the base-case model in the 2011 assessment (Taylor and Wetzel 2011).

### Fishing Mortality

Yelloweye rockfish are caught coastwide in all sectors of the fishery. Yelloweye are particularly vulnerable to hook-and-line gears, which are effective in the high relief habitats yelloweye reside. The current non-trawl RCA and the recreational depth closures are primarily configured based on yelloweye distribution and projected impacts in these hook-and-line fisheries. Small footrope trawls, including selective flatfish trawls, do not have the rollers and anti-chafing protection needed to fish in the high relief habitats yelloweye are found. Mandating these gears for trawl efforts on the shelf shoreward of the trawl RCA, the configuration of the trawl RCA, and a small IFQ allocation of yelloweye are the primary strategies currently used to minimize trawl impacts on yelloweye. Yelloweye are also a bycatch species in the Pacific halibut fishery (Love, *et al.* 2002).

Yelloweye rockfish are mostly encountered north of 36° N lat. Yelloweye occur in depths from 25 to 475 m and are most commonly found at depths from 91 to 180 m (Love, *et al.* 2002).

Fishing mortality rates estimated in the 2011 assessment have been in excess of the current  $F_{MSY}$  harvest rate for rockfish (SPR = 50 percent) from 1976 through 1999 (Figure 1-22). Relative exploitation rates (catch/biomass of age-8 and older fish) are estimated to have peaked at 12.7 percent in 1992, but have been at or less than 1.1 percent after 2001. The  $F_{MSY}$  exploitation rate assuming the proxy SPR of 50 percent is 2.2 percent. Annual yelloweye harvest rates in the 1976-1999 period averaged over five times the estimated  $F_{MSY}$  and spawning biomass declined rapidly during that period.

The commercial RCAs substantially reduce yelloweye impacts. North of 40°10' N lat., the highest bycatch rates of yelloweye rockfish occur in waters less than 100 fm. Yelloweye rockfish have a patchy distribution and as such, using fleetwide bycatch rates over a large area (north and south of 40°10' N lat.) may misrepresent actual catch rates. North of Cape Alava, yelloweye bycatch rates are lowest inside of the 60 fm line; bycatch rates would increase substantially if shoreward RCAs were moved from the 60 fm line to the 75 fm line. The seaward boundary of the non-trawl RCA extends out to 150 fm year round south of

40°10' N lat. The seaward boundary of the non-trawl RCA north of 40°10' N lat. is at 100 fm year round with a few exceptions where the seaward boundary is at 125 fm. Between 45°03.83' to 43° N lat. the seaward is at 125 fm year round.

Area closures and a prohibition on retention are the main strategies used to minimize recreational yelloweye impacts. The California recreational fishery is subject to depth restrictions that are more restrictive in the northern management areas where yelloweye are more prevalent. CDFG evaluated and has available four potential YRCAs which include habitat in both state and Federal waters where high yelloweye encounter rates have been documented. If implemented, YRCAs are anticipated to reduce yelloweye impacts during the open fishing seasons in both the Northern Groundfish Management Area and the North-Central North of Pt. Arena Groundfish Management Area, possibly allowing for a longer fishing season. To date, these YRCAs have not been implemented but would remain available management measures that can be routinely implemented inseason if needed. Depth management is the main tool used for controlling yelloweye rockfish fishing mortality in the Washington and Oregon recreational fisheries.




Figure 1-22. Time series of estimated relative exploitation rates (catch/biomass of age 8+ fish) of yelloweye rockfish, 1916-2010 (Taylor and Wetzel 2011).

Catch monitoring uncertainty is high given the relatively small contribution of yelloweye to rockfish market categories and the relatively large scale of recreational removals. In addition, since 2001, management restrictions have required nearly all yelloweye rockfish caught by recreational and commercial fishermen to be discarded at sea. Precisely tracking recreational catch inseason, especially in the California recreational fishery, has been a challenge.

### Rebuilding Duration and Probabilities

Rebuilding under the SPR harvest rate specified in the rebuilding plan has a predicted  $P_{MAX}$  of 72.9 percent and a probability of rebuilding by the target year of 2074 of 62.1 percent.

## 1.1.4 Non-Overfished Groundfish Stocks

### 1.1.4.1 Arrowtooth Flounder

#### Distribution and Life History

Arrowtooth flounder (*Atheresthes stomias*) range from the southern coast of Kamchatka to the northwest Bering Sea and Aleutian Islands to San Simeon, California. Arrowtooth flounder is the dominant flounder species on the outer continental shelf from the western Gulf of Alaska to Oregon. They are members of the family Pleuronectidae, the right eyed flounders. Arrowtooth reach sizes of nearly 90 cm and can live to 27 years. Eggs and larvae are pelagic; juveniles and adults are demersal (Garrison and Miller 1982; NOAA 1990). Juveniles and adults are most commonly found on sand or sandy gravel substrates, but occasionally occur over low-relief rock-sponge bottoms. Arrowtooth flounder exhibit a strong migration from shallow water summer feeding grounds on the continental shelf to deep water spawning grounds over the continental slope (NOAA 1990). Depth distribution may vary from as little as 50 m in summer to more than 500 m in the winter (Garrison and Miller 1982; NOAA 1990; Rickey 1995).

Arrowtooth flounder are oviparous with external fertilization and eggs are about 2.5 mm in diameter. Spawning may occur deeper than 500 m off Washington (Rickey 1995). Arrowtooth are batch spawners (Rickey 1995). They spawn in the deeper continental shelf waters (>200 m) in the late fall through early spring and appear to move inshore during the summer (Zimmerman and Goddard 1996). The larvae spend approximately four weeks in the upper 100 m of the water column (Fargo and Starr 2001) and settle to the bottom in the late winter and early spring. Larvae eat copepods, their eggs, and copepod nauplii (Yang 1995; Yang and Livingston 1985). Juveniles and adults feed on crustaceans (mainly ocean pink shrimp and krill) and fish (mainly gadids, herring, and pollock) (Hart 1988; NOAA 1990).

Arrowtooth flounder exhibit two feeding peaks, at noon and midnight. Arrowtooth are piscivorous, but they also eat shrimp, worms, and euphausiids (Love 1996). Buckley et al. (1999) analyzed 380 arrowtooth stomachs that were collected in 1989 and 1992 from Oregon and Washington and found that hake (*Merluccius productus*) and unidentified gadids dominate their stomach contents (45 percent and 22 percent respectively) followed by herring (19 percent; *Clupea pallasii*), mesopelagics (0.5 percent), rex sole (1 percent; *Glyptocephalus zachirus*), slender sole (*Lyopsetta exilis*) and other small flatfish (3 percent), other arrowtooth (1.5 percent), other unidentified flatfish (1 percent), pandalid shrimp (~3 percent), and euphausiids (3 percent). Yang (1995) analyzed 1,144 stomachs from arrowtooth collected in the Gulf of Alaska, and found that walleye pollock (*Theragra chalcogramma*) composed 66 percent of the arrowtooth diet, although arrowtooth smaller than 40 cm primarily feed on capelin (*Mallotus villosus*), herring, and shrimp. Gotshall (1969) examined 425 arrowtooth stomachs from northern California throughout the 1960s and found that pandalid shrimp made up nearly 40 percent of the prey by volume, along with other shrimps, crabs, euphausiids, Pacific sanddabs (*Citharichthys sordidus*), and slender sole. However, Gotshall's samples were taken directly from shrimp beds, so higher concentrations of shrimp would be expected. It is clear that arrowtooth have a broad diet, consuming most of the common fish and invertebrates found on soft bottom substrate and in the water column.

Predators of juvenile arrowtooth include skates, dogfish, shortspine thornyhead, halibut, coastal sharks, orcas, toothed whales, and harbor seals (Field, et al. 2006). Adult arrowtooth are likely to be vulnerable only to the largest of these predators.

Female arrowtooth off Oregon reach 50 percent maturity at 8 years of age, and males at four years (Hosie 1976). Rickey (1995) found that the arrowtooth reach 50 percent maturity at lengths of 36.8 cm for females and 28 cm for males off Washington, and 44 cm for females and 29 cm for males off Oregon. As a



comparison, female length at 50 percent maturity is 47 cm in the Gulf of Alaska (Turnock, *et al.* 2005) and 38 cm in British Columbia (Fargo and Starr 2001).

### **Stock Status and Management History**

Arrowtooth are commonly caught by trawl fleets off Washington and Oregon, but they are frequently discarded due to low flesh quality. For this reason, the market for arrowtooth has been fairly limited over the last 50 years. It is likely that the stock off the U.S. west coast is linked to the population off British Columbia and, possibly, to the stock in the Gulf of Alaska. However, for assessment purposes it is assumed that the U.S. west coast population is a unit stock.

The west coast stock of arrowtooth flounder was assessed in 1993 (Rickey 1993), and a full stock assessment was done in 2007 (Kaplan and Helser 2008). Three components of the arrowtooth fishery were used in modeling: the mink food fishery in the 1950s-1970s; a targeted fillet/headed-and-gutted fishery that began around 1981; and a “bycatch fleet” that represents west coast trawl effort with arrowtooth bycatch, but no landings. Estimates of historical catch are highly uncertain. The model contains assumed fixed values for natural mortality and steepness of the stock-recruitment relationship. Likelihood profiles suggest that the estimates of biomass and depletion are not sensitive to values of steepness. Assumed values of natural mortality have a small effect on estimated depletion, but strongly influence the estimates of absolute biomass.

The base model shows a period of moderate depletion through the 1950s and 1960s, followed by a rebuilding of the stock beginning in the late 1970s. Strong year classes, in particular the 1999 year class, have led to an increase in the stock since the late 1990s. The spawning biomass at the beginning of 2007 was estimated to be 63,302 mt and 79 percent of the estimated unfished spawning biomass. Total biomass at the start of 2007 was estimated to be 85,175 mt. The 2007 stock assessment estimated that the arrowtooth stock has never fallen below the overfished threshold.

A new data-moderate assessment of arrowtooth flounder is planned for 2015 since the 2007 assessment is now considered out of date.

### **Stock Productivity**

Arrowtooth flounder are a very productive stock with high growth rates, high natural mortality rates, and a high stock-recruitment steepness. A mean flatfish steepness of 0.8 was determined in a 2010 meta-analysis conducted by the SSC and described in the 2011-2012 specifications FEIS (PFMC and NMFS 2011). A steepness of 0.902 was assumed in the 2007 arrowtooth flounder assessment based on a flatfish meta-analysis conducted by Dorn (2002a). Arrowtooth received a relatively high productivity score of 1.95 in the PSA analysis (Table 1-2).

The 2007 assessment estimated strong recruitments for most years between 1998 and 2007, with a particularly strong recruitment of the 1999 year class. That year class has dominated the population and fishery for the last ten years but is now diminished through high natural mortality. However, the 2007 assessment projects a very healthy stock through 2018 under catch streams much higher than has been realized since then.

### **Fishing Mortality**

The target  $F_{MSY}$  SPR harvest rate for arrowtooth is 30 percent. The 2007 assessment estimated annual SPR harvest rates between 1997 and 2006 of 49-75 percent, substantially lower than the target. The arrowtooth ACL/OY has never been exceeded.

Arrowtooth flounder are a trawl-dominant species and are not particularly valuable. Given that arrowtooth are caught on the northern shelf where Pacific halibut, darkblotched rockfish, and yelloweye rockfish are caught incidentally to arrowtooth, this is not a species with a high attainment since valuable quota for these highly constraining species would have to be invested to target arrowtooth. About 26% of the arrowtooth quota was attained in the 2012 fishery (see <https://www.webapps.nwfsc.noaa.gov/ifq/>). Management uncertainty is low with the 100 percent observer coverage for the trawl fleet under trawl rationalization. The PSA vulnerability score of 1.21 indicates a low concern of overfishing.

#### 1.1.4.2 Black Rockfish off California and Oregon

##### **Distribution and Life History**

Black rockfish (*Sebastes melanops*) are found from Southern California (San Miguel Island) to the Aleutian Islands (Amchitka Island) and they occur most commonly from San Francisco northward (Hart 1988; Miller and Lea 1972; Phillips 1957; Stein and Hassler 1989). Black rockfish occur from the surface to greater than 366 m; however, they are most abundant at depths less than 54 m (Stein and Hassler 1989). Off California, black rockfish are found along with the blue, olive, kelp, black-and-yellow, and gopher rockfishes (Hallacher and Roberts 1985). The abundance of black rockfish in shallow water declines in the winter and increases in the summer (Stein and Hassler 1989). Densities of black rockfish decrease with depth during both the upwelling and non-upwelling seasons (Hallacher and Roberts 1985). Off Oregon, larger fish seem to be found in deeper water (20 m to 50 m) (Stein and Hassler 1989). Black rockfish off the northern Washington coast and outer Strait of Juan de Fuca exhibit no significant movement. However, fish appear to move from the central Washington coast southward to the Columbia River, but not into waters off Oregon. Movement displayed by black rockfish off the northern Oregon coast is primarily northward to the Columbia River (Culver 1986). Black rockfish form mixed sex, midwater schools, especially in shallow water (Hart 1988; Stein and Hassler 1989). Black rockfish larvae and young juveniles (<40 mm to 50 mm) are pelagic, but are benthic at larger sizes (Laroche and Richardson 1980).

Black rockfish have internal fertilization and annual spawning (Stein and Hassler 1989). Parturition occurs from February through April off British Columbia, January through March off Oregon, and January through May off California (Stein and Hassler 1989). Spawning areas are unknown, but spawning may occur in offshore waters because gravid (egg-carrying) females have been caught well offshore (Dunn and Hitz 1969; Hart 1988; Stein and Hassler 1989). Black rockfish can live to be more than 20 years in age. The maximum length attained by the black rockfish is 60 cm (Hart 1988; Stein and Hassler 1989). Off Oregon, black rockfish primarily prey on pelagic nekton (anchovies and smelt) and zooplankton such as salps, mysids, and crab megalops. Off Central California, juveniles eat copepods and zoea, while adults prey on juvenile rockfish, euphausiids, and amphipods during upwelling periods. During periods without upwelling they primarily consume invertebrates. Black rockfish feed almost exclusively in the water column (Culver 1986). Black rockfish are known to be eaten by lingcod and yelloweye rockfish (Stein and Hassler 1989).

##### **Stock Status and Management History**

A black rockfish assessment was completed in 2003 and pertained to the portion of the coastwide stock occurring off the coasts of Oregon and California (Ralston and Dick 2003) or the southern stock unit. Alternative harvest levels in the 2003 assessment were ranged to capture the major uncertainty of historical landings prior to 1978. Black rockfish catches prior to 1945 were assumed to be zero in the assessment. Many gaps in historical landings of black rockfish since 1945 were evident, and these landings were reconstructed using a variety of data sources. The base model assumed cumulative landings of black rockfish from all fisheries was 17,100 mt from 1945 to 1977. The 2003 assessment concluded the southern

California-Oregon stock of black rockfish was in healthy condition with a 2002 spawning output estimated to be at 49 percent of its unexploited level.

The southern stock of black rockfish was again assessed in 2007 (Sampson 2008) using a similar approach and structure as the 2003 assessment, but included historical catch series that extended back to 1916 with relatively large catches of black rockfish in California during World War II. The 2007 assessment estimated the southern stock was at 70 percent of its unfished level at the start of 2007. The 2007 assessment was structured into six fisheries: a set of trawl, commercial non-trawl, and recreational fisheries for Oregon and California, respectively. The fisheries for each state were based on fish capture location rather than where they were landed and therefore represented separate geographic areas. The model in the 2007 assessment did not include any underlying spatial structure in the population dynamics. Like the previous southern stock assessment, abundance indices for tuning the assessment were based on recreational CPUE data with two independent indices available for each state. The standard research trawl surveys along the U.S. west coast do not operate in shallow enough water to catch appreciable numbers of black rockfish and therefore do not provide any fishery independent index of stock biomass for black rockfish. The 2007 assessment had two additional abundance indices that were not available for the previous assessment: a black rockfish pre-recruit index for 2001-2006 and estimates from a tag-recapture study of exploitable black rockfish abundance off Newport, Oregon for 2003-2005. The 2007 assessment for the southern stock of black rockfish used the same sex- and age-specific formulation for natural mortality (M) that was used in the assessment for northern black rockfish, but there is little evidence to confirm that the assumed formulation is correct. The 2003 assessment for southern black rockfish used much smaller values for M that were more consistent with observed values for the maximum age of southern black rockfish.

A new full assessment of black rockfish is planned for 2015 since the 2007 assessment is now considered out of date.

### **Stock Productivity**

The 2007 southern black rockfish assessment assumed a steepness of 0.6 based on the Dorn meta-analysis of rockfish steepness done at that time. The revised rockfish steepness meta-analysis now predicts a mean steepness of 0.779. The PSA productivity score of 1.33 indicates a stock of moderate productivity.

The 2007 assessment estimated above-average recruitments in the 1990s (with particularly strong recruitments in 1994 and 1999), 2000, 2001, and 2007; and below-average recruitments during 2002-2006. These recruitments are projected to keep the stock healthy under the 1,000 mt constant catch strategy implemented in 2009.

### **Fishing Mortality**

The nearshore commercial and recreational fisheries that take black rockfish are managed well in California and Oregon, and ACLs/OYs have not been exceeded. The PSA vulnerability score of 1.94 indicates a stock of medium concern for overfishing.

Over most of the stock's history the fishing rate has been less than the 50% SPR target fishing rate. The estimated spawning output has been above the target level during all years except 1991 to 1998, and has never dropped below the overfished level. The southern stock of black rockfish is estimated to be well above the overfished level.

### 1.1.4.3 Black Rockfish off Washington

#### **Distribution and Life History**

See the description of black rockfish distribution and life history in section 1.1.4.2.

#### **Stock Status and Management History**

The black rockfish stock found between Cape Falcon, Oregon and the U.S. Canadian border was first assessed in 1994 (Wallace and Tagart 1994). Estimated biomass was 60 percent of the unfished level and female egg production was estimated to be 43 percent of the unfished level. A harvest guideline of 517 mt for this area was specified beginning in 1995 based on assessment results. Catches remained well below the harvest guideline in the years subsequent to the assessment.

The 1999 assessment of the black rockfish stock north of Cape Falcon, Oregon determined the stock was at 45 percent of the unfished level (Wallace, *et al.* 1999). The population was regarded as healthy and stock abundance was estimated to be slightly increasing after a period of low abundance in the late 1980s and early 1990s.

The most recent assessment of the northern stock was done in 2007, which estimated a depletion of 53.4 percent of the unfished level (Wallace, *et al.* 2008). The base model for the 2007 assessment assumed a female natural mortality rate to be age-specific using age at first and full maturity for inflections (10 and 15). A constant natural mortality rate of 0.16 was assumed for males and young females (< 10 years of age), and a rate of 0.2 was assumed for old females ( $\geq 15$  years of age). Model sensitivity analysis showed that model configurations using higher natural mortality for older females provided better overall fits to the data. In the model, spawning biomass and age 3+ biomass reached the lowest levels in 1995, following poor recruitment and intense fishing in the late 1980s. The population trajectory remained just above minimum stock size threshold, and the model indicated that the stock is currently well above the management target of  $B_{40\%}$ .

A new full assessment of black rockfish is planned for 2015 since the 2007 assessment is now considered out of date.

#### **Stock Productivity**

The 2007 assessment assumed a steepness 0.6 in the stock-recruitment relationship of the northern black rockfish stock based on the Dorn prior (as was done in the southern black rockfish assessment). Steepness may be even higher based on the revised prior of 0.779. The PSA productivity score of 1.33 indicates a stock of moderate productivity.

The 2007 assessment estimated strong recruitments in the 1990s (including strong recruitments in 1994 and 1999 as also estimated in the southern assessment) and above-average recruitments from 2002-2006.

#### **Fishing Mortality**

Total mortality of black rockfish off Washington has consistently been well below established ACLs/OYs. The stock is targeted in the Washington recreational fishery; however, that fishery is tightly regulated to minimize canary and yelloweye rockfish impacts. There is also a relatively low tribal take of black rockfish off Washington. There are no commercial nearshore fisheries off Washington.

Exploitation of black rockfish reached a peak in 1988 of 13 percent of the age 3+ biomass and remained near that level for 7 years, dropping precipitously between 1995 and 2000. In recent years exploitation has been relatively low (4-6 percent). Exploitation rate relative to spawning biomass indicate that harvest rates exceeded management targets between the mid-1980s through the mid-1990s for the northern stock of black rockfish.

The PSA vulnerability score of 1.94 indicates a stock of medium concern for overfishing.

#### 1.1.4.4 Cabezon off California

##### **Distribution and Life History**

Cabezon (*Scorpaenichthys marmoratus*) are distributed along the entire west coast of the continental United States. They range from central Baja California north to Sitka, Alaska (Love 1996; Miller and Lea 1972). Cabezon are primarily a nearshore species found intertidally and among jetty rocks, out to depths of greater than 100 m (Love 1996; Miller and Lea 1972).

Cabezon are known to spawn in recesses of natural and manmade objects, and males are reported to show nest-guarding behavior (Garrison and Miller 1982). Spawning is protracted, and there appears to be a seasonal progression of spawning that begins off California in winter and proceeds northward to Washington by spring. Spawning off California peaks in January and February (O'Connell 1953) while spawning in Puget Sound (Washington State) occurs for up to 10 months (November-August), peaking in March-April (Lauth 1987). Laid eggs are sticky and adhere to the surface where deposited. After hatching, the young of the year spend 3-4 months as pelagic larvae and juveniles. Settlement takes place after the young fish have attained 3-5 cm in length (Lauth 1987; O'Connell 1953). It is apparent that females lay multiple batches in different nests, but whether these eggs are temporally distinct enough to qualify for separate spawning events is not understood (Lauth 1987; O'Connell 1953).

##### **Stock Status and Management History**

Cabezon in California waters was first assessed in 2003; depletion was estimated at 34.7 percent at the start of 2003 (Cope, *et al.* 2004). The assessment delineated two stocks (north and south) at the Oregon-California border, a distinction based on differences in the catch history, CPUE trends and biological parameters (mainly growth) between the two areas. Due to the lack of data for the northern population, the assessment focused on only the southern population. As with most nearshore groundfish stocks, this assessment lacked a fishery-independent index of abundance, and consequently relied on recreational CPUE indices and information about larval abundance.

The 2005 assessment modeled two California substocks north and south of Point Conception (Cope and Punt 2006). Historically, the recreational fishery had been the primary source of removals of cabezon in California; however, commercial catches had become a major source of removals in the ten years preceding the assessment because of the developing live-fish fishery. Removals were reconstructed back to 1916, when the commercial fishery began. The estimated stock depletions of the northern and southern substocks of cabezon at the start of 2005 were 40.1 percent and 28.3 percent, respectively.

The most recent cabezon assessment for cabezon occurring in waters off California, done in 2009, estimated a stock depletion of 48.3 percent of unfished biomass at the start of 2009 (Cope and Key 2009). The 2009 assessment modeled two California substocks, and also evaluated the population as a coastwide California stock. The SSC recommended combining the results of the area models for the two California substocks of cabezon for use in deciding statewide harvest specifications.

## **Stock Productivity**

The 2009 cabezon assessment assumed a steepness of 0.7 for all models. The PSA productivity score of 1.72 indicates a stock of relatively high productivity.

Recruitment deviations were estimated from 1970-2006 for both of the assessed substocks. Recruitment patterns are distinctly different for the substocks occurring north and south of Pt. Conception at 34°27' N lat. Large recruitment events in the 1970s and 1990s in the north and the south have increased spawning biomass to healthy levels. Interannual variation in recruitment is greater in the north. The large increase in biomass in the south was driven by a large 1999 recruitment, the largest seen in the time series. Large recruitments in the southern substock were estimated immediately after major El Niño events (e.g., 1984 and 1994 recruitments). Recruitment events for the northern substock appear to lag large recruitments in the south by a year.

## **Fishing Mortality**

Exploitation of the southern cabezon substock began in the 1960s and caused a substantial decline in stock biomass. The large recruitments discussed above and a reduction in exploitation rates in the late 1990s and 2000s caused the substock to rebound to healthy levels. Exploitation in the north also increased in the 1960s, although fishing pressure was not as great. The spawning biomass of the northern substock declined, although not as dramatically as in the south. The stock rebounded with good recruitment and a reduction in fishing pressure.

The cabezon stock(s) off California were first assessed in 2003, and OYs were first specified in 2004. Specified OYs were exceeded in each year through 2006, but a reduction in cumulative landing limits adequately reduced fishing mortality starting in 2007. The percent of OY attainment ranged from 56 to 74 percent in the 2007-2010 period.

The PSA vulnerability score of 1.68 indicates a low risk of overfishing.

### **1.1.4.5 Cabezon off Oregon**

#### **Distribution and Life History**

See the description of cabezon distribution and life history in section 1.1.4.4.

#### **Stock Status and Management History**

The 2009 assessment of the Oregon substock of cabezon (Cope and Key 2009) was the first for cabezon in Oregon waters; the assessment indicated a healthy stock status for Oregon cabezon at 52.4 percent depletion at the start of 2009. Only one index of abundance was used for modeling the Oregon cabezon substock (the Oregon Recreational Boat Survey or ORBS CPUE index). The Oregon model was robust to almost all data and parameter manipulation trials except the removal of the ORBS survey. Removal of the only abundance index causes the population to drop sharply below the overfished level and absolute biomass to be much smaller than in the base case. Unlike the assessments for the California substocks, the assessment of the Oregon cabezon substock does not show recent increases in spawning biomass. While the uncertainty in the estimated depletion level of the Oregon substock is generally low, uncertainty in the estimated spawning biomass is high.

## **Stock Productivity**

Steepness in the 2009 assessment of the Oregon substock of cabezon was assumed to be 0.7. Recruitment in the Oregon substock of cabezon was estimated to be less dynamic than that for the California substocks. The PSA productivity score of 1.72 indicates a stock of relatively high productivity.

The assessment estimates large recruitments in 1999 and 2004. Uncertainty in estimating recruitment for the Oregon substock is less than the uncertainty in recruitment estimation for the California substocks.

## **Fishing Mortality**

Cabezon exploitation in Oregon started in the 1970s and caused the biomass to decline. However, exploitation was not excessive and the estimated spawning biomass has always been above the  $B_{MSY}$  target.

The PSA vulnerability score of 1.68 indicates a low risk of overfishing.

### **1.1.4.6 California Scorpionfish**

#### **Distribution and Life History**

California scorpionfish (*Scorpaena guttata*), also known locally as sculpin, is a generally benthic species found from central California to the Gulf of California in depths between the inter-tidal and about 170 m (Eschmeyer, *et al.* 1983; Love, *et al.* 1987). California scorpionfish generally inhabits rocky reefs, but in certain areas and seasons they aggregate over sandy or muddy substrate (Frey 1971; Love, *et al.* 1987). Catch rate analysis and tagging studies show that most, but not all, California scorpionfish migrate to deeper water to spawn during May-September (Love, *et al.* 1987). Tagging data suggest that they return to the same spawning site (Love, *et al.* 1987), but information is not available on non-spawning season site fidelity. California scorpionfish are quite mobile and may not be permanently tied to a particular reef (Love, *et al.* 1987).

California scorpionfish spawn from May through August, peaking in July (Love, *et al.* 1987). The species is oviparous, producing floating, gelatinous egg masses in which the eggs are embedded in a single layer (Orton 1955). California scorpionfish utilize the “explosive breeding assemblage” reproductive mode in which fish migrate to, and aggregate at traditional spawning sites for brief periods (Love, *et al.* 1987). These spawning aggregations have been targeted by fishermen. Few California scorpionfish are mature at one year of age, but over 50 percent are mature by age two and most are mature by age three (Love, *et al.* 1987).

The species feeds on a wide variety of foods, including crabs, fishes, octopi, isopods and shrimp, but juvenile Cancer crabs are the most important prey (Limbaugh 1955; Love, *et al.* 1987).

#### **Stock Status and Management History**

California scorpionfish were assessed in 2005 (Maunder, *et al.* 2006) in the southern California Bight south of Point Conception at 34°27' N lat. to the U.S.-Mexico border. The stock assessment indicated the California scorpionfish stock was healthy with an estimated spawning stock biomass of 79.8 percent of its initial, unfished biomass in 2005.

In most years, 99 percent or more of the landings occur in the southern California ports. The California nearshore FMP includes California scorpionfish. The stock is managed by the state under provisions for improved fishery monitoring and research data collection.

## Stock Productivity

A steepness value of 0.7 was assumed for California scorpionfish in the 2005 assessment. The PSA productivity score of 1.83 indicates a stock of relatively high productivity, especially for a rockfish.

The assessment noted a high recruitment variation in the stock and recruitments in the 1990s and early 2000s were estimated to be substantially above average. Relatively large recruitment events were estimated starting in 1984.

## Fishing Mortality

A substantial but unknown portion of the stock occurs in Mexican waters. The exploitation of the stock in Mexican waters is unknown and the connectivity of that stock with the U.S. stock in the Southern California Bight is also unknown.

Commercial catch records for scorpionfish were available beginning in 1928. Commercial catches were the dominant removals until the 1990s when the recreational catch became dominant. High catches and low recruitments in the 1950s and 1960s precipitated a decline in biomass. Stock biomass has been on an increasing trend since the mid-1970s.

The PSA vulnerability score of 1.41 indicates a low risk of overfishing.

### 1.1.4.7 Chilipepper Rockfish South of 40°10' N Lat.

## Distribution and Life History

Chilipepper rockfish (*Sebastes goodei*) are found from Magdalena Bay, Baja California, Mexico, to as far north as the northwest coast of Vancouver Island, British Columbia (Allen 1982; Hart 1988; Miller and Lea 1972). The region of greatest abundance is found between Point Conception and Cape Mendocino, California. Chilipepper have been taken as deep as 425 m, but nearly all in survey catches were taken between 50 and 350 m (Allen and Smith 1988). Adults and older juveniles usually occur over the shelf and slope; larvae and small juveniles are generally found near the surface. In California, chilipepper are most commonly found associated with deep, high relief rocky areas and along cliff drop-offs (Love, *et al.* 1990), as well as on sand and mud bottoms (MBC 1987). They are occasionally found over flat, hard substrates (Love, *et al.* 1990). Love (1996) does not consider this to be a migratory species. Chilipepper may travel as far as 45 m off the bottom during the day to feed (Love 1996). Chilipepper rockfish are described as an elongate fish with reduced head spines similar in appearance to both shortbelly rockfish (at smaller sizes, although shortbelly tend to be slimmer) and bocaccio rockfish (bocaccio tend to have larger mouths).

Chilipeppers are ovoviviparous and eggs are fertilized internally (Reilly, *et al.* 1992). Chilipepper school by sex just prior to spawning (MBC 1987). In California, fertilization of eggs begins in October and spawning occurs from September to April (Oda 1992) with the peak occurring during December to January (Love, *et al.* 2002). Chilipepper may spawn multiple broods in a single season (Love, *et al.* 2002). Females of the species are significantly larger, reaching lengths of up to 56 cm (Hart 1988). Males are usually smaller than 40 cm (Dark and Wilkins 1994). Males mature at two years to six years of age, and 50 percent are mature at three years to four years. Females mature at two years to five years with 50 percent mature at three years to four years (MBC 1987). Females may attain an age of about 27 years, whereas the maximum age for males is about 12 years (MBC 1987).



Larval and juvenile chilipepper eat all life stages of copepods and euphausiids, and are considered to be somewhat opportunistic feeders (Reilly, *et al.* 1992). In California, adults prey on large euphausiids, squid, and small fishes such as anchovies, lanternfish, and young Pacific whiting (Hart 1988; Love, *et al.* 2002). Chilipepper are found with widow rockfish, greenspotted rockfish, and swordspine rockfish (Love, *et al.* 2002). Juvenile chilipepper compete for food with bocaccio, yellowtail rockfish, and shortbelly rockfish (Reilly, *et al.* 1992). Pelagic juveniles are preyed upon by a wide range of predators, including seabirds, salmon, lingcod and marine mammals. Larger piscivorous fishes, marine mammals, and in recent years jumbo squid are among the predators of larger adults.

### **Stock Status and Management History**

Chilipepper have been one of the most important commercial target species in California waters since the 1880s and were historically an important recreational target in Southern California waters. With the exception of excluding foreign fishing effort from the U.S. EEZ in the late 1970s, management actions were modest (and usually general to all rockfish and other groundfish) prior to the implementation of the Groundfish FMP in 1982. When the FMP was implemented, management for the groundfish trawl fishery was based on individual vessel trip limits, which were set at 40,000 lbs per trip on the *Sebastes* (all rockfish species) complex. These limits were maintained until 1991, when they were reduced to 25,000; in 1993 the trip limit system was revised from daily to biweekly trip limits, which were set at 50,000 lbs (south of Cape Mendocino). The trip limit regime continued to evolve in its absolute amounts and temporal duration (monthly, bimonthly) throughout the 1990s, with a general trend towards lower limits as conservation concerns arose for other rockfish species (particularly bocaccio rockfish in the region south of Mendocino). The chilipepper catch in the bottom trawl fishery has been managed under an IFQ system since 2011.

Chilipepper rockfish were assessed in 1998 (Ralston, *et al.* 1998), at which time the stock south of 40°10' N lat. was estimated to be at 46 percent to 61 percent of unfished biomass.

A full chilipepper assessment was conducted in 2007 (Field 2008). The 2007 assessment estimated a substantial increase in the spawning biomass of chilipepper rockfish in recent years, due to a strong 1999 year class as well as greatly reduced harvest rates in commercial and recreational fisheries. The 2007 assessment's base model result suggests a spawning biomass of 23,889 tons in 2006, corresponding to approximately 70 percent of the unfished spawning biomass of 33,390 tons and representing a near tripling of spawning biomass from the estimated low of 8,696 mt (26 percent of unfished) in 1999. The strong 1999 year class represents the largest estimated historical recruitment, and is the primary cause for the current population trajectory. There are no obvious signs of strong year classes since 1999, and coastwide pelagic juvenile surveys suggested average to low recruitment in years immediately preceding the assessment, suggesting that the stock may dip slightly in the near term.

The 2007 assessment was first used in 2008 to decide 2009 and 2010 chilipepper harvest specifications. The Council consideration for 2011 and 2012 was whether or not to remove chilipepper rockfish from the Shelf Rockfish North complex and manage it coastwide. Chilipepper rockfish are predominantly found south of 40°10' N lat. Prior to 2007 they were only assessed in the area south of 40°10' N lat. To date, chilipepper rockfish have been managed with stock-specific harvest specifications south of 40°10' N lat. and within the Shelf Rockfish North complex north of 40°10' N lat. When the stock assessment area was extended for the 2007 chilipepper stock assessment, it was extended to the stock's entire west coast range through waters off Oregon (chilipepper rockfish are not believed to occur in waters off Washington). However, it was decided to continue to manage chilipepper rockfish south of 40°10' N lat. with stock-specific harvest specifications and as part of the Shelf Rockfish complex north of 40°10' N lat.

An update of the 2007 assessment of chilipepper rockfish south of 40°10' N lat. is planned for 2015.

## Stock Productivity

Steepness in the 2007 assessment was fixed at 0.57, which was the mean of the prior probability distribution in the base model. Since steepness was thought to be poorly specified in the model, this parameter was chosen as the major axis of uncertainty. The decision table projected outcomes for a low productivity and a high productivity model using steepness values of 0.34 and 0.81, respectively. The PSA productivity score of 1.83 indicates a stock of relatively high productivity, especially for a rockfish.

There have been strong recruitments estimated for the stock in the late 1960s, early 1970s, and very strong recruitments in 1984 and 1999. The 1999 year class was the biggest recruitment event in the assessment time series, causing spawning biomass to increase substantially in the ten years preceding the assessment.

## Fishing Mortality

Chilipepper rockfish have been one of the most important commercial target species in California since the late 1800s and was also a recreational target in southern California waters. Catches and exploitation rate have declined substantially since the early 1990s. While chilipepper has always been an important target species in California, the exploitation rate has rarely exceeded the  $F_{MSY}$  target of a 50 percent SPR. Exploitation rates declined substantially since the late 1990s with the implementation of more restrictive management measures to rebuild depleted stocks.

Throughout most of the past three decades, domestic landings have ranged between approximately 2,000 and 3,000 mt; however, since 2002 landings have averaged less than 100 mt per year. The highest exploitation rates occurred from the late 1980s through the mid-1990s, when they were above target levels and the stock was approaching its lowest estimated historical levels. From the late 1990s through the present, exploitation rates have been declining significantly down to incidental levels, as a result of management measures implemented to rebuild co-occurring depleted rockfish species (particularly bocaccio, but including canary, widow, cowcod and yelloweye). Discards are assumed to be negligible in the historical period; however, regulatory discards have been substantial in recent years, more than doubling the total catch relative to landings since 2002.

The PSA vulnerability score of 1.35 indicates a low risk of overfishing.

### 1.1.4.8 Dover Sole

#### Distribution and Life History

Dover sole (*Microstomus pacificus*) are distributed from the Navarin Canyon in the northwest Bering Sea and westernmost Aleutian Islands to San Cristobal Bay, Baja California, Mexico (Hagerman 1952; Hart 1988; NOAA 1990). Dover sole are a dominant flatfish on the continental shelf and slope from Washington to Southern California. Adults are demersal and are found from 9 m to 1,450 m, with highest abundance below 200 m to 300 m (Allen and Smith 1988). Adults and juveniles show a high affinity toward soft bottoms of fine sand and mud. Juveniles are often found in deep nearshore waters. Dover sole are considered to be a migratory species. In the summer and fall, mature adults and juveniles can be found in shallow feeding grounds, as shallow as 55 m off British Columbia (Westrheim and Morgan 1963). By late fall, Dover sole begin moving offshore into deep waters (400 m or more) to spawn. Although there is an inshore-offshore seasonal migration, little north-south coastal migration occurs (Westrheim and Morgan 1963).

Spawning occurs from November through April off Oregon and California in waters 80 m to 550 m depth at or near the bottom (Hagerman 1952; Hart 1988; NOAA 1990; Percy, *et al.* 1977). Dover sole are

oviparous and fertilization is external. Larvae are planktonic and are transported to offshore nursery areas by ocean currents and winds for up to two years. Settlement to benthic living occurs mid-autumn to early spring off Oregon, and February through July off California (Markle, *et al.* 1992). Juvenile fish move into deeper water with age and begin seasonal spawning and feeding migrations upon reaching maturity.

Dover sole larvae eat copepods, eggs, and nauplii, as well as other plankton. Juveniles and adults eat polychaetes, bivalves, brittle stars, and small benthic crustaceans. Dover sole feed diurnally by sight and smell (Dark and Wilkins 1994; Gabriel and Pearcy 1981; Hart 1988; NOAA 1990). Dover sole larvae are eaten by pelagic fishes like albacore, jack mackerel and tuna, as well as sea birds. Juveniles and adults are preyed upon by sharks, demersally feeding marine mammals, and to some extent by sablefish (NOAA 1990). Dover sole compete with various eelpout species, rex sole, English sole, and other fishes of the mixed species flatfish assemblage (NOAA 1990).

### **Stock Status and Management History**

Dover sole have been the target of trawl operations along the west coast of North America since World War II and were almost certainly caught prior to the war as incidental take in directed fisheries for English sole and petrale sole. Almost all of the harvests have been taken by groundfish trawl, and in particular as part of the Dover sole, shortspine thornyhead, longspine thornyhead, and sablefish (DTS) trawl fishery. Annual landings from U.S. waters averaged 6,700 mt during the 1960s, 12,800 mt during the 1970s, 18,400 mt during the 1980s, 12,400 mt during the 1990s, and 7,200 mt since 2000.

The 1997 Dover sole stock assessment (Brodziak, *et al.* 1997) treated the entire population from the Monterey area through the U.S.-Vancouver area as a single stock based on research addressing the genetic structure of the population. Under a range of harvest policies and recruitment scenarios, the 1997 model projected that spawning biomass would increase from the estimated year-end level in 1997 through the year 2000 due to growth of the exceptionally large 1991 year class and to the lower catches observed in the fishery since 1991.

Dover sole were next assessed in 2001, resulting in an estimated spawning stock size of 29 percent of the unexploited biomass (Sampson and Wood 2001). The unexploited spawning stock biomass was estimated to be 176,500 mt and the stock steadily declined from the 1950s until the mid-1990s with little subsequent variation. The 1991 year class was the last strong one, consistent with the 1997 assessment.

The 2005 Dover sole assessment indicated the stock was above target levels and had an increasing abundance and biomass trend since the late 1990s (Sampson 2005). The final base model estimated the unexploited spawning stock biomass to be slightly less than 300,000 mt and spawning biomass at the start of 2005 was estimated to be about 189,000 mt, equivalent to 63 percent of the unexploited level. Spawning biomass and age 5+ biomass (roughly corresponding to the exploitable biomass) were estimated to have reached their lowest points in the mid-1990s and rose steadily since. The estimated increases in biomass since the mid-1990s were due primarily to strong year classes in 1990 and 1991, and exceptionally strong year classes in 1997 and 2000.

A new Dover sole assessment was done in 2011, which indicated the stock was healthy with a 2011 spawning stock biomass depletion of 83.7 percent of unfished biomass (Hicks and Wetzel 2011).

### **Stock Productivity**

Steepness in the 2011 Dover sole assessment was fixed at 0.8, the mean steepness estimated in the SSC's 2010 meta-analysis of flatfish productivity (PFMC and NMFS 2011). While the 2011 assessment was considered data-rich, estimates of steepness are uncertain partly because the stock has not been fished to

low levels to understand potential recruitment at low spawning biomass. The PSA productivity score of 1.8 indicates a stock of relatively high productivity.

There is little information regarding recruitment prior to 1960. Estimates of recruitment appear to oscillate between periods of low recruitment and periods of high recruitment. The five largest recruitments were predicted in the years 2000, 1992, 1988, 1965, and 1991. The five smallest recruitments were predicted in 2003, 2002, 2004, 2006, and 1974.

Larger than average recruitments in the early 1960s resulted in an increase in the Dover sole spawning biomass. A period of smaller than average recruitments in the late 1970s and early 1980s, along with the highest catches on record caused a decline in spawning biomass throughout the 1980s. More recently, spawning biomass has been increasing. However, a recent increase in Dover sole catches and low estimated recruitment in the early 2000s seem to be resulting in a slight downward trend in spawning biomass.

### **Fishing Mortality**

The spawning biomass of Dover sole reached a low in the mid-1990s before beginning to increase throughout the last decade. The estimated depletion has remained above the 25 percent biomass target and it is unlikely that the stock has ever fallen below this threshold. Throughout the 1970s, 1980s, and 1990s the exploitation rate and SPR generally increased, but never exceeded the SPR 30 percent  $F_{MSY}$  target. Recent exploitation rates on Dover sole have been much lower than  $F_{MSY}$ , even after management increased catch levels in 2007.

Given the productivity of the stock and constraints on fishing, projections assuming a 25,000 mt constant annual catch predict the stock would remain above the target  $B_{MSY}$  level for the next ten years even under the more pessimistic and less likely low state of nature in the assessment decision table. Higher ACLs than the preferred No Action ACL were initially considered but rejected from more detailed analysis since the current market is projected to limit the take of Dover sole in the next management cycle to less than 25,000 mt. Higher ACLs are predicted to be sustainable; future mortalities as high as the OFL (above the allowable ACL maximum of the ABC) would maintain the stock above the target level of  $B_{25\%}$  under the most likely base case model in the 2011 assessment.

Sablefish quota pounds are needed to target Dover sole and the other DTS species using trawl gear. Sablefish IFQ quota is also used in a single-species target fishery using fixed gears. The competition and price for sablefish quota are affected by Asian sablefish demand and supply from north Pacific fisheries outside the west coast EEZ (e.g., BC and the Gulf of Alaska fisheries). It may be the case that the supply and demand of west coast Dover sole will remain limited until there is an increased harvestable surplus of sablefish above recent levels.

Dover sole are caught primarily by bottom trawls and are managed using IFQs in the rationalized fishery. Despite Dover sole being an important target species, only 35 percent of the 2011 quota was attained in the IFQ fishery.

The PSA vulnerability score of 1.54 indicates a low risk of overfishing.

#### **1.1.4.9 English Sole**

### **Distribution and Life History**

English sole (*Parophrys vetulus*) are found from Nunivak Island in the southeast Bering Sea and Agattu Island in the Aleutian Islands, to San Cristobal Bay, Baja California Sur, Mexico (Allen and Smith 1988).

In research survey data, nearly all occurred at depths greater than 250 m (Allen and Smith 1988). Adults and juveniles prefer soft bottoms composed of fine sands and mud (Ketchen 1956), but also occur in eelgrass habitats (Pearson and Owen 1992). English sole use nearshore coastal and estuarine waters as nursery areas (Krygier and Percy 1986; Rogers, *et al.* 1988). Adults make limited migrations. Those off Washington show a northward post-spawning migration in the spring on their way to summer feeding grounds and a southerly movement in the fall (Garrison and Miller 1982). Tagging studies have identified separate stocks based on this species' limited movements and meristic characteristics (Jow 1969).

Spawning occurs over soft-bottom mud substrates (Ketchen 1956) from winter to early spring, depending on the stock. Eggs are neritic and buoyant, but sink just before hatching (Hart 1988); juveniles and adults are demersal (Garrison and Miller 1982). Small juveniles settle in the estuarine and shallow nearshore areas all along the coast, but are less common in southerly areas, particularly south of Point Conception. Large juveniles commonly occur up to depths of 150 m. Although many post larvae may settle outside of estuaries, most will enter estuaries during some part of their first year of life (Gunderson, *et al.* 1990). Some females mature as three-year-olds (26 cm), but all females over 35 cm long are mature. Males mature at two years (21 cm). Females attain much larger sizes than males. Landings by the fishery are composed primarily of female fish, but at-sea discards of small fish include large numbers of male English sole.

Larvae are planktivorous. Juveniles and adults are carnivorous, eating copepods, amphipods, cumaceans, mysids, polychaetes, small bivalves, clam siphons, and other benthic invertebrates (Allen 1982; Becker 1984; Hogue and Carey 1982; Simenstad, *et al.* 1979). English sole feed primarily by day, using sight and smell, and sometimes dig for prey (Allen 1982; Hulberg and Oliver 1979). A juvenile English sole's main predators are piscivorous birds such as great blue heron (*Ardia herodias*), larger fishes, and marine mammals. Adults may be eaten by marine mammals, sharks, and other large fishes.

### **Stock Status and Management History**

English sole have been captured by the bottom trawl fishery operating off the western coast of North America for over a century. Stewart (2006) found that peak catches from the southern area occurred in the 1920s with a maximum of 3,976 mt of English sole landed in 1929, and peak catches from the northern area occurred in the 1940s to the 1960s with a maximum of 4,008 mt landed in 1948. Landings from both areas have generally declined since the mid-1960s and have been at nearly historical lows in recent years

The most recent stock assessment of English sole prior the current 2005 assessment was performed in 1993 (Sampson and Stewart 1993). That assessment considered the female portion of the stock off Oregon and Washington during the years 1977-1993 because the landings were dominated by females (greater than 90% by weight). The English sole spawning biomass was found to be increasing and it was concluded that the fishery was sustainable at (then) contemporary harvest levels.

The 2005 assessment of English sole (Stewart 2006) modeled a single coastwide stock, although both commercial and fishery independent data sources were treated separately for a southern (INPFC Conception and Monterey) and a northern (INPFC Eureka, Columbia and U.S. Vancouver) area. The assessment found that English sole spawning biomass had increased rapidly over the last decade after a period of poor recruitments from the mid-1970s to the mid-1990s, which left the stock at nearly historically low levels. Strong year classes were estimated for 1995, 1996, and 1999. The data indicated that the 1999 year class may be the largest in the time-series. There was substantial uncertainty related to certain parameters in the assessment, specifically biomass, recruitment, and relative depletion, as indicated by the wide confidence intervals for those parameters. Nevertheless, sensitivity analyses indicated that the conclusion that current spawning biomass exceeds the target level ( $B_{40\%}$ ) was robust to all three of these sources of uncertainty. The spawning biomass at the beginning of 2005 was estimated to be 31,379 mt, which corresponds to 91.5

percent of the unexploited equilibrium level. Total catches for 2004 were estimated to be 1,341 mt, of which 950 mt were landed.

The 2007 update assessment (Stewart 2008c) confirmed the magnitude of increased biomass through a large quantity of age data through 2006, which became available. The 2007 assessment also included data on fishery length and age (primarily from Washington) that was previously unavailable. These new data provided substantially improved information regarding recent year class strengths and current stock status. The spawning biomass at the beginning of 2007 was estimated to be 41,906 mt, which corresponded to 116 percent of the unexploited equilibrium level.

Cope et al. (2014) assessed English sole using the data-moderate exSSS model platform. The English sole assessment was conducted for a coastwide stock and stock depletion was estimated to be 88% at the start of 2013 (Table 1-4). The current spawning biomass was estimated to be 25,719 mt and the projected 2015 and 2016 OFLs are 10,792 mt and 7,890 mt, respectively. Since the new English sole assessment was conducting using data-moderate methods, the stock was downgraded from a category 1 to a category 2 stock.

### **Stock Productivity**

There is little evidence for a strong stock-recruitment relationship, with some of the largest recruitments occurring at moderate levels of spawning biomass. This corresponds to the relatively high estimate of steepness of 0.87 in the assessment. In general, recruitment deviations are well-informed by the data between 1940 and 2000.

Following two decades of low recruitments, strong year classes were estimated for 1995, 1998-2000, and 2002. The data indicate that the 1999 year class was the largest in the time-series.

The PSA productivity score of 2.25 indicates a very productive stock, which is true for most nearshore and shelf flatfishes.

### **Fishing Mortality**

The estimated SPR for English sole has never been below the proxy target of 30 percent for flatfish. Exploitation rates were highest from the late 1940s to the early 1990s. Since 1992, the intensity of exploitation has been substantially less, resulting in higher SPR levels. This corresponds to a relative exploitation rate (catch/biomass of age 3 and older fish) history that is high from the late 1940s to the early 1990s, and steadily declining to very low levels over the last 15 years.

English sole are primarily caught by groundfish bottom trawls. Management uncertainty is low with the 100 percent observer coverage for the groundfish trawl fleet under trawl rationalization. Very small amounts of English sole were landed in the 2011 IFQ fishery with only 1 percent of the quota attained. This is due to low trawl effort on the shelf since such efforts require investment of limited quota for Pacific halibut, darkblotched rockfish, and yelloweye rockfish.

The PSA vulnerability score of 1.19 shows a very low concern of overfishing on the stock.

#### 1.1.4.10 Lingcod North and South of 40°10' N Lat.

##### **Distribution and Life History**

Lingcod (*Ophiodon elongatus*), a top order predator of the family *Hexagrammidae*, ranges from Baja California, Mexico, to Kodiak Island in the Gulf of Alaska. Lingcod are demersal at all life stages (Allen and Smith 1988; NOAA 1990; Shaw and Hassler 1989). Adult lingcod prefer two main habitat types: slopes of submerged banks 10 m to 70 m below the surface with seaweed, kelp, and eelgrass beds and channels with swift currents that flow around rocky reefs (Emmett, *et al.* 1991; Giorgi and Congleton 1984; NOAA 1990; Shaw and Hassler 1989). Juveniles prefer sandy substrates in estuaries and shallow subtidal zones (Emmett, *et al.* 1991; Hart 1988; NOAA 1990). As the juveniles grow they move to deeper waters. Adult lingcod are considered a relatively sedentary species, but there are reports of migrations of greater than 100 km by sexually immature fish (Jagiello 1990; Mathews and LaRiviere 1987; Matthews 1992; Smith, *et al.* 1990).

Mature females live in deeper water than males and move from deep water to shallow water in the winter to spawn (Forrester 1969; Hart 1988; Jagiello 1990; LaRiviere, *et al.* 1980; Mathews and LaRiviere 1987; Matthews 1992; Smith, *et al.* 1990). Mature males may live their whole lives associated with a single rock reef, possibly out of fidelity to a prime spawning or feeding area (Allen and Smith 1988; LaRiviere, *et al.* 1980; Shaw and Hassler 1989). Spawning generally occurs over rocky reefs in areas of swift current (Adams 1986; Adams and Hardwick 1992; Giorgi and Congleton 1984; LaRiviere, *et al.* 1980). After the females leave the spawning grounds, the males remain in nearshore areas to guard the nests until the eggs hatch. Hatching occurs in April off Washington, but as early as January and as late as June at the geographic extremes of the lingcod range. Males begin maturing at about two years (50 cm), whereas females mature at three plus years (76 cm). In the northern extent of their range, fish mature at an older age and larger size (Emmett, *et al.* 1991; Adams, 1992 #438; Hart 1988; Mathews and LaRiviere 1987; Miller and Geibel 1973; Shaw and Hassler 1989). The maximum age for lingcod is about 20 years (Adams and Hardwick 1992).

Lingcod are a visual predator, feeding primarily by day. Larvae are zooplanktivores (NOAA 1990). Small demersal juveniles prey upon copepods, shrimps, and other small crustaceans. Larger juveniles shift to clupeids and other small fishes (Emmett, *et al.* 1991; NOAA 1990). Adults feed primarily on demersal fishes (including smaller lingcod), squids, octopi, and crabs (Hart 1988; Miller and Geibel 1973; Shaw and Hassler 1989). Lingcod eggs are eaten by gastropods, crabs, echinoderms, spiny dogfish, and cabezon. Juveniles and adults are eaten by marine mammals, sharks, and larger lingcod (Miller and Geibel 1973; NOAA 1990).

##### **Stock Status and Management History**

Lingcod have been a target of commercial fisheries since the early 1900s in California, and since the late 1930s in Oregon and Washington waters. Recreational fishermen have targeted lingcod since the 1920s in California. A smaller recreational fishery has taken place in Washington and Oregon since at least the 1970s. Although historically the catches of lingcod have been greater in the commercial sector than in the recreational sector, this pattern has been reversed since the late 1990s.

In 1997, Jagiello, *et al.* (1997) assessed the size and condition of the portion of the stock in the Columbia and Vancouver areas (including the Canadian portion of the Vancouver management area), and concluded the stock had fallen to below ten percent of its unfished size at 8.8 percent of its unfished biomass. The Council responded by imposing substantial harvest reductions coastwide, reducing the harvest targets for the Eureka, Monterey, and Conception areas by the same percentage as in the north.

In 1999, Adams, et al. (1999) assessed the southern portion of the stock and concluded the condition of the southern stock was similar to the northern stock with a depletion of  $B_{15\%}$ , thus confirming the Council had taken appropriate action to reduce harvest coastwide. Based on these assessments, the lingcod stock was declared overfished in 1999. A rebuilding plan establishing a target year of 2009 and harvest rates of  $F = 0.0531$  and  $F = 0.0610$  for fisheries in the northern and southern areas, respectively was adopted and implemented in 2000.

Jagiello et al. (2000) conducted a coastwide lingcod assessment and determined the total biomass increased from 6,500 mt in the mid-1990s to about 8,900 mt in 2000. In the south, the population had also increased slightly from 5,600 mt in 1998 to 6,200 mt in 2000. In addition, the assessment concluded previous aging methods portrayed an older population; whereas new aging efforts showed the stock to be younger and more productive. Therefore, the ABC and OY were increased in 2001 on the basis of the new assessment. A revised rebuilding analysis of coastwide lingcod (Jagiello and Hastie 2001) confirmed the major conclusions of the 2000 assessment and rebuilding analysis, but slightly modified recruitment projections to stay on the rebuilding trajectory to reach target biomass in 2009.

The lingcod rebuilding plan was formally adopted by the Council and incorporated into the FMP under Amendment 16-2. The rebuilding plan established a target rebuilding year of 2009 and the harvest control rule of  $F = 0.0531$  for fisheries in the northern areas and  $F = 0.0610$  for fisheries in the southern areas (with a  $P_{MAX}$  of 60 percent). Depth-based restrictions and a winter season fishing closure to protect nest-guarding males were also implemented as part of the rebuilding plan.

Jagiello et al. (2004) conducted a coastwide assessment for lingcod in 2003 that indicated the lingcod stock had achieved the rebuilding objective of  $B_{40\%}$  in the north with a 68 percent depletion, but was at a 31 percent depletion in the south. The Council's SSC, working in concert with the lead assessment author, recalculated the coastwide lingcod stock status in March 2004 using actual 2003 harvests (the assessment, which was completed during 2003, assumed harvest would be equal to the specified OY in 2003). Their calculations indicated that the spawning biomass at the start of 2004 was within 99.3 percent of  $B_{MSY}$  ( $B_{40\%}$ ) on a coastwide basis. The harvest control rule was recalculated to be  $F = 0.17$  for fisheries in the northern areas and  $F = 0.15$  for fisheries in the southern areas.

The 2005 coastwide assessment (Jagiello and Wallace 2006) again modeled two populations of lingcod north and south of  $40^{\circ}10'$  N. lat. On a coastwide basis, the lingcod population was concluded to be fully rebuilt, with the spawning biomass in 2005 estimated to be 64 percent of its unfished level. Within the separate area models current biomass was estimated to be closer to unfished biomass in the north ( $B_{87\%}$ ) than in the south ( $B_{24\%}$ ). Given that the lingcod stock is managed on a coastwide basis, the Council announced the lingcod stock to be fully rebuilt in 2005, which is four years earlier than the target rebuilding year established in the rebuilding plan.

The 2009 lingcod assessment modeled two populations north and south of the California-Oregon border at  $42^{\circ}$  N. lat. (Hamel, *et al.* 2009). Both populations were healthy with stock depletion estimated at 62 and 74 percent for the north and south, respectively.

The Council and NMFS elected to maintain the management line for lingcod at  $40^{\circ}10'$  N lat. by specifying separate ACLs north and south of that line. This action was intended to not overly encumber the commercial fishing industry, which is required to fish within a single management area within one trip. Specifying the lingcod management line at  $42^{\circ}$  N lat. would create two management areas stratified at  $40^{\circ}10'$  N lat. and  $42^{\circ}$  N lat. This would especially burden vessels home ported out of Brookings, Crescent City, Eureka, and Ft. Bragg, since they would have to restructure their current fishing practices to avoid a violation of the management line crossover provisions. It is stated in the 2009 assessment that a management break at Cape Mendocino would be likely more biologically accurate than stratifying the



assessment north and south of 42° N lat. In general, given the crossover provisions and the other regulations that foster area management strategies, the fewer latitudinal management lines there are, the less burdened the offshore commercial fishery will be. Two major biogeographic breaks occur on the west coast at Pt. Conception at 34°27' N lat. and Cape Mendocino approximately at 40°10' N lat., and many stocks show differences north and south of these latitudes. These biogeographic breaks are probably the more appropriate latitudes to specify management lines, given how north-south physical processes such as current patterns tend to be different, creating stock differences for species affected by these different physical processes.

The lingcod STAT evaluated the swept area biomass estimates calculated annually (2003-2010) from the NMFS NWFSC trawl survey, which indicated that 48 percent of the lingcod biomass for the stock south of 42° N lat. occurred between 40°10' N lat. and 42° N lat. Therefore, 48 percent of the 2013 and 2014 OFLs projected in the 2009 lingcod assessment for the southern lingcod stock were added to OFLs proposed for the stock north of 40°10' N lat. Likewise, 48 percent of the projected OFLs for the southern stock were subtracted from the OFLs proposed for the stock south of 40°10' N lat. Given that the trawl survey is the main fishery-independent tuning index of biomass in the assessment, using swept area biomass from the trawl survey to estimate relative biomass north and south of 40°10' N lat. was considered appropriate.

### **Stock Productivity**

Steepness was fixed at 0.8 in the 2009 assessment. The PSA productivity score of 1.75 indicates a stock of relatively high productivity.

Recruitments in the north were estimated from 1928-2007, with bias correction ramping in from 1950 to 1964 as data became informative. The base model indicated a very strong recruitment event in 1964, a secondary event in 1970, and recent relatively strong recruitments in 1999-2002, with fairly high recruitment in 2006 as well. Recruitments in the south were estimated from 1928-2007, with bias correction ramping in from 1960 to 1974 as data became informative. The base model indicated relatively strong recruitment events in 1976, 1983, and 1999-2003, similar to the period of increased recruitment in the north, with a very high but uncertain recruitment in 2007.

### **Fishing Mortality**

Lingcod exploitation coastwide was above the target rate for most of the 1970s through the 1990s, driving the stock below the MSST and into an overfished condition. The stock was successfully rebuilt by 2006 based on good recruitments and very low fishing mortality rates. The SPR for northern lingcod has been above the proxy target of 45 percent since 1998, and in recent years has been far above that level. The SPR for the southern lingcod stock has been above the proxy target of 45 percent since 2001, and in recent years has been far above that level.

The PSA vulnerability score for lingcod is 1.55, indicating a low risk of overfishing of the stock.

#### **1.1.4.11 Longnose Skate**

### **Distribution and Life History**

Skates are the largest and most widely distributed group of batoid fish with approximately 245 species ascribed to two families (Ebert and Compagno 2007; McEachran 1990). Skates are benthic fish that are found in all coastal waters but are most common in cold temperatures and polar waters (Ebert and Compagno 2007).

There are about eleven species of skates from either of three genera (*Amblyraja*, *Bathyraja*, and *Raja*) present in the Northeast Pacific Ocean off California, Oregon and Washington (Ebert 2003). Of that number, just three species (longnose skate *Raja rhina*, big skate *Raja binoculata*, and sandpaper skate *Bathyraja interrupta*) make up over 95 percent of survey catches in terms of biomass and numbers, with the longnose skate leading in both categories (62% of biomass and 56% of numbers). Species compositions of fishery landings also show that longnose skate are the predominant skates in commercial catches. On average, longnose skate represents 75 percent of total skate landings in Oregon for the last 20 years and 45 percent in Washington for the last 10 years. There are no species composition data available for commercial landings in California, but anecdotal evidence suggests that the majority of skates landed there are longnose skates.

The distribution of the longnose skate is limited to the eastern Pacific Ocean. It is found from the southeastern Bering Sea to just below Punta San Juanico, southern Baja California, and Gulf of California at depths of 9-1,069 m (Love, *et al.* 2005). Longnose skates do not exhibit a size-specific pattern in distribution relative to bottom depth; average fish size does not vary greatly with depth.

Currently, there is no information available that indicates the existence of multiple breeding units in the Northeast Pacific Ocean. Several tagging studies have found that elasmobranchs, such as sharks and skates, can undertake extensive migrations within their geographic range (Martin and Zorzi 1993; McFarlane and King 2003). This behavior suggests the likelihood that there is a high degree of genetic mixing within the population, across its range. As a result, the longnose skate population off California, Oregon and Washington is modeled in this assessment as a single stock.

The life history of skates is characterized by late maturity, low fecundity and slow growth to large body size (King and McFarlane 2003; Moyle and Cech 1996; Walker and Hislop 1998). Skates invest considerable energy in developing a few large, well-protected embryos. These characteristics are associated with a K-type reproductive strategy, as opposed to r-type strategy, wherein reproductive success is achieved by high productivity and early maturity (Hoenig and Gruber 1990).

The longnose skate is oviparous. After fertilization, the female forms tough, but permeable egg cases that surround eggs and then deposits these egg cases onto the sea floor at daily to weekly intervals for a period of several months or longer (Hamlett and Koob 1999). The eggs within egg cases incubate for several months in a benthic habitat. Inside the egg cases, the embryos develop with nourishment provided by yolk. The longnose skate is known to have only a single embryo per egg case (David Ebert, Moss Landing Marine Laboratories, pers. com. as cited by Gertseva and Schirripa (2008)). When the yolk is depleted and the juvenile is fully formed, it exits in the egg case. Once hatched, the young skate is similar in appearance to an adult, but smaller in size. Upon reaching maturity, skates enter the reproductive stage, which lasts for the remainder of their lives (Frisk, *et al.* 2002; Pratt and Casey 1990). On average off the continental US Pacific Coast, female longnose skates mature between 11-18 years, which corresponds to 75-125 cm in total length (Thompson 2006). The life span of the longnose skate is not well known, although individuals up to 23 years of age have been found (Thompson 2006). Longnose skates attain a maximum length of about 145 cm, although individuals as large as 180 cm have been reported off the U.S. west coast (Thompson 2006).

The reproductive cycle of oviparous skates has been observed for a few species but not for longnose skate. These studies indicate that egg production generally occurs throughout the year although there have been some instances where seasonality in egg laying was observed (Hamlett and Koob 1999). Information on fecundity of longnose skate is extremely limited. Holden (1974) found that species of the family *Rajidae* are the most fecund of all elasmobranchs and can lay 100 egg cases per year, although eggs may not be produced every year. Frisk *et al.* (2002) estimated that annual fecundity for skates similar in size with longnose may be less than 50 eggs per year; however, those eggs exhibit high survival rates due to the large

parental investment. Overall, little is known about breeding frequency, egg survival, hatching success and other early life history characteristics of longnose skate.

### **Stock Status and Management History**

Longnose skate was managed in a complex of dissimilar species, the Other Fish complex, from 1982, when the Groundfish FMP was implemented through 2008. In 2009, longnose skate was removed from the Other Fish complex and managed with stock-specific harvest specifications.

Gertseva and Schirripa (2008) assessed the west coast longnose skate stock in 2007. The spawning stock biomass was estimated to be at 66 percent of its unfished biomass at the start of 2007. Based on that assessment, a constant catch strategy (OY = 1,349 mt) was implemented in 2009 based on a 50 percent increase in the average 2004-2006 landings and discard mortality. The constant catch strategy was revised in 2013 by implementing an ACL of 2,000 mt to provide greater access to the stock and to limit disruption of current fisheries. This level of harvest was projected to maintain the population at a healthy level as projected in the 10-year forecast for longnose skate in the 2007 assessment (Gertseva and Schirripa 2008).

The SSC recommended changing the proxy  $F_{MSY}$  rate for longnose skate and other elasmobranchs from an SPR of 45% to an SPR of 50% beginning in 2015. This recommendation, driven primarily by conservation concerns for spiny dogfish (see section 1.1.4.18), was heeded by the Council when they adopted 2015 and 2016 OFLs consistent with this lower harvest rate.

### **Stock Productivity**

Steepness of the stock-recruitment curve was fixed at a value of 0.4 in the 2007 assessment to reflect the K-type reproductive strategy of the longnose skate. Recruitments were deterministically provided using this steepness value and a Beverton-Holt stock-recruitment relationship since the data in the 2007 assessment was not informative of relative year-class strength. In general, elasmobranchs have relatively low productivity given the K-type reproductive strategy of producing few eggs per female with a significant parental energy investment to increase survival of those few eggs (e.g., production of egg cases and relatively large yolk masses).

### **Fishing Mortality**

Historically, skates in general, and longnose skate in particular, have not been high-priced fishery products. They are taken mostly as bycatch in other commercially important fisheries (Bonfil 1994). Although skates are caught in almost all demersal fisheries and areas off the U.S. west coast, the vast majority (almost 97%) are caught with trawl gear.

Landing records indicate that skates have been retained on the U.S. Pacific Coast at least since 1916 (Martin and Zorzi 1993). Little is known about the species composition of west coast skate fisheries, particularly prior to 1990. With few exceptions, longnose skate landings have been reported, along with other skate species, under the market category “unspecified skates”, until 2009 when a sorting requirement for longnose skate was required.

Historically, only the skinned pectoral fins or “wings” were sold, although a small portion of catch would be marketed in the round (whole). The wings were cut onboard the boat and the remainder discarded. Currently, west coast skates are marketed both whole and as wings. Skates wings are sold fresh or fresh-frozen, as well as dried or salted and dehydrated, for sale predominantly in Asian markets (Bonfil 1994; Martin and Zorzi 1993). It appears that the demand for whole skates did increase greatly during the mid-1990s, as evidenced by the increase in the number of trips where skates were landed. While skates were

encountered predominantly as bycatch previously, landings data from this period reveal greater targeting of skates by some vessels. After a few years, the whole-skate market cooled due to downturns in Asian financial markets (Peter Leipzig, Fishermen's Marketing Association, pers. com. as cited by Gertseva and Schirripa (2008).

Historically, the exploitation rate for the longnose skate has been low. It reached its maximum level of 4.02% in 1981 (Gertseva and Schirripa 2008). An exploitation rate of 1.25% was estimated in 2006.

A vulnerability score of 1.68 indicates a low concern for overfishing the stock.

#### 1.1.4.12 Longspine Thornyhead

##### **Distribution and Life History**

Longspine thornyhead occur from the southern tip of Baja, California, to the Aleutian Islands (Jacobson and Vetter 1996; Orr, *et al.* 1998). There appears to be no distinct geographic breaks in stock abundance along the west coast (Fay 2006; Rogers, *et al.* 1997). Adult longspine thornyhead are bottom dwellers, and inhabit the deep waters of the continental slope throughout their range.

Longspine occur at depths greater between 201 and 1,756 m, most typically between 500 and 1,300 m (Love, *et al.* 2002), and a peak in abundance and spawning biomass in the oxygen minimum zone (OMZ) at about 1,000 m depth (Jacobson and Vetter 1996; Wakefield 1990). Longspine are better adapted to deep water than shortspine (Siebenaller 1978; Siebenaller and Somero 1982). Wakefield (Wakefield 1990) estimated that in Central California, 83% of the longspine population resides within an area of the continental slope bounded by 600 and 1,000 m depth.

Unlike shortspine thornyhead, the mean size of longspines is similar throughout the depth range of the species (Jacobson and Vetter 1996). Camera sled observations indicate that longspines do not school or aggregate, and are distributed relatively evenly over soft sediments (Wakefield 1990). Differences in density of individuals at depth do occur with latitude, with higher densities of longspine in deep water (1,000-1,400 m) off Oregon than off central California (Jacobson and Vetter 1996).

The strong relationship between depth and size found in shortspine thornyhead (Jacobson and Vetter 1996) is not observed for longspines, with the distribution of longspines being relatively uniform with depth (Rogers, *et al.* 1997). Unlike shortspines, longspine do not undergo an ontogenetic migration to deeper waters (Wakefield 1990).

Longspine thornyheads prefer muddy or soft sand bottoms in deep-water environments characterized by high pressure and low oxygen concentrations. These are low productivity (Vetter and Lynn 1997) and low diversity (Haigh and Schnute 2003) habitats where food availability is limited. Longspines have adapted to this environment with an extremely slow metabolism that allows it to wait up to 180 days between feedings (Vetter and Lynn 1997). They are not territorial, and do not school. They have no swim bladders; instead oil in the bones and spines provides floatation. Video observations from submersibles and ROVs indicate that thornyhead are sit-and-wait predators that rest on the bottom and remain motionless for extended periods (John Butler, NOAA Fisheries, Southwest Fisheries Science Center, CA, as cited in Jacobson and Vetter (1996)).

The spawning season for longspine thornyheads appears to be extended, and occurs over several months during February, March and April (Best 1964; Moser 1974; Pearcy 1962; Wakefield and Smith 1990). Both thornyhead species produce a bi-lobed jellied egg mass that is fertilized at depth and which then floats to the surface where final development and hatching occur (Pearcy 1962). An extended larval and pelagic

juvenile phase follows, which is thought to be 18-20 months long (Jacobson and Vetter 1996; Moser 1974; Wakefield 1990). Juvenile longspine settle on the continental slope at depths between 600 and 1,200 m (Wakefield 1990). Moser (1974) reports a mean length at settlement of 4.2-6.0 cm, although pelagic juveniles up to 69 mm in length have been collected in midwater trawls off Oregon (J. Siebenaller unpublished data, as cited in Wakefield and Smith (1990)).

Following settlement, longspine thornyhead are strictly benthic (Jacobson and Vetter 1996). No apparent pulse in recruitment during the year was observed by Wakefield and Smith (1990), perhaps due to the long (4-5 months) spawning season, variation in growth rates, and variation in the duration of the pelagic period (Wakefield and Smith 1990). There is potential for cannibalism because juveniles settle directly on to the adult habitat (Jacobson and Vetter 1996).

Adult females release between 20,000 and 450,000 eggs over a 4-5 month period (Best 1964; Moser 1974). Wakefield (1990) and Cooper et al. (2005) both found linear relationships between fecundity and somatic weight. The data analyzed by Cooper et al. (2005) indicated that fecundity of longspine between 20 and 30 cm in length ranged from 20,000 to 50,000 eggs.

There is considerable uncertainty regarding age and growth of thornyheads (Jacobson and Vetter 1996), although data indicate that longspine thornyhead are long lived. Age estimates of over 40 years have been obtained from otoliths using thin-section and break- and-burn techniques (Ianelli, *et al.* 1994). High frequencies of large longspine thornyheads may be due to a strongly asymptotic growth pattern, with accumulation of many age groups in the largest size-classes (Jacobson and Vetter 1996).

Size-at-age data (Ianelli, *et al.* 1994) indicate that longspine grow to a maximum size of about 30 cm TL at ages of about 25-45 years, with little or no sexual dimorphism in length at age – longspines in British Columbia, Canada also display no sexual dimorphism (Starr and Haigh 2000). Orr et al. (1998) report a maximum length for longspines of 38 cm, although individuals of this size are rare in both trawl surveys and commercial landings. Growth increments on otoliths suggest that juveniles reach 80 mm after 1 year of life as demersal juveniles (Wakefield unpublished data, as cited in (Wakefield and Smith 1990)), which would correspond to an age of 2.5 - 3 years old.

Longspine thornyhead are ambush predators (Jacobson and Vetter 1996). They consume fish fragments, crustaceans, bivalves, and polychaetes and occupy a tertiary consumer level in the food web. Pelagic juveniles prey largely on herbivorous euphausiids and occupy a secondary consumer level in the food web (Love 1996; Smith and Brown 1983). Sablefish and shortspine thornyhead commonly prey on longspine thornyhead (Buckley, *et al.* 1999).

### **Stock Status and Management History**

Longspine thornyhead are exploited in the limited entry deep-water trawl fishery operating on the continental slope that also targets shortspine thornyhead, Dover sole and sablefish (i.e., the DTS fishery). A very small proportion of longspine landings is due to non-trawl gears (gillnets, hook and line). Longspine and shortspine thornyhead make up a single market category; however, they have been managed under separate harvest specifications since 1992. Beginning in 2011, trawl catches of longspine north of 34°27' N lat. have been managed using individual fishing quotas.

The thornyhead fishery developed in Northern California during the 1960s. The fishery then expanded north and south, and the majority of the landings of longspine thornyhead have since been in the Monterey, Eureka, and Columbia INPFC areas, with some increase in landings from the Conception (southern CA) and Vancouver (northern WA) INPFC areas in recent years (Fay 2006).

The most recent stock assessment of west coast longspine thornyhead was done in 2013. This was the fifth assessment done for longspines, but only the second in which it was assessed individually (earlier assessments were of longspine and shortspine thornyheads in combination). Previous assessments were conducted by Jacobson (Jacobson 1990; 1991), Ianelli et al. (1994), Rogers et al. (1997), and Fay (2006). The 1990 and 1991 assessments were very similar. Important features included reviews of available biological data, and analyses of trends in mean lengths from port samples and catch rates calculated from logbook data. Swept-area and video biomass estimates were used to estimate average biomass levels and exploitation rates in the Monterey to US-Vancouver management areas. The available data were used to conduct per-recruit analyses of yield, revenue, and spawning biomass, and to develop estimates of the then target level of  $F_{35\%}$ .

Ianelli et al. (1994) assessed the coastwide abundance of longspine and shortspine thornyheads based on slope survey data, an updated analysis of the logbook data, and fishery length-composition data to estimate the parameters of length-based Stock Synthesis models, under different assumptions regarding discarding practices.

The Rogers et al. (1997) assessment used a length-based version of Stock Synthesis 1 to fit an age-structured model to data for the Monterey, Eureka, Columbia and Vancouver INPFC areas. Models were fitted to biomass estimates and length data from the AFSC slope surveys (1988-1996), a logbook CPUE index, discarded proportions by year, and length composition data from California and Oregon. Sensitivity to discard rates based on changes in prices and minimum size were explored.

The 2005 assessment of longspine thornyhead estimated spawning biomass in 2005 was approximately 71 percent of unfished spawning biomass (Fay 2006). The model assumed one coastwide stock with one coastwide trawl fishery. Results from the base model suggested that the length compositions from the slope surveys were influencing recruitment in the model, such that the model estimated slightly higher recruitment in the early 1990s, which then declined in the mid to late 1990s.

The 2013 longspine thornyhead assessment indicated a stock depletion of 75 percent at the start of 2013 (Stephens and Taylor 2013).

### **Stock Productivity**

Stephens and Taylor (2013) estimated annual longspine recruitment using a Beverton-Holt stock-recruitment function and assuming a steepness value of 0.6. Most 2013 rockfish assessments used a steepness prior of 0.779, estimated from a meta-analysis of rockfish assessment results. This value might be expected in the 2013 longspine assessment; however, rockfish ecology and reproduction are quite different from those of thornyheads, which (for example) do not give birth to live young but rather spawn floating egg masses.

Steepness in the shortspine thornyhead assessment was fixed at 0.6 both in the 2005 and 2013 models (Hamel 2006c; Taylor and Stephens 2013). This value was justified based on consistency between the modeling approach and management targets, in addition to being within a range of biologically reasonable values. For consistency, therefore, steepness for the longspine model was also fixed at 0.6.

Annual deviations about this stock-recruitment curve were estimated for the years 1944 through 2012. Estimated recruitments do not show high variability, and the uncertainty in each estimate is greater than the variability between estimates. The 2013 longspine assessment is relatively uninformative of relative year class strength since ages were not used in the model (thornyheads are notoriously difficult to age). Therefore, a length-based assessment with an assumed steepness is used to determine recruitment.

## **Fishing Mortality**

The estimated exploitation rate of longspine thornyheads was above the current  $F_{MSY}$  harvest rate through much of the 1990s and, in hindsight, given the current target harvest rate, overfishing was occurring. However, stock biomass was estimated to have never dropped below the target  $B_{MSY}$  level. There is very little risk of overexploitation of longspines given their deep distribution beyond the 700 fm limit to west coast bottom trawling implemented under Amendment 19.

The PSA vulnerability score of 1.54 for longspine thornyheads also indicates a low concern for potential overfishing of the stock.

### **1.1.4.13 Pacific Cod**

#### **Distribution and Life History**

Pacific cod (*Gadus macrocephalus*) are widely distributed in the coastal north Pacific, from the Bering Sea to Southern California in the east, and to the Sea of Japan in the west. Adult Pacific cod occur as deep as 875 m (Allen and Smith 1988), but the vast majority occurs between 50 m and 300 m (Allen and Smith 1988; Love 1996; NOAA 1990). Along the west coast, Pacific cod prefer shallow, soft-bottom habitats in marine and estuarine environments (Garrison and Miller 1982), although adults have been found associated with coarse sand and gravel substrates (Garrison and Miller 1982; Palsson 1990). Larvae and small juveniles are pelagic; large juveniles and adults are parademersal (Dunn and Matarese 1987)(NOAA 1990). Adult Pacific cod are not considered to be a migratory species. There is, however, a seasonal bathymetric movement from deep spawning areas of the outer shelf and upper slope in fall and winter to shallow middle-upper shelf feeding grounds in the spring (Dunn and Matarese 1987).

Pacific cod have external fertilization (Hart 1988; NOAA 1990) with spawning occurring from late fall to early spring. Their eggs are demersal. Larvae may be transported to nursery areas by tidal currents (Garrison and Miller 1982). Half of females are mature by three years (55 cm) and half of males are mature by two years (45 cm) (Dunn and Matarese 1987). Juveniles and adults are carnivorous and feed at night (Allen and Smith 1988; Palsson 1990) with the main part of the adult Pacific cod diet being whatever prey species is most abundant (Kihara and Shimada 1988; Klovach, *et al.* 1995). Larval feeding is poorly understood. Pelagic fish and sea birds eat Pacific cod larvae, while juveniles are eaten by larger demersal fish, including Pacific cod. Adults are preyed upon by toothed whales, Pacific halibut, salmon shark, and larger Pacific cod (Hart 1988; Love 1996; NOAA 1990; Palsson 1990). The closest competitor of the Pacific cod for resources is the sablefish (Allen 1982).

#### **Stock Status and Management History**

The west coast population of Pacific cod has never been formally assessed. Targetable amounts of Pacific cod occur off northern Washington infrequently since the west coast EEZ is at the southern limit of their distribution. The Pacific cod OFL has been set at the highest annual historical catch observed for the stock and ACLs/OYs have been set at half that amount.

Pacific cod is the only unassessed, data-poor groundfish stock currently managed with stock-specific harvest specifications on the west coast.

## Stock Productivity

The PSA productivity score of 2.11 indicates a relatively high productivity and the vulnerability score of 1.34 for Pacific cod indicates a low concern for potential overfishing of the stock.

## Fishing Mortality

Pacific cod occur periodically in targetable amounts off northern Washington. In some years they are targeted because the abundance of this fringe population (in the context of the species' distribution off the west coast) is large enough to be targeted and in some years they are not available. The annual total mortality of Pacific cod has ranged from 39 mt (2008) to 1,415 mt (2004) during 2002-2012. The ACL of 1,600 mt has never been exceeded.

### 1.1.4.14 Pacific Whiting

#### Distribution and Life History

Pacific whiting (*Merluccius productus*), also referred to as Pacific hake, is a semi-pelagic schooling species distributed along the west coast of North America generally ranging from 25° N lat. to 55° N lat. It is among 18 species of hake from four genera (being the majority of the family *Merluccidae*), which are found in both hemispheres of the Atlantic and Pacific oceans (Alheit and Pitcher 1995; Lloris, *et al.* 2005). The coastal stock of Pacific whiting is currently the most abundant groundfish population in the California Current system. Smaller populations of this species occur in the major inlets of the Northeast Pacific Ocean, including the Strait of Georgia, Puget Sound, and the Gulf of California. Genetic studies indicate that the Strait of Georgia and the Puget Sound populations are genetically distinct from the coastal population (Iwamoto, *et al.* 2004; King, *et al.* 2012). Genetic differences have also been found between the coastal population and hake off the west coast of Baja California (Vrooman and Paloma 1977). The coastal stock is also distinguished from the inshore populations by larger body size and seasonal migratory behavior.

The coastal stock of Pacific whiting typically ranges from the waters off southern California to northern British Columbia and in some years to southern Alaska, with the northern boundary related to fluctuations in annual migration. In spring, adult Pacific whiting migrate onshore and northward to feed along the continental shelf and slope from northern California to Vancouver Island. In summer, Pacific whiting often form extensive mid-water aggregations in association with the continental shelf break, with highest densities located over bottom depths of 200–300 m (Dorn and Methot 1991; Dorn and Methot 1992).

Older Pacific whiting exhibit the greatest northward migration each season, with two- and three-year old fish rarely observed in Canadian waters north of southern Vancouver Island. During El Niño events (warm ocean conditions, such as occurred in 1998), a larger proportion of the stock migrates into Canadian waters, apparently due to intensified northward transport during the period of active migration (Agostini, *et al.* 2006; Dorn 1995). In contrast, La Niña conditions (colder water, such as occurred in 2001) result in a southward shift in the stock's distribution, with a much smaller proportion of the population found in Canadian waters, as seen in the 2001 survey.

Spawning occurs from December through March, peaking in late January (Smith 1995). Pacific whiting are oviparous with external fertilization. Eggs of the Pacific whiting are neritic and float to neutral buoyancy (Bailey 1982; Bailey, *et al.* 1982; NOAA 1990). Hatching occurs in five days to six days, and within three months to four months juveniles are typically 35 mm (Hollowed 1992). Juveniles move to deeper water as they get older (NOAA 1990). Females mature at three years to four years (34 cm to 40 cm) and nearly all males are mature by three years (28 cm). Females grow more rapidly than males after four years; growth ceases for both sexes at 10 years to 13 years (Bailey, *et al.* 1982).



All life stages feed near the surface late at night and early in the morning (Sumida and Moser 1984). Larvae eat calanoid copepods, as well as their eggs and nauplii (McFarlane and Beamish 1986; Sumida and Moser 1984). Juveniles and small adults feed chiefly on euphausiids (NOAA 1990). Large adults also eat amphipods, squid, herring, smelt, crabs, and sometimes juvenile whiting (Bailey 1982; Dark and Wilkins 1994; McFarlane and Beamish 1986). Eggs and larvae of Pacific whiting are eaten by pollock, herring, invertebrates, and sometimes Pacific whiting. Juveniles are eaten by lingcod, Pacific cod, and rockfish species. Adults are preyed on by sablefish, albacore, pollock, Pacific cod, marine mammals, soupfin sharks, and spiny dogfish (Fiscus 1979; McFarlane and Beamish 1986).

### **Stock Status and Management History**

The history of the coastal whiting fishery is characterized by rapid changes brought about by the development of foreign fisheries in 1966, joint-venture fisheries in the early 1980s, and domestic fisheries in 1990s. Since implementation of the Magnuson-Stevens Fishery Conservation and Management Act in the U.S. and the declaration of a 200 mile fishery conservation zone in the U.S. and Canada in the late 1970s, annual quotas (or catch targets) have been used to limit the catch of Pacific hake in both zones by foreign and domestic fisheries. Scientists from both countries historically collaborated through the Technical Subcommittee of the Canada-U.S. Groundfish Committee (TSC), and there were informal agreements on the adoption of annual fishing policies. During the 1990s, however, disagreements between the U.S. and Canada on the allotment of the catch limits between U.S. and Canadian fisheries led to quota overruns; 1991-1992 quotas summed to 128% of the limit, while the 1993-1999 combined quotas were 107% of the limit on average. In 2003, a bilateral Pacific whiting management agreement was signed by both countries that created formal allocations of the harvestable surplus, as well as an international process for assessing and managing the stock. This international process was fully implemented in 2012.

Pacific whiting is managed consistent with the Agreement with Canada on Pacific Hake/Whiting. Annual catch limits, now called TACs (total allowable catches), for Pacific whiting are adopted on an annual basis after a stock assessment is completed by a Joint Technical Committee (JTC) and reviewed in February by an international Scientific Review Group (SRG). In March the JTC and SRG present the assessment to the Joint Management Committee (JMC), the international decision-making body. The JMC presents their TAC recommendations to their respective government officials before these TACs are implemented in regulations. The coastwide TAC for the U.S. West Coast and Canada is allocated 26.12% to Canada and 73.88% to the U.S. under Article III (2) of the Agreement.

The most recent Pacific whiting assessment estimated female spawning biomass to be 1.72 million mt, with a depletion ratio of 81.8% of unfished equilibrium levels at the start of 2014 (Taylor, *et al.* 2014). The base model estimates indicate that since the 1960s, Pacific Hake female spawning biomass has ranged from well below to near the unfished equilibrium biomass. The model estimates that the stock was below the unfished equilibrium in the 1960s and 1970s, increased toward the unfished equilibrium after two or more large recruitments occurred in the early 1980s, and then declined steadily through the 1990s to a low in 2000. This long period of decline was followed by a brief peak in 2003 as the large 1999 year class matured and subsequently supported the fishery for several years. Estimated female spawning biomass declined to an all-time low of 0.479 million mt in 2009 because of low recruitment between 2000 and 2007, along with a declining 1999 year class. Spawning biomass estimates have increased since 2009 on the strength of a large 2010 cohort and above average 2008 and 2009 cohorts.

## Stock Productivity


Pacific whiting have high relative productivity as evidenced by fast growth, a high natural mortality rate ( $M$ ), and high steepness in the Beverton-Holt stock-recruitment function. The prior for steepness in the 2014 Pacific whiting assessment is based on the median (0.79), 20th (0.67) and 80th (0.87) percentiles from the Myers et al. (1999) meta-analysis of the family *Gadidae*, and has been used in previous U.S. assessments since 2007.

Pacific whiting exhibit low average recruitment with occasional large year-classes. Very large year classes in 1980, 1984, and 1999 supported much of the commercial catch from the 1980s to the early 2000s. In the last decade, estimated recruitment has been at some of the lowest values in the time series (1966-present) as well as some of the highest. The 2014 assessment estimates a strong 2010 year class, second only to the 1980 year class in the time series, comprising 67% of the 2013 commercial catch.

The PSA productivity score for Pacific whiting ( $P = 2.00$ ) is relatively high and the low vulnerability score ( $V = 1.69$ ) indicates a low concern for potential overfishing.

## Fishing Mortality

Taylor et al. (Taylor, *et al.* 2014) estimated fishing intensity on the stock was consistently below the  $F_{40\%}$  target until recently when the target was likely exceeded in 2008, 2010 and 2011 (Figure 1-23). The exploitation fraction does not necessarily correspond to fishing intensity because fishing intensity accounts for the age-structure: for example, fishing intensity remained nearly constant and above target from 2010 to 2011 but exploitation fraction declined in these years because of high estimated abundances of 1 year old fish (Figure 1-24).



**Figure 1-23. Trend in median fishing intensity (relative to the SPR management target) of Pacific whiting through 2013 with 95% posterior credibility intervals. The management target defined in the Agreement is shown as a horizontal line at 1.0.**




Figure 1-24. Trend in median exploitation fraction through 2013 with 95% posterior credibility intervals.

#### 1.1.4.15 Sablefish

##### Distribution and Life History

Sablefish, or black cod, (*Anoplopoma fimbria*) are distributed in the northeastern Pacific ocean from the southern tip of Baja California, northward to the north-central Bering Sea and in the Northwestern Pacific ocean from Kamchatka, southward to the northeastern coast of Japan. Although few studies have critically evaluated issues regarding the stock structure of this species, it appears there may exist at least three different stocks of sablefish along the west coast of North America: (1) a stock that exhibits relatively slow growth and small maximum size that is found south of Monterey Bay (Cailliet, *et al.* 1988; Phillips and Inamura 1954); (2) a stock that is characterized by moderately fast growth and large maximum size that occurs from northern California to Washington; and (3) a stock that grows very quickly and contains individuals that reach the largest maximum size of all sablefish in the northeastern Pacific ocean, distributed off British Columbia, Canada and in the Gulf of Alaska (Mason, *et al.* 1983; McFarlane and Beamish 1983a). Large adults are uncommon south of Point Conception (Hart 1988; Love 1996; McFarlane and Beamish 1983b; NOAA 1990). Adults are found as deep as 1,900 m, but are most abundant between 200 m and 1,000 m (Beamish and McFarlane 1988; Kendall and Matarese 1987; Mason, *et al.* 1983). Off southern California, sablefish are abundant to depths of 1,500 m (MBC 1987). Adults and large juveniles commonly occur over sand and mud (McFarlane and Beamish 1983a; NOAA 1990) in deep marine waters. They were also reported on hard-packed mud and clay bottoms in the vicinity of submarine canyons (MBC 1987).

Spawning occurs annually in the late fall through winter in waters greater than 300 m (Hart 1988; NOAA 1990). Sablefish are oviparous with external fertilization (NOAA 1990). Eggs hatch in about 15 days (Mason, *et al.* 1983; NOAA 1990) and are demersal until the yolk sac is absorbed (Mason, *et al.* 1983). Age-zero juveniles become pelagic after the yolk sac is absorbed. Older juveniles and adults are benthopelagic. Larvae and small juveniles move inshore after spawning and may rear for up to four years (Boehlert and Yoklavich 1985; Mason, *et al.* 1983). Older juveniles and adults inhabit progressively deeper

waters. Estimates indicate that 50 percent of females are mature at five years to six years (24 inches) and 50 percent of males are mature at five years (20 inches).

Sablefish larvae prey on copepods and copepod nauplii. Pelagic juveniles feed on small fishes and cephalopods—mainly squids (Hart 1988; Mason, *et al.* 1983). Demersal juveniles eat small demersal fishes, amphipods, and krill (NOAA 1990). Adult sablefish feed on fishes like rockfishes and octopus (Hart 1988; McFarlane and Beamish 1983a). Larvae and pelagic juvenile sablefish are heavily preyed upon by seabirds and pelagic fishes. Juveniles are eaten by Pacific cod, Pacific halibut, lingcod, spiny dogfish, and marine mammals, such as Orca whales (Cailliet, *et al.* 1988; Hart 1988; Love 1996; Mason, *et al.* 1983; NOAA 1990). Sablefish compete with many other co-occurring species for food, mainly Pacific cod and spiny dogfish (Allen 1982).

### **Stock Status and Management History**

Formal stock assessments of sablefish began in 1984. The first coastwide assessment established regulations on the sablefish fishery off the U.S. Pacific coast which were implemented as trip limits in October 1982. Since 1982, the sablefish fishery has been managed intensively, with limited entry and open access programs used in various manners to limit catches.

In 2001, two assessments were completed and reviewed by a STAR Panel: one by NMFS (Schirripa and Methot 2001) and one by the Pacific Groundfish Conservation Trust (Hilborn, *et al.* 2001). The two assessments were in agreement, and the Council adopted the NMFS assessment for management purposes. Schirripa and Methot (2001) focused on evaluating the sensitivity of the model and the outcomes to changes in the survey data. These changes included the combining of the AFSC slope survey data and the NWFSC Industry Co-operative Survey data using a statistical Generalized Linear Models (GLM) procedure. This analysis made it possible to extend the southern boundary of the assessment south to Point Conception at 34°27' N lat. rather than 36° N lat., used in previous assessments. The assessment indicated a normal decline in biomass since the late 1970s due to the fishing down of the unfished stock and an unexpected decline in recruitment during the early 1990s. It introduced for the first time, the possibility that sablefish recruitment may be linked to environmental factors. A seemingly meaningful relationship was demonstrated between changes in northern and southern copepod abundances and sablefish recruitment. Conditions and projections in the model considered two competing “states of nature” to calculate the mean virgin recruitment: a “density-dependent” state that used the average of 1975-1991 recruitments, and a “regime shift” state that used the 1975-2000 recruitments. To account for this uncertainty, the Council adopted a 2002 ABC based on the proxy harvest rate ( $F_{45\%}$ ) adjusted to reflect the distribution north and south of 36° N latitude. This was done because a plan amendment would be needed to change the management area since Groundfish FMP Amendment 14 specified only the area north of 36° N lat.

The Council also wanted to verify industry reports of a large abundance of juvenile sablefish, an observation that was confirmed to some extent by preliminary results from the 2001 NMFS slope survey. Based on these considerations, the Council recommended a new expedited assessment be done in 2002. This update assessment (Schirripa 2002), by definition, sought to document changes in the estimates of the status of the stock by only considering newly available data for 2001 while not considering any new changes in the model structure or model assumptions. The expedited assessment confirmed fishermen’s anecdotal reports of a large 1999 year class, which was also apparent in the preliminary results of the 2001 slope survey.

The 2005 sablefish assessment estimated stock depletion at 34.3 percent of unfished biomass (Schirripa and Colbert 2006). The assessment fit a relationship between sea level and recruitment deviations for the period 1973-2003 and used that relationship to hindcast recruitment variability back to 1925. The 2005 assessment found that spawning stock biomass had steadily declined since 1900 and suggested that there was little evidence that recruitment from 2001-2005 was as high as that for the strong 1999 and 2000 year classes.

As a result, the assessment's biomass projections indicate a short-term increase, followed by a continued decline.

The 2007 updated sablefish assessment estimated spawning depletion to be 38.3 percent of unfished biomass at the start of 2007 (Schirripa 2008). This increase from 2005 was attributed in part to the continued recruitment of the strong 1999 and 2000 year classes into the spawning stock biomass. The assessment also estimated a series of poor recruitments in the mid- to late-1990s, and if fished at the full OY level, depletion was forecasted to decrease for the next five years.

The 2011 sablefish assessment estimated spawning stock biomass to be at 33 percent of its unfished biomass at the beginning of 2011 (Stewart, *et al.* 2011). The resource was modeled as a single stock; however, there is some dispersal to and from offshore seamounts and along the coastal waters of the continental U.S., Canada, Alaska, and across the Aleutian Islands to the western Pacific which was not explicitly accounted for in this analysis. Environmental time-series including both sea-surface height (used in previous sablefish assessments) and zooplankton abundance were also investigated. These environmental indices were not used in the 2011 assessment in the interest of parsimony since they did not affect results.

An update of the 2011 sablefish assessment is planned for 2015.

### **Stock Productivity**

It was not possible to estimate the steepness parameter of the Beverton-Holt stock-recruitment relationship in the 2011 sablefish assessment, so this quantity was fixed at a value of 0.6 and explored via sensitivity analyses.

Stewart et al. (2011) estimated sablefish were exploited at a modest level through the first half of the 20th century. Following a period of above-average recruitments, the spawning stock biomass increased to nearly unexploited levels. Large harvests in the 1970s and 1980s were believed to have caused the stock biomass to decline. Estimates of the stock's productivity were highly uncertain due to lack of information on mortality, absolute stock size, and productivity. Sablefish recruitment was estimated to be variable over the historical record, with substantial uncertainty in individual recruitment events. Recruitments during the 1980s were, on average, roughly an order of magnitude higher than the very poor recent cohorts estimated between 2002 and 2007.

### **Fishing Mortality**

The sablefish fishery has been managed with a rich history of seasons, size-limits, trip-limits, and a complex permit system. Coastwide yield targets have been divided among the different gears (hook-and-line, pot and trawl), fishery sectors (including both limited entry and open access) as well as north and south of 36° N lat. Peak catches occurred in the late 1970s just prior to the imposition of the first catch limits. Since 2001, the total estimated dead catch has been only 79 percent of the sum of the OFLs (ABCs at the time) and 87 percent of the ACLs (OYs at the time). In only one year of the last 10, 2008, did the estimated dead catch exceed the ACL (and OFL) by 5% (3%).

The PSA vulnerability score of 1.64 indicates a relatively low concern for potential overfishing.

#### 1.1.4.16 Shortbelly Rockfish

##### **Distribution and Life History**

Shortbelly rockfish (*Sebastes jordani*) range from Punta Baja in Baja California (Klingbeil 1976) as far north as La Perouse Bank off of British Columbia, and as far west as the Cobb seamount off the southern Washington coast (Pearson, *et al.* 1993). However, they are most abundant along the continental shelf break between the northern end of Monterey Bay and Point Reyes, California (particularly in the regions of Ascension Canyon and the Farallon Islands), and around the Channel Islands in the Southern California Bight (Love, *et al.* 2002; Moser, *et al.* 2000; Pearson, *et al.* 1991; Phillips 1964). Although stock structure is poorly understood, genetic analysis of fish collected between San Diego and Cape Mendocino suggests a single coastwide stock, with slight differences in allele frequencies across Point Conception (Constable 2006). The shortbelly rockfish is one of the most abundant rockfish species in the California Current and is a key forage species for many piscivorous fish, birds, and marine mammals.

Shortbelly rockfish feed primarily on juvenile and adult euphausiids, and are an important prey item to a wide range of piscivorous fishes, seabirds and marine mammals (Chess, *et al.* 1988; Lowry and Carretta 1999; Sydeman, *et al.* 2001). Merkel (1957) reported that juvenile shortbelly rockfish were important prey of Chinook salmon along the central California coast in late spring and summer, accounting for more than 60% of those identified to species. For many breeding California seabirds, as much as 90% of their diet is comprised of pelagic stages of juvenile (age 0) rockfish during the late spring and early summer breeding seasons, and unexploited species (such as shortbelly) generally account for more than two thirds of the juvenile rockfish identified (Ainley, *et al.* 1993; Miller and Sydeman 2004; Sydeman, *et al.* 2001). However there is considerable interannual and interdecadal variability in the frequency of rockfish in seabird diets. Throughout the 1990s, foraging rates on juvenile rockfish by central California seabirds declined for both exploited and unexploited rockfish species primarily in response to changes in ocean conditions associated with poor recruitment for rockfish (Miller and Sydeman 2004; Mills, *et al.* 2007; Sydeman, *et al.* 2001). Although rockfish have rarely been identified to the species level in the diets of many California Current marine mammals (Antonelis and Fiscus 1980; Morejohn, *et al.* 1978; Perez and Bigg 1986; Stroud, *et al.* 1981), shortbelly were among the five most significant prey items for California sea lion (*Zalophus californianus*) in the Channel Islands (Lowry and Carretta 1999) and are frequently encountered in sea lion food habits samples off of Central California (Weise and Harvey 2005). Shortbelly rockfish are also described as important prey to thresher sharks (Preti, *et al.* 2004), longnose skate (Robinson, *et al.* 2007), and jumbo squid (Field, *et al.* 2007), among others. Consequently, shortbelly rockfish are an important forage species to a wide range of predators throughout the California Current ecosystem, and generally have a trophic position and life history traits more similar to forage fishes than most other *Sebastes*.

##### **Stock Status and Management History**

The expectation of eventual development of a domestic commercial fishery (Kato 1981) led to past efforts to estimate stock abundance and productivity (Lenarz 1980, Pearson *et al.* 1989, Pearson *et al.* 1991) as well as evaluations of commercial potential. The first ABC for shortbelly rockfish was set by the Council at 10,000 mt for 1983 through 1989. A stock assessment by Pearson *et al.* (Pearson, *et al.* 1991) estimated that allowable catches for shortbelly might range from 13,900 to 47,000 mt per year, based on life history data and hydroacoustic survey estimates of abundance. Subsequently, the Council established an ABC of 23,500 mt, which was reduced to 13,900 mt in 2001 based on observations of poor recruitment throughout the 1990s and the continued lack of a targeted fishery. Yet despite several attempts to develop a commercial fishery for shortbelly, domestic fishery landings have never exceeded 80 mt per year along the west coast.

A shortbelly rockfish assessment was done as an academic exercise in 2007 to understand the potential environmental determinants of fluctuations in the recruitment and abundance of an unexploited rockfish population in the California Current ecosystem (Field, *et al.* 2008). The results of the assessment indicated the shortbelly stock was healthy with an estimated spawning stock biomass of 67 percent of its unfished biomass in 2005.

Shortbelly rockfish is an abundant species that is not targeted in any commercial or recreational fisheries or caught in substantial amounts. However, shortbelly rockfish is a valuable forage fish species in the California Current ecosystem with fluctuations in stock recruitment and biomass driven by environmental conditions. The consequence of fisheries, including high and low estimates of plausible discards, were estimated to be negligible ( $P < 0.01$ ) in all years with the exception of the foreign fisheries of the mid-1960s (Field, *et al.* 2008). Shortbelly rockfish were initially considered for an Ecosystem Component (EC) species categorization under Amendment 23. Rather than classifying shortbelly rockfish as an EC species, the Council chose to recommend a very restrictive ACL of 50 mt for the 2011-2012 and the 20-13-2014 management cycles. The ACL is proposed to be increased to 500 mt beginning in 2015 to prevent unavoidable bycatch from prematurely shutting down emerging midwater trawl fisheries targeting yellowtail and widow rockfish. The 500 mt ACL is still less than 10% of the ABC and is a level of harvest meant to accommodate unavoidable incidental bycatch of shortbelly rockfish while allowing most of the harvestable surplus of the stock to be available as forage for species in the California Current ecosystem. Such ecological considerations are made when setting ACLs for west coast groundfish species.

### **Stock Productivity**

Field *et al.* (Field, *et al.* 2008) assumed a steepness of 0.65 in a Mace-Doonan stock-recruitment relationship (Mace and Doonan 1988) in the 2007 shortbelly assessment. The data in the assessment model were insufficient for estimating steepness; therefore, an assumed value was used based on the Dorn (2002b) meta-analysis of rockfish steepness available at the time the assessment was conducted.

Recruitment deviations of shortbelly from 1960-2005 were estimated in the 2007 assessment; however, there was greater confidence in relative year class strength from 1975-2005. The model suggested a long period of poor recruitment through most of the 1990s, associated with a significant decline in biomass. The interesting conclusion of the 2007 shortbelly assessment was how apparent environmental determinants of shortbelly recruitment and not fishing mortality affected biomass and stock status.

### **Fishing Mortality**

Shortbelly rockfish are not targeted in any west coast fisheries and are incidentally caught in very small amounts. Love *et al.* (2002) reported that shortbelly rockfish were commonly caught incidentally with trawl gear in the San Francisco-Monterey region during the development of the trawl fishery in the 1930s and 1940s when they were often referred to as steamer rockcod, as they tended to be common in the steamer lanes south of San Francisco. However, as a result of the small size and poor marketability, only modest domestic landings (1 to 65 mt per year) have been reported in the last 25 years. Historical landings were almost certainly less. Phillips (1939) reported that *S. jordani* accounted for 1 lb. out of 332,630 lbs examined in Monterey wholesale fish markets between 1937 and 1938. Nitsos (1965) reported trace amounts (approximately 1,000 lbs out of 1,920,000 lbs landed) of *S. jordani* landed in Monterey ports from trawlers in 1962-1963, but none were reported from ports other than Monterey. There was historically a short period in which large numbers of shortbelly were caught during the foreign fisheries of the 1960s and 1970s (Rogers 2003b). These landings (nearly 15,000 mt through 1976, over half of which was taken in 1966) were presumably incidental to the targeting of other rockfish and Pacific hake. Only in the early days of the foreign fisheries (the mid-1960s) were Pacific hake pursued in large numbers south of Cape Mendocino, which is when the bulk of documented historical landings of shortbelly occurred. Since the

early 1970s the Pacific hake fishery has been prosecuted primarily off of Oregon and Washington, and to a lesser extent off of Northern California (generally north of Cape Mendocino).

The available data for historical bycatch rates of shortbelly rockfish are extremely sparse. Shortbelly have been caught incidentally, at times in large numbers, by trawlers targeting other semi-pelagic rockfish (usually chilipepper and widow rockfish). As large hauls of shortbelly are not marketable but occasionally foul the mesh of typical groundfish trawls, more experienced fishermen generally recognize shortbelly sign (as well as habitat preferences) on their acoustics, and work to actively avoid schools. Bycatch monitoring programs conducted north of Cape Mendocino in the mid-1980s suggested very negligible levels of bycatch, such that shortbelly were less than 0.25% of total catches in all fishing strategies (which included nearshore flatfish, bottom rockfish, midwater rockfish and whiting, shrimp and the deep water complex), including less than 0.05% for midwater trawl whiting and rockfish (Pikitch 1988). Very little contemporary information is available for the region south of Mendocino. However, all of these data were collected far north of the usual range of shortbelly. Data processed from the West Coast Groundfish Observer Program suggests that approximately one mt of shortbelly rockfish were caught and discarded in trawl fisheries south of Mendocino. As regulatory measures have closed the vast majority of habitat optimal to adult shortbelly, such trace landings are to be expected in recent years, and comparable data prior to these closures does not exist.

Field et al. (2007) acknowledged the uncertain historical estimates of shortbelly bycatch and therefore explored higher and lower bycatch streams in the 2007 assessment. Varying the historical catch assumptions in the assessment did not result in meaningful deviations from the base model results; therefore, they concluded it was unlikely fishing mortality had any substantive impact on the stock since the days of the foreign fisheries.

#### 1.1.4.17 Shortspine Thornyhead

##### **Distribution and Life History**

Shortspine thornyhead (*Sebastolobus alascanus*) are found in the waters off of the West Coast of the United States from northern Baja California to the Bering Sea. They are found from 20 to over 1,500 m in depth. The majority of the spawning biomass occurs in the oxygen minimum zone between 600 and 1,400 m, where longspine thornyheads are most abundant (Bradburn, *et al.* 2011; Jacobson and Vetter 1996). The distribution of the smallest shortspine thornyheads suggests that they tend to settle at around 100–400 m and are believed to have ontogenetic migration down the slope, although large individuals are found across the depth range.

Shortspine thornyhead do not appear to be distributed evenly across the west coast, with higher densities of thornyheads in shallower areas (under 500 m) off of Oregon and Washington, and higher densities in deeper areas off of California. The mean latitude of the largest shortspine is slightly further north than of the medium sizes, suggesting the possibility of either a J-shaped migration, differential patterns of recruitment, or regional differences in exploitation history.

Although their densities vary, shortspine thornyheads are present in almost all trawlable areas below 500 m. They are caught in 91% of the trawl survey hauls below 500 m and 94% of the commercial bottom trawl hauls below 500 m. In camera tows, thornyheads are seen to be spaced randomly across the sea floor (Wakefield 1990), indicating a lack both of schooling and territoriality.

Genetic studies of stock structure do not suggest separate stocks along the west coast. Siebenaller (1978) and Stepien (1995) found few genetic differences among shortspine thornyheads along the Pacific coast. Stepien (1995), however, did suggest that there may be a separate population of shortspine thornyhead in



the isolated area around Cortes Bank off San Diego, California. Stepien (1995) also suggested that juvenile dispersion might be limited in the area where the Alaska and California currents split. This occurs towards the northern boundary of the assessment area, near 48° N lat.

Stepien et al. (2000), using a more discerning genetic material (mtDNA), found evidence of a pattern of genetic divergence corresponding to geographic distance. However, this study, which included samples collected from southern California to Alaska, did not identify a clear difference between stocks even at the extremes of the range. No such pattern was seen in longspine thornyhead, which suggests that the shorter pelagic stage (~1 yr vs. ~2 yrs) of shortspine may contribute to an increased genetic separation with distance.

Shortspine thornyheads along the west coast spawn pelagic, gelatinous masses between December and May (Erickson and Pikitch 1993; Pearson and Gunderson 2003; Wakefield 1990). Juveniles settle at around 1 year of age (22- 27 mm in length), likely in the range of 100-200 m (Vetter and Lynn 1997), and migrate down the slope with age and size, although large individuals are found across the depth range.

Shortspine thornyhead grow very slowly, but may continue growing throughout their lives, reaching maximum lengths of over 70 cm. Females appear to reach larger sizes than do males. Maturity in females has been estimated as occurring near 18 cm, at 8-10 years of age (Pearson and Gunderson 2003), although new information suggests that patterns of maturity may be more complex.

Shortspine and longspine thornyheads have historically been caught with each other and with Dover sole and sablefish, making up the DTS fishery. Other groundfish species that frequently co-occur in these deep waters include a complex of slope rockfishes, rex sole, longnose skate, rougtail skate, Pacific grenadier, giant grenadier, Pacific flatnose as well as non-groundfish species such as Pacific hagfish and a diverse complex of eelpouts. Shortspine thornyheads typically occur in shallower water than the shallowest longspine thornyheads, and migrate to deeper water as they age. When shortspines have reached a depth where they overlap with longspines, they are typically larger than the largest longspines. Shortspine thornyhead stomachs have been found to include longspine thornyheads, suggesting a predator-prey linkage between the two species.

Thornyheads spawn gelatinous masses of eggs which float to the surface. This may represent a significant portion of the upward movement of organic carbon from the deep ocean (Wakefield 1990). Thornyheads have been observed in towed cameras beyond the 1,280 m limit of the current fishery and survey, but their distribution, abundance, and ecosystem interactions in these deep waters are relatively unknown.

### **Stock Status and Management History**

Beginning in 1989, both thornyhead species were managed as part of the deep water complex with sablefish and Dover sole (DTS). In 1991, the Council first adopted separate ABC levels for thornyheads and catch limits were imposed on the thornyhead group. Harvest guidelines (HGs) were instituted in 1992 along with an increase in the minimum mesh size for bottom trawl fisheries. In 1995 separate landing limits were placed on shortspine and longspine thornyheads and trip limits became more restrictive. Trip limits (predominantly 2-month limits on cumulative vessel landings) have often been adjusted during the year since 1995 in order to not exceed the HG or OY for that year. At first, the HG for shortspine thornyhead was set higher than the ABC (1,500 vs. 1,000 mt in 1995-1997) in order to allow a greater catch of longspine thornyhead, which was considered a relatively underutilized and healthy stock. In 1999 the OY was set at less than 1,000 mt and remained close to that level through 2006. As a result of the 2005 shortspine assessment, catch limits increased to about 2,000 mt per year and have remained near that level to the present.

Since early 2011, trawl harvest of each thornyhead species has been managed under the PFMC's catch share, or individual fishing quota (IFQ), program. Whereas the trip limits previously used to limit harvest restricted only the amount of fish each vessel could land, individual vessels fishing under the catch-share program are now held accountable for all of the quota-share species they catch.

Ianelli et al. (Ianelli, *et al.* 1994) assessed the coastwide abundance of longspine and shortspine thornyheads based on slope survey data, an updated analysis of the logbook data, and fishery length-composition data to estimate the parameters of length-based Stock Synthesis models, under different assumptions regarding discarding practices.

The assessment of thornyheads in 1997 covered the area from Central California at 36° N lat. to the U.S.-Canada border (Rogers, *et al.* 1997). The STAR Panel expressed concern that management requires more detailed information on thornyheads than could be obtained from the available data. In 1998, two separate stock assessments covering the area north of 36° N lat. were prepared and accepted by the Council (NMFS and OT 1998; Rogers, *et al.* 1998). A synthesis of these two assessments was used to set the harvest specifications 1999 and 2000. Given that the synthesis estimated 1999 depletion at 32 percent of virgin biomass, the Council used the precautionary 40-10 policy to set the OYs for those two years.

There were a range of uncertainties in the 2001 assessment of shortspine thornyhead, in 2001, not the least of which was the estimated biomass (Piner and Methot 2001). The assessment was extended south to Point Conception (in contrast to past surveys, which were limited to stocks north of the 36° N latitude management area boundary). The authors concluded the 2001 spawning biomass ranged between 25 percent and 50 percent of unexploited spawning biomass. As was also the case in the 1998 assessment, the uncertainty in abundance largely revolved around the uncertainty in recruitment and survey  $q$ , or catchability, of shortspine thornyhead in slope surveys. The authors also concluded that the trend in stock biomass was increasing and the stock was not depleted. Based on estimated biomass and application of the GMT-recommended  $F=0.75M$  principle (which approximated an  $F_{50\%}$  proxy harvest rate for shortspine thornyhead), the assessment authors and GMT recommended a slight increase in the ABC and OY for 2002. They also recommended that the harvest specifications be set for two areas divided by Point Conception at 34°27' N lat., rather than the previous policy to separate the management areas at the Conception-Monterey border (36° N lat.). Despite the uncertainty in biomass estimates and determination of whether shortspine thornyhead should be treated as a "precautionary zone" stock, these recommendations did treat the stock as such by applying the 40-10 adjustment.

The 2005 stock assessment estimated the shortspine thornyhead spawning stock biomass to be at 62.9 percent of its initial, unfished biomass in 2005 (Hamel 2006c). The 2005 assessment extended the southern border of the assessment area from Point Conception to the Mexican border (32.5° N latitude). Including the entire Conception area resulted in a larger basis for unfished biomass, given that this area was estimated to contain nearly half of the stock's total west coast biomass. It was noted that there could be regional management concerns with this stock because while the assessment OY was coastwide, there are differences in historic exploitation rates north and south of Point Conception. It was also noted the biomass estimate south of Pt. Conception was more uncertain than that in the north.

The 2013 stock assessment estimated the shortspine thornyhead spawning stock biomass to be at 74.2 percent of its initial, unfished biomass in 2013 (Taylor and Stephens 2013). A longer time series of the coastwide NWFSC trawl survey biomass estimates were included in this assessment relative to the 2005 assessment. Therefore, the STAT concluded there was no greater uncertainty in the biomass south of Pt. Conception relative to estimates for the rest of the coast. As in the previous assessment, no age data were used in the 2013 assessment and growth parameters were fixed at the same values used in 2005.

## Stock Productivity

Taylor and Stephens (2013) estimated annual shortspine recruitment using a Beverton-Holt stock-recruitment function and assuming a steepness value of 0.6. Most 2013 rockfish assessments used a steepness prior of 0.779, estimated from a meta-analysis of rockfish assessment results. This value might be expected in the 2013 longspine assessment; however, rockfish ecology and reproduction are quite different from those of thornyheads, which (for example) do not give birth to live young but rather spawn floating egg masses.

Steepness in the shortspine thornyhead assessment was fixed at 0.6 both in the 2005 and 2013 models (Hamel 2006c; Taylor and Stephens 2013). This value was justified based on consistency between the modeling approach and management targets, in addition to being within a range of biologically reasonable values.

Annual deviations about this stock-recruitment curve were estimated for the years 1944 through 2012. Estimated recruitments do not show high variability, and the uncertainty in each estimate is greater than the variability between estimates. The 2013 shortspine assessment is relatively uninformative of relative year class strength since ages were not used in the model (thornyheads are notoriously difficult to age). Therefore, a length-based assessment with an assumed steepness is used to determine recruitment.

## Fishing Mortality

Landings of shortspine were estimated to have risen to a peak of 4,815 mt in 1989, followed by a sharp decline during a period of trip limits and other management measures imposed in the 1990s. Since the institution of separate trip limits for shortspine and longspine thornyheads, the fishery had more moderate removals of between 1,000 and 2,000 mt per year from 1995 through 1998. Landings fell below 1,000 mt per year from 1999 through 2006, then rose to 1,531 in 2009 and have declined since that time.

Exploitation rates in terms of spawning potential ratio indicates that the exploitation slightly exceeded the  $F_{MSY}$  target for a single year in 1985 and then for the period 1989-1994. However, the stock status is estimated to have never fallen below the  $B_{40\%}$  management target.

### 1.1.4.18 Spiny Dogfish

#### Distribution and Life History

In the Northeast Pacific, spiny dogfish (*Squalus suckleyi*) occur from the Gulf of Alaska, with isolated individuals found in the Bering Sea, southward to San Martin Island, in southern Baja California. They are extremely abundant in waters off British Columbia and Washington, but decline in abundance southward along the Oregon and California coasts (Ebert 2003; Ebert, *et al.* 2010).

The U.S. west coast spiny dogfish stock likely has interaction and overlap with dogfish observed off British Columbia. About 1,300 dogfish were tagged along the coast of Washington from 1942-1946, during the period of the strong directed fishery for dogfish. Only 50 of these fish were recaptured and had tags returned (4%), of which 54% were recaptured within U.S. coastal waters, while 32% were recaptured in coastal Canada and 12% in the inside waters of Puget Sound and the Strait of Georgia. One fish was recaptured in coastal Japanese waters (7 years after being tagged). Because many of the releases were close to the U.S.-Canada border, and the fractions do not take into account the relative fishing pressure within each area, this study is of limited use in providing reliable information about dogfish movement rates.

A spatial population dynamics model (Taylor 2008), which included these tagging data (along with much larger tagging experiments conducted in Canada and inside U.S. waters of Puget Sound) estimated movement rates of about 5% per year between the U.S. coastal sub-population of dogfish and that found along the west coast of Vancouver Island in Canada. The model also estimated movement rates of less than 1% per year between the U.S. coastal sub-population of dogfish and that in the Puget Sound.

These sharks appear to prefer areas in which the water temperature ranges from 5 to 15° C, often making latitudinal and depth migrations to follow this optimal temperature gradient (Brodeur, *et al.* 2009). There is also evidence of seasonal movement along the coast based on both tagging data and timing of historical fisheries (Ketchen 1986). One estimate of the seasonal movement along the Pacific coast is a North-South shift of about 600 km from winter to summer (Taylor 2008). This seasonal pattern is not as extreme as that found among spiny dogfish in Atlantic waters of the U.S., which are likely due to larger fluctuations in temperature. Dogfish have also been captured in high-seas salmon gillnets across the North Pacific between about 40° and 50° N lat. (Nakano and Nagasawa 1996), but the extent of these wide-ranging pelagic movements is poorly understood.

The biology and life history of spiny dogfish are relatively well studied (Campana, *et al.* 2009; Di Giacomo, *et al.* 2009; Taylor 2008; Tribuzio 2009; Tribuzio, *et al.* 2009; Tribuzio, *et al.* 2010; Vega, *et al.* 2009). This species is an opportunistic feeder that consumes a wide range of prey (whatever is abundant). Schooling pelagic fish, such as herring, make up the majority of its diet. They also feed on invertebrates such as shrimp, crab and squid. In turn, dogfish are preyed upon by larger cod, hake, and other spiny dogfish (Beamish, *et al.* 1992; Brodeur, *et al.* 2009; Tanasichuk, *et al.* 1991). Larger species of sharks, as well as seals and killer whales, also feed on dogfish.

Spiny dogfish have internal fertilization and ovoviviparous development. The internal development takes place over 22-24 months, the longest gestation period known for sharks. The number of pups in each litter ranges between 5 and 15 individuals depending on the size of the female (larger females bearing more pups). The size at birth is generally between 20 and 30 cm for both genders. Male spiny dogfish are reported to grow faster than females, but females reach larger sizes. This species is the latest maturing (with 50% female maturity reported at 35.5 years) and longest lived of all elasmobranchs (Cortes 2002; Saunders and McFarlane 1993; Smith, *et al.* 1998; Taylor 2008). Life history traits of spiny dogfish make the species highly susceptible to overfishing and slow to recover from stock depletion since its slow growth, late maturation, and low fecundity are directly related to recruitment and spawning stock biomass (Holden 1974; King and McFarlane 2003).

### **Stock Status and Management History**

Spiny dogfish on the U.S. west coast have been utilized for almost a thousand years, with those in Puget Sound first used by Native Americans (Bargmann 2009). The exploitation of spiny dogfish in coastal waters, however, started in the 20th century. Even though the history of spiny dogfish utilization on the U.S. west coast included a brief but intense commercial fishery in the 1940s, in general this species is not highly prized and is mostly taken as bycatch in other fisheries.

Prior to 1936, coastal catches of spiny dogfish were extremely minimal, but in 1936, shortly after it was discovered that livers of spiny dogfish have high level of vitamin A, a large scale fishery for dogfish developed in the Pacific Northwest. Before World War II, Northeast Pacific dogfish livers could not compete with the cheaper and more potent sources of vitamin A from Europe. But when World War II started and European supplies were cut, dogfish shark livers became the major source of vitamin A in the United States, and the spiny dogfish fishery grew rapidly along the Pacific coast. The processed liver oils were used in pharmaceuticals, food processing, and animal feed (Bargmann 2009; Ketchen 1986).

During the liver fishery, dogfish were targeted by three major gear groups, including setlines, set nets, and bottom trawls. The timing of the dogfish liver fishery coincided with the development of bottom trawling in the U.S. Northwest, and though at the onset of the fishery the catches by trawl were low, by the mid-1940s trawling was the dominant type of fishing for dogfish.

In 1945, a sharp decline in spiny dogfish catches began. This decline occurred despite continued strong demand for vitamin A and high prices for dogfish livers, but because of decreased availability of the species in the Northeast Pacific (Bargmann 2009; Ketchen 1986). In 1950, with the advent of synthetic vitamins, demand for spiny dogfish livers declined and catches in the Northeast Pacific virtually ended.

Between 1950 and 1974, the landings of spiny dogfish remained minimal. By the late 1950s it was reported that species availability had increased. Also, in the late 1950s-early 1960s, dogfish earned a bad reputation among fishermen. They were blamed for driving off commercially valuable species such as herring and mackerel, while consuming large numbers of them. Spiny dogfish have also been observed biting through nets to get to their fish prey, releasing many of them and damaging fishing gear in the process. They were also reported damaging gear when become entangled in commercial nets. As a result, fishermen were trying to avoid areas with higher chances of dogfish catches (such as soft bottoms, for example) to prevent encountering dogfish and potentially damaging their gear.

A market opportunity for dogfish developed in the mid-1970s. In Europe, spiny dogfish has long been used an inexpensive source of human food, for fish and chips in particular. A decline in the European dogfish supply provided an opportunity for developing an export dogfish food fishery on the U.S. west coast. Also, during the late 1970s, shark cartilage started to be used in cancer treatment, and a portion of spiny dogfish catches have since been sold for medical research and treatment (Gregory Lippert, WDFW, pers. com. as cited by Gertseva and Taylor (2011)). As before, three types of gear were involved in catching dogfish (bottom trawl, setlines, and sunken gill nets), but since the mid-1980s catches by gillnets have been minimal.

Spiny dogfish is a common bycatch species, often caught in other fisheries and largely discarded. For instance, it has long been incidentally caught in the hake fishery, which is almost exclusively conducted with mid-water trawls. Large-scale harvesting of Pacific hake in the U.S. began in 1966, when factory trawlers from the Soviet Union and other countries began targeting this stock. After the 200-mile U.S. EEZ was declared in 1977, a joint-venture fishery was initiated between U.S. trawlers and Soviet factory trawlers acting as motherships (larger, slower ships for fish processing and storage while at sea). By 1989 the U.S. fleet capacity had grown to a level sufficient to harvest the entire quota, and no further foreign fishing was allowed. The Pacific hake fishery is currently 100% observed at sea and data on bycatch species, including spiny dogfish, is being routinely collected.

Spiny dogfish on the U.S. west coast has been managed under the Other Fish complex since implementation of the Groundfish FMP by the Council in 1982. In 2005, reduction in the Other Fish ABC was implemented due to removal of the California substock of cabezon from the Other Fish complex. The same year, a 50% precautionary OY reduction was implemented to accommodate uncertainty associated with managing unassessed stocks. In 2006, a trip limit for spiny dogfish was imposed for U.S. west coast waters which varied between 45 and 91 mt per two months for all gears. In 2009, another ABC reduction was implemented due to removal of longnose skate from the Other Fish complex and the 50% OY reduction was maintained.

In 2011, reduction in the Other Fish OFL was implemented due to removal of the Oregon substock of cabezon from the Other Fish complex. The 50% precautionary reduction to the ACL was maintained; however, a scientific uncertainty buffer was specified as an ABC of 7,742 mt under the Amendment 23 framework.

Gertseva and Taylor (2011) estimated the spawning stock output of spiny dogfish to be 44,660 thousands of fish (95% confidence interval: 8,937-80,383), which represents 63% of the unfished spawning output level. While this depletion level indicates the stock is currently healthy, fishing at the target SPR of 45% is expected to severely reduce the spawning output over the long term because of the extremely low productivity and other reproductive characteristics of the stock. The Council partially addressed this by setting a more conservative spiny dogfish ABC for 2013 by specifying a  $P^*$  of 0.3.

The Council further decided to manage spiny dogfish with stock-specific harvest specifications beginning in 2015. The SSC also investigated establishing a more conservative  $F_{MSY}$  harvest rate for spiny dogfish and other elasmobranchs in recognition of their lower productivity. The SSC recommended and the Council adopted a more conservative proxy 50% SPR harvest rate as an interim measure for elasmobranchs. The 50% SPR was based on an SSC meta-analysis of *Chondrichthyes* species using the posterior distribution for  $F_{MSY}/M$  values as reported by Zhou et al. (2012). The SSC said they may further investigate sustainable harvest rates for Council-managed elasmobranchs as more information becomes available in the future.

### **Stock Productivity**

Spiny dogfish have a relatively low stock productivity due to slow growth, late maturation, and low fecundity. The fecundity of dogfish in the Northeast Pacific Ocean has been well studied, with pregnant females having relatively few pups per litter (5 to 15), and with relatively little variability among individuals. Unlike fish producing millions of eggs, the low fecundity of dogfish suggests both low productivity in general and a more direct connection between spawning output and recruitment than for many species.

Gertseva and Taylor (2011) modeled the spiny dogfish spawner-recruit relationship using a new functional form that was recently added to the Stock Synthesis platform, which allowed a more explicit modeling of pre-recruit survival between the stage during which embryos can be counted in pregnant females to their recruitment as age 0 dogfish. This new method may be useful for a variety of low fecund species, as well as providing additional flexibility in the spawner-recruit relationship that may be explored for any stock. The method is an expansion and improvement on similar approaches previously applied to dogfish (Taylor 2008; Wood, *et al.* 1979), which assumed a linear decline in age 0 survival as a function of population density. While steepness was not estimated or assumed in the conventional sense of a Beverton-Holt stock-recruitment relationship, a value for steepness can be calculated using a formula provided by Gertseva and Taylor (2011). The calculated value of steepness is 0.28, indicating a great degree of compensation or density-dependent recruitment.

### **Fishing Mortality**

During the last 10 years, relative exploitation rates (catch/summary biomass) are estimated to have hovered around 1% and SPR is estimated to be well above pre-2015 management target of SPR 45%. The 2011 assessment identified a period during the vitamin A fishery in the 1940s when the exploitation rate exceeded the current  $F_{MSY}$  proxy harvest rate.

#### **1.1.4.19 Splitnose Rockfish South of 40°10' N Lat.**

### **Distribution and Life History**

Splitnose rockfish (*Sebastes diploproa*) are distributed from the northern Gulf of Alaska (Prince William Sound) to central Baja California and occur at depths between 91-795 meters. Adults are the most abundant between British Columbia and southern California at depths from 215 to 350 meters (Alverson, *et al.* 1964;

Gunderson and Sample 1980; Love, *et al.* 2002). The species is distinguished by having a deeply notched upper jaw, which inspired its Greek name *diploproa*, meaning “double prow”. Splitnose rockfish are commonly seen on low-relief mud fields of the continental shelf and upper slope, often near isolated rock, cobble or shell debris. Solitary individuals are often found resting on the seafloor, although they occasionally form schools that move more than 100 meters in the water column (Love, *et al.* 2002; Rogers 1994).

Splitnose rockfish co-occur with an assemblage of slope rockfish, including Pacific ocean perch (*Sebastes alutus*), darkblotched rockfish (*Sebastes crameri*), yellowmouth rockfish (*Sebastes reedi*), and sharpchin rockfish (*Sebastes zacentrus*) off Washington and Oregon, and striptail rockfish (*Sebastes saxicola*), darkblotched rockfish and shortspine thornyhead (*Sebastolobus alascanus*) off central California. Pacific ocean perch and darkblotched rockfish are the most abundant members of that assemblage off the coasts of Oregon and Washington, but splitnose rockfish and darkblotched rockfish dominate off the northern coast of California. Lesser amounts of splitnose have also been noted in the deep water DTS assemblage and with shrimp catch (Rogers 1994; Rogers and Pikitch 1992; Weinberg 1994).

There are no clear stock delineations for splitnose rockfish in the U.S. waters. No molecular markers have yet been developed for this species, and no genetic data are currently available to suggest the presence of several stocks (Waples, *et al.* 2008). No distinct breaks are seen in the fishery landings and catch distributions. Survey catches imply a continuous distribution. The spatial dynamic cluster analysis of the Northwest Fisheries Science Center (NWFS) survey abundance indices (Cope and Punt 2009) provided no evidence of spatial stock structure for splitnose rockfish off Washington, Oregon, and California.

Splitnose rockfish are documented in the literature to live to at least 86 years (Bennett, *et al.* 1982), although a fish encountered in a NMFS survey was aged at 103 years old. This is a small species – the maximum size reported in the literature is 46 cm (Love, *et al.* 2002); the vast majority of individuals caught in NMFS surveys were under 44 cm in fork length, although a few fish larger than this were caught.

Splitnose rockfish exhibit sexual dimorphism in growth. Although the males grow to their maximum lengths earlier than females, females reach larger sizes than males (Boehlert 1980; Love, *et al.* 2002). It was hypothesized that life history characteristics may vary with latitude, but that is uncertain. Boehlert and Kappenman (1980) detected greater size-at-age with increasing latitude and suggested more rapid growth of fish in the northern end of their range. Analysis of the NWFS shelf-slope survey data did not show a distinct gradient in growth rate between north and south, although the asymptotic length ( $L_{inf}$ ) exhibits a latitudinal gradient (Gertseva, *et al.* 2009). Growth of splitnose rockfish was found to correlate with climate and environmental variables, including sea surface temperature, the El Niño Southern Oscillation (ENSO) index, and the Pacific Decadal Oscillation (PDO) (Black 2009; Black, *et al.* 2008); more information is needed to develop climate-growth relationships for stock assessment purposes.

Female splitnose rockfish off California mature at 6-9 years old (18-23 cm long) (Echeverria 1987), and their fecundity increases with size (Phillips 1964). Splitnose rockfish mature somewhat later off British Columbia - both males and females reach 50% maturity at size of 27 cm (Westrheim 1975). Like other rockfishes, splitnose utilize internal fertilization and bear live young (Love, *et al.* 2002). This species can exhibit a long reproductive season, with young larvae found in all months off southern California, from January to September off central California, from March to September in Oregon, and in July off Washington (Love, *et al.* 2002; Moser, *et al.* 2000).

Young juveniles live at the surface for several months, then go through a transitory midwater residence, and finally settle to benthic habitats near the end of their first year of life (Love, *et al.* 2002). During their first year, splitnose have been found living among drifting vegetation in Puget Sound and southern

California, and under floating objects in Queen Charlotte Sound, British Columbia (Shaffer, *et al.* 1995). Pelagic juvenile splitnose feed on calanoid copepods and amphipods (Shaffer, *et al.* 1995), while benthic juveniles and adults eat krill, copepods, sergestid shrimps and amphipods. Splitnose are prey of Steller sea lions and other pinnipeds (Love, *et al.* 2002).

Size-composition data for splitnose rockfish show a strong gradient of body size with depth, with smaller fish in shallow waters, suggesting ontogenetic movements of splitnose rockfish to deeper waters with increasing size and age, a common phenomenon in the genus *Sebastes* (Boehlert 1980).

### **Stock Status and Management History**

Limits on domestic rockfish catches were first instituted in 1983, with splitnose rockfish managed as a part of the *Sebastes* complex, which included around 50 species. The ABC for the *Sebastes* complex was estimated for each International North Pacific Fisheries Council (INPFC) area along the coast based on historic landings. In 1994, the *Sebastes* complex was divided into southern and northern management areas, and harvest guidelines were established for the complex in each area. The southern area included Conception, Monterey and Eureka INPFC areas, and the northern area included Columbia and U.S.-Vancouver INPFC areas.

In response to a concern that deep water species off Oregon and Washington might have been overharvested, Rogers (1994) conducted a preliminary assessment of splitnose rockfish, which focused on compiling and reviewing the available data. However, since the data were sparse and no evident trends in biomass or mean size were detected, the results were inconclusive. In 1996 the status of several rockfish species, which were part of the *Sebastes* complex, were assessed (Rogers, *et al.* 1996), and ABCs for splitnose rockfish in the southern area were calculated to be 868 mt for the southern management area and 274 mt for the northern management area. These amounts were not specified individually, but included in the total ABCs for the *Sebastes* complex.

In 1998, unusually high splitnose rockfish landings drove *Sebastes* complex harvests in the southern management area sharply upward. In 1999, for the first time, splitnose rockfish were individually separated from the southern *Sebastes* complex. Individual ABCs and OYs for splitnose rockfish in that area have been specified along with splitnose-specific trip limits since then. The ABC for the southern management area was set at 868 mt, as estimated in the 1996 assessment of the remaining rockfish in the *Sebastes* complex (Rogers, *et al.* 1996).

Additionally in 1999, the general *Sebastes* complex was divided into nearshore, shelf, and slope assemblages, and the dividing line between the northern and southern management areas was shifted southward to 40°10' N lat., near Cape Mendocino. Since that time, in the northern area, splitnose has been managed under trip limits for minor slope rockfish. In 2000, harvest specifications for splitnose rockfish were set for the Conception and Monterey areas only, and 48 mt for the Eureka area were added to the northern minor rockfish ABC. Also, a precautionary adjustment of the OY (reduced from the ABC by 25%) was specified to account for the limited nature of the assessment. In 2000, the ABC and OY for splitnose rockfish south of 40°10' N lat. were reduced based on the revised  $F_{MSY}$  harvest rate policy. During the last 10 years, the coastwide landings and total catch of splitnose rockfish were relatively low, and the limits established for the area south of 40°10' N lat. have not been exceeded.

Gertseva *et al.* (2009) assessed splitnose rockfish coastwide and determined the stock was healthy with a depletion of 66 percent at the start of 2009. Since 1999, the splitnose spawning output was estimated to have been increasing in response to below-average removals and above-average recruitment during the last decade. At the beginning of 2009 the estimated spawning stock output was 8,426 million eggs. Uncertainty in the model was explored through asymptotic variance estimates and sensitivity analyses. Asymptotic



confidence intervals were estimated within the model and reported throughout the assessment for key model parameters and management quantities. Uncertainty in recent recruitment was used to define alternative states of nature and develop the decision table.

### **Stock Productivity**

Steepness of the stock-recruitment curve was fixed at a value of 0.58 in the 2009 splitnose rockfish assessment, as estimated by a meta-analysis for unassessed rockfish. Recruitment deviations were estimated for each year between 1960 and 2006, which was the period best informed by the data based on evaluation of the variance of the recruitment deviations. Prior to 1960 and after 2006, recruits were taken deterministically from the stock-recruit curve. The model estimated above-average recruitments in the most recent years beginning 1999 (), which along with low catches during the last decade determine a population increase in recent and early forecast years. Uncertainty in recent recruitment was used to define alternative states of nature and develop the decision table.

### **Fishing Mortality**

Splitnose rockfish have been taken incidentally in fisheries such as the trawl fisheries targeting POP, mixed slope rockfish, and other deep water targets, but have not been a commercial target species. Splitnose rockfish were lightly exploited until the 1940s, when the trawl fishery for rockfish first became important. With the development of the POP fishery (a species with which splitnose rockfish co-occur), spawning output of splitnose rockfish began to decline. A sharp drop in the 1960s was associated with large harvests of POP by foreign trawl fleets operating in the U.S. EEZ. Another drop occurred in 1998 when the increased availability of splitnose rockfish led to high removals off California. Since 1999, the splitnose spawning output was estimated to have been increasing in response to below-average removals and above-average recruitment during the last decade.

It was decided to continue management of splitnose rockfish with stock-specific specifications south of 40°10' N lat. and under the Slope Rockfish complex north of 40°10' N lat. when the coastwide splitnose rockfish assessment was first used to inform management in 2011. A north-south apportionment based on the average 1916-2008 assessed area catch resulting in 64.2 percent stock-specific specification in the southern area and 35.8 percent for the contribution of splitnose rockfish to the Slope Rockfish North complex was used to apportion harvest specifications since 2011. The Council recommended continuing this management strategy largely due to the implications of determining the uncertain catch history by trawl permit to initially allocate trawl splitnose quota shares (QS) under Amendment 20. Since splitnose rockfish are not targeted and predominantly discarded at sea, little data would be available to determine catch history.

#### **1.1.4.20 Starry Flounder**

### **Distribution and Life History**

Starry flounder (*Platichthys stellatus*) have a very broad geographic distribution around the rim of the North Pacific Ocean and have been recorded from Los Angeles to the Aleutian Islands, although they are rare south of Point Conception (Kramer and O'Connell 1995; Orcutt 1950). Off the U.S. west coast starry flounder are found commonly in nearshore waters, especially in the vicinity of estuaries (Baxter 1999; Kimmerer 2002; NOAA 1990; Orcutt 1950; Pearson 1989; Sopher 1974). It has quite a shallow bathymetric distribution, with most individuals occurring in waters less than 80 m, although specimens have been collected off the continental shelf in excess of 350 m (Kramer and O'Connell 1995; Orcutt 1950). They are most often found on gravel, clean shifting sand, hard stable sand, and mud substrates.

Spawning occurs primarily during the winter months of December and January, at least in central California (Orcutt 1950); it may occur somewhat later in the year (February-April) off British Columbia and Washington (Hart 1988; Love 1996). Egg/larval development apparently takes about 2-3 months to occur. Offspring principally remain within the estuaries until age two, when many have migrated to the adjacent ocean habitats (Baxter 1999; Kimmerer 2002; Orcutt 1950). Reproductive maturity occurs at age two years for males and age three years for females, when the fish are 28 cm and 35 cm, respectively. Tagging studies have shown that fish are relatively sedentary and move little during their adult lives (Love 1996); however, there is little information on regional variation in stock structure.

Starry flounder consume crabs, shrimps, worms, clams and clam siphons, other small mollusks, small fish, nemertean worms, and brittle stars (Hart 1988).

### **Stock Status and Management History**

The U.S. west coast starry flounder stock was assessed in 2005 (Ralston 2006). The assessment was based on the assumption of separate biological populations north and south of the California-Oregon border. The assessment used catch data, relative abundance indices derived from trawl logbook data, and an index of age-1 abundance from trawl surveys in the San Francisco Bay and Sacramento-San Joaquin River estuary. Unlike most other groundfish stock assessments, no age- or length-composition data were directly used in the assessment. Both the northern and southern populations were estimated to be above the target level of 40 percent of virgin spawning biomass (44 percent in Washington-Oregon and 62 percent in California), although the status of this data-poor species remained fairly uncertain compared to that of many other groundfish species. One of the most significant areas of uncertainty in the assessment was the estimate of natural mortality rate, which was quite high (0.30 for females and 0.45 for males).

Starry flounder were managed in the Other Flatfish complex until 2007, when the stock was removed from the complex and managed with stock-specific specifications determined from the assessment. Starry flounder have never been overfished or subject to overfishing.

### **Stock Productivity**

Recruitment deviations were estimated in both the northern and southern starry flounder assessment models, although selectivity patterns were fixed external to the model after analysis of trawl length composition information from the PacFIN-BDS data base and sport length composition information from the RecFIN data base. Growth and other life history parameters were also fixed, largely based on a detailed study of starry flounder by Orcutt (1950). Finally, spawner-recruit steepness ( $h = 0.80$ ) and recruitment variability ( $\sigma_r = 1.00$ ) were also held constant.

Starry flounder is a relatively productive stock with a PSA productivity score of 2.15. They are also not vulnerable to potential overfishing ( $V = 1.04$ ).

### **Fishing Mortality**

Starry flounder are mostly caught in nearshore recreational fisheries. Historically, they were also caught in nearshore trawl efforts; however, this catch is rare today given that Washington and California have closed their state nearshore waters to trawling. Both the northern and southern stocks were estimated to be well above the  $B_{25\%}$   $B_{MSY}$  threshold ( $B_{44\%}$  in Washington-Oregon and  $B_{62\%}$  in California). In addition, recent exploitation rates have been well below the  $F_{MSY}$  proxy for flatfish. Recent landings in both areas have been less than 20% of the calculated ABC/OFL.

#### 1.1.4.21 Widow Rockfish

##### **Distribution and Life History**

Widow rockfish (*Sebastes entomelas*) range from Albatross Bank off Kodiak Island to Todos Santos Bay, Baja California, Mexico (Eschmeyer, *et al.* 1983; Miller and Lea 1972; NOAA 1990). They occur over hard bottoms along the continental shelf (NOAA 1990) and prefer rocky banks, seamounts, ridges near canyons, headlands, and muddy bottoms near rocks. Large widow rockfish concentrations occur off headlands such as Cape Blanco, Cape Mendocino, Point Reyes, and Point Sur. Adults form dense, irregular, midwater and semi-demersal schools deeper than 100 m at night and disperse during the day (Eschmeyer, *et al.* 1983; NOAA 1990; Wilkins 1986). All life stages are pelagic, but older juveniles and adults are often associated with the bottom (NOAA 1990). All life stages are fairly common from Washington to California (NOAA 1990). Pelagic larvae and juveniles co-occur with yellowtail rockfish, chilipepper, shortbelly rockfish, and bocaccio larvae and juveniles off Central California (Reilly, *et al.* 1992).

Widow rockfish are ovoviviparous, have internal fertilization, and brood their eggs until released as larvae (NOAA 1990; Reilly, *et al.* 1992). Mating occurs from late fall-early winter. Larval release occurs from December through February off California, and from February through March off Oregon. Juveniles are 21 mm to 31 mm at metamorphosis, and they grow to 25 cm to 26 cm over three years. Age and size at sexual maturity varies by region and sex, generally increasing northward and at older ages and larger sizes for females. Some mature in three years (25 cm to 26 cm), 50 percent are mature by four years to five years (25 cm to 35 cm), and most are mature in eight years (39 cm to 40 cm) (NOAA 1990). The maximum age of widow rockfish is 28 years, but rarely over 20 years for females and 15 years for males (NOAA 1990). The largest size is 53 cm and about 2.1 kg (Eschmeyer, *et al.* 1983; NOAA 1990).

Widow rockfish are carnivorous. Adults feed on small pelagic crustaceans, midwater fishes (such as age-one or younger Pacific whiting), salps, caridean shrimp, and small squids (Adams 1987; NOAA 1990). During spring, the most important prey item is salps, during the fall fish are more important, and during the winter widow rockfish primarily eat sergestid shrimp (Adams 1987). Feeding is most intense in the spring after spawning (NOAA 1990). Pelagic juveniles are opportunistic feeders, and their prey consists of various life stages of calanoid copepods, and euphausiids (Reilly, *et al.* 1992).

##### **Stock Status and Management History**

Widow rockfish are an important commercial species from British Columbia to central California, particularly since 1979, when Oregon trawl fisherman demonstrated the ability to make large catches at night using midwater trawl gear. Many additional participants entered the fishery resulting in a rapid increase in landings of widow rockfish (Love, *et al.* 2002). Widow rockfish are a minor component of the recreational groundfish fisheries.

Williams *et al.* (2000) assessed the coastwide stock of widow rockfish in 2000. The spawning output level (8,223 mt eggs), based on that assessment and a revised rebuilding analysis (Punt and MacCall 2002) adopted by the Council in June 2001, was at 23.6 percent of the unfished level (33,490 mt eggs) in 1999. The widow rockfish stock was declared overfished in 2001 based on this assessment result.

It was concluded in the 2003 assessment (He, *et al.* 2003) that the widow rockfish stock size was at 24.7 percent of the unfished biomass and that stock productivity was considerably lower than previously thought. Results from the 2003 widow rockfish rebuilding analysis were used to develop the first widow rockfish rebuilding plan, which was adopted in April 2004 under Amendment 16-3 to the groundfish FMP. The rebuilding plan established a target rebuilding year of 2038 and a harvest control rule of  $F = 0.0093$ .

A full assessment was completed in 2005 for widow rockfish (He, *et al.* 2006a). The base model estimated that spawning biomass declined steadily since the early 1980s and that spawning output in 2004 was 31 percent of the unexploited level, above the Council's overfished threshold. Further, spawning output in the base model was estimated to have never dropped below the 25 percent overfished threshold. The 2005 rebuilding analysis indicated that the stock was much closer to reaching a rebuilt biomass than previously estimated: under the 2005 rebuilding analysis (He, *et al.* 2006b),  $T_{MIN}$  was estimated to be 2013, compared to a  $T_{MIN}$  of 2026 in the 2003 analysis (He, *et al.* 2003). This rebuilding analysis was used to modify the widow rockfish rebuilding plan, which was adopted under Amendment 16-4 in 2006. The target rebuilding year under the modified rebuilding plan was 2015 and the harvest control rule was an SPR harvest rate of 95 percent.

An updated assessment was done in 2007 (He, *et al.* 2008) using the same age-based model (written in ADMB) and data compiling procedures used in the previous assessment. The estimated total biomass in 2006 was 120,132 mt and the estimated depletion rate was 35.5 percent of the unfished spawning output. The population was projected to recover to the target in 2009, which was six years earlier than the target year in the rebuilding plan. Based on these results, the SSC recommended no changes to the rebuilding plan.

A full assessment of widow rockfish was conducted in 2011 (He, *et al.* 2011), which indicated the spawning stock biomass was successfully rebuilt with a depletion of 51 percent at the start of 2011. However, there was considerable uncertainty regarding the stock assessment's finding that the stock has rebuilt. Productivity and status of this stock are highly uncertain because the available biomass indices are not informative. Nonetheless, the SSC considered the base model of the new widow rockfish assessment to be the best available science.

Assessment of widow rockfish is hampered by a lack of reliable assessment information. Future research needs include reliable abundance indices; continuation of the long-term midwater juvenile trawl survey used to infer year class strengths; better understanding of the relationship between recruitment and environmental conditions in the California Current Ecosystem, improved short-term forecasts of productivity, biomass levels and allowable catches from stock assessments; incorporation of new discard data; evaluation of the utility of acoustic surveys; increased age-collection programs to increase sample size; and determination of age-composition for the triennial survey.


A new full assessment of widow rockfish is planned for 2015.

### **Stock Productivity**

The major axis of uncertainty in the new widow rockfish assessment is steepness, which defines the relative productivity of the stock. The SSC recommended fixing the steepness parameter at 0.76 in the assessment, due to the lack of information to reliably estimate steepness. The steepness parameter of 0.76 is the median value in the distribution of steepness parameters of assessed rockfish species in the Dorn (2002b) meta-analysis. The decision table in the assessment was developed to bracket model uncertainty in widow rockfish productivity with alternative values of steepness. The 12.5 percent and 87.5 percent quantiles from the prior distribution on  $h$  translate into steepness values of 0.54 and 0.95, respectively. This range was considered reasonable to account for the uncertainty associated with steepness. It was, however, agreed by the STAT and the SSC to shift this range to a lower steepness value to (a) take account of the data which, while not greatly informative, did provide some evidence for a lower steepness value, and (b) provide continuity by considering the value of steepness used in the 2009 assessment (0.41). As a result, steepness values of 0.41 and 0.90 were used for the low and high states of nature in the assessment decision table.

The high uncertainty in the steepness of the stock-recruitment relationship and the lack of recent strong recruitments motivates a precautionary approach to managing widow rockfish. If the pessimistic state of nature is correct ( $h = 0.41$ ), then annual constant catches of up to 1,500 mt are projected to maintain spawning stock biomass above the MSST during the 10-year projection period (i.e., 2013-2022).

The base model in the 2011 widow assessment estimated a time series of recruitment of age-0 fish from 1948 to 2009. The highest recruitment occurred in 1970 (Figure 1-25). Recruitments remained generally low in the early 1990s and have been very low since 2001, as compared to the long-term average. As in the past widow assessments, uncertainties in estimation of recruitment remain high.



**Figure 1-25. Time series of estimated recruitments from the base model in the 2011 widow rockfish assessment.**

### **Fishing Mortality**

Widow rockfish are caught mostly in midwater trawls used to target Pacific whiting and, before 2002, used to target widow and yellowtail rockfish. The exploitation rate was above the target SPR of 50 percent (i.e.,  $F < F_{MSY}$ ) until the late 1970s when trawl catches in the target midwater fishery increased to rates beyond the target. This continued until the stock was declared overfished and managed under a rebuilding plan. Harvest declined dramatically and the estimated SPR harvest rates increased rapidly above target  $F_{MSY}$ . The increase in biomass during the past decade was the result of reduced catches rather than strong year-classes.

Lower OYs specified in 2005-2010 were not exceeded as the fishery was managed to avoid widow bycatch and the percent of OY attainment decreased with time during that period. The percent attainment of the 2011 IFQ allocation was 40 percent. The at-sea whiting sectors have been better able to avoid widow rockfish in recent years with the lowest bycatch rates (widow catch/whiting catch) observed in the past couple of years.

Management uncertainty is low since widow rockfish is a trawl-dominant species and there is mandatory 100 percent observer coverage in trawl fisheries.

#### 1.1.4.22 Yellowtail Rockfish North of 40°10' N lat.

##### **Distribution and Life History**

Yellowtail rockfish (*Sebastes flavidus*) range from San Diego, California, to Kodiak Island, Alaska (Fraidenburg 1980; Gotshall 1981; Lorz, *et al.* 1983; Love, *et al.* 2002; Miller and Lea 1972; Norton and MacFarlane 1995). The center of yellowtail rockfish abundance is from Oregon to British Columbia (Fraidenburg 1980). Yellowtail rockfish are a common species abundant over the middle shelf (Carlson and Haight 1972; Fraidenburg 1980; Tagart 1991; Weinberg 1994). Yellowtail rockfish are most common near the bottom, but not on the bottom (Love, *et al.* 2002; Stanley, *et al.* 1994). Yellowtail rockfish adults are considered semi-pelagic (Stanley, *et al.* 1994; Stein, *et al.* 1992) or pelagic, which allows them to range over wider areas than benthic rockfish (Percy 1992). Adult yellowtail rockfish occur along steeply sloping shores or above rocky reefs (Love, *et al.* 2002). They can be found above mud with cobble, boulder and rock ridges, and sand habitats; they are not, however, found on mud, mud with boulder, or flat rock (Love, *et al.* 2002; Stein, *et al.* 1992). Yellowtail rockfish form large (sometimes greater than 1,000 fish) schools and can be found alone or in association with other rockfishes (Love, *et al.* 2002; Percy 1992; Rosenthal, *et al.* 1982; Stein, *et al.* 1992; Tagart 1991). These schools may persist at the same location for many years (Percy 1992).

Yellowtail rockfish are viviparous (Norton and MacFarlane 1995) and mate from October to December. Parturition peaks in February and March and from November to March off California (Westrheim 1975). Young-of-the-year pelagic juveniles often appear in kelp beds beginning in April and live in and around kelp in midwater during the day, descending to the bottom at night (Love, *et al.* 2002; Tagart 1991). Male yellowtail rockfish are 34 cm to 41 cm in length (five years to nine years) at 50 percent maturity, females are 37 cm to 45 cm (six years to ten years) (Tagart 1991). Yellowtail rockfish are long-lived and slow-growing; the oldest recorded individual was 64 years old (Fraidenburg 1980; Tagart 1991). Yellowtail rockfish have a high growth rate relative to other rockfish species (Tagart 1991). They reach a maximum size of about 55 cm in approximately 15 years (Tagart 1991). Yellowtail rockfish feed mainly on pelagic animals, but are opportunistic, occasionally eating benthic animals as well (Lorz, *et al.* 1983). Large juveniles and adults eat fish (small Pacific whiting, Pacific herring, smelt, anchovies, lanternfishes, and others), along with squid, krill, and other planktonic organisms (euphausiids, salps, and pyrosomes) (Love, *et al.* 2002; Phillips 1964; Rosenthal, *et al.* 1982; Tagart 1991).

##### **Stock Status and Management History**

Until the late 1990s, yellowtail rockfish were harvested as part of a directed midwater trawl fishery. Yellowtail rockfish are common in both commercial and recreational fisheries throughout its range, and commonly occur with canary and widow rockfishes (Cope and Haltuch 2012). Despite its popularity in commercial and recreational fisheries, its association with those highly regulated species has greatly decreased removals over the last decade. From the end of 2002 through 2010, implementation of the RCAs and small landings limits designed to only accommodate incidental bycatch eliminated directed mid-water fishing opportunities for yellowtail rockfish in non-tribal trawl fisheries. A limited opportunity to target

yellowtail rockfish in the trawl fishery has been available since 2011 under the trawl rationalization program, yet low quotas for widow rockfish, canary rockfish, and for other constraining stocks has continued to limit mid-water targeting of yellowtail rockfish.

Yellowtail rockfish are currently managed with stock-specific harvest specifications north of 40°10' N lat. and within the southern Shelf Rockfish complex south of 40°10' N lat. There has never been an assessment of the southern stock and the OFL contribution of yellowtail rockfish to the southern Shelf Rockfish complex is based on a DB-SRA estimate.

Yellowtail rockfish on the U.S. west coast north of 40°10' N lat. were assessed in 1984 (Weinberg, *et al.* 1984), 1986 (Coleman 1986), 1988 (Tagart 1988), 1993 (Tagart 1993), 1996 (Tagart and Wallace 1996), and 1997 (Tagart, *et al.* 1997) to determine harvest specifications for the stock. A full assessment in 2000 (Tagart, *et al.* 2000) was the first that estimated stock status with an estimated depletion of 60.5% at the start of 2000. Lai *et al.* (2003) updated the 2000 assessment and estimated stock depletion was 46% at the start of 2003. Another assessment update was prepared in 2005 (Wallace and Lai 2006) with an estimated depletion of 55% at the start of 2005.

A new data-moderate assessment of yellowtail rockfish north of 40°10' N lat. was conducted in 2013 (Cope, *et al.* 2014). The estimated depletion at the start of 2013 was 68% and the spawning biomass was estimated to be 50,043 mt. This was a large biomass increase relative to previous estimates and can be attributed to the low removals in the last 10 years.

### **Stock Productivity**

The posterior median estimate of steepness in the 2013 yellowtail rockfish assessment is 0.78, equal to the mean prior steepness used in the assessment and indicative of a relatively productive stock. However, this estimate may not be as informative of relative stock productivity. Due to the low susceptibility of yellowtail rockfish to fisheries removals, the vulnerability to overfishing of yellowtail rockfish is relatively low ( $V = 1.88$ ), though the productivity of this species is also relatively low ( $P = 1.33$ ) based on other life history traits, including a longevity to almost 70 years.

### **Fishing Mortality**

Fishing mortality of yellowtail rockfish north of 40°10' N lat. was relatively high in the 1980s and 1990s with direct targeting by mid-water trawl gear of yellowtail and widow rockfish. The elimination of that fishery in 2003 to reduce impacts on widow rockfish (and canary rockfish to some degree), coupled with RCA implementation, significantly reduced fishing mortality of yellowtail rockfish. The decision table in the 2013 assessment predicts the stock will keep building under the average annual catch estimated in the assessment (1,376 mt) and would remain at a healthy level in the next 10 years (i.e., above  $B_{MSY}$ ) at catch levels over 4 times that amount.

#### **1.1.5 Groundfish Stock Complexes**

There are eight stock complexes for which ACLs were specified through the 2015-2016 management cycle. These complexes are the Nearshore, Shelf, and Slope Rockfish complexes north and south of 40°10' N lat., the Other Flatfish, and the Other Fish complexes.

Most of the component stocks comprising the stock complexes are unassessed category 3 stocks with OFLs that are determined using data-poor methods such as DB-SRA, DCAC, or average historical catch (see Section 1.1.1). In cases where assessments were used to inform OFLs for component stocks managed in

stock complexes, the OFLs were projected from those assessments using proxy  $F_{MSY}$  harvest rates. A more detailed description of the assessed stocks managed in stock complexes follows.

#### 1.1.5.1 Nearshore Rockfish North and South of 40°10' N Lat.

The nearshore rockfish complexes north and south of 40°10' N lat. are comprised of both assessed and unassessed species. Of the stocks managed in the nearshore rockfish complexes, only blue rockfish in California north of Pt. Conception, brown rockfish, China rockfish, copper rockfish, and gopher rockfish in California north of Pt. Conception have been assessed. The following section defines these complexes in terms of their component stocks and provides further detail on those component stocks that have been assessed.

The Nearshore Rockfish complex north of 40°10' N lat. is composed of the following species: black and yellow rockfish (*Sebastes chrysomelas*); blue rockfish (*S. mystinus*); brown rockfish (*S. auriculatus*); calico rockfish (*S. dalli*); China rockfish (*S. nebulosus*); copper rockfish (*S. caurinus*); gopher rockfish (*S. carnatus*); grass rockfish (*S. rastrelliger*); kelp rockfish (*S. atrovirens*); olive rockfish (*S. serranoides*); quillback rockfish (*S. maliger*); and treefish (*S. serriceps*). These stocks are all unassessed with the exception of blue rockfish in California, brown rockfish, China rockfish, copper rockfish, and gopher rockfish in California.

The Nearshore Rockfish complex south of 40°10' N lat. is further subdivided into the following management categories: 1) shallow nearshore rockfish [comprised of black and yellow rockfish (*Sebastes chrysomelas*), China rockfish (*S. nebulosus*), gopher rockfish (*S. carnatus*), grass rockfish (*S. rastrelliger*), and kelp rockfish (*S. atrovirens*)], and 2) deeper nearshore rockfish [comprised of black rockfish (*S. melanops*), blue rockfish (*S. mystinus*), brown rockfish (*S. auriculatus*), calico rockfish (*S. dalli*), copper rockfish (*S. caurinus*), olive rockfish (*S. serranoides*), quillback rockfish (*S. maliger*), and treefish (*S. serriceps*)]. With the exception of the blue rockfish stock occurring in waters off California north of Point Conception (i.e., 34°27' N lat. to 40°10' N lat.), brown rockfish, China rockfish, copper rockfish, and gopher rockfish north of Point Conception (34°27' N lat.), all of the Nearshore Rockfish South stocks are unassessed.

### **Blue Rockfish in California**

#### **Distribution and Life History**

Blue rockfish (*Sebastes mystinus*) range from the Gulf of Alaska to northern Baja California, although they are most commonly found between Oregon and central California (Love, *et al.* 2002). They inhabit kelp forests and rocky reefs in relatively shallow depths usually to about 90 meters (50 fm) (Miller and Lea 1972; Reilly 2001), but have been landed as deep as 549 meters (300 fm) (Love, *et al.* 2002). Blue rockfish are residential, with their movements restricted to a small area, usually near the kelp canopy or pinnacles for shelter and spatial orientation (Jorgensen, *et al.* 2006; Lea, *et al.* 1999; Miller and Geibel 1973). Genetic evidence suggests distinct subpopulations of blue rockfish with a biogeographic barrier at Cape Mendocino, California (Cope 2004). More recently, evidence suggests the presence of two genetically distinct cryptic species in central California (Burford, *et al.* 2011).

Blue rockfish are primarily “selective opportunity” planktivores (Gotshall, *et al.* 1965; Love and Ebeling 1978). As juveniles, they feed on planktonic crustacea, hydroids, and algae (Miller and Geibel 1973). Adults also consume fish, squid, tunicates, scyphozoids, bull kelp nori, and pelagic gastropods (Hobson, *et al.* 1996; Lea, *et al.* 1999; Love, *et al.* 2002). Many of these prey items are made available from the relaxation of upwelling or southerly winds, explaining high blue rockfish numbers in the summer off central and northern California, where these conditions are well developed (Hobson and Chess 1988; Love, *et al.*



2002). Due to their great abundance in kelp forests, blue rockfish juveniles are recognized as a key species in the piscivore trophic web of these ecosystems (Hallacher and Roberts 1985).

### **Stock Status and Management History**

The blue rockfish stock in California waters north of Pt. Conception was assessed in 2007 and the stock's depletion was estimated to be 29.7% of its unfished spawning output at the start of 2007 (Key, *et al.* 2008); therefore, the stock is considered to be in the precautionary zone. Blue rockfish were not a highly sought species historically, but an increase in catches in the 1970s resulted in a continuous decline in spawning biomass through the early 1990s. The abundance of blue rockfish was at the management target (SB<sub>40%</sub>) in 1980 and at the overfished threshold in 1982. Spawning biomass reached a minimum (10% of unexploited) in 1994 and 1995; however, there has been a constant increase since then.

During the 2009 and 2010 biennial specifications process, the Council contemplated removing blue rockfish from the Nearshore Rockfish complexes. Blue rockfish was managed within the Nearshore Rockfish complexes because of scientific uncertainty and management needs, given the interaction of blue rockfish with other nearshore species. When blue rockfish occur offshore they can be targeted separately from other nearshore rockfish, but those that occur inshore mix with other nearshore rockfish stocks. Blue rockfish are managed under California's Nearshore Fishery Management Plan which has mandatory sorting requirements for landed catch. Landings are routinely tracked and monitored, thereby reducing management uncertainty. For more efficient state management, blue rockfish remains a component stock within the Nearshore Rockfish complexes. The OFL contribution of blue rockfish is projected from the 2007 assessment using the proxy  $F_{50\%}$   $F_{MSY}$  harvest rate and apportioning 87.3 percent of the OFL based on average catches of the assessed stock south of 40°10' N lat. (Key, *et al.* 2008). The OFL contribution of blue rockfish south of 34°27' N lat. is based on DCAC. The assessed portion of the blue rockfish stock is categorized as a category 2 stock, and the unassessed portion south of 34°27' N lat. is categorized as a category 3 stock. The Council has implemented precautionary management of the California population of blue rockfish since 2009 by setting a harvest guideline for California fisheries based on the sum of the 40-10 adjusted ACL contribution north of Pt. Conception and the ABC contribution south of Pt. Conception. This HG has not been exceeded since then.

### **Stock Productivity**

A Beverton-Holt steepness of 0.58 was assumed in the 2007 blue rockfish assessment based on the median steepness in the rockfish meta-analysis done at that point in time. The GMT's PSA analysis indicates a relatively high vulnerability to potential overfishing ( $V = 2.01$ ) due partly to a relatively low relative productivity ( $P = 1.22$ ) (Table 1-2).

Recruitment is variable and highly uncertain for blue rockfish. There was little information other than the pre-recruitment index in the recent years to inform the 2007 assessment model about recruitments. Recruitment appeared to be high in the 1960s, and more recently strong year classes appeared in 1993 and 1998. The late 1970s showed all time low recruitment, with 2006 among the 3 lowest recruitment years estimated.

### **Fishing Mortality**

Blue rockfish have been an important part of the recreational fishery in California since the late 1950s (Mason 1998; Reilly, *et al.* 1993; Wilson-Vandenberg, *et al.* 1996). Commonly taken by Commercial

Passenger Fishing Vessels (CPFVs, aka partyboats), skiffs, and divers, it is among the most frequently caught species north of Point Conception (Karpov, *et al.* 1995). However, since the mid-1980s the California recreational catch has declined significantly, especially in the south. This may be a result of overfishing from the more heavily populated southern coast (Love, *et al.* 1998), where there is more angling opportunity due to more favorable access and ocean conditions (Bennett, *et al.* 2004); poor recruitment resulting from a long-term shift away from preferred cold, productive waters (Jarvis, *et al.* 2004; Love, *et al.* 2002); or the effect of increasingly strict fishing regulations. Fishing mortality exceeded current target levels from the mid-1970s through the late 1990s, but has been close to target levels since 2000.

The California blue rockfish catch has played a relatively minor role in the commercial fishery compared to the recreational fishery. This has remained true, even with the advent of the live-fish fishery in the late 1980s, although the contribution of blue rockfish has been increasing in recent years. Since the preferred dinner plate-sized catch for this fishery results in immature fish being targeted in many cases, there is concern over the potential implications of the increasing effort in this fishery. Selection of younger, smaller individuals has led to lower lifetime egg production and consequently, threatened population viability (O'Farrell and Botsford 2005; O'Farrell and Botsford 2006).

## **Brown Rockfish**

### **Distribution and Life History**

Brown rockfish (*Sebastes auriculatus*) are distributed from Prince William Sound to southern Baja California in Mexico, but are most abundant on the U.S. west coast south of Bodega Bay, California (Love, *et al.* 2002). They occur from very shallow inshore waters out to 135 m (74 fm). Brown rockfish are a sedentary rockfish found in shallow water and bays (Eschmeyer, *et al.* 1983), among sheltering weed-covered rocks or around pilings (Lamb and Edgell 1986). Brown rockfish show distinct genetic differentiation by distance in coastal populations off California (Buonaccorsi, *et al.* 2005), though no distinct break is obvious to define substocks. Life history information is not spatially resolved. While coastwide populations may be subject to localized depletion because of reef-specific associations and small home ranges, no subpopulations have been distinguished.

Brown rockfish have been aged to 34 years (Love, *et al.* 2002).

### **Stock Status and Management History**

Brown rockfish are managed in the northern and southern Nearshore Rockfish complexes. A single coastwide data-moderate assessment of brown rockfish was conducted in 2013 (Cope, *et al.* 2014). The assessment estimated the brown rockfish stock to be healthy with a depletion of 42% of its unfished biomass at the start of 2013. The brown rockfish assessment used two CPUE indices of the California recreational fisheries derived from dockside intercept surveys during 1980-2003 (north and south of Point Conception). The assessment also used two observer-based recreational CPUE indices from California Party Fishing Vessels (CPFV; i.e., charter boats) during 1999-2011 south of Point Conception and during 1988-2011 between Point Conception and Cape Mendocino. No indices were constructed for north of 40°10' N lat. since this is a rare species north of Cape Mendocino. While coastwide landings were used in the assessment, only about 1% of the cumulative coastwide landings of brown rockfish were from fisheries north of 40°10' N lat. It was assumed that the population in the north followed the same trends as the southern population.

## Stock Productivity

Brown rockfish has a notably elevated vulnerability to overfishing ( $V = 1.99$ ) but a relatively high productivity score for a rockfish ( $P = 1.72$ ) in the GMT's PSA analysis (Table 1-2).

## Fishing Mortality

Estimated exploitation rates for brown rockfish were at or above the MSY harvest level during most years between 1973 and 2003, but have remained below the MSY harvest level since then (Cope, *et al.* 2014). Median spawning biomass declined rapidly during the 1970s and 1980s but has shown an increasing trend since the mid-1990s. The fishing mortality rate in 2012 was estimated to be 63% of  $F_{MSY}$ .

## China Rockfish

### Distribution and Life History

China rockfish (*Sebastes nebulosus*) range from Kachemak Bay in the Gulf of Alaska to Redondo Beach and Nicholas Island in the Southern California Bight but are most abundant from Prince William Sound to northern California (Love, *et al.* 2002). They occur primarily in nearshore and shelf waters in depths ranging from 3 to 128 m. China rockfish are a solitary species associated with high relief habitats, especially boulder fields with many crevices. They are territorial and a study off Vancouver Island indicates that individuals are likely to move 10 m or less within their territories.

China rockfish are long-lived with the oldest age reported at 79 years (Love, *et al.* 2002). Males and females mature at about the same size and age with some fish mature at 26 cm and all fish mature at 30 cm. The maximum size is reported to be 45 cm. Larval release occurs off California from January to June peaking in January. Larvae are released later in the season in the Gulf of Alaska during April to August with peak release in May.

Chinas prey on benthic organisms including brittle stars, crabs, shrimps, chitons, and small fishes. Nudibranchs, octopi, snails, and red abalone were observed prey for China rockfish off central and northern California.

### Stock Status and Management History

China rockfish are managed in the northern and southern Nearshore Rockfish complexes. Separate data-moderate assessments of China rockfish north and south of 40°10' N lat.<sup>7</sup> were conducted in 2013 (Cope, *et al.* 2014). The China rockfish population south of 40°10' N lat. was estimated to be healthy with a depletion of 66% of its unfished biomass at the start of 2013. However, the population north of 40°10' N lat. was estimated to be more depleted and in the precautionary zone with a depletion ratio of 37% at the start of 2013. The southern China rockfish assessment used a CPUE index of the California recreational fisheries derived from dockside intercept surveys during 1980-2003, as well as an observer-based recreational CPUE index from CPFVs during 1988-2011 as indices of abundance. The northern China rockfish assessment used a CPUE index of the Oregon and northern California recreational fisheries derived

---

<sup>7</sup> Separate China rockfish data-moderate assessments were also conducted north and south of the California-Oregon border at 42° N lat. at the Council's request. The SSC recommended the Council's choice of a management line for China rockfish should dictate which assessments should be used to set harvest specifications. The Council's decision to continue to manage the stock within the Nearshore Rockfish complexes north and south of 40°10' N lat. in 2015 and beyond rendered the second set of assessments stratified at 42° N lat. moot.

from dockside intercept surveys during 1980-2003 and an Oregon onboard charter boat index during 2001-2012 as indices of abundance and assumed the population off Washington followed the same trends.

The Council decided to continue to manage China rockfish in the Nearshore Rockfish complexes in 2015-2016. The Council decided a full assessment of China rockfish should be conducted in 2015 with greater focus on modeling state-specific indices of abundance. There was also the suggestion of exploring an Oregon nearshore commercial index which can be theoretically constructed given Oregon's logbook program (an effort metric can be derived from logbook data). Available age and length composition data may also help inform some of the China rockfish dynamics.


### Stock Productivity

The productivity score for China rockfish is relatively low ( $P = 1.33$ ) and there is a major vulnerability to potential overfishing ( $V = 2.23$ ).

### Fishing Mortality

China rockfish are an important species in the nearshore recreational and commercial fisheries on the west coast. They are particularly valuable in the commercial live-fish fishery where their unique coloration and high quality flesh commands the highest prices for rockfish delivered as a live product on the west coast. California and Oregon allow nearshore commercial fisheries while Washington does not.

Fishing mortality of China rockfish south of  $40^{\circ}10'$  N lat. has been well below the  $F_{MSY}$  harvest rate. The XDB-SRA base model estimates that median MSY for southern China rockfish is 32 mt per year (Table 1-6), and the fishing mortality rate in 2012 was 28% of  $F_{MSY}$ . However, that is not the case for the population north  $40^{\circ}10'$  N lat. The XDB-SRA base model estimates that median MSY for northern China rockfish is 9 mt per year, and the fishing mortality rate in 2012 was 191% of  $F_{MSY}$ . Figure 1-26 depicts total estimated catch of China rockfish north of  $40^{\circ}10'$  N lat. relative to the 2015 OFL and ACL contributions to the complex. The cumulative 2004-2012 total estimated catch of China rockfish north of  $40^{\circ}10'$  N lat. was 191% and 221% of the cumulative 2015 OFL and ACL contributions, respectively. Maintaining these catch levels is predicted to lead to continued stock decline.



**Figure 1-26. Estimated total catch of China rockfish north of 40°10' N lat. in 2004-2012 relative to the preferred 2015 OFL contribution and ACL contribution.**

## **Copper Rockfish**

### **Distribution and Life History**

Copper rockfish (*Sebastes caurinus*) are a deep bodied and spiny rockfish that range from Kachemak Bay in the Gulf of Alaska to Isla San Benito in central Baja California and are common from Port Valdez, Alaska to northern Baja California (Love, *et al.* 2002). They range in depth from the subtidal zone to 183 m. Subadult and adult copper rockfish are found primarily in boulder fields and over high relief rocks, although they also inhabit low relief rock substrata. They perch on the substrata or hover a few meters above the bottom in aggregations and as solitary individuals (Love, *et al.* 2002). Depending on the habitat and the geographic location, coppers are often found with vermilion, brown, black, dusky, silvergray, yelloweye, quillback, or tiger rockfishes. Coppers have small home ranges in high relief habitats (<10 m<sup>2</sup>) and large home ranges in low relief habitats (<4,000 m<sup>2</sup>).

### **Stock Status and Management History**

Copper rockfish are managed in the northern and southern Nearshore Rockfish complexes. Separate data-moderate assessments of copper rockfish north and south of 34°27' N lat. were conducted in 2013 (Cope, *et al.* 2014). Both copper rockfish populations were estimated to be healthy with depletions of 76% and 48% of unfished biomass at the start of 2013 for the southern and northern populations, respectively. The southern copper rockfish assessment used a CPUE index of the California recreational fisheries derived from dockside intercept surveys during 1980-2003, as well as an observer-based recreational CPUE index from CPFVs during 1999-2011 as indices of abundance. The northern copper rockfish assessment used a CPUE index of the California recreational fisheries derived from dockside intercept surveys during 1980-2003, an observer-based recreational CPUE index from CPFVs during 1988-2011, and an Oregon onboard charter boat index during 2001-2012 as indices of abundance. The northern copper rockfish assessment assumed the population off Washington followed the same trends.

### **Stock Productivity**

The PSA productivity score of 1.95 for copper rockfish indicates a high relative productivity among rockfish species. There is a relatively major vulnerability of potential overfishing ( $V = 2.27$ ) for the stock, which ranks as the highest vulnerability score in the GMT's analysis of species managed under the Groundfish FMP (tied with roughey rockfish; Table 1-2).

### **Fishing Mortality**

Estimated exploitation rates for copper rockfish south of Point Conception were at or above the MSY harvest level during most of the 1980s. North of Point Conception, harvest rates exceeded  $F_{MSY}$  from the late 1970s through the early 1990s (Cope, *et al.* 2014). Median spawning biomass in the northern region declined rapidly from the 1960s to early 1980s, but has shown an increasing trend since the mid-1990s. Biomass south of Point Conception has been increasing at a faster rate than in the north, but follows a similar pattern of decline and recovery. The median exploitation rate in 2012 was estimated to be 35% of proxy ( $B_{40\%}$ )  $F_{MSY}$  north of Point Conception and 33% of proxy  $F_{MSY}$  south of Point Conception.

## Gopher Rockfish

### Distribution and Life History

Gopher rockfish (*Sebastes carnatus*) range from Eureka, California, to San Roque, central Baja California (Miller and Lea 1972), but are most common from Mendocino County to Santa Monica Bay, California (Love 1996). Gopher rockfish is a residential and demersal species, associated with kelp beds or rocky reefs, from the intertidal to about 264 ft. (80 m), most commonly between 30 and 120 ft. (9-37 m) (Eschmeyer, *et al.* 1983; Love 1996; Love, *et al.* 2002). One tagging study off central California (Lea, *et al.* 1999) revealed that gopher rockfish exhibit minor patterns of movement (<1.5 nm, 2.8 km) with all fish being recaptured on the same reef system where they were tagged. Another study, conducted by Matthews (1986), reported movements up to 1.2 km (0.65 nm) by gopher rockfish that traveled from a low-relief natural reef to a high-relief artificial reef. The change in substrate type may have been a factor in the movement in the Matthews study.

Gopher rockfish settle out of the plankton as large larvae (2 cm. or less in length) primarily in the canopies of giant and bull kelp (*Macrocystis pyrifera* and *Nereocystis luetkeana*, respectively) where they remain close to the fronds (Love, *et al.* 2002). Settlement occurs primarily in June and July. With growth, older individuals move down the kelp stipes to the bottom where they take up residence in rocks and crevices. They are largely territorial with home ranges of 10-12 m<sup>2</sup> (Love, *et al.* 2002).

Gopher rockfish are closely related to black-and-yellow rockfish (*Sebastes chrysomelas*) and kelp rockfish (*Sebastes atrovirens*). Gopher and black-and-yellow rockfish are distinct morphologically by color and inhabit different depth ranges (gopher have a deeper depth range), but cannot be distinguished genetically (Love, *et al.* 2002). This presents an interesting phenomenon in how speciation in rockfish may occur. There are theories that interbreeding may be lessened by individuals only breeding with others of the same color. If it is determined the two species are actually one, then the name *S. carnatus* will prevail since it was described first (Love, *et al.* 2002).

### Stock Status and Management History

Gopher rockfish was assessed for the first time in 2005 and estimated stock depletion under the base model was 97% of its unfished biomass at the start of 2005 (Key, *et al.* 2006). Although the distribution of gopher rockfish extends south into the Southern California Bight, the assessment was restricted to the stock north of Point Conception. The assessment is based on landings and length composition data from commercial and recreational fisheries (primarily hook and line gear) and an index of relative abundance (CPUE) from the CPFV Sportfish Survey database. These data sources were used to estimate population trends from 1965 to 2004. There are no fishery-independent indices of stock biomass for gopher rockfish. Assessment results indicate an upward trend in gopher rockfish biomass since the 1980s and estimates of 2005 abundance ranged between 60 percent and 110 percent of average unfished stock size; this range of depletion levels is the result of alternative emphases in the model given to the CPFV in the CPUE index, a data element identified as a major source of uncertainty.

During the 2007-2008 biennial specifications process, the Council decided to continue managing gopher rockfish within the Nearshore Rockfish South complex since there was adequate resource protection under the California Nearshore Fishery Management Plan and managing gopher rockfish with stock-specific harvest specifications could disrupt that plan. The OFL contribution of gopher rockfish north of 34°27' N lat. is projected from the 2005 assessment (Key, *et al.* 2006) using the proxy F<sub>50%</sub> F<sub>MSY</sub> harvest rate. The OFL contribution of gopher rockfish south of 34°27' N lat. is based on DCAC. The assessed portion of the

gopher rockfish stock is categorized as a category 1 stock and the unassessed portion south of 34°27' N lat. is categorized as a category 3 stock.

### **Stock Productivity**

Recruitments were modeled in the 2005 assessment assuming a Beverton-Holt relationship, with steepness fixed at  $h=0.65$  and recruitment variability fixed at  $\sigma_r = 0.5$ . Recruitment deviations were estimated for the period 1965-2000. This stock showed evidence of weak recruitment in the 1970s, with peaks in the mid-1980s and mid-1990s. Recruitment estimates in the 1970s are not reliable since length information was not available until the 1980s. Overall, recruitment has been variable throughout the entire time series.

The PSA productivity score of 1.56 for gopher rockfish indicates a moderate relative productivity among rockfish species. There is a relatively low vulnerability of potential overfishing ( $V = 1.76$ ) for the stock.

### **Fishing Mortality**

Gopher rockfish have been a minor component of the commercial and recreational rockfish fishery since at least the late 1960s. In 1980, an estimated 63 mt of gopher rockfish were landed commercially north of Point Conception, with a decrease in landings in the mid-1980s. Landings then began to increase, with a peak in the fishery occurring in 1992 when approximately 74 mt were landed. Since then, landings have slightly decreased over time. Lower recent landings in 2003 and 2004 (13 and 15 mt, respectively) are in part due to more restrictive federal limits placed on rockfishes. Hook-and-line gears have been the dominant gear type used during the 1969 to 2004 period accounting for 98% of commercial landings.

The recreational gopher rockfish fishery for California ports north of Point Conception peaked during a five-year period in the early 1990s, with 2001 and 2003 also being productive years. Since 1983, anglers caught the greatest proportion of gopher rockfish from private and rental boats (71%), followed next by party and charter boats (27%). However, in more recent years (1997 to 2004) these proportions have changed, with the private and rental boats taking 59% of gopher rockfish in the recreational fishery and 41% by the party and charter boats. Also since 1983, gopher rockfish have ranked 25th in northern California recreational fishery landings, accounting for approximately 1% of the total harvest for all recreationally caught fishes. However, gopher rockfish made up approximately 50% of the estimated take of the shallow nearshore rockfishes and 6% of all nearshore rockfish species combined. Additionally, recent catches have been influenced by size and bag limits.

Starting in the late 1980s the premium quality live- fish market developed (Larson and Wilson-Vandenberg 2001). Currently, nearly all gopher rockfish are landed in this condition due to a more lucrative high-demand market. As a result of the increasing demand for live- fish the average price per pound has risen steadily from a low of less than \$2.00/lb. at the inception of the live- fish market to approximately \$6.15/lb. in 2004 (unadjusted for inflation).

Recent exploitation rates are estimated to have been well below the  $F_{MSY}$  proxy for rockfish.

#### **1.1.5.2 Shelf Rockfish North and South of 40°10' N Lat.**

The shelf rockfish complexes north and south of 40°10' N lat. are comprised of both assessed and unassessed species. Of the stocks managed in the shelf rockfish complexes, chilipepper rockfish north of 40°10' N lat. (the assessment for the northern stock only covers the area from 40°10' N lat. to Cape Blanco, OR at 43° N lat. – see section 1.1.4.7 for more details), greenspotted rockfish, greenstriped rockfish, and

stripetail rockfish have been assessed. The following section defines these complexes in terms of their component stocks and provides further detail on those component stocks that have been assessed.

The Shelf Rockfish complex north of 40°10' N lat. is comprised of the following species: bronzespotted rockfish (*Sebastes gilli*); bocaccio (*Sebastes paucispinis*); chameleon rockfish (*S. phillipsi*); cowcod (*S. levis*); dusky rockfish (*S. ciliatus*); dwarf-red rockfish (*S. rufianus*); flag rockfish (*S. rubrivinctus*); freckled rockfish (*S. lentiginosus*); greenblotched rockfish (*S. rosenblatti*); greenspotted rockfish (*S. chlorostictus*); greenstriped rockfish (*S. elongatus*); halfbanded rockfish (*S. semicinctus*); harlequin rockfish (*S. variegatus*); honeycomb rockfish (*S. umbrosus*); Mexican rockfish (*S. macdonaldi*); pink rockfish (*S. eos*); pinkrose rockfish (*S. simulator*); pygmy rockfish (*S. wilsoni*); redstripe rockfish (*S. proriger*); rosethorn rockfish (*S. helvomaculatus*); rosy rockfish (*S. rosaceus*); silvergray rockfish (*S. brevispinis*); speckled rockfish (*S. ovalis*); squarespot rockfish (*S. hopkinsi*); sunset rockfish (*S. crocotulus*); starry rockfish (*S. constellatus*); stripetail rockfish (*S. saxicola*); swordspine rockfish (*S. ensifer*); tiger rockfish (*S. nigrocinctus*); and vermilion rockfish (*S. miniatus*).

The Shelf Rockfish complex south of 40°10' N lat. is composed of the following species: bronzespotted rockfish (*Sebastes gilli*); chameleon rockfish (*S. phillipsi*); dusky rockfish (*S. ciliatus*); dwarf-red rockfish (*S. rufianus*); flag rockfish (*S. rubrivinctus*); freckled rockfish (*S. lentiginosus*); greenblotched rockfish (*S. rosenblatti*); greenspotted rockfish (*S. chlorostictus*); greenstriped rockfish (*S. elongatus*); halfbanded rockfish (*S. semicinctus*); harlequin rockfish (*S. variegatus*); honeycomb rockfish (*S. umbrosus*); Mexican rockfish (*S. macdonaldi*); pink rockfish (*S. eos*); pinkrose rockfish (*S. simulator*); pygmy rockfish (*S. wilsoni*); redstripe rockfish (*S. proriger*); rosethorn rockfish (*S. helvomaculatus*); rosy rockfish (*S. rosaceus*); silvergray rockfish (*S. brevispinis*); speckled rockfish (*S. ovalis*); squarespot rockfish (*S. hopkinsi*); sunset rockfish (*S. crocotulus*); starry rockfish (*S. constellatus*); stripetail rockfish (*S. saxicola*); swordspine rockfish (*S. ensifer*); tiger rockfish (*S. nigrocinctus*); vermilion rockfish (*S. miniatus*); and yellowtail rockfish (*S. flavidus*).

## **Greenspotted Rockfish**

### **Distribution and Life History**

Greenspotted rockfish (*Sebastes chlorostictus*) are found in waters off the west coast of North America, ranging from Copalis Head, Washington to Isla Cedros, Baja California (approximately 25° to 47° N lat.). Abundance of this species is greatest from northern Baja California to Mendocino County in California. Greenspotted rockfish associate with several benthic habitat types between depths of 30-363 m, although adults are most common between 60 and 240 m (Love, *et al.* 2002).

Greenspotted rockfish are a long-lived and slow growing species, with sedentary adults associating with a wide variety of benthic habitats. Maximum reported age is 51 years (Benet, *et al.* 2009). Estimates of maximum length for greenspotted rockfish are in the vicinity of 50 cm. Benet *et al.* (2009) report maximum fork length as 48 cm for central California. Miller and Gotshall (1965) report 51 cm total length for the same area, but did not attempt to distinguish between greenspotted rockfish and pink rockfish (*Sebastes eos*), which grow to 56 cm (Love, *et al.* 2002). Commercial port samplers in California have reported individuals larger than 50 cm fork length (up to 57 cm), although fish of this size appear to be rare (CALCOM, 2011). In southern California, Love *et al.* (1990) report maximum length as 50 cm total length. Sexual dimorphism is not apparent in greenspotted rockfish (Benet, *et al.* 2009; Lenarz and Wyllie Echeverria 1991; Mason 1998), although latitudinal differences in weight-at-length, length-at-age, and size-at-maturity have been observed.

Seasonal maturation and size at maturity vary with latitude, a trend commonly seen in rockfishes (Benet, *et al.* 2009; Love, *et al.* 1990). In central and northern California, spawning months have been reported from



March to September, with peak parturition from April to June (Benet, *et al.* 2009; Wyllie Echeverria 1987). In southern California spawning months begin in February and extend through July, with peak parturition in April (Love, *et al.* 1990). Benet *et al.* (2009) estimate length at 50% maturity for female greenspotted as 26 cm, consistent with a previous estimate of 27 cm (Wyllie Echeverria 1987) based on females from the same area. In southern California, Love *et al.* (1990) report length at 50% maturity as 22 cm (converted to fork length from total length). Love *et al.* (1990) detected evidence of multiple broods in females from southern California (ovaries containing eyed larvae and large numbers of fertilized or unfertilized eggs). No evidence of multiple broods was found in studies of greenspotted rockfish north of Point Conception (Benet, *et al.* 2009; Wyllie Echeverria 1987).

Several studies have reported on habitat associations for greenspotted rockfish. Yoklavich *et al.* (2000) quantified deep, rocky habitat in Monterey Bay. They observed smaller greenspotted rockfish in shallow depths (75-174 m), and reported strong associations with heterogeneous habitats (cobble-mud, mud-boulder, rock-mud, and rock-ridge). Laidig *et al.* (2009) studied habitat associations of demersal fishes from a manned submersible in central California, observing 809 greenspotted rockfish. They mainly encountered immature individuals (86% of greenspotted were <25 cm), identifying positive associations with all habitat types (boulder, brachiopod beds, cobble) other than mud. The predominance of juvenile rockfish in the study area suggests that the areas and depths surveyed may be nursery grounds for juvenile rockfish and/or transitional zones as individuals move toward adult habitats (Laidig, *et al.* 2009). Juvenile greenspotted rockfish are commonly seen in traps targeting spot prawn in Monterey Bay, usually in low-relief habitats (Dick, *et al.* 2011).

Adult greenspotted rockfish are generally sedentary, and associate with a wide range of habitat types. Yoklavich *et al.* (2000) observed 426 greenspotted rockfish (fourth highest abundance of observed species) in Monterey Bay, noting that adults were common near rocky outcrops, ridges, caves, and overhangs. Anderson *et al.* (2009) described greenspotted rockfish as characteristic of transition zones between hard and soft sediments, based on in situ observations across Cordell Bank in central California. They classified habitat for greenspotted rockfish over a range of spatial scales. At the finest scale (1-10s of m), greenspotted were found to have weak associations with four of five possible categories: mud, boulders, cobbles, and rock (sand being the fifth category). At intermediate scales (10-100s of m) Anderson *et al.* (2009) characterized greenspotted habitat as depths between 100-300 m and soft and mixed sediment types.

Movements of greenspotted rockfish have been monitored using acoustic tagging experiments. Starr *et al.* (2002) implanted acoustic tags in six adults in Monterey Bay, finding that adults exhibit limited horizontal movement and almost no vertical movement. They also identified two movement patterns. In the first pattern, 94% of time was spent within a 0.58 km<sup>2</sup> area. The second pattern involved larger movements, with excursions up to 3 km, but 60% of time was spent within the 1.6 km<sup>2</sup> study area. Lowe *et al.* (2009) monitored 4 adult greenspotted rockfish near oil platforms in southern California using acoustic tags. Probabilities of detection near the release sites dropped by 14% in one year of monitoring. Two individuals returned to their release sites after a 7-month absence.

Williams and Ralston (2002) studied the distribution and co-occurrence of rockfishes over continental shelf and slope habitats using fishery-independent trawl survey data. Greenspotted rockfish were consistently caught (>80% co-occurrence) with bocaccio, chilipepper, stripetail (*S. saxicola*), and shortbelly rockfish. Williams and Ralston (2002) proposed species assemblages for management purposes, including greenspotted in a “southern shelf” assemblage along with bocaccio, chilipepper, shortbelly, stripetail, greenstriped, and cowcod. Since greenspotted rockfish is not a primary target of commercial fisheries, its association with other desirable shelf rockfish species (e.g., bocaccio and chilipepper) is likely a driving force behind historical exploitation of this species.

Molecular systematic studies (Hyde and Vetter 2007) report that greenspotted rockfish are closely related to pink rockfish and greenblotched rockfish (*S. rosenblatti*). Greenspotted rockfish can be distinguished from pink and greenblotched rockfishes by a smooth lower jaw, lacking scales found on the lower mandibles of the other two species (Love, *et al.* 2002).

### **Stock Status and Management History**

The 2011 greenspotted rockfish assessment conducted for the portion of the stock off California was modeled as two area assessments north and south of Point Conception at 34°27' N lat. The assessment indicates the stock is in the precautionary zone with spawning biomass depletions of 30.6 percent and 37.4 percent for the stocks north and south of Point Conception, respectively. The stocks have shown substantial biomass increases since implementation of the RCAs in 2003. Shelf rockfish are particularly well protected by the RCAs, and greenspotted rockfish catches have been negligible since 2003. The Council recommended continuing to manage greenspotted rockfish within the Shelf Rockfish complexes since catch histories were too uncertain to allocate QS in the IFQ fishery. The OFL contribution of greenspotted rockfish to the Shelf Rockfish North complex was based on apportioning 22.2 percent of the projected OFLs from the assessment for the stock north of Point Conception, which is the average estimated catch proportion in the assessment for the stock occurring in the area between 40°10' N lat. and the California-Oregon border at 42° N lat. The OFL contribution for the portion of the stock occurring north of 42° N lat. was derived using DB-SRA. The SSC categorized the assessed portion of the stock as a category 2 stock since recruitments were not estimated. The unassessed portion of the stock was categorized as a category 3 stock.

### **Stock Productivity**

Length and age composition data available for the 2011 greenspotted rockfish assessment contained insufficient information to reliably resolve year-class strength. Both base models assumed that recruitment followed a deterministic Beverton-Holt stock-recruitment relationship, so trends in recruitment reflected trends in estimated spawning output.

While the productivity score for greenspotted rockfish is relatively low ( $P = 1.39$ ), the susceptibility score is sufficiently low to estimate a medium vulnerability to potential overfishing ( $V = 1.98$ ).

### **Fishing Mortality**

Greenspotted rockfish are not usually a primary target of commercial or recreational fisheries. Regulations affecting this species are typically intended to alter fishing mortality of primary targets and/or overfished species. For example, implementation of RCAs statewide and CCAs in southern California has greatly reduced fishing mortality for greenspotted rockfish in the past decade.

Historical harvest rates for greenspotted rockfish peaked in the mid-1980s in southern California, but continued to rise in northern California until about a decade later. SPR harvest rates exceeded the current proxy MSY value in northern California from 1973-2000, and from 1969-1998 in southern California. Biomass in both regions is currently below target (<40% unfished spawning output), but above the MSST, and equilibrium SPR harvest rates have been below the proxy MSY level since 2001 in the north and since 1999 in the south.

## Greenstriped Rockfish

### Distribution and Life History

Greenstriped rockfish (*Sebastes elongatus*) can be found in abundance from British Columbia to Northern Baja California, but range from Chirikof Island in the Aleutian Islands (Gulf of Alaska) to central Baja California (Love, *et al.* 2002). Adults may inhabit depths between 12 and 500 meters, but are more commonly found between 100 and 250 m, and adults typically move to deeper water as they mature (Love, *et al.* 2002; Shaw and Gunderson 2006). This species of rockfish is found with other congeners or alone in a wide range of habitats, which include rocky outcroppings. However, unlike most other species of rockfish they seem to prefer mud or sand bottoms (Love, *et al.* 2002; Shaw and Gunderson 2006).

A genetic study of greenstriped rockfish was recently undertaken by Jon Hess (pers. comm., NWFSC, NOAA as cited in by Hicks *et al.* (2009)) to study the stock structure of greenstriped rockfish. The genetic variability was remarkably low and showed less variability than most other rockfish species, even when including samples from Puget Sound. However, latitudinal differences in life-history traits have been observed.

Typical of other species of the genus *Sebastes*, greenstriped rockfish are long-lived with maximum observed ages greater than 50 years (Love, *et al.* 2002). Females grow larger than males, but typically mature at about the same length, between 18 and 24 cm, which corresponds to an age between 7 and 10 years. A latitudinal cline in maturity has been observed with fish maturing at a smaller size in the southern areas (Wyllie Echeverria 1987).

Greenstriped rockfish give birth to live young and the fecundity of a 0.5 kilogram female is on average around 200,000 eggs (Dick 2009), although a wide range of fecundity has been reported (Love, *et al.* 2002). The reproductive development of males and females is slightly offset with mating occurring in December through February, fertilization occurring in early spring, and parturition occurring about a month later in late spring (Shaw and Gunderson 2006). Females have the ability to store sperm during the time between copulation and fertilization to ensure the availability of spermatozoa when oocyte maturation has occurred (Shaw and Gunderson 2006). However, in southern latitudes, parturition may occur from January to July and females in Southern California may release two broods during this time (Love, *et al.* 2002). Juveniles settle to the bottom at about 3 cm in length in autumn and are commonly found along the interface of fine sand and clay. Maturing adults typically move to deeper water (Love, *et al.* 2002).

A wide range of prey items make up the diet of greenstriped rockfish. They will feed from the water column or the bottom on such things as fish, krill, shrimps, copepods, amphipods, and squid. Other fish species may prey on greenstriped rockfish. They have been found in the stomachs of king salmon (Love, *et al.* 2002). Reefs with small numbers of piscivorous rockfish had much higher numbers of small rockfish, such as greenstriped rockfish, than reefs with high numbers of piscivorous rockfish (PFMC 2006).

### Stock Status and Management History

Greenstriped rockfish are a bycatch species with little market value mainly due to its small size, and it has been reported that fillets from this species have a short shelf life (Love, *et al.* 2002). As a result, there has not been a long-term directed fishery for this species. However, greenstriped rockfish are often observed in landings from various fisheries, although in small proportions. The most common occurrence of greenstriped rockfish is in trawl fisheries, but they are often caught in recreational fisheries, especially when fishing vessels drift off of the rocks.

After many attempts to start trawl fisheries off the west coast of the United States in the late 1800s, the availability of the otter trawl and the diesel engine in the mid-1920s helped the trawl fisheries expand (Douglas 1998). The trawl fisheries really became established during World War II when demand increased for shark livers and bottomfish. A mink food fishery also developed during World War II (Jones and Harry 1960). Foreign fleets began fishing for rockfish in the mid-1960s until the EEZ was implemented in 1977 (Rogers 2003b). Since 1977, landings of rockfish were high until management restrictions were implemented in 2000.

Greenstriped rockfish are often caught in bottom trawls, but a long-term directed fishery has not occurred for this species and historical discarding rates are not well known. There have been many reports of greenstriped rockfish occurring in various fisheries, even as early as 1884 (Goode 1884). Fishermen report that greenstriped rockfish are ubiquitous, but are rarely if ever caught in great numbers.

A coastwide assessment of greenstriped rockfish was done in 2009, which indicated stock depletion was at 81% of its unfished biomass at the start of 2009 (Hicks, *et al.* 2009). The coastwide greenstriped harvest specifications were apportioned beginning in 2011 using the mean of the 2003-2008 swept area biomass estimates north of 40°10' N lat. (84.5 percent) from the NMFS trawl survey. This stock has continued to be managed within the Shelf Rockfish complexes due to the complications associated with managing this species with IFQs. Species pulled out of a complex managed with IFQs must be converted into an IFQ management unit under the Amendment 20 rules. Greenstriped rockfish is a trawl-dominant bycatch species that is rarely landed due to their diminutive size and low market desirability. An initial allocation of quota share for greenstriped would be less than straightforward given the unreliable catch history. The SSC rated the greenstriped stock as category 2 on the basis of the very uncertain catch history in the 2009 assessment that prevented the estimation of discrete year classes.

### **Stock Productivity**

Recruitment deviations were estimated in the 2009 assessment starting in 1970. The estimates showed that recruitment was highly variable for greenstriped rockfish with high values in 1971, 1984, 1993, and 1998, and low estimates of recruitment in the 1990s, early 1970s, and 2006. The age data from the NWFSC trawl survey were very consistent with these estimates and precisely showed a very strong 1993 cohort.

While the greenstriped productivity score is relatively low ( $P = 1.28$ ), the susceptibility to high exploitation was also low leading to a medium vulnerability to potential overfishing ( $V = 1.88$ ).

### **Fishing Mortality**

The spawning output of greenstriped rockfish reached a low in the late 1990s before beginning to increase throughout the last decade. The estimated depletion has remained above the 40% of unfished spawning output target and it is unlikely that the stock has ever fallen below this threshold. Throughout the 1970s, 1980s, and 1990s the exploitation rate and SPR have generally increased and occasionally exceeded current estimates of the harvest rate limit (SPR = 50%). Recent exploitation rates on greenstriped rockfish have been very small, which is primarily due to management actions in the late 1990s and early 2000s to rebuild other species.

## **Stripetail Rockfish**

### **Distribution and Life History**

Stripetail rockfish (*Sebastes saxicola*) are found from Yakutat Bay in the eastern Gulf of Alaska to Bahia Sebastian Vizcaino in central Baja California, but are more common from coastal British Columbia to southern California (Love, *et al.* 2002). They occur in depths ranging from 25 to 547 m but are most abundant between 100 and 200 m. Adult stripetail are benthically oriented and are most often associated mud, sand, and other low relief habitats. Stripetails are found in the same habitats as splitnose rockfish, greenstriped rockfish, Dover sole, and thornyheads.

Stripetail rockfish live at least 38 years and females grow faster (after reaching maturity) and achieve a larger size than males. Stripetail rockfish are relatively small-sized rockfish with a maximum size of 41 cm and 1 kg (Love, *et al.* 2002). Female stripetails along the California coast are mature by 18 cm or about 9 years of age. Off California, larval release occurs from November to March with peak release occurring off central and northern California in February and in December in the Southern California Bight (Love, *et al.* 2002). Females produce between 15,000 and 230,000 eggs.

Stripetails are primarily water column planktivores feeding mainly on krill and copepods. They are preyed on by a number of predators including Chinook salmon.

### **Stock Status and Management History**

Stripetail rockfish are managed in the northern and southern Shelf Rockfish complexes. They are a relatively minor component stock to these complexes since stripetail are not targeted nor landed in large amounts.

A new data-moderate assessment of stripetail rockfish was conducted in 2013, which indicated the stock was healthy with a depletion exceeding 77.5% (Cope, *et al.* 2014). The 2013 assessment did not produce a reliable estimate of the scale of the stock's biomass; therefore, the SSC did not recommend using the OFL estimates in the assessment. However, the SSC did recommend the available data in the assessment provided strong evidence that the stock was well above the target  $B_{MSY}$  and that the assessment results could be used for status determination. Given that the assessment-based OFLs were not endorsed by the SSC, the OFL continues to be based on a DB-SRA methodology and the stock is therefore categorized as a category 3 stock.

### **Stock Productivity**

Two recruitment events reported in trawl studies off California from 1973-1993 occurred during El Niños (Love, *et al.* 2002). It is not clear from the literature whether this is a representative recruitment pattern for the stock.

The PSA productivity score of 1.39 for stripetail rockfish indicates a relatively low productivity among rockfish species. There is a moderate vulnerability of potential overfishing ( $V = 1.8$ ) for the stock.

## Fishing Mortality

Stripetail rockfish are not targeted in commercial or recreational fisheries due to their small size. However, they are caught incidentally in bottom trawl fisheries due to their occurrence in low relief, trawlable habitats. They are rarely landed in current trawl fisheries although they were frequently landed and sold for animal food in the 1950s and 1960s. The stock has never experienced overfishing with the exploitation rate remaining well below the proxy  $SPR = 50\% F_{MSY}$  harvest rate for rockfish.

### 1.1.5.3 Slope Rockfish North and South of 40°10' N Lat.

The slope rockfish complexes north and south of 40°10' N lat. are comprised of both assessed and unassessed species. Of the stocks managed in the slope rockfish complexes, aurora rockfish, blackgill rockfish south of 40°10' N lat., rougheye rockfish (and blackspotted rockfish), and sharpchin rockfish have been assessed. There is an older assessment of bank rockfish that was done in 2000 (Piner, *et al.* 2000) that was limited in area and is not used in current management. The following section defines these complexes in terms of their component stocks and provides further detail on those component stocks that have been assessed.

The Slope Rockfish complex north of 40°10' N lat. is comprised of the following species: aurora rockfish (*Sebastes aurora*); bank rockfish (*S. rufus*); blackgill rockfish (*S. melanostomus*); blackspotted rockfish (*S. melanostictus*); redbanded rockfish (*S. babcocki*); rougheye rockfish (*S. aleutianus*); sharpchin rockfish (*S. zacentrus*); shortraker rockfish (*S. borealis*); splitnose rockfish (*S. diploproa*); and yellowmouth rockfish (*S. reedi*).

The Slope Rockfish complex south of 40°10' N lat. is composed of the following species: aurora rockfish (*Sebastes aurora*), bank rockfish (*S. rufus*), blackgill rockfish (*S. melanostomus*), Pacific ocean perch (*S. alutus*), redbanded rockfish (*S. babcocki*), rougheye rockfish (*S. aleutianus*), sharpchin rockfish (*S. zacentrus*), shortraker rockfish (*S. borealis*), and yellowmouth rockfish (*S. reedi*).

## Aurora Rockfish

### Distribution and Life History

Aurora rockfish (*Sebastes aurora*) are encountered between the Queen Charlotte Islands (British Columbia, Canada) south to mid-Baja California (Mexico). Off of the United States, they are common from northern Oregon to southern California, and are most abundant in the area around Point Conception, California. They occur at depths from 200 to 700 m (~100 to 400 fm) with the median depth increasing to the south, such that they are most abundant from 350 to 550 m in the north and 400 to 600 m in the south.

While there are areas of greater abundance off of northern Oregon and especially off of Point Conception, California, the population appears continuous over the entire coast, so that there is no clear point for stock delineation. Survey catches exhibit a continuous distribution along the entire coast, though with areas of higher and lower abundances along the coast.

Aurora rockfish is a long-lived rockfish species, with maximum observed age of 125 years on the U.S. west coast based upon otoliths aged in the 2013 assessment (Hamel, *et al.* 2013). This is slightly greater than the maximum of 118 years seen by Thompson and Hannah (2010) and consistent with a maximum age greater than 75 as reported by Love *et al.* (2002). As with many rockfish species, aurora rockfish exhibit both spatially varying and sexually dimorphic growth, with females reaching a slightly larger size than males. Off of Oregon, females reached an asymptotic length of 36.9 cm, while males reached only 33.6 cm (Thompson and Hannah 2010). Asymptotic size and size at age decreases with latitude, and since the

bulk of the stock is south of Oregon, the average asymptotic lengths are quite a bit lower than those reported above.

Thompson and Hannah (2010) found the age at 50% maturity for female aurora rockfish to be 12.56 years and the length at 50% maturity to be 25.54 cm. Maturity data collected coastwide during the 2012 NWFSC trawl survey found similar values, though with more evidence of atresia in older and larger fish than observed in the Thomson and Hannah study.

Aurora rockfish larvae have been collected off of California in months ranging from November to August, with abundance peaking in May and June, corresponding to the observation of females with developed embryos from March to May off of California and in May in Oregon (Love, *et al.* 2002). Thompson and Hannah (2010) also found that parturition peaked in May off of Oregon. Auroras settle on the bottom when they reach a length of about 3.3 cm (Love, *et al.* 2002).

Aurora rockfish display ontogenetic movement, with smaller fish found in shallower waters (below 400-450 m). They are distributed over both hard and soft substrates (Love, *et al.* 2002).

Aurora rockfish co-occurs with many prominent groundfish targets such as Dover sole, sablefish, thornyheads, and hake, though are most reported in the catch of splitnose rockfish. Aurora rockfish contributes to the overall California Current ecosystem as both predator on crustaceans and small fishes, and as prey to larger fishes, marine mammals, and large squid. Juvenile aurora rockfishes are preyed on by salmon, birds, and other fishes (Love 2011).

Several aspects of aurora rockfish population biology are affected by the ecosystem. The recruitment of many species of rockfish appears to be high in 1999, suggesting that environmental conditions influence the spawning success and survival of larvae and juvenile rockfish, including aurora rockfish. The mechanism behind this observation is not well understood, but zooplankton abundance, changes in water temperature and currents, distribution of prey and predators, and amount and timing of upwelling are all possible linkages. Changes in the environment may also directly influence age-at-maturity, fecundity, growth, and survival, which can affect stock status determination and its susceptibility to fishing. Thompson and Hannah (2010) found variations in growth corresponding to individual years based upon dendrochronological techniques and otoliths, and found a correlation between an observed growth anomaly in otoliths and sea level in individual years.

### **Stock Status and Management History**


Aurora rockfish reside in deep waters below 200 m. The primary gear type that has been used to catch aurora rockfish and other deep water rockfish has been trawl gear. The use of trawls off the west coast of the United States dates to the late 1800s, though there was little fishery expansion until the availability of the otter trawl and the diesel engine in the mid-1920s (Douglas 1998). Trawl fisheries were mainly conducted on the shelf and became more established during World War II when demand increased for groundfish. Mink farms were also a major destination of groundfish removals in the 1940s and 1950s (Jones and Harry 1960). Foreign fleets began fishing for rockfish, including deeper waters of the slope, in the mid-1960s, with declining participation until the 200-mile EEZ was implemented in 1977 (Rogers 2003b). Peaks in the foreign catch have typically been seen in the mid-1960s for rockfishes, but for aurora rockfish, the largest catches were taken in the early 1970s. Foreign fishing was limited in the northern regions by 1970, shifting effort southward and more into aurora rockfish habitat. After 1977, domestic landings of rockfish increased rapidly until about 1990. Subsequent declines in rockfish landings were driven by declining biomass levels and implementation of new, more restrictive management practices, particularly between 1997 and 2002.

Documented and estimated removals of aurora rockfish do not reach consistently large levels until the 1980s. Aurora rockfish are and have been historically most commonly taken from central California to Oregon, tightly coupled with catches of splitnose rockfish. The term “rosefish” was often used to describe either splitnose or aurora rockfish and has been used as a reporting category in California since 1982. Aurora rockfish remains largely a non-targeted member of the slope rockfish complexes.

Limits on select rockfishes, which included the co-occurring species splitnose, were established in 1982. The first imposed catch limits on a coastwide *Sebastes* complex (aurora being one of the 50 rockfishes in the complex) were instituted in 1983. This complex was divided into two management areas north and south of 43° N lat. (separating the Eureka and Columbia INPFC areas) in 1994. Ongoing concern that shelf and slope rockfishes may be undergoing overfishing led the attempt by Rogers et al. (1996) to describe the status of most rockfishes contained in the *Sebastes* complex. Aurora rockfish information content was low, so only estimates of exploitation rates were provided, indicating the stock was undergoing very high exploitation rates relative to biomass estimates in both management areas.

The *Sebastes* complex was subsequently divided into nearshore, shelf, and slope complexes effective in the year 2000 and the dividing line between the northern and southern management areas was shifted to 40°10' N. lat. Aurora rockfish has been managed under trip limits for the minor slope rockfish complex in both the north and south management areas from 2000-2010. Beginning in 2011, bottom trawl catches of slope rockfish north and south of 40°10' N lat. have been managed under an IF Q system.

The first assessment of the west coast stock of aurora rockfish was conducted in 2013 (Hamel, *et al.* 2013); the assessment estimated stock depletion was at 64 percent of its unfished equilibrium at the start of 2013 and had never dropped below its  $B_{MSY}$  target (Figure 1-27). The assessment was an age-based full assessment with natural mortality identified as the major axis of uncertainty. The SSC categorized aurora rockfish as a category 1 stock based on the assessment. However, the uncertainty in estimated biomass in the 2013 assessment was greater than for other category 1 assessments resulting in a higher sigma value for defining the ABC buffer (see section 1.3.2 for more details).



**Figure 1-27. Time series of estimated spawning biomass and depletion of aurora rockfish, 1916-2013.**



## Stock Productivity

Steepness was fixed to the mean of the most recent rockfish steepness prior ( $h = 0.779$ ; Thorson, 2013) in the 2013 assessment. Recruitment deviations were estimated from 1916 (the beginning of the modeling period), with a ramp towards bias correction beginning in 1962, full-bias adjustment beginning in 1970 and ending in 2008, and a ramping back down to no bias correction in 2012. Two of the largest contemporary recruitment events are found in 1999 and 2007 (Figure 1-28). Despite the inclusion of estimated ageing error, discerning individual year classes remains difficult and significant correlation exists between the estimated strength of adjacent year classes, which may be primarily due to ageing error rather than actual correlation in recruitment strength.




Figure 1-28. Time series of estimated age-0 recruits of aurora rockfish on the U.S. west coast, 1970-2013.

## Fishing Mortality

The 2013 estimates that exploitation of aurora rockfish has been relatively low, with total catch estimated to have exceeded the current management harvest-rate limits in 7 years (1983, 1988-1990, and 1992-1994), during the early peak in trawl catch (Figure 1-29). Recent levels of removals have remained moderate. There seems to be very low risk that current removals are causing overfishing.

While stock-specific OFLs/ABCs were not historically set for aurora rockfish specifically, the reauthorized Magnuson-Stevens Act of 2006 and FMP Amendment 23 required OFLs for all species in a management plan, including those managed in stock complexes. The first OFL contributions were calculated using DB-SRA and provided in 2011. The 2015 and beyond OFLs are projected from the 2013 assessment. Recent catches since 2002 have been below the new 2015 OFL and ABC (Figure 1-30).




Figure 1-29. Time series of estimated relative spawning potential ratio (1-SPR/1-SPR(Target=0.50)) for aurora rockfish, 1916-2012. Values of relative SPR above 1.0 reflect harvests in excess of the current overfishing proxy.




Figure 1-30. Estimated coastwide total annual catch of aurora rockfish in 2002-2012 relative to the proposed 2015 OFLs and 2015 ABCs (summed north and south of 40°10' N lat. to compare to coastwide catches).

## Blackgill Rockfish

### Distribution and Life History

Blackgill rockfish (*Sebastes melanostomus*), also known at times as blackmouth rockfish or deepsea rockfish, range from at least central Vancouver Island to central Baja California (Love, *et al.* 2002). However, the species is relatively uncommon north of Cape Mendocino and occurs in the greatest densities in the Southern California Bight (SCB). The name very accurately describes the most identifying characteristic of adult blackgill rockfish, in that they have black pigmentation on the rear edge of their gill cover, as well as in the fold above the upper jaw and inside of the mouth. The rest of the fish appears pink

with brown and white blotches underwater, or reddish with distinct brown saddles upon capture. It is a medium-sized (to about 62 cm maximum length) and deep bodied species. Additional descriptions and meristics can be found in Love et al. (2002) for adults and Moser (1996) for larvae and juveniles.

Hyde and Vetter (2007) did not find any evidence for close molecular or evolutionary relationships between blackgill and other rockfish species. Blackgill were found to be moderately related with several other slope or deep shelf species (*S. aurora*, *S. phillipsi*, *S. gilli*, *S. diploproa*, and *S. melanosema*) as well to a suite of mostly rare and poorly known species from the Gulf of California (*S. sinensis*, *S. peduncularis*, and *S. cortezi*) or southern California.

Blackgill are a slope rockfish species, and are generally rare in waters less than 100 meters and most abundant in waters between 300 and 500 meters depth. Love et al. (2002) report a depth distribution of 87 to 768 meters; however, from ten years of data from the NWFSC combined trawl survey, only one haul greater than 600 meters encountered blackgill (that tow was at 647 meters) and the shallowest fish was encountered at 133 meters. Survey data suggest that smaller fish tend to be encountered in shallower water and larger fish in deeper water; survey data also suggest few small fish in waters north of Cape Mendocino. Juveniles are often seen over soft bottom habitats with low relief. Adults are usually associated with high relief rocky outcrops, canyons or deep rock pinnacles, although fishermen often report taking them in midwater (Kronman 1999; Love, *et al.* 2002).

Little is known about the population structure of blackgill rockfish. Like most rockfish, larvae and juveniles circulate in the plankton for 3-4 months. Love et al. (2002) report that some juveniles may be pelagic for up to 7 months; however, this may be atypical. Thus, like most shelf and slope species, blackgill likely disperse over fairly long distances before settling to the bottom. Abundance south of the U.S./Mexico border is uncertain, but there appear to be substantial numbers and catches of blackgill in many areas, and pelagic juveniles have been found as far south as Punta Abreojos, in southern Baja California (Moser and Ahlstrom 1978). The CalCOFI Ichthyoplankton survey has been used to develop or explore indices of relative abundance for several rockfish species for which larvae can be morphologically identified to species (Moser, *et al.* 2000), and such indices have been used as relative abundance indices for assessments of bocaccio (Field, *et al.* 2009) and shortbelly rockfish (Field, *et al.* 2008) as well as northern anchovy (Jacobson and Lo 1994), Pacific sardine (Hill, *et al.* 2008), and California sheephead (Alonzo, *et al.* 2008). Unfortunately, blackgill rockfish is not among the species that have been historically sorted to the species level using morphological methods, although recent developments have led to the potential to use genetic methods to identify historical and contemporary *Sebastes* from the ichthyoplankton archives (e.g., (Taylor, *et al.* 2004), J. Hyde, FRD/SWFSC, unpublished data). Thus, it is possible that these collections could provide relative abundance information from past and contemporary monitoring programs.

Moser and Ahlstrom also found that blackgill represented approximately 16% of the total number of rockfish specimens encountered in a series of midwater trawls for late larvae and juvenile stage rockfish done in the early 1970s (prior to most historical exploitation). By contrast, from ongoing pelagic juvenile surveys run by the Fisheries Ecology Division used to develop juvenile (pre-recruit) indices for some species, blackgill rockfish comprised only about 3% of juveniles collected from the southern California region from 2004 through 2010 (K. Sakuma and J. Field, unpublished data as cited in Field and Pearson (2011)). However, these results are not likely to be comparable unless seasonal and depth of survey efforts are accounted for; the Moser and Ahlstrom (1978) study in particular fished depths ranging from 0 to 600 meters using an Isaacs-Kidd midwater trawl, while the FED survey uses a considerably larger (modified Cobb) midwater trawl and typically only fishes at 30 meters headrope depth. There is at least some potential to consider relative abundance indices of age-0 juveniles from the FED/SWFSC survey in the future, although given the very slow growth and difficulty in ageing of blackgill rockfish, it is unlikely that validation of survey indices or improved understandings of high frequency variation in year class strength will be of substantial near term benefit to the model.

Nearly 2/3rds of all U.S. landings are from waters south of Point Conception, for which blackgill accounted for as much as 20 to 30% of total *Sebastes* landings in the SCB during the 1980s, when deep water fixed gear fisheries rapidly expanded (more details in catch history section). Nearly all of the remaining landings took place between Conception and Cape Mendocino, such that less than 1.3% of historical California landings have come from waters north of Cape Mendocino. Landings in Oregon waters are even less, and only trace landings of blackgill are reported from Washington waters. Trawl survey abundance data (discussed later in the document) are consistent with these results, although they represent the period following the greatest extent of exploitation: surveys that took place from the 1970s through the late 1990s had virtually no coverage in southern waters where blackgill are the most abundant.

Blackgill rockfish have among the deepest distribution of all of the California Current *Sebastes* (although the three *Sebastobus* species are common at considerably greater depths), and live at the edge of the low oxygen (hypoxic) conditions that characterize the slope waters of the California Current. Below these depths, species diversity declines to a smaller suite of species that have adapted to cope with low oxygen waters, notably the DTS complex species (Dover sole, thornyheads and sablefish), which have evolved a range of adaptive strategies including metabolic suppression, slow growth rates, late ages at maturity, and ambush (rather than active searching) predation methods (Childress and Seibel 1998; Jacobson and Vetter 1996; Koslow, *et al.* 2000; Vetter and Lynn 1997). These low oxygen waters, known as the oxygen minimum zone (OMZ), are a natural feature of the Eastern Pacific Rim and other regions characterized by high surface productivity and/or the upwelling of oxygen-poor source waters (Helly and Levin 2004). The California Current has a relatively deeper OMZ than the Equatorial Eastern Tropical Pacific (ETP) or the Humboldt Current (Helly and Levin 2004), with the zone starting at approximately 500 to 600 meters depth in the waters off of southern and central California. The observation that blackgill are likely the most deeply distributed medium-size *Sebastes* (at least in southern California Current waters) suggests that they have adapted to live on the edge of the OMZ, where oxygen availability is rapidly declining relative to shelf waters, although no *Sebastes* species appears able to tolerate the very low oxygen conditions within the OMZ itself.

Seibel (2011) describes two oxygen thresholds that are temperature dependent (as opposed to species or situation-specific), one in which virtually all species are capable of physiologically adjusting or adapting to declining oxygen availability, and a second for which no further adjustment or adaptation in aerobic O<sub>2</sub> utilization is possible. Seibel (2011) describes this latter threshold as one at which “organisms that are not specifically adapted to low O<sub>2</sub> will suffer physiological stress and eventual death.” Importantly, this threshold falls just below the currently observed oxygen levels throughout the slope waters of much of the California Current, implying that any expansion of the OMZ in this region is likely to have tremendous impacts on the vertical distribution of populations and the species composition of ecosystems. Equally importantly, there is already some evidence of a shoaling (shallowing) of the depth of the OMZ throughout the California Current (Bograd, *et al.* 2008; Whitney, *et al.* 2007), with Bograd *et al.* (2008) reporting oxygen declines of 20-30% at depths of approximately 300 to 500 meters in the waters of the Southern California Bight, the region in which most of the blackgill biomass resides. A shoaling of the OMZ has been predicted to be a likely or plausible response to global climate change due to the fact that oxygen is less soluble in warmer waters, and warming is also expected to increase stratification in the upper ocean, which will both reduce oxygen supply and increase oxygen demand at depth (Keeling, *et al.* 2010; Sarmiento, *et al.* 1998; Seibel 2011).

For blackgill rockfish, it is the shoaling of the OMZ at depth that is likely to be the greatest long-term threat, as such a shoaling would likely represent a severe compression of the available habitat for this species. McClatchie *et al.* (2010) evaluated potential scenarios for hypoxia to impact the habitat of cowcod (*Sebastes levis*), a rebuilding shelf species that is a focus of management in the SCB. They found that as much as 37% of deep (240-350 m) cowcod habitat is currently affected by hypoxia, but that if the current trends of

a shoaling OMZ continue for 20 years, this could increase to 55% of deep habitat, as well as an additional 18% of habitat in the 180 to 240 m depth range. These numbers would presumably differ substantially for blackgill rockfish, which have a very different (considerably deeper) distribution; due to their proximity to the OMZ, they may be at considerably greater risk to the longer-term impacts of shoaling. Moreover, changes in the characteristics and dynamics of the OMZ could lead to changes in the forage base for blackgill, which are described as foraging primarily on mesopelagic fishes which undergo diel migrations from the edge of the OMZ to surface waters in order to feed.

As previously mentioned, blackgill have been described as having a strong affinity for deep water habitat, particularly around offshore banks, canyons and areas of high depth gradients. They have been described as feeding on small mesopelagic fishes, such as myctophids and bathylagids (Love, *et al.* 2002). Isaacs and Schwartzlose (1965), Genin *et al.* (1988), Koslow (2000) and Genin (2004) describe the mechanisms by which vertical migrants, such as zooplankton and mesopelagic fishes, become trapped by topographic features. High densities of deep water adapted resident species are consequently found in the relatively small, confined areas where these diurnally-migrating prey become aggregated. Such observations are consistent with the reports by fishermen of isolated deep banks, pinnacles or other habitat features often hosting very large numbers of fish over a relatively small spatial range, such that vertical hook and line gear (which can be more precisely targeted at small habitat features) is the gear of choice for targeting these species (as opposed to horizontal, or set, hook and line gear often used to target species in deeper slope waters, such as sablefish and thornyheads, which tend to be more widely dispersed).

With respect to predators and predation mortality, it is likely that sablefish and shortspine thornyheads are among the most important predators of blackgill rockfish. Both species are large (up to 100 and 75 cm, respectively, although individuals greater than 80 or 65 cm of either species are uncommon) and largely piscivorous ambush predators that are typically (along with longspine thornyhead and Dover sole) the most abundant and commercially important groundfish in the continental slope ecosystem (Lauth 2000). Food habits information for adult sablefish found that *Sebastolobus* and *Sebastes* species, particularly *Sebastolobus altivelis*, are key prey items, representing 15% to 30% of total prey by volume (Buckley, *et al.* 1999; Laidig, *et al.* 1997). Similarly, shortspine thornyhead preyed heavily on *S. altivelis*, unidentified *Sebastes*, and other fishes (Buckley, *et al.* 1999). Although no *S. melanostomus* were conclusively identified in either study, other slope rockfish species (*S. crameri*, *S. diploproa*, and *S. alutus*) were. The lack of specimens is likely due to both studies' focused sampling in northern California, Oregon and Washington slope waters, rather than the south-central and southern California waters in which *S. melanostomus* are most abundant.

### **Stock Status and Management History**

Blackgill rockfish have historically represented a minor part of California rockfish landings north of Point Conception, but a substantial fraction of landings occur south of Conception. Based on consultations with fishery participants, Butler *et al.* (1999a) and Kronman (1999) defined the southern California targeted fishery for blackgill rockfish as being a relatively recent phenomenon. Although longline fishing had long been the primary means of catching rockfish in southern California waters, increased participation and declines in the catches of many highly desired shelf species (such as vermilion rockfish and cowcod) contributed to a gradual shift in effort towards deeper and more offshore waters. Moreover, improvements in technology and gear (such as LORAN, affordable acoustic systems, electric line haulers) helped ease the difficulties of fishing (and relocating good fishing sites) in deeper waters. Additionally, set nets (gillnets) also began to be deployed at a larger scale in southern California in the 1970s and 1980s, often targeting deep reefs for large bocaccio, cowcod, blackgill, bank and other rockfish species.

Such developments seem to have been associated with a geographic expansion of the regions fished, such that fishing locations were sequentially depleted and new fishing locations discovered and developed over time. The first stock assessment for blackgill rockfish (Butler, *et al.* 1999a) noted that there was significant evidence for sequential depletion of blackgill rockfish in localized areas. This included reports from fishery participants that many pinnacles or other fishing sites that routinely yielded 20,000 pounds of blackgill per trip in the early days of the fishery were now only yielding 500 or so pounds per trip and were often covered with lost gear. Similarly, in a review of historical southern California fisheries, Kronman (1999) also documented the rapid growth and development of the blackgill fishery specifically as one in which fishermen would often “completely decimate” rockfish spots with deep fishing vertical line gear, based on the accounts of the participants themselves. Consequently, there was an ongoing shift to newer fishing spots, generally further offshore and to greater depths, as well as greater experimentation with alternative gears and target species.

These observations suggest the potential for a situation in which the stock may have undergone the “sequential depletion” of biomass from available habitat patches. If so, this would suggest that a traditional (non-spatial) stock assessment assumption of evenly distributed fishing mortality across space is substantially flawed. In fact, if the fishery were sequentially depleting specific areas, the length frequency information would not be likely to suggest a shift to smaller fish over time as the length frequencies could essentially reflect “unfished” population structure for the duration over which the new habitats were discovered and exploited. The consequences of failing to recognize such patterns can lead to overexploitation and collapse, and such processes have been described for several marine invertebrate populations (Karpov *et al.* 2000, Orensanz *et al.* 2000) as well as temperate water reef fishes (Epperly and Dodrill 1995, Rudershausen *et al.* 2008). Ongoing efforts to analyze historical block summary data have the potential to identify such shifts and consider whether such factors are likely to be important for west coast groundfish species such as blackgill, as well as to determine whether there is sufficient data to estimate spatial effects or develop spatially-explicit models more capable of accounting for such factors.

Management of blackgill rockfish has generally not been to the species level, but rather as part of the “*Sebastes* complex” in the Pacific Fishery Management Council era (prior to which management was under the direction of the California Department of Fish and Game). The PFMC allowable biological catches (ABC) of blackgill have historically been grouped together with eleven other species of minor rockfishes called “remaining rockfish” and all “other” rockfish. The PFMC historically used trip limits, and later cumulative trip limits (over set time periods), to slow the pace of harvest based on allowable biological catch and to promote a year-round fishery. For all commercial gear types, the limits were initiated in 1983 when the PFMC imposed a monthly limit of 40,000 pounds per trip for the entire coastwide *Sebastes* complex, a limit that stayed in place through 1990. After recognizing the differential spatial distribution of the remaining rockfishes and the fisheries that target them, harvest limits on both open access and limited entry fisheries were divided between the northern and southern *Sebastes* complexes, and trip limits began to be implemented at variable levels over both time (month and year) and space (north and south of Mendocino), often with species-specific limits in addition to the overall limit on *Sebastes* catches. Although early limits applied to both trawl and fixed gears, beginning in 1995 fixed gear limits (hook and line and pot, primarily, as gill nets were phased out through the 1990s) were set to 10,000 lbs of *Sebastes* per trip, which persisted through the 1990s.

Consequently, prior to 1999 cumulative trip limits had been historically high relative to landings of blackgill rockfish from individual trips, and unlikely to have impacted fishing for blackgill and catches. Limits were dramatically reduced in 1999 for the southern *Sebastes* complex; 2-month cumulative limit of 3,500 pounds for limited entry and 3,600 pounds per month for open access. Since 2000, blackgill has been managed as part of the Minor Slope Rockfish sub-group, with limits ranging from 3,000-50,000 pounds per 2 months; Tables 1-3 show the trip limits implemented since 2000 for this complex for the limited entry trawl, limited entry fixed gear and open access fixed gear fisheries.

In 2001 the Cowcod Conservation area was established outside of 20 fathoms and directly excludes directed groundfish fishing from an expansive area in the Conception and southern Monterey INPFC areas.<sup>8</sup> This regulation has had a tremendous impact on the southern fixed gear fleet that targets blackgill, as the deep offshore banks and features that characterize the CCAs in deep water are optimal habitat for this species. By contrast, the shelf closures (rockfish conservation areas) implemented to protect rebuilding shelf species (such as bocaccio, cowcod, canary and widow rockfish) have presumably had a negligible direct effect, as the depths closed in the RCAs do not encompass the depths at which most blackgill are encountered. Such measures may have had an indirect effect, by virtue of shifting trawl effort to deeper waters, although for much of California the overall effect has been a sharp decline in active participation in the trawl fishery more generally.

The first assessment for blackgill rockfish was conducted in 1998 and estimated stock depletion was between 40 and 54 percent of its unfished equilibrium at the start of 1998 (Butler, *et al.* 1999a). That assessment assumed a unit stock in southern and central California (Conception INPFC area) and was based on a stock reduction analysis assuming constant recruitment. The dynamics of the simple model were tuned to average mortality rates from catch curves and landings data. Fishery selectivity was assumed to mirror maturity at size/age; trends in fishable/mature biomass were then estimated.

A second blackgill rockfish stock assessment was completed in 2005 indicating a stock depletion of 52 percent (Helser 2006). This assessment expanded the geographic range of that in Butler *et al.* (1999a), including both the Monterey and Conception INPFC areas, where over 90 percent of the landings have occurred. The assessment was based on catch and length composition data from commercial fisheries and indices of relative abundance and size composition from the AFSC shelf trawl survey and the AFSC slope survey. The modeling approach included fishery and survey length compositions to explicitly estimate selectivity. The assumed natural mortality rate was identified as a key axis of uncertainty for this stock.

The most recent blackgill rockfish assessment, conducted in 2011 for the stock south of 40°10' N lat. (Field and Pearson 2011), estimated the stock was below target with a depletion of 30 percent of its unfished biomass at the start of 2011 (Figure 1-31). The spawning output of blackgill rockfish was at high levels in the mid-1970s, but began to decline steeply in the late 1970s through the 1980s, consistent with the rapid development and growth of the targeted fishery. The biomass reached a low of approximately 18 percent of the unfished level in the mid-1990s. Since that time, catches have declined and spawning output has increased. The estimated depletion level in 2011 is 30.2 percent.

Catch data used in the assessment are generally reliable throughout the time period, although there is a lot of uncertainty in catch data prior to the early 1980s. Ageing is very difficult for this species, which appears to have highly variable size at age, as well as apparent regional differences in growth rates and potentially other life history traits. The lack of a reliable, long-term, fishery-independent survey index that reflects abundance from the entire range of the stock is problematic. In general, natural mortality and growth parameters comprised the greatest contribution to model uncertainty.

---

**8** As the current trawl survey also excludes this region from trawl gear impacts, the area of the CCAs is shown in later maps of survey CPUE for blackgill rockfish, in Figure 13




Figure 1-31. Time series of estimated spawning output and depletion of blackgill rockfish south of 40°10' N lat., 1950-2011.

### Stock Productivity

In the 2013 assessment, the Beverton-Holt model was used to describe the stock-recruitment relationship. The log of the unexploited recruitment level was treated as an estimated parameter; recruits were taken deterministically from the stock-recruit curve. Recruitment deviations were not estimated, as the lack of obvious cohorts in either age or length data and the high degree of ageing uncertainty make plausible estimates unlikely. The estimated recruitment is projected to be at relatively high levels due to the fixed value of steepness ( $h = 0.76$ ); this trend, however, is consistent with the trends from the survey data.

Blackgill rockfish have a relatively high potential vulnerability to overfishing ( $V = 2.08$ ) driven by a combination of low productivity ( $P = 1.22$ ) and relatively high susceptibility to being caught in the fishery (Table 1-2). The low productivity is due to the stock being long-lived (max. age = 87 yrs; (Love, *et al.* 2002)), with late maturation, and relatively low natural mortality (Table 1-5).

### Fishing Mortality

Catches of blackgill rockfish primarily occur in the Southern California Bight south of Point Conception (34°27' N. lat.) where the species is caught in both directed fixed gear (hook-and-line) and historically, gillnet fisheries. Landings of this species are estimated to have risen slowly from very low levels (approximately 20-30 mt) in the 1950s, and then climbed rapidly in the 1970s and 1980s as improvements in technology and declines in other target species led fishermen to target blackgill rockfish in deeper and more offshore waters. Landings peaked in the mid-1980s at just over 1,000 mt, but have declined to approximately 100 mt to 150 mt in recent years.

The 2011 depletion estimate indicated the stock was in the precautionary zone compelling the Council to reduce impacts to prevent overfishing and allow the stock to rebuild back to its target biomass. The Council and NMFS implemented stringent harvest guidelines of 106 and 110 mt for 2013 and 2014 respectively, corresponding to calculated 40-10 reductions. Specifying HGs created a sorting requirement for the stock to allow better inseason catch monitoring. The Council further established an apportionment of the non-



trawl allocation of 60 percent to limited entry and 40 percent to open access fixed gears, which reflects the historical distribution of catch between the limited entry and open access fixed gear sectors from 2005-2010. Non-trawl landing limits for blackgill south of 40°10' N lat. were reduced beginning in 2013 to prevent targeting of the stock. Cumulative landing limits for blackgill south of 40°10' N lat. were reduced from 40,000 lbs/2 months for slope rockfish including blackgill to 40,000 lbs/ 2 months with a sublimit of 1,375 lbs/2 months for blackgill for the limited entry fixed gear sector. Open access cumulative landing limits south of 40°10' N lat. were reduced from 10,000 lbs/2 months for slope rockfish including blackgill to 10,000 lbs/2 months with a sublimit 475 lbs/2 months for blackgill. While final catch accounting of groundfish in 2013 groundfish fisheries will not be available until the end of 2014, landed catch of blackgill has been reduced significantly and fishermen report blackgill targeting is no longer occurring.




Figure 1-32. Time series of estimated summary harvest rate for the blackgill rockfish south of 40°10' N lat., 1950-2011. The dashed line is the harvest rate at the overfishing  $F_{MSY}$  proxy.



Figure 1-33. Estimated coastwide total annual catch of blackgill rockfish in 2002-2012 relative to the proposed 2015 OFLs, 2015 ABCs, and 2015 ACLs (summed north and south of 40°10' N lat. to compare to coastwide catches).

## Rougheye/Blackspotted Rockfish

### Distribution and Life History

Rougheye rockfish (*Sebastes aleutianus*) and blackspotted rockfish (*S. melanostictus*) are slope rockfish that share broad overlap in their depth and geographic distributions from the Eastern Aleutian Islands along the North American continental margin to southern Oregon, with blackspotted rockfish's range extending east beyond the Aleutian chain to the Pacific Coast of Japan (Gharrett, *et al.* 2005; Hawkins, *et al.* 2005; Orr and Hawkins 2008). It is very difficult to visually distinguish between the two species and they have been persistently confused in surveys and catches. Off the U.S. west coast the two species have been reported as rougheye rockfish or in an even more generic rockfish category. It has only been from recent genetic studies in the early 2000s that the two separate species have been identified and described (Orr and Hawkins 2008).

Both species are encountered at depths shallower than 100 m to at least 439 m, however, blackspotted rockfish tend to be more prevalent in deeper waters (Hawkins *et al.* 2005, Orr and Hawkins 2008). Genetic information is not available to provide positive species identification in historical survey and landings information, but these data indicate that density of the nominal rougheye rockfish complex decreases sharply south of the Oregon-California border at 42° N lat. Studies suggest that rougheye rockfish account for a greater proportion of the species complex along the coast of Washington and Oregon than in Alaskan waters (Gharrett, *et al.* 2005; Hawkins, *et al.* 2005; Orr and Hawkins 2008). Recent discussions with port samplers in southern Oregon suggest that both rougheye and blackspotted rockfish are encountered with some regularity in the commercial trawl and fixed-gear landings in Charleston, Port Orford, and Brookings, with blackspotted rockfish composing approximately one third to one half of identified specimens (C. Good and N. Wilsman, ODFW, pers. comm. as cited in Hicks *et al.* 2013).

The west coast of the U.S. is the southern portion of the range of rougheye rockfish, and it is likely that the population north of the U.S.-Canada border is not a separate stock. The connectivity of rougheye populations throughout its range is unknown.

Compared with other rockfish species on the west coast of the U.S., rougheye rockfish life-history is poorly described and the recent resurrection of the two species classification (rougheye and blackspotted rockfishes) has further complicated the understanding of life-history characteristics. Rougheye rockfish are often associated with boulders and steep habitats, and are typically found alone or in small aggregations (Love, *et al.* 2002). Younger fish may school and are often found in shallower waters on the shelf, and larger fish may form larger aggregations in the Pacific Northwest during the autumn and winter.

Rougheye rockfish give birth to live young with larvae released between February and June and at lengths between 4.5-5.3 mm (Love, *et al.* 2002). There are no studies on the fecundity of rougheye rockfish on the west coast of the U.S.


A wide range of prey items make up the diet of rougheye rockfish. Crangid and pandalid shrimps make up the majority of their diets, and larger individuals, greater than 30 cm, feeding upon other fishes (Love 2011). They are also known to feed upon gammarid amphipods; mysids, crabs, polychaetes, and octopuses (Love 2011; Love, *et al.* 2002).

## Stock Status and Management History

Rougheye and blackspotted rockfish (henceforth denoted as rougheye) are landed as part of the minor slope rockfish complexes north and south of 40°10' N lat.; however, they are rarely caught in the south. The historical reconstruction of landings for rougheye rockfish suggests that fixed gear fisheries have caught rougheye rockfish since the turn of the 20th century and landings in the trawl fishery are estimated to have increased into the 1940s. Landings remained relatively constant throughout the 1950s and into the 1960s before the foreign trawl fleet increased catches into the 1970s. The declaration of the EEZ resulted in the buildup of a domestic fleet and landings increased rapidly into the late 1980s and early 1990s. Subsequently, landings declined in the late 1990s and have been between 100 and 200 mt in recent years. Trawl, longline, and Pacific whiting at-sea trawl fisheries make up the majority of the catch.

Rougheye rockfish are a desirable market species and discarding has been low, historically. However, management restrictions (e.g., trip limits) have resulted in increased discarding since 2000. Trawl rationalization was introduced in 2011, and since then very little discarding of rougheye rockfish has occurred.

Hicks et al. (2013) conducted the first assessment of the U.S. west coast stock of rougheye and blackspotted rockfish as a complex of two species. The coastwide population was modeled assuming parameters for combined sexes (a single-sex model) and assuming removals beginning in 1916. The predicted spawning biomass from the base model generally showed a slight decline over the entire time series with a period of steeper decline during the 1980s and 1990s. Since 2000, the spawning biomass has stabilized and possibly increased because of reduced catches and above average recruitment in 1999. The 2013 spawning biomass relative to unfished equilibrium spawning biomass was estimated to be 47 percent of its unfished equilibrium at the start of 2013. The stock has been estimated to be healthy throughout the time series in the new assessment (Figure 1-34).




**Figure 1-34. Time series of estimated spawning biomass and depletion of rougheye/blackspotted rockfish, 1916-2013 (from Hicks et al. 2013).**

## Stock Productivity

The parameter for steepness of the spawner-recruit relationship was fixed at 0.779 in the 2013 assessment based on a steepness meta-analysis for west coast rockfishes (Jim Thorson, NWFSC). There is little information regarding recruitment prior to 1980, and the uncertainty in these estimates is expressed in the assessment. Estimates of recruitment appear to oscillate between periods of low and high recruitment. The four largest recruitments were estimated in 1999, 1998, 2001, and 1988, and the four smallest recruitments were estimated in 2002, 2006, 2005, and 1995 (Figure 1-35).

Rougheye rockfish have the highest potential vulnerability to overfishing ( $V = 2.27$ ) driven by a combination of low productivity ( $P = 1.17$ ) and relatively high susceptibility to being caught in the fishery (Table 1-2). Despite this, the 2013 assessment estimated the stock to be above the  $B_{40\%}$  spawning biomass target. The low productivity is due to the stock being long-lived (max. age = 205 yrs; (Love, *et al.* 2002)), with late maturation, and relatively low natural mortality (Table 1-5).



**Figure 1-35. Time series of estimated age-0 recruits of rougheye/blackspotted rockfish on the U.S. west coast, 1980-2013(from Hicks et al. 2013).**

## Fishing Mortality


Rougheye rockfish are not often targeted by a specific fishery, but are desirable and marketable, thus are typically retained when captured. They are often captured in bottom trawl, mid-water trawl, and longline fisheries. Small numbers have been observed in pot, shrimp, and recreational fisheries.

After many attempts to start trawl fisheries off the west coast of the United States in the late 1800s, the availability of the otter trawl and the diesel engine in the mid-1920s helped the trawl fisheries expand (Douglas 1998). Trawl fisheries really became established during World War II when demand increased for shark livers and bottomfish. A mink food fishery also developed during World War II (Jones and Harry 1960). Foreign fleets began fishing for rockfish in the mid-1960s until the EEZ was implemented in 1977 (Rogers 2003b). Since 1977, landings of rockfish were high until management restrictions were implemented in 2000. Longline catches of rougheye rockfish are present from the turn of the century and continue in recent years, targeting sablefish and halibut.


A long-term directed fishery has not occurred for roughey rockfish and historical discarding practices are not well known. Roughey rockfish inhabit deeper water as adults, which were fished less often historically.

Throughout the 1980s and 1990s exploitation rates (1-SPR) were mostly above target levels (Figure 1-36). Recent exploitation rates on roughey rockfish were predicted to be near target levels.

While stock-specific OFLs/ABCs were not historically set for roughey rockfish specifically, the reauthorized Magnuson-Stevens Act of 2006 and FMP Amendment 23 required OFLs for all species in a management plan, including those managed in stock complexes. The first OFL contributions were calculated using DB-SRA and provided in 2011. The 2015 and beyond OFLs are projected from the 2013 assessment. Recent catches since 2002 have been above the new 2015 OFL since 2008 (Figure 1-37).



**Figure 1-36. Time-series of estimated summary harvest rate for the west coast stocks of roughey and blackspotted rockfish, 1916-2012. The dotted line is the harvest rate at the overfishing  $F_{MSY}$  proxy.**



**Figure 1-37. Estimated coastwide total annual catch of rougheye and blackspotted rockfish in 2002-2012 relative to the proposed 2015 OFLs and 2015 ABCs (summed north and south of 40°10' N lat. to compare to coastwide catches).**

## Sharpchin Rockfish

### Distribution and Life History

Sharpchin rockfish (*Sebastes zacentrus*) range from the western Aleutian Islands (Attu Is.) to Southern California, though the core range is northern California to the Gulf of Alaska in waters between 100 m and 300 m (Love, *et al.* 2002). There is no indication of population structure in sharpchin rockfish. Sharpchin rockfish is a smaller-sized rockfish that inhabits waters up to 500 m, typically over muddy-rock habitats.

Mitochondrial DNA analyses indicate sharpchins are related mostly to harlequin, Puget Sound, and pygmy rockfishes (Love, *et al.* 2002).

Sharpchin rockfishes live to at least 58 years (Love, *et al.* 2002). Females attain a larger size than males with a reported maximum size of 45 cm (Love, *et al.* 2002). Off Oregon and Washington, the size at 50% maturity for females is 22cm with all females being mature at 30 cm. The size at 50% maturity is larger for samples farther north with 25 cm and 28 cm reported off British Columbia and the Gulf of Alaska, respectively. Larval releases occur from March to June off California and Oregon and during July off British Columbia.

Sharpchin eat a variety of prey including krill, shrimps, gammarid amphipods, copepods, and small fishes.

### Stock Status and Management History

Sharpchin rockfish are managed in the northern and southern Slope Rockfish complexes.

A new data-moderate assessment of sharpchin rockfish was conducted in 2013, which indicated the stock was healthy with a depletion of 68% at the start of 2013 (Cope, *et al.* 2014). The SSC recommended the 2013 assessment be used for setting harvest specifications and upgraded the stock from a category 3 to a

category 2 stock. The 2015 and 2016 OFLs are 416 and 404 mt, respectively. The coastwide OFLs were apportioned 80% to the north of 40°10' N lat. and 20% to the south to determine the OFL contributions to the Slope Rockfish complexes based on swept area biomass estimates from the triennial survey.

## **Stock Productivity**

A high steepness of 0.77 was estimated in the 2013 sharpchin rockfish assessment, near the prior used in the assessment.

Sharpchin have a relatively low productivity ( $P = 1.36$ ) and a relatively high vulnerability ( $V = 2.05$ ) to overfishing based on the PSA scores derived prior to the 2013 assessment (Table 1-2).

## **Fishing Mortality**

Sharpchin are not a major commercial target, though they are taken in large numbers and commonly seen in trawls that target Pacific ocean perch. They are taken most commonly off Oregon and Washington with POP, darkblotched, splitnose, and yellowmouth rockfish. While they are common in west coast bottom trawl catches, their smaller size makes them less valuable than the larger rockfish species. They are rarely taken in recreational fisheries.

### **1.1.5.4 Other Flatfish**

The Other Flatfish complex contains most of the flatfish species managed in the Groundfish FMP (with the exception of arrowtooth flounder, Dover sole, English sole, petrale sole, and starry flounder). These species include butter sole (*Isopsetta isolepis*), curlfin sole (*Pleuronichthys decurrens*), flathead sole (*Hippoglossoides elassodon*), Pacific sanddab (*Citharichthys sordidus*), rex sole (*Glyptocephalus zachirus*), rock sole (*Lepidopsetta bilineata*), and sand sole (*Psettichthys melanostictus*).

## **Pacific Sanddabs**

### **Distribution and Life History**

Pacific sanddab (*Citharichthys sordidus*) is a left-eyed flounder of the family *Paralichthyidae* and is widely distributed along the Pacific west coast from the Bering Sea to Cabo San Lucas, Baja California (Arora 1951; Hart 1988; Kramer and O'Connell 1995; Love, *et al.* 2005; Miller and Lea 1972; Rackowski and Pikitch 1989). Early studies reported that the species is the most abundant in the north-central portion of California from Eureka to San Francisco, but were also fairly common in southern California (Rackowski and Pikitch 1989). Early studies also reported that the species is usually found at depths between 18m and 275m and most commonly found at depths between 35m and 95m (Arora 1951; Demory 1971; Hart 1988; Miller and Lea 1972; Roedel 1953). On Oregon's continental shelf, Pacific sanddab is the most abundant small flatfish on sandy-bottom in the depths between 74 and 102 m (Pearcy 1978). Young Pacific sanddab (ages 0 and 1) are also found to be concentrated in the same depth range (Donohoe 2000). Pacific sanddab was also found to be relatively more abundant in shallow waters at higher latitudes (Chamberlain 1979).

Pacific sanddab are generally not considered a primary target for commercial fisheries along the U.S. west coast, but they are nevertheless highly prized by the commercial and recreational fisheries for their excellent edibility (CDFG 2001), and have long been an important component of the nearshore flatfish fishery, commanding a high price in fresh fish markets (Arora 1951). Commercial catches of Pacific sanddab were mostly from bottom trawl fisheries, and there is a long history of catches. Recreational catches of Pacific

sanddab are from the hook and line fishery and most of this catch is from southern California waters. Some recreational anglers target Pacific sanddab in southern California, mostly from small boats and CPFVs (CDFG 2001).

Pacific sanddabs can grow to 35cm in length. They are sexually dimorphic, with females attaining larger sizes than males. Analysis of growth rates for both sexes between the southern and northern areas (divided at the California-Oregon border at 42° N lat.) showed no significant difference in growth rates for both sexes between the two areas.

There are no genetic or tagging studies informing stock structure of Pacific sanddab along the U.S. Pacific coast. Bottom trawl surveys in recent years (both NWFSC and triennial surveys) showed that Pacific sanddab are commonly caught along the coastal areas of all U.S. waters.

Pacific sanddabs play an important role in the coastal ecosystems in the U.S. waters, particularly because they are a relatively abundant species and are important prey items to a wide range of marine predators, including piscivorous fishes, sea mammals, and sea birds (Field, *et al.* 2006; Levin, *et al.* 2006).

### **Stock Status and Management History**

Pacific sanddabs have been under federal management since the implementation of the groundfish FMP in 1982 and managed within the Other Flatfish complex of unassessed flatfish species. The management performance in recent years for Pacific sanddab has been good; the average 2005-2012 total annual catch has been about 23% of the stock's ACL/OY contribution to the Other Flatfish complex.

A coastwide assessment of Pacific sanddab was done in 2013 indicating the stock was at 95.5% of its unfished biomass (He, *et al.* 2013). The SSC recommended in 2013 that this assessment not be used for deciding harvest specifications since the scale of the stock's biomass could not be adequately estimated. However, the status estimate was precise enough to conclude the stock was well above the  $B_{MSY}$  proxy of  $B_{25\%}$ . The SSC recommended the stock continue to be categorized as a category 3 stock given the OFL estimate from the assessment depends on the biomass estimate, which was not estimated with adequate precision. The OFL estimate is therefore based on the DB-SRA method used since 2011. The 2015 and 2016 OFL contribution of Pacific sanddab to the Other Flatfish complex is 4,801 mt.

### **Stock Productivity**

A steepness prior of 0.8 was used in the 2013 assessment. Annual recruitment deviations were estimated between 1966 and 2011. Annual recruitment deviations were treated in a log-normal distribution with  $\sigma_R$  fixed at 0.45. Low recruitments occurred from the early 2000s to the mid-2000s. Recruitments in recent years have been at or above the long term average, with a strong recruitment in 2010.

The PSA productivity score of 2.4 indicates a very high relative productivity of Pacific sanddabs. This leads to a very low vulnerability ( $V = 1.25$ ) of potential overfishing for the stock.

### **Fishing Mortality**

There is a long history of commercial catches on Pacific sanddab (Barss 1976). Sette and Fiedler (1928) reported that landings of flatfish in California waters were first reported in 1892. The first available landings of Pacific sanddab in Oregon waters were in 1942 (Karnowski, *et al.* 2012). There were also



commercial catches for mink foods in both California and Oregon waters in the 1950s and 1960s (Best 1959; Best 1961; Nitsos and Reed 1965). Reported total catches of Pacific sanddab were high in the late 1920s. And there was an increasing trend from the 1960s and reached the highest catch level in the late 1990s. Discards of Pacific sanddab in commercial trawl fisheries were high, primarily due to its small size (Sampson 2002). Catches of the species in recent years were in the range of 200 mt and 400 mt, well below the OFL contribution of the stock to the Other Flatfish complex of 4,801 mt.

## **Rex Sole**

### **Distribution and Life History**

Rex sole (*Glyptocephalus zachirus*) is a right-eyed flounder of the family *Pleuronectidae* ranging from central Baja California to the Aleutian Islands and the western Bering Sea. They are common from southern California to the Aleutian Islands. They are distributed over mud and sand bottom habitat in deeper depths, are commonly found in waters up to at least 500 m, and range down to more than 1,100 m.

Rex sole grow slowly and are relatively long-lived for a flatfish with a maximum age of 29 years (Cope, *et al.* 2014). Females grow faster and attain a larger size than males.

### **Stock Status and Management History**

Rex sole are currently managed in the Other Flatfish complex.

A new data-moderate assessment of rex sole using the exSSS model was conducted in 2013, which indicated the stock was healthy with a depletion of 80% at the start of 2013 (Cope, *et al.* 2014). The SSC recommended the 2013 assessment be used for setting harvest specifications and upgraded the stock from a category 3 to a category 2 stock. The 2015 and 2016 OFL contributions of rex sole to the Other Flatfish complex are 5,764 and 3,956 mt, respectively.

### **Stock Productivity**

A steepness prior of 0.89 was estimated in the 2013 assessment, higher than the mean prior value.

The PSA productivity score of 2.05 indicates a high relative productivity of rex sole. This leads to a low vulnerability ( $V = 1.28$ ) of potential overfishing for the stock.

### **Fishing Mortality**

Rex sole are a very commonly occurring species in the fishery-independent trawl surveys and are very accessible to trawl fisheries. Targeting for rex sole in commercial fisheries has varied over the years, with major removals occurring in the mid-1900s to provide feed for mink farms. They have not been targeted heavily in the last few decades. While their flesh is tasty and of high quality, they are small fish with very thin fillets and therefore less desired in commercial markets.

Females are larger than males and are most commonly caught. Males are small enough to escape the minimum mesh size of west coast bottom trawls.

### 1.1.5.5 Other Fish

Beginning in 2015, the Other Fish stock complex is comprised of three species managed as five populations: kelp greenling (*Hexagrammos decagrammus*) (separate populations off WA, OR, and CA), cabezon (*Scorpaenichthys marmoratus*) (off WA), and leopard shark (*Triakis semifasciata*) (primarily off CA). Of these five populations, only kelp greenling off CA, cabezon off WA, and leopard shark have OFL contributions to the Other Fish complex.

The other species managed in the Other Fish complex prior to 2015 are proposed to be managed under stock-specific specifications (spiny dogfish, see section 1.1.4.18) or designated as Ecosystem Component species (see section 1.1.6) beginning in 2015.

A full assessment of the population of kelp greenling occurring in Oregon waters is planned for 2015.

### 1.1.6 Ecosystem Component Species

The Council proposed the following species to be designated Ecosystem Component (EC) species under FMP Amendment 24: big skate, California skate, all other endemic skates, soupfin shark, finescale codling, Pacific grenadier, all other endemic grenadier species, and spotted ratfish (Table 1-8). An EC species can be so designated if it is not targeted, is not subject to overfishing or being overfished in the absence of conservation measures, and not generally retained for sale or personal use. No harvest specifications or management reference points are required for EC species; however, there is a monitoring requirement to determine potential changes in their status or their vulnerability to the fishery. An unexpected increasing catch trend infers an EC species' vulnerability to overfishing may have increased, compelling a consideration to reclassify the stock as "in the fishery". Any designation of a species as an EC species or a change from an EC designation to a species considered to be "in the fishery" requires an FMP amendment. The GMT analyzed historical data to determine those species in the Other Fish complex (previous to the restructuring of the complex proposed for implementation in 2015) that were in and out of the fishery to inform the decision to designate EC species (see [Agenda Item H.4.b, GMT Report 2, November 2013](#)).

**Table 1-8. Groundfish species proposed for designation as Ecosystem Component Species under FMP Amendment 24.**

Common Name	Scientific Name
Aleutian skate	<i>Bathyraja aleutica</i>
Bering/sandpaper skate	<i>B. interrupta</i>
Big skate	<i>Raja binoculata</i>
California skate	<i>R. inornata</i>
Roughtail/black skate	<i>Bathyraja trachura</i>
All other skates	Endemic species in the family <i>Arhynchobatidae</i>
Pacific grenadier	<i>Coryphaenoides acrolepis</i>
Giant grenadier	<i>Albatrossia pectoralis</i>
All other grenadiers	Endemic species in the family <i>Macrouridae</i>
Finescale codling (aka Pacific flatnose)	<i>Antimora microlepis</i>
Ratfish	<i>Hydrolagus colliei</i>
Soupfin shark	<i>Galeorhinus zyopterus</i>

## 1.2 Discard Mortality Rates Used to Manage West Coast Groundfish Stocks

Some groundfish species caught in the west coast groundfish fishery are discarded at sea because they are incidentally caught and are not marketable (market-induced discards) or they are caught in excess of allowable cumulative landing limits or are not of a legal size to keep (regulatory discards). The SSC recommended the discard mortality rates by gear type that were modeled in approved stock assessments be used to manage the fishery. Table 1-9 shows the discard mortality rates by commercial gear type used in the most recent assessments for lingcod, longnose skate, sablefish, and spiny dogfish. These discard mortality rates are also applied to estimated discards of these species to estimate total discard mortality when reconciling total mortality in west coast groundfish fisheries.

The GMT recommended using the 50% mortality rate for lingcod discarded in west coast bottom trawl fisheries based on a study that evaluated tow duration and time on deck of trawl-caught lingcod that were ultimately discarded (Parker, *et al.* 2003). The 2009 lingcod stock assessment also modeled the 50% discard mortality rate for discarded lingcod in trawl fisheries (Hamel, *et al.* 2009). The GMT recommended a 7% lingcod discard mortality rate be used for commercial fixed gear fisheries (Table 1-9), as well as discards in recreational fisheries (Table 1-10) based on a study off California evaluating immediate and delayed mortality of lingcod caught using these gears (Albin and Karpov 1996). However Hamel *et al.* (2009) assumed a 5% discard mortality rate in recreational fisheries in the 2009 assessment. This difference has yet to be reconciled.

No studies had been conducted to estimate the mortality of discarded longnose skate or any other skate at the time of the 2007 longnose skate assessment. In tagging studies conducted in Canada (Gordon McFarlane, Pacific Biological Station, Fisheries and Oceans Canada, pers. com. as cited in (Gertseva and Schirripa 2008)), tagged skates were recovered several times in trawl surveys, indicating that skates can survive trawl capture and on-deck sorting time. Anecdotal evidence from commercial fisheries also indicates that skates are generally durable, and can handle capture and release well. However, many factors, such as trawl time, handling techniques, and time spent on the deck certainly affect skate survival. Gertseva and Schirripa (2008) assumed that 50% of commercially-discarded skates die in the 2007 longnose skate assessment.

Sablefish discard mortality rates have been the subject of numerous research studies and analyses supporting historical sablefish stock assessments. Sablefish, lacking a swim-bladder (and therefore the propensity for severe barotrauma), have a very good chance of survival after capture depending on the specific conditions they experience during the process. Generally warmer water results in higher mortality, as the physiological stress of transitioning from very cold bottom temperatures to warmer surface water and air temperatures can be great (Davis, *et al.* 2001). Further, some gears, such as pot and hook-and-line gear are less physically damaging to sablefish than, for example, spending an extended period of hours in a trawl cod-end with a large catch volume. Treatment and handling of captured fish, including time-on-deck is also important for subsequent survival. The GMT reviewed the research studies informing sablefish discard mortality and recommended the mortality rates of 50% for trawl discards and 20% for fixed gear discards as shown in Table 1-9. Stewart *et al.* (2011) assumed the same discard mortality rates by gear type in the 2011 sablefish assessment.

There have been no studies performed on discard mortality of spiny dogfish in the Northeast Pacific Ocean for neither the bottom trawl nor the hook-and-line fleet. In spiny dogfish assessments conducted elsewhere, different values of discard mortality were assumed, from 5% to 50% for bottom trawl and from 6% to 75% for hook-and-line gears, but all sources noted considerable uncertainty in these estimates. Gertseva and Taylor (2011) assumed trawl discard mortality to be 100% and hook-and-line discard mortality to be 50%.

**Table 1-9. Discard mortality rates by commercial gear type used to manage west coast groundfish.**

<b>Stock</b>	<b>Discard Mortality Rate (# dead/# discarded)</b>	<b>Gear Type</b>
Lingcod	50%	Trawl
Lingcod	7%	H&L fixed gear
Longnose skate	50%	Trawl and fixed gear
Sablefish	50%	Trawl
Sablefish	20%	Fixed gear
Spiny dogfish	50%	H&L fixed gear

The GMT analyzed the disposition of observed discards of groundfish species released at the surface of recreational charter fishing efforts off California and Oregon to determine depth-based discard mortality rates using recreational hook-and-line gear. The GMT considered “surface” mortality (i.e., mortality that is observable when a fish is brought to the surface, handled on deck, and thrown back) from charter observations. The GMT then considered short-term, below-surface mortality that has been documented in research trials to a limited extent using underwater cameras or divers. The GMT used a guild-based Generalized Linear Model (GLM) analysis comparing mortality rates of species with similar depth-distributions and vertical orientation in the water column to determine mortality rates for species with limited discard observations. The GMT calculated the upper 95% confidence intervals of surface mortality rates to illustrate the uncertainty associated with GLM predictions. Since upper 95% confidence limits for surface mortality approach 100% at depths greater than 30 fm, mortality beyond this depth was assumed to be 100%. The two exceptions to this approach were yellowtail and black rockfish, given their relatively low mortality rates. The depth-based discard mortality matrix developed by the GMT shows a wide variation in rockfish mortality rates by depth reflecting the diversity of rockfish adaptations to barotrauma (Table 1-10). Yellowtail and black rockfish, which are more pelagic than most of the other rockfish, tend to suffer less barotrauma and therefore exhibit lower surface-release mortality rates. Estimates of surface release discard mortalities for groundfish species that lack a swim bladder (e.g., lingcod and flatfishes) were based on research efforts off California (Albin and Karpov 1996). The resulting depth-based surface release mortality rates for various groundfish species released using recreational hook-and-line gears were implemented in 2009 to determine discard mortalities (Table 1-10).

**Table 1-10. Discard mortality rates by depth of groundfish species released at the surface in west coast recreational fisheries using hook-and-line gear.**

Species Group	Species	Depth Bin			
		0-10 fm	11-20 fm	21-30 fm	>30 fm
Rockfish	Black Rockfish	11%	20%	29%	63%
	Black and Yellow Rockfish	13%	24%	37%	100%
	Blue Rockfish	18%	30%	43%	100%
	Bocaccio	19%	32%	46%	100%
	Brown Rockfish	12%	22%	33%	100%
	Calico Rockfish	24%	43%	60%	100%
	Canary Rockfish	21%	37%	53%	100%
	China Rockfish	13%	24%	37%	100%
	Copper Rockfish	19%	33%	48%	100%
	Gopher Rockfish	19%	34%	49%	100%
	Grass Rockfish	23%	45%	63%	100%
	Kelp Rockfish	11%	19%	29%	100%
	Olive Rockfish	34%	45%	57%	100%
	Quillback Rockfish	21%	35%	52%	100%
	Tiger Rockfish	20%	35%	51%	100%
	Treefish	14%	25%	39%	100%
	Vermilion Rockfish	20%	34%	50%	100%
	Widow Rockfish	21%	36%	52%	100%
	Yelloweye Rockfish	22%	39%	56%	100%
Yellowtail Rockfish	10%	17%	25%	50%	
Other Fish	Cabezon	7%	7%	7%	7%
	California scorpionfish	7%	7%	7%	7%
	Kelp Greenling	7%	7%	7%	7%
	Lingcod	7%	7%	7%	7%
	Pacific Cod	5%	32%	53%	97%
General Cat.	Flatfish	7%	7%	7%	7%
	Sharks and Skates	7%	7%	7%	7%
	Dogfish	7%	7%	7%	7%

The GMT also determined surface-release discard mortality rates by depth for rockfish species caught in nearshore commercial fisheries. The GMT assumed the same discard mortality rates by depth and species for nearshore commercial fisheries using recreational hook-and-line gear (i.e., rod and reel gear) as recommended for recreational fisheries (Table 1-10). The GMT also assumed a 100% discard mortality rate for all rockfish species and depths in nearshore commercial fisheries using non-recreational gears (e.g., longline, dinglebar, etc.). The GMT then weighted the discard rates by depth bin for recreational and non-recreational gear types by the proportion of these gears types deployed in the Oregon nearshore commercial fishery using 2004-2006 Oregon logbook data (Table 1-11). The combined weighted discard rates for all nearshore commercial gears by rockfish species and depth bin were implemented in 2009 (Table 1-12).

**Table 1-11. Proportion of recreational and non-recreational gears used in 2004-2006 Oregon nearshore commercial fisheries based on logbook data.**

Gear Type	Depth Bin		
	0-10 fm	11-20 fm	>20 fm
Recreational	86.6%	72.3%	60.7%
Non-recreational	13.4%	27.7%	39.3%

**Table 1-12. Discard mortality rates by depth of rockfish species released at the surface in west coast nearshore commercial fisheries.**

Species	Depth Bin		
	0-10 fm	11-20 fm	>20 fm
Black Rockfish	23%	42%	90%
Blue Rockfish	29%	49%	100%
Bocaccio	30%	51%	100%
Canary Rockfish	32%	54%	100%
Widow Rockfish	32%	54%	100%
Yelloweye Rockfish	32%	56%	100%
Shallow Nearshore Rockfish	25%	49%	100%
Deeper Nearshore Rockfish	23%	48%	100%
Other Nearshore Rockfish	24%	48%	100%

Recent research has shown the effects of barotrauma can be mitigated in physoclistous (i.e., the swim bladder is not connected to the alimentary canal via a pneumatic duct) fish such as rockfish by releasing them at depth by using descending devices (Hannah, *et al.* 2012; Jarvis and Lowe 2008; Parker, *et al.* 2006; Pribyl, *et al.* 2012). The Council tasked the GMT to investigate the research on rockfish barotrauma mitigation for canary rockfish, cowcod, and yelloweye rockfish and propose depth-based mortality rates associated with release using descending devices. The GMT determined depth-based discard mortality rates associated with the use of descending devices for these stocks in a Bayesian Hierarchical model that considered the uncertainty of using other species as a proxy for these three, as well as the uncertainty associated with missing observations in one or more depth bins. They also calculated the upper 60 percent, 75 percent, 90 percent, and 95 percent confidence intervals so the Council could choose their preferred level of risk tolerance, given the uncertainties characterized in the analysis. The Council decided to recommend the mortality estimates calculated for these three stocks at the 90 percent upper confidence interval (Table 1-13). The Council also explained that these rates may be revised in the future as more research emerges informing survival of rockfish released at sea using descending devices. The Council also asked that these rates be applied retrospectively in estimating total mortality of these species for those samples where adequate sampling information exists.

**Table 1-13. Discard mortality rates by depth of canary rockfish, cowcod, and yelloweye rockfish released in west coast recreational fisheries using descending devices.**

<b>Depth (fm)</b>	<b>Canary</b>	<b>Cowcod</b>	<b>Yelloweye</b>
0-10	21%	21%	22%
10-20	25%	35%	26%
20-30	25%	52%	26%
30-50	48%	57%	27%
50-100	57%	57%	57%
>100	100%	100%	100%

### **1.3 The Groundfish Harvest Specification Framework and Harvest Specifications for Fisheries in 2015 and Beyond**

West coast groundfish stocks are managed under a harvest specification framework that considers scientific and management uncertainties. The first specification is the overfishing limit (OFL), which is the maximum sustainable yield (MSY) estimated for the stock and the legal harvest limit beyond which constitutes overfishing. The OFL is determined either by applying the harvest rate estimated to result in a biomass capable of sustaining MSY (i.e.,  $F_{MSY}$ ) recommended by the Council’s Scientific and Statistical Committee (SSC) to an estimate of exploitable biomass in the case of assessed stocks or through an approved data-poor method (e.g., depletion-corrected average catch (DCAC) or depletion-based stock reduction analysis (DB-SRA)) in the case of unassessed stocks. Regardless of the method or data informing the calculation of an OFL, there is scientific uncertainty in the estimation of an OFL. The Pacific Coast Groundfish Fishery Management Plan (FMP) mandates a precautionary buffer to account for this uncertainty by prescribing an acceptable biological catch (ABC) harvest level that is less than the OFL. A further reduction from the ABC can be specified when setting an annual catch limit (ACL) that accounts for management uncertainty, socioeconomic considerations, ecological considerations, conservation objectives, and/or other considerations the Council and NMFS wish to address. Since the ACL can be set equal to the ABC, the ABC is the highest harvest level that can be specified for west coast groundfish stocks.

The following sections describe in detail the science informing 2015 and 2016 harvest specifications. Table 1-14 summarizes the 2014, 2015, and 2016 harvest specifications for west coast groundfish stocks.

**Table 1-14. Status quo 2014 harvest specifications and final preferred overfishing limits (OFLs in mt), acceptable biological catches (ABCs in mt), annual catch limits (ACLs in mt), and ecosystem component species designations for west coast groundfish stocks and stock complexes in 2015 and 2016 (stocks with new assessments in bold).**

Stock	2014			2015			2016		
	OFL	ABC	ACL	OFL	ABC	ACL	OFL	ABC	ACL
<b>OVERFISHED STOCKS</b>									
<b>BOCACCIO S. of 40°10'</b>	<b>881</b>	<b>842</b>	<b>337</b>	<b>1,444</b>	<b>1,380</b>	<b>349</b>	<b>1,351</b>	<b>1,291</b>	<b>362</b>
CANARY	741	709	119	733	701	122	729	697	125
<b>COWCOD S. of 40°10'</b>	<b>12</b>	<b>9</b>	<b>3</b>	<b>67</b>	<b>60</b>	<b>10</b>	<b>66</b>	<b>59</b>	<b>10</b>
<b>DARKBLOTCHED</b>	<b>553</b>	<b>529</b>	<b>330</b>	<b>574</b>	<b>549</b>	<b>338</b>	<b>580</b>	<b>554</b>	<b>346</b>
PACIFIC OCEAN PERCH	838	801	153	842	805	158	850	813	164
<b>PETRALE SOLE</b>	<b>2,774</b>	<b>2,652</b>	<b>2,652</b>	<b>2,946</b>	<b>2,816</b>	<b>2,816</b>	<b>3,044</b>	<b>2,910</b>	<b>2,910</b>
YELLOWEYE	51	43	18	52	47	18	52	47	19
<b>NON-OVERFISHED STOCKS</b>									
Arrowtooth Flounder	6,912	5,758	5,758	6,599	5,497	5,497	6,396	5,328	5,328
Black Rockfish (OR-CA)	1,166	1,115	1,000	1,176	1,124	1,000	1,183	1,131	1,000
Black Rockfish (WA)	428	409	409	421	402	402	423	404	404
Cabazon (CA)	165	158	158	161	154	154	158	151	151
Cabazon (OR)	49	47	47	49	47	47	49	47	47
California scorpionfish	122	117	117	119	114	114	117	111	111
Chilipepper S. of 40°10'	1,722	1,647	1,647	1,703	1,628	1,628	1,694	1,619	1,619
Dover Sole	77,774	74,352	25,000	66,871	63,929	50,000	59,221	56,615	50,000
<b>English Sole</b>	<b>5,906</b>	<b>5,646</b>	<b>5,646</b>	<b>10,792</b>	<b>9,853</b>	<b>9,853</b>	<b>7,890</b>	<b>7,204</b>	<b>7,204</b>
Lingcod N of 40°10'	3,162	2,878	2,878	3,010	2,830	2,830	2,891	2,719	2,719
Lingcod S. of 40°10'	1,276	1,063	1,063	1,205	1,004	1,004	1,136	946	946
Longnose skate	2,816	2,692	2,000	2,449	2,341	2,000	2,405	2,299	2,000
<b>Longspine Thornyhead (coastwide)</b>	<b>3,304</b>	<b>2,752</b>	NA	<b>5,007</b>	<b>4,171</b>	NA	<b>4,763</b>	<b>3,968</b>	NA
<b>Longspine Thornyhead N of 34°27'</b>	NA	NA	<b>1,958</b>	NA	NA	<b>3,170</b>	NA	NA	<b>3,015</b>
<b>Longspine Thornyhead S. of 34°27'</b>	NA	NA	<b>347</b>	NA	NA	<b>1,001</b>	NA	NA	<b>952</b>
Pacific Cod	3,200	2,221	1,600	3,200	2,221	1,600	3,200	2,221	1,600



Stock	2014			2015			2016		
	OFL	ABC	ACL	OFL	ABC	ACL	OFL	ABC	ACL
Sablefish (coastwide)	7,158	6,535	NA	7,857	7,173	NA	8,526	7,784	NA
Sablefish N of 36°	NA	NA	4,349	NA	NA	4,793	NA	NA	5,241
Sablefish S. of 36°	NA	NA	1,560	NA	NA	1,719	NA	NA	1,880
Shortbelly	6,950	5,789	50	6,950	5,789	500	6,950	5,789	500
<b>Shortspine Thornyhead (coastwide)</b>	<b>2,310</b>	<b>2,208</b>	<b>NA</b>	<b>3,203</b>	<b>2,668</b>	<b>NA</b>	<b>3,169</b>	<b>2,640</b>	<b>NA</b>
<b>Shortspine Thornyhead N of 34°27'</b>	<b>NA</b>	<b>NA</b>	<b>1,525</b>	<b>NA</b>	<b>NA</b>	<b>1,745</b>	<b>NA</b>	<b>NA</b>	<b>1,726</b>
<b>Shortspine Thornyhead S. of 34°27'</b>	<b>NA</b>	<b>NA</b>	<b>393</b>	<b>NA</b>	<b>NA</b>	<b>923</b>	<b>NA</b>	<b>NA</b>	<b>913</b>
Spiny dogfish	2,950	2,024	NA	2,523	1,912	1,912	2,503	1,897	1,897
Splitnose S. of 40°10'	1,747	1,670	1,670	1,794	1,715	1,715	1,826	1,746	1,746
Starry Flounder	1,834	1,528	1,528	1,841	1,534	1,534	1,847	1,539	1,539
Widow	4,435	4,212	1,500	4,137	3,929	2,000	3,990	3,790	2,000
<b>Yellowtail N of 40°10'</b>	<b>4,584</b>	<b>4,382</b>	<b>4,382</b>	<b>7,218</b>	<b>6,590</b>	<b>6,590</b>	<b>6,949</b>	<b>6,344</b>	<b>6,344</b>
<b>STOCK COMPLEXES</b>									
Nearshore Rockfish North	110	94	94	90	79	69	90	79	69
Shelf Rockfish North	2,195	1,932	968	2,208	1,944	1,944	2,217	1,953	1,952
Slope Rockfish North	1,553	1,414	1,160	1,831	1,693	1,693	1,844	1,706	1,706
Nearshore Rockfish South	1,160	1,001	990	1,309	1,165	1,114	1,317	1,163	1,006
Shelf Rockfish South	1,913	1,620	714	1,914	1,625	1,624	1,915	1,626	1,625
Slope Rockfish South	685	622	622	813	705	693	814	705	695
Other Flatfish	10,060	6,982	4,884	11,453	8,749	8,749	9,645	7,243	7,243
Other Fish	6,802	4,697	4,697						
<i>Cabezon (WA)</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>4.5</i>	<i>3.7</i>	<i>3.7</i>	<i>4.8</i>	<i>4.0</i>	<i>4.0</i>
<i>Kelp greenling (CA)</i>	<i>118.9</i>	<i>82.5</i>	<i>NA</i>	<i>118.9</i>	<i>99.2</i>	<i>99.2</i>	<i>118.9</i>	<i>99.2</i>	<i>99.2</i>
<i>Kelp greenling (OR)</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>
<i>Kelp greenling (WA)</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>	<i>a/</i>
<i>Leopard shark</i>	<i>167.1</i>	<i>139.4</i>	<i>139.4</i>	<i>167.1</i>	<i>139.4</i>	<i>139.4</i>	<i>167.1</i>	<i>139.4</i>	<i>139.4</i>

Stock	2014			2015			2016		
	OFL	ABC	ACL	OFL	ABC	ACL	OFL	ABC	ACL
<b>ECOSYSTEM COMPONENT SPECIES</b>									
Big skate	458.0	317.9	NA	No harvest specifications for an EC species					
California skate	86.0	59.7	NA						
Aleutian skate	b/	b/	b/						
Roughtail/black skate	b/	b/	b/						
Bering/sandpaper skate	b/	b/	b/						
All other skates	b/	b/	b/						
Pacific grenadier	1,519.0	1,054.2	NA						
Giant grenadier	b/	b/	b/						
All other grenadiers	b/	b/	b/						
Ratfish	1,441.0	1,000.1	NA						
Soupfin shark	61.6	42.8	NA						
Finescale codling	a/	a/	a/						


a/ No OFL or ABC contribution for these stocks given the lack of an approved method.

b/ No harvest specifications adopted since these species are not currently managed in the FMP.

### 1.3.1 Overfishing Limits

The OFL is the MSY harvest level associated with the current stock abundance and is the estimated or proxy MSY harvest level, which is the harvest threshold above which overfishing occurs. The methods for determining OFL are based on the best available science and the recommendation of the SSC; therefore, alternatives are not developed for this reference point.

The OFL is calculated by applying a deterministic or proxy MSY harvest rate (denoted  $F_{MSY}$ ) to the estimated exploitable biomass of a managed stock. The  $F_{MSY}$  harvest rate may be converted to a Spawning Potential Ratio (SPR). For ease of comparison among stocks and to standardize the basis of rebuilding calculations, it is useful to express any specific fishing mortality rate in terms of its effect on Spawning Potential Ratio (SPR = spawning per recruit at the current population level relative to that at the stock's unfished condition). Given fishery selectivity patterns and basic life history parameters, there is a direct inverse relationship between  $F$  and SPR (Figure 1-38). When there is no fishing, each new female recruit is expected to achieve 100 percent of its spawning potential. As fishing intensity increases, expected lifetime reproduction declines due to this added source of mortality. Conversion of  $F$  into the equivalent SPR has the benefit of standardizing for differences in growth, maturity, fecundity, natural mortality, and fishery selectivity patterns and, as a consequence, the Council's SSC recommends that it be used routinely.



**Figure 1-38. Relationship between SPR and instantaneous fishing mortality rate (F) for a hypothetical rockfish.**

Amendment 23, which was adopted in December 2010 and implemented in 2011, revised the descriptions of species categories used in the development of harvest specifications. The first category (category 1) includes species with relatively data-rich quantitative stock assessments that are developed on the basis of catch-at-age, catch-at-length, or other data. Recruitments are estimated for category 1 stocks. OFLs and overfished/rebuilding thresholds can generally be calculated for these species. The second category (category 2) includes species for which some biological indicators are available yet data informing an assessment are limited. Category 2 assessments include a new class of data-moderate assessments where catch data and one or more indices of abundance inform the status and biomass of the stock, but age and length compositional data are excluded. This type of assessment allows for a more expeditious assessment

review than the category 1 full assessments, which require a rigorous review process<sup>9</sup>, thus enabling more stocks to be assessed in an assessment cycle. Two data-moderate assessment models were approved for the 2013 assessment cycle (which informs management decision-making for 2015 and beyond): extended depletion-based stock reduction analysis (XDB-SRA) and extended simple stock synthesis (exSSS). The third assessment category (category 3) includes minor species for which are caught and where the only available information is catch. When setting the 2015 and 2016 OFLs for category 1 or 2 species, the  $F_{MSY}$  harvest rate or a proxy was applied to the estimated exploitable biomass. A policy of using a default harvest rate as a proxy for the fishing mortality rate that is expected to achieve MSY is also referred to as the  $F_{MSY}$  control rule or maximum fishing mortality threshold (MFMT) harvest rate. Catch-based methods are generally used to determine the OFL for category 3 species.

New stock assessments and rebuilding analyses recommended by the SSC for use in setting biennial harvest specifications were approved by the Council for setting the 2015 and 2016 biennial harvest specifications. Eight full stock assessments, eight data-moderate stock assessments, and one stock assessment update were prepared to inform the 2015 and 2016 harvest specifications. Full stock assessments included those for aurora rockfish, cowcod south of 34°27' N lat., darkblotched rockfish, longspine thornyhead, petrale sole, Pacific sanddabs, rougheye/blackspotted rockfish (analyzed as a complex of two stocks), and shortspine thornyhead. Aurora rockfish, rougheye/blackspotted rockfish, and Pacific sanddabs were assessed for the first time in 2013. OFLs were estimated for seven of these eight stocks; the assessment for Pacific sanddabs was recommended to inform stock status but not current stock biomass or OFL. Eight stocks were assessed using the data-moderate methods adopted by the Council. These stocks included brown rockfish, China rockfish, copper rockfish, English sole, rex sole, sharpchin rockfish, stripetail rockfish, and yellowtail rockfish north of 40°10' N lat. OFLs were estimated for seven of the eight data-moderate stocks; the stripetail rockfish assessment was recommended to inform stock status but not current stock biomass or OFL. A stock assessment update, which incorporates new data but does not change model structure or assumptions, was prepared for bocaccio. For stocks that were not assessed in 2013, the Council used 2015-2016 OFLs calculated using the most recent stock assessments or update or historical landings data.

Two data-poor methods, depletion-corrected average catch (DCAC) and depletion-based stock reduction analysis (DB-SRA) have been used to determine most of the category 3 OFLs since 2011. These methods were recommended for determining 2015 and 2016 OFLs for unassessed stocks, for which there are enough relevant data. The average historical catch approach was used to determine OFLs for stocks where the historical catches were too sparse to use DCAC or DB-SRA.

For 2015 and 2016, default harvest rates were used as a proxy for the fishing mortality rate that is expected to achieve the MSY ( $F_{MSY}$ ). A proxy is used because there is insufficient information for most Pacific Coast groundfish stocks to establish a species-specific  $F_{MSY}$ . In 2015 and 2016, the following default harvest rate proxies, based on SSC recommendations, were used:  $F_{30\%}$  for assessed flatfish,  $F_{40\%}$  for Pacific whiting,  $F_{50\%}$  for rockfish (including thornyheads),  $F_{50\%}$  for spiny dogfish and longnose skate, and  $F_{45\%}$  for other groundfish such as sablefish and lingcod. The FMP allows default harvest rate proxies to be modified as scientific knowledge improves for a particular species.

Table 1-15 compares the 2015 and 2016 OFLs with the 2014 OFL for stocks managed with stock-specific harvest specifications. The OFLs are specified for all the stocks and stock complexes actively managed in the fishery, as required by the FMP. The 2014 OFLs in Table 1-15 were projected from stock assessments done in 2011 or earlier. The 2015 and 2016 OFLs in Table 1-15 include the results of stock assessments conducted in 2013. The OFL contributions for the cowcod stock south of 40°10' N lat. are shown as area-

---

<sup>9</sup> The review process for new full assessments includes a Stock Assessment Review (STAR) panel review and a subsequent review by the Council's Scientific and Statistical Committee (SSC). Only those assessments that are endorsed by the SSC are considered for formal adoption in the Council process.

specific OFL contributions because they were derived using different methods. The Conception area OFLs for cowcod were projected from the 2013 rebuilding analysis (Dick and MacCall 2014) and the Monterey area OFLs were derived using DB-SRA. Although the area-specific OFL contributions for cowcod are displayed in Table 1-15, the OFL is specified for the entire stock south of 40°10' N lat. and not for each area. The 2014 OFL and 2015 and 2016 OFL contributions of individual stocks within the six Rockfish complexes, the Other Flatfish complex, and the Other Fish complex are shown in italics in Table 1-15. The OFL contributions for the individual stocks were summed to determine the complex OFLs.

The Council recommended restructuring the status quo Other Fish complex by removing spiny dogfish and managing that stock with stock-specific harvest specifications (Table 1-15), designating the skates, grenadiers, soupfin shark, finescale codling, and ratfish stocks as Ecosystem Component species, and managing kelp greenling, the Washington stock of cabezon, and leopard shark in a reconfigured Other Fish complex.

The preferred 2015 and 2016 OFLs for west coast groundfish stocks and stock complexes used the same policies (e.g.,  $F_{MSY}$  harvest rates and methodologies) used to determine the 2014 OFLs (i.e., No Action) with the following exceptions:

- The spiny dogfish  $F_{MSY}$  proxy harvest rate was changed from  $F_{45\%}$  to  $F_{50\%}$ ;
- Spiny dogfish is recommended to be removed from the Other Fish complex and managed with stock-specific harvest specifications;
- The skates, Pacific grenadier, finescale codling, soupfin shark, and ratfish stocks are removed from the Other Fish complex and designated as Ecosystem Component species;
- Those endemic skate and grenadier species not previously managed in the FMP are recommended to be added to the FMP and designated as Ecosystem Component species; and
- Kelp greenling, the Washington stock of cabezon, and leopard shark are recommended for management in a new Shallow Roundfish complex (there is also an alternative to manage these stocks individually with stock-specific harvest specifications).

**Table 1-15. Specified 2014 OFLs (mt) and preferred 2015 and 2016 OFLs (mt) for west coast groundfish stocks stock complexes (overfished stocks in CAPS, stocks with new assessments in bold, substock contributions to a stock OFL in *italics*).**

Stock	2014 OFL	Category	2015 OFL	2016 OFL
<b>OVERFISHED STOCKS</b>				
<b>BOCACCIO S. of 40°10' N. lat.</b>	<b>881</b>	<b>1</b>	<b>1,444</b>	<b>1,351</b>
CANARY	741	1	733	729
COWCOD S. of 40°10' N. lat.	12		<b>66.6</b>	<b>68.4</b>
<i>COWCOD (Conception)</i>	<b>7</b>	<b>2</b>	<b>55.0</b>	<b>56.4</b>
<i>COWCOD (Monterey)</i>	5	3	<i>11.6</i>	<i>12.0</i>
<b>DARKBLOTCHED</b>	<b>553</b>	<b>1</b>	<b>574</b>	<b>580</b>
PACIFIC OCEAN PERCH	838	1	842	850
<b>PETRALE SOLE</b>	<b>2,774</b>	<b>1</b>	<b>2,946</b>	<b>3,044</b>
YELLOWEYE	51	2	52	52
<b>NON-DEPLETED STOCKS</b>				
Arrowtooth Flounder	6,912	2	6,599	6,396
Black Rockfish (OR-CA)	1,166	1	1,176	1,183
Black Rockfish (WA)	428	1	421	423
Cabazon (CA)	165	1	161	158
Cabazon (OR)	49	1	49	49
California scorpionfish	122	1	119	117
Chilipepper S. of 40°10' N. lat.	1,722	1	1,703	1,694
Dover Sole	77,774	1	66,871	59,221
<b>English Sole</b>	<b>5,906</b>	<b>2</b>	<b>10,792</b>	<b>7,890</b>
Lingcod N. of 40°10' N. lat.	3,162	1	3,010	2,891
Lingcod S. of 40°10' N. lat.	1,276	2	1,205	1,136
Longnose skate	2,816	1	2,449	2,405
<b>Longspine Thornyhead (coastwide)</b>	<b>3,304</b>	<b>2</b>	<b>5,007</b>	<b>4,763</b>
Pacific Cod	3,200	3	3,200	3,200
Sablefish (coastwide)	7,158	1	7,857	8,526
Shortbelly	6,950	2	6,950	6,950
<b>Shortspine Thornyhead (coastwide)</b>	<b>2,310</b>	<b>2</b>	<b>3,203</b>	<b>3,169</b>
Spiny dogfish	2,950	2	2,523	2,503
Splitnose S. of 40°10' N. lat.	1,747	1	1,794	1,826
Starry Flounder	1,834	2	1,841	1,847
Widow	4,435	1	4,137	3,990
<b>Yellowtail N. of 40°10' N. lat.</b>	<b>4,584</b>	<b>2</b>	<b>7,218</b>	<b>6,949</b>
<b>STOCK COMPLEXES</b>				
Nearshore Rockfish North	110		88	88
<i>Black and yellow</i>	<i>0.01</i>	3	<i>0.01</i>	<i>0.01</i>
<i>Blue (CA)</i>	<i>27.4</i>	2	<i>27.4</i>	<i>27.7</i>
<i>Blue (OR &amp; WA)</i>	<i>32.3</i>	3	<i>32.3</i>	<i>32.3</i>
<b>Brown</b>	<b>5.5</b>	<b>2</b>	<b>1.9</b>	<b>1.9</b>
<i>Calico</i>	-	3	-	-
<b>China</b>	<b>9.8</b>	<b>2</b>	<b>7.2</b>	<b>7.4</b>
<b>Copper</b>	<b>26.0</b>	<b>2</b>	<b>10.6</b>	<b>10.3</b>
<i>Gopher</i>	-	3	-	-
<i>Grass</i>	<i>0.7</i>	3	<i>0.7</i>	<i>0.7</i>
<i>Kelp</i>	<i>0.01</i>	3	<i>0.01</i>	<i>0.01</i>
<i>Olive</i>	<i>0.3</i>	3	<i>0.3</i>	<i>0.3</i>

Stock	2014 OFL	Category	2015 OFL	2016 OFL
<i>Quillback</i>	7.4	3	7.4	7.4
<i>Treefish</i>	0.2	3	0.2	0.2
Shelf Rockfish North	2,195		2,209	2,218
<i>Bronzespotted</i>	-	3	-	-
<i>Bocaccio</i>	284.0	3	284.0	284.0
<i>Chameleon</i>	-	3	-	-
<i>Chilipepper</i>	129.6	3	128.2	127.5
<i>Cowcod</i>	-	3	0.4	0.4
<i>Flag</i>	0.1	3	0.1	0.1
<i>Freckled</i>	-	3	-	-
<i>Greenblotched</i>	1.3	3	1.3	1.3
<i>Greenspotted 40°10' to 42° N. lat.</i>	9.4	2	10.0	9.9
<i>Greenspotted N. of 42 N. lat. (OR &amp; WA)</i>	6.1	3	6.1	6.1
<i>Greenstriped</i>	1,268.3	2	1,281.9	1,292.0
<i>Halfbanded</i>	-	3	-	-
<i>Harlequin</i>	-	3	-	-
<i>Honeycomb</i>	-	3	-	-
<i>Mexican</i>	-	3	-	-
<i>Pink</i>	0.004	3	0.004	0.004
<i>Pinkrose</i>	-	3	-	-
<i>Puget Sound</i>	-	3	-	-
<i>Pygmy</i>	-	3	-	-
<i>Redstripe</i>	269.9	3	269.9	269.9
<i>Rosethorn</i>	12.9	3	12.9	12.9
<i>Rosy</i>	3.0	3	3.0	3.0
<i>Silvergray</i>	159.4	3	159.4	159.4
<i>Speckled</i>	0.2	3	0.2	0.2
<i>Squarespot</i>	0.2	3	0.2	0.2
<i>Starry</i>	0.004	3	0.004	0.004
<b>Stripetail</b>	40.4	3	40.4	40.4
<i>Swordspine</i>	0.0001	3	0.0001	0.0001
<i>Tiger</i>	1.0	3	1.0	1.0
<i>Vermilion</i>	9.7	3	9.7	9.7
Slope Rockfish North	1,553		1,831	1,844
<b>Aurora</b>	<b>15.4</b>	<b>1</b>	<b>17.4</b>	<b>17.5</b>
<i>Bank</i>	17.2	3	17.2	17.2
<i>Blackgill</i>	4.7	3	4.7	4.7
<i>Redbanded</i>	45.3	3	45.3	45.3
<b>Rougheye/Blackspotted</b>	<b>71.1</b>	<b>2</b>	<b>201.9</b>	<b>206.8</b>
<b>Sharpchin</b>	<b>214.5</b>	<b>2</b>	<b>332.8</b>	<b>323.2</b>
<i>Shortraker</i>	18.7	3	18.7	18.7
<i>Splitnose</i>	974.1	1	1,000.6	1,018.2
<i>Yellowmouth</i>	192.4	3	192.4	192.4
Nearshore Rockfish South	1,160		1,313	1,288
<i>Shallow Nearshore Species</i>	NA	NA	NA	NA
<i>Black and yellow</i>	27.5	3	27.5	27.5
<b>China</b>	<b>16.6</b>	<b>2</b>	<b>55.2</b>	<b>52.7</b>
<i>Gopher (N of Pt. Conception)</i>	153.0	1	148.0	144.0
<i>Gopher (S of Pt. Conception)</i>	25.6	3	25.6	25.6
<i>Grass</i>	59.6	3	59.6	59.6

Stock	2014 OFL	Category	2015 OFL	2016 OFL
<i>Kelp</i>	27.7	3	27.7	27.7
<i>Deeper Nearshore Species</i>	NA	NA	NA	NA
<i>Blue (assessed area)</i>	187.8	2	188.6	190.3
<i>Blue (S of 34°27' N. lat.)</i>	72.9	3	72.9	72.9
<b>Brown</b>	<b>204.6</b>	<b>2</b>	<b>163.8</b>	<b>160.2</b>
<i>Calico</i>	-	3	-	-
<b>Copper</b>	<b>141.5</b>	<b>2</b>	<b>301.1</b>	<b>284.3</b>
<i>Olive</i>	224.6	3	224.6	224.6
<i>Quillback</i>	5.4	3	5.4	5.4
<i>Treefish</i>	13.2	3	13.2	13.2
Shelf Rockfish South	1,912.9		1,917.9	1,918.9
<i>Bronzespotted</i>	3.6	3	3.6	3.6
<i>Chameleon</i>	-	3	-	-
<i>Flag</i>	23.4	3	23.4	23.4
<i>Freckled</i>	-	3	-	-
<i>Greenblotched</i>	23.1	3	23.1	23.1
<i>Greenspotted</i>	80.3	2	82.8	82.0
<i>Greenstriped</i>	232.7	2	235.1	237.0
<i>Halfbanded</i>	-	3	-	-
<i>Harlequin</i>	-	3	-	-
<i>Honeycomb</i>	9.9	3	9.9	9.9
<i>Mexican</i>	5.1	3	5.1	5.1
<i>Pink</i>	2.5	3	2.5	2.5
<i>Pinkrose</i>	-	3	-	-
<i>Pygmy</i>	-	3	-	-
<i>Redstripe</i>	0.5	3	0.5	0.5
<i>Rosethorn</i>	2.1	3	2.1	2.1
<i>Rosy</i>	44.5	3	44.5	44.5
<i>Silvergray</i>	0.5	3	0.5	0.5
<i>Speckled</i>	39.4	3	39.4	39.4
<i>Squarespot</i>	11.1	3	11.1	11.1
<i>Starry</i>	62.6	3	62.6	62.6
<b>Stripetail</b>	<b>23.6</b>	<b>3</b>	<b>23.6</b>	<b>23.6</b>
<i>Swordspine</i>	14.2	3	14.2	14.2
<i>Tiger</i>	0.04	3	0.04	0.04
<i>Vermilion</i>	269.3	3	269.3	269.3
<i>Yellowtail</i>	1,064.4	3	1,064.4	1,064.4
Slope Rockfish South	685		813	814
<b>Aurora</b>	<b>26.1</b>	<b>1</b>	<b>74.3</b>	<b>74.3</b>
<i>Bank</i>	503.2	3	503.2	503.2
<i>Blackgill</i>	134.0	2	137.0	140.0
<i>Pacific ocean perch</i>	-	3	-	-
<i>Redbanded</i>	10.4	3	10.4	10.4
<b>Rougheye/Blackspotted</b>	<b>0.4</b>	<b>2</b>	<b>4.1</b>	<b>4.2</b>
<b>Sharpchin</b>	<b>9.8</b>	<b>2</b>	<b>83.2</b>	<b>80.8</b>
<i>Shortraker</i>	0.1	3	0.1	0.1
<i>Yellowmouth</i>	0.8	3	0.8	0.8
Other Flatfish	10,060		11,453	9,645
<i>Butter sole</i>	4.6	3	4.6	4.6
<i>Curlfin sole</i>	8.2	3	8.2	8.2



Stock	2014 OFL	Category	2015 OFL	2016 OFL
<i>Flathead sole</i>	35.0	3	35.0	35.0
<b><i>Pacific sanddab</i></b>	4,801.0	<b>3</b>	4,801.0	4,801.0
<b><i>Rex sole</i></b>	<b>4,371.5</b>	<b>2</b>	<b>5,764.0</b>	<b>3,956.0</b>
<i>Rock sole</i>	66.7	3	66.7	66.7
<i>Sand sole</i>	773.2	3	773.2	773.2
Other Fish a/	6,802	3	291	291
<i>Cabezon (WA)</i>	b/	3	4.5	4.8
<i>Kelp greenling (CA)</i>	118.9	3	118.9	118.9
<i>Kelp greenling (OR)</i>	b/	3	b/	b/
<i>Kelp greenling (WA)</i>	b/	3	b/	b/
<i>Leopard shark</i>	167.1	3	167.1	167.1

a/ Values for these specifications are the sum of known contributions of component stocks. The 2014 OFL is not comparable to the 2015 and 2016 OFLs since the Other Fish complex was restructured by designating most of the component species as Ecosystem Component species.

b/ No OFL contribution for these stocks given the lack of an approved method.

### 1.3.2 Acceptable Biological Catches

The 2014, 2015, and 2016 ABCs are annual catch specifications that are the stock or stock complex's OFL reduced by an amount associated with the scientific uncertainty in estimating the OFL. Under the FMP harvest specification framework, scientific advice that is relatively more uncertain will result in ABCs that are relatively lower, all other things being equal (i.e., a precautionary reduction in catch will occur due purely to scientific uncertainty in estimating the OFL). The ABC is an SSC-recommended catch level that ACLs may not exceed. As explained in more detail below, the SSC developed a two-step approach referred to as the P\* approach for determining ABCs. In the P\* approach, the SSC determines the amount of scientific uncertainty associated with estimating the OFL in stock assessments, referred to as the sigma ( $\sigma$ ) value. The Council then chooses its preferred level of risk of overfishing, a policy decision, which is designated as the overfishing probability (P\*) (see Section 1.3.1). The SSC then applies the P\* value to the sigma value to determine the amount by which the OFL is reduced to establish the ABC. The SSC's recommendations for sigma and the reductions from OFL associated with different P\* values are science-based recommendations; therefore, alternatives to these values are not analyzed.

The SSC assigned each species in the groundfish fishery to one of three categories based on the level of information available about the species. Table 1-16 shows the criteria used by the SSC to categorize stocks. The SSC's recommended sigma value for category 1 stocks is based on a statistical analysis of the variance within and among stock assessments. The variance in estimating biomass is the metric used for determining sigma since the OFL is estimated by applying the harvest rate estimated or assumed to produce MSY (i.e.,  $F_{MSY}$ ) to the exploitable biomass, and uncertainty in exploitable biomass was considered the major contributor to overall uncertainty. The meta-analysis used stock assessments from 17 data-rich stocks to determine the proxy sigma value for category 1 stocks (Ralston, *et al.* 2011). The general methodology used by the SSC subcommittees to assess among-assessment uncertainty was to compare previous stock assessments and stock assessment updates, and consider the logarithms of the ratios of the biomass estimates for each pair of assessments and their reciprocals using the last 20 years from an assessment. This provides a distribution of stock size differences in log-space and, if this variation is averaged over species, provides a general view of total biomass variation (represented as sigma -  $\sigma$ ) that emerges among repeat assessments of stocks, while embracing a wide range of factors that affect variability in results. The SSC

indicated that biomass is most likely the dominant source of uncertainty; however, it is anticipated that other factors will need to be considered in the future.

**Table 1-16. Criteria used by the SSC to categorize stocks based on the quantity and quality of data informing the estimate of OFL. Stock categories are used in deciding 2015 and 2016 ABCs that accommodate the uncertainty in estimating OFLs.**

Category	Sub-category	Criteria
Category 1 - Data rich stocks. OFL based on $F_{MSY}$ or $F_{MSY}$ proxy from model output. ABC based on $P^*$ buffer.		
1	a	Reliable compositional (age and/or size) data sufficient to resolve year-class strength and growth characteristics. Only fishery-dependent trend information available. Age/size structured assessment model.
1	b	As in 1a, but trend information also available from surveys. Age/size structured assessment model.
1	c	Age/size structured assessment model with reliable estimation of the stock-recruit relationship.
Category 2 - Data moderate. OFL derived from model output (or natural mortality).		
2	a	$M^*$ survey biomass assessment (as in Rogers 1996).
2	b	Historical catches, fishery-dependent trend information only. An aggregate population model is fit to the available information.
2	c	Historical catches, survey trend information, or at least one absolute abundance estimate. An aggregate population model is fit to the available information.
2	d	Full age-structured assessment, but results are substantially more uncertain than assessments used in the calculation of the $P^*$ buffer. The SSC will provide a rationale for each stock placed in this category. Reasons could include that assessment results are very sensitive to model and data assumptions, or that the assessment has not been updated for many years.
Category 3 - Data poor. OFL derived from data-poor methods using historical catch.		
3	a	No reliable catch history. No basis for establishing OFL.
3	b	Reliable catch estimates only for recent years. OFL is average catch during a period when stock is considered to be stable and close to $B_{MSY}$ equilibrium on the basis of expert judgment.
3	c	Reliable aggregate catches during period of fishery development and approximate values for natural mortality. Default analytical approach DCAC.
3	d	Reliable annual historical catches and approximate values for natural mortality and age at 50% maturity. Default analytical approach DB-SRA.

Based on this analysis, the SSC recommended using the biomass variance statistic of  $\sigma = 0.36$  as a default for category 1 stocks. In cases where the stock biomass estimated in the most recent assessment has a variance for ending biomass greater than the variance estimated for that stock's category, the assessment's estimated variance is used instead. The logic behind this approach is that the uncertainty in any particular assessment is at least as large as the uncertainty that is estimated in the assessment. The stock biomass estimated in the 2011 widow rockfish assessment was judged to have a greater variance than the sigma of

0.36 used for other category 1 stocks. In this case, the SSC recommended using a sigma value of 0.41 for deciding the widow rockfish ABC. Likewise, the 2013 assessment for aurora rockfish also indicated a greater variance than the sigma of 0.36 used for other category 1 stocks. In that case, a sigma value of 0.39 was chosen for deciding the aurora rockfish ABC. The Council then recommends an appropriate P\* value and that P\* is mapped to its corresponding buffer fraction. When the P\* approach is used, the upper limit of P\* allowed by the FMP is 0.45.

Since there is greater scientific uncertainty for category 2 and 3 stocks relative to category 1 stocks, the scientific uncertainty buffer is generally greater than that recommended for category 1 stocks. The SSC recommended sigma values for category 2 and 3 stocks of 0.72 and 1.44, respectively (i.e., two and four times the sigma for category 1 stocks). The specific values of 0.72 and 1.44 were recommended by the SSC and considered to be the best available scientific information; however, the values are not based on a formal analysis of assessment outcomes and should be considered placeholders that could change substantially when the SSC reviews additional analyses in future management cycles. Table 1-17 shows the relationship between the values for sigma and the buffer for a range of values for P\*.

### 1.3.2.1 Considerations for Deciding the Overfishing Probability (P\*) When Specifying an Acceptable Biological Catch

The overfishing probability metric (P\*) is technically defined as the probability of overfishing a stock based on the scientific uncertainty in estimating the OFL. Interpretation of this definition has generated much discussion in the Council's harvest specification decision-making process. Either P\* is interpreted narrowly as the actual probability of overfishing, or P\* is considered more broadly as the Council's level of tolerance towards the risk that the OFL will be exceeded. Both viewpoints have merit but the latter view has more utility in the Council process, and is a more accurate representation of how the P\* value is decided.

The one problem with the literal definition of P\* is that the SSC has recommended a default value of sigma (0.36) for category 1 stocks, which are stocks that have assessments with estimated recruitment deviations (i.e., the strength of individual year classes is estimated). Nevertheless, category 1 assessments vary greatly both in the degree of uncertainty and how that uncertainty is characterized in the assessment model. It is common that one or more parameters are either estimated outside the model or assumed based on the assessment scientist's best judgment. In such cases, the uncertainty associated with that parameter is also not estimated nor characterized in any way within the assessment. For example, the 2011 sablefish assessment (Stewart, *et al.* 2011) appears to estimate current biomass with significant uncertainty. However, within that assessment many of the key parameters that affect the estimated biomass such as growth and natural mortality are explicitly estimated within the model<sup>10</sup>. The confidence interval associated with the ending year biomass estimate appears quite large relative to other assessments since the uncertainties associated with estimated growth and natural mortality are included within the overall assessment uncertainty. This compares to many other assessments, such as splitnose rockfish in 2009 (Gertseva, *et al.* 2009) or longspine thornyhead in 2013 (Stephens and Taylor 2013) where many parameters are assumed and fixed (e.g., natural mortality and steepness) because there is insufficient information to estimate these parameters in the assessment. In these cases, the biomass variances tend to underestimate the actual uncertainty.

The spectrum of assessment approaches vary between fully Bayesian models with most key parameters estimated (e.g., sablefish in 2011) to deterministic models with most parameters fixed (e.g., longspine thornyhead in 2013). Within the spectrum are parameter estimations using informed or diffuse priors. Given this variety of approaches and the degree to which uncertainty is characterized, it is hard to pursue a

---

<sup>10</sup> Stock-recruitment steepness (h), another parameter that affects the estimate of biomass, is fixed at an assumed 0.6 in the 2011 sablefish assessment.

formulaic approach where the P\* decision hinges on the scientific uncertainty associated with estimating the OFL. For the most part, the relative uncertainty in estimating the OFL is addressed with the SSC's sigma specification, which is only intended to broadly distinguish between assessments along the data-poor to data-rich continuum. The Council's P\* decision is therefore most appropriately considered as a risk assessment given many sources of uncertainty regarding the true state of nature for a stock.

### 1.3.2.2 Preferred 2015 and 2016 Acceptable Biological Catches

The ABCs for actively-managed stock complexes were determined by summing ABC values of the component stocks. Table 1-18 and Table 1-19 depict the potential alternative 2015 and 2016 ABCs, respectively for stocks and stock complexes across a range of P\* values from 0.25 to 0.45. Table 1-20 depicts the preferred 2015 and 2016 ABCs for stocks and stock complexes as well as the P\* values chosen to determine these ABCs.

The Council selected a P\* value of 0.45 for most category 1 stocks. With a P\* value of 0.45, a sigma value of 0.36 corresponds with a reduction of 4.4 percent from the OFL when deriving the ABC. For sablefish, the thornyheads, and assessed flatfish stocks, the Council selected a P\* value of 0.4. The preferred 2015 and 2016 ABCs used the same policies (i.e., stock categories, sigma and P\* values) used to determine the 2014 No Action ABCs with the following exceptions:

- Aurora rockfish was changed from a category 3 to a category 1 stock based on the new benchmark stock assessment adopted in 2013 (Hamel, *et al.* 2013). Therefore, the sigma of 1.44 for category 3 stocks was used to determine the 2014 ABC and a stock-specific sigma of 0.39 estimated for aurora rockfish was used to determine the 2015 and 2016 ABCs. The same P\* of 0.45 was used to determine 2014, 2015, and 2016 ABCs;
- Rougheye/blackspotted rockfish was changed from a category 3 (for rougheye alone) to a category 2 stock based on the new benchmark stock assessment adopted in 2013 (Hicks, *et al.* 2013). The SSC decided to designate the rougheye/blackspotted assemblage of stocks as category 2 since the assessment was for the complex of these two hard to distinguish stocks;
- English sole and yellowtail rockfish north of 40°10' N lat. were changed from category 1 to category 2 stocks based on their new data-moderate assessments;
- Brown rockfish, China rockfish, copper rockfish, rex sole, and sharpchin rockfish were changed from category 3 stocks to category 2 stocks based on new data-moderate assessments for these stocks in 2013; and
- Shortspine thornyhead was changed from a category 1 stock to a category 2 stock based on the lack of age data in the new benchmark assessment for this stock in 2013 (Taylor and Stephens 2013). The same P\* value of 0.4 was used to determine the 2014, 2015, and 2016 ABCs.

**Table 1-17. Relationship between P\* and the percent reduction of the OFL for deciding the 2015 and 2016 ABCs for category 1, aurora rockfish, widow rockfish, category 2, and category 3 stocks based on  $\sigma$  values of 0.36, 0.39, 0.41, 0.72, and 1.44, respectively.**

P*	Assessment Uncertainty ( $\sigma$ )				
	Cat. 1 0.36	Aurora 0.39	Widow 0.41	Cat. 2 0.72	Cat. 3 1.44
0.5	0	0	0	0	0
0.45	4.4%	4.8%	5.0%	8.7%	16.6%
0.44	5.3%		6.0%	10.3%	19.5%
0.43	6.2%		7.0%	11.9%	22.4%
0.42	7.0%		7.9%	13.5%	25.2%
0.41	7.9%		8.9%	15.1%	27.9%
0.4	8.7%	9.4%	9.9%	16.7%	30.6%
0.39	9.6%		10.8%	18.2%	33.1%
0.38	10.4%		11.8%	19.7%	35.6%
0.37	11.3%		12.7%	21.3%	38.0%
0.36	12.1%		13.7%	22.7%	40.3%
0.35	13.0%	14.0%	14.6%	24.2%	42.6%
0.34	13.8%		15.6%	25.7%	44.8%
0.33	14.6%		16.5%	27.1%	46.9%
0.32	15.5%		17.4%	28.6%	49.0%
0.31	16.3%		18.4%	30.0%	51.0%
0.3	17.2%	18.5%	19.3%	31.4%	53.0%
0.29	18.1%		20.3%	32.9%	54.9%
0.28	18.9%		21.3%	34.3%	56.8%
0.27	19.8%		22.2%	35.7%	58.6%
0.26	20.7%		23.2%	37.1%	60.4%
0.25	21.6%	23.1%	24.2%	38.5%	62.1%
0.24	22.5%		25.1%	39.9%	63.8%
0.23	23.4%		26.1%	41.3%	65.5%
0.22	24.3%		27.1%	42.6%	67.1%
0.21	25.2%		28.2%	44.0%	68.7%
0.2	26.1%	28.0%	29.2%	45.4%	70.2%
0.19	27.1%		30.2%	46.9%	71.8%
0.18	28.1%		31.3%	48.3%	73.2%
0.17	29.1%		32.4%	49.7%	74.7%
0.16	30.1%		33.5%	51.1%	76.1%
0.15	31.1%	33.2%	34.6%	52.6%	77.5%
0.14	32.2%		35.8%	54.1%	78.9%
0.13	33.3%		37.0%	55.6%	80.2%
0.12	34.5%		38.2%	57.1%	81.6%
0.11	35.7%		39.5%	58.7%	82.9%
0.1	37.0%	39.3%	40.9%	60.3%	84.2%
0.09	38.3%		42.3%	61.9%	85.5%
0.08	39.7%		43.8%	63.6%	86.8%
0.07	41.2%		45.4%	65.4%	88.1%
0.06	42.9%		47.1%	67.4%	89.3%
0.05	44.7%	47.3%	49.1%	69.4%	90.6%

**Table 1-18. 2014 ABCs (mt) and a range of alternative 2015 ABCs (mt) varied by the probability of overfishing (P\*) for west coast groundfish stocks (overfished stocks in CAPS; stocks with new assessments in bold; component stocks in stock complexes in italics).**

Stock	Status Quo 2014 ABC	2015 OFL	Cat.	Range of Alternative 2015 ABCs				
				Overfishing Probability (P*)				
				0.45	0.40	0.35	0.30	0.25
<b>OVERFISHED STOCKS</b>								
<b>BOCACCI</b> S. of 40°10' N. lat.	842	<b>1,444</b>	<b>1</b>	<b>1,380</b>	<b>1,318</b>	<b>1,256</b>	<b>1,195</b>	<b>1,132</b>
CANARY	709	733	1	701	669	638	607	575
COWCOD S. of 40°10' N. lat.	9	67		60	54	48	43	38
<i>COWCOD (Conception)</i>	<b>6</b>	<b>55</b>	<b>2</b>	<b>50</b>	<b>46</b>	<b>42</b>	<b>38</b>	<b>34</b>
<i>COWCOD (Monterey)</i>	3	12	3	10	8	7	5	4
<b>DARKBLOTCHED</b>	529	<b>574</b>	<b>1</b>	<b>549</b>	<b>524</b>	<b>499</b>	<b>475</b>	<b>450</b>
PACIFIC OCEAN PERCH	801	842	1	805	769	733	697	660
<b>PETRALE SOLE</b>	2,652	<b>2,946</b>	<b>1</b>	<b>2,816</b>	<b>2,690</b>	<b>2,563</b>	<b>2,439</b>	<b>2,310</b>
YELLOWEYE	43	52	2	47	43	39	35	32
<b>NON-DEPLETED STOCKS</b>								
Arrowtooth Flounder	5,758	6,599	2	6,025	5,497	5,002	4,527	4,058
Black Rockfish (OR-CA)	1,115	1,176	1	1,124	1,074	1,023	974	922
Black Rockfish (WA)	409	421	1	402	384	366	349	330
Cabazon (CA)	158	161	1	154	147	140	133	126
Cabazon (OR)	47	49	1	47	45	43	41	38
California scorpionfish	117	119	1	114	109	104	99	93
Chilipepper S. of 40°10' N. lat.	1,647	1,703	1	1,628	1,555	1,482	1,410	1,335
Dover Sole	74,352	66,871	1	63,929	61,053	58,178	55,369	52,427
<b>English Sole</b>	<b>5,646</b>	<b>10,792</b>	<b>2</b>	<b>9,853</b>	<b>8,990</b>	<b>8,180</b>	<b>7,403</b>	<b>6,637</b>
Lingcod N. of 40°10' N. lat.	2,878	3,010	1	2,830	2,659	2,494	2,334	2,172
Lingcod S. of 40°10' N. lat.	1,063	1,205	2	1,100	1,004	913	827	741
Longnose skate	2,692	2,449	1	2,341	2,236	2,130	2,027	1,920
<b>Longspine Thornyhead (coastwide)</b>	<b>2,752</b>	<b>5,007</b>	<b>2</b>	<b>4,571</b>	<b>4,171</b>	<b>3,795</b>	<b>3,435</b>	<b>3,079</b>
Pacific Cod	2,221	3,200	3	2,669	2,221	1,837	1,504	1,213
Sablefish (coastwide)	6,535	7,857	1	7,511	7,173	6,836	6,506	6,160
Shortbelly	5,789	6,950	2	6,345	5,789	5,268	4,768	4,274
<b>Shortspine Thornyhead (coastwide)</b>	<b>2,208</b>	<b>3,203</b>	<b>2</b>	<b>2,924</b>	<b>2,668</b>	<b>2,428</b>	<b>2,197</b>	<b>1,970</b>
Spiny dogfish	2,024	2,523	2	2,303	2,101	1,912	1,731	1,551
Splitnose S. of 40°10' N. lat.	1,670	1,794	1	1,715	1,638	1,561	1,485	1,406
Starry Flounder	1,528	1,841	2	1,681	1,534	1,395	1,263	1,132
Widow	4,212	4,137	1	3,929	3,729	3,532	3,337	3,138
<b>Yellowtail N. of 40°10' N. lat.</b>	<b>4,382</b>	<b>7,218</b>	<b>2</b>	<b>6,590</b>	<b>6,013</b>	<b>5,471</b>	<b>4,952</b>	<b>4,439</b>
<b>STOCK COMPLEXES</b>								
Nearshore Rockfish North	94	88		77	68	59	52	45
<i>Black and yellow</i>	<i>0.0</i>	<i>0.014</i>	3	<i>0.011</i>	<i>0.009</i>	<i>0.008</i>	<i>0.006</i>	<i>0.005</i>
<i>Blue (CA)</i>	<i>25.0</i>	<i>27.4</i>	2	<i>25.0</i>	<i>22.9</i>	<i>20.8</i>	<i>18.8</i>	<i>16.9</i>
<i>Blue (OR &amp; WA)</i>	<i>26.9</i>	<i>32.3</i>	3	<i>26.9</i>	<i>22.4</i>	<i>18.5</i>	<i>15.2</i>	<i>12.2</i>
<b>Brown</b>	<b>4.6</b>	<b>1.9</b>	<b>2</b>	<b>1.7</b>	<b>1.6</b>	<b>1.4</b>	<b>1.3</b>	<b>1.2</b>
<i>Calico</i>	<i>0.0</i>	-	3	-	-	-	-	-
<b>China</b>	<b>8.2</b>	<b>7.2</b>	<b>2</b>	<b>6.6</b>	<b>6.0</b>	<b>5.5</b>	<b>4.9</b>	<b>4.4</b>

Stock	Status Quo 2014 ABC	2015 OFL	Cat.	Range of Alternative 2015 ABCs				
				Overfishing Probability (P*)				
				0.45	0.40	0.35	0.30	0.25
<b>Copper</b>	<b>21.6</b>	<b>10.6</b>	<b>2</b>	<b>9.7</b>	<b>8.9</b>	<b>8.1</b>	<b>7.3</b>	<b>6.5</b>
<i>Gopher</i>	0.0	-	3	-	-	-	-	-
<i>Grass</i>	0.5	0.7	3	0.5	0.5	0.4	0.3	0.2
<i>Kelp</i>	0.0	0.009	3	0.008	0.006	0.005	0.004	0.003
<i>Olive</i>	0.3	0.3	3	0.3	0.2	0.2	0.1	0.1
<i>Quillback</i>	6.2	7.4	3	6.2	5.1	4.2	3.5	2.8
<i>Treefish</i>	0.2	0.2	3	0.2	0.2	0.1	0.1	0.1
Shelf Rockfish North	1,932	2,209		1,944	1,712	1,505	1,317	1,142
<i>Bronzespotted</i>	0.0	-	3	-	-	-	-	-
<i>Bocaccio</i>	236.9	284.0	3	236.9	197.1	163.0	133.5	107.6
<i>Chameleon</i>	0.0	-	3	-	-	-	-	-
<i>Chilipepper</i>	108.1	128.2	3	106.9	88.9	73.6	60.2	48.6
<i>Cowcod</i>	0.0	0.4	3	0.3	0.3	0.3	0.3	0.3
<i>Flag</i>	0.1	0.07	3	0.06	0.05	0.04	0.03	0.03
<i>Freckled</i>	0.0	-	3	-	-	-	-	-
<i>Greenblotched</i>	1.1	1.3	3	1.1	0.9	0.7	0.6	0.5
<i>Greenspotted 40°10' to 42° N. lat.</i>	9	10.0	2	9.1	8.3	7.5	6.8	6.1
<i>Greenspotted N. of 42 N. lat. (OR &amp; WA)</i>	5.1	6.1	3	5.1	4.2	3.5	2.9	2.3
<i>Greenstriped</i>	1,158	1,281.9	2	1,170.3	1,067.8	971.7	879.4	788.3
<i>Halfbanded</i>	0.0	-	3	-	-	-	-	-
<i>Harlequin</i>	0.0	-	3	-	-	-	-	-
<i>Honeycomb</i>	0.0	-	3	-	-	-	-	-
<i>Mexican</i>	0.0	-	3	-	-	-	-	-
<i>Pink</i>	0.0	0.004	3	0.003	0.003	0.002	0.002	0.001
<i>Pinkrose</i>	0.0	-	3	-	-	-	-	-
<i>Puget Sound</i>	0.0	-	3	-	-	-	-	-
<i>Pygmy</i>	0.0	-	3	-	-	-	-	-
<i>Redstripe</i>	225.1	269.9	3	225.1	187.3	154.9	126.9	102.3
<i>Rosethorn</i>	10.8	12.9	3	10.8	9.0	7.4	6.1	4.9
<i>Rosy</i>	2.5	3.0	3	2.5	2.1	1.7	1.4	1.1
<i>Silvergray</i>	133.0	159.4	3	133.0	110.6	91.5	74.9	60.4
<i>Speckled</i>	0.1	0.17	3	0.14	0.12	0.10	0.08	0.06
<i>Squarespot</i>	0.1	0.17	3	0.14	0.12	0.10	0.08	0.07
<i>Starry</i>	0.0	0.00	3	0.003	0.003	0.002	0.002	0.001
<b>Stripetail</b>	<b>33.7</b>	<b>40.4</b>	<b>3</b>	<b>33.7</b>	<b>28.0</b>	<b>23.2</b>	<b>19.0</b>	<b>15.3</b>
<i>Swordspine</i>	0.0	0.0001	3	0.00008	0.00007	0.00006	0.00005	0.00004
<i>Tiger</i>	0.8	1.0	3	0.8	0.7	0.6	0.5	0.4
<i>Vermilion</i>	8.1	9.7	3	8.1	6.7	5.6	4.6	3.7
Slope Rockfish North	1,414	1,831		1,693	1,568	1,451	1,340	1,232
<b>Aurora</b>	<b>12.8</b>	<b>17.4</b>	<b>1</b>	<b>16.6</b>	<b>15.8</b>	<b>15.0</b>	<b>14.2</b>	<b>13.4</b>
<i>Bank</i>	14.4	17.2	3	14.4	12.0	9.9	8.1	6.5
<i>Blackgill</i>	3.9	4.7	3	3.9	3.3	2.7	2.2	1.8
<i>Redbanded</i>	37.7	45.3	3	37.7	31.4	26.0	21.3	17.2
<b>Rougheye/Blackspotted</b>	<b>59.3</b>	<b>201.9</b>	<b>2</b>	<b>184</b>	<b>168</b>	<b>153</b>	<b>138</b>	<b>124</b>



Stock	Status Quo 2014 ABC	2015 OFL	Cat.	Range of Alternative 2015 ABCs				
				Overfishing Probability (P*)				
				0.45	0.40	0.35	0.30	0.25
<b>Sharpchin</b>	<b>178.9</b>	<b>332.8</b>	<b>2</b>	<b>303.8</b>	<b>277.2</b>	<b>252.3</b>	<b>228.3</b>	<b>204.7</b>
<i>Shorotraker</i>	15.6	18.7	3	15.6	13.0	10.7	8.8	7.1
<i>Splitnose</i>	931.3	1,000.6	1	956.6	913.6	870.5	828.5	784.5
<i>Yellowmouth</i>	160.5	192.4	3	160.5	133.6	110.5	90.4	72.9
Nearshore Rockfish South	1,001	1,313		1,169	1,042	928	823	725
<i>Shallow Nearshore Species</i>	NA	NA	NA	NA	NA	NA	NA	NA
<i>Black and yellow</i>	23.0	28	3	23.0	19.1	15.8	12.9	10.4
<b>China</b>	<b>13.8</b>	<b>55</b>	<b>2</b>	<b>50.4</b>	<b>46.0</b>	<b>41.9</b>	<b>37.9</b>	<b>34.0</b>
<i>Gopher (N of Pt. Conception)</i>	146.3	148	1	141.5	135.1	128.8	122.5	116.0
<i>Gopher (S of Pt. Conception)</i>	21.4	26	3	21.4	17.8	14.7	12.0	9.7
<i>Grass</i>	49.7	60	3	49.7	41.4	34.2	28.0	22.6
<i>Kelp</i>	23.1	28	3	23.1	19.2	15.9	13.0	10.5
<i>Deeper Nearshore Species</i>	NA	NA	NA	NA	NA	NA	NA	NA
<i>Blue (assessed area)</i>	171.4	189	2	172.2	157.1	142.9	129.4	116.0
<i>Blue (S of 34°27' N. lat.)</i>	60.8	73	3	60.8	50.6	41.8	34.3	27.6
<b>Brown</b>	<b>170.6</b>	<b>164</b>	<b>2</b>	<b>149.5</b>	<b>136.4</b>	<b>124.1</b>	<b>112.3</b>	<b>100.7</b>
<i>Calico</i>	0.0	-	3	-	-	-	-	-
<b>Copper</b>	<b>118.0</b>	<b>301</b>	<b>2</b>	<b>274.9</b>	<b>250.8</b>	<b>228.2</b>	<b>206.6</b>	<b>185.2</b>
<i>Olive</i>	187.4	225	3	187.4	155.9	128.9	105.6	85.1
<i>Quillback</i>	4.5	5	3	4.5	3.7	3.1	2.5	2.0
<i>Treefish</i>	11.0	13	3	11.0	9.2	7.6	6.2	5.0
Shelf Rockfish South	1,620	1,918		1,625	1,375	1,159	970	802
<i>Bronzespotted</i>	3.0	3.6	3	3.0	2.5	2.1	1.7	1.4
<i>Chameleon</i>	0.0	-	3	-	-	-	-	-
<i>Flag</i>	19.5	23.4	3	19.5	16.3	13.4	11.0	8.9
<i>Freckled</i>	0.0	-	3	-	-	-	-	-
<i>Greenblotched</i>	19.3	23.1	3	19.3	16.1	13.3	10.9	8.8
<i>Greenspotted</i>	73.3	82.8	2	75.6	69.0	62.8	56.8	50.9
<i>Greenstriped</i>	212.4	235.1	2	214.7	195.9	178.2	161.3	144.6
<i>Halfbanded</i>	0.0	-	3	-	-	-	-	-
<i>Harlequin</i>	0.0	-	3	-	-	-	-	-
<i>Honeycomb</i>	8.2	9.9	3	8.2	6.8	5.7	4.6	3.7
<i>Mexican</i>	4.2	5.1	3	4.2	3.5	2.9	2.4	1.9
<i>Pink</i>	2.1	2.5	3	2.1	1.8	1.5	1.2	1.0
<i>Pinkrose</i>	0.0	-	3	-	-	-	-	-
<i>Pygmy</i>	0.0	-	3	-	-	-	-	-
<i>Redstripe</i>	0.4	0.5	3	0.4	0.3	0.3	0.2	0.2
<i>Rosethorn</i>	1.8	2.1	3	1.8	1.5	1.2	1.0	0.8
<i>Rosy</i>	37.1	44.5	3	37.1	30.9	25.5	20.9	16.9
<i>Silvergray</i>	0.4	0.5	3	0.4	0.4	0.3	0.3	0.2
<i>Speckled</i>	32.8	39.4	3	32.8	27.3	22.6	18.5	14.9
<i>Squarespot</i>	9.2	11.1	3	9.2	7.7	6.4	5.2	4.2
<i>Starry</i>	52.2	62.6	3	52.2	43.4	35.9	29.4	23.7
<b>Stripetail</b>	<b>19.7</b>	<b>23.6</b>	<b>3</b>	<b>19.7</b>	<b>16.4</b>	<b>13.6</b>	<b>11.1</b>	<b>9.0</b>

Stock	Status Quo 2014 ABC	2015 OFL	Cat.	Range of Alternative 2015 ABCs				
				Overfishing Probability (P*)				
				0.45	0.40	0.35	0.30	0.25
<i>Swordspine</i>	11.9	14.2	3	11.9	9.9	8.2	6.7	5.4
<i>Tiger</i>	0.0	0.04	3	0.03	0.03	0.02	0.02	0.02
<i>Vermilion</i>	224.6	269.3	3	224.6	186.9	154.6	126.6	102.1
<i>Yellowtail</i>	887.7	1,064.4	3	887.7	738.7	611.0	500.3	403.4
Slope Rockfish South	622	813		705	611	529	456	390
<b><i>Aurora</i></b>	<b>21.7</b>	<b>74.3</b>	<b>1</b>	<b>70.7</b>	<b>67.3</b>	<b>63.9</b>	<b>60.6</b>	<b>57.1</b>
<i>Bank</i>	459.4	503.2	3	419.7	349.2	288.8	236.5	190.7
<i>Blackgill</i>	122.3	137.0	2	125.1	114.1	103.8	94.0	84.3
<i>Pacific ocean perch</i>	0.0	-	3	-	-	-	-	-
<i>Redbanded</i>	8.7	10.4	3	8.7	7.2	6.0	4.9	3.9
<b><i>Rougheye/Blackspotted</i></b>	<b>0.3</b>	<b>4.1</b>	<b>2</b>	<b>3.8</b>	<b>3.4</b>	<b>3.1</b>	<b>2.8</b>	<b>2.5</b>
<b><i>Sharpchin</i></b>	<b>8.2</b>	<b>83.2</b>	<b>2</b>	<b>76.0</b>	<b>69.3</b>	<b>63.1</b>	<b>57.1</b>	<b>51.2</b>
<i>Shortraker</i>	0.1	0.10	3	0.09	0.07	0.06	0.05	0.04
<i>Yellowmouth</i>	0.7	0.8	3	0.7	0.6	0.5	0.4	0.3
Other Flatfish	6,982	11,453		10,007	8,749	7,634	6,628	5,701
<i>Butter sole</i>	3.2	4.6	3	3.9	3.2	2.7	2.2	1.8
<i>Curlfin sole</i>	5.7	8.2	3	6.9	5.7	4.7	3.9	3.1
<i>Flathead sole</i>	24.3	35.0	3	29.2	24.3	20.1	16.5	13.3
<b><i>Pacific sanddab</i></b>	<b>3,331.9</b>	<b>4,801.0</b>	<b>3</b>	<b>4,004.0</b>	<b>3,331.9</b>	<b>2,755.8</b>	<b>2,256.5</b>	<b>1,819.6</b>
<b><i>Rex sole</i></b>	<b>3,033.8</b>	<b>5,764.0</b>	<b>2</b>	<b>5,262.5</b>	<b>4,801.4</b>	<b>4,369.1</b>	<b>3,954.1</b>	<b>3,544.9</b>
<i>Rock sole</i>	46.3	66.7	3	55.6	46.3	38.3	31.3	25.3
<i>Sand sole</i>	536.6	773.2	3	644.8	536.6	443.8	363.4	293.0
Other Fish	NA	290.5	3	242.3	201.6	166.7	136.5	110.1
<i>Cabezon (WA)</i>	a/	4.5	3	3.8	3.1	2.6	2.1	1.7
<i>Kelp greenling (CA)</i>	82.5	118.9	3	99.2	82.5	68.2	55.9	45.1
<i>Kelp greenling (OR)</i>	a/	b/	3	a/	a/	a/	a/	a/
<i>Kelp greenling (WA)</i>	a/	b/	3	a/	a/	a/	a/	a/
<i>Leopard shark</i>	1,054.2	167.1	3	139.4	116.0	95.9	78.5	63.3

a/ No OFL or ABC contribution for these stocks given the lack of an approved method for estimating the OFL.

**Table 1-19. 2014 ABCs (mt) and a range of alternative 2016 ABCs (mt) varied by the probability of overfishing (P\*) for west coast groundfish stocks (overfished stocks in CAPS; stocks with new assessments in bold; component stocks in stock complexes in *italics*).**

Stock	Status Quo 2014 ABC	2016 OFL	Range of Alternative 2016 ABCs				
			Overfishing Probability (P*)				
			0.45	0.40	0.35	0.30	0.25
<b>DEPLETED STOCKS</b>							
<b>BOCACCIO S. of 40°10' N. lat.</b>	842	<b>1,351</b>	<b>1,291</b>	<b>1,233</b>	<b>1,175</b>	<b>1,118</b>	<b>1,059</b>
CANARY	709	729	697	666	634	604	572
COWCOD S. of 40°10' N. lat.	9	68	62	55	50	44	39
<i><b>COWCOD (Conception)</b></i>	<b>6</b>	<b>56</b>	<b>51</b>	<b>47</b>	<b>43</b>	<b>39</b>	<b>35</b>
<i>COWCOD (Monterey)</i>	3	12	10	8	7	6	5
<b>DARKBLOTCHED</b>	529	<b>580</b>	<b>554</b>	<b>530</b>	<b>505</b>	<b>480</b>	<b>455</b>
PACIFIC OCEAN PERCH	801	850	813	776	740	704	666
<b>PETRALE SOLE</b>	2,652	<b>3,044</b>	<b>2,910</b>	<b>2,779</b>	<b>2,648</b>	<b>2,520</b>	<b>2,386</b>
YELLOWEYE	43	52	47	43	39	35	32
<b>NON-DEPLETED STOCKS</b>							
Arrowtooth Flounder	5,758	6,396	5,840	5,328	4,848	4,388	3,934
Black Rockfish (OR-CA)	1,115	1,183	1,131	1,080	1,029	980	927
Black Rockfish (WA)	409	423	404	386	368	350	332
Cabazon (CA)	158	158	151	144	137	131	124
Cabazon (OR)	47	49	47	45	43	41	38
California scorpionfish	117	117	111	106	101	97	91
Chilipepper S. of 40°10' N. lat.	1,647	1,694	1,619	1,547	1,474	1,403	1,328
Dover Sole	74,352	59,221	56,615	54,069	51,522	49,035	46,429
<b>English Sole</b>	<b>5,646</b>	<b>7,890</b>	<b>7,204</b>	<b>6,572</b>	<b>5,981</b>	<b>5,413</b>	<b>4,852</b>
Lingcod N. of 40°10' N. lat.	2,878	2,891	2,719	2,555	2,398	2,245	2,089
Lingcod S. of 40°10' N. lat.	1,063	1,136	1,037	946	861	779	699
Longnose skate	2,692	2,405	2,299	2,196	2,092	1,991	1,885
<b>Longspine Thornyhead (coastwide)</b>	<b>2,752</b>	<b>4,763</b>	<b>4,349</b>	<b>3,968</b>	<b>3,610</b>	<b>3,267</b>	<b>2,929</b>
Pacific Cod	2,221	3,200	2,669	2,221	1,837	1,504	1,213
Sablefish (coastwide)	6,535	8,526	8,151	7,784	7,418	7,060	6,684
Shortbelly	5,789	6,950	6,345	5,789	5,268	4,768	4,274
<b>Shortspine Thornyhead (coastwide)</b>	2,208	<b>3,169</b>	<b>2,893</b>	<b>2,640</b>	<b>2,402</b>	<b>2,174</b>	<b>1,949</b>
Spiny dogfish	2,024	2,503	2,285	2,085	1,897	1,717	1,540
Splitnose S. of 40°10' N. lat.	1,670	1,826	1,746	1,667	1,589	1,512	1,432
Starry Flounder	1,528	1,847	1,686	1,539	1,400	1,267	1,136
Widow	4,212	3,990	3,790	3,596	3,407	3,218	3,026
<b>Yellowtail N. of 40°10' N. lat.</b>	4,382	<b>6,949</b>	<b>6,344</b>	<b>5,789</b>	<b>5,267</b>	<b>4,767</b>	<b>4,274</b>
<b>STOCK COMPLEXES</b>							
Nearshore Rockfish North	94	88	77	68	59	52	45
<i>Black and yellow</i>	<i>0.0</i>	<i>0.014</i>	<i>0.011</i>	<i>0.009</i>	<i>0.008</i>	<i>0.006</i>	<i>0.005</i>
<i>Blue (CA)</i>	<i>25.0</i>	<i>27.7</i>	<i>25.3</i>	<i>23.1</i>	<i>21.0</i>	<i>19.0</i>	<i>17.0</i>
<i>Blue (OR &amp; WA)</i>	<i>26.9</i>	<i>32.3</i>	<i>26.9</i>	<i>22.4</i>	<i>18.5</i>	<i>15.2</i>	<i>12.2</i>
<b>Brown</b>	<b>4.6</b>	<b>1.9</b>	<b>1.7</b>	<b>1.6</b>	<b>1.4</b>	<b>1.3</b>	<b>1.2</b>
<i>Calico</i>	<i>0.0</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>
<b>China</b>	<b>8.2</b>	<b>7.4</b>	<b>6.8</b>	<b>6.2</b>	<b>5.6</b>	<b>5.1</b>	<b>4.6</b>

Stock	Status Quo 2014 ABC	2016 OFL	Range of Alternative 2016 ABCs				
			Overfishing Probability (P*)				
			0.45	0.40	0.35	0.30	0.25
<b>Copper</b>	<b>21.6</b>	<b>10.3</b>	<b>9.4</b>	<b>8.6</b>	<b>7.8</b>	<b>7.1</b>	<b>6.4</b>
<i>Gopher</i>	0.0	-	-	-	-	-	-
<i>Grass</i>	0.5	0.7	0.5	0.5	0.4	0.3	0.2
<i>Kelp</i>	0.0	0.009	0.008	0.006	0.005	0.004	0.003
<i>Olive</i>	0.3	0.32	0.26	0.22	0.18	0.15	0.12
<i>Quillback</i>	6.2	7.4	6.2	5.1	4.2	3.5	2.8
<i>Treefish</i>	0.2	0.22	0.18	0.15	0.12	0.10	0.08
Shelf Rockfish North	1,932	2,218	1,953	1,720	1,513	1,324	1,148
<i>Bronzespotted</i>	0.0	-	-	-	-	-	-
<i>Bocaccio</i>	236.9	284.0	236.9	197.1	163.0	133.5	107.6
<i>Chameleon</i>	0.0	-	-	-	-	-	-
<i>Chilipepper</i>	108.1	127.5	106.4	88.5	73.2	59.9	48.3
<i>Cowcod</i>	0.0	0.4	0.3	0.3	0.3	0.3	0.3
<i>Flag</i>	0.1	0.07	0.06	0.05	0.04	0.03	0.03
<i>Freckled</i>	0.0	-	-	-	-	-	-
<i>Greenblotched</i>	1.1	1.3	1.1	0.9	0.7	0.6	0.5
<i>Greenspotted 40°10' to 42° N. lat.</i>	9	9.9	9.0	8.2	7.5	6.8	6.1
<i>Greenspotted N. of 42 N. lat. (OR &amp; WA)</i>	5.1	6.1	5.1	4.2	3.5	2.9	2.3
<i>Greenstriped</i>	1,158	1,292.0	1,179.6	1,076.2	979.3	886.3	794.6
<i>Halfbanded</i>	0.0	-	-	-	-	-	-
<i>Harlequin</i>	0.0	-	-	-	-	-	-
<i>Honeycomb</i>	0.0	-	-	-	-	-	-
<i>Mexican</i>	0.0	-	-	-	-	-	-
<i>Pink</i>	0.0	0.004	0.003	0.003	0.002	0.002	0.001
<i>Pinkrose</i>	0.0	-	-	-	-	-	-
<i>Puget Sound</i>	0.0	-	-	-	-	-	-
<i>Pygmy</i>	0.0	-	-	-	-	-	-
<i>Redstripe</i>	225.1	269.9	225.1	187.3	154.9	126.9	102.3
<i>Rosethorn</i>	10.8	12.9	10.8	9.0	7.4	6.1	4.9
<i>Rosy</i>	2.5	3.0	2.5	2.1	1.7	1.4	1.1
<i>Silvergray</i>	133.0	159.4	133.0	110.6	91.5	74.9	60.4
<i>Speckled</i>	0.1	0.17	0.14	0.12	0.10	0.08	0.06
<i>Squarespot</i>	0.1	0.17	0.14	0.12	0.10	0.08	0.07
<i>Starry</i>	0.0	0.004	0.003	0.003	0.002	0.002	0.001
<b>Stripetail</b>	<b>33.7</b>	<b>40.4</b>	<b>33.7</b>	<b>28.0</b>	<b>23.2</b>	<b>19.0</b>	<b>15.3</b>
<i>Swordspine</i>	0.00008	0.00010	0.00008	0.00007	0.00006	0.00005	0.00004
<i>Tiger</i>	0.8	1.0	0.8	0.7	0.6	0.5	0.4
<i>Vermilion</i>	8.1	9.7	8.1	6.7	5.6	4.6	3.7
Slope Rockfish North	1,414	1,844	1,706	1,580	1,462	1,352	1,243
<b>Aurora</b>	<b>12.8</b>	<b>17.5</b>	<b>16.7</b>	<b>15.9</b>	<b>15.1</b>	<b>14.3</b>	<b>13.5</b>
<i>Bank</i>	14.4	17.2	14.4	12.0	9.9	8.1	6.5
<i>Blackgill</i>	3.9	4.7	3.9	3.3	2.7	2.2	1.8
<i>Redbanded</i>	37.7	45.3	37.7	31.4	26.0	21.3	17.2
<b>Rougheye/Blackspotted</b>	<b>59.3</b>	<b>206.8</b>	<b>189</b>	<b>172</b>	<b>157</b>	<b>142</b>	<b>127</b>

Stock	Status Quo 2014 ABC	2016 OFL	Range of Alternative 2016 ABCs				
			Overfishing Probability (P*)				
			0.45	0.40	0.35	0.30	0.25
<b>Sharpchin</b>	<b>178.9</b>	<b>323.2</b>	<b>295.1</b>	<b>269.2</b>	<b>245.0</b>	<b>221.7</b>	<b>198.8</b>
<i>Shorotraker</i>	15.6	18.7	15.6	13.0	10.7	8.8	7.1
<i>Splimose</i>	931.3	1,018.2	973.4	929.6	885.8	843.0	798.2
<i>Yellowmouth</i>	160.5	192.4	160.5	133.6	110.5	90.4	72.9
Nearshore Rockfish South	1,001	1,291	1,148	1,023	910	807	710
<i>Shallow Nearshore Species</i>	NA	NA	NA	NA	NA	NA	NA
<i>Black and yellow</i>	23.0	27.5	23.0	19.1	15.8	12.9	10.4
<b>China</b>	<b>13.8</b>	<b>55.2</b>	<b>50.4</b>	<b>46.0</b>	<b>41.9</b>	<b>37.9</b>	<b>34.0</b>
<i>Gopher (N of Pt. Conception)</i>	146.3	144.0	137.7	131.5	125.3	119.2	112.9
<i>Gopher (S of Pt. Conception)</i>	21.4	25.6	21.4	17.8	14.7	12.0	9.7
<i>Grass</i>	49.7	59.6	49.7	41.4	34.2	28.0	22.6
<i>Kelp</i>	23.1	27.7	23.1	19.2	15.9	13.0	10.5
<i>Deeper Nearshore Species</i>	NA	NA	NA	NA	NA	NA	NA
<i>Blue (assessed area)</i>	171.4	190.3	173.8	158.5	144.3	130.6	117.0
<i>Blue (S of 34°27' N. lat.)</i>	60.8	72.9	60.8	50.6	41.8	34.3	27.6
<b>Brown</b>	<b>170.6</b>	<b>160.2</b>	<b>146.3</b>	<b>133.5</b>	<b>121.4</b>	<b>109.9</b>	<b>98.5</b>
<i>Calico</i>	0.0	-	-	-	-	-	-
<b>Copper</b>	<b>118.0</b>	<b>284.3</b>	<b>259.6</b>	<b>236.9</b>	<b>215.5</b>	<b>195.1</b>	<b>174.9</b>
<i>Olive</i>	187.4	224.6	187.4	155.9	128.9	105.6	85.1
<i>Quillback</i>	4.5	5.4	4.5	3.7	3.1	2.5	2.0
<i>Treefish</i>	11.0	13.2	11.0	9.2	7.6	6.2	5.0
Shelf Rockfish South	1,620	1,919	1,626	1,376	1,160	971	803
<i>Bronzespotted</i>	3.0	3.6	3.0	2.5	2.1	1.7	1.4
<i>Chameleon</i>	0.0	-	-	-	-	-	-
<i>Flag</i>	19.5	23.4	19.5	16.3	13.4	11.0	8.9
<i>Freckled</i>	0.0	-	-	-	-	-	-
<i>Greenblotched</i>	19.3	23.1	19.3	16.1	13.3	10.9	8.8
<i>Greenspotted</i>	73.3	82.0	74.9	68.3	62.2	56.3	50.4
<i>Greenstriped</i>	212.4	237.0	216.4	197.4	179.6	162.6	145.8
<i>Halfbanded</i>	0.0	-	-	-	-	-	-
<i>Harlequin</i>	0.0	-	-	-	-	-	-
<i>Honeycomb</i>	8.2	9.9	8.2	6.8	5.7	4.6	3.7
<i>Mexican</i>	4.2	5.1	4.2	3.5	2.9	2.4	1.9
<i>Pink</i>	2.1	2.5	2.1	1.8	1.5	1.2	1.0
<i>Pinkrose</i>	0.0	-	-	-	-	-	-
<i>Pygmy</i>	0.0	-	-	-	-	-	-
<i>Redstripe</i>	0.4	0.5	0.4	0.3	0.3	0.2	0.2
<i>Rosethorn</i>	1.8	2.1	1.8	1.5	1.2	1.0	0.8
<i>Rosy</i>	37.1	44.5	37.1	30.9	25.5	20.9	16.9
<i>Silvergray</i>	0.4	0.5	0.4	0.4	0.3	0.3	0.2
<i>Speckled</i>	32.8	39.4	32.8	27.3	22.6	18.5	14.9
<i>Squarespot</i>	9.2	11.1	9.2	7.7	6.4	5.2	4.2
<i>Starry</i>	52.2	62.6	52.2	43.4	35.9	29.4	23.7
<b>Stripetail</b>	<b>19.7</b>	<b>23.6</b>	<b>19.7</b>	<b>16.4</b>	<b>13.6</b>	<b>11.1</b>	<b>9.0</b>

Stock	Status Quo 2014 ABC	2016 OFL	Range of Alternative 2016 ABCs				
			Overfishing Probability (P*)				
			0.45	0.40	0.35	0.30	0.25
<i>Swordspine</i>	11.9	14.2	11.9	9.9	8.2	6.7	5.4
<i>Tiger</i>	0.0	0.04	0.03	0.03	0.02	0.02	0.02
<i>Vermilion</i>	224.6	269.3	224.6	186.9	154.6	126.6	102.1
<i>Yellowtail</i>	887.7	1,064.4	887.7	738.7	611.0	500.3	403.4
Slope Rockfish South	622	814	705	612	530	457	391
<b><i>Aurora</i></b>	<b>21.7</b>	<b>74.3</b>	<b>70.7</b>	<b>67.3</b>	<b>63.9</b>	<b>60.6</b>	<b>57.1</b>
<i>Bank</i>	459.4	503.2	419.7	349.2	288.8	236.5	190.7
<i>Blackgill</i>	122.3	140.0	127.8	116.6	106.1	96.0	86.1
<i>Pacific ocean perch</i>	0.0	-	-	-	-	-	-
<i>Redbanded</i>	8.7	10.4	8.7	7.2	6.0	4.9	3.9
<b><i>Rougheye/Blackspotted</i></b>	<b>0.3</b>	<b>4.2</b>	<b>3.9</b>	<b>3.5</b>	<b>3.2</b>	<b>2.9</b>	<b>2.6</b>
<b><i>Sharpchin</i></b>	<b>8.2</b>	<b>80.8</b>	<b>73.8</b>	<b>67.3</b>	<b>61.2</b>	<b>55.4</b>	<b>49.7</b>
<i>Shortraker</i>	0.1	0.10	0.09	0.07	0.06	0.05	0.04
<i>Yellowmouth</i>	0.7	0.8	0.7	0.6	0.5	0.4	0.3
Other Flatfish	6,982	9,645	8,356	7,243	6,264	5,388	4,589
<i>Butter sole</i>	3.2	4.6	3.9	3.2	2.7	2.2	1.8
<i>Curlfin sole</i>	5.7	8.2	6.9	5.7	4.7	3.9	3.1
<i>Flathead sole</i>	24.3	35.0	29.2	24.3	20.1	16.5	13.3
<b><i>Pacific sanddab</i></b>	<b>3,331.9</b>	<b>4,801.0</b>	<b>4,004.0</b>	<b>3,331.9</b>	<b>2,755.8</b>	<b>2,256.5</b>	<b>1,819.6</b>
<b><i>Rex sole</i></b>	<b>3,033.8</b>	<b>3,956.0</b>	<b>3,611.8</b>	<b>3,295.3</b>	<b>2,998.6</b>	<b>2,713.8</b>	<b>2,432.9</b>
<i>Rock sole</i>	46.3	66.7	55.6	46.3	38.3	31.3	25.3
<i>Sand sole</i>	536.6	773.2	644.8	536.6	443.8	363.4	293.0
Other Fish	NA	290.8	242.5	201.8	166.9	136.7	110.2
<i>Cabezon (WA)</i>	a/	4.8	4.0	3.3	2.8	2.3	1.8
<i>Kelp greenling (CA)</i>	82.5	118.9	99.2	82.5	68.2	55.9	45.1
<i>Kelp greenling (OR)</i>	a/	a/	a/	a/	a/	a/	a/
<i>Kelp greenling (WA)</i>	a/	a/	a/	a/	a/	a/	a/
<i>Leopard shark</i>	116.0	167.1	139.4	116.0	95.9	78.5	63.3

a/ No ABC contribution for these stocks given the lack of an approved method for estimating the OFL.

**Table 1-20. Preferred 2015 and 2016 ABCs (mt) for west coast groundfish stocks and stock complexes and the associated ABC harvest control rule (P\* = overfishing probability) (overfished stocks in CAPS; stocks with new assessments in bold; component stocks in stock complexes in *italics*).**

Stock	Preferred ABCs		
	2015	2016	ABC Harvest Control Rule
<b>OVERFISHED STOCKS</b>			
<b>BOCACCIO S. of 40°10' N. lat.</b>	<b>1,380</b>	<b>1,291</b>	<b>P* = 0.45</b>
CANARY	701	697	P* = 0.45
COWCOD S. of 40°10' N. lat.	<b>60</b>	62	<b>P* = 0.45</b>
<i>COWCOD (Conception)</i>	<b>50</b>	<b>51</b>	<b>P* = 0.45</b>
<i>COWCOD (Monterey)</i>	10	10	P* = 0.45
<b>DARKBLOTCHED</b>	<b>549</b>	<b>554</b>	<b>P* = 0.45</b>
PACIFIC OCEAN PERCH	805	813	P* = 0.45
<b>PETRALE SOLE</b>	<b>2,816</b>	<b>2,910</b>	<b>P* = 0.45</b>
YELLOWEYE	43	43	P* = 0.4
<b>NON-DEPLETED STOCKS</b>			
Arrowtooth Flounder	5,497	5,328	P* = 0.4
Black Rockfish (OR-CA)	1,124	1,131	P* = 0.45
Black Rockfish (WA)	402	404	P* = 0.45
Cabezon (CA)	154	151	P* = 0.45
Cabezon (OR)	47	47	P* = 0.45
California scorpionfish	114	111	P* = 0.45
Chilipepper S. of 40°10' N. lat.	1,628	1,619	P* = 0.45
Dover Sole	63,929	56,615	P* = 0.45
<b>English Sole</b>	<b>9,853</b>	<b>7,204</b>	<b>P* = 0.45</b>
Lingcod N. of 40°10' N. lat.	2,830	2,719	P* = 0.45
Lingcod S. of 40°10' N. lat.	1,004	946	P* = 0.4
Longnose skate	2,341	2,299	P* = 0.45
<b>Longspine Thornyhead (coastwide)</b>	<b>4,171</b>	<b>3,968</b>	<b>P* = 0.4</b>
Pacific Cod	2,221	2,221	P* = 0.4
Sablefish (coastwide)	7,173	7,784	P* = 0.4
Shortbelly	5,789	5,789	P* = 0.4
<b>Shortspine Thornyhead (coastwide)</b>	<b>2,668</b>	<b>2,640</b>	<b>P* = 0.4</b>
Spiny dogfish	2,101	2,085.2	P* = 0.4
Splitnose S. of 40°10' N. lat.	1,715	1,746	P* = 0.45
Starry Flounder	1,681	1,686	P* = 0.4
Widow	3,929	3,790	P* = 0.45
<b>Yellowtail N. of 40°10' N. lat.</b>	<b>6,590</b>	<b>6,344</b>	<b>P* = 0.45</b>
<b>STOCK COMPLEXES</b>			
Nearshore Rockfish North	77	77	P* = 0.45
<i>Black and yellow</i>	0.0	0.0	P* = 0.45
<i>Blue (CA)</i>	25.0	25.3	P* = 0.45
<i>Blue (OR &amp; WA)</i>	26.9	26.9	P* = 0.45
<b>Brown</b>	<b>1.7</b>	<b>1.7</b>	<b>P* = 0.45</b>
<i>Calico</i>	-	-	P* = 0.45
<b>China</b>	<b>6.6</b>	<b>6.8</b>	<b>P* = 0.45</b>

Stock	Preferred ABCs		
	2015	2016	ABC Harvest Control Rule
<i>Copper</i>	9.7	9.4	<b>P* = 0.45</b>
<i>Gopher</i>	-	-	P* = 0.45
<i>Grass</i>	0.5	0.5	P* = 0.45
<i>Kelp</i>	0.0	0.0	P* = 0.45
<i>Olive</i>	0.3	0.3	P* = 0.45
<i>Quillback</i>	6.2	6.2	P* = 0.45
<i>Treefish</i>	0.2	0.2	P* = 0.45
Shelf Rockfish North	1,944	1,953	P* = 0.45
<i>Bronzespotted</i>	-	-	P* = 0.45
<i>Bocaccio</i>	236.9	236.9	P* = 0.45
<i>Chameleon</i>	-	-	P* = 0.45
<i>Chilipepper</i>	106.9	106.4	P* = 0.45
<i>Cowcod</i>	0.3	0.3	P* = 0.45
<i>Flag</i>	0.1	0.06	P* = 0.45
<i>Freckled</i>	-	-	P* = 0.45
<i>Greenblotched</i>	1.1	1.1	P* = 0.45
<i>Greenspotted 40°10' to 42° N. lat.</i>	9.1	9.0	P* = 0.45
<i>Greenspotted N. of 42 N. lat. (OR &amp; WA)</i>	5.1	5.1	P* = 0.45
<i>Greenstriped</i>	1,170.3	1,179.6	P* = 0.45
<i>Halfbanded</i>	-	-	P* = 0.45
<i>Harlequin</i>	-	-	P* = 0.45
<i>Honeycomb</i>	-	-	P* = 0.45
<i>Mexican</i>	-	-	P* = 0.45
<i>Pink</i>	0.0	0.003	P* = 0.45
<i>Pinkrose</i>	-	-	P* = 0.45
<i>Puget Sound</i>	-	-	P* = 0.45
<i>Pygmy</i>	-	-	P* = 0.45
<i>Redstripe</i>	225.1	225.1	P* = 0.45
<i>Rosethorn</i>	10.8	10.8	P* = 0.45
<i>Rosy</i>	2.5	2.5	P* = 0.45
<i>Silvergray</i>	133.0	133.0	P* = 0.45
<i>Speckled</i>	0.1	0.14	P* = 0.45
<i>Squarespot</i>	0.1	0.14	P* = 0.45
<i>Starry</i>	0.0	0.003	P* = 0.45
<i>Stripetail</i>	33.7	33.7	<b>P* = 0.45</b>
<i>Swordspine</i>	0.0	0.0	P* = 0.45
<i>Tiger</i>	0.8	0.8	P* = 0.45
<i>Vermilion</i>	8.1	8.1	P* = 0.45
Slope Rockfish North	1,693	1,706	P* = 0.45
<i>Aurora</i>	16.6	16.7	<b>P* = 0.45</b>
<i>Bank</i>	14.4	14.4	P* = 0.45
<i>Blackgill</i>	3.9	3.9	P* = 0.45
<i>Redbanded</i>	37.7	37.7	P* = 0.45
<i>Rougheye/Blackspotted</i>	184.3	189	<b>P* = 0.45</b>



Stock	Preferred ABCs		
	2015	2016	ABC Harvest Control Rule
<b>Sharpchin</b>	<b>303.8</b>	<b>295.1</b>	<b>P* = 0.45</b>
<i>Shorthead</i>	15.6	15.6	P* = 0.45
<i>Splitnose</i>	956.6	973.4	P* = 0.45
<i>Yellowmouth</i>	160.5	160.5	P* = 0.45
Nearshore Rockfish South	1,169	1,148	P* = 0.45
<i>Shallow Nearshore Species</i>	NA	NA	P* = 0.45
<i>Black and yellow</i>	23.0	23.0	P* = 0.45
<b>China</b>	<b>50.4</b>	<b>50</b>	<b>P* = 0.45</b>
<i>Gopher (N of Pt. Conception)</i>	141.5	137.7	P* = 0.45
<i>Gopher (S of Pt. Conception)</i>	21.4	21.4	P* = 0.45
<i>Grass</i>	49.7	49.7	P* = 0.45
<i>Kelp</i>	23.1	23.1	P* = 0.45
<i>Deeper Nearshore Species</i>	NA	NA	P* = 0.45
<i>Blue (assessed area)</i>	172.2	173.8	P* = 0.45
<i>Blue (S of 34°27' N. lat.)</i>	60.8	60.8	P* = 0.45
<b>Brown</b>	<b>149.5</b>	<b>146</b>	<b>P* = 0.45</b>
<i>Calico</i>	-	-	P* = 0.45
<b>Copper</b>	<b>274.9</b>	<b>260</b>	<b>P* = 0.45</b>
<i>Olive</i>	187.4	187.4	P* = 0.45
<i>Quillback</i>	4.5	4.5	P* = 0.45
<i>Treefish</i>	11.0	11.0	P* = 0.45
Shelf Rockfish South	1,625	1,626	P* = 0.45
<i>Bronzespotted</i>	3.0	3.0	P* = 0.45
<i>Chameleon</i>	-	-	P* = 0.45
<i>Flag</i>	19.5	19.5	P* = 0.45
<i>Freckled</i>	-	-	P* = 0.45
<i>Greenblotched</i>	19.3	19.3	P* = 0.45
<i>Greenspotted</i>	75.6	74.9	P* = 0.45
<i>Greenstriped</i>	214.7	216.4	P* = 0.45
<i>Halfbanded</i>	-	-	P* = 0.45
<i>Harlequin</i>	-	-	P* = 0.45
<i>Honeycomb</i>	8.2	8.2	P* = 0.45
<i>Mexican</i>	4.2	4.2	P* = 0.45
<i>Pink</i>	2.1	2.1	P* = 0.45
<i>Pinkrose</i>	-	-	P* = 0.45
<i>Pygmy</i>	-	-	P* = 0.45
<i>Redstripe</i>	0.4	0.4	P* = 0.45
<i>Rosethorn</i>	1.8	1.8	P* = 0.45
<i>Rosy</i>	37.1	37.1	P* = 0.45
<i>Silvergray</i>	0.4	0.4	P* = 0.45
<i>Speckled</i>	32.8	32.8	P* = 0.45
<i>Squarespot</i>	9.2	9.2	P* = 0.45
<i>Starry</i>	52.2	52.2	P* = 0.45
<b>Stripetail</b>	<b>19.7</b>	<b>19.7</b>	<b>P* = 0.45</b>

Stock	Preferred ABCs		
	2015	2016	ABC Harvest Control Rule
<i>Swordspine</i>	11.9	11.9	P* = 0.45
<i>Tiger</i>	0.0	0.03	P* = 0.45
<i>Vermilion</i>	224.6	224.6	P* = 0.45
<i>Yellowtail</i>	887.7	887.7	P* = 0.45
Slope Rockfish South	704.7	705	P* = 0.45
<b><i>Aurora</i></b>	<b>70.7</b>	<b>70.7</b>	<b>P* = 0.45</b>
<i>Bank</i>	419.7	419.7	P* = 0.45
<i>Blackgill</i>	125.1	127.8	P* = 0.45
<i>Pacific ocean perch</i>	-	-	P* = 0.45
<i>Redbanded</i>	8.7	8.7	P* = 0.45
<b><i>Rougheye/Blackspotted</i></b>	<b>3.8</b>	<b>4</b>	<b>P* = 0.45</b>
<b><i>Sharpchin</i></b>	<b>76.0</b>	<b>73.8</b>	<b>P* = 0.45</b>
<i>Shortraker</i>	0.1	0.09	P* = 0.45
<i>Yellowmouth</i>	0.7	0.7	P* = 0.45
Other Flatfish	8,749	7,243	P* = 0.4
<i>Butter sole</i>	3.2	3.2	P* = 0.4
<i>Curlfin sole</i>	5.7	5.7	P* = 0.4
<i>Flathead sole</i>	24.3	24.3	P* = 0.4
<b><i>Pacific sanddab</i></b>	<b>3,331.9</b>	<b>3,331.9</b>	<b>P* = 0.4</b>
<b><i>Rex sole</i></b>	<b>4,801.4</b>	<b>3,295.3</b>	<b>P* = 0.4</b>
<i>Rock sole</i>	46.3	46.3	P* = 0.4
<i>Sand sole</i>	536.6	536.6	P* = 0.4
Other Fish	242	243	P* = 0.45
<i>Cabezon (WA)</i>	3.8	4.0	P* = 0.45
<i>Kelp greenling (CA)</i>	99.2	99.2	P* = 0.45
<i>Kelp greenling (OR)</i>	a/	a/	P* = 0.45
<i>Kelp greenling (WA)</i>	a/	a/	P* = 0.45
<i>Leopard shark</i>	139.4	139.4	P* = 0.45

a/ No OFL or ABC contribution for these stocks given the lack of an approved method for estimating the OFL.

### 1.3.3 Annual Catch Limits

Annual catch limits (ACLs) are specified for each stock and stock complex that is “in the fishery” as specified under the FMP framework. An ACL is a harvest specification set equal to the ABC or below the ABC in consideration of conservation objectives, management uncertainty, socioeconomic considerations, ecological considerations, and other factors (e.g. rebuilding considerations) needed to meet management objectives. Sector-specific ACLs may be specified in cases where a sector has a formal, long-term allocation of the harvestable surplus of a stock or stock complex. The ACL counts all sources of fishing-related mortality including landed catch, discard mortalities, research catches, and set-asides for exempted fishing permits (EFPs).

Under the FMP, the biomass level that produces MSY ( $B_{MSY}$ ) is defined as the precautionary threshold. When the biomass for an assessed category 1 or 2 stock falls below the precautionary threshold, the harvest rate will be reduced to help the stock return to the  $B_{MSY}$  level, which is the management target for groundfish stocks. If a stock biomass is larger than  $B_{MSY}$ , the ACL may be set equal to or less than ABC. Because  $B_{MSY}$  is a long-term average, the true biomass could be below  $B_{MSY}$  in some years and above  $B_{MSY}$  in other years. Even in the absence of overfishing, biomass may decline to levels below  $B_{MSY}$  due to natural fluctuations in recruitment. The minimum stock size threshold (MSST) is the biomass threshold for declaring a stock overfished. When spawning stock biomass falls below the MSST, a rebuilding plan must be developed that determines the strategy for rebuilding the stock in the shortest time possible while considering impacts to fishing-dependent communities and other factors. When spawning stock biomass is below  $B_{MSY}$  yet above the MSST, the stock is considered to be in the precautionary zone. The current proxy  $B_{MSY}$  and MSST reference points for west coast groundfish stocks are as follows:


- Assessed flatfish stocks:  $B_{MSY} = 25$  percent of initial biomass or  $B_{25\%}$ ; MSST = 12.5 percent of initial biomass or  $B_{12.5\%}$  (PFMC and NMFS 2011); and
- All other assessed groundfish stocks:  $B_{MSY} = 40$  percent of initial biomass or  $B_{40\%}$ ; MSST = 25 percent of initial biomass or  $B_{25\%}$ .

These reference points are only used to manage assessed stocks since they require estimates of spawning stock biomass.

West coast groundfish stocks are managed with harvest control rules that calculate ACLs below the ABCs when spawning biomass is estimated to be in the precautionary zone. These harvest control rules are designed to prevent a stock from becoming overfished. The FMP defines the 40-10 harvest control rule for stocks with a  $B_{MSY}$  proxy of  $B_{40\%}$  that are in the precautionary zone. The analogous harvest control rule for assessed flatfish stocks is the 25-5 harvest control rule. Both ACL harvest control rules are applied after the ABC deduction is made. The further the stock biomass is below the precautionary threshold, the greater the reduction in ACL relative to the ABC, until at  $B_{10\%}$  for a stock with a  $B_{MSY}$  proxy of  $B_{40\%}$  or  $B_{5\%}$  for a stock with a  $B_{MSY}$  proxy of  $B_{25\%}$ , the ACL would be set at zero<sup>11</sup> (Figure 1-39). These harvest policies foster a quicker return to the  $B_{MSY}$  level and serve as an interim rebuilding policy for stocks that are below the MSST. The Council may recommend setting the ACL higher than what the default ACL harvest control rule specifies as long as the ACL does not exceed the ABC, complies with the requirements of the MSA, and is consistent with the FMP and National Standard Guidelines. Additional precautionary adjustments may be made to an ACL if necessary to address management uncertainty, conservation concerns, socioeconomic concerns, ecological considerations, and the other factors that are considered when setting ACLs.

---

<sup>11</sup> The lower  $B_{10\%}$  and  $B_{5\%}$  thresholds in the precautionary ACL harvest control rules are used to establish the slope of the ACL curve in Figure 36. These precautionary ACL control rules only apply for stocks in the precautionary zone ( $B_{MSY} > B_{CURRENT} > MSST$ ). A rebuilding plan governs the ACL harvest control rule for any stock that falls below the MSST and is designated as overfished.



**Figure 1-39. Conceptual diagram of the 25-5 and 40-10 ACL harvest control rules used to manage assessed west coast flatfish and other groundfish species, respectively, that are in the precautionary zone.**

The ACL serves as the basis for invoking accountability measures (AMs), which are management measures or mechanisms used to address any management uncertainty that may result in exceeding an ACL. If ACLs are exceeded more often than 1 in 4 years, then AMs, such as catch monitoring and inseason adjustments to fisheries, need to improve or additional AMs may need to be implemented. Additional AMs may include setting an annual catch target (ACT), which is a specified level of harvest below the ACL. The use of ACTs may be especially important for a stock subject to highly uncertain inseason catch monitoring. A sector-specific ACT may serve as a harvest guideline (HG) for a sector or may be used strategically in a rebuilding plan to attempt to reduce mortality of an overfished stock more than the rebuilding plan limits prescribe.

The Council has the discretion to adjust the ACLs for uncertainty on a case-by-case basis. In cases where there is a high degree of uncertainty about the condition of the stock or stocks, the ACL may be reduced accordingly. Most category 3 species are managed in a stock complex (such as the minor rockfish complexes and the Other Flatfish complex) where harvest specifications are set for the complex in its

entirety. For stock complexes, the ACL will be less than or equal to the sum of the individual component ABCs. The ACL may be adjusted below the sum of component ABCs as appropriate.

For most stocks and stock complexes, the Council elected to use the same general policies for deciding 2015 and 2016 ACLs as were used for deciding the 2014 ACLs (No Action) (Table 1-21). The No Action ACLs are the 2014 ACLs specified in Federal regulations.

Section 4.6.3 of the FMP states the Council's general policies on rebuilding overfished stocks. Section 4.6.3.1 of the FMP specifies the overall goals of rebuilding programs are to (1) achieve the population size and structure that will support the MSY within a specified time period that is as short as possible, taking into account the status and biology of the stock, the needs of fishing communities, and the interaction of the stock of fish within the marine ecosystem; (2) minimize, to the extent practicable, the adverse social and economic impacts associated with rebuilding, including adverse impacts on fishing communities; (3) fairly and equitably distribute both the conservation burdens (overfishing restrictions) and recovery benefits among commercial, recreational, and charter fishing sectors; (4) protect the quantity and quality of habitat necessary to support the stock at healthy levels in the future; and (5) promote widespread public awareness, understanding and support for the rebuilding program. These overall goals are derived from and consistent with the requirements of the MSA. The first goal embodies MSA National Standard 1 (NS1) and the requirements for rebuilding overfished stocks found at MSA section 304(e)(4)(A). The third goal is required by MSA section 304(e)(4)(B). The fourth and fifth goals represent additional policy preferences of the Council that recognize the importance of habitat protection to the rebuilding of some fish stocks and the desire for public outreach and education on the complexities—biological, economic, and social issues—involved with rebuilding overfished stocks. Overfished groundfish species are those with spawning biomasses that have dropped below the Council's MSST (i.e., 25 percent of initial spawning biomass or  $B_{25\%}$  for all groundfish species other than flatfish where the MSST is  $B_{12.5\%}$ ). The FMP requires these stocks to be rebuilt to a target biomass that supports MSY (i.e.,  $B_{MSY}$  or  $B_{40\%}$  for all groundfish species other than flatfish where the target is  $B_{25\%}$ ).

Rebuilding plans are in place for six overfished rockfish species, as well as petrale sole, where assessments have indicated spawning biomass has declined to below the MSST. New full and updated assessments and rebuilding analyses were done in 2013 inform the 2015 and 2016 harvest specifications for many of the overfished species. New full assessments were conducted for cowcod, darkblotched rockfish, and petrale sole in 2013; however, a new rebuilding analysis was only prepared for cowcod. The results of the new assessments for darkblotched rockfish and petrale sole indicated those stocks would be rebuilt by 2015 and 2014, respectively. The SSC did not recommend new rebuilding analyses for these two stocks given their imminent rebuilding expectation. An update assessment for bocaccio was prepared in 2013. Like darkblotched, the stock is predicted to rebuild by 2015 and the SSC therefore recommended no new rebuilding analysis be prepared. Catch reports for canary rockfish, Pacific ocean perch, and yelloweye rockfish were prepared in 2013. These catch reports indicated total catches were within limits prescribed in these stocks' respective rebuilding plans.

**Table 1-21. The status quo 2014 and preferred 2015 and 2016 annual catch limits (ACLs in mt) for west coast groundfish stocks and stock complexes (stocks with new assessments in bold).**

Stock	Status Quo ACL	Preferred ACLs		
	2014	2015	2016	ACL Harvest Control Rule
<b>OVERFISHED STOCKS</b>				
<b>BOCACIO S. of 40°10'</b>	337	<b>349</b>	<b>362</b>	SPR = 77.7%
CANARY	119	122	125	SPR = 88.7%
<b>COWCOD S. of 40°10'</b>	3	<b>10</b>	<b>10</b>	SPR = 82.7% (F = 0.007); ACT = 4 mt
<b>DARKBLOTCHED</b>	330	<b>338</b>	<b>346</b>	SPR = 64.9%
PACIFIC OCEAN PERCH	153	158	164	SPR = 86.4%
<b>PETRALE SOLE</b>	2,652	<b>2,816</b>	<b>2,910</b>	25-5 rule
YELLOWEYE	18	18	19	SPR = 76.0%
<b>NON-DEPLETED STOCKS</b>				
Arrowtooth Flounder	5,758	5,497	5,328	ACL = ABC (P* = 0.4)
Black Rockfish (OR-CA)	1,000	1,000	1,000	Constant catch strategy
Black Rockfish (WA)	409	402	404	ACL = ABC (P* = 0.45)
Cabazon (CA)	158	154	151	ACL = ABC (P* = 0.45)
Cabazon (OR)	47	47	47	ACL = ABC (P* = 0.45)
California scorpionfish	117	114	111	ACL = ABC (P* = 0.45)
Chilipepper S. of 40°10'	1,647	1,628	1,619	ACL = ABC (P* = 0.45)
Dover Sole	25,000	50,000	50,000	Constant catch strategy; alt. ACL = 50,000 mt
<b>English Sole</b>	5,646	<b>9,853</b>	<b>7,204</b>	ACL = ABC (P* = 0.45)
Lingcod N. of 40°10'	2,878	2,830	2,719	ACL = ABC (P* = 0.45)
Lingcod S. of 40°10'	1,063	1,004	946	ACL = ABC (P* = 0.4)
Longnose skate	2,000	2,000	2,000	Constant catch strategy
<b>Longspine Thornyhead N. of 34°27'</b>	1,958	<b>3,170</b>	<b>3,015</b>	ACL = 76% of coastwide ABC (P* = 0.4)
<b>Longspine Thornyhead S. of 34°27'</b>	347	<b>1,001</b>	<b>952</b>	ACL = 24% of coastwide ABC (P* = 0.4)
Pacific Cod	1,600	1,600	1,600	ACL = 50% of OFL
Sablefish N. of 36°	4,349	4,793	5,241	40-10 rule applied to 73.6% of coastwide ABC (P* = 0.4)
Sablefish S. of 36°	1,560	1,719	1,880	40-10 rule applied to 26.4% of coastwide ABC (P* = 0.4)
Shortbelly	50	500	500	ACL increased from No Action to accommodate incidental bycatch in emerging midwater trawl fisheries; harvestable surplus to be available as forage
<b>Shortspine Thornyhead N. of 34°27'</b>	1,525	<b>1,745</b>	<b>1,726</b>	ACL = 65.4% of coastwide ABC (P* = 0.4)

Stock	Status Quo ACL	Preferred ACLs		
	2014	2015	2016	ACL Harvest Control Rule
<b>Shortspine Thornyhead S. of 34°27'</b>	393	<b>923</b>	<b>913</b>	ACL = 34.6% of coastwide ABC (P* = 0.4)
Spiny dogfish	NA	2,101	2,085	ACL = ABC (P* = 0.4)
Splitnose S. of 40°10'	1,670	1,715	1,746	ACL = ABC (P* = 0.45)
Starry Flounder	1,528	1,534	1,539	ACL = ABC (P* = 0.4)
Widow	1,500	2,000	2,000	Constant catch strategy; ACL = 2,000 mt
<b>Yellowtail N. of 40°10'</b>	4,382	<b>6,590</b>	<b>6,344</b>	ACL = ABC (P* = 0.45)
<b>STOCK COMPLEXES</b>				
Nearshore Rockfish North	94	69	69	ACL = ABC (P* = 0.45); 40-10 adj. ACL contrib. for blue RF in CA and China RF
Shelf Rockfish North	968	1,944	1,952	ACL = ABC (P* = 0.45); 40-10 adj. ACL contrib. for greenspotted RF in CA
Slope Rockfish North	1,160	1,693	1,706	ACL = ABC (P* = 0.45)
Nearshore Rockfish South	990	1,114	1,006	ACL = ABC (P* = 0.45); 40-10 adj. ACL contrib. for blue RF N of 34°27' N lat.
Shelf Rockfish South	714	1,624	1,625	ACL = ABC (P* = 0.45); 40-10 adj. ACL contrib. for greenspotted RF in CA
Slope Rockfish South	622	693	695	ACL = ABC (P* = 0.45); 40-10 adj. ACL contrib. for blackgill RF
Other Flatfish	4,884	8,749	7,243	ACL = ABC (P* = 0.4)
Other Fish	4,697	242	243	Sum of component species ACLs (ACLs = ABCs (P* = 0.45))
<i>Cabazon (WA)</i>	NA	3.7	4.0	ACL = ABC (P* = 0.45)
<i>Kelp greenling (CA)</i>	NA	99.2	99.2	ACL = ABC (P* = 0.45)
<i>Kelp greenling (OR) a/</i>	NA	NA	NA	No approved method for calculating the OFL contribution
<i>Kelp greenling (WA) a/</i>	NA	NA	NA	No approved method for calculating the OFL contribution
<i>Leopard shark</i>	NA	139.4	139.4	ACL = ABC (P* = 0.45)

Stock	Status Quo ACL	Preferred ACLs		
	2014	2015	2016	ACL Harvest Control Rule
<b>ECOSYSTEM COMPONENT SPECIES</b>				
Big skate	No harvest specifications for an EC species			
California skate				
Aleutian skate				
Roughtail/black skate				
Bering/sandpaper skate				
All other skates				
Pacific grenadier				
Giant grenadier				
All other grenadiers				
Ratfish				
Soupfin shark				
Finescale codling				

a/ No approved method for calculating the OFL contributions for these stocks in 2015-2016.



## CHAPTER 2      **SOCIOECONOMICS OF THE GROUND FISH FISHERY**

---

### **2.1      Socioeconomic Summary**

The following tables were compiled using data downloaded in April 2013 from the PacFIN vdrfd table and the NorPAC 4900 table for at-sea Pacific whiting sectors. Groundfish fisheries are defined within the vdrfd table in the “dahl\_sector” column. Nominal dollar values adjusted for inflation using Bureau of Economic Analysis Table 1.1.9 “Implicit Price Deflators for Gross Domestic Product, 1995=100” dated March 28, 2013.

A description of the dahl\_sector coding is included at the end of this section.

**Table 2-1. Landings (shoreside commercial and tribal) by species group (mt), 1981-2012.**

Year	CPS	CRAB	Groundfish	HMS	Other	Salmon	Shellfish	Shrimp	Total	Pct of Ann Average
1981	139,770	9,652	103,344	26,883	14,787	7,972	10,813	19,186	332,406	107%
1982	126,970	8,235	119,356	17,272	12,173	8,823	3,623	13,433	309,884	99%
1983	69,346	7,862	98,978	55,062	11,011	2,935	3,288	6,879	255,360	82%
1984	63,822	7,177	89,804	43,992	10,653	2,178	3,661	5,618	226,904	73%
1985	68,174	8,559	90,923	14,249	13,603	5,046	2,447	13,609	216,610	69%
1986	84,042	8,390	82,480	13,011	19,753	7,377	508	26,893	242,454	78%
1987	90,441	9,373	91,982	12,881	24,553	9,410	458	31,514	270,611	87%
1988	108,364	17,509	92,248	15,005	27,920	12,515	344	32,832	306,737	98%
1989	113,412	16,814	99,372	7,711	30,985	6,869	425	36,398	311,985	100%
1990	88,510	14,507	94,539	9,886	29,739	4,685	320	25,632	267,817	86%
1991	90,592	7,027	105,547	11,007	25,043	3,734	262	20,157	263,368	84%
1992	60,596	15,882	132,555	13,608	19,227	2,048	328	36,422	280,667	90%
1993	79,172	18,076	116,394	16,959	16,155	2,213	537	23,496	273,002	88%
1994	85,674	18,167	135,677	16,065	14,598	1,802	336	15,752	288,071	92%
1995	128,069	17,468	134,493	14,046	13,706	4,755	300	12,312	325,149	104%
1996	135,920	25,139	146,452	25,957	12,779	3,306	158	14,919	364,629	117%
1997	151,057	13,013	143,571	23,202	11,678	3,697	98	18,881	365,196	117%
1998	74,372	12,388	131,011	16,322	7,682	1,850	57	5,662	249,344	80%
1999	171,035	16,191	125,883	11,878	9,188	2,711	45	14,226	351,157	113%
2000	225,774	13,566	123,031	10,955	9,816	3,704	114	16,283	403,242	129%
2001	195,827	11,850	103,556	12,708	8,989	3,363	93	18,599	354,985	114%
2002	182,843	16,113	75,056	10,834	9,666	5,109	168	26,246	326,035	105%
2003	125,368	34,013	82,905	17,648	7,932	6,015	108	14,594	288,583	93%
2004	143,398	28,537	122,303	15,190	8,284	5,663	191	9,688	333,254	107%
2005	157,885	25,098	135,454	10,050	8,208	4,296	113	11,404	352,509	113%
2006	159,783	35,707	151,260	13,503	7,716	1,190	137	8,914	378,211	121%
2007	195,044	20,722	117,495	12,519	8,590	1,451	148	11,604	367,573	118%
2008	145,498	17,373	97,667	11,610	10,163	264	177	15,835	298,586	96%
2009	171,619	23,441	82,194	13,246	9,618	476	240	14,952	315,785	101%
2010	201,475	24,863	94,045	11,926	8,926	1,031	259	20,763	363,287	117%
2011	174,232	26,792	128,585	11,776	9,317	1,185	200	30,051	382,137	123%
2012	208,618	20,784	90,384	14,132	9,755	2,330	179	29,866	376,049	121%
<b>Total</b>	<b>4,216,702</b>	<b>550,284</b>	<b>3,538,542</b>	<b>531,092</b>	<b>442,212</b>	<b>130,002</b>	<b>30,131</b>	<b>602,619</b>	<b>10,041,584</b>	
Pct of Total	42%	5%	35%	5%	4%	1%	0%	6%	100%	
Ann Average	129,293	17,081	111,231	16,676	13,950	4,118	966	18,476	311,791	

**Table 2-2. Ex-vessel revenue (shoreside commercial and tribal) by species group in current (2012) dollars, \$1,000s, 1981-2012.**

Year	CPS	CRAB	Groundfish	HMS	Other	Salmon	Shellfish	Shrimp	Total	Pct of Ann Average
1981	\$59,680	\$43,250	\$99,360	\$96,687	\$23,753	\$70,120	\$27,104	\$51,459	\$471,413	130%
1982	\$58,446	\$41,171	\$124,177	\$50,687	\$20,059	\$77,789	\$14,583	\$36,922	\$423,833	116%
1983	\$50,204	\$49,450	\$104,735	\$118,259	\$20,092	\$18,183	\$9,867	\$25,767	\$396,557	109%
1984	\$25,646	\$45,052	\$93,393	\$106,863	\$20,511	\$20,716	\$11,676	\$19,136	\$342,992	94%
1985	\$36,476	\$48,946	\$104,769	\$49,319	\$25,310	\$39,028	\$11,401	\$24,801	\$340,050	93%
1986	\$35,178	\$45,031	\$103,392	\$46,522	\$35,713	\$46,099	\$3,973	\$61,303	\$377,212	104%
1987	\$37,000	\$48,619	\$127,638	\$53,329	\$38,634	\$81,970	\$4,707	\$86,593	\$478,491	132%
1988	\$42,311	\$77,072	\$117,690	\$54,284	\$50,513	\$117,051	\$3,747	\$53,704	\$516,373	142%
1989	\$37,472	\$68,464	\$114,014	\$29,611	\$57,754	\$44,329	\$5,551	\$50,467	\$407,661	112%
1990	\$35,479	\$75,863	\$102,573	\$29,671	\$64,952	\$35,040	\$4,338	\$46,585	\$394,502	108%
1991	\$35,493	\$35,797	\$112,096	\$26,281	\$73,952	\$21,900	\$3,146	\$40,474	\$349,139	96%
1992	\$28,189	\$61,027	\$107,278	\$38,097	\$62,280	\$13,960	\$5,146	\$44,558	\$360,535	99%
1993	\$24,753	\$65,077	\$94,081	\$43,503	\$55,447	\$13,160	\$6,666	\$29,036	\$331,725	91%
1994	\$31,226	\$77,503	\$97,997	\$43,526	\$52,800	\$10,478	\$4,210	\$33,694	\$351,434	97%
1995	\$53,118	\$91,621	\$122,590	\$31,052	\$48,845	\$21,833	\$4,532	\$31,633	\$405,223	111%
1996	\$59,598	\$105,157	\$113,988	\$57,171	\$42,620	\$12,954	\$3,184	\$33,171	\$427,843	118%
1997	\$60,609	\$72,798	\$110,234	\$48,246	\$39,954	\$13,776	\$1,533	\$31,298	\$378,448	104%
1998	\$13,257	\$64,549	\$71,877	\$34,640	\$26,015	\$7,700	\$94	\$18,571	\$236,704	65%
1999	\$57,182	\$90,813	\$76,162	\$31,446	\$31,957	\$12,876	\$65	\$26,692	\$327,194	90%
2000	\$54,390	\$82,109	\$79,651	\$29,679	\$34,940	\$18,152	\$208	\$26,679	\$325,808	90%
2001	\$40,900	\$67,313	\$63,837	\$30,808	\$30,163	\$13,454	\$226	\$21,293	\$267,995	74%
2002	\$40,665	\$74,472	\$52,936	\$21,548	\$29,262	\$17,951	\$457	\$26,757	\$264,047	73%
2003	\$42,829	\$144,262	\$58,633	\$34,541	\$25,910	\$25,773	\$165	\$14,026	\$346,140	95%
2004	\$38,551	\$122,800	\$55,574	\$35,187	\$26,173	\$36,791	\$558	\$13,095	\$328,729	90%
2005	\$49,789	\$96,980	\$63,237	\$26,521	\$23,793	\$27,808	\$323	\$16,234	\$304,685	84%
2006	\$45,409	\$146,062	\$67,733	\$29,978	\$24,588	\$11,351	\$400	\$12,196	\$337,716	93%
2007	\$48,641	\$113,854	\$63,983	\$27,507	\$23,641	\$13,835	\$391	\$16,597	\$308,449	85%
2008	\$47,496	\$95,833	\$74,119	\$33,028	\$29,010	\$2,150	\$521	\$24,660	\$306,817	84%
2009	\$77,109	\$109,774	\$69,427	\$31,752	\$27,298	\$2,632	\$711	\$15,854	\$334,556	92%
2010	\$87,805	\$116,459	\$70,256	\$32,118	\$30,545	\$9,318	\$716	\$20,971	\$368,188	101%
2011	\$79,681	\$159,591	\$94,146	\$47,467	\$35,898	\$11,138	\$522	\$39,210	\$467,653	129%
2012	\$86,956	\$151,408	\$73,448	\$46,790	\$35,593	\$21,970	\$489	\$37,964	\$454,618	125%
<b>Total</b>	<b>\$1,521,538</b>	<b>\$2,688,177</b>	<b>\$2,885,025</b>	<b>\$1,416,120</b>	<b>\$1,167,978</b>	<b>\$891,284</b>	<b>\$131,211</b>	<b>\$1,031,398</b>	<b>\$11,732,730</b>	
Pct of Total	13%	23%	25%	12%	10%	8%	1%	9%	100%	
Ann Average	\$46,277	\$81,831	\$90,696	\$44,172	\$36,529	\$28,042	\$4,217	\$32,046	\$363,810	

**Table 2-3. Groundfish landings (shoreside commercial and tribal) by species or species group (mt), 1981-2012.**

Year	P. Whiting	Sablefish	Lingcod	P. Cod	Other Roundfish	Rockfish	Thornyheads	Arrowtooth Flounder	Dover Sole	English Sole	Petrale Sole	Other Flatfish	Other Groundfish	Total	Pct of Ann Average
1981	839	11,419	3,304	1,237	41	57,779	1,801	1,074	16,468	2,711	2,041	3,672	960	103,344	93%
1982	1,027	18,627	3,840	908	46	59,316	2,158	2,351	21,000	2,793	2,630	3,931	728	119,356	108%
1983	1,051	14,652	4,252	597	16	46,289	1,749	2,077	20,084	2,356	2,214	3,001	639	98,978	90%
1984	2,721	14,015	4,029	585	25	36,819	3,189	2,379	19,307	1,721	1,739	2,660	615	89,804	81%
1985	3,894	14,132	3,839	411	18	33,263	4,069	2,679	20,616	1,929	1,840	3,460	772	90,923	82%
1986	3,465	13,150	1,891	331	37	33,387	3,610	2,230	17,396	2,039	1,750	2,761	433	82,480	75%
1987	4,795	12,602	2,587	2,281	38	36,526	3,747	2,830	18,489	2,482	2,205	2,915	485	91,982	83%
1988	6,868	10,744	2,767	3,345	41	35,193	5,663	1,946	18,185	2,103	2,149	2,731	514	92,248	83%
1989	7,414	10,285	3,563	2,189	43	37,238	8,085	3,553	18,881	2,412	2,153	2,969	589	99,372	90%
1990	9,633	9,065	2,907	1,064	22	33,179	10,084	5,824	15,753	1,912	1,765	2,505	828	94,539	85%
1991	23,970	9,501	3,167	1,796	24	28,737	6,515	4,945	18,274	2,185	1,927	3,239	1,267	105,547	95%
1992	56,128	9,361	1,888	1,778	37	28,121	8,873	3,573	16,074	1,626	1,554	2,018	1,525	132,555	120%
1993	42,108	8,147	2,210	1,370	23	29,005	9,224	2,713	14,371	1,603	1,503	1,938	2,180	116,394	105%
1994	73,617	7,579	1,907	866	50	23,539	8,048	3,249	9,389	1,124	1,375	2,439	2,497	135,677	123%
1995	74,963	7,915	1,469	505	102	21,980	7,555	2,321	10,593	1,133	1,659	2,559	1,740	134,493	122%
1996	85,129	8,317	1,559	445	124	21,669	6,532	2,192	12,187	1,154	1,829	1,999	3,318	146,452	132%
1997	87,417	7,943	1,569	595	175	18,245	5,504	2,344	10,126	1,505	1,948	2,310	3,891	143,571	130%
1998	87,857	4,384	350	413	221	16,298	3,526	3,169	8,023	1,140	1,463	1,700	2,468	131,011	118%
1999	83,471	6,648	358	280	191	10,815	2,648	5,285	9,141	913	1,498	2,015	2,621	125,883	114%
2000	85,855	6,281	146	279	193	8,887	2,378	3,276	8,780	769	1,893	1,622	2,672	123,031	111%
2001	73,412	5,637	156	324	424	5,673	1,761	2,465	6,890	993	1,845	1,717	2,258	103,556	94%
2002	45,708	3,798	206	752	3,876	2,962	2,716	2,085	6,301	1,175	1,797	1,724	1,957	75,056	68%
2003	55,336	5,420	165	1,250	338	1,668	2,402	2,327	7,356	931	2,070	1,575	2,068	82,905	75%
2004	96,504	5,755	178	1,403	163	2,112	1,477	2,327	6,746	952	1,962	1,394	1,328	122,303	111%
2005	109,053	6,208	203	851	297	1,900	1,312	2,240	6,903	929	2,734	1,237	1,587	135,455	122%
2006	127,166	6,199	260	367	68	1,463	1,460	1,922	5,970	912	2,610	1,215	1,649	151,260	137%
2007	91,442	5,241	268	89	68	1,493	1,822	2,262	9,279	690	2,253	972	1,617	117,495	106%
2008	67,761	5,871	285	38	72	1,524	2,679	2,668	11,217	363	2,220	811	2,159	97,667	88%
2009	49,223	7,198	233	236	84	2,073	2,703	3,844	11,753	357	1,767	971	1,751	82,194	74%
2010	64,654	6,829	173	345	68	2,280	2,712	3,228	10,391	221	803	745	1,595	94,045	85%
2011	103,190	6,407	376	604	89	2,552	1,904	2,292	7,745	170	932	686	1,638	128,585	116%
2012	66,369	5,246	484	631	84	2,937	1,773	2,243	7,066	189	1,094	690	1,577	90,384	82%
<b>Total</b>	<b>1,692,039</b>	<b>274,576</b>	<b>50,588</b>	<b>28,160</b>	<b>7,097</b>	<b>644,922</b>	<b>129,679</b>	<b>89,912</b>	<b>400,752</b>	<b>43,492</b>	<b>59,223</b>	<b>66,177</b>	<b>51,926</b>	<b>3,538,543</b>	
Pct of Total	48%	8%	1%	1%	0%	18%	4%	3%	11%	1%	2%	2%	1%	100%	
Ann. Average	52,876	8,580	1,581	880	222	20,154	4,052	2,810	12,524	1,359	1,851	2,068	1,623	110,579	

**Table 2-4. Groundfish ex-vessel revenue (shoreside commercial and tribal) by species or species group in current (2012) dollars, \$1,000s, 1981-2012.**

Year	P. Whiting	Sablefish	Lingcod	P. Cod	Other Roundfish	Rockfish	Thornyheads	Arrowtooth Flounder	Dover Sole	English Sole	Petrale Sole	Other Flatfish	Other Groundfish	Total	Pct of Ann Average
1981	\$311	\$11,605	\$3,669	\$1,128	\$31	\$47,223	\$1,960	\$499	\$17,528	\$3,905	\$5,084	\$5,724	\$691	\$99,358	110%
1982	\$377	\$21,576	\$4,380	\$926	\$40	\$52,974	\$2,228	\$1,180	\$22,179	\$4,068	\$7,305	\$6,296	\$649	\$124,175	137%
1983	\$376	\$15,379	\$4,683	\$622	\$34	\$45,757	\$1,789	\$915	\$19,748	\$3,344	\$6,669	\$4,841	\$572	\$104,731	115%
1984	\$781	\$12,857	\$4,207	\$579	\$25	\$39,359	\$3,254	\$978	\$18,850	\$2,351	\$5,235	\$4,178	\$731	\$93,387	103%
1985	\$1,067	\$19,950	\$4,188	\$416	\$17	\$39,234	\$4,194	\$1,093	\$20,294	\$2,651	\$5,578	\$5,327	\$760	\$104,769	116%
1986	\$830	\$20,157	\$2,434	\$366	\$24	\$42,888	\$4,067	\$918	\$18,065	\$2,960	\$5,483	\$4,600	\$597	\$103,389	114%
1987	\$1,184	\$23,907	\$3,852	\$2,944	\$32	\$50,592	\$4,765	\$1,627	\$22,072	\$3,929	\$7,044	\$5,048	\$644	\$127,641	141%
1988	\$1,933	\$21,798	\$3,867	\$3,361	\$46	\$42,721	\$7,554	\$873	\$20,907	\$3,137	\$6,645	\$4,260	\$585	\$117,686	130%
1989	\$1,753	\$17,664	\$4,605	\$2,073	\$41	\$42,201	\$10,936	\$1,294	\$18,934	\$3,232	\$6,440	\$4,242	\$600	\$114,015	126%
1990	\$2,335	\$15,489	\$3,657	\$1,014	\$29	\$38,141	\$13,665	\$2,157	\$14,810	\$2,212	\$5,126	\$3,320	\$615	\$102,568	113%
1991	\$5,986	\$22,154	\$3,795	\$1,837	\$42	\$34,553	\$10,095	\$1,935	\$18,707	\$2,565	\$5,410	\$4,255	\$765	\$112,098	124%
1992	\$8,931	\$20,558	\$2,440	\$1,927	\$104	\$34,097	\$13,524	\$1,262	\$15,044	\$1,808	\$4,161	\$2,620	\$800	\$107,277	118%
1993	\$4,295	\$14,753	\$2,749	\$1,444	\$196	\$33,853	\$14,080	\$867	\$12,761	\$1,661	\$3,844	\$2,572	\$1,010	\$94,083	104%
1994	\$7,278	\$19,869	\$2,519	\$927	\$432	\$29,423	\$18,594	\$1,023	\$8,838	\$1,227	\$3,684	\$2,988	\$1,195	\$97,995	108%
1995	\$11,119	\$33,232	\$2,108	\$612	\$971	\$29,075	\$23,769	\$809	\$10,760	\$1,314	\$4,929	\$2,954	\$936	\$122,588	135%
1996	\$7,156	\$35,753	\$2,234	\$540	\$1,209	\$27,032	\$17,461	\$687	\$11,514	\$1,268	\$5,127	\$2,337	\$1,673	\$113,990	126%
1997	\$11,131	\$38,094	\$2,262	\$701	\$1,380	\$22,863	\$13,014	\$691	\$8,933	\$1,473	\$5,275	\$2,450	\$1,970	\$110,237	122%
1998	\$6,423	\$15,257	\$770	\$538	\$1,937	\$21,947	\$7,547	\$964	\$8,119	\$1,186	\$4,107	\$1,796	\$1,286	\$71,877	79%
1999	\$9,088	\$22,672	\$833	\$351	\$1,891	\$15,891	\$6,756	\$1,525	\$8,805	\$897	\$4,171	\$2,085	\$1,197	\$76,162	84%
2000	\$9,907	\$26,350	\$449	\$376	\$2,203	\$14,394	\$6,797	\$1,055	\$8,704	\$765	\$5,490	\$1,826	\$1,336	\$79,651	88%
2001	\$6,632	\$22,159	\$491	\$455	\$1,810	\$10,263	\$5,064	\$805	\$6,809	\$979	\$5,155	\$2,051	\$1,162	\$63,837	70%
2002	\$5,465	\$14,577	\$650	\$1,053	\$2,050	\$6,395	\$7,320	\$594	\$6,246	\$1,110	\$4,508	\$2,054	\$915	\$52,938	58%
2003	\$6,487	\$22,351	\$499	\$1,731	\$1,234	\$3,887	\$5,441	\$646	\$7,135	\$853	\$5,487	\$1,858	\$1,024	\$58,632	65%
2004	\$8,800	\$20,047	\$516	\$1,692	\$1,149	\$4,689	\$3,370	\$641	\$6,412	\$860	\$5,196	\$1,570	\$633	\$55,574	61%
2005	\$13,814	\$23,353	\$531	\$1,005	\$944	\$4,302	\$3,081	\$597	\$6,405	\$767	\$6,344	\$1,360	\$733	\$63,236	70%
2006	\$17,776	\$25,503	\$643	\$450	\$761	\$3,910	\$3,613	\$511	\$5,433	\$723	\$6,465	\$1,214	\$733	\$67,735	75%
2007	\$15,216	\$22,554	\$718	\$117	\$753	\$4,223	\$3,850	\$543	\$8,309	\$532	\$5,391	\$903	\$876	\$63,984	71%
2008	\$16,114	\$28,825	\$781	\$58	\$804	\$4,479	\$5,108	\$623	\$9,787	\$282	\$5,270	\$745	\$1,243	\$74,120	82%
2009	\$6,652	\$36,186	\$634	\$260	\$717	\$4,890	\$4,445	\$874	\$9,109	\$259	\$3,741	\$872	\$786	\$69,427	77%
2010	\$10,328	\$37,274	\$494	\$358	\$681	\$4,712	\$4,716	\$714	\$7,237	\$160	\$2,083	\$718	\$780	\$70,255	77%
2011	\$24,137	\$45,539	\$878	\$715	\$861	\$5,321	\$4,218	\$501	\$7,089	\$124	\$2,940	\$771	\$1,052	\$94,146	104%
2012	\$20,499	\$28,033	\$1,114	\$788	\$846	\$5,721	\$3,662	\$607	\$6,496	\$145	\$3,550	\$795	\$1,192	\$73,448	
<b>Total</b>	<b>\$244,182</b>	<b>\$755,473</b>	<b>\$67,651</b>	<b>\$31,363</b>	<b>\$23,294</b>	<b>\$767,010</b>	<b>\$239,937</b>	<b>\$30,008</b>	<b>\$392,041</b>	<b>\$52,747</b>	<b>\$162,932</b>	<b>\$88,629</b>	<b>\$29,742</b>	<b>\$2,885,009</b>	
Pct of Total	8%	26%	2%	1%	1%	27%	8%	1%	14%	2%	6%	3%	1%	100%	
Ann. Average	\$7,631	\$23,609	\$2,114	\$980	\$728	\$23,969	\$7,498	\$938	\$12,251	\$1,648	\$5,092	\$2,770	\$929	\$90,157	

**Table 2-5. Groundfish landings (shoreside commercial and tribal) by gear type (mt), 1991-2012.**

Year	Dredge	Hook-and-line	Miscellaneous	Net	Pot	Shrimp Trawl	Trawl	Troll	Total
1981		4,308	2	1,770	3,961	1,846	90,949	508	103,344
1982		5,016	3	2,249	6,550	1,395	103,575	567	119,356
1983		3,990	6	3,203	5,989	1,221	84,141	430	98,978
1984		3,210	9	4,326	4,448	497	77,020	296	89,804
1985		5,361	1	5,499	3,938	522	75,261	340	90,923
1986		6,579	2	5,777	3,049	1,600	65,211	262	82,480
1987		7,576	11	4,650	2,139	1,622	75,789	196	91,982
1988		6,440	3	3,054	2,277	1,310	78,952	213	92,248
1989		6,680	11	3,248	2,125	1,232	85,800	276	99,372
1990		6,627	28	3,163	1,705	966	81,693	358	94,539
1991		8,293	1	1,898	1,086	873	93,282	115	105,547
1992		9,151	6	1,794	827	926	119,693	159	132,555
1993		7,559	2	1,287	871	1,597	104,844	235	116,394
1994		6,461	1	757	1,404	890	125,793	373	135,677
1995		6,396	2	792	1,118	754	125,214	217	134,493
1996		7,474	2	328	861	890	136,675	222	146,452
1997		7,100	<0.5	322	662	393	134,678	415	143,570
1998		4,661	2	387	546	386	124,751	279	131,011
1999		4,647	<0.5	140	821	427	119,755	93	125,883
2000		4,110	1	94	939	311	117,541	35	123,031
2001		3,697	1	87	734	241	98,760	37	103,556
2002	*	3,191	3	73	520	89	71,156	23	75,056
2003		3,489	1	80	842	32	78,438	24	82,905
2004	*	3,704	*	65	850	27	117,616	39	122,303
2005		3,932	2	56	1,023	16	130,383	42	135,454
2006		3,663	<0.5	63	1,079	19	146,397	39	151,260
2007		3,157	1	47	714	25	113,530	23	117,495
2008		3,588	<0.5	34	704	14	93,313	15	97,667
2009		4,391	*	13	890	16	76,872	13	82,195
2010		4,367	<0.5	8	934	14	88,704	18	94,045
2011		4,299	<0.5	9	1,484	70	122,708	16	128,585
2012	*	3,548	1	10	1,219	20	85,564	23	90,384
<b>Total</b>	<b>*</b>	<b>166,665</b>	<b>102</b>	<b>45,280</b>	<b>56,303</b>	<b>20,236</b>	<b>3,244,054</b>	<b>5,901</b>	<b>3,538,542</b>
Pct of Total	<1%	4%	<1%	<1%	1%	<1%	95%	<1%	100%
Ann. Average	*	5,268	3	1,460	1,777	652	101,879	191	111,231

**Table 2-6. Groundfish ex-vessel revenue (shoreside commercial and tribal) by gear type in current (2012) dollars, \$1,000s, 1981-2012.**

Year	Dredge	Hook-and-line	Miscellaneous	Net	Pot	Shrimp Trawl	Trawl	Troll	Total
1981		\$8,464	\$7	\$3,457	\$4,353	\$1,404	\$81,027	\$649	<b>\$99,360</b>
1982		\$9,895	\$8	\$3,835	\$10,844	\$1,215	\$97,594	\$786	<b>\$124,177</b>
1983		\$6,961	\$10	\$4,593	\$7,790	\$1,117	\$83,720	\$546	<b>\$104,737</b>
1984		\$6,007	\$14	\$6,283	\$4,792	\$484	\$75,370	\$446	<b>\$93,394</b>
1985		\$11,595	\$28	\$8,641	\$6,439	\$515	\$77,051	\$500	<b>\$104,769</b>
1986		\$14,217	\$27	\$8,939	\$5,382	\$1,716	\$72,668	\$443	<b>\$103,392</b>
1987		\$18,572	\$34	\$7,719	\$4,424	\$2,044	\$94,499	\$347	<b>\$127,638</b>
1988		\$16,924	\$38	\$4,970	\$4,791	\$1,153	\$89,465	\$348	<b>\$117,690</b>
1989		\$15,332	\$17	\$4,967	\$3,644	\$1,098	\$88,509	\$448	<b>\$114,014</b>
1990		\$15,213	\$37	\$4,942	\$2,932	\$867	\$77,958	\$624	<b>\$102,573</b>
1991		\$22,746	\$3	\$2,839	\$2,615	\$874	\$82,825	\$193	<b>\$112,096</b>
1992		\$21,603	\$14	\$2,647	\$1,957	\$941	\$79,865	\$252	<b>\$107,278</b>
1993		\$16,864	\$3	\$2,003	\$1,792	\$1,489	\$71,614	\$317	<b>\$94,081</b>
1994		\$16,686	\$3	\$1,185	\$4,232	\$1,114	\$74,195	\$582	<b>\$97,997</b>
1995		\$23,698	\$4	\$1,280	\$5,215	\$1,066	\$90,950	\$376	<b>\$122,588</b>
1996		\$27,070	\$4	\$566	\$4,354	\$1,171	\$80,462	\$362	<b>\$113,990</b>
1997		\$30,217	<\$1	\$517	\$3,970	\$564	\$74,336	\$630	<b>\$110,234</b>
1998		\$15,378	\$3	\$578	\$2,400	\$572	\$52,538	\$410	<b>\$71,878</b>
1999		\$18,574	\$1	\$235	\$3,613	\$614	\$52,963	\$162	<b>\$76,162</b>
2000		\$19,408	\$9	\$155	\$5,175	\$605	\$54,215	\$85	<b>\$79,651</b>
2001		\$17,065	\$6	\$160	\$3,630	\$374	\$42,525	\$78	<b>\$63,838</b>
2002	*	\$13,878	\$33	\$134	\$2,678	\$176	\$35,983	\$53	<b>\$52,936</b>
2003		\$15,977	\$10	\$137	\$4,316	\$82	\$38,056	\$54	<b>\$58,632</b>
2004	*	\$15,917	*	\$148	\$3,515	\$72	\$35,842	\$77	<b>\$55,574</b>
2005		\$17,558	\$7	\$96	\$4,356	\$69	\$41,062	\$88	<b>\$63,236</b>
2006		\$17,729	\$1	\$130	\$5,006	\$60	\$44,727	\$79	<b>\$67,733</b>
2007		\$16,180	\$10	\$108	\$3,383	\$51	\$44,180	\$72	<b>\$63,983</b>
2008		\$19,093	\$3	\$60	\$3,987	\$26	\$50,915	\$36	<b>\$74,119</b>
2009		\$24,129	*	\$16	\$4,966	\$24	\$40,270	\$19	<b>\$69,427</b>
2010		\$26,148	\$2	\$14	\$5,676	\$30	\$38,345	\$42	<b>\$70,256</b>
2011		\$31,589	\$2	\$15	\$10,734	\$101	\$51,646	\$58	<b>\$94,145</b>
2012		\$21,543	\$4	\$18	\$6,452	\$42	\$45,319	\$70	<b>\$73,448</b>
<b>Total</b>	<b>*</b>	<b>\$572,229</b>	<b>\$347</b>	<b>\$71,384</b>	<b>\$149,411</b>	<b>\$21,731</b>	<b>\$2,060,695</b>	<b>\$9,230</b>	<b>\$2,885,028</b>
Pct of Total	<0.5%	34%	<0.5%	<0.5%	11%	<0.5%	55%	<0.5%	100%
Ann. Average	*	\$17,452	\$11	\$2,262	\$4,532	\$687	\$63,861	\$290	\$89,096

**Table 2-7. Shoreside IFQ - Trawl (whiting and non-whiting) by groundfish species or species group (mt), 2003-2012.**

Fishery	P. Whiting	Sablefish	Lingcod	P. Cod	Other Roundfish	Rockfish	Thorny heads	Arrowtooth Flounder	Dover Sole	English Sole	Petrale Sole	Other Flatfish	Other Groundfish	Nonground fish	Total
<b>Whiting Total</b>	<b>717,634</b>	<b>361</b>	<b>35</b>	<b>10</b>	<b>178</b>	<b>2,706</b>	<b>24</b>	<b>59</b>	<b>2</b>	<b>2</b>	<b>1</b>	<b>15</b>	<b>804</b>	<b>2,241</b>	<b>724,072</b>
2003	51,183	40	<0.5	<0.5	*	68	<0.5	<0.5			<0.5	<0.5	4	88	51,385
2004	89,641	131	4	1	10	176	1	1			<0.5	<0.5	33	205	90,203
2005	97,559	22	6	1	165	289	<0.5	1			<0.5	<0.5	97	280	98,420
2006	97,267	11	6	1	<0.5	226	<0.5	2			<0.5	<0.5	38	71	97,622
2007	73,277	9	5	<0.5	*	319	<0.5	3			<0.5	1	52	197	73,864
2008	50,760	<0.5	3	<0.5		151	<0.5	2			<0.5	<0.5	60	880	51,857
2009	40,294	49	1	<0.5		211	<0.5	4			<0.5	<0.5	21	24	40,605
2010	62,655	21	2	<0.5	1	333	12	10			1	8	155	177	63,377
2011	89,826	30	5	7		530	2	13			<0.5	1	182	78	90,674
2012	65,171	47	4	<0.5		401	8	25			<0.5	4	162	242	66,065
<b>Nonwhiting Total</b>	<b>104</b>	<b>22,969</b>	<b>1,273</b>	<b>3,487</b>	<b>31</b>	<b>7,744</b>	<b>17,897</b>	<b>21,991</b>	<b>82,302</b>	<b>4,804</b>	<b>17,351</b>	<b>9,255</b>	<b>12,013</b>	<b>398</b>	<b>201,618</b>
2003	30	2,097	48	720	21	708	2,171	936	6,872	666	1,659	1,390	1,320	93	18,730
2004	12	2,183	53	825	4	979	1,279	1,246	6,555	817	1,749	1,280	784	38	17,803
2005	*	2,315	74	724	5	500	1,134	2,076	6,748	859	2,701	1,114	1,090	31	19,372
2006	*	2,467	115	330	<0.5	515	1,261	1,716	5,740	868	2,581	1,101	1,199	50	17,946
2007	2	2,428	119	43	*	576	1,605	2,025	8,951	622	2,207	884	1,090	40	20,592
2008	1	2,871	107	12	<0.5	603	2,438	2,635	10,970	327	2,175	743	1,391	43	24,313
2009	<0.5	3,009	109	87	*	757	2,459	3,823	11,611	265	1,696	887	1,441	49	26,192
2010	9	2,511	73	100	*	825	2,428	3,211	10,326	158	770	684	1,308	23	22,425
2011	26	1,663	239	252	*	928	1,585	2,167	7,586	108	792	585	1,177	12	17,120
2012	19	1,427	337	395	<0.5	1,354	1,538	2,158	6,944	115	1,021	587	1,212	19	17,126
<b>Grand Total</b>	<b>717,738</b>	<b>23,331</b>	<b>1,308</b>	<b>3,497</b>	<b>208</b>	<b>10,449</b>	<b>17,921</b>	<b>22,051</b>	<b>82,305</b>	<b>4,805</b>	<b>17,352</b>	<b>9,271</b>	<b>12,818</b>	<b>2,639</b>	<b>925,690</b>

**Table 2-8. Shoreside IFQ - Trawl (whiting and non-whiting) ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012.**

Fishery	P. Whiting	Sablefish	Lingcod	P. Cod	Other Roundfish	Rockfish	Thorny heads	Arrowtooth Flounder	Dover Sole	English Sole	Petrale Sole	Other Flatfish	Other Groundfish	Nonground fish	Total
<b>Whiting Total</b>	<b>123,210</b>	<b>663</b>	<b>32</b>	<b>4</b>	<b>18</b>	<b>2,358</b>	<b>11</b>	<b>9</b>	<b>&lt;\$1</b>	<b>&lt;\$1</b>	<b>2</b>	<b>3</b>	<b>143</b>	<b>185</b>	<b>126,639</b>
2003	5,966	49	<\$1	<\$1	*	66	<\$1	<\$1			<\$1	<\$1	<\$1	17	6,100
2004	8,271	67	4	1	2	174	<\$1	<\$1			<\$1	<\$1	10	27	8,558
2005	12,413	43	6	1	15	274	<\$1	<\$1			<\$1	<\$1	38	23	12,813
2006	14,018	21	5	<\$1	<\$1	207	<\$1	<\$1			<\$1	<\$1	12	11	14,276
2007	12,305	14	4	<\$1	*	246	<\$1	<\$1			<\$1	<\$1	16	40	12,627
2008	12,305	<\$1	2	<\$1		109	<\$1	<\$1			<\$1	<\$1	4	24	12,444
2009	5,590	8	<\$1	<\$1		151	<\$1	<\$1			<\$1	<\$1	<\$1	<\$1	5,751
2010	10,116	79	1	<\$1	<\$1	226	2	<\$1			1	<\$1	3	6	10,436
2011	22,027	189	5	<\$1		531	2	2			<\$1	<\$1	46	12	22,814
2012	20,199	194	3	<\$1		375	6	4			<\$1	1	13	24	20,820
<b>Nonwhiting Total</b>	<b>29</b>	<b>87,160</b>	<b>2,166</b>	<b>4,394</b>	<b>25</b>	<b>9,535</b>	<b>2,485</b>	<b>5,427</b>	<b>71,485</b>	<b>3,923</b>	<b>4,347</b>	<b>9,121</b>	<b>6,440</b>	<b>1,229</b>	<b>269,266</b>
2003	5	6,885	97	1,021	21	805	3,922	289	6,669	617	4,374	1,586	610	341	27,243
2004	2	5,984	93	1,000	2	1,186	2,007	371	6,241	735	4,617	1,340	312	140	24,029
2005	*	6,516	119	860	2	567	1,767	553	6,267	704	6,260	1,146	430	113	25,302
2006	*	7,997	184	402	<\$1	582	2,171	463	5,224	684	6,390	1,026	447	116	25,687
2007	<\$1	8,556	216	57	*	747	2,311	487	8,016	474	5,273	783	584	136	27,639
2008	1	12,073	177	15	<\$1	853	3,319	615	9,565	251	5,154	655	865	109	33,652
2009	<\$1	13,096	184	94	*	976	2,726	870	8,989	186	3,557	770	647	97	32,193
2010	3	11,152	138	102	*	1,025	2,638	713	7,182	111	2,000	607	690	86	26,445
2011	9	9,340	402	322	*	1,152	1,927	476	6,952	76	2,538	604	827	40	24,666
2012	8	5,562	556	520	<\$1	1,642	2,072	86	6,380	86	3,312	605	1,028	51	22,410
<b>Grand Total</b>	<b>123,239</b>	<b>87,823</b>	<b>2,199</b>	<b>4,398</b>	<b>43</b>	<b>11,894</b>	<b>2,486</b>	<b>5,436</b>	<b>71,485</b>	<b>3,923</b>	<b>4,347</b>	<b>9,124</b>	<b>6,583</b>	<b>1,414</b>	<b>395,904</b>



**Table 2-9. Shoreside IFQ – Non-trawl by groundfish species or species group (mt), 2011-2012.**

<b>Species</b>	<b>2011</b>	<b>2012</b>
Sablefish	1,116	923
Rougheye Rockfish	7	15
Blackgill Rockfish	2	6
Other Slope Rockfish	3	3
Shelf Rockfish	<0.5	<0.5
Thornyheads	23	13
Other Roundfish	3	2
Other Rockfish	1	<0.5
Flatfish	44	3
Other Groundfish	4	6
Nongroundfish	<0.5	<0.5
<b>Total</b>	<b>1203</b>	<b>971</b>

**Table 2-10. Shoreside IFQ - Non-trawl ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2011-2012.**

<b>Species</b>	<b>2011</b>	<b>2012</b>
Sablefish	\$7,611	\$4,896
Rougheye Rockfish	\$8	\$17
Blackgill Rockfish	\$8	\$19
Other Slope Rockfish	\$4	\$4
Shelf Rockfish	<\$1	<\$1
Thornyheads	\$146	\$29
Other Roundfish	\$6	\$4
Other Rockfish	\$1	<\$1
Flatfish	\$37	\$9
Other Groundfish	\$2	\$5
Nongroundfish	<\$1	<\$1
<b>Total</b>	<b>\$7,822</b>	<b>\$4,983</b>

Spiny dogfish landings excluded due to data confidentiality rules.

**Table 2-11. Limited entry fixed gear landings by groundfish species or species group (mt), 2003-2012.**

Year	Sablefish	Rougeye Rockfish	Spiny Dogfish	Blackgill Rockfish	Other Slope Rockfish	Shelf Rockfish	Thorny heads	Other Roundfish	Other Rockfish	Flatfish	Other Groundfish	Nonground fish	Total
2003	1,890	10	105	72	23	6	178	18	22	6	61	49	2,441
2004	2,136	19	90	42	25	12	166	22	19	5	37	54	2,627
2005	2,188	25	230	24	30	15	161	20	21	6	42	47	2,808
2006	2,209	40	131	38	23	14	172	21	23	4	36	50	2,760
2007	1,782	37	196	15	21	14	174	22	28	4	45	40	2,377
2008	1,844	39	181	20	21	13	199	32	38	7	66	31	2,491
2009	2,434	67	24	48	22	5	200	27	30	8	50	12	2,928
2010	2,474	48	8	40	29	3	225	24	32	8	57	4	2,953
2011	2,416	38	8	80	22	2	247	22	22	9	59	4	2,929
2012	1,845	34	1	47	36	6	177	20	18	10	84	8	2,286
<b>Total</b>	<b>21,218</b>	<b>356</b>	<b>974</b>	<b>425</b>	<b>250</b>	<b>91</b>	<b>1,901</b>	<b>226</b>	<b>252</b>	<b>67</b>	<b>539</b>	<b>299</b>	<b>26,599</b>

**Table 2-12. Limited entry fixed gear ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012.**

Year	Sablefish	Rougeye Rockfish	Spiny Dogfish	Blackgill Rockfish	Other Slope Rockfish	Shelf Rockfish	Thorny heads	Other Roundfish	Other Rockfish	Flatfish	Other Groundfish	Nonground fish	Total
2003	\$9,620	\$12	\$56	\$244	\$41	\$15	\$1,404	\$98	\$119	\$6	\$33	\$300	\$11,948
2004	\$8,976	\$20	\$47	\$152	\$38	\$53	\$1,307	\$88	\$87	\$7	\$17	\$320	\$11,112
2005	\$10,226	\$28	\$133	\$71	\$35	\$60	\$1,274	\$72	\$87	\$5	\$22	\$262	\$12,274
2006	\$11,143	\$43	\$75	\$109	\$33	\$55	\$1,382	\$74	\$107	\$3	\$18	\$373	\$13,414
2007	\$9,244	\$42	\$115	\$60	\$27	\$51	\$1,452	\$99	\$160	\$2	\$23	\$325	\$11,600
2008	\$10,716	\$46	\$94	\$79	\$32	\$57	\$1,708	\$150	\$208	\$9	\$36	\$239	\$13,377
2009	\$14,285	\$80	\$10	\$135	\$37	\$18	\$1,610	\$125	\$155	\$8	\$27	\$67	\$16,556
2010	\$16,056	\$68	\$4	\$109	\$44	\$11	\$1,846	\$111	\$131	\$17	\$37	\$33	\$18,466
2011	\$19,603	\$61	\$3	\$232	\$38	\$8	\$1,994	\$102	\$83	\$19	\$37	\$32	\$22,213
2012	\$11,620	\$60	<0.5	\$165	\$74	\$26	\$1,418	\$99	\$99	\$34	\$67	\$56	\$13,718
<b>Total</b>	<b>\$121,488</b>	<b>\$461</b>	<b>\$538</b>	<b>\$1,356</b>	<b>\$399</b>	<b>\$354</b>	<b>\$15,396</b>	<b>\$1,018</b>	<b>\$1,237</b>	<b>\$110</b>	<b>\$316</b>	<b>\$2,007</b>	<b>\$144,678</b>

**Table 2-13. Open access landings (other than by fixed gear) by groundfish species or species group (mt), 2003-2012.**

Year	Sablefish	Black Rockfish	Brown Rockfish	Lingcod	Cabezon	Nearshore Rockfish	Thorny heads	Other Rockfish	Flatfish	Other Groundfish	Nonground fish	Total
2003	4	<0.5		2		<0.5	1	27	<0.5	27	8	69
2004	4	<0.5		2		<0.5	1	24	7	20	6	64
2005	4	1		2		<0.5	*	13	<0.5	22	8	50
2006	3	<0.5		3		<0.5	*	19	<0.5	20	5	52
2007	7	<0.5		5		1	*	12	<0.5	12	5	43
2008	3	*	*	1	*	*		2	<0.5	8	7	22
2009	3	*		<0.5		*	*	1	*	4	1	16
2010		*		1		*	*	1	*	3	1	7
2011	3	<0.5		1		*	*	<0.5	<0.5	4	2	11
2012	6	<0.5		1		<0.5	<0.5	3	<0.5	5	2	16
<b>Total</b>	<b>36</b>	<b>2</b>	<b>&lt;0.5</b>	<b>18</b>	<b>&lt;0.5</b>	<b>2</b>	<b>6</b>	<b>101</b>	<b>13</b>	<b>125</b>	<b>45</b>	<b>347</b>

Spiny dogfish, kelp greenling, and other roundfish excluded for data confidentiality.

**Table 2-14. Open access ex-vessel revenue (other than by fixed gear) by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012.**

Year	Sablefish	Black Rockfish	Brown Rockfish	Lingcod	Cabezon	Nearshore Rockfish	Thorny heads	Other Rockfish	Flatfish	Other Groundfish	Nonground fish	Total
2003	\$9	\$1		\$7		\$1	\$4	\$64	<\$1	\$37	\$42	\$165
2004	\$11	\$1		\$11		\$2	\$2	\$62	\$8	\$33	\$29	\$161
2005	\$13	\$1		\$13		\$1	*	\$36	<\$1	\$31	\$47	\$144
2006	\$9	\$1		\$16		\$2	*	\$47	<\$1	\$34	\$30	\$139
2007	\$24	\$1		\$28		\$4	*	\$45	<\$1	\$20	\$33	\$155
2008	\$17	*	*	\$6	*	*		\$13	\$1	\$11	\$48	\$100
2009	\$19	*		\$1		*	*	\$2	*	\$6	\$8	\$31
2010		*		\$4		*	*	\$3	*	\$3	\$4	\$28
2011	\$17	<\$1		\$6		*	*	\$2	\$3	\$5	\$12	\$47
2012	\$28	<\$1		\$3		<\$1	*	\$9	\$1	\$4	\$14	\$61
<b>Total</b>	<b>\$146</b>	<b>\$6</b>	<b>\$3</b>	<b>\$95</b>	<b>\$3</b>	<b>\$13</b>	<b>\$15</b>	<b>\$282</b>	<b>\$17</b>	<b>\$184</b>	<b>\$267</b>	<b>\$1,031</b>

**Table 2-15. Non-nearshore fixed gear landings by groundfish species or species group (mt), 2003-2012.**

Year	Sablefish	Rougheye Rockfish	Spiny Dogfish	Blackgill Rockfish	Other Slope Rockfish	Shelf Rockfish	Thorny heads	Other Roundfish	Other Rockfish	Flatfish	Other Groundfish	Nonground fish	Total
2003	2,445	11	149	80	26	4	159	9	2	7	104	55	3,051
2004	2,604	20	99	47	27	3	161	14	4	5	81	58	3,122
2005	3,085	30	135	26	36	5	158	15	6	7	80	52	3,637
2006	3,007	41	101	48	26	6	170	16	9	5	52	57	3,537
2007	2,243	38	107	16	23	4	152	12	8	5	55	50	2,712
2008	2,419	41	125	24	22	3	155	20	8	8	96	45	2,965
2009	3,421	69	29	58	26	2	150	17	9	10	60	17	3,867
2010	3,521	49	10	67	32	1	172	15	15	8	73	6	3,969
2011	3,024	39	10	114	24	1	192	12	17	8	71	6	3,518
2012	2,247	36	2	80	37	4	146	11	8	8	91	11	2,680
<b>Total</b>	<b>28,016</b>	<b>374</b>	<b>766</b>	<b>560</b>	<b>278</b>	<b>34</b>	<b>1,615</b>	<b>140</b>	<b>84</b>	<b>70</b>	<b>764</b>	<b>359</b>	<b>33,058</b>

**Table 2-16. Non-nearshore fixed gear ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012.**

Year	Sablefish	Rougheye Rockfish	Spiny Dogfish	Blackgill Rockfish	Other Slope Rockfish	Shelf Rockfish	Thorny heads	Other Roundfish	Other Rockfish	Flatfish	Other Groundfish	Nonground fish	Total
2003	\$11,895	\$13	\$77	\$256	\$48	\$7	\$1,221	\$17	\$2	\$6	\$58	\$352	\$13,953
2004	\$10,651	\$20	\$51	\$162	\$43	\$7	\$1,252	\$20	\$6	\$6	\$37	\$355	\$12,611
2005	\$13,572	\$36	\$77	\$77	\$46	\$9	\$1,246	\$27	\$8	\$6	\$44	\$302	\$15,450
2006	\$14,335	\$45	\$59	\$137	\$40	\$15	\$1,361	\$29	\$13	\$4	\$26	\$428	\$16,494
2007	\$11,134	\$43	\$62	\$46	\$29	\$4	\$1,268	\$24	\$11	\$3	\$29	\$446	\$13,100
2008	\$13,528	\$49	\$64	\$75	\$30	\$6	\$1,318	\$49	\$13	\$5	\$52	\$414	\$15,603
2009	\$18,946	\$82	\$12	\$154	\$42	\$3	\$1,187	\$41	\$15	\$6	\$32	\$152	\$20,671
2010	\$21,343	\$71	\$5	\$179	\$48	\$2	\$1,362	\$36	\$28	\$5	\$48	\$56	\$23,183
2011	\$23,699	\$61	\$4	\$323	\$42	\$5	\$1,540	\$29	\$37	\$7	\$45	\$57	\$25,847
2012	\$13,956	\$63	\$1	\$251	\$73	\$13	\$1,168	\$29	\$18	\$13	\$71	\$79	\$15,735
<b>Total</b>	<b>\$153,058</b>	<b>\$483</b>	<b>\$413</b>	<b>\$1,660</b>	<b>\$441</b>	<b>\$72</b>	<b>\$12,923</b>	<b>\$301</b>	<b>\$151</b>	<b>\$63</b>	<b>\$441</b>	<b>\$2,642</b>	<b>\$172,647</b>

**Table 2-17. Nearshore fixed gear landings by groundfish species or species group (mt), 2003-2012.**

Year	Black Rockfish	Cabazon	Lingcod	Other Nearshore Rockfish	Brown Rockfish	Kelp Greenling	Gopher Rockfish	Other Rockfish	Blue Rockfish	Nonground fish	Flatfish	Other Groundfish	Thorny heads	Total
2003	172	64	55	45	20	25	13	13	13	4	<0.5	<0.5		424
2004	182	75	63	50	24	25	16	24	18	5	<0.5	<0.5		482
2005	170	58	52	50	22	23	18	24	23	3	<0.5	1		444
2006	153	49	52	53	21	16	15	26	24	4	<0.5	2		415
2007	184	46	54	59	22	20	19	26	17	3	<0.5	1	*	451
2008	181	47	56	57	24	23	24	23	30	4	2	1		470
2009	225	47	45	50	24	22	23	21	11	4	1	1		473
2010	152	45	38	39	26	20	27	14	9	4	1	1		375
2011	123	60	51	44	28	23	30	17	14	4	1	<0.5		395
2012	119	57	57	45	26	24	22	18	11	4	3	1		386
<b>Total</b>	<b>1,661</b>	<b>548</b>	<b>521</b>	<b>491</b>	<b>236</b>	<b>220</b>	<b>206</b>	<b>205</b>	<b>169</b>	<b>39</b>	<b>9</b>	<b>8</b>	<b>&lt;0.5</b>	<b>4,314</b>

**Table 2-18. Nearshore fixed gear ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012.**

Year	Black Rockfish	Blue Rockfish	Brown Rockfish	Cabazon	Flatfish	Gopher Rockfish	Kelp Greenling	Lingcod	Nonground fish	Other Groundfish	Other Nearshore Rockfish	Other Rockfish	Thorny heads	Total
2003	\$583	\$43	\$249	\$723	\$1	\$188	\$279	\$248	\$31	<\$1	\$723	\$71		\$3,139
2004	\$632	\$64	\$345	\$821	\$2	\$243	\$283	\$271	\$38	\$1	\$809	\$135		\$3,645
2005	\$621	\$77	\$327	\$639	\$2	\$299	\$267	\$226	\$20	\$2	\$820	\$132		\$3,433
2006	\$652	\$87	\$303	\$550	\$2	\$268	\$189	\$237	\$21	\$3	\$918	\$143		\$3,373
2007	\$810	\$71	\$319	\$515	\$2	\$328	\$217	\$256	\$14	\$2	\$1,004	\$152	*	\$3,691
2008	\$796	\$123	\$355	\$515	\$7	\$412	\$273	\$281	\$24	\$1	\$953	\$133		\$3,876
2009	\$959	\$44	\$346	\$467	\$4	\$365	\$236	\$230	\$26	\$1	\$770	\$123		\$3,570
2010	\$668	\$32	\$346	\$451	\$8	\$404	\$210	\$192	\$33	\$1	\$625	\$88		\$3,062
2011	\$569	\$49	\$382	\$604	\$10	\$466	\$233	\$263	\$32	\$1	\$687	\$107		\$3,403
2012	\$589	\$37	\$359	\$557	\$21	\$348	\$260	\$311	\$22	\$1	\$699	\$127		\$3,331
<b>Total</b>	<b>\$6,879</b>	<b>\$628</b>	<b>\$3,330</b>	<b>\$5,843</b>	<b>\$61</b>	<b>\$3,321</b>	<b>\$2,447</b>	<b>\$2,514</b>	<b>\$261</b>	<b>\$14</b>	<b>\$8,009</b>	<b>\$1,211</b>	<b>\$4</b>	<b>\$34,522</b>

Other roundfish excluded for data confidentiality.

**Table 2-19. Incidental open access landings by groundfish species or species group (mt), 2003-2012.**

Year	Sablefish	Roundfish	Rockfish	Flatfish	Other Groundfish	CPS	Crab	HMS	Other	Salmon	Shrimp	Total
2003	78	230	262	51	72	18	61	49	248	228	8	1,305
2004	72	219	303	49	64	21	45	42	254	258	6	1,333
2005	52	179	296	40	59	16	35	36	226	253	3	1,195
2006	64	141	199	41	63	27	42	37	202	144	4	964
2007	50	132	193	40	69	23	42	43	206	140	6	944
2008	46	96	101	33	46	25	33	32	149	15	8	584
2009	41	98	91	33	40	18	36	18	149	23	6	553
2010	21	120	132	22	31	10	30	24	152	81	5	628
2011	35	160	126	55	47	13	36	20	180	89	5	766
2012	38	194	177	62	36	15	37	25	197	110	4	895
<b>Total</b>	<b>497</b>	<b>1,569</b>	<b>1,880</b>	<b>426</b>	<b>527</b>	<b>186</b>	<b>397</b>	<b>326</b>	<b>1,963</b>	<b>1,341</b>	<b>55</b>	<b>9,167</b>

**Table 2-20. Incidental open access ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012.**

Year	Sablefish	Roundfish	Rockfish	Flatfish	Other Groundfish	CPS	Crab	HMS	Other	Salmon	Shrimp	Total
2003	\$99	\$44	\$33	\$2	\$21	\$4	\$181	\$72	\$1,169	\$934	\$260	\$2,820
2004	\$72	\$44	\$38	\$4	\$32	\$1	\$85	\$51	\$1,267	\$1,267	\$104	\$2,965
2005	\$83	\$24	\$42	\$1	\$18	\$1	\$36	\$27	\$1,246	\$1,125	\$74	\$2,677
2006	\$85	\$27	\$23	\$2	\$23	\$39	\$58	\$55	\$1,345	\$510	\$158	\$2,325
2007	\$101	\$27	\$17	\$2	\$25	\$685	\$67	\$59	\$1,452	\$389	\$133	\$2,958
2008	\$64	\$23	\$15	<\$1	\$10	\$1,622	\$52	\$48	\$868	\$40	\$64	\$2,805
2009	\$57	\$18	\$13	\$1	\$5	\$695	\$50	\$14	\$651	\$98	\$43	\$1,644
2010	\$18	\$28	\$27	\$1	\$7	\$624	\$40	\$23	\$843	\$363	\$76	\$2,049
2011	\$48	\$36	\$35	\$1	\$5	\$389	\$92	\$84	\$1,212	\$415	\$53	\$2,370
2012	\$32	\$44	\$38	\$3	\$5	\$528	\$96	\$24	\$1,125	\$619	\$46	\$2,560
<b>Total</b>	<b>\$658</b>	<b>\$315</b>	<b>\$281</b>	<b>\$18</b>	<b>\$152</b>	<b>\$4,588</b>	<b>\$756</b>	<b>\$456</b>	<b>\$11,179</b>	<b>\$5,760</b>	<b>\$1,009</b>	<b>\$25,174</b>

**Table 2-21. Non-nearshore non-sablefish landings by groundfish species or species group (mt), 2003-2012.**

Year	Blackgill Rockfish	Rougeye Rockfish	Thorny heads	Spiny Dogfish	Other Slope Rockfish	Shelf Rockfish	Other Rockfish	Other Roundfish	Flatfish	Other Groundfish	Nonground fish	Total
2003	42	<0.5	21		2	3	2	11	2	13	1	97
2004	19	*	7	78	3	14	2	13	4	14	*	155
2005	9	*	4	98	4	12	1	17	1	8	*	155
2006	6	<0.5	4	90	1	17	5	17	2	4	2	148
2007	7		23	91	1	19	4	20	3	6	2	177
2008	12	<0.5	45	67	2	16	3	21	2	3	1	172
2009	21	1	57	<0.5	6	11	2	14	2	2	2	116
2010	21	1	67		1	9	1	10	3	1	1	114
2011	16	1	65	*	1	13	2	11	4	2	1	116
2012	26	1	40	*	3	15	2	17	5	7	2	115
<b>Total</b>	<b>177</b>	<b>4</b>	<b>332</b>	<b>425</b>	<b>23</b>	<b>128</b>	<b>23</b>	<b>151</b>	<b>28</b>	<b>59</b>	<b>14</b>	<b>1,364</b>

**Table 2-22. Non-nearshore non-sablefish ex-vessel revenue by groundfish species or species group in current (2012) dollars, \$1,000s, 2003-2012.**

Year	Blackgill Rockfish	Rougeye Rockfish	Thorny heads	Spiny Dogfish	Other Slope Rockfish	Shelf Rockfish	Other Rockfish	Other Roundfish	Flatfish	Other Groundfish	Nonground fish	Total
2003	\$141	<\$1	\$195		\$7	\$10	\$6	\$44	\$15	\$15	\$6	\$439
2004	\$66	*	\$64	\$42	\$8	\$66	\$6	\$52	\$19	\$17	\$11	\$351
2005	\$35	*	\$37	\$57	\$13	\$55	\$6	\$67	\$5	\$12	\$5	\$290
2006	\$20	<\$1	\$37	\$51	\$4	\$76	\$20	\$72	\$15	\$9	\$11	\$315
2007	\$39		\$196	\$55	\$4	\$97	\$17	\$86	\$22	\$10	\$11	\$537
2008	\$55	<\$1	\$400	\$36	\$7	\$82	\$13	\$113	\$14	\$3	\$10	\$732
2009	\$62	\$2	\$471	<\$1	\$18	\$63	\$7	\$78	\$9	\$3	\$9	\$724
2010	\$60	\$1	\$580		\$2	\$50	\$3	\$54	\$20	\$2	\$6	\$779
2011	\$48	\$3	\$549	*	\$2	\$66	\$8	\$61	\$24	\$2	\$9	\$773
2012	\$83	\$2	\$334	*	\$9	\$80	\$9	\$98	\$32	\$7	\$15	\$669
<b>Total</b>	<b>\$609</b>	<b>\$8</b>	<b>\$2,862</b>	<b>\$241</b>	<b>\$76</b>	<b>\$645</b>	<b>\$96</b>	<b>\$725</b>	<b>\$174</b>	<b>\$79</b>	<b>\$93</b>	<b>\$5,607</b>

Table 2-23. Groundfish landings by shoreside commercial fishery sectors (mt), 2003-2012.

Year	Shoreside IFQ Trawl (Nonwhiting)	Shoreside IFQ Trawl (Whiting)	Shoreside IFQ Nonrawl	Non Nearshore Fixed Gear	Nearshore Fixed Gear	Non Fixed Gear Open Access	Incidental Open Access	Exempted trawl, EFP/Research, Misc.	Grand Total
2003	18,638	51,297		3,092	420	69	54	3,747	77,317
2004	17,765	89,999		3,217	476	62	53	1,982	113,554
2005	19,342	98,141		3,739	441	48	52	293	122,054
2006	17,896	97,552		3,625	411	52	44	121	119,701
2007	20,552	73,667		2,837	448	38	49	116	97,707
2008	24,270	50,977		3,090	466	28	26	156	79,013
2009	26,143	40,580		3,964	469	14	25	136	71,331
2010	22,401	63,200		4,076	371	7	17	304	90,375
2011	17,108	90,596	1,203	3,626	391	9	20	1,850	114,804
2012	17,107	65,823	971	2,783	382	14	23	786	87,890
<b>Total</b>	<b>201,220</b>	<b>721,831</b>	<b>2,175</b>	<b>34,049</b>	<b>4,275</b>	<b>341</b>	<b>363</b>	<b>9,490</b>	<b>973,745</b>
Pct of total	21%	74%	0%	3%	0%	0%	0%	1%	100%
Ann Average	20,122	72,183	1,087	3,405	428	34	36	949	97,375



Table 2-24. Groundfish ex-vessel revenue in current (2012) dollars, \$1,000s, by shoreside commercial fishery sectors, 2003-2012.

Year	Shoreside IFQ Trawl (Nonwhiting)	Shoreside IFQ Trawl (Whiting)	Shoreside IFQ Nonrawl	Non Nearshore Fixed Gear	Nearshore Fixed Gear	Non Fixed Gear Open Access	Incidental Open Access	Exempted trawl, EFP/Research, Misc.	Grand Total
2003	\$26,902	\$6,083		\$14,033	\$3,108	\$137	\$200	\$3,731	\$54,194
2004	\$23,889	\$8,531		\$12,597	\$3,607	\$142	\$190	\$1,766	\$50,721
2005	\$25,189	\$12,790		\$15,433	\$3,413	\$103	\$169	\$615	\$57,712
2006	\$25,571	\$14,264		\$16,369	\$3,353	\$127	\$160	\$333	\$60,177
2007	\$27,503	\$12,587		\$13,179	\$3,678	\$125	\$171	\$243	\$57,485
2008	\$33,544	\$12,420		\$15,912	\$3,850	\$64	\$111	\$324	\$66,225
2009	\$32,096	\$5,751		\$21,235	\$3,543	\$23	\$93	\$421	\$63,162
2010	\$26,359	\$10,430		\$23,901	\$3,027	\$25	\$80	\$1,132	\$64,955
2011	\$24,626	\$22,802	\$7,821	\$26,553	\$3,372	\$35	\$126	\$904	\$86,239
2012	\$22,359	\$20,796	\$4,982	\$16,308	\$3,309	\$47	\$123	\$616	\$68,538
<b>Total</b>	<b>\$268,037</b>	<b>\$126,454</b>	<b>\$12,803</b>	<b>\$175,519</b>	<b>\$34,259</b>	<b>\$827</b>	<b>\$1,423</b>	<b>\$10,084</b>	<b>\$629,407</b>
Pct of total	43%	20%	2%	28%	5%	0%	0%	2%	100%
Ann Average	\$26,803.8	\$12,645.4	\$6,401.5	\$17,552.0	\$3,426.0	\$82.8	\$142.3	\$1,008.5	\$62,941

**Table 2-25. Treaty non-whiting groundfish sector landings (groundfish only) by gear group (mt), 2003-2012.**

Year	Hook-and-Line	Pot	Shrimp Trawl	Other Trawl	Total
2003	629		10	758	1,397
2004	754		16	1,071	1,841
2005	679		25	1,242	1,946
2006	654	<0.5	30	865	1,548
2007	535		11	935	1,481
2008	669		13	726	1,408
2009	759		2	1,046	1,807
2010	598	34	12	1,050	1,693
2011	557	20	7	1,431	2,014
2012	552	47	10	1,273	1,881
<b>Total</b>	<b>6,386</b>	<b>101</b>	<b>135</b>	<b>10,395</b>	<b>17,017</b>

**Table 2-26. Treaty non-whiting groundfish sector ex-vessel revenue (groundfish only), current (2012) dollars, \$1,000s, 2003-2012.**

Year	Hook-and-Line	Pot	Shrimp Trawl	Other Trawl	Total
2003	\$2,652		\$13	\$1,101	\$3,766
2004	\$2,938		\$21	\$1,341	\$4,300
2005	\$2,620		\$35	\$1,452	\$4,107
2006	\$2,754	<\$1	\$47	\$932	\$3,733
2007	\$2,483		\$18	\$998	\$3,499
2008	\$3,014		\$21	\$912	\$3,947
2009	\$3,902		\$3	\$1,250	\$5,155
2010	\$3,609	\$214	\$15	\$1,254	\$5,092
2011	\$4,235	\$140	\$9	\$1,814	\$6,198
2012	\$3,048	\$164	\$16	\$1,550	\$4,777
<b>Total</b>	<b>\$31,255</b>	<b>\$518</b>	<b>\$197</b>	<b>\$12,603</b>	<b>\$44,574</b>

**Table 2-27. Landings (mt) by whiting sectors, 2003-2012.**

Sector / Species Group	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	Grand Total
<b>Catcher-Processor Subtotal</b>	<b>41,434</b>	<b>71,004</b>	<b>79,333</b>	<b>79,096</b>	<b>74,304</b>	<b>109,134</b>	<b>38,748</b>	<b>54,787</b>	<b>72,759</b>	<b>55,669</b>	<b>676,267</b>
P. Whiting	41,215	69,412	78,890	78,865	73,264	108,240	34,801	54,292	71,680	55,264	665,922
Other Groundfish	101	401	204	97	211	718	49	219	835	290	3,125
Nongroundfish	118	1,192	239	134	829	175	3,898	277	244	114	7,220
<b>Mothership Subtotal</b>	<b>26,040</b>	<b>24,163</b>	<b>49,295</b>	<b>55,601</b>	<b>47,986</b>	<b>57,687</b>	<b>24,297</b>	<b>35,935</b>	<b>50,331</b>	<b>38,604</b>	<b>409,939</b>
P. Whiting	26,022	24,102	48,597	55,355	47,811	57,498	24,091	35,714	50,051	38,442	407,683
Other Groundfish	5	53	123	176	157	162	199	175	192	108	1,350
Nongroundfish	13	8	575	70	18	26	7	47	88	54	906
<b>Shoreside Whiting Trawl Subtotal</b>	<b>51,385</b>	<b>90,204</b>	<b>98,420</b>	<b>97,622</b>	<b>73,864</b>	<b>51,857</b>	<b>40,605</b>	<b>63,377</b>	<b>90,674</b>	<b>66,065</b>	<b>724,072</b>
P. Whiting	51,183	89,641	97,559	97,267	73,277	50,760	40,294	62,655	89,826	65,171	717,634
Other Groundfish	115	358	582	285	390	217	287	545	770	651	4,198
Nongroundfish	88	205	280	71	197	880	24	177	78	242	2,241
<b>Treaty Shoreside Whiting Trawl Subtotal</b>	<b>4,196</b>	<b>6,909</b>	<b>11,457</b>	<b>30,026</b>	<b>18,321</b>	<b>17,516</b>	<b>9,158</b>	<b>1,977</b>	<b>11,766</b>	<b>613</b>	<b>111,938</b>
P. Whiting	4,079	6,848	11,422	29,896	18,158	16,972	8,929	1,968	11,756	613	110,642
Other Groundfish	113	61	32	115	149	275	127	9	10		889
Nongroundfish	5	<0.5	2	15	14	269	102	<0.5	<0.5		407
<b>Treaty Mothership Subtotal</b>	<b>20,684</b>	<b>23,950</b>	<b>24,356</b>	<b>5,661</b>	<b>5,275</b>	<b>15,152</b>	<b>14,107</b>	<b>16,530</b>	<b>6,438</b>	<b>33</b>	<b>132,186</b>
P. Whiting	19,376	23,459	23,582	5,568	5,167	14,944	13,458	16,309	6,344	31	128,239
Other Groundfish	1,270	470	746	91	82	205	142	218	89	2	3,316
Nongroundfish	38	20	28	2	26	3	507	3	5	<0.5	631
<b>Grand Total</b>	<b>143,739</b>	<b>216,230</b>	<b>262,861</b>	<b>268,006</b>	<b>219,750</b>	<b>251,346</b>	<b>126,915</b>	<b>172,606</b>	<b>231,968</b>	<b>160,984</b>	<b>2,054,402</b>

Table 2-28. Ex-vessel revenue, current (2012) dollars, \$1,000s, by whiting sectors, 2003-2012.

Sector / Species Group	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	Grand Total
<b>Catcher-Processor Subtotal</b>	<b>\$6,542</b>	<b>\$11,760</b>	<b>\$9,987</b>	<b>\$10,670</b>	<b>\$11,910</b>	<b>\$26,610</b>	<b>\$4,178</b>	<b>\$9,995</b>	<b>\$16,263</b>	<b>\$16,376</b>	<b>\$124,291</b>
P. Whiting	\$6,372	\$11,567	\$9,823	\$10,582	\$11,775	\$26,299	\$4,163	\$9,937	\$16,022	\$16,259	\$122,799
Other Groundfish	\$131	\$85	\$96	\$42	\$121	\$264	\$13	\$46	\$160	\$86	\$1,044
Nongroundfish	\$38	\$109	\$68	\$46	\$14	\$47	\$3	\$12	\$81	\$31	\$448
<b>Mothership Subtotal</b>	<b>\$6,041</b>	<b>\$3,185</b>	<b>\$6,057</b>	<b>\$7,271</b>	<b>\$7,423</b>	<b>\$17,355</b>	<b>\$2,959</b>	<b>\$6,437</b>	<b>\$12,331</b>	<b>\$11,016</b>	<b>\$80,075</b>
P. Whiting	\$6,039	\$3,133	\$5,948	\$7,199	\$7,356	\$17,274	\$2,818	\$6,356	\$12,275	\$10,974	\$79,373
Other Groundfish	\$1	\$47	\$79	\$67	\$64	\$76	\$139	\$81	\$35	\$38	\$627
Nongroundfish	<1\$	\$4	\$31	\$5	\$3	\$4	\$2	<1\$	\$21	\$4	\$76
<b>Shoreside Whiting Trawl Subtotal</b>	<b>\$6,100</b>	<b>\$8,558</b>	<b>\$12,813</b>	<b>\$14,276</b>	<b>\$12,627</b>	<b>\$12,444</b>	<b>\$5,751</b>	<b>\$10,436</b>	<b>\$22,814</b>	<b>\$20,820</b>	<b>\$126,639</b>
P. Whiting	\$5,966	\$8,271	\$12,413	\$14,018	\$12,305	\$12,305	\$5,590	\$10,116	\$22,027	\$20,199	\$123,210
Other Groundfish	\$116	\$260	\$378	\$247	\$282	\$115	\$161	\$314	\$776	\$597	\$3,245
Nongroundfish	\$17	\$27	\$23	\$11	\$40	\$24	<1\$	\$6	\$12	\$24	\$185
<b>Treaty Shoreside Whiting Trawl Subtotal</b>	<b>\$680</b>	<b>\$554</b>	<b>\$1,418</b>	<b>\$3,823</b>	<b>\$3,000</b>	<b>\$4,010</b>	<b>\$1,132</b>	<b>\$209</b>	<b>\$1,708</b>	<b>\$133</b>	<b>\$16,668</b>
P. Whiting	\$507	\$524	\$1,389	\$3,756	\$2,908	\$3,799	\$1,061	\$205	\$1,701	\$133	\$15,985
Other Groundfish	\$165	\$29	\$29	\$67	\$91	\$148	\$48	\$4	\$7		\$588
Nongroundfish	\$7	<1\$	<1\$	<1\$	\$2	\$62	\$23	<1\$	<1\$		\$95
<b>Treaty Mothership Subtotal</b>	<b>\$2,860</b>	<b>\$2,123</b>	<b>\$3,119</b>	<b>\$834</b>	<b>\$921</b>	<b>\$3,523</b>	<b>\$1,308</b>	<b>\$1,841</b>	<b>\$1,512</b>	<b>\$10</b>	<b>\$18,051</b>
P. Whiting	\$2,237	\$2,036	\$3,026	\$779	\$911	\$3,477	\$1,296	\$1,738	\$1,479	\$10	\$16,989
Other Groundfish	\$624	\$87	\$93	\$55	\$10	\$46	\$12	\$98	\$30	<1\$	\$1,055
Nongroundfish	<1\$	<1\$	<1\$	<1\$	<1\$	<1\$	<1\$	\$4	\$3	<1\$	\$7
<b>Grand Total</b>	<b>\$22,223</b>	<b>\$26,180</b>	<b>\$33,394</b>	<b>\$36,874</b>	<b>\$35,881</b>	<b>\$63,942</b>	<b>\$15,328</b>	<b>\$28,918</b>	<b>\$54,628</b>	<b>\$48,355</b>	<b>\$365,724</b>

Table 2-29. Average monthly landings (mt) by commercial fishery sectors, 5 years, 2008-2012 (except as noted).

Sector /Species	January	February	March	April	May	June	July	August	September	October	November	December
<b>At-Sea Catch-Processor (Whiting)</b>					15,132	4,226	56	4,860	9,904	11,412	12,914	6,773
P. Whiting					15,074	4,190	56	4,845	9,872	11,386	12,833	6,599
Other Groundfish					58	36	<0.5	15	32	26	82	174
<b>At-Sea Mothership (Whiting)</b>					16,346	6,871	1,624	1,820	2,395	8,696	3,259	316
P. Whiting					16,253	6,846	1,621	1,816	2,385	8,679	3,250	310
Other Groundfish					93	25	3	5	11	17	9	6
<b>Shoreside IFQ Trawl (Whiting)</b>					1,829	10,966	11,276	17,855	7,008	8,611	4,062	628
P. Whiting					1,815	10,888	11,167	17,719	6,947	8,576	4,011	618
Other Groundfish					14	78	109	136	61	34	51	11
<b>Shoreside IFQ Trawl (Nonwhiting)</b>	1,303	1,897	2,003	2,287	2,131	1,796	1,805	1,816	1,674	1,745	1,508	1,440
P. Whiting	<0.5	<0.5	1	1	2	1	1	1	2	1	1	1
Other Groundfish	1,302	1,897	2,001	2,286	2,130	1,795	1,805	1,816	1,672	1,745	1,508	1,439
<b>Non Nearshore Fixed Gear</b>	97	104	131	339	414	423	406	443	517	352	164	118
P. Whiting	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5
Other Groundfish	97	104	131	339	414	423	406	443	517	352	164	118
<b>Nearshore Fixed Gear</b>	19	16	11	23	58	53	57	53	56	34	21	14
Other Groundfish	19	16	11	23	58	53	57	53	56	34	21	14
<b>Non Fixed Gear Open Access</b>	1	1	<0.5	1	1	1	1	5	2	1	<0.5	<0.5
Other Groundfish	1	1	<0.5	1	1	1	1	5	2	1	<0.5	<0.5
<b>Incidental Open Access</b>	1	1	<0.5	1	2	7	6	1	1	1	1	1
P. Whiting		*					*					
Other Groundfish	1	1	<0.5	1	2	7	6	1	1	1	1	1
<b>Exempted trawl, EFP/Research, Misc.</b>	7	6	5	6	19	179	117	64	192	40	6	5
P. Whiting		*	<0.5		*	136	87	25	*	28	*	
Other Groundfish	7	6	5	6	19	43	30	39	*	12	6	5
<b>Shoreside IFQ Nontrawl**</b>	*	*	33	64	6	57	69	95	265	281	128	77
Other Groundfish	*	*	33	64	6	57	69	95	265	281	128	77

Table 2-30. Average monthly ex-vessel revenue in nominal dollars, \$1,000s, by commercial fishery sectors, 5 years, 2008-2012 (except as noted).

Sector /Species	January	February	March	April	May	June	July	August	September	October	November	December
<b>At-Sea Catch-Processor (Whiting)</b>					\$3,434	\$1,362	\$9	\$1,040	\$2,045	\$2,576	\$2,756	939
P. Whiting					\$3,405	\$1,337	\$9	\$1,039	\$2,038	\$2,569	\$2,744	910
Other Groundfish					\$29	\$25	<1\$	\$1	\$7	\$7	\$11	29
<b>At-Sea Mothership (Whiting)</b>					\$2,936	\$1,954	\$534	\$525	\$617	\$2,242	\$836	43
P. Whiting					\$2,886	\$1,944	\$534	\$524	\$616	\$2,238	\$832	41
Other Groundfish					\$50	\$10	<1\$	\$1	\$1	\$4	\$4	\$1
<b>Shoreside IFQ Trawl (Whiting)</b>					\$338	\$1,965	\$2,651	\$4,222	\$1,619	\$2,131	\$1,056	\$93
P. Whiting					\$333	\$1,908	\$2,533	\$4,107	\$1,574	\$2,113	\$1,032	\$90
Other Groundfish					\$5	\$57	\$118	\$116	\$45	\$17	\$24	\$2
<b>Shoreside IFQ Trawl (Nonwhiting)</b>	\$1,797	\$2,285	\$2,238	\$2,420	\$2,472	\$2,266	\$2,409	\$2,314	\$2,226	\$2,251	\$2,074	\$2,036
P. Whiting	<1\$	<1\$	\$1	<1\$	<1\$	<1\$	<1\$	<1\$	\$1	\$1	<1\$	<1\$
Other Groundfish	\$1,797	\$2,285	\$2,238	\$2,420	\$2,471	\$2,266	\$2,409	\$2,314	\$2,224	\$2,251	\$2,074	\$2,036
<b>Non Nearshore Fixed Gear</b>	\$494	\$511	\$685	\$1,835	\$2,452	\$2,523	\$2,438	\$2,643	\$2,965	\$2,052	\$879	\$626
P. Whiting	<1\$	<1\$	<1\$	<1\$	<1\$	<1\$	<1\$	<1\$	<1\$	<1\$	<1\$	<1\$
Other Groundfish	\$494	\$511	\$685	\$1,835	\$2,452	\$2,523	\$2,438	\$2,643	\$2,965	\$2,052	\$879	\$626
<b>Nearshore Fixed Gear</b>	\$200	\$160	\$71	\$130	\$432	\$393	\$448	\$398	\$426	\$282	\$207	\$157
Other Groundfish	\$200	\$160	\$71	\$130	\$432	\$393	\$448	\$398	\$426	\$282	\$207	\$157
<b>Non Fixed Gear Open Access</b>	\$2	\$2	\$1	\$1	\$3	\$3	\$5	\$8	\$5	\$5	\$1	<1\$
Other Groundfish	\$2	\$2	\$1	\$1	\$3	\$3	\$5	\$8	\$5	\$5	\$1	<1\$
<b>Incidental Open Access</b>	\$7	\$4	\$1	\$2	\$9	\$28	\$26	\$8	\$4	\$5	\$4	\$5
P. Whiting		*					*					
Other Groundfish	\$7	\$4	\$1	\$2	\$9	\$28	\$26	\$8	\$4	\$5	\$4	\$5
<b>Exempted trawl, EFP/Research, Misc.</b>	\$16	\$13	\$8	\$18	\$45	\$116	\$131	\$110	\$110	\$47	\$26	\$20
P. Whiting		*	<1\$		*	\$30	\$26	\$8	*	\$7	*	
Other Groundfish	\$16	\$13	\$8	\$18	\$45	\$85	\$105	\$102	*	\$40	\$26	\$20
<b>Shoreside IFQ Nontrawl**</b>	*	*	\$131	\$382	\$40	\$174	\$294	\$565	\$1,649	\$1,841	\$764	\$475
Other Groundfish	*	*	\$131	\$382	\$40	\$174	\$294	\$565	\$1,649	\$1,841	\$764	\$475

**Table 2-31. Average monthly landings (mt) by Treaty fishery sectors, 5 years, 2008-2012.**

Sector/ Species	January	February	March	April	May	June	July	August	September	October	November	December
<b>Treaty Whiting Mothership</b>						6,426	10,541	5,769	3,312	5,013	1,869	630
P. Whiting						6,063	10,222	5,618	3,267	4,859	1,824	625
Other Groundfish						362	319	151	44	154	45	6
<b>Treaty Shoreside Whiting Trawl</b>					17	4	2,125	1,269	1,102	2,090	990	535
P. Whiting					14	4	2,112	1,266	1,098	2,070	965	520
Other Groundfish					3		14	3	5	20	25	14
<b>Treaty Shoreside Nonwhiting Groundfish</b>	40	47	107	247	236	204	165	146	202	198	93	77
Other Groundfish	40	47	107	247	236	204	165	146	202	198	93	77

**Table 2-32. Average monthly ex-vessel revenue in nominal dollars, \$1,000s, by Treaty fishery sectors, 5 years, 2008-2012.**

Sector/ Species	January	February	March	April	May	June	July	August	September	October	November	December
<b>Treaty Whiting Mothership</b>						\$676	\$1,086	\$659	\$524	\$778	\$252	\$22
P. Whiting						\$571	\$1,024	\$607	\$492	\$732	\$249	\$21
Other Groundfish						\$106	\$62	\$52	\$32	\$46	\$2	\$1
<b>Treaty Shoreside Whiting Trawl</b>					\$6	\$1	\$282	\$175	\$250	\$417	\$148	\$77
P. Whiting					\$3	\$1	\$277	\$173	\$248	\$408	\$138	\$70
Other Groundfish					\$3		\$5	\$2	\$3	\$9	\$10	\$7
<b>Treaty Shoreside Nonwhiting Groundfish</b>	\$46	\$53	\$315	\$931	\$649	\$601	\$438	\$290	\$503	\$559	\$280	\$209
Other Groundfish	\$46	\$53	\$315	\$931	\$649	\$601	\$438	\$290	\$503	\$559	\$280	\$209

**Table 2-33. Average monthly groundfish landings (mt) by species and species groups, 5 years, 2008-2012.**

Month	P. Whiting	Sablefish	P. Cod	Other Roundfish	Rockfish	Thorny heads	Arrowtooth Flounder	Dover Sole	English Sole	Lingcod	Other Flatfish	Other Groundfish	Petrale Sole
January	<0.5	186	6	6	73	132	118	562	11	6	33	82	255
February	1	225	12	4	78	192	254	864	18	13	41	130	244
March	2	312	14	3	97	175	257	1,062	18	20	42	127	143
April	1	645	37	5	123	235	398	1,196	19	11	56	155	50
May	1,831	661	61	11	228	249	420	877	23	36	79	162	73
June	11,029	683	59	8	260	220	277	672	35	40	108	181	85
July	13,366	687	58	11	289	194	260	670	38	48	102	177	86
August	19,010	669	46	8	295	192	235	769	35	41	110	197	83
September	8,209	821	22	10	311	215	152	776	20	38	72	150	62
October	10,675	687	38	6	238	227	189	806	16	25	64	163	52
November	4,977	418	5	4	143	184	141	743	13	14	43	136	77
December	1,139	318	13	3	141	140	155	638	14	18	32	85	154

**Table 2-34. Average monthly groundfish ex-vessel revenue in nominal dollars, \$1,000s, by species and species groups, 5 years, 2008-2012.**

Month	P. Whiting	Sablefish	P. Cod	Other Roundfish	Rockfish	Thorny heads	Arrowtooth Flounder	Dover Sole	English Sole	Lingcod	Other Flatfish	Other Groundfish	Petrale Sole
January	<\$1	\$838	\$7	\$63	\$230	\$295	\$26	\$444	\$8	\$9	\$31	\$48	\$564
February	<\$1	\$1,019	\$13	\$46	\$212	\$326	\$56	\$654	\$13	\$24	\$39	\$73	\$559
March	<\$1	\$1,481	\$16	\$29	\$160	\$336	\$58	\$815	\$13	\$34	\$37	\$68	\$323
April	<\$1	\$3,461	\$43	\$41	\$237	\$378	\$93	\$943	\$14	\$21	\$52	\$85	\$124
May	\$336	\$3,708	\$69	\$112	\$515	\$400	\$96	\$694	\$16	\$101	\$75	\$96	\$203
June	\$1,939	\$3,782	\$69	\$82	\$553	\$372	\$62	\$530	\$24	\$106	\$109	\$113	\$224
July	\$2,837	\$3,772	\$65	\$83	\$633	\$367	\$58	\$548	\$27	\$115	\$105	\$102	\$234
August	\$4,288	\$3,767	\$56	\$77	\$597	\$339	\$52	\$641	\$25	\$104	\$110	\$111	\$226
September	\$1,865	\$4,616	\$24	\$98	\$649	\$394	\$36	\$644	\$15	\$99	\$68	\$87	\$176
October	\$2,529	\$3,926	\$45	\$58	\$469	\$399	\$41	\$644	\$12	\$70	\$57	\$87	\$150
November	\$1,170	\$2,102	\$6	\$37	\$317	\$364	\$30	\$586	\$10	\$44	\$41	\$59	\$214
December	\$161	\$1,551	\$14	\$31	\$294	\$302	\$34	\$513	\$10	\$29	\$30	\$49	\$394



Table 2-35. Groundfish landings (mt) by "IOPAC port groups", 2003-2012.

Port	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>Washington Subtotal</b>	<b>23,997</b>	<b>39,341</b>	<b>50,471</b>	<b>65,430</b>	<b>50,139</b>	<b>39,386</b>	<b>24,516</b>	<b>35,645</b>	<b>39,393</b>	<b>22,311</b>
Puget Sound	3,580	3,465	2,983	1,959	1,462	1,138	1,565	1,403	706	804
North WA coast	2,284	2,128	2,438	1,891	1,592	1,522	1,829	1,642	1,833	1,819
South and central WA coast	18,133	33,748	45,049	61,580	47,086	36,726	21,123	32,600	36,855	19,688
<b>Oregon Subtotal</b>	<b>48,266</b>	<b>70,680</b>	<b>73,807</b>	<b>73,743</b>	<b>56,807</b>	<b>45,077</b>	<b>47,330</b>	<b>48,236</b>	<b>81,822</b>	<b>61,727</b>
Astoria	19,136	21,556	23,145	31,489	25,734	18,441	22,662	21,567	49,935	30,672
Tillamook	107	88	46	73	49	41	62	37	30	30
Newport	22,679	41,380	43,304	33,647	24,066	19,074	17,347	20,702	26,459	27,211
Coos Bay	5,110	6,857	6,203	7,458	5,570	5,850	5,515	4,148	3,962	2,377
Brookings	1,234	799	1,109	1,075	1,389	1,671	1,743	1,783	1,436	1,437
<b>California Subtotal</b>	<b>10,643</b>	<b>12,282</b>	<b>11,176</b>	<b>12,087</b>	<b>10,549</b>	<b>13,204</b>	<b>10,349</b>	<b>10,164</b>	<b>7,368</b>	<b>6,342</b>
Crescent City	1,024	2,177	1,568	2,140	1,642	4,284	2,666	2,746	386	178
Eureka	3,817	4,950	5,166	6,223	5,213	4,738	3,152	2,726	2,229	1,935
Fort Bragg	1,597	1,616	1,902	1,446	1,470	1,758	1,956	1,857	1,670	1,484
Bodega Bay	212	41	13	62	95	118	82	75	73	53
San Francisco	1,031	1,208	690	714	1,119	1,027	774	723	521	373
Monterey	1,585	1,009	989	916	457	541	476	566	585	618
Morro Bay	956	893	507	244	223	393	874	937	1,275	1,162
Santa Barbara	97	100	92	107	91	80	126	266	332	257
Los Angeles	200	221	160	111	165	145	144	151	181	146
San Diego	123	67	89	122	76	122	100	119	117	136
<b>Grand Total</b>	<b>82,905</b>	<b>122,303</b>	<b>135,454</b>	<b>151,260</b>	<b>117,495</b>	<b>97,667</b>	<b>82,195</b>	<b>94,045</b>	<b>128,583</b>	<b>90,380</b>

Table 2-36. Groundfish ex-vessel revenue in current (2012) dollars, \$1,000s, by "TOPAC port groups", 2003-2012.

Port	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<b>Washington Subtotal</b>	<b>\$14,862</b>	<b>\$14,787</b>	<b>\$18,601</b>	<b>\$20,137</b>	<b>\$17,254</b>	<b>\$17,949</b>	<b>\$14,786</b>	<b>\$16,529</b>	<b>\$23,224</b>	<b>\$17,007</b>
Puget Sound	\$4,801	\$4,599	\$5,071	\$4,545	\$3,033	\$2,594	\$2,838	\$2,011	\$2,030	\$1,605
North WA coast	\$4,840	\$4,662	\$4,614	\$4,099	\$3,852	\$3,844	\$4,822	\$4,659	\$5,706	\$4,377
South and central WA coast	\$5,221	\$5,526	\$8,916	\$11,492	\$10,369	\$11,511	\$7,125	\$9,859	\$15,487	\$11,025
<b>Oregon Subtotal</b>	<b>\$25,349</b>	<b>\$24,362</b>	<b>\$28,649</b>	<b>\$31,188</b>	<b>\$29,321</b>	<b>\$35,882</b>	<b>\$33,590</b>	<b>\$32,309</b>	<b>\$45,760</b>	<b>\$38,443</b>
Astoria	\$8,813	\$9,257	\$10,741	\$12,470	\$11,317	\$12,840	\$11,577	\$9,675	\$20,217	\$17,527
Tillamook	\$265	\$239	\$140	\$236	\$201	\$180	\$198	\$144	\$171	\$150
Newport	\$7,542	\$8,316	\$9,536	\$9,593	\$8,734	\$11,478	\$10,607	\$10,902	\$14,237	\$12,829
Coos Bay	\$5,663	\$4,397	\$5,222	\$5,989	\$5,797	\$7,193	\$6,416	\$6,962	\$6,221	\$4,260
Brookings	\$3,065	\$2,152	\$3,011	\$2,900	\$3,273	\$4,191	\$4,792	\$4,626	\$4,914	\$3,678
<b>California Subtotal</b>	<b>\$18,421</b>	<b>\$16,425</b>	<b>\$15,986</b>	<b>\$16,407</b>	<b>\$17,408</b>	<b>\$20,288</b>	<b>\$21,050</b>	<b>\$21,418</b>	<b>\$25,151</b>	<b>\$17,993</b>
Crescent City	\$2,108	\$1,291	\$1,661	\$1,863	\$1,903	\$2,708	\$2,520	\$1,296	\$897	\$563
Eureka	\$3,692	\$3,303	\$3,616	\$4,278	\$4,826	\$5,403	\$4,505	\$4,494	\$4,314	\$3,284
Fort Bragg	\$2,933	\$2,897	\$3,400	\$2,730	\$2,987	\$3,637	\$4,356	\$4,113	\$4,834	\$3,330
Bodega Bay	\$385	\$151	\$103	\$155	\$261	\$325	\$271	\$380	\$444	\$358
San Francisco	\$1,865	\$2,070	\$1,451	\$1,703	\$2,161	\$2,055	\$1,538	\$1,484	\$1,991	\$1,096
Monterey	\$2,847	\$2,115	\$2,067	\$2,080	\$1,434	\$1,556	\$1,364	\$1,569	\$1,899	\$1,428
Morro Bay	\$2,362	\$2,411	\$1,675	\$1,502	\$1,597	\$2,086	\$3,932	\$4,669	\$6,796	\$4,305
Santa Barbara	\$489	\$538	\$469	\$560	\$617	\$563	\$790	\$1,531	\$2,111	\$1,883
Los Angeles	\$1,090	\$1,350	\$967	\$688	\$1,059	\$982	\$1,027	\$1,063	\$1,158	\$924
San Diego	\$651	\$300	\$579	\$848	\$564	\$972	\$746	\$819	\$705	\$822
<b>Grand Total</b>	<b>\$58,633</b>	<b>\$55,574</b>	<b>\$63,236</b>	<b>\$67,732</b>	<b>\$63,983</b>	<b>\$74,120</b>	<b>\$69,427</b>	<b>\$70,256</b>	<b>\$94,135</b>	<b>\$73,443</b>

Table 2-37. Landings (mt) ex-vessel revenue in current (2012) dollars, \$1,000s, by "IOPAC port groups" and species and species groups, 2010-2012.

	P. Whiting		Sablefish		Other Roundfish		Rockfish		Flatfish		Other Groundfish	
	mt	\$000s	mt	\$000s	mt	\$000s	mt	\$000s	mt	\$000s	mt	\$000s
<b>2010</b>												
<b>Washington Subtotal</b>	<b>30,688</b>	<b>\$4,270</b>	<b>1,473</b>	<b>\$9,653</b>	<b>340</b>	<b>\$379</b>	<b>984</b>	<b>\$1,092</b>	<b>1,946</b>	<b>\$1,077</b>	<b>215</b>	<b>\$57</b>
Puget Sound			192	\$1,275	47	\$51	74	\$79	1,035	\$579	55	\$28
North WA coast			497	\$3,324	287	\$322	649	\$806	175	\$194	34	\$12
South and central WA coast	30,688	\$4,270	784	\$5,054	6	\$6	261	\$206	736	\$304	125	\$18
<b>Oregon Subtotal</b>	<b>31,539</b>	<b>\$5,628</b>	<b>2,858</b>	<b>\$15,665</b>	<b>175</b>	<b>\$672</b>	<b>2,194</b>	<b>\$2,618</b>	<b>10,365</b>	<b>\$7,133</b>	<b>1,105</b>	<b>\$593</b>
Astoria	14,099	\$2,176	681	\$2,895	83	\$106	853	\$829	5,392	\$3,436	458	\$233
Tillamook			2	\$13	6	\$39	20	\$78	*	*	*	*
Newport	17,304	\$3,439	926	\$5,482	13	\$52	572	\$586	1,590	\$1,186	295	\$158
Coos Bay	135	\$13	761	\$4,453	17	\$55	463	\$458	2,474	\$1,806	297	\$177
Brookings			487	\$2,822	55	\$421	287	\$667	901	\$691	53	\$25
<b>California Subtotal</b>	<b>2,427</b>	<b>\$430</b>	<b>2,498</b>	<b>\$11,956</b>	<b>71</b>	<b>\$481</b>	<b>1,814</b>	<b>\$5,717</b>	<b>3,077</b>	<b>\$2,704</b>	<b>276</b>	<b>\$129</b>
Crescent City	2,343	\$419	86	\$395	8	\$43	134	\$298	163	\$136	12	\$4
Eureka	84	\$11	523	\$2,629	4	\$11	402	\$560	1,598	\$1,247	115	\$36
Fort Bragg			504	\$2,588	23	\$109	546	\$786	752	\$620	32	\$10
Bodega Bay			41	\$262	2	\$10	17	\$65	*	*	*	*
San Francisco			109	\$660	5	\$23	170	\$349	400	\$433	39	\$19
Monterey			212	\$896	6	\$48	159	\$436	127	\$146	63	\$42
Morro Bay	*	*	773	\$3,288	19	\$161	134	\$1,200	7	\$15	5	\$5
Santa Barbara	<0.5	<\$1	174	\$794	5	\$68	80	\$655	4	\$11	3	\$4
Los Angeles	*	*	53	\$311	<0.5	\$2	80	\$692	11	\$52	6	\$6
San Diego			25	\$133	1	\$6	93	\$677	*	*	1	\$3
<b>2011</b>												
<b>Washington Subtotal</b>	<b>34,482</b>	<b>\$7,319</b>	<b>1,535</b>	<b>\$12,457</b>	<b>505</b>	<b>\$663</b>	<b>1,063</b>	<b>\$1,204</b>	<b>1,496</b>	<b>\$1,387</b>	<b>314</b>	<b>\$195</b>
Puget Sound	*	*	159	\$1,446	76	\$135	32	\$38	407	\$390	30	\$21
North WA coast			486	\$4,027	392	\$472	530	\$661	375	\$525	49	\$22
South and central WA coast	34,480	\$7,319	889	\$6,984	37	\$57	501	\$505	714	\$471	235	\$152
<b>Oregon Subtotal</b>	<b>68,704</b>	<b>\$16,813</b>	<b>2,305</b>	<b>\$17,662</b>	<b>497</b>	<b>\$1,217</b>	<b>1,812</b>	<b>\$2,517</b>	<b>7,478</b>	<b>\$6,901</b>	<b>1,027</b>	<b>\$651</b>

	P. Whiting		Sablefish		Other Roundfish		Rockfish		Flatfish		Other Groundfish	
	mt	\$000s	mt	\$000s	mt	\$000s	mt	\$000s	mt	\$000s	mt	\$000s
Astoria	42,444	\$10,373	590	\$3,592	391	\$539	1,085	\$1,190	4,700	\$4,037	726	\$486
Tillamook			7	\$48	8	\$50	15	\$72				
Newport	24,722	\$6,095	785	\$7,103	15	\$60	224	\$259	638	\$681	76	\$40
Coos Bay	1,538	\$345	522	\$3,906	12	\$54	247	\$299	1,473	\$1,522	169	\$94
Brookings	<0.5	<\$1	401	\$3,012	71	\$514	240	\$696	667	\$661	56	\$31
<b>California Subtotal</b>	<b>5</b>	<b>\$5</b>	<b>2,566</b>	<b>\$15,411</b>	<b>67</b>	<b>\$574</b>	<b>1,581</b>	<b>\$5,818</b>	<b>2,852</b>	<b>\$3,136</b>	<b>296</b>	<b>\$207</b>
Crescent City			72	\$457	7	\$41	54	\$160	*	*	11	\$9
Eureka	<0.5	<\$1	363	\$2,391	3	\$9	282	\$417	1,459	\$1,404	123	\$92
Fort Bragg	*	*	462	\$3,185	14	\$95	503	\$796	636	\$720	51	\$33
Bodega Bay			67	\$412	1	\$5	4	\$26	1	\$1	*	*
San Francisco	*	*	175	\$1,341	4	\$28	77	\$234	248	\$375	18	\$13
Monterey	*	*	221	\$1,223	5	\$55	121	\$370	176	\$220	62	\$31
Morro Bay			875	\$4,618	28	\$254	282	\$1,796	69	\$113	22	\$15
Santa Barbara	<0.5	<\$1	244	\$1,318	6	\$78	73	\$699	6	\$11	4	\$5
Los Angeles	<0.5	<\$1	58	\$316	<0.5	\$4	102	\$772	16	\$61	6	\$6
San Diego			30	\$150	1	\$5	85	\$548	*	*	*	*
<b>2012</b>												
<b>Washington Subtotal</b>	<b>17,535</b>	<b>\$5,882</b>	<b>1,316</b>	<b>\$7,514</b>	<b>406</b>	<b>\$558</b>	<b>1,276</b>	<b>\$1,492</b>	<b>1,511</b>	<b>\$1,385</b>	<b>269</b>	<b>\$175</b>
Puget Sound	*	*	142	\$911	31	\$57	170	\$205	400	\$380	62	\$52
North WA coast			483	\$2,766	285	\$355	620	\$783	344	\$419	87	\$55
South and central WA coast	17,535	\$5,882	691	\$3,837	89	\$147	486	\$504	767	\$587	120	\$69
<b>Oregon Subtotal</b>	<b>48,831</b>	<b>\$14,611</b>	<b>2,152</b>	<b>\$11,529</b>	<b>709</b>	<b>\$1,550</b>	<b>1,857</b>	<b>\$2,660</b>	<b>7,188</b>	<b>\$7,315</b>	<b>990</b>	<b>\$779</b>
Astoria	23,518	\$7,558	596	\$3,150	598	\$841	1,107	\$1,225	4,189	\$4,171	665	\$582
Tillamook			*	*	12	\$66	18	\$81				
Newport	25,312	\$7,053	741	\$4,512	17	\$69	246	\$295	774	\$829	121	\$71
Coos Bay	1	<\$1	464	\$2,159	13	\$58	231	\$328	1,510	\$1,618	158	\$98
Brookings	<0.5	<\$1	352	\$1,705	70	\$515	255	\$731	714	\$699	46	\$28

	P. Whiting		Sablefish		Other Roundfish		Rockfish		Flatfish		Other Groundfish	
	mt	\$000s	mt	\$000s	mt	\$000s	mt	\$000s	mt	\$000s	mt	\$000s
<b>California Subtotal</b>	<b>4</b>	<b>\$6</b>	<b>1,778</b>	<b>\$8,989</b>	<b>83</b>	<b>\$639</b>	<b>1,577</b>	<b>\$5,231</b>	<b>2,582</b>	<b>\$2,891</b>	<b>318</b>	<b>\$238</b>
Crescent City	*	*	63	\$318	6	\$38	60	\$162	*	*	6	\$2
Eureka	<0.5	<\$1	318	\$1,515	6	\$19	291	\$413	1,198	\$1,241	122	\$97
Fort Bragg	2	\$2	392	\$1,845	18	\$86	475	\$728	546	\$634	52	\$36
Bodega Bay			*	*	1	\$12	5	\$50	*	*	*	*
San Francisco	*	*	98	\$498	6	\$35	90	\$261	170	\$294	11	\$8
Monterey	*	*	164	\$728	8	\$79	98	\$345	284	\$242	66	\$35
Morro Bay	2	\$2	426	\$1,982	30	\$271	340	\$1,656	318	\$352	47	\$43
Santa Barbara			169	\$1,121	7	\$85	64	\$649	10	\$18	7	\$10
Los Angeles	<0.5	\$2	47	\$277	1	\$8	79	\$568	12	\$64	7	\$6
San Diego			57	\$411	1	\$6	76	\$400	1	\$3	1	\$2

**Table 2-38. Landings (mt) ex-vessel revenue in current (2012) dollars, \$1,000s, by "IOPAC port groups" and commercial fishery sectors, 2010-2012 (Washington and Oregon).**

	2010				2011				2012			
	P. Whiting		Other Groundfish		P. Whiting		Other Groundfish		P. Whiting		Other Groundfish	
	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s
<b>Puget Sound</b>			1,399	1,979	*	*	703	2,022	*	*	804	1,604
Shoreside IFQ Trawl (Nonwhiting)			1,250	1,016	*	*	551	698	*	*	*	*
Shoreside IFQ Non-trawl							*	*			*	*
Non Nearshore Fixed Gear			142	941			131	1,141			108	626
Incidental Open Access			*	*							*	*
Exempted trawl, EFP/Research, Misc.			*	*			*	*			*	*
<b>North WA coast</b>			172	947			154	1,124			154	846
Shoreside IFQ Trawl (Nonwhiting)			*	*								
Non Nearshore Fixed Gear			159	927			153	1,121			150	829
Non Fixed Gear Open Access			*	*							2	13
Incidental Open Access			1	2			1	3			1	4
<b>South and central WA coast</b>	28,721	4,065	1,684	4,237	22,724	5,617	2,032	6,563	16,922	5,749	1,940	3,899
Shoreside IFQ Trawl (Whiting)	28,720	4,065	254	96	22,063	5,456	219	179	16,813	5,715	157	98
Shoreside IFQ Trawl (Nonwhiting)	<0.5	<\$1	866	570	*	*	1,172	1,867	*	*	1,346	1,524
Shoreside IFQ Non-trawl							297	1,814			183	857
Non Nearshore Fixed Gear			*	*			332	2,677			249	1,399
Non Fixed Gear Open Access			1	7			*	*			1	4
Incidental Open Access			2	6			*	*			3	12
Exempted trawl, EFP/Research, Misc.			*	*	*	*	10	12	*	*	3	6
<b>Astoria</b>	14,099	2,176	7,468	7,499	42,444	10,373	7,492	9,843	23,518	7,558	7,154	9,969
Shoreside IFQ Trawl (Whiting)	14,077	2,171	102	106	41,631	10,160	415	344	23,117	7,440	297	229
Shoreside IFQ Trawl (Nonwhiting)	*	*	7,331	7,190	18	3	6,805	8,188	14	3	6,433	7,635
Shoreside IFQ Non-trawl							45	389			257	1,591
Non Nearshore Fixed Gear			31	195			90	759			64	390

Nearshore Fixed Gear							*	*				
Non Fixed Gear Open Access			*	*								
Incidental Open Access			*	*			2	9			3	8
Exempted trawl, EFP/Research, Misc.	*	*	4	6	795	210	135	155	387	115	100	116
<b>Tillamook</b>			<b>37</b>	<b>144</b>			<b>30</b>	<b>171</b>			<b>30</b>	<b>150</b>
Shoreside IFQ Trawl (Nonwhiting)			*	*								
Non Nearshore Fixed Gear			4	19			9	60			4	21
Nearshore Fixed Gear			24	109			21	110			25	126
Non Fixed Gear Open Access			<0.5	1			*	*			*	*
Incidental Open Access			*	*			<0.5	<\$1			<0.5	2
Exempted trawl, EFP/Research, Misc.							*	*			*	*
<b>Newport</b>	<b>17,304</b>	<b>3,439</b>	<b>3,397</b>	<b>7,464</b>	<b>24,722</b>	<b>6,095</b>	<b>1,737</b>	<b>8,143</b>	<b>25,312</b>	<b>7,052</b>	<b>1,899</b>	<b>5,776</b>
Shoreside IFQ Trawl (Whiting)	*	*	157	102	24,595	6,066	132	250	25,242	7,044	198	271
Shoreside IFQ Trawl (Nonwhiting)	*	*	2,723	3,808	<0.5	<\$1	890	1,435	<0.5	<\$1	1,052	1,532
Shoreside IFQ Non-trawl							260	2,291			216	1,174
Non Nearshore Fixed Gear			500	3,486			409	4,035			377	2,656
Nearshore Fixed Gear			13	47			7	30			11	49
Non Fixed Gear Open Access			<0.5	<\$1			<0.5	3			<0.5	1
Incidental Open Access			2	9			2	15			3	14
Exempted trawl, EFP/Research, Misc.			3	11	*	*	37	82	*	*	41	80
<b>Coos Bay</b>	<b>135</b>	<b>13</b>	<b>4,012</b>	<b>6,949</b>	<b>1,538</b>	<b>345</b>	<b>2,423</b>	<b>5,876</b>	<b>1</b>	<b>&lt;\$1</b>	<b>2,377</b>	<b>4,260</b>
Shoreside IFQ Trawl (Whiting)	*	*	1	<\$1	*	*	*	*				
Shoreside IFQ Trawl (Nonwhiting)	*	*	3,616	4,247	1	<\$1	2,094	3,101	*	*	2,138	2,836
Shoreside IFQ Non-trawl							20	132			*	*
Non Nearshore Fixed Gear			388	2,656	*	*	286	2,541			207	1,269
Nearshore Fixed Gear			5	37			8	57			11	81
Non Fixed Gear Open Access			<0.5	1			*	*			*	*
Incidental Open Access			1	2			3	15			1	5
Exempted trawl, EFP/Research, Misc.			1	5			8	25	*	*	9	12
<b>Brookings</b>			<b>1,783</b>	<b>4,626</b>	<b>&lt;0.5</b>	<b>&lt;\$1</b>	<b>1,436</b>	<b>4,914</b>	<b>&lt;0.5</b>	<b>&lt;\$1</b>	<b>1,437</b>	<b>3,678</b>

Shoreside IFQ Trawl (Nonwhiting)			1,321	1,867	<0.5	<\$1	967	1,621	<0.5	<\$1	1,070	1,527
Non Nearshore Fixed Gear			322	1,912	*	*	285	2,240			207	1,124
Nearshore Fixed Gear			133	835			163	1,012			154	1,014
Non Fixed Gear Open Access							*	*			<0.5	<\$1
Incidental Open Access			<0.5	<\$1			*	*			1	6
Exempted trawl, EFP/Research, Misc.			6	12	*	*	21	39			5	7

**Table 2-39. Landings (mt) ex-vessel revenue in current (2012) dollars, \$1,000s, by "IOPAC port groups" and commercial fishery sectors, 2010-2012 (California).**

	2010				2011				2012			
	P. Whiting		Other Groundfish		P. Whiting		Other Groundfish		P. Whiting		Other Groundfish	
	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s
<b>Crescent City</b>	<b>2,343</b>	<b>419</b>	<b>403</b>	<b>876</b>			<b>386</b>	<b>897</b>	*	*	<b>178</b>	<b>563</b>
Shoreside IFQ Trawl (Whiting)	*	*	29	7								
Shoreside IFQ Trawl (Nonwhiting)			259	403			*	*	*	*	*	*
Non Nearshore Fixed Gear			40	181			52	344			36	202
Nearshore Fixed Gear			59	272			36	165			32	155
Incidental Open Access			*	*			<0.5	1			<0.5	1
Exempted trawl, EFP/Research, Misc.	*	*	*	*			*	*			*	*
<b>Eureka</b>	<b>84</b>	<b>11</b>	<b>2,641</b>	<b>4,483</b>	<b>&lt;0.5</b>	<b>&lt;\$1</b>	<b>2,229</b>	<b>4,314</b>	<b>&lt;0.5</b>	<b>&lt;\$1</b>	<b>1,935</b>	<b>3,284</b>
Shoreside IFQ Trawl (Whiting)	79	9	2	3								
Shoreside IFQ Trawl (Nonwhiting)	6	2	2,441	3,450	<0.5	<\$1	2,120	3,415	<0.5	<\$1	1,855	2,722
Non Nearshore Fixed Gear			194	1,009			105	874			77	540
Nearshore Fixed Gear			4	20			4	17			3	20
Non Fixed Gear Open Access			*	*			*	*			*	*
Incidental Open Access			*	*			*	*			<0.5	<\$1
<b>Fort Bragg</b>			<b>1,857</b>	<b>4,113</b>	<b>*</b>	<b>*</b>	<b>1,665</b>	<b>4,829</b>	<b>2</b>	<b>2</b>	<b>1,482</b>	<b>3,329</b>
Shoreside IFQ Trawl (Nonwhiting)			1,572	2,338	*	*	1,331	2,401	2	2	1,195	1,843
Shoreside IFQ Non-trawl							*	*			*	*
Non Nearshore Fixed Gear			268	1,586			286	2,004			254	1,306
Nearshore Fixed Gear			15	178			17	210			10	104
Non Fixed Gear Open Access							*	*			<0.5	2



	2010				2011				2012			
	P. Whiting		Other Groundfish		P. Whiting		Other Groundfish		P. Whiting		Other Groundfish	
	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s
Incidental Open Access			*	*			<0.5	1			*	*
Exempted trawl, EFP/Research, Misc.			*	*			<0.5	2			1	3
<b>Bodega Bay</b>			<b>75</b>	<b>380</b>			<b>73</b>	<b>444</b>			<b>53</b>	<b>358</b>
Shoreside IFQ Trawl (Nonwhiting)			*	*			*	*			*	*
Non Nearshore Fixed Gear			41	265			68	420			47	312
Nearshore Fixed Gear			3	48			1	18			3	43
Non Fixed Gear Open Access							<0.5	2				
Incidental Open Access			<0.5	1			<0.5	1			<0.5	<\$1
Exempted trawl, EFP/Research, Misc.			*	*							*	*
<b>San Francisco</b>			<b>722</b>	<b>1,484</b>	*	*	<b>521</b>	<b>1,991</b>	*	*	<b>373</b>	<b>1,096</b>
Shoreside IFQ Trawl (Nonwhiting)			625	820	*	*	335	557	*	*	251	420
Shoreside IFQ Non-trawl							54	323			27	49
Non Nearshore Fixed Gear			72	509			107	935			60	381
Nearshore Fixed Gear			10	119			11	136			12	140
Non Fixed Gear Open Access			*	*			<0.5	5			1	4
Incidental Open Access			*	*			<0.5	4			1	5
Exempted trawl, EFP/Research, Misc.			15	27			14	31			23	97
<b>Monterey</b>			<b>566</b>	<b>1,569</b>	*	*	<b>584</b>	<b>1,899</b>	*	*	<b>618</b>	<b>1,428</b>
Shoreside IFQ Trawl (Nonwhiting)			*	*			*	*			*	*
Shoreside IFQ Non-trawl							*	*			14	60
Non Nearshore Fixed Gear			214	865	*	*	195	928	*	*	169	682
Nearshore Fixed Gear			13	147			12	149			14	173
Non Fixed Gear Open Access			*	*			*	*			3	9
Incidental Open Access			<0.5	1			1	4			<0.5	2
Exempted trawl, EFP/Research, Misc.			*	*			5	15			*	*

	2010				2011				2012			
	P. Whiting		Other Groundfish		P. Whiting		Other Groundfish		P. Whiting		Other Groundfish	
	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s	mt	\$1,000s
<b>Morro Bay</b>	*	*	937	4,668			1,275	6,796	2	2	1,161	4,303
Shoreside IFQ Trawl (Nonwhiting)							*	*	*	*	*	*
Shoreside IFQ Non-trawl							454	2,336			209	964
Non Nearshore Fixed Gear	*	*	655	2,764			549	2,890	*	*	296	1,374
Nearshore Fixed Gear			74	945			96	1,243			86	1,124
Non Fixed Gear Open Access			*	*			*	*			*	*
Incidental Open Access			1	5			1	4			1	6
Exempted trawl, EFP/Research, Misc.			207	955			8	37			11	41
<b>Santa Barbara</b>	<0.5	<\$1	266	1,531	*	*	332	2,111			257	1,883
Non Nearshore Fixed Gear	<0.5	<\$1	239	1,241	*	*	300	1,827			222	1,584
Nearshore Fixed Gear			14	229			10	178			14	212
Non Fixed Gear Open Access			1	2			3	9			*	*
Incidental Open Access			4	29			4	37			*	*
Exempted trawl, EFP/Research, Misc.			9	29			15	61			17	50
<b>Los Angeles</b>	*	*	151	1,063	*	*	181	1,158	<0.5	2	146	922
Non Nearshore Fixed Gear	*	*	132	991	*	*	158	1,065	<0.5	2	125	810
Nearshore Fixed Gear			4	33			5	40			7	65
Non Fixed Gear Open Access			3	3			3	3			3	3
Incidental Open Access			3	5			3	6			3	9
Exempted trawl, EFP/Research, Misc.			9	31			13	43			8	36
<b>San Diego</b>			119	819			117	705			136	821
Non Nearshore Fixed Gear			116	800			114	690			131	798
Nearshore Fixed Gear			1	8			*	*			1	5
Non Fixed Gear Open Access			*	*			1	1			1	4
Incidental Open Access			1	9			1	8			2	11
Exempted trawl, EFP/Research, Misc.			*	*			*	*			1	3

**Table 2-40. Number of vessels making at least one groundfish landing by "IOPAC port groups" and commercial fishery sectors, 5 years 2008-2012.**

Fishery / Port	Vessels	Fishery / Port	Vessels	Fishery / Port	Vessels
<b>Shoreside IFQ Trawl (Whiting)</b>		<b>Nearshore Fixed Gear</b>		<b>Exempted trawl, EFP/Research, Misc.</b>	
<b>Washington</b>	<b>21</b>	<b>Oregon</b>	<b>216</b>	<b>Washington</b>	<b>19</b>
South and central WA coast	21	Astoria	1	Puget Sound	4
<b>Oregon</b>	<b>32</b>	Tillamook	42	South and central WA coast	16
Astoria	26	Newport	35	<b>Oregon</b>	<b>76</b>
Newport	20	Coos Bay	36	Astoria	23
Coos Bay	4	Brookings	132	Tillamook	2
<b>California</b>	<b>12</b>	<b>California</b>	<b>382</b>	Newport	28
Crescent City	11	Crescent City	27	Coos Bay	16
Eureka	7	Eureka	18	Brookings	29
<b>Coastwide</b>	<b>41</b>	Fort Bragg	36	<b>California</b>	<b>133</b>
<b>Shoreside IFQ Trawl (Nonwhiting)</b>		Bodega Bay	20	Crescent City	5
<b>Washington</b>	<b>20</b>	San Francisco	49	Eureka	2
Puget Sound	8	Monterey	56	Fort Bragg	17
North WA coast	3	Morro Bay	113	Bodega Bay	2
South and central WA coast	13	Santa Barbara	61	San Francisco	23
<b>Oregon</b>	<b>84</b>	Los Angeles	23	Monterey	8
Astoria	42	San Diego	13	Morro Bay	35
Tillamook	3	<b>Coastwide</b>	<b>597</b>	Santa Barbara	38
Newport	29	<b>Non Fixed Gear Open Access</b>		Los Angeles	24
Coos Bay	29	<b>Washington</b>	<b>18</b>	San Diego	8
Brookings	14	North WA coast	6	<b>Coastwide</b>	<b>218</b>
<b>California</b>	<b>42</b>	South and central WA coast	12		
Crescent City	13	<b>Oregon</b>	<b>44</b>		
Eureka	16	Astoria	3		
Fort Bragg	7	Tillamook	5		
Bodega Bay	3	Newport	11		
San Francisco	13	Coos Bay	19		
Monterey	3	Brookings	7		

Fishery / Port	Vessels	Fishery / Port	Vessels	Fishery / Port	Vessels
Morro Bay	2	<b>California</b>	<b>88</b>		
Santa Barbara	1	Eureka	3		
<b>Coastwide</b>	<b>125</b>	Fort Bragg	7		
<b>Shoreside IFQ Non-trawl</b>		Bodega Bay	3		
<b>Washington</b>	<b>10</b>	San Francisco	12		
Puget Sound	3	Monterey	10		
South and central WA coast	10	Morro Bay	7		
<b>Oregon</b>	<b>17</b>	Santa Barbara	18		
Astoria	7	Los Angeles	26		
Newport	8	San Diego	6		
Coos Bay	4	<b>Coastwide</b>	<b>150</b>		
<b>California</b>	<b>22</b>	<b>Incidental Open Access</b>			
Fort Bragg	1	<b>Washington</b>	<b>46</b>		
San Francisco	4	Puget Sound	2		
Monterey	4	North WA coast	22		
Morro Bay	15	South and central WA coast	34		
<b>Coastwide</b>	<b>40</b>	<b>Oregon</b>	<b>200</b>		
<b>Non Nearshore Fixed Gear</b>		Astoria	25		
<b>Washington</b>	<b>124</b>	Tillamook	27		
Puget Sound	23	Newport	59		
North WA coast	40	Coos Bay	76		
South and central WA coast	87	Brookings	29		
<b>Oregon</b>	<b>317</b>	<b>California</b>	<b>367</b>		
Astoria	33	Crescent City	11		
Tillamook	43	Eureka	10		
Newport	123	Fort Bragg	16		
Coos Bay	97	Bodega Bay	20		
Brookings	64	San Francisco	51		
<b>California</b>	<b>722</b>	Monterey	49		
Crescent City	21	Morro Bay	70		
Eureka	61	Santa Barbara	64		

Fishery / Port	Vessels	Fishery / Port	Vessels	Fishery / Port	Vessels
Fort Bragg	91	Los Angeles	62		
Bodega Bay	32	San Diego	46		
San Francisco	126	<b>Coastwide</b>	<b>604</b>		
Monterey	142				
Morro Bay	182				
Santa Barbara	85				
Los Angeles	75				
San Diego	61				
<b>Coastwide</b>	<b>1129</b>				

**Table 2-41. Number of vessels making at least one groundfish landing by commercial fishery sector and length category (feet), 5 years 2008-2012.**

Fishery	Length category)				
	<= 40	41-50	51-60	61-70	71-150
Shoreside IFQ Trawl (Whiting)			2	8	35
Shoreside IFQ Trawl (Nonwhiting)	1	15	34	34	50
Shoreside IFQ Nontrawl	9	10	7	12	5
Non Nearshore Fixed Gear	831	211	65	26	9
Nearshore Fixed Gear	563	30	2		1
Non Fixed Gear Open Access	102	41	6		1
Incidental Open Access	437	124	27	8	10
Exempted trawl, EFP/Research, Misc.	109	47	27	15	22

**Table 2-42. Engagement (groundfish ex-vessel revenue in port as percent of ex-vessel coastwide revenue) and dependence (groundfish ex-vessel revenue in port as percent of total ex-vessel revenue in port), using current (2012) dollars, 2003-2012.**

	Engagement	Dependence
Puget Sound	5%	44%
North WA coast	7%	45%
South and central WA coast	14%	14%
<b>Washington</b>	<b>25%</b>	<b>20%</b>
Astoria	18%	37%
Tillamook	0%	5%
Newport	15%	30%
Coos Bay	8%	22%
Brookings	5%	32%
<b>Oregon</b>	<b>47%</b>	<b>30%</b>
Crescent City	2%	10%
Eureka	6%	26%
Fort Bragg	5%	36%
Bodega Bay	0%	4%
San Francisco	3%	9%
Monterey	3%	16%
Morro Bay	5%	65%
Santa Barbara	1%	3%
Los Angeles	1%	3%
San Diego	1%	10%
<b>California</b>	<b>28%</b>	<b>12%</b>
<b>Coastwide</b>		<b>19%</b>

**Table 2-43. Groundfish landings (mt) on the west coast (Washington-Oregon-California) from inside and outside the Pacific Council management area.**

Year	Area		Council Area		Area	
	metric tons	percent	metric tons	percent	metric tons	percent
1981	103,344	85%	8,254	7%	9,827	8%
1982	119,356	85%	10,051	7%	10,579	8%
1983	98,978	80%	10,114	8%	14,352	12%
1984	89,804	73%	18,483	15%	14,353	12%
1985	90,923	77%	8,969	8%	17,760	15%
1986	82,480	76%	5,185	5%	20,488	19%
1987	91,982	77%	553	<1%	26,633	22%
1988	92,248	85%	270	<1%	16,123	15%
1989	99,372	84%	131	<1%	18,456	16%
1990	94,539	86%	1,755	2%	13,416	12%
1991	105,547	83%	3,711	3%	17,425	14%
1992	132,555	85%	379	<1%	22,179	14%
1993	116,394	87%	437	<1%	17,125	13%
1994	135,677	91%	569	<1%	12,093	8%
1995	134,493	92%	485	<1%	10,523	7%
1996	146,452	92%	459	<1%	12,956	8%
1997	143,571	93%	346	<1%	10,488	7%
1998	131,011	93%	363	<1%	9,729	7%
1999	125,883	95%	251	<1%	6,364	5%
2000	123,031	97%	191	<1%	3,586	3%
2001	103,556	97%	93	<1%	3,154	3%
2002	75,056	97%	94	<1%	2,058	3%
2003	82,905	97%	91	<1%	2,532	3%
2004	122,303	97%	87	<1%	3,301	3%
2005	135,454	98%	28	<1%	2,898	2%
2006	151,260	100%	39	<1%	484	<1%
2007	117,495	100%	48	<1%	356	<1%
2008	97,667	100%	31	<1%	158	<1%
2009	82,194	100%	23	<1%	125	<1%
2010	94,045	100%	23	<1%	79	<1%
2011	128,585	100%	18	<1%	136	<1%
2012	90,384	98%	15	<1%	1,732	2%

## 2.2 PacFIN Groundfish Sector Criteria

The field “dahl\_sector” in the vessel daily summary file (vdrfd) contains numeric codes identifying groundfish “sectors.” These sectors are meant to identify landings according to fishery components, or sectors, used in management. Sectors are defined through a combination of species composition of landings, gear type, and permit status, among other factors. It should be noted that the results of this coding may differ from other categorizations of landings, such as that developed by the West Coast Groundfish Observer Program in their total mortality reports, because of differences in the underlying data structure (e.g., use of fish ticket data versus vessel summary data) and the procedures used to code the data. The record level for coding data is:

- Sector codes assigned at vessel-day-gear level
- Species composition criteria used in sector coding are calculated on a vessel-day-gear basis (agid+drvid+tdate+grgroup)

Sector codes 1, 2, and 16 are not available in the PacFIN Explorer tool, because the necessary landings data reside in a separate database (NORPAC).

The PFMC trawl rationalization program was implemented in 2011. This introduced individual fishing quota (IFQ) management for trawl-endorsed groundfish limited access permits. The program also allows vessels with these permits to use any legal groundfish gear. To account for this change a new sector was created (sector 20) for the shoreside non-trawl IFQ sector and the procedure for updating the sector field was updated accordingly.

The criteria used to code landings by sector are described below.

### Global criteria

- Year  $\geq$  1994
- Catch is from a PFMC area
- Sectors 1-15 PARGRP=C
- Sectors 16-19 PARGRP=I
- Sectors 01, 02, and 16 AGID = N (as noted)
- Sectors 03-15, 17-19 AGID = W, O, C

### 1: Whiting Catcher Processor Sector

- AGID =N
- Gear Group is TWL
- Valid trawl endorsement
- DRVID=PROC

### 2: Whiting Mothership Sector

- AGID =N
- Gear Group is TWL
- Valid trawl endorsement
- DRVID not equal to PROC

### 3: Shoreside Whiting Sector (Shoreside IFQ Trawl)

- Removal type (ftl.removal\_type) not R (research)
- Whiting (PWHT) RWT  $\geq$  50% total vessel-day-gear RWT
- Gear Group is TWL
- Valid trawl endorsement



#### 4: Shoreside Nonwhiting Trawl Sector (Shoreside IFQ Trawl)

- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Whiting (PWHT) RWT < 50% total vessel-day-gear RWT
- Nonwhiting groundfish (sp.mgrp=GRND and sp.spid <> PWHT) RWT >= 50% total vessel-day-gear RWT
- Groundfish (sp.mgrp=GRND) RWT > California halibut (CHLB, CHL1) RWT
- Pink shrimp, ridgeback prawn, or spot prawn (PHSP, RPRW, SPRW) RWT < 100 lbs
- Gear Group is TWL
- Valid trawl endorsement

#### 5: Nearshore Sector (Limited Entry)

- Gear Group is HKL or POT
- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Sum of nearshore species<sup>12</sup> vessel-day-gear RWT >0
- Sablefish RWT = 0
- Groundfish (sp.mgrp=GRND) >= 50% of total vessel-day-gear RWT.<sup>13</sup>
- Valid longline endorsement or pot gear endorsement.
- IFQ landing flag (ifq\_landing) = 'N'<sup>14</sup>

#### 6: Nearshore Sector (Open Access)

- Gear Group = HKL or POT
- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Sum of nearshore species vessel-day-gear RWT >0
- Sablefish RWT = 0
- Groundfish (sp.mgrp=GRND) >= 50% of total vessel-day-gear RWT
- No valid longline endorsement or pot gear endorsement
- IFQ landing flag (ifq\_landing) = 'N'

#### 7: Non Nearshore Sector (Limited Entry)

- Gear Group is HKL or POT
- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Sum of nearshore species RWT =0
- Sablefish RWT >0
- Groundfish (sp.mgrp=GRND) >= 50% of total vessel-day-gear RWT
- Valid longline endorsement or pot gear endorsement
- IFQ landing flag (ifq\_landing) = 'N'

#### 8: Non Nearshore Sector (Open Access)

- Gear Group = HKL or POT
- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Sum of nearshore species RWT =0
- Sum of sablefish vessel-day-gear RWT >0
- Groundfish (sp.mgrp=GRND) >= 50% of total vessel-day-gear RWT

---

<sup>12</sup> Nearshore rockfish species as defined by WCGOP spid list, see Table 2.

<sup>13</sup> This is equivalent to Groundfish (sp.mgrp=GRND) >= non-groundfish (all other sp.mgrp) by weight.

<sup>14</sup> The ifq\_landing code is assigned at the fish ticket level (ft table). This means that species not managed under the Groundfish Shoreside IFQ Program are flagged if landed with IFQ species.

- No valid longline endorsement or pot gear endorsement
- IFQ landing flag (ifq\_landing) = 'N'

#### 9: Non Nearshore Non Sablefish Sector (Limited Entry)

- Gear Group is HKL or POT
- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Sum of nearshore species RWT =0
- Sablefish RWT =0
- Groundfish (sp.mgrp=GRND) >= 50% of total vessel-day-gear RWT
- Valid longline endorsement or pot gear endorsement
- IFQ landing flag (ifq\_landing) = 'N'

#### 10: Non Nearshore Non Sablefish Sector (Open Access)

- Gear Group = HKL or POT
- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Sum of nearshore species RWT =0
- Sablefish =0
- Groundfish (sp.mgrp=GRND) >= 50% of total vessel-day-gear RWT
- No valid longline endorsement or pot gear endorsement
- IFQ landing flag (ifq\_landing) = 'N'

#### 11: Non Fixed Gear Directed Open Access Sector

- Gear Group = NET or TLS
- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Groundfish (sp.mgrp=GRND) RWT >= 50% of total vessel-day-gear RWT
- IFQ landing flag (ifq\_landing) = 'N'

#### 12: Incidental Open Access Sector

- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Groundfish (sp.mgrp=GRND) RWT > 0 and <50% of total vessel-day-gear RWT
- Gear Group = HKL or POT or NET or TLS
- No valid longline endorsement or pot gear endorsement
- IFQ landing flag (ifq\_landing) = 'N'

#### 13: Exempted Trawl Sector (With Groundfish Landings)

- Removal type (ftl.removal\_type) not E or R (EFP, research)
- Gear group is TWL or TWS
- Groundfish (sp.mgrp=GRND) RWT > 0
- California halibut (CHLB, CHL1) RWT > 0 or pink shrimp, ridgeback prawn, or spot prawn (PSHP, RPRW, SPRW) RWT => 100 lbs
- No valid trawl endorsement

#### 14: EFP and Miscellaneous Sector

- Groundfish (sp.mgrp=GRND) RWT > 0
- *Landings not accounted for in other sectors*

#### 15: Commercial Non-groundfish Sector

- groundfish (sp.mgrp=GRND) RWT = 0

- IFQ landing flag (ifq\_landing) = 'N'

16: Treaty Mothership Whiting Sector

- AGID =N
- Whiting (PWHT) RWT >= 50% total vessel-day-gear RWT
- Gear Group is TWL
- PROC is a mothership

17: Treaty Shoreside Whiting Sector

- Whiting (PWHT) RWT >= 50% total vessel-day-gear RWT
- Gear Group is TWL

18: Treaty Shoreside Nonwhiting Groundfish Sector

- Whiting (PWHT) RWT < 50% total vessel-day-gear RWT
- Groundfish (sp.mgrp=GRND) RWT > 0

19: Treaty Non-groundfish Sector

- Groundfish (sp.mgrp=GRND) RWT = 0

20: Shoreside IFQ Non-trawl Sector<sup>15</sup>

- Gear Group not TWL
- Removal type (ftl.removal\_type) not E or R (EFP, research)
- IFQ landing flag (ifq\_landing) = 'Y'

XXX

- Remaining records not accounted for in sectors

---

<sup>15</sup> The vdrfd.dahl\_sector column is coded by means of a SQL searched CASE expression. In a CASE expression the conditions are evaluated in order. When a condition evaluates to TRUE, the corresponding result is returned, and the execution of CASE logic ends. Therefore the order of each clause in the expression is relevant. The clause for Sector 20 is placed between those for Sector 12 and Sector 13.

**Table 2-44. Sector codes.**

Code	Sector Name
1	Whiting Catcher Processor
2	Whiting Mothership
3	Shoreside Whiting (Shoreside IFQ Trawl)
4	Shoreside Nonwhiting Trawl (Shoreside IFQ Trawl)
5	Nearshore (Limited Entry)
6	Nearshore (Open Access)
7	Non Nearshore (Limited Entry)
8	Non Nearshore (Open Access)
9	Non Nearshore Non Sablefish (Limited Entry)
10	Non Nearshore Non Sablefish (Open Access)
11	Non Fixed Gear Directed Open Access
12	Incidental Open Access
13	Exempted Trawl (With Groundfish Landings)
14	EFP and Miscellaneous
15	Commercial Non-groundfish
16	Treaty Mothership Whiting
17	Treaty Shoreside Whiting
18	Treaty Shoreside Nonwhiting Groundfish
19	Treaty Non-groundfish
20	Shoreside IFQ Non-trawl Sector

**Table 2-45. Nearshore species list. (Source: M. Bellman, Heery, E., Jannot, J., and Majewski, J. 2010. *Explicit Retrieval and Processing of PacFIN Data Used in Total Mortality Estimation*. FRAM, NWFSC, NMFS.)**

SPID	Common Name	SPID	Common Name
BISC	BROWN IRISH SCULPIN	KLPG	KELP GREENLING
BLCK	BLACK ROCKFISH	KLPR	KELP ROCKFISH
BLK1	NOM. BLACK ROCKFISH	NSHR	NORTHERN NEAR-SHORE ROCKFISH
BLU1	NOM. BLUE ROCKFISH	NUSR	NOR. UNSP. NEAR-SHORE ROCKFISH
BLUR	BLUE ROCKFISH	OLV1	NOM. OLIVE ROCKFISH
BRW1	NOM. BROWN ROCKFISH	OLVE	OLIVE ROCKFISH
BRWN	BROWN ROCKFISH	QLB1	NOM. QUILLBACK ROCKFISH
BSCL	BUFFALO SCULPIN	QLBK	QUILLBACK ROCKFISH
BYEL	BLACK-AND-YELLOW ROCKFISH	RCK2	UNSP. BOLINA RCKFSH
BYL1	NOM. BLACK-AND-YELLOW ROCKFISH	RCK7	UNSP. GOPHER RCKFSH
CBZ1	NOM. CABEZON	RCK9	BLACK+BLUE ROCKFISH
CBZN	CABEZON	RSCL	RED IRISH SCULPIN
CHN1	NOM. CHINA ROCKFISH	SCOR	CALIFORNIA SCORPIONFISH
CHNA	CHINA ROCKFISH	SCR1	NOM. CALIF. SCORPIONFISH
CLC1	NOM. CALICO ROCKFISH	SHP1	NOM. CALIFORNIA SHEEPHEAD
CLCO	CALICO ROCKFISH	SHPD	CALIFORNIA SHEEPHEAD
COP1	NOM. COPPER ROCKFISH	SSHR	SOUTHERN NEAR-SHORE ROCKFISH
COPP	COPPER ROCKFISH	SSRD	Deep So. Near-shore RF
GPH1	NOM. GOPHER ROCKFISH	SSRS	Shallow So. Near-shore RF
GPHR	GOPHER ROCKFISH	SUSR	SOU. UNSP. NEAR-SHORE ROCKFISH
GRAS	GRASS ROCKFISH	TRE1	NOM. TREEFISH
GRS1	NOM. GRASS ROCKFISH	TREE	TREEFISH
KGL1	NOM. KELP GREENLING	UDNR	UNSP. DEEP NEAR-SHORE RF
KLP1	NOM. KELP ROCKFISH	UGLG	UNSP. GREENLING
		USHR	UNSP. NEAR-SHORE ROCKFISH

## CHAPTER 3      **ESSENTIAL FISH HABITAT**

---

Essential fish habitat (EFH) for groundfish was first established in 1998 under FMP Amendment 11, and in response to the Magnuson Act reauthorization of 1996. EFH was revised significantly and finalized in 2006 as part of [Amendment 19 to the groundfish FMP](#).

The EFH regulations call for a review of EFH elements at least every five years. The current review was initiated in December 2010, and consists of three phases. Phase 1 consisted of a literature survey and a subsequent synthesis report by NMFS. The [Phase 1 Report](#) summarized new and newly-available information regarding physical and biogenic habitats, habitat models, trophic interactions, and fishing and non-fishing activities that may affect groundfish EFH (PFMC 2012). The [NMFS synthesis report](#) included comparisons of habitat, substrate, and fishing effort maps, pre-2005 and post-2005 (NMFS 2013).

The second phase consisted of a further examination of the new and newly-available information compiled during the first phase, and also consisted of a request for proposals to generate potential changes to existing EFH elements. The Council received eight proposals, of which two were subsequently withdrawn from consideration. The Essential Fish Habitat Review Committee (EFHRC) managed the first two phases, which concluded with a final [Phase 2 report](#) in March 2014 (PFMC 2014).

The Council will consider changes to existing EFH provisions, based on best available science, the proposals received in response to the RFP, and AB and public input. Any major changes to EFH would be contained in an FMP amendment, while more minor changes may be made outside of the amendment process.

## CHAPTER 4 SAFETY AT SEA

---

National Standard 10 (NS10) guidelines interpreting the MSA state, “Conservation and management measures shall, to the extent practicable, promote the safety of human life at sea”. During preparation of any FMP, FMP amendment, or regulation that might affect safety of human life at sea, the Council should consult with the United States Coast Guard (USCG) and the fishing industry as to the nature and extent of any adverse impacts. This consultation may be done through a Council advisory panel, committee, or other review of the FMP, FMP amendment, or regulations.

There are many ways in which an FMP may avoid or provide alternative measures to reduce potential impacts on safety of human life at sea. The following is a list of some factors that could be considered when management measures are developed:

- 1) Setting seasons to avoid hazardous weather.
- 2) Providing for seasonal or trip flexibility to account for bad weather (weather days).
- 3) Allowing for pre- and post-season “soak time” to deploy and pick up fixed gear, so as to avoid overloading vessels with fixed gear.
- 4) Tailoring gear requirements to provide for smaller or lighter gear for smaller vessels.
- 5) Avoiding management measures that require hazardous at-sea inspections or enforcement if other comparable enforcement could be accomplished as effectively.
- 6) Limiting the number of participants in the fishery.
- 7) Spreading effort over time and area to avoid potential gear and/or vessel conflicts.
- 8) Implementing management measures that reduce the race for fish and the resulting incentives for fishermen to take additional risks with respect to vessel safety.

The Council consults with the USCG on safety-at-sea considerations through a non-voting USCG seat on the Council and through the Council’s enforcement advisory body, the Enforcement Consultants. The Council also has considered safety-at-sea factors when deciding groundfish management measures. For example, the sablefish fishery for limited entry fixed gear permit holders with a sablefish endorsement fish their tier limits any time during the April to October primary season, which allows fishermen to fish when weather conditions are amenable to fishing safely. Likewise, the rationalized trawl fishery, managed using a system of IFQs and harvest cooperatives, has reduced the propensity to race for fish in that fishery and enhanced safety-at-sea. In general, most of the groundfish fishery has also limited participation through a limited entry system.

The National Institute for Occupational Safety and Health (NIOSH) is the federal government agency responsible for conducting research and making recommendations for the prevention of work-related injury and illness. NIOSH recently completed an in-depth study of commercial fishing fatalities in the United States during 2000-2009. The purpose of the study was to identify the most hazardous fisheries around the country and to describe the unique safety issues in each. NIOSH published a report on fatal occupational injuries in the west coast commercial fishing industry available at [http://www.cdc.gov/niosh/docs/2011-104/pdfs/WC CFID Summary EV.pdf](http://www.cdc.gov/niosh/docs/2011-104/pdfs/WC_CFID_Summary_EV.pdf). During 2000-2009, 86 commercial fishing deaths occurred off the US. west coast, an average of 9 per year. Almost 70% of the deaths were caused by drowning following a vessel disaster (e.g., sinking, capsizing, fire, etc.) in which the crew was forced to abandon ship. During two years (2001 and 2006) vessel disasters were the sole cause of commercial fishing fatalities. About one-quarter (24%) of fatalities were the result of falls overboard. The remaining fatalities were due to traumatic injuries sustained on-board, while diving or on-shore. The NIOSH report identified the highest fatality rate in the Dungeness crab fishery followed by the Columbia River Tribal salmon fishery.

Vessel disasters often result in multiple fatalities. The 58 deaths due to vessel disasters during 2000-2009 took place in 32 separate incidents. Vessel disasters were usually caused by a sequence of events, starting with an initiating event. The most common initiating events were: flooding, being struck by a large wave, and crossing a river bar during hazardous conditions. In addition, severe weather conditions contributed to 78% of vessel disasters. During 2000-2009, 21 Dungeness crab fishermen died in 10 separate vessel disasters. There were also 16 other vessel disasters in which all the fishermen survived. Crossing a bar in hazardous conditions led to 40% of fatal vessel disasters. None of the non-fatal vessel disasters involved crossing a bar. Vessel instability led to both fatal and non-fatal disasters, but was slightly more likely to be involved in fatal disasters. Several initiating events only resulted in non-fatal vessel disasters, such as flooding and striking rocks.

Falls overboard accounted for 24% of all fatalities in the west coast commercial fishing industry during 2000-2009. Falls overboard were caused most often by tripping or slipping on deck and by entanglement in fishing gear. Factors that contributed to falls overboard were: working alone on deck (52%), using alcohol or drugs (19%), and poor weather conditions (14%). None of the victims of falls overboard were wearing a Personal Flotation Device (PFD).

The NIOSH report recommended the following to prevent or mitigate injuries and fatalities from vessel disasters, falls overboard, or on-board injuries:

### **Vessel Disasters**

- Take a marine safety class at least once every 5 years - Safety training for fishermen is available, affordable, and saves lives. All fishermen should learn and know how to use basic lifesaving equipment like immersion suits, life rafts, EPIRBs, and fire extinguishers.
- Do monthly drills: Abandon ship, Flooding, Fire - Safety training equips fishermen with survival skills and knowledge. Monthly drills give fishermen an opportunity to practice and re-enforce those skills.
- Test immersion suit for leaks - When watertight, immersion suits provide thermal protection and flotation in cold water. If an immersion suit has leaks, it will provide less protection from cold water. Instructions for inflation testing immersion suits are available at [www.amsea.org](http://www.amsea.org).
- Heed weather forecasts and avoid fishing in severe sea conditions - Hazardous weather conditions contributed to nearly 80% of vessel disasters off the west coast during 2000-2009, and the deaths of 52 fishermen. Make the decision to stay in port when the seas are too rough for your vessel to operate in. Keep track of forecasts and seek shelter before the storm arrives or intensifies beyond the safe operating limits of your vessel.
- Maintain watertight integrity - Flooding is the most common initiating event for vessel disasters on the west coast. Inspect and maintain the hull of your vessel and all through-hull fittings. When seas are rough, ensure that watertight doors and hatches are sealed. Inspect and test high water alarms regularly.

### **Falls Overboard**

- Wear a PFD on deck - Falls overboard occur without warning or time to prepare. A PFD stowed away onboard will not help float a fisherman who has fallen overboard. Wearing a PFD on deck is the single most important thing a fisherman can do to increase survivability following a fall overboard. There are many new styles of PFDs which have been evaluated by fishermen in real working conditions and are comfortable to work in on deck. Results of the NIOSH PFD study are available at [www.cdc.gov/niosh/topics/fishing](http://www.cdc.gov/niosh/topics/fishing).



- Utilize a man overboard alarm system - Man overboard alarms are devices which alert others instantly to a fall overboard emergency, even if the fall was not witnessed. Systems vary in features and cost, but even the most inexpensive and basic system can save lives by immediately sounding an alarm if a fisherman falls overboard. Some of these systems can also benefit fishermen who work alone on small vessels by shutting down the engine if the sole operator falls overboard. This gives the fisherman, especially one prepared by wearing a PFD, a chance to get back to the vessel and re-board it.
- Conduct monthly man-overboard drills - If you fell overboard, would you want it to be the first time your crewmates tried to recover a man-overboard? Practicing man-overboard recovery procedures is essential for a crew to perform well in an actual emergency.
- Install bulwarks or rails at a minimum height of 36 to 39 inches.

### **On-Board Injuries**

- Install emergency stop (e-stop) devices on deck machinery - Deck machinery, especially deck winches, are particularly hazardous and result in many fatal and non-fatal injuries. Emergency-stop buttons have been developed specifically for deck machinery on fishing vessels and can be adapted and retrofitted onto any winch or other machinery. More information about e-stops for fishing vessels can be found at [www.cdc.gov/niosh/topics/fishing](http://www.cdc.gov/niosh/topics/fishing).
- The Coast Guard recommends vessels carry all required safety and survival equipment as required for the vessels operating parameters. You can generate a checklist for a particular vessel by using the “Checklist Generator” at <http://www.uscg.mil/d13/cfvs/default.asp>. Then have a Coast Guard dockside examiner on board to inspect the condition of each of the required items.

## LITERATURE CITED

- Adams, P. B. 1986. Status of lingcod (*Ophiodon elongatus*) stocks off the coast of Washington, Oregon and California. Pages 60 in Status of Pacific coast groundfish fishery in 1986 and recommended acceptable biological catches for 1987. Pacific Fisheries Management Council, Portland, OR.
- Adams, P. B. 1987. The diet of widow rockfish *Sebastes entomelas* in northern California. W. H. Lenarz and D. R. Gunderson, editors. Widow rockfish, proceedings of a workshop. NOAA Tech. Rep. NMFS 48, Tiburon, CA.
- Adams, P. B. and J. E. Hardwick. 1992. Lingcod. Pages 161-164 in L. W.S, C. M. Dewees, and C. W. Haugen, editors. California's Living Marine Resources and Their Utilization. California Sea Grant Program, Davis, CA.
- Adams, P. B., E. H. Williams, K. R. Silberberg, and T. E. Laidig. 1999. Southern lingcod stock assessment in 1999. in Appendix to Status of the Pacific Coast groundfish fishery through 1999 and recommended acceptable biological catches for 2000 (SAFE Report). Pacific Fishery Management Council, Portland.
- Agostini, V. N., R. C. Francis, A. B. Hollowed, S. D. Pierce, C. Wilson, and A. N. Hendrix. 2006. The relationship between Pacific hake (*Merluccius productus*) distribution and poleward subsurface flow in the California Current system. Canadian Journal of Fisheries and Aquatic Sciences 63:2648-2659.
- Ainley, D. G., W. H. Sydeman, R. H. Parrish, and W. H. Lenarz. 1993. Oceanic factors influencing distribution of young rockfish (*Sebastes*) in Central California: a predator's perspective. CalCOFI Reports 34(133-139).
- Albin, D. and K. A. Karpov. 1996. Mortality of lingcod, *Ophiodon elongatus*, related to capture by hook and line. Marine Fisheries Review 60(3):29-34.
- Alderdice, D. F. and C. R. Forrester. 1971. Effects of salinity and temperature on embryonic development of the petrale sole (*Eopsetta jordani*). J.Fish.Res.Board Canada 28:727-744.
- Alheit, J. and T. J. Pitcher. 1995. Hake: biology, fisheries, and markets. Chapman and Hall, London.
- Allen, L. G., D. J. Pondella II, and M. H. Hom. 2006. The ecology of marine fishes: California and adjacent waters. University of California Press, Los Angeles, CA.
- Allen, M. J. 1982. Functional structure of soft-bottom fish communities of the southern California shelf. Ph.D Dissertation. University of California.
- Allen, M. J. and G. B. Smith. 1988. Atlas and zoogeography of common fishes in the Bering Sea and northeastern Pacific.
- Alonzo, S. H., T. Ish, M. Key, A. D. MacCall, and M. Mangel. 2008. The importance of incorporating protogynous sex change into stock assessments. Bulletin of Marine Science 83:163-179.
- Alverson, D. L. and B. M. Chatwin. 1957. Results from tagging experiments on a spawning stock of petrale sole, *Eopsetta jordani* (Lockington). J.Fish.Res.Board Canada 14:953-974.

- Alverson, D. L., A. T. Pruter, and L. L. Ronholt. 1964. A study of demersal fishes and fisheries of the Northeastern Pacific Ocean. Institute of Fisheries, University of British Columbia, Vancouver, British Columbia.
- Anderson, T. J., C. Syms, D. A. Roberts, and D. F. Howard. 2009. Multi-scale fish-habitat associations and the use of habitat surrogates to predict the organization and abundance of deep-water fish assemblages. *Journal of Experimental Marine Biology and Ecology* 379:34-42.
- Antonelis, G. A. and C. H. Fiscus. 1980. The pinnipeds of the California Current. *CalCOFI Reports* 21:68-78.
- Archibald, C., P. D. Fournier, and B. M. Leaman. 1983. Reconstruct of stock history and development of rehabilitation strategies for Pacific ocean perch in Queen Charlotte Sound, Canada. *N.Amer.J.Fish.Mgmt.* 3:283-294.
- Arora, H. L. 1951. An investigation of the California sand dab, (*Citharichthys sordidus*) (Girard). *California Fish and Game* 37:3-42.
- Bailey, K. M. 1982. The early life history of the Pacific hake, *Merluccius productus* *Fish.Bull.* 80:589-598.
- Bailey, K. M., R. C. Francis, and P. R. Stevens. 1982. The life history and fishery of Pacific whiting, *Merluccius productus* *Calif.Coop.Oceanic Fish.Invest.Rep.* 23:81-98.
- Bargmann, G. G. 2009. A history of the fisheries for spiny dogfish along the Pacific Coast from California to Washington. *in* V. F. Gallucci, G. McFarlane, and G. G. Bargmann, editors. *Biology and Management of Dogfish Sharks*. American Fisheries Society.
- Barss, W. H. 1976. The Pacific sanddab. Oregon Department of Fish and Wildlife Informational Report 76-5.
- Baxter, R. 1999. Pleuronectiformes. Pages 369-415 *in* E. Interagency Ecological Program for the Sacramento-San Joaquin, editor. Report on the 1980-1995 fish, shrimp, and crab sampling in the San Francisco Estuary, California, volume Technical Report 63. California Department of Fish and Game, Stockton, CA.
- Beamish, R. J. 1979. New information on the longevity of Pacific ocean perch (*Sebastes alutus*). *J.Fish.Res.Board Canada* 36:1395-1400.
- Beamish, R. J. and G. A. McFarlane. 1988. Resident and dispersal behavior of adult sablefish (*Anoplopoma fimbria*) in the slope waters off Canada's West Coast. *Can.J.Fish.Aquat.Sci.* 45:152-164.
- Beamish, R. J., G. A. McFarlane, and A. Benson. 2006. Longevity overfishing. *Progress in Oceanography* 68:289-302.
- Beamish, R. J., B. L. Thomson, and G. A. McFarlane. 1992. Spiny dogfish predation on chinook and coho salmon and the potential effects on hatchery-produced salmon. *Trans.Am.Fish.Soc.* 121(4):444-455.
- Becker, D. S. 1984. Resource partitioning by small-mouthed pleuronectids in Puget Sound, Washington. Ph.D Dissertation. University of Washington.

- Bence, J. R. and J. E. Hightower. 1990. Status of bocaccio in the Conception/Monterey/Eureka INPFC areas in 1990. *in* Appendix to Status of the Pacific Coast groundfish fishery through 1990 and recommended acceptable biological catches for 1991 (SAFE Report). Pacific Fishery Management Council, Portland.
- Benet, D. L., E. J. Dick, and D. E. Pearson. 2009. Life history aspects of greenspotted rockfish (*Sebastes chlorostictus*) from central California. NOAA Technical Memorandum NMFS NOAA-TM-NMFS-SWFSC-466:43 pgs.
- Bennett, J. T., G. W. Boehlert, and K. K. Turekian. 1982. Confirmation of longevity in *Sebastes diploproa* (Pisces: Scorpaenidae) from  $^{210}\text{Pb}/^{226}\text{Ra}$  measurements in otoliths. *Marine Biology* 71:209-215.
- Bennett, W. A., K. Roinestad, L. Rogers-Bennett, L. Kaufman, D. Wilson-Vandenberg, and B. Heneman. 2004. Inverse regional responses to climate change and fishing intensity by the recreational rockfish (*Sebastes* spp.) fishery in California. *Canadian Journal of Fisheries and Aquatic Sciences* 61:2499-2510.
- Berkeley, S. A. 2004. Maternal age as a determinant of larval growth and survival in a marine fish, *Sebastes melanops* *Ecology* 85(5):1258-1264.
- Berkeley, S. A., M. A. Hixon, R. J. Larson, and M. S. Love. 2004. Fisheries sustainability via protection of age structure and spatial distribution of fish populations. *Fisheries* 29(8):23-32.
- Best, E. A. 1959. Status of the animal food fishery in northern California, 1956 and 1957. *California Fish and Game* 45(1):5-18.
- Best, E. A. 1960. Petrale Sole. Pages 79 *in* California ocean fisheries resources to the year 1960. California Department of Fish and Game.
- Best, E. A. 1961. The California animal food fishery, 1958-1960. *Bulletin of the Pacific Marine Fisheries Commission* 5:5-15.
- Best, E. A. 1963. Movements of petrale sole, *Eopsetta jordani*, tagged off of California. *Pacific Marine Fisheries Commission Bulletin* 6:24-38.
- Best, E. A. 1964. Spawning of longspine channel rockfish, *Sebastes altivelis*. *California Fish and Game* 50:265-267.
- Beverton, R. J. H. 1992. Patterns of reproductive strategy parameters in some marine teleost fishes. *Journal of Fish Biology* 42(Supplement B):137-160.
- Birtwell, I. K., M. Wood, and D. K. Gordon. 1984. Fish diets and benthic invertebrates in the estuary of the Somass River, Port Alberni, British Columbia. *Canadian Manuscript Report of Fisheries and Aquatic Sciences* 1799:49.
- Black, B. A. 2009. Climate-driven synchrony across tree, bivalve, and rockfish growth-increment chronologies of the northeast Pacific. *Marine Ecology Progress Series* 378:37-46.
- Black, B. A., G. W. Boehlert, and M. M. Yoklavich. 2008. Using tree-ring crossdating techniques to validate annual growth increments in long-lived fishes. *Canadian Journal of Fisheries and Aquatic Sciences* 62:2277-2284.

- Bobko, S. J. and S. A. Berkeley. 2004. Maturity, ovarian cycle, fecundity, and age-specific parturition of black rockfish, *Sebastes melanops*. Fish.Bull. 102(3):418-429.
- Boehlert, G. W. 1980. Size composition, age composition, and growth of canary rockfish, *Sebastes pinniger*, and splitnose rockfish, *S. diploproa* from the 1977 rockfish survey. Mar.Fish.Rev. 42:57-63.
- Boehlert, G. W. and M. Y. Yoklavich. 1985. Larval and juvenile growth of sablefish *Anoplopoma fimbria* as determined from otolith increments. Fish.Bull. 83:475-481.
- Bograd, S. J., C. G. Castro, E. D. Lorenzo, D. M. Palacios, H. Bailey, W. Gilly, and coauthors. 2008. Oxygen declines and the shoaling of the hypoxic boundary in the California Current. Geophysical Research Letters 35:L12607.
- Bond, C. E. 1979. Biology of fishes. Saunders College Publishing, Philadelphia.
- Bonfil, R. 1994. Overview of world elasmobranch fisheries. FAO Fisheries Technical Paper No. 341.
- Bradburn, M. J., A. Keller, and B. H. Horness. 2011. The 2003 to 2008 U.S. West Coast bottom trawl surveys of groundfish resources off Washington, Oregon, and California: Estimates of distribution, abundance, length, and age composition. NOAA Technical Memorandum NMFS-NWFSC-114:323 pgs.
- Brodeur, R. D., I. A. Fleming, J. M. Bennett, and M. A. Campbell. 2009. Summer distribution and feeding of spiny dogfish off the Washington and Oregon coasts. *in* V. F. Gallucci, G. A. McFarlane, and G. G. Bargmann, editors. Biology and Management of Dogfish Sharks. American Fisheries Society.
- Brodeur, R. D. and W. G. Percy. 1984. Food habits and dietary overlap of some shelf rockfishes (genus *Sebastes*) from the northeastern Pacific Ocean. Fishery Bulletin 82:262-294.
- Brodziak, J., L. Jacobson, R. Lauth, and M. Wilkins. 1997. Assessment of the Dover sole stock for 1997. *in* Appendix to Status of the Pacific Coast groundfish fishery through 1997 and recommended biological catches for 1998 (Stock assessment and fishery evaluation). Pacific Fishery Management Council, Portland, OR.
- Buckley, T. W., G. E. Tyler, D. M. Smith, and P. A. Livingston (O. National and A. Atmospheric). 1999. Food habits of some commercially important groundfish off the coasts of California, Oregon, Washington, and British Columbia.
- Buonaccoursi, V. P., C. A. Kimbrel, E. A. Lynn, and R. D. Vetter. 2005. Limited realized dispersal and introgressive hybridization influence genetic structure and conservation strategies for brown rockfish, *Sebastes auriculatus*. Conservation Genetics 6:697-713.
- Burford, M. O., M. H. Carr, and G. Bernardi. 2011. Age-structured genetic analysis reveals temporal and geographic variation within and between two cryptic rockfish species. Marine Ecology Progress Series 442:201-215.
- Butler, J. L., T. Barnes, P. Crone, and R. Conser. 2003. Cowcod rebuilding review. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches

- for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Butler, J. L., L. D. Jacobson, and J. T. Barnes. 1999a. Stock assessment for blackgill rockfish. *in* Appendix to the status of the Pacific coast groundfish fishery through 1998 and recommended acceptable biological catches for 1999: stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Butler, J. L., L. D. Jacobson, J. T. Barnes, H. G. Moser, and R. Collins. 1999b. Stock assessment of cowcod. *in* Appendix to Status of the Pacific Coast groundfish fishery through 1998 and recommended acceptable biological catches for 1999 (SAFE Report). Pacific Fishery Management Council, Portland, OR.
- Cailliet, G. M., E. K. Osada, and M. Moser. 1988. Ecological studies of sablefish in Monterey Bay. *Calif.Dept.Fish and Game* 74:133-153.
- Campana, S. E., W. Joyce, and D. W. Kulka. 2009. Growth and reproduction of spiny dogfish off the Eastern Coast of Canada, including inferences on stock structure. *in* V. F. Gallucci, G. A. McFarlane, and G. G. Bargmann, editors. *Biology and Management of Dogfish Sharks*. American Fisheries Society.
- Carlson, H. R. and R. E. Haight. 1972. Evidence for a home site and homing of adult yellowtail rockfish, *Sebastes flavidus* *J.Fish.Res.Bd.Canada* 29:1011-1014.
- Casillas, E., L. Crockett, Y. deReynier, J. Glock, M. Helvey, B. Meyer, and coauthors. 1998. Essential Fish Habitat, West Coast Groundfish. *in* Appendix to Amendment 11 of the Pacific Coast Groundfish Plan, Fishery Management Plan Environmental Impact Statement for the California, Oregon Washington Groundfish Fishery. National Marine Fisheries Service, Seattle.
- Castillo, G. C. 1992. Fluctuations of year-class strength of petrale sole (*Eopsetta jordani*) and their relation to environmental factors. Oregon State University, Corvallis, OR.
- Castillo, G. C. 1995. Latitudinal patterns in reproductive life history traits of Northeast Pacific flatfish. Pages 51-72 *in* International Symposium on North Pacific Flatfish. University of Alaska Sea Grant College Program, Fairbanks, AK, 1995.
- Castillo, G. C., H. W. Li, and J. T. Golden. 1993. Environmental induced recruitment variation in petrale sole, *Eopsetta jordani*. *Fisheries Bulletin* 92:481-493.
- CDFG. 2001. Sanddabs. Pages 201-202 *in* California's Marine Living Resources: A Status Report. California Department of Fish and Game.
- Chamberlain, D. W. 1979. Histology of the reproductive systems and comparison of selected morphological characters in four Eastern Pacific species of *Citharichys* (*Pisces: Bothidae*). Ph.D. University of Southern California, Los Angeles, CA.
- Chess, J. R., S. E. Smith, and P. C. Fischer. 1988. Trophic relationships of the shortbelly rockfish, *Sebastes jordani*, off central California. *CalCOFI Reports* 29(129-136).
- Childress, J. J. and B. A. Seibel. 1998. Life at stable low oxygen levels: Adaptations of animals to oceanic oxygen minimum layers. *Journal of Experimental Biology* 201:1223-1232.

- Clark, W. G. 1991. Groundfish exploitation rates based on life history parameters. *Canadian Journal of Fisheries and Aquatic Sciences* 48:734-750.
- Clark, W. G. 2002. F35% revisited ten years later. *North American Journal of Fisheries Management* 22:251-257.
- Coleman, B. A. 1986. The 1980 Pacific West Coast bottom trawl survey of groundfish resources: estimates of distribution, abundance, length and age composition.
- Constable, H. 2006. Population genetics of juvenile *Sebastes jordani*, shortbelly rockfish, along the California coast. San Francisco State University, San Francisco, CA.
- Cooper, D. W., K. E. Pearson, and D. R. Gunderson. 2005. Fecundity of shortspine thornyhead (*Sebastolobus alascanus*) and longspine thornyhead (*S. altivelis*) (Scorpaenidae) from the northeastern Pacific Ocean, determined by sterological and gravimetric techniques. *Fishery Bulletin* 103:15-22.
- Cope, J., E. J. Dick, A. MacCall, M. Monk, B. Soper, and C. Wetzel. 2014. Data-moderate stock assessments for brown, China, copper, sharpchin, stripetail, and yellowtail rockfishes and English and rex soles in 2013. Pacific Fishery Management Council, Portland, OR.
- Cope, J. M. 2004. Population genetics and phylogeography of the blue rockfish (*Sebastes mystinus*) from Washington to California. *Canadian Journal of Fisheries and Aquatic Sciences* 61:332-342.
- Cope, J. M. 2013. Implementing a statistical catch-at-age model (Stock Synthesis) as a tool for deriving overfishing limits in data-limited situations. *Fish. Res.* 142:3-14.
- Cope, J. M., J. DeVore, E. J. Dick, K. Ames, J. Budrick, D. L. Erickson, and coauthors. 2011. An approach to defining stock complexes for U.S. West Coast groundfishes using vulnerabilities and ecological distributions. *N.Amer.J.Fish.Mgmt.* 31(4):589-604.
- Cope, J. M. and M. A. Haltuch. 2012. Temporal and spatial summer groundfish assemblages in trawlable habitat off the west coast of the USA, 1977 to 2009. *Marine Ecology Progress Series* 451:187-200.
- Cope, J. M. and M. Key. 2009. Status of Cabezon ( *Scorpaenichthys marmoratus* ) in California and Oregon waters as assessed in 2009. Pacific Fishery Management Council, Portland, OR.
- Cope, J. M., K. Piner, C. V. Minte-Vera, and A. E. Punt. 2004. Status and future prospects for the cabezon (*Scorpaenichthys marmoratus*) as assessed in 2003. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2004 and recommended acceptable biological catches for 2005-2006 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Cope, J. M. and A. Punt. 2006. Status of Cabezon ( *Scorpaenichthys marmoratus* ) in California Waters as Assessed in 2005. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Cope, J. M. and A. E. Punt. 2009. Drawing the lines: resolving fishery management units with simple fisheries data. *Canadian Journal of Fisheries and Aquatic Sciences* 66(8):1256-1273.

- Cope, J. M., J. T. Thorson, J. R. Wetzel, and J. DeVore. In Press. Evaluating a prior on relative stock status using simplified age-structured models. Fisheries Research.
- Cortes, E. 2002. Incorporating uncertainty into demographic modeling: Application to shark populations and their conservation. Conservation Biology 16:1048-1062.
- Crone, P. R., R. D. Methot, R. J. Conser, and T. L. Builder. 1999. Status of the canary rockfish resource off Oregon and Washington in 1999. *in* Status of the Pacific Coast groundfish fishery through 1998 and recommended acceptable biological catches for 1999 (SAFE Report). Pacific Fishery Management Council, Portland, OR.
- Culver, B. N. 1986. Results of tagging black rockfish (*Sebastes melanops*) off the Washington and northern Oregon coast. Alaska Sea Grant College Program, Anchorage, Alaska, 1986.
- Dark, T. A. and M. E. Wilkins. 1994. Distribution, abundance, and biological characteristics of groundfish off the coast of Washington, Oregon and California, 1977-1986. NOAA.
- Davis, M. W., B. L. Olla, and C. B. Schreck. 2001. Stress induced by hooking, net towing, elevated sea water temperature and air in sablefish: lack of concordance between mortality and physiological measures of stress. Journal of Fish Biology 58:1-15.
- Demory, R. L. 1971. Depth distribution of some small flatfishes off the northern Oregon-Washington coast. Research Report, Fish Commission of Oregon 3:44-48.
- Demory, R. L. 1984. Appendix 12: Progress report on the status of petrale sole in the INPFC Columbia-Vancouver areas in 1984. *in* Status of the Pacific coast groundfish fishery and recommendations for management in 1985. Pacific Fishery Management Council, Portland, OR.
- Di Giacomo, E. E., M. R. Perier, and M. Coller. 2009. Reproduction of spiny dogfish in San Matias Gulf, Patagonia. *in* V. F. Gallucci, G. A. McFarlane, and G. G. Bargmann, editors. Biology and Management of Dogfish Sharks. American Fisheries Society.
- Dick, E. J. and A. D. MacCall. 2011. Depletion-based stock reduction analysis: a catch-based method for determining sustainable yields for data-poor fish stocks. Fisheries Research 110:331-341.
- Dick, E. J. and A. D. MacCall. 2013. Status and productivity of cowcod, *Sebastes levis*, in the Southern California Bight, 2013. Pacific Fishery Management Council, Portland, OR.
- Dick, E. J. and A. D. MacCall. 2014. Cowcod Rebuilding Analysis. Pacific Fishery Management Council, Portland, OR.
- Dick, E. J., D. Pearson, and S. Ralston. 2011. Status of greenspotted rockfish, *Sebastes chlorostictus*, in U.S. waters off California. Pacific Fishery Management Council, Portland, OR.
- Dick, E. J. and S. Ralston. 2007. Cowcod rebuilding analysis. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation.
- Dick, E. J., S. Ralston, and D. Pearson. 2007. Status of cowcod, *Sebastes levis*, in the southern California Bight. Pacific Fishery Management Council, Portland, OR.



- Dick, E. J., S. Ralston, D. Pearson, and J. Wiedenmann. 2009. Updated status of cowcod, *Sebastes levis*, in the southern Californian night bight. Portland, OR.
- Donohoe, C. J. 2000. Metamorphosis, growth, and settlement of Pacific sanddab (*Citharichthys sordidus*) to a continental shelf nursery, inferred from otolith microstructure. Ph.D. Oregon State University, Corvallis, OR.
- Dorn, M. 1995. Effects of age composition and oceanographic conditions on the annual migration of Pacific whiting, *Merluccius productus* Calif.Coop.Oceanic Fish.Invest.Rep. 36:97-105.
- Dorn, M., S. A. Berkeley, R. J. Conser, O. S. Hamel, R. Mohn, K. Piner, and coauthors. 2006. Petrale Sole STAR Panel Meeting Report [September 2005]. in Volume 2: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Dorn, M. W. 2002a. Advice on west coast rockfish harvest rates from Bayesian meta-analysis of stock-recruit relationships. Pages 22:290-300 in. N. Am J. fish. Manage.
- Dorn, M. W. 2002b. Advice on west coast rockfish harvest rates from Bayesian meta-analysis of stock-recruit relationships. N. Am J. Fish. Manage. 22:290-300.
- Dorn, M. W. and R. D. Methot. 1991. Status of the coastal Pacific whiting resource in 1990. NOAA Tech.Memo NMFS F/AFSC-47.
- Dorn, M. W. and R. D. Methot. 1992. Status of the coastal Pacific whiting resource in 1992. in. Pacific Fishery Management Council, Portland, Oregon.
- Douglas, D. A. 1998. Species composition of rockfish in catches by Oregon trawlers, 1963-93.
- Dunn, J. R. and C. R. Hitz. 1969. Oceanic occurrence of black rockfish (*Sebastes melanops*) in the central north Pacific. J.Fish.Res.Bd.Canada 26:3094-3097.
- Dunn, J. R. and A. C. Matarese. 1987. A review of early life history of northeast Pacific gadoid fishes. Fish. Res. 5:163-184.
- Ebert, D. A. 2003. Sharks, Rays and Chimaeras of California. University of California Press, Berkeley, CA.
- Ebert, D. A. and L. J. V. Compagno. 2007. Biodiversity and systematics of skates (Chondrichthyes: Rajiformes: Rajoidei). Environ.Biol.Fish. 80(2-3):111-124.
- Ebert, D. D., W. T. White, K. J. Goldman, L. J. V. Compagno, T. S. Daly-Engel, and R. D. Ward. 2010. Resurrection and redescription of *Squalus suckleyi* (Girard, 1854) from the North Pacific, with comments on the *Squalus acanthias* subgroup (Squaliformes: Squalidae). Zootaxa 2612:22-40.
- Echeverria, T. W. 1987. Thirty-four species of California rockfishes: maturity and seasonality of reproduction. Fish.Bull., U.S. 85(2):229-250.
- Emmett, R. L., S. L. Stone, S. A. Hinton, and M. E. Monaco. 1991. Distribution and abundance of fishes and invertebrates in West Coast estuaries, Volume II: Species life history summaries. NOAA/NOS Strategic Environmental Assessments Division, Rockville, Maryland.

- Erickson, D. L. and E. K. Pikitch. 1993. A histological description of shortspine thornyhead, *Sebastes alascanus*, ovaries: Structures associated with the production of gelatinous egg masses. *Environ.Biol.Fish.* 36:273-282.
- Eschmeyer, W. N., E. S. Herald, and H. Hammon. 1983. *A Field Guide to Pacific Coast Fishes of North America*. Houghton Mifflin, Boston.
- Fargo, J. and P. J. Starr. 2001. Turbot stock assessment for 2001 and recommendation for management in 2002. Canadian Science Advisory Secretariat.
- Fay, G. 2006. Stock Assessment and Status of Longspine Thornyhead ( *Sebastes altivelis* ) off California, Oregon and Washington in 2005. *in* Volume 4: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Field, J. C. 2008. Status of the Chilipepper rockfish, *Sebastes goodei*, in 2007. Pacific Fishery Management Council, Portland, Oregon.
- Field, J. C. 2011a. Rebuilding analysis for bocaccio in 2011. Pacific Fishery Management Council, Portland, OR.
- Field, J. C. 2011b. Status of bocaccio, *Sebastes paucispinis*, in the Conception, Monterey and Eureka INPFC areas as evaluated for 2011. Pacific Fishery Management Council, Portland, OR.
- Field, J. C. 2013. Status of bocaccio, *Sebastes paucispinis*, in the Conception, Monterey and Eureka INPFC areas as evaluated for 2013. Pacific Fishery Management Council, Portland, OR.
- Field, J. C., K. Baltz, A. J. Phillips, and W. A. Walker. 2007. Range expansion and trophic interactions of the jumbo squid, *Dosidicus gigas*, in the California Current. *CalCOFI Reports* 48:131-146.
- Field, J. C., E. J. Dick, and A. D. MacCall. 2008. Stock Assessment Model for the Shortbelly Rockfish, *Sebastes Jordani*, in the California Current. Pacific Fishery Management Council, Portland, Oregon.
- Field, J. C., E. J. Dick, D. Pearson, and A. D. MacCall. 2009. Status of bocaccio, *Sebastes paucispinis*, in the Conception, Monterey and Eureka INPFC areas for 2009. Pacific Fishery Management Council, Portland, OR.
- Field, J. C., R. C. Francis, and K. Aydin. 2006. Top-down modeling and bottom-up dynamics: linking a fisheries-based ecosystem model with climate hypotheses in the Northern California Current. *Progress in Oceanography* 68:238-270.
- Field, J. C. and X. He. 2009. Bocaccio rebuilding analysis for 2009. Pacific Fishery Management Council, Portland, OR.
- Field, J. C. and D. Pearson. 2011. Status of the blackgill rockfish, *Sebastes melanostomus*, in the Conception and Monterey INPFC areas for 2011. Pacific Fishery Management Council, Portland, OR.
- Fiscus, C. H. 1979. Interactions of marine mammals and Pacific hake. *Mar.Fish.Rev.* 41:1-9.

- Ford, R. F. 1965. Distribution, population dynamics and behavior of a Bothid flatfish, *Citharichthys stigmaeus*. Ph.D. University of California San Diego.
- Forrest, R. E., M. K. McAllister, M. W. Dorn, S. J. D. Martell, and R. D. Stanley. 2010. Hierarchical Bayesian estimation of recruitment parameters and reference points for Pacific rockfishes (*Sebastes* spp.) under alternative assumptions about the stock-recruit function. *Can.J.Fish.Aquat.Sci.* 67:1611-1634.
- Forrester, C. R. 1969. Life history information on some groundfish species.
- Fraidenburg, M. E. 1980. Yellowtail rockfish, *Sebastes flavidus*, length and age composition off California, Oregon, and Washington in 1977. *Mar.Fish.Rev.* 42:54-56.
- Francis, R. C. 1986. Two fisheries biology problems in West coast groundfish management. *North American Journal of Fisheries Management* 6:453-462.
- Frey, H. W. 1971. California's living marine resources and their utilization. Calif. Dept. Fish Game.
- Frisk, M. G., T. J. Miller, and M. J. Fogarty. 2002. The population dynamics of little skate *Leucoraja erinacea*, winter skate *Leucoraja ocellata*, and barndoor skate *Dipturus leavis*: predicting exploitation limits using matrix analysis. *ICES J.of Mar.Sci.* 59:576-586.
- Gabriel, W. L. and W. G. Pearcy. 1981. Feeding selectivity of Dover sole, *Microstomus pacificus* *Fish.Bull.* 79:749-763.
- Garrison, K. J. and B. S. Miller. 1982. Review of the early life history of Puget Sound fishes. University of Washington Fish. Res. Inst., Seattle, Washington.
- Genin, A. 2004. Bio-physical coupling in the formation of zooplankton and fish aggregations over abrupt topographies. *Journal of Marine Systems* 50:3-20.
- Genin, A., L. Hauray, and P. Greenblatt. 1988. Interactions of migrating zooplankton with shallow topography: predation by rockfishes and intensification of patchiness. *Deep Sea Research* 35:151-175.
- Gertseva, V., J. M. Cope, and D. E. Pearson. 2009. Status of the U.S. splitnose rockfish (*Sebastes diploproa*) resource in 2009. Pacific Fishery Management Council, Portland, OR.
- Gertseva, V. and I. G. Taylor. 2011. Status of the spiny dogfish shark resource off the continental U.S. Pacific Coast in 2011. Pacific Fishery Management Council, Portland, OR.
- Gertseva, V. V. and M. J. Schirripa. 2008. Status of the Longnose Skate (*Raja rhina*) off the continental US Pacific Coast in 2007. Pacific Fishery Management Council, Portland, Oregon.
- Gertseva, V. V. and J. T. Thorson. 2013. Status of the darkblotched rockfish resource off the continental U.S. Pacific Coast in 2013. Pacific Fishery Management Council, Portland, OR.
- Gharrett, A. J., A. P. Matala, E. L. Peterson, A. K. Gray, and Z. Li. 2005. Two genetically distinct forms of rougheyeye rockfish are different species. *Transactions of the American Fisheries Society* 134:242-260.

- Giorgi, A. E. and J. L. Congleton. 1984. Effects of current velocity on the development and survival of lingcod, *Ophiodon elongatus*, embryos. *Env.Bio.Fish.* 10:15-27.
- Golden, J. T. and R. L. Demory. 1984. A progress report on the status of canary rockfish (*Sebastes pinniger*) in the INPFC Vancouver, Columbia and Eureka areas in 1984, Appendix 6. *in* Status of the Pacific coast groundfish fishery through 1990 and recommended acceptable biological catches for 1991: stock assessment and fishery evaluation. Pacific Fishery Management Council. Pacific Fishery Management Council, Portland, OR.
- Golden, J. T. and C. Wood. 1990. Status of canary rockfish in the INPFC Vancouver, Columbia and Eureka areas and recommended ABC for 1991. Pages Appendix H *in*, Portland, Oregon.
- Goode, G. B. 1884. The fishery and fisheries industries of the United States. Washington Government Printing Office.
- Gotshall, D. W. 1969. Stomach contents of Pacific hake and arrowtooth flounder from northern California. *Calif.Fish Game* 55(1):75-82.
- Gotshall, D. W. 1981. Pacific Coast Inshore Fishes. Sea Challengers and Western Marine Enterprises Publication, Los Osos California.
- Gotshall, D. W., J. G. Smith, and A. Holbert. 1965. Food of the blue rockfish *Sebastes mystinus*. California Department of Fish and Game 51(3):147-162.
- Gregory, P. A. and T. Jow. 1976. The validity of otoliths as indicators of age of petrale sole from California. California Department of Fish and Game 62(2):132-140.
- Gunderson, D. R. 1971. Reproductive patterns of Pacific ocean perch (*Sebastes alutus*) off Washington and British Columbia and their relation to bathymetric distribution and seasonal abundance. *J.Fish.Res.Board Canada* 28:417-425.
- Gunderson, D. R. 1977. Population biology of Pacific ocean perch, *Sebastes alutus*, stocks in the Washington-Queen Charlotte Sound region, and their response to fishing. *Fish.Bull.* 75(2):369-403.
- Gunderson, D. R. 1979. Results of cohort analysis for Pacific ocean perch stocks off British Columbia, Washington, and Oregon and an evaluation of alternative rebuilding strategies for these stocks. Unpublished report prepared for the Pacific Fishery Management Council, Portland OR.
- Gunderson, D. R. 1981. An updated cohort analysis for Pacific ocean perch stocks off Washington and Oregon. Unpublished report prepared for the Pacific Fishery Management Council, Portland, OR.
- Gunderson, D. R. 1984. The great widow rockfish hunt of 1980-82. *North American Journal of Fisheries Management* 4:465-468.
- Gunderson, D. R. and T. M. Sample. 1980. Distribution and abundance of rockfish off Washington, Oregon, and California during 1977. *U.S.Natl.Mar.Fish.Serv., Mar.Fish.Rev.* 42(3-4):2-16.
- Hagerman, F. B. 1952. The biology of the Dover sole, *Microstomus pacificus* (Lockington). *Calif.Dept.Fish and Game, Fish Bull.* 85:48 p.

- Haigh, R. and J. T. Schnute. 2003. The longspine thornyhead fishery along the west coast of Vancouver Island, British Columbia, Canada: portrait of a developing fishery. *N.Amer.J.Fish.Mgmt.* 23:120-140.
- Hallacher, L. E. and D. A. Roberts. 1985. Differential utilization of space and food by the inshore rockfishes (*Scorpaenidae: Sebastes*) of Carmel Bay, California. *Environ.Biol.Fish.* 12:91-110.
- Haltuch, M. 2011. 2011 petrale sole rebuilding analysis. NMFS, NWFSC, Seattle, WA, November 23, 2011.
- Haltuch, M. and A. Hicks. 2009a. 2009 petrale sole rebuilding analysis. Pacific Fishery Management Council, Portland, OR.
- Haltuch, M., A. Hicks, and K. See. 2011. Status of the U.S. petrale sole resource in 2010, Seattle, WA, November 23, 2011.
- Haltuch, M. A. and A. Hicks. 2009b. Status of the U.S. petrale sole resource in 2008. Pacific Fishery Management Council, Portland, Oregon.
- Haltuch, M. A., K. Ono, and J. Valero. 2013. Status of the U.S. petrale sole resource in 2012. Pacific Fishery Management Council, Portland, OR.
- Hamel, O. 2011. Pacific Ocean Perch Rebuilding Analysis 2011. NMFS, NWFSC, Seattle, WA, December 13, 2011.
- Hamel, O. and K. Ono. 2011. Stock Assessment of Pacific Ocean Perch in Waters off of the U.S. West Coast in 2011. NWFSC, Seattle, WA, September 20, 2011.
- Hamel, O. S. 2006a. Rebuilding update for Pacific ocean perch. *in* Volume 3: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Hamel, O. S. 2006b. Status and future prospects for the Pacific ocean perch resource in waters off Washington and Oregon as assessed in 2005. *in* Volume 3: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Hamel, O. S. 2006c. Status and Future Prospects for the Shortspine Thornyhead Resource in Waters off Washington, Oregon, and California as Assessed in 2005. *in* Volume 4: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Hamel, O. S. 2008a. Rebuilding update for darkblotched rockfish. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation.
- Hamel, O. S. 2008b. Rebuilding update for Pacific ocean perch. Pacific Fishery Management Council, Portland, OR.
- Hamel, O. S. 2008c. Status and future prospects for the darkblotched rockfish resource in waters off Washington, Oregon, and California as assessed in 2007. *in* Status of the Pacific Coast groundfish

- fishery through 2007, Stock assessment and fishery evaluation. Portland Fishery Management Council, Portland, OR.
- Hamel, O. S. 2008d. Status and future prospects for the Pacific ocean perch resource in waters off Washington and Oregon as assessed in 2007. Pacific Fishery Management Council, Portland, OR.
- Hamel, O. S. 2009a. Rebuilding update for Pacific ocean perch in 2009. Pacific Fishery Management Council, Portland, OR.
- Hamel, O. S. 2009b. Status and Future Prospects for the Pacific Ocean Perch Resource in Waters off Washington and Oregon as Assessed in 2009. Pacific Fishery Management Council, Portland, Oregon.
- Hamel, O. S., J. M. Cope, and S. Matson. 2013. Stock Assessment of Aurora Rockfish in 2013. Pacific Fishery Management Council, Portland, OR.
- Hamel, O. S., S. A. Sethi, and T. F. Wadsworth. 2009. Status and future prospects for Lingcod in waters off Washington, Oregon, and California as assessed in 2009. Pacific Fishery Management Council, Portland, OR.
- Hamel, O. S., I. J. Stewart, and A. E. Punt. 2003. Status and future prospects for the Pacific ocean perch resource in waters off Washington and Oregon as assessed in 2003. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Hamlett, W. C. and T. J. Koob. 1999. Female reproductive system. Pages 398-443 *in* W. C. Hamlett, editor. *In Sharks, Skates and Rays*. The Johns Hopkins University Press, Baltimore, MD.
- Hannah, R. W., P. S. Rankin, and M. T. O. Blume. 2012. Use of a novel cage system to measure postrecompression survival of Northeast Pacific rockfish. *Marine and Coastal Fisheries* 4(1):46-56.
- Harry, G. Y. 1959. Time of spawning, length at maturity, and fecundity of the English, petrale, and Dover soles (*Parophrys vetulus*, *Eopsetta jordani*, and *Mircrostomus pacificus*, respectively). *Fisheries Commission of Oregon, Research Briefs* 7(1):5-13.
- Hart, J. L. 1988. Pacific Fishes of Canada. *Bull.Fish.Res.Bd.Canada* 180:1-730.
- Hawkins, S. L., J. Heifetz, C. M. Kondzela, J. E. Pohl, R. L. Wilmot, O. N. Katugin, and coauthors. 2005. Genetic variation of roughey rockfish (*Sebastes aleutianus*) and shortraker rockfish (*S. borealis*) inferred from allozymes. *Fishery Bulletin* 103:524-535.
- He, X., D. Pearson, E. J. Dick, J. Field, S. Ralston, and A. D. MacCall. 2006a. Status of the Widow Rockfish Resource in 2005. *in* Volume 3: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- He, X., D. Pearson, E. J. Dick, J. C. Field, S. Ralston, and A. D. MacCall. 2008. Status of the widow rockfish resource in 2007: An Update. PFMC, Portland, Oregon.

- He, X., D. E. Pearson, E. J. Dick, J. C. Field, S. Ralston, and A. D. MacCall. 2011. Status of the widow rockfish resources in 2011. Pacific Fishery Management Council, Portland, OR.
- He, X., D. E. Pearson, J. C. Field, L. Lefebvre, and M. Key. 2013. Status of the Pacific sanddab resource in 2013. Pacific Fishery Management Council, Portland, OR.
- He, X., A. Punt, A. D. MacCall, and S. Ralston. 2006b. Rebuilding Analysis for Widow Rockfish in 2005. *in* Volume 3: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- He, X., S. V. Ralston, A. D. MacCall, D. E. Pearson, and E. J. Dick. 2003. Status of the widow rockfish resource in 2003. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Heifetz, J., J. N. Ianelli, D. M. Clausen, D. L. Courtney, and J. T. Fujioka. 2000. Slope rockfish. *in* Stock assessment fishery evaluation report for the groundfish resources of the Gulf of Alaska as projected for 2001. North Pacific Fishery Management Council, Anchorage.
- Helly, J. J. and L. A. Levin. 2004. Global distribution of naturally occurring marine hypoxia on continental margins. *Deep Sea Research* 51:1159-1168.
- Helser, T. 2006. Stock assessment of the blackgill rockfish (*Sebastes melanostomus*) population off the west coast of the United States in 2005. *in* Volume 5: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Hess, J., P. Chittaro, A. Elz, E. Gilbert-Horvath, V. Simon, and J. Garza. 2014. Cryptic population structure in the severely depleted cowcod, *Sebastes levis*. *Canadian Journal of Fisheries and Aquatic Sciences* 71(1):81-92.
- Hicks, A. C., M. A. Haltuch, and C. Wetzel. 2009. Status of greenstriped rockfish (*Sebastes elongatus*) along the outer coast of California, Oregon, and Washington. Pacific Fishery Management Council, Portland, OR.
- Hicks, A. C. and C. Wetzel. 2011. The Status of Dover Sole (*Microstomus pacificus*) along the U.S. West Coast in 2011. Pacific Fishery Management Council, Portland, OR.
- Hicks, A. C., C. Wetzell, and J. Harms. 2013. The Status of Rougheye Rockfish (*Sebastes aleutianus*) and Blackspotted Rockfish (*S. melanostictus*) as a Complex Along the U.S. West Coast in 2013. Pacific Fishery Management Council, Portland, OR.
- Hilborn, R., J. L. Valero, and M. Maunder. 2001. Status of the sablefish resource off the U.S. Pacific coast in 2001. *in* Appendix to the status of the Pacific Coast groundfish fishery through 2001 and acceptable biological catches for 2002 (Stock assessment and fishery evaluation). Pacific Fishery Management Council, Portland, OR.
- Hill, K. T., E. Dorval, N. C. H. Lo, B. J. Macewicz, C. Show, and R. Felix-Uraga. 2008. Assessment of the Pacific sardine resource in 2007 for U.S. management in 2008. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-413.

- Hobson, E. S. and J. R. Chess. 1988. Trophic relations of the blue rockfish, *Sebastes mystinus*, in a coastal upwelling system off northern California. *Fishery Bulletin* 86(4):715-743.
- Hobson, E. S., J. R. Chess, and D. F. Howard. 1996. Zooplankters consumed by blue rockfish during brief access to a current off California's Sonoma coast. *California Department of Fish and Game* 82(2):87-92.
- Hoenig, J. M. and S. H. Gruber. 1990. Life-history pattern in the elasmobranchs: implications for fisheries management. Pages 1-16 in H. L. J. Pratt, S. H. Gruber, and T. Taniuchi, editors. *Elasmobranchs as Living Resources: Advances in the Biology, Ecology, Systematic, and the Status of the Fisheries*. Proceedings of the Second United States-Japan Workshop. NOAA Technical Report, Honolulu, HI.
- Hogue, E. W. and A. G. Carey. 1982. Feeding ecology of 0-age flatfishes at a nursery ground on the Oregon coast. *Fish.Bull.* 80:555-565.
- Holden, M. J. 1974. Problems in the rational exploitation of elasmobranch populations and some suggested solutions. Pages 117-137 in *Sea Fisheries Research*. John Wiley & Son, New York, NY.
- Hollowed, A. B. 1992. Spatial and temporal distribution of Pacific hake, *Merluccius productus*, larvae and estimates of survival during early life stages. *Calif.Coop.Oceanic Fish.Invest.Rep.* 33:100-123.
- Hosie, M. J. 1976. The arrowtooth flounder. Oregon Department of Fish and Wildlife.
- Hulberg, L. W. and J. S. Oliver. 1979. Prey availability and the diets of two co-occurring flatfish. Pages 29-36 in S. J. Lipovsky and C. A. Simenstad, editors. *Washington Sea Grant University of Washington, Seattle, 1979*.
- Hyde, J. R. and R. D. Vetter. 2007. The origin, evolution, and diversification of rockfishes of the genus *Sebastes* (Cuvier). *Molecular Phylogenetics and Evolution* 44:790-811.
- Ianelli, J. N., R. Lauth, and L. D. Jacobson. 1994. Status of the thornyhead resource in 1994. in *Status of the Pacific groundfish fishery through 1994 and recommended acceptable biological catches for 1995*. Pacific Fishery Management Council, Portland, Oregon.
- Ianelli, J. N., M. Wilkins, and S. Harley. 2000. Status and future prospects for the Pacific ocean perch resource in waters off Washington and Oregon as assessed in 2000. in *Appendix to Status of the Pacific coast groundfish fishery through 2000 and recommended Acceptable Biological Catches for 2001 (Stock Assessment and Fishery Evaluation)*. Pacific Fishery Management Council, Portland, OR.
- Ianelli, J. N. and M. Zimmerman. 1998. Status and future prospects for the Pacific ocean perch resource in waters off Washington and Oregon as assessed in 1998. Pacific Fishery Management Council, Portland, OR.
- Isaacs, J. D. and R. A. Schwartzlose. 1965. Migrant sound scatterers: interaction with the sea floor. *Science* 150:1810-1813.
- Ito, D. H., D. K. Kimura, and M. E. Wilkins. 1986. Appendix 3: Current status and future prospects for the Pacific ocean perch resource in waters off Washington and Oregon. in *Status of the Pacific coast*



- groundfish fishery through 1986 and recommended acceptable biological catches for 1987. Pacific Fishery Management Council, Portland, OR.
- Iwamoto, E., M. J. Ford, and R. G. Gustafson. 2004. Genetic population structure of Pacific hake, *Merluccius productus*, in the Pacific Northwest. *Environmental Biology of Fishes* 69:187-199.
- Jacobson, L. D. 1990. Thornyheads stock assessment for 1990. Appendix D. *in* Status of the Pacific Coast Groundfish Fishery Through 1990 and Recommended Acceptable Biological Catches for 1991. Pacific Fishery Management Council, Portland, OR.
- Jacobson, L. D. 1991. Thornyheads stock assessment for 1991. Appendix D. *in* Status of the Pacific Coast Groundfish Fishery Through 1991 and Recommended Acceptable Biological Catches for 1992. Pacific Fishery Management Council, Portland, OR.
- Jacobson, L. D. and N. C. H. Lo. 1994. Spawning biomass of the northern anchovy in 1994, La Jolla, CA.
- Jacobson, L. D. and R. D. Vetter. 1996. Bathymetric demography and niche separation of thornyhead rockfish: *Sebastolobus alascanus* and *Sebastolobus altivelis*. *Can.J.Fish.Aquat.Sci.* 53:600-609.
- Jagiello, T. H. 1990. Movement of tagged lingcod, *Ophiodon elongatus*, at Neah Bay, Washington. *Fish.Bull.* 88:815-820.
- Jagiello, T. H., P. Adams, M. Peoples, S. Rosenfield, K. R. Silberberg, and T. E. Laidig. 1997. Assessment of lingcod in 1997. Pacific Fishery Management Council, Portland, OR.
- Jagiello, T. H. and J. Hastie. 2001. Updated rebuilding analysis for lingcod. Unpublished report prepared for the Pacific Fishery Management Council, Portland, OR.
- Jagiello, T. H. and F. Wallace. 2006. Assessment of Lingcod (*Ophiodon elongatus*) for the Pacific Fishery Management Council in 2005 *in* Volume 5: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Jagiello, T. H., F. R. Wallace, and Y. W. Cheng. 2004. Assessment of lingcod (*Ophiodon elongatus*) for the Pacific Fishery Management Council in 2003. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2004 and recommended acceptable biological catches for 2005-2006 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Jagiello, T. H., D. Wilson-Vandenberg, J. Sneva, S. Rosenfield, and F. Wallace. 2000. Assessment of lingcod (*Ophiodon elongatus*) for the Pacific Fishery Management Council in 2000. *in* Appendix to Status of the Pacific coast groundfish fishery through 2000 and recommended acceptable biological catches for 2001 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Jarvis, E. T., M. J. Allen, and R. W. Smith. 2004. Comparison of recreational fish catch trends to environment-species relationships and fishery-independent data in the Southern California Bight, 1980-2000. *CalCOFI Reports* 45:167-179.
- Jarvis, E. T. and C. G. Lowe. 2008. The effects of barotrauma on the catch-and-release survival of southern California nearshore and shelf rockfish (*Scorpaenidae*, *Sebastes* spp.). *Canadian Journal of Fisheries and Aquatic Sciences* 65(7):1286-1296.

- Johnson, S. L., W. H. Barss, and R. L. Demory. 1982. Rockfish assessment studies on Heceta Bank, Oregon, 1980-81. Oregon Department of Fish and Wildlife Project Annual Report.
- Jones, W. G. and G. Y. Harry, Jr. 1960. The Oregon trawl fishery for mink food 1948-1957. Fish Commission of Oregon Research Briefs 8:14-30.
- Jorgensen, S. J., D. M. Kaplan, A. P. Klimley, S. G. Morgan, M. R. O'Farrell, and L. W. Botsford. 2006. Limited movement in blue rockfish *Sebastes mystinus*: internal structure of home range. Marine Ecological Progress Series 327:157-170.
- Jow, T. 1969. Results of English sole tagging off California. Pac.Mar.Fish.Comm.Bull. 7:16-33.
- Kaplan, I. C. and T. E. Helser. 2008. Stock Assessment of the Arrowtooth flounder ( *Atheresthes stomias* ) Population off the West Coast of the United States in 2007. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation Stock Assessments and Rebuilding Analyses.
- Karnowski, M., V. Gertseva, and A. Stephens. 2012. Historical reconstruction of Oregon's commercial fisheries landings. Oregon Department of Fish and Wildlife.
- Karpov, K. A., D. P. Albin, and W. H. Van Buskirk. 1995. The marine recreational fishery in northern and central California. California Department of Fish and Game Bulletin 176:192 pgs.
- Keeling, R. F., A. Körtzinger, and N. Gruber. 2010. Ocean deoxygenation in a warming world. Annual Reviews in Marine Science 2:199-229.
- Kendall, A. W., Jr. and W. H. Lenarz. 1986. Status of early life history studies of northeast Pacific rockfishes. Pages 99-128 in. Alaska Sea Grant College Program, Anchorage, Alaska, 1986.
- Kendall, A. W., Jr. and A. C. Matarese. 1987. Biology of eggs, larvae, and epipelagic juveniles of sablefish, *Anoplopoma fimbria* , in relation to their potential use in management. Marine Fisheries Review 49:1-13.
- Ketchen, K. S. 1956. Factors influencing the survival of the lemon sole (*Parophrys vetulus*) in Hecate Strait, British Columbia. Fish.Res.Bd.Canada 13:647-694.
- Ketchen, K. S. 1986. The spiny dogfish (*Squalus acanthias*) in the northeast Pacific and a history of its utilization. Canadian Special Publication of Fisheries and Aquatic Sciences 88.
- Ketchen, K. S. and C. R. Forrester. 1966. Population dynamics of the petrale sole, *Eopsetta jordani*, in waters off western Canada. Bull.Fish.Res.Bd.Canada 153:1-195.
- Key, M., A. D. MacCall, T. Bishop, and B. Leos. 2006. Stock assessment of the gopher rockfish (*Sebastes carnatus*) in Volume 5: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Key, M., A. D. MacCall, D. B. Field, D. Aseltine-Neilson, and K. Lynn. 2008. The 2007 Assessment of Blue Rockfish ( *Sebastes mystinus* ) in California. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation Stock Assessments and Rebuilding Analyses.

- Kihara, K. and A. M. Shimada. 1988. Prey-predator interactions of the Pacific cod, *Gadus macrocephalus*, and water temperature. *Bull.Jpn.Soc.Sci.Fish.* 54:2085-2088.
- Kimmerer, W. J. 2002. Effects of freshwater flow on abundance of estuarine organisms: physical effects or trophic linkages? *Mar.Ecol.Prog.Ser.* 243:39-55.
- King, J. R., G. A. MacFarlane, S. R. M. Jones, S. R. Gilmore, and C. L. Abbott. 2012. Stock delineation of migratory and resident Pacific hake in Canadian waters. *Fisheries Research* 114:19-30.
- King, J. R. and G. A. McFarlane. 2003. Marine fish life history strategies: applications to fishery management. *Fisheries Management and Ecology* 10:249-264.
- Klingbeil, R. A. 1976. Southern range extensions of the blue rockfish, *Sebastes mystinus*; the flag rockfish, *S. rubrivinctus*; and the shortbelly rockfish, *S. jordani*. *California Fish and Game* 62:160.
- Klovach, N. V., O. A. Rovnina, and D. V. Kol'stov. 1995. Biology and exploitation of Pacific cod, *Gadus macrocephalus*, in the Anadyr-Navarin region of the Bering Sea. *J.Ichthy.* 35:9-17.
- Koslow, J. A., G. W. Boehlert, J. D. M. Gordon, R. L. Haedrick, P. Lorance, and N. Parin. 2000. Continental slope and deep-sea fisheries: implications for a fragile ecosystem. *ICES Journal of Marine Science* 57:548-557.
- Kramer, D. E. and V. M. O'Connell. 1995. Guide to Northeast Pacific rockfishes. Genera *Sebastes* and *Sebastolobus*. Alaska Sea Grant, Marine Advisory Bulletin 25.
- Kravitz, M. J., W. G. Pearcy, and M. P. Guin. 1977. Food of five species of co-occurring flatfishes on Oregon's continental shelf. *Fishery Bulletin* 74(4):984-990.
- Kronman, M. (S. B. M. H. Museum). 1999. Santa Barbara's hook-and-line fisheries, Santa Barbara, CA.
- Krygier, E. E. and W. G. Pearcy. 1986. The role of estuarine and offshore nursery areas for young English sole, *Parophrys vetulus*, Girard, off Oregon. *Fish.Bull.* 84:119-132.
- Lai, H. L., M. Haltuch, A. Punt, and J. M. Cope. 2006. Stock assessment of petrale sole: 2004. *in* Volume 2: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, Oregon.
- Lai, H. L., J. V. Tagart, J. N. Ianelli, and F. R. Wallace. 2003. Status of the yellowtail rockfish resource in 2003. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Laidig, T. E., P. B. Adams, and W. M. Samiere (M. E. Wilkins and M. W. Saunders). 1997. Feeding habits of Sablefish, *Anoplopoma fimbria*, off the coast of Oregon and California. National Oceanic and Atmospheric Administration.
- Laidig, T. E., D. L. Watters, and M. M. Yoklavich. 2009. Demersal fish and habitat associations from visual surveys on the central California shelf. *Estuarine, Coastal, and Shelf Science* 83:629-637.

- Lamb, A. and P. Edgell. 1986. Coastal fishes of the Pacific northwest. Harbour Publishing Co. Ltd., Madeira Park, British Columbia.
- LaRiviere, M. G., D. D. Jessup, and S. B. Mathews. 1980. Lingcod, *Ophiodon elongatus*, spawning and nesting in San Juan Channel, Washington. *Calif. Fish and Game* 67:231-239.
- Laroche, W. A. and S. L. Richardson. 1980. Development and occurrence of larvae and juveniles of the rockfishes *Sebastes flavidus* and *Sebastes melanops* (Scorpaenidae) off Oregon. *Fish. Bull.* 77:901-923.
- Larson, R. J. and D. Wilson-Vandenberg. 2001. Other nearshore rockfishes. Pages 185-188 in W. S. Leet, R. Klingbeil, and E. J. Larson, editors. *California's Living Marine Resources: A status report*, volume 1. California Department of Fish and Game.
- Lauth, R. 1987. Spawning ecology and nesting behavior of the Cabezon, *Scorpaenichthys mamoratus* in Puget Sound, Washington. Master's Thesis. University of Washington.
- Lauth, R. 2000. The 1999 Pacific west coast upper continental slope trawl survey of groundfish resources off Washington, Oregon, and California: estimates of distribution, abundance, and length composition. National Oceanic and Atmospheric Administration.
- Lea, R. N., R. D. McAllister, and D. A. VanTresca. 1999. Biological aspects of nearshore rockfishes of the genus *Sebastes* from central California. California Department of Fish and Game Fish Bulletin 177
- Leaman, B. M. and R. J. Beamish. 1984. Ecological and management implications of longevity in some northeast Pacific groundfish. *Bulletin of the International North Pacific Fisheries Commission* 42:85-97.
- Lee, Y. W. 2002. Oceanographic effect on the dynamics of food habits and growth conditions of some groundfish species of the Pacific Northwest. Ph.D Dissertation. Oregon State University.
- Lenarz, W. H. 1993. An initial examination of the status of the darkblotched rockfish fishery off the coasts of California, Oregon, and Washington. in Appendix C in Appendices to the status of the Pacific Coast groundfish through 1993 and recommended acceptable biological catches for 1994.
- Lenarz, W. H. and T. Wyllie Echeverria. 1991. Sexual dimorphism in *Sebastes*. *Environmental Biology of Fishes* 30:71-80.
- Levin, P. S., E. E. Holmes, K. R. Piner, and C. J. Harvey. 2006. Shifts in a Pacific Ocean fish assemblage: The potential influence of exploitation. *Conservation Biology* 20:1181-1190.
- Limbaugh, C. 1955. Fish life in the kelp beds and the effects of kelp harvesting. University of California, Institute of Marine Research 55-9:1-158.
- Lloris, D., J. E. Matallanas, and P. Oliver. 2005. Hakes of the World (Family *Merlucciidae*): an annotated and illustrated catalogue of hake species known to date. FAO species catalogue for fisheries purposes, Rome.
- Lorz, H. V., W. G. Pearcy, and M. Fraidenburg. 1983. Notes on the feeding habits of the yellowtail rockfish, *Sebastes flavidus*, off Washington and in Queen Charlotte Sound. *Calif. Fish. Game* 69:33-38.

- Love, M. S. 1996. Probably more than you want to know about the fishes of the Pacific Coast. Really Big Press, Santa Barbara, California.
- Love, M. S. 2011. Certainly more than you want to know about the fishes of the Pacific Coast. A postmodern experience. Really Big Press, Santa Barbara, CA.
- Love, M. S., B. Axell, P. Morris, R. Collins, and A. Brooks. 1987. Life history and fishery of the California scorpionfish, *Scorpaena guttata*, within the Southern California Bight. *Fishery Bulletin* 85:90-116.
- Love, M. S., J. E. Caselle, and W. Van Buskirk. 1998. A severe decline in the commercial passenger fishing vessel rockfish (*Sebastes* spp.) catch in the southern California bight, 1980-1996. *CalCOFI Reports* 39:180-195.
- Love, M. S. and A. W. Ebeling. 1978. Food and habitat of three switch-feeding fishes in the kelp forests off Santa Barbara, California. *Fishery Bulletin* 76(1):257-271.
- Love, M. S., C. W. Mecklenburg, T. A. Mecklenburg, and L. K. Thorsteinson. 2005. Resource Inventory of Marine and Estuarine Fishes of the West Coast and Alaska: A Checklist of North Pacific and Arctic Ocean Species from Baja California to the Alaska-Yukon Border. US Department of the Interior, US Geological Survey. National Biological Information Infrastructure.
- Love, M. S., P. Morris, M. McCrae, and R. Collins. 1990. Life history aspects of 19 rockfish species (*Scorpaenidae*: *Sebastes*) from the southern California bight.
- Love, M. S. and M. Yoklavich. 2008. Habitat characteristics of juvenile cowcod, *Sebastes levis* (*Scorpaenidae*), in Southern California. *Environmental Biology of Fishes* (82):195-202.
- Love, M. S., M. Yoklavich, and L. Thorsteinson. 2002. The rockfishes of the northeast Pacific. University of California Press, Berkeley, California.
- Lowe, C. G., K. M. Anthony, E. T. Jarvis, L. F. Bellquist, and M. S. Love. 2009. Site fidelity and movement patterns of groundfish associated with offshore petroleum platforms in the Santa Barbara Channel. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* 1:71-89.
- Lowry, M. S. and J. V. Carretta. 1999. Market squid (*Loligo opalescens*) in the diet of California sea lions (*Zalophus californianus*) in southern California (1981-1995). *CalCOFI Reports* 40:196-207.
- MacCall, A. D. 2002. Status of bocaccio off California in 2002. *in* Volume 1 Status of the Pacific Coast groundfish fishery through 2002 and recommended acceptable biological catches for 2003 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- MacCall, A. D. 2003a. Bocaccio rebuilding analysis for 2003. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- MacCall, A. D. 2003b. Status of bocaccio off California in 2003. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.

- MacCall, A. D. 2006a. Bocaccio Rebuilding Analysis for 2005. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- MacCall, A. D. 2006b. Status of Bocaccio off California in 2005. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- MacCall, A. D. 2008a. Bocaccio rebuilding analysis for 2007. Pacific Fishery Management Council, Portland, OR.
- MacCall, A. D. 2008b. Status of bocaccio off California in 2007. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation Stock Assessments and Rebuilding Analyses.
- MacCall, A. D. 2009. Depletion-corrected average catch: a simple formula for estimating sustainable yields in data-poor situations. *ICES J.of Mar.Sci.* 66:2267-2271.
- MacCall, A. D. and X. He. 2002. Bocaccio rebuilding analysis for 2002. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2002 and recommended acceptable biological catches for 2003 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- MacCall, A. D., S. Ralston, D. Pearson, and E. Williams. 1999. Status of bocaccio off California in 1999 and outlook for the next millennium. *in* Appendix to Status of the Pacific Coast groundfish fishery through 1999 and recommended acceptable biological catches for 2000 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Mace, P. M., N. W. Bartoo, A. B. Hollowed, P. Kleiber, R. D. Methot, S. A. Murawski, and coauthors. 2001. Marine Fisheries Stock Assessment Improvement Plan. NOAA Technical Memorandum NMFS-F/SPO-56:69.
- Mace, P. M. and I. J. Doonan. 1988. A generalized bioeconomic simulation model for fish population dynamics. *New Zealand Fishery Assessment Research Document* 88(4).
- MacGregor, J. S. 1986. Relative abundance of four species of *Sebastes* off California and Baja California. *Calif.Coop.Oceanic Fish.Invest.Rep.* 27:121-135.
- Mantua, N. J., S. R. Hare, Y. Zhang, J. M. Wallace, and R. C. Francis. 1997. A Pacific interdecadal climate oscillation with impacts on salmon production. *Bulletin of the American Meteorological Society* 78(6):1069-1079.
- Markle, D. F., P. M. Harris, and C. L. Toole. 1992. Metamorphosis and an overview of early life history stages in Dover sole, *Microstomus pacificus* *Fish.Bull.* 90:285-301.
- Martin, L. and G. D. Zorzi. 1993. Status and review of the California skate fishery. Pages 39-52 *in* S. Branstetter, editor. Conservation biology of elasmobranchs, volume 115. NOAA Technical Report NMFS.
- Mason, J. C., R. J. Beamish, and G. A. McFarlane. 1983. Sexual maturity, fecundity, spawning, and early life history of sablefish (*Anoplopoma fimbria*) in waters off the Pacific coast of Canada. *Canadian Journal of Fisheries and Aquatic Sciences* 40:2621-2134.

- Mason, J. E. 1998. Declining rockfish lengths in the Monterey Bay, California, recreational fishery, 1959-94. *Marine Fisheries Review* 60(3):15-28.
- Mathews, S. B. and M. LaRiviere. 1987. Movement of tagged lingcod, *Ophiodon elongatus*, in the Pacific Northwest. *Fish Bull.* 85:153-159.
- Matthews, K. R. 1986. Movement of two nearshore, territorial rockfishes previously reported as non-movers and implications to management. *California Department of Fish and Game Bulletin* 72(2):103-109.
- Matthews, K. R. 1992. A telemetric study of the home ranges and homing routes of lingcod, *Ophiodon elongatus*, on shallow rocky reefs off Vancouver Island, British Columbia. *Fish.Bull.* 90:784-790.
- Maunder, M., J. T. Barnes, D. Aseltine-Neilson, and A. D. MacCall. 2006. The Status of California Scorpionfish (*Scorpaena guttata*) off Southern California in 2004. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- MBC. 1987. Ecology of important fisheries species offshore California. Minerals Management Service, Pacific Outer Continental Shelf Region, Washington, D.C.
- McClatchie, S., R. Goerick, R. Rich-Cosgrove, and R. Vetter. 2010. Oxygen in the Southern California Bight: multidecadal trends and implications for demersal fisheries. *Geophysical Research Letters* 37(19).
- McEachran, J. D. 1990. Diversity of rays: why are there so many species? *Chondros* 5(2):1-6.
- McFarlane, G. A. and R. Beamish. 1983a. Preliminary observations on the juvenile biology of sablefish (*Anoplopoma fimbria*) off the west coast of Canada. *in* Proceedings of the International Sablefish Symposium. Alaska Sea Grant Report 83-3.
- McFarlane, G. A. and R. J. Beamish. 1983b. Biology of adult sablefish (*Anoplopoma fimbria*) in waters off western Canada. Alaska Sea Grant College Program, University of Alaska, Anchorage, 1983.
- McFarlane, G. A. and R. J. Beamish. 1986. Biology and fishery of Pacific hake, *Merluccius productus*, in the Strait of Georgia. *Int.N.Pac.Fish.Comm.Bull.* 50:365-392.
- McFarlane, G. A. and J. R. King. 2003. Migration patterns of spiny dogfish (*Squalus acanthias*) in the North Pacific Ocean. *Fishery Bulletin* 101:358-367.
- Merkel, T. J. 1957. Food habits of king salmon, *Oncorhynchus tshawytscha* (Walbaum), in the vicinity of San Francisco, California. *California Department of Fish and Game* 43:249-270.
- Method, R. D. 2000. Rebuilding analysis for canary rockfish. Unpublished report prepared for the Pacific Fishery Management Council, Portland, OR.
- Method, R. D. 2006. Updated Rebuilding Analysis for Canary Rockfish Based on Stock Assessment in 2005. *in* Volume 6: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.

- Methot, R. D. and K. Piner. 2002a. Rebuilding analysis for yelloweye rockfish: update to incorporate results of coastwide assessment in 2002. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Methot, R. D. and K. Piner. 2002b. Status of the canary rockfish resource off California, Oregon and Washington in 2001. *in* Volume 1 Status of the Pacific Coast groundfish fishery through 2002 and recommended acceptable biological catches for 2003 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Methot, R. D. and J. Rogers. 2001. Rebuilding analysis for darkblotched rockfish. Unpublished report prepared for the Pacific Fishery Management Council, Portland, OR.
- Methot, R. D. and I. J. Stewart. 2006. Status of the U.S. canary rockfish resource in 2005. *in* Volume 6: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Methot, R. D., F. Wallace, and K. Piner. 2003. Status of yelloweye rockfish off the U.S. West Coast in 2002. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Miller, A. K. and W. Sydeman. 2004. Rockfish response to low-frequency ocean climate change as revealed by the diet of a marine bird over multiple time scales. *Marine Ecology Progress Series* 281:207-216.
- Miller, D. J. and J. J. Geibel. 1973. Summary of blue rockfish and lingcod life histories; a reef ecology study and giant kelp *Macrocystis pyrifera*, experiments in Monterey Bay, California. *Calif.Dept.Fish and Game, Fish Bull.* 158:131.
- Miller, D. J. and D. Gotshall. 1965. Ocean sportfish catch and effort from Oregon to Point Arguello, California: July 1, 1957-June 30, 1961. *California Department of Fish and Game Fishery Bulletin* 130:135 pgs.
- Miller, D. J. and R. N. Lea. 1972. Guide to the coastal marine fishes of California. *Calif.Dept.Fish and Game, Fish.Bull.* 157:249.
- Mills, K. L., T. Laidig, S. Ralston, and W. J. Sydeman. 2007. Diets of top predators indicate pelagic juvenile rockfish (*Sebastes* spp.) abundance in the California Current System. *Fisheries Oceanography* 16(3):273-283.
- Morejohn, G. V., J. T. Harvey, and L. T. Krasnow. 1978. The importance of *Loligo opalescens* in the food web of marine vertebrates in Monterey Bay, California. *California Department of Fish and Game Bulletin* 169:67-98.
- Moser, H. G. 1974. Development and distribution of larvae and juveniles of *Sebastolobus* (Pisces: Family Scorpaenidae). *Fishery Bulletin* 72:865-884.



- Moser, H. G. and E. H. Ahlstrom. 1978. Larvae and pelagic juveniles of blackgill rockfish, *Sebastes melanostomus*, taken in midwater trawls off Southern California and Baja California. Journal of the Fisheries Research Board of Canada 35:981-996.
- Moser, H. G., R. L. Charter, W. Watson, D. A. Ambrose, J. Butler, S. R. Charter, and coauthors. 2000. Abundance and distribution of rockfish (*Sebastes*) larvae in the Southern California Bight in relation to environmental conditions and fishery exploitation. CalCOFI Rep. 41:132-147.
- Moser, H. G. e. 1996. The early stages of fishes in the California Current region. California Cooperative Oceanic Fisheries Investigations, Atlas No. 33. Allen Press, Inc., Lawrence, KS.
- Moyle, P. B. and J. J. J. Cech. 1996. Fishes, An Introduction to Ichthyology, 3rd edition. Prentice-Hall, Upper Saddle River, NJ.
- Mulligan, T. J. and B. M. Leaman. 1992. Length-at-age analysis: Can you get what you see? Can.J.Fish.Aquat.Sci. 49:632-643.
- Myers, R. A., K. G. Bowen, and N. Barrowman. 1999. Maximum reproductive rate of fish at low population sizes. Canadian Journal of Fisheries and Aquatic Sciences (56):2404-2419.
- Nakano, H. and K. Nagasawa. 1996. Distribution of pelagic elasmobranchs caught by salmon research gillnets in the North Pacific. Fisheries Science 62:860-865.
- Nichol, D. 1990. Life history examination of darkblotched rockfish ( *Sebastes crameri* ) off the Oregon coast. M.S. Thesis. Oregon State University.
- Nichol, D. G. and E. K. Pikitch. 1994. Reproduction of darkblotched rockfish off the Oregon coast. Transactions of the American Fisheries Society 123:469-481.
- Nitsos, R. J. 1965. Species composition of rockfish (family *Scorpaenidae*) landed by California otter trawl vessels, 1962-1963. Pacific Marine Fisheries Commission, Portland, OR.
- Nitsos, R. J. and P. H. Reed. 1965. The animal food fishery in California, 1961-1962. California Fish and Game 51(1):16-27.
- NMFS (National Marine Fisheries Service). 2013. Groundfish Essential Fish Habitat Synthesis: A Report to the Pacific Fishery Management Council. NOAA NMFS Northwest Fisheries Science Center, Seattle, April 2013.
- NMFS, S. and S. OT (S. National Marine Fisheries). 1998. Status of the shortspine thornyhead resource off the U.S. Pacific coast in 1998. In: Appendix to Status of the Pacific coast groundfish fishery through 1998 and recommended acceptable biological catches for 1999. Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- NOAA (N. O. a. A. Administration). 1990. West coast of North America coastal and ocean zones strategic assessment: Data atlas. OMA/NOS, Ocean Assessments Division, Strategic Assessment Branch, NOAA.
- Norton, E. C. and R. B. MacFarlane. 1995. Nutritional dynamics of reproduction in viviparous yellowtail rockfish, *Sebastes flavidus* Fish.Bull. 93:299-307.

- O'Connell, C. P. 1953. The life history of the cabezon *Scorpaenichthys marmoratus*
- O'Connell, V. M. and D. W. Carlile. 1993. Habitat-specific density of adult yelloweye rockfish *Sebastes ruberrimus* in the eastern Gulf of Alaska. *Fish.Bull.* 91:304-309.
- O'Connell, V. M. and F. C. Funk. 1986. Age and growth of yelloweye rockfish (*Sebastes ruberrimus*) landed in southeastern Alaska. Pages 171-185 *in*, volume 87-2. Alaska Sea Grant College Program, Anchorage, Alaska, 1986.
- O'Farrell, M. R. and L. W. Botsford. 2005. Estimation of change in lifetime egg production from length frequency data. *Canadian Journal of Fisheries and Aquatic Sciences* 62:1626-1639.
- O'Farrell, M. R. and L. W. Botsford. 2006. Estimating the status of nearshore rockfish (*Sebastes* spp.) populations with length frequency data. *Ecological Applications* 16(3):977-986.
- Oda, K. T. 1992. Chilipepper. Pages 122 *in* W. S. Leet, C. M. Dewees, and C. W. Haugen, editors. *California's Living Marine Resources and Their Utilization*. California Sea Grant Program, Davis, CA.
- Orcutt, H. G. 1950. The life history of the starry flounder (*Platichthys stellatus*). *Calif.Dept.Fish and Game, Fish.Bull.* 78.
- Orr, J. W., M. A. Brown, and D. C. Baker. 1998. Guide to Rockfishes (*Scorpaenidae*) of the Genera *Sebastes*, *Sebastolobus*, and *Adelosebastes* of the Northeast Pacific Ocean.
- Orr, J. W. and S. Hawkins. 2008. Species of the rougheye rockfish complex: resurrection of *Sebastes melanosticus* (Matsubara, 1934) and a redescription of *Sebastes aleutianus* (Jordan and Evermann, 1898) (Teleostei: *Scorpaeniformes*). *Fishery Bulletin* 106:111-134.
- Orton, G. 1955. Early developmental stages of the California scorpionfish, *Scorpaena guttata* Copeia 1955:210-214.
- Palsson, W. A. 1990. Pacific cod in Puget Sound and adjacent waters: Biology and stock assessment. *Wash.Dept.Fish.Tech.Rep.* 112:137.
- Parker, S. J., H. I. McElderry, P. S. Rankin, and R. W. Hannah. 2006. Buoyancy regulation and barotrauma in two species of nearshore rockfish. *Transactions of the American Fisheries Society* 135(5):1213-1223.
- Parker, S. J., P. S. Rankin, R. W. Hannah, and C. B. Schreck. 2003. Discard mortality of trawl-caught lingcod in relation to tow duration and time on deck. *North American Journal of Fisheries Management* 23:530-542.
- Patrick, W. S., P. Spence, O. Osmeth, J. M. Cope, J. Field, D. Kobayashi, and coauthors. 2009. Use of productivity and susceptibility indices to determine stock vulnerability, with example applications to six U.S. fisheries. U.S. Dep. Commerce.
- Pearcy, W. G. 1962. Egg masses and early developmental stages of the scorpaneid fish, *Sebastolobus*. *J.Fish.Res.Bd.Canada* 1169-1173.

- Pearcy, W. G. 1978. Distribution and abundance of small flatfishes and other demersal fishes in a region of diverse sediments and bathymetry off Oregon. *Fishery Bulletin* 78:629-643.
- Pearcy, W. G. 1992. Movements of acoustically-tagged yellowtail rockfish *Sebastes flavidus* on Heceta Bank, Oregon. *Fish.Bull.* 90:726-735.
- Pearcy, W. G., M. J. Hosie, and S. L. Richardson. 1977. Distribution and duration of pelagic life of larvae of Dover sole, *Microstomus pacificus*; rex sole, *Glyptocephalus zachirus*; and petrale sole, *Eopsetta jordani*, in waters off Oregon. *Fish.Bull.* 75:173-183.
- Pearsall, I. A. and J. J. Fargo. 2007. Diet composition and habitat fidelity for groundfish in Hecate Strait, British Columbia. Canadian Technical Report of Fisheries and Aquatic Sciences 2692:141.
- Pearson, D. 1989. Survey of fishes and water properties of South San Francisco Bay, California, 1973-82. U.S. Dept. Comm.
- Pearson, D. E., D. A. Douglas, and B. Barss. 1993. Biological observation from the Cobb Seamount rockfish fishery. *Fishery Bulletin* 91:573-576.
- Pearson, D. E., J. E. Hightower, and J. T. H. Chan. 1991. Appendix D: Age, growth, year class strength, and potential yield for shortbelly rockfish. *in* Status of the Pacific coast groundfish fishery through 1989 and recommended acceptable biological catches for 1990. Pacific Fishery Management Council, Portland, OR.
- Pearson, D. E. and S. L. Owen. 1992. English sole. Pages 99-100 *in* W. S. Leet, C. M. Dewees, and C. W. Haugen, editors. California's Living Marine Resources and Their Utilization. California Sea Grant Program, Davis, CA.
- Pearson, K. E. and D. R. Gunderson. 2003. Reproductive biology and ecology of shortspine thornyhead rockfish, *Sebastolobus alascanus*, and longspine thornyhead rockfish, *S. altivelis*, from the northeastern Pacific Ocean. *Environ.Biol.Fish.* 67:117-136.
- Pedersen, M. G. 1975. Movements and growth of petrale sole tagged off Washington and southwest Vancouver Island. *J.Fish.Res.Bd.Canada* 32:2169-2177.
- Pederson, M. G. 1975. Movements and growth of petrale sole (*Eopsetta jordani*) tagged off Washington and southwest Vancouver Island. *Fishery Research Board of Canada Progress Report* 32(1):2169-2177.
- Perez, M. A. and M. A. Bigg. 1986. Diet of northern fur seals, *Callorhinus ursinus*, off Western North America. *Fishery Bulletin* 84:957-970.
- Perry, R. I., M. Stocker, and J. Fargo. 1994. Environmental effects on the distribution of groundfish in Hecate Strait, British Columbia. *Can.J.Fish.Aquat.Sci.* 51:1401-1409.
- PFMC. 2004. Amendment 16-3 to the Pacific Groundfish Fishery Management Plan; rebuilding plans for bocaccio, cowcod, widow rockfish, and yelloweye rockfish. Draft Environmental Impact Statement. Pacific Fishery Management Council, Portland, OR.
- PFMC. 2006. Final environmental impact statement for the proposed groundfish acceptable biological catch and optimum yield specifications and management measures: 2007-2008 Pacific coast groundfish

- fishery and Amendment 16-4: Rebuilding plans for seven depleted Pacific coast groundfish species. Pacific Fishery Management Council, Portland, OR.
- PFMC. 2012. Pacific coast groundfish 5-year review of essential fish habitat. Report to the Pacific Fishery Management Council. Phase 1: New information. Pacific Fishery Management Council, Portland, OR.
- PFMC. 2014. Review of Pacific Coast groundfish essential fish habitat. Phase 2 report to the Pacific Fishery Management Council. Pacific Fishery Management Council, Portland, OR.
- PFMC and NMFS (Pacific Fishery Management Council and National Marine Fisheries Service). 2011. Proposed Harvest Specifications and Management Measures for the 2011-2012 Pacific Coast Groundfish Fishery and Amendment 16-5 to the Pacific Coast Groundfish Fishery Management Plan to Update Existing Rebuilding Plans and Adopt a Rebuilding Plan for Petrale Sole, Final Environmental Impact Statement. National Marine Fisheries Service, Northwest Region, Seattle, February 2011.
- Phillips, J. B. 1939. The rockfish of the Monterey wholesale fish markets. *California Fish and Game* 25:214-225.
- Phillips, J. B. 1957. A review of the rockfishes of California (Family Scorpaenidae). *Calif. Dep. Fish and Game, Fish Bull.* 104:158.
- Phillips, J. B. 1964. Life history studies on ten species of rockfishes (genus *Sebastes*). *Calif. Dep. Fish and Game, Fish Bull.* 126:70.
- Phillips, J. B. and S. Inamura. 1954. The sablefish fishery of California. *Pac. Mar. Fish. Comm. Bull.* 3:5-38.
- Piner, K. 2006. Cowcod Rebuilding Analysis 2005 Analysis of the Progress towards Rebuilding in the Southern California Bight. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Piner, K., E. J. Dick, and J. Field. 2006. 2005 Stock Status of Cowcod in the Southern California Bight and Future Prospects. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Piner, K. and R. Methot. 2001. Stock status of shortspine thornyhead off the Pacific west coast of the United States 2001. *in* Appendix to Status of the Pacific Coast Groundfish Fishery Through 2001 and Acceptable Biological Catches for 2002 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Piner, K., M. Schirripa, T. Builder, J. Rogers, and R. Methot. 2000. Bank rockfish (*Sebastes rufus*) stock assessment for Eureka, Monterey, and Conception INPFC Areas north of Pt. Conception, California. Pacific Fishery Management Council, Portland, OR.
- Porter, P. 1964. Notes on fecundity, spawning, and early life history of petrale sole (*Eopsetta jordani*), with descriptions of flatfish larvae collected in the Pacific Ocean off Humboldt Bay, California. Humboldt State College, Arcata, CA.

- Pratt, H. L. J. and J. G. Casey. 1990. Shark reproductive strategies as a limiting factor in directed fisheries, with a review of Holden's method of estimating growth parameters Pages 97-110 in H. L. J. Pratt, S. H. Gruber, and T. Taniuchi, editors. Elasmobranchs as Living Resources: Advances in the Biology, Ecology, Systematic, and the Status of the Fisheries. Proceedings of the Second United States-Japan Workshop. NOAA Technical Report NMFS, Honolulu, HI.
- Preti, A., S. E. Smith, and D. A. Ramon. 2004. Diet differences in the thresher shark (*Alopias vulpinus*) during transition from a warm-water regime to a cool-water regime off California-Oregon, 1988-2000. CalCOFI Reports 45:118-125.
- Pribyl, A. L., C. B. Schreck, M. L. Kent, M. Kelley, and S. J. Parker. 2012. Recovery potential of black rockfish, *Sebastes melanops* Girard, recompressed following barotrauma. Journal of Fish Diseases 35(4):275-286.
- SSC default rebuilding analysis (Version 2.8a, April 2005). 2005. University of Washington, Seattle.
- Punt, A. E., O. S. Hamel, and I. J. Stewart. 2003. Rebuilding analysis for Pacific ocean perch for 2003. in Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Punt, A. E. and J. N. Ianelli. 2001. Revised rebuilding analysis for Pacific ocean perch. Unpublished report to the Pacific Fishery Management Council, Portland, OR.
- Punt, A. E. and A. D. MacCall. 2002. Revised rebuilding analysis for widow rockfish for 2002. Unpublished report to the Pacific Fishery management Council, Portland, OR.
- Rackowski, J. P. and E. K. Pikitch. 1989. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest), Pacific and speckled sanddabs. Biological Report 82(11.107). Fish and Wildlife Service, U.S. Department of the Interior.
- Ralston, S. 2006. An Assessment of Starry Flounder off California, Oregon, and Washington (2005). in Volume 2: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Ralston, S. and E. J. Dick. 2003. The status of black rockfish ( *Sebastes melanops* ) off Oregon and northern California in 2003. in Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Ralston, S. and J. N. Ianelli. 1998. When lengths are better than ages: The complex case of bocaccio. Pages 451-468 in F. Funk and T. J. Quinn II and J. Heifetz and J. Ianelli and J. E. Powers and J. F. Schweigert, and coeditors, editors. Fishery Stock Assessment Models. University of Alaska, Alaska Sea Grant College Program Report No. AK-SG-98-01, Fairbanks, Alaska.
- Ralston, S., J. N. Ianelli, D. E. Pearson, M. E. Wilkins, R. A. Miller, and D. Thomas. 1996. Status of bocaccio in the Conception/Monterey/Eureka INPFC areas in 1996 and recommendations for management in 1997. in Appendix Vol. 1: Status of the Pacific Coast groundfish fishery through 1996 and recommended acceptable biological catches for 1997 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.

- Ralston, S., D. Pearson, and J. Reynolds. 1998. Status of the chilipepper rockfish stock in 1998. *in* Status of the Pacific coast groundfish fishery through 1998 and recommended acceptable biologic catches for 1999: stock assessment and fishery evaluation, appendix. Pacific Fishery Management Council, Portland, OR.
- Ralston, S., D. E. Pearson, J. C. Field, and M. Key. 2010. Documentation of the California catch reconstruction project. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-461.
- Ralston, S., A. E. Punt, O. S. Hamel, J. D. DeVore, and R. J. Conser. 2011. A meta-analytic approach to quantifying scientific uncertainty in stock assessments. *Fishery Bulletin* 109(2):217-213.
- Reilly, C. A., T. W. Wyllie-Echeverria, and S. Ralston. 1992. Interannual variation and overlap in the diets of pelagic juvenile rockfish (Genus: *Sebastes*) off central California. *Fish.Bull.* 90:505-515.
- Reilly, P. 2001. Blue Rockfish. Pages 165-167 *in* W. S. Leet, C. M. Dewees, R. Klingbeil, and E. J. Larson, editors. California's Living Marine Resources: A Status Report. The Resources Agency, California Department of Fish and Game.
- Reilly, P. N., D. Wilson-Vandenberg, R. N. Lea, C. Wilson, and M. Sullivan. 1994. Recreational angler's guide to the common nearshore fishes of Northern and Central California.
- Reilly, P. N., D. Wilson-Vandenberg, D. L. Watters, J. E. Hardwick, and D. Short. 1993. On board sampling of the rockfish and lingcod commercial passenger fishing vessel industry in northern and central California, May 1987 to December 1991. Marine Research Division Administrative Report:242 pgs.
- Richardson, S. L. and W. A. Laroche. 1979. Development and occurrence of larvae and juveniles of the rockfishes *Sebastes crameri*, *Sebastes pinniger*, and *Sebastes helvomaculatus* (Family Scorpaenidae) off Oregon. *Fish.Bull.* 77:1-46.
- Rickey, M. H. 1993. Status of the coastal arrowtooth flounder resource in 1993. *in* Appendices to the Status of the Pacific Coast Groundfish Fishery through 1993 and Recommended Acceptable Biological Catches for 1994: Stock Assessment and Fishery Evaluation. Pacific Fishery Management Council, Portland, OR.
- Rickey, M. H. 1995. Maturity, spawning, and seasonal movements of arrowtooth flounder, *Atheresthes stomias*, off Washington. *Fish.Bull.* 93:127-138.
- Robinson, H. J., G. M. Cailliet, and D. A. Ebert. 2007. Food habits of the longnose skate, *Raja rhina* (Jordan and Gilbert, 1880), in central California waters. *Environmental Biology of Fishes* 80:165-179.
- Roedel, P. M. 1953. Common ocean fishes of the California coast. *California Fish and Game Bulletin* 91:184 pgs.
- Rogers, C. W., D. R. Gunderson, and D. A. Armstrong. 1988. Utilization of a Washington estuary by juvenile English sole, *Parophrys vetulus* *Fish.Bull.* 86:823-831.
- Rogers, J. B. 1994. Preliminary status of the splitnose rockfish stock in 1994. *in* Appendix H to the Status of the Pacific coast groundfish fishery through 1994 and recommended acceptable biological catches for 1995. Pacific Fishery Management Council, Portland, OR.

- Rogers, J. B. 2003a. Darkblotched rockfish (*Sebastes crameri*) 2003 stock status and rebuilding update. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2003 and recommended acceptable biological catches for 2004 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Rogers, J. B. 2003b. Species allocation of *Sebastes* and *Sebastolobus* sp. caught by foreign countries from 1965 through 1976 off Washington, Oregon, and California, U.S.A. U.S. Department of Commerce.
- Rogers, J. B. 2005a. Status of the darkblotched rockfish (*Sebastes crameri*) resource in 2005. *in* Volume 3: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Rogers, J. B. 2005b. Update of darkblotched rockfish (*Sebastes crameri*) rebuilding analyses. *in* Volume 3: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Rogers, J. B., T. L. Builder, P. R. Crone, J. Brodziak, R. D. Methot, R. J. Conser, and coauthors. 1998. Status of the shortspine thornyhead ( *Sebastolobus alascanus* ) resource in 1998. *in* Appendix to Status of the Pacific coast groundfish fishery through 1998 and recommended acceptable biological catches for 1999 (Stock assessment and fishery evaluation). Pacific Fishery Management Council, Portland, OR.
- Rogers, J. B., L. D. Jacobson, R. Lauth, J. N. Ianelli, and M. Wilkins. 1997. Status of the thornyhead resource in 1997. *in* Appendix to Status of the Pacific coast groundfish fishery through 1997 and recommended acceptable biological catches for 1998 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Rogers, J. B., R. D. Methot, T. L. Builder, K. Piner, and M. Wilkins. 2000. Status of the darkblotched rockfish ( *Sebastes crameri* ) resource in 2000. *in* Appendix to Status of the Pacific coast groundfish fishery through 2000 and recommended acceptable biological catches for 2001 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Rogers, J. B. and E. K. Pikitch. 1992. Numerical definition of groundfish assemblages caught off the coasts of Oregon and Washington using commercial fishing strategies. *Canadian Journal of Fisheries and Aquatic Sciences* 49:2648-2656.
- Rogers, J. B., M. Wilkins, D. Kamakawa, F. Wallace, T. Builder, M. Zimmerman, and coauthors. 1996. Status of the remaining rockfish in the *Sebastes* complex in 1996 and recommendations for management in 1997. *in* Appendix to Status of the Pacific coast groundfish fishery through 1996 and recommended acceptable biological catches for 1997 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Rosenthal, R. J., L. Haldorson, L. J. Field, V. Moran-O'Connell, M. G. LaRiviere, J. Underwood, and coauthors. 1982. Inshore and shallow offshore bottomfish resources in the southeastern Gulf of Alaska (1981-1982). Alaska Dept. Fish and Game, Juneau, Alaska.

- Sakuma, K. M. and S. Ralston. 1995. Distribution patterns of late larval groundfish off central California in relation to hydrographic features during 1992 and 1993. *Calif.Coop.Oceanic Fish.Invest.Rep.* 36:179-192.
- Sampson, D. and Y. W. Lee. 1999. An Assessment of the Stocks of Petrale Sole off Washington, Oregon and Northern California in 1998. *in* Status of the Pacific Coast groundfish fishery through 1998 and recommended biological catches for 1994: stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Sampson, D. B. 1996. Appendix C: Stock status of canary rockfish off Oregon and Washington in 1996. *in* C. Pacific Fishery Management, editor. Status of the Pacific coast groundfish fishery through 1996 and recommended acceptable biological catches for 1997: stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Sampson, D. B. 2002. Analysis of data from the at-sea data collection project. Coastal Oregon Marine Experiment Station, Oregon State University.
- Sampson, D. B. 2005. The status of Dover sole off the U.S. west coast in 2005. Pacific Fishery Management Council, Portland, OR, In press.
- Sampson, D. B. 2008. The Status of Black Rockfish off Oregon and California in 2007. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation Stock Assessments and Rebuilding Analyses.
- Sampson, D. B. and E. M. Stewart. 1993. An assessment of the English sole stock off Oregon and Washington. *in* Appendix H-1 to status of the Pacific coast groundfish fishery through 1993 and recommended acceptable biological catches for 1994 (SAFE Report). Pacific Fishery Management Council, Portland, OR.
- Sampson, D. B. and E. M. Stewart. 1994. Appendix G: Status of the canary rockfish resource off Oregon and Washington in 1994. *in* Status of the Pacific coast groundfish fishery through 1994 and recommended acceptable biological catches for 1995: stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Sampson, D. B. and C. Wood. 2001. Stock status of Dover sole off the west coast in 2000. *in* Appendix to the Status of the Pacific Coast Groundfish Fishery Through 2001 and Acceptable Biological Catches for 2002 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Sarmiento, J. L., T. M. Hughes, R. J. Stouffer, and S. Manabe. 1998. Simulated response of the ocean carbon cycle to anthropogenic climate warming. *Nature* 393:245-249.
- Saunders, M. W. and G. A. McFarlane. 1993. Age and length at maturity of the female spiny dogfish, *Squalus acanthias*, in the Strait of Georgia, British Columbia, Canada. *Environ.Biol.Fish.* 38:49-57.
- Schirripa, M. and J. J. Colbert. 2006. Status of the sablefish resource off the continental U.S. Pacific coasts in 2005. *in* Volume 4: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.



- Schirripa, M. J. 2002. Status of the sablefish resource off the continental U.S. Pacific coast in 2002. *in* Volume 1: Status of the Pacific Coast groundfish fishery through 2002 and recommended acceptable biological catches for 2003 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Schirripa, M. J. 2008. Status of the sablefish resource off the continental U.S. Pacific coast in 2007. Status of the Pacific Coast groundfish fishery through 2007. Stock assessment and fishery evaluation. Stock assessments and rebuilding analyses.
- Schirripa, M. J. and R. Methot. 2001. Status of the sablefish resource off the U.S. Pacific coast in 2001. *in* Appendix to the Status of the Pacific Coast Groundfish Fishery Through 2001 and Acceptable Biological Catches for 2002 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Seibel, B. A. 2011. Critical oxygen levels and metabolic suppression in oceanic oxygen minimum zones. *The Journal of Experimental Biology* 214:326-336.
- Sette, O. E. and R. H. Fielder. 1928. Fishery industries of the United States, 1927. *in* Report of the United States Commissioner of Fisheries for the Fiscal Year 1928. U.S. Department of Commerce.
- Shaffer, J. A., D. C. Doty, R. M. Buckley, and J. E. West. 1995. Crustacean community composition and trophic use of the drift vegetation habitat by juvenile splitnose rockfish *Sebastes diploproa* *Mar.Ecol.Prog.Ser.* 123:13-21.
- Shaw, F. R. and D. R. Gunderson. 2006. Life history traits of the greenstriped rockfish, *Sebastes elongatus*. *California Fish and Game* 92:1-23.
- Shaw, W. N. and T. J. Hassler. 1989. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (Pacific Northwest) -- lingcod. Army Corps of Engineers TR EL-82-4.
- Siebenaller, J. F. 1978. Genetic variability in deep-sea fishes of the genus *sebastolobus*. Pages 95-122 *in* B. Battaglia and J. Beardmore, editors. *Marine Organisms*, New York, NY.
- Siebenaller, J. F. and G. N. Somero. 1982. The Maintenance of Different Enzyme Activity Levels in Congeneric Fishes Living at Different Depths. *Physiological Zoology* 55:171-179.
- Simenstad, C. A., B. S. Miller, C. F. Nybalde, K. Thornburgh, and L. J. Bledsoe. 1979. Food web relationships of northern Puget Sound and the Strait of Juan de Fuca. US Interagency (NOAA, EPA) Energy/Environ. Res. Dev. Prog. Rep., Washington, D.C.
- Smith, B. D., G. A. McFarlane, and A. J. Cass. 1990. Movements and mortality of tagged male and female lingcod in the Strait of Georgia, British Columbia. *Trans.Am.Fish.Soc.* 119:813-824.
- Smith, K. L. and N. O. Brown. 1983. Oxygen consumption of pelagic juveniles and demersal adults of the deep-sea fish *Sebastes altivelis*, measured by depth. *Mar.Biol.* 76:325-332.
- Smith, P. E. 1995. Development of the population biology of the Pacific hake, *Merluccius productus* *Calif.Coop.Oceanic Fish.Invest.Rep* 36:144-152.

- Smith, S. E., D. W. Au, and C. Show. 1998. Intrinsic rebound potentials of 26 species of Pacific sharks. *Marine and Freshwater Research* 49:663-678.
- Sopher, T. R. 1974. A trawl survey of the fishes of Arcata Bay, California. Humbolt State University.
- Stanley, R. D., B. M. Leaman, L. Haldorson, and V. M. O'Connell. 1994. Movements of tagged adult yellowtail rockfish, *Sebastes flavidus*, off the West Coast of North America. *Fish.Bull.* 92:655-663.
- Starr, P. J. and J. Fargo. 2004. Petrale sole stock assessment for 2003 and recommendations for management in 2004. CSAS Res. Doc. 2004/036:92.
- Starr, P. J. and R. Haigh. 2000. Assessment of the Canadian longspine thornyhead (*Sebastolobus altivelis*) for 2000. Canadian Stock Assessment Secretariat, Nanaimo, British Columbia.
- Starr, R. M., J. H. Heine, J. M. Felton, and G. M. Cailliet. 2002. Movements of bocaccio (*Sebastes paucispinis*) and greenspotted (*Sebastes chlorostictus*) rockfishes in a Monterey submarine canyon: implications for the design of marine reserves. *Fishery Bulletin* 100:324-337.
- Stein, D. and T. J. Hassler. 1989. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (Pacific southwest): Brown rockfish, copper rockfish, black rockfish. U.S. Fish and Wildlife Service.
- Stein, D. L., B. N. Tissot, M. A. Hixon, and W. Barss. 1992. Fish-habitat associations on a deep reef at the edge of the Oregon continental shelf. *Fish.Bull.* 90:540-551.
- Steiner, R. 1978. Food habits and species composition of neritic reef fishes off Depoe Bay, Oregon. M.S. Thesis. Oregon State University.
- Stephens, A. 2011. 2011 rebuilding analysis for darkblotched rockfish. Pacific Fishery Management Council, Portland, OR, November, 2011.
- Stephens, A., O. Hamel, I. Taylor, and C. Wetzel. 2011. Status and future prospects for the darkblotched rockfish resource in waters off Washington, Oregon, and California in 2011. Pacific Fishery Management Council, Portland, OR, November, 2011.
- Stephens, A. and I. G. Taylor. 2013. Stock assessment and status of longspine thornyhead (*Sebastolobus altivelis*) off California, Oregon and Washington in 2013. Pacific Fishery Management Council, Portland, OR.
- Stepien, C. A. 1995. Population genetic divergence and geographic patterns from DNA sequences: examples from marine and freshwater fishes. Pages 263-287 *in* Evolution and the Aquatic Ecosystem Defining Unique Units in Population Conservation, volume 17. American Fisheries Society.
- Stepien, C. A., A. K. Dillon, and A. K. Patterson. 2000. Population genetics, phylogeography, and systematics of the thornyhead rockfishes (*Sebastolobus*) along the deep continental slopes of the North Pacific Ocean. *Can.J.Fish.Aquat.Sci.* 57:1701-1717.
- Stewart, I. J. 2006. Status of the U.S. English sole resource in 2005. *in* Volume 2: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.

- Stewart, I. J. 2008a. Rebuilding analysis for canary rockfish based on the 2007 stock assessment. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation.
- Stewart, I. J. 2008b. Status of the U.S. canary rockfish resource in 2007. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation Stock Assessments and Rebuilding Analyses.
- Stewart, I. J. 2008c. Updated U.S. English sole stock assessment: Status of the resource in 2007. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation Stock Assessments and Rebuilding Analyses.
- Stewart, I. J. 2009a. Rebuilding analysis for canary rockfish based on the 2009 updated stock assessment. Pacific Fishery Management Council, Portland, OR.
- Stewart, I. J. 2009b. Rebuilding analysis for yelloweye rockfish based on the 2009 stock assessment. Pacific Fishery Management Council, Portland, OR.
- Stewart, I. J. 2009c. Status of the U.S. canary rockfish resource in 2009 (Update of 2007 assessment model). Pacific Fishery Management Council, Portland, OR.
- Stewart, I. J., J. T. Thorson, and C. Wetzel. 2011. Status of the U.S. sablefish resource in 2011. Pacific Fishery Management Council, Portland, OR.
- Stewart, I. J., J. R. Wallace, and C. McGilliard. 2009. Status of the U.S. yelloweye rockfish resource in 2009. Pacific Fishery Management Council, Portland, OR.
- Stroud, R. K., G. H. Fiscus, and H. Kajimura. 1981. Food of the Pacific white-sided dolphin, *Lagenorhynchus obliquidens*, Dall's porpoise, *Phocoenoides dalli*, and northern fur seal, *Callorhinus ursinus*, off California and Washington. Fishery Bulletin 78(4):951-959.
- Sumida, B. Y. and H. G. Moser. 1984. Food and feeding of Bocaccio and comparison with Pacific hake larvae in the California current. Calif.Coop.Oceanic Fish.Invest.Rep. 25:112-118.
- Sydeman, W., M. M. Hester, J. A. Thayer, F. Gress, P. Martin, and J. Buffa. 2001. Climate change, reproductive performance and diet composition of marine birds in the Southern California Current System, 1969 - 1997. Progress in Oceanography 49:309-329.
- Tagart, J. V. 1988. Appendix D: Status of the yellowtail rockfish stocks in the International North Pacific Fishery Commission Vancouver and Columbia areas. *in* Status of the Pacific coast groundfish fishery through 1988 and recommended acceptable biological catches for 1989. Pacific Fishery Management Council, Portland, OR.
- Tagart, J. V. 1991. Population dynamics of yellowtail rockfish (*Sebastes flavidus*) stocks in the northern California to Vancouver Island region. Ph.D. Dissertation. University of Washington.
- Tagart, J. V. 1993. Status of the yellowtail rockfish resource in 1993. *in* Stock assessment and fishery evaluation: Appendices to the status of the Pacific Coast groundfish fishery through 1993 and recommended acceptable biological catches for 1994. Pacific Fishery Management Council, Portland, Oregon.

- Tagart, J. V., J. T. Golden, D. K. Kimura, and R. L. Demory. 1980. Evaluation of alternative trip limits for Pacific ocean perch. Unpublished report prepared for the Pacific Fishery Management Council, Portland, OR.
- Tagart, J. V., J. N. Ianelli, A. Hoffman, and F. R. Wallace. 1997. Status of the yellowtail rockfish resource in 1997. *in* Stock assessment and fishery evaluation: Status of the Pacific Coast groundfish fishery through 1997 and recommended acceptable biological catches for 1998. Pacific Fishery Management Council, Portland, Oregon.
- Tagart, J. V. and F. R. Wallace. 1996. Status of the yellowtail rockfish resource in 1996. *in* Stock assessment and fishery evaluation: Appendix volume 2 to the status of the Pacific Coast groundfish fishery through 1996 and recommended acceptable biological catches for 1997. Pacific Fishery Management Council, Portland, Oregon.
- Tagart, J. V., F. R. Wallace, and J. N. Ianelli. 2000. Status of the yellowtail rockfish resource in 2000. *in* Appendix to Status of the Pacific coast groundfish fishery through 2000 and recommended acceptable biological catches for 2001 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Tanasichuk, R. W., D. M. Ware, W. Shaw, and G. A. McFarlane. 1991. Variations in diet, daily ration, and feeding periodicity of pacific hake (*Merluccius productus*) and spiny dogfish (*Squalus acanthias*) off the Lower West Coast of Vancouver Island. *Can.J.Fish.Aquat.Sci.* 48(11):2118-2128.
- Taylor, C. A., W. Watson, T. Chereskin, J. Hyde, and R. Vetter. 2004. Retention of larval rockfishes, *Sebastes* near natal habitat in the Southern California Bight, as indicated by molecular identification methods. *CalCOFI Rep.* 45:152-166.
- Taylor, I. G. 2008. Modeling spiny dogfish population dynamics in the Northeast Pacific. University of Washington, Seattle, WA.
- Taylor, I. G. 2011. Rebuilding analysis for yelloweye rockfish based on the 2011 update stock assessment. NMFS, Seattle, WA, October 13, 2011.
- Taylor, I. G. and A. Stephens. 2013. Stock Assessment of Shortpine Thornyhead in 2013. Pacific Fishery Management Council, Portland, OR.
- Taylor, I. G. and C. Wetzel. 2011. Status of the U.S. yelloweye rockfish resource in 2011 (Update of 2009 assessment model). National Marine Fisheries Service, Northwest Fisheries Science Center, Seattle.
- Taylor, N., A. C. Hicks, I. G. Taylor, C. Grandin, and S. Cox. 2014. Status of the Pacific hake (whiting) stock in U.S. and Canadian waters in 2014 with a management strategy evaluation.
- Thomas, D. H. and A. D. MacCall. 2001. Bocaccio. Pages 162-164 *in* W. S. Leet, C. M. Dewees, R. Klingbeil, and E. J. Larson, editors. California's Living Marine Resources: A Status Report. Calif. Dept. Fish and Game.
- Thompson, J. E. 2006. Age, growth and maturity of the longnose skate (*Raja rhina*) for the US west coast and sensitivity to fishing impacts. Oregon State University, Corvallis, OR.

- Thompson, J. E. and R. W. Hannah. 2010. Using cross-dating techniques to validate ages of aurora rockfish (*Sebastes aurora*): estimates of age, growth and female maturity. *Environmental Biology of Fishes* 88:377-388.
- Tribuzio, C. 2009. Life history, ecology and population demographics of spiny dogfish in the Gulf of Alaska. University of Alaska, Fairbanks, AK.
- Tribuzio, C. A., V. F. Gallucci, and G. G. Bargmann. 2009. Reproductive biology and management implications for spiny dogfish in Puget Sound, Washington. *in* V. F. Gallucci, G. A. McFarlane, and G. G. Bargmann, editors. *Biology and Management of Dogfish Sharks*. American Fisheries Society.
- Tribuzio, C. A., G. H. Kruse, and J. T. Fujioka. 2010. Age and growth of spiny dogfish (*Squalus acanthias*) in the Gulf of Alaska: analysis of alternative growth models. *Fishery Bulletin* 108(2):119-135.
- Trumble, S. J. 1995. Abundance, movements, dive behavior, food habits, and mother – pup interactions of harbor seals (*Phoca vitulina richardsi*) near Monterey Bay, California. M.S. California State University Fresno.
- Tsou, T. and F. R. Wallace. 2006. Updated Rebuilding Analysis for Yelloweye Rockfish Based on Stock Assessment in 2006. *in* Volume 6: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Turnock, B. J., T. K. Wilderbuer, and E. S. Brown (T. Gulf of Alaska Groundfish Plan). 2005. Arrowtooth Flounder. North Pacific Fishery Management Council, P. O. Box 103136, Anchorage, AK 99510.
- Turnock, J., M. Wilkins, M. Saelens, and C. Wood. 1993. Status of West Coast petrale sole in 1993. *in* Appendix G. *In*: Status of the Pacific Coast groundfish fishery through 1993 and recommended biological catches for 1994: stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland, OR.
- Vega, N. M., V. F. Gallucci, L. Hauser, and J. Franks. 2009. Differences in growth in the spiny dogfish over a latitudinal gradient in the Northeast Pacific. *in* V. F. Gallucci, G. A. McFarlane, and G. G. Bargmann, editors. *Biology and Management of Dogfish Sharks*. American Fisheries Society.
- Vetter, R. D. and E. A. Lynn. 1997. Bathymetric demography, enzyme activity patterns, and bioenergetics of deep-living scorpaenid fishes: paradigms revisited. *Mar. Ecol. Prog. Ser.* 155:173-188.
- Vrooman, A. M. and P. A. Paloma. 1977. Dwarf hake off the coast of Baja California, Mexico. *California Cooperative Oceanic Fisheries Investigations Report* 19:67-72.
- Wakefield, W. W. 1990. Patterns in the distribution of demersal fishes on the upper continental slope off central California with studies on the role of ontogenetic vertical migration in particle flux. University of California, San Diego, CA.
- Wakefield, W. W. and K. L. J. Smith. 1990. Ontogenetic vertical migration in *Sebastes altivelis* as a mechanism for transport of particulate organic matter at continental slope depths. *Limnology and Oceanography* 35:1314-1328.

- Walker, P. A. and R. G. Hislop. 1998. Sensitive skates or resilient rays? Spatial and temporal shifts in ray species composition in the central and north-western North Sea between 1930 and the present day. *ICES J. of Mar. Sci.* 55:392-402.
- Wallace, F. R. 2002. Status of the yelloweye rockfish resource in 2001 for northern California and Oregon waters. *in* Appendix to the Status of the Pacific Coast Groundfish Fishery Through 2001 and Acceptable Biological Catches for 2002 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Wallace, F. R., Y. W. Cheng, and T. S. Tsou. 2008. Status of the black rockfish resource north of Cape Falcon, Oregon to the U.S.-Canadian border in 2006. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation Stock Assessments and Rebuilding Analyses.
- Wallace, F. R., A. Hoffman, and J. A. Tagart. 1999. Status of the black rockfish resource in 1999. *in* Appendix to the Status of the Pacific Coast Groundfish Fishery Through 1999 and Recommended Acceptable Biological Catches for 2000. Pacific Fishery Management Council, Portland, OR.
- Wallace, F. R. and J. V. Tagart. 1994. Status of coastal black rockfish stocks in Washington and northern Oregon in 1994. *in* Appendix F: Status of the Pacific Coast Groundfish Fishery Through 1994 and Recommended Acceptable Biological Catches for 1995. Pacific Fisheries Management Council, Portland, Oregon.
- Wallace, F. R., T. Tsou, T. Jagielo, and Y. W. Cheng. 2006. Status of yelloweye rockfish (*Sebastes ruberrimus*) off the U.S. West Coast in 2006. *in* Volume 6: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, OR.
- Wallace, J. and H. L. Lai. 2006. Status of the yellowtail rockfish in 2004. *in* Volume 3: Status of the Pacific Coast groundfish fishery through 2005, Stock assessment and fishery evaluation: Stock assessments and rebuilding analyses. Pacific Fishery Management Council, Portland, Oregon.
- Wallace, J. R. 2008a. Update to the status of yelloweye rockfish ( *Sebastes ruberrimus* ) off the U.S. West Coast in 2007. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation Stock Assessments and Rebuilding Analyses.
- Wallace, J. R. 2008b. Updated Rebuilding Analysis for Yelloweye Rockfish Based on the Stock Assessment Update in 2007. Status of the Pacific Coast groundfish fishery through 2007, Stock assessment and fishery evaluation.
- Wallace, J. R. 2009. 2009 darkblotched rockfish rebuilding analysis. Pacific Fishery Management Council, Portland, OR.
- Wallace, J. R. 2011. Rebuilding analysis for canary rockfish based on the 2011 updated stock assessment. Pacific Fishery Management Council, Portland, OR.
- Wallace, J. R. and J. M. Cope. 2011. Status update of the U.S. canary rockfish resource in 2011. National Marine Fisheries Service, Northwest Fisheries Science Center, Seattle.

- Wallace, J. R. and O. S. Hamel. 2009. Status and Future Prospects for the Darkblotched Rockfish Resource in Waters off Washington, Oregon, and California as Updated in 2009. Pacific Fishery Management Council, Portland, OR.
- Waples, R. S., A. E. Punt, and J. M. Cope. 2008. Integrating genetic data in marine resource management: how can we do it better? *Fish and Fisheries* 9:423-449.
- Weinberg, K. L. 1994. Rockfish assemblages of the middle shelf and upper slope off Oregon and Washington. *Fish.Bull.* 92:620-632.
- Weinberg, K. L., M. E. Wilkins, and T. A. Dark. 1984. The 1983 Pacific West Coast bottom trawl survey of groundfish resources: estimates of distribution, abundance, age and length composition. U.S. Dept. of Commerce.
- Weise, M. J. and J. T. Harvey. 2005. Impact of the California sea lion (*Zalophus californianus*) on salmon fisheries in Monterey Bay, California. *Fishery Bulletin* 103:685-696.
- Westrheim, S. J. 1975. Reproduction, maturation, and identification of larvae of some *Sebastes* (*Scorpaenidae*) species in the northeast Pacific Ocean. *J.Fish.Res.Board Canada* 32:2399-2411.
- Westrheim, S. J. and A. R. Morgan. 1963. Results from tagging a spawning stock of Dover sole, *Microstomus pacificus* *Pac.Mar.Fish.Comm.Bull.* 6:13-21.
- Wetzel, C. R. and A. E. Punt. In Press. Management strategy evaluation of the performance of data-moderate and data-limited assessment methods for U.S. west coast groundfish. *Fisheries Research*.
- Whitney, F. A., H. J. Freeland, and M. Robert. 2007. Persistently declining oxygen levels in the interior waters of the eastern subarctic Pacific. *Progress in Oceanography* 75:179-199.
- Wilkins, M. E. 1986. Development and evaluation of methodologies for assessing and monitoring the abundance of widow rockfish, *Sebastes entomelas* *Fish.Bull.* 84:287-310.
- Williams, E. H., A. D. MacCall, S. Ralston, and D. E. Pearson. 2000. Status of the widow rockfish resource in Y2K. *in* Appendix to Status of the Pacific coast groundfish fishery through 2000 and recommended acceptable biological catches for 2001 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Williams, E. H. and S. Ralston. 2002. Distribution and co-occurrence of rockfishes (family: *Sebastidae*) over trawlable shelf and slope habitats of California and southern Oregon. *Fisheries Bulletin* 100:836-855.
- Williams, E. H., S. Ralston, A. D. MacCall, D. Woodbury, and D. E. Pearson. 1999. Stock assessment of the canary rockfish resource in the waters off southern Oregon and California in 1999. *in* Status of the Pacific coast groundfish fishery through 1999 and recommended acceptable biological catches for 2000 (Stock Assessment and Fishery Evaluation). Pacific Fishery Management Council, Portland, OR.
- Wilson-Vandenberg, D., P. N. Reilly, and C. E. Wilson. 1996. Onboard sampling of the rockfish and lingcod commercial passenger fishing vessel industry in northern and central California, January through December 1994. *Marine Research Division Administrative Report*:96 pgs.

- Wood, C. C., K. S. Ketchen, and R. J. Beamish. 1979. Population dynamics of the spiny dogfish (*Squalus acanthias*) in British Columbia waters. *Journal of the Fisheries Research Board of Canada* 36:647-656.
- Wyllie Echeverria, T. 1987. Thirty-four species of California rockfishes: Maturity and seasonality of reproduction. *Fish.Bull.* 85:229-240.
- Yang, M. S. 1995. Food habits and diet overlap of arrowtooth flounder ( *Atheresthes stomias* ) and Pacific halibut ( *Hippoglossus stenolepis* ) in the Gulf of Alaska. Pages 205-223 *in*. Alaska Sea Grant College Program, University of Alaska, Anchorage, Alaska, 1995.
- Yang, M. S. and P. A. Livingston. 1985. Food habits and diet overlap of two congeneric species, *Atheresthes stomias* and *A. evermanni* , in the eastern Bering Sea. *Fish.Bull.* 84:615-623.
- Yoklavich, M. M., H. G. Greene, G. M. Cailliet, D. E. Sullivan, R. N. Lea, and M. S. Love. 2000. Habitat associations of deep-water rockfishes in a submarine canyon: an example of a natural refuge. *Fishery Bulletin* 98:625-641.
- Yoklavich, M. M., M. S. Love, and K. A. Forney. 2007. A fishery independent assessment of cowcod ( *Sebastes levis* ) in southern California's Cowcod Conservation Areas using direct observations from an occupied submersible. *Canadian Journal of Fisheries and Aquatic Sciences* 64(12):1795-1804.
- Zhou, S., S. Yin, J. T. Thorson, A. D. M. Smith, and M. Fuller. 2012. Linking fishing mortality reference points to life history traits: an empirical study. *Can.J.Fish.Aquat.Sci.* 69:1292-1301.
- Zimmerman, M. and P. Goddard. 1996. Biology and distribution of arrowtooth flounder, *Atheresthes stomias* , and Kamchatka flounders ( *A. evermanni* ) in Alaskan waters. *Fishery Bulletin* 94:358-370.