

# Evaluating potential responses of salmon populations exposed to acetylcholinesterase inhibiting insecticides



April 24, 2009  
Scott Hecht, NMFS

## Office of Protected Resources Pesticide Team

Angela Somma	Division Chief
Pamela Lawrence	NOAA GC
Paige Doelling	Ecotoxicologist
Kira Goetschius	Fisheries biologist
Tony Hawkes	Ecotoxicologist
Scott Hecht	Ecotoxicologist
Peter Johnsen	Fisheries biologist
Arlene Pangelinan	Coordinator

## NWFSC's Ecotoxicology and Environmental Fish Health Program



Tracy Collier	Director: Environmental Conservation Division
Nathaniel Scholz	Program manager
David Baldwin	
Julann Spromberg	
Kate Macneale	
Cathy Laetz	

## Objective:

Provide an overview of population models used in NMFS' biological opinions on the effects of acetylcholinesterase- inhibiting insecticides.

# Overview

Introduction

Conceptual framework

Life history and species


Model 1. Survival of subyearling salmon: acute lethality

Model 2. Growth of sub-yearling salmon: direct and indirect effects


Applications within opinions

Discussion

# Framework for Assessing Effects




# Framework for Assessing Effects (continued)



# Salmonid Lifecycle

Key Lifestage and Model  
Parameter:  
Subyearling survival




# What do these insecticides do?

## NEUROTOXICANTS


### Mode of action:

- disrupt neurotransmission
- inhibit an enzyme, acetylcholinesterase (AChE), by binding to it
- Nerve cells continue to fire





# Linking data across biological scales using models



# Two Models:

Acetylcholinesterase (AChE)  
Inhibition

Prey Availability



**Death of subyearling salmon  
from acute exposure**

Salmon Growth  
(Size)


First Year Survival  
(S1)

Lambda (Intrinsic population  
growth rates)

Population  
model



# Chinook Salmon Life-History



- Stream and ocean-type life-history
- Maximum Age 5 years
- Earliest female Reproduction Age 3
- Density Independent vital rates
- Slight changes in survival and reproduction could alter lambda and age distribution
- Impacts to first year survival produce the largest change in lambda




WDFW

# Transition matrix for life-history graph of Chinook salmon

Chinook Salmon

*O. tshawytscha*




$$A = \begin{bmatrix}
 0 & 0 & R3=a_{13} & R4=a_{14} & R5=a_{15} \\
 S1=a_{21} & 0 & 0 & 0 & 0 \\
 0 & S2=a_{32} & 0 & 0 & 0 \\
 0 & 0 & S3=a_{43} & 0 & 0 \\
 0 & 0 & 0 & S4=a_{54} & 0
 \end{bmatrix}$$

# Life-History Modeling


## Coho Salmon

*Oncorhynchus kisutch*




## Sockeye Salmon

*O. nerka*



## Chinook Salmon

*O. tshawytscha*



# Control Populations

Four “control” populations: Coho, Ocean- and stream type Chinook, sockeye

Chum and steelhead not modeled

Represent a population that we can compare to an impacted population

Comments:

Hatchery fish


Data taken from field studies

# Model 1:

Death of subyearling salmonids  
from acute exposure (96 hours)

# Model 1: Acute lethality

Death of subyearling ocean type Chinook salmon: LC50 and slope





# Model 1 Information

- Ocean- and stream-type Chinook, chum, coho, and sockeye
- All subyearlings exposed at sometime during their first year of survival
- Model repeats the lambda calculation 1000 times to integrate variability in transition values
- Lowest 96 hour salmonid LC<sub>50</sub> selected
- Sigmoid slope of 3.63 used (probit slope of 4.5)
- Input a range of concentrations to bracket dose-response

# Model 1: Inputs

Insecticide	96 h LC50 ug/L	96 h LC50 Slope
Carbaryl	250	3.63
Carbofuran	164	3.63
Methomyl	560	3.63

96 hour exposure concentration (ug/L):  
Ran multiple concentrations to bracket the LC50 to determine a population's response

# Output of models

- Percent change in Lambda between control and impacted population
- Percent mortality in population
- Survival rates of subyearling salmon ( $S_1$ ) in control and impacted populations

When is a change in lambda significant?

# Example of population model results from death of subyearling salmon

Table 74. Modeled output for Ocean-type Chinook salmon exposed to 4 d exposures of carbaryl, carbofuran, and methomyl reporting the impacted factors of survival as percent dead, lambda and standard deviation, and percent change in lambda compared to an unexposed population.

<u>Carbaryl</u>	0 µg/L	50 µg/L	100 µg/L	200 µg/L	250 µg/L	350 µg/L	500 µg/L	750 µg/L
% dead	0	0	3	31	50	77	93	98
Lambda (STD)	1.09 (0.1)	1.08 (0.1)	1.08 (0.1)	0.98 (0.09)	0.89 (0.08)	0.71 (0.06)	.53 (0.05)	0.36 (0.03)
% change in lambda	NA	NS	NS (-1)	-10	-18	-34	-52	-67
Threshold for significant change in lambda	-9.1 % ~ 190 µg/L							

# Model 1: Mixtures

- Used an additivity model based on dose-addition
- 4 day exposure
- Three scenarios:

Scenario	Carbamate	Application	Concentration ug/L
1 PRZM- EXAMS	Carbaryl	2 lbs/acre, 4 apps,	19
	Carbofuran	apples	35
	Methomyl	2 lb/acre, artichoke 0.9 lb/acre, 10 apps	88
2 GENEEC	Carbaryl	Corn	229
	Carbofuran	Corn	53
	Methomyl	Corn	49
3 Off-channel habitat	Carbaryl	5 lb/acre	335
	Carbofuran	1 lb/acre	67
	Methomyl	0.9 lbs/acre	17.1

<u>Scenario 1:</u> PRZM-EXAMS 24-h averages	Ocean-type Chinook	Stream-type Chinook	Sockeye	Coho
% dead	5	5	5	5
Lambda (STD)	1.07 (0.10)	0.99 (0.03)	1.00 (0.06)	1.01 (0.05)
% change in lambda	NS(-1)	NS(-1)	NS(-1)	NS(-2)
<u>Scenario 2:</u> GENEEC 90-d averages	Ocean-type Chinook	Stream-type Chinook	Sockeye	Coho
% dead	74	74	74	74
Lambda (STD)	0.74 (0.07)	0.72 (0.02)	0.74 (0.04)	0.66 (0.04)
% change in lambda	-32	-28	-27	-36
<u>Scenario 3:</u> Offchannel habitats 0.5 m deep	Ocean-type Chinook	Stream-type Chinook	Sockeye	Coho
% dead	89	89	89	89
Lambda (STD)	0.59 (0.05)	0.58 (0.02)	0.60 (0.03)	0.49 (0.03)
% change in lambda	-46	-42	-41	-52


NA denotes non applicable; NS denotes values less than one standard deviation of lambda expressed as the percent of lambda. (Calculated value, omitted when less than or equal to one)

## Model 2:

Effects to growth and subsequent size of subyearling salmon from:

1. Reduced feeding  
and
2. Reduced abundance of salmon prey

# Linking reductions in juvenile size to reductions in individual survival for ocean-type Chinook







# Model 2:

Acetylcholinesterase  
(AChE) inhibition of  
subyearlings

Reduction in prey  
availability

Salmon growth  
(Size )

First Year Survival  
(S1)

Population  
model

$$A = \begin{bmatrix} 0 & 0 & R3=a_{13} & R4=a_{14} & R5=a_{15} \\ S1=a_{21} & 0 & 0 & 0 & 0 \\ 0 & S2=a_{32} & 0 & 0 & 0 \\ 0 & 0 & S3=a_{43} & 0 & 0 \\ 0 & 0 & 0 & S4=a_{54} & 0 \end{bmatrix}$$

Lambda (Intrinsic  
population growth rates)

# Growth model control values

**Table 2. Species specific control parameters to model organismal growth and survival rates. Growth period and survival rate are determined from the literature data listed for each species.  $G_c$  and  $\alpha$  were calculated to make the basic model produce the appropriate size and survival values from the literature.**

	Chinook Stream-type <sup>1</sup>	Chinook Ocean-type <sup>2</sup>	Coho <sup>3</sup>	Sockeye <sup>4</sup>
days to run organismal growth model	184	140	184	168
growth rate % body wt/day ( $G_c$ )	1.28	1.30	0.90	1.183
$\alpha$ from equation S	-0.33	-1.99	-0.802	-0.871
Control Survival $\phi$	0.418	0.169	0.310	0.295


<sup>1</sup> Values from data in Healy and Heard 1984, Fast et al., 1988, Beckman et al., 2000, Knudsen et al., 2006

<sup>2</sup> Values from data in Healey and Heard 1984, Howell et al., 1985, Roni and Quinn 1995, Ratner et al., 1997, PSCCTC 2002, Green and Beechie, 2004, Johnson et al., 2007


<sup>3</sup> Values from data in Pess et al., 2002, Knudsen et al., 2002

<sup>4</sup> Values from data in Pauley et al., 1989, Gustafson et al., 1997, McGurk 2000


# Impact of chlorpyrifos on the AChE activity and feeding behavior of coho





# Impact of chlorpyrifos on the AChE activity of coho links to feeding behavior



# Ration (food uptake) can be linked to growth rate



# Creating a model to link AChE inhibition to reduced growth





## Step 1: Define...

- pesticide exposure
- magnitude of AChE inhibition
- timecourse for effect and recovery


# Creating a model to link AChE inhibition to reduced growth

On any given day...




Step 2: Connect the dots...

- AChE inhibition
- reduced feeding
- reduced ration











- Freshwater exposure to insecticides reduces abundances of salmonid prey
- Reductions of prey potentially affects growth and size which impacts survival


# Recovery: Abundances of 16 salmonid prey taxa





Compiled data supplied by van den Brink et al. 1996. ET&C (15):1143-1153

# Model 2: Modeling reductions in prey abundance






# Consequences of reduced growth on final weight of subyearling Chinook salmon




# Linking reductions in juvenile size to reductions in individual survival for ocean-type Chinook





**Noteworthy input parameters for Model 2**

# How we selected salmon prey sensitivity (LC50)










# Selection of a community prey slope

96 hour static bioassays with carbaryl on several salmonid prey taxa collected from PNW streams

Species	Survival EC50 ug/L	95% CI ug/L	Probit slope	Sigmoid slope
Stonefly- <i>Calineuria californica</i>	17.3	14.06-20.2	8.24	6.0
Mayfly- <i>Cinygma sp.</i>	11.1	7.7-13.9	4.10	3.0
Mayfly- <i>Ameletus sp.</i>	20.4	na	5.34	3.9
Caddisfly- <i>Brachycentrus americanus</i>	41.2	37.6-50.5	15.0	10.9
Caddisfly- <i>Psychoglypha sp. early instar</i>	30.3	25.0-40.4	9.10	6.6
Caddisfly- <i>Psychoglypha sp. Late instar</i>	61	55.6-68.54	7.50	5.5
Caddisfly- <i>Lepidostoma unicolor</i>	29	19.5-37.0	4.80	3.5

# Dose-response relationship for Ocean-type Chinook



**Figure 44.** Percent change in lambda for Ocean-type Chinook salmon following 4 d, 21 d, and 60 d exposures to carbaryl, carbofuran, and methomyl. Open symbols denote a percent change in lambda of less than one standard deviation from control population. Closed symbols represent a percent change in lambda of more than one standard deviation from control population.

**Table 80. Multiple application scenarios for carbaryl and methomyl and predicted percent change in lambdas for salmon populations**

	Carbaryl	Methomyl
Crop examples	Almonds, chestnuts, pecans, filberts, walnuts, pistachios	Sweet corn
Application rate	5 lbs a.i./acre	0.45 lbs a.i./acre
Number of applications/yr	4	10
Application interval	14 days	3 days
Method of application	Aerial (fine-medium droplet distribution)	aerial (fine-medium droplet distribution)
No-application Buffer	none	100 ft
Off-channel habitat characteristics	water depth = 0.5 m Initial average concentration 335 µg/L; 24 h exposure	water depth = 0.5 m Initial average concentration 8.55 µg/L; 96 h exposure
% change in Lambda		
Ocean-type Chinook	-19%	-8%
Stream-type Chinook	-15%	-6%
Sockeye	-16%	-7%
Coho	-18%	-8%

# Application of population modeling results

Comparison of change in lambdas with  
ESA-listed independent populations

Relationship of results to exposure

- Pesticides
- Subyearling salmonids

# Population modeling results are one line of evidence

- Results address risk hypotheses related to juvenile growth, survival, and prey availability
- Results of other non-modeled risk hypotheses also evaluated at the population level including:
  - survival of adults
  - swimming ability
  - olfaction-mediated behaviors (carbofuran)
  - starvation

# Risk at the Species level (ESU/DPS)

## Integration and Synthesis

- Status of the Species
- Environmental Baseline
- Cumulative Effects



The background of the image is a light beige or tan color with a subtle, mottled texture. It is covered with a repeating pattern of small, stylized fish. The fish are drawn in a simple, sketchy style using a brownish-gold color. They are arranged in a grid-like pattern, with two rows of fish in each horizontal section. The top row of fish in each section is oriented vertically, while the bottom row is oriented horizontally. The fish are small and appear to be swimming in various directions.

**Thank you**