

**Seasonal Management Areas
to Reduce Ship Strikes
of Northern Right Whales
in the Gulf of Maine**

by

Richard L. Merrick

December 2005

Recent Issues in This Series

- 05-01 **Results from the 2004 Cooperative Survey of Atlantic Surfclams.** By J.R. Weinberg, E.N. Powell, C. Pickett, V.A. Nordahl, Jr., and L.D. Jacobson. February 2005.
- 05-02 **Proceedings of a Workshop to Review and Evaluate the Design and Utility of Fish Mark - Recapture Projects in the Northeastern United States; October 19-21, 2004; Nonantum Resort, Kennebunkport, Maine.** By Workshop Organizing Committee (S. Tallack, editor, P. Rago, chairperson, T. Brawn, workshop coordinator, and (alphabetically) S. Cadrin, J. Hoey, and L. Taylor Singer. March 2005.
- 05-03 **Description of the 2004 Oceanographic Conditions on the Northeast Continental Shelf.** By M.H. Taylor, C. Bascuñán, and J.P. Manning. April 2005.
- 05-04 **40th SAW Assessment Report.** By 40th Northeast Regional Stock Assessment Workshop. April 2005.
- 05-05 **Effectiveness of a Square-Mesh Escape Panel in Reducing Finfish Bycatch in a Small-Mesh Bottom Trawl Used in the Longfin Inshore Squid (*Loligo pealeii*) Fishery.** By L. Hendrickson. June 2005.
- 05-06 **Use of the Historic Area Remediation Site by Black Sea Bass and Summer Flounder.** By M.C. Fabrizio, J.P. Pessutti, J.P. Manderson, A.F. Drohan, and B.A. Phelan. June 2005.
- 05-07 **Benthic Macrofauna and Associated Hydrographic Observations Collected in Newark Bay, New Jersey, between June 1993 and March 1994.** By L.L. Stehlik, S.J. Wilk, R.A. Pikanowski, D.G. McMillan, and E.M. MacHaffie. July 2005.
- 05-08 **Mortality and Serious Injury Determinations for Large Whale Stocks along the Eastern Seaboard of the United States, 1999-2003.** By T.V.N. Cole, D.L. Hartley, and R.L. Merrick. July 2005.
- 05-09 **NEFSC Bycatch Estimation Methodology: Allocation, Precision, and Accuracy.** By P.J. Rago, S.E. Wigley, and M.J. Fogarty. August 2005.
- 05-10 **41st SAW Assessment Summary Report.** By 41st Northeast Regional Stock Assessment Workshop. August 2005.
- 05-11 **Expanding Opportunities in Oceanic and Atmospheric Sciences III: Proceedings of the Third National Conference to Strengthen the Links among HBMSCUs, NOAA, Business, and Graduate Studies in Marine and Atmospheric Sciences.** By A. Jearld, Jr., and D. Peloquin, compilers. August 2005.
- 05-12 **Total Bycatch Estimate of Loggerhead Turtles (*Caretta caretta*) in the 2004 Atlantic Sea Scallop (*Placopecten magellanicus*) Fishery.** By K.T. Murray. August 2005.
- 05-13 **Assessment of 19 Northeast Groundfish Stocks through 2004: 2005 Groundfish Assessment Review Meeting (2005 GARM), Northeast Fisheries Science Center, Woods Hole, Massachusetts, 15-19 August 2005.** By R.K. Mayo and M. Terceiro, editors. September 2005.
- 05-14 **41st SAW Assessment Report.** By the 41st Northeast Regional Stock Assessment Workshop. September 2005.
- 05-15 **Economic Analysis of Alternative Harvest Strategies for Eastern Georges Bank Haddock [5Zjm; 551, 552, 561, 562].** By E.M. Thunberg, D.S.K. Liew, C.M. Fulcher, and J.K.T. Brodziak. October 2005.
- 05-16 **Northeast Fisheries Science Center Publications, Reports, and Abstracts for Calendar Year 2004.** By L. Garner. October 2005.
- 05-17 **Modification of Defined Medium ASP₁₂ for Picoplankter *Aureococcus anophagefferens*, with Limited Comparison of Physiological Requirements of New York and New Jersey Isolates.** By J.B. Mahoney. November 2005.

Seasonal Management Areas to Reduce Ship Strikes of Northern Right Whales in the Gulf of Maine

by

Richard L. Merrick

Postal Address: National Marine Fisheries Serv., Woods Hole Lab., 166 Water St., Woods Hole, MA 02543
E-Mail Address: Richard.Merrick@noaa.gov

**U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts**

December 2005

Northeast Fisheries Science Center Reference Documents

This series is a secondary scientific series designed to assure the long-term documentation and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review but no technical or copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

All documents issued in this series since April 2001, and several documents issued prior to that date, have been copublished in both paper and electronic versions. To access the electronic version of a document in this series, go to <http://www.nefsc.noaa.gov/nefsc/publications/series/crdlist.htm>. The electronic version will be available in PDF format to permit printing of a paper copy directly from the Internet. If you do not have Internet access, or if a desired document is one of the pre-April 2001 documents available only in the paper version, you can obtain a paper copy by contacting the senior Center author of the desired document. Refer to the title page of the desired document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate (*i.e.*, non-individualized) authorship, then contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

This document's publication history is as follows: manuscript submitted for review--October 31, 2005; manuscript accepted through technical review--November 9, 2005; manuscript accepted through policy review--December 8, 2005; and final copy submitted for publication--December 8, 2005. This document may be cited as:

Merrick, R.L. 2005. Seasonal management areas to reduce ship strikes of northern right whales in the Gulf of Maine. *U.S. Dep. Commer., Northeast Fish. Sci. Cent. Ref. Doc.* 05-19; 18 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.

TABLE OF CONTENTS

ABSTRACT.....	v
INTRODUCTION	1
METHODS	1
Datasets	1
Analysis.....	2
RESULTS.....	4
Seasonal Management Areas	4
Traffic Separation Scheme (TSS) Analyses	6
DISCUSSION	6
LITERATURE CITED	7
ACKNOWLEDGMENTS	7

List of Tables

Table 1.	Right, humpback, and fin whales sightings through 2003 in the original and two alternative configurations of the part of the Boston Traffic Separation Scheme	8
----------	---	---

List of Figures

Figure 1.	Northeast ALWTRP Management Areas through 2005	9
Figure 2.	Northeast ALWTRP Management Areas proposed as of 2005	9
Figure 3.	Right whale sightings in Northeast US waters, 1960-2003	10
Figure 4.	Right whale groups (2+ animals) sighted in Northeast US waters during March-July, 1999-2003.....	11
Figure 5.	Right whale groups sighted and DAM buffer zones for Northeast US waters, March-July, 1999-2003	12
Figure 6.	Gulf of Maine combined seasonal management area, right whale groups, and DAM buffer zones for NortheastUS waters, March-July, 1999-2003.....	13
Figure 7.	Seasonal management areas and right whale sightings for Northeast US waters April-July, 1999-2003	14
Figure 8.	Seasonal management areas and right whale sighting densities for Northeast US waters March-July, 1999-2003.....	15
Figure 9.	Seasonal management areas and right whale sightings densities for Northeast US waters June, 1999-2003.....	16
Figure 10.	Existing Boston Traffic Separation Scheme (TSS) and associated right whale sightings.....	17
Figure 11.	Alternative Boston Traffic Separation Scheme (TSS) and associated right whale sightings.....	18

ABSTRACT

Northeast Fisheries Science Center staff evaluated the spatial and temporal distribution of northern right whales (*Eubalena glacialis*) within the Gulf of Maine to develop potential management areas to reduce ship strikes of these whales. The following management areas coupled with routing and speed restrictions should decrease the risk of right whale ship strikes in the Gulf of Maine:

1. A Cape Cod Bay Seasonal Management Area (SMA) to be coterminous with the existing Cape Cod Bay critical habitat (January 1- May 15).
2. A Gulf of Maine SMA extending eastward from Massachusetts Bay (70.50° W) to the Hague Line, and southward from 42.50° N to the Cape Cod Bay SMA and the southern edge of the Great South Channel critical habitat area. Time period for this SMA would be March 1 - July 31. If further temporal structuring of the SMA is desired then it could be split as follows:
 - a. A Western Seasonal Management Area extending from Massachusetts Bay (70.50° W) eastward to 69.40° W, and from 42.50° N southward to the Cape Cod Bay SMA and Great South Channel critical habitat areas (March 1 - April 30)
 - b. An Eastern Seasonal Management Area extending from 69.75° W eastward to the Hague line, and from 42.50° N southward to the southern edge of the Great South Channel critical habitat area (April 1 - July 31)
3. A modified Boston Traffic Separation Scheme (TSS) which includes moving the existing East-West leg of the lanes further north, and adding a new North-South TSS as a spur from the existing TSS. Realignment of the existing Boston TSS has the potential to reduce interactions between ships and large whales along the northern portion of the TSS by 50% or more by directing shipping traffic away from seasonal large whale concentrations within the existing TSS.

INTRODUCTION

NOAA Fisheries has used seasonal and spatial gear restrictions to reduce interactions between northern right whales (*Eubalaena glacialis*) and commercial fixed gear fisheries as part of the Atlantic Large Whale Take Reduction Plan (ALWTRP; <http://www.nero.noaa.gov/whaletrp/>). The most recent application of these restrictions was the Seasonal Area Management (SAM) and Dynamic Area Management (DAM) zones implemented in March 2002. Design of these zones was based in part on analyses conducted by the Northeast Fisheries Science Center (Clapham and Pace 2001, Merrick et al. 2001). Based on these analyses, Management Areas (MAs) were established to regulate fishery gear deployed in the Northwest Atlantic Ocean in areas occupied seasonally by right whales (Figure 1).

Since implementation of the MAs to reduce fishing gear interactions with right whales, a strategy has also been developed to reduce ship strikes of right whales in the same waters. In theory, MAs to reduce gear and ship strike interactions could be similar in time and space. Nonetheless, it would be prudent to review the design of the existing MAs before extending their application under the ship strike strategy. It may be necessary to modify the original fishing gear based MAs to meet the needs of ship strike reduction.

The goal of this paper is to evaluate Seasonal Management Areas for reducing ship strikes of right whales. The original MAs are updated with additional data, and then modified to accommodate shipping characteristics. An initial analysis is also provided of an alternative alignment of the Boston, MA Traffic Separation Scheme that allows shipping to continue but provides increased protection to right whales.

METHODS

Datasets

Right whale sightings data were obtained from the North Atlantic Right Whale database (NARW) curated at the University of Rhode Island (URI), which includes survey data from a variety of sources back to the 1960s. This dataset comprises all right whale sighting information from reliable sources collected on the East Coast of the US since at least the 1960s, and includes observations from fine- and broad-scale aerial and vessel surveys conducted in a variety of projects. Major sources of data for New England waters include the Cetacean and Turtle Assessment Program (1978-1982), the South Channel Ocean Productivity Experiment (1988-1989), and recent NOAA surveys (1999-2003). The NARW dataset is continually updated. The NARW dataset used in this analysis was distributed in December 2004, and includes sighting records through fall 2003. Areas surveyed (including track lines) can be found at <http://www.marinegis.org/series1/index.html>.

Right whale sighting records from ~1960 through 2003 collected within Northeast waters between latitude 40° and 45° N, and eastward of 71° W to the Hague Line were analyzed.

Analysis of the Boston Traffic Separation Scheme also included humpback (*Megaptera novaeangliae*) and fin whale (*Balaenoptera musculus*) sightings.

Recent NOAA Fisheries Service data

The only recent unbiased, broad-scale survey observations included in the NARW database are those from NOAA aerial surveys. Directed surveys for North Atlantic right whales were flown during March-November, 1999-2003 in the area from south of Nantucket, MA northward to the Bay of Fundy, and from the New England coast eastward to the Hague Line. Surveys were accomplished using two high-wing, twin-engine fixed wing aircraft. One was a NOAA DeHaviland Twin Otter, and the other was a chartered amphibious Grumman Widgeon (1999-2001) or Goose (2002-2003). Surveys were flown at ca. 230 m (1999-2000) or 310 m (2001-2003) altitude, and 100 kts (185 km/hr) by at least one aircraft on virtually all days when sighting conditions were Beaufort 3 or better. Flights typically were not conducted when visibility was obscured (snow, rain, or fog).

During 1999-2000, flights were either flown along tracklines developed for SCOPEX in the Great South Channel area, or directed towards areas of reported or suspected right whale concentrations. During 2001-2003, one aircraft flew a monthly series of east-west track lines over the entire study area, while the second aircraft focused on the Great South Channel and other offshore areas.

Analysis

Defining Seasonal Management Areas

The MAs developed in Merrick et al (2001) provided the baseline for the ship strike MAs, and the additional analyses focused on refining the temporal and spatial extent of the East and West Seasonal MAs (Figure 1) previously developed for the ALWTRP (Merrick et al. 2001). The Cape Cod Bay MA was not analyzed further, but was assumed to be coterminous with the existing Cape Cod Bay critical habitat management area defined as part of the northern right whale recovery plan and the ALWTRP.

Additional analyses¹ of right whale sightings prompted a potential spatial adjustment of the western Seasonal MA (Figure 2) to better reflect recent data on right whale seasonal distribution. This report provides a further update using the full 2003 dataset.

Analysis of DAM actions during 1999-2003: The predictability of right whale concentrations distinguish Dynamic from Seasonal MAs. Dynamic MAs were developed to address unpredictable concentrations (which could be because of a lack of knowledge or because the whales truly do represent an unusual aggregation), while seasonal MAs were designed to cover areas of known right whale concentrations. The Cape Cod Bay and Great South Channel

¹ Memo from Richard Merrick to Mary Colligan dated 31 October 2003. Avail. Protected Species Branch, NEFSC, 166 Water Street, Woods Hole, MA 02543

Management Areas are examples of Seasonal MAs, because they were drawn based on historical sightings of right whales in these areas.

Dynamic MAs may be candidates for Seasonal MAs if whales are observed in the same area during the same season during three or more years; the area would then be considered to have predictable concentrations of whales. If, however, whales were observed in 1-2 years or in different months in different years, then the area would not be considered as a candidate Seasonal MA. For analysis, this means that Dynamic MAs needed to be created for each sighting of 2+ whales during each analysis year, and boundaries drawn around applicable zones for the aggregate of years. In summary, the process was to:

1. Identify right whale sightings which met the trigger criteria (a sighting with 2+ whales)
2. Define 15 nm radius buffer around the core area
3. Join, as necessary, the circular buffer zones

Dynamic event were triggered whenever there was a sighting of 2+ right whales sufficiently close to each other to achieve a density of 0.04 right whales/nm² (Clapham and Pace 2001). Revised Seasonal MAs were then drawn to encapsulate all Dynamic areas exhibiting such whale concentrations during the same season for three or more years.

Analyses focused on sightings during March-July, because this is the time period when whales move out of Cape Cod Bay into offshore areas.

Gridded Analysis: Ideally, right whale sightings in an area would be corrected for sightings effort to evaluate spatial and temporal distributions. However, this is difficult given the varying levels and quality of the sightings survey efforts; at best, adjusting for effort would entail a long and complex process that would eliminate from consideration any sightings without associated effort data. GIS analyses by New England Aquarium staff (<http://www.marinegis.org/series1/index.html>) using only “good survey effort and sightings data” disclosed only a single sighting of one animal north and west of Cape Cod Bay during the month of May from 1978 through 1999. In contrast, during this same time period the URI right whale database contains a total of 19 sightings (mostly from surveys without associated effort data) representing 25 individual animals.

As an alternative, one can grid the data and then normalize the sightings within each grid cell (dividing by the cell’s area). The normalized values (“densities”) can then be compared between cells to evaluate the overall distribution of animals. Comparing relatively close cells should compensate, to a limited degree, for the lack of uniform survey effort over the entire Gulf of Maine. This assumes that sighting effort is relatively consistent for the area. For this reason, analyses were restricted to the March-July 1999-2003 period, the time interval when the NEFSC conducted systematic right whale surveys of the Gulf of Maine complementing the systematic surveys of Cape Cod Bay conducted by the Provincetown Center for Coastal Studies.

Using these datasets, ARCVIEW was used to generate a monthly series of “density” plots in the SAM West area:

1. A rectangular grid was generated for the area between 40° and 45° N latitude from the Hague Line westward to the New England coast or 73° W longitude. Cells throughout the area were defined as 0.25 degree longitude and latitude squares.
2. Counts of individual right whales sighted in surveys conducted during 1999-2003 were then summed for the March-July period and by month for each grid cell.
3. Of the total 1,994 cells, 983 had no right whale observations during the 5-year, 5-month period. Four cells had very high right whale densities (e.g., Cape Cod Bay in March-April) that skewed the distribution. The remaining cells were all below 100 animals per cell for the entire period. Cells were grouped binned into six bins based on right whale densities: 0, 1, 2-4, 5-14, 15-99, and >99 sightings per cell for the period.
4. Using these categories, the data were plotted on a gridded, ARCVIEW base map of the Gulf of Maine using shaded values to distinguish between ranges of values.

The resulting monthly plots were then compared to the Seasonal MAs prepared as part of the DAM analysis to provide an alternative assessment of the area.

Traffic Separation Scheme (TSS) Analyses

The existing TSS was designed prior to the availability of much of the data included in the analysis presented in this paper. Nonetheless, the existing TSS was evaluated, and an alternative configuration provided that reduces the potential for large whale ship strikes. This configuration was designed to address two issues: (a) to move the TSS to an area with the lowest densities of whales, and (b) to design a spur or “cutout” from the TSS to allow vessels to exit and enter the TSS from the north.

The analysis involved moving the lanes northward until the number of potential interactions (defined as the number of sightings which fell within the TSS) was minimized. This is only meant as a preliminary evaluation of the routing, as alternative evaluations are currently underway by staff of the Stellwagen Banks National Marine Sanctuary (pers. comm., David Wiley, 12 October 2005), and by the US Coast Guard as part of Port Access Route Study for the Boston area requested by NOAA.

RESULTS

Seasonal Management Areas

Simulated Dynamic MAs

The URI database includes 7,698 right whale sighting events (13,118 animals; Figure 3) recorded during 1960-2003 in Northeast waters. During March-July 1999-2003, there were 2,396 right

whale sighting events in Northeast U.S. waters, with 1,301 events recorded outside of Cape Cod Bay. Of these events, there were 155 sightings of groups of 2+ animals observed outside of Cape Cod Bay (Figure 4).

The DAM zones drawn using the five seasons of data are shown in Figure 5. Most of the zones, (particularly the overlapping zones) occur from Cape Cod Bay eastward across the Great South Channel and on the northern edge of Georges Bank. Temporally isolated (less than 3 years) sightings of 2+ animal groups occurred southwest of the Great South Channel ($n = 2$), west of the Cape Cod Bay critical habitat area ($n = 3$), and north of Wilkinson Basin ($n = 2$). A cluster of repeated sightings occurred in the Cashes Ledge area ($n = 11$); 9 of these occurred in 2000 and 2 in 2003. Because these sightings do not suggest a pattern of repeated inter-annual use of the area, they do not qualify for considerations as a Seasonal MA.

Consolidating the DAMs to produce single zones which included 3 or more seasons of overlapping buffers resulted in the rectilinear shape shown in Figure 6. This area encompasses 137 (88%) of the Gulf of Maine 2+ animal sightings during March-June 1999-2003. It also includes 1,095 (84%) of the 1,301 Gulf of Maine right whale sighting events outside of Cape Cod Bay for the March-July 1999-2003 period.

The rectilinear Gulf of Maine Seasonal MA includes the existing Great South Channel right whale critical habitat area. This MA expands protection for right whales during March-July in two general areas:

1. North (Stellwagen Bank) and east (Wilkinson Basin) of the Cape Cod Bay critical habitat and,
2. North and east (northern edge of Georges Bank) of the Great South Channel MA.

Sighting data were also analyzed for the months of April through July (Figure 7), the period after most right whales have moved out of Cape Cod Bay but still remain in the Gulf of Maine prior to moving to the Bay of Fundy. The analyses suggest that whales occur throughout the Gulf of Maine Seasonal MA during this time; alternatively, the zone could be split into a West zone during March through April (eastward to 69.40° W) and an East zone during April – July (eastward from 69.75° W).

Gridded Data

The gridded results for the March-July ($n = 2,396$ sightings; Figure 8) suggest that the Cape Cod Bay and Gulf of Maine Seasonal Management Areas encompass most of the animals sighted. The only obvious discrepancy is to the north in the central Gulf of Maine where right whale sightings occurred in June 2000 and 2003 (Figure 9). Otherwise, the distributional patterns generally follow the known seasonal movements of right whales through the Gulf of Maine. During March, animals are located at relatively high densities only within Cape Cod Bay, though scattered sightings exist outside the Bay. Animals begin moving out of the Bay in April, with densities still high there, but also moderately high to the north of the Bay, and in isolated areas of the Great South Channel. By May, most of the activity has shifted out of the Bay and into the

Great South Channel. Densities remain high there through June (Fig. 9), and then in July animals depart the Bay, reaching temporarily high densities north of Cape Cod Bay in July, before animals move to the Bay of Fundy.

Traffic Separation Scheme (TSS) Analyses

The current or original TSS has the unfortunate property of routing vessels through high whale density areas north of Race Point, MA (Figures 7 and 10). However, by moving the East-West portion of the TSS somewhat north, it is possible to avoid many of these animals (Figure 11). Analyses of the sighting data available through 2003 from the URI dataset suggest that encounter rates could be decreased by 46%, 58%, and 66% for right, humpback and fin whales, respectively (Table 1), by such a move. Also shown in Table 1 and Figure 11 is an alternative alignment of the East-West portion of the TSS proposed by the Stellwagen Bank National Marine Sanctuary (“NOS alternative”). Both alignments show the potential for significant reduction of ship strike risk for large whales.

DISCUSSION

Extension of the existing gear management areas (Cape Cod Bay, SAM West, SAM East, and the Great South Channel) in time and space would provide greater protection for right whales. Such an extension would also protect the concentrations of right whales observed in the Gulf of Maine during spring-summer in recent years. This includes groups of animals observed on Stellwagen Bank in April, in Wilkinson Basin in April-May, and in the northern Great South Channel in April-May. Right whale concentrations to the north of SAM East appear to be less predictable, but occur in areas with a lower likelihood of gear interactions due to the limited fishing there.

The simulated DAM and gridded density results both suggested that SAM West (Figure 1) should be expanded to the west and south to provide greater protection for right whales. The SAM East’s southern boundary should also be modified to better encompass the known distribution of right whales on Georges Bank (e.g., Figure 8). The seasonal division on May 1 between the original SAM East and West zones is a significant problem in that aggregations of right whales have been repeatedly sighted in the eastern part of SAM West (Wilkinson Basin), after it has been reopened. This problem could be remedied by extending the SAM West Area to the end of May, or by extending the western boundary of SAM East to 69.75°W.

Finally, the existing Boston Traffic Separation Scheme could be realigned to pass through areas with lower concentrations of right, humpback and fin whales (Figure 11). Such a realignment would significantly reduce the likelihood of ship strikes in the area. Additional analyses being conducted cooperatively between Stellwagen Bank National Marine Sanctuary and NEFSC staff, and by the US Coast Guard as part of a current Boston Port Access Route Study, will further refine this alignment.

LITERATURE CITED

Clapham, P. J., and R. M. Pace, III. 2001. Defining Triggers for Temporary Area Closures to Protect Right Whales from Entanglements: Issues and Options. Northeast Fish. Sci. Cent. Ref. Doc. 01-06; 28 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.

Merrick, R. L., P. J. Clapham, T. V. N. Cole, P. Gerrior, and R. M. Pace, III. 2001. Identification of Seasonal Area Management Zones for North Atlantic Right Whale Conservation. Northeast Fish. Sci. Cent. Ref. Doc. 01-14; 18 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.

ACKNOWLEDGMENTS

The efforts of the survey teams and pilots (NOAA, Provincetown Center for Coastal studies, and commercial pilots) are gratefully acknowledged. Without their efforts over the past three years this report would not have been possible to produce. Recent surveys by NOAA were authorized under MMPA/ESA permit 775-1600. I would also like to thank Dr. Fred Serchuk, Dr. Greg Silber, David Wiley, and David Gouveia for reviews of earlier versions of this report.

Table 1. Right, humpback, and fin whales sightings through 2003 in the original and two alternative configurations of the part of the Boston Traffic Separation Scheme which begins due east of Race Point and is aligned first northwest-southeast and then west-east. Original refers to the existing Traffic Separation Scheme (Fig. 10), NOS Alt. Refers to alternative routing suggested by NOS², and NMFS Alt. refers to NMFS alternative Traffic Separation Scheme (Fig. 11).

Lanes	Right whale			Humpback whale			Fin whale		
	Original	NOS Alt	NMFS Alt.	Original	NOS Alt.	NMFS Alt.	Original	NOS Alt.	NMFS Alt.
East-West	221	134	121	227	111	95	225	87	76
North-South ³	>>0	5	5	>>0	4	4	>>0	8	8
Total	221+	139	126	227+	115	99	225+	95	84

² NOS Alternative as provided by staff of Stellwagen Bank National Marine Sanctuary

³ Note that because under the present alignment of the TSS, commercial vessels can move North-South across the breadth of the Georges Bank, all of these vessels have the potential to pass through areas seasonally occupied by large whales. Thus, the potential for vessel interactions, while significantly greater than 0, is not easily calculated




Figure 1. Northeast ALWTRP Management Areas through 2005.




Figure 2. Northeast ALWTRP Management Areas proposed as of 2005.

Right Whales Sighted in Northeast US Waters 1960-2003




Figure 3. Right whale sightings in Northeast US waters, 1960-2003

Right Whale Groups Sighted in Northeast US Waters March-July, 1999-2003




Figure 4. Right whale groups (2+ animals) sighted in Northeast US waters during March-July, 1999-2003.

Right Whale Groups Sighted and DAM Buffer Zones for Northeast US Waters March-July, 1999-2003




Figure 5. Right whale groups sighted and DAM buffer zones for Northeast US waters, March-July, 1999-2003.

**Seasonal Management Areas,
Right Whale Groups Sighted and DAM Buffer Zones for
Northeast US Waters March-July, 1999-2003**




Figure 6. Gulf of Maine combined seasonal management area, right whale groups, and DAM buffer zones for Northeast US waters, March-July, 1999-2003.

Seasonal Management Areas, and Right Whale Sightings for Northeast US Waters April-July, 1999-2003




Figure 7. Seasonal management areas and right whale sightings for Northeast US waters April-July, 1999-2003.




Figure 8. Seasonal management areas and right whale sightings densities for Northeast US waters, March-July, 1999-2003.

Seasonal Management Areas, and Right Whale Sightings Densities for Northeast US Waters June, 1999-2003




Figure 9. Seasonal management areas and right whale sightings densities for Northeast US waters, June, 1999-2003.

Current Boston Traffic Separation Schemes (TSS) & Associated Right Whale Sightings (March-July; 1960-2003)




Figure 10. Existing Boston Traffic Separation Scheme (TSS) and associated right whale sightings.

Alternative Boston Traffic Separation Schemes (TSS) & Associated Right Whale Sightings (March-July; 1960-2003)




Figure 11. Alternative Boston Traffic Separation Scheme (TSS) and associated right whale sightings.

Procedures for Issuing Manuscripts in the *Northeast Fisheries Science Center Reference Document (CRD) Series*

Clearance: All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If any author is not a federal employee, he/she will be required to sign an "NEFSC Release-of-Copyright Form." If your manuscript includes material lifted from another work which has been copyrighted, then you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

Organization: Manuscripts must have an abstract and table of contents, and — if applicable — lists of figures and tables. As much as possible, use traditional scientific manuscript organization for sections: "Introduction," "Study Area"/"Experimental Apparatus," "Methods," "Results," "Discussion" and/or "Conclusions," "Acknowledgments," and "Literature/References Cited."

Style: The CRD series is obligated to conform with the style contained in the current edition of the *United States Government Printing Office Style Manual*. That style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the *CBE Style Manual*. Manuscripts should be prepared to conform with these style manuals.

The CRD series uses the American Fisheries Society's guides to names of fishes, mollusks, and decapod crustaceans, the Society for Marine Mammalogy's guide to names of marine mammals, the Biosciences Information Service's guide to serial title abbreviations, and the International Standardization Organization's guide to statistical terms.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of cited works. Personal communications must include date, full name, and full mailing address of the contact.

Preparation: Type a clean/neat, single-spaced version of the document. The document must be paginated continuously from beginning to end and must have a "Table of Contents." Begin the preliminary pages of the document — always the "Table of Contents" — with page "iii." Begin the body of the document — normally the "Introduction" — with page "1," and continuously paginate all pages including tables, figures, appendices, and indices. You can insert blank pages as appropriate throughout the document, but account for them in your pagination (*e.g.*, if your last figure ends on an odd-numbered/right-hand page such as "75," and if your next page is the first page of an appendix, then you would normally insert a blank page after the last figure, and paginate the first page of the appendix as "77" to make it begin on an odd-numbered/right-hand page also). Forward the final version to the Editorial Office as both a paper copy and electronically (*i.e.*, e-mail attachment, 3.5-inch floppy disk, high-density zip disk, or CD). For purposes of publishing the CRD series only, the use of Microsoft Word is preferable to the use of Corel WordPerfect.

Production and Distribution: The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages (pages "i" and "ii") of the document, then combine those covers and preliminary pages with the text that you have supplied. The document will then be issued online.

Paper copies of the four covers and two preliminary pages will be sent to the sole/senior NEFSC author should he/she wish to prepare some paper copies of the overall document as well. The Editorial Office will only produce three paper copies (*i.e.*, two copies for the NEFSC's libraries and one copy for its own archives) of the overall document.

A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the online version of the document. The sole/senior NEFSC author of the document will receive a list of those so notified.

Research Communications Branch
Northeast Fisheries Science Center
National Marine Fisheries Service, NOAA
166 Water St.
Woods Hole, MA 02543-1026

**MEDIA
MAIL**

Publications and Reports of the Northeast Fisheries Science Center

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of living marine resources for the benefit of the nation through their science-based conservation and management and promotion of the health of their environment." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (*e.g.*, anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media. Currently, there are three such media:

NOAA Technical Memorandum NMFS-NE -- This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document -- This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review, but no technical or copy editing.

Resource Survey Report (formerly *Fishermen's Report*) -- This information report is a quick-turnaround report on the distribution and relative abundance of selected living marine resources as derived from each of the NEFSC's periodic research vessel surveys of the Northeast's continental shelf. There is no scientific review, nor any technical or copy editing, of this report.

OBTAINING A COPY: To obtain a copy of a *NOAA Technical Memorandum NMFS-NE* or a *Northeast Fisheries Science Center Reference Document*, or to subscribe to the *Resource Survey Report*, either contact the NEFSC Editorial Office (166 Water St., Woods Hole, MA 02543-1026; 508-495-2228) or consult the NEFSC webpage on "Reports and Publications" (<http://www.nefsc.noaa.gov/nefsc/publications/>).

ANY USE OF TRADE OR BRAND NAMES IN ANY NEFSC PUBLICATION OR REPORT DOES NOT IMPLY ENDORSEMENT.