

Contaminants Report Number: R6&R8/725C/11



**U.S. FISH & WILDLIFE SERVICE
REGION 6**

ENVIRONMENTAL CONTAMINANTS PROGRAM



**Characterizing Contaminant Exposure of Mountain Plovers
on Wintering Grounds in California and Breeding
Grounds in Colorado, Wyoming, and Montana**



Photo Credit – Fritz Knopf

By
Kimberly Dickerson, Karen J. Nelson, and Catherine Zeeman, Ph.D

FFS# 6N56 and 6F51
DEQ Project ID: 200660002.3

July 2011

Abstract

The mountain plover (*Charadrius montanus*) is a shorebird that primarily inhabits dry upland sites, such as shortgrass prairie and shrub-steppe landscapes during both wintering and breeding seasons. Nesting occurs primarily in Montana, Wyoming, Colorado, and New Mexico. The birds winter in the Central and Imperial Valleys of California as well as Arizona, Texas, and Mexico. Conversion of the mountain plovers' native grassland habitat to agricultural lands in both breeding and wintering areas has resulted in these birds using farmlands for feeding and roosting. Both the Central and Imperial Valleys report year round agriculture with over 809,400 ha (2 million acres) in production; but, these habitat preferences expose wintering mountain plovers to contaminants associated with agricultural practices, including current use pesticides and their adjuvants, persistent chemicals from historic applications, inorganic elements from irrigation drainwater, and chemicals present in the irrigation water. To evaluate the exposure of mountain plover to contaminants, we collected soil and food items at both their wintering and breeding grounds in 2006. We also collected eggs (2006-2008) to determine exposure of developing embryos to contaminants transferred by adults.

Some elevated levels of inorganic elements were found in soil and terrestrial insect food items of the mountain plover at both wintering and breeding sites but none of these was elevated in the eggs. Organic contaminants in soil and insects collected from breeding grounds were very low. However, several organic contaminants were elevated in soil and insect samples from the wintering grounds, which were also elevated in the eggs. These elevated levels of contaminants in eggs were likely the result of accumulation in adults ingesting insects and soils at the wintering grounds and then transferring them to the embryo when egg-laying occurred.

Specifically, residues of DDT (o,p' and p,p') and its metabolites DDD (o,p' and p,p') and DDE (o,p' and p,p') were detected in virtually all of the soil samples from the wintering grounds. In the Imperial Valley soils, concentrations of total DDT ranged from slightly below to more than ten-times above the Ecological Soil Screening Level of 22 ng/g dry weight (dw) for protection of avian species. Dieldrin and aldrin were also detected but were below level established for the protection of avian species. DDE was detected in all invertebrate samples, reflecting the widespread occurrence of DDT in general. The DDE concentration in invertebrates tended to be between 0.5 to 5.0 times the concentration in corresponding soil samples and total DDT concentrations were higher in invertebrates than the soil collected from the same location demonstrating the potential for these contaminants to bioaccumulate in the tissues of soil invertebrates, which are then ingested by mountain plover.

DDE was detected in all eggs collected from breeding sites in 2006. Concentrations ranged from 5 ng/g to 5,400 ng/g wet weight (ww) for eggs from Montana (n = 7), from 10 to 930 ng/g ww for eggs from Wyoming (n = 7), and from 11.0 - 11,500 ng/g ww for eggs from Colorado (n = 5). Furthermore, two eggs from Montana had DDE concentrations (3,000 and 5,400 ng/g ww) that were at least as high as those found to adversely affect productivity in pelicans. Two eggs from Colorado were near this concentration as well (2,700 and 2,900 ng/g ww) with an

additional egg having a concentration (11,500 ng/g ww) approaching that known to affect bald eagle reproduction.

In 2007, mountain plover eggs were collected from breeding sites in Colorado and Wyoming. All eggs had detectable levels of p,p'-DDE (range = 35 – 51,200 ng/g ww, n = 6), with the highest concentration found in an egg from Colorado. This concentration (51,200 ng/g ww) is well above the 300 ng/g shown to cause colony failure in brown pelicans and above the 16 ug/g ww level shown to cause complete nest failure in eggs of bald eagles. This egg also had detectable concentrations of o,p'-DDT, dieldrin, heptachlor epoxide, oxychlorane, and PCBs, which, if acting additively, increases the potential for adverse effects.

In 2008, eggs were collected from BLM lands, Phillips County, Montana (n = 8). DDT or one of its metabolites was detected in seven of the eight mountain plover eggs. DDE concentrations in two eggs (47,500 ng/g ww and 53,200 ng/g ww) were very high. It is possible these two eggs were not viable because of the DDE concentrations and the likelihood of either additive or synergistic effects from the other detected organochlorine residues.

Archived mountain plover eggs (n = 13) collected in 2003 were submitted for organic chemical analyses to demonstrate the presence of organic compounds and that exposure to mountain plover is ongoing. DDT was detected in three eggs and DDE was detected in all eggs with concentrations ranging from 192 ng/g dw to 292,000 ng/g dw. The egg with the highest concentration (192,000 ng/g dw or 80,000 ng/g ww) also had many other organic contaminants detected that could contribute additive toxicity to the embryo. It is unknown if this egg was viable when it was collected, but it is unlikely the egg would have hatched.

Data from this study provides information on the contaminant exposure of young-of-the-year birds (as developing embryos and as nestlings) prior to their migration to the presumed more contaminated wintering grounds. Although, we did not specifically investigate nest success or embryo survival, it is possible that the elevated levels of some organic contaminants detected would affect chick survival or cause reproductive failure. A future study investigating reproductive success is warranted to answer this question.

TABLE OF CONTENTS

| | |
|--|------|
| Abstract | ii |
| List of Figures | vi |
| List of Tables | vii |
| List of Appendices | viii |
| List of Acronyms and Abbreviations | x |
| Introduction | 1 |
| Methods..... | 4 |
| Study Sites | 4 |
| Wintering Grounds | 4 |
| Kern NWR..... | 6 |
| Sonny Bono NWR..... | 7 |
| Imperial Valley | 8 |
| Breeding Grounds | 9 |
| Wyoming | 10 |
| Archived Mountain Plover Egg Sampling Locations..... | 11 |
| Colorado | 13 |
| Montana..... | 14 |
| Southern Phillips County (CMR NWR, BLM, State, and Private lands)..... | 14 |
| Lake Mason NWR..... | 16 |
| Hailstone NWR | 17 |
| Sample Collection and Processing..... | 18 |
| Wintering Grounds | 18 |
| Breeding Grounds..... | 20 |
| Chemical Analyses..... | 22 |
| Data Analysis | 24 |
| Egg Data Analysis | 24 |
| Mixtures..... | 25 |
| Screening for Risk | 25 |
| Screening for Background..... | 27 |
| Quality Assurance..... | 27 |

| | |
|--|----|
| Results and Discussion | 28 |
| Wintering Grounds | 28 |
| Soils-Inorganic Elements..... | 28 |
| Soils-Organic Compounds..... | 32 |
| Terrestrial Invertebrates-Inorganic Elements | 39 |
| Terrestrial Invertebrates-Organic Compounds | 45 |
| Breeding Grounds | 50 |
| Soils-Inorganic Elements..... | 50 |
| Soils-Organic Compounds..... | 55 |
| Terrestrial Invertebrates-Inorganic Elements | 55 |
| Terrestrial Invertebrates-Organic Compounds | 59 |
| Mountain Plover Eggs-Inorganic Elements-2006 | 59 |
| Mountain Plover Eggs-Organic Compounds-2006 | 65 |
| Mountain Plover Eggs-Organic Compounds-2007 | 69 |
| Mountain Plover Eggs-Organic Compounds-2008 | 72 |
| Mountain Plover Eggs-Organic Compounds-2003 | 77 |
| Conclusions..... | 79 |
| Management Recommendations..... | 82 |
| Acknowledgements..... | 83 |
| Literature Cited | 85 |
| Appendices..... | 96 |

Figures

| | |
|---|----|
| Figure 1. General location of mountain plover wintering grounds in California and breeding grounds in Montana, Wyoming, and Colorado. | 2 |
| Figure 2. Sampling locations in California. | 5 |
| Figure 3. Kern National Wildlife Refuge sampling locations..... | 7 |
| Figure 4. SBSS NWR and Imperial Valley sample locations..... | 8 |
| Figure 5. Sampling locations in Wyoming. | 9 |
| Figure 6. Thunder Basin National Grassland sampling locations..... | 10 |
| Figure 7. Foote Creek Rim sampling locations..... | 11 |
| Figure 8. Archived mountain plover egg sampling locations..... | 12 |
| Figure 9. Sampling locations in Colorado..... | 13 |
| Figure 10. Sampling locations in Montana. | 14 |
| Figure 11. Charles M. Russell NWR, BLM, and State sampling locations..... | 15 |
| Figure 12. Lake Mason NWR sampling locations..... | 16 |
| Figure 13. Hailstone NWR sampling locations..... | 18 |
| Figure 14. Pitfall trap design..... | 19 |
| Figure 15. DDT isomer concentrations in soils from California (1985 and 2006)..... | 34 |
| Figure 16. Percentage of DDT isomers comprising total DDT in California soils (1985 and 2006)..... | 34 |
| Figure 17. DDT isomer p,p'DDE concentrations in soils and invertebrates collected in California..... | 47 |
| Figure 18. Total DDT concentrations in soils and invertebrates collected in California..... | 47 |
| Figure 19. Trifluralin concentrations in soils and invertebrates collected in California..... | 48 |
| Figure 20. Concentration of total PCBs in soil and invertebrate samples collected in California..... | 50 |
| Figure 21. Correlation between p,p'DDE concentrations in eggs and eggshell thickness..... | 69 |
| Figure 22. Mountain plover egg data compared to p,p'-DDE effects levels..... | 81 |

Tables

| | |
|---|----|
| Table 1. Eco-SSLs (ug/g dry weight in soil) showing the concentrations of contaminants in soil that are protective of ecological receptors which commonly come into contact with and/or consume biota living in or on soil..... | 26 |
| Table 2. Arithmetic mean and range of concentrations of select inorganics (ug/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006, for comparison with geometric mean and range of inorganics in soils from the western United States as reported by Shacklett and Boerngen (1984)..... | 36 |
| Table 3. Arithmetic mean and range of concentrations of selected organic analytes (ng/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006..... | 38 |
| Table 4. Arithmetic mean and range of concentrations of select inorganics (ug/g dry weight) in invertebrate samples collected from mountain plover wintering sites in California, 2006..... | 42 |
| Table 5. Arithmetic mean and range of concentrations of the most frequently detected organics (ng/g dry weight) in invertebrate samples collected from mountain plover wintering sites in California, 2006. | 46 |
| Table 6. Average and range of concentrations of select inorganics (ug/g dry weight) in soil samples collected from mountain plover breeding sites in Montana and Wyoming, 2006, for comparison with average and range of inorganics in soils from the western United States as reported by Shacklett and Boerngen (1984) | 52 |
| Table 7. Average and range of concentrations of select inorganics (ug/g dry weight) in invertebrate samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006 | 58 |
| Table 8. Average and range of concentrations of inorganics (ug/g dry weight) in egg samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006..... | 60 |
| Table 9. Barium concentrations and developmental stage of mountain plover eggs collected from breeding sites in Colorado, Montana, and Wyoming in 2006..... | 62 |
| Table 10. Average and range of concentrations of most frequently detected organochlorine compounds (ng/g dry weight) in egg samples collected from mountain plover breeding sites in Colorado, Montana and Wyoming, 2006..... | 66 |
| Table 11. Concentrations of detected organochlorine pesticides and total PCBs (ng/g wet weight) in egg samples collected from mountain plover breeding sites in Colorado and Wyoming, 2007..... | 70 |
| Table 12. Concentrations of detected organic analytes (ng/g dry weight) in mountain plover eggs collected from BLM lands, Montana, 2008 | 73 |
| Table 13. Concentrations of DDE in mountain plover eggs compared to burrowing owl eggs collected in the Central and Imperial Valley of California | 80 |

Appendices

| | |
|--|-----|
| Appendix 1. Concentrations of inorganics (ug/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006..... | 97 |
| Appendix 2. Concentrations of organic pesticides and trifluralin (ng/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006..... | 100 |
| Appendix 3. Concentrations of PCBs (ng/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006..... | 104 |
| Appendix 4. Concentrations of inorganics (ug/g dry weight) in terrestrial invertebrate samples collected from mountain plover wintering sites in California, 2006..... | 117 |
| Appendix 5. Concentrations of organic pesticides and trifluralin (ng/g dry weight) in terrestrial invertebrate samples collected from mountain plover wintering sites in California, 2006..... | 119 |
| Appendix 6. Concentrations of PCBs (ng/g dry weight) in terrestrial invertebrate samples collected from mountain plover wintering sites in California, 2006..... | 122 |
| Appendix 7. Concentrations of inorganics (ug/g dry weight) in soil samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006. | 129 |
| Appendix 8. Organochlorine pesticides, organophosphate pesticides, and total PCBs (ng/g dry weight) were not detected in soil samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006..... | 131 |
| Appendix 9. Concentrations of inorganics (ug/g dry weight) in terrestrial invertebrate samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006. | 132 |
| Appendix 10. Organochlorine pesticides, organophosphate pesticides, and total PCBs (ng/g dry weight) were not detected in terrestrial invertebrate samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006..... | 135 |
| Appendix 11. Concentrations of inorganics (ug/g dry weight) in eggs samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006. | 136 |
| Appendix 12. Concentrations of organic compounds (ng/g dry weight) in eggs samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006..... | 139 |
| Appendix 13. Mountain plover egg content data from breeding sites in Colorado, Montana, and Wyoming, 2006..... | 140 |
| Appendix 14. Mountain plover egg measurement data from breeding sites in Colorado, Montana, and Wyoming, 2006..... | 141 |
| Appendix 15. Mountain plover egg thickness data from breeding sites in Colorado, Montana, and Wyoming, 2006..... | 142 |

| | |
|--|-----|
| Appendix 16. Concentrations of organic pesticides, PCBs, and trifluralin (ng/g dry weight) in eggs samples collected from mountain plover breeding sites in Colorado and Wyoming, 2007..... | 143 |
| Appendix 17. Mountain plover egg content data from breeding sites in Colorado and Wyoming, 2007. | 144 |
| Appendix 18. Mountain plover egg measurement data from breeding sites in Colorado and Wyoming, 2007..... | 144 |
| Appendix 19. Mountain plover eggshell thickness data from breeding sites in Colorado and Wyoming, 2007..... | 145 |
| Appendix 20. Concentrations of organic pesticides, PCBs, and trifluralin (ng/g dry weight) in eggs samples collected from mountain plover breeding sites in Montana, 2008. | 146 |
| Appendix 21. Mountain plover egg content data from breeding sites in Montana, 2008. | 148 |
| Appendix 22. Mountain plover egg measurement data from breeding sites in Montana, 2008..... | 148 |
| Appendix 23. Mountain plover egg thickness data from breeding sites in Montana, 2008..... | 149 |
| Appendix 24. Mountain plover egg content data from breeding sites in Wyoming, 2003 | 149 |
| Appendix 25. Concentrations of organic pesticides, PCBs, and trifluralin (ng/g dry weight) in eggs samples collected from mountain plover breeding sites in Wyoming, 2003 | 150 |
| Appendix 26. Concentrations of organic pesticides (ng/g dry weight) in egg samples collected from mountain plover breeding sites in Montana and Colorado, 2003 | 153 |

List of Acronyms and Abbreviations

Measurements

cm – centimeter
dw – dry weight
ft – foot
ha – hectare
km – kilometer
m – meter
mg/kg – milligram per kilogram
mm – millimeter
ng/g – nanogram
ppb – part per billion
ppm – part per million
µg/g – microgram per gram
µg/kg - microgram per kilogram
uL – microliter
ww – wet weight

Agencies

ATSDR – Agency for Toxic Substances and Disease Registry
BLM – Bureau of Land Management
USDOI – U.S. Department of Interior
USEPA – United State Environmental Protection Agency
USFS – U.S. Forest Service
USFWS – U.S. Fish and Wildlife Service
USGS-BRD – U.S. Geological Survey – Biological Research Division

Locations

ACF – Analytical Control Facility
ACM – Antelope Coal Mine
CMR – Charles M. Russell National Wildlife Refuge
CRWO – Carlsbad Fish and Wildlife Office
FCR – Foote Creek Rim
LMU – Lake Mason Unit
NWR – National Wildlife Refuge
PNG – Pawnee National Grassland
SBSS – Sony Bono Salton Sea
TBNG – Thunder Basin National Grassland

Chemicals and related terms

BHC – Benzene hexachloride
BW – Body Weight
DDD - dichlorodiphenyldichloroethanes

DDE – dichlorodiphenyldichloroethylenes
DDT – dichlorodiphenyltrichloroethanes
ECO-SSLs – Ecological Soil Screening Levels
HCB – Hexachlorobenzene
HCH – Hexachlorocyclohexane
ICP-MS – Inductively Coupled Plasma Emission Mass Spectrometry
LOECs – Low Observed Effect Concentration
NOAEL – No Observed Adverse Effect Level
NOEC – No Observed Effect Concentration
OCs – organochlorine compounds
OPs – Organophosphate chemicals
PCBs – Polychlorinated biphenyls
QA/QC – Quality Assurance/Quality Control
RED – Reregistration eligibility decision
TOC – Total Organic Carbon
TRV – Toxicity Reference Value

INTRODUCTION

The mountain plover (*Charadrius montanus*) is a shorebird that primarily inhabits dry upland sites, such as shortgrass prairie and shrub-steppe landscapes during both wintering and breeding seasons. This plover prefers and often breeds in black-tailed prairie dog (*Cynomys ludovicianus*) towns, especially in areas where taller grasses surround the dog town, but they will also breed in fallow fields and recently plowed ground (Knopf 1996). Nesting occurs primarily in Montana (including three different National Wildlife Refuges (NWRs), Charles M. Russell, Lake Mason, and Hailstone), Wyoming, and Colorado; although new data suggest that New Mexico may have comparable numbers. The birds winter in the Central and Imperial Valleys of California, Arizona, Texas, and Mexico (Figure 1; NatureServe 2010). Approximately half the entire United States population winters in California (Andres and Stone 2009; USFWS 2011), with the Imperial Valley becoming the increasingly favored location. Conversion of the mountain plovers' native grassland habitat to agricultural lands in both wintering and breeding areas has resulted in the birds using farmlands for feeding and roosting (Knopf and Rupert 1995), particularly in both the Central and Imperial Valleys where over 809,400 ha (2 million acres) are in production all year. This intensive production results in increasing dependence on agricultural chemicals and potential increased exposure of mountain plovers to applied pesticides and fertilizers.

In 1982, the U.S. Fish and Wildlife Service (Service) designated the mountain plover as a category 2 candidate species, meaning that more information was necessary to determine the status of the species. In 1990, a status report suggested Federal listing under the Endangered Species Act may be warranted, and in November of 1994, the status of the mountain plover was elevated to a category 1 candidate species (USFWS 1999a). In 1999, the Service proposed to list the mountain plover as a threatened species, but in 2003, determined that the listing was not warranted (USFWS 2003a). On November 16, 2006, Forest Guardians and the Biological Conservation Alliance filed a complaint in the District Court for the Southern District of California challenging the withdrawal of the proposal to list the mountain plover (Forest Guardians, et al. v Ken Salazar et al., Case No. 3:06-cv-02560-MMA-BLM).




Figure 1. General location of mountain plover wintering grounds in California and breeding grounds in Montana, Wyoming, and Colorado.

A settlement agreement between the plaintiffs and the Federal defendants was filed on August 8, 2009. As part of the settlement, the Service agreed to reconsider its 2003 decision to withdraw the proposed listing of the mountain plover and to submit to the Federal Register by July 31, 2010, a notice re-opening the proposal to list the mountain plover and provide for public comment. It was agreed that upon publication of the notice, the 2003 withdrawal of the proposed

rule would be vacated. After a thorough review of all available scientific and commercial information, the Service determined that the mountain plover is not threatened or endangered throughout all or a significant portion of its range as of May 12, 2011.

Although the mountain plover has never been a listed species, it is protected under the Migratory Bird Treaty Act (16 U.S.C. 703) and is on the Service's Birds of Conservation Concern list (USFWS 2008). Current mountain plover population estimates approximately 20,000 birds (USFWS 2011). Based on data from the Breeding Bird Survey, plover population declined by 67% from 1966 to 1993 (Knopf 1996) but this decline has currently moderated (USFWS 2011). Because contaminant concerns have been identified as potentially affecting the mountain plover and potentially contributing to its decline (USFWS 1999a; USFWS 2002), this investigation was designed to address this gap.

Specifically, while wintering in the Imperial Valley of California, mountain plovers spend the early winter in harvested or fallow alfalfa fields before switching over to Bermuda grass fields after the fields are burned in late winter (F. Knopf, USGS-BRD, Pers. Comm. 2/05). These habitat preferences expose wintering mountain plovers to contaminants associated with agricultural practices, including current use pesticides and their adjuvants (e.g., organophosphates and carbamate pesticides, and herbicides), persistent chemicals from historic applications (e.g., organochlorine pesticides), trace elements from irrigation drainwater (e.g., selenium), chemicals present in the irrigation water (e.g., perchlorate), and fertilizers. Although the potential for direct exposure to organophosphate and carbamate pesticides may be limited because certain applications occur when plovers are not using the fields (Wunder and Knopf 2003, USFWS 2003a), some of these pesticides (along with adjuvants and carriers) are classified as moderately persistent and residues can remain in the fields and be accumulated by wintering plovers. Numerous pesticides are applied to alfalfa and Bermuda grass fields in the Imperial Valley that can adversely affect the long-term health of birds or their offspring. Some of these applied pesticides are known immunosuppressors in mammals (effects to birds have not been studied) and some are known endocrine disruptors.

Mountain plover exposure to environmental contaminants is not restricted to the wintering grounds. Although more limited, plovers on their breeding grounds may be exposed to insecticides used for grasshopper control, as well as DeltaDust, a synthetic pyrethroid used in some prairie dog towns to control the primary vector (fleas) of the plague bacterium *Yersinia pestis*.

To evaluate plover exposure on both wintering and breeding grounds, this study assessed (1) contaminant concentrations in likely food items consumed by mountain plovers at their wintering grounds; (2) exposure of developing embryos to contaminants transferred by adults to eggs; and, (3) contaminant levels in likely food items of nestling and adult plovers while on their breeding grounds. Data from objectives two and three will provide information on contaminant exposure by young-of-the-year birds (as developing embryos) before migrating to the presumed more contaminated wintering grounds.

METHODS

Overall, samples of soil and soil invertebrate samples were collected from five wintering locations and seven breeding areas. Presumed viable plover eggs were collected in 2006 and 2007 from breeding areas in Colorado and Wyoming and abandoned or failed-to-hatch eggs were collected in 2006 and 2008 in Montana. Additional eggs collected from Wyoming and archived in 2003 were also analyzed for contaminants as part of this study.

Study Sites

Wintering grounds

Sampling sites for wintering mountain plovers included several locations in the Imperial and Central Valleys of California (Figure 2) with one on-refuge site at the Kern NWR (K15) in the Central Valley and four sites in the Imperial Valley. Imperial Valley sites included one on-refuge field within the boundaries of the Sony Bono Salton Sea (SBSS) NWR (SB420) and three off-refuge fields nearby (J20, J22, K18). The two on-refuge sites, one on SBSS NWR (Imperial Valley) and one at Kern NWR (Central Valley) were selected based on recommendations by



Mountain Plover Sample Locations - California



Figure 2. Sampling locations in California.

NWR managers. The three off-refuge sites in Imperial Valley were selected based on reports of where plovers were found foraging prior to spring migration and where permission to sample could be obtained from property owners and managers.

Kern NWR

Kern NWR is located in the historic Tulare Lake Basin in an area once covered by an inland lake and wetland complex totaling over 252,938 ha (625,000 acres). The refuge was established on November 18, 1960, and covers 4,297 ha (10,618 acres). It is located approximately 30.6 km (19 miles) west of the city of Delano in the San Joaquin Valley, and all of the land within the Refuge boundary is controlled by the Service. Habitats within the refuge include seasonally flooded marshes, grasslands, alkali scrub/playas, and riparian lands. The refuge contains one of the few remaining wetlands left in the area and is intensively managed to produce habitat for migrating and wintering waterfowl and other waterbirds. This includes the creation of marsh conditions and other diverse habitat production to ensure the lands will provide a continuing food source for the birds (USFWS 2005).

The sampling location on Kern NWR is one of the few parcels managed for upland endangered species, primarily reptiles and small mammals, but it also may attract mountain plovers (Figure 3). Vegetation in the field that was sampled included mixed grasses, with patches of scrub such as iodine bush (*Allenrolfea occidentalis*), suaeda (*Suaeda* sp.), and alkali heath (*Frankenia salina*). According to refuge management, the field had not been farmed for many years with grazing used to control non-native grasses. Pixley NWR, established on November 17, 1959, is located approximately 30.6 km (19 miles) south of the city of Tulare in the San Joaquin Valley. The NWR is located within the Tulare Basin, of which Tulare Lake is the lowest portion. This refuge has agricultural fields with annual grasses that are used by mountain plovers, but the site could not be sampled due to concerns about potential impacts on the federally endangered blunt-nosed leopard lizard (*Gambelia silus*).

Imperial Valley – Off-Refuge Fields

The three off-refuge sites in the Imperial Valley were selected based on reports of where plovers were found to forage prior to spring migration and where permission to sample could be obtained from property owners and managers (Figure 4).




Figure 4. SBSS NWR and Imperial Valley sample locations.

The three off-refuge sites (J20, J22, and K18) were active agricultural fields. All three fields were subjected to post-harvest burns prior to use by the plovers. Between March, when the plovers left, and the first of May, when samples were taken, the fields were tilled and grasses had re-grown to a height of approximately 15.2 cm. Records of pesticide use for individual fields were not readily available. However, two fields (J20 and J22) were replanted with Bermuda grass, assumed to have been treated with the fumigants (e.g. sulfur) and pesticides typically

employed for Bermuda grass crops in Imperial Valley (Pesticide Action Network 2010). The third agricultural field (K18) had a mix of grasses indicating that it may have gone fallow that season. As such, K18 may not have been treated with products used for Bermuda grass cultivation prior to sampling.

Breeding Grounds

Several known breeding grounds in Montana, Wyoming, and Colorado were selected as study sites. This included public lands administered by the Bureau of Land Management (BLM), U.S. Forest Service (USFS), and the Service's NWR system.




Figure 5. Sampling locations in Wyoming.

Wyoming

Study sites in Wyoming included the Thunder Basin National Grassland (TBNG) with an adjacent area near the Antelope Coal Mine (ACM) and Foote Creek Rim (FCR) in southern Wyoming (Figure 5).

TBNG, managed by the Douglas Ranger District of the USFS, is located in northeastern Wyoming in the Powder River Basin with the Big Horn Mountains to the west and the Black Hills to the east (Figure 6). The area consists primarily of shortgrass prairie and sagebrush, which provides habitat for white-tailed deer (*Odocoileus virginianus*), mule deer (*Odocoileus hemionus*), pronghorn antelope (*Antilocapra americana*), numerous small mammals, and a variety of raptors and songbirds (USFS 2010).




Figure 6. Thunder Basin National Grassland sampling locations.

An additional breeding area, located on Foote Creek Rim (FCR) near the Arlington Wind Plant in southeastern Wyoming was also selected as a study area (Figure 7). FCR is a remote, treeless plateau consisting of shortgrass prairie. Located between the towns of Laramie and Rawlins, the area is considered one of the windiest places in America. It is important habitat for white-tailed deer, mule deer, pronghorn antelope, elk (*Cervus elaphus*), and numerous species of small mammals, raptors, and passerines (BLM 2010).




Figure 7. Foote Creek Rim sampling locations.

Archived Mountain Plover Egg Sampling Locations

Archived mountain plover eggs collected in 2003 were submitted for organic chemical analysis using funds from this investigation. This information was used to demonstrate the ongoing

exposure of mountain plover to organic compounds. Sites sampled in 2003 included a site in Albany County and another in the Shirley Basin located in Carbon County in southern Wyoming (Figure 8). These high elevation grasslands consist primarily of mixed-grass prairie and sagebrush, which supports pronghorn antelope, domestic cattle, and black-tailed prairie dogs (Plumb 2004). A third site, located in Park County in the northwestern Wyoming, was also sampled (Figure 8). This area is a desert-shrub dominated by greasewood (*Sarcobatus vermiculatus*) with some sagebrush and mixed grass prairie interspersed. Pronghorn antelope, domestic sheep and cattle, and white-tailed prairie dogs (*Cynomys leucurus*) are the primary inhabitants (Plumb 2004).




Figure 8. Archived mountain plover egg sampling locations.

Colorado

Mountain plovers nest across the eastern plateau region of Colorado, including a breeding stronghold in Weld County (Knopf and Rupert 1996). Samples collected for this investigation were primarily collected in Weld County on the Pawnee National Grassland (PNG). PNG is a shortgrass prairie managed by the USFS and is known internationally for the birding opportunities. Additionally, pronghorn antelope, mule deer, and many other small mammal species inhabit this grassland.




Figure 9. Sampling locations in Colorado.

Montana

Several known breeding grounds in Montana were selected as study sites. These included public lands administered by the BLM and Charles M. Russell, Lake Mason, and Hailstone NWRs in north-central and central Montana (Figure 10).




Figure 10. Sampling locations in Montana.

Southern Phillips County (CMR NWR, BLM, State and Private lands)

Southern Phillips County in Montana is thought to contain not only the largest mountain plover breeding populations in Montana but also one of the largest breeding populations in North America (Dinsmore 2001). Study sites located in southern Phillips County included Charles M. Russell (CMR) NWR and the adjacent BLM and Montana State School Trust properties. These

areas consist of rolling uplands interspersed with steeply eroded ravines. Sagebrush and interspersed shortgrass prairie dominate the habitat (USFWS 2010). Mountain plovers nest almost exclusively on active black-tailed prairie dog towns in this part of Montana. Vegetation in these colonies is sparse, but includes such species as fringed sagewort (*Atemisia frigida*), plains prickly pear (*Opuntia polyacantha*), blue grama (*Bouteloua gracilis*), needle-and-thread grass (*Stipa comata*), and Sandberg bluegrass (*Poa secunda*). Prairie dog towns were mapped entirely in 2003, which showed that the towns covered more than 2,954 ha (7,300 acres) at CMR NWR (Jo Ann L. D. Dullum, Wildlife Biologist, CMR NWR, Pers. Comm. 2010). In 2002, prairie dog towns mapped on BLM, State, and private lands north of the refuge totaled 9022 ha (22,292 acres; Figure 11).




Figure 11. Charles M. Russell NWR, BLM, and State sampling locations.

Lake Mason NWR

Lake Mason NWR is located in central Montana approximately 12.9 km (8 miles) northwest of the town of Roundup (population 4,000) in Musselshell County. The refuge includes 6730 ha (16,630 acres) in fee title and 2227 ha (5,502 acres) in refuge easements. It is divided into three units: Lake Mason, Willow Creek Unit, and the North Unit. Lake Mason and the Willow Creek Units were the only units sampled during this investigation (Figure 12).




Figure 12. Lake Mason National Wildlife Refuge in Montana. Units from north to south are the North Unit, Willow Creek Unit, and Lake Mason.

Lake Mason Unit (LMU) is a 526 ha (1,300 acre) enhanced wetland surrounded by rolling shortgrass prairie hills. The lake is fed and drained by South Willow Creek. Water in the creek

is mainly from runoff and agricultural return water is minimal. During this investigation, Lake Mason was dry and contained active prairie dog town located on the western portion of the lake. This colony was mapped in 2002 and at that time it covered 193 ha (478 acres; Jo Ann L. D. Dullum, Wildlife Biologist, CMR NWR, Pers. Comm. 2010).

The Willow Creek Unit of Lake Mason NWR is located approximately 19 km northwest of the LMU, is 874 ha (2,160 acres) in size and is managed to maintain a healthy shortgrass prairie community. Mountain plovers are attracted to this area as are many other migratory birds typical of shortgrass prairie ecosystems. The prairie dog town on this unit was mapped in 2002 and covered 190 ha (469 acres; Jo Ann L. D. Dullum, Wildlife Biologist, CMR NWR, Pers. Comm. 2010).

Hailstone NWR

Hailstone NWR, located in Stillwater County, approximately 5 km northeast of the town Rapelje and is part of a large closed basin located in south-central Montana. The refuge was created in 1942 by an Executive Order of President Franklin D. Roosevelt as an easement refuge of 1112 ha (2,748 acres). This contains approximately 51 ha (126 acres) of prairie dog towns based on mapping completed in 2002 (Jo Ann L. D. Dullum, Wildlife Biologist, CMR NWR, Pers. Comm. 2010) (Figure 13).




Figure 13. Hailstone NWR sampling locations

Sample Collection and Processing

Wintering Grounds

Samples of soil and soil invertebrates were collected in early May 2005, by environmental contaminant biologists from the Carlsbad Fish and Wildlife Office (CFWO) and by refuge biologists. As indicated previously, samples were collected from three off-refuge fields in Imperial Valley, one on-refuge field in Imperial Valley, and one on-refuge field in Central Valley (Kern NWR).

Composite surface soil and insect samples were collected from five stations in each field. Soil samples were collected using stainless steel equipment that was either pre-cleaned (dedicated) or cleaned between stations using standard protocols (USEPA/ERT 1994) requiring phosphate-free detergent, dilute nitric acid, acetone, and distilled water. Each station was defined by a circle with a radius of one meter and twelve pitfall traps were positioned in each circle using the layout in Figure 14. A labeled stake was placed in the center to help relocate the station. Holes were dug using a bulb planter and trowels for pitfall trap placement.



Figure 14. Pitfall trap design (arrow depicts 1 m radius).

Soils removed for the pitfall traps were placed in pre-cleaned stainless steel bowls and homogenized using stainless steel spoons to produce one composite soil sample for each station. A portion of the composite soil sample was placed in a WhirlPak® bag for total organic carbon (TOC) and grain size. The remaining soil was placed in an I-Chem® jar for inorganic and organic chemical analysis.

Pitfall traps were 11.3 cm (4.5 inch) high 528 mL (16 fluid ounce) cups. These traps tended to be narrow, which made it difficult for large insects to escape once they fell into the cup. To avoid introduction of sampling artifacts, no water or solvent was added to the traps to immobilize insects once they had fallen in the trap.

Pitfall traps were left in place over night. The next day, all insects in each of the 12 cups at a station were composited into one sample and some sweep netting was performed to augment the sample. However, because the traps had soil in them, which made it almost impossible to

separate insects from the soil in the field, the contents from pitfall traps and net sweeps were put into labeled sample bags and placed on ice for further processing at the lab at CFWO. Once at the lab, insects were separated from any clumps of soil that may have fallen into the traps, transferred to cleaned glass jars, and placed in the freezer. Samples consisted primarily of spiders and to lesser extent beetles. Sample weights averaged 1 gram, limiting the number of analyses (inorganic and organic) that could be performed. Once processed, all samples were stored frozen until they were shipped to laboratories designated by the Service's Analytical Control Facility (ACF) for chemical analyses.

Soil samples from Imperial and Kern were analyzed for metals, organochlorine compounds (OCs) including polychlorinated biphenyls (PCBs) and pesticides, organophosphates (OPs), triazine herbicides, sulfur, trifluralin, and perchlorate. All insect samples were analyzed for metals and trifluralin. Because of the small sample sizes, only a subset of the insect samples was analyzed for OCs.

Breeding Grounds

Samples collected on the breeding grounds included soil, terrestrial insects, and mountain plover eggs. These matrices were collected from the study area locations in Montana, Wyoming, and Colorado (Figures 6, 7, 8, 9, 11, 12, and 13). The number of eggs collected from the different locations depended on egg availability. Eggs were collected by hand with no more than one egg collected from any nest. In Montana, eggs were collected in May and June of 2006 and 2008 from southern Phillips County by graduate students under the direction of mountain plover researcher Dr. Stephen Dinsmore. These eggs were collected from abandoned or failed nests, and all remaining eggs from these nests were collected. Many of the eggs collected from failed nests in Montana had cracked eggshells. These eggs were only dissected into jars to reduce handling. Wet weight concentrations of contaminants of these eggs are likely elevated from possible desiccation. Dry weight concentrations should be comparable to literature values. Mountain plover researcher, Dr. Fritz Knopf, collected mountain plover eggs from Wyoming and Colorado sites in May and June 2006 and 2007.

Eggs collected by environment contaminant biologists were processed using standard protocols (Hudson River Natural Resource Trustees 2002). All eggs were placed in cartons and transported immediately to the lab for processing by biologists using standard protocols adopted for studies on the impacts of PCBs in the Hudson River (Hudson River Natural Resources Trustees 2002).

Intact eggs were gently cleaned with distilled water, weighed, and breadth and width were measured to the nearest 0.1 mm using a dial caliper. The volume was measured as the weight of water displaced by the egg. A scalpel or curved scissors was pre-rinsed with dilute nitric acid followed by distilled water and used to cut the shell around the equator. Contents were aged and examined for abnormalities and malpositioning. If an embryo was present, it was further evaluated for evidence of conspicuous malformations (e.g., missing limbs, malformed beaks). To minimize potential loss of the sample and prevent cross-contamination that could result from physical manipulation, embryos were not evaluated for subtle malformations that required measurements. Egg contents that were in various stages of decomposition made the assessment of some embryos difficult if not impossible. Once observations were recorded, contents were placed in certified cleaned amber glass jars and frozen until they could be submitted for chemical analyses by a laboratory under contract with ACF. Eggs were analyzed for metals, OCs, and total PCBs in 2006. In 2007 and 2008, eggs were analyzed for OCs and trifluralin.

Eggshells were placed in cartons to dry at room temperature for a minimum of 30 days. Once dry, the thickness of each eggshell (shell + shell membrane) was measured at four points around the girth with a Mitutoyo micrometer, which was fitted with ball attachments and accurate to 0.01 mm. Some eggshells had hardened yolk residue that could not be removed even at the time the contents were harvested. This was generally confined to one side of the shell and was avoided when taking measurements for shell thickness. Eggshell thickness was measured by methods described in Bunck et al. (1985) and Ratcliffe (1970).

Archived mountain plover eggs collected in 2003, that were not collected specifically as part of this investigation, were submitted for organic chemical analysis using funds from this

investigation. These eggs were collected by University of Wyoming graduate student, R.E. Plumb. The number of eggs collected from the different locations depended on egg availability. Eggs were collected in June and July of 2003 and no more than one egg was collected from any nest (Plumb 2004). Service personnel dissected the eggs according to the methods previously described. Eggs were analyzed for OCs, PCBs, and trifluralin to demonstrate the presence of organic compounds and that exposure to mountain plover is ongoing.

Soil and terrestrial insect samples were collected in May and June of 2006. In Montana this was completed by the Service's Montana Field Office personnel and in Wyoming and Colorado by the Service's Wyoming Field Office personnel. Insect samples were collected using the same methods described in the wintering grounds sampling. Traps were left out overnight. In Montana, sweep netting was also used to augment the pitfall trap insects. All trap and net samples were composited into WhirlPak® bags and frozen until contents could be sorted with chemically-cleaned Teflon® tweezers. All insects were placed in chemically-cleaned amber glass jars and frozen.

Because foraging birds remain close to their nesting areas throughout the season (Knopf and Wunder 2006), sampling for insects was limited to one event and focused in areas where the birds were observed foraging. Samples of soil and insects were analyzed for metals, OPs, OCs, and total PCBs by the same laboratories listed above.

Chemical Analyses

Samples of soil and invertebrates from California sites, and soil, terrestrial insects, and mountain plover eggs samples collected on the breeding grounds were shipped to ACF designated laboratories for analyses, under catalog numbers 1040078 and 1040079, for California, 6070089 and 6070096 for Montana, and 6030187 and 6030192 for Wyoming and Colorado.

Samples were analyzed for organic contaminants, soil TOC and soil grain size by Geochemical and Environmental Research Group in College Station, Texas. Analyses for inorganic constituents were conducted by Trace Element Research Laboratory in College Station, Texas.

The percent TOC was determined by measuring carbon dioxide produced on ignition of acidified, freeze-dried sample, using an infrared detector. Grain size distribution was determined as percent sand (>0.0063 mm), silt (0.0039 mm to 0.063 mm) and clay (<0.0039 mm) using sieves and settling columns after sample had been treated with hydrogen peroxide and a dispersing agent. For metals, samples were acid extracted, then analyzed using hydride generation atomic absorption spectrometry (selenium and arsenic), cold vapor atomic absorption spectrometry (for mercury) or Inductively Coupled Plasma Emission (ICP) spectrometry (other metals).

To determine concentrations of the organic compounds (pesticides, total PCBs, and trifluralin), samples were first extracted with methylene chloride and cleaned using gel-permeation chromatography. Silica gel chromatography was used for additional cleaning and separation of PCBs. Samples were then analyzed by gas chromatography with an electron capture detector (pesticides and PCBs) or high resolution gas chromatography-mass spectrometry in Selective Ion Monitoring mode (for certain PCB congeners).

The target analytes and method detection limits reported by the contract labs follow. Target analytes for soil samples included the following parameters: % moisture, % TOC, and grain size distribution. Target analytes for soil, invertebrate, and egg samples for organic compounds included: PCB-total¹ (polychlorinated biphenyls as Aroclors), aldrin², alpha-, beta-, gamma, and delta-BHC² (s), alpha- and gamma-chlordanes², chlorpyrifos², dieldrin², endosulfan II², endrin², heptachlor², heptachlor epoxide², HCB² (hexachlorobenzene), HCH² (hexachlorocyclohexane), mirex², cis- and trans-nonachlor², oxychlordane², o,p'- and p,p'-DDD² (dichlorodiphenyldichloroethanes), o,p'- and p,p'-DDE² (dichlorodiphenyldichloroethylenes), o,p'- and p,p'-DDT² (dichlorodiphenyltrichloroethanes), pentachloro-anisole², 1,2,3,4- and 1,2,4,5-TeCB² (tetrachlorobenzenes), toxaphene¹, tetrachlorobenzenes² (1,2,3,4- and 1,2,4,5-),

and trifluralin². Results of analyses for organic contaminants are reported as ng/g or parts per billion (ppb). The dry weight detection limits are as follows: ¹1.43 as µg/kg or ppb; ²0.143 as µg/kg or ppb.

The target analytes for soil, invertebrate, and egg samples for inorganics included: aluminum¹, arsenic², barium², beryllium³, boron¹, cadmium³, chromium-total⁴, copper⁴, iron¹, lead⁵, magnesium⁵, manganese⁷, mercury⁶, molybdenum⁵, nickel⁵, selenium², strontium⁴, vanadium⁴, and zinc⁷. Results of analyses for inorganic contaminants are reported as µg/g or parts per million (ppm). The dry weight detection limits are as follows: ¹10.2 as mg/kg or ppm; ²0.5 as mg/kg or ppm; ³0.2 mg/kg or ppm; ⁴1.0 as mg/kg or ppm; ⁵5.0 as mg/kg or ppm; ⁶0.1 mg/kg or ppm; ⁷0.5 mg/kg or ppm.

Data analysis

Analytical results were provided by contract labs as both wet weight (ww) and dry weight (dw) based concentrations. To be consistent with the various screening values, it was necessary to consider both ww-based and dw-based contaminant levels because dw-based concentrations are used to evaluate contaminant levels in soils, whereas either ww or dw-based concentrations may be used to evaluate contaminant levels in biota.

The percent moisture reported for each sample is provided to enable conversion between ww and dw concentrations as follows.

Wet weight concentration = dry weight concentration x (f_{solids}), or

Dry weight concentration = wet weight concentration / (f_{solids}), where

$$f_{\text{solids}} = 1 - (\% \text{moisture}/100)$$

Egg Data Analysis

Shell thickness for each egg was reported as the mean of four individual measurements. The mean, standard deviation, and 95% confidence interval was then computed for each species.

Contaminant levels in eggs were evaluated as ww concentrations. This was done to allow for adjustment for moisture loss that can occur due to the delay in time between when the egg was laid and when it was collected. Contaminant levels measured in bird eggs were adjusted accordingly using equations from the published literature (e.g. Stickel et al. 1983; Hoyt 1979) and from direct measurements with intact eggs.

Mixtures

DDT and its metabolites are a family of compounds (isomers) that occur as mixtures. Individual isomers (e.g. p,p'-DDE) can be evaluated individually but it is also desirable to evaluate the distribution and potential risks posed by concentrations of mixtures as a whole. In general, the total DDT concentration was computed as the sums of the six individual compounds (o,p and p,p'-DDD, DDE and DDT) with ½ the detection limit used as a surrogate for those that were below the limit of detection.

Screening for risk

To screen for ecological risks, dry weight concentrations of contaminants measured in soils were compared with U.S. Environmental Protection Agency (USEPA) ecological soil screening levels (Eco-SSLs; Table 1). The USEPA adopted a protocol for deriving Eco-SSLs in 2000 and has since developed Eco-SSLs for 15 of the target analytes in this study. The Eco-SSLs are ecological risk-based soil screening levels derived in a standardized manner and are based on contaminant risks to soil microorganisms, soil invertebrates, plants, birds and mammals. Separate values were derived for each of the receptors for which data of adequate quantity and quality were available. The Eco-SSLs for invertebrates and plants are the means of concentrations that affect 10% to 20% of exposed organisms (10-20% effect levels), or are maximum acceptable concentrations computed from No Observed Effect Concentrations (NOECs) and Low Observed Effect Concentrations (LOECs). The Eco-SSLs for wildlife are calculated using literature-based toxicity reference values (TRVs), which are high end No Observed Adverse Effect Levels (NOAELs as daily dose rate), combined with bioaccumulation factors for highly exposed forage and prey species. The Eco-SSLs were used preferentially over

other currently available ecological risk-based screening levels. Most Eco-SSLs were in draft form in 2006, and were finalized in subsequent years.

Table 1. Eco-SSLs¹ (ug/g dry weight in soil) showing show the concentrations of contaminants in soil that are protective of ecological receptors which commonly come into contact with and/or consume biota living in or on soil.

| Trace Element | Plants | Soil Invertebrates | Wildlife | |
|------------------------|--------|--------------------|-------------------------|----------------------------|
| | | | Avian | Mammalian |
| Arsenic | 18 | NA ² | 43 | 46 |
| Barium | NA | 330 | NA | 2,000 |
| Beryllium | NA | 40 | NA | 21 |
| Cadmium | 32 | 140 | 0.77 | 0.36 |
| Chromium | NA | NA | Cr III - 26 Cr VI-NA | Cr III – 34 Cr VI - 130 |
| Cobalt | 13 | NA | 120 | 230 |
| Copper | 70 | 80 | 28 | 49 |
| Lead | 120 | 1,700 | 11 | 56 |
| Manganese | 220 | 450 | 4,300 | 4,000 |
| Nickel | 38 | 280 | 210 | 130 |
| Selenium | 0.52 | 4.1 | 1.2 | 0.63 |
| Vanadium | NA | NA | 7.8 | 280 |
| Zinc | 160 | 120 | 46 | 79 |
| Organochlorines | | | | |
| Dieldrin | NA | NA | 0.022 | 0.0049 |
| DDT & metabolites | NA | NA | 0.093 | 0.021 |

¹Eco-SSLs - USEPA 2000a and b; USEPA 2005a-f; USEPA 2007a-h; USEPA 2008

²NA = Not Available.

Concentrations of trace elements and two pesticides in soil samples collected from mountain plover wintering and breeding areas were compared to the Eco-SSL values to determine if concentrations detected could potentially adversely affect these birds directly by exposure through diet and incidental ingestion of soil or indirectly by impacting their food source (toxicity to soil invertebrates) or habitat (toxicity to plants).

Screening for Background

Because metals are naturally occurring constituents of soils, data on metals were also compared with background concentrations for surface soils to help managers with decisions regarding feasibility of actions to manage risk.

Data from large-scale studies and/or compilations for nation-wide application were used to identify ranges of concentrations that may be considered typical for undisturbed soils. The available data offer rough estimates at best of background concentrations because soil characteristics are highly variable from region to region, state to state, and even locale to locale. Data from Shacklette and Boerngen (1984) for surface soils from throughout the conterminous United States, specifically western states, and from Bradford et al. (1996) for soils from California were used to evaluate concentrations of naturally occurring inorganic analytes in soil samples from wintering grounds.

Soil samples from Imperial and Kern were analyzed for metals, organochlorine compounds (OCs) including polychlorinated biphenyls (PCBs) and pesticides, organophosphates (OPs), triazine herbicides, sulfur, trifluralin, and perchlorate. All insect samples were analyzed for metals and trifluralin. Because of the small sample sizes, only a subset of the insect samples was analyzed for OCs.

In breeding areas, all soil and invertebrate samples were analyzed for metals and OCs (including PCBs and pesticides). In Wyoming and Colorado, these samples were also analyzed for OPs. Eggs collected in 2006 were analyzed for metals and OCs (including PCBs and pesticides). For 2003 archived eggs and eggs collected in 2008, analyses included OCs and trifluralin.

Quality Assurance

Laboratories used analytical methods and followed quality assurance (QA) and quality control (QC) protocols in accordance with requirements by ACF. A rigorous program of methods standardization and QA/QC assessment is maintained by ACF for all contract laboratories.

Procedural blanks, duplicates, spiked samples, and analysis of standard reference materials were used routinely with each batch of samples to evaluate and maintain QA/QC. For digestion of all matrices for the ICP-MS semi-quantitative scan, quality control included digestion blanks, reference solutions and materials, samples replicate, and sample spikes. Quality control parameters for sample analysis by ICP-MS scan included running the single calibration standard as a sample, a measurement of precision by repeated runs of a reference solution, the analysis of independent-source laboratory control samples, and within-run monitoring of changes in the internal standards. For samples analyzed by flow injection atomic absorption (selenium), pre-digestion quality control included digestion blanks, replicates, spikes, and reference solutions/materials. Analytical quality control for selenium included calibration verification solutions and analysis spikes. For analysis of mercury in tissue by thermal combustion, amalgamation, and atomic absorption spectroscopy (DMA-80), quality control included calibration verification checks, reference tissues, replicates, method spikes, and blanks.

RESULTS AND DISCUSSION

Wintering Grounds

Soils-Inorganic Elements

Summary statistics for trace element concentrations in soil samples collected from two refuges in California and three sites in the Imperial Valley are contained in Table 2; full analytical results are listed in Appendix 1. The averages and ranges of inorganic constituents in soil samples from wintering areas in California were mostly within the western average and range of soil in the western U.S. (Shacklette and Boerngen 1984), but some constituents were above the Eco-SSLs that are protective of birds and/or soil invertebrates. Average concentrations of beryllium, cobalt, manganese, molybdenum, and nickel in soil samples were slightly higher than the average reported for western soils but fell at the lower range of concentrations reported for western U.S. soils and were at or below mean background concentrations in California soils (Bradford et al. 1996). Only manganese in soil samples exceeded the Eco-SSL for plants and invertebrates.

Average concentrations of arsenic, barium, mercury, lead, and vanadium in samples were below the geometric mean for background in western U.S. soils (Shacklette and Boerngen 1984). Average concentrations of arsenic, cobalt, manganese, nickel and selenium in soil were lower than Eco-SSLs for the protection of birds. There are no established Eco-SSLs for barium, beryllium, boron, and mercury for the protection of birds. Concentrations of barium and beryllium were below Eco-SSLs for protection of soil invertebrates.

Individual boron concentrations in soil samples taken from Kern NWR were higher (range = 34.3 – 72 ug/g dw) than the averages reported for western U.S. soils, but were still at the lower end of the overall range of <20 – 300 ug/g dw (Shacklette and Boerngen 1984). Boron concentrations in soils from Kern NWR were all >15 ug/g dw considered to be a threshold for observable impacts on growth of sensitive plant species (Miwa et al. 2007). However, the concentrations of boron observed in these samples were considered to be below levels of concern for mountain plovers because their direct exposure via incidental ingestion of soil is only fraction of their diet.

Average concentrations of selenium in soil samples were within the ranges reported by Shacklette and Boerngen (1984), but some were elevated relative to background concentrations for soils specific to California (Bradford et al. 1996). Selenium concentrations measured in soil samples from Imperial Valley were high (Table 2), particularly in the actively cultivated fields where even the minimum concentrations (≥ 0.55 ug/g dw) were greater than the maximum reported by Bradford et al. (1996) for background soils in California. All of the off-refuge fields sampled exceeded the Eco-SSL for protection of plants, but invertebrates and avian ecological receptor levels were not exceeded. While the elevated selenium levels observed in the actively cultivated fields from Imperial Valley are not based on geology alone (USDOI 1998), conditions in the Imperial Valley favor the accumulation of selenium introduced by irrigation with water from the Colorado River, which has elevated levels of selenium (USDOI 1998). Selenium concentrations in soil from Kern NWR were consistent with background concentrations reported by Bradford et al. (1996) for California soils (mean = 0.058 ug/g dw; range = 0.015 – 0.430 ug/g dw).

Average strontium concentrations measured in fields from California ranged from 224 ug/g dw to 441 ug/g dw (range = 216 – 504 ug/g dw), all of which exceeded the average background concentration for California soils (128 ug/g dw) and some were above the range for background (20 – 271 ug/g dw). Strontium is ubiquitous in the environment and present in nearly all types of rocks and soils (ATSDR 2004). Elevated concentrations in soils may result from atmospheric deposition of particles from coal fired power plants, pyrotechnic devices, and dust from areas where phosphate fertilizers have been applied. Releases directly to soil are primarily from application of solid waste and the use of phosphate fertilizers (ATSDR 2004). The toxicological implications for mountain plovers of elevated strontium concentrations in Imperial Valley soils are not known.

Average lead concentrations (14.0 – 18.9 ug/g dw) in soil samples collected from Imperial Valley were slightly above the Eco-SSL for the protection of birds; but it is unlikely that the lead concentrations in the Imperial Valley soil samples pose a significant threat to mountain plover (Eisler 1988b). Average concentrations of vanadium in soils from all of the California (34.8 – 42.0 ug/g dw) were greatly above the Eco-SSL of 7.8 ug/g dw for the protection of birds, but interpreting the risk posed by vanadium at measured concentrations in soil samples is difficult because information on vanadium's toxicity to wild terrestrial birds is scarce. In studies with domestic chickens (*Gallus domesticus*), adverse effects depended on the chemical form of vanadium and ranged from impaired growth in newly hatched to reduced egg production to mortality (Rattner et al. 2006).

No background cadmium values were reported by Shacklette and Boerngen (1984). However, concentrations measured in soils from actively cultivated fields in Imperial Valley exceeded both the geometric mean and the maximum reported for background in soils from California (Bradford et al. 1996). In addition, all of the samples collected from agricultural fields in Imperial Valley had cadmium concentrations exceeding the Eco-SSL for birds (0.77 ug/g dw) with averages ranging from 1.93 ug/g dw to 3.30 ug/g dw. In comparison, cadmium concentrations in soils from more passively cultivated fields in Imperial Valley (SBSS NWR)

and the Central Valley (Kern NWR) were below 0.55 ug/g dw (Table 2) and were below levels of concern (the Eco-SSL) for avian wildlife (USEPA 2005d).

Average copper concentrations in soils from fields were between 17.5 ug/g dw and 33.7 ug/g dw, which are higher than the geometric mean, but well within the range of copper concentration reported for western U.S. soils (mean = 2.07 ug/g dw; range = 2 – 300 ug/g dw). These concentrations were only slightly higher than the range of copper concentrations reported for non-contaminated sites (Kabata-Pendias and Pendias 1992), and either did not or only slightly exceeded the Eco-SSL for avian wildlife (28 ug/kg dw). Copper deficiency or excess can be harmful to organisms, although reported toxic effects from excess dietary exposure in birds in field conditions are rare (Eisler 1997; USDOJ 1998).

Average phosphorus concentrations measured in soil samples (803 – 3,316 ug/g dw) were elevated compared to the geometric mean concentration of 320 ug/g but were within the range of 40 ug/g dw to 4,500 ug/g dw for background soils in the western U.S. (Shacklette and Boerngen 1984). Phosphorus occurs naturally and is a primary ingredient in fertilizers (UNEP 2001). Birds require a specific level of phosphorus and the availability of phosphorus depends on the soil's pH. Lower pH conditions allow aluminum to become soluble, which if taken up by birds, competes with phosphorus and calcium needed for eggshell structure and proper growth during embryo development (Schwarzbach et al. 2006). Soils in the western U.S. are typically alkaline that readily alleviates this concern.

Zinc concentrations in soil samples from actively cultivated fields in Imperial Valley (range = 392 – 9,310 ug/g dw) were much higher than the geometric mean background concentration of 55 ug/g reported for the western U.S. (Shacklette and Boerngen 1984) and all samples greatly exceeded the respective Eco-SSLs for protection of birds (46 ug/kg dw) and soil invertebrates (120 ug/g dw). Zinc concentrations in soil samples from refuges were much lower, although several samples exceeded the Eco-SSL for protection of birds. Zinc occurs naturally but elevated concentrations in soil may be the result of application of certain fertilizers, atmospheric

deposition from smelting operations of galvanized metal, or application of domestic and industrial biosolids (Eisler 1993).

Although not tested for statistical significance, it appears that concentrations of arsenic, barium, beryllium, cobalt, copper, lead, magnesium, manganese and selenium were higher in soils from Imperial Valley fields than in the field from the Kern NWR. Conversely, concentrations of sodium, boron, and potassium were higher in soil from the field at Kern NWR. These differences may reflect regional variation in conditions and influences. For example, soils at Kern NWR were generally saltier than those in Imperial Valley, while soils in the Imperial Valley had somewhat higher concentrations of selenium, possibly the result of irrigation with Colorado River water (USDOI 1998). Within Imperial Valley, soils from actively cultivated fields had higher concentrations of calcium, cadmium, phosphorus, sulfur, and zinc than soils from the more passively managed field on SBSS NWR. These differences appear to reflect the influence of land management practices. For example, calcium, phosphorus, sulfur, and zinc are among the nutrients added as fertilizers to support the growth of Bermuda grass crops. Sulfur is also the most commonly used fumigant on Bermuda grass crops, and cadmium may be present at elevated levels if biosolids are used on the fields.

Soils-Organic Compounds

Most organic pesticides were either not detected or were at very low concentrations in soil samples (Appendix 2; Table 3). The herbicide trifluralin is the third most heavily used pesticide in Imperial Valley for alfalfa and Bermuda grass crops (PAN 2010). Trifluralin was detected in all soil samples collected from Imperial Valley fields, including at SBSS NWR (Table 3). However, concentrations (range = 0.514 – 1.3 ng/g dw) were below mean concentrations detected in soils collected in 1972 for the National Soils Monitoring Program (<10 ng/g; Carey et al. 1979) and well below concentrations reported for soils the same year from fields in California (650 – 6,450 ng/g dw; Carey et al. 1979). Concentrations measured in soils from SBSS NWR were the same as concentrations measured in actively farmed fields nearby. This suggests that concentrations measured in the actively cultivated fields were not from recent applications. Instead, trifluralin concentrations measured on those fields and on SBSS NWR appear to reflect

residual trifluralin transported by from area-wide non-point sources such as drift, and/or irrigation water.

Residues of DDT (o,p' and p,p') and its metabolites DDD (o,p' and p,p') and DDE (o,p' and p,p') were detected in virtually all soil samples, with higher concentrations occurring in soil samples from the Imperial Valley, including SBSS NWR (Table 3, Figure 15) than from the Central Valley. DDT was banned for use in the U.S. in 1972 but it is still transported via air, water, and animal movements from countries that continue to use the insecticide. DDE is a breakdown product of DDT and residues in soil may also be the result of historical agricultural use because DDE is extremely persistent (USDOI 1998). Twelve years after DDT was banned, total DDT concentrations in soils from agricultural fields in California were between 1.0 ng/g dw and 31,273 ng/g dw, which are orders of magnitude higher than concentrations measured in this study. However, total DDT concentrations in samples collected in 1984 from Imperial Valley were between 37 ng/g dw and 495 ng/g dw (average = 228 ng/g dw; Mischke et al. 1985), which is not greatly different from concentrations measured in Imperial Valley soils collected for this study in 2006 (average = 140 ng/g dw). The fraction of total DDT that is the parent compound (p,p'-DDT) has declined from approximately 20% in samples collected in 1984 (Mischke et al. 1985) to approximately 7% in samples collected from Imperial Valley in 2006 (Figure 16 and Table 3). While there was an approximately 2-fold decline in total DDT concentrations between 1984 and 2006, there was an approximately 3-fold decline in concentrations of parent compound (DDT), which is consistent with an approximately 15 year half-life for DDT in soil. Although concentrations of total DDT in Imperial Valley soils have declined over time, they still ranged from slightly below to more than ten-times the Eco-SSL of 22 ng/g dw for protection of avian species.

Dieldrin was detected in all soil samples from fields. Aldrin, which is readily converted to dieldrin in the environment, was detected in approximately half of the samples. Although frequently detected, concentrations of dieldrin and aldrin were less than the Eco-SSL of 22 ng/g (0.022 ug/g dw) for avian species. The Eco-SSLs for DDT and dieldrin are designed to address risk posed by exposure to these contaminants, which is primarily through bioaccumulation in




Figure 15. DDT isomer concentrations in soils from California (1985 and 2006)




Figure 16. Percentage of DDT isomers comprising total DDT in California

food web organisms and incidental ingestion of soil. There are no Eco-SSLs for other potentially bioaccumulative contaminants detected in soil samples. These include substances such as chlordanes and HCHs that were detected at low levels. While concentrations of OC pesticides other than dieldrin and DDT may appear low in the soils, they may still pose a risk to plovers by bioaccumulating in the tissues of the soil invertebrates that plovers consume.

PCB concentrations in soils were low (arithmetic mean concentrations <110 ng/g dw; Table 3 and Appendix 3) and below levels associated with soils at contaminated sites (ATSDR 2000). Concentrations, as the sum of the congeners, in the most actively worked fields (Imperial Valley off-refuge) were between 36 ng/g dw and 175 ng/g dw and appeared to be significantly higher than concentrations measured in less intensively worked fields on refuge lands nearby (range = 15.3 – 20.3 ng/g dw). The field on Kern NWR, which has not been active for decades, had the lowest concentration (range = 4.75 – 8.84 ng/g dw). Atmospheric deposition contributes to PCB concentrations observed in environmental media. However, there may be additional sources for actively cultivated fields in Imperial Valley such as routine use of heavy farm equipment.

Table 2. Arithmetic mean and range of concentrations of select inorganics¹ (ug/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006, for comparison with geometric mean and range of inorganics in soils from the western United States as reported by Shacklett and Boerngen (1984).

| Location and Samples | | Al | As | B | Ba | Be | Ca | Cd | Co | Cr | Cu | Fe | Hg | K |
|--|---------|----------|-------|------|------|-------|---------|-------|------|------|------|----------|-------|--------|
| Imperial Valley Off-refuge J20A - E (N=5) | ave | 19,900 | 6.56 | 7.42 | 313 | 1.11 | 81,400 | 1.93 | 7.82 | 30.9 | 31.2 | 17,520 | 0.016 | 4,424 |
| | min | 17,100 | 6.07 | 4.71 | 307 | 1.06 | 78,600 | 1.48 | 7.30 | 27.7 | 29.5 | 16,300 | 0.015 | 3,880 |
| | max | 23,500 | 7.05 | 10.4 | 324 | 1.20 | 84,800 | 2.54 | 8.41 | 33.4 | 32.8 | 18,900 | 0.016 | 5,050 |
| Imperial Valley Off-refuge J22A - E (N=5) | ave | 19,280 | 5.90 | 9.49 | 317 | 1.142 | 83,480 | 3.30 | 7.54 | 41.3 | 33.7 | 17,280 | 0.018 | 4,812 |
| | min | 17,900 | 5.74 | 7.73 | 309 | 1.050 | 77,000 | 1.87 | 7.17 | 27.5 | 28.5 | 16,900 | 0.017 | 4,100 |
| | max | 20,000 | 6.05 | 11.2 | 326 | 1.240 | 92,800 | 4.66 | 8.04 | 53.0 | 38.7 | 17,700 | 0.019 | 5,250 |
| Imperial Valley Off-refuge K18A-E (N=5) | ave | 23,180 | 5.72 | 6.82 | 292 | 1.25 | 77,540 | 2.65 | 8.24 | 35.9 | 29.2 | 19,220 | 0.016 | 4,812 |
| | min | 19,700 | 5.00 | 5.81 | 266 | 1.19 | 69,000 | 1.91 | 7.61 | 30.9 | 24.6 | 17,800 | 0.015 | 4,050 |
| | max | 24,900 | 6.25 | 8.92 | 313 | 1.30 | 91,600 | 3.78 | 8.50 | 42.0 | 34.3 | 19,800 | 0.017 | 5,170 |
| Imperial Valley On-refuge; SBSS NWR SB420A - E (N=5) | ave | 24,420 | 7.33 | 6.94 | 241 | 1.24 | 50,560 | 0.389 | 9.7 | 22.8 | 27.2 | 20,980 | 0.018 | 5,202 |
| | min | 22,500 | 6.57 | 5.21 | 234 | 1.17 | 50,000 | 0.370 | 9.27 | 21.5 | 25.3 | 20,000 | 0.018 | 4,690 |
| | max | 29,000 | 8.25 | 10.0 | 249 | 1.41 | 51,000 | 0.435 | 10.5 | 25.6 | 29.6 | 22,900 | 0.019 | 5,960 |
| Central Valley On-refuge; Kern NWR K15A - E (N=5) | ave | 17,180 | 2.92 | 48.1 | 126 | 0.486 | 36,460 | 0.404 | 5.64 | 26.1 | 17.5 | 15,880 | 0.014 | 5,766 |
| | min | 12,800 | 2.03 | 34.3 | 110 | 0.363 | 29,600 | 0.349 | 3.85 | 22.0 | 10.6 | 12,100 | 0.009 | 4,580 |
| | max | 23,600 | 4.23 | 72.0 | 152 | 0.684 | 45,300 | 0.543 | 8.61 | 31.8 | 28.4 | 21,800 | 0.027 | 7,620 |
| western U.S. background soils | geomean | 58,000 | 5.5 | 23 | 580 | 0.68 | 18,000 | 0.26* | 7.1 | 41 | 2.07 | 21,000 | 0.46 | 18,000 |
| | min | 5,000 | <0.10 | <20 | 70 | <1 | 600 | 0.05* | <3 | 3 | 2 | 1,000 | <0.01 | 1,900 |
| | max | >100,000 | 97 | 300 | 5000 | 15 | 320,000 | 1.7* | 50 | 2000 | 300 | >100,000 | 4.6 | 63,000 |

¹ See Appendix 1 for complete list of inorganic constituent concentrations by sample

* California background (Bradford et al. 1996)

Table 2. cont.

| Location and Samples | | Mg | Mn | Mo | Na | Ni | P | Pb | S | Se | Sr | Zn |
|--|---------|-----------|-----------|-----------|-----------|-----------|----------|-----------|----------|-----------|-----------|-----------|
| Imperial Valley Off-refuge J20A - E (N=5) | ave | 14,240 | 485 | 0.91 | 1,222 | 18.5 | 1,872 | 14.0 | 1,420 | 0.68 | 407 | 2,564 |
| | min | 13,200 | 461 | 0.51 | 804 | 17.7 | 1,370 | 13.4 | 1,190 | 0.56 | 392 | 1,130 |
| | max | 14,900 | 506 | 1.22 | 2,640 | 19.8 | 2,260 | 14.9 | 1,610 | 0.76 | 418 | 3,900 |
| Imperial Valley Off-refuge J22A - E (N=5) | ave | 14,040 | 500 | 1.049 | 1,275 | 19.5 | 3,136 | 14.1 | 2,094 | 0.915 | 441 | 5,708 |
| | min | 12,900 | 482 | 0.505 | 995 | 18.5 | 2,740 | 13.4 | 1,920 | 0.717 | 399 | 1,170 |
| | max | 15,000 | 548 | 1.390 | 1,460 | 21.3 | 3,650 | 14.9 | 2,330 | 1.11 | 504 | 9,310 |
| Imperial Valley Off-refuge K18A-E (N=5) | ave | 14,100 | 460 | 0.507 | 958 | 19.8 | 2,488 | 16.2 | 1,184 | 0.605 | 391 | 1,432 |
| | min | 12,700 | 436 | 0.505 | 804 | 19.0 | 1,840 | 14.4 | 1,080 | 0.550 | 338 | 392 |
| | max | 14,700 | 478 | 0.510 | 1,140 | 20.1 | 3,060 | 17.6 | 1,300 | 0.716 | 482 | 2,700 |
| Imperial Valley On-refuge; SBSS NWR SB420A - E (N=5) | ave | 14,920 | 499 | 1.22 | 1,428 | 21.9 | 803 | 18.9 | 814 | 0.392 | 224 | 78.7 |
| | min | 14,400 | 478 | 1.10 | 1,190 | 20.4 | 781 | 17.6 | 646 | 0.366 | 216 | 75.2 |
| | max | 16,000 | 534 | 1.29 | 1,700 | 23.3 | 822 | 21.0 | 1,180 | 0.418 | 233 | 84.3 |
| Central Valley On-refuge; Kern NWR K15A - E (N=5) | ave | 12,980 | 398 | 1.24 | 5,484 | 21.8 | 1,848 | 6.52 | 754 | 0.087 | 314 | 56.7 |
| | min | 9,600 | 317 | 0.51 | 3,180 | 17.4 | 1,410 | 4.94 | 387 | 0.075 | 235 | 44.4 |
| | max | 18,600 | 531 | 3.08 | 8,950 | 28.7 | 2,120 | 8.27 | 1,340 | 0.114 | 394 | 79.4 |
| western U.S. background soils | geomean | 7,400 | 380 | 0.85 | 9,700 | 15 | 320 | 17 | 1,300 | 0.23 | 200 | 55 |
| | min | 300 | 30 | <3 | 500 | <5 | 40 | <10 | <800 | <0.1 | 10 | 10 |
| | max | >100,000 | 5,000 | 7 | 100,000 | 700 | 4,500 | 700 | 48,000 | 4.3 | 3000 | 2100 |

Table 3. Arithmetic mean and range of concentrations of selected organic analytes (ng/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006.

| Location and samples | | DDD (o,p' + p,p') | DDE (o,p' + p,p') | DDT (o,p' + p,p') | DDTs- total ² | Chlordanes ³ | HCHs ⁴ | Dieldrin & Aldrin ⁵ | PCBs - Aroclors | PCBs congeners | Trifluralin | Chlorpyrifos |
|--|-----|-------------------|-------------------|-------------------|--------------------------|-------------------------|-------------------|--------------------------------|-----------------|----------------|-------------|--------------|
| Imperial Valley Off-refuge J20A - E (N=5) | ave | 11.1 | 232 | 14.2 | 258 | 5.56 | 0.112 | 1.94 | 42.2 | 50.0 | 0.97 | 0.798 |
| | min | 6.83 | 134 | 8.51 | 155 | 4.09 | 0.093 | 1.44 | 27.3 | 36.0 | 0.549 | 0.266 |
| | max | 14.7 | 348 | 23.7 | 376 | 7.60 | 0.132 | 2.81 | 55.8 | 66.4 | 1.30 | 1.07 |
| Imperial Valley Off-refuge J22A - E (N=5) | ave | 14.3 | 152 | 9.23 | 176 | 4.76 | 0.105 | 2.08 | 37.5 | 114 | 1.02 | 0.707 |
| | min | 4.40 | 63 | 3.01 | 70.0 | 1.43 | 0.077 | 0.64 | 14.5 | 74.2 | 0.850 | 0.464 |
| | max | 24.7 | 253 | 16.8 | 292 | 7.63 | 0.126 | 3.45 | 59.4 | 175 | 1.20 | 1.05 |
| Imperial Valley Off-refuge K18A-E (N=5) | ave | 1.72 | 18.6 | 1.81 | 22.1 | 0.360 | 0.086 | 0.175 | 12.0 | 64.3 | 0.737 | 0.264 |
| | min | 1.20 | 11.3 | 1.19 | 14.4 | 0.301 | 0.077 | 0.139 | 8.87 | 45.4 | 0.573 | 0.177 |
| | max | 2.76 | 22.3 | 2.43 | 26.3 | 0.400 | 0.093 | 0.231 | 15.3 | 85.1 | 1.12 | 0.325 |
| Imperial Valley On-refuge; SBSS NWR SB420A - E (N=5) | ave | 4.51 | 86.9 | 11 | 103 | 0.301 | 0.378 | 0.276 | 12.5 | 17.5 | 0.829 | 0.619 |
| | min | 4.17 | 60.5 | 9.44 | 74.2 | 0.269 | 0.305 | 0.248 | 10.9 | 15.3 | 0.514 | 0.489 |
| | max | 5.02 | 111 | 13 | 129 | 0.336 | 0.410 | 0.289 | 13.7 | 20.3 | 1.30 | 0.798 |
| Central Valley On-refuge; Kern NWR K15A - E (N=5) | ave | 0.488 | 1.95 | 1.93 | 4.37 | 0.483 | 0.076 | 0.127 | 6.67 | 6.61 | <0.260 | 0.058 |
| | min | 0.077 | 0.55 | 0.151 | 0.78 | 0.333 | 0.065 | 0.079 | 4.76 | 4.75 | <0.252 | <0.026 |
| | max | 0.85 | 3.74 | 4.53 | 9.12 | 0.70 | 0.105 | 0.240 | 8.91 | 8.84 | <0.266 | 0.110 |

¹ See Appendix 1 for complete list of organic constituent concentrations by sample

² DDTs - total = [o,p'-DDD] + [p,p'-DDD] + [o,p'-DDE] + [p,p'-DDE] + [o,p'-DDT] + [p,p'-DDT]

³ Chlordanes = [cis-chlordane] + [trans chlordane] + [cis-nonachlor] + [trans-nonachlor] + [oxychlordane]

⁴ HCHs = [alpha HCH] + [beta HCH] + [delta HCH]

⁵ Dieldrin & Aldrin combined because Arden is readily converted to Dieldrin in the environment.

Terrestrial Invertebrates-Inorganic Elements

All California sampling sites had at least one invertebrate sample exceeding 1,000 ug/g dw of aluminum (Table 4; Appendix 4) and concentrations in those from SBSS NWR averaged 7,150 ng/g dw (Table 4). A dietary concentration above 1,000 ug/g dw is considered toxic to birds (Sample et al. 1996; Sparling and Lowe 1996). High levels of aluminum in the diet can affect the metabolism of calcium and phosphorus, which results in loss of eggshell quality, reduced growth in chicks, and/or decreased egg-laying in female birds (Sparling and Lowe 1996). Even so, a debate still remains on the risk to birds associated with elevated aluminum concentrations in the diet because bioavailability is not well known.

Arsenic is a common ingredient in herbicides and insecticides, which represent the largest source in agricultural fields (Eisler 1988a); but average concentrations of arsenic in terrestrial invertebrate samples from all wintering areas were below levels of concern for avian species (Table 4). The dietary NOAEL for arsenic, as a daily dose, is estimated to be 2.46 ug/g/day as copper acetoarsenite for the brown-headed cowbird (*Molothrus ater*), which is considered a sensitive species to arsenic (Sample et al. 1996). Mallards (*Anas platyrhynchos*) had an estimated NOAEL as daily dose of 5.14 ug/g/d with sodium arsenite. One individual invertebrate sample from SBSS NWR had an arsenic concentration of 4.5 ug/g dw. Using a body weight and a daily food ingestion rate more specific to plovers (Dunning 1984, Nagy 2001), the NOAEL for cowbirds translates into a dietary concentration for mountain plovers of approximately 30 ug/g dw. The maximum concentration observed in samples from California wintering grounds was 4.5 ug/g dw.

Barium concentrations in three individual invertebrate samples (47.8 ug/g, 33.8 ug/g, and 137 ug/g dw) were higher than any of the remaining 21 samples (Appendix 4), but these concentrations were well below an estimated NOAEL of 2,000 ug/g (Sample et al. 1996). In addition, beryllium was very low or below the detection limit for invertebrate samples (Wiemeyer et al. 2004).

Most research on dietary ingestion of boron by birds has occurred in waterfowl as a result of agricultural irrigation drainwater concerns (USDOI 1998); but, in laboratory studies where adult domestic chickens consumed 875 ug/g of boron as boric acid for six days, the only ill effect was egg production cessation, which returned to normal 14 days after the boron diet was removed (Eisler 1990b). However, boron is teratogenic if injected into domestic chicken eggs (Eisler 1990b). Boron concentrations in all but one of the invertebrate samples collected at California wintering grounds were below 30 ug/g (Table 4; Appendix 4) shown to affect growth in mallard ducklings (Sample et al. 1996; USDOI 1998).

Cadmium concentrations measured in invertebrate samples were all <1.2 ug/g dw. Laboratory studies suggest that dietary intake of <1 ug/g should not lead to toxicological effects in birds (Furness 1996). Chromium concentrations in soil invertebrates were <3.5 ug/g in all samples but one (19 ug/g dw), but this concentration does not represent a risk to birds even if considered in its most toxic hexavalent form (Eisler 1986).

Average copper concentrations in terrestrial invertebrates varied from 41.2 ug/g dw to 89.1 ug/g dw (Table 4). Little information exists on the toxicity of excess dietary copper to wild birds, although information based on poultry studies indicate that a diet of 300 ug/g is the maximum tolerable limit, which is well below copper concentrations in collected invertebrate samples. Typically, copper deficiency is more of a concern in birds' diet because it is a necessary nutrient (Eisler 1997; USDOI 1998).

Lead was detected infrequently, and in all but two samples, concentrations ranged from not detected (mean detection limit = 1.8 ug/g dw) to 2.6 ug/g dw. Concentrations in two samples from Imperial Valley fields were noticeably higher, at 7.2 ug/g dw in one off-refuge field and 11 ug/g dw in the on-refuge field at SBSS NWR (Appendix 4). Toxicological benchmarks indicate that dietary concentration of 10 ug/g as lead acetate has no effect on egg hatching success in Japanese quail (*Coturnix japonica*), whereas a concentration of 100 ug/g of lead impaired reproduction. In other bird species such as domestic chickens and pheasants (*Phasianus colchicus*), the growth of precocial hatchlings exposed to dietary lead levels of 500 ug/g was not

affected (Hoffman et al. 1985). Dietary ingestion from food containing biologically incorporated lead is unlikely to result in toxicity, particularly at the level measured in invertebrates from fields in Imperial Valley (Beyer et al. 1988; Eisler 1988b).

Terrestrial invertebrates from SBSS NWR were higher in iron and magnesium than at other wintering sites where invertebrates were collected. These elements are important nutrients and required by living organisms in large amounts (Hulse et al. 1980). Manganese is also an important nutrient but is required in much smaller amounts and can be toxic at high concentrations. Manganese was highest in terrestrial invertebrates (maximum concentration = 226 ug/g dw) from SBSS NWR (Table 4; Appendix 4). With an estimated dry weight-based food ingestion rate of 0.180 g/g_{BW}-d for mountain plovers, the maximum observed manganese concentration in invertebrate samples translates into a daily dose rate for mountain plovers of approximately 41 ug/g_{BW}-d, which is much less than the daily dose of 977 ug/g/day shown to be the NOAEL for Japanese quail (Sample et al. 1996).

Mercury concentrations in invertebrates were below the detection limit (0.12 ug/g dw), but this is approximately ten-times greater than concentrations measured in co-located soils. Additionally, the wet weight-based detection limit for mercury in invertebrate samples was 0.06 ug/g, which exceeds dietary screening levels considered by USFWS (2003b) for the protection of western snowy plover (*Charadrius alexandrinus nivosus*) in California (0.009 ug/g or 0.026 ug/g depending on the uncertainty factor). However, the dry weight-based detection limit for mercury in invertebrates (0.12 ug/g) translates into a potential ingestion rate of 0.022 ug/g/day for mountain plovers, which is below the lowest NOAEL (0.030 ug/g/day) identified by USEPA (1995) for chronic exposure by avian species.

Molybdenum concentrations were low (Eisler 1989; USDOJ 1998) in terrestrial invertebrate samples. Information available on nickel indicates that concentrations measured in terrestrial invertebrates were low, but whether they were below levels of concern for wild birds is not known (Weber and Reid 1968; Cain and Pafford 1981).

Table 4. Arithmetic mean and range of concentrations of select inorganics¹ (ug/g dry weight) in invertebrate samples collected from mountain plover wintering sites in California, 2006.

| Location and Samples | Statistical Function | Al ug/g dw | As ug/g dw | B ug/g dw | Ba ug/g dw | Be ug/g dw | Cd ug/g dw | Cr ug/g dw | Cu ug/g dw | Hg ug/g dw | Mn ug/g dw |
|--|-----------------------------|-----------------------|-----------------------|----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Imperial Valley Off-refuge J20A - E (N=5) | average | 990 | 0.54 | 11 | 13.5 | <0.1 | 0.528 | 1.08 | 56.3 | <0.12 | 33.4 |
| | minimum | 150 | <0.3 | <10 | 1.8 | <0.1 | 0.2 | <0.5 | 25 | <0.12 | 13 |
| | maximum | 3120 | 1.4 | 20 | 47.8 | <0.1 | 0.84 | 3.5 | 94.9 | <0.12 | 68 |
| Imperial Valley Off-refuge J22A - E (N=5) | average | 925.4 | 1.02 | 12.1 | 11.2 | <0.1 | 0.684 | 1.44 | 89.1 | <0.12 | 36.6 |
| | minimum | 343 | <0.6 | <7 | 4.8 | <0.1 | 0.45 | 0.6 | 32.9 | <0.12 | 25 |
| | maximum | 1810 | 2.1 | 37 | 18 | <0.1 | 0.9 | 2.2 | 208 | <0.12 | 49 |
| Imperial Valley Off-refuge K18A-E (N=5) | average | 1344 | 0.42 | 4.3 | 12.8 | <0.1 | 0.786 | 2.4 | 48.3 | <0.12 | 43.8 |
| | minimum | 1190 | <0.5 | 6 | 10 | <0.1 | 0.5 | 1.5 | 26 | <0.12 | 35 |
| | maximum | 1540 | 0.75 | 6 | 15 | <0.1 | 1.1 | 2.8 | 78.8 | <0.12 | 50 |
| Imperial Valley On-refuge; SBSS NWR SB420A - E (N=5) | average | 7150 | 1.59 | 7.5 | 43.8 | 0.152 | 0.582 | 5.42 | 41.2 | <0.12 | 91.7 |
| | minimum | 3620 | 0.6 | <4 | 13 | <0.1 | <0.1 | 1 | 24 | <0.12 | 46 |
| | maximum | 24500 | 4.5 | 18 | 137 | 0.56 | 1.2 | 19 | 63.7 | <0.12 | 226 |
| Central Valley On-refuge; Kern NWR K15A - E (N=5) | average | 1585 | 0.896 | 8.2 | 39.9 | <0.1 | 0.916 | 1.96 | 59.9 | <0.12 | 49.9 |
| | minimum | 855 | <0.6 | <6 | 16 | <0.1 | 0.6 | 1 | 31 | <0.12 | 34 |
| | maximum | 2010 | 2.3 | 10 | 68.7 | <0.1 | 1.2 | 2.7 | 102 | <0.12 | 64.2 |

Table 4, cont.

| Location and Samples | Statistical Function | Mo ug/g dw | Ni ug/g dw | Pb ug/g dw | Se ug/g dw | Sr ug/g dw | V ug/g dw | Zn ug/g dw |
|--|-----------------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|----------------------|-----------------------|
| Imperial Valley Off-refuge J20A - E (N=5) | average | 1.4 | 1.18 | 2.02 | 2.38 | 71.1 | 1.99 | 413 |
| | minimum | <2 | 0.6 | <0.8 | 1.4 | 9 | <0.5 | 194 |
| | maximum | 3 | 2.4 | 7.2 | 3 | 301 | 6.1 | 592 |
| Imperial Valley Off-refuge J22A - E (N=5) | average | 2 | 1.19 | 0.42 | 2.92 | 25.6 | 2.08 | 550 |
| | minimum | <2 | <0.5 | <0.5 | 1.6 | 20 | 0.7 | 427 |
| | maximum | 6 | 1.8 | 0.5 | 4.4 | 32.7 | 3.3 | 747 |
| Imperial Valley Off-refuge K18A-E (N=5) | average | 1 | 0.6 | 0.763 | 1.29 | 27.4 | 2.68 | 405 |
| | minimum | <0.5 | <0.5 | <0.7 | 0.93 | 22.3 | 2.2 | 273 |
| | maximum | 1 | 1 | 2 | 1.6 | 38.8 | 3.2 | 606 |
| Imperial Valley On-refuge; SBSS NWR SB420A - E (N=5) | average | 1.4 | 3.88 | 3.32 | 1.40 | 85.8 | 11.5 | 181 |
| | minimum | 1 | 1.5 | 1 | 0.78 | 27 | 3.4 | 138 |
| | maximum | 2 | 11 | 11 | 2.8 | 298 | 36 | 215 |
| Central Valley On-refuge; Kern NWR K15A - E (N=5) | average | 1.2 | 2.5 | 0.62 | 0.608 | 212 | 4.2 | 159 |
| | minimum | 1 | 2.2 | <0.6 | 0.5 | 43.4 | 2.7 | 118 |
| | maximum | 2 | 2.7 | 1 | 0.91 | 413 | 5.4 | 213 |

¹See Appendix 4 for complete list of inorganic constituent concentrations by sample.

Average concentrations of selenium in invertebrate samples (0.608 – 2.92 ug/g dw) were consistently higher than average concentrations of selenium in co-located soil samples (0.087 – 0.915 ug/g dw). Birds may suffer reproductive impairment at a dietary range of 3-8 ug/g (USDOJ 1998); and a range of 10 –15 ug/g dietary selenium exposure is the toxicity threshold for non-breeding birds exposed to winter stress (USDOJ 1998). However, the selenium in invertebrate samples indicate that it is likely available for uptake and accumulation by organisms that constitute the mountain plover food web but overall does not present a risk to the birds.

Strontium concentrations were below 50 ug/g dw in most of the invertebrate samples collected. However, noticeably higher concentrations were observed with invertebrates from Kern NWR (four out of five samples = 127 – 413 ug/g dw), and in two samples from Imperial Valley (301 ug/g dw from an off-refuge field and 298 ug/g dw from SBSS NWR). Strontium is associated with calcium or potassium rich rock with weathering and atmospheric deposition being the primary input (Mora 2003). It is considered to have low toxicity (Wennig and Kirsch 1950), but it is unknown if the strontium levels measured in invertebrates pose a threat to mountain plover.

Little information exists on the toxicity of vanadium to wild birds, particularly terrestrial species. Research on acute toxicity and chronic toxicity of vanadium compounds to mallards and Canada geese (*Branta canadensis*) indicate that dietary effect levels may be greater than 250 ug/g, which is much higher than concentrations measured in invertebrates samples collected at the wintering grounds (Rattner et al. 2006).

Because zinc concentrations in soil samples from the Imperial Valley were higher than those from other wintering sites, it is not unexpected that terrestrial invertebrate samples from this area also had higher zinc concentrations. Zinc is an essential element but it can bioaccumulate in organisms and cause adverse effects at both high and low concentrations (USDOJ 1998). Concentrations of zinc in invertebrates from the Imperial Valley sites were above levels shown to cause immunosuppression in young domestic chicks that were fed zinc at 178 ug/g for three weeks, although growth was not affected (Eisler 1993). Dietary concentrations of zinc that may affect the mountain plover are unknown.

Terrestrial Invertebrates-Organic Compounds

A few OCs were detected in terrestrial invertebrate samples (Table 5; Appendix 5). As with the soil samples, higher OC concentrations occurred samples from Imperial Valley, including SBSSNWR. DDE and two PCB congeners (#28 and #175) were detected in all of the invertebrate samples, reflecting the wide-spread occurrence of DDT and PCBs in general. PCB congeners #28 and #175 are not one of the more toxic co-planer PCB congeners. DDE concentrations in individual invertebrate samples from Imperial Valley ranged from approximately 48 ng/g dw to 1,310 ng/g dw, and averages ranged from 120 ng/g dw to 732 ng/g dw. The DDE concentration in invertebrates tended to be between 0.5 to 5.0 times the concentrations in corresponding soil samples (Figure 17). Total DDT concentrations were also higher in invertebrates than the soil collected from the same location (Figure 18).

Trifluralin, a dinitroaniline herbicide used to control annual grasses and broadleaf weeds, is widely used in the Imperial Valley. This herbicide is known to be persistent in soil, and current product labeling allows several applications during the growing season (USEPA 1996).

Trifluralin was detected in most of the insect samples (13 out of 15) from actively cultivated fields in Imperial Valley, less frequently (1 out of 5) in the Imperial Valley field on SBSS NW, and not at all in the field at Kern NWR. Concentrations of trifluralin in invertebrates were 10 to 100 times the concentrations in corresponding soil samples (Figure 19).

According to the reregistration eligibility decision (RED) for trifluralin, the chronic level of concern was exceeded for birds. Additionally, trifluralin application rates specified on the label may adversely affect reproduction (USEPA 1996). As part of the RED, USEPA requires avian reproduction studies. These studies used northern bobwhite (*Colinus virginianus*) and mallard duck and revealed that cracked eggs were the common endpoint affected (USEPA 1996).

Data from invertebrate samples demonstrate the potential for DDE and trifluralin to bioaccumulate in the tissues of soil invertebrates and be introduced into the diet of mountain plovers while wintering in Imperial Valley.

Table 5. Arithmetic mean and range of concentrations of the most frequently detected organics¹ (ng/g dry weight) in invertebrate samples collected from mountain plover wintering sites in California, 2006.

| Location and samples | | PCB# 28 | PCB# 72 | PCB# 138/160 | PCB# 172 | PCB# 175 | PCB# 180 | PCB - total (Σ congeners) | p,p'-DDE | Total DDT | Trifluralin |
|--|-----|---------|---------|--------------|----------|----------|----------|--------------------------------------|----------|-----------|-------------|
| Imperial Valley Off-refuge J20A - E (N=5) | ave | 12.4 | 19 | 17.2 | 6.97 | 20.2 | 5.51 | 222 | 322 | 359 | 98.3 |
| | min | 8.76 | 2.35 | 8.62 | 0.85 | 10.6 | 3.06 | 45.9 | 137 | 158 | <16.0 |
| | max | 20.8 | 32 | 26.2 | 12.0 | 32.1 | 11.7 | 767 | 478 | 537 | 256 |
| Imperial Valley Off-refuge J22A - E (N=5) | ave | 18.0 | 39 | 17.3 | 15.4 | 16.8 | 3.18 | 129 | 732 | 759 | 83.4 |
| | min | 11.6 | 1.02 | 7.66 | 5.09 | 7.57 | 1.09 | 92.3 | 222 | 264 | <11.3 |
| | max | 34.1 | 65 | 23.8 | 22.0 | 28.7 | 8.35 | 152 | 1,310 | 1,337 | 252 |
| Imperial Valley Off-refuge K18A-E (N=5) | ave | 5.47 | 7.83 | 6.50 | 5.30 | 4.64 | 1.57 | 62.5 | 120 | 130 | 69.1 |
| | min | 4.21 | 0.320 | 4.53 | 2.19 | 1.77 | 0.63 | 38.9 | 47.8 | 62.3 | 35.30 |
| | max | 6.81 | 17 | 8.54 | 7.13 | 7.80 | 2.50 | 86.3 | 310 | 320 | 157 |
| | | | | | | | | | | | |
| Imperial Valley On-refuge; SBSS NWR SB420A - E (N=5) | ave | 5.93 | 6.81 | 6.84 | 4.44 | 6.05 | 1.24 | 64.3 | 306 | 324 | <6.11 |
| | min | 3.92 | 0.371 | 5.18 | 2.56 | 4.52 | 0.417 | 34.1 | 142 | 151 | <4.17 |
| | max | 7.58 | 22.9 | 8.36 | 6.19 | 8.80 | 2.94 | 95.1 | 547 | 575 | 9.13 |
| | | | | | | | | | | | |
| Central Valley On-refuge; Kern NWR K15A - E (N=5) | ave | 22.1 | nd | 11.6 | 3.40 | 14.2 | 3.44 | 178 | 10.2 | 38.5 | <11.6 |
| | min | 10.5 | nd | 2.10 | 0.62 | 7.77 | 0.62 | 45.9 | 3.10 | 18.6 | <6.08 |
| | max | 39.8 | nd | 20.7 | 8.95 | 22.7 | 6.55 | 604 | 17.2 | 63.0 | <22.4 |

¹See Appendix 5 for complete list of organic constituent concentrations by sample.




Figure 17. DDT isomer p,p'-DDE concentrations in soils and invertebrates collected in California.




Figure 18. Total DDT concentrations in soils and invertebrates collected in California.




Figure 19. Trifluralin concentrations in soils and invertebrates collected in California.

Endosulfan II and endrin were both detected in one invertebrate sample from Kern NWR at levels considerably higher than the other Kern NWR invertebrates samples (all below non-detect) and those from Imperial Valley, including SBSS NWR (Appendix 5). Endosulfan does not accumulate in warm blooded animals indicating a low risk to the mountain plover (Wiemeyer 1996), however endrin is considered one of the most acutely toxic OCs (Blus 1995). Endrin was detected at a low concentration of 0.203 ng/g dw in the corresponding soil sample but the invertebrate sample contained 1,020 ng/g dw indicating significant bioaccumulation and a potential risk to mountain plovers consuming these invertebrates.

The total PCB concentrations in invertebrates ranged from 6 ng/g dw to 768 ng/g dw (Appendix 6). The median for all 25 samples was 138 ng/g dw and all but two samples had concentrations greater than 155 ng/g dw. These concentrations fell below dietary levels associated with most

adverse reproductive effects in gallinaceous birds, which tend to be among the most sensitive of the avian receptors to PCB toxicity (USEPA 1995). These concentrations were also below 2,000 ng/g identified by USEPA (1995) as a NOAEL for multiple reproductive endpoints in gallinaceous species. Additionally, USEPA (1995) noted that 2,000 ng/g dw is a LOAEL for the most sensitive effect (reduced chick growth) among chickens exposed to most toxic PCB mixtures (Aroclors 1248 and 1254). Using an uncertainty factor of 10, the corresponding estimated NOAEL would be 200 ng/g dw, which is greater than concentrations observed in all but two samples of invertebrates.

Total PCB concentrations in the invertebrate samples were routinely higher than total PCB concentrations in co-located soil samples. However, the variation among samples was high so that site-specific differences or correlations with total PCB concentrations in soils, if present, could not be detected. Some of the variability observed with PCBs in invertebrate samples may be due to the nature of the samples, as each had different proportions of spiders, beetles, ants, and larvae. As a result, samples with more spiders may have had a higher concentration of PCBs, as some bioconcentration of PCBs within the invertebrate community had likely already occurred.




Figure 20. Concentration of total PCBs in soil and invertebrate samples collected in California.

Breeding Grounds

Soil-Inorganic Elements

Soil samples were collected from three refuges in Montana, three locations in Wyoming, and one location in Colorado. Soil trace element concentrations for all sites were lower than the Eco-SSLs for arsenic, beryllium, chromium, cobalt, and manganese. In addition, the average boron, barium, beryllium, chromium, mercury, molybdenum, and strontium in soil samples from breeding areas were either below or near the western U.S. average for background soils (Shacklette and Boerngen 1984). One composite soil sample from CMR NWR was the only site that had elevated concentrations of several trace elements in comparison to other sites sampled in the breeding area. Summary statistics of these inorganic element concentrations are contained in Table 6; full analytical results are listed in Appendix 7.

In soil samples from Montana sites, arsenic concentrations (range = 8.2 – 12.4 ug/g dw) were generally higher than average (5.5 ug/g dw) but within the range (<0.10 – 97 ug/g dw) for western U.S. background soils (Shacklette and Boerngen 1984) and did not exceed the Eco-SSL of 43 ug/g for the protection of birds or soil invertebrates (Table 6). Arsenic concentrations from Wyoming soil samples (range = 2.9 – 5.3 ug/g dw) centered around the western U.S. average (Shacklette and Boerngen 1984).

The average background concentration for cobalt in western U.S. soils is 7.1 (range = <3 – 50 ug/g dw). Six of the nine composite soil samples taken, primarily in Montana, were above this average (range = 7.4 – 23.2 ug/g dw) but all fell below the Eco-SSL of 120 ug/g for the protection of birds. Cobalt makes up part of vitamin B₁₂ and is an essential micronutrient for animals including birds. Cobalt deficiency is typically more of a problem and studies on the effects of excess concentrations on birds are few (Nelson et al. 1966). However, comparison with the Eco-SSL suggests that risks posed to mountain plovers by cobalt concentrations in soils are likely to be low.

While Shacklette and Boerngen (1984) provide no values for background concentrations of cadmium in U.S. soils, the cadmium concentrations measured in soils from plover breeding sites (range = 0.13 – 1.1 ug/g dw) fell within the range of mean concentrations reported by Baudo et al. (1990) for soils uninfluenced by obvious sources (0.01 – 2 ug/g dw). Only one soil sample exceeded the Eco-SSL for avian wildlife (0.77 mg/kg dw), which is based on the woodcock, a ground insectivore (USEPA 2005d). The composite soil sample collected from North and South Buckskin Ridge on CMR NWR contained 1.09 mg/kg dw. Laboratory studies suggest that dietary intake of <1 mg/kg cadmium in food would not lead to any toxicological effect in birds (Furness 1996).

Copper concentrations in soils from breeding sites ranged from 6.7 ug/g dw to 30 ug/g dw, which were higher than the average of 2.07 ug/g, but within the range (2 – 300 ug/g dw) reported by Shacklette and Boerngen (1984) for western U.S. soils. These concentrations, with the exception of a soil sample from ACM study area in Wyoming (30 ug/g dw), also fell below the Eco-SSL of 28 ug/g for protection of avian species and all were below the Eco-SSL of 80 ug/g for protection

Table 6. Average and range of concentrations of select inorganics¹ (ug/g dry weight) in soil samples collected from mountain plover breeding sites² in Montana and Wyoming, 2006, for comparison with average and range of inorganics in soils from the western United States as reported by Shacklett and Boerngen (1984).

| | | As | B | Ba | Be | Cd | Co | Cr | Cu | Hg | Mn | Mo | Ni | P |
|----------------------------------|---------|-----------|------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|----------|
| WY - Soil (n=3) | average | 4.29 | 4.96 | 112 | 0.60 | 0.32 | 6.61 | 15.9 | 23.4 | 0.04 | 179 | BDL | 19.81 | 358 |
| | minimum | 2.9 | BDL ³ | 88.4 | 0.393 | 0.27 | 5.0 | 8.58 | 18.1 | 0.014 | 97.8 | BDL | 9.12 | 272 |
| | maximum | 5.32 | 4.96 | 148 | 0.972 | 0.394 | 8.15 | 22.1 | 30 | 0.07 | 288 | BDL | 33.7 | 402 |
| | | | | | | | | | | | | | | |
| MT - Soil (n=5) | average | 10.0 | 12.42 | 229 | 0.79 | 0.48 | 12.23 | 23.1 | 18.7 | 0.02 | 826 | 0.58 | 23.94 | 704 |
| | minimum | 8.18 | 9.41 | 115 | 0.626 | 0.274 | 7.4 | 21.1 | 16.5 | 0.020 | 426 | BDL | 17.8 | 591 |
| | maximum | 12.4 | 14.2 | 363 | 0.988 | 1.09 | 23.2 | 25.6 | 20.6 | 0.028 | 2290 | 0.99 | 40.8 | 863 |
| | | | | | | | | | | | | | | |
| western U.S. background soils | average | 5.5 | 23 | 580 | 0.68 | No value | 7.1 | 41 | 2.07 | 0.46 | 380 | 0.85 | 15 | 320 |
| | minimum | <0.10 | <20 | 70 | <1 | | <3 | 3 | 2 | <0.01 | 30 | <3 | <5 | 40 |
| | maximum | 97 | 300 | 5000 | 15 | | 50 | 2000 | 300 | 4.6 | 5000 | 7 | 700 | 4500 |

| | | Pb | Se | Sr | V | Zn |
|----------------------------------|---------|-----------|-----------|-----------|----------|-----------|
| WY - Soil (n=3) | average | 11.51 | 0.56 | 45.97 | 31.13 | 54.17 |
| | minimum | 8.14 | 0.204 | 17.9 | 22.7 | 38.1 |
| | maximum | 17.1 | 1.22 | 95.2 | 37.3 | 80.6 |
| | | | | | | |
| MT - Soil (n=5) | average | 13.68 | 0.46 | 51.20 | 55.26 | 77.16 |
| | minimum | 12.4 | 0.262 | 24 | 41.1 | 65.1 |
| | maximum | 15.1 | 0.589 | 109 | 65.6 | 84.9 |
| | | | | | | |
| western U.S. background soils | average | 17 | 0.23 | 200 | 70 | 55 |
| | minimum | <10 | <0.1 | 10 | 7 | 10 |
| | maximum | 700 | 4.3 | 3000 | 500 | 2100 |

¹See Appendix 7 for complete list of inorganic constituent concentrations by sample.

²Only one soil sample collected from the Colorado mountain plover breeding site so data are not presented in this table.

³BDL = Below Detection Limit

of soil invertebrates. Copper is an essential nutrient and deficiency or excess can be harmful to organisms. Reports of toxic effects from excess dietary exposure in birds in field conditions are rare (Eisler 1997; USDOJ 1998).

The average lead concentrations in soil samples from breeding areas were below the average background concentration of 17 ug/g for western U.S. soils (Shacklette and Boerngen 1984), but most individual samples exceeded the Eco-SSL of 11 ug/g (Table 1). The likelihood of poisoning from the ingestion of lead in forms other than spent ammunition or where soil is extremely contaminated from mining operations is slim (Eisler 1988b).

Manganese in soil samples from Montana were above the average of 380 ug/g for western U.S. soils with one on-refuge soil sample having a high value of 2,290 ug/g dw (Table 6; Appendix 7). Even so, this high concentration is below the Eco-SSL 4,300 ug/g and is of little concern (Table 1).

Nickel concentrations (range = 16.6 – 40.8 ug/g dw) were slightly above the average (15 ug/g) for western U.S. soils in seven of nine soil samples from breeding grounds but below the Eco-SSLs for birds (210 ug/g) and soil invertebrates (280 ug/g). Nickel does not biomagnify and even in areas near smelters where nickel is significantly elevated in soils, it is unlikely that higher trophic level species are affected (Phipps et al. 2002).

Phosphorus was above the average (320 ug/g) for western U.S. soils for all soil samples in the breeding areas except one sample from TBNG in Wyoming (Table 6; Appendix 7). The availability of phosphorus depends on a soil's pH, with a lower pH increasing aluminum's solubility. The aluminum then readily binds to phosphorus making it unavailable to birds. Phosphorus is an essential element for eggshell development and bone growth (Miles et al. 1993). Western U.S. soils are primarily alkaline thereby reducing the availability of phosphorus to birds (Miles et al. 1993).

Selenium concentrations in breeding area soil samples (range = 0.2 – 1.22 ug/g dw; average = 0.459) were mostly above the average of 0.23 ug/g dw for background western U.S. soils

(Shacklette and Boerngen 1984), but appear to be typical for native rangeland soils sampled in parts of Wyoming and Montana (0.061 – 3.3 ug/g; Tidball 1976). Because selenium is very mobile in alkaline soils, geological formations in the western U.S. readily contribute selenium to the soils (USDOI 1998). Although, selenium is an essential mineral for organisms, there is a small margin between what is required nutritionally and what is considered toxic. These background concentrations are slightly lower than the avian insectivore Eco-SSL of 1.2 mg/kg dw in soil. Only one soil sample from ACM exceeded this level with a concentration of 1.22 mg/kg dw. Because the Eco-SSL is based on a no-effect level, dietary ingestion of selenium from soils is not likely to cause any impacts to mountain plovers on their breeding grounds based on the samples collected as a part of this investigation, or to soil invertebrates, which plovers rely on for food (Eco-SSL 4.1 ug/g dw; Table 1).

Vanadium concentrations in all soil samples were below the average background of 70 ug/g dw for western U.S. soils (Shacklette and Boerngen 1984), but above the Eco-SSL of 7.8 ug/g dw for the protection of birds. Whether the exceedances of Eco-SSLs translate into significant risk for mountain plovers is uncertain because the avian Eco-SSL is based on results from studies on chickens, and data on vanadium toxicity to wild species are lacking. Results of one study on waterfowl demonstrated that sensitivity to vanadium toxicity depends on the form of the vanadium and the species of the bird (Rattner et al. 2006).

Concentrations of zinc in soil samples (range = 65.1 – 84.9 ug/g dw) from Montana and one soil sample (80.6 ug/g dw) from Wyoming were above the average (55 ug /g dw) for background in western U.S. soils (Shacklette and Boerngen 1984), but were within the range of means reported by Kabata-Pendias and Pendias (1992) for uncontaminated sites (17 – 125 ug/g dw and rarely >200 ug/g dw). All Montana soils and the ACM in Wyoming sample exceeded the Eco-SSL for an avian insectivore (46 mg/kg dw). However, the extent to which the Eco-SSL was exceeded was small. The maximum zinc concentration of 85 ug/g dw was 1.8-fold greater than the Eco-SSL, which accounts for exposure through ingestion of soil and soil invertebrates that have accumulated zinc in their tissues. Zinc is an essential element and while it may be bioaccumulated by exposed invertebrates, the zinc concentrations measured in soils from breeding sites were not elevated enough to be considered harmful by either direct (incidental

ingestion) or indirect (dietary) exposure (Eisler 1993). Zinc is a required nutrient and bird diets should contain 25 – 38 mg/kg dw in feed to prevent the effects of zinc deficiency and 93 – 120 mg/kg dw for optimal growth (Eisler 1993).

Soils-Organic Compounds

OCs, OPs, and total PCBs (ng/g dw) were not detected in soil samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming in 2006. The specific compounds (and detection limits) for which all samples were analyzed are listed in Appendix 8.

Terrestrial Invertebrates-Inorganic Elements

Concentrations of inorganics in terrestrial invertebrates from breeding sites were typically lower than those found in terrestrial invertebrate samples collected from wintering sites in California (Table 7). Among breeding sites, one composite invertebrate sample from CMR NWR had higher concentrations of inorganic compounds than samples from the other sites. This is not surprising given that most of the inorganic elements in soil sampled from this same site were also higher than soil sampled from other sites (Appendices 7 and 9).

Aluminum concentrations in all but one of the terrestrial invertebrate samples (range = approximately 52 – 387 ug/g dw) were much lower than those collected from wintering grounds in California. The one exception is the composite sample from CMR NWR with an aluminum concentration of 2,180 ug/g dw. This concentration is far below the aluminum concentration of 5,000 ug/g considered to be the adverse dietary effect level in waterfowl (Sparling and Lowe 1996), but is above the 1,000 ug/g NOAEL developed for ring doves (*Streptopelia risoria*) (Sample et al. 1996).

Arsenic concentrations measured in invertebrates ranged from 0.07 ug/g dw to 2.15 ug/g dw. Dietary effects level of arsenic to birds depends on the arsenical compound and the bird species (USDOI 1998). For example, female mallard ducklings exhibited reduced growth when fed 30 ug/g dw as As⁺⁵ over a 10 week period (Eisler 1988a); whereas 99.8 ug/g dw of dietary sodium arsenate was lethal to 50 percent of the brown-headed cowbirds tested over an 11-day period

(Camardese et al. 1990). Concentrations measured in invertebrates consumed by plovers were all well below concentrations associated with adverse effects.

Barium concentrations in all but one of the terrestrial invertebrate samples from breeding grounds were below the estimated dietary NOAEL-based benchmark for avian wildlife of 20.8 ug/g (Sample et al. 1996). Beryllium was below detection limits for all samples except one in Montana and that concentration was very low (0.132 ug/g dw). Additionally, concentrations of boron, cadmium, chromium, and molybdenum observed in terrestrial invertebrates were low (Eisler 1985a, 1986, and 1989; Sample et al. 1996; USDOJ 1998).

Cobalt concentrations were low in invertebrates collected (range = <0.45 – 0.84 ug/g dw) with the exception of one sample in Montana that had the maximum concentration 2.12 ug/g dw. It is unclear if this cobalt concentration could affect mountain plover because effects of dietary cobalt on birds are not well studied.

Copper concentrations in invertebrate samples ranged from approximately 14 ug/g dw to 48 ug/g dw. Unfortunately, little information exists on the dietary concentration of copper that may cause toxicity to wild birds. Information on poultry indicate that a diet with 300 ug/g of copper is the maximum tolerable limit, but concentrations found in terrestrial invertebrates from all mountain plover breeding sites were well below this level (Eisler 1997; USDOJ 1998).

Mercury was detected in invertebrate samples but concentrations were low (range = <0.0047 – 0.017 ug/g dw) compared with concentrations reported in reviews such as Eisler (1987). Nickel concentrations ranged from <0.49 ug/g dw to 1.77 ug/g dw for all terrestrial invertebrate samples collected at breeding sites except the one sample from CMR NWR site (94.94 ug/g dw). There is little information on dietary nickel concentrations required for adverse effects in wild birds. However, the high concentration measured in the sample from CMR NWR was below the nickel concentration found to affect domestic chickens (<700 ug/g dw; Weber and Reid 1968) and mallard ducklings (<800 ug/g dw; Cain and Pafford 1981). Like nickel, the maximum concentration of lead was in the sample from CMR NWR (3.15 ug/g dw versus 0.135 – 0.866 ug/g dw), but this concentration is well below dietary effects levels for birds reported by Eisler

(1988b) at 50 ug/g inorganic lead and Hoffman et al. (1985) at >25 ug/g body weight (BW) or approximately 139 ug/g mountain plover diet. The selenium concentration (2.56 ug/g dw) in the invertebrate sample at the CMR NWR site was also higher than from invertebrates collected at the other sites (range 0.613 – 1.76 ug/g dw), but were below dietary effects levels for birds (>3 ug/g dw; USDOJ 1998).

Strontium concentrations measured in invertebrates from plover breeding grounds were between 0.438 ug/g dw and 2.56 ug/g dw. Strontium is considered to have low toxicity to animals (Wennig and Kirsch 1950), although it is unknown if strontium levels measured in invertebrates at this site pose a threat to mountain plover. Dietary concentrations between 8 ug/g dw and 117 ug/g dw have been related to elevated levels in egg contents and shells of passerine birds in Arizona (Mora 2003) but samples from the mountain plovers' breeding sites were at the lower end of this range.

Titanium concentrations ranged from 5.8 ug/g dw to 27.3 ug/g dw. The effect of titanium in the diet of birds is unknown. Similarly, information on the toxicity of dietary vanadium to terrestrial birds is limited. Results of studies by Rattner et al. (2006) on mallards and Canada geese indicate that dietary effect levels vary, depending on the form of the vanadium, but also that dietary concentrations of 250 ug/g caused little known toxicity. Vanadium concentrations measured in invertebrates (maximum = 7.7 ug/g dw) are well below published adverse effect levels for mallards and Canada geese. Concentrations of zinc ranged from 85.8 ug/g dw to 169 ug/g dw, which are below zinc dietary levels of 178 ug/g that cause immunosuppression in young domestic chickens following subchronic exposure of three weeks (Eisler 1993).

Table 7. Average and range of concentrations of select inorganics¹ (ug/g dry weight) in invertebrate samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006.

| | | Al | As | Ba | Co | Cu | Hg | Ni | Pb | Se | Sr | Ti | V | Zn |
|-----------------------|----------------------|-----------|------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|----------|-----------|
| WY - Insects (n=5) | average | 108.8 | 0.18 | 8.57 | 0.48 | 23.96 | 0.006 | 1.32 | 0.25 | 0.66 | 9.4 | 7.2 | 0.399 | 125.4 |
| | minimum | 52.8 | 0.07 | 5.85 | 0.46 | 18.3 | BDL | 0.83 | 0.14 | 0.44 | 5.8 | 5.9 | BDL | 111 |
| | maximum | 183 | 0.41 | 11.8 | 0.51 | 28 | 0.007 | 1.77 | 0.34 | 0.75 | 17.8 | 9.8 | 0.682 | 137 |
| | | | | | | | | | | | | | | |
| MT - Insects (n=5) | average | 631.5 | 0.88 | 18.1 | 0.81 | 30.44 | 0.011 | 1.9 | 0.99 | 1.39 | 20.9 | 15.5 | 2.31 | 153.6 |
| | minimum | 90.3 | BDL ² | 9.68 | BDL | 17.1 | 0.007 | 0.83 | 0.21 | 0.68 | 10.1 | 2.9 | BDL | 137 |
| | maximum | 2180 | 2.15 | 42.6 | 2.12 | 47.5 | 0.017 | 4.94 | 3.15 | 2.56 | 31 | 43 | 7.77 | 169 |
| | | | | | | | | | | | | | | |
| CO - Insects (n=2) | average ³ | | | | | | | | | | | | | |
| | minimum | 101 | 0.09 | 6.93 | 0.47 | 14.1 | BDL | BDL | 0.23 | 0.61 | 6.2 | 6.7 | BDL | 85.5 |
| | maximum | 134 | 0.11 | 9.18 | 0.49 | 23.6 | BDL | 0.62 | 0.33 | 0.68 | 7.5 | 9.4 | BDL | 104 |
| | | | | | | | | | | | | | | |

¹See Appendix 9 for complete list of inorganic constituent concentrations by sample.

²BDL = Below Detection Limit

³Average not calculable as n=2

Terrestrial Invertebrates-Organic Compounds

None of the target organic analytes (OCs, OPs, and total PCBs) was detected in terrestrial invertebrate samples collected from mountain plover breeding sites in 2006 from Colorado, Montana, or Wyoming. Specific compounds analyzed in terrestrial invertebrate samples are listed in Appendix 10.

Mountain Plover Eggs-Inorganic Elements-2006

Generally, concentrations of most inorganic constituents were low in mountain plover eggs. Boron, beryllium, cobalt, chromium, molybdenum, and vanadium were not detected in any of the mountain plover eggs collected in 2006. Silver was not analyzed in eggs collected in Montana and was not detected in any eggs collected from Colorado or Wyoming. Arsenic was detected in more than half the eggs collected (Table 8; Appendix 11), but at concentrations below the tentative screening level of 1.3 ug/g dw for no effects in birds (USDOJ 1998).

Barium concentrations measured in mountain plover eggs (range = 3.15 ug/g – 19.6 ug/g dw; Table 9) were above levels found in the yellow-breasted chat (*Icteria virens*) (2.5+2 ug/g dw) and the endangered southwestern willow flycatcher (*Empidonax traillii extimus*) (0.5 + 0.1 ug/g dw) from Arizona (Mora 2003). In eggs of the California clapper rail (*Rallus longirostris obsoletus*) collected in the south and north sections of San Francisco Bay averaged 0.34 ug/g dw (max 0.59 ug/g dw) and 0.45 ug/g dw (max 4.13 ug/g dw), respectively (Schwarzbach et al. 2006). Three of the clapper rail embryos exhibited deformed feet where barium concentrations ranged from 2.2 ug/g dw to 4.1 ug/g dw. Such high barium concentrations may possibly be associated with malpositioning in late stage embryos (Schwarzbach et al. 2006). Of the 20 mountain plover eggs collected, only one embryo (age 25 days) from Wyoming was deformed. This embryo exhibited an abnormally small head and an upper bill that twisted out and up. The barium concentration in that egg was 13.1 ug/g dw, which is greater than the barium concentration found in deformed California clapper rail embryos. However, whether the deformed mountain plover is the result of the elevated barium concentration is unknown as three eggs from PNG had higher barium concentrations than the egg with the deformed chick and these did not appear to have deformities when examined.

Table 8. Average and range of concentrations of inorganics^{1,2} (ug/g dry weight) in egg samples³ collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006.

| Location | Parameter | Moisture % | Al ug/g dw | As ug/g dw | Ba ug/g dw | Ca ug/g dw | Cu ug/g dw | Fe ug/g dw | Hg ug/g dw |
|---|-----------|------------|------------|------------|------------|------------|------------|------------|------------|
| Pawnee National Grasslands, CO (n=5) | average | 72.66 | 1.15 | 0.13 | 14.60 | 5538.00 | 3.05 | 111.46 | 0.037 |
| | minimum | 71 | 0.68 | 0.05 | 10.5 | 2680 | 2.23 | 96.3 | 0.026 |
| | maximum | 74.7 | 1.47 | 0.22 | 19.6 | 9900 | 4.29 | 136 | 0.051 |
| BLM Lands, Phillips County, MT (n=7) | average | 72.61 | 0.71 | 0.18 | 8.12 | 7394.29 | 3.18 | 106.11 | 0.035 |
| | minimum | 70.3 | 0.54 | 0.18 | 5.75 | 2680 | 2.66 | 84.9 | 0.025 |
| | maximum | 75.9 | 0.89 | 0.19 | 9.72 | 19200 | 4.13 | 126 | 0.049 |
| Foote Creek Rim, WY (n=5) | average | 72.6 | 0.82 | 0.11 | 4.86 | 5898 | 2.824 | 88.3 | 0.052 |
| | minimum | 71.6 | 0.62 | 0.05 | 3.15 | 2750 | 2.65 | 77.4 | 0.039 |
| | maximum | 74.3 | 1.19 | 0.29 | 5.85 | 13500 | 3 | 97.7 | 0.075 |

Table 8. cont.

| Location | Parameter | K ug/g dw | Mg ug/g dw | Mn ug/g dw | Na ug/g dw | Ni ⁴ ug/g dw | P ug/g dw | Pb ug/g dw | S ug/g dw |
|---|-----------|-----------|------------|------------|------------|-------------------------|-----------|------------------|-----------|
| Pawnee National Grasslands, CO (n=5) | average | 5300 | 481 | 1.55 | 5254 | 22.06 | 8958 | 0.06 | 6294 |
| | minimum | 4490 | 371 | 0.96 | 4490 | 1.15 | 7920 | BDL ⁵ | 5420 |
| | maximum | 6310 | 590 | 2.14 | 6190 | 55.7 | 9810 | 0.19 | 6840 |
| BLM Lands, Phillips County, MT (n=7) | average | 5582.86 | 559.71 | 1.27 | 5928.57 | NC ⁶ | 8588.57 | 0.06 | 7596 |
| | minimum | 4700 | 403 | 0.67 | 5200 | BDL | 7380 | BDL | 6670 |
| | maximum | 6350 | 922 | 2.05 | 7310 | BDL | 10900 | 0.13 | 9070 |
| Foote Creek Rim, WY (n=5) | average | 5328 | 507.8 | 1.29 | 5352 | NC | 8612 | 0.03 | 6744 |
| | minimum | 4620 | 401 | 0.907 | 4520 | BDL | 7810 | BDL | 5920 |
| | maximum | 5730.00 | 748.00 | 1.83 | 6570.00 | BDL | 10300 | 0.05 | 7650 |

Table 8. cont.

| Location | Parameter | Se ug/g dw | Si ug/g dw | Sr ug/g dw | Ti ug/g dw | Zn ug/g dw |
|---|-----------|---------------|---------------|---------------|---------------|---------------|
| Pawnee National Grasslands, CO (n=5) | average | 2.89 | 11.59 | 7.87 | 0.54 | 51.88 |
| | minimum | 2.19 | 9.77 | 3.81 | BDL | 39.7 |
| | maximum | 3.84 | 12.9 | 13.5 | 1.06 | 60 |
| BLM Lands, Phillips County, MT (n=7) | average | 4.87 | NA | 8.49 | NC | 45.37 |
| | minimum | 3.64 | NA | 2.74 | BDL | 36.5 |
| | maximum | 7.32 | NA | 16.60 | BDL | 52.9 |
| Foote Creek Rim, WY (n=5) | average | 2.64 | 8.96 | 7.75 | 0.61 | 52.5 |
| | minimum | 2.16 | 8.03 | 3.63 | BDL | 43 |
| | maximum | 3.1 | 9.93 | 17.1 | 1.78 | 60.2 |

¹See Appendix 11 for complete list of inorganic constituent concentrations by sample.

²One sample at Pawnee National Grasslands, CO had a Cd concentration of 0.021 ug/g dw; the other four samples were below the detection limit. All BLM Lands, MT samples and all Foote Creek Rim, WY samples were below the detection limit for Cd.

³One egg collected from Charles M. Russell NWR, MT and two egg samples collected from the Antelope Coal Mine site, WY are not included in this table but are addressed in the text if the inorganic element concentration is elevated.

⁴One sample at Foote Creek Rim, WY had a Ni concentration of 0.67 ug/g dw; the other four samples were below the detection limit. All BLM Lands, MT samples were below the detection limit for Ni.

⁵BDL = Below Detection Limit.

⁶NC = Not calculable.

Table 9. Barium concentrations and developmental stage of mountain plover eggs collected from breeding sites in Colorado, Montana, and Wyoming in 2006.

| Location | Sample | Barium ug/g dw | Fertile | Age days | Abnormal |
|--------------------------------|-------------|-------------------|---------|-------------|----------|
| Pawnee National Grasslands, CO | MPWCPE01 | 10.8 | Yes | 8 | No |
| Pawnee National Grasslands, CO | MPWCPE02 | 13.7 | Yes | 4 | No |
| Pawnee National Grasslands, CO | MPWCPE03 | 18.4 | Yes | 24 | No |
| Pawnee National Grasslands, CO | MPWCPE04 | 19.6 | Yes | 12 | No |
| Pawnee National Grasslands, CO | MPWCPE05 | 10.5 | Yes | 17 | No |
| | | | | | |
| BLM Lands, Phillips County, MT | BLM025MPE | 7.83 | Yes | 10 | No |
| BLM Lands, Phillips County, MT | BLM081-2MPE | 8.83 | * | * | * |
| BLM Lands, Phillips County, MT | BLM081MPE | 9.37 | * | * | * |
| BLM Lands, Phillips County, MT | BLM189MPE | 5.75 | * | * | * |
| BLM Lands, Phillips County, MT | BLMB026MPE | 6.71 | Yes | 15 | No |
| BLM Lands, Phillips County, MT | BLMS014MPE | 8.65 | Yes | 15 | No |
| BLM Lands, Phillips County, MT | BLMS020MPE | 9.72 | * | * | * |
| Charles M. Russell NWR, MT | CMRNMCMPE06 | 10 | ** | Fresh | No |
| | | | | | |
| Near Antelope Coal Mine, WY | MPAMPE01 | 5.2 | Yes | 9 | No |
| Near Antelope Coal Mine, WY | MPAMPE02 | 13.1 | Yes | 25 | Yes |
| | | | | | |
| Foot Creek Rim, WY | MPFCPE01 | 4.86 | Yes | 21 | No |
| Foot Creek Rim, WY | MPFCPE02 | 4.97 | Yes | 21 | No |
| Foot Creek Rim, WY | MPFCPE03 | 3.15 | No | 0 | No |
| Foot Creek Rim, WY | MPFCPE04 | 5.85 | Yes | 25 | No |
| Foot Creek Rim, WY | MPFCPE05 | 5.49 | Yes | 10 | No |

* Egg collected from abandoned nest, and yolk in advanced state of decay

** Fertility not noted

Cadmium was not detected in most eggs except one egg from the ACM site (0.065 ug/g dw) and one egg from PNG (0.021 ug/g dw; Appendix 11). According to Ohlendorf (1993), non-toxic concentrations of cadmium in avian eggs are <0.5 ug/g. The toxic effects most commonly associated with cadmium exposure include impaired growth, anemia, and testicular damage (Eisler 1985a). The toxic effects of cadmium in avian species have also included altered avoidance response, reduced egg production, eggshell thinning, kidney damage, damage to the gut epithelium, altered energy metabolism, impaired bone marrow function, heart abnormalities, and adrenal abnormalities (Eisler 1985a; Furness 1996). These effects, which were observed in

laboratory tests, have been related to cadmium levels in the diet, the liver, or the kidney. Benchmarks relating adverse effects of cadmium concentrations in eggs are lacking (Ohlendorf 1993) and the likelihood of cadmium reaching these adverse levels in eggs is considered to be very low (Furness 1996).

Copper was detected in eggs, but all below the tentative screening level of 5.5 ug/g dw eggs (USDOI 1998). Similarly, mercury was detected in all egg samples, with a maximum of 0.075 ug/g dw (or 0.021 ug/g ww for that sample). The level of concern for mercury in eggs is between 0.2-1.0 ug/g ww (USDOI 1998) and a level of 0.5 ug/g ww in eggs (or about 2.5 ug/g dw assuming 80% egg moisture) does not appear to have any adverse effects on reproduction (Thompson 1996).

Lead was detected in nine of 20 eggs. Concentrations in all of the eggs were less than 0.5 ug/g dw, which is consistent with concentrations of <0.5 ug/g dw reported for eggs of yellow-breasted chat and willow flycatchers (Mora 2003), and what is considered background (<1.0 ug/g dw; Ohlendorf 1993; Gochfeld and Burger 1998).

Nickel was below the detection limit in most eggs from Colorado, Montana, and Wyoming. One egg collected at FCR, Wyoming, had a detectable nickel concentration of 0.67 ug/g dw. Interestingly, all eggs collected at PNG had detectable nickel concentrations (1.2, 4.1, 14.4, 35, and 55.7 ug/g dw). It is unknown what the source of the elevated nickel concentrations is because the corresponding soil sample and invertebrate diet sample analyzed from this site and from the California wintering sites did not show elevated nickel concentrations. Nickel is toxic and teratogenic in domestic chickens at concentrations in eggs as low as 0.02 mg/egg (approximately 2.0 ug/g dw assuming a 55 g/egg with 80% moisture; Eisler 1998). Studies where nickel concentrations of 700 ug/g as nickel mesotetraphenylporphine in 1 uL of crude oil (= 0.7 ug per egg or ~0.06 ug/g dw assuming 60 g/egg and 80% moisture) injected into mallard duck eggs did not affect embryo survival significantly compared to eggs where just 1 uL of crude oil was injected, although the embryonic weight was significantly lower, and the percentage of abnormalities in post hatch survivors was increased (Hoffman 1979). The potential for toxicity depends on the form that the nickel is in and the species of the exposed organism. Nickel

concentrations measured in mountain plover eggs from PNG (average = 22 ug/g dw) are greater than concentrations associated with adverse effects in other species, and as such, may be at levels of concern for the plover. No abnormalities were noted, however.

Bird eggs collected from study sites not impacted by selenium usually contain ≤ 3 ug/g dw with a maximum of 5 ug/g dw (USDOI 1998). The average selenium concentrations in eggs from PNG (2.89 ug/g dw), FCR (2.64 ug/g dw), and the one egg collected near the ACM site (3.03 ug/g dw) were at or below the mean background concentration. Selenium concentrations in one egg from CMR NWR (3.63 ug/g dw) and most of the eggs collected from BLM lands in Montana were above average background, but within the range of background samples (Table 8). Three eggs collected on the BLM lands were >5 ug/g dw but still below the embryotoxic threshold for selenium in bird eggs of 10 ug/g dw (Heinz 1996). The reasons as to why selenium concentrations in eggs from Montana were higher than in eggs from Colorado and Wyoming are not clear because selenium concentrations in co-located soil and invertebrate samples from the Montana site tended to be lower in selenium than concentrations in soils and invertebrates from Wyoming and from wintering sites in the Imperial Valley (Tables 2, 4, 6, and 7). Because ingested selenium is readily transferred from the female bird to the egg (Eisler 1985c), the apparent higher levels in eggs from the Montana sites may be the result of incidental soil ingestion (USDOI 1998). The suggested embryotoxic threshold for selenium in bird eggs is estimated to be 10 ug/g (Heinz 1996; USDOI 1998).

Concentrations of sulfur and silicon are reported for mountain plover egg contents, but the significance of the concentrations is not known. Research on domestic chickens indicates that a laying hen will deposit sulfate in the egg and that some of this will be converted into the amino acid cystine (Machlin et al. 1953). Silicon is considered an essential nutrient for animals in small quantities (Ohlendorf 1993). However, concentrations that may be associated with adverse effects have not been identified.

Strontium concentrations in plover eggs did not appear elevated when compared with levels in eggs of the yellow-breasted chats (mean = 23.9 ± 12.7 ug/g dw) and willow flycatchers (mean = 35.1 ± 58 ug/g dw) in Arizona (Mora 2003) or with normal embryos of California clapper rail

(mean = 66.1 ug/g dw; range 30.2 – 94.6 ug/g dw; Schwarzbach et al. 2006). Titanium concentrations in plover eggs were low or below the detection limit. The significance of titanium in avian egg contents is unknown.

Average zinc concentrations in mountain plover eggs from PNG (51.98 ug/g dw; range 39.7 – 60.0 ug/g dw) and FCR (52.5 ug/g dw; range 43.0 – 60.2 ug/g dw) were slightly above what is considered background for avian eggs in general (50 ug/g dw; USDOJ 1998). The one egg from ACM had the highest zinc concentration (64.1 ug/g dw) of any of the eggs collected from the breeding areas, and eggs from Montana (range 36.5 – 52.9 ug/g dw) were less than- to slightly higher than background. The significance of observed zinc concentrations on the development of mountain plover is not known, but zinc is an essential element whose mobilization and transfer from storage sites to the developing embryo is regulated during development (Richards 1997).

Because of the generally overall low concentrations of inorganic elements in egg samples from 2006, and to control costs, eggs collected in 2007 and 2008, and archived eggs collected in 2003 were analyzed for organic constituents only.

Mountain Plover Eggs-Organic Compounds-2006

Overall, very few organic compounds were detected in any of the mountain plover eggs sampled from Colorado, Montana, and Wyoming (Appendix 12). Hexachlorobenzene (HCB) was detected at low concentrations in mountain plover eggs with the highest concentration (32.5 ng/g dw) detected in an egg from PNG (Table 10; Appendix 12). HCB was used as a fungicide to treat crop seeds but banned in 1965 in the U.S because of its carcinogenic properties. In 1995, it was banned globally at the Stockholm Convention because, as a persistent organic pollutant, it is observed to have long-range transport, significant human health and environmental effects, biomagnifies up the food chain, and degrades very slowly (half-life of 3 to 6 years in soil; ATSDR 2002). HCB is lipophilic and is primarily acquired by ingesting contaminated food items. In birds, females pass HCB directly into the egg (primarily the yolk) as shown in studies with Japanese and bobwhite quail (Ingebrigsten et al. 1981; Breslin et al. 1983). Little information exists on the effects of these concentrations on embryo survival and health but a

levels of 80 ug/g (80,000 ng/g) or higher was necessary before reproductive failure occurred in Japanese quail (Bechard 1981). Therefore, the concentrations of HCB found in the mountain plover eggs are unlikely to affect the embryos.

Table 10. Average and range of concentrations of most frequently detected organochlorine compounds¹ (ng/g dry weight) in egg samples collected from mountain plover breeding sites in Colorado, Montana² and Wyoming, 2006.

| Location | Parameter | HCB ng/g dw | oxychlordane ng/g dw | p,p'-DDE ng/g dw |
|---|-----------|----------------|-------------------------|---------------------|
| Pawnee National Grasslands, CO (n=5) | average | 15.4 | 18.4 | 13,062 |
| | minimum | 7.52 | BDL | 46.8 |
| | maximum | 32.5 | 48.8 | 43,200 |
| BLM Lands, Phillips County, MT (n=7) | average | 8.24 | 6.92 | 3,990 |
| | minimum | BDL | BDL | 134 |
| | maximum | 16.7 | 25.1 | 18,600 |
| Foote Creek Rim, WY (n=5) | average | 5.6 | NC | 1,328 |
| | minimum | BDL | BDL | 45 |
| | maximum | 8.4 | BDL | 4,090 |

¹See Appendix 12 for complete list of organic constituent concentrations by sample including those constituents not detected in any mountain plover egg sample.

²One egg collected from Charles M. Russell NWR, MT and two egg samples collected from the Antelope Coal Mine site, WY are not included in this table but are addressed in the text if the organic element concentration is elevated.

³BDL = Below Detection Limit.

⁴NC = Not calculable.

Oxychlordane was detected primarily in eggs collected from Colorado (Table 10; Appendix 12) with the highest concentration (48.4 ng/g dw) in the same egg that exhibited the highest concentration of HCB. Oxychlordane is a metabolite of chlordane, which was used primarily for the control of various crop pests and termites. All uses of chlordane were banned in the U.S. in 1988. Oxychlordane is more lipophilic, toxic, and persistent than chlordane (Eisler 1990a). Unfortunately, there is little information on the levels of oxychlordane in eggs that might be associated with adverse effects, and consequently, the implications of concentrations observed in plover eggs are not known.

DDE was detected in all eggs collected from all breeding sites (Table 10; Appendix 12). DDE is a major metabolite of the insecticide DDT, which was banned in 1972 in the U.S. DDE is still

commonly found in the environment because of its persistence as well as atmospheric transport and deposition from other countries which continue to use DDT (DOI 1998). Although no DDT or DDT metabolites were found in any soil or invertebrate samples collected on the breeding grounds in Montana, Wyoming, or Colorado, it was still detected in both soil and invertebrate samples collected in plover wintering grounds. Since little to no exposure to DDE occurs on mountain plover breeding grounds sampled as part of this investigation, it appears that DDE deposited in eggs of mountain plovers was from exposures on their wintering grounds.

DDE, specifically p,p'-DDE, is one of the metabolites responsible for deleterious effects in birds, with the most sensitive and most documented being eggshell thinning and subsequent breakage (Blus 1996). Much of the research on DDE's deleterious effect in terrestrial birds has focused on tissue residues (primarily brain and liver; Blus 1996), or concentrations of DDE found in the diet of these birds (USDOI 1998) rather than concentrations in eggs (with the exception of raptors), making it difficult to determine exactly what effects the concentrations of DDE observed in mountain plover eggs collected for this study may have. Even so, the concentrations of DDE in some of the individual plover eggs collected from this study were higher than 3 ug/g ww (3,000 ng/g ww) shown to cause almost complete colony failure due to egg breakage in the brown pelican (*Pelecanus occidentalis*), considered one of the most sensitive avian species (Blus 1995). For bald eagles (*Haliaeetus leucocephalus*) concentrations ≤ 3 ug/g ww in eggs appears to have no effects but almost complete nest failure occurred when eggs contained ≥ 16 ug/g ww (16,000 ng/g; Wiemeyer et al. 1984). DDE concentrations measured in mourning doves eggs from the Colorado River delta in Mexico, ranged from 4.3 ng/g ww to 58.6 ng/g ww, which were below the calculated NOAEL of 2.2 ug/g ww (2,200 ng/g ww) for terrestrial birds (Clark et al. 1995; Garcia-Hernandez et al. 2006). The DDE concentrations in mountain plover eggs collected in 2006 for this study ranged from 5 ng/g ww to 5,400 ng/g ww for eggs from Montana, from 10 ng/g ww to 930 ng/g ww for eggs from Wyoming, and from 11.0 ng/g ww to 11,500 ng/g ww for eggs from Colorado (Appendix 12). Furthermore, two eggs from Montana had DDE concentrations (3,000 ng/g ww and 5,400 ng/g ww) that were at least as high as those found to adversely affect productivity in pelicans. Two eggs from Colorado were near this concentration (2,700 ng/g ww and 2,900 ng/g ww) with an additional egg having a concentration (11,500 ng/g ww) approaching that known to affect bald eagle reproduction (Blus 1995).

Information on eggshell thickness of mountain plover eggs is lacking but the concentration of DDE associated with significant eggshell thinning (e.g. >20% thinner compared with pre-DDT eggshell thicknesses) ranged from 0.1 ug/g ww for the brown pelican (Blus 1984) to 2 ug/g ww for the peregrine falcon (Blus 1996). Concentrations of DDE in individual mountain plover eggs collected in Montana equaled or exceeded the threshold for significant thinning in brown pelicans in four of the seven eggs (range 0.1 – 5.4 ug/g ww) and DDE concentrations (3 ug/g ww and 5.4 ug/g ww) in two of these eggs exceeded the threshold for the peregrine falcon (Appendix 12). In Wyoming, DDE concentrations (0.9 ug/g ww and 0.93 ug/g/ww) in two of the seven eggs exceeded the threshold for significant thinning for brown pelicans but none exceeded the threshold for the peregrine falcon. Three of the five eggs collected from Colorado had DDE concentrations (range = 2.7 – 11.5 ug/g ww) that exceeded the threshold for significant thinning in sensitive species with one of these eggs (11.5 ug/g ww) exceeding the threshold for peregrine falcon.

Shell thickness of mountain plover eggs has not been reported prior to this report (Knopf and Wunder 2006; Appendices 13 – 15). Although our sample size is small and we did not attempt to locate archived mountain plover eggshells, correlations between DDE concentrations and eggshell thickness were still investigated. According to Blus (1984), egg residues are more accurate for indicating nest success rather than eggshell thickness and because the amount of eggshell thinning resulting from each incremental increase in DDE concentration is greater at lower residue levels (USDOI 1998). Additionally, researchers have shown that eggshell thinning of 5 – 7% is statistically significant but that it probably is not biologically significant. Specifically, field studies have shown that an average thinning of 10% is seldom associated with egg breakage or population decline (USDOI 1998). As an example, 1 ug DDE/g may result in 5-10% thinning, whereas 59 ug/g can result in 44% thinning in brown pelicans (USDOI 1998).

Data on the eggs collected in 2006 and 2007 were used to determine if a correlation existed between eggshell thickness using the Ratcliffe Index (Ratcliffe 1970) and p'p'-DDE concentrations. Statistical analysis revealed a weak negative association between DDE residue and egg index thickness (Spearman $r = -0.27$), however the two variables were not statistically correlated ($p = 0.185$; Figure 21).




Figure 21. Correlation between p'p' DDE concentrations in eggs and eggshell thickness

Mountain Plover Eggs-Organic Compounds-2007

Eggs collected from sites in Wyoming and Colorado in 2007 did not have detectable levels of trifluralin or its metabolites. Trifluralin is one of the more heavily used pesticides on forage grass fields in Imperial County, California, and was present in invertebrate samples at concentrations that were many times higher than concentrations in soil. The absence of trifluralin in plover eggs could be for a variety of factors: plovers are not ingesting trifluralin-contaminated prey; plovers are exposed but the trifluralin is not accumulated in plover tissues; plovers ingest trifluralin but it is not transferred to the eggs; or plover ingest trifluralin but depurate any residues before arriving on breeding grounds.

Furthermore, only a few of the target organic analytes were detected in mountain plover eggs from breeding sites in Colorado and Wyoming (Table 11; Appendix 16 – 19). With the exception of DDT (primarily p,p'-DDE, but including DDDs and DDTs), PCBs, and oxychlordanes, the organochlorine compounds were detected infrequently and generally at low

Table 11. Concentrations of detected organochlorine pesticides and total PCBs (**ng/g wet weight**) in egg samples collected from mountain plover breeding sites in Colorado and Wyoming, 2007.

| Location | Sample | % lipid | alpha BHC | 1,2,3,4-Tetrachlorobenzene | dieldrin | HCB | trans-nonachlor | heptachlor epoxide | oxychlorodane | o,p'-DDT | p,p'-DDE | PCB-TOTAL |
|--------------------------------|---------------|----------------|------------------|-----------------------------------|-----------------|------------|------------------------|---------------------------|----------------------|-----------------|-----------------|------------------|
| Pawnee National Grasslands, CO | MPWCPE06 | 0.83 | <0.435 | <0.435 | 3.37 | <0.435 | <0.435 | 0.97 | 1.27 | 1.68 | 51,200 | 33.6 |
| Pawnee National Grasslands, CO | MPWCPE07 | 7.8 | 0.566 | <0.424 | <0.424 | <0.424 | <0.424 | 1.02 | 2.72 | <0.42 | 783 | 13.8 |
| Pawnee National Grasslands, CO | MPWCPE08 | 0.82 | 1.04 | 4.07 | <0.446 | <0.446 | 0.659 | 0.85 | 0.633 | <0. | 60.2 | 29.1 |
| Pawnee National Grasslands, CO | MPWCPE09 | 1.20 | 0.697 | <0.49 | <0.49 | <0.49 | <0.49 | <0.49 | 0.834 | <0.49 | 213 | 13.3 |
| Pawnee National Grasslands, CO | MPWCPE10 | 0.95 | <0.459 | <0.459 | <0.459 | <0.459 | <0.459 | <0.46 | 3.0 | <0.459 | 35 | 9.17 |
| | | | | | | | | | | | | |
| Foote Creek Rim, WY | MPFCPE06 | 8.97 | <0.495 | <0.495 | <0.495 | 1.77 | <0.495 | <0.495 | 2.63 | <0.495 | 74.6 | 39.6 |

concentrations. Alpha-BHC, also known as hexachlorocyclohexane (HCH) was detected in three eggs from Colorado (1.97, 13.2, and 21.2 ng/g dw), at concentrations that were not much greater than the respective detection limits for the samples. That BHC was detected, albeit at low concentrations attests to the persistence of this substance, which has not been used in the U.S. since 1978 (Blus 1995; Wiemeyer 1996). The 1,2,3,4 isomer of tetrachlorobenzene was detected in one egg from Colorado but at a very low concentration (83.1 ng/g dw). This isomer is currently used in dielectric mixtures and is also a metabolite of γ HCH (lindane; Blus 1995; Handel et al. 2006).

Dieldrin was detected in one egg from Colorado at a concentration of 70.4 ng/g dw (3.37 ug/g ww). There are few data on what levels of dieldrin in eggs cause adverse effects. Brown pelicans, considered a sensitive species to OCs, had lower productivity when dieldrin concentrations in eggs were ≥ 1 ug/g ww ($\geq 1,000$ ng/g ww; Blus 1995), although this effect has not been definitely established (Peakall 1996). In addition, researchers recommend caution with the interpretation of effects of individual OCs because many OC compounds occur with other OCs, and effects, which are compound specific (e.g., lethality caused by dieldrin, or eggshell thinning caused by DDE) may or may not be additive (Blus 1995). This may be the case with this particular egg because other OCs (heptachlor epoxide, o,p'-DDT, p,p'-DDE, and oxychlorodane) were detected along with the dieldrin.

Hexachlorobenzene was detected at a concentration of 9.24 ug/g dw (1.77 ug/g ww) in the single egg collected from FCR, Wyoming. The detection does not appear to be an anomaly as HCB was detected at roughly similar concentrations (max = 8.4 ug/g dw) in multiple eggs collected from the same site in 2006 (Table 10). Unfortunately, data relating adverse effects to HCB levels in terrestrial birds are lacking. The lipid weight-based concentration detected in the plover egg from Colorado (~20 ng/g lipid) would be considered low, based on other studies of OCs in terrestrial species (Jaspers et al. 2005). The detected concentration is also orders of magnitude below levels identified as NOAEL for reproductive effects in seabirds (Boersma et al. 1986) and waterfowl (Blus et al. 1984).

Trans-nonachlor, heptachlor epoxide, and oxychlordanes detected in mountain plover eggs are all constituents of technical chlordane, with heptachlor epoxide and oxychlordanes being the more toxic products (Wiemeyer 1996). Trans-nonachlor is deemed relatively non-toxic to birds but it is possible it is metabolized to oxychlordanes (Stickel et al. 1983). While there have been a few studies on levels in the diet that are toxic, data relating chlordane concentrations, or its metabolite oxychlordanes, in eggs to reproductive effects are lacking (Wiemeyer 1996). Concentrations of heptachlor epoxide detected in plover eggs are orders of magnitude lower than concentrations that appear to be associated with adverse effects in American kestrels (*Falco sparverius*; >1,500 ng/g), Japanese quail (>10,000 ng/g), and gray partridge (*Perdix perdix*; ~3,000 – 7,000 ng/g ww; Wiemeyer 1996).

As in 2006, all eggs collected in 2007 had detectable levels of p,p'-DDE (range = 35 – 51,200 ng/g ww; Table 11). The highest concentration of DDE was from an egg collected in Colorado. The DDE residue was 51,200 ng/g ww (51.2 ug/g ww), which is well above the 3 ug/g shown to cause almost colony failure in brown pelicans (Blus 1995) and above the 16 ug/g ww level shown to cause complete nest failure in eggs of bald eagles (Wiemeyer et al. 1984). This egg also had detectable concentrations of o,p'-DDT (1.68 ng/g ww), dieldrin, heptachlor epoxide, oxychlordanes, and PCBs, which, if acting additively, increases the potential for adverse effects (Wiemeyer 1996; USDOJ 1998).

PCBs were the second most prevalent OC detected in mountain plover eggs, with concentrations ranging from 9.2 – 42 ng/g ww. PCBs were used primarily in electrical transformers and capacitors for their insulating and cooling properties. The manufacturing of PCBs was banned in the U.S. by 1977. Because of their persistence, PCBs are found worldwide (Rice and O'Keefe 1995), so it is not surprising that mountain plover eggs contained detectable total PCBs. PCBs are known to affect survival and reproductive success in birds but at levels much higher than those detected in this study with mountain plovers (Hoffman et al. 1996).

Mountain Plover Eggs-Organic Compounds-2008

In 2008, eggs were collected from BLM lands, Phillips County, Montana. Like eggs collected from Colorado and Wyoming in 2007, these had no detectable levels of trifluralin or its

metabolites, confirming the previous observation that either the plovers are not ingesting trifluralin-contaminated prey, or more probably, that they are experiencing dietary exposure, but the trifluralin is not accumulated by the mountain plovers and subsequently transferred to the eggs. Unlike the eggs collected from Colorado and Wyoming in 2007, most of the target OC compounds were detected in one or more of the eggs collected from Montana in 2008. Of the detected analytes, only nine were detected frequently (i.e., in >four out of eight samples) and three (p,p'-DDE, HCB, and PCBs) were detected in nearly all, if not all, of the eggs that were collected (Table 12; Appendices 20 – 23).

Table 12. Concentration of detected organic analytes (ng/g dw) in mountain plover eggs collected from BLM lands, Montana in 2008.

| Analyte | minimum | maximum | Detects (out of 8) |
|-----------------------------|----------------|----------------|-------------------------------|
| alpha BHC | <1.60 | 4.22 | 5 |
| alpha chlordane | <1.60 | 1.79 | 1 |
| beta BHC | <1.60 | 1.43 | 2 |
| chlorpyrifos | <1.60 | 12.8 | 4 |
| dieldrin | <1.60 | 130 | 4 |
| endosulfan II | <1.60 | 6.18 | 1 |
| endrin | <1.60 | 139 | 4 |
| gamma chlordane | <1.60 | 2.5 | 1 |
| HCB | 4.43 | 32.4 | 8 |
| heptachlor | <1.60 | 2.63 | 1 |
| heptachlor epoxide | <1.60 | 15.4 | 2 |
| mirex | <1.60 | 12.6 | 3 |
| o,p'-DDD | <1.60 | 2.44 | 1 |
| o,p'-DDE | <1.60 | 33.2 | 1 |
| o,p'-DDT | <1.60 | 6.67 | 1 |
| oxychlordane | <1.60 | 111 | 5 |
| p,p'-DDD | <1.60 | 320 | 3 |
| p,p'-DDE | 16.4 | 185,000 | 7 |
| p,p'-DDT | <1.60 | 57.6 | 3 |
| pentachloro-anisole | <1.60 | 3.54 | 4 |
| 1,2,3,4-Tetrachloro-benzene | <1.60 | 9.38 | 4 |
| 1,2,4,5-Tetrachloro-benzene | <1.60 | 20.4 | 3 |
| trans-nonachlor | <1.60 | 2.99 | 1 |
| PCB TOTAL | 75.9 | 2,760 | 8 |

Alpha BHC, alpha chlordane, beta-BHC, endosulfan II, gamma chlordane, heptachlor, pentachloro-anisole, and trans-nonachlor were detected in one or more mountain plover eggs collected from Montana but at low concentrations (Blus 1995; Wiemeyer 1996; Appendix 20). Mirex was found in three eggs at concentrations not likely to affect embryo survival (Wiemeyer 1996). Although mirex was banned in 1978, it is considered extremely persistent (Eisler 1985b) and one of its breakdown products is HCB, which was detected in all of the eggs. The HCB was detected at low concentrations (Table 12), but its presence reflects its persistence in the environment as it has not been used in the U.S. since 1978 (Blus 1995; Wiemeyer 1996). The maximum HCB concentration observed in the eggs from Montana (32.4 ug/g dw) is comparable to the highest level (32.5 ng/g dw) detected in eggs from PNG in 2006 (Table 10; Appendix 12). Although, female birds pass HCB directly into the egg (primarily the yolk), little information exists on the effects of these concentrations on embryo survival and health (Ingebrigsten et al. 1981; Breslin et al. 1983). One study on Japanese quail indicated that levels of 80 ug/g ww (8,000 ng/g) or higher were necessary before reproductive failure occurred (as cited in Bechard 1981). It is unlikely that the concentrations of HCB found in these eggs would adversely affect the mountain plover embryos.

Heptachlor, which was used primarily to control soil insect pests, was detected in only one plover egg at a low concentration (Table 12). Heptachlor, if ingested by vertebrate animals, is readily metabolized to heptachlor epoxide and like other OCs can be passed from the female bird to the egg (Henny et al. 1983; Henny et al. 1984). Heptachlor epoxide as well as trans-nonachlor and oxychlordane were contained in the insecticide chlordane but neither heptachlor nor chlordane is used in the U.S. anymore (Wiemeyer 1996). These other chlordane-related compounds were detected infrequently and at low concentrations in eggs collected from Montana. Trans-nonachlor was found in one plover egg at a low concentration (2.99 ng/g dw). This same egg also contained heptachlor epoxide and oxychlordane at low concentrations (9.84 ng/g dw and 18.7 ng/g dw, respectively). An additional egg also contained these latter two OCs at low concentrations (15.4 ng/g dw and 10.1 ng/g dw, respectively) but not trans-nonachlor. One egg contained a high concentration of oxychlordane (111 ng/g dw) but did not have any trans-nonachlor or heptachlor epoxide residues. Both heptachlor epoxide and oxychlordane are considered toxic to birds; whereas, trans-nonachlor is considered relatively non-toxic to birds. It

is possible, however, for trans-nonachlor to be metabolized to oxychlordan explaining the absence of the specific metabolite in the third egg (Stickel et al. 1983).

Dieldrin and endrin were detected in one egg at low concentrations (5.33 ng/g dw and 1.84 ng/g dw, respectively) but two other eggs had much higher concentrations of both these compounds (egg # S10P001E1 had a dieldrin and endrin residue of 130 ng/g dw and 88.5 ng/g dw, respectively; egg #S10P001E2 had a dieldrin and endrin residue of 125 ng/g dw and 139 ng/g dw, respectively). Both dieldrin and endrin are cyclodiene pesticides. Dieldrin is very persistent while endrin is less so but endrin is considered highly toxic (Peakall 1996). Blus (1995) and Peakall (1996) identified a concentration of 0.5 ug/g as the critical level for endrin in brown pelican eggs, with higher levels leading to possible impairment. In owls, reproductive impairment occurred when endrin residues in eggs were 0.27 ug/g ww (Fleming et al. 1982; Peakall 1996). The individual concentrations of dieldrin and endrin are below critical levels for pelicans and owls. But when individual OCs co-occur with other OCs there is the potential for additive effects (Blus 1995). Both of these eggs contain residues of other OCs (e.g. DDE, oxychlordan) that may increase the odds that the embryos may have been adversely affected.

DDT or one of its metabolites was detected in seven of the eight mountain plover eggs, although only p, p'-DDD, p,p'-DDE, p,p'-DDT have been shown to cause adverse effects (Blus 1996). DDT and its metabolites are the dominant class of organochlorine contaminants in plover eggs, with concentrations that are orders of magnitude greater than concentrations of the other OCs. Although at low concentrations compared with metabolites (especially DDE), DDT was detected in some eggs, even though the use of the parent compound in the U.S. was banned in 1972. The source of the DDT may be residual from past uses or it may have been transported into U.S. via wind from parts of the world where DDT is still in use (USDOI 1998). DDT was detected in most soil samples and some invertebrate samples from wintering areas in Imperial Valley. Given the proximity to Mexico, where DDT is still in use, the source of the DDT in Imperial Valley soils and invertebrates, may well be drift from Mexico. In addition, the three mountain plover eggs with detectable levels of p,p'-DDT may have overwintered in Mexico. DDD was also detected in three eggs but concentrations were not elevated and it is not associated with significant eggshell thinning like DDE (USDOI 1998). DDE concentrations in two eggs (47.5

ug/g ww and 53.2 ug/g ww) were very high and above the 3 ug/g shown to cause almost complete colony failure due to egg breakage in brown pelicans (Blus 1995). These two eggs also had concentrations greater than 16 ug/g ww, which has been found to cause complete failure in eggs of bald eagles (Wiemeyer et al. 1984). It is very possible these two eggs were not viable because of these DDE concentrations and the likelihood of either additive or synergistic effects from the other detected OC residues (Wiemeyer 1996, USDOJ 1998).

Four eggs had detectable concentrations of 1,2,3,4-tetrachlorobenzene (range 6.17 – 9.38 ng/g dw) and two eggs had detectable levels of 1,2,4,5-tetrachlorobenzene (19.0 ng/g dw and 20.4 ng/g dw; Appendix 20). These compounds may be present in the system as metabolites of the pesticide lindane (Menzie 1980). They were also used in DDT mixtures and dielectric fluids (Blus 1995; Handel et al. 2006). The concentrations observed in plover eggs are considered low (Blus 1995).

Total PCBs were detected in all eggs (75.9 – 2,760 ng/g dw; Appendix 20) at concentrations second only to DDT and its metabolites. Although banned in 1997, PCBs are found worldwide (Rice and O'Keefe 1995). Concentrations of PCBs measured in mountain plover eggs were lower than what is known to affect survival and reproductive success in other birds (Hoffman et al. 1996).

Chlorpyrifos was detected at concentrations between 3.01 ng/g dw and 12.8 ng/g dw in four of the eight eggs collected from Montana. Chlorpyrifos is a broad spectrum OP insecticide and although it is no longer used to kill mosquitoes, it is still registered for terrestrial systems (PMEP 2011). Unlike OCs, OPs are typically short-lived in the environment. The primary route of toxicity is through dietary ingestion or dermal application (Odenkirchen and Eisler 1988). These pesticides are also metabolized or excreted rapidly by most animals and do not bioaccumulate or biomagnify through the food chain (Hill 1995). OPs are not transferred from a mother bird to an egg in amounts that are biologically important (Hill 1995), although a mother can transfer the pesticide via her feathers while incubating the eggs or the egg can become contaminated from direct pesticide application (Hill 1995). No known chlorpyrifos applications occurred in the

Montana study area and it was not detected in soil or terrestrial invertebrate samples. Therefore, it is unknown as to why this insecticide was found in the mountain plover eggs.

Mountain Plover Eggs-Organic Compounds-2003

All of the eggs collected in 2003 had detectable levels of alpha-BHC, alpha-chlordane, cis-nonachlor, HCB, pentachloroanisole, trans-nonachlor, but at concentrations that are considered low (Appendices 24 – 26; Blus 1995; Wiemeyer 1996). Beta-BHC was also detected in eggs collected in 2003, with one egg from Shirley Basin, Wyoming, exhibiting a concentration (77.5 ng/g dw) that was notably higher than the other eggs. Little if anything is known about the toxicity of beta-BHC, which is the main BHC isomer retained and accumulated in animal tissues. In general, BHC is considered not very toxic to avian embryos (Wiemeyer 1996). There are limited data on γ -BHC (lindane), which is the most active as an insecticide of the BHC isomers. In two studies, no effects were observed even when concentrations of γ -BHC in eggs were as high as 10,000 ng/g for ring-necked pheasant (Blus et al. 1984) and 5,500 ng/g for American kestrel (review by Wiemeyer 1996). Beta-BHC concentrations observed in mountain plover eggs (maximum 77.5 ng/g dw) are orders of magnitude lower than no effect levels for gamma-BHC in other species and therefore are of minor concern at this time (IPCS 2010).

Dieldrin and endrin concentrations observed in mountain plover eggs were below levels of concern based on studies with brown pelicans and owls (Fleming et al. 1982; Blus 1995; Peakall 1996). However, because other OCs are present (e.g. p,p'-DDE) and the toxicity of individual OCs may be additive, there is the potential for negative effects of combined OCs on mountain plover embryos, particularly in the one egg (sample # MPSBPE04) collected from Shirley Basin, which has a very high DDE residue (Blus 1995).

Heptachlor epoxide and oxychlordane are metabolites of chlordane and heptachlor epoxide is also a metabolite of heptachlor. Most of the toxicity data on these compounds is for heptachlor epoxide for which concentrations in eggs associated with adverse effects appear to be >1,500 ng/g in American kestrels, >10,000 ng/g Japanese quail, and 3,000 – 7,000 ng/g in gray partridge (Wiemeyer 1996). Concentrations of heptachlor epoxide and oxychlordane in eggs were all below 30 ng/g dw and are orders of magnitude lower than thresholds for other species. While

concentrations of heptachlor epoxide and oxychlorodane are below levels of concern individually, there is still concern about the potential for these substances to add to the toxicity of other OCs that are present, most notably DDE (Blus 1995; Wiemeyer 1996).

Both 1,2,3,4- and 1,2,4,5-tetrachlorobenzenewere detected in eggs from Park County but not from Albany County or Shirley Basin. These products are metabolites of the insecticide lindane. Overall, concentrations were low (Appendix 25), but one egg (sample # MPCDPE05) from Park County had higher concentrations than any of the other eggs (40.8 ng/g dw and 61.6 ng/g dw, respectively). This egg also had a DDE concentration of 6,000 ng/g ww, which is above the concentration known to cause colony failure in brown pelicans (Blus 1995). It is unknown if levels of 1,2,3,4- and 1,2,4,5-tetrachlorobenzene measured in egg #MPCDPE05 are at concentrations to affect survival of mountain plover embryos. However, as stated above, these OCs may act additively with other OCs that are present (Blus 1995).

DDT was detected in three eggs (all <4.0 ng/g dw), and DDD was detected in one egg (31.6 ug/g dw) at concentrations that are considered to be low (Blus 1996; USDOJ 1998). DDE was detected in all eggs with concentrations ranging from 192 ng/g dw to 292,000 ng/g dw (Appendix 25). The egg with the highest concentration (192,000 ng/g dw or 80,000 ng/g ww) is the same egg (sample # MPSBPE04) from Shirley Basin in which many other OCs were detected and could contribute additive toxicity to the embryo (Blus 1995). At 80,000 ng/g ww (80 ug/g ww), this DDE level is significantly greater than the 3 ug/g shown to cause almost complete colony failure due to egg breakage in brown pelicans (Blus 1995) and above the 16 ug/g ww found to cause complete failure in bald eagle eggs (Wiemeyer et al. 1984). It is unknown if this egg was viable when it was collected but because of the DDE concentration it is very likely that the egg would not have hatched (Wiemeyer 1996, USDOJ 1998).

As with eggs collected in 2006 – 2008, PCBs were detected in all eggs except two eggs from Shirley Basin (Appendix 25). PCBs were the second most abundant of the OCs, after DDE, with detected concentrations that ranged from approximately 40 ng/g dw to 400 ng/g dw. Even so, concentrations were lower than what is known to affect survival and reproductive success in other birds (Hoffman et al. 1996).

CONCLUSIONS

Mountain plover on their wintering grounds in California are exposed to elevated levels of trace elements and organic contaminants. The only contaminants from the wintering grounds that appear in eggs at levels of concern are OCs. Organic contaminants were below the limits of detection in soil and invertebrate samples collected on the plovers' breeding grounds, so detection of the OC in eggs is likely linked to exposures that occur to adults on their wintering grounds in California, Arizona, Texas, and Mexico. This study only has data on wintering grounds in California, but nearly half of the United States population of mountain plover winter in the state (USFWS 2011)

DDT and its metabolites are still present in agricultural fields in the Imperial and Central Valleys of California (USDOI 1998; Kegley et al. 1999; Gervais et al. 2000), although DDT was banned in 1972. One of the most significant findings of this study was the high DDE levels found in soils and invertebrates collected in the Imperial Valley, and in eggs collected in Colorado, Montana, and Wyoming. Similar results were found in studies investigating burrowing owls using the Central and Imperial Valleys in California. Residues of p,p'-DDE in burrowing owl eggs in the Imperial Valley were documented in 1996 and 2002, and during the six years between collections, p,p'-DDE did not decline. In addition, eggshells measured in 1996 were 22% thinner than eggs collected prior to 1937 (Gervais et al. 2000; Gervais and Catlin 2004). Previous research also found mountain plover eggs contained a range of p,p'-DDE from 0.05 to 17 ug/g ww (Archuleta, USFWS, unpubl. data). Additionally, p,p'-DDT was detected in seven out of 25 adult mountain plover carcasses at concentrations from 0.02 ug/g dw to 0.36 ug/g dw (Archuleta, USFWS, unpubl.data), indicating exposure to parent product is still occurring. Table 13 compares concentration of DDE collected in this study to p,p'-DDE levels found in burrowing owl studies mentioned previously (Gervais et al. 2000; Gervais and Catlin 2004).

Mountain plover egg residues of p,p'-DDE levels were also compared to literature effects levels. Many eggs were well below any effects levels but several eggs exceeded the effects levels (Gervais et al. 2000; Figure 22). Understanding where breeding adults spent their winters may

Table 13. Concentrations of DDE in mountain plover eggs collected in Colorado, Montana, and Wyoming, compared to burrowing owl eggs collected in the Central and Imperial Valleys of California.

| Location | Species of Bird Egg collected and number (n) | Mean p,p'-DDE ug/g (ww) |
|--|---|------------------------------------|
| Pawnee National Grasslands, CO (2006) | Mountain Plover (5) | 3.44 |
| BLM Lands and CMR, Phillips County, MT (2006) | Mountain Plover (8) | 1.32 |
| Near Antelope Coal Mine and Foote Creek Rim, WY (2006) | Mountain Plover (7) | 0.324 |
| SBSS NWR (1996) | Burrowing Owl (5) | 1.19 |
| Near Pixley NWR (1996) | Burrowing Owl (7) | 0.62 |

help with future management actions. Adults spending winters in Mexico may be exposed to much higher levels of p,p'-DDE than those wintering in California.

Trifluralin was not detected in eggs collected in this investigation, although exposures likely occur to plovers through soil ingestion and invertebrate food items on wintering grounds. The RED on trifluralin identified cracked eggshells as a common endpoint in northern bobwhite quail and mallard duck exposed to the herbicide (USEPA 1996). All mountain plover eggs collected in Montana in 2008 had cracked shells. The studies cited in the RED did not report concentrations in eggs (USEPA 1996) and did not go into the reasons for decreased eggshell integrity. Because mountain plovers are exposed to trifluralin in addition to p,p'-DDE levels, they may be at increased risk of compromised eggshells from both of these contaminants.




Figure 22. Mountain plover egg data compared to p,p'-DDE effects levels

MANAGEMENT RECOMMENDATIONS

Mountain plovers winter in the Central and Imperial Valleys of California, Arizona, Texas, and Mexico. In fact, approximately half of the entire United States population winters in California (USFWS 2011), with the Imperial Valley becoming the increasingly favored location. Both the Central and Imperial Valleys report year round agriculture with over 809,400 ha (2 million acres) in production. While wintering in the Imperial Valley, mountain plovers spend the early winter in harvested or fallow alfalfa fields before switching over to Bermuda grass fields after the fields are burned in late winter (F. Knopf, pers. comm. 2/05). Because of the documented potential exposures to mountain plovers on their wintering grounds, any actions to create suitable habitat with lower contaminant concentrations on refuge grounds would be beneficial.

Although the Kern NWR is located within an intensive agricultural area, studies in 1986-1987 found that pesticide residues and trace element concentrations were comparatively low (USFWS 2000). The low concentrations measured at the Kern NWR suggest these contaminants pose little threat to wildlife; however, impounded subsurface drainage and evaporation ponds from agricultural irrigation do pose a potential threat to wildlife (USFWS 2000). Results of samples collected in 2006 also revealed the lowest levels of contaminants of any of the study sites on the wintering grounds. In the 1960's mountain plover numbers on Kern NWR ranged from 1,000 to 10,000, but no sightings have been recorded since 1985 (USFWS 2005). Similarly, another nearby refuge, Pixley NWR, not sampled in this investigation because of the presence of the endangered blunt-nosed leopard lizard, has also experienced declining plover numbers. In 1993, 109 plovers were documented, decreasing to 21 in 1994, and 15 in 1997 when the last sighting was made (USFWS 2005). As with Kern NWR, Pixley NWR is located in a highly agricultural area but studies in 1986-1987 found that pesticide residues and trace element levels were low (USFWS 1999b). Attempts to provide habitats on Kern and Pixley NWRs to attract mountain plovers could potentially reduce contaminant exposures.

In the Imperial Valley, SBSSNWR had lower contaminant concentrations than other samples collected, but still higher than samples collected on Kern NWR. Plover use on the refuge is minimal but could perhaps be managed to increase plover use in this specific area. The Refuge

has approximately 352 ha (869 acres) of croplands with many fields cooperatively farmed by local farmers. Giving preference to those farmers that plant and manage fields that attract plovers may help lower exposure to the more contaminated Imperial Valley fields.

Acknowledgements –

Funding for this project was provided by the Department of Interior's USFWS Department of Environmental Quality, Contaminants Investigation Program. Biologists from Charles M. Russell NWR provided mapping support and located potential breeding sites on the refuge as well as housing on UL Bend NWR for sampling events on CMR NWR (Randy Matchett and Jo Ann L. Dullum). Steven Dinsmore, professor, and Paul D. B. Skrade, graduate assistant with the Department of Natural Resource Ecology & Management at Iowa State University, collected all plover eggs in Montana and provided information on mountain plover breeding areas on CMR. Craig Knowles and Pamela Knowles with FaunaWest Wildlife Consultants provided information on mountain plover siting on Lake Mason NWR. Nancy Golden, Environmental Contaminant Specialist from the USFWS Washington Office, assisted with field collections of soil and invertebrate samples in Montana. Tyler Fox, biological technician for USFWS, Fritz Knopf of U.S. Geological Survey, and Regan Plumb, graduate student at the University of Wyoming, provided valuable support with sample collections in Wyoming and Colorado. Jim Lindstrom of the USFWS's Wyoming Ecological Field office developed many of the GIS maps in this report.

The following people provided assistance with the implementation of the wintering grounds portion of this study. Fritz Knopf, U.S. Geological Survey, provided technical support in the design and implementation of the wintering grounds portion of study. Christian Schoneman, refuge manager at SBSS NWR, and David Hardt, refuge manager at Kern NWR were kind enough to provide expertise and access to fields and facilities for sampling on the refuges. Vance Russell, Audubon California, and Bruce Wilcox, Imperial Irrigation District, provided invaluable information on potential off-refuge locations and contacts for requesting access. Al Kalin with the Imperial County Farm Bureau provided critical assistance with identification and initial contacts with land owners willing to let the Service collect samples for this study. We would like to thank Pat Dockstader of the Golden Eagle Hay Company and Lt. Raymond Din at

the Calipatria State Prison for allowing access to the off-refuge fields. Finally, sampling on wintering ground fields could not have been physically completed without assistance from Scott Sobiech and Carol Roberts, of the USFWS CFWO (Imperial Valley fields) and Jihadda Govan, Kern NWR biologist (Central Valley, Kern NWR field). Peter Plage of USFWS Denver Regional Office and Fritz Knopf provided helpful reviews of this report.

LITERATURE CITED

- Agency for Toxic Substances and Disease Registry (ATSDR). 2002. Toxicological profile for hexachlorobenzene. ATSDR, U.S. Public Health Service, Atlanta, Georgia. 403 pp.
- Agency for Toxic Substances and Disease Registry (ATSDR). 2000. Toxicological profile for polychlorinated biphenyls (PCBs). ATSDR, U.S. Public Health Service, Atlanta, Georgia. 948 pp.
- Agency for Toxic Substances and Disease Registry (ATSDR). 2004. Toxicological profile for strontium. ATSDR, U.S. Public Health Service, Atlanta, Georgia. 387 pp.
- Andres, B.A. and K.L. Stone. 2009. Conservation Plan for the Mountain Plover (*Charadrius montanus*), Version 1.0. Manomet Center for Conservation Sciences, Manomet, Massachusetts.
- Baudo, R., J. Giesy and H. Muntau, eds. 1990. Sediments; Chemistry and toxicity of in-place pollutants. Lewis Publishers, Inc. Boca Raton, Florida. 405 pp.
- Bechard, M. 1981. DDT and hexachlorobenzene residues in southeastern Washington Swainson's hawks (*Buteo swainsoni*). Bull Environ. Contam. Toxicol. 26: 248-253.
- Beyer, W.N., J.W. Spann, L. Sileo, and J.C. Franson. 1988. Lead poisoning in six captive avian species. Arch. Environ. Contam. Toxicol. 17: 121-130.
- Blus, L.J. 1984. DDE in birds' eggs: comparison of two methods for estimating critical levels. Wilson Bull. 96: 268-276.
- Blus, L.J. 1996. DDT, DDD, and DDE in birds. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*, W.N. Beyer, G.H. Heinz, and A.W. Redmon-Norwood, eds. CRC Press, Boca Raton, Florida. Pp. 49-71.
- Blus, L.J. 1995. Organochlorine pesticides. In: *Handbook of Ecotoxicology*, Hoffman, D.J., B.A. Rattner, G.A. Burton, Jr., and J. Cairns, Jr., eds. Lewis Publishers. New York, New York. Pp. 275-300.
- Boersma, D.C., J.A. Ellenton, and A. Yagminas. 1986. Investigation of the hepatic mixed-function oxidase system in herring gull embryos in relation to environmental contaminants. Environ. Toxicol. Chem. 5: 309-318
- Bradford, G.R., A.C. Chang, A.L. Page, D. Bakhtar, J.A. Frampton and H. Wright. 1996. Background concentrations of trace and major elements in California soils. Kearny Foundation of Soil Science, University of California, Riverside, California.

- Breslin, W.J., M.R. Bleavins, and R.K. Ringer. 1983. Distribution and excretion of hexachlorobenzene in bobwhite (*Colinus virginianus*). *J. Toxicol. Environ. Health.* 11: 885-896.
- Bunck, C.M., J.W. Spann, O.H. Pattee, and W.J. Fleming. 1985. Changes in eggshell thickness during incubation: Implications for evaluating the impact of organochlorine contaminants on productivity. *Bull. Environ. Contam. Toxicol.* 35: 173-182.
- Bureau of Reclamation (BLM). 2010. Foote Creek Rim Wind Energy Project. Accessed May 2010 at http://www.blm.gov/wy/st/en/field_offices/Rawlins.html
- Cain, B.W. and E.A. Pafford. 1981. Effects of dietary nickel on survival and growth of mallard ducklings. *Arch. Environ. Contam. Toxicol.* 10: 737-745.
- Camardese, M.B., D.J. Hoffman, L.J. LeCaptain, and G.W. Pendleton. 1990. Effects of arsenate on growth and physiology in mallard ducklings. *Environ. Toxicol. Chem.* 9: 785-795.
- Carey, A.E., J.A. Gowen, H.Tai, W.G. Mitchell and B. Wiersma. 1979. Pesticide residue levels in soils and crops from 37 states 1972 – National Soils Monitoring Program (IV). *Pest. Monit. J.* 12(4): 209-229.
- Clark, D.R., Jr., E.L. Flinckinger, D.H. Whit, R.L. Hothem, and A.A. Belisle. 1995. Dicolof and DDT residues in lizard carcasses and bird eggs from Texas, Florida, and California. *Bull. Environ. Contam. Toxicol.* 54: 817-824.
- Dinsmore, S.J. 2001. Population biology of mountain plovers in southern Phillips County, Montana. Dissertation. Dept. of Fishery and Wildlife Biology. Colorado State University.
- Dunning, J.B., Jr. 1984. Body weights of 686 species of North American birds. *Western Bird Banding Assoc. Monograph No. 1.* 41 pp.
- Eisler, R. 1985a. Cadmium hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 2. U.S. Fish and Wildlife Service, Laurel, Maryland. 46 pp.
- Eisler, R. 1985b. Mirex hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 1. U.S. Fish and Wildlife Service, Laurel, Maryland. 42 pp.
- Eisler, R. 1985c. Selenium hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 5. U.S. Fish and Wildlife Service, Laurel, Maryland. 41 pp.

- Eisler, R. 1986. Chromium hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 6. U.S. Fish and Wildlife Service, Laurel, Maryland. 60 pp.
- Eisler, R. 1987. Mercury hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 10. U.S. Fish and Wildlife Service, Laurel, Maryland. 90 pp.
- Eisler, R. 1988a. Arsenic hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No.85. U.S. Fish and Wildlife Service, Laurel, Maryland. 92 pp.
- Eisler, R. 1988b. Lead hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 14. U.S. Fish and Wildlife Service, Laurel, Maryland. 134 pp.
- Eisler, R. 1989. Molybdenum hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 85. U.S. Fish and Wildlife Service, Laurel, Maryland. 61 pp.
- Eisler, R. 1990a. Chlordane hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 21. U.S. Fish and Wildlife Service, Laurel, Maryland. 49 pp.
- Eisler, R. 1990b. Boron hazards to fish, wildlife, and invertebrates: a synoptic review. U.S. Fish Wildlife Service Biological Report. 85(1.20). 32 pp.
- Eisler, R. 1993. Zinc hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 26. U.S. Fish and Wildlife Service, Laurel, Maryland. 106 pp.
- Eisler, R. 1997. Copper hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 26. U.S. Fish and Wildlife Service, Laurel, Maryland. 98 pp.
- Eisler, R. 1998. Nickel hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 34. U.S. Fish and Wildlife Service, Laurel, Maryland. 76 pp.
- Fleming, W.J., M.A.R. McLane, and E. Cromartie. 1982. Endrin decreases screech owl productivity. J. Wildl. Manage. 46: 462-468.
- Furness, R.W. 1996. Cadmium in Birds. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*, W.N. Beyer, G.H. Heinz, and A.W. Redmon-Norwood, eds. CRC Press, Boca Raton, Florida. Pp. 389-404.

- Garcia-Hernandez, J.Y., V. Sapozhnikova, D. Schlenk, A.Z. Mason, O. Hinojosa-Huerta, J.J. Rivera-Dias, N.A. Ramos-Delgado, and G. Sanchez-Bon. 2006. Concentration of contaminants in breeding bird eggs from the Colorado River Delta, Mexico. *Environ. Toxicol. Chem.* 25: 1640-1647.
- Gervais, J.A., D.K. Rosenberg, D.M. Fry, L. Trulio, and K. K. Sturm. 2000. Burrowing owls and agricultural pesticides: evaluation of residues and risks from three populations in California, USA. *Environ. Toxicol. Chem.* 19: 337-343.
- Gervais, J.A. and D.H. Catlin. 2004. Temporal patterns of DDE in burrowing owl eggs from the Imperial Valley, California. *Southw. Assoc. Natural.* 49: 509-512.
- Gochfeld, M and J. Burger. 1998. Temporal trends in metal levels in eggs of endangered Roseate tern (*Sterna dougallii*) in New York. *Environ. Res.* 77: 36-42.
- Handel, C.M., L.M. Pajot, S.M. Matsuoka, K.A. Trust, J.M. Stotts, J. Terenzi, and S.L. Talbot. 2006. Potential role of environmental contaminants in the pathology of beak deformities among black-capped chickadees in south-central Alaska. Unpublished final report. Project ID: 1130-7F22. U.S. Geological Survey, Alaska Science Center, Anchorage, Alaska.
- Heinz, G.H. 1996. Selenium in Birds. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*, W.N. Beyer, G.H. Heinz, and A.W. Redmon-Norwood, eds. CRC Press, Boca Raton, Florida. Pp. 447-471.
- Henny, C., L.J. Blus, and C.J. Stafford. 1983. Effects of heptachlor on American kestrel in the Columbia Basin, Oregon. *J. Wildl. Manage.* 47: 1080-1087.
- Henny, C., L.J. Blus, and T.E. Kaiser. 1984. Heptachlor seed treatment contaminates hawks, owls, and eagles of Columbia Basin, Oregon. *Raptor Res.* 18: 41-48.
- Hill, E.F. 1995. Organophosphorus and carbamate pesticides. In: *Handbook of Ecotoxicology*, Hoffman, D.J., B.A. Rattner, G.A. Burton, Jr., and J. Cairns, Jr., eds. Lewis Publishers. New York, New York. Pp. 242-273.
- Hoffman, D.J. 1979. Embryotoxic effects of crude oil containing nickel and vanadium in mallards. *Bull. Environ. Contam. Toxicol.* 23: 203-206.
- Hoffman, D.J., C.P. Rice, and T.J. Kubiak. 1996. PCBs and dioxins in birds. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*, W.N. Beyer, G.H. Heinz, and A.W. Redmon-Norwood, eds. CRC Press, Boca Raton, Florida. Pp. 165-207.
- Hoffman, D.J., J.C. Franson, O.H. Pattee, C.M. Bunck, and A. Anderson. 1985. Survival, growth, and accumulation of ingested lead in nestling American kestrels (*Falco sparverius*). *Arch. Environ. Contam. Toxicol.* 14: 89-94.

- Hoyt, D.F. 1979. Practical methods of estimating volume and fresh weight of bird eggs. *The Auk*. 96: 73-77.
- Hudson River Natural Resource Trustees. 2002. Work plan for the collection of eggs from spotted sandpipers, American woodcock, belted kingfisher, American robin, red-winged blackbird, and eastern phoebe associated with the Hudson River from Hudson Falls to Schodack Island, New York. March 2002. U.S. Department of Commerce, Silver Spring, Maryland. 60 pp.
- Hulse, M., J.S. Mahoney, G.D. Schroder, C.S. Hacker, and S.M. Pier. 1980. Environmentally acquired lead, cadmium, and manganese in the cattle egret (*Bubulcus ibis*) and the laughing gull (*Larus atricilla*). *Arch. Environ. Contam. Toxicol.* 9: 63-78.
- Ingebrigsten, K., E.M. Brevik, and I. Nafstad. 1981. Distribution and elimination of hexachlorobenzene (HCB) after single oral exposure in Japanese quail (*Coturnix coturnix japonica*). *J. Toxicol. Environ. Health.* 8: 845-856.
- International Programme on Chemical Safety. 2011. Environmental Health Criteria 123, Alpha - and Beta-Hexachlorocyclohexanes. Accessed April 2011 at <http://www.inchem.org/documents/ehc/ehc/ehc123.htm#PartNumber:2>
- International Programme on Chemical Safety (IPCS). 2010. Benzene hexachloride. Accessed April 2011 at <http://www.eea.europa.eu/themes/chemicals/links/international-organisations-conventions-and-agreements/international-programme-on-chemical-safety-ipcs>
- Jaspers, V., A. Covaci, J. Maervoet, T. Dauwe, S. Voorspoels, P. Schepens and M. Eens. 2005. Brominated flame retardants and organochlorine pollutants in eggs of little owls (*Athene noctua*) from Belgium. *Environ. Pollut.* 136: 81-88
- Kabata-Pendias, A. and H. Pendias. 1992. Trace elements in soils and plants, 2nd ed. CRC Press. Boca Raton, Florida. 365 pp.
- Kegley, S., L. Neumeister, and T. Martin. 1999. Disrupting the balance: ecological impacts of pesticides in California. Pesticide Action Network. San Francisco, California. 99 pp.
- Knopf, Fritz L. and M. B. Wunder. 2006. Mountain Plover (*Charadrius montanus*), *The Birds of North America Online* (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the *Birds of North America Online*: <http://bna.birds.cornell.edu/bna/species/211>.
- Knopf, F.L. 1996. Mountain Plover (*Charadrius montanus*). In: *The Birds of North America*, No. 211, A. Poole and F. Gill, eds. The Academy of Natural Sciences. Philadelphia, PA., and The American Ornithologists' Union, Washington, D.C. 16 pp.
- Knopf, F.L., and J.R. Rupert. 1995. Habits and habitats of mountain plovers in California. *Condor.* 97: 743-751.

- Knopf, F.L., and J.R. Rupert. 1996. Productivity and movements of mountain plovers breeding in Colorado. *Wilson Bull.* 108: 28-35.
- Machlin, L.J., P.B. Pearson, C.A. Denton, and H.R. Bird. 1953. The utilization of sulfate sulfur by the laying hen and its incorporation into cystine. *J. Biolog. Chem.* 205: 213-219.
- Menzie, C.M. 1980. Metabolism of Pesticides - Update III. Special Scientific Report - Wildlife No. 232. U.S. Department of the Interior, Fish and Wildlife Service, Washington, D.C. 75 pp.
- Miles, A.K., C.E. Grue, G.W. Pendleton, and J.H. Soares, Jr. 1993. Effects of dietary aluminum, calcium, and phosphorus on egg and bone of European starlings. *Arch. Environ. Contam. Toxicol.* 24: 206-212.
- Mischke, T., K. Brunetti, V. Acosta, D. Weaver, and M. Brown. 1985. Agricultural sources of DDT residues in California's environment. Report for the Environmental Hazards Assessment Program, California Department of Foods and Agriculture, Sacramento, California.
- Miwa, K., J. Takano, H. Omori, M. Seki, K. Shinozaki and T. Fujiwara. 2007. Plants tolerant of high boron levels. *Science.* 318: 1417
- Mora, M.A. 2003. Heavy metals and metalloids in egg contents and eggshells of passerine birds from Arizona. *Environ. Pollut.* 125: 393-400.
- Nagy, K.A. 2001. Food requirements of wild animals: Predictive equations for free-living mammals, reptiles, and birds. *Nutrition Abstracts and Reviews, Series B.* 71: 21R-31R.
- NatureServe. 2010. InfoNatura: Animals and Ecosystems of Latin America [web application]. 2007. Version 5.0. Arlington, Virginia (USA). Accessed May 2010 at: <http://www.natureserve.org/infonatura>.
- Nelson, M.M., J.B. Moyle, and A.T. Farnham. 1966. Cobalt levels in foods and livers of pheasants. *J. Wildl. Manage.* 30(2): 423-425.
- Odenkirchen, E.W. and R. Eisler. 1988. Chlorpyrifos hazards to fish, wildlife, and invertebrates: A synoptic review. Contaminant hazards reviews, Report No. 13. U.S. Fish and Wildlife Service, Laurel, Maryland. 34 pp.
- Ohlendorf, H.M. 1993. Marine birds and trace elements in the temperate North Pacific. In: *The Status, Ecology, and Conservation of Marine Birds of the North Pacific*, K. Vermeer, K.T. Briggs, K.H. Morgan, and D. Siegel-Causey, eds. Special Publication, Canadian Wildlife Service, Ottawa. Pp. 232-240.

- Peakall, D.B. 1996. Dieldrin and other cyclodiene pesticides in wildlife. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*, W.N. Beyer, G.H. Heinz, and A.W. Redmon-Norwood, eds. CRC Press, Boca Raton, Florida. Pp. 73-97.
- Pesticide Action Network (PAN). 2010. Available: <http://www.panna.org/>
- Pesticide Management Education Program (PMEP). 2011. Pesticide Active Ingredient Information. Accessed May 2011 at: <http://pmez.cce.cornell.edu/profiles/insect-mite/cadusafos-cyromazine/carbaryl/index.html>
- Phipps, T., S.L. Tank, J. Wirtz, L. Brewer, A. Coyner, L.S. Ortego, and A. Fairbrother. 2002. Essentiality of nickel and homeostatic mechanisms for its regulation in terrestrial organisms. *Environ. Rev.* 10 (4): 209-261.
- Plumb, R.E. 2004. Minimum population size and concentration areas of mountain plovers breeding in Wyoming. Master's thesis, University of Wyoming.
- Rattcliffe, D.A. 1970. Changes attributable to pesticides in egg breakage frequency and eggshell thickness in some British birds. *J. Appl. Ecol.* 7: 67-115.
- Rattner, B.A., M.A. McKernan, K.M. Eisenreich, W.A. Link, G.H. Olsen, D.J. Hoffman, K.A. Knowles, P.C. McGowan. 2006. Toxicity and hazard of vanadium to mallard ducks (*Anas platyrhynchos*) and Canada geese (*Branta canadensis*). *J. Toxicol. Environ. Health, Part A.* 69: 331-351.
- Rice, C.P. and P. O'Keefe. 1995. Sources, pathways, and effects of PCBs, Dioxins, and dibenzofurans. In: *Handbook of Ecotoxicology*, Hoffman, D.J., B.A. Rattner, G.A. Burton, Jr., and J. Cairns, Jr., eds. Lewis Publishers. New York, New York. Pp. 424-468.
- Richards, M.P. 1997. Trace mineral metabolism in the avian embryo. *Poultry Sci.* 76: 152-164.
- Sample, B.E., D.M. Opresko, and G.W. Suter II. 1996. Toxicological benchmarks for wildlife: 1996 revision. Oak Ridge National Laboratory, Health Sciences Division, Oak Ridge, Tennessee. Rept. ES/ER/TM-86/R3. Prepared for the United States Department of Energy, Office of Environmental Restoration and Waste Management. 217 pp.
- Schwarzbach, S.E., J.D. Albertson, and C.M. Thomas. 2006. Effects of predation, flooding, and contamination on reproductive success of California clapper rails (*Rallus longirostris obsoletus*) in San Francisco Bay. *The Auk*: 123(1): 45-60.
- Shacklette, H.T., and J.G. Boerngen. 1984. Element concentrations in soils and other surficial materials of the conterminous United States. U.S. Geological Survey Professional Paper 1270. 105 pp.

- Sparling, D.W. and T.P. Lowe. 1996. Environmental hazards of aluminum to plants, invertebrates, fish, and wildlife. *Rev. Environ. Contam. Toxicol.* 145: 1-127.
- Stickel, L.F., W.H. Stickel, and R.A. Dyrland. 1983. Oxychlorane, HCS-3260, and nonachlor in birds: lethal residues and loss rates. *J. Toxicol. Environ. Health.* 12: 611-622.
- Thompson, D.R. 1996. Mercury in birds and terrestrial mammals. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*, W.N. Beyer, G.H. Heinz, and A.W. Redmon-Norwood, eds. CRC Press, Boca Raton, Florida. Pp. 341-356.
- Tidball, R.R. and R.J. Ebens. 1976. Regional geochemical baselines in soils of the Powder River Basin, Montana-Wyoming. U.S. Geological Survey - Twenty-eighth annual field conference. Wyoming Geological Association Guidebook. Denver, Colorado.
- United Nations Environment Programme (UNEP). 2001. Environmental Aspects of Phosphate and Potash Mining. Paris, France. 61 pp.
- U.S. Department of the Interior (USDOI). 1998. Guidelines for the interpretation of the biological effects of selected constituents in biota, water, and sediment. National Irrigation Water Quality Program. Information Report 3. Bureau of Reclamation, Denver, Colorado. 198 pp.
- U.S. Environmental Protection Agency Environmental Response Team (USEPA/ERT). 1994. SOP# 2006. Sampling equipment decontamination. U.S. Environmental Protection Agency Environmental Response Team. Edison, New Jersey.
- U. S. Environmental Protection Agency (USEPA). 1995. Great Lakes water quality initiative criteria documents for the protection of wildlife - DDT, mercury, 2,3,7,8-TCDD and PCBs. EPA-820-B-95-008. U.S. Environmental Protection Agency, Office of Water, Washington, D.C. 82 pp.
- U. S. Environmental Protection Agency (USEPA). 1996. Reregistration Eligibility Decision (RED) for trifluralin. Office of Prevention, Pesticides and Toxic Substances, EPA 738-R-95-040.
- U.S. Environmental Protection Agency (USEPA). 2000. Ecological soil screening level guidance, July 10, 2000 Draft. U.S. Environmental Protection Agency, Office of Emergency and Remedial Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2000a. Ecological soil screening level guidance, July 10, 2000 Draft, Exhibit 5-1 Review of background concentrations for metals. U.S. Environmental Protection Agency, Office of Emergency and Remedial Response. Washington, D.C.

- U.S. Environmental Protection Agency (USEPA). 2005a. Ecological soil screening levels for arsenic, Interim final. OSWER Directive 9285.7-62. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2005b. Ecological soil screening levels for barium, Interim final. OSWER Directive 9285.7-63. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2005c. Ecological soil screening levels for beryllium, Interim final. OSWER Directive 9285.7-64. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2005d. Ecological soil screening levels for cadmium, Interim final. OSWER Directive 9285.7-65. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2005e. Ecological soil screening levels for lead, Interim final. OSWER Directive 9285.7-70. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2005f. Ecological soil screening levels for vanadium, Interim final. OSWER Directive 9285.7-75. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2007a. Ecological soil screening levels for copper, Interim final. OSWER Directive 9285.7-68, February 2007 Revision. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2007b. Ecological soil screening levels for DDT and metabolites. OSWER Directive 9285.7-57. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2007c. Ecological soil screening levels for dieldrin, Interim final. OSWER Directive 9285.7-56, April 2007 Revision. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2007d. Ecological soil screening levels for manganese, Interim final. OSWER Directive 9285.7-71. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2007e. Ecological soil screening levels for nickel, Interim final. OSWER Directive 9285.7-76. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.

- U.S. Environmental Protection Agency (USEPA). 2007f. Ecological soil screening levels for polycyclic aromatic hydrocarbons (PAHs), Interim final. OSWER Directive 9285.7-78. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2007g. Ecological soil screening levels for selenium, Interim final. OSWER Directive 9285.7-72. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2007h. Ecological soil screening levels for zinc, Interim final. OSWER Directive 9285.7-73. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 2008. Ecological soil screening levels for chromium, Interim final. OSWER Directive 9285.7-66, April 2008 Revision. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. Washington, D.C.
- U.S. Fish and Wildlife Service (USFWS). 1999a. Endangered and threatened wildlife and plants; threatened status and special regulation for the mountain plover. Federal Register. 64(30): 7587-7601.
- U.S. Fish and Wildlife Service (USFWS). 1999b. Contaminant Assessment Process, Pixley National Wildlife Refuge. Accessed May 2011 at: https://ecos.fws.gov/cap/report/complete?study_id=533762
- U.S. Fish and Wildlife Service (USFWS). 2000. Contaminant Assessment Process, Kern National Wildlife Refuge. Accessed May 2011 at: https://ecos.fws.gov/cap/report/complete?study_id=519997.
- U.S. Fish and Wildlife Service (USFWS). 2002. Birds of Conservation Concern, Division of Migratory Bird Management. Arlington Virginia. 105 pp.
- U.S. Fish and Wildlife Service (USFWS). 2003a. Endangered and threatened wildlife and plants: Withdrawal of the proposed rule to list the mountain plover as threatened. Federal Register. 68(174): 53083-53101.
- U.S. Fish and Wildlife Service (USFWS). 2003b. Evaluation of the clean water act section 304(a) human health criterion for methyl mercury: protectiveness for threatened and endangered wildlife in California. U.S. Fish and Wildlife Service, Sacramento Fish and Wildlife Office, Environmental Contaminants Division. Sacramento, California. 96 pp + appendix.
- U.S. Fish and Wildlife Service (USFWS). 2005. Kern and Pixley National Wildlife Refuges Final Comprehensive Conservation Plan. 103pp.

- U.S. Fish and Wildlife Service (USFWS). 2002. Birds of Conservation Concern, Division of Migratory Bird Management. Arlington, Virginia. 87 pp.
- U.S. Fish and Wildlife Service (USFWS). 2010. Charles M. Russell NWR website. Available at: <http://www.fws.gov/cmr/wildlife.html>
- U.S. Fish and Wildlife Service (USFWS). 2011. Endangered and Threatened Wildlife and Plants: Withdrawal of the Proposed Rule To List the Mountain Plover as Threatened. Federal Register. 76(92): 27756-27799.
- U.S. Forest Service (USFS). 2010. Thunder Basin National Grassland Land and Resource Management Plan. Accessed May 2010 at: <http://www.fs.usda.gov/>
- Weber, C.W. and B.L. Reid. 1968. Nickel toxicity in growing chicks. J. Nutrit. 95: 612.
- Wennig, R. and N. Kirsch. 1950. Strontium. In: *Handbook on Toxicity of Inorganic Compounds*, H.G. Seiler and H. Sigel, eds. Marcel Dekker, Inc., New York. pp. 631-638.
- Wiemeyer, S.N. 1996. Other organochlorine pesticides in birds. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*, W.N. Beyer, G.H. Heinz, and A.W. Redmon-Norwood, eds. CRC Press, Boca Raton, Florida. Pp. 99-115.
- Wiemeyer, S.N., T.G. Lamont, C.M. Bunck, C.R. Sindelar, F.J. Grandich, J.D. Fraser, and M.A. Byrd. 1984. Organochlorine pesticide, polychlorobiphenyl, and mercury residues in bald eagle eggs, 1969-79, and their relationships to shell thinning and reproduction. Arch. Environ. Contam. Toxicol. 13: 529-549.
- Wiemeyer, S.N., P.L. Tuttle, and D.K. Higgins. 2004. Assessment of wildlife hazards associated with mine pit lakes. Environmental Contaminants Report 1F34. U.S. Fish and Wildlife Service. Nevada Field Office. 24 pp.
- Wunder, M.B. and F.L. Knopf. 2003. The Imperial Valley of California is critical to wintering mountain plovers. J. Field Ornithol. 74(1): 74-80.

Appendices

Appendix 1. Concentrations of inorganics* (ug/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006.

| Location | Sample | Al ug/g dw | As ug/g dw | B ug/g dw | Ba ug/g dw | Be ug/g dw | Ca ug/g dw | Cd ug/g dw | Co ug/g dw | Cr ug/g dw | Cu ug/g dw |
|---------------------|---------------|----------------------|----------------------|---------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| Imperial Valley J20 | S-IV-J20A | 21400 | 7.05 | 9.28 | 309 | 1.12 | 81300 | 1.48 | 8 | 27.7 | 29.5 |
| Imperial Valley J20 | S-IV-J20B | 17100 | 6.25 | 5.48 | 311 | 1.06 | 80200 | 2.54 | 7.3 | 33.4 | 30.8 |
| Imperial Valley J20 | S-IV-J20C | 20000 | 7.05 | 7.24 | 307 | 1.09 | 84800 | 1.54 | 7.79 | 28.5 | 30.8 |
| Imperial Valley J20 | S-IV-J20D | 17500 | 6.07 | 4.71 | 315 | 1.1 | 82100 | 2.13 | 7.58 | 32.3 | 32.8 |
| Imperial Valley J20 | S-IV-J20E | 23500 | 6.39 | 10.4 | 324 | 1.2 | 78600 | 1.94 | 8.41 | 32.5 | 32.1 |
| Imperial Valley J22 | S-IV-J22A | 20000 | 6.05 | 9.07 | 314 | 1.24 | 77000 | 3.26 | 7.4 | 53 | 31.3 |
| Imperial Valley J22 | S-IV-J22B | 19700 | 5.93 | 11.2 | 321 | 1.11 | 80300 | 2.56 | 7.6 | 31.3 | 34.1 |
| Imperial Valley J22 | S-IV-J22C | 19800 | 5.74 | 10.4 | 314 | 1.05 | 85700 | 1.87 | 8.04 | 27.5 | 38.7 |
| Imperial Valley J22 | S-IV-J22D | 19000 | 5.95 | 9.04 | 326 | 1.17 | 92800 | 4.66 | 7.49 | 49.2 | 35.8 |
| Imperial Valley J22 | S-IV-J22E | 17900 | 5.85 | 7.73 | 309 | 1.14 | 81600 | 4.16 | 7.17 | 45.7 | 28.5 |
| Imperial Valley K18 | S-IV-K18A | 19700 | 5 | 5.81 | 266 | 1.19 | 69000 | 2.36 | 7.61 | 36.1 | 24.6 |
| Imperial Valley K18 | S-IV-K18B | 22800 | 5.56 | 6.22 | 292 | 1.23 | 72400 | 2.34 | 8.43 | 33.3 | 28.8 |
| Imperial Valley K18 | S-IV-K18C | 24900 | 6.25 | 8.92 | 291 | 1.26 | 75800 | 1.91 | 8.5 | 30.9 | 28.7 |
| Imperial Valley K18 | S-IV-K18D | 24600 | 6 | 6.44 | 296 | 1.28 | 78900 | 2.88 | 8.45 | 37.1 | 29.7 |
| Imperial Valley K18 | S-IV-K18E | 23900 | 5.81 | 6.7 | 313 | 1.3 | 91600 | 3.78 | 8.23 | 42 | 34.3 |
| Sonny Bono NWR | S-SB-420A | 22800 | 7.08 | 6.37 | 234 | 1.17 | 51000 | 0.388 | 9.27 | 22.1 | 25.7 |
| Sonny Bono NWR | S-SB-420B | 22900 | 8.25 | 6.91 | 239 | 1.18 | 50000 | 0.435 | 9.35 | 21.7 | 26.4 |
| Sonny Bono NWR | S-SB-420C | 22500 | 6.57 | 6.27 | 242 | 1.17 | 50700 | 0.37 | 9.34 | 21.5 | 25.3 |
| Sonny Bono NWR | S-SB-420D | 24900 | 7.38 | 5.21 | 239 | 1.29 | 50500 | 0.373 | 10.2 | 23.2 | 29.6 |
| Sonny Bono NWR | S-SB-420E | 29000 | 7.39 | 9.95 | 249 | 1.41 | 50600 | 0.378 | 10.5 | 25.6 | 29.1 |
| | | | | | | | | | | | |
| Kern NWR | S-K-15A | 12800 | 3.3 | 35.8 | 152 | 0.363 | 36000 | 0.371 | 4.07 | 22 | 10.6 |
| Kern NWR | S-K-15B | 23600 | 4.23 | 72 | 131 | 0.684 | 45300 | 0.543 | 8.61 | 31.8 | 25.1 |
| Kern NWR | S-K-15C | 13100 | 2.03 | 42.7 | 110 | 0.373 | 33800 | 0.349 | 3.85 | 22 | 11.2 |
| Kern NWR | S-K-15D | 18400 | 2.93 | 55.6 | 112 | 0.52 | 37600 | 0.405 | 5.8 | 26.1 | 28.4 |
| Kern NWR | S-K-15E | 18000 | 2.12 | 34.3 | 123 | 0.49 | 29600 | 0.354 | 5.89 | 28.5 | 12.1 |

Appendix 1. cont.

| Location | Sample | Hg ug/g dw | K ug/g dw | Mg ug/g dw | Mn ug/g dw | Mo ug/g dw | Na ug/g dw | Ni ug/g dw | P ug/g dw | Pb ug/g dw | S ug/g dw |
|---------------------|---------------|----------------------|---------------------|----------------------|----------------------|----------------------|----------------------|----------------------|---------------------|----------------------|---------------------|
| Imperial Valley J20 | S-IV-J20A | 0.0153 | 4810 | 14700 | 461 | 1.18 | 2640 | 18.4 | 1370 | 14.9 | 1610 |
| Imperial Valley J20 | S-IV-J20B | 0.0155 | 3880 | 13200 | 483 | <1.02 | 847 | 17.7 | 2200 | 13.4 | 1560 |
| Imperial Valley J20 | S-IV-J20C | 0.0153 | 4480 | 14500 | 471 | <1.01 | 937 | 18.1 | 1760 | 13.8 | 1190 |
| Imperial Valley J20 | S-IV-J20D | 0.0156 | 3900 | 13900 | 503 | 1.22 | 883 | 18.5 | 2260 | 13.8 | 1500 |
| Imperial Valley J20 | S-IV-J20E | 0.0162 | 5050 | 14900 | 506 | 1.11 | 804 | 19.8 | 1770 | 14.3 | 1240 |
| Imperial Valley J22 | S-IV-J22A | 0.0182 | 4700 | 13300 | 489 | 1.39 | 995 | 20.2 | 2810 | 14.2 | 2080 |
| Imperial Valley J22 | S-IV-J22B | 0.0185 | 5240 | 15000 | 482 | 1.07 | 1460 | 18.6 | 3440 | 13.4 | 2050 |
| Imperial Valley J22 | S-IV-J22C | 0.0171 | 5250 | 14900 | 487 | <1.01 | 1310 | 18.5 | 3040 | 13.9 | 1920 |
| Imperial Valley J22 | S-IV-J22D | 0.0168 | 4770 | 14100 | 548 | 1.22 | 1370 | 21.3 | 3650 | 14.9 | 2330 |
| Imperial Valley J22 | S-IV-J22E | 0.0182 | 4100 | 12900 | 494 | 1.06 | 1240 | 19 | 2740 | 14.2 | 2090 |
| Imperial Valley K18 | S-IV-K18A | 0.0161 | 4050 | 12700 | 436 | <1.02 | 827 | 19 | 2390 | 14.4 | 1180 |
| Imperial Valley K18 | S-IV-K18B | 0.0156 | 4970 | 14300 | 460 | <1.01 | 1140 | 20.1 | 2370 | 15.8 | 1300 |
| Imperial Valley K18 | S-IV-K18C | 0.0152 | 5170 | 14700 | 449 | <1.02 | 1130 | 20.1 | 1840 | 17.6 | 1160 |
| Imperial Valley K18 | S-IV-K18D | 0.0162 | 5010 | 14400 | 476 | <1.01 | 887 | 19.9 | 2780 | 16.7 | 1080 |
| Imperial Valley K18 | S-IV-K18E | 0.0169 | 4860 | 14400 | 478 | <1.01 | 804 | 20 | 3060 | 16.5 | 1200 |
| Sonny Bono NWR | S-SB-420A | 0.0181 | 4920 | 14400 | 479 | 1.26 | 1190 | 21.5 | 781 | 18.3 | 646 |
| Sonny Bono NWR | S-SB-420B | 0.0175 | 4890 | 14500 | 478 | 1.29 | 1400 | 21.5 | 801 | 21 | 798 |
| Sonny Bono NWR | S-SB-420C | 0.0189 | 4690 | 14400 | 483 | 1.1 | 1390 | 20.4 | 795 | 17.6 | 696 |
| Sonny Bono NWR | S-SB-420D | 0.0179 | 5550 | 15300 | 519 | 1.28 | 1700 | 22.6 | 822 | 18.5 | 1180 |
| Sonny Bono NWR | S-SB-420E | 0.018 | 5960 | 16000 | 534 | 1.18 | 1460 | 23.3 | 818 | 19.2 | 748 |
| | | | | | | | | | | | |
| Kern NWR | S-K-15A | 0.00985 | 4580 | 9600 | 341 | <1.01 | 3180 | 17.4 | 2040 | 5.94 | 572 |
| Kern NWR | S-K-15B | 0.011 | 7620 | 18600 | 531 | 3.08 | 8950 | 28.7 | 2080 | 8.27 | 1340 |
| Kern NWR | S-K-15C | 0.0266 | 4670 | 10900 | 317 | <1.01 | 4110 | 17.5 | 2120 | 4.94 | 574 |
| Kern NWR | S-K-15D | 0.0108 | 6000 | 13600 | 410 | 1.03 | 7690 | 22.4 | 1590 | 6.61 | 897 |
| Kern NWR | S-K-15E | 0.00939 | 5960 | 12200 | 392 | 1.07 | 3490 | 23.1 | 1410 | 6.84 | 387 |

Appendix 1. cont.

| Location | Sample | Se ug/g dw | Sr ug/g dw | V ug/g dw | Zn ug/g dw |
|---------------------|---------------|----------------------|----------------------|---------------------|----------------------|
| Imperial Valley J20 | S-IV-J20A | 0.56 | 392 | 37.5 | 1130 |
| Imperial Valley J20 | S-IV-J20B | 0.76 | 413 | 31.3 | 3900 |
| Imperial Valley J20 | S-IV-J20C | 0.59 | 417 | 35.6 | 2060 |
| Imperial Valley J20 | S-IV-J20D | 0.74 | 418 | 31.7 | 3690 |
| Imperial Valley J20 | S-IV-J20E | 0.73 | 393 | 38 | 2040 |
| Imperial Valley J22 | S-IV-J22A | 1.11 | 399 | 36.2 | 9310 |
| Imperial Valley J22 | S-IV-J22B | 0.88 | 443 | 36.7 | 4310 |
| Imperial Valley J22 | S-IV-J22C | 0.72 | 454 | 36.1 | 1170 |
| Imperial Valley J22 | S-IV-J22D | 1.03 | 504 | 36.1 | 6230 |
| Imperial Valley J22 | S-IV-J22E | 0.84 | 405 | 35.1 | 7520 |
| Imperial Valley K18 | S-IV-K18A | 0.72 | 338 | 37.9 | 2700 |
| Imperial Valley K18 | S-IV-K18B | 0.56 | 361 | 40.2 | 1750 |
| Imperial Valley K18 | S-IV-K18C | 0.55 | 375 | 42.5 | 392 |
| Imperial Valley K18 | S-IV-K18D | 0.59 | 400 | 42.6 | 1030 |
| Imperial Valley K18 | S-IV-K18E | 0.6 | 482 | 43.4 | 1290 |
| Sonny Bono NWR | S-SB-420A | 0.38 | 216 | 40.5 | 75.2 |
| Sonny Bono NWR | S-SB-420B | 0.37 | 218 | 40.1 | 78 |
| Sonny Bono NWR | S-SB-420C | 0.39 | 221 | 39.8 | 75.8 |
| Sonny Bono NWR | S-SB-420D | 0.41 | 233 | 42.6 | 80.4 |
| Sonny Bono NWR | S-SB-420E | 0.42 | 231 | 47.1 | 84.3 |
| | | | | | |
| Kern NWR | S-K-15A | 0.08 | 284 | 25.8 | 44.9 |
| Kern NWR | S-K-15B | 0.11 | 394 | 51 | 79.4 |
| Kern NWR | S-K-15C | 0.08 | 323 | 27.4 | 44.4 |
| Kern NWR | S-K-15D | 0.08 | 332 | 36.3 | 57.4 |
| Kern NWR | S-K-15E | 0.07 | 235 | 34.7 | 57.3 |

*Perchlorate and silver not detected in any soil samples.

Appendix 2. Concentrations of organic pesticides* and trifluralin (ng/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006.

| Location | Sample | 1,2,3,4-tetra-chlorobenzene ng/g dw | 1,2,4,5-tetra-chlorobenzene ng/g dw | aldrin ng/g dw | alpha BHC ng/g dw | alpha-chlordane ng/g dw | beta BHC ng/g dw | chlorpyrifos ng/g dw | cis-nonachlor ng/g dw |
|---------------------|-----------|--|--|-------------------|----------------------|----------------------------|---------------------|-------------------------|--------------------------|
| Imperial Valley J20 | S-IV-J20A | <0.0316 | 0.092 | <0.0316 | 0.046 | 0.305 | <0.0316 | 0.266 | 1.19 |
| Imperial Valley J20 | S-IV-J20B | 0.0523 | 0.184 | 0.0591 | 0.0716 | 0.606 | <0.0283 | 0.843 | 1.48 |
| Imperial Valley J20 | S-IV-J20C | <0.0278 | 0.264 | 0.0552 | 0.0744 | 0.856 | <0.0278 | 0.91 | 2.18 |
| Imperial Valley J20 | S-IV-J20D | 0.0343 | 1.31 | 0.072 | 0.0753 | 0.589 | <0.0274 | 0.9 | 1.45 |
| Imperial Valley J20 | S-IV-J20E | <0.029 | 0.652 | 0.0762 | 0.0621 | 0.657 | <0.029 | 1.07 | 1.32 |
| Imperial Valley J22 | S-IV-J22A | <0.0277 | 1.29 | 0.0839 | 0.047 | 0.977 | <0.0277 | 0.464 | 0.853 |
| Imperial Valley J22 | S-IV-J22B | <0.0291 | 1.21 | 0.152 | 0.0436 | 1.59 | <0.0291 | 0.931 | 1.7 |
| Imperial Valley J22 | S-IV-J22C | <0.0284 | 0.199 | 0.0928 | 0.0699 | 1.51 | <0.0284 | 1.05 | 1.76 |
| Imperial Valley J22 | S-IV-J22D | <0.029 | 0.214 | 0.105 | 0.0536 | 0.585 | <0.029 | 0.589 | 0.762 |
| Imperial Valley J22 | S-IV-J22E | <0.0276 | 0.0707 | <0.0276 | 0.0354 | 0.235 | <0.0276 | 0.503 | 0.343 |
| Imperial Valley K18 | S-IV-K18A | 0.077 | 0.321 | <0.0274 | 0.0517 | 0.0363 | <0.0274 | 0.177 | 0.103 |
| Imperial Valley K18 | S-IV-K18B | <0.0275 | 0.313 | 0.0725 | 0.0457 | <0.0275 | <0.0275 | 0.284 | 0.105 |
| Imperial Valley K18 | S-IV-K18C | <0.028 | 0.48 | 0.0676 | 0.0394 | 0.036 | <0.028 | 0.243 | 0.0822 |
| Imperial Valley K18 | S-IV-K18D | 0.0898 | 0.602 | 0.0875 | 0.0505 | 0.0572 | <0.028 | 0.325 | 0.0696 |
| Imperial Valley K18 | S-IV-K18E | <0.0286 | 0.341 | 0.145 | 0.0338 | 0.0502 | <0.0286 | 0.293 | 0.0945 |
| Sonny Bono NWR | S-SB-420A | <0.0256 | 0.346 | <0.0256 | 0.178 | 0.0323 | 0.0428 | 0.489 | 0.165 |
| Sonny Bono NWR | S-SB-420B | <0.0257 | 0.318 | <0.0257 | 0.222 | <0.0257 | 0.0433 | 0.651 | 0.161 |
| Sonny Bono NWR | S-SB-420C | <0.0259 | 0.348 | <0.0259 | 0.241 | 0.0272 | 0.0418 | 0.798 | 0.17 |
| Sonny Bono NWR | S-SB-420D | <0.0259 | 0.218 | <0.0259 | 0.215 | 0.0315 | 0.0462 | 0.609 | 0.185 |
| Sonny Bono NWR | S-SB-420E | <0.026 | 0.455 | <0.026 | 0.221 | 0.0419 | 0.0765 | 0.55 | 0.164 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.0369 | 0.175 | <0.0252 | 0.0461 | 0.181 | 0.0338 | 0.11 | 0.0615 |
| Kern NWR | S-K-15B | <0.0267 | 0.148 | <0.0267 | 0.0334 | 0.121 | <0.0267 | 0.0971 | 0.0852 |
| Kern NWR | S-K-15C | <0.0261 | 0.139 | <0.0261 | 0.0295 | 0.145 | <0.0261 | 0.0411 | 0.0538 |
| Kern NWR | S-K-15D | <0.0261 | 0.147 | <0.0261 | <0.0261 | 0.0794 | 0.0307 | <0.0261 | 0.0529 |
| Kern NWR | S-K-15E | <0.0261 | 0.13 | <0.0261 | 0.0263 | 0.0727 | <0.0261 | 0.0306 | <0.0261 |

Appendix 2. cont.

| Location | Sample | delta BHC ng/g dw | dieldrin ng/g dw | endosulfan II ng/g dw | endrin ng/g dw | gamma BHC ng/g dw | gamma chlordane ng/g dw | HCB ng/g dw |
|---------------------|---------------|-----------------------------|----------------------------|---------------------------------|--------------------------|-----------------------------|-----------------------------------|-----------------------|
| Imperial Valley J20 | S-IV-J20A | <0.0316 | 1.42 | 0.133 | 0.536 | <0.0316 | 0.217 | 0.443 |
| Imperial Valley J20 | S-IV-J20B | <0.0283 | 1.58 | 0.112 | 0.55 | <0.0283 | 0.294 | 0.481 |
| Imperial Valley J20 | S-IV-J20C | 0.0293 | 2.75 | 0.246 | 0.79 | <0.0278 | 0.46 | 0.61 |
| Imperial Valley J20 | S-IV-J20D | <0.0274 | 1.71 | 0.117 | 0.589 | <0.0274 | 0.264 | 0.585 |
| Imperial Valley J20 | S-IV-J20E | <0.029 | 1.98 | 0.159 | 1 | <0.029 | 0.383 | 0.78 |
| Imperial Valley J22 | S-IV-J22A | 0.0403 | 1.85 | 0.0973 | 0.491 | <0.0277 | 0.424 | 0.933 |
| Imperial Valley J22 | S-IV-J22B | 0.053 | 3 | 0.14 | 1.02 | <0.0291 | 0.821 | 1.55 |
| Imperial Valley J22 | S-IV-J22C | <0.0284 | 3.36 | 0.121 | 1.04 | <0.0284 | 0.671 | 1.84 |
| Imperial Valley J22 | S-IV-J22D | <0.029 | 1.13 | 0.0758 | 0.6 | <0.029 | 0.3 | 0.91 |
| Imperial Valley J22 | S-IV-J22E | <0.0276 | 0.627 | <0.0276 | 0.255 | <0.0276 | 0.228 | 0.473 |
| Imperial Valley K18 | S-IV-K18A | <0.0274 | 0.165 | <0.0274 | 0.0968 | <0.0274 | 0.0275 | 0.396 |
| Imperial Valley K18 | S-IV-K18B | <0.0275 | 0.0814 | 0.0424 | 0.0825 | <0.0275 | 0.0368 | 0.566 |
| Imperial Valley K18 | S-IV-K18C | <0.028 | 0.071 | <0.028 | 0.169 | <0.028 | <0.028 | 0.47 |
| Imperial Valley K18 | S-IV-K18D | <0.028 | 0.083 | <0.028 | 0.0887 | <0.028 | <0.028 | 0.345 |
| Imperial Valley K18 | S-IV-K18E | <0.0286 | 0.0864 | <0.0286 | 0.126 | <0.0286 | <0.0286 | 0.334 |
| Sonny Bono NWR | S-SB-420A | 0.0448 | 0.275 | 0.206 | 1.51 | 0.0396 | <0.0256 | 0.277 |
| Sonny Bono NWR | S-SB-420B | 0.0485 | 0.255 | 0.132 | 1.3 | 0.0712 | <0.0257 | 0.279 |
| Sonny Bono NWR | S-SB-420C | 0.0648 | 0.272 | 0.109 | 1.28 | 0.0627 | <0.0259 | 0.29 |
| Sonny Bono NWR | S-SB-420D | 0.0682 | 0.235 | 0.15 | 1.28 | 0.0692 | <0.0259 | 0.237 |
| Sonny Bono NWR | S-SB-420E | 0.0346 | 0.276 | 0.0314 | 1.15 | 0.0566 | <0.026 | 0.256 |
| | | | | | | | | |
| Kern NWR | S-K-15A | <0.0252 | 0.227 | 0.283 | 0.203 | <0.0252 | 0.167 | 0.233 |
| Kern NWR | S-K-15B | <0.0267 | 0.105 | <0.0267 | <0.0267 | <0.0267 | 0.0788 | 0.275 |
| Kern NWR | S-K-15C | <0.0261 | 0.1 | <0.0261 | 0.167 | <0.0261 | 0.127 | 0.192 |
| Kern NWR | S-K-15D | <0.0261 | 0.0656 | <0.0261 | 0.0815 | <0.0261 | 0.103 | 0.111 |
| Kern NWR | S-K-15E | <0.0261 | 0.0727 | 0.0727 | 0.0917 | <0.0261 | 0.058 | 0.0674 |

Appendix 2. cont.

| Location | Sample | heptachlor ng/g dw | hepta- chlorepoide ng/g dw | mirex ng/g dw | oxychlorane ng/g dw | o,p'-DDD ng/g dw | o,p'-DDE ng/g dw | o,p'-DDT ng/g dw |
|---------------------|---------------|------------------------------|--|-------------------------|-------------------------------|----------------------------|----------------------------|----------------------------|
| Imperial Valley J20 | S-IV-J20A | <0.0316 | <0.0316 | 0.49 | 0.363 | 4.56 | 1.42 | 2.29 |
| Imperial Valley J20 | S-IV-J20B | 0.168 | 0.0432 | 0.855 | 0.381 | 6.66 | 1.83 | 2.99 |
| Imperial Valley J20 | S-IV-J20C | 0.149 | 0.0564 | 1.01 | 0.591 | 9 | 2.8 | 3.91 |
| Imperial Valley J20 | S-IV-J20D | 0.0554 | 0.0598 | 0.687 | 0.446 | 6.49 | 1.84 | 3.22 |
| Imperial Valley J20 | S-IV-J20E | 0.0563 | <0.029 | 1.22 | 0.368 | 8.84 | 2.48 | 5.32 |
| Imperial Valley J22 | S-IV-J22A | 0.194 | 0.0291 | 0.131 | 0.251 | 5.77 | 1.45 | 1.34 |
| Imperial Valley J22 | S-IV-J22B | 0.0966 | 0.0601 | 0.547 | 0.375 | 16.4 | 3.13 | 3.7 |
| Imperial Valley J22 | S-IV-J22C | 0.296 | 0.0916 | 0.714 | 0.419 | 15.5 | 3.05 | 4.31 |
| Imperial Valley J22 | S-IV-J22D | 0.338 | <0.029 | 0.261 | 0.216 | 5.79 | 1.59 | 1.89 |
| Imperial Valley J22 | S-IV-J22E | 0.175 | 0.0332 | 0.0718 | 0.0497 | 2.38 | 0.612 | 0.914 |
| Imperial Valley K18 | S-IV-K18A | 0.177 | <0.0274 | 0.0572 | 0.0473 | 1.21 | 0.205 | 0.536 |
| Imperial Valley K18 | S-IV-K18B | 0.157 | 0.0591 | 0.0468 | <0.0275 | 0.609 | 0.0636 | 0.278 |
| Imperial Valley K18 | S-IV-K18C | 0.195 | <0.028 | <0.028 | <0.028 | 0.458 | 0.0327 | 0.238 |
| Imperial Valley K18 | S-IV-K18D | 0.166 | <0.028 | <0.028 | <0.028 | 0.451 | 0.147 | 0.397 |
| Imperial Valley K18 | S-IV-K18E | 0.0945 | <0.0286 | <0.0286 | <0.0286 | 0.639 | 0.224 | 0.47 |
| Sonny Bono NWR | S-SB-420A | 0.121 | <0.0256 | 0.35 | <0.0256 | 1.94 | 0.859 | 3.05 |
| Sonny Bono NWR | S-SB-420B | <0.0257 | <0.0257 | 0.243 | <0.0257 | 1.94 | 0.781 | 2.8 |
| Sonny Bono NWR | S-SB-420C | <0.0259 | <0.0259 | 0.246 | <0.0259 | 2.04 | 0.844 | 3.16 |
| Sonny Bono NWR | S-SB-420D | <0.0259 | <0.0259 | 0.176 | <0.0259 | 1.64 | 0.524 | 2.46 |
| Sonny Bono NWR | S-SB-420E | <0.026 | <0.026 | 0.156 | <0.026 | 1.74 | 0.469 | 2.24 |
| | | | | | | | | |
| Kern NWR | S-K-15A | 0.103 | 0.0256 | 0.0943 | <0.0252 | 0.04 | 0.293 | 0.3 |
| Kern NWR | S-K-15B | 0.0399 | <0.0267 | 0.0464 | <0.0267 | 0.0464 | 0.0938 | 0.119 |
| Kern NWR | S-K-15C | 0.0475 | <0.0261 | 0.0464 | <0.0261 | 0.0264 | 0.131 | 0.229 |
| Kern NWR | S-K-15D | <0.0261 | <0.0261 | 0.0709 | <0.0261 | 0.0646 | 0.261 | 0.291 |
| Kern NWR | S-K-15E | 0.0495 | <0.0261 | 0.039 | <0.0261 | <0.0261 | 0.0927 | 0.0548 |

Appendix 2. cont.

| Location | Sample | p,p'-DDD ng/g dw | p,p'-DDE ng/g dw | p,p'-DDT ng/g dw | pentachloro- anisole ng/g dw | propham ng/g dw | trans- nonachlor ng/g dw | trifluralin ng/g dw |
|---------------------|-----------|---------------------|---------------------|---------------------|------------------------------------|--------------------|--------------------------------|------------------------|
| Imperial Valley J20 | S-IV-J20A | 2.27 | 195 | 6.22 | 0.124 | <0.127 | 2.01 | 0.549 |
| Imperial Valley J20 | S-IV-J20B | 3.22 | 132 | 8.37 | 0.144 | <0.113 | 2.62 | 1.22 |
| Imperial Valley J20 | S-IV-J20C | 4.71 | 345 | 10.8 | 0.105 | <0.112 | 3.51 | 0.835 |
| Imperial Valley J20 | S-IV-J20D | 3.62 | 163 | 9.7 | 0.416 | 3.73 | 2.62 | 1.3 |
| Imperial Valley J20 | S-IV-J20E | 5.89 | 317 | 18.4 | 0.0844 | <0.115 | 2.66 | 0.938 |
| Imperial Valley J22 | S-IV-J22A | 4.54 | 119 | 3.89 | 0.267 | <0.111 | 1.7 | 0.85 |
| Imperial Valley J22 | S-IV-J22B | 8.27 | 250 | 10.2 | 0.1 | <0.117 | 2.96 | 0.966 |
| Imperial Valley J22 | S-IV-J22C | 7.36 | 217 | 12.5 | 0.147 | <0.112 | 3.27 | 0.916 |
| Imperial Valley J22 | S-IV-J22D | 3.49 | 103 | 5.31 | 0.163 | 2.39 | 1.24 | 1.2 |
| Imperial Valley J22 | S-IV-J22E | 2.02 | 62 | 2.1 | 0.139 | 3.25 | 0.577 | 1.18 |
| Imperial Valley K18 | S-IV-K18A | 1.55 | 21.1 | 1.69 | 0.0627 | 4.3 | 0.165 | 0.66 |
| Imperial Valley K18 | S-IV-K18B | 1.02 | 22.2 | 1.03 | 0.0569 | 2.6 | 0.231 | 0.736 |
| Imperial Valley K18 | S-IV-K18C | 0.742 | 17.2 | 0.947 | 0.0586 | 2.28 | 0.2 | 0.597 |
| Imperial Valley K18 | S-IV-K18D | 0.763 | 11.2 | 1.48 | 0.0797 | 4.47 | 0.146 | 0.573 |
| Imperial Valley K18 | S-IV-K18E | 1.15 | 20.7 | 1.96 | 0.0583 | <0.116 | 0.198 | 1.12 |
| Sonny Bono NWR | S-SB-420A | 2.62 | 102 | 9.28 | <0.0256 | <0.104 | 0.0605 | 0.73 |
| Sonny Bono NWR | S-SB-420B | 2.55 | 92.8 | 8.82 | 0.032 | 2.47 | 0.0691 | 1.3 |
| Sonny Bono NWR | S-SB-420C | 2.98 | 110 | 9.63 | <0.0259 | <0.103 | 0.0867 | 0.7 |
| Sonny Bono NWR | S-SB-420D | 2.53 | 66.2 | 7.79 | <0.0259 | <0.104 | 0.0934 | 0.514 |
| Sonny Bono NWR | S-SB-420E | 2.59 | 60 | 7.2 | 0.0293 | <0.103 | 0.0765 | 0.901 |
| | | | | | | | | |
| Kern NWR | S-K-15A | 0.562 | 2.76 | 1.72 | 0.1 | 3.44 | 0.282 | <0.0252 |
| Kern NWR | S-K-15B | 0.291 | 0.943 | 0.858 | 0.0378 | 2.28 | 0.177 | <0.0267 |
| Kern NWR | S-K-15C | 0.545 | 1.23 | 1.74 | <0.0261 | <0.104 | 0.182 | <0.0261 |
| Kern NWR | S-K-15D | 0.786 | 3.48 | 4.24 | <0.0261 | <0.105 | 0.134 | <0.0261 |
| Kern NWR | S-K-15E | 0.0643 | 0.456 | 0.0959 | <0.0261 | <0.104 | 0.176 | <0.0261 |

*Aldicarb, azinophosmethyl, barban, bendiocarb, carbaryl, carbofuran, coumaphos, demeton, diazinon, dichlorvos/naled dimethoate, disulfoton, EPN, ethoprop, fensultothion, fenthion, malathion, merphos/tribufos, methiocarb, methomyl, methylparathion, mevinphos, monocrotophos, naled, oxamyl, parathion, phorate, protothiofos, runnel, sulfotepp, sulprofos, tetrachlorvinphos, tetraethylpyrophosphate, trichloronate, and toxaphene were not detected in any soil samples.

Appendix 3. Concentrations of PCBs* (ng/g dry weight) in soil samples collected from mountain plover wintering sites in California, 2006.

| Location | Sample | Cl3-PCB ng/g dw | Cl4-PCB ng/g dw | Cl5-PCB ng/g dw | Cl6-PCB ng/g dw | Cl7-PCB ng/g dw | Cl8-PCB ng/g dw | Cl9-PCB ng/g dw | PCB# 1 ng/g dw |
|---------------------|-----------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|-------------------|
| Imperial Valley J20 | S-IV-J20A | 6.63 | 1.99 | 6.69 | 5.7 | 5.83 | <0.632 | <0.632 | <0.00632 |
| Imperial Valley J20 | S-IV-J20B | 8.12 | 2.65 | 7.44 | 7.76 | 8.26 | <0.566 | <0.566 | 0.119 |
| Imperial Valley J20 | S-IV-J20C | 11.2 | 4.23 | 12.8 | 11.3 | 12.1 | 0.741 | 0>.557 | <0.00557 |
| Imperial Valley J20 | S-IV-J20D | 8.31 | 3.09 | 9.79 | 7.43 | 9.58 | 0.568 | >0.549 | 0.0609 |
| Imperial Valley J20 | S-IV-J20E | 16.8 | 4.01 | 11.1 | 10.1 | 11.9 | 0.954 | >0.579 | 0.0879 |
| Imperial Valley J22 | S-IV-J22A | 4.93 | 2.91 | 7.47 | 6.65 | 6.87 | <0.554 | >0.554 | 0.094 |
| Imperial Valley J22 | S-IV-J22B | 14.8 | 5.76 | 12.8 | 10.9 | 13.3 | 0.804 | <0.582 | 0.0436 |
| Imperial Valley J22 | S-IV-J22C | 13 | 5.29 | 13.2 | 10.7 | 12.8 | 0.94 | 0.607 | <0.00568 |
| Imperial Valley J22 | S-IV-J22D | 4.53 | 2.97 | 6.06 | 5.49 | 6.47 | <0.579 | <0.579 | 0.0583 |
| Imperial Valley J22 | S-IV-J22E | 1.63 | 1.68 | 2.84 | 4.09 | 3.2 | <0.552 | <0.552 | 0.0995 |
| Imperial Valley K18 | S-IV-K18A | 5.37 | 1.03 | 1.65 | 1.87 | 1.84 | <0.548 | <0.548 | 0.0198 |
| Imperial Valley K18 | S-IV-K18B | 8.93 | 1.2 | 0.737 | 1.79 | 1.63 | <0.551 | <0.551 | 0.0123 |
| Imperial Valley K18 | S-IV-K18C | 6.44 | 1.32 | 0.619 | 1.34 | 1.44 | <0.561 | <0.561 | 0.0743 |
| Imperial Valley K18 | S-IV-K18D | 3.34 | 0.914 | 0.937 | 1.48 | 1.45 | <0.561 | <0.561 | <0.00561 |
| Imperial Valley K18 | S-IV-K18E | 4.95 | 1.17 | 0.928 | 1.61 | 1.97 | <0.572 | <0.572 | 0.014 |
| Sonny Bono NWR | S-SB-420A | 0.892 | 1.18 | 2.7 | 2.71 | 5.15 | 0.577 | <0.512 | 0.0323 |
| Sonny Bono NWR | S-SB-420B | 0.998 | 1.37 | 2.84 | 2.63 | 4.95 | 0.564 | <0.514 | <0.00514 |
| Sonny Bono NWR | S-SB-420C | 1.11 | 1.25 | 2.64 | 2.55 | 4.76 | 0.598 | <0.518 | 0.134 |
| Sonny Bono NWR | S-SB-420D | 1.1 | 1.01 | 2.17 | 1.97 | 4.1 | 0.618 | <0.518 | <0.00518 |
| Sonny Bono NWR | S-SB-420E | 1.07 | 1.08 | 2.35 | 1.95 | 3.79 | 0.548 | <.521 | <0.00521 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 1.54 | 1.37 | 1.99 | 2.43 | 1.01 | <0.504 | <0.504 | <0.00504 |
| Kern NWR | S-K-15B | 1.06 | 1.28 | 0.981 | 1.87 | 0.955 | <0.534 | <0.534 | <0.00534 |
| Kern NWR | S-K-15C | 0.967 | 0.701 | 1.47 | 2.25 | 0.902 | <0.522 | <0.522 | <0.00522 |
| Kern NWR | S-K-15D | 0.583 | <0.523 | 1.04 | 1.78 | 0.605 | <0.523 | <0.523 | 0.019 |
| Kern NWR | S-K-15E | 2.49 | 0.556 | 0.826 | 1.34 | <0.521 | <0.521 | <0.521 | <0.00521 |

Appendix 3. cont.

| Location | Sample | PCB# 101/90 ng/g dw | PCB# 105 ng/g dw | PCB# 107 ng/g dw | PCB# 110 ng/g dw | PCB# 118 ng/g dw | PCB# 119 ng/g dw | PCB# 128 ng/g dw | PCB# 129 ng/g dw |
|---------------------|-----------|---------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| Imperial Valley J20 | S-IV-J20A | 0.218 | 0.0243 | 0.126 | 0.691 | 0.00766 | 0.0306 | 0.567 | 0.0192 |
| Imperial Valley J20 | S-IV-J20B | 0.307 | 0.0296 | 0.12 | 0.773 | <0.00566 | 0.145 | 0.639 | <0.00566 |
| Imperial Valley J20 | S-IV-J20C | 0.446 | 0.0406 | 0.172 | 0.855 | <0.00557 | 0.198 | 0.919 | <0.00557 |
| Imperial Valley J20 | S-IV-J20D | 0.266 | 0.0255 | 0.0952 | 1.1 | <0.00549 | 0.166 | 0.864 | <0.00549 |
| Imperial Valley J20 | S-IV-J20E | 0.279 | 0.0176 | 0.122 | 0.809 | <0.00579 | 0.14 | 1.01 | <0.00579 |
| Imperial Valley J22 | S-IV-J22A | 0.438 | 0.0459 | 0.0649 | 0.445 | 0.0358 | <0.00554 | 0.936 | 0.00671 |
| Imperial Valley J22 | S-IV-J22B | 0.422 | 0.0353 | 0.0648 | 0.628 | 0.0554 | 0.135 | 1.18 | 0.148 |
| Imperial Valley J22 | S-IV-J22C | 0.499 | 0.0619 | 0.0928 | 0.805 | 0.0573 | 0.144 | 1.39 | 0.0321 |
| Imperial Valley J22 | S-IV-J22D | 0.26 | 0.0163 | 0.0723 | 0.499 | 0.0664 | 0.0256 | 0.818 | 0.0175 |
| Imperial Valley J22 | S-IV-J22E | 0.0309 | 0.0155 | <0.00552 | 0.275 | 0.14 | 0.0652 | 0.485 | 0.103 |
| Imperial Valley K18 | S-IV-K18A | 0.0253 | <0.00548 | <0.00548 | 0.136 | <0.00548 | 0.0286 | 0.262 | 0.0396 |
| Imperial Valley K18 | S-IV-K18B | 0.0368 | 0.01 | <0.00551 | 0.0401 | <0.00551 | <0.00551 | 0.295 | 0.103 |
| Imperial Valley K18 | S-IV-K18C | 0.0282 | 0.00788 | <0.00561 | 0.0473 | <0.00561 | <0.00561 | 0.342 | 0.0845 |
| Imperial Valley K18 | S-IV-K18D | 0.0393 | <0.00561 | <0.00561 | 0.055 | 0.0101 | <0.00561 | 0.312 | 0.0236 |
| Imperial Valley K18 | S-IV-K18E | <0.00572 | <0.00572 | <0.00572 | 0.042 | 0.00933 | <0.00572 | 0.322 | <0.00572 |
| Sonny Bono NWR | S-SB-420A | <0.00512 | <0.00512 | 0.0229 | 0.191 | <0.00512 | <0.00512 | 0.297 | 0.0803 |
| Sonny Bono NWR | S-SB-420B | 0.0681 | 0.00826 | 0.0268 | 0.275 | 0.0361 | <0.00514 | 0.296 | 0.0795 |
| Sonny Bono NWR | S-SB-420C | 0.0345 | 0.0094 | 0.0334 | 0.161 | <0.00518 | <0.00518 | 0.296 | 0.0324 |
| Sonny Bono NWR | S-SB-420D | 0.0493 | 0.00944 | 0.0136 | 0.0766 | <0.00518 | <0.00518 | 0.26 | 0.0357 |
| Sonny Bono NWR | S-SB-420E | 0.0461 | 0.00524 | 0.0199 | 0.169 | 0.00524 | <0.00521 | 0.235 | 0.0231 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.129 | 0.0205 | 0.0502 | 0.0195 | 0.0143 | 0.0676 | 0.297 | <0.00504 |
| Kern NWR | S-K-15B | 0.161 | 0.0173 | 0.0205 | 0.027 | 0.00647 | 0.0226 | 0.254 | 0.00534 |
| Kern NWR | S-K-15C | 0.133 | 0.00949 | 0.0179 | 0.0179 | 0.00633 | 0.0274 | 0.401 | 0.0548 |
| Kern NWR | S-K-15D | 0.0571 | 0.0127 | 0.0116 | 0.0571 | 0.00529 | 0.019 | 0.297 | 0.00847 |
| Kern NWR | S-K-15E | 0.101 | 0.02 | 0.0158 | 0.0148 | <0.00521 | 0.0148 | 0.217 | 0.00521 |

Appendix 3. cont.

| Location | Sample | PCB# 130 ng/g dw | PCB# 135 ng/g dw | PCB# 136 ng/g dw | PCB# 138/160 ng/g dw | PCB# 141/179 ng/g dw | PCB# 149/123 ng/g dw | PCB# 15 ng/g dw | PCB# 151 ng/g dw |
|---------------------|-----------|---------------------|---------------------|---------------------|----------------------------|----------------------------|----------------------------|--------------------|---------------------|
| Imperial Valley J20 | S-IV-J20A | 2.02 | 0.24 | 0.171 | 1.24 | 0.364 | 0.378 | <0.00632 | <0.00632 |
| Imperial Valley J20 | S-IV-J20B | 2.7 | 0.282 | 0.186 | 1.66 | 0.457 | 0.513 | <0.00566 | 0.45 |
| Imperial Valley J20 | S-IV-J20C | 4.3 | 0.564 | 0.198 | 2.57 | 0.575 | 0.74 | <0.00557 | <0.00557 |
| Imperial Valley J20 | S-IV-J20D | 2.55 | 0.301 | 0.172 | 1.69 | 0.416 | 0.483 | <0.00549 | <0.00549 |
| Imperial Valley J20 | S-IV-J20E | 2.98 | 0.395 | 0.205 | 1.99 | 0.491 | 0.572 | <0.00579 | 0.694 |
| Imperial Valley J22 | S-IV-J22A | 2.17 | 0.469 | 0.22 | 0.718 | 0.424 | 0.674 | <0.00554 | <0.00554 |
| Imperial Valley J22 | S-IV-J22B | 3.84 | 0.812 | 0.188 | 1.41 | 0.285 | 1.02 | <0.00582 | 0.481 |
| Imperial Valley J22 | S-IV-J22C | 3.9 | 0.743 | 0.126 | 1.51 | 0.298 | 0.969 | <0.00568 | <0.00568 |
| Imperial Valley J22 | S-IV-J22D | 1.81 | 0.321 | 0.0909 | 0.732 | 0.293 | 0.478 | <0.00579 | <0.00579 |
| Imperial Valley J22 | S-IV-J22E | 1.24 | 0.112 | 0.0508 | 0.568 | 0.239 | 0.389 | <0.00552 | <0.00552 |
| Imperial Valley K18 | S-IV-K18A | 0.701 | 0.0231 | 0.0891 | 0.293 | 0.151 | 0.135 | <0.00548 | <0.00548 |
| Imperial Valley K18 | S-IV-K18B | 0.488 | <0.00551 | 0.0201 | 0.249 | 0.173 | 0.111 | <0.00551 | <0.00551 |
| Imperial Valley K18 | S-IV-K18C | 0.354 | <0.00561 | 0.0146 | 0.164 | 0.133 | 0.0912 | <0.00561 | <0.00561 |
| Imperial Valley K18 | S-IV-K18D | 0.486 | 0.0539 | 0.018 | 0.263 | 0.129 | 0.102 | <0.00561 | <0.00561 |
| Imperial Valley K18 | S-IV-K18E | 0.539 | 0.049 | 0.0163 | 0.242 | 0.148 | 0.127 | <0.00572 | <0.00572 |
| Sonny Bono NWR | S-SB-420A | 0.663 | 0.0469 | 0.0657 | 0.423 | 0.131 | 0.126 | <0.00512 | 0.212 |
| Sonny Bono NWR | S-SB-420B | 0.601 | 0.0464 | 0.064 | 0.393 | 0.169 | 0.136 | <0.00514 | 0.227 |
| Sonny Bono NWR | S-SB-420C | 0.625 | 0.0449 | 0.0585 | 0.339 | 0.126 | 0.116 | 0.0115 | 0.222 |
| Sonny Bono NWR | S-SB-420D | 0.531 | 0.022 | 0.0367 | 0.301 | 0.084 | 0.107 | 0.00735 | 0.147 |
| Sonny Bono NWR | S-SB-420E | 0.48 | 0.0587 | 0.0493 | 0.297 | 0.107 | 0.123 | <0.00521 | 0.151 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.701 | 0.0471 | 0.00512 | 0.503 | 0.219 | 0.264 | 0.0707 | <0.00504 |
| Kern NWR | S-K-15B | 0.432 | 0.0809 | 0.00539 | 0.299 | 0.352 | 0.152 | 0.0529 | <0.00534 |
| Kern NWR | S-K-15C | 0.568 | 0.179 | 0.00527 | 0.36 | 0.151 | 0.204 | 0.0696 | 0.0243 |
| Kern NWR | S-K-15D | 0.414 | 0.141 | 0.00529 | 0.326 | 0.14 | 0.142 | 0.0201 | 0.0434 |
| Kern NWR | S-K-15E | 0.35 | 0.0284 | 0.00738 | 0.227 | 0.159 | 0.13 | 0.0306 | 0.0348 |

Appendix 3. cont.

| Location | Sample | PCB# 153/132 ng/g dw | PCB# 156 ng/g dw | PCB# 157/173/201 ng/g dw | PCB# 158 ng/g dw | PCB# 16/32 ng/g dw | PCB# 166 ng/g dw | PCB# 167 ng/g dw | PCB# 170/190 ng/g dw |
|---------------------|-----------|----------------------------|---------------------|--------------------------------|---------------------|--------------------------|---------------------|---------------------|----------------------------|
| Imperial Valley J20 | S-IV-J20A | 0.416 | <0.00632 | 0.0434 | 0.0843 | 0.0588 | 0.0115 | 0.175 | 0.0358 |
| Imperial Valley J20 | S-IV-J20B | 0.474 | <0.00566 | <0.00566 | 0.118 | <0.00566 | <0.00566 | 0.238 | 0.0489 |
| Imperial Valley J20 | S-IV-J20C | 0.805 | <0.00557 | <0.00557 | 0.141 | 0.235 | <0.00557 | 0.353 | 0.144 |
| Imperial Valley J20 | S-IV-J20D | 0.495 | <0.00549 | <0.00549 | 0.117 | <0.00549 | <0.00549 | 0.32 | 0.0631 |
| Imperial Valley J20 | S-IV-J20E | 0.735 | <0.00579 | <0.00579 | 0.121 | 0.236 | 0.0328 | 0.347 | 0.217 |
| Imperial Valley J22 | S-IV-J22A | 0.415 | 0.0112 | 0.0257 | 0.0716 | 0.246 | 0.0112 | 0.523 | 0.123 |
| Imperial Valley J22 | S-IV-J22B | 0.66 | 0.164 | 0.0471 | 0.23 | 0.408 | 0.00589 | 0.463 | 0.152 |
| Imperial Valley J22 | S-IV-J22C | 0.777 | 0.079 | 0.0527 | 0.111 | 0.129 | 0.00573 | 0.788 | 0.186 |
| Imperial Valley J22 | S-IV-J22D | 0.3 | 0.0373 | 0.0326 | 0.0699 | 0.255 | <0.00579 | 0.517 | 0.155 |
| Imperial Valley J22 | S-IV-J22E | 0.313 | 0.259 | 0.0409 | 0.108 | 0.356 | <0.00552 | 0.134 | 0.0862 |
| Imperial Valley K18 | S-IV-K18A | 0.0814 | <0.00548 | 0.0209 | 0.0374 | 0.228 | <0.00548 | 0.0627 | 0.0715 |
| Imperial Valley K18 | S-IV-K18B | 0.0903 | 0.128 | 0.0312 | 0.0346 | 0.298 | <0.00551 | 0.068 | 0.146 |
| Imperial Valley K18 | S-IV-K18C | 0.0721 | <0.00561 | 0.0158 | 0.0169 | 0.259 | <0.00561 | 0.0338 | 0.106 |
| Imperial Valley K18 | S-IV-K18D | 0.0875 | <0.00561 | 0.0314 | <0.00561 | 0.169 | <0.00561 | <0.00561 | 0.0965 |
| Imperial Valley K18 | S-IV-K18E | 0.0712 | <0.00572 | 0.0233 | 0.007 | 0.3 | <0.00572 | 0.0898 | 0.167 |
| Sonny Bono NWR | S-SB-420A | 0.109 | <0.00512 | 0.0282 | 0.0626 | 0.128 | <0.00512 | 0.363 | 0.0354 |
| Sonny Bono NWR | S-SB-420B | 0.185 | <0.00514 | 0.0248 | 0.0475 | 0.0836 | <0.00514 | 0.312 | 0.0351 |
| Sonny Bono NWR | S-SB-420C | 0.168 | 0.0178 | 0.0366 | 0.0261 | 0.15 | 0.0125 | 0.359 | 0.0491 |
| Sonny Bono NWR | S-SB-420D | 0.139 | 0.0178 | 0.0273 | 0.0241 | 0.151 | 0.0063 | 0.256 | 0.0315 |
| Sonny Bono NWR | S-SB-420E | 0.132 | 0.00839 | 0.0335 | 0.0262 | 0.135 | 0.00839 | 0.254 | 0.0252 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.272 | 0.0379 | <0.00504 | 0.0082 | 0.965 | 0.0123 | 0.0646 | 0.00615 |
| Kern NWR | S-K-15B | 0.174 | <0.00534 | 0.0432 | <0.00534 | 0.565 | <0.00534 | 0.0636 | 0.0388 |
| Kern NWR | S-K-15C | 0.178 | 0.00527 | 0.00633 | 0.00633 | 0.637 | 0.0148 | 0.0981 | 0.0137 |
| Kern NWR | S-K-15D | 0.161 | 0.00741 | 0.00847 | 0.00529 | 0.276 | <0.00523 | 0.0698 | 0.0127 |
| Kern NWR | S-K-15E | 0.124 | <0.00521 | 0.0126 | 0.00738 | 2 | <0.00521 | 0.0527 | 0.00948 |

Appendix 3. cont.

| Location | Sample | PCB# 171/202 ng/g dw | PCB# 172 ng/g dw | PCB# 174 ng/g dw | PCB# 175 ng/g dw | PCB# 176/137 ng/g dw | PCB# 177 ng/g dw | PCB# 178 ng/g dw | PCB# 18/17 ng/g dw |
|---------------------|-----------|-------------------------|---------------------|---------------------|---------------------|-------------------------|---------------------|---------------------|-----------------------|
| Imperial Valley J20 | S-IV-J20A | <0.00632 | <0.00632 | 0.0549 | 0.147 | 0.163 | 0.105 | 0.548 | <0.00632 |
| Imperial Valley J20 | S-IV-J20B | 0.0375 | 0.058 | 0.0852 | 0.175 | 0.289 | 0.144 | 0.753 | <0.00566 |
| Imperial Valley J20 | S-IV-J20C | 0.139 | 0.118 | 0.0924 | 0.284 | 0.319 | <0.00557 | 1.03 | <0.00557 |
| Imperial Valley J20 | S-IV-J20D | 0.185 | 0.156 | 0.0831 | 0.189 | 0.496 | <0.00549 | 0.75 | <0.00549 |
| Imperial Valley J20 | S-IV-J20E | 0.0363 | 0.147 | 0.0973 | 0.297 | 0.477 | <0.00579 | 0.965 | 0.0973 |
| Imperial Valley J22 | S-IV-J22A | 0.0336 | 0.0492 | 0.0895 | 0.164 | 0.175 | 0.104 | 0.642 | 0.101 |
| Imperial Valley J22 | S-IV-J22B | 0.0306 | 0.0224 | 0.105 | 0.282 | 0.346 | 0.0919 | 1.48 | 0.053 |
| Imperial Valley J22 | S-IV-J22C | 0.00573 | 0.047 | 0.0939 | 0.273 | 0.328 | 0.108 | 1.31 | 0.0596 |
| Imperial Valley J22 | S-IV-J22D | 0.0256 | 0.0245 | 0.0629 | 0.15 | 0.223 | 0.118 | 0.603 | 0.0944 |
| Imperial Valley J22 | S-IV-J22E | 0.00773 | 0.0387 | 0.0608 | 0.0983 | 0.169 | 0.0774 | 0.269 | <0.00552 |
| Imperial Valley K18 | S-IV-K18A | <0.00548 | 0.0099 | 0.0132 | 0.066 | 0.163 | 0.0803 | 0.107 | 0.109 |
| Imperial Valley K18 | S-IV-K18B | 0.0078 | 0.0279 | 0.0223 | 0.0636 | 0.192 | 0.0847 | 0.0925 | 0.201 |
| Imperial Valley K18 | S-IV-K18C | 0.00676 | 0.0158 | 0.0135 | 0.0574 | 0.167 | 0.0957 | 0.071 | 0.0372 |
| Imperial Valley K18 | S-IV-K18D | 0.0314 | 0.00786 | 0.0707 | 0.0539 | 0.139 | 0.104 | 0.073 | <0.00561 |
| Imperial Valley K18 | S-IV-K18E | 0.0187 | 0.0128 | 0.0513 | 0.0642 | 0.247 | 0.0642 | 0.106 | <0.00572 |
| Sonny Bono NWR | S-SB-420A | 0.0282 | 0.0167 | 0.073 | 0.103 | 0.0949 | 0.072 | 0.36 | 0.0751 |
| Sonny Bono NWR | S-SB-420B | 0.033 | 0.0103 | 0.0712 | 0.0929 | 0.0836 | 0.0898 | 0.291 | 0.0846 |
| Sonny Bono NWR | S-SB-420C | 0.0408 | 0.0104 | 0.0679 | 0.115 | 0.00731 | 0.094 | 0.311 | 0.0554 |
| Sonny Bono NWR | S-SB-420D | 0.0493 | 0.0168 | 0.0724 | 0.0913 | 0.0556 | 0.0986 | 0.221 | 0.129 |
| Sonny Bono NWR | S-SB-420E | 0.0367 | 0.0105 | 0.0734 | 0.0828 | 0.0356 | 0.124 | 0.207 | 0.0398 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.0215 | 0.041 | 0.0369 | 0.0768 | 0.0973 | 0.204 | 0.0133 | 0.0471 |
| Kern NWR | S-K-15B | 0.0291 | 0.041 | 0.0205 | 0.0561 | 0.0442 | 0.42 | 0.00539 | 0.0313 |
| Kern NWR | S-K-15C | 0.00949 | 0.0422 | 0.019 | 0.0707 | 0.0844 | 0.121 | 0.0707 | 0.0232 |
| Kern NWR | S-K-15D | 0.00847 | 0.0148 | 0.0212 | 0.0709 | 0.0635 | 0.0487 | 0.00529 | 0.0222 |
| Kern NWR | S-K-15E | 0.0158 | 0.0432 | 0.0148 | 0.0548 | 0.0327 | 0.0864 | <0.00521 | 0.0179 |

Appendix 3. cont.

| Location | Sample | PCB# 180 ng/g dw | PCB# 183 ng/g dw | PCB# 185 ng/g dw | PCB# 187 ng/g dw | PCB# 189 ng/g dw | PCB# 191 ng/g dw | PCB# 193 ng/g dw | PCB# 194 ng/g dw |
|---------------------|-----------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| Imperial Valley J20 | S-IV-J20A | 0.0255 | 3.94 | 0.307 | 0.354 | <0.00632 | <0.00632 | 0.139 | <0.00632 |
| Imperial Valley J20 | S-IV-J20B | 0.0989 | 6.03 | 0.433 | 0.0102 | <0.00566 | 0.0204 | 0.0841 | <0.00566 |
| Imperial Valley J20 | S-IV-J20C | 0.0552 | 8.88 | 0.684 | 0.0169 | 0.0564 | 0.0609 | 0.165 | <0.00557 |
| Imperial Valley J20 | S-IV-J20D | 0.0277 | 6.71 | 0.495 | 0.00997 | 0.0321 | 0.0244 | 0.364 | <0.00549 |
| Imperial Valley J20 | S-IV-J20E | 0.0492 | 8.37 | 0.496 | 0.0352 | 0.0586 | 0.0199 | 0.614 | 0.135 |
| Imperial Valley J22 | S-IV-J22A | 0.0805 | 4.42 | 0.619 | 0.301 | 0.00783 | <0.00554 | 0.0548 | 0.0548 |
| Imperial Valley J22 | S-IV-J22B | 0.053 | 9.18 | 0.631 | 0.733 | 0.0495 | <0.00582 | 0.101 | 0.066 |
| Imperial Valley J22 | S-IV-J22C | 0.0561 | 8.62 | 0.747 | 0.821 | 0.0412 | <0.00568 | 0.12 | 0.215 |
| Imperial Valley J22 | S-IV-J22D | 0.0664 | 4.24 | 0.425 | 0.307 | 0.00583 | <0.00579 | 0.0618 | 0.0291 |
| Imperial Valley J22 | S-IV-J22E | 0.135 | 1.85 | 0.211 | 0.16 | 0.00663 | 0.00663 | 0.0309 | 0.0497 |
| Imperial Valley K18 | S-IV-K18A | 0.0671 | 0.998 | 0.0891 | 0.077 | 0.0099 | 0.0066 | 0.077 | 0.0121 |
| Imperial Valley K18 | S-IV-K18B | 0.0223 | 0.715 | 0.0847 | 0.0691 | 0.00557 | 0.00892 | 0.0847 | 0.0279 |
| Imperial Valley K18 | S-IV-K18C | 0.0248 | 0.636 | 0.08 | 0.0642 | 0.0203 | 0.00788 | 0.0777 | 0.0124 |
| Imperial Valley K18 | S-IV-K18D | 0.0247 | 0.64 | 0.0651 | 0.0415 | 0.00898 | <0.00561 | 0.0954 | 0.00561 |
| Imperial Valley K18 | S-IV-K18E | 0.0408 | 0.782 | 0.0898 | 0.107 | 0.0257 | <0.00572 | 0.189 | 0.0338 |
| Sonny Bono NWR | S-SB-420A | 0.0188 | 3.98 | 0.105 | 0.166 | 0.0229 | <0.00512 | 0.074 | <0.00512 |
| Sonny Bono NWR | S-SB-420B | 0.0402 | 3.88 | 0.096 | 0.142 | 0.0279 | <0.00514 | 0.0568 | <0.00514 |
| Sonny Bono NWR | S-SB-420C | 0.0136 | 3.71 | 0.112 | 0.152 | 0.0146 | <0.00518 | 0.0616 | <0.00518 |
| Sonny Bono NWR | S-SB-420D | 0.0304 | 3.15 | 0.0871 | 0.116 | 0.0189 | <0.00518 | 0.0504 | <0.00518 |
| Sonny Bono NWR | S-SB-420E | 0.0157 | 2.94 | 0.0849 | 0.086 | 0.0178 | <0.00521 | 0.0472 | <0.00521 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.0768 | 0.106 | 0.112 | 0.145 | 0.0123 | <0.00504 | 0.0564 | 0.0266 |
| Kern NWR | S-K-15B | 0.0755 | 0.0626 | 0.0518 | 0.0529 | 0.00755 | 0.00539 | 0.0464 | 0.0647 |
| Kern NWR | S-K-15C | 0.057 | 0.148 | 0.077 | 0.111 | 0.0106 | <0.00522 | 0.0654 | 0.0232 |
| Kern NWR | S-K-15D | 0.0656 | 0.101 | 0.0582 | 0.0762 | 0.00952 | <0.00523 | 0.0487 | 0.00952 |
| Kern NWR | S-K-15E | 0.0485 | 0.0316 | 0.0622 | 0.0601 | <0.00521 | 0.00527 | 0.019 | 0.0158 |

Appendix 3. cont.

| Location | Sample | PCB# 195/208 ng/g dw | PCB# 196 ng/g dw | PCB# 197 ng/g dw | PCB# 199 ng/g dw | PCB# 200 ng/g dw | PCB# 205 ng/g dw | PCB# 206 ng/g dw | PCB# 207 ng/g dw |
|---------------------|-----------|-------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| Imperial Valley J20 | S-IV-J20A | 0.0179 | 0.0447 | 0.0192 | 0.166 | <0.00632 | 0.0422 | 0.0881 | 0.103 |
| Imperial Valley J20 | S-IV-J20B | <0.00566 | 0.142 | 0.0216 | 0.249 | <0.00566 | 0.108 | 0.0989 | 0.147 |
| Imperial Valley J20 | S-IV-J20C | <0.00557 | 0.257 | 0.0406 | 0.371 | <0.00557 | 0.0733 | 0.266 | 0.216 |
| Imperial Valley J20 | S-IV-J20D | <0.00549 | 0.192 | 0.0831 | 0.24 | <0.00549 | 0.0532 | 0.184 | 0.157 |
| Imperial Valley J20 | S-IV-J20E | 0.00586 | 0.286 | 0.0293 | 0.397 | 0.0434 | 0.0574 | 0.236 | 0.209 |
| Imperial Valley J22 | S-IV-J22A | 0.0447 | 0.0212 | <0.00554 | 0.197 | 0.0392 | 0.0302 | 0.0447 | 0.0101 |
| Imperial Valley J22 | S-IV-J22B | 0.0554 | 0.0424 | 0.0471 | 0.45 | 0.0165 | 0.0801 | 0.309 | 0.252 |
| Imperial Valley J22 | S-IV-J22C | 0.0458 | 0.0756 | 0.0321 | 0.432 | 0.0114 | 0.0768 | 0.275 | 0.332 |
| Imperial Valley J22 | S-IV-J22D | 0.0676 | 0.0198 | <0.00579 | 0.235 | <0.00579 | 0.0525 | 0.0466 | 0.0268 |
| Imperial Valley J22 | S-IV-J22E | <0.00552 | 0.108 | 0.0177 | 0.122 | <0.00552 | 0.0508 | 0.255 | 0.0365 |
| Imperial Valley K18 | S-IV-K18A | <0.00548 | 0.0583 | 0.0132 | 0.228 | <0.00548 | 0.0253 | 0.0649 | <0.00548 |
| Imperial Valley K18 | S-IV-K18B | <0.00551 | 0.019 | 0.0223 | 0.402 | 0.00669 | 0.0178 | 0.167 | <0.00551 |
| Imperial Valley K18 | S-IV-K18C | <0.00561 | 0.0293 | 0.0191 | 0.375 | 0.00676 | 0.0135 | 0.16 | <0.00561 |
| Imperial Valley K18 | S-IV-K18D | <0.00561 | <0.00561 | 0.00898 | 0.239 | <0.00561 | 0.0449 | 0.0606 | 0.0101 |
| Imperial Valley K18 | S-IV-K18E | <0.00572 | <0.00572 | 0.00933 | 0.238 | <0.00572 | 0.028 | 0.132 | 0.00933 |
| Sonny Bono NWR | S-SB-420A | <0.00512 | 0.0438 | 0.0667 | 0.424 | <0.00512 | 0.0115 | 0.0709 | 0.0521 |
| Sonny Bono NWR | S-SB-420B | <0.00514 | 0.0485 | 0.0733 | 0.4 | <0.00514 | 0.0175 | 0.031 | <0.00514 |
| Sonny Bono NWR | S-SB-420C | 0.00627 | 0.0491 | 0.0732 | 0.398 | 0.0104 | 0.023 | <0.00518 | <0.00518 |
| Sonny Bono NWR | S-SB-420D | 0.00735 | 0.0588 | 0.0598 | 0.437 | 0.0126 | 0.0136 | 0.0063 | <0.00518 |
| Sonny Bono NWR | S-SB-420E | 0.00521 | 0.0388 | 0.065 | 0.382 | 0.0126 | 0.0115 | <0.00521 | <0.00521 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.0082 | 0.0891 | 0.0102 | 0.0256 | 0.0164 | 0.11 | 0.04 | <0.00504 |
| Kern NWR | S-K-15B | 0.00863 | 0.0464 | 0.0119 | 0.125 | 0.0162 | 0.147 | 0.0895 | <0.00534 |
| Kern NWR | S-K-15C | 0.0158 | 0.0422 | 0.0222 | 0.0137 | 0.0116 | 0.0939 | 0.0369 | <0.00522 |
| Kern NWR | S-K-15D | 0.00847 | 0.019 | 0.00635 | 0.018 | <0.00523 | 0.0339 | 0.0148 | 0.018 |
| Kern NWR | S-K-15E | 0.00632 | 0.0664 | 0.00738 | 0.0158 | 0.00632 | 0.0432 | 0.0169 | <0.00521 |

Appendix 3. cont.

| Location | Sample | PCB# 209 ng/g dw | PCB# 22/51 ng/g dw | PCB# 24/27 ng/g dw | PCB# 25 ng/g dw | PCB# 26 ng/g dw | PCB# 28 ng/g dw | PCB# 29 ng/g dw | PCB# 30 ng/g dw |
|---------------------|-----------|---------------------|-----------------------|-----------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| Imperial Valley J20 | S-IV-J20A | <0.00632 | <0.00632 | <0.00632 | <0.00632 | 0.198 | 0.0907 | 5.76 | <0.00632 |
| Imperial Valley J20 | S-IV-J20B | 0.0148 | 0.0307 | 0.0114 | <0.00566 | 0.241 | 0.589 | 6.74 | <0.00566 |
| Imperial Valley J20 | S-IV-J20C | 0.0519 | 0.0406 | 0.0361 | <0.00557 | <0.00557 | 0.445 | 10.2 | 0.0541 |
| Imperial Valley J20 | S-IV-J20D | 0.0487 | 0.00886 | 0.00886 | <0.00549 | 0.0775 | 0.698 | 7.48 | <0.00549 |
| Imperial Valley J20 | S-IV-J20E | 0.0879 | 0.0516 | 0.0316 | <0.00579 | 0.149 | 0.198 | 16 | <0.00579 |
| Imperial Valley J22 | S-IV-J22A | 0.0962 | 0.103 | <0.00554 | 0.0705 | 0.372 | 0.661 | 2.26 | 0.372 |
| Imperial Valley J22 | S-IV-J22B | 0.0978 | 0.174 | <0.00582 | 0.0294 | 0.15 | 0.224 | 12.1 | 0.0907 |
| Imperial Valley J22 | S-IV-J22C | 0.166 | 0.32 | <0.00568 | 0.0596 | 0.307 | 0.255 | 10.9 | 0.15 |
| Imperial Valley J22 | S-IV-J22D | 0.035 | 0.0792 | 0.0454 | 0.0373 | 0.0723 | 0.279 | 2.65 | 0.507 |
| Imperial Valley J22 | S-IV-J22E | 0.0398 | 0.0354 | 0.011 | <0.00552 | <0.00552 | 0.134 | 0.864 | 0.0243 |
| Imperial Valley K18 | S-IV-K18A | 0.0198 | 0.0099 | 0.0561 | 0.022 | 0.108 | 0.29 | 3.89 | 0.0187 |
| Imperial Valley K18 | S-IV-K18B | 0.0123 | 0.0424 | 0.0201 | <0.00551 | 0.106 | 0.184 | 7.51 | <0.00551 |
| Imperial Valley K18 | S-IV-K18C | 0.0214 | 0.0124 | 0.036 | <0.00561 | 0.101 | 0.233 | 5.29 | <0.00561 |
| Imperial Valley K18 | S-IV-K18D | 0.118 | 0.00898 | <0.00561 | <0.00561 | 0.0584 | 0.15 | 2.62 | <0.00561 |
| Imperial Valley K18 | S-IV-K18E | 0.0653 | 0.0128 | <0.00572 | <0.00572 | 0.0432 | 0.261 | 3.86 | <0.00572 |
| Sonny Bono NWR | S-SB-420A | 0.0073 | 0.0198 | <0.00512 | <0.00512 | 0.025 | 0.0365 | 0.593 | <0.00512 |
| Sonny Bono NWR | S-SB-420B | 0.0134 | 0.0155 | <0.00514 | <0.00514 | 0.0351 | 0.0248 | 0.695 | <0.00514 |
| Sonny Bono NWR | S-SB-420C | 0.0125 | 0.00627 | <0.00518 | <0.00518 | 0.0428 | 0.0251 | 0.717 | <0.00518 |
| Sonny Bono NWR | S-SB-420D | 0.00944 | 0.00839 | <0.00518 | <0.00518 | 0.022 | 0.0556 | 0.717 | <0.00518 |
| Sonny Bono NWR | S-SB-420E | 0.0115 | 0.0136 | 0.00629 | <0.00521 | 0.0451 | 0.0126 | 0.77 | <0.00521 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.0441 | <0.00504 | 0.0133 | <0.00504 | 0.0195 | <0.00504 | 0.395 | 0.00922 |
| Kern NWR | S-K-15B | 0.0496 | 0.0658 | 0.00755 | <0.00534 | <0.00534 | 0.0421 | 0.175 | <0.00534 |
| Kern NWR | S-K-15C | 0.0338 | 0.0316 | 0.0158 | <0.00522 | 0.00844 | <0.00522 | 0.152 | <0.00522 |
| Kern NWR | S-K-15D | 0.0127 | <0.00523 | 0.00741 | <0.00523 | <0.00523 | 0.0328 | 0.114 | <0.00523 |
| Kern NWR | S-K-15E | 0.0116 | 0.04 | <0.00521 | <0.00521 | <0.00521 | 0.0179 | 0.337 | <0.00521 |

Appendix 3. cont.

| Location | Sample | PCB# 31 ng/g dw | PCB# 33/20 ng/g dw | PCB# 39 ng/g dw | PCB# 40 ng/g dw | PCB# 41/64 ng/g dw | PCB# 42/59/37 ng/g dw | PCB# 44 ng/g dw | PCB# 45 ng/g dw |
|---------------------|-----------|--------------------|-----------------------|--------------------|--------------------|-----------------------|-----------------------------|--------------------|--------------------|
| Imperial Valley J20 | S-IV-J20A | 0.525 | <0.00632 | <0.00632 | 0.0332 | 8.66 | 0.0511 | 0.155 | <0.00632 |
| Imperial Valley J20 | S-IV-J20B | 0.484 | 0.0239 | <0.00566 | 0.0352 | 5.01 | <0.00566 | 0.215 | 0.0602 |
| Imperial Valley J20 | S-IV-J20C | <0.00557 | 0.143 | <0.00557 | 0.105 | 6.28 | <0.00557 | 0.205 | 0.0496 |
| Imperial Valley J20 | S-IV-J20D | <0.00549 | 0.0365 | <0.00549 | 0.0299 | 9.21 | <0.00549 | 0.22 | 0.0532 |
| Imperial Valley J20 | S-IV-J20E | <0.00579 | 0.0551 | <0.00579 | 0.0598 | 11.3 | <0.00579 | 0.204 | 0.0516 |
| Imperial Valley J22 | S-IV-J22A | 0.63 | 0.0168 | 0.0906 | <0.00554 | 145 | <0.00554 | 0.15 | <0.00554 |
| Imperial Valley J22 | S-IV-J22B | 1.49 | 0.0353 | 0.0436 | 0.284 | 70.7 | <0.00582 | 0.266 | 0.0766 |
| Imperial Valley J22 | S-IV-J22C | 0.656 | 0.0252 | 0.12 | 0.0596 | 58.6 | <0.00568 | 0.249 | 0.055 |
| Imperial Valley J22 | S-IV-J22D | 0.383 | 0.0268 | 0.103 | <0.00579 | 47.8 | <0.00579 | 0.212 | 0.0653 |
| Imperial Valley J22 | S-IV-J22E | 0.137 | 0.0199 | 0.0541 | <0.00552 | 63 | 0.0155 | 0.218 | 0.0486 |
| Imperial Valley K18 | S-IV-K18A | 0.568 | 0.0187 | 0.0528 | <0.00548 | 43.7 | 0.0099 | 0.199 | 0.0374 |
| Imperial Valley K18 | S-IV-K18B | 0.537 | 0.0312 | <0.00551 | <0.00551 | 69.2 | 0.0245 | 0.178 | <0.00551 |
| Imperial Valley K18 | S-IV-K18C | 0.413 | 0.0304 | 0.0349 | <0.00561 | 72.9 | 0.306 | 0.13 | 0.0439 |
| Imperial Valley K18 | S-IV-K18D | 0.311 | 0.0258 | <0.00561 | <0.00561 | 36.4 | <0.00561 | 0.159 | 0.0494 |
| Imperial Valley K18 | S-IV-K18E | 0.445 | 0.028 | <0.00572 | <0.00572 | 38.6 | <0.00572 | 0.194 | 0.028 |
| Sonny Bono NWR | S-SB-420A | <0.00512 | 0.0136 | <0.00512 | 0.072 | 6.74 | <0.00512 | 0.181 | <0.00512 |
| Sonny Bono NWR | S-SB-420B | 0.0588 | <0.00514 | <0.00514 | 0.0134 | 4.72 | 0.0237 | 0.184 | <0.00514 |
| Sonny Bono NWR | S-SB-420C | 0.0742 | 0.0324 | <0.00518 | 0.024 | 4.02 | <0.00518 | 0.191 | 0.00522 |
| Sonny Bono NWR | S-SB-420D | <0.00518 | 0.0189 | <0.00518 | 0.0199 | 4.23 | <0.00518 | 0.173 | 0.0346 |
| Sonny Bono NWR | S-SB-420E | 0.0314 | 0.0168 | <0.00521 | 0.0314 | 5.25 | <0.00521 | 0.17 | 0.0335 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.0266 | 0.0482 | 0.0154 | 0.0533 | <0.00504 | 0.0174 | 0.292 | 0.0318 |
| Kern NWR | S-K-15B | 0.124 | 0.0194 | 0.0259 | <0.00534 | <0.00534 | 0.00863 | 0.277 | 0.0216 |
| Kern NWR | S-K-15C | 0.0443 | 0.0338 | 0.02 | 0.0264 | <0.00522 | 0.00633 | 0.276 | 0.0264 |
| Kern NWR | S-K-15D | 0.116 | 0.00847 | <0.00523 | 0.0127 | <0.00523 | 0.00529 | 0.201 | <0.00523 |
| Kern NWR | S-K-15E | 0.0327 | 0.0348 | <0.00521 | <0.00521 | <0.00521 | 0.0126 | 0.23 | 0.0348 |

Appendix 3. cont.

| Location | Sample | PCB# 46 ng/g dw | PCB# 47/75 ng/g dw | PCB# 48 ng/g dw | PCB# 49 ng/g dw | PCB# 52 ng/g dw | PCB# 53 ng/g dw | PCB# 60/56 ng/g dw | PCB# 63 ng/g dw |
|---------------------|-----------|--------------------|-----------------------|--------------------|--------------------|--------------------|--------------------|-----------------------|--------------------|
| Imperial Valley J20 | S-IV-J20A | 0.313 | 0.12 | <0.00632 | 0.0115 | 0.16 | <0.00632 | <0.00632 | <0.00632 |
| Imperial Valley J20 | S-IV-J20B | 0.255 | <0.00566 | <0.00566 | 0.1 | 0.292 | <0.00566 | <0.00566 | <0.00566 |
| Imperial Valley J20 | S-IV-J20C | 0.541 | 0.219 | <0.00557 | <0.00557 | 0.309 | <0.00557 | <0.00557 | <0.00557 |
| Imperial Valley J20 | S-IV-J20D | 0.466 | <0.00549 | <0.00549 | 0.0997 | 0.279 | <0.00549 | <0.00549 | <0.00549 |
| Imperial Valley J20 | S-IV-J20E | 0.986 | 0.109 | 0.0129 | <0.00579 | 0.369 | <0.00579 | <0.00579 | <0.00579 |
| Imperial Valley J22 | S-IV-J22A | 0.254 | <0.00554 | 0.00554 | 0.182 | 0.0492 | <0.00554 | 0.31 | <0.00554 |
| Imperial Valley J22 | S-IV-J22B | 0.816 | 0.0189 | 0.00582 | 0.231 | 0.551 | <0.00582 | 0.371 | <0.00582 |
| Imperial Valley J22 | S-IV-J22C | 0.717 | 0.0527 | 0.00568 | 0.197 | 0.433 | <0.00568 | 0.314 | <0.00568 |
| Imperial Valley J22 | S-IV-J22D | 0.178 | 0.0571 | 0.00579 | 0.177 | 0.282 | <0.00579 | 0.432 | <0.00579 |
| Imperial Valley J22 | S-IV-J22E | 0.0873 | <0.00552 | 0.00552 | 0.0332 | 0.0729 | <0.00552 | <0.00552 | <0.00552 |
| Imperial Valley K18 | S-IV-K18A | 0.0594 | <0.00548 | 0.00548 | 0.169 | 0.0506 | <0.00548 | <0.00548 | <0.00548 |
| Imperial Valley K18 | S-IV-K18B | 0.117 | 0.0178 | <0.00551 | 0.0758 | 0.162 | <0.00551 | <0.00551 | <0.00551 |
| Imperial Valley K18 | S-IV-K18C | 0.14 | <0.00561 | <0.00561 | 0.0394 | 0.0372 | <0.00561 | <0.00561 | <0.00561 |
| Imperial Valley K18 | S-IV-K18D | 0.0673 | <0.00561 | <0.00561 | 0.0191 | 0.0348 | <0.00561 | <0.00561 | <0.00561 |
| Imperial Valley K18 | S-IV-K18E | 0.0875 | 0.0338 | <0.00572 | 0.0152 | 0.042 | <0.00572 | <0.00572 | <0.00572 |
| Sonny Bono NWR | S-SB-420A | 0.122 | <0.00512 | <0.00512 | 0.0125 | 0.0469 | <0.00512 | <0.00512 | <0.00512 |
| Sonny Bono NWR | S-SB-420B | 0.0557 | <0.00514 | <0.00514 | 0.0165 | 0.115 | <0.00514 | <0.00514 | <0.00514 |
| Sonny Bono NWR | S-SB-420C | 0.101 | 0.00522 | <0.00518 | 0.0125 | 0.0418 | <0.00518 | <0.00518 | 0.00836 |
| Sonny Bono NWR | S-SB-420D | 0.0934 | 0.00735 | <0.00518 | 0.0168 | 0.0346 | <0.00518 | <0.00518 | 0.00839 |
| Sonny Bono NWR | S-SB-420E | 0.0702 | 0.0178 | <0.00521 | 0.022 | 0.043 | <0.00521 | <0.00521 | 0.00629 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.0164 | 0.118 | 0.121 | 0.0338 | 0.04 | <0.00504 | 0.127 | 0.0307 |
| Kern NWR | S-K-15B | 0.0151 | 0.0281 | <0.00534 | 0.0173 | 0.0723 | <0.00534 | 0.134 | 0.00863 |
| Kern NWR | S-K-15C | 0.00633 | 0.0232 | 0.0401 | 0.0169 | 0.0179 | <0.00522 | <0.00522 | 0.0148 |
| Kern NWR | S-K-15D | 0.0106 | 0.0222 | 0.0328 | 0.0254 | 0.00741 | 0.0138 | 0.0286 | <0.00523 |
| Kern NWR | S-K-15E | 0.00527 | 0.0316 | <0.00521 | 0.0274 | 0.00948 | <0.00521 | 0.04 | <0.00521 |

Appendix 3. cont.

| Location | Sample | PCB# 66 ng/g dw | PCB# 67 ng/g dw | PCB# 7/9 ng/g dw | PCB# 70 ng/g dw | PCB# 72 ng/g dw | PCB# 74/61 ng/g dw | PCB# 8/5 ng/g dw | PCB# 82 ng/g dw |
|---------------------|-----------|--------------------|--------------------|---------------------|--------------------|--------------------|-----------------------|---------------------|--------------------|
| Imperial Valley J20 | S-IV-J20A | 0.0677 | 0.0613 | <0.00632 | 0.105 | 0.752 | 0.0358 | 0.0128 | 1.77 |
| Imperial Valley J20 | S-IV-J20B | 0.126 | 0.293 | <0.00566 | 0.18 | 0.928 | 0.05 | 0.233 | 2.48 |
| Imperial Valley J20 | S-IV-J20C | 0.331 | 0.493 | <0.00557 | 0.374 | 1.04 | 0.159 | 0.177 | 3.69 |
| Imperial Valley J20 | S-IV-J20D | 0.255 | 0.279 | <0.00549 | 0.235 | 0.96 | 0.0709 | 0.295 | 2.45 |
| Imperial Valley J20 | S-IV-J20E | <0.00579 | 0.182 | <0.00579 | 0.318 | 1.41 | 0.0903 | 0.281 | 2.92 |
| Imperial Valley J22 | S-IV-J22A | 0.3 | 0.276 | 0.123 | 0.152 | 0.978 | 0.116 | 0.336 | 1.88 |
| Imperial Valley J22 | S-IV-J22B | 0.37 | 0.391 | <0.00582 | 0.29 | 1.52 | 0.233 | 0.355 | 3.57 |
| Imperial Valley J22 | S-IV-J22C | 0.43 | 0.521 | <0.00568 | 0.299 | 1.49 | 0.235 | 0.389 | 3.62 |
| Imperial Valley J22 | S-IV-J22D | 0.307 | 0.284 | <0.00579 | 0.119 | 0.674 | 0.0851 | 0.302 | 1.67 |
| Imperial Valley J22 | S-IV-J22E | 0.169 | 0.032 | <0.00552 | 0.11 | 0.791 | 0.0409 | 0.197 | 0.93 |
| Imperial Valley K18 | S-IV-K18A | 0.0583 | <0.00548 | <0.00548 | 0.0968 | 0.294 | 0.055 | 0.239 | 0.437 |
| Imperial Valley K18 | S-IV-K18B | 0.0903 | 0.0156 | <0.00551 | 0.101 | 0.372 | 0.0424 | 0.243 | 0.314 |
| Imperial Valley K18 | S-IV-K18C | 0.0957 | 0.0146 | <0.00561 | 0.106 | 0.37 | 0.0394 | 0.253 | 0.309 |
| Imperial Valley K18 | S-IV-K18D | 0.204 | <0.00561 | <0.00561 | 0.00673 | 0.323 | 0.0483 | 0.221 | 0.276 |
| Imperial Valley K18 | S-IV-K18E | 0.0688 | <0.00572 | <0.00572 | 0.0187 | 0.63 | 0.049 | 0.215 | 0.386 |
| Sonny Bono NWR | S-SB-420A | 0.0605 | <0.00512 | <0.00512 | 0.0156 | 0.575 | 0.0949 | 0.234 | 0.791 |
| Sonny Bono NWR | S-SB-420B | 0.114 | <0.00514 | <0.00514 | 0.031 | 0.709 | 0.109 | 0.209 | 0.771 |
| Sonny Bono NWR | S-SB-420C | 0.0648 | 0.00731 | <0.00518 | 0.0199 | 0.702 | 0.0711 | 0.107 | 0.771 |
| Sonny Bono NWR | S-SB-420D | 0.0682 | 0.0063 | <0.00518 | 0.0315 | 0.426 | 0.0787 | 0.084 | 0.573 |
| Sonny Bono NWR | S-SB-420E | 0.0786 | 0.00839 | <0.00521 | 0.021 | 0.527 | 0.0493 | 0.108 | 0.601 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.0984 | 0.0656 | <0.00504 | 0.0891 | 0.133 | 0.0686 | 0.0994 | 0.75 |
| Kern NWR | S-K-15B | 0.0561 | 0.0647 | <0.00534 | 0.0485 | 0.166 | 0.159 | 0.0906 | 0.29 |
| Kern NWR | S-K-15C | 0.0454 | 0.0443 | <0.00522 | 0.0327 | 0.0464 | 0.0422 | 0.096 | 0.543 |
| Kern NWR | S-K-15D | 0.0243 | 0.0529 | <0.00523 | 0.019 | <0.00523 | 0.00952 | 0.0656 | 0.462 |
| Kern NWR | S-K-15E | 0.019 | 0.0432 | <0.00521 | 0.0337 | 0.00843 | 0.058 | 0.0527 | 0.322 |

Appendix 3. cont.

| Location | Sample | PCB# 83 ng/g dw | PCB# 84 ng/g dw | PCB# 85 ng/g dw | PCB# 87/115 ng/g dw | PCB# 92 ng/g dw | PCB# 95/80 ng/g dw | PCB# 97 ng/g dw | PCB# 99 ng/g dw |
|---------------------|-----------|--------------------|--------------------|--------------------|------------------------|--------------------|-----------------------|--------------------|--------------------|
| Imperial Valley J20 | S-IV-J20A | 0.0524 | 1.6 | 0.0358 | 0.158 | 0.926 | 0.221 | 0.718 | 0.105 |
| Imperial Valley J20 | S-IV-J20B | 0.177 | 1.8 | 0.0955 | 0.201 | <0.00566 | <0.00566 | 0.984 | 0.324 |
| Imperial Valley J20 | S-IV-J20C | 0.156 | 2.76 | 0.0676 | 0.295 | 2.18 | <0.00557 | 1.46 | 0.446 |
| Imperial Valley J20 | S-IV-J20D | 0.196 | 1.73 | 0.062 | 0.212 | 1.55 | 0.529 | 1.01 | 0.401 |
| Imperial Valley J20 | S-IV-J20E | 0.25 | 1.69 | 0.0574 | 0.26 | 2.9 | <0.00579 | 1.28 | 0.428 |
| Imperial Valley J22 | S-IV-J22A | 0.00783 | 1.2 | 0.122 | 0.202 | 0.508 | 0.34 | 1.51 | 0.668 |
| Imperial Valley J22 | S-IV-J22B | 0.119 | 1.95 | 0.105 | 0.305 | 2.24 | 0.649 | 2.11 | 0.371 |
| Imperial Valley J22 | S-IV-J22C | 0.136 | 2.28 | 0.147 | 0.292 | 2.2 | 0.536 | 1.97 | 0.409 |
| Imperial Valley J22 | S-IV-J22D | 0.0373 | 0.904 | 0.0758 | 0.154 | 0.781 | 0.558 | 0.881 | 0.0688 |
| Imperial Valley J22 | S-IV-J22E | 0.074 | 0.256 | 0.0995 | 0.191 | <0.00552 | <0.00552 | 0.681 | 0.0751 |
| Imperial Valley K18 | S-IV-K18A | 0.0429 | 0.0528 | 0.0176 | 0.0858 | 0.244 | 0.12 | 0.372 | 0.0847 |
| Imperial Valley K18 | S-IV-K18B | <0.00551 | 0.0814 | 0.039 | 0.0424 | <0.00551 | <0.00551 | 0.144 | 0.0279 |
| Imperial Valley K18 | S-IV-K18C | 0.018 | 0.054 | <0.00561 | 0.0439 | <0.00561 | <0.00561 | 0.0856 | 0.027 |
| Imperial Valley K18 | S-IV-K18D | 0.0337 | 0.055 | 0.00786 | 0.0505 | 0.242 | <0.00561 | 0.119 | 0.0438 |
| Imperial Valley K18 | S-IV-K18E | 0.0292 | 0.0502 | 0.00817 | 0.0665 | 0.123 | <0.00572 | 0.179 | 0.0315 |
| Sonny Bono NWR | S-SB-420A | 0.049 | <0.00512 | 0.00938 | 0.0688 | 1.16 | <0.00512 | 0.332 | 0.0772 |
| Sonny Bono NWR | S-SB-420B | 0.0495 | <0.00514 | 0.00929 | 0.0867 | 1.06 | 0.0836 | 0.287 | 0.0846 |
| Sonny Bono NWR | S-SB-420C | 0.047 | <0.00518 | 0.00627 | 0.0711 | 1.2 | <0.00518 | 0.304 | <0.00518 |
| Sonny Bono NWR | S-SB-420D | 0.0346 | 0.0818 | <0.00518 | 0.0556 | 0.977 | <0.00518 | 0.242 | 0.0504 |
| Sonny Bono NWR | S-SB-420E | 0.0377 | 0.102 | 0.00629 | 0.0566 | 1.01 | <0.00521 | 0.245 | 0.0451 |
| | | | | | | | | | |
| Kern NWR | S-K-15A | 0.193 | 0.409 | 0.0256 | 0.154 | <0.00504 | 0.0533 | <0.00504 | 0.0471 |
| Kern NWR | S-K-15B | 0.12 | 0.184 | 0.0281 | 0.0367 | <0.00534 | 0.0281 | <0.00534 | 0.0259 |
| Kern NWR | S-K-15C | 0.141 | 0.258 | 0.02 | 0.149 | <0.00522 | 0.039 | <0.00522 | 0.0802 |
| Kern NWR | S-K-15D | 0.0529 | 0.135 | 0.00847 | 0.101 | 0.054 | 0.0148 | <0.00523 | 0.0381 |
| Kern NWR | S-K-15E | 0.0379 | 0.227 | 0.00948 | 0.0337 | <0.00521 | 0.0158 | <0.00521 | <0.00521 |

Appendix 3. cont.

| Location | Sample | PCB-1268 ng/g dw | PCB- TOTAL ng/g dw |
|---------------------|---------------|-----------------------------|-----------------------------------|
| Imperial Valley J20 | S-IV-J20A | 27.4 | 27.3 |
| Imperial Valley J20 | S-IV-J20B | 35.4 | 35.3 |
| Imperial Valley J20 | S-IV-J20C | 53 | 53 |
| Imperial Valley J20 | S-IV-J20D | 39.5 | 39.5 |
| Imperial Valley J20 | S-IV-J20E | 55.7 | 55.8 |
| Imperial Valley J22 | S-IV-J22A | 29.9 | 30 |
| Imperial Valley J22 | S-IV-J22B | 59.4 | 59.4 |
| Imperial Valley J22 | S-IV-J22C | 57.1 | 57.2 |
| Imperial Valley J22 | S-IV-J22D | 26.4 | 26.5 |
| Imperial Valley J22 | S-IV-J22E | 14.5 | 14.5 |
| Imperial Valley K18 | S-IV-K18A | 12.5 | 12.4 |
| Imperial Valley K18 | S-IV-K18B | 15.2 | 15.3 |
| Imperial Valley K18 | S-IV-K18C | 12.2 | 12.2 |
| Imperial Valley K18 | S-IV-K18D | 8.87 | 8.87 |
| Imperial Valley K18 | S-IV-K18E | 11.4 | 11.4 |
| Sonny Bono NWR | S-SB-420A | 13.6 | 13.7 |
| Sonny Bono NWR | S-SB-420B | 13.6 | 13.6 |
| Sonny Bono NWR | S-SB-420C | 13.2 | 13.2 |
| Sonny Bono NWR | S-SB-420D | 11.1 | 11.1 |
| Sonny Bono NWR | S-SB-420E | 10.9 | 10.9 |
| | | | |
| Kern NWR | S-K-15A | 8.88 | 8.91 |
| Kern NWR | S-K-15B | 6.89 | 6.9 |
| Kern NWR | S-K-15C | 6.76 | 6.75 |
| Kern NWR | S-K-15D | 4.76 | 4.76 |
| Kern NWR | S-K-15E | 5.99 | 6.01 |

* C11-PCB, C110-PCB, C12-PCB, PCB#114, PCB#126, PCB#146, PCB#169, PCB#69, PCB#77, PCB #81, PCB-1242, PCB-1248, PCB#1254, and PCB 1260 not detected in any soil samples.

Appendix 4. Concentrations of inorganics* (ug/g dry weight) in terrestrial invertebrate samples collected from mountain plover wintering sites in California, 2006.

| Location | Sample | Moisture % | Al ug/g dw | As ug/g dw | B ug/g dw | Ba ug/g dw | Be ug/g dw | Cd ug/g dw | Cr ug/g dw | Cu ug/g dw |
|---------------------|-----------|------------|------------|------------|-----------|------------|------------|------------|------------|------------|
| Imperial Valley J20 | I-IV-J20A | 54.8 | 190 | <0.3 | 20 | 1.8 | <0.1 | 0.2 | <0.5 | 25 |
| Imperial Valley J20 | I-IV-J20B | 56.2 | 3120 | 1.4 | 10 | 47.8 | <0.1 | 0.84 | 3.5 | 94.9 |
| Imperial Valley J20 | I-IV-J20C | 57 | 480 | <0.8 | <20 | 5.6 | <0.1 | 0.6 | <0.7 | 57 |
| Imperial Valley J20 | I-IV-J20D | 59.3 | 1010 | 0.4 | <10 | 10 | <0.1 | 0.5 | 1 | 62.8 |
| Imperial Valley J20 | I-IV-J20E | 42.3 | 150 | <0.7 | <20 | 2.4 | <0.1 | 0.5 | <0.6 | 42 |
| Imperial Valley J22 | I-IV-J22A | 61.6 | 1810 | 1 | <10 | 18 | <0.1 | 0.82 | 2.2 | 67.7 |
| Imperial Valley J22 | I-IV-J22B | 65.3 | 603 | 2.1 | <10 | 7.1 | <0.1 | 0.9 | 1 | 67 |
| Imperial Valley J22 | I-IV-J22C | 68 | 343 | 0.91 | <7 | 4.8 | <0.1 | 0.45 | 0.6 | 69.8 |
| Imperial Valley J22 | I-IV-J22D | 65.4 | 851 | <0.6 | 37 | 14 | <0.1 | 0.8 | 1.7 | 208 |
| Imperial Valley J22 | I-IV-J22E | 59.6 | 1020 | 0.81 | 10 | 12 | <0.1 | 0.45 | 1.7 | 32.9 |
| Imperial Valley K18 | I-IV-K18A | 69.5 | 1220 | 0.75 | 6 | 10 | <0.1 | 0.5 | 1.5 | 48.6 |
| Imperial Valley K18 | I-IV-K18B | 70.6 | 1540 | 0.5 | <7 | 14 | <0.1 | 1.1 | 2.8 | 78.8 |
| Imperial Valley K18 | I-IV-K18C | 68 | 1250 | 0.3 | <7 | 13 | <0.1 | 1 | 2.8 | 50.3 |
| Imperial Valley K18 | I-IV-K18D | 63.8 | 1190 | <0.5 | <10 | 12 | <0.1 | 0.65 | 2.4 | 38 |
| Imperial Valley K18 | I-IV-K18E | 64.4 | 1520 | 0.3 | <7 | 15 | <0.1 | 0.68 | 2.5 | 26 |
| Sonny Bono NWR | I-SB-420A | 65.1 | 2200 | 0.8 | <7 | 19 | <0.1 | <0.1 | 1.6 | 24 |
| Sonny Bono NWR | I-SB-420B | 70.8 | 1810 | 0.6 | <4 | 13 | <0.1 | 0.51 | 1 | 49.3 |
| Sonny Bono NWR | I-SB-420C | 65.8 | 2270 | 0.96 | 10 | 16 | <0.1 | 0.43 | 1.8 | 32 |
| Sonny Bono NWR | I-SB-420D | 68.4 | 4970 | 1.1 | <8 | 33.8 | <0.1 | 0.72 | 3.7 | 37 |
| Sonny Bono NWR | I-SB-420E | 54.1 | 24500 | 4.5 | 18 | 137 | 0.56 | 1.2 | 19 | 63.7 |
| | | | | | | | | | | |
| Kern NWR | I-K-15A | 62.4 | 1230 | 0.88 | <6 | 68.7 | <0.1 | 1.2 | 1.7 | 102 |
| Kern NWR | I-K-15B | 53.7 | 1970 | 0.5 | 8 | 41.9 | <0.1 | 1 | 2.2 | 51.3 |
| Kern NWR | I-K-15C | 59.6 | 855 | 2.3 | 10 | 22.7 | <0.1 | 0.6 | 1 | 31 |
| Kern NWR | I-K-15D | 55.2 | 2010 | <0.6 | <20 | 16 | <0.1 | 0.97 | 2.2 | 59 |
| Kern NWR | I-K-15E | 54.4 | 1860 | 0.5 | 10 | 50.4 | <0.1 | 0.81 | 2.7 | 56 |

Appendix 4. cont.

| Location | Sample | Fe ug/g dw | Mg ug/g dw | Mn ug/g dw | Mo ug/g dw | Ni ug/g dw | Pb ug/g dw | Se ug/g dw | Sr ug/g dw | V ug/g dw | Zn ug/g dw |
|---------------------|-----------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--------------|---------------|
| Imperial Valley J20 | I-IV-J20A | 170 | 1270 | 29 | 3 | 0.6 | <0.8 | 1.4 | 9.8 | 0.8 | 194 |
| Imperial Valley J20 | I-IV-J20B | 2160 | 3300 | 68 | <2 | 2.4 | 7.2 | 3 | 301 | 6.1 | 592 |
| Imperial Valley J20 | I-IV-J20C | 374 | 1720 | 22 | <2 | 0.9 | <2 | 2.8 | 15 | 1 | 468 |
| Imperial Valley J20 | I-IV-J20D | 686 | 1930 | 35 | <2 | 1 | <1 | 2 | 20.8 | 1.8 | 409 |
| Imperial Valley J20 | I-IV-J20E | 160 | 1360 | 13 | <2 | 1 | <2 | 2.7 | 9 | <0.5 | 401 |
| Imperial Valley J22 | I-IV-J22A | 1220 | 2230 | 49 | <2 | 1.8 | <1 | 2.2 | 32.7 | 3.3 | 640 |
| Imperial Valley J22 | I-IV-J22B | 458 | 2060 | 42 | <2 | 1.5 | <1 | 3 | 23.2 | 1 | 446 |
| Imperial Valley J22 | I-IV-J22C | 298 | 1850 | 25 | <2 | <0.5 | <0.7 | 3.4 | 20 | 0.7 | 491 |
| Imperial Valley J22 | I-IV-J22D | 673 | 1770 | 40 | 6 | 1.5 | <1 | 4.4 | 29 | 3.2 | 747 |
| Imperial Valley J22 | I-IV-J22E | 745 | 1400 | 27 | <2 | 0.9 | <0.5 | 1.6 | 23.2 | 2.2 | 427 |
| Imperial Valley K18 | I-IV-K18A | 772 | 1860 | 50 | <2 | 1 | <0.6 | 1.3 | 24 | 2.2 | 420 |
| Imperial Valley K18 | I-IV-K18B | 922 | 2280 | 49 | <2 | 1 | 2 | 1.5 | 28 | 3.2 | 606 |
| Imperial Valley K18 | I-IV-K18C | 822 | 2000 | 49 | <2 | <0.5 | <0.7 | 1.1 | 24 | 2.8 | 359 |
| Imperial Valley K18 | I-IV-K18D | 755 | 1720 | 36 | <2 | <0.5 | <1 | 1.6 | 22.3 | 2.3 | 368 |
| Imperial Valley K18 | I-IV-K18E | 891 | 1700 | 35 | <2 | 0.5 | <0.7 | 0.93 | 38.8 | 2.9 | 273 |
| Sonny Bono NWR | I-SB-420A | 1700 | 2090 | 48 | <2 | 1.9 | 1 | 0.78 | 27.3 | 4.7 | 171 |
| Sonny Bono NWR | I-SB-420B | 1260 | 2280 | 46 | 2 | 1.7 | 1 | 1 | 27 | 3.4 | 215 |
| Sonny Bono NWR | I-SB-420C | 1580 | 2230 | 51.8 | <2 | 1.5 | 1 | 1.3 | 30.1 | 4.4 | 206 |
| Sonny Bono NWR | I-SB-420D | 3240 | 4040 | 86.9 | 2 | 3.3 | 2.6 | 1.1 | 46.6 | 8.9 | 176 |
| Sonny Bono NWR | I-SB-420E | 12300 | 7880 | 226 | <2 | 11 | 11 | 2.8 | 298 | 36 | 138 |
| | | | | | | | | | | | |
| Kern NWR | I-K-15A | 1160 | 2630 | 34 | <2 | 2.4 | <0.6 | 0.63 | 413 | 3.3 | 160 |
| Kern NWR | I-K-15B | 1970 | 2750 | 64.2 | 2 | 2.7 | <0.8 | 0.5 | 259 | 5.1 | 140 |
| Kern NWR | I-K-15C | 909 | 2330 | 36 | <2 | 2.2 | <0.8 | 0.5 | 127 | 2.7 | 118 |
| Kern NWR | I-K-15D | 1950 | 2300 | 55.5 | <2 | 2.6 | <2 | 0.5 | 43.4 | 5.4 | 164 |
| Kern NWR | I-K-15E | 1840 | 2430 | 59.9 | <2 | 2.6 | 1 | 0.91 | 217 | 4.5 | 213 |

*Mercury not detected in any of the terrestrial invertebrate samples.

Appendix 5. Concentrations of organic pesticides* and trifluralin (ng/g dry weight) in terrestrial invertebrate samples collected from mountain plover wintering sites in California, 2006.

| Location | Sample | Moisture % | beta BHC ng/g dw | dieldrin ng/g dw | endosulfan II ng/g dw | endrin ng/g dw | gamma chlordane ng/g dw | HCB ug/g dw |
|---------------------|-------------|------------|------------------|------------------|-----------------------|----------------|-------------------------|-------------|
| Imperial Valley J20 | I-IV-J20A | 58.1 | <8.52 | <8.52 | <8.52 | <8.52 | <8.52 | <8.52 |
| Imperial Valley J20 | I-IV-J20B | 58.5 | <8.03 | 23.4 | 30.5 | 126 | <8.03 | <8.03 |
| Imperial Valley J20 | I-IV-J20C | 56.8 | <16.5 | <16.5 | <16.5 | <16.5 | <16.5 | <16.5 |
| Imperial Valley J20 | I-IV-J20D | 61.8 | <16.4 | <16.4 | <16.4 | <16.4 | <16.4 | <16.4 |
| Imperial Valley J20 | I-IV-J20E | 57.4 | <23.5 | <23.5 | <23.5 | <23.5 | <23.5 | <23.5 |
| Imperial Valley J22 | I-IV-J22A | 62.5 | <16.7 | <16.7 | <16.7 | <16.7 | <16.7 | <16.7 |
| Imperial Valley J22 | I-IV-J22B | 68.5 | <11.4 | <11.4 | <11.4 | <11.4 | <11.4 | <11.4 |
| Imperial Valley J22 | I-IV-J22C | 64.5 | <5.22 | <5.22 | <5.22 | <5.22 | <5.22 | 7.91 |
| Imperial Valley J22 | I-IV-J22D | 69.5 | <10.9 | <10.9 | <10.9 | <10.9 | <10.9 | <10.9 |
| Imperial Valley J22 | I-IV-J22E | 55.6 | <10.2 | 28.1 | <10.2 | 31.5 | <10.2 | <10.2 |
| Imperial Valley K18 | I-IV-K18A** | 71.6 | <4.29 | 32.8 | 11.1 | 21.1 | <4.29 | <4.29 |
| Imperial Valley K18 | I-IV-K18B** | 68.8 | <3.2 | <3.2 | <3.2 | 7.13 | 3.56 | <3.2 |
| Imperial Valley K18 | I-IV-K18C | 69.2 | <5.8 | <5.8 | <5.8 | <5.8 | <5.8 | <5.8 |
| Imperial Valley K18 | I-IV-K18D | 63.5 | 4.22 | <4.03 | <4.03 | 9.6 | 6.84 | <4.03 |
| Imperial Valley K18 | I-IV-K18E** | 61.3 | <3.01 | <3.01 | <3.01 | 11.7 | <3.01 | <3.01 |
| Sonny Bono NWR | I-SB-420A** | 66.3 | <3.71 | <3.71 | <3.71 | <3.71 | <3.71 | <3.71 |
| Sonny Bono NWR | I-SB-420B | 70.4 | <7.03 | <7.03 | <7.03 | <7.03 | <7.03 | <7.03 |
| Sonny Bono NWR | I-SB-420C | 64.9 | 7.86 | <5.7 | <5.7 | 8.65 | <5.7 | <5.7 |
| Sonny Bono NWR | I-SB-420D | 69.9 | <4.37 | <4.37 | <4.37 | 4.75 | <4.37 | <4.37 |
| Sonny Bono NWR | I-SB-420E** | 62.5 | <4.17 | <4.17 | <4.17 | 29.7 | 6.32 | <4.17 |
| | | | | | | | | |
| Kern NWR | I-K-15A | 62.4 | <10.2 | <10.2 | 98 | 1,020 | <10.2 | <10.2 |
| Kern NWR | I-K-15B | 52.5 | <21 | <21 | <21 | <21 | <21 | <21 |
| Kern NWR | I-K-15C | 59.7 | <6.2 | <6.2 | <6.2 | <6.2 | <6.2 | <6.2 |
| Kern NWR | I-K-15D | 55.6 | <11.3 | <11.3 | <11.3 | <11.3 | <11.3 | <11.3 |
| Kern NWR | I-K-15E | 52.3 | <8.06 | <8.06 | <8.06 | <8.06 | <8.06 | <8.06 |

Appendix 5. cont.

| Location | Sample | oxychlorane ng/g dw | p,p'-DDE ng/g dw | p,p'-DDT ng/g dw | 1,2,4,5- Tetrachlorobenzene ng/g dw | Trifluralin ng/g dw |
|---------------------|-----------|------------------------|---------------------|---------------------|---|------------------------|
| Imperial Valley J20 | I-IV-J20A | <8.52 | 137 | <8.52 | 25.9 | 256 |
| Imperial Valley J20 | I-IV-J20B | 11.6 | 283 | <8.03 | 90.9 | 82.5 |
| Imperial Valley J20 | I-IV-J20C | <16.5 | 312 | <16.5 | 42.1 | 58.4 |
| Imperial Valley J20 | I-IV-J20D | <16.4 | 401 | <16.4 | <16.4 | <32 |
| Imperial Valley J20 | I-IV-J20E | <23.5 | 478 | <23.5 | 123 | 78.6 |
| Imperial Valley J22 | I-IV-J22A | <16.7 | 222 | <16.7 | <16.7 | 49.3 |
| Imperial Valley J22 | I-IV-J22B | <11.4 | 342 | <11.4 | <11.4 | <22.5 |
| Imperial Valley J22 | I-IV-J22C | <5.22 | 547 | <5.22 | 50.9 | 32.6 |
| Imperial Valley J22 | I-IV-J22D | <10.9 | 1,310 | <10.9 | <10.9 | 252 |
| Imperial Valley J22 | I-IV-J22E | <10.2 | 1,240 | <10.2 | <10.2 | 71.8 |
| Imperial Valley K18 | I-IV-K18A | <4.29 | 56.6 | <4.29 | <4.29 | 51.3 |
| Imperial Valley K18 | I-IV-K18B | <3.2 | 54.4 | <3.2 | 5.43 | 49.7 |
| Imperial Valley K18 | I-IV-K18C | <5.8 | 47.8 | <5.8 | 18.8 | 52.4 |
| Imperial Valley K18 | I-IV-K18D | <4.03 | 310 | <4.03 | 29.9 | 35.3 |
| Imperial Valley K18 | I-IV-K18E | <3.01 | 130 | <3.01 | <3.01 | 157 |
| Sonny Bono NWR | I-SB-420A | <3.71 | 142 | <3.71 | <3.71 | 9.13 |
| Sonny Bono NWR | I-SB-420B | <7.03 | 225 | <7.03 | 21.4 | <14.4 |
| Sonny Bono NWR | I-SB-420C | <5.7 | 259 | <5.7 | 27.9 | <11.3 |
| Sonny Bono NWR | I-SB-420D | <4.37 | 358 | 10.9 | <4.37 | <8.78 |
| Sonny Bono NWR | I-SB-420E | <4.17 | 547 | 19.5 | <4.17 | <8.33 |
| | | | | | | |
| Kern NWR | I-K-15A | <10.2 | 14.4 | <10.2 | 21.4 | <20.9 |
| Kern NWR | I-K-15B | <21 | <21 | <21 | <21 | <44.8 |
| Kern NWR | I-K-15C | <6.2 | <6.2 | <6.2 | 8.93 | <12.2 |
| Kern NWR | I-K-15D | <11.3 | <11.3 | <11.3 | 14.7 | <22.3 |
| Kern NWR | I-K-15E | <8.06 | 17.2 | <8.06 | <8.06 | <16.1 |

Appendix 5. cont.

*Aldrin, alpha BHC, alpha chlordane, chlorpyrifos, cis-nonachlor, delta BHC, gamma BHC, heptachlor, heptachlor epoxide, mirex, o,p'-DDD, o,p'-DDE, o,p'-DDT, p,p'-DDD, pentachloro- anisole, toxaphene, trans-nonachlor, and 1,2,3,4-tetrachlorobenzene were analyzed for in all samples but were not detected.

** The following organophosphates aldicarb, azinphos-methyl, barban, bendiocarb, carbaryl, carbofuran, coumaphos demeton, diazinon, dichlorvos/naled, dimethoate, disulfoton, EPN, ethoprop, fensulfothion, fenthion, malathion, merphos/tribufos, methiocarb, methomyl, methyl parathion, mevinphos, monocrotophos, naled, oxamyl, parathion, phorate, propham, protothiofos, ronnel, sulfotepp, sulprofos, tetrachlorvinphos, tetraethylpyrophosphate, trichloronate were analyzed for in these five terrestrial invertebrate samples but were not detected.

Appendix 6. Concentrations of PCBs* (ng/g dry weight) in terrestrial invertebrate samples collected from mountain plover wintering sites in California, 2006.

| Location | Sample | Moisture % | C14-PCB ng/g dw | C15-PCB ng/g dw | C16-PCB ng/g dw | PCB# 1 ng/g dw | PCB# 105 ng/g dw | PCB# 107 ng/g dw |
|---------------------|-----------|------------|-----------------|-----------------|-----------------|----------------|------------------|------------------|
| Imperial Valley J20 | I-IV-J20A | 58.1 | <170 | <170 | <170 | <1.7 | <1.7 | <1.7 |
| Imperial Valley J20 | I-IV-J20B | 58.5 | 314 | 306 | <161 | <1.61 | <1.61 | 36.8 |
| Imperial Valley J20 | I-IV-J20C | 56.8 | <331 | <331 | <331 | <3.31 | <3.31 | <3.31 |
| Imperial Valley J20 | I-IV-J20D | 61.8 | <328 | <328 | <328 | <3.28 | <3.28 | <3.28 |
| Imperial Valley J20 | I-IV-J20E | 57.4 | <469 | <469 | <469 | <4.69 | <4.69 | <4.69 |
| Imperial Valley J22 | I-IV-J22A | 62.5 | <333 | <333 | <333 | <3.33 | <3.33 | <3.33 |
| Imperial Valley J22 | I-IV-J22B | 68.5 | <227 | <227 | <227 | <2.27 | <2.27 | <2.27 |
| Imperial Valley J22 | I-IV-J22C | 64.5 | <104 | <104 | <104 | <1.04 | <1.04 | <1.04 |
| Imperial Valley J22 | I-IV-J22D | 69.5 | <218 | <218 | <218 | <2.18 | <2.18 | <2.18 |
| Imperial Valley J22 | I-IV-J22E | 55.6 | <205 | <205 | <205 | <2.05 | <2.05 | <2.05 |
| Imperial Valley K18 | I-IV-K18A | 71.6 | <85.8 | <85.8 | <85.8 | <0.858 | <0.858 | <0.858 |
| Imperial Valley K18 | I-IV-K18B | 68.8 | <64 | <64 | <64 | <0.64 | <0.64 | <0.64 |
| Imperial Valley K18 | I-IV-K18C | 69.2 | <116 | <116 | <116 | <1.16 | <1.16 | <1.16 |
| Imperial Valley K18 | I-IV-K18D | 63.5 | <80.6 | <80.6 | <80.6 | <0.806 | <0.806 | <0.806 |
| Imperial Valley K18 | I-IV-K18E | 61.3 | <60.1 | <60.1 | <60.1 | 14.4 | <0.601 | <0.601 |
| Sonny Bono NWR | I-SB-420A | 66.3 | <74.1 | <74.1 | <74.1 | <0.741 | <0.741 | <0.741 |
| Sonny Bono NWR | I-SB-420B | 70.4 | <141 | <141 | <141 | <1.41 | <1.41 | <1.41 |
| Sonny Bono NWR | I-SB-420C | 64.9 | <114 | <114 | <114 | <1.14 | <1.14 | <1.14 |
| Sonny Bono NWR | I-SB-420D | 69.9 | <87.3 | <87.3 | <87.3 | <0.873 | <0.873 | <0.873 |
| Sonny Bono NWR | I-SB-420E | 62.5 | <83.3 | <83.3 | <83.3 | <0.833 | 5.57 | <0.833 |
| Kern NWR | I-K-15A | 62.4 | <204 | <204 | 301 | <2.04 | <2.04 | <2.04 |
| Kern NWR | I-K-15B | 52.5 | <421 | <421 | <421 | <4.21 | <4.21 | <4.21 |
| Kern NWR | I-K-15C | 59.7 | <124 | <124 | <124 | <1.24 | <1.24 | <1.24 |
| Kern NWR | I-K-15D | 55.6 | <225 | <225 | <225 | <2.25 | <2.25 | <2.25 |
| Kern NWR | I-K-15E | 52.3 | <161 | <161 | <161 | <1.61 | <1.61 | <1.61 |

Appendix 6. cont.

| Location | Sample | PCB# 119 ng/g dw | PCB# 135 ng/g dw | PCB# 138/160 ng/g dw | PCB# 141/179 ng/g dw | PCB# 149/123 ng/g dw | PCB# 151 ng/g dw | PCB# 172 ng/g dw |
|---------------------|-----------|---------------------|---------------------|----------------------------|----------------------------|----------------------------|---------------------|---------------------|
| Imperial Valley J20 | I-IV-J20A | <1.7 | <1.7 | 13 | <1.7 | <1.7 | <1.7 | <1.7 |
| Imperial Valley J20 | I-IV-J20B | <1.61 | 24.5 | 8.62 | 31.2 | 3.84 | 43.9 | 2.26 |
| Imperial Valley J20 | I-IV-J20C | <3.31 | <3.31 | 20.1 | <3.31 | <3.31 | <3.31 | 8.55 |
| Imperial Valley J20 | I-IV-J20D | <3.28 | <3.28 | 18.3 | <3.28 | <3.28 | <3.28 | 12 |
| Imperial Valley J20 | I-IV-J20E | <4.69 | <4.69 | 26.2 | <4.69 | <4.69 | <4.69 | 11.2 |
| Imperial Valley J22 | I-IV-J22A | <3.33 | <3.33 | 23.8 | <3.33 | <3.33 | <3.33 | 22 |
| Imperial Valley J22 | I-IV-J22B | <2.27 | <2.27 | 21 | <2.27 | <2.27 | <2.27 | 14.3 |
| Imperial Valley J22 | I-IV-J22C | <1.04 | <1.04 | 7.66 | <1.04 | <1.04 | <1.04 | 5.09 |
| Imperial Valley J22 | I-IV-J22D | <2.18 | <2.18 | 15 | <2.18 | <2.18 | <2.18 | 14 |
| Imperial Valley J22 | I-IV-J22E | <2.05 | <2.05 | 19.1 | <2.05 | 6.97 | <2.05 | 21.5 |
| Imperial Valley K18 | I-IV-K18A | 1.38 | <0.858 | 5.74 | 11.3 | <0.858 | <0.858 | 5.19 |
| Imperial Valley K18 | I-IV-K18B | <0.64 | <0.64 | 6.63 | <0.64 | <0.64 | <0.64 | 5.05 |
| Imperial Valley K18 | I-IV-K18C | <1.16 | <1.16 | 7.06 | 16.4 | 1.16 | <1.16 | 7.13 |
| Imperial Valley K18 | I-IV-K18D | <0.806 | <0.806 | 8.54 | <0.806 | <0.806 | <0.806 | 6.95 |
| Imperial Valley K18 | I-IV-K18E | <0.601 | <0.601 | 4.53 | <0.601 | <0.601 | <0.601 | 2.19 |
| Sonny Bono NWR | I-SB-420A | <0.741 | <0.741 | 5.19 | <0.741 | <0.741 | <0.741 | 2.56 |
| Sonny Bono NWR | I-SB-420B | <1.41 | <1.41 | 8.36 | <1.41 | <1.41 | <1.41 | 6.19 |
| Sonny Bono NWR | I-SB-420C | <1.14 | <1.14 | 7.29 | 6.86 | <1.14 | <1.14 | 4.23 |
| Sonny Bono NWR | I-SB-420D | <0.873 | <0.873 | 8.19 | 3.28 | <0.873 | <0.873 | 6.03 |
| Sonny Bono NWR | I-SB-420E | 7.38 | <0.833 | 5.18 | 3.12 | <0.833 | <0.833 | 3.19 |
| | | | | | | | | |
| Kern NWR | I-K-15A | <2.04 | 193 | 16.8 | <2.04 | <2.04 | 90.4 | 8.95 |
| Kern NWR | I-K-15B | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 |
| Kern NWR | I-K-15C | <1.24 | <1.24 | 7.13 | <1.24 | <1.24 | <1.24 | <1.24 |
| Kern NWR | I-K-15D | <2.25 | <2.25 | 20.7 | <2.25 | <2.25 | <2.25 | <2.25 |
| Kern NWR | I-K-15E | <1.61 | <1.61 | 11.2 | <1.61 | <1.61 | <1.61 | 4.21 |

Appendix 6. cont.

| Location | Sample | PCB# 175 ng/g dw | PCB# 177 ng/g dw | PCB# 180 ng/g dw | PCB# 193 ng/g dw | PCB# 196 ng/g dw | PCB# 197 ng/g dw | PCB# 206 ng/g dw |
|---------------------|-----------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| Imperial Valley J20 | I-IV-J20A | 10.7 | <1.7 | 3.41 | <1.7 | <1.7 | <1.7 | <1.7 |
| Imperial Valley J20 | I-IV-J20B | 10.6 | 24.5 | 3.06 | 6.48 | <1.61 | <1.61 | <1.61 |
| Imperial Valley J20 | I-IV-J20C | 23.5 | <3.31 | 3.63 | <3.31 | <3.31 | <3.31 | <3.31 |
| Imperial Valley J20 | I-IV-J20D | 24.1 | <3.28 | 5.73 | <3.28 | <3.28 | <3.28 | <3.28 |
| Imperial Valley J20 | I-IV-J20E | 32.1 | <4.69 | 11.7 | <4.69 | <4.69 | <4.69 | <4.69 |
| Imperial Valley J22 | I-IV-J22A | 28.7 | <3.33 | <3.33 | <3.33 | <3.33 | <3.33 | <3.33 |
| Imperial Valley J22 | I-IV-J22B | 18.3 | <2.27 | <2.27 | <2.27 | <2.27 | <2.27 | <2.27 |
| Imperial Valley J22 | I-IV-J22C | 7.57 | <1.04 | 3.65 | 4.06 | <1.04 | <1.04 | <1.04 |
| Imperial Valley J22 | I-IV-J22D | 16.9 | <2.18 | <2.18 | <2.18 | <2.18 | <2.18 | <2.18 |
| Imperial Valley J22 | I-IV-J22E | 12.4 | <2.05 | 8.35 | <2.05 | <2.05 | <2.05 | <2.05 |
| Imperial Valley K18 | I-IV-K18A | 5.52 | 4.5 | 1.18 | <0.858 | 3.46 | <0.858 | <0.858 |
| Imperial Valley K18 | I-IV-K18B | 3.4 | <0.64 | 1.23 | <0.64 | 3.91 | <0.64 | <0.64 |
| Imperial Valley K18 | I-IV-K18C | 7.8 | <1.16 | 2.5 | <1.16 | 3.6 | <1.16 | <1.16 |
| Imperial Valley K18 | I-IV-K18D | 4.69 | <0.806 | 2.3 | <0.806 | 4.3 | 4.01 | <0.806 |
| Imperial Valley K18 | I-IV-K18E | 1.77 | <0.601 | 0.633 | <0.601 | 1.26 | <0.601 | <0.601 |
| Sonny Bono NWR | I-SB-420A | 4.52 | <0.741 | 1.72 | <0.741 | <0.741 | <0.741 | 3.66 |
| Sonny Bono NWR | I-SB-420B | 8.8 | <1.41 | <1.41 | <1.41 | <1.41 | <1.41 | <1.41 |
| Sonny Bono NWR | I-SB-420C | 6.82 | <1.14 | 2.94 | <1.14 | <1.14 | <1.14 | <1.14 |
| Sonny Bono NWR | I-SB-420D | 5.05 | <0.873 | <0.873 | <0.873 | <0.873 | <0.873 | <0.873 |
| Sonny Bono NWR | I-SB-420E | 5.05 | <0.833 | <0.833 | <0.833 | <0.833 | <0.833 | <0.833 |
| | | | | | | | | |
| Kern NWR | I-K-15A | 17 | 193 | 3.67 | <2.04 | <2.04 | <2.04 | <1.41 |
| Kern NWR | I-K-15B | 22.7 | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 | <1.14 |
| Kern NWR | I-K-15C | 7.77 | <1.24 | <1.24 | <1.24 | <1.24 | <1.24 | <0.873 |
| Kern NWR | I-K-15D | 13.5 | <2.25 | 6.55 | <2.25 | <2.25 | <2.25 | <0.833 |
| Kern NWR | I-K-15E | 9.81 | <1.61 | 4.26 | <1.61 | <1.61 | <1.61 | <1.61 |

Appendix 6. cont.

| Location | Sample | PCB# 207 ng/g dw | PCB# 209 ng/g dw | PCB# 28 ng/g dw | PCB# 31 ng/g dw | PCB# 41/64 ng/g dw | PCB# 47/75 ng/g dw | PCB# 49 ng/g dw |
|---------------------|-----------|---------------------|---------------------|--------------------|--------------------|-----------------------|-----------------------|--------------------|
| Imperial Valley J20 | I-IV-J20A | <1.7 | <1.7 | 9.35 | <1.7 | <1.7 | <1.7 | <1.7 |
| Imperial Valley J20 | I-IV-J20B | <1.61 | 4.08 | 8.76 | <1.61 | <1.61 | <1.61 | <1.61 |
| Imperial Valley J20 | I-IV-J20C | <3.31 | 11.9 | 12.6 | <3.31 | <3.31 | <3.31 | <3.31 |
| Imperial Valley J20 | I-IV-J20D | <3.28 | <3.28 | 10.4 | <3.28 | <3.28 | <3.28 | <3.28 |
| Imperial Valley J20 | I-IV-J20E | <4.69 | <4.69 | 20.8 | <4.69 | <4.69 | <4.69 | <4.69 |
| Imperial Valley J22 | I-IV-J22A | <3.33 | <3.33 | 34.1 | <3.33 | 4.39 | <3.33 | <3.33 |
| Imperial Valley J22 | I-IV-J22B | <2.27 | <2.27 | 16 | <2.27 | 8.98 | <2.27 | <2.27 |
| Imperial Valley J22 | I-IV-J22C | <1.04 | <1.04 | 11.6 | 1.12 | 2.34 | <1.04 | <1.04 |
| Imperial Valley J22 | I-IV-J22D | <2.18 | <2.18 | 14.7 | <2.18 | <2.18 | <2.18 | <2.18 |
| Imperial Valley J22 | I-IV-J22E | <2.05 | <2.05 | 13.4 | <2.05 | <2.05 | <2.05 | <2.05 |
| Imperial Valley K18 | I-IV-K18A | <0.858 | <0.858 | 5.5 | <0.858 | <0.858 | <0.858 | <0.858 |
| Imperial Valley K18 | I-IV-K18B | <0.64 | 1.29 | 5.6 | <0.64 | 2.81 | <0.64 | <0.64 |
| Imperial Valley K18 | I-IV-K18C | <1.16 | 1.9 | 6.81 | <1.16 | <1.16 | <1.16 | <1.16 |
| Imperial Valley K18 | I-IV-K18D | <0.806 | 2.61 | 5.25 | <0.806 | <0.806 | <0.806 | <0.806 |
| Imperial Valley K18 | I-IV-K18E | <0.601 | 2.42 | 4.21 | <0.601 | <0.601 | <0.601 | <0.601 |
| Sonny Bono NWR | I-SB-420A | 2.05 | 5.65 | 3.92 | <0.741 | <0.741 | <0.741 | <0.741 |
| Sonny Bono NWR | I-SB-420B | <1.41 | <1.41 | 7.58 | <1.41 | <1.41 | <1.41 | <1.41 |
| Sonny Bono NWR | I-SB-420C | <1.14 | <1.14 | 5.95 | <1.14 | <1.14 | <1.14 | <1.14 |
| Sonny Bono NWR | I-SB-420D | <0.873 | <0.873 | 5.78 | <0.873 | <0.873 | <0.873 | <0.873 |
| Sonny Bono NWR | I-SB-420E | <0.833 | 1.19 | 6.4 | <0.833 | <0.833 | <0.833 | <0.833 |
| | | | | | | | | |
| Kern NWR | I-K-15A | <1.41 | <2.04 | 17.1 | <2.04 | <2.04 | <2.04 | <2.04 |
| Kern NWR | I-K-15B | <1.14 | <4.21 | 39.8 | <4.21 | <4.21 | <4.21 | <4.21 |
| Kern NWR | I-K-15C | <0.873 | <1.24 | 10.5 | <1.24 | <1.24 | 59.8 | 13.8 |
| Kern NWR | I-K-15D | <0.833 | <2.25 | 26.6 | <2.25 | <2.25 | <2.25 | <2.25 |
| Kern NWR | I-K-15E | <1.61 | <1.61 | 16.4 | <1.61 | <1.61 | <1.61 | <1.61 |

Appendix 6. cont.

| Location | Sample | PCB# 52 ng/g dw | PCB# 53 ng/g dw | PCB# 60/56 ng/g dw | PCB# 63 ng/g dw | PCB# 66 ng/g dw | PCB# 7/9 ng/g dw | PCB# 70 ng/g dw |
|---------------------|-----------|--------------------|--------------------|-----------------------|--------------------|--------------------|---------------------|--------------------|
| Imperial Valley J20 | I-IV-J20A | <1.7 | <1.7 | <1.7 | <1.7 | <1.7 | <1.7 | <1.7 |
| Imperial Valley J20 | I-IV-J20B | <1.61 | <1.61 | <1.61 | 218 | 10.6 | <1.61 | 16.2 |
| Imperial Valley J20 | I-IV-J20C | <3.31 | <3.31 | <3.31 | <3.31 | <3.31 | <3.31 | <3.31 |
| Imperial Valley J20 | I-IV-J20D | <3.28 | <3.28 | <3.28 | <3.28 | <3.28 | <3.28 | <3.28 |
| Imperial Valley J20 | I-IV-J20E | <4.69 | <4.69 | <4.69 | <4.69 | <4.69 | <4.69 | <4.69 |
| Imperial Valley J22 | I-IV-J22A | <3.33 | <3.33 | <3.33 | <3.33 | <3.33 | <3.33 | <3.33 |
| Imperial Valley J22 | I-IV-J22B | <2.27 | <2.27 | <2.27 | <2.27 | <2.27 | <2.27 | <2.27 |
| Imperial Valley J22 | I-IV-J22C | 11.3 | <1.04 | <1.04 | <1.04 | <1.04 | <1.04 | <1.04 |
| Imperial Valley J22 | I-IV-J22D | <2.18 | <2.18 | <2.18 | <2.18 | <2.18 | <2.18 | <2.18 |
| Imperial Valley J22 | I-IV-J22E | <2.05 | <2.05 | <2.05 | <2.05 | <2.05 | <2.05 | <2.05 |
| Imperial Valley K18 | I-IV-K18A | <0.858 | <0.858 | <0.858 | <0.858 | 5.54 | <0.858 | <0.858 |
| Imperial Valley K18 | I-IV-K18B | <0.64 | <0.64 | <0.64 | <0.64 | <0.64 | <0.64 | <0.64 |
| Imperial Valley K18 | I-IV-K18C | <1.16 | <1.16 | <1.16 | <1.16 | <1.16 | <1.16 | <1.16 |
| Imperial Valley K18 | I-IV-K18D | <0.806 | 7.12 | 12 | <0.806 | <0.806 | 2.38 | <0.806 |
| Imperial Valley K18 | I-IV-K18E | <0.601 | <0.601 | <0.601 | <0.601 | <0.601 | <0.601 | <0.601 |
| Sonny Bono NWR | I-SB-420A | <0.741 | <0.741 | <0.741 | <0.741 | <0.741 | <0.741 | <0.741 |
| Sonny Bono NWR | I-SB-420B | <1.41 | <1.41 | <1.41 | <1.41 | <1.41 | <1.41 | <1.41 |
| Sonny Bono NWR | I-SB-420C | <1.14 | <1.14 | <1.14 | <1.14 | <1.14 | <1.14 | <1.14 |
| Sonny Bono NWR | I-SB-420D | <0.873 | <0.873 | <0.873 | <0.873 | <0.873 | <0.873 | <0.873 |
| Sonny Bono NWR | I-SB-420E | <0.833 | <0.833 | <0.833 | 9.27 | <0.833 | <0.833 | <0.833 |
| | | | | | | | | |
| Kern NWR | I-K-15A | <2.04 | <2.04 | <2.04 | <2.04 | <2.04 | <2.04 | <2.04 |
| Kern NWR | I-K-15B | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 |
| Kern NWR | I-K-15C | <1.24 | <1.24 | <1.24 | <1.24 | <1.24 | <1.24 | <1.24 |
| Kern NWR | I-K-15D | <2.25 | <2.25 | <2.25 | <2.25 | <2.25 | <2.25 | <2.25 |
| Kern NWR | I-K-15E | <1.61 | <1.61 | <1.61 | <1.61 | <1.61 | <1.61 | <1.61 |

Appendix 6 . cont.

| Location | Sample | PCB# 72 ng/g dw | PCB# 74/61 ng/g dw | PCB# 82 ng/g dw | PCB# 84 ng/g dw | PCB# 87/115 ng/g dw | PCB# 92 ng/g dw | PCB# 95/80 ng/g dw |
|---------------------|-----------|--------------------|-----------------------|--------------------|--------------------|------------------------|--------------------|-----------------------|
| Imperial Valley J20 | I-IV-J20A | 10.7 | <1.7 | <1.7 | <1.7 | <1.7 | 14.4 | <1.7 |
| Imperial Valley J20 | I-IV-J20B | 22.2 | 46.7 | <1.61 | 4.67 | <1.61 | 4.5 | <1.61 |
| Imperial Valley J20 | I-IV-J20C | 26.2 | <3.31 | <3.31 | <3.31 | <3.31 | <3.31 | <3.31 |
| Imperial Valley J20 | I-IV-J20D | 32.4 | <3.28 | <3.28 | 22.2 | <3.28 | 5.98 | <3.28 |
| Imperial Valley J20 | I-IV-J20E | <4.69 | <4.69 | <4.69 | <4.69 | <4.69 | <4.69 | <4.69 |
| Imperial Valley J22 | I-IV-J22A | 38.7 | <3.33 | <3.33 | <3.33 | <3.33 | <3.33 | <3.33 |
| Imperial Valley J22 | I-IV-J22B | 65.1 | <2.27 | <2.27 | <2.27 | <2.27 | <2.27 | <2.27 |
| Imperial Valley J22 | I-IV-J22C | 37.9 | <1.04 | <1.04 | <1.04 | <1.04 | <1.04 | <1.04 |
| Imperial Valley J22 | I-IV-J22D | 50 | <2.18 | <2.18 | <2.18 | <2.18 | <2.18 | <2.18 |
| Imperial Valley J22 | I-IV-J22E | <2.05 | <2.05 | <2.05 | 27.4 | <2.05 | 36.1 | <2.05 |
| Imperial Valley K18 | I-IV-K18A | 4.92 | <0.858 | <0.858 | 3.78 | <0.858 | 10.7 | <0.858 |
| Imperial Valley K18 | I-IV-K18B | <0.64 | <0.64 | <0.64 | 10.1 | <0.64 | 1.65 | <0.64 |
| Imperial Valley K18 | I-IV-K18C | 7.89 | <1.16 | <1.16 | <1.16 | <1.16 | 15.7 | <1.16 |
| Imperial Valley K18 | I-IV-K18D | 17 | <0.806 | <0.806 | 5.58 | <0.806 | 1.13 | <0.806 |
| Imperial Valley K18 | I-IV-K18E | 9.04 | <0.601 | <0.601 | <0.601 | <0.601 | <0.601 | <0.601 |
| Sonny Bono NWR | I-SB-420A | <0.741 | <0.741 | <0.741 | <0.741 | <0.741 | <0.741 | <0.741 |
| Sonny Bono NWR | I-SB-420B | 9.78 | <1.41 | <1.41 | 12.5 | <1.41 | 14.8 | <1.41 |
| Sonny Bono NWR | I-SB-420C | <1.14 | <1.14 | <1.14 | <1.14 | <1.14 | <1.14 | <1.14 |
| Sonny Bono NWR | I-SB-420D | 22.9 | <0.873 | <0.873 | <0.873 | <0.873 | 5.85 | <0.873 |
| Sonny Bono NWR | I-SB-420E | <0.833 | <0.833 | 4.7 | <0.833 | 6.83 | <0.833 | 3.69 |
| | | | | | | | | |
| Kern NWR | I-K-15A | <2.04 | <2.04 | <2.04 | <2.04 | <2.04 | <2.04 | <2.04 |
| Kern NWR | I-K-15B | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 | <4.21 |
| Kern NWR | I-K-15C | <1.24 | <1.24 | <1.24 | <1.24 | <1.24 | <1.24 | <1.24 |
| Kern NWR | I-K-15D | <2.25 | <2.25 | <2.25 | <2.25 | <2.25 | <2.25 | <2.25 |
| Kern NWR | I-K-15E | <1.61 | <1.61 | <1.61 | <1.61 | <1.61 | <1.61 | <1.61 |

Appendix 6. cont.

| Location | Sample | PCB# 99 ng/g dw | PCB-1268 ng/g dw | PCB-TOTAL ng/g dw |
|---------------------|---------------|----------------------------|-----------------------------|------------------------------|
| Imperial Valley J20 | I-IV-J20A | <1.7 | <170 | <170 |
| Imperial Valley J20 | I-IV-J20B | 260 | 768 | 768 |
| Imperial Valley J20 | I-IV-J20C | <3.31 | <331 | <331 |
| Imperial Valley J20 | I-IV-J20D | <3.28 | <328 | <328 |
| Imperial Valley J20 | I-IV-J20E | <4.69 | <469 | <469 |
| Imperial Valley J22 | I-IV-J22A | <3.33 | <333 | <333 |
| Imperial Valley J22 | I-IV-J22B | <2.27 | <227 | <227 |
| Imperial Valley J22 | I-IV-J22C | <1.04 | <104 | <104 |
| Imperial Valley J22 | I-IV-J22D | <2.18 | <218 | <218 |
| Imperial Valley J22 | I-IV-J22E | <2.05 | <205 | <205 |
| Imperial Valley K18 | I-IV-K18A | <0.858 | <85.8 | <85.8 |
| Imperial Valley K18 | I-IV-K18B | <0.64 | <64 | <64 |
| Imperial Valley K18 | I-IV-K18C | <1.16 | <116 | <116 |
| Imperial Valley K18 | I-IV-K18D | <0.806 | 86.7 | 86.8 |
| Imperial Valley K18 | I-IV-K18E | <0.601 | <60.1 | 60.1 |
| Sonny Bono NWR | I-SB-420A | <0.741 | <74.1 | 74.1 |
| Sonny Bono NWR | I-SB-420B | <1.41 | <141 | 141 |
| Sonny Bono NWR | I-SB-420C | <1.14 | <114 | 114 |
| Sonny Bono NWR | I-SB-420D | <0.873 | <87.3 | 87.3 |
| Sonny Bono NWR | I-SB-420E | 33.5 | 95.4 | 95.5 |
| | | | | |
| Kern NWR | I-K-15A | 191 | 605 | 605 |
| Kern NWR | I-K-15B | <4.21 | <421 | <421 |
| Kern NWR | I-K-15C | <1.24 | <124 | <124 |
| Kern NWR | I-K-15D | <2.25 | <225 | <225 |
| Kern NWR | I-K-15E | <1.61 | <161 | <161 |

Appendix 6. cont.

* C11# PCB, C110# PCB, C12# PCB, C13# PCB, C17# PCB, C18# PCB, C19# PCB, PCB# 101/90, PCB# 110, PCB#114, PCB# 118, PCB# 126, PCB# 128, PCB# 129, PCB# 130, PCB# 136, PCB# 146, PCB# 15, PCB# 153/132, PCB# 156, PCB# 157/173/201, PCB# 158, PCB# 16/32, PCB# 166, PCB# 167, PCB# 169, PCB# 170/190, PCB# 171/202, PCB# 174, PCB# 176/137, PCB# 178, PCB# 18/17, PCB# 183, PCB# 185, PCB# 187, PCB# 189, PCB# 191, PCB# 194, PCB# 195/208, PCB# 199, PCB# 200, PCB# 205, PCB# 22/51, PCB# 24/27, PCB# 25, PCB# 26, PCB# 29, PCB# 30, PCB# 33/20, PCB# 39, PCB# 40, PCB# 42/59/37, PCB# 44, PCB# 45, PCB# 46, PCB# 48, PCB#67, PCB#69, PCB#77, PCB# 8/5, PCB# 81, PCB# 83, PCB# 85, PCB# 97, PCB-1242, PCB-1248, PCB-1254, and PCB 1260 not detected in any terrestrial invertebrate samples.

Appendix 7. Concentrations of inorganics (ug/g dry weight) in soil samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006.

| Location | Sample | Ag ug/g dw | Al ug/g dw | As ug/g dw | B ug/g dw | Ba ug/g dw | Be ug/g dw | Ca ug/g dw | Cd ug/g dw | Co ug/g dw |
|---------------------------------------|--------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|---------------|---------------|
| Pawnee National Grasslands, CO | MPPGSL01 | <0.192 | 11200 | 2.26 | 4.08 | 123 | 0.526 | 6270 | 0.128 | 3.66 |
| Charles M. Russell NWR, MT | CMRDLWLDS06 | <0.194 | 20100 | 12.4 | 12.1 | 272 | 0.799 | 2830 | 0.31 | 12.6 |
| Charles M. Russell NWR, MT | CMRNMCRBWS06 | <0.192 | 14500 | 9.97 | 9.41 | 115 | 0.626 | 1940 | 0.274 | 9.67 |
| Charles M. Russell NWR, MT | CMRNSBSRS06 | 0.258 | 21500 | 10 | 14.2 | 363 | 0.743 | 5060 | 1.09 | 23.2 |
| Hailstone NWR, MT | HailS060506 | <0.19 | 21100 | 9.47 | 12.5 | 187 | 0.79 | 4340 | 0.354 | 7.4 |
| Lake Mason NWR, MT | LMS072606 | <0.193 | 22700 | 8.18 | 13.9 | 206 | 0.988 | 27900 | 0.392 | 8.27 |
| Antelope Coal Mine, WY | MPAMSL01 | <0.196 | 11100 | 5.32 | <1.96 | 148 | 0.972 | 6000 | 0.394 | 8.15 |
| Foote Creek Rim, WY | MPFCSL01 | <0.193 | 11500 | 4.66 | 4.96 | 101 | 0.441 | 4230 | 0.27 | 6.69 |
| Thunder Basin National Grasslands, WY | MPTBSL01 | <0.192 | 4620 | 2.9 | <1.92 | 88.4 | 0.393 | 1370 | 0.281 | 5 |

Appendix 7. cont.

| Location | Sample | Cr ug/g dw | Cu ug/g dw | Fe ug/g dw | Hg ug/g dw | K ug/g dw | Mg ug/g dw | Mn ug/g dw | Mo ug/g dw | Na ug/g dw |
|---------------------------------------|--------------|---------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|---------------|
| Pawnee National Grasslands, CO | MPPGSL01 | 8.91 | 6.74 | 11500 | 0.0101 | 3440 | 3320 | 228 | <0.960 | <192 |
| Charles M. Russell NWR, MT | CMRDLWLDS06 | 24 | 20.6 | 27000 | 0.0259 | 4000 | 5510 | 458 | <0.97 | 560 |
| Charles M. Russell NWR, MT | CMRNMCRBWS06 | 21.1 | 16.5 | 22400 | 0.0257 | 3160 | 3720 | 468 | <0.96 | 536 |
| Charles M. Russell NWR, MT | CMRNSBSRS06 | 25.6 | 18.3 | 24700 | 0.02 | 4810 | 6630 | 2290 | <0.96 | 356 |
| Hailstone NWR, MT | HailS060506 | 21.9 | 18.1 | 22300 | 0.0281 | 6000 | 6780 | 426 | 0.99 | <190 |
| Lake Mason NWR, MT | LMS072606 | 23 | 19.9 | 22700 | 0.0233 | 7330 | 9150 | 487 | <0.96 | <193 |
| Antelope Coal Mine, WY | MPAMSL01 | 16.9 | 30 | 16800 | 0.07 | 2540 | 3120 | 141 | <0.980 | 507 |
| Foote Creek Rim, WY | MPFCSL01 | 22.1 | 22 | 14900 | 0.0142 | 2780 | 2990 | 288 | <0.970 | <193 |
| Thunder Basin National Grasslands, WY | MPTBSL01 | 8.58 | 18.1 | 8700 | 0.0344 | 1280 | 840 | 97.8 | <0.960 | <192 |

Appendix 7. cont.

| Location | Sample | Ni ug/g dw | P ug/g dw | Pb ug/g dw | S ug/g dw | Se ug/g dw | Sr ug/g dw | V ug/g dw | Zn ug/g dw |
|---------------------------------------|--------------|---------------|--------------|---------------|--------------|---------------|---------------|--------------|---------------|
| Pawnee National Grasslands, CO | MPPGSL01 | 4.97 | 387 | 7.86 | 108 | 0.143 | 37.8 | 21.5 | 30.3 |
| Charles M. Russell NWR, MT | CMRDLWLDS06 | 23.1 | 638 | 14.3 | 690 | 0.589 | 51.2 | 65.6 | 78 |
| Charles M. Russell NWR, MT | CMRNMCRBWS06 | 17.8 | 591 | 12.4 | 408 | 0.561 | 30.5 | 59.3 | 65.1 |
| Charles M. Russell NWR, MT | CMRNSBSRS06 | 40.8 | 602 | 13 | 521 | 0.391 | 41.3 | 63.8 | 74.9 |
| Hailstone NWR, MT | HailS060506 | 18 | 828 | 15.1 | 337 | 0.499 | 24 | 46.5 | 82.9 |
| Lake Mason NWR, MT | LMS072606 | 20 | 863 | 13.6 | 464 | 0.262 | 109 | 41.1 | 84.9 |
| Antelope Coal Mine, WY | MPAMSL01 | 16.6 | 402 | 17.1 | 2200 | 1.22 | 95.2 | 37.3 | 80.6 |
| Foote Creek Rim, WY | MPFCSL01 | 33.7 | 401 | 8.14 | 305 | 0.204 | 17.9 | 33.4 | 38.1 |
| Thunder Basin National Grasslands, WY | MPTBSL01 | 9.12 | 272 | 9.29 | 188 | 0.27 | 24.8 | 22.7 | 43.8 |

Appendix 8. Organochlorine pesticides,* organophosphate pesticides,** and total PCBs (ng/g dry weight) were not detected in soil samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006.

*Organochlorine pesticides analyzed for in soil samples from Colorado, Montana, and Wyoming included: alpha BHC, alpha chlordane, beta BHC, cis-nonachlor, delta BHC, dieldrin, endrin, gamma BHC, gamma chlordane, HCB, heptachlor epoxide, mirex, o,p'-DDD, o,p'-DDE, o,p'-DDT, oxychlordane, p,p'-DDD, p,p'-DDE, p,p'-DDT, PCB Total, toxaphene, and trans-nonachlor.

**Organophosphate pesticides analyzed for in soil samples from Colorado and Wyoming included: alachlor, aldicarb, azinphosmethyl, carbaryl, carbofuran, chlorpyrifos, coumaphos, demeton, diazinon, dichlorvos, dimethoate, disulfoton, ethoprop, famphur, fensulfothion, fenthion, malathion, methiocarb, methomyl, methylparathion, mevinphos, oxamyl, parathion, phorate, terbufos, 1-naphthol, aldicarb, sulfone, aldicarb sulfoxide, baygon, carbofuran 3-OH, and EPN. These pesticides were not analyzed for in soil samples from Montana.

Appendix 9. Concentrations of inorganics* (ug/g dry weight) in terrestrial invertebrate samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006.

| Location | Sample | Moisture % | Al ug/g dw | As ug/g dw | B ug/g dw | Ba ug/g dw | Be ug/g dw | Ca ug/g dw | Cd ug/g dw | Co ug/g dw |
|---------------------------------------|-------------|------------|---------------|---------------|--------------|---------------|---------------|---------------|---------------|---------------|
| Pawnee National Grasslands, CO | MPPGTI01 | 65.6 | 134 | 0.112 | 4.56 | 9.18 | <0.0468 | 1440 | 0.226 | 0.468 |
| Pawnee National Grasslands, CO | MPPGTI03 | 65.7 | 101 | 0.0895 | 3.61 | 6.93 | <0.0486 | 1530 | 0.348 | 0.486 |
| Charles M. Russell NWR, MT | CMRDLWLDI06 | 60.9 | 381 | 0.951 | 5.11 | 12.5 | <0.0468 | 1600 | 0.439 | 0.835 |
| Charles M. Russell NWR, MT | CMRNMCBWI06 | 54.3 | 2180 | 2.15 | 6.64 | 42.6 | 0.132 | 2810 | 0.333 | 2.12 |
| Charles M. Russell NWR, MT | CMRNSBSRI06 | 64.3 | 387 | 1.01 | 4.38 | 14.3 | <0.0449 | 2920 | 0.385 | 0.627 |
| Hailstone NWR, MT | HailI060506 | 66.2 | 119 | 0.202 | 18.8 | 11.4 | <0.0452 | 1640 | 0.306 | <0.452 |
| Lake Mason NWR, MT | LMI072606 | 63.6 | 90.3 | <0.178 | 17.8 | 9.68 | <0.045 | 1870 | 0.221 | <0.45 |
| Antelope Coal Mine, WY | MPAMTI01 | 66.9 | 79 | 0.102 | 5.24 | 7.19 | <0.0476 | 1140 | 0.188 | 0.476 |
| Foote Creek Rim, WY | MPFCTI01 | 68.5 | 115 | 0.105 | 3.24 | 5.85 | <0.0461 | 2250 | 0.0941 | 0.461 |
| Foote Creek Rim, WY | MPFCTI04 | 66.7 | 183 | 0.407 | 4.65 | 11.8 | <0.0457 | 8200 | 0.164 | 0.457 |
| Thunder Basin National Grasslands, WY | MPTBTI03 | 66.5 | 52.8 | 0.0745 | 3.86 | 6.51 | <0.0514 | 857 | 0.309 | 0.514 |
| Thunder Basin National Grasslands, WY | MPTBTI04 | 67.2 | 114 | 0.212 | 13.1 | 11.5 | <0.0494 | 1120 | 0.285 | 0.494 |

Appendix 9. cont.

| Location | Sample | Cr ug/g dw | Cu ug/g dw | Fe ug/g dw | Hg ug/g dw | K ug/g dw | Mg ug/g dw | Mn ug/g dw | Mo ug/g dw | Na ug/g dw |
|--|--------------|---------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|---------------|
| Pawnee National Grasslands, CO | MPPGTI01 | 0.547 | 14.1 | 198 | <0.0048 | 8150 | 881 | 15.4 | <0.936 | 910 |
| Pawnee National Grasslands, CO | MPPGTI03 | <0.486 | 23.6 | 132 | <0.0047 | 8870 | 879 | 12.1 | <0.972 | 1000 |
| Charles M. Russell NWR, MT | CMRDLWLDI06 | 1.69 | 19.8 | 771 | 0.0103 | 6450 | 1710 | 79.9 | <0.936 | 2010 |
| Charles M. Russell NWR, MT | CMRNMCRBWI06 | 3.71 | 25.5 | 4300 | 0.0166 | 9230 | 2070 | 165 | 1.03 | 920 |
| Charles M. Russell NWR, MT | CMRNSBSRI06 | 1.52 | 17.1 | 1260 | 0.00659 | 5740 | 1540 | 71.2 | <0.898 | 2220 |
| Hailstone NWR, MT | HailI060506 | 0.565 | 42.3 | 225 | 0.0106 | 9370 | 1880 | 20.5 | <0.904 | 1800 |
| Lake Mason NWR, MT | LMI072606 | <0.45 | 47.5 | 161 | 0.0121 | 9840 | 1970 | 15.8 | 1.04 | 1540 |
| Antelope Coal Mine, WY | MPAMTI01 | <0.476 | 28 | 163 | <0.0049 | 9060 | 920 | 11.7 | <0.953 | 1110 |
| Foote Creek Rim, WY | MPFCTI01 | 0.702 | 27.8 | 176 | <0.0048 | 8890 | 874 | 17.8 | <0.922 | 700 |
| Foote Creek Rim, WY | MPFCTI04 | 0.692 | 20.3 | 242 | 0.0068 | 8470 | 989 | 22.2 | <0.913 | 661 |
| Thunder Basin National Grasslands, WY | MPTBTI03 | 0.592 | 18.3 | 136 | 0.0067 | 8230 | 1000 | 14.6 | <1.03 | 985 |
| Thunder Basin National Grasslands, WY | MPTBTI04 | 0.666 | 25.4 | 358 | 0.0101 | 7940 | 977 | 20.4 | <0.988 | 954 |

Appendix 9. cont.

| Location | Sample | Ni ug/g dw | P ug/g dw | Pb ug/g dw | S ug/g dw | Se ug/g dw | Si ug/g dw | Sr ug/g dw | Ti ug/g dw | V ug/g dw | Zn ug/g dw |
|--|--------------|---------------|--------------|---------------|--------------|---------------|---------------|---------------|---------------|--------------|---------------|
| Pawnee National Grasslands, CO | MPPGTI01 | 0.62 | 5990 | 0.328 | 4240 | 0.678 | 179 | 6.2 | 9.35 | <0.468 | 85.5 |
| Pawnee National Grasslands, CO | MPPGTI03 | <0.486 | 6480 | 0.226 | 4560 | 0.613 | 207 | 7.51 | 6.65 | <0.486 | 104 |
| Charles M. Russell NWR, MT | CMRDLWLDI06 | 1.64 | 6110 | 0.866 | 5240 | 1.11 | NA** | 10.1 | 12.6 | 1.47 | 168 |
| Charles M. Russell NWR, MT | CMRNMCRBWI06 | 4.94 | 7180 | 3.15 | 4570 | 1.76 | NA | 19.2 | 43 | 7.77 | 147 |
| Charles M. Russell NWR, MT | CMRNSBSRI06 | 1.3 | 5680 | 0.443 | 4980 | 2.56 | NA | 17 | 14.2 | 1.85 | 169 |
| Hailstone NWR, MT | HailI060506 | 0.831 | 6740 | 0.263 | 5270 | 0.853 | NA | 27.3 | 4.72 | <0.452 | 147 |
| Lake Mason NWR, MT | LMI072606 | 0.916 | 7090 | 0.212 | 4790 | 0.676 | NA | 31 | 2.93 | <0.45 | 137 |
| Antelope Coal Mine, WY | MPAMTI01 | 0.829 | 7120 | 0.202 | 4690 | 0.734 | 150 | 8.16 | 5.93 | <0.476 | 137 |
| Foote Creek Rim, WY | MPFCTI01 | 1.12 | 7290 | 0.34 | 4340 | 0.473 | 155 | 5.82 | 5.98 | <0.461 | 116 |
| Foote Creek Rim, WY | MPFCTI04 | 1.66 | 7030 | 0.244 | 4160 | 0.438 | 199 | 17.8 | 9.79 | 0.588 | 111 |
| Thunder Basin National Grasslands, WY | MPTBTI03 | 1.2 | 6820 | 0.135 | 4320 | 0.92 | 131 | 6.44 | 6.67 | <0.514 | 126 |
| Thunder Basin National Grasslands, WY | MPTBTI04 | 1.77 | 6600 | 0.34 | 4340 | 0.754 | 141 | 8.8 | 7.58 | 0.682 | 137 |

*Silver not detected in any terrestrial invertebrate samples.

**NA = Not analyzed.

Appendix 10. Organochlorine pesticides,* organophosphate pesticides,** and total PCBs (ng/g dry weight) were not detected in terrestrial invertebrate samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006.

*Organochlorine pesticides analyzed for in terrestrial invertebrate samples from Colorado, Montana, and Wyoming included: alpha BHC, alpha chlordane, beta BHC, cis-nonachlor, delta BHC, dieldrin, endrin, gamma BHC, gamma chlordane, HCB, heptachlor epoxide, mirex, o,p'-DDD, o,p'-DDE, o,p'-DDT, oxychlordane, p,p'-DDD, p,p'-DDE, p,p'-DDT, PCB Total, toxaphene, and trans-nonachlor.

**Organophosphate pesticides analyzed for in terrestrial invertebrate samples from Colorado and Wyoming included: alachlor, aldicarb, azinphosmethyl, carbaryl, carbofuran, chlorpyrifos, coumaphos, diazinon, dichlorvos, dimethoate, ethoprop, famphur, fensulfothion, fenthion, malathion, methiocarb, methomyl, methylparathion, mevinphos, oxamyl, parathion, phorate, terbufos, 1-naphthol, aldicarb, sulfone, aldicarb sulfoxide, baygon, carbofuran 3-OH, and EPN. These pesticides were not analyzed for in invertebrate samples from Montana.

Appendix 11. Concentrations of inorganics* (ug/g dry weight) in eggs samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006.

| Location | Sample | Moisture % | Al ug/g dw | As ug/g dw | Ba ug/g dw | Ca ug/g dw | Cd ug/g dw | Cu ug/g dw | Fe ug/g dw |
|--------------------------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Pawnee National Grasslands, CO | MPWCPE01 | 72.1 | 1.47 | 0.0593 | 10.8 | 2850 | <0.0192 | 2.65 | 104 |
| Pawnee National Grasslands, CO | MPWCPE02 | 71.7 | 0.909 | <0.0483 | 13.7 | 2680 | <0.0193 | 2.71 | 105 |
| Pawnee National Grasslands, CO | MPWCPE03 | 73.8 | 1.34 | 0.224 | 18.4 | 9900 | <0.02 | 3.37 | 116 |
| Pawnee National Grasslands, CO | MPWCPE04 | 71 | 0.677 | 0.0507 | 19.6 | 3140 | <0.0188 | 2.23 | 96.3 |
| Pawnee National Grasslands, CO | MPWCPE05 | 74.7 | 1.33 | 0.193 | 10.5 | 9120 | 0.0207 | 4.29 | 136 |
| | | | | | | | | | |
| BLM Lands, Phillips County, MT | BLM025MPE | 73.4 | <0.443 | 0.175 | 7.83 | 4130 | <0.0175 | 2.66 | 97.9 |
| BLM Lands, Phillips County, MT | BLM081-2MPE | 70.7 | <0.432 | <0.171 | 8.83 | 2680 | <0.0171 | 2.8 | 105 |
| BLM Lands, Phillips County, MT | BLM081MPE | 70.3 | 0.893 | <0.176 | 9.37 | 2980 | <0.0176 | 2.77 | 102 |
| BLM Lands, Phillips County, MT | BLM189MPE | 70.8 | <0.429 | <0.17 | 5.75 | 4740 | <0.017 | 3.33 | 84.9 |
| BLM Lands, Phillips County, MT | BLMB026MPE | 74.6 | 0.599 | 0.188 | 6.71 | 13400 | <0.0188 | 3.92 | 126 |
| BLM Lands, Phillips County, MT | BLMS014MPE | 75.9 | 0.538 | 0.182 | 8.65 | 19200 | <0.0182 | 4.13 | 125 |
| BLM Lands, Phillips County, MT | BLMS020MPE | 72.6 | 0.818 | <0.177 | 9.72 | 4630 | <0.177 | 2.68 | 102 |
| Charles M. Russell NWR, MT | CMRNMCMPE06 | 74.2 | 1.73 | <0.178 | 10 | 2510 | <0.178 | 2.98 | 111 |
| | | | | | | | | | |
| Near Antelope Coal Mine, WY | MPAMPE01 | 72.6 | 0.732 | <0.0483 | 5.2 | 2800 | <0.0193 | 2.63 | 103 |
| Near Antelope Coal Mine, WY | MPAMPE02 | 72 | 1.65 | 0.274 | 13.1 | 14100 | 0.0645 | 3.58 | 115 |
| Foote Creek Rim, WY | MPFCPE01 | 71.6 | 0.658 | 0.0631 | 4.86 | 4260 | <0.0204 | 2.83 | 77.4 |
| Foote Creek Rim, WY | MPFCPE02 | 72 | 0.729 | 0.113 | 4.97 | 5960 | <0.0176 | 2.65 | 92.8 |
| Foote Creek Rim, WY | MPFCPE03 | 72.1 | 0.619 | 0.0492 | 3.15 | 2750 | <0.019 | 2.77 | 82.9 |
| Foote Creek Rim, WY | MPFCPE04 | 74.3 | 1.19 | 0.292 | 5.85 | 13500 | <0.0186 | 3 | 97.7 |
| Foote Creek Rim, WY | MPFCPE05 | 73 | 0.909 | 0.0547 | 5.49 | 3020 | <0.0177 | 2.87 | 90.7 |

Appendix 11. Cont.

| Location | Sample | Hg ug/g dw | K ug/g dw | Mg ug/g dw | Mn ug/g dw | Na ug/g dw | Ni ug/g dw | P ug/g dw | Pb ug/g dw |
|--------------------------------|-------------|---------------|--------------|---------------|---------------|---------------|---------------|--------------|---------------|
| Pawnee National Grasslands, CO | MPWCPE01 | 0.0514 | 4490 | 371 | 0.962 | 5380 | 14.4 | 8690 | 0.0593 |
| Pawnee National Grasslands, CO | MPWCPE02 | 0.0365 | 4800 | 433 | 1.46 | 4610 | 4.06 | 7920 | <0.0483 |
| Pawnee National Grasslands, CO | MPWCPE03 | 0.0349 | 6310 | 586 | 1.24 | 5600 | 35 | 9810 | 0.185 |
| Pawnee National Grasslands, CO | MPWCPE04 | 0.0262 | 5190 | 425 | 1.95 | 4490 | 1.15 | 8670 | <0.0471 |
| Pawnee National Grasslands, CO | MPWCPE05 | 0.0344 | 5710 | 590 | 2.14 | 6190 | 55.7 | 9700 | <0.0487 |
| | | | | | | | | | |
| BLM Lands, Phillips County, MT | BLM025MPE | 0.0267 | 5340 | 471 | 1.06 | 5200 | <0.443 | 8470 | <0.0438 |
| BLM Lands, Phillips County, MT | BLM081-2MPE | 0.0488 | 5140 | 403 | 1.51 | 5410 | <0.432 | 7800 | <0.0428 |
| BLM Lands, Phillips County, MT | BLM081MPE | 0.0327 | 4700 | 434 | 1.39 | 5410 | <0.444 | 7880 | 0.0951 |
| BLM Lands, Phillips County, MT | BLM189MPE | 0.0409 | 5910 | 448 | 0.671 | 5200 | <0.429 | 7380 | <0.0425 |
| BLM Lands, Phillips County, MT | BLMB026MPE | 0.0322 | 6330 | 742 | 1.26 | 6950 | <0.475 | 9470 | 0.131 |
| BLM Lands, Phillips County, MT | BLMS014MPE | 0.0359 | 6350 | 922 | 2.05 | 7310 | <0.46 | 10900 | 0.0812 |
| BLM Lands, Phillips County, MT | BLMS020MPE | 0.0252 | 5310 | 498 | 0.979 | 6020 | <0.447 | 8220 | <0.0443 |
| Charles M. Russell NWR, MT | CMRNMCMPE06 | 0.0307 | 5480 | 448 | 1.43 | 5020 | <0.448 | 8240 | 0.0468 |
| | | | | | | | | | |
| Near Antelope Coal Mine, WY | MPAMPE01 | 0.0225 | 4910 | 480 | 1.2 | 5020 | <0.483 | 8080 | 0.0501 |
| Near Antelope Coal Mine, WY | MPAMPE02 | 0.0707 | 5420 | 679 | 2.61 | 5720 | <0.482 | 10800 | 0.068 |
| Foote Creek Rim, WY | MPFCPE01 | 0.0419 | 5730 | 484 | 1.08 | 4940 | 0.669 | 8510 | <0.0509 |
| Foote Creek Rim, WY | MPFCPE02 | 0.075 | 5240 | 486 | 1.83 | 5580 | <0.44 | 7810 | <0.044 |
| Foote Creek Rim, WY | MPFCPE03 | 0.0505 | 4620 | 401 | 0.907 | 4520 | <0.476 | 8290 | <0.0476 |
| Foote Creek Rim, WY | MPFCPE04 | 0.0518 | 5730 | 748 | 1.52 | 6570 | <0.465 | 10300 | <0.0465 |
| Foote Creek Rim, WY | MPFCPE05 | 0.0385 | 5320 | 420 | 1.1 | 5150 | <0.443 | 8150 | 0.046 |

Appendix 11. Cont.

| Location | Sample | S ug/g dw | Se ug/g dw | Si ug/g dw | Sr ug/g dw | Ti ug/g dw | Zn ug/g dw |
|--------------------------------|-------------|--------------|---------------|---------------|---------------|---------------|---------------|
| Pawnee National Grasslands, CO | MPWCPE01 | 5420 | 3.84 | 9.77 | 4.3 | <0.481 | 48.4 |
| Pawnee National Grasslands, CO | MPWCPE02 | 6740 | 2.66 | 12.9 | 3.81 | <0.483 | 39.7 |
| Pawnee National Grasslands, CO | MPWCPE03 | 6840 | 3 | 11.8 | 11.5 | 1.06 | 60 |
| Pawnee National Grasslands, CO | MPWCPE04 | 5950 | 2.19 | 11.6 | 6.22 | <0.471 | 55.3 |
| Pawnee National Grasslands, CO | MPWCPE05 | 6520 | 2.75 | 11.9 | 13.5 | 0.946 | 56 |
| | | | | | | | |
| BLM Lands, Phillips County, MT | BLM025MPE | 7160 | 3.85 | NA** | 6.21 | <0.0443 | 43.6 |
| BLM Lands, Phillips County, MT | BLM081-2MPE | 6840 | 5.15 | NA | 4.34 | <0.432 | 36.9 |
| BLM Lands, Phillips County, MT | BLM081MPE | 6670 | 4.93 | NA | 5.07 | <0.444 | 36.5 |
| BLM Lands, Phillips County, MT | BLM189MPE | 7040 | 4.01 | NA | 2.74 | <0.429 | 43.3 |
| BLM Lands, Phillips County, MT | BLMB026MPE | 9040 | 5.22 | NA | 15.5 | <0.475 | 52.9 |
| BLM Lands, Phillips County, MT | BLMS014MPE | 9070 | 7.32 | NA | 16.6 | <0.46 | 51.5 |
| BLM Lands, Phillips County, MT | BLMS020MPE | 7350 | 3.64 | NA | 8.98 | <0.447 | 52.9 |
| Charles M. Russell NWR, MT | CMRNMCMPE06 | 6530 | 3.63 | NA | 6.88 | <0.448 | 40.6 |
| | | | | | | | |
| Near Antelope Coal Mine, WY | MPAMPE01 | 6580 | 3.03 | 10.1 | 4.35 | <0.483 | 43 |
| Near Antelope Coal Mine, WY | MPAMPE02 | 6610 | 3.67 | 12.9 | 22 | 2.08 | 64.1 |
| Foote Creek Rim, WY | MPFCPE01 | 6680 | 2.53 | 8.03 | 5.68 | <0.509 | 55 |
| Foote Creek Rim, WY | MPFCPE02 | 7020 | 3.1 | 8.24 | 7.92 | <0.44 | 44.3 |
| Foote Creek Rim, WY | MPFCPE03 | 5920 | 2.68 | 9.38 | 3.63 | <0.476 | 43 |
| Foote Creek Rim, WY | MPFCPE04 | 7650 | 2.73 | 9.23 | 17.1 | 1.78 | 60.2 |
| Foote Creek Rim, WY | MPFCPE05 | 6450 | 2.16 | 9.93 | 4.42 | 0.564 | 60 |

*Silver, boron, beryllium, cobalt, chromium, molybdenum, and vanadium not detected in any mountain plover egg samples.

**NA = Not analyzed.

Appendix 12. Concentrations of organic compounds¹ (ng/g dry weight²) in eggs samples collected from mountain plover breeding sites in Colorado, Montana, and Wyoming, 2006.

| Location | Sample | Moisture % | HCB ng/g dw | oxychlordane ng/g dw | p,p'-DDE ng/g dw | p,p'-DDE ng/g ww |
|--------------------------------|--------------|------------|-------------|----------------------|------------------|------------------|
| Pawnee National Grasslands, CO | MPWCPE01 | 75.4 | 32.5 | 48.8 | 11,800 | 2,900 |
| Pawnee National Grasslands, CO | MPWCPE02 | 73.4 | 7.52 | 11.3 | 43,200 | 11,500 |
| Pawnee National Grasslands, CO | MPWCPE03 | 72.8 | 14.7 | 18.4 | 9,930 | 2,700 |
| Pawnee National Grasslands, CO | MPWCPE04 | 78.9 | 9.48 | 9.48 | 332 | 70 |
| Pawnee National Grasslands, CO | MPWCPE05 | 76.5 | 12.8 | <8.51 | 46.8 | 11 |
| BLM Lands, Phillips County, MT | BLM025MPE | 76.3 | <8.44 | <8.44 | 1,730 | 410 |
| BLM Lands, Phillips County, MT | BLM081-2MPE | 71 | 10.3 | <6.9 | 18,600 | 5,400 |
| BLM Lands, Phillips County, MT | BLM081MPE | 71.6 | <7.04 | <7.04 | 10,600 | 3,000 |
| BLM Lands, Phillips County, MT | BLM189MPE | 73.8 | <7.63 | <7.63 | 382 | 100 |
| BLM Lands, Phillips County, MT | BLMB026MPE | 76.6 | 12.8 | <8.55 | 303 | 71 |
| BLM Lands, Phillips County, MT | BLMS014MPE | 76.1 | 16.7 | 25.1 | 134 | 32 |
| BLM Lands, Phillips County, MT | BLMS020MPE | 71.6 | 10.6 | <7.04 | 151 | 43 |
| Charles M. Russell NWR, MT | CMRNMCMPPE06 | 74.8 | <7.94 | <7.94 | 19.8 | 5 |
| Near Antelope Coal Mine, WY | MPAMPE01 | 76.7 | 8.58 | <8.58 | 1,160 | 270 |
| Near Antelope Coal Mine, WY | MPAMPE02 | 77.4 | <8.85 | 8.85 | 111 | 25 |
| Foote Creek Rim, WY | MPFCPE01 | 76.2 | 8.4 | <8.40 | 298 | 71 |
| Foote Creek Rim, WY | MPFCPE02 | 78 | <9.09 | <9.09 | 4,090 | 900 |
| Foote Creek Rim, WY | MPFCPE03 | 71.6 | <7.04 | <7.04 | 3,270 | 930 |
| Foote Creek Rim, WY | MPFCPE04 | 77.8 | <9.01 | <9.01 | 45 | 10 |
| Foote Creek Rim, WY | MPFCPE05 | 80.8 | <10.4 | <10.4 | 323 | 62 |

¹ alpha BHC, alpha chlordane, beta BHC, cis-nonachlor, delta BHC, dieldrin, endrin, gamma BHC, gamma chlordane, heptachlor epoxide, mirex, o,p'-DDD, o,p'-DDE, o,p'-DDT, p,p'-DDD, p,p'-DDT, toxaphene, trans -nonachlor, and total PCBs not detected in any mountain plover egg samples.

² wet weight (ww) results given for DDE for ease of comparison among other studys' results in the discussion section of this report.

Appendix 13. Mountain plover egg content data from breeding sites in Colorado, Montana, and Wyoming, 2006.

| Location | Sample | Whole Egg Weight (g) | Egg Contents Weight (g) | Egg Volume (cc) | Nest Status | Contents Condition | Egg Fertile | Embryo Age (days) |
|--------------------------------|-------------|----------------------|-------------------------|-----------------|-------------|--------------------|-------------|-------------------|
| Pawnee National Grasslands, CO | MPWCPE01 | 16.1 | 14.3 | 22 | Active | Fresh | Yes | 8 |
| Pawnee National Grasslands, CO | MPWCPE02 | 14.9 | 13 | 10 | Active | Fresh | Yes | 4 |
| Pawnee National Grasslands, CO | MPWCPE03 | 14 | 12.3 | 16 | Active | Fresh | Yes | 24 |
| Pawnee National Grasslands, CO | MPWCPE04 | 14.4 | 12.7 | 15.5 | Active | Fresh | Yes | 14 |
| Pawnee National Grasslands, CO | MPWCPE05 | 13.8 | 11.8 | 15.4 | Active | Fresh | Yes | 17 |
| | | | | | | | | |
| BLM Lands, Phillips County, MT | BLM025MPE | 14.56 | 13.04 | 10 | Abandoned | Addled | Yes | 10 |
| BLM Lands, Phillips County, MT | BLM081-2MPE | 13.96 | 11.75 | 10 | Abandoned | Rotten | Unknown | Unknown |
| BLM Lands, Phillips County, MT | BLM081MPE | 14.42 | 11.8 | 10 | Abandoned | Rotten | Unknown | Unknown |
| BLM Lands, Phillips County, MT | BLM189MPE | 12.87 | 10.41 | 10 | Abandoned | Rotten | Unknown | Unknown |
| BLM Lands, Phillips County, MT | BLMB026MPE | 13.55 | 11.8 | 11 | Abandoned | Addled | Yes | 15 |
| BLM Lands, Phillips County, MT | BLMS014MPE | 12.65 | 10.72 | 10 | Abandoned | Addled | Yes | 15 |
| BLM Lands, Phillips County, MT | BLMS020MPE | 14.48 | 10.49 | 10 | Abandoned | Rotten | Unknown | Unknown |
| Charles M. Russell NWR, MT | CMRNMCMPE06 | 15.47 | 13.39 | 11 | Abandoned | Fresh | * | 0 |
| | | | | | | | | |
| Near Antelope Coal Mine, WY | MPAMPE01 | 14.7 | 13.2 | 15 | Active | Fresh | Yes | 9 |
| Near Antelope Coal Mine, WY | MPAMPE02 | 13.7 | 11.9 | 13.5 | Active | Fresh | Yes | 25** |
| Foote Creek Rim, WY | MPFCPE01 | 13.6 | 11.8 | 17 | Active | Fresh | Yes | 21 |
| Foote Creek Rim, WY | MPFCPE02 | 15.1 | 13.2 | 15.5 | Active | Fresh | Yes | 21 |
| Foote Creek Rim, WY | MPFCPE03 | 14.8 | 11.9 | 18 | Active | Fresh | No | 0 |
| Foote Creek Rim, WY | MPFCPE04 | 13.9 | 12.5 | 14 | Active | Fresh | Yes | 25 |
| Foote Creek Rim, WY | MPFCPE05 | 15.3 | 13.4 | 17 | Active | Fresh | Yes | 10 |

*Fertility not noted

**Embryo abnormal – head was abnormally small; upper bill twisted out and up.

Appendix 14. Mountain plover egg measurement data from breeding sites in Colorado, Montana, and Wyoming, 2006.

| Location | Sample | Egg Length (mm) | Egg Width (mm) | Dry Shell Weight (mg) | Thickness Index* |
|--------------------------------|--------------|---------------------------|---------------------------|-----------------------|------------------|
| | | Average of 3 measurements | Average of 3 measurements | | |
| Pawnee National Grasslands, CO | MPWCPE01 | 38.2 | 29.1 | 1300 | 1.17 |
| Pawnee National Grasslands, CO | MPWCPE02 | 36.2 | 28.3 | 1200 | 1.17 |
| Pawnee National Grasslands, CO | MPWCPE03 | 37.8 | 28.1 | 1300 | 1.23 |
| Pawnee National Grasslands, CO | MPWCPE04 | 37.7 | 27.8 | 1400 | 1.33 |
| Pawnee National Grasslands, CO | MPWCPE05 | 36.2 | 27.9 | 1700 | 1.69 |
| | | | | | |
| BLM Lands, Phillips County, MT | BLM025MPE | 38.2 | 27.6 | 1150 | 1.09 |
| BLM Lands, Phillips County, MT | BLM081-2MPE | 38.3 | 28.0 | 1230 | 1.15 |
| BLM Lands, Phillips County, MT | BLM081MPE | 39.6 | 25.6 | 1310 | 1.3 |
| BLM Lands, Phillips County, MT | BLM189MPE | 35.3 | 27.9 | 1250 | 1.27 |
| BLM Lands, Phillips County, MT | BLMB026MPE | 38.8 | 27.2 | 1120 | 1.06 |
| BLM Lands, Phillips County, MT | BLMS014MPE | 37.3 | 27.4 | 1210 | 1.18 |
| BLM Lands, Phillips County, MT | BLMS020MPE | 37.4 | 29.8 | 1390 | 1.25 |
| Charles M. Russell NWR, MT | CMRNMCMPPE06 | 37.5 | 28.1 | 1280 | 1.22 |
| | | | | | |
| Near Antelope Coal Mine, WY | MPAMPE01 | 36.4 | 28.7 | 1200 | 1.15 |
| Near Antelope Coal Mine, WY | MPAMPE02 | 39.4 | 27.5 | 1400 | 1.29 |
| Foote Creek Rim, WY | MPFCPE01 | 36.6 | 27.6 | 1500 | 1.48 |
| Foote Creek Rim, WY | MPFCPE02 | 37.7 | 29.1 | 1600 | 1.46 |
| Foote Creek Rim, WY | MPFCPE03 | 37.6 | 27.7 | 1100 | 1.06 |
| Foote Creek Rim, WY | MPFCPE04 | 37.6 | 28.5 | 1400 | 1.31 |
| Foote Creek Rim, WY | MPFCPE05 | 37.5 | 28.5 | 1600 | 1.5 |

* Thickness Index = eggshell weight (mg) / [length (mm) x width (mm)] (Ratcliffe 1970)

Appendix 15. Mountain plover egg thickness data from breeding sites in Colorado, Montana, and Wyoming, 2006.

| Location | Sample | 1 st Half Eggshell Thickness (um) | 2 nd Eggshell Half Thickness (um) | Overall Thickness (um) | Thickness Index* |
|--------------------------------|--------------|--|--|------------------------|------------------|
| | | Average of 3 measurements | Average of 3 measurements | Average Thickness | |
| Pawnee National Grasslands, CO | MPWCPE01 | 22.8 | 24.5 | 23.63 | 1.17 |
| Pawnee National Grasslands, CO | MPWCPE02 | 25.3 | 22 | 23.63 | 1.17 |
| Pawnee National Grasslands, CO | MPWCPE03 | 27.3 | 24 | 25.63 | 1.23 |
| Pawnee National Grasslands, CO | MPWCPE04 | 24.3 | 23 | 23.63 | 1.33 |
| Pawnee National Grasslands, CO | MPWCPE05 | 22.3 | 22.3 | 22.25 | 1.69 |
| | | | | | |
| BLM Lands, Phillips County, MT | BLM025MPE | 22 | 21.5 | 21.75 | 1.09 |
| BLM Lands, Phillips County, MT | BLM081-2MPE | 22.5 | 21.5 | 22 | 1.15 |
| BLM Lands, Phillips County, MT | BLM081MPE | 21.5 | 21.8 | 21.63 | 1.3 |
| BLM Lands, Phillips County, MT | BLM189MPE | 25 | 24.5 | 24.75 | 1.27 |
| BLM Lands, Phillips County, MT | BLMB026MPE | 23.5 | 21.5 | 22.5 | 1.06 |
| BLM Lands, Phillips County, MT | BLMS014MPE | 23.3 | 24.3 | 23.75 | 1.18 |
| BLM Lands, Phillips County, MT | BLMS020MPE | 23.8 | 25 | 24.38 | 1.25 |
| Charles M. Russell NWR, MT | CMRNMCMPPE06 | 21.5 | 21.3 | 21.38 | 1.22 |
| | | | | | |
| Near Antelope Coal Mine, WY | MPAMPE01 | 24.8 | 24.5 | 24.6 | 1.15 |
| Near Antelope Coal Mine, WY | MPAMPE02 | 20.3 | 20.5 | 20.38 | 1.29 |
| Foote Creek Rim, WY | MPFCPE01 | 22.8 | 22.5 | 22.63 | 1.48 |
| Foote Creek Rim, WY | MPFCPE02 | 23 | 23 | 23 | 1.46 |
| Foote Creek Rim, WY | MPFCPE03 | 20.5 | 24.5 | 22.5 | 1.06 |
| Foote Creek Rim, WY | MPFCPE04 | 23.3 | 21.8 | 22.5 | 1.31 |
| Foote Creek Rim, WY | MPFCPE05 | 23.3 | 24.5 | 23.88 | 1.5 |

* Thickness Index = eggshell weight (mg) / [length (mm) x width (mm)] (Ratcliffe 1970)

Appendix 16. Concentrations of organic pesticides, PCBs, and trifluralin* (ng/g dry weight) in eggs samples collected from mountain plover breeding sites in Colorado and Wyoming, 2007.

| Location | Sample | Moisture % | alpha BHC ng/g dw | dieldrin ng/g dw | HCB ng/g dw | heptachlor epoxide ng/g dw | o,p'-DDT ng/g dw | oxychlorane ng/g dw | p,p'-DDE ng/g dw |
|--------------------------------|----------|------------|-------------------|------------------|-------------|----------------------------|------------------|---------------------|------------------|
| Pawnee National Grasslands, CO | MPWCPE06 | 95.2 | <9.08 | 70.4 | <9.08 | 20.2 | 35.2 | 26.5 | 1,070,000 |
| Pawnee National Grasslands, CO | MPWCPE07 | 71.3 | 1.97 | <1.47 | <1.47 | 3.56 | <1.47 | 9.48 | 2,730 |
| Pawnee National Grasslands, CO | MPWCPE08 | 95.1 | 21.2 | <9.13 | <9.13 | 17.4 | <9.13 | 12.9 | 1,230 |
| Pawnee National Grasslands, CO | MPWCPE09 | 94.7 | 13.2 | <9.27 | <9.27 | <9.27 | <9.27 | 15.8 | 4,030 |
| Pawnee National Grasslands, CO | MPWCPE10 | 94.5 | <8.36 | <8.36 | <8.36 | <8.36 | <8.36 | 54.7 | 638 |
| | | | | | | | | | |
| Foote Creek Rim, WY | MPFCPE06 | 71.5 | <1.74 | <1.74 | 6.23 | <1.74 | <1.74 | 9.24 | 262 |

DRY WEIGHT / FRESH WEIGHT CONVERSION FOR ADDLED EGGS WITH >75% MOISTURE = 0.325 (fw concentrations will be approximately 6x higher than wet weight concentrations for eggs with 95% moisture - assuming the high moisture content is due to the addition of rinse water)

Appendix 16. cont.

| Location | Sample | 1,2,3,4-Tetrachlorobenzene ng/g dw | trans-nonachlor ng/g dw | Cl4-PCB ng/g dw | Cl6-PCB ng/g dw | Cl7-PCB ng/g dw | PCB-TOTAL ng/g dw |
|--------------------------------|----------|------------------------------------|-------------------------|-----------------|-----------------|-----------------|-------------------|
| Pawnee National Grasslands, CO | MPWCPE06 | <9.08 | <9.08 | <182 | 300 | <182 | 701 |
| Pawnee National Grasslands, CO | MPWCPE07 | <1.47 | <1.47 | 38.1 | <29.5 | <29.5 | 48 |
| Pawnee National Grasslands, CO | MPWCPE08 | 83.1 | 13.5 | 217 | <183 | <183 | 595 |
| Pawnee National Grasslands, CO | MPWCPE09 | <9.27 | <9.27 | 202 | <185 | <185 | 251 |
| Pawnee National Grasslands, CO | MPWCPE10 | <8.36 | <8.36 | <167 | <167 | <167 | <167 |
| | | | | | | | |
| Foote Creek Rim, WY | MPFCPE06 | <1.74 | <1.74 | <34.8 | 50.8 | 58.7 | 139 |

*aldrin, alpha chlordane, beta BHC, chlorpyrifos, cis-nonachlor, delta BHC, endosulfan II, endrin, gamma BHC, gamma chlordane, heptachlor, mirex, o,p'-DDD, o,p'-DDE, p,p'-DDD, p,p'-DDT, pentachloro-anisole, toxaphene, trans-nonachlor, Cl11-PCB, Cl10-PCB, Cl2-PCB, Cl3-PCB, Cl5-PCB, Cl8-PCB, Cl9-PCB, 1,2,4,5-tetrachlorobenzene, trifluralin, trifluralin #10, trifluralin #11, trifluralin #12, trifluralin #13, trifluralin #14, trifluralin #15, trifluralin #17, trifluralin #2, trifluralin #20, trifluralin #21, trifluralin #3, trifluralin #4, trifluralin #5, and trifluralin #7 not detected in any plover egg samples.

Appendix 17. Mountain plover egg content data from breeding sites in Colorado and Wyoming, 2007.

| Location | Sample | Whole Egg Weight (g) | Egg Contents Weight (g) | Egg Volume (cc) | Nest Status | Contents Condition | Egg Fertile | Embryo Age (days) |
|--------------------------------|----------|----------------------|-------------------------|-----------------|-------------|--------------------|-------------|-------------------|
| Pawnee National Grasslands, CO | MPWCPE06 | 15.1 | 12.8 | 14 | Active | Fresh | Yes | 1 |
| Pawnee National Grasslands, CO | MPWCPE07 | 14.5 | 10.8 | 17 | Active | Fresh | No | 0 |
| Pawnee National Grasslands, CO | MPWCPE08 | 13.2 | 11.7 | 16 | Active | Fresh | Yes | 8 |
| Pawnee National Grasslands, CO | MPWCPE09 | 14.8 | 11.4 | 15 | Active | Fresh | Yes | 7 |
| Pawnee National Grasslands, CO | MPWCPE10 | 15.8 | 11.9 | 15 | Active | Fresh | Yes | 7 |
| | | | | | | | | |
| Foote Creek Rim, WY | MPFCPE06 | 14.7 | 11.8 | 14 | Active | Fresh | No | 0 |

Appendix 18. Mountain plover egg measurement data from breeding sites in Colorado and Wyoming, 2007.

| Location | Sample | Egg Length (mm) | Egg Width (mm) | Overall Thickness (um) | Thickness Index* |
|--------------------------------|----------|---------------------------|---------------------------|------------------------|------------------|
| | | Average of 3 measurements | Average of 3 measurements | Average Thickness | |
| Pawnee National Grasslands, CO | MPWCPE06 | 37.8 | 28.2 | 23.75 | 1.31 |
| Pawnee National Grasslands, CO | MPWCPE07 | 37.7 | 28.2 | 21.19 | 1.22 |
| Pawnee National Grasslands, CO | MPWCPE08 | 36.4 | 27.1 | 19.38 | 0.913 |
| Pawnee National Grasslands, CO | MPWCPE09 | 37.7 | 28.3 | 22.63 | 1.13 |
| Pawnee National Grasslands, CO | MPWCPE10 | 38.3 | 29.0 | 21.44 | 1.08 |
| | | | | | |
| Foote Creek Rim, WY | MPFCPE06 | 37.2 | 28.2 | 20.69 | 1.24 |

* Thickness Index = eggshell weight (mg) / [length (mm) x width (mm)] (Ratcliffe 1970)

Appendix 19. Mountain plover egg thickness data from breeding sites in Colorado and Wyoming, 2007.

| Location | Sample | 1 st Half Eggshell Thickness (um) | 2 nd Eggshell Half Thickness (um) | Overall Thickness (um) | Thickness Index* |
|--------------------------------|----------|--|--|------------------------|------------------|
| | | Average of 3 measurements | Average of 3 measurements | Average Thickness | |
| Pawnee National Grasslands, CO | MPWCP06 | 24.5 | 23 | 23.75 | 1.31 |
| Pawnee National Grasslands, CO | MPWCPE07 | 21.6 | 20.8 | 21.19 | 1.33 |
| Pawnee National Grasslands, CO | MPWCPE08 | 18.9 | 19.9 | 19.38 | 0.913 |
| Pawnee National Grasslands, CO | MPWCPE09 | 23.9 | 21.4 | 22.63 | 1.13 |
| Pawnee National Grasslands, CO | MPWCPE10 | 22.4 | 20.5 | 21.44 | 1.08 |
| | | | | | |
| Foote Creek Rim, WY | MPFCPE06 | 20.1 | 21.3 | 21.69 | 1.24 |

* Thickness Index = eggshell weight (mg) / [length (mm) x width (mm)] (Ratcliffe 1970)

Appendix 20. Concentrations of organic pesticides, PCBs, and trifluralin* (ng/g dry weight) in eggs samples collected from mountain plover breeding sites in Montana, 2008.

| Location | Sample | Moisture % | alpha BHC ng/g dw | alpha chlordane ng/g dw | beta BHC ng/g dw | chlorpyrifos ng/g dw | dieldrin ng/g dw | endosulfan II ng/g dw |
|--------------------------------|------------|------------|-------------------|-------------------------|------------------|----------------------|------------------|-----------------------|
| BLM Lands, Phillips County, MT | B148P033E1 | 70.8 | 3.68 | <1.65 | <1.65 | <1.65 | <1.65 | <1.65 |
| BLM Lands, Phillips County, MT | B41P025E1 | 71 | 4.22 | <1.67 | <1.67 | <1.67 | <1.67 | <1.67 |
| BLM Lands, Phillips County, MT | B41P050E1 | 69.6 | 2.07 | <1.46 | <1.46 | <1.46 | <1.46 | <1.46 |
| BLM Lands, Phillips County, MT | B45P009E1 | 71.9 | 2.33 | <1.62 | <1.62 | <1.62 | <1.62 | <1.62 |
| BLM Lands, Phillips County, MT | B72P017E1 | 64.9 | 2.07 | 1.79 | <1.38 | 3.01 | 7.59 | 6.18 |
| BLM Lands, Phillips County, MT | S1001E1 | 71.5 | <1.72 | <1.72 | <1.72 | 3.15 | 5.33 | <1.72 |
| BLM Lands, Phillips County, MT | S10P001E1 | 66.1 | <1.54 | <1.54 | 1.43 | 10.4 | 130 | <1.54 |
| BLM Lands, Phillips County, MT | S10P001E2 | 71.2 | <1.59 | <1.59 | 1.28 | 12.8 | 125 | <1.59 |

Appendix 20. cont.

| Location | Sample | endrin ng/g dw | gamma chlordane ng/g dw | HCB ng/g dw | heptachlor ng/g dw | heptachlor epoxide ng/g dw | mirex ng/g dw | o,p'-DDD ng/g dw |
|---------------------------------|------------|----------------|-------------------------|-------------|--------------------|----------------------------|---------------|------------------|
| BLM Lands, Phillips County, MT | B148P033E1 | <1.65 | <1.65 | 14.4 | <1.65 | <1.65 | 2.04 | <1.65 |
| BLM Lands, Phillips County, MT | B41P025E1 | <1.67 | <1.67 | 6.58 | <1.67 | <1.67 | <1.67 | <1.67 |
| BLM Lands, Phillips County, MT | B41P050E1 | <1.46 | <1.46 | 4.43 | <1.46 | <1.46 | <1.46 | <1.46 |
| 4BLM Lands, Phillips County, MT | B45P009E1 | <1.62 | <1.62 | 6.46 | 2.63 | <1.62 | <1.62 | <1.62 |
| BLM Lands, Phillips County, MT | B72P017E1 | 3.53 | <1.38 | 5.93 | <1.38 | <1.38 | <1.38 | 2.44 |
| BLM Lands, Phillips County, MT | S1001E1 | 1.84 | <1.72 | 7.84 | <1.72 | <1.72 | <1.72 | <1.72 |
| BLM Lands, Phillips County, MT | S10P001E1 | 88.5 | 2.5 | 32.4 | <1.54 | 9.84 | 12.6 | <1.54 |
| BLM Lands, Phillips County, MT | S10P001E2 | 139 | <1.59 | 19.8 | <1.59 | 15.4 | 4.43 | <1.59 |

Appendix 20. cont.

| Location | Sample | o,p'-DDE ng/g dw | o,p'-DDT ng/g dw | oxychlorane ng/g dw | p,p'-DDD ng/g dw | p,p'-DDE ng/g dw | p,p'-DDT ng/g dw | pentachloro- anisole ng/g dw |
|--------------------------------|------------|---------------------|---------------------|------------------------|---------------------|---------------------|---------------------|------------------------------------|
| BLM Lands, Phillips County, MT | B148P033E1 | <1.65 | <1.65 | 7.69 | <1.65 | 110 | <1.65 | <1.65 |
| BLM Lands, Phillips County, MT | B41P025E1 | <1.67 | <1.67 | <1.67 | <1.67 | <1.67 | <1.67 | 1.72 |
| BLM Lands, Phillips County, MT | B41P050E1 | <1.46 | <1.46 | 1.79 | <1.46 | 220 | <1.46 | 1.52 |
| BLM Lands, Phillips County, MT | B45P009E1 | <1.62 | <1.62 | <1.62 | <1.62 | 58.2 | <1.62 | 3.54 |
| BLM Lands, Phillips County, MT | B72P017E1 | <1.38 | <1.38 | <1.38 | <1.38 | 16.4 | 3.3 | <1.38 |
| BLM Lands, Phillips County, MT | S1001E1 | <1.72 | <1.72 | 111 | 2.64 | 261 | <1.72 | <1.72 |
| BLM Lands, Phillips County, MT | S10P001E1 | 33.2 | <1.54 | 18.7 | 320 | 140,000 | 57.6 | <1.54 |
| BLM Lands, Phillips County, MT | S10P001E2 | <1.59 | 6.67 | 10.1 | 300 | 185,000 | 26 | 2.31 |

Appendix 20. cont.

| Location | Sample | 1,2,3,4- Tetrachloro- benzene ng/g dw | 1,2,4,5- Tetrachloro- benzene ng/g dw | trans- nonachlor ng/g dw | C14- PCB ng/g dw | C15- PCB ng/g dw | C16- PCB ng/g dw | C17- PCB ng/g dw | PCB TOTAL ng/g dw |
|--------------------------------|------------|--|--|--------------------------------|------------------------|------------------------|------------------------|------------------------|-------------------------|
| BLM Lands, Phillips County, MT | B148P033E1 | <1.65 | 15.7 | <1.65 | <32.9 | 43 | <32.9 | 389 | 477 |
| BLM Lands, Phillips County, MT | B41P025E1 | <1.67 | <1.67 | <1.67 | <33.5 | 42.4 | <33.5 | 365 | 424 |
| BLM Lands, Phillips County, MT | B41P050E1 | 7.59 | 19.3 | <1.46 | <29.1 | 41.2 | <29.1 | 413 | 490 |
| BLM Lands, Phillips County, MT | B45P009E1 | 9.38 | <1.62 | <1.62 | <32.4 | <32.4 | <32.4 | <32.4 | 75.9 |
| BLM Lands, Phillips County, MT | B72P017E1 | <1.38 | 20.4 | <1.38 | <27.7 | 28 | <27.7 | <27.7 | 96.9 |
| BLM Lands, Phillips County, MT | S1001E1 | <1.72 | <1.72 | <1.72 | <34.4 | <34.4 | 39.4 | <34.4 | 83.8 |
| BLM Lands, Phillips County, MT | S10P001E1 | 8.87 | <1.54 | 2.99 | 133 | 225 | 517 | 188 | 1,110 |
| BLM Lands, Phillips County, MT | S10P001E2 | 6.17 | <1.59 | <1.59 | 66 | 2,020 | 459 | 193 | 2,760 |

*aldrin, cis-nonachlor, delta BHC, gamma BHC, toxaphene, C11-PCB, C110-PCB, C12-PCB, C13-PCB, C18-PCB, C19-PCB, trifluralin, trifluralin #10, trifluralin #11, trifluralin #12, trifluralin #13, trifluralin #14, trifluralin #15, trifluralin #17, trifluralin #2, trifluralin #20, trifluralin #21, trifluralin #3, trifluralin #4, trifluralin #5, and trifluralin #7 not detected in any plover egg samples.

Appendix 21. Mountain plover egg content data from breeding sites in Montana, 2008.

| Location | Sample | Whole Egg Weight (g) | Egg Contents Weight (g) | Egg Volume (cc) | Nest Status | Contents Condition | Egg Fertile | Embryo Age (days) |
|--------------------------------|------------|----------------------|-------------------------|-----------------|-------------|--------------------|-------------|-------------------|
| BLM Lands, Phillips County, MT | B148P033E1 | * | 7.77 | | Abandoned | Rotten | unknown | unknown |
| BLM Lands, Phillips County, MT | B41P025E1 | * | 10.92 | | Abandoned | Rotten | unknown | unknown |
| BLM Lands, Phillips County, MT | B41P050E1 | * | 11 | | Abandoned | Rotten | unknown | unknown |
| BLM Lands, Phillips County, MT | B45P009E1 | * | 11.43 | | Abandoned | Rotten | unknown | unknown |
| BLM Lands, Phillips County, MT | B72P017E1 | * | 6.81 | | Abandoned | Rotten | unknown | unknown |
| BLM Lands, Phillips County, MT | S1001E1 | * | ** | | Abandoned | Rotten | unknown | unknown |
| BLM Lands, Phillips County, MT | S10P001E1 | * | ** | | Abandoned | Rotten | Yes | 15 |
| BLM Lands, Phillips County, MT | S10P001E2 | * | ** | | Abandoned | Rotten | Yes | 15 |

*Egg collected cracked, no measurements possible without danger of breaking the egg

**weight entered onto data sheet incorrectly

Appendix 22. Mountain plover egg measurement data from breeding sites in Montana, 2008.

| Location | Sample | Egg Length (mm) | Egg Width (mm) | Overall Thickness (um) | Thickness Index+ |
|--------------------------------|------------|---------------------------|---------------------------|------------------------|------------------|
| | | Average of 3 measurements | Average of 3 measurements | Average Thickness | |
| BLM Lands, Phillips County, MT | B148P033E1 | * | * | 23.125 | * |
| BLM Lands, Phillips County, MT | B41P025E1 | * | * | 24 | * |
| BLM Lands, Phillips County, MT | B41P050E1 | * | * | 20.375 | * |
| BLM Lands, Phillips County, MT | B45P009E1 | * | * | 23.5 | * |
| BLM Lands, Phillips County, MT | B72P017E1 | * | * | 19.375 | * |
| BLM Lands, Phillips County, MT | S1001E1 | * | * | 21.875 | * |
| BLM Lands, Phillips County, MT | S10P001E1 | * | * | 20.625 | * |
| BLM Lands, Phillips County, MT | S10P001E2 | * | * | 21.75 | * |

+ Thickness Index = eggshell weight (mg) / [length (mm) x width (mm)] (Ratcliffe 1970)

*Egg collected cracked, no measurements possible without danger of breaking the egg

Appendix 23. Mountain plover egg thickness data from breeding sites in Montana, 2008.

| Location | Sample | 1 st Half Eggshell Thickness (um) | 2 nd Eggshell Half Thickness (um) | Overall Thickness (um) | Thickness Index+ |
|-----------------------------|------------|--|--|------------------------|------------------|
| | | Average of 3 measurements | Average of 3 measurements | Average Thickness | |
| BLM Lands, Phillips County, | B148P033E1 | 24 | 22.25 | 23.125 | * |
| BLM Lands, Phillips County, | B41P025E1 | 24.75 | 23.25 | 24 | * |
| BLM Lands, Phillips County, | B41P050E1 | 21 | 19.75 | 20.375 | * |
| BLM Lands, Phillips County, | B45P009E1 | 24 | 23 | 23.5 | * |
| BLM Lands, Phillips County, | B72P017E1 | 20.25 | 18.5 | 19.375 | * |
| BLM Lands, Phillips County, | S1001E1 | 22 | 21.75 | 21.875 | * |
| BLM Lands, Phillips County, | S10P001E1 | 20.25 | 21 | 20.625 | * |
| BLM Lands, Phillips County, | S10P001E2 | 22 | 21.5 | 21.75 | * |

+ Thickness Index = eggshell weight (mg) / [length (mm) x width (mm)] (Ratcliffe 1970)

*Egg collected cracked, no measurements possible without danger of breaking the egg

Appendix 24. Mountain plover egg content data from breeding sites in Wyoming, 2003.

| Location | Sample | Egg Contents Weight (g) | Egg Volume (cc) | Contents Condition | Egg Fertile |
|----------------------------------|----------|-------------------------|-----------------|---------------------------------|-------------|
| Albany County, WY | MPACPE01 | 11.2 | 10 | Egg cracked, embryonated, early | Yes |
| Park County, WY | MPCDPE01 | 17.3 | 16 | Egg good, embryonated, early | Yes |
| Park County, WY | MPCDPE02 | 16.2 | 15 | Egg good, embryonated, early | Yes |
| Park County, WY | MPCDPE03 | 17.8 | 16 | Egg rotten, embryonated, early | Yes |
| Park County, WY | MPCDPE04 | 17.9 | 15 | Egg rotten | Unknown |
| Park County, WY | MPCDPE05 | 15.6 | 13 | Egg rotten | Unknown |
| Park County, WY | MPCDPE06 | 16.3 | 14 | Egg rotten | Unknown |
| Park County, WY | MPCDPE07 | 15 | 12 | Egg cracked, embryonated, late | Yes |
| Shirley Basin, Carbon County, WY | MPSBPE01 | 9.2 | 11 | Egg good, embryonated, late | Yes |
| Shirley Basin, Carbon County, WY | MPSBPE02 | 11.6 | 13 | Egg good, embryonated, late | Yes |
| Shirley Basin, Carbon County, WY | MPSBPE03 | 11.2 | 13 | Egg good, embryonated, late | Yes |
| Shirley Basin, Carbon County, WY | MPSBPE04 | 17.7 | 18 | Egg good, embryonated, mid | Yes |
| Shirley Basin, Carbon County, WY | MPSBPE05 | 13.1 | 13 | Egg good, infertile | No |

Appendix 25. Concentrations of organic pesticides, PCBs, and trifluralin* (ng/g dry weight) in eggs samples collected from mountain plover breeding sites in Wyoming, 2003.

| Location | Sample | Moisture % | alpha BHC ng/g dw | alpha chlordane ng/g dw | beta BHC ng/g dw | cis-nonachlor ng/g dw | dieldrin ng/g dw | endrin ng/g dw | HCB ng/g dw |
|----------------------------------|----------|------------|-------------------|-------------------------|------------------|-----------------------|------------------|----------------|-------------|
| Albany County, WY | MPACPE01 | 69.4 | 3.29 | <1.54 | <1.54 | <1.54 | <1.54 | <1.54 | 6.26 |
| Park County, WY | MPCDPE01 | 70.3 | 1.94 | <1.59 | <1.59 | <1.59 | <1.59 | <1.59 | 7.33 |
| Park County, WY | MPCDPE02 | 71.3 | 1.7 | <1.61 | <1.61 | <1.61 | 17.3 | <1.61 | 7.24 |
| Park County, WY | MPCDPE03 | 70 | 2.35 | 2.11 | 1.96 | <1.52 | 5.67 | <1.52 | 6.45 |
| Park County, WY | MPCDPE04 | 70 | 1.8 | <1.62 | <1.62 | <1.62 | 6.05 | <1.62 | 6.32 |
| Park County, WY | MPCDPE05 | 65.5 | <1.41 | <1.41 | <1.41 | <1.41 | 31.4 | <1.41 | 15.6 |
| Park County, WY | MPCDPE06 | 67.2 | <1.25 | <1.25 | <1.25 | 2.08 | 19.5 | <1.25 | 8.83 |
| Park County, WY | MPCDPE07 | 67.5 | <1.25 | <1.25 | <1.25 | <1.25 | 4.37 | <1.25 | 6.23 |
| Shirley Basin, Carbon County, WY | MPSBPE01 | 74.4 | 1.78 | <1.7 | <1.7 | <1.7 | <1.7 | 3.85 | 5.77 |
| Shirley Basin, Carbon County, WY | MPSBPE02 | 73.1 | <1.84 | <1.84 | <1.84 | <1.84 | <1.84 | <1.84 | 5.83 |
| Shirley Basin, Carbon County, WY | MPSBPE03 | 73.7 | 2.22 | <1.78 | <1.78 | <1.78 | <1.78 | 2.09 | 6.44 |
| Shirley Basin, Carbon County, WY | MPSBPE04 | 72.5 | 4.45 | 4.06 | 77.5 | <1.71 | 23.3 | 12.7 | 9.9 |
| Shirley Basin, Carbon County, WY | MPSBPE05 | 70.4 | 2.61 | <1.67 | <1.67 | <1.67 | 3.1 | <1.67 | 4.29 |

Appendix 25. cont.

| Location | Sample | heptachlor epoxide ng/g dw | o,p'-DDT ng/g dw | oxychlordane ng/g dw | p,p'-DDD ng/g dw | p,p'-DDE ng/g dw | p,p'-DDT ng/g dw | pentachloro-anisole ng/g dw | 1,2,3,4-Tetrachloro-benzene ng/g dw |
|----------------------------------|----------|-------------------------------|---------------------|-------------------------|---------------------|---------------------|---------------------|--------------------------------|--|
| Albany County, WY | MPACPE01 | 2.53 | <1.54 | 12.8 | <1.54 | 1,960 | <1.54 | <1.54 | <1.54 |
| Park County, WY | MPCDPE01 | 14.4 | <1.59 | 17.7 | <1.59 | 4,810 | <1.59 | 2.06 | 3.58 |
| Park County, WY | MPCDPE02 | 19.6 | <1.61 | 12.3 | <1.61 | 10,800 | <1.61 | <1.61 | 18.4 |
| Park County, WY | MPCDPE03 | 8.75 | <1.52 | 11 | <1.52 | 4,150 | 3.86 | <1.52 | 7.92 |
| Park County, WY | MPCDPE04 | 8.28 | <1.62 | 14.4 | <1.62 | 4,220 | 4.05 | <1.62 | 8.38 |
| Park County, WY | MPCDPE05 | 48 | <1.41 | 28.5 | <1.41 | 17,400 | <1.41 | <1.41 | 40.8 |
| Park County, WY | MPCDPE06 | 26.8 | <1.25 | 20.1 | <1.25 | 10,000 | <1.25 | 1.85 | 9.17 |
| Park County, WY | MPCDPE07 | 2.37 | <1.25 | 12.4 | <1.25 | 2,210 | <1.25 | <1.25 | 3.63 |
| Shirley Basin, Carbon County, WY | MPSBPE01 | 3.27 | <1.7 | 6.79 | <1.7 | 149 | <1.7 | <1.7 | <1.7 |
| Shirley Basin, Carbon County, WY | MPSBPE02 | 2.79 | <1.84 | 6.55 | <1.84 | 1,330 | <1.84 | 2.49 | <1.84 |
| Shirley Basin, Carbon County, WY | MPSBPE03 | 2.09 | <1.78 | 6.67 | <1.78 | 139 | <1.78 | <1.78 | <1.78 |
| Shirley Basin, Carbon County, WY | MPSBPE04 | 6.88 | 8.41 | 24.1 | 31.6 | 292,000 | 3.79 | <1.71 | <1.71 |
| Shirley Basin, Carbon County, WY | MPSBPE05 | 1.87 | <1.67 | 10.4 | <1.67 | 1,830 | <1.67 | <1.67 | <1.67 |

Appendix 25. cont.

| Location | Sample | 1,2,4,5-Tetrachloro-benzene ng/g dw | trans-nonachlor ng/g dw | C14-PCB ng/g dw | C15-PCB ng/g dw | C16-PCB ng/g dw | C17-PCB ng/g dw | C19-PCB ng/g dw | PCB-TOTAL ng/g dw |
|----------------------------------|----------|--|----------------------------|--------------------|--------------------|--------------------|--------------------|--------------------|----------------------|
| Albany County, WY | MPACPE01 | <1.54 | <1.54 | <30.9 | <30.9 | <30.9 | <30.9 | <30.9 | 53.3 |
| Park County, WY | MPCDPE01 | 6.6 | <1.59 | 73.8 | <31.7 | <31.7 | <31.7 | <31.7 | 139 |
| Park County, WY | MPCDPE02 | 28.2 | 2.13 | <32.2 | <32.2 | 67.5 | 49.9 | 36.3 | 180 |
| Park County, WY | MPCDPE03 | 11.1 | <1.52 | 31.9 | <30.3 | 35.9 | 32.3 | <30.3 | 119 |
| Park County, WY | MPCDPE04 | 12.7 | <1.62 | 49.5 | <32.3 | 43.9 | 36.4 | <32.3 | 156 |
| Park County, WY | MPCDPE05 | 61.6 | 2.61 | <28.1 | <28.1 | 171 | 134 | 43.5 | 400 |
| Park County, WY | MPCDPE06 | 25.7 | 1.35 | 107 | <25 | 74.8 | 65.1 | <25 | 269 |
| Park County, WY | MPCDPE07 | <1.25 | 1.66 | <25 | <25 | <25 | <25 | <25 | 51.6 |
| Shirley Basin, Carbon County, WY | MPSBPE01 | <1.7 | <1.7 | <33.9 | <33.9 | <33.9 | <33.9 | <33.9 | <33.9 |
| Shirley Basin, Carbon County, WY | MPSBPE02 | <1.84 | <1.84 | <36.8 | <36.8 | <36.8 | <36.8 | <36.8 | <36.8 |
| Shirley Basin, Carbon County, WY | MPSBPE03 | <1.78 | <1.78 | <35.5 | <35.5 | <35.5 | <35.5 | <35.5 | 41.4 |
| Shirley Basin, Carbon County, WY | MPSBPE04 | <1.71 | 3.53 | 39.3 | 45.8 | 154 | 49.9 | <34.3 | 299 |
| Shirley Basin, Carbon County, WY | MPSBPE05 | <1.67 | <1.67 | <33.4 | <33.4 | <33.4 | <33.4 | <33.4 | 39.8 |

*aldrin, chlorpyrifos, delta BHC, endosulfan II, gamma BHC, gamma chlordane, heptachlor, mirex, o,p'-DDD, o,p'-DDE, toxaphene, trans-nonachlor, C11-PCB, C10-PCB, C12-PCB, C13-PCB, C18-PCB, trifluralin, trifluralin #10, trifluralin #11, trifluralin #12, trifluralin #13, trifluralin #14, trifluralin #15, trifluralin #17, trifluralin #2, trifluralin #20, trifluralin #21, trifluralin #3, trifluralin #4, trifluralin #5, and trifluralin #7 not detected in any plover egg samples.

Appendix 26. Concentrations of organic pesticides (ng/g dry weight) in egg samples* collected from mountain plover breeding sites in Montana and Colorado, 2003

| Location | Sample | Lipid % | Moisture % | o,p'-DDD ng/g dw | o,p'-DDE ng/g dw | o,p'-DDT ng/g dw | p,p'-DDD ng/g dw | p,p'-DDE ng/g dw | p,p'-DDT ng/g dw |
|------------------------|----------|---------|------------|------------------|------------------|------------------|------------------|------------------|------------------|
| Park County (MD2), CO | COPA0201 | 17.1 | 62.6 | <13.4 | <14.6 | <14.6 | <13.4 | 115,000 | <13.4 |
| Park County (HS7), CO | COPA0701 | 14.1 | 69.1 | <16.2 | <13.6 | <13.6 | <16.2 | 110 | <16.2 |
| Park County (MD8), CO | COPA0801 | 13.5 | 67.5 | <15.4 | <15.2 | <15.2 | <15.4 | 3,660 | <15.4 |
| Park County (BN9), CO | COPA0901 | 10.8 | 69.7 | <16.5 | <18 | <18 | <16.5 | 1,620 | <16.5 |
| Park County (BS10), CO | COPA1001 | 23.1 | 60.4 | <12.6 | <16.7 | <16.7 | <12.6 | 2,580 | <12.6 |
| Park County (HS12), CO | COPA1201 | 15.6 | 65.8 | <14.6 | <14.5 | <14.5 | <14.6 | 439 | <14.6 |
| Park County (HS14), CO | COPA1401 | 18.1 | 63.1 | <13.6 | <17.4 | <17.4 | <13.6 | 325 | <13.6 |
| Park County (FM20), CO | COPA2001 | 16.9 | 67.1 | <15.2 | <18.3 | <18.3 | <15.2 | 116 | <15.2 |
| Park County (FM3), CO | COPAM301 | 11.4 | 72.2 | <18 | <15.7 | <15.7 | <18 | 201 | <18 |
| Park County (HS3), CO | COPAS301 | 12.8 | 70.1 | <16.7 | <14 | <14 | <16.7 | 177 | <16.7 |
| South Park County, CO | COSP0100 | 16 | 65.4 | <14.5 | <14.7 | <14.7 | <14.5 | 636 | <14.5 |
| South Park County, CO | COSP1020 | 11.4 | 71.3 | <17.4 | <14.4 | <14.4 | <17.4 | 143 | <17.4 |
| South Park County, CO | COSP2700 | 7.96 | 72.7 | <18.3 | <15 | <15 | <18.3 | 117 | <18.3 |
| South Park County, CO | COSP3700 | 15.1 | 68.2 | <15.7 | <15.5 | <15.5 | <15.7 | 50.3 | <15.7 |
| South Park County, CO | COSP4100 | 14.9 | 64.2 | <14 | <18.9 | <18.9 | <14 | 20,500 | <14 |
| South Park County, CO | COSP4400 | 14.8 | 66 | <14.7 | <14.6 | <14.6 | <14.7 | 7,320 | <14.7 |
| Weld County, CO | COWE0600 | 14.1 | 65.2 | <14.4 | <13.6 | <13.6 | <14.4 | 8,070 | <14.4 |
| Weld County, CO | COWE2002 | 16.2 | 66.6 | <15 | <15.2 | <15.2 | <15 | 111 | <15 |
| Weld County, CO | COWE3702 | 13.5 | 67.8 | <15.5 | <18 | <18 | <15.5 | 106 | <15.5 |
| Weld County, CO | COWE5002 | 13.1 | 73.6 | <18.9 | <16.7 | <16.7 | <18.9 | 30.3 | <18.9 |
| | | | | | | | | | |
| Phillips County, MT | MTPH2099 | 14.9 | 69 | <16.1 | <16.1 | <16.1 | <16.1 | 11,100 | <16.1 |
| Phillips County, MT | MTPH3100 | 12.9 | 72.7 | <18.3 | <18.3 | <18.3 | <18.3 | 22 | <18.3 |
| Phillips County, MT | MTPH3799 | 10.1 | 73.7 | <19 | <19 | <19 | <19 | <19 | <19 |
| Phillips County, MT | MTPH6300 | 12.9 | 71.2 | <17.4 | <17.4 | <17.4 | <17.4 | 2,260 | <17.4 |

*Contents of all eggs were fresh but the fertility and age of the embryo were unknown. Eggs were not collected as part of this study and the quality of the sampling effort cannot be validated.