

**ENDANGERED, THREATENED, PROPOSED, AND CANDIDATE SPECIES**  
**U.S. Fish and Wildlife Service**  
**Region 3 (Illinois, Indiana, Iowa, Michigan, Missouri, Minnesota, Ohio, Wisconsin)**  
**October 1, 2016**

**MAMMALS**

Canada lynx (*Lynx canadensis*) Lead Region 6

Status: Threatened

Critical Habitat designated

Habitat: Mix of evergreens and hardwoods, such as maple and birch; interspersed of mature and young forest

Range: CO, ID, ME, MI, MN, MT, NH, NY, OR, UT, VT, WA, WI, WY

Gray bat (*Myotis grisescens*) Lead: Region 3

Status: Endangered

Habitat: Caves

Range: Alabama, Arkansas, Florida, Georgia, Illinois, Indiana, Kansas, Kentucky, Mississippi, Missouri, North Carolina, Oklahoma, Tennessee, Virginia, West Virginia

Gray wolf (*Canis lupus*) Lead: Region 3

Status:

Western Great Lakes DPS - - Relisted Dec. 19, 2014: Threatened in Minnesota and Endangered in the remainder of the DPS

Minnesota, Wisconsin, and Michigan and portions of North Dakota, South Dakota, Iowa, Illinois, Indiana, and Ohio. Specifically the part of North Dakota north and east of the Missouri River upstream as far as Lake Sakakawea and east of U.S. Highway 83 to the Canadian border; the part of South Dakota north and east of the Missouri River; the parts of Iowa, Illinois and Indiana that are north of Interstate 80; and the part of Ohio north of I-80 and west of the Maumee River at Toledo.

Outside of Western Great Lakes DPS - - Endangered, except Minnesota (Threatened) and Idaho, Montana, eastern Oregon and eastern Washington where it has been delisted

<http://www.fws.gov/mountain-prairie/species/mammals/wolf/>

Indiana bat (*Myotis sodalis*) Lead: Region 3

Status: Endangered

Habitat: Summer habitat includes small to medium river and stream corridors with well developed riparian woods; woodlots within 1 to 3 miles of small to medium rivers and streams; and upland forests. Caves and mines as hibernacula. **[Critical habitat has been designated for hibernacula: Blackball Mine in LaSalle County, Illinois; Big Wyandotte Cave in Crawford County and Ray=s Cave in Greene County, Indiana; Onyx Cave in Crawford**

**County, Cooper Hollow Sink Cave and Bear (Mud Sink) Cave in Franklin County, Pilot Knob Mine in Iron County, Great Scott Cave in Washington County, and Bat Cave in Shannon County, Missouri]**

Range: Alabama, Arkansas, Florida, Georgia, Illinois, Indiana, Iowa, Kentucky, Maryland, Michigan, Mississippi, Missouri, New Jersey, New York, North Carolina, Ohio, Oklahoma, Pennsylvania, Tennessee, Vermont, Virginia, west Virginia

Northern long-eared Bat  
Status: Threatened

Lead: Region 3

Habitat: Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. Roosts and forages in upland forests during spring and summer.

Range: Alabama , Arkansas , Connecticut , Delaware , District of Columbia , Florid , Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Vermont, Virginia, West Virginia, Wisconsin, Wyoming

Ozark big-eared bat (*Plecotus townsendii ingens*)  
Status: Endangered

Lead: Region 2

Habitat: Caves in limestone karst regions dominated by mature hardwood forest  
Range in Region 3: Missouri

**BIRDS**

Kirtland's warbler (*Setophaga kirtlandii =Dendroica kirtlandii*)  
Status: Endangered

Lead: Region 3

Habitat: Breeding in jack pine  
Range: Breeding habitat in Michigan and Wisconsin

Least tern (*Sterna antillarum*) Interior population

Lead: Region 4

Status: Endangered  
Habitat: Bare alluvial islands and dredged spoil islands  
Range: Alabama, Colorado, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Mississippi, Missouri, Montana, Nebraska, New Mexico, North Dakota, Oklahoma, South Dakota, Tennessee, Texas

Piping plover (*Charadrius melodus*) - Threatened and Endangered  
**Great Lakes population**

Lead: Region 3

Status: Endangered  
Habitat: beaches along shorelines of the Great Lakes

**Critical Habitat designated in:**

Illinois - Lake county

Indiana - Porter county

Michigan - Alger, Benzie, Charlevoix, Cheboygan, Chippewa, Emmet, Iosco, Leelanau, Luce, Mackinac, Mason, Muskegon, Presque Isle, and Schoolcraft counties

Minnesota - St. Louis county

Ohio - Erie and Lake counties

Wisconsin - Ashland, Douglas, Manitowoc, and Marinette counties

Range: Illinois, Indiana, Michigan, Minnesota, New York, Ohio, Pennsylvania, Wisconsin

***Great Plains population***

Status: Threatened

Habitat: Bare alluvial and dredged spoil islands; sand and gravel areas around fly ash ponds

Range: Alabama, Arkansas, Colorado, Connecticut, Florida, Iowa, Kansas, Louisiana, Maine, Massachusetts, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Oklahoma, Rhode Island, South Carolina, South Dakota, Texas, Virginia

Rufa Red Knot (*Calidris canutus rufa*)

Lead Region: Region 5

Status: Threatened, Dec. 11, 2014

Range: The knot migrates between breeding grounds in the Canadian Arctic and several wintering regions: southeast United States, northwest Gulf of Mexico, northern Brazil and Tierra del Fuego at the southern tip of South America. During migrations, groups of a few individuals to thousands of knots can be found anywhere along the coastal and inland U.S. migration corridors from Argentina to Canada. It is an irregular migrant through most the Region 3 states but regularly uses the Lake Michigan shoreline, in Michigan, and the Lake Erie shoreline, in Ohio.

Whooping Crane (*Grus americana*)

Lead: Region 2

Status: nonessential, experimental population

Habitat: open wetlands and lakeshores

Range: Iowa, Illinois, Indiana, Michigan, Minnesota, Ohio, and Wisconsin are within the Nonessential, Experimental Population area

## REPTILES

Copperbelly water snake (*Nerodia erythrogaster neglecta*) Lead: Region 3  
[northern population - north of 40E north latitude; approximately Indianapolis, Indiana]

Status: Threatened

Habitat: Wooded and permanently wet areas such as oxbows, sloughs, brushy ditches and floodplain woods

Region 3 Lead Office: East Lansing Michigan Field Office

Range: Indiana, Michigan, Ohio

Eastern massasauga (*Sistrurus catenatus*) Lead: Region 3

Status: Threatened

Habitat: Open to forested wetlands and adjacent upland areas

Region 3 Lead Office: Chicago Illinois Field Office

Range: Illinois, Indiana, Iowa, Michigan, Minnesota, Ohio, Wisconsin

Lake Erie water snake (*Nerodia sipedon insularum*) Lead: Region 3

Status: Delisted

Habitat: Shorelines of islands in western Lake Erie

Region 3 Lead Office: Columbus Ohio Field Office

Range: Ohio

## AMPHIBIANS

Ozark hellbender (*Cryptobranchus alleganiensis bishopi*) Lead: Region 3

Status: Endangered

Habitat: Streams of the Ozark plateau in southern Missouri and northern Arkansas.

Region 3 Lead Office: Columbia Missouri Field Office

Range: Arkansas and Missouri

## FISHES

Arkansas darter (*Etheostoma cragini*) Lead Region 6

Status: Candidate

Range: AR, CO, KS, MO, OK

Region 3 Lead Office: Columbia Missouri Field Office

Listing Priority: 11

Listed as Candidate: 1991

Grotto sculpin (*Cottus specus*) Lead: Region 3

Status: Endangered

Habitat: Streams in cave systems, occupying pools and riffles with moderate stream flow and low to moderate stream depth. Restricted to two karst (limestone regions characterized by sink holes, abrupt ridges, caves and underground streams) areas, the Central Perryville Karst and Mystery-Rimstone Karst in Perry County, southeast Missouri.

Region 3 Lead Office: Columbia Missouri Field Office

Range: Perry County, Missouri

Neosho madtom (*Noturus placidus*) Lead: Region 6  
Status: Threatened  
Habitat: Medium to large rivers in shallow riffles over gravel bottoms  
Range: Kansas, Missouri, Oklahoma

Niangua darter (*Etheostoma nianguae*) Lead: Region 3  
Status: Threatened  
Habitat: Clear, medium-sized streams that run off hilly areas underlain by chert and dolomite.  
**Critical habitat has been designated in Big Tavern Creek, Miller County; Brush Creek, Cedar and St. Clair Counties; Niangua River, Dallas County; and Pomme de Terre River, Greene County.**  
Region: Missouri

Ozark cavefish (*Amblyopsis rosae*) Lead: Region 4  
Status: Threatened  
Habitat: Caves in the Boone and Burlington limestone formations of the Ozark Mountains  
Range: Arkansas, Missouri, Oklahoma

Pallid sturgeon (*Scaphirhynchus albus*) Lead: Region 6  
Status: Endangered  
Habitat: Mississippi River downstream of its confluence with the Missouri River; Ohio River below Dam #53; Missouri River  
Range: Arkansas , Illinois , Iowa , Kansas , Kentucky , Louisiana , Mississippi , Missouri , Montana , Nebraska , North Dakota , South Dakota , Tennessee

Scioto madtom (*Noturus trautmani*) Lead: Region 3  
Status: Endangered  
Habitat: Stream riffles of moderate flow over sandy gravel bottom; may be extinct (Ohio Division of Wildlife will not consider it extinct until after the year 2000)  
Range: Ohio

Topeka shiner (*Notropis topeka*) Lead: Region 6  
Status: Endangered - Critical Habitat designated in Minnesota and Iowa  
Habitat: Small prairie streams  
Range: Iowa, Kansas, Minnesota, Missouri, Nebraska, South Dakota

## MUSSELS

Clubshell (*Pleurobema clava*) Lead: Region 5  
Status: Endangered  
Habitat: Found in coarse sand and gravel areas of runs and riffles in streams and small rivers  
Range: Illinois, Indiana, Kentucky, Michigan, New York, Ohio, Pennsylvania, Tennessee, West Virginia

- Curtis' pearlymussel (*Epioblasma florentina curtisi*) Lead: Region 3  
 Status: Endangered  
 Habitat: Restricted to Little Black River in Ripley County, Missouri  
 Range: Missouri
- Fanshell (*Cyprogenia stegaria*) Lead: Region 4  
 Status: Endangered  
 Habitat: Found in areas of packed sand and gravel at locations in a good current  
 Range: Alabama, Illinois, Indiana, Kentucky, Ohio, Tennessee, Virginia, West Virginia
- Fat pocketbook (*Potamilus capax*) (= *Proptera c.*) Lead: Region 4  
 Status: Endangered  
 Habitat: Slow-flowing water in large rivers  
 Range: Arkansas, Illinois, Indiana, Kentucky, Louisiana, Mississippi, Missouri
- Higgins eye pearly mussel (*Lampsilis higginsii*) Lead: Region 3  
 Status: Endangered  
 Habitat: Mississippi River and some of its larger northern tributaries (i.e., St. Croix and Wisconsin Rivers) in gravel or sand  
 Range: Illinois, Iowa, Minnesota, Missouri, South Dakota, Wisconsin
- Neosho mucket (*Lampsilis rafinesqueana*) Lead: Region 4  
 Status: Endangered - - Critical Habitat  
 Habitat: rivers  
 Range: Arkansas, Kansas, Missouri, and Oklahoma
- Northern riffleshell (*Epioblasma torulosa rangiana*) Lead: Region 5  
 Status: Endangered  
 Habitat: Large streams and small rivers in firm sand of riffle areas; also occurs in Lake Erie  
 Range: Indiana, Kentucky, Michigan, Ohio, Pennsylvania, West Virginia
- Orange-foot pimpleback (*Plethobasus cooperianus*) Lead: Region 4  
 Status: Endangered  
 Habitat: Gravel bars with strong currents in large rivers  
 Range: Alabama, Illinois, Indiana (possibly extirpated), Kentucky, Tennessee
- Pink mucket pearlymussel (*Lampsilis abrupta*) (= *L. orbiculata*) Lead: Region 4  
 Status: Endangered  
 Habitat: The lower Mississippi and Ohio Rivers and their larger tributaries  
 Range: Alabama, Arkansas, Illinois, Indiana, Kentucky, Louisiana, Missouri, Ohio, Tennessee, Virginia, West Virginia

- Purple cat's paw pearl mussel (*Epioblasma obliquata obliquata*) Lead: Region 3  
 Status: Endangered  
 Habitat: Gravel riffles of medium to large rivers  
 Range: Ohio
- Rabbitsfoot (*Quadrula cylindrica cylindrica*) Lead Region: 4  
 Status: Threatened - - Critical Habitat  
 Habitat: Rivers  
 Range: AL, AR, IL, IN, KS, KY, LA, MO, MS, OH, OK, PA, TN
- Rayed Bean (*Villosa fabalis*) Lead: Region 3  
 Status: Endangered  
 Habitat: Smaller, headwater creeks, but sometimes found in large rivers and Lake Erie  
 Range: Indiana, Michigan, New York, Ohio, Pennsylvania, Virginia
- Rough pigtoe (*Pleurobema plenum*) Lead: Region 4  
 Status: Endangered  
 Habitat: Medium to large rivers in sand or gravel  
 Range: Alabama, Indiana, Kentucky, Tennessee, Virginia
- Scaleshell (*Leptodea leptodon*) Lead: Region 3  
 Status: Endangered  
 Habitat: medium-sized and large rivers with stable channels and good water quality  
 Range: Arkansas, Illinois, Missouri, Nebraska, Oklahoma, South Dakota
- Sheepnose (*Plethobasus cyphus*) Lead: Region 3  
 Status: Endangered  
 Habitat: Shallow areas in larger rivers and streams  
 Range: Alabama, Illinois, Indiana, Iowa, Kentucky, Minnesota, Mississippi, Missouri, Ohio, Pennsylvania, Tennessee, Virginia, Wisconsin
- Snuffbox (*Epioblasma triquetra*) Lead: Region 3  
 Status: Endangered  
 Habitat: Small to medium-sized creeks in areas with a swift current and some larger rivers, and Lake Erie  
 Region 3 Lead Office: Ohio Field Office  
 Range: Alabama, Arkansas, Illinois, Indiana, Kentucky, Michigan, Missouri, Ohio, Pennsylvania, Tennessee, Virginia, and Wisconsin

Spectaclecase (*Cumberlandia monodonta*) Lead: Region 3  
Status: Endangered  
Habitat: Large rivers in areas sheltered from the main force of the current  
Range: Alabama, Arkansas, Illinois, Iowa, Kansas, Kentucky, Minnesota, Missouri, Tennessee, Virginia, West Virginia, Wisconsin

White catspaw (*Epioblasma obliquata perobliqua*) Lead: Region 3  
Status: Endangered  
Habitat: Firm sand or gravel riffles in small streams and medium to large rivers  
Range: Ohio, Indiana (possibly extirpated)

Winged mapleleaf (*Quadrula fragosa*) Lead: Region 3  
Status: Endangered  
Habitat: Medium to large rivers in mud, sand, or gravel  
Range: Arkansas, Minnesota, Missouri, Oklahoma, Wisconsin

## SNAILS

Iowa Pleistocene snail (*Discus macclintocki*) Lead: Region 3  
Status: Endangered  
Habitat: North-facing algific talus slopes  
Range: Illinois, Iowa

Tumbling Creek cavesnail (*Antrobia culveri*) Lead: Region 3  
Status: Endangered  
Habitat: Tumbling Creek in Tumbling Creek cave in Taney County  
Range: Missouri

## INSECTS

American burying beetle (*Nicrophorus americanus*) Lead: Region 5  
Status: Endangered  
Habitat:  
Range: Arkansas, Kansas, Massachusetts, Missouri, Nebraska, Ohio, Oklahoma, Rhode Island, South Dakota, Texas

Dakota skipper (*Hesperia dacotae*) Lead: Region 3  
Status: Threatened with 4(d) Rule - - Critical Habitat  
Habitat: high-quality tallgrass and mixed grass prairie  
Range: MN, SD, ND  
Critical Habitat: MN, SD, ND



- Hines emerald dragonfly (*Somatochlora hineana*) Lead: Region 3  
 Status: Endangered - - Critical Habitat  
 Habitat: Spring fed wetlands, wet meadows and marshes; calcareous streams & associated wetlands overlying dolomite bedrock  
 Range: Illinois, Michigan, Missouri, Wisconsin
- Hungerford's crawling water beetle (*Brychius hungerfordi*) Lead: Region 3  
 Status: Endangered  
 Habitat: Cool riffles of clean, slightly alkaline streams; known to occur in only 3 isolated locations  
 Range: Michigan
- Karner blue butterfly (*Lycaeides melissa samuelis*) Lead: Region 3  
 Status: Endangered  
 Habitat: Pine barrens and oak savannas on sandy soils and containing wild lupines (*Lupinus perennis*), the only known food plant of larvae.  
 Range: Illinois, Indiana, Michigan, Minnesota (extirpated), New Hampshire, New York, Ohio, Wisconsin
- Mitchell's satyr butterfly (*Noenympha mitchelli mitchelli*) Lead: Region 3  
 Status: Endangered  
 Habitat: Fens; wetlands characterized by calcareous soils which are fed by carbonate-rich water from seeps and springs  
 Range: Alabama, Indiana, Michigan, Mississippi, Ohio, Virginia
- Poweshiek skipperling (*Oarisma poweshiek*) Lead: Region 3  
 Status: Endangered - - Critical Habitat  
 Habitat: Remnants of tallgrass prairie and, in Michigan, fens  
 Lead Region: Region 3  
 Region 3 Lead Office: Twin Cities, Minnesota Field Office  
 Range: Michigan, Wisconsin  
 Critical Habitat: Iowa, Michigan, Minnesota, North Dakota, South Dakota, Wisconsin
- Rattlesnake-master Borer Moth (*Papaipema eryngii*) Lead: Region 3  
 Status: Candidate (Added 2013)  
 Listing Priority Number: 8  
 Habitat: Prairie with the plant, rattlesnake-master  
 Region 3 Lead Office: Rock Island Field Office  
 Current Range: AR, IL, KY, NC, OK
- Rusty patched bumble bee (*Bombus affinis*) Lead: Region 3  
 Status: Proposed as Endangered  
 Habitat: Prairie with the plant, rattlesnake-master

Region 3 Lead Office: Twin Cities Field Office

Current Range: IL, IN, IA, ME, MD, MA, MN, NC, OH, PA, TN, VA, WI – and Ontario, Canada

## CRUSTACEANS

- Cave crayfish, no common name (*Cambarus aculabrum*) Lead: Region 4  
Status: Endangered  
Habitat: Caves  
Range: Arkansas, Missouri
- Illinois cave amphipod (*Gammarus acherondytes*) Lead: Region 3  
Status: Endangered  
Habitat: cave streams  
Range: Illinois (currently found in 3 caves)

## PLANTS

- American hart's-tongue fern (*Asplenium scolopendrium* var. *americanum*) Lead: Region 4  
(=*Phyllitis japonica* ssp. *a.*)  
Status: Threatened  
Habitat: Cool limestone sinkholes in mature hardwood forest  
Range: Alabama, Michigan, New York, Tennessee
- Decurrent false aster (*Boltonia decurrens*) Lead: Region 3  
Status: Threatened  
Habitat: Disturbed alluvial soils (Mississippi and Illinois River alluvial floodplain);  
Range: Illinois, Missouri
- Dwarf lake iris (*Iris lacustris*) Lead: Region 3  
Status: Threatened  
Habitat: Partially shaded sandy-gravelly soils on lakeshores  
Range: Michigan, Wisconsin
- Eastern prairie fringed orchid (*Platanthera leucophaea*) Lead: Region 3  
Status: Threatened  
Habitat: Mesic to wet prairies and meadows  
Range: Illinois, Indiana, Iowa, Maine, Michigan, Missouri, Ohio, Oklahoma, Virginia,  
Wisconsin
- Fassett's locoweed (*Oxytropis campestris* var. *chartaceae*) Lead: Region 3  
Status: Threatened  
Habitat: Open sandy lakeshores  
Range: Wisconsin
- Geocarpon (no common name) (*Geocarpon minimum*) Lead: Region 4

- Status: Threatened  
Habitat: Moist soils in exposed sandstone glades  
Range: Arkansas, Louisiana, Missouri, Texas
- Houghton's goldenrod (*Solidago houghtonii*) Lead: Region 3  
Status: Threatened  
Habitat: Sandy flats along Great Lakes shores  
Range: Michigan, New York
- Lakeside daisy (*Hymenoxys herbacea*)(=*H. acaulis* var. *glabra*) Lead: Region 3  
Status: Threatened  
Habitat: Dry rocky prairies; limestone rock surfaces including outcrops and quarries  
Range: Illinois, Michigan, Ohio
- Leafy prairie clover (*Dalea foliosa*) Lead: Region 4  
Status: Endangered  
Habitat: Prairie remnants on thin soil over limestone (Des Plaine River floodplain)  
Range: Alabama, Illinois, Tennessee
- Leedy's roseroot (*Sedum integrifolium* ssp. *leedyi*) Lead: Region 3  
Status: Threatened  
Habitat: Cool, wet groundwater-fed limestone cliffs  
Range: driftless area of southeastern Minnesota, New York
- Mead's milkweed (*Asclepias meadii*) Lead: Region 3  
Status: Threatened  
Habitat: Prairies; and rhyolite glades in Missouri  
Range: Illinois, Indiana, Iowa, Kansas, Missouri, Wisconsin
- Michigan monkey-flower (*Mimulus glabratus* var. *michiganensis*) Lead: Region 3  
Status: Endangered  
Habitat: Soils saturated with cold flowing spring water; found along seepages, streams and lakeshores  
Range: Michigan
- Minnesota dwarf trout lily (*Erythronium propullans*) Lead: Region 3  
Status: Endangered  
Habitat: North facing slopes and floodplains in deciduous forests  
Range: Minnesota
- Missouri bladderpod (*Physaria filiformis*) Lead: Region 3  
Status: Threatened  
Habitat: Open glades in shallow limestone soils

Range: Missouri

Northern wild monkshood (*Aconitum noveboracense*) Lead: Region 3

Status: Threatened

Habitat: Cool, moist, shaded cliff faces or talus slopes in wooded ravines, near water seeps

Range: Iowa, New York, Ohio, Wisconsin

Pitcher's thistle (*Cirsium pitcheri*) Lead: Region 3

Status: Threatened

Habitat: Stabilized dunes and blowout areas

Range: Illinois, Indiana, Michigan, Wisconsin

Pondberry (*Lindera melissifolia*) Lead: Region 4

Status: Endangered

Habitat: Bottomland hardwood forest, poorly drained depressions, and margins of limestone sinks

Range: Alabama, Arkansas, Georgia, Mississippi, Missouri, North Carolina, South Carolina

Prairie bush-clover (*Lespedeza leptostachya*) Lead: Region 3

Status: Threatened

Habitat: Dry to mesic prairies with gravelly soils

Range: Illinois, Iowa, Minnesota, Wisconsin

Price's potato-bean (*Apios priceana*) Lead: Region 4

Status: Threatened

Habitat: Wet floodplain forests, shrubby swamps

Range: Alabama, Illinois (possibly extirpated), Kentucky, Mississippi, Tennessee

Running buffalo clover (*Trifolium stoloniferum*) Lead: Region 3

Status: Endangered

Habitat: Disturbed bottomland meadows; disturbed sites that have shade during part of each day

Range: Arkansas, Indiana, Kentucky, Missouri, Ohio, West Virginia

Short's bladderpod (*Physaria globosa*) Lead: Region 4

Status: Endangered - - Critical Habitat

Habitat: Steep, rocky wooded slopes and talus areas, occurs along cliff tops and bases and cliff ledges. Found adjacent to rivers or streams and on south to west facing slopes. The Indiana population is found within the Shawnee Hills section of the Interior Low Plateau Physiographic Province.

Range: Indiana, Kentucky, Tennessee

- Short's goldenrod (*Solidago shortii*) Lead: Region 4  
Status: Endangered  
Habitat: Cedar glades and openings in oak and hickory forests  
Range: Indiana, Kentucky
- Small whorled pogonia (*Isotria medeoloides*) Lead: Region 5  
Status: Threatened  
Habitat: Dry woodland; upland sites in mixed forests (second or third growth stage)  
Range: Connecticut, Delaware, Georgia, Illinois, Maine, Massachusetts, Michigan, New Hampshire, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, South Carolina, Tennessee, Virginia, West Virginia
- Virginia sneezeweed (*Helenium virginicum*) Lead: Region 5  
Status: Threatened  
Habitat: Sinkhole ponds  
Range: Missouri, Virginia
- Virginia spiraea (*Spiraea virginiana*) Lead: Region 5  
Status: Threatened  
Habitat: Stream banks and floodplains  
Range: Georgia, Kentucky, North Carolina, Ohio, Pennsylvania, Tennessee, Virginia, West Virginia
- Western prairie fringed orchid (*Platanthera praeclara*) Lead: Region 3  
Status: Threatened  
Habitat: Wet prairies & sedge meadows.  
Range: Iowa, Kansas, Minnesota, Missouri, Nebraska, North Dakota, Oklahoma, South Dakota
-

## Regional Office Addresses

### **Region 1 - Pacific**

U.S. Fish and Wildlife Service  
Eastside Federal Complex  
911 N.E. 11th Avenue  
Portland, Oregon 97232-4181

### **Region 2 - Southwest**

U.S. Fish and Wildlife Service  
Box 1306  
Albuquerque, New Mexico 87103

### **Region 3 - Great Lakes-Big Rivers**

U.S. Fish and Wildlife Service  
5600 American Blvd. West, Ste. 990  
Bloomington, Minnesota 55437-1458

### **Region 4 - Southeast**

U.S. Fish and Wildlife Service  
1875 Century Boulevard  
Atlanta, Georgia 30345

### **Region 5 - Northeast**

U.S. Fish and Wildlife Service  
300 Westgate Center Drive  
Hadley, Massachusetts 01035-9589

### **Region 6 - Mountain-Prairie**

U.S. Fish and Wildlife Service  
Box 25486  
Denver Federal Center  
Denver, Colorado 80225

### **Region 7 - Alaska**

U.S. Fish and Wildlife Service  
1011 East Tudor Road  
Anchorage, Alaska 99503