

# An Introduction to lowa Soil Surveys



Introduction 3
Soil Horizons5
How Soil Develops 7
General Soil Maps 9
Soil Properties12
Using Soil Maps16
Locating Information







properties of the soil and predict soil behavior for a host of uses. These predictions, called soil interpretations, are developed to help users of soils manage the resource. Soil surveys of each Iowa county are available in three formats: as publications, on CDs, and online.

The publications represent a snapshot in time. They contain information that was current at the time of printing. But the text, tables and soil maps may have been updated since publication. The most current information about the soil survey program in Iowa is available at www.ia.nrcs.usda.gov/soils.html.

The Natural Resources Conservation Service (NRCS) also has several web sites devoted to the soil survey program on a national basis. The primary one is <a href="http://websoilsurvey.nrcs.usda.gov/app/">http://websoilsurvey.nrcs.usda.gov/app/</a>.


To gather information for a soil survey, soil scientists walk the landscapes, dig holes with soil augers, shovels and probes, and examine cross-sections of soil profiles. They observe soil textures (the ratio of sand, silt and clay), soil color, structure and thickness of the different soil horizons. These and many other soil properties are studied in the field, while others are deter-

mined through laboratory tests. The intent of this publication is to acquaint users and potential users of soil survey reports with the content of those reports, and to help them extract useful soils data. To be proficient in using soil survey data, it is imperative that users have a basic understanding of the concepts of soil development and of soil-land-scape relationships. These topics are covered briefly in the next two sections of this guide.

Photo: NRCS soil scientist Rick Bednarek (right) evaluates a soil probe drawn from a continuous no-till field in Adair County with former District Conservationist Marvin Lundstedt.

### **Soil Horizons**

**Soils develop in layers.** These layers, called horizons, can be seen where roads have been cut through hills and other areas where the soil is exposed. The presence and thickness of each horizon varies with location. Under disturbed conditions, such as intensive agriculture, building sites, or where there is severe erosion, some horizons may not be present. The photos on Page 6 are profiles of Iowa soils with visible horizons labeled.



#### **O** Horizon

O horizons are present in soils that have significant amounts of decomposed plant and animal materials at the soil surface. The layer is black in color, and is most commonly present in shallow, depressional areas of north-central Iowa. Because of the high levels of organic content, O horizons feel much lighter than mineral horizons.

#### **A Horizon**

All soils have an A horizon. It is the layer of mineral soil at the soil surface or immediately below the O horizon. Commonly referred to as topsoil, the A horizon is typically darker and more fertile than lower horizons.

#### **E** Horizon

Soils in eastern Iowa that formed under forest vegetation commonly have a subsurface layer called an E horizon. Soil in the E horizon is silty and light colored. As water moves through this horizon, soluble minerals, such as iron, are leached to lower horizons. This horizon will usually have less clay than the horizons above and below.

#### **B** Horizon

Most soils have a B horizon, where soil development has taken place since the original soil parent material was formed. B horizons are lighter colored than the organically enriched A and O horizons and have developed good soil structure. The B horizon typically has more clay and different soil pH than the layers directly above and below.

#### **C** Horizon

The C horizon is the undeveloped material from which the soil is formed. It is also called soil parent material. Common parent materials in Iowa include glacial till, which is the material left by glaciers that passed through the state tens of thousands of years ago; wind blown material called loess: and water deposited material from flooding events called alluvium. The C horizon is usually deep in the soil profile, but can also be near the soil surface where soils have not been able to develop, such as on the Missouri or Mississippi River bluffs.

#### **R** Horizon

The lowest horizon, R horizon, is bedrock. In Iowa, this horizon is typically very deep. However, there are areas in northeast Iowa where the bedrock plateau was so high that the glaciers could not pass over the bedrock, and in central Iowa where bedrock is exposed due to geologic erosion.

### **lowa Soil Horizons**













- A. Colo soil is found in bottomlands throughout lowa. The topsoil (A Horizon) is one meter or more thick due to the low lying landscape position.
- B. Sperry soil is found in southern lowa. The subsurface layer (E Horizon) in this model has been stripped of organic coatings through frequent wetting and drying, and appears bleached.
- C. Clarion soil occurs in north-central lowa. Soils in this region of the state were formed from glacial deposits where stones and rocks are common in the parent material (C Horizon).
- D. Tama soil occurs in eastern lowa and has been designated as lowa's State Soil. It formed under prairie vegetation that resulted in a thick topsoil (A Horizon) and a well-developed subsoil (B Horizon). The parent material (C Horizon) has wind deposited sediments, called loess.
- E. Racine soil occurs in northeast lowa. It formed under oak-savannah vegetation. The parent material (C Horizon) was the result of significant geologic erosion.
- Allamakee soil formed in eastern lowa under forest vegetation. The surface layer (A Horizon) is much thinner in forested soils than in prairie soils. Most of the organic matter is stored in trees, as opposed to being stored in many fine roots of grasses.

	Summit	Midslope	Footslope
	Weakly Developed Soil	<b>Moderately Developed Soil</b>	Well Developed Soil
12"	A - TOPSOIL	A - TOPSOIL	A - TOPSOIL
		B - SUBSOIL	
24"	Poil	1.000	B - SUBSOIL
36"	C-PARENT MATERIAL	C-PARENT MATERIAL	C-PARENT MATERIAL

### **How Soil Develops**

The three boxes in the diagram represent either a soil in the same spot at different times in history or a soil at three different points on a hill slope. During that time, climate, living organisms and topography changed the soil. Notice in this example how the parent material is broken down, and the soil's structure changes from weakly developed to well-developed. Not all soils undergo this exact type of change, and the time for change differs with location. These differences occur because the climate differs from one location to another, as do the number and types of organisms living in the soil.

#### **Parent Material**

Parent material (C Horizon) consists of the loosely arranged mineral and organic matter from which soils are developed. Parent material can be ash from volcanoes, sediments moved and

deposited by wind and water, or sand and rock deposited by glaciers. Broken down bedrock is another example of parent material. In Iowa, Alluvial soils deposited by water are the youngest and least developed.

#### **Climate**

Climate helps change parent material into subsoil, subsurface soil and topsoil. Freezing, thawing, wetting and drying breaks parent material apart. Rain dissolves some minerals and transports them deeper into the soil. Soils in northwest Iowa develop slower because it is drier and cooler than other parts of the state.


#### **Living Organisms**

Plants and animals change the weathered parent material into subsoil and topsoil. Leaves, twigs, stems and bark from plants fall onto the soil, and are broken down by fungi, bacteria, insects and other animals that live in

the soil. Insects and earthworms burrowing through the soil eat and break down organic matter and minerals into simpler compounds. Plants and animals help make the soil rich in nutrients as they die and decay. Living organisms are directly related to the amount of vegetation produced, which is directly related to soil moisture.

#### **Topography**

Topography refers to the lay of the land. Steeper slopes may have eroded topsoil, exposing subsoil or parent material. This makes the soil look like the weakly developed soil in the diagram (above). Steeper slopes take longer to develop because moisture runs off the slope instead of infiltrating the soil. Soils in low areas and depressions tend to be wetter and more developed. In contrast, soils in more sloping areas tend to be drier and less developed.



#### SOIL ASSOCIATIONS

Mahaska-Tainter association; Nearly level and gently sloping, somewhat poorly drained and poorly drained soils that have a subsoil of silty clay loam or silty clay; on uplands

Otley-Ladega-Nira association: Gently sloping to strongly sloping, moderately well drained soils that have a subsoil of sity clay loam; on uplands

Clinton-Lindley-Gara association: Gently sloping to steep, well drained and moderately well drained soils that have a subsoil of silty clay learn or clay learn; on uplands

Ladega-Clinton-Gesport association: Gently sloping to steep, moderately well drained soils that have a subsoil of silty clay learn or silty clay; on uplands

Colo-Nodaway-Zook association: Nearly level and gently sleping, poor by drained to mederately well drained sells that are deminantly silty clay leam and silt leam throughout or that have a subsell of silty clay leam or silty clay; on bettem lands

Clinten-Chelsea association: Mederately sloping to steep, mederately well drained and excessively drained soils that have a subsoil of silty class of that have a subsoil of silty class of the sand throughout; on uplands

Pershing-Grundy-Haig association: Moderately sloping to nearly level, moderately well drained to poorly drained soils that have a subsoil of sitty clay and sitty clay leam; on uplands

Compiled 1975

#### U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE

IOWA AGRICULTURE AND HOME ECONOMICS EXPERIMENT STATION COOPERATIVE EXTENSION SERVICE, IOWA STATE UNIVERSITY DEPARTMENT OF SOIL CONSERVATION, STATE OF IOWA

#### GENERAL SOIL MAP

MAHASKA COUNTY, IOWA

	5	TOWNSHIP						
٧	6	5	4	3	2			
	7		9	10	11	1		
<b>^</b>	18	17	16	15	14	1		
	19	20	21	22	23	2		
	30	29	28	27	26	2		
	31	32	33	34	35	3		

Such area outlined on this map cannists of more than one kind of soil. The map is thus maged for general planning syther than a bas her decisions on the use of specific tracks.

8


### **General Soil Maps**


A General Soil Map is a small-scale map of the survey area. It is color-coded to show major soil associations. Descriptions of each of the soil associations are near the front of the soil survey report. This section of the report is labeled *General Soil Map Units* in the Table of Contents of a Soil Survey. A map of Iowa Soil Regions is also available online at www.ia.nrcs.usda. gov/soils.html.

Each color-coded area on the General Soil Map has a corresponding description. In the example General Soil Map of Mahaska County (Page 8), green designates the Pershing-Grundy-Haig soil association. As the name of the map implies, Pershing, Grundy and Haig are the major soils that occupy the landscapes of this section of Mahaska County. Likewise, the description of soil association 7 gives general information about this section of the county.

Iowa soil survey reports contain a number of three-dimensional drawings, called block diagrams, depicting the relationships of soils, parent material and topography for the major soil associations. The illustration below shows the Pershing-Grundy-Haig the relationship of parent material (loess, glacial till, shale and alluvium) and the topography to the different soils. Refer to the *How Soil Develops* section of this guide (Page 7) for additional insights on these relationships.

The General Soil Map can be used to compare the suitability of large areas for general land uses. Because of the small scale, it is not intended to be used to make management decisions on specific sites. Soils within a soil association may vary greatly in slope, depth, drainage and other characteristics that affect management.





Natural Resources Conservation Service photo illustration

#### **Soil Capability Classes**

Each soil is grouped by capability class, and in most cases by subclass. These groupings reveal the suitability of soils for field crops and pasture, and indicate associated risks and responses to conservation treatment and management.

There are eight capability classes, which represent progressively greater limitations and narrower choices for practical land use. These classes are designated by Roman numerals I through VIII.

#### Classes:

- I have few limitations that restrict their use. Class I has no subclasses because these soils have few limitations.
- II have moderate limitations that restrict the choice of plants or that require moderate conservation practices.
- **III** have severe limitations that reduce the choices of plants or

- that require special conservation practices, or both.
- IV have very severe limitations that reduce the choice of plants or that require very careful management, or both.
- V are not likely to erode but have other limitations, impractical to remove, that limit their use.

- VI have severe limitations that make them generally unsuitable for cultivation.
- VII have very severe limitations that make them unsuitable for cultivation.
- VIII soils and miscellaneous areas have limitations that nearly preclude their use for commercial crop production.


Capability subclasses are noted with an "e," "w" or "s" following the capability class numbers. For example, in 2e, the "e" indicates the soil is erosive. A "w" would signify a wetness limitation. An "s" would denote a shallow, droughty or stony soil.

#### **How To Use Soil Capability Classes**

On the whole, producers make rational use of their farmland. Almost all United States land in Capability Class I—with soils that are level, deep, well-drained and easy to work—is used to grow crops. The more extensive capability classes of II and III comprise the majority of cultivated cropland, although that land generally requires conservation measures.

Most of Iowa's land is very well suited to cropland. More than 84 percent of the state's 36 million acres are in Class I, II or III, and only 8 percent is in Class VI, VII or VIII.

More than two-thirds of Iowa is cultivated cropland, and there is an additional 1 million acres that is in non-cultivated hay or horticultural crops. About 3.5 million acres are grazed pastures and a little more than 2 million acres in Iowa are forested.



### **Soil Properties**

All soils possess properties that impact their suitability for specific uses. Although some properties, such as slope, may be altered to some extent to make a site more suitable, it is usually impractical to modify these properties extensively. Following is a list of major soil properties that affect the suitability of soils for a number of specific uses. The table on Page 15 provides examples of various land uses that are affected by specific soil properties.

#### **Texture**

Soil particles vary in size. From coarsest to finest, they are sand, silt or clay. The percentage of each in any given soil determines its texture.

Texture is an important soil property because it is closely related to many aspects of soil behavior. It influences ease of tillage and root development. It also affects the availability of nutrients in the soil, the amount of water that a soil will hold and how water moves through the soil.



Relative sizes of sand, silt and clay

Sand causes little restriction of air, water and root movement through the soil, whereas clay often reduces movement.

#### **Structure**

Two soils with the same texture may have distinctly different physical properties because of the arrangement of soil particles. This arrangement is called soil structure. Soil structure forms when individual grains of sand, silt and clay are bound together physically or chemically. Plant roots, organic matter and clay particles all provide physical and chemical binding agents. These bound particles form larger units called peds. A ped is a single unit of soil structure. The shapes of peds, which range in size, determine the structure type

(granular, platy, blocky or prismatic).

Soil structure is constantly changing. Although the process is too slow for you to notice, a soil's structure becomes stronger and more distinct over time.

#### Color

Color is one of the most noticeable soil properties. Color is affected by organic matter content, climate, soil drainage and minerals. Most soil minerals are naturally white or light gray. However, humus and iron compounds change the outer color of the soil particles much like a coat of paint.

Soil color gives clues about the nature of the root zone, which is the normal depth that roots penetrate into the soil. Dark colors reflect favorable amounts of humus. Gray colors suggest wetness. Brown and red colors indicate favorable air-water relationships.

### Depth to High Water Table

A saturated soil layer, or water table, will fluctuate throughout the year. How high the water table rises, and how long it stays at that

level, will affect the use of the soil.

#### **Depth to Bedrock**

Bedrock is the solid rock under the soil and parent material. Sometimes it is exposed at the surface. The depth from the soil surface to bedrock influences the soil's potential and limitations.

#### **Depth Classes**

#### **Very Shallow:**

less than 10 inches to bedrock

#### **Shallow:**

10 to 20 inches to bedrock

#### **Moderately Deep:**

20 to 40 inches to bedrock

#### Deep:

40 to 60 inches to bedrock

#### Very Deep:

More than 60 inches to bedrock

### Permeability and Saturated Hydraulic Conductivity

Permeability and Saturated Hydraulic Conductivity both refer to how easily water moves through soil. Water movement is influenced by texture, structure, compaction and the presence or absence of cementing agents.



Permeability is measured as the inches per hour that water moves downward and laterally through saturated soil. (See the illustration at the bottom of this page.)


Saturated hydraulic conductivity is the measurement of this movement of water through the soil. It is measured in micrometers per second.

#### **Rock Fragments**

The number of rock fragments on the surface and the percentage of fragments in the soil profile are important to land-use planning. Excessive rock fragments reduce the available water capacity of the soil.

#### Slope

Perfectly level land has a slope of 0 percent. All other land rises or falls to some degree. Slope is an important factor, and impacts water erosion and surface drainage.



### Available Water Capacity

Available Water Capacity is a measure of how much water a soil can store for use by plants. The Available Water Capacity is influenced by soil texture, organic matter content and pore space. It is measured in inches of water per inch of soil.

#### **Flooding and Ponding**

Periodic floodwater from overflowing streams or runoff from nearby slopes makes sites risky for some uses. If floodwaters persist, they may delay planting and harvesting.


#### Permeability and Saturated Hydraulic Conductivity-How Water Moves Through Soil



Granular



**Subangular Blocky** 



**Columnar or Prismatic** 



**Platy** (closely resembles clay and compacted soils)

(SOIL STRUCTURE TYPES)

### Soil Property – Land Use Table

Indicates a particular land use that could be affected by a specific soil property.		SOIL PROPERTY									
		Wetness	Perme- ability	Depth to Bedrock	Slope	Surface Texture	Sub- surface Texture	Small Stones	Large Stones	Flooding	
SELECTED LAND USE	Tank Absorption Fields	X	X	X	X		X		X	X	
	Sanitary Landfill (Trench)	X	X	X	X		X		X	X	
	Sanitary Landfill (Area)	X	X		X					X	
	Dwellings Without Basements	X		X	X	X	X		X	X	
	Dwellings With Basements	X		X	X		X		X	X	
	Small Commercial Buildings	X		X	×		X		X	X	
	Local Roads and Streets	X		X	X	X	X		X	X	
	Lawns and Landscaping	X		X	x	X		x	X	X	
	Camp Areas	X	X		X	X		X	X	X	
	Picnic Areas	X			X	X		x	X	X	
	Playgrounds	X	x	X	X	X		x	X	X	
	Paths and Trails	X			X	x		x	X	X	
	Golf Fairways	X		×	X	X		X	X	X	


## Using Soil Maps

The General Soil Map, Index to Map Sheets and Detailed Soil Map Sheets are located at the back of the soil survey, or as separate maps in a folder. To obtain information about a particular plot of land, locate the area on the Index to Map Sheets, select the appropriate map sheet number, and locate the map sheet. Map sheets are in numerical order in the soil survey. Then locate a specific parcel of land on the map sheet by noting imagery or section numbers, roads, streams and towns.

The symbols on the map sheets are listed on the back of the Index to Map Sheets with the names of each soil that they represent. You can obtain detailed information about each soil area of interest by referring to the map unit description located in the text or interpretive tables located in the back of the soil survey publication. The map unit description or tables can be located easily by using the Table of Contents in the front of the publication.

Individual soil symbols on the soil maps may contain more than one soil. They generally include areas of similar soils, and possibly small areas of contrasting soils. The composition of soils is explained in each

map unit description. The soil maps provide sufficient information for developing resource plans, but an onsite investigation should be conducted to plan intensive uses in small areas.



# Locating Information

Every soil type has dozens of specific soil properties and interpretations that aid users in making decisions on the land. These properties and interpretations are listed by soil map unit number in various tables located throughout the soil survey report or online at: http://soildatamart.nrcs.usda.gov/Survey.aspx?State=IA.

Below is a list of tables and specific information they contain that are available in most soil survey reports.



**Temperature and Precipitation** average daily minimum and maximum temperatures, growing degree days, snowfall, spring and fall freeze dates, length of growing

season

**Cropland Management** Considerations

cropping considerations for each type such as 'acid soil', 'excessive permeability', or 'water erosion'

**Crop and Pasture Yield Tables** crop and pasture yields, land capability class, subsoil potassium and

phosphorus levels

**Prime Farmland Lists** 

includes additional lands of statewide and local importance

Windbreaks and Environmental Plantings, Forestland **Productivity** 

specific trees that are suitable for each soil type and the height they will grow in 20 years, site index and volume of wood fiber is also available in areas where there are forests

**Recreational Development** 

soil suitability for camp areas, picnic areas, playgrounds, paths and trails, motorcycle trails, and golf fairways

Wildlife Habitat

openland wildlife, woodland wildlife and wetland wildlife suitability

**Building Site Development** 

soil suitability for dwelling with or without basements, small commercial buildings, local roads and streets, shallow excavations and lawns and landscaping

**Sanitary Facilities** 

soil considerations for septic tank absorption fields, sewage lagoons and landfills

**Construction Materials** 

soils as a potential source of gravel, sand, topsoil, roadfill or reclamation material

Water Management

soil suitability for ponds, embankments, dikes and levees

Soil Survey of Allamakee County, Iowa

Agricultural Waste Management soil suitability for the application

of manure or sewage sludge or for wastewater treatment

**Engineering Index Properties** 

Unified and AASHTO classifications, rock fragments, percent passing sieves, liquid limit and plasticity index

Physical Properties of the Soil

soil bulk density, clay content, available water capacity, shrink-swell potential, organic matter content, erosion factors

Chemical Properties of the Soil

cation exchange capacity, soil pH, calcium carbonate content


Water Features

depth to saturated soil for each month, frequency and duration of ponding and flooding for each month for each soil type

Soil Features

depth to bedrock, potential for soil subsidence and frost heaving, risk of corrosion to steel and concrete

Soil interpretations are constantly being developed and tested. The Department of Homeland Security uses soil maps to determine suitable sites for animal burial and composting, or to identify soils that can sequester radioactive materials. The Department of Defense looks at soil surveys to determine areas vehicle trafficability, helicopter landing areas, and even strategic fighting positions.





This publication was produced and printed by:
United States Department of Agriculture
Natural Resources Conservation Service
210 Walnut Street, Room 693
Des Moines, IA 50309

April 2007

For questions about information contained in this publication or in specific soil surveys, contact the lowa NRCS Soils Staff at the above address, or phone (515) 284-4135.

For information about NRCS and its programs in lowa, visit us online at www.ia.nrcs.usda.gov



The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact

USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.





www.ia.nrcs.usda.gov

Helping People Help the Land in lowa