

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	ATLANTIC COD								TOTAL	
	NORTH ATLANTIC		MID-ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----										
1981. . .	15,182	22	2,542	16	0	0	0	0	17,725	19
1982. . .	12,464	26	9,070	34	0	0	0	0	21,534	21
1983. . .	13,112	22	1,472	38	0	0	0	0	14,584	20
1984. . .	5,425	12	515	26	0	0	0	0	5,940	11
1985. . .	22,525	58	554	26	0	0	0	0	23,078	56
1986. . .	7,237	34	290	22	0	0	0	0	7,527	33
1987. . .	8,135	20	852	26	0	0	0	0	8,987	18
1988. . .	8,186	20	3,919	22	0	0	0	0	12,105	15
1989. . .	5,155	12	1,955	21	0	0	0	0	7,110	10
1990. . .	6,790	17	1,121	18	0	0	0	0	7,911	15
1991. . .	8,844	17	463	23	0	0	0	0	9,307	17
1992. . .	1,906	18	430	26	0	0	0	0	2,335	16
1993. . .	5,190	17	1,949	20	0	0	0	0	7,139	14
1994. . .	4,083	25	203	27	0	0	0	0	4,285	24
1995. . .	3,437	20	1,164	17	0	0	0	0	4,601	16
1996. . .	3,260	21	72	27	0	0	0	0	3,332	20

YEAR	ATLANTIC CROAKER								TOTAL	
	NORTH ATLANTIC		MID-ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----										
1981. . .	0	0	538	19	809	17	1,735	19	3,082	12
1982. . .	0	0	526	58	1,132	17	3,143	20	4,800	15
1983. . .	0	0	518	25	946	21	1,828	14	3,292	11
1984. . .	0	0	721	17	2,684	19	1,419	18	4,824	12
1985. . .	0	0	861	13	1,076	15	1,228	15	3,164	8
1986. . .	0	0	2,153	11	3,105	29	2,806	14	8,064	13
1987. . .	0	0	1,498	9	1,437	18	1,193	22	4,128	9
1988. . .	0	0	3,318	13	1,327	14	1,643	39	6,288	13
1989. . .	0	0	1,349	8	1,029	10	449	19	2,827	6
1990. . .	0	0	913	13	804	10	771	29	2,488	10
1991. . .	0	0	1,860	11	688	11	1,888	44	4,437	19
1992. . .	0	0	1,826	10	802	7	569	13	3,197	7
1993. . .	0	0	2,481	9	537	8	593	10	3,611	6
1994. . .	0	0	4,019	6	851	9	602	17	5,472	5
1995. . .	0	0	3,374	8	661	18	392	18	4,426	7
1996. . .	0	0	3,815	8	413	9	433	12	4,661	7

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	ATLANTIC MACKEREL								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----										
1981. . .	1,911	22	5,166	21	0	0	0	0	7,077	16
1982. . .	844	31	1,781	28	0	0	0	0	2,625	22
1983. . .	1,093	19	5,487	34	38	72	0	0	6,618	28
1984. . .	1,726	17	3,387	27	0	0	0	0	5,113	19
1985. . .	4,197	31	1,784	76	0	0	0	0	5,982	32
1986. . .	2,128	32	7,183	31	0	0	0	0	9,311	25
1987. . .	2,623	23	6,265	23	0	0	0	0	8,889	17
1988. . .	5,669	21	1,511	42	18	0	0	0	7,198	18
1989. . .	2,444	12	1,495	24	0	0	0	0	3,939	12
1990. . .	2,638	14	1,478	26	0	0	0	0	4,116	13
1991. . .	3,576	17	2,080	21	0	0	0	0	5,657	13
1992. . .	530	14	96	35	0	0	0	0	629	13
1993. . .	1,255	12	66	44	0	0	0	0	1,326	12
1994. . .	3,675	16	85	52	0	0	0	0	3,760	16
1995. . .	1,847	13	907	36	0	0	0	0	2,754	15
1996. . .	2,699	12	422	25	2	59	0	0	3,123	11
YEAR	BLACK DRUM								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----										
1981. . .	0	0	95	0	122	21	1,065	18	1,283	15
1982. . .	0	0	0	0	276	26	2,600	18	2,876	16
1983. . .	0	0	1,378	21	397	22	3,020	19	4,796	14
1984. . .	0	0	39	0	586	23	1,385	22	2,011	16
1985. . .	0	0	345	36	572	21	1,506	23	2,424	16
1986. . .	0	0	722	20	469	21	2,817	19	4,008	14
1987. . .	0	0	463	32	388	17	3,799	35	4,651	29
1988. . .	0	0	36	0	428	20	2,222	12	2,686	11
1989. . .	0	0	247	29	222	37	1,903	16	2,372	14
1990. . .	0	0	11	79	181	36	832	22	1,023	19
1991. . .	0	0	84	1	549	20	1,219	22	1,852	16
1992. . .	0	0	238	39	566	15	1,549	8	2,352	8
1993. . .	0	0	2	42	428	13	1,478	13	1,908	11
1994. . .	0	0	3	74	693	13	1,177	20	1,873	13
1995. . .	0	0	155	42	658	16	1,307	9	2,121	8
1996. . .	0	0	99	67	568	13	1,159	8	1,826	8

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	BLACK SEA BASS								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	143	15	1,060	16	759	19	265	75	2,229	13
1982. . .	319	26	9,568	36	1,872	17	404	20	12,163	28
1983. . .	294	23	3,760	21	907	21	128	26	5,088	16
1984. . .	66	31	1,165	14	2,333	16	197	24	3,761	11
1985. . .	48	37	2,036	14	1,328	16	1,834	34	5,246	14
1986. . .	639	22	11,745	22	475	14	1,177	19	14,037	18
1987. . .	141	39	1,777	14	1,043	14	812	19	3,773	9
1988. . .	245	28	2,610	16	1,472	33	1,361	12	5,687	12
1989. . .	98	20	3,182	7	1,155	21	1,765	14	6,200	7
1990. . .	47	24	2,705	9	546	21	607	18	3,905	7
1991. . .	63	42	4,070	11	853	15	488	14	5,475	8
1992. . .	30	18	2,570	8	727	10	417	10	3,744	6
1993. . .	49	16	4,776	22	560	16	509	16	5,894	18
1994. . .	37	53	2,890	13	623	17	531	11	4,081	10
1995. . .	64	27	6,099	14	588	12	288	11	7,040	12
1996. . .	85	19	5,812	15	654	13	249	11	6,801	13

YEAR	BLUEFISH								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	29,632	22	59,148	7	6,507	21	580	34	95,867	8
1982. . .	41,226	19	34,978	13	6,802	18	418	27	83,424	11
1983. . .	33,988	22	38,431	15	16,703	32	3,589	32	92,711	12
1984. . .	18,985	13	38,066	12	10,402	17	880	78	68,333	8
1985. . .	17,205	21	28,339	13	6,970	22	665	32	53,180	10
1986. . .	38,718	9	48,710	10	5,459	13	856	25	93,742	6
1987. . .	20,930	10	48,407	6	7,316	12	551	18	77,204	5
1988. . .	11,726	7	27,996	7	8,501	14	1,171	13	49,393	5
1989. . .	11,332	10	23,183	5	4,744	13	1,724	25	40,984	5
1990. . .	10,487	9	16,697	5	3,372	8	356	18	30,913	4
1991. . .	12,199	8	17,672	6	3,127	10	969	28	33,966	4
1992. . .	8,414	8	13,365	6	2,496	10	506	15	24,782	4
1993. . .	9,209	6	8,735	7	2,348	7	349	22	20,641	4
1994. . .	7,996	8	6,284	8	1,261	6	451	21	15,992	5
1995. . .	5,992	8	6,973	9	1,209	9	600	26	14,775	5
1996. . .	6,644	14	7,050	10	1,040	8	354	14	15,089	8

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	DOLPHIN								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	0	0	4,918	23	550	20	5,469	21
1982. . .	0	0	1	0	3,788	14	3,249	38	7,038	19
1983. . .	0	0	51	38	6,152	17	665	24	6,868	15
1984. . .	0	0	0	0	3,423	19	983	30	4,406	16
1985. . .	0	0	79	31	5,537	23	1,280	30	6,895	20
1986. . .	0	0	183	37	6,784	17	2,984	22	9,952	14
1987. . .	0	0	73	32	4,350	11	2,739	22	7,162	11
1988. . .	1	0	145	0	6,302	14	1,158	18	7,606	12
1989. . .	0	0	806	24	9,810	9	2,882	20	13,499	8
1990. . .	0	0	349	32	7,331	13	5,096	30	12,777	14
1991. . .	8	0	555	18	11,198	8	5,448	15	17,209	7
1992. . .	0	0	692	29	5,154	8	3,945	23	9,791	10
1993. . .	143	21	1,783	40	5,368	6	3,524	13	10,818	9
1994. . .	0	0	393	30	9,619	8	2,765	15	12,778	7
1995. . .	0	0	754	27	12,142	5	6,896	9	19,792	5
1996. . .	0	0	522	49	7,387	6	4,640	12	12,549	6

YEAR	GAG								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	0	0	86	50	332	22	418	20
1982. . .	0	0	0	0	3	0	1,308	20	1,311	20
1983. . .	0	0	0	0	91	59	1,483	18	1,574	17
1984. . .	0	0	0	0	406	32	2,642	62	3,048	54
1985. . .	0	0	0	0	161	36	1,667	22	1,828	21
1986. . .	0	0	0	0	45	49	627	22	672	21
1987. . .	0	0	0	0	504	36	978	15	1,482	16
1988. . .	0	0	0	0	222	22	1,410	18	1,632	16
1989. . .	0	0	0	0	430	26	960	16	1,391	14
1990. . .	0	0	0	0	349	46	1,432	28	1,781	24
1991. . .	0	0	0	0	220	23	2,114	16	2,334	15
1992. . .	0	0	0	0	476	16	1,729	9	2,205	8
1993. . .	0	0	0	0	544	22	2,282	8	2,827	8
1994. . .	0	0	0	0	561	21	1,947	9	2,507	9
1995. . .	0	0	0	0	303	19	2,628	8	2,930	8
1996. . .	0	0	0	0	278	21	1,964	8	2,242	8

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	GRAY SNAPPER								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	0	0	601	52	671	27	1,272	28
1982. . .	0	0	0	0	140	20	844	17	983	15
1983. . .	0	0	0	0	282	24	945	21	1,227	17
1984. . .	0	0	0	0	219	20	3,623	30	3,843	28
1985. . .	0	0	0	0	733	30	1,538	24	2,272	19
1986. . .	0	0	0	0	172	20	1,148	17	1,320	15
1987. . .	0	0	0	0	281	22	1,161	37	1,442	30
1988. . .	0	0	0	0	342	15	746	16	1,089	12
1989. . .	0	0	0	0	234	15	1,356	12	1,590	11
1990. . .	0	0	0	0	245	19	1,144	19	1,389	16
1991. . .	0	0	0	0	366	16	1,690	12	2,056	10
1992. . .	0	0	0	0	369	12	1,091	7	1,460	6
1993. . .	0	0	0	0	245	13	1,124	8	1,369	7
1994. . .	0	0	0	0	281	10	1,136	9	1,417	8
1995. . .	0	0	0	0	320	14	1,058	8	1,378	7
1996. . .	0	0	0	0	231	13	955	8	1,187	7

YEAR	GRAY TRIGGERFISH								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	0	0	82	38	648	39	730	35
1982. . .	0	0	1	0	139	26	1,010	24	1,151	22
1983. . .	0	0	12	72	64	43	239	18	315	17
1984. . .	6	0	0	0	78	21	112	21	196	15
1985. . .	0	0	24	82	74	29	183	28	281	21
1986. . .	7	0	17	38	49	26	1,338	25	1,411	23
1987. . .	0	0	69	40	36	25	1,666	33	1,772	31
1988. . .	0	0	6	78	72	24	1,366	15	1,445	14
1989. . .	3	0	119	34	257	35	1,422	15	1,800	13
1990. . .	0	0	52	23	208	28	2,095	18	2,355	16
1991. . .	3	53	137	26	165	21	2,121	14	2,426	13
1992. . .	0	0	32	25	273	17	1,325	9	1,631	8
1993. . .	1	0	106	25	177	17	1,182	11	1,465	9
1994. . .	2	78	39	32	142	13	1,165	11	1,348	10
1995. . .	0	0	65	32	97	15	1,017	9	1,179	8
1996. . .	1	65	155	43	183	15	496	11	836	11

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.
 NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	GREATER AMBERJACK								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----										
1981. . .	0	0	0	0	1,344	17	568	43	1,912	17
1982. . .	0	0	0	0	494	48	3,380	20	3,874	18
1983. . .	0	0	0	0	207	25	2,015	18	2,222	17
1984. . .	0	0	335	0	1,439	26	955	27	2,730	17
1985. . .	0	0	0	0	1,347	25	1,323	20	2,670	16
1986. . .	0	0	664	30	1,568	21	7,540	20	9,773	16
1987. . .	0	0	242	60	2,641	42	5,240	20	8,123	19
1988. . .	0	0	68	0	1,787	24	2,501	23	4,356	16
1989. . .	0	0	207	39	1,626	29	4,563	22	6,395	17
1990. . .	0	0	357	51	981	22	690	33	2,027	18
1991. . .	0	0	20	52	1,042	20	2,891	19	3,953	15
1992. . .	0	0	25	63	1,102	13	2,659	12	3,787	9
1993. . .	0	0	254	23	578	21	3,498	17	4,330	14
1994. . .	0	0	0	0	1,626	19	1,562	16	3,188	12
1995. . .	0	0	0	0	623	21	811	15	1,434	13
1996. . .	0	0	6	57	970	14	994	10	1,970	9

YEAR	KING MACKEREL								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----										
1981. . .	0	0	10	0	4,394	30	1,622	8	6,026	22
1982. . .	0	0	0	0	6,807	12	5,755	71	12,562	33
1983. . .	0	0	0	0	7,438	21	2,077	32	9,515	18
1984. . .	0	0	0	0	6,612	16	3,133	27	9,744	14
1985. . .	0	0	6	0	7,562	24	1,104	19	8,671	21
1986. . .	0	0	109	34	6,206	10	1,723	19	8,037	9
1987. . .	0	0	44	41	5,402	11	2,878	11	8,324	8
1988. . .	0	0	137	29	5,810	11	2,919	12	8,866	8
1989. . .	0	0	67	20	3,917	10	2,630	16	6,614	9
1990. . .	0	0	30	0	4,225	8	3,168	14	7,422	8
1991. . .	2	0	53	31	5,183	7	4,325	13	9,563	7
1992. . .	0	0	58	30	7,017	9	3,135	9	10,210	7
1993. . .	0	0	141	50	4,285	7	4,119	8	8,545	5
1994. . .	0	0	11	41	4,176	7	4,634	8	8,821	5
1995. . .	0	0	15	0	5,008	7	4,697	7	9,719	5
1996. . .	0	0	7	71	4,314	6	5,700	7	10,020	5

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	RED DRUM								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	352	60	409	38	3,127	14	3,888	13
1982. . .	0	0	0	0	911	20	6,415	18	7,326	16
1983. . .	0	0	54	44	1,075	16	6,878	13	8,008	11
1984. . .	0	0	1	0	2,615	41	6,879	15	9,496	15
1985. . .	0	0	0	0	2,203	32	5,435	14	7,638	13
1986. . .	0	0	900	57	1,065	13	5,189	9	7,154	10
1987. . .	0	0	44	90	1,592	12	4,569	11	6,206	9
1988. . .	0	0	9	71	1,900	16	2,932	8	4,841	8
1989. . .	0	0	30	34	885	12	5,741	8	6,655	7
1990. . .	0	0	3	100	1,355	29	4,053	9	5,411	10
1991. . .	0	0	36	32	1,247	12	5,362	9	6,645	8
1992. . .	0	0	55	44	1,116	10	7,801	4	8,972	4
1993. . .	0	0	46	11	1,168	8	8,710	5	9,924	5
1994. . .	0	0	4	0	1,439	9	7,528	5	8,970	4
1995. . .	0	0	66	0	1,661	11	11,939	4	13,665	4
1996. . .	0	0	2	0	1,333	8	11,708	5	13,043	4

YEAR	RED SNAPPER								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	0	0	383	33	3,999	26	4,382	24
1982. . .	0	0	0	0	158	27	2,198	19	2,356	18
1983. . .	0	0	0	0	199	27	4,128	20	4,327	20
1984. . .	0	0	0	0	454	17	1,426	23	1,880	18
1985. . .	0	0	0	0	1,336	29	2,457	22	3,792	18
1986. . .	0	0	0	0	114	27	2,425	17	2,538	17
1987. . .	0	0	0	0	134	20	1,647	27	1,780	25
1988. . .	0	0	0	0	225	23	1,899	22	2,124	19
1989. . .	0	0	0	0	269	28	1,814	22	2,083	19
1990. . .	0	0	0	0	115	8	1,060	16	1,175	14
1991. . .	0	0	0	0	132	34	1,503	13	1,635	13
1992. . .	0	0	0	0	618	38	2,613	8	3,231	10
1993. . .	0	0	0	0	142	27	4,136	7	4,278	7
1994. . .	0	0	0	0	201	36	3,734	8	3,935	8
1995. . .	0	0	0	0	67	20	2,876	8	2,943	8
1996. . .	0	0	0	0	102	35	2,617	8	2,719	8

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.
 NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	SAND SEATROUT								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	0	0	0	0	4,073	41	4,073	41
1982. . .	0	0	0	0	0	0	1,249	18	1,249	18
1983. . .	0	0	0	0	0	0	2,486	22	2,486	22
1984. . .	0	0	0	0	0	0	3,032	27	3,032	27
1985. . .	0	0	0	0	0	0	5,094	20	5,094	20
1986. . .	0	0	0	0	0	0	3,047	9	3,047	9
1987. . .	0	0	0	0	0	0	2,014	10	2,014	10
1988. . .	0	0	0	0	0	0	1,561	11	1,561	11
1989. . .	0	0	0	0	0	0	1,455	18	1,455	18
1990. . .	0	0	0	0	0	0	1,835	14	1,835	14
1991. . .	0	0	0	0	0	0	2,709	13	2,709	13
1992. . .	0	0	0	0	0	0	1,643	10	1,643	10
1993. . .	0	0	0	0	0	0	2,090	10	2,090	10
1994. . .	0	0	0	0	0	0	2,884	6	2,884	6
1995. . .	0	0	0	0	0	0	2,002	8	2,002	8
1996. . .	0	0	0	0	0	0	1,889	8	1,889	8

YEAR	SCUP								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	2,586	15	3,225	44	3	1	0	0	5,815	25
1982. . .	4,281	14	924	24	2	62	0	0	5,206	12
1983. . .	2,502	14	3,750	21	0	69	0	0	6,252	14
1984. . .	1,185	16	1,230	27	0	0	0	0	2,416	16
1985. . .	3,243	19	2,850	18	3	78	0	0	6,096	13
1986. . .	7,525	13	4,080	14	5	0	0	0	11,610	10
1987. . .	4,853	17	1,343	20	1	0	0	0	6,197	14
1988. . .	3,383	11	883	25	1	9	0	0	4,268	10
1989. . .	3,564	11	1,990	9	4	21	0	0	5,558	8
1990. . .	1,746	14	2,394	9	1	0	0	0	4,140	8
1991. . .	5,003	8	3,082	9	3	35	0	0	8,088	6
1992. . .	2,300	10	2,102	11	10	21	0	0	4,412	8
1993. . .	2,275	13	920	12	4	18	0	0	3,198	10
1994. . .	1,542	16	1,076	16	10	57	0	0	2,629	12
1995. . .	1,013	13	299	29	1	54	0	0	1,313	12
1996. . .	1,618	10	621	14	0	0	0	0	2,239	8

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	SHEEPSHEAD								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	17	0	480	22	1,031	24	1,528	18
1982. . .	0	0	0	0	1,875	17	1,962	13	3,837	11
1983. . .	0	0	2	0	735	21	3,288	20	4,026	17
1984. . .	0	0	0	0	1,857	14	2,812	15	4,669	10
1985. . .	0	0	0	0	888	23	2,939	23	3,827	18
1986. . .	0	0	0	0	1,402	14	2,279	30	3,680	19
1987. . .	0	0	76	73	1,891	16	1,762	12	3,730	10
1988. . .	0	0	2	0	1,569	13	4,382	10	5,953	8
1989. . .	0	0	0	0	921	12	4,953	12	5,874	10
1990. . .	0	0	0	0	1,243	13	2,478	14	3,721	10
1991. . .	0	0	2	0	2,019	11	3,068	14	5,089	10
1992. . .	0	0	0	0	2,466	10	4,897	6	7,363	5
1993. . .	0	0	0	0	1,879	13	4,577	6	6,456	6
1994. . .	0	0	0	0	2,902	10	2,734	8	5,636	6
1995. . .	0	0	9	77	2,597	11	4,901	6	7,507	6
1996. . .	0	0	5	58	1,702	8	3,348	6	5,055	5

YEAR	SOUTHERN FLOUNDER								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	0	0	160	30	822	22	982	19
1982. . .	0	0	106	72	827	59	827	15	1,760	29
1983. . .	0	0	0	0	393	17	3,055	39	3,448	35
1984. . .	0	0	3	0	421	12	359	15	783	9
1985. . .	0	0	4	65	560	13	837	22	1,401	14
1986. . .	0	0	1	0	538	15	2,619	21	3,159	18
1987. . .	0	0	0	0	289	11	415	12	706	8
1988. . .	0	0	12	36	475	14	861	10	1,348	8
1989. . .	0	0	0	0	522	21	515	18	1,038	14
1990. . .	0	0	0	0	503	12	855	15	1,358	11
1991. . .	0	0	2	0	645	12	932	10	1,579	7
1992. . .	0	0	4	60	818	9	814	8	1,637	6
1993. . .	0	0	0	0	693	8	655	9	1,348	6
1994. . .	0	0	0	0	1,132	7	664	8	1,796	5
1995. . .	0	0	0	0	924	9	715	8	1,639	6
1996. . .	0	0	1	0	478	11	780	8	1,258	7

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	SPANISH MACKEREL								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----										
1981. . .	0	0	0	0	1,290	20	1,733	15	3,023	12
1982. . .	0	0	0	0	1,209	24	2,794	20	4,004	16
1983. . .	0	0	0	0	227	29	2,646	16	2,873	15
1984. . .	0	0	0	0	1,398	31	1,061	22	2,458	20
1985. . .	0	0	0	0	698	21	1,197	12	1,895	11
1986. . .	0	0	13	58	1,195	14	6,789	15	7,997	13
1987. . .	0	0	19	48	1,685	7	2,925	10	4,630	7
1988. . .	0	0	160	39	2,939	7	2,052	9	5,151	6
1989. . .	1	0	120	23	1,386	7	1,560	8	3,066	6
1990. . .	0	0	95	24	1,630	7	2,367	8	4,092	6
1991. . .	44	34	189	15	2,240	7	2,520	8	4,992	5
1992. . .	0	0	107	17	1,841	6	3,423	6	5,371	4
1993. . .	1	0	175	18	1,262	7	1,941	7	3,378	5
1994. . .	0	0	214	14	1,158	6	1,762	6	3,135	4
1995. . .	0	0	145	29	672	8	1,630	9	2,447	6
1996. . .	0	0	81	46	986	9	1,637	7	2,704	6

YEAR	SPOT								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----										
1981. . .	0	0	5,216	15	1,700	18	42	53	6,957	12
1982. . .	0	0	2,324	17	1,662	10	16	27	4,002	11
1983. . .	0	0	2,879	21	2,119	19	30	48	5,029	15
1984. . .	0	0	776	18	1,024	13	6	5	1,806	11
1985. . .	0	0	2,156	9	3,788	14	7	69	5,952	9
1986. . .	0	0	2,352	10	1,041	16	33	29	3,426	8
1987. . .	0	0	2,882	10	965	9	54	39	3,901	8
1988. . .	0	0	888	13	1,634	14	30	62	2,553	10
1989. . .	0	0	2,045	7	1,249	8	0	0	3,294	6
1990. . .	0	0	2,912	9	672	9	54	83	3,639	7
1991. . .	0	0	3,432	8	1,082	8	26	56	4,540	7
1992. . .	0	0	3,048	11	976	8	43	47	4,067	9
1993. . .	0	0	1,591	11	1,780	7	108	56	3,479	7
1994. . .	0	0	1,956	6	2,372	8	55	42	4,383	5
1995. . .	0	0	1,571	10	1,536	7	50	42	3,158	6
1996. . .	0	0	832	12	1,214	12	5	51	2,051	9

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	SPOTTED SEATRUT								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	0	0	1,184	33	4,652	14	5,836	13
1982. . .	0	0	0	0	1,547	16	11,440	10	12,988	9
1983. . .	0	0	0	0	1,343	16	11,161	11	12,505	10
1984. . .	0	0	0	0	1,163	13	8,606	20	9,770	17
1985. . .	0	0	0	0	2,065	14	6,471	12	8,536	10
1986. . .	0	0	69	21	2,052	12	16,966	18	19,087	16
1987. . .	0	0	61	31	2,334	10	11,865	5	14,260	5
1988. . .	0	0	497	24	1,829	10	12,610	5	14,936	4
1989. . .	0	0	297	15	1,474	9	12,798	8	14,569	7
1990. . .	0	0	160	32	1,357	15	4,847	7	6,364	6
1991. . .	0	0	157	19	3,085	10	12,052	6	15,294	5
1992. . .	0	0	64	23	1,710	7	9,545	5	11,320	4
1993. . .	0	0	214	22	1,554	7	8,295	5	10,064	4
1994. . .	0	0	202	14	1,633	6	9,342	4	11,178	3
1995. . .	0	0	178	18	2,030	8	10,361	5	12,569	4
1996. . .	0	0	78	22	803	8	9,665	5	10,545	4
YEAR	STRIPED BASS								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	215	28	931	26	807	53	0	0	1,953	25
1982. . .	2,134	52	388	33	0	0	1	0	2,523	44
1983. . .	596	31	2,087	71	15	99	0	0	2,698	55
1984. . .	138	50	1,139	43	5	7	1	0	1,284	38
1985. . .	407	54	414	36	25	50	3	0	849	31
1986. . .	418	31	686	39	4	43	20	49	1,128	26
1987. . .	356	31	499	20	2	40	5	59	862	17
1988. . .	591	20	665	27	28	33	38	34	1,321	16
1989. . .	470	22	262	26	8	0	1	0	742	17
1990. . .	657	24	1,569	14	0	0	69	27	2,295	12
1991. . .	1,128	17	2,513	11	4	0	62	28	3,706	9
1992. . .	1,524	18	2,494	13	21	15	28	36	4,067	10
1993. . .	1,846	10	3,803	12	16	38	1	0	5,666	9
1994. . .	2,284	10	4,443	9	77	31	9	37	6,813	7
1995. . .	2,657	8	8,365	9	237	15	5	17	11,264	7
1996. . .	3,441	9	10,885	6	288	13	6	36	14,619	5

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	SUMMER FLOUNDER								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	389	34	9,221	9	488	24	0	0	10,098	8
1982. . .	2,310	16	14,314	32	1,640	23	0	0	18,264	25
1983. . .	1,322	17	26,100	8	586	21	0	0	28,008	7
1984. . .	900	16	16,559	8	1,378	29	0	0	18,837	8
1985. . .	744	34	10,883	12	863	21	0	0	12,490	11
1986. . .	5,880	19	10,817	9	1,177	44	0	0	17,885	9
1987. . .	1,338	20	10,634	10	257	11	0	0	12,228	9
1988. . .	712	12	13,302	4	644	9	0	0	14,658	4
1989. . .	316	17	2,559	7	301	13	0	0	3,176	6
1990. . .	235	21	4,376	5	531	11	0	0	5,142	4
1991. . .	356	14	7,370	4	432	16	0	0	8,158	4
1992. . .	430	13	6,459	5	267	8	0	0	7,157	4
1993. . .	552	11	7,812	5	479	7	0	0	8,844	4
1994. . .	980	9	7,886	4	481	7	0	0	9,347	4
1995. . .	858	11	4,394	5	251	23	0	0	5,503	5
1996. . .	1,232	8	8,777	4	407	7	0	0	10,416	3

YEAR	TAUTOG								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	1,698	26	2,417	24	1	0	0	0	4,116	18
1982. . .	4,615	15	3,707	14	16	67	0	0	8,338	10
1983. . .	2,911	14	2,818	19	21	67	0	0	5,751	12
1984. . .	3,277	16	2,104	20	0	0	0	0	5,381	12
1985. . .	1,077	16	3,221	17	7	65	0	0	4,305	14
1986. . .	10,744	14	6,159	13	4	60	0	0	16,906	10
1987. . .	3,365	17	5,515	12	8	0	0	0	8,889	10
1988. . .	3,478	18	5,819	17	5	33	0	0	9,302	13
1989. . .	2,413	11	3,935	9	31	35	0	0	6,379	7
1990. . .	1,485	11	3,669	9	3	31	0	0	5,156	7
1991. . .	2,459	11	5,622	7	25	54	0	0	8,105	6
1992. . .	3,374	14	4,285	10	13	35	0	0	7,671	8
1993. . .	1,673	10	4,244	11	10	64	0	0	5,927	8
1994. . .	1,119	14	2,346	19	3	0	0	0	3,468	13
1995. . .	959	16	3,641	10	3	33	0	0	4,604	9
1996. . .	918	13	2,336	13	12	31	0	0	3,266	10

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	VERMILION SNAPPER								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	0	0	0	0	5	43	86	21	91	20
1982. . .	0	0	0	0	233	34	271	20	504	19
1983. . .	0	0	0	0	416	45	64	19	480	39
1984. . .	0	0	0	0	210	17	155	21	365	13
1985. . .	0	0	0	0	435	26	246	64	681	29
1986. . .	0	0	0	0	11	31	932	40	944	39
1987. . .	0	0	0	0	207	41	491	30	699	24
1988. . .	0	0	0	0	135	28	467	17	602	14
1989. . .	0	0	0	0	106	23	411	20	517	17
1990. . .	0	0	0	0	121	28	518	26	639	22
1991. . .	0	0	0	0	105	26	829	22	934	20
1992. . .	0	0	0	0	118	22	602	10	720	9
1993. . .	0	0	0	0	98	23	759	11	857	10
1994. . .	0	0	0	0	73	19	608	13	681	12
1995. . .	0	0	0	0	44	21	564	11	608	10
1996. . .	0	0	0	0	79	45	229	14	309	15

YEAR	WEAKFISH								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	281	33	15,724	16	99	27	0	0	16,105	16
1982. . .	155	50	7,792	16	339	43	0	0	8,285	15
1983. . .	624	57	10,403	13	703	21	0	0	11,731	12
1984. . .	28	68	6,424	23	562	47	0	0	7,014	21
1985. . .	149	57	5,025	11	315	28	0	0	5,489	11
1986. . .	102	54	9,464	10	575	19	0	0	10,142	9
1987. . .	4	79	5,957	16	789	23	0	0	6,750	14
1988. . .	0	0	5,881	12	450	14	0	0	6,332	11
1989. . .	0	0	1,912	9	265	14	0	0	2,177	8
1990. . .	1	63	1,215	9	131	14	0	0	1,347	8
1991. . .	0	0	1,919	8	212	20	0	0	2,131	7
1992. . .	21	110	1,258	9	120	16	0	0	1,399	9
1993. . .	7	59	887	11	209	13	0	0	1,102	9
1994. . .	0	0	1,471	11	324	11	0	0	1,796	10
1995. . .	0	0	1,689	8	163	18	0	0	1,852	7
1996. . .	0	0	2,849	7	102	10	0	0	2,951	7

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.

U.S. MARINE RECREATIONAL FISHERIES

ESTIMATED WEIGHT (LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES, 1981-1996 FOR THE ATLANTIC AND GULF COASTS

YEAR	WINTER FLOUNDER								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	6,751	15	5,673	14	0	0	0	0	12,424	10
1982. . .	11,865	37	4,553	12	0	0	0	0	16,417	27
1983. . .	3,781	11	5,857	14	2	0	0	0	9,640	10
1984. . .	5,012	11	10,145	15	0	0	0	0	15,157	11
1985. . .	6,608	15	8,765	24	0	0	0	0	15,373	15
1986. . .	4,039	19	3,596	9	0	0	0	0	7,635	11
1987. . .	5,663	15	5,305	13	0	0	0	0	10,967	10
1988. . .	3,243	17	5,537	13	0	0	0	0	8,780	10
1989. . .	3,180	18	2,184	15	0	0	0	0	5,363	12
1990. . .	1,651	14	1,505	10	0	0	0	0	3,156	9
1991. . .	769	13	2,131	13	0	0	0	0	2,899	10
1992. . .	430	12	641	11	0	0	0	0	1,072	8
1993. . .	559	12	1,570	30	0	0	0	0	2,130	22
1994. . .	393	12	1,104	17	0	0	0	0	1,497	13
1995. . .	389	23	1,161	13	0	0	0	0	1,549	11
1996. . .	328	12	1,383	14	0	0	0	0	1,712	11
YEAR	ALL FISHES								TOTAL	
	NORTH ATLANTIC		MID ATLANTIC		SOUTH ATLANTIC		GULF OF MEXICO (1)			
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
	----- THOUSANDS OF POUNDS -----									
1981. . .	68,790	11	118,575	5	37,873	7	52,978	6	278,215	4
1982. . .	85,706	11	105,417	8	48,533	4	75,703	7	315,360	4
1983. . .	68,382	12	124,645	6	65,202	10	80,920	5	339,150	4
1984. . .	39,631	7	100,531	6	50,050	5	71,753	7	261,966	3
1985. . .	59,426	23	79,398	7	59,956	7	65,446	5	264,227	6
1986. . .	81,972	6	135,525	5	53,561	5	96,564	5	367,623	3
1987. . .	55,168	6	116,717	4	51,555	5	66,536	4	289,976	2
1988. . .	39,726	6	85,891	4	54,851	4	70,845	3	251,314	2
1989. . .	33,100	5	76,967	4	46,353	3	66,904	3	223,324	2
1990. . .	28,887	8	56,797	4	35,772	4	51,548	4	173,004	2
1991. . .	35,631	6	65,186	3	47,656	3	79,768	3	228,241	2
1992. . .	21,169	5	47,334	3	44,967	3	68,931	2	182,401	2
1993. . .	24,298	5	55,082	4	37,346	2	68,519	2	185,244	2
1994. . .	23,915	6	45,859	3	50,089	3	63,572	2	183,436	1
1995. . .	19,788	6	58,870	8	50,444	2	73,055	2	202,157	3
1996. . .	21,286	6	55,744	3	43,757	3	64,567	2	185,354	2

(1) GULF OF MEXICO ESTIMATES DO NOT INCLUDE TEXAS DATA.

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR.