

## KILLER WHALE (*Orcinus orca*): Western North Atlantic Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE

Killer whales are characterized as uncommon or rare in waters of the U.S. Atlantic Exclusive Economic Zone (EEZ) (Katona et al. 1988). The 12 killer whale sightings constituted 0.1% of the 11,156 cetacean sightings in the 1978-81 CETAP surveys (CETAP 1982). The same may be true for eastern Canadian waters, where the species has been described as relatively uncommon and numerically few (Mitchell and Reeves 1988). Lawson and Stevens (2013) reported on eastern Canada killer whale sighting events from 1758 to 2012, and found that sightings were most common from June to September and especially more frequent over the last ten years. This is possibly due to increased public awareness of this species, and more boats, people and cameras on the water during those months. In eastern Canada 17.4% and in the U.S. Gulf of Maine 9.3% of humpbacks had scars on their flukes caused by non-fatal predatory interactions with killer whales. This may be due to migration patterns or may reflect dietary differences and relative distributions of different ecotypes of killer whales (McCordic et al. 2013). Killer whale distribution extends from the Arctic ice edge to the West Indies. They are normally found in small groups, although 40 animals were reported from the southern Gulf of Maine in September 1979, and 29 animals in Massachusetts Bay in August 1986 (Katona et al. 1988). In the U.S. Atlantic EEZ, while their occurrence is unpredictable, they do occur in fishing areas, perhaps coincident with tuna, in warm seasons (Katona et al. 1988; NMFS unpublished data). In an extensive analysis of historical whaling records, Reeves and Mitchell (1988) plotted the distribution of killer whales in offshore and mid-ocean areas. Their results suggest that the offshore areas need to be considered in present-day distribution, movements, and stock relationships.

Stock and ecotype definitions are largely unknown. Results from other areas (e.g., the Pacific Northwest and Norway) suggest that social structure and territoriality may be important.

### POPULATION SIZE

The total number of killer whales off the eastern U.S. coast is unknown.

#### Minimum Population Estimate


Present data are insufficient to calculate a minimum population estimate.

#### Current Population Trend

There are insufficient data to determine the population trends for this species.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are not known for this stock. The maximum net productivity rate was assumed to be 0.04 for purposes of this assessment. This value is based on theoretical calculations showing that cetacean populations may not generally grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow et al. 1995).



**Figure 1.** Distribution of killer whale sightings from NEFSC and SEFSC shipboard and aerial surveys during the summer in 1998, 1999, 2002, 2004, 2006 and 2011. Isobaths are the 100m, 1,000m, and 4,000m depth contours.

## POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (Wade and Angliss 1997). The minimum population size is unknown. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5 because this stock is of unknown. PBR for the western North Atlantic killer whale is unknown because the minimum population size cannot be determined.

## ANNUAL HUMAN-CAUSED MORTALITY

In 1994, one killer whale was caught in the New England multispecies sink gillnet fishery but released alive. Known mortality events in eastern Canada (DFO, unpublished data; Lawson and Stevens 2013), for the last 40 years equate to at least one killer whale death every 2 years.

## Fishery Information

Data on current incidental takes in U.S. fisheries are available from several sources. In 1986, NMFS established a mandatory self-reported fishery information system for large pelagic fisheries. Data files are maintained at the Southeast Fisheries Science Center (SEFSC). The Northeast Fisheries Science Center (NEFSC) Sea Sampling Observer Program was initiated in 1989, and since that year several fisheries have been covered by the program. In late 1992 and in 1993, the SEFSC provided observer coverage of pelagic longline vessels fishing off the Grand Banks (Tail of the Banks) and provides observer coverage of vessels fishing south of Cape Hatteras.

There have been no observed mortalities or serious injuries by NMFS Sea Samplers in the pelagic drift gillnet, pelagic longline, pelagic pair trawl, New England multispecies sink gillnet, mid-Atlantic coastal sink gillnet, and North Atlantic bottom trawl fisheries.

## STATUS OF STOCK

The status of killer whales relative to OSP in U.S. Atlantic EEZ is unknown. Because there are no observed mortalities or serious injury between 2008 and 2012, the total U.S. fishery-related mortality and serious injury for this stock is considered insignificant and approaching zero mortality and serious injury rate. The species is not listed as threatened or endangered under the Endangered Species Act. In Canada, the Cetacean Protection Regulations of 1982, promulgated under the standing Fisheries Act, prohibit the catching or harassment of all cetacean species. There are insufficient data to determine the population trends for this species. This is not a strategic stock because, although PBR could not be calculated, there is no evidence of human-induced mortality.

## REFERENCES

- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade 1995. U.S. Marine Mammal Stock Assessments: Guidelines for Preparation, Background, and a Summary of the 1995 Assessments. NOAA Tech. Memo. NMFS-OPR-6, 73 pp.
- CETAP (Cetacean and Turtle Assessment Program, University of Rhode Island). 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf, final report. #AA 551-C T8-48. Bureau of Land Management, Washington, DC, 538 pp.
- Katona, S.K., J.A. Beard, P.E. Girton and F. Wenzel 1988. Killer whales (*Orcinus orca*) from the Bay of Fundy to the Equator, including the Gulf of Mexico. Rit. Fiskideild. 9: 205-224.
- Lawson, J.W. and T.S. Stevens 2013. Historic and current distribution patterns, and minimum abundance, of killer whales (*Orcinus orca*) in the north-west Atlantic. J. Mar. Biol. Assoc. U.K. DOI: <http://dx.doi.org/10.1017/S0025315413001409>
- McCordic, J.A., S.K. Todd and P.T. Stevick. 2013. Differential rates of killer whale attacks on humpback whales in the North Atlantic as determined by scarification. J. Mar. Biol. Assoc. U.K. DOI: <http://dx.doi.org/10.1017/S0025315413001008>
- Mitchell, E. and R.R. Reeves 1988. Records of killer whales in the western North Atlantic, with emphasis on eastern Canadian waters. Rit. Fiskideild. 9: 161-193.
- Reeves, R.R. and E. Mitchell. 1988. Killer whale sightings and takes by American pelagic whalers in the North Atlantic. Rit. Fiskideild. 9: 7-23.

Wade P.R., and R.P. Angliss.1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12, 93 pp.