

## SPERM WHALE (*Physeter macrocephalus*): North Pacific Stock


### STOCK DEFINITION AND GEOGRAPHIC RANGE

The sperm whale is one of the most widely distributed marine mammal species, perhaps exceeded in its global range only by the killer whale (Rice 1989). In the North Pacific, sperm whales were depleted by extensive commercial whaling over a period of more than a hundred years, and the species was the primary target of illegal Soviet whaling in the second half of the 20th century (Ivashchenko et al. 2013, 2014).

Sperm whales feed primarily on medium-sized to large-sized squids but also take substantial quantities of large demersal and mesopelagic sharks, skates, and fishes (Rice 1989). In the North Pacific, sperm whales are distributed widely (Fig. 1), with the northernmost boundary extending from Cape Navarin (62°N) to the Pribilof Islands (Omura 1955). Although females and young sperm whales were thought to remain in tropical and temperate waters year-round, Mizroch and Rice (2006) and Ivashchenko et al. (2014) showed that there were extensive catches of female sperm whales above 50°N, in the western Bering Sea and in the western Aleutian Islands. Mizroch and Rice (2013) also showed female movements into the Gulf of Alaska and western Aleutians. Males are found in the summer in the Gulf of Alaska, Bering Sea, and waters around the Aleutian Islands (Kasuya and Miyashita 1988, Mizroch and Rice 2013, Ivashchenko et al. 2014). Sighting surveys conducted by the Alaska Fisheries Science Center's National Marine Mammal Laboratory in the summer months between 2001 and 2010 have found sperm whales to be the most frequently sighted large cetacean in the coastal waters around the central and western Aleutian Islands (NMML, unpubl. data). Acoustic surveys detected the presence of sperm whales year-round in the Gulf of Alaska although they appear to be more common in summer than in winter (Mellinger et al. 2004). These seasonal detections are consistent with the hypothesis that sperm whales move to higher latitudes in summer and to lower latitudes in winter (Whitehead and Arnobom 1987).

Mizroch and Rice (2013) examined 261 Discovery mark recoveries from the days of commercial whaling and found extensive movements from U.S. and Canadian coastal waters into the Gulf of Alaska and Bering Sea. The U.S. marked 176 sperm whales from 1962 to 1969 off southern California and northern Baja California (Mizroch and Rice 2013). Seven of those marked whales were recovered in locations ranging from offshore California, Oregon, and British Columbia waters to the western Gulf of Alaska. A male whale marked by Canadian researchers moved from near Vancouver Island, British Columbia, to the Aleutian Islands near Adak. A whale marked by Soviet researchers moved from coastal Michoacán, mainland Mexico, to a location about 1,300 km offshore of Washington State. Similar extensive movements have also been demonstrated by recent satellite-tagging studies (Straley et al. 2014). Three adult males satellite-tagged off southeastern Alaska moved far south, one to coastal Baja California, one into the north-central Gulf of California, and the third to a location near the Mexico-Guatemala border (Straley et al. 2014). Marking data show extensive movements throughout the North Pacific and along the U.S. west coast into the Gulf of Alaska and Bering Sea/Aleutian Islands region.

Mizroch and Rice (2013) also analyzed whaling data and found that males and females concentrated seasonally along oceanic frontal zones, for example, in the subtropical frontal zone (ca. 28-34°N) and the subarctic frontal zones (ca. 40-43°N). Males also concentrated seasonally near the Aleutian Islands and along the Bering Sea shelf edge. Their analyses of marking and whaling data indicate that there are no apparent divisions between separate demes or stocks within the North Pacific. Analysis of Soviet catch data by Ivashchenko et al. (2014)



**Figure 1.** The approximate distribution of sperm whales in the North Pacific includes deep waters south of 62°N to the equator.

showed broad agreement with these results, although a sharp division was evident at Amchitka Pass in the Aleutians, with mature males to the east and males and family groups to the west, including in the Commander Islands. There were four main areas of concentration in the Soviet catches: a large pelagic area (30-50°N) in the eastern North Pacific, including the Gulf of Alaska and western coast of North America; the northeastern and southwestern central North Pacific; and the southern Kuril Islands. Some of the catch distribution was similar to that of 19th century Yankee whaling catches plotted by Townsend (1935), notably in the “Japan Ground” (in the pelagic western Pacific) and the “Coast of Japan Ground.” Many females were caught in Olyutorsky Bay (western Bering Sea) and around the Commander Islands.

The following information was considered in classifying stock structure based on the Dizon et al. (1992) phylogeographic approach: 1) Distributional data: no apparent discontinuities based on whale marking data; 2) Population response data: unknown; 3) Phenotypic data: unknown; and 4) Genotypic data: genetics studies indicate the possibility of a “somewhat” discrete U.S. coastal stock (Mesnick et al. 2011). For management purposes, the International Whaling Commission (IWC) recognizes two management units of sperm whales in the North Pacific (eastern and western). However, the IWC has not reviewed its sperm whale stock boundaries in recent years (Donovan 1991). For management purposes, three stocks of sperm whales are currently recognized in U.S. waters: 1) Alaska (North Pacific stock), 2) California/Washington/Oregon, and 3) Hawaii. New information from Mizroch and Rice (2013) suggests that this structure should be reviewed and updated to reflect current data. The California/Oregon/Washington and Hawaii sperm whale stocks are reported separately in the Stock Assessment Reports for the U.S. Pacific Region.

### **POPULATION SIZE**

Current and historical estimates of the abundance of sperm whales in the North Pacific are considered unreliable, and caution should be exercised in interpreting published estimates. The abundance of sperm whales in the North Pacific was reported to be 1,260,000 prior to exploitation, which by the late 1970s was estimated to have been reduced to 930,000 whales (Rice 1989). Confidence intervals for these estimates were not provided. These estimates include whales from the California/Oregon/Washington stock, for which a separate abundance estimate is currently available (see Stock Assessment Reports for the U.S. Pacific Region). Estimates for a large area of the eastern temperate North Pacific were produced from line-transect and acoustic survey data by Barlow and Taylor (2005), but no recent estimate exists for other areas, including for the central or western North Pacific.

Although Kato and Miyashita (1998) believe their estimate to be positively biased, their analysis suggested 102,112 (CV = 0.155) sperm whales in the western North Pacific. The number of sperm whales occurring within Alaska waters is unknown.

As the data used in estimating the abundance of sperm whales in the entire North Pacific are more than 8 years old at this time, and there are no available estimates for numbers of sperm whales in Alaska waters, a reliable estimate of abundance for the North Pacific stock is not available.

### **Minimum Population Estimate**

At this time, it is not possible to produce a reliable estimate of minimum abundance for this stock, as a current estimate of abundance is not available.

### **Current Population Trend**

No current estimate of abundance exists for this stock; therefore, reliable information on trends in abundance for this stock is currently not available (Braham 1992).

### **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

A reliable estimate of the maximum net productivity rate is not currently available for the North Pacific stock of sperm whales. Hence, until additional data become available, it is recommended that the cetacean maximum net productivity rate ( $R_{MAX}$ ) of 4% be employed for this stock at this time (Wade and Angliss 1997).

## **POTENTIAL BIOLOGICAL REMOVAL**

Under the 1994 reauthorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor:  $PBR = N_{MIN} \times 0.5R_{MAX} \times F_R$ . The recovery factor ( $F_R$ ) for this stock is 0.1, the value for cetacean stocks which are classified as “endangered” (Wade and Angliss 1997). However, because a reliable estimate of minimum abundance ( $N_{MIN}$ ) is currently not available, the PBR for this stock is unknown.

## **ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY**

### **Fisheries Information**

Detailed information (including observer programs, observer coverage, and observed incidental takes of marine mammals) for federally-managed and state-managed U.S. commercial fisheries in Alaska waters is presented in Appendices 3-6 of the Alaska Stock Assessment Reports.

Between 2009 and 2013, there were four serious injuries of sperm whales observed in the Gulf of Alaska sablefish longline fishery (two each in 2012 and 2013), resulting in an average annual observed mortality and serious injury of 0.8 sperm whales in U.S. commercial groundfish fisheries in 2009-2013 (Helker et al. 2015). Extrapolations based on observer effort are not available at this time.

### **Alaska Native Subsistence/Harvest Information**

Sperm whales have never been reported to be taken by subsistence hunters (Rice 1989).

### **Other Mortality**

Sperm whales were the dominant species killed by the commercial whaling industry as it developed in the North Pacific in the years after the Second World War (Mizroch and Rice 2006, Ivashchenko et al. 2014). Between 1946 and 1967, most of the sperm whales were caught in waters near Japan and in the Bering Sea/Aleutian Islands (BSAI) region. The BSAI catches were dominated by males. After 1967, whalers moved out of the BSAI region and began to catch even larger numbers of sperm whales further south in the North Pacific between 30° and 50°N (Mizroch and Rice 2006: Figs. 7-9). The reported catch of sperm whales taken by commercial whalers operating in the North Pacific between 1912 and 2006 was 261,148 sperm whales, of which, 259,120 were taken between 1946 and 1987 (IWC, Bureau of International Whaling Statistics (BIWS) catch data, February 2008 version, unpubl.). This value underestimates the actual kill in the North Pacific as a result of under-reporting by U.S.S.R. pelagic whaling operations. Berzin (2008) described extreme under-reporting and misreporting of Soviet sperm whale catches from the mid-1960s into the early 1970s, including enormous (and under-reported) whaling pressure on female sperm whales in the latter years of whaling. More recently, Ivashchenko et al. (2013, 2014) estimate that 157,680 sperm whales were killed by the U.S.S.R. in the North Pacific between 1948 and 1979, of which 25,175 were unreported; the Soviets also extensively misreported the sex and length of catches. In addition, new information indicates that Japanese land-based whaling operations also misreported the number and sex of sperm whale catches during the post-World War II era (Kasuya 1999). The last year that the U.S.S.R. reported catches of sperm whales was in 1979 and the last year that Japan reported substantial catches was in 1987, but Japanese whalers reported catches of 48 sperm whales between 2000 and 2009 (IWC, BIWS catch data, October 2010 version, unpubl.). It should be noted that the reliability of data concerning large pelagic catches of sperm whales by Japan in the North Pacific is unknown, but analysis of length distribution data suggests at least some degree of systematic misreporting (Cooke et al. 1983). Thus, studies that use Japanese data to assess the North Pacific distribution of this species, including by sex, should be interpreted with caution.

From 2009 to 2013, one suspected human-related sperm whale mortality was reported to the NMFS Alaska Region stranding database (Helker et al. 2015). A beachcast sperm whale was found in 2012 on a beach near Yakutat with a net from an unknown fishery wrapped around its lower jaw. However, due to the advanced decomposition of this whale, the cause of death could not be determined.

### **Other Issues**

NMFS observers aboard longline vessels targeting both sablefish and halibut have documented sperm whales feeding off longline gear in the Gulf of Alaska (Hill and Mitchell 1998, Hill et al. 1999, Perez 2006, Sigler et al. 2008). Fishery observers recorded several instances during 1995-1997 in which sperm whales were deterred by fishermen (i.e., yelling at the whales or throwing seal bombs in the water).

Annual longline surveys have been recording sperm whale predation on catch since 1998 (Hanselman et al. 2008). Sperm whale depredation in the sablefish longline fishery is widespread in the central and eastern Gulf of

Alaska but rarely observed in the Bering Sea; the majority of interactions occur in the West Yakutat and East Yakutat/Southeast areas (Perez 2006, Hanselman et al. 2008). Sigler et al. (2008) analyzed catch data from 1998 to 2004 and found that catch rates were about 2% less at locations where depredation occurred, but the effect was not significant ( $p = 0.34$ ). Hill et al. (1999) analyzed data collected by fisheries observers in Alaska waters and also found no significant effect on catch. A small, significant effect on catch rates was found in a study using data collected in Southeast Alaska, in which longline fishery catches in sets with sperm whales present were compared to catches in sets with sperm whales absent (3% reduction, t-test, 95% CI of 0.4-5.5%,  $p = 0.02$ ; Straley et al. 2005). Undamaged catches may also occur when sperm whales are present; in these cases, sperm whales apparently feed off the discard.

## STATUS OF STOCK

Sperm whales are listed as “endangered” under the Endangered Species Act of 1973, and therefore designated as “depleted” under the MMPA. As a result, this stock is classified as a strategic stock. However, on the basis of total abundance, current distribution, and regulatory measures that are currently in place, it is unlikely that this stock is in danger of extinction (Braham 1992). Reliable estimates of the minimum population, population trends, PBR, and status of the stock relative to its Optimum Sustainable Population are currently not available, although the estimated annual rate of human-caused mortality and serious injury seems minimal for this stock. Because the PBR is unknown, the level of annual U.S. commercial fishery-related mortality and serious injury that can be considered insignificant and approaching zero mortality and serious injury rate is unknown.

## HABITAT CONCERNS

There are no known habitat issues that are of particular concern for this stock. However, potential concerns include elevated levels of sound from anthropogenic sources (e.g., shipping, military sonars), possible changes in prey distribution with climate change, entanglement in fishing gear, ship strikes due to increased vessel traffic (e.g., from increased shipping in higher latitudes), and oil and gas activities.

## CITATIONS

- Barlow J., and B. L. Taylor. 2005. Estimates of sperm whale abundance in the northeastern temperate Pacific from a combined acoustic and visual survey. *Mar. Mammal Sci.* 21:429-445.
- Berzin A. A. 2008. The truth about Soviet whaling: a memoir. *Mar. Fish. Rev.* 70(2):4-59.
- Braham, H. 1992. Endangered whales: status update. Working document presented at A Workshop on the Status of California Cetacean Stocks (SOCCS/14). 35 pp. + tables. Available from NMML, Alaska Fisheries Science Center, NMFS, 7600 Sand Point Way NE, Seattle, WA 98115.
- Cooke, J. G., W. K. de la Mare, and J. R. Beddington. 1983. Some aspects of the reliability of the length data for the western North Pacific stock of sperm whales. *Rep. Int. Whal. Comm.* 33:265-267.
- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. *Conserv. Biol.* 6:24-36.
- Donovan, G. P. 1991. A review of IWC stock boundaries. *Rep. Int. Whal. Comm. (Special Issue 13)*:39-68.
- Hanselman, D. H., C. R. Lunsford, J. T. Fujioka, and C. J. Rodgveller. 2008. Assessment of the sablefish stock in Alaska. Section 3, Pp. 303-420 *In* Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska. North Pacific Fisheries Management Council, Anchorage, AK.
- Helker, V. T., B. M. Allen, and L. A. Jemison. 2015. Human-caused injury and mortality of NMFS-managed Alaska marine mammal stocks, 2009-2013. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-300, 94 p.
- Hill, P. S., and E. Mitchell. 1998. Sperm whale interactions with longline vessels in Alaska waters during 1997. Unpubl. doc. submitted to Fish. Bull., U.S. Available from S. Mizroch, NMML, Alaska Fisheries Science Center, NMFS, 7600 Sand Point Way NE, Seattle, WA 98115.
- Hill, P. S., J. L. Laake, and E. Mitchell. 1999. Results of a pilot program to document interactions between sperm whales and longline vessels in Alaska waters. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-108, 42 p.
- Ivashchenko, Y. V., P. J. Clapham, and R. L. Brownell, Jr. 2013. Soviet catches of whales in the North Pacific: revised totals. *J. Cetacean Res. Manage.* 13(1):59-71.
- Ivashchenko, Y. V., R. L. Brownell, Jr., and P. J. Clapham. 2014. Distribution of Soviet catches of sperm whales *Physeter macrocephalus* in the North Pacific. *Endang. Species Res.* 25:249-263.
- Kasuya, T. 1999. Examination of the reliability of catch statistics in the Japanese coastal sperm whale fishery. *J. Cetacean Res. Manage.* 1:109-122.

- Kasuya T., and T. Miyashita. 1988. Distribution of sperm whale stocks in the North Pacific. *Sci. Rep. Whales Res. Inst.* 39: 31-75.
- Kato, H., and T. Miyashita. 1998. Current status of North Pacific sperm whales and its preliminary abundance estimates. Unpubl. doc. submitted to Int. Whal. Comm. (SC/50/CAWS/52). 6 pp.
- Mellinger, D. K., K. M. Stafford, and C. G. Fox. 2004. Seasonal occurrence of sperm whale (*Physeter macrocephalus*) sounds in the Gulf of Alaska, 1999-2001. *Mar. Mammal Sci.* 20(1):48-62.
- Mesnick S. L., B. L. Taylor, F. I. Archer, K. K. Martien, S. Escorza Treviño, B. L. Hancock-Hanser, S. C. Moreno Medina, V. L. Pease, K. M. Robertson, J. M. Straley, R. W. Baird, J. Calambokidis, G. S. Schorr, P. Wade, V. Burkanov, C. R. Lunsford, L. Rendell, and P. A. Morin 2011. Sperm whale population structure in the eastern and central North Pacific inferred by the use of single nucleotide polymorphisms, microsatellites and mitochondrial DNA. *Mol. Ecol. Res.* 11(Suppl. 1):278-298.
- Mizroch S. A., and D. W. Rice. 2006. Have North Pacific killer whales switched prey species in response to depletion of the great whale populations? *Mar. Ecol. Prog. Ser.* 310:235-246.
- Mizroch S. A., and D. W. Rice. 2013. Ocean nomads: distribution and movements of sperm whales in the North Pacific shown by whaling data and Discovery marks. *Mar. Mammal Sci.* 29(2):E136-E165. DOI: 10.1111/j.1748-7692.2012.00601.x.
- Omura, H. 1955. Whales in the northern part of the North Pacific. *Nor. Hvalfangst-tidende* 44(6):323-345.
- Perez, M. A. 2006. Analysis of marine mammal bycatch data from the trawl, longline, and pot groundfish fisheries of Alaska, 1998-2004, defined by geographic area, gear type, and target groundfish catch species. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-167, 194 p.
- Rice, D. W. 1989. Sperm whale, *Physeter macrocephalus*. Pp. 177-233 In S. H. Ridgway and R. Harrison (eds.), *Handbook of Marine Mammals*. Vol. 4. River Dolphins and the Larger Toothed Whales. Academic Press, New York.
- Sigler, M. F., C. R. Lunsford, J. M. Straley, and J. B. Liddle. 2008. Sperm whale depredation of sablefish longline gear in the northeast Pacific Ocean. *Mar. Mammal Sci.* 24(1):16-27.
- Straley, J., T. O'Connell, S. Mesnick, L. Behnken, and J. Liddle. 2005. Sperm whale and longline fisheries interactions in the Gulf of Alaska. North Pacific Research Board R0309 Final Report. 15 pp.
- Straley, J. M., G. S. Schorr, A. M. Thode, J. Calambokidis, C. R. Lunsford, E. M. Chenoweth, V. M. O'Connell, and R. D. Andrews. 2014. Depredating sperm whales in the Gulf of Alaska: local habitat use and long distance movements across putative population boundaries. *Endang. Species Res.* 24:125-135.
- Townsend, C. 1935. The distribution of certain whales as shown by logbook records of American whalships. *Zoologica* 19:1-50.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 p.
- Whitehead, H., and T. Arnbohm. 1987. Social organization of sperm whale off the Galapagos Island, February-April 1985. *Can. J. Zool.* 65(4):913-919.