

DWARF SPERM WHALE (*Kogia sima*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

The dwarf sperm whale (*Kogia sima*) is distributed worldwide in temperate to tropical waters (Caldwell and Caldwell 1989; McAlpine 2002). Sightings of these animals in the western North Atlantic occur in oceanic waters (Figure 1; Mullin and Fulling 2003; NMFS unpublished data). Stranding records exist from Florida to Maine, but there are no stranding records for the east Canadian coast (Willis and Baird 1998). Dwarf sperm whales and pygmy sperm whales (*K. breviceps*) are difficult to differentiate at sea (Caldwell and Caldwell 1989, Wursig *et al.* 2000), and sightings of either species are often categorized as *Kogia* sp. Diagnostic morphological characters have been useful in distinguishing the two *Kogia* species (Barros and Duffield 2003), thus enabling researchers to use stranding data in distributional and ecological studies. Specifically, the distance from the snout to the center of the blowhole in proportion to the animal's total length, as well as the height of the dorsal fin in proportion to the animal's total length, can be used to differentiate between the two *Kogia* species when such measurements are obtainable (Barros and Duffield 2003; Handley 1966). Duffield *et al.* (2003) proposed using the molecular weights of myoglobin and hemoglobin, as determined by blood or muscle tissues of stranded animals, as a quick and robust way to provide species confirmation.

Using hematological as well as stable-isotope data, Barros *et al.* (1998) speculated that dwarf sperm whales may have a more pelagic distribution than pygmy sperm whales, and/or dive deeper during feeding bouts. This behavior may result in differential exposure to marine debris, collision with vessels and other anthropogenic activities between the two *Kogia* species.

The western North Atlantic dwarf sperm whale population is being considered a separate stock for management purposes, although there is currently no information to differentiate this stock from the northern Gulf of Mexico stock(s). Additional morphological, genetic and/or behavioral data are needed to provide further information on stock delineation.

POPULATION SIZE

Abundance estimates from selected regions of the dwarf sperm whale habitat exist for select time periods. Because *K. sima* and *K. breviceps* are difficult to differentiate at sea, the reported abundance estimates are for both species of *Kogia*. The best estimate for *Kogia* spp. is 3,785 (CV=0.47; Table 1). This estimate is from summer 2011 surveys covering waters from central Florida to the lower Bay of Fundy. This estimate is almost certainly negatively biased. One component of line transect estimates is $g(0)$, the probability of seeing an animal on the transect line. Estimating $g(0)$ is difficult because it consists of accounting for both perception bias (i.e., at the surface but missed) and availability bias (i.e., below the surface while in range of the observers), and many uncertainties (e.g., group size and diving behavior) can confound both




Figure 1. Distribution of *Kogia* spp. sightings from NEFSC and SEFSC shipboard and aerial surveys during the summers in 2004 and 2011. Isobaths are the 100-m, 1,000-m and 4,000-m depth contours.

(Marsh and Sinclair 1989; Barlow 1999). The best estimate was corrected for perception bias (see below) but not availability bias and a corrected estimate could be 2-4 times larger.

Earlier abundance estimates

Please see Appendix IV for a summary of abundance estimates, including earlier estimates and survey descriptions.

Recent surveys and abundance estimates

An abundance estimate of 1,783 (CV=0.62) *Kogia* spp. was generated from aerial and shipboard surveys conducted during June-August 2011 between central Virginia and the lower Bay of Fundy. The aerial portion covered 6,850 km of tracklines over waters north of New Jersey between the coastline and the 100-m depth contour through the U.S. and Canadian Gulf of Maine, and up to and including the lower Bay of Fundy. The shipboard portion covered 3,811 km of tracklines between central Virginia and Massachusetts in waters deeper than the 100-m depth contour out to beyond the U.S. EEZ. Both sighting platforms used a double-platform data collection procedure, which allows estimation of abundance corrected for perception bias of the detected species (Laake and Borchers 2004). Estimation of the abundance was based on the independent observer approach assuming point independence (Laake and Borchers 2004) and calculated using the mark-recapture distance sampling option in the computer program Distance (version 6.0, release 2, Thomas *et al.* 2009).

An abundance estimate of 2,002 (CV=0.69) *Kogia* spp. was generated from a shipboard survey conducted concurrently (June-August 2011) in waters between central Virginia and central Florida. This shipboard survey included shelf-break and inner continental slope waters deeper than the 50-m depth contour within the U.S. EEZ. The survey employed two independent visual teams searching with 25x bigeye binoculars. A total of 4,445 km of tracklines were surveyed, yielding 290 cetacean sightings. The majority of sightings occurred along the continental shelf break with generally lower sighting rates over the continental slope. Estimation of the abundance was based on the independent observer approach assuming point independence (Laake and Borchers 2004) and calculated using the mark-recapture distance sampling option in the computer program Distance (version 6.0, release 2, Thomas *et al.* 2009).

Table 1. Summary of abundance estimates for the western North Atlantic <i>Kogia</i> spp. Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV).			
Month/Year	Area	N_{best}	CV
Jun-Aug 2011	central Virginia to lower Bay of Fundy	1,783	0.62
Jun-Aug 2011	central Florida to central Virginia	2,002	0.69
Jun-Aug 2011	central Florida to lower Bay of Fundy (COMBINED)	3,785	0.47

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for *Kogia* spp. is 3,785 (CV=0.47). The minimum population estimate for *Kogia* spp. is 2,598 animals.

Current Population Trend

A trend analysis has not been conducted for this stock. The statistical power to detect a trend in abundance for this stock is poor due to the relatively imprecise abundance estimates and long survey interval. For example, the power to detect a precipitous decline in abundance (i.e., 50% decrease in 15 years) with estimates of low precision (e.g., CV > 0.30) remains below 80% (alpha = 0.30) unless surveys are conducted on an annual basis (Taylor *et al.* 2007).

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size for *Kogia* spp. is 2,598. The maximum productivity rate is 0.04, the default value for cetaceans. The recovery factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5 because this stock is of unknown status. PBR for western North Atlantic *Kogia* spp. is 26.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

The estimated annual average fishery-related mortality or serious injury for *Kogia* sp. during 2007-2011 was 3.4 (CV=1.0; Table 2).

New Serious Injury Guidelines

NMFS updated its serious injury designation and reporting process, which uses guidance from previous serious injury workshops, expert opinion, and analysis of historic injury cases to develop new criteria for distinguishing serious from non-serious injury (Angliss and DeMaster 1998; Andersen *et al.* 2008; NOAA 2012). NMFS defines serious injury as an “injury that is more likely than not to result in mortality”. Injury determinations for stock assessments revised in 2013 or later incorporate the new serious injury guidelines, based on the most recent 5-year period for which data are available.

Fishery Information

The commercial fishery that could potentially interact with this stock in the Atlantic Ocean is the Atlantic Ocean, Caribbean, Gulf of Mexico large pelagic longline fishery (Appendix III). Pelagic swordfish, tunas and billfish are the targets of the longline fishery. Total estimated annual average fishery-related mortality and serious injury during 2007-2011 was unknown for dwarf sperm whales because species-specific mortality estimates could not be made. However, there was 1 report of a *Kogia* sp. seriously injured by the pelagic longline fishery during quarter 4 of 2011. Estimated serious injuries of *Kogia* sp. attributable to the pelagic longline fishery in the mid-Atlantic Bight region during quarter 4 of 2011 were 17.0 (CV=1.0; Garrison and Stokes 2012). The annual average serious injury and mortality attributable to the Atlantic pelagic longline fishery for the 5-year period from 2007 to 2011 was 3.4 animals (CV=1.0; Table 2).

Table 2. Summary of the incidental mortality and serious injury of Atlantic Ocean *Kogia* sp. by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the observed mortalities and serious injuries recorded by on-board observers, the estimated annual mortality and serious injury, the combined annual estimates of mortality and serious injury (Estimated Combined Mortality), the estimated CV of the combined estimates (Estimated CVs) and the mean of the combined estimates (CV in parentheses).

Fishery	Years	Vessels ^a	Data Type ^b	Observer Coverage	Observed Serious Injury	Observed Mortality	Estimated Serious Injury	Estimated Mortality	Estimated Combined Mortality	Est. CVs	Mean Annual Mortality
Pelagic Longline	07-11	74,78, 75,79, 83	Obs. Data Logbook	.07, .07, .10, .08, .09	0,0,0,0,1	0,0,0,0,0	0,0,0,0,17	0,0,0,0,0	0,0,0,0,17	NA, NA, NA, NA, 1.00	3.4 (1.0)
TOTAL											3.4 (1.0)

^a Number of vessels in the fishery is based on vessels reporting effort to the pelagic longline logbook.

^b Observer data (Obs. Data) are used to measure bycatch rates, and the data are collected within the Northeast Fisheries Observer Program. Mandatory logbook data were used to measure total effort for the longline fishery. These data are collected at the Southeast Fisheries Science Center (SEFSC).

Earlier Interactions

Between 1992 and 2006, 1 *Kogia* sp. was hooked, released alive and considered seriously injured in 2000 (in the Florida East coast fishing area) (Yeung 2001).

Other Mortality

From 2007-2011, at least 35 dwarf sperm whales were reported stranded along the U.S. Atlantic coast and Puerto Rico (Table 3; NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 13 September 2012 (SER) and 9 November 2012 (NER)). In addition, there were 6 records of unidentified stranded *Kogia*.

Table 3. Dwarf and pygmy sperm whale (*Kogia sima* (Ks), *Kogia breviceps* (Kb) and *Kogia* sp. (Sp)) strandings along the Atlantic coast, 2007-2011. Strandings that were not reported to species have been reported as *Kogia* sp. The level of technical expertise among stranding network personnel varies, and given the potential difficulty in correctly identifying stranded *Kogia* whales to species, reports to specific species should be viewed with caution.

STATE	2007			2008			2009			2010			2011			TOTALS		
	Ks	Kb	Sp	Ks	Kb	Sp	Ks	Kb	Sp	Ks	Kb	Sp	Ks	Kb	Sp	Ks	Kb	Sp
Maine	0	2	0	0	0	0	0	1	0	0	0	0	0	0	0	0	3	0
Massachusetts	0	1	1	0	2	0	1	4	0	1	0	0	0	0	0	2	7	1
New York	0	1	0	0	0	0	0	1	0	0	2	0	0	1	0	0	5	0
New Jersey	0	1	0	0	1	0	0	2	0	0	3	0	1	1	0	1	8	0
Delaware	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0
Maryland	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0
Virginia	2	4	0	2	0	0	0	0	0	0	0	2	1	1	0	5	5	2
North Carolina	5	7	0	1	4	1	1	6	0	3	5	0	2	10	0	12	32	1
South Carolina	1	3	0	0	5	0	1	6	0	1	6	0	1	2	0	4	22	0
Georgia	0	1	0	0	3	0	0	2	0	0	2	1	0	4	0	0	12	1
Florida	1	5	0	3	6	0	1	6	0	3	17	0	2	14	1	10	48	1
Puerto Rico	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	1	0
TOTALS	9	27	1	6	21	1	4	28	0	8	36	3	8	34	1	35	146	6

There were three documented strandings of dwarf sperm whales along the U.S. Atlantic coast during 2007-2011 that were classified as human interactions. One was a 2007 stranding in Virginia that was classified as a fishery interaction. The second was a whale stranded in Florida during 2010 whose flukes were cut off by a public person on the beach. Finally, plastic was found in the stomach of an animal that stranded in New Jersey during 2011.

Historical stranding records (1883-1988) of dwarf sperm whales in the southeastern U.S. (Credle 1988), and strandings recorded during 1988-1997 (Barros *et al.* 1998) indicate that this species accounts for about 17% of all *Kogia* strandings in the entire southeastern U.S. waters. During the period 1990-October 1998, 3 dwarf sperm whale strandings occurred in the northeastern U.S. (Maryland, Massachusetts, and Rhode Island), whereas 43 strandings were documented along the U.S. Atlantic coast between North Carolina and the Florida Keys in the same period.

Stranding data probably underestimate the extent of human-related mortality and serious injury because all of the marine mammals that die or are seriously injured may not wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other human interactions. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of human interactions.

Rehabilitation challenges for *Kogia* sp. are numerous due to limited knowledge regarding even the basic biology of these species. Advances in recent rehabilitation success have potential implications for future release and tracking of animals at sea to potentially provide information on distribution, movements and habitat use of these species (Manire *et al.* 2004).

STATUS OF STOCK

Dwarf sperm whales are not listed as threatened or endangered under the Endangered Species Act, and the western North Atlantic stock is not considered strategic under the Marine Mammal Protection Act. Total U.S. fishery-related mortality and serious injury for *Kogia* sp. is not less than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. The status of dwarf sperm whales in the western U.S. Atlantic EEZ relative to OSP is unknown. There are insufficient data to determine population trends for this species.

REFERENCES CITED

- Andersen, M.S., K.A. Forney, T.V.N. Cole, T. Eagle, R. Angliss, K. Long, L. Barre, L. Van Atta, D. Borggaard, T. Rowles, B. Norberg, J. Whaley and L. Engleby. 2008. Differentiating serious and non-serious injury of marine mammals: report of the serious injury technical workshop, 10-13 September 2007, Seattle, WA. NOAA Tech. Memo. NMFS-OPR-39. 94 pp.
- Angliss, R.P. and D.P. DeMaster. 1998. Differentiating serious and non-serious injury of marine mammals taken incidental to commercial fishing operations: Report of the serious injury workshop, 1-2 April 1997, Silver Spring, MD. NOAA Tech. Memo. NMFS-OPR-13. 48 pp.
- Barlow, J. 1999. Trackline detection probability for long-diving whales. Pages 209-221 *in*: Garner *et al.* (eds.) Marine mammal survey and assessment methods. Balkema, Rotterdam. 287 pp.
- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. Marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6, 73pp. Available from NMFS, Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.
- Barros, N.B. and D.A. Duffield. 2003. Unraveling the mysteries of Pygmy and Dwarf sperm whales. Strandings Newsletter of the Southeast U.S. Marine Mammal Stranding Network. December 2003. NOAA Tech. Memo NMFS-SEFSC-521, 11 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Drive, Miami, FL 33149.
- Barros, N.B., D.A. Duffield, P.H. Ostrom, D.K. Odell and V.R. Cornish. 1998. Nearshore vs. offshore ecotype differentiation of *Kogia breviceps* and *K. simus* based on hemoglobin, morphometric and dietary analyses. World Marine Mammal Science Conference Abstracts. Monaco. 20-24 January.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, J.L. Laake, D.L. Borchers and L. Thomas. 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press, New York, 432 pp.
- Caldwell, D.K. and M.C. Caldwell. 1989. Pygmy sperm whale *Kogia breviceps* (de Blainville 1838): dwarf sperm whale *Kogia simus* Owen, 1866. Pages 235-260 *in*: S.H. Ridgway and R. Harrison (eds.) Handbook of marine mammals, Vol. 4: River dolphins and the larger toothed whales. Academic Press, San Diego. 442 pp.
- Credle, V.R. 1988. Magnetite and magnetoreception in dwarf and pygmy sperm whales, *Kogia simus* and *Kogia breviceps*. MSc. Thesis. University of Miami. Coral Gables, FL.
- Duffield, D.A., N.B. Barros, E.O. Espinoza, S. Ploen, F.M.D. Gulland, and J.E. Heyning. 2003. Identifying pygmy and dwarf sperm whales (Genus *Kogia*) using electrospray ionization mass spectrometry of myoglobin and hemoglobin. Mar. Mamm. Sci. 19(2):395-399.
- Garrison, L.P. and L. Stokes. 2012. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2011. NOAA Tech. Memo. NMFS-SEFSC-632, 61 pp.
- Handley, C.O. 1966. A synopsis of the genus *Kogia* (pygmy sperm whales) *in* Norris, K.S. (ed) Whales, dolphins, and porpoises. U. of CA Press, xv + 789 pp.
- Laake, J.L. and D.L. Borchers. 2004. Methods for incomplete detection at distance zero, *In*: Advanced distance sampling, edited by S. T. Buckland, D. R. Andersen, K. P. Burnham, J. L. Laake, and L. Thomas, pp. 108–189, Oxford University Press, New York.
- Manire, C.A., H.L. Rhinehart, N.B. Barros, L. Byrd and P. Cunningham-Smith. 2004. An approach to the rehabilitation of *Kogia* sp. Aquatic Mamm. 30(2):257-270.
- Marsh, H. and D.F. Sinclair. 1989. Correcting for visibility bias in strip transect surveys of aquatic fauna. J. Wildl. Manage. 53:1017-1024.
- McAlpine, D.F. 2002. Pygmy and Dwarf Sperm whales. Pages 1007-1009 *in*: W.F. Perrin, B. Wursig, and J.G.M. Thewissen (eds.) Encyclopedia of Marine Mammals. Academic Press, San Diego, CA.
- Mullin, K.D. and G.L. Fulling. 2003. Abundance and distribution of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. Fish. Bull., U.S. 101:603-613.
- NOAA. 2012. Federal Register 77:3233. National policy for distinguishing serious from non-serious injuries of marine mammals. Available from: <http://www.nmfs.noaa.gov/op/pds/documents/02/238/02-238-01.pdf>
- Taylor, B.L., M. Martinez, T. Gerrodette, J. Barlow and Y.N. Hrovat. 2007. Lessons from monitoring trends in abundance in marine mammals. Mar. Mamm. Sci. 23(1): 157-175.
- Thomas, L., J.L. Laake, E. Rexstad, S. Strindberg, F.F.C. Marques, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, M.L. Burt, S.L. Hedley, J.H. Pollard, J.R.B. Bishop and T.A. Marques. 2009. Distance 6.0.

- Release 2. [Internet]. University of St. Andrews (UK): Research Unit for Wildlife Population Assessment. Available from: <http://www.ruwpa.st-and.ac.uk/distance/>
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Würsig, B., T.A. Jefferson and D.J. Schmidly. 2000. The marine mammals of the Gulf of Mexico. Texas A&M University Press, College Station, TX, 256 pp.
- Willis, P.M., and R.W. Baird. 1998. Status of the dwarf sperm whale (*Kogia simus*) in Canada. Can. Fld-Nat. 112:114-125.
- Wursig, B., T.A. Jefferson and D.J. Schmidly. 2000. The marine mammals of the Gulf of Mexico. Texas A&M University Press, College Station, TX, 256 pp.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo. NOAA-TM-SEFSC-467, 42 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.