

Findings for the OMPS  
Nadir Mapper 1<sup>st</sup> Guess Total Column Ozone (INCTO)  
&  
Nadir Mapper Total Column Ozone EDR (OOTCO)  
in  
Support of Promotion to Provisional Maturity

Compiled by L. Flynn from  
JPSS and NPP OMPS Teams  
Last Updated January 16, 2013

# Outline

- Provisional Definition
- Performance Requirements
- OMPS Background
- SOMTC Caveats
- INCTO/OOTCO Performance
  - Internal Evaluation
    - Flags – Error Code, QF1, QF2, Scene Condition
    - INCTO/OOTCO comparisons
    - Cross-Track Dependence
    - Geolocation of Small FOV
  - External Accuracy Evaluation
    - Comparisons to OMPS V8TOZ
    - Comparisons to SBUV/2, OMI and GOME-2 V8TOZ
    - Comparisons to ground-based Dobson and Brewer
  - Known Deficiencies (repeated)
- Summary of Findings and Recommendations (repeated)
  - Promotion to Provisional
  - Monitoring plots
  - [http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ\\_OOTCO.php](http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ_OOTCO.php)
  - [http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ\\_INCTO.php](http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ_INCTO.php)
  - Upgrade to V8TOZ

Provisional Definition	Artifacts (Deliverables)
Product quality may not be optimal	Product accuracy is determined for a broader (but still limited) set of conditions. No requirement to demonstrate compliance with specifications.
Incremental product improvements are still occurring	Narrative, listing and discussing known errors. All DRs are identified and prioritized (1-5). Provisional readiness will address priorities 1-2. Pathway towards algorithm improvements to meet specifications is demonstrated.
Version control is in affect	Description of the development environment, algorithm version (IDPS build number), and LUTs/PCTs versions used to generate the product validation materials. ATBDs are accurate, up-to-date and consistent with the product running.
General research community is encouraged to participate in the QA and validation of the product, but need to be aware that product validation and QA are ongoing	ADP STAR will request feedback from appropriate users for the product. The notification letter will include a Provisional Maturity disclaimer. DPA will send request to Project Science to post Provisional Maturity disclaimer on CLASS. DPA will submit readme document to CLASS.
Users are urged to consult the EDR product status document prior to use of the data in publications	Warning of potential non-reproducibility of results due to continuing calibration and code changes. Identify known deficiencies regarding product quality.
May be replaced in the archive when the validated product becomes available	Technical evaluation of limited data reprocessing is presented.
Ready for operational evaluation	Key NOAA and non-NOAA end users are identified and feedback requested.

**Table 4.2.3 - Ozone Total Column (OMPS-N)**

EDR Attribute	Threshold	Objective
<b>Ozone TC Applicable Conditions:</b> 1. Threshold requirements only apply under daytime conditions with Solar Zenith Angles (SZA) up to 80 degrees. 2. The EDR shall be delivered for all SZA.		
a. Horizontal Cell Size	50 x 50 km <sup>2</sup> @ nadir	10 x 10 km <sup>2</sup>
b. Vertical Cell Size	0 -60 km	0 -60 km
c. Mapping Uncertainty, 1 Sigma	5 km at Nadir	5 km
d. Measurement Range	50 -650 milli-atm-cm	50-650 milli-atm-cm
e. Measurement Precision		
1. $X < 0.25$ atm-cm	6.0 milli-atm-cm	1.0 milli-atm-cm
2. $0.25 < X < 0.45$ atm-cm	7.7 milli-atm-cm	1.0 milli-atm-cm
3. $X > 0.45$ atm-cm	2.8 milli-atm-cm + 1.1%	1.0 milli-atm-cm
f. Measurement Accuracy		
1. $X < 0.25$ atm-cm	9.5 milli-atm-cm	5.0 milli-atm-cm
2. $0.25 < X < 0.45$ atm-cm	13.0 milli-atm-cm	5.0 milli-atm-cm
3. $X > 0.45$ atm-cm	16.0 milli-atm-cm	5.0 milli-atm-cm
g. Refresh	At least 90% coverage of the globe every 24 hours (monthly average)	24 hrs.
		v2.0, 9/22/12

# INCTO/OOTCO Summary of Findings and Recommendations

- OMPS NM EDR (OOTCO) Status

The OOTCO product is producing reasonable values for total column ozone and effective reflectivity. Sulfur Dioxide (SO<sub>2</sub>) Index and Aerosol Index values show large variations with cross-track view angle and latitude due to inter-channel biases. Most error flags are functioning as designed. The OMPS NM EDR product (OOCTO) and IP product (INCTO) use the same algorithm and measurements. Differences are present because of the use of external data, e.g., CrIS Temperature Profiles and NRT VIIRS Snow/Ice fields are used in OOTCO in place of climatologies, monthly tilings and forecasts. We are replacing the VIIRS cloud top pressure initially with the UV climatology (CCR #736) and eventually with OMPS-measurement-based estimates. We have turned off the use of the VIIRS cloud fraction in the algorithm as previously applied for OOCTO. The inputs and corrections to the EDR and SDR algorithms and input data sets (e.g., VIIRS snow/ice fields) needed to further improve the product are known.

The OMPS Team recommends that the OOTCO Product be promoted to Provisional Maturity.

- OMPS NM First Guess IP (INCTO)

The INCTO product is producing reasonable values for total column ozone and effective reflectivity.

The OMPS Team recommends that the INCTO Product be promoted to Provisional Maturity.

- Monitoring Figures are available at

[http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ\\_INCTO.php](http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ_INCTO.php)

- Upgrade to V8TOZ

The team is investigating an upgrade from the OMPS EDR multiple triplet algorithm to the V8TOZ algorithm currently in use with SBUV/2, OMI and GOME-2 measurements for climate data records and operational products. As of January 1, 2013, the V8TOZ algorithm has been used to process the first year of OMPS data independently at NOAA STAR and the NASA Ozone PEATE. The OMPS NPP Science Team is adapting OMI algorithms to create better cloud top pressures, aerosol indices and SO<sub>2</sub> indices from the OMPS measurements for use with the V8 algorithm.

# INCTO/OOTCO Known Product Deficiencies

- Problems with fixes in the pipeline or changed over the course of the study
  - Preliminary Day 1 Solar (CCR #411 **Implemented the middle of June 11, 2012<sup>^</sup>**)
  - EDR RT LUT – corrected bandpasses (CCR #343)
  - Cloud Top Pressure – new UV climatology (CCR #385, **CCR #736 Mx7.0**)
    - Replace IR climatology in INCTO **and VIIRS cloud top pressure in OOTCO**
  - Partial Cloud Logic – consistent surface reflectivity (CCR #419)
    - Switched from VIIRS Cloud Fraction to measurement-based, partial-cloud estimates
  - SDR weekly dark current updates (DR #4750, etc. Started 12/22/2012)
  - Adjustment to Solar spectra for varying Earth/Sun distance (DR #4798, CCR #481 in Mx6.6 early 2013)
- Problems under investigation / in preparation
  - **VIIRS Snow/Ice Data** (DR #4678). New tilings under evaluation. Work on NRT product by VIIRS team.
  - Ozone values out of range (infrequent TOZ > 650 DU) (DR #4692)
  - Broken Ascending/Descending Flag (DR #4804)
  - Radiance/Irradiance Calibration/Adjustments (DR #5047)
  - Stray Light correction (DR #4907)
  - Incorrect SO2 Coefficients (DR #4918). This is a table change and can go in out-of-build.
- Longer term refinements and improvements
  - Soft internal calibration/adjustments, e.g., cross track bias (DR #5047)
  - Wavelength Scale and Adjustments
    - Day 1 Scale (**Preliminary update May 7, 2012<sup>^</sup>**) and trending
    - Radiance/Irradiance Doppler Shift adjustment
    - Intra-orbit scale drift

**<sup>^</sup> Both of these created large discontinuities in the product performance. Similar effects can be expected as further changes enter the system.**


# OMPS Fundamentals

**NOAA, through the Joint Polar Satellite System (JPSS) program, in partnership with National Aeronautical Space Administration (NASA), launched the Suomi National Polar-orbiting Partnership (Suomi NPP) satellite on October 28, 2011. The Ozone Mapping and Profiler Suite (OMPS) consists of two telescopes feeding three detectors measuring solar radiance scattered by the Earth's atmosphere and solar irradiance by using diffusers. The measurements are used to generate estimates of total column ozone and vertical ozone profiles.**

**The nadir mapper (total column) sensor uses a single grating monochromator and a CCD array detector to make measurements every 0.42 nm from 300 nm to 380 nm with 1.0-nm resolution. It has a 110° cross-track FOV and 0.27° along-track slit width FOV. The measurements are currently combined into 35 cross-track bins: 3.35° (50 km) at nadir, and 2.84° at ±55°. The resolution is 50 km along-track at nadir, with a 7.6-second reporting period. The instrument is capable of making measurements with much better horizontal resolution.**

**The nadir profiler sensor uses a double monochromator and a CCD array detector to make measurements every 0.42 nm from 250 nm to 310 nm with 1.0-nm resolution. It has a 16.6° cross-track FOV, 0.26° along-track slit width. The current reporting period is 38 seconds giving it a 250 km x 250 km cell size collocated with the five central total column cells.**

**The limb profiler sensor is a prism spectrometer with spectral coverage from 290 nm to 1000 nm. It has three slits separated by 4.25° with a 19-second reporting period that equates to 125 km along-track motion. The slits have 112 km (1.95°) vertical FOVs equating to 0 to 60 km coverage at the limb, plus offsets for pointing uncertainty, orbital variation, and Earth oblateness. The CCD array detector provides measurements every 1.1 km with 2.1-km vertical resolution. The products for the Limb Profiler are not discussed here.**



Instrument Fields of View. Schematic from Ball Aerospace and Technology Corporation.



# Nadir Mapper & Profiler







Diagram from Ball Aerospace and Technology Corporation



Each instrument can view the Earth or either of two solar diffusers; a working and a reference.

The instruments measure radiance scattered from the Earth's atmosphere and surface. They also make solar measurements using pairs of diffusers. Judicious operation of working and reference diffusers allows analysts to track the diffuser degradation. The solar measurements also provide checks on the wavelength scale and bandpass. The instruments have completed multiple passes through their internal dark and nonlinearity calibration sequences and are beginning to make regular solar measurements. (See information on the OMPS SDRs.)






## Ozone Absorption Cross Sections:

Ozone has four main absorption bands in the ultraviolet, visible and near-infrared as follows: the Hartley bands from 200 nm to 310 nm, the Huggins bands from 310 nm to 380 nm, the Chappuis bands from 400 nm to 650 nm, and the Wulf bands from 600 nm to 1100 nm. The OMPS nadir telescope directs photons to two spectrometers, one with a wide, cross-track field-of-view (FOV) and spectral coverage in the Huggins ozone absorption bands, and the other with a smaller, nadir FOV and spectral coverage in the Hartley ozone absorption bands. Figures (a) and (b) show the ozone absorption cross-sections at a nominal atmospheric temperature for parts of these bands. These cross-sections are for  $-50^{\circ}\text{C}$  as estimated from a quadratic fit in temperature of the Brion-Daumont-Malicet data set.

# OMPS Nadir Mapper Spectra

- The plot at the top of the following slide shows a sample OMPS Nadir Mapper solar spectrum measured in January. The initial calibration, goniometry and wavelengths scales have been applied. Notice the Fraunhofer lines, e.g., a deep one near 360 nm.
- The plot in the middle shows a sample spectrum for the Earth View data for the nadir field-of view.
- The plot on the bottom shows the ratio of the first two spectra. Notice that much of the structure in the solar spectrum cancels out in the ration. Also notice the variations between 320 and 330 nm produced by differential ozone absorption with wavelength as illustrated in the Figure (a) from two slides earlier.

# Typical spectra from 310 to 380 nm for OMPS Nadir Mapper



# Caveats for OMPS NM Earth-View SDR SOMTC

- The OMPS NM and NP dark will be updated weekly. This can result in small but systematic positive bias (higher than truth) in SDR. For NM, the bias is negligible for most wavelength. In rare cases where signals are extreme weak, such as near the terminator and for wavelength shorter than 305 nm, the bias can reach 0.2%. For NP, the bias generally decreases with wavelength, from as much as 0.02% for wavelengths shorter than 255 nm and down to 0.002% for wavelengths longer than 285 nm.
- The spectral solar irradiances have been updated with on-orbit measurements for both NM and NP. Further analysis and update of Day One solar irradiance may be provided in future. These preliminary and additional updates of Day One solar have been planned before launch as part of normal calibration update.
- The wavelength scales for the OMPS NM and NP for both Earth and solar spectra have been updated with on-orbit measurements. The adjustments were somehow larger than expected from pre-launch thermal analysis. We will monitor and re-evaluate periodically.
- Out-of-band stray light was expected before launch and confirmed after launch for the OMPS NM, especially for wavelength less than 305 nm. The impact can be severe, up to 5%, but limited to wavelength less than 310 nm that are less critical for ozone retrieval. Stray light correction will be a continuing effort in the coming months.

# SDR Requirements and Performance

Requirements	Specification or Prediction	On-Orbit Performance
Non-linearity Knowledge	< <b>0.5%</b>	~ <b>0.1%</b>
Non-linearity	< <b>2% full well</b>	< <b>0.46%</b>
On-orbit Wavelength Calibration	< <b>0.01 nm</b>	<b>average ~0.01 nm RMS</b>
Stray Light NM	≤ <b>2</b>	<b>average ~± 2%</b>
Intra-Orbit Wavelength Stability	< <b>0.02 nm</b>	< <b>0.013 nm</b>
SNR	> <b>1000</b>	> <b>1000 from SV and EV</b>
Inter-Orbital Thermal Wavelength Shift	< <b>0.02 nm</b>	< <b>0.013 nm</b>
CCD Read Noise	< <b>60 -e RMS</b>	< <b>25 -e RMS</b>
Detector Gain	> <b>46</b>	~ <b>42</b>
Absolute Irradiance Calibration Accuracy	< <b>7%</b>	<b>5%</b>
Absolute Radiance Calibration Accuracy	< <b>8%</b>	< <b>5%</b>

# SDR Table 3: Open DRs

DR #	Short Description
4978	TC EDR fails for Sample Table with extra pixel column
4951	Large error in TLE geo calculation
4927	FT document 474-00181 need update
4920	Indicate Graceful Degradation mode when using TLE geolocation
4914	OMPS TC and NP GEO
4907	TC needs Stray light correction
4879	NP and TC Darks need to be updated
4823	The NP stray light needs to be corrected.
4818	Smears in error - Smears show negative and unexpected values
4799	Inconsistent structure between OAD and EV SDR
4749	OMPS darks have negative values
4693	Cal SDR strategy study
4676	Radiance error associated with aggregation
4673	Correction for different linearity slope $T_{up}$ for CCD2
4672	Linearity correction update for xml file
4671	OMPS Data quality threshold tables non existence for SDR
4627	Quantization introduced by linearity correction error - Cal SDR
4615	Transient filter issue
4602	Spatial pixel mismatch in Cal SDR

# Total Column Ozone\* Products

The spectral measurements from the OMPS Nadir Mapper\* of the radiances scattered by the Earth's atmosphere are used to generate estimates of the total column ozone. The algorithm uses ratios of Earth radiance to Solar irradiance at triplets of wavelengths to obtain estimates of the total column ozone, effective reflectivity, and the wavelength dependence of the reflectivity. Table values computed for a set of standard profiles, cloud heights, latitudes and solar zenith angles are interpolated and compared to the measured top-of-atmosphere albedos. The triplets combine an ozone insensitive wavelength channel (at 364, 367, 372 or 377 nm) to obtain cloud fraction and reflectivity information, with a pair of measurements at shorter wavelengths. The pairs are selected to have one "weak" and one "strong" ozone absorption channel. The hyperspectral capabilities of the sensor are used to select multiple sets of triplets to balance ozone sensitivity across the range of expected ozone column amounts and solar zenith angles. The "strong" ozone channels are placed at 308.5, 310.5, 312.0, 312.5, 314.0, 315.0, 316.0, 317.0, 318.0, 320.0, 322.5, 325.0, 328.0, or 331.0 nm. They are paired with a longer "weak" channel at 321.0, 329.0, 332.0, or 336.0 nm. The ozone absorption cross-sections decrease from  $3.0 \text{ (atm. cm)}^{-1}$  to  $0.3 \text{ (atm. cm)}^{-1}$  over the range of "strong" wavelengths. Typical ozone columns range from 100 DU or 0.1 atm-cm to 600 DU or 0.6 atm-cm.

\*There is sometimes confusion on what to call the OMPS instruments and products. The **OMPS Nadir Mapper (NM)** makes the principal measurements that are used to create the **Total Column Ozone (TC or TOZ)** Products.



# The 1<sup>st</sup> Guess Total Ozone Product INCTO

The Multiple Triplet algorithm described in the previous slide is applied twice for each FOV. This was done to resolve the “Who goes first?” problem created by the desires to use information from other sensors in the retrieval algorithms, e.g., OMPS wanted to use the CrIS temperature profile, and CrIS wanted to use the OMPS ozone estimates. The “1<sup>st</sup> Guess” OMPS products (**INCTO**) use climatological or forecast fields for surface reflectivity and pressure, snow/ice coverage, cloud optical centroid depth, and atmospheric temperature. They use internally calculated estimates of cloud fractions and effective reflectivity from measurements at non-ozone absorbing UV wavelengths. As we will show, this application of the algorithm is performing well. This product is sometimes called the Total Ozone Intermediate Product (TOZ IP).

REFERENCES – Additional information is in the OMPS Total Column Algorithm Theoretical Basis and Operational Algorithm Description Documents, and a volume of the Common Data Format Control Book:

Available at <http://npp.gsfc.nasa.gov/documents.html>

OMPS Total Column Ozone ATBD [474-00029 Rev-Baseline.pdf](#)

OMPS Total Column Ozone OAD [474-00066 OAD-OMPS-TC-EDR-SW RevA 20120127.pdf](#)

Atmospheric EDRs CDFCB [474-0001-04-02 Rev-Baseline.pdf](#)

# The 2<sup>nd</sup> Pass Total Ozone Product, OOTCO

The “2<sup>nd</sup> Pass or EDR” OMPS products (**OOTCO**) use cloud top pressures (soon to be replaced with the UV climatology) and snow/ice coverage from VIIRS near-real-time products and temperature profiles from CrIS products. Recent studies have found that the estimates cloud top pressure from an infrared or visible sensor do not provide the correct quantities for use with UV radiances, e.g., thin cirrus may be optically opaque in the IR but optically thin in the UV. CCR #736 implements a switch to a UV-based cloud top climatology. The products use internally calculated estimates of cloud fractions and effective reflectivity from measurements at non-ozone absorbing UV wavelengths. As we will show, this application of the algorithm is performing well. This product is sometimes called the Total Ozone Environmental Data Record (TOZ EDR). The INCTO and OOTCO products use identical sets of measurements from the OMPS Nadir Mapper. The INCTO final ozone estimate is included as a parameter in the OOTCO output files.

REFERENCES – Additional information for this product is available in the documents listed for INCTO on the previous slide.

# INCTO/OOTCO Error and Quality Flags

- Error Flag
  - Bit1 0 good, 1 large residual; Bit2 1 large SO2 Index; Bit3 1 triplet inconsistency; Bit4 1 ozone out of range; Bit5 1 surface reflectivity out of range
- Quality Flag 1
  - Bit1/Bit2 Quality 0 no retrieval, 1 low, 2 medium, 3 high; Bit3 1 input data quality is not good; Bit4 1 triplet selection is not consistent; Bit5 1 inconsistent residuals; Bit6/Bit7 0 SZA<80, 1 80<SZA<88, 2 SZA>88
- Quality Flag 2 – Duplicates other flags or information
  - Bit1 1 snow/ice present; Bit2 1 sun glint geometry over open water; Bit3 1 solar eclipse in FOV; Bit4 1 TOZ<50 or TOZ>650; Bit5/6 0 TOZ> 450\*, 1 250<TOZ<450, 2 TOZ<250, 3 not used; Bit7 1 Aerosol Index Too Large, AI> 0.5; Bit8 Spare
- South Atlantic Anomaly Flag – Climatological intensity
  - 0 0-10%, 1 10-20%, 2 20-30%, 3 30-40%, 4 40-50%, 5 50-60%, 6 60-70%, 7 70-80%, 8 >80%
- Scene Condition Flag
  - Bit1 0 Descending, 1 Ascending^; Bit2 1 Snow/Ice present #; Bit3 1 Tropospheric Aerosols present; Bit4 1 Snow/Ice Fraction > 0; Bit5 1 Solar Zenith Angle (>80°); Bit6 1 Surface Reflectivity (>1.2 or <-0.05)

See references on the previous slide for more details on these flags.

\*QF2 B5/6 can be 0 when the condition is not checked in addition to when the total column ozone is greater than 450 DU.

^SCF B1 is not currently set properly according to the orbital path, i.e., by checking the changes in latitude during a measurement. It is almost always set to 1, except for the first and last granules in a sequence of measurements as processed at IDPS. A fix has been tested in ADL.


# SCF B2 is not currently set properly. It is set to snow/ice present almost everywhere. It is inconsistent with SCF B4 Snow/Ice Fraction > 0.

# Sun Glint Flags


OMPS/TC SunGlintflag at03/05/2012



VIIRS Blue Marble  
January 23, 2012



OMPS/DOFCD SunGlint at 01/02/2013




SOMTC\_npp\_d20120305\*  
GOTCD\_npp\_d20120305\*




generated by aitoff.pro

**This figure gives the location of pixels with the OMPS NM Sun Glint flag set to 1 – viewing geometry has the potential for sun glint and the location is over open water – for the OMPS Nadir Mapper for March 5, 2012. The code is consistent in assigning this condition. (The VIIRS image in the upper left corner shows similar locations for Sun Glint but shifted to the South as the image is for Sun angles in January.) This flag is passed through from the SDR to both the TOZ IP and EDR products.**

**INCTO Eclipse Flag & Aerosol Index:**  
These two figures compare the occurrences of Eclipse Flags and elevated Aerosol Index values. The figure below looks at the Eclipse Flag for the recent Solar Eclipse on May 20/21, 2012. The flag is activated for the proper times and locations. The reduced sunlight in eclipse conditions leads to poor values for the Aerosol Index as shown on the Left.




↑ Map showing locations of high Aerosol Index values for May 21, 2012. The large values in the Northwestern Pacific are present because the algorithm does not account for the low radiances in the Moon's shadow




↑ Location of Eclipse Flags for OMPS Nadir Mapper First Guess Total Column Ozone Product for May 21, 2012. The map for OOTCO is identical.

# Scene Condition Flag: Ascending/Descending



Nadir FOV locations for May 15, 2012. Each orbit starts near  $60^{\circ}\text{S}$  and ascends to  $80^{\circ}\text{N}$  and then descends to  $70^{\circ}\text{N}$ . The **Green** FOVs are where the Scene Condition Flag is set to Ascending, and the **Red** FOVs are where the flag is set to Descending. This flag erroneously identifies most descending orbit locations as ascending and erroneously identifies the first ascending location as descending. For all of the descending, it only correctly identifies the last location as descending. This may not be set at all and just have zeros from initialization. It is supposed to be passed through from SDR. DR #4804 is open on this issue.


# Scene Condition Flag: Ascending/Descending






Nadir FOV locations for January 2, 2013. Each orbit starts near  $60^{\circ}\text{S}$  and descends to  $80^{\circ}\text{S}$  and then ascends to  $70^{\circ}\text{N}$ . The **Green** FOVs are where the Scene Condition Flag is set to Ascending, and the **Red** FOVs are where the flag is set to Descending. This flag erroneously identifies most descending orbit locations as ascending and erroneously identifies the first ascending location as descending. For all of the descending, it only correctly identifies the last location as descending. This may not be set at all and just have zeros from initialization. It is supposed to be passed through from the SDR. DR #4804 is open on this issue.

# Time Series of Percent Good for OOTCO Error Flag

OOTCO Percent-Good in 01/2013







**SO2 Index** values for INCTO versus Latitude in Degrees North for two days – May 5th and May 10<sup>th</sup>, and for January 2, 2013. Notice that the distribution has shifted down so that there are fewer values above the Error Flag threshold of 6 DU. The algorithm has a cutoff in values at -12 DU. Much of the variation in this index is caused by deficiencies in the current calibration, not real atmospheric SO2 content. Similar changes in the product will occur as the day 1 solar spectra, wavelength scales and radiance calibration are adjusted during the coming months.

Another update (in the middle of the day on June 11, 2012), this time for the Day 1 solar spectra, reduced the percent good for the INCTO Error Flag from 99% down to approximately 50%. The new solar spectra had  $\pm 5\%$  variations relative to the prelaunch values. The decrease in good error flags was primarily due to increased SO<sub>2</sub> flagging. Until definitive inter-channel calibration values are determined and in the system, users can expect similar shifts in the product behavior.

## Daily Maps of Error Flags for OMPS

INCTO for June 11<sup>th</sup>:

**Purple** 1 Large Residual;


**Orange** 2 SO<sub>2</sub> Index;

**Green** 16 Surface Reflectivity out of range.


Since SO<sub>2</sub> is produced from residuals, both of the increased flag frequencies from May to June are related to residuals.

The first eight orbits (Eastern Hemisphere) have low occurrences of non-zero error flags.


OMPS/INCTO ErrorFlags at 06/11/2012



DMPS/INTCO ErrorFlags at 03/05/2012



DMPS/INTCO ErrorFlags at 05/18/2012



## Daily Maps of Error Flags for OMPS

INCTO for March 5<sup>th</sup>  
and May 18<sup>th</sup>, 2012:

**Purple** 1 Large  
Residual;


**Orange** 2 SO<sub>2</sub> Index;

**Green** 16 Surface  
Reflectivity out of  
range.


Since SO<sub>2</sub> is produce  
from residuals, both of  
the reduced flag  
frequencies from  
March to May are  
related to residuals.



OMPS/OOTCO ErrorFlags at 03/05/2012



OMPS/OOTCO ErrorFlags at 05/18/2012



## Daily Maps of Error Flags for OMPS

OOTCO for March 5<sup>th</sup> and May 18<sup>th</sup>, 2012:


**Purple** 1 Large Residual;

**Orange** 2 SO<sub>2</sub> Index;


**Green** 16 Surface Reflectivity out of range.

The OOTCO product in this time period is suffering from problems associated with the use of VIIRS cloud fraction estimates.

OMPS/INCTO ErrorFlags at 01/02/2013




OMPS/OOTCO ErrorFlags at 01/02/2013




Daily Maps of **Error Flags** for OMPS INCTO (Top) and OOTCO Bottom for January 2, 2013 : **Purple** 1 Large Residual; **Orange** 2 SO2 Index; **Green** 16 Surface Reflectivity out of range.

The OOTCO product is suffering from problems associated with the use of VIIRS cloud fraction estimates. The plan is to remove this dependence in future processing.

OMPS/INTCO ErrorFlags = 8,16 at 03/05/2012




OMPS/INTCO ErrorFlags = 8,16 at 05/18/2012




Daily Maps of **Error Flags** for OMPS INCTO for March 5<sup>th</sup> and May 18<sup>th</sup>, 2012: Since they were hard to distinguish in the previous plots, two of the less frequent flags are re-plotted here.  
**Red** 8 Ozone out of range (This flag is not activated for May 18<sup>th</sup>.);  
**Green** 16 Surface Reflectivity out of range.



OMPS/ODTCO ErrorFlags=8,16 at 03/05/2012




OMPS/ODTCO ErrorFlags=8,16 at 05/18/2012




Daily Maps of **Error Flags** for OMPS OOTCO for March 5<sup>th</sup> and May 18<sup>th</sup>, 2012: Since they were hard to distinguish in the previous plots, two of the less frequent flags are plotted here. **Red** 8 Ozone out of range, and **Green** 16 Surface Reflectivity out of range are replotted here.

OMPS/INCTO ErrorFlags=8,16 at 01/02/2013




OMPS/ODTDO ErrorFlags=8,16 at 01/02/2013




Daily Maps of **Error Flags** for OMPS INCTO (Top) and OOTCO (Bottom) for January 2, 2013: Since they were hard to distinguish in the previous plots, two of the less frequent flags are plotted here. **Red** 8 Ozone out of range, and **Green** 16 Surface Reflectivity out of range are replotted here.



OMPS/INTCO QF1 at 03/05/2012



OMPS/INTCO QF1 at 05/18/2012



## INCTO Quality

Flag 1 for March 5<sup>th</sup>  
(top) and May 18<sup>th</sup>  
(bottom), 2012:

**Orange** is QF1=4  
(input data quality is  
not good),


**Blue** is QF1=16  
(Residuals are not  
consistent),

**Green** is QF1=32 (SO<sub>2</sub>  
index  $\geq 6$  DU).


There are no QF1=8  
(O<sub>3</sub> triplet selection  
is not consistent)  
values.

Again the change in  
frequency is for  
residual related flags.

OMP5/OOTCO QF1 at 03/05/2012




OMP5/OOTCO QF1 at 05/18/2012




**OOTCO Quality Flag 1** for March 5<sup>th</sup> (top) and May 18<sup>th</sup> (bottom), 2012:  
**Orange** is QF1=4 (input data quality is not good),  
**Blue** is QF1=16 (Residuals are not consistent),  
**Green** is QF1=32 (SO<sub>2</sub> index  $\geq$  6 DU).  
The inconsistent effective reflectivity calculations are producing widespread errors.

OMPS/INCTO QF1 at 01/02/2013



OMPS/OOTCO QF1 at 01/02/2013



INCTO (Top) and  
OOTCO (Bottom)

Quality Flag 1 for  
January 2, 2013:

**Orange** is QF1=4 (input  
data quality is not  
good),

**Blue** is QF1=16  
(Residuals are not  
consistent),


**Green** is QF1=32 (SO<sub>2</sub>  
index  $\geq 6$  DU).

There are no QF1=8 (O<sub>3</sub>  
triplet selection is not  
consistent) values.


Again the change in  
frequency is for residual  
related flags. This day  
had an orbit that  
continued taking data  
on the night-side



OMPS/INTCO QF1\_TCO at 03/05/2012



OMPS/INTCO QF1\_TCO at 05/18/2012



## INCTO Quality Flag 1;


Total Column Quality:

**Orange** – 3 High Quality,  
**Blue** – 2 Medium Quality (large residue, input quality, triplet, SZA>80, SO<sub>2</sub>, SAA, or bad surface reflectivity), and  
**Purple** – 1 Poor Quality (Sun Glint, Eclipse, or Bad TOZ).


Poor quality is frequently due to Sun Glint, and Medium quality is frequently due to SAA flagging or Solar Zenith Angles greater than 80°. The white areas are regions where there is no sunlight for the FOV, so measurements can not be made. Two sets of observations are made at high latitudes for May 18<sup>th</sup> in the North so parts of earlier orbits are covered up by the results of later ones.



OMPS/OOTCO QF1\_TCO at 03/05/2012



OMPS/OOTCO QF1\_TCO at 05/18/2012



## OOTCO QualityFlag 1;

Total Column Quality:

**Orange** – 3 High Quality,

**Blue** – 2 Medium Quality


(large residue, input quality, triplet, SZA>80, SO<sub>2</sub>, SAA, or bad surface reflectivity), and

**Purple** – 1 Poor Quality (Sun Glint, Eclipse, High SO<sub>2</sub>, or Bad TOZ).


Poor quality is frequently due to Sun Glint or SO<sub>2</sub>, and Medium Quality flags are now present over much of the world. Two sets of observations are made at high latitudes for May 18<sup>th</sup> in the North so parts of earlier orbits are covered up by the results of later ones.



OMPS/INTCO QF1\_TCO at 01/02/2013




OMPS/OOTCO QF1\_TCO at 01/02/2013




January 2, 2013 INTCO (Top) and OOTCO (Bottom) Quality Flag 1; Total Column Quality: **Orange** – 3 High Quality, **Blue** – 2 Medium Quality (large residue, input quality, triplet, SZA>80, SO<sub>2</sub>, SAA, or bad surface reflectivity), and **Purple** – 1 Poor Quality (Sun Glint, Eclipse, Bad SO<sub>2</sub> or Aerosol Index, or Bad TOZ). Poor quality is frequently due to SO<sub>2</sub>, and Medium quality is frequently due to SAA flagging or Solar Zenith Angles greater than 80°. The white areas are regions where there is no sunlight for the FOV, so measurements can not be made. Two sets of observations are made at high latitudes for May 18<sup>th</sup> in the North so parts of earlier orbits are covered up by the results of later ones.

OMPS/INTCO Snowice at 03/05/2012




OMPS/INTCO Snowice at 05/18/2012




INCTO Quality  
Flag 2, Bit1,  
**Purple** Snow/Ice  
These are the values  
provided in the  
initial pre-launch  
data file. They are  
static; they are not  
being properly  
updated. The  
differences between  
the two figures are  
produced solely by  
changes in the  
OMPS coverage.

DMPS/DOTCD SnowIce at 03/05/2012




DMPS/DOTCD SnowIce at 05/18/2012




**OOTCO Quality**  
Flag 2, Bit1,  
**Purple** Snow/Ice.  
The snow/ice data is  
not correct. This  
problem is under  
investigation.



OMPS/INCTO SnowIce at 01/02/2013




DMPS/DOFCO SnowIce at 01/02/2013




**INCTO (Top) and OOTCO (Bottom) Quality Flag 2, Bit1, Purple Snow/Ice.** The snow/ice for INCTO is probably the November VIIRS tiling. **The snow/ice data is not correct in the OOTCO using the NRT VIIRS, e.g., in the Amazon. This problem is under investigation.**

OMPS/INTCO Aerosol at 03/05/2012




OMPS/INTCO Aerosol at 05/18/2012




**INCTO Quality Flag 2, Bit7, Red**  
 Aerosol Index limit exceeded. These are set consistent with the Aerosol Index values. The regions above the Sahara Desert and Arabian Peninsula are from elevated dust. The regions in the Equatorial Pacific are due to cross-track differences in the biases between longer wavelengths and effects of sun glint.

OMPS/ODTDO Aerosol at 03/05/2012




OMPS/ODTDO Aerosol at 05/18/2012




## OOTCO Quality Flag 2, Bit7, Red

Aerosol Index limit exceeded. These are set consistently with the Aerosol Index values. The Aerosol Index is sensitive to discrepancies in the effective reflectivity and to Sun glint. The blocked region in the lower left on the lower map is where the VIIRS cloud fraction was not available and the algorithm set the cloud fraction to zero.

OMPS/INCO Aerosol at 01/02/2013




OMPS/OOTCO Aerosol at 01/02/2013




January 2, 2013 INCTO (Top) and OOTCO (Bottom) Quality Flag 2, Bit 7, Red

Aerosol Index limit exceeded. These are set consistently with the Aerosol Index values. The Aerosol Index is sensitive to discrepancies in the effective reflectivity and to Sun glint. The blocked region in the lower left on the upper map is where the VIIRS cloud fraction was not available and the algorithm set the cloud fraction to zero.



November 23, 2012  
OOTCO Ozone (Top)  
and Input Data Quality  
Flag (Bottom) during  
VIIRS roll maneuver.  
Satellite View Angles exceed  
70 degrees,

DO QF1(2,input data quality is not good) at 11/23/2012



# Typical Distribution of Non-Fill Effective Reflectivity


The two figures on the next slide show the distribution of non-fill effective reflectivity values for the OOTCO (top) and INCTO (bottom) for April 6, 2012 with latitude. The two figures on the slide following it show the histograms of effective reflectivity for the two products for the same day.

The OOTCO estimates are a mixture of calculations produced by the surface reflectivity (adjusted for snow/ice fractional coverage from auxiliary data values times 95%), the VIIRS cloud fraction (times a minimum of 80%), and partial cloud calculations if no VIIRS cloud fraction is reported. Notice the plateaus of values at 80% (for cloud fractions of 1) and 95% (for snow/ice fractions of 1). The algorithm logic does not currently allow it to adjust the values downward. The IR cloud fractions are frequently much larger than those modeled for UV measurements, and the snow/ice tiles are not currently updated. These factors produce reflectivity errors that lead to large errors in the current OOTCO ozone products.


The INCTO estimates are derived directly from the OMPS UV measurements for selected channels between 330 and 380 nm. Fewer than 1% of the 120000 values computed in the INCTO product for this day are greater than 100 and only four values are less than 0. 2.5 % of the effective reflectivity have fill values with most of these occurring for  $SZA \geq 88^\circ$  and all are for cases with N-values containing fill – no other error flags are set by the algorithm for fill N-value cases. The algorithm does not currently adjust the surface reflectivity properly for low reflectivity cases but this logic is included in the next build scheduled for implementation in Fall of 2012.




OOTCO Reflectivity for 4/6/2012




INCTO Reflectivity for 4/6/2012



OOTCO Reflectivity for 4/6/2012



INCTO Reflectivity for 4/6/2012






# OOTCO/INCTO Differences on Next Slide

- Top Left – Cloud Top Pressure differences (UV climatology versus VIIRS IR estimates)
- Bottom Right – Snow/Ice differences (Monthly tiles versus near-real time both from VIIRS)
- Top Left – Cloud Fraction differences (Most are produced by Snow/Ice differences and how the pressure is handled for cloudy scenes over snow)
- Bottom Right – Total Ozone differences (These are a result of the different cloud heights and interacting with the assumed below cloud ozone column and reflectivity contributions to the ozone sensitive channel)

# Cloud Top Pressure Differences

ootco & into cloud top pressure difference on 11/12/2012




Atmosphere

0.00 0.08 0.17 0.25 0.33 0.42 0.50

# Cloud Fraction Differences

ootco & into cloud fraction difference on 11/12/2012




Fraction

0.00 0.03 0.05 0.08 0.10 0.13 0.15

# Total Ozone Differences

ootco & into ozone difference on 11/12/2012




Dobson Unit

0.00 2.50 5.00 7.50 10.00 12.50 15.00

# Snow/Ice Differences



ootco & into o3 diff at snow/ice different points on 11/12/2012




Dobson Unit

0.00 1.67 3.33 5.00 6.67 8.33 10.00


# Ozone differences between OOTCO and INCTO versus CTP difference                      Cloud Fraction difference



## INCTO Cloud Fraction



## CTP diff \* CF(INCTO)



# OMPS NM OOTCO Minimum

## Reflectivity Cross-Track Dependence

The lines in the figure on the next slide show the weekly, one-percentile effective reflectivity values for the months of November and December for all the data in a latitude/longitude box in the Equatorial Pacific versus cross-track view position. (17 is the nadir position and 0 and 34 are the extreme viewing angles.) The December values are given by the lower four lines and are offset -2% relative to the y-axis scale on the left. We expect the one-percentile effective reflectivity values to be approximately 4% for this region of the globe from climatological measurements made by other instruments. The larger values observed here are due to inaccuracies in the radiance calibration. Some of the changes between the two months are due to a lack of adjustment for the Earth/Sun distance in the solar spectrum. This particular, annually-varying bias will be rectified with changes in the code scheduled for implementation in early 2013.




# OMPS NM OOTCO/INCTO Aerosol Index Cross-Track Dependence

The lines in the figure on the next slide show the weekly, average aerosol index values for the months of November for all the data in a latitude/longitude box in the Equatorial Pacific versus cross-track view position. (17 is the nadir position and 0 and 34 are the extreme viewing angles.) The INCTO values are given by the upper four lines and are offset +1% relative to the y-axis scale on the left. The cross-track variation is probably related to calibration uncertainty for the radiances relative to the irradiances with some contributions from sun glint.




# OOTCO/INCTO Aerosol Index Cross-Track Dependence for November 2012



Weekly average Aerosol Index values, for November for a latitude / longitude box in the Equatorial Pacific versus cross track pixel..

# OOTCO Cross-Track Ozone Dependence



November weekly-mean total column ozone (OMPS Nadir Mapper EDR Multiple-Triplet algorithm retrievals) as a function of the cross-track view angle for a Latitude/Longitude in the Equatorial Pacific. The persistent cross-track bias is caused by deficiencies in the initial calibration/solar data values. The INCTO results are shifted down 20 DU relative to the scale on the left vertical axis.





# Total Ozone Maps for INCTO and OOTCO

The false color daily maps (October 15, 2012) of Total Column Ozone on the next two slides show the expected distribution of global ozone values for October including the Antarctic Ozone Hole. The OOTCO map (first) has greatly varying quality depending on the consistency of the VIIRS cloud fraction for the first half of the day followed by a transition to the use of a measurement-based calculation for the second half of the day (the Western Hemisphere).


# Sample OOTCO Total Ozone Map

OMPS OOTCO Total Ozone for 20121015



# Sample INCTO Total Ozone Map

OMPS INCTO Total Ozone for 20121015





# Total Ozone Maps

## OOTCO, INCTO, OMI, GOME-2 & V8TOZ

The false color daily maps (Nov 29, 2012) of Total Column Ozone on the next five slides show the expected distribution of global ozone values for late November. The OOTCO map (first) has varying quality depending on the consistency of the VIIRS cloud top pressure and NRT snow/ice imputed reflectivity values as compared with the UV measurements used for INCTO (second). Both maps show some “scalloping” due to the cross-track bias about each of the 14 orbital tracks. The distributions of ozone reported by the INCTO/ OOTCO are similar to that seen in the EOS Aura OMI map and MetOp-A GOME-2 total ozone products for the same day on the third and fourth slides, and the OMPS Version 8 retrievals on the fifth slide.


# Sample OOTCO Total Ozone Map

OMPS OOTCO Total Ozone for 20121129



# Sample INCTO Total Ozone Map


OMPS INCTO Total Ozone for 20121129





# Sample EOS Aura OMI Total Ozone Map

OMI Total Ozone for 20121129



100

200

300



400

500





# Sample MetOp-A GOME-2 Total Ozone Map

GOME-2 Total Ozone for 20121129



# Sample OMPS V8TOZ Total Ozone Map

OMPS V8 Total Ozone for 20121129





(29/29)

# Ozone: OMPS vs CrIMSS-EDR

OMPS nadir

mapper beta EDR

CrIMSS Mx6.4 O3-IP




Total column ozone (integral of CrIMSS ozone IP profile)

Quick Look slides and analysis by OMPS team


Slides courtesy of OMPS team (P.I. Larry Flynn, Jianguo Niu, Eric Beach)

# Time series of daily zonal mean ozone for Pacific Box for 2012

OMPS N18N19 SBUV2 GOME-2 Zonal Mean Total O3 1.2012-12.2012 20S20N/100W180W




# Time series of daily zonal mean ozone Pacific Box for October







# Effective Reflectivity Map for INCTO

The false color daily map (Nov 29, 2012) of Effective Reflectivity on the next slide shows the expected distribution of global values for November with high values over Greenland and low values over desert regions. Bands of clouds are also evident. For example, there is a feature in the western Atlantic running parallel to the US coastline. The OMI UV Cloud Fraction for the same day is given to the right. Clouds are found in consistent locations between the two maps.





# Sample INCTO Effective Reflectivity Map

OMPS INCTO Reflectivity for 20121129




# Sample OOTCO Effective Reflectivity Map

OMPS OOTCO Reflectivity for 20121129



# Aerosol Index Map for INCTO


The false color daily map (Nov 29, 2012) of Aerosol Index on the next slide shows the expected distribution of global aerosol with high values over Northern Africa and the Middle East due to desert dust. The North-South bands evident along the the eastern side of each orbit are caused by cross-track bias across the 14 orbital tracks. The OMI Aerosol Index for the same day is given to the right.






# Sample INCTO Aerosol Index Map

OMPS INCTO Aerosol Index for 20121129



# Sample OOCTO Aerosol Index Map

OMPS OOCTO Aerosol Index for 20121129






# Aerosol Index Map for OOTCO

The false color daily map (May 12, 2012) of Aerosol Index on the next slide shows a very variable product with reasonable values in regions with small cloud fractions reported by VIIRS (e.g., the high Aerosol Index values over Northern Africa and the Middle East, due to desert dust, where the VIIRS cloud fraction and UV cloud fraction are both close to zero) but poor performance over most of the globe where the computed effective reflectivity does not match the UV measurements. This changes starting on October 15, 2012.


# Sample OOTCO Aerosol Index Map

OMPS OOTCO Aerosol Index for 20120512





## ← 331-nm Channel Radiances

for the first eight orbits of OMPS Nadir Mapper measurement (end of 1/26/2012 and start of 1/27/2012). This image shows the expected range of values and variations across the orbital track and with solar zenith angles at the times of the measurements. The white circle around the North Pole is the region of polar night during the Northern Hemisphere Winter. The OMPS needs scattered sunlight to make its measurements, so there are no data there.



## Effective Reflectivity →

from the multiple triplet retrieval algorithm in IDPS for the same eight orbits. The quantity represents the UV reflectivity of the clouds and surface in each Field-of-View. Again, the range of values from bright clouds to dark open ocean scenes is as expected.




## ← Total Ozone

from the multiple triplet retrieval algorithm first guess product in IDPS for the same eight orbits for the first pass ozone retrieval (IP product without CrIS or VIIRS information). The values show some cross track variations and are offset approximately 5% from another satellite ozone product. These uncertainty levels for preliminary products are consistent with the use of prelaunch calibration parameters and tables in the initial operational system.

**First Light Products are Beta Quality**



# Geolocation Verification




The image on the left above shows the OMPS-calculated UV effective reflectivity over the Persian Gulf region for 30 January 2012. The image on the right is an Aqua MODIS RGB image for the same day. Comparison of the two indicate the capability of the OMPS TC sensor to identify surface features and characteristics of 5 km or less.

# Comparisons to MLO Dobson Station

- The next slide has a time series showing the matchup overpass data from the OOTCO for Mauna Loa compared to the Dobson station measurements. The OMPS values are a weighted average of all measurements within a  $1 \times 1.06$  degree<sup>2</sup> latitude x longitude box centered at the station.
- The slide following it shows a scatter plot comparing the Dobson values for Direct Sun measurement to the matchup data from OMPS (wavg) and from OMI


# March to December 2012

OOTCO is averaging ~3% above the Dobson Satellite sees to sea level, MLO misses ~7DU





# OMP and OMI vs MLO Dobson, daily TO3 DU, Direct Sun



# INCTO/OOTCO Known Product Deficiencies

- Problems with fixes in the pipeline or changed over the course of the study
  - Preliminary Day 1 Solar (CCR #411 **Implemented the middle of June 11, 2012<sup>^</sup>**)
  - EDR RT LUT – corrected bandpasses (CCR #343)
  - Cloud Top Pressure – new UV climatology (CCR #385, **CCR #736 Mx7.0**)
    - Replace IR climatology in INCTO **and VIIRS cloud top pressure in OOTCO**
  - Partial Cloud Logic – consistent surface reflectivity (CCR #419)
    - Switched from VIIRS Cloud Fraction to measurement-based, partial-cloud estimates
  - SDR weekly dark current updates (DR #4750, etc. Started 12/22/2012)
  - Adjustment to Solar spectra for varying Earth/Sun distance (DR #4798, CCR #481 in Mx6.6)
- Problems under investigation / in preparation
  - **VIIRS Snow/Ice Data** (DR #4678). New tilings under evaluation. Work on NRT product by VIIRS team.
  - Ozone values out of range (infrequent TOZ > 650 DU) (DR #4692)
  - Broken Ascending/Descending Flag (DR #4804)
  - Definitive Day One Solar & Radiance/Irradiance Calibration/Adjustments (DR #5047)
  - Stray Light correction (DR #4907)
  - Incorrect SO2 Coefficients (DR #4918). This is a table change and can go in out-of-build.
- Longer term refinements and improvements
  - Soft internal calibration/adjustments, e.g., cross track bias (DR #5047)
  - Wavelength Scale and Adjustments
    - Day 1 Scale (**Preliminary update May 7, 2012<sup>^</sup>**) and trending
    - Radiance/Irradiance Doppler Shift adjustment
    - Intra-orbit scale drift

**<sup>^</sup> Both of these created large discontinuities in the product performance. Similar effects can be expected as further changes enter the system.**

# Timeline of major changes in OMPS Nadir Products

- Problem with wavelength scale in NM February
  - Problem with wavelength scale in NP February (DOY error) Final data at CLASS – No.
  - May 7, 2012 – New OMPS NM and OMPS NP Wavelength Scales CCR #389
  - June 11, 2012 – New OMPS NM Day 1 Solar Irradiance CCR #411
  - July 17, 2012 – OMPS NP Day 1 Solar Irradiance CCR #458
  - August 10, 2012 – New NM RT LUT CCR #343 (Mx6.1)
  - August 10, 2012 – CTP to UV for INCTO CCR #385 (Mx6.2)
  - October 15, 2012 – OMPS NM E/S distance CCR #481 (Mx6.3)
  - October 15, 2012 – Partial cloud and VIIRS CF CCR #419 (Mx6.3)
  - TLE in use for Ephemeris Mx6.3 – Mx6.4
  - Updates to VIIRS Snow/Ice Tiles xxx
- Future
- February 2013 – IMOPO Surface Pressure limit too restrictive CCR #595 (Mx6.6)
  - April 2013 – CTP for OOTCO CCR #736 (Mx7.0)
  - April 2013 – Profile and TOZ flags switched in IMOPO PCR (Mx7.0)

# INCTO/OOTCO Summary of Findings and Recommendations

- OMPS NM First Guess IP (INCTO)

The INCTO product is producing reasonable values for total column ozone, effective reflectivity and aerosol index values. Most error flags are functioning as designed. Several problems with the product have been identified and there is progress on implementing corrections and adjustments.

The OMPS Team recommends that the INCTO Product be promoted to Beta Maturity.

- Monitoring Figures are available at

[http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ\\_INCTO.php](http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ_INCTO.php)

[http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ\\_OOTCO.php](http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.TOZ_OOTCO.php)

<http://www.star.nesdis.noaa.gov/icvs/PROD/proOMPSbeta.php>

## OMPS NM EDR (OOTCO) Status

The OMPS NM EDR product (OOTCO) uses the same algorithm and measurements as the INCTO product. Differences are present because of the use of external data, e.g., CrIS Temperature Profiles and VIIRS Cloud Top Pressures are used in OOTCO in place of forecasts or climatologies. We plan to replace the VIIRS cloud top pressure initially with the UV climatology and eventually with OMPS-measurement-based estimates. We have turned off the use of the VIIRS cloud fraction in the algorithm as applied for OOTCO as of October 15, 2012.

The OMPS Team recommends that the OOTCO Product be promoted to Beta Maturity. The inputs and corrections needed to improve the product further are known.

- Upgrade to V8TOZ

The team is investigating an upgrade from the OMPS EDR multiple triplet algorithm to the V8TOZ algorithm currently in use with SBUV/2, OMI and GOME-2 measurements for climate data records and operational products. As of July 1, 2012, the V8TOZ algorithm has been used to process the first five months of OMPS data independently at NOAA STAR and the NASA Ozone PEATE. The OMPS NPP Science Team is adapting OMI algorithms to create better cloud top pressures, aerosol indices and SO<sub>2</sub> indices from the OMPS measurements for use with the V8 algorithm.



Table 2.1.3 - Ozone Total Column		
EDR Attribute	Threshold (1,2)	Objective
a. Horizontal Cell Size	50 x 50 km <sup>2</sup> @ nadir (10)	10 x 10 km <sup>2</sup> (10)
b. Vertical Cell Size	0 - 60 km	0 - 60 km
c. Mapping Uncertainty, 1 Sigma (3)	5 km at Nadir (3)	5 km
d. Measurement Range	50 - 650 milli-atm-cm	50-650 milli-atm-cm
e. Measurement Precision (4)		
1. X < 0.25 atm-cm	6.0 milli-atm-cm (4,5)	1.0 milli-atm-cm
2. 0.25 < X < 0.45 atm-cm	7.7 milli-atm-cm (4,5)	1.0 milli-atm-cm
3. X > 0.45 atm-cm	2.8 milli-atm-cm + 1.1% (4,5)	1.0 milli-atm-cm
f. Measurement Accuracy (6)		
1. X < 0.25 atm-cm	9.5 milli-atm-cm (6,5)	5.0 milli-atm-cm
2. 0.25 < X < 0.45 atm-cm	13.0 milli-atm-cm (6,5)	5.0 milli-atm-cm
3. X > 0.45 atm-cm	16.0 milli-atm-cm (6,5)	5.0 milli-atm-cm
g. Latency	120 min. (7)	15 min
h. Refresh	At least 90% coverage of the globe every 24 hours (monthly average) (8)	24 hrs. (8)
i. Long-term Stability (9)	1% over 7 years	0.5% over 7 years
		v1.4.2, 7/29/11

**Notes:**

1. The OMPS Limb Profiler instrument does not fly on JPSS-1. Thus, only the Ozone Total Column elements are shown in this Table.
2. The loss of the OMPS Limb Profiler has had a small effect on the total column performance as the estimates of the profile shape and the tropospheric ozone are poorer, so the corrections are also poorer. There is new information that the OMPS algorithm use of the IR cloud top pressures will lead to errors as the IR values tend to be higher than the UV ones that should be used. A Discrepancy Report has been submitted and this error contribution is being carried as a risk with known mitigation in the form of improved climatology and measurement-based UV pressure estimates. The values in brackets are the performance requirements prior to OMPS Limb Profiler demanifestation.
3. The IORD-II Mapping "Accuracy" of 5 km at nadir was changed to "Uncertainty, 1 sigma" in accordance with user desires as expressed by the OATS and JARG.
4. The IORD-II required TC Measurement Precision of 3.0 milli-atm-cm + 0.5% of Measured Ozone. However, the TC Measurement Precision attributes are driven by the limitations of the nadir instrument when the contributions of the limb instrument are not available. The Government agrees with the contractor that the revised Threshold specifications represent realistic performance targets based on TOMS-EP heritage and EDR performance assessments.
5. The performance requirements with the contribution of an OMPS Limb Profiler would be as follows: Attribute e(1): 2.5 + 0.5% of measured ozone; Attribute e(2): 2.75 + 0.5%; Attribute e(3): 3.0 + 0.5%; Attribute f(1): 9.0; Attribute f(2): 12; Attribute f(3): 15.
6. The IORD-II TC Measurement Accuracy requirement is 15.0 milli-atm-cm. The TC Measurement Accuracy of 16 milli-atm-cm for Total Column > 450 milli-atm-cm is driven by the limitations of the nadir instrument when the contributions of the limb instrument are not available. The Government agrees with the contractor that the revised specifications represent realistic performance targets based on TOMS-EP heritage and EDR performance assessments.
7. Relaxed IORD-II Threshold requirement.
8. The IORD-II included Refresh threshold and objective requirements of 24 hrs which accommodate the stated requirement that the product be delivered under clear, daytime conditions only. This interpretation of the IORD-II Refresh requirement is consistent with the baseline OMPS Cross-track Swath Width design of ~ 2800 km (110° FOV) for a single orbit plane. All OMPS measurements require sunlight, so there is no coverage in polar night areas. This EDR is currently measured by the SBUV and High Resolution Infrared Sounder (HIRS) on POES, but the EDR produced by this legacy system does not satisfy JPSS threshold requirements.



**Table 2.1.3 - Ozone Total Column**

EDR Attribute	Threshold (1,2)	Objective
a. Horizontal Cell Size	50 x 50 km <sup>2</sup> @ nadir (10)	10 x 10 km <sup>2</sup> (10)
b. Vertical Cell Size	0 - 60 km	0 - 60 km
c. Mapping Uncertainty, 1 Sigma (3)	5 km at Nadir (3)	5 km
d. Measurement Range	50 - 650 milli-atm-cm	50-650 milli-atm-cm
e. Measurement Precision (4)		
1. X < 0.25 atm-cm	6.0 milli-atm-cm (4,5)	1.0 milli-atm-cm
2. 0.25 < X < 0.45 atm-cm	7.7 milli-atm-cm (4,5)	1.0 milli-atm-cm
3. X > 0.45 atm-cm	2.8 milli-atm-cm + 1.1% (4,5)	1.0 milli-atm-cm
f. Measurement Accuracy (6)		
1. X < 0.25 atm-cm	9.5 milli-atm-cm (6,5)	5.0 milli-atm-cm
2. 0.25 < X < 0.45 atm-cm	13.0 milli-atm-cm (6,5)	5.0 milli-atm-cm
3. X > 0.45 atm-cm	16.0 milli-atm-cm (6,5)	5.0 milli-atm-cm
g. Latency	120 min. (7)	15 min
h. Refresh	At least 90% coverage of the globe every 24 hours (monthly average) (8)	24 hrs. (8)
i. Long-term Stability (9)	1% over 7 years	0.5% over 7 years
		v1.4.2, 7/29/11

**Notes:**




- The OMPS Limb Profiler instrument does not fly on JPSS-1. Thus, only the Ozone Total Column elements are shown in this Table.
- The loss of the OMPS Limb Profiler has had a small effect on the total column performance as the estimates of the profile shape and the tropospheric ozone are poorer, so the corrections are also poorer. There is new information that the OMPS algorithm use of the IR cloud top pressures will lead to errors as the IR values tend to be higher than the UV ones that should be used. A Discrepancy Report has

# Northrop Grumman Algorithms and Data Products

**OMPS Team Task Status**


**Oct 10, 2012**

# Mx6.3 Preview Data Analysis: Comparison of INCTO and OOTCO




The difference in total column ozone retrieval between the IP and EDR (top left panel) appears to correlate with cloud fraction (top right panel). The cloud fraction difference between the IP and EDR (bottom right panel) does not show the same pattern, so this is not the primary cause of the differences. (M. Novicki and B. Sen, NGAS)

# Mx6.3 Preview Analysis: Cause of difference in cloud fraction



Most of the differences in the cloud fraction between the IP and EDR can be explained by differences in the snow/ice fraction used by the algorithm. The snow/ice fraction used by the IP comes from the Snow/Ice Rolling Tile, while the snow/ice fraction used by the EDR is derived from the VIIRS Snow and Ice EDRs. (M. Novicki and B. Sen, NGAS)

# Mx6.3 Preview Analysis: Comparison of difference in total column ozone



While the differences in total column retrievals between the IP and EDR appear to be correlated (see slide 1), the comparison shown above does not qualitatively show a correlation. Further statistical analysis needs to be done to determine if any correlation exists. This analysis is based on only two orbits of data and may show a different trend when examined over a whole day. (M. Novicki and B. Sen, NGAS)