

ESTIMATED WEIGHT (IN LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES AND FISHING AREA, 1981-1996 FOR THE ATLANTIC AND GULF COASTS.

YEAR	ATLANTIC COD						ATLANTIC CROAKER					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	1,019	36	4,719	35	11,986	24	1,636	18	1,270	17	176	34
1982.	701	97	3,320	48	17,514	24	1,414	15	3,194	22	192	30
1983.	159	35	2,177	22	12,247	24	1,756	15	1,353	16	183	27
1984.	162	50	1,573	24	4,204	13	3,211	17	1,322	13	290	23
1985.	639	60	1,738	46	20,701	63	1,887	11	1,174	14	103	21
1986.	194	56	1,313	31	6,020	40	6,386	15	1,515	17	164	29
1987.	54	39	2,576	41	6,358	20	2,613	11	1,192	20	323	26
1988.	330	30	1,394	26	10,381	17	3,579	9	1,749	37	960	34
1989.	1,269	24	600	35	5,241	12	2,113	7	602	12	111	18
1990.	280	32	1,182	17	6,449	18	1,962	8	515	41	10	31
1991.	516	37	728	25	8,063	19	3,862	21	560	56	15	29
1992.	25	37	360	30	1,951	18	2,681	7	431	19	85	29
1993.	106	39	467	21	6,566	15	3,177	7	328	11	105	32
1994.	304	30	553	22	3,428	30	4,595	6	756	15	121	27
1995.	87	47	196	24	4,318	17	3,800	7	339	12	288	41
1996.	105	80	220	25	3,007	22	3,629	8	525	11	507	18

YEAR	ATLANTIC MACKEREL						BLACK DRUM					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	620	44	4,082	22	2,375	27	677	23	449	24	156	20
1982.	742	33	67	28	1,816	28	1,159	25	1,687	22	30	42
1983.	165	23	132	37	6,321	29	2,081	15	1,799	25	916	40
1984.	109	25	895	19	4,109	23	754	22	1,186	24	70	51
1985.	1,044	76	1,932	51	3,006	47	866	18	1,377	25	180	48
1986.	612	45	625	43	8,073	29	3,079	17	888	24	41	32
1987.	395	39	3,712	23	4,782	27	3,238	40	874	22	539	53
1988.	1,313	63	2,163	26	3,722	24	1,666	14	815	19	206	27
1989.	387	23	2,141	15	1,410	24	1,687	14	578	36	108	31
1990.	650	35	1,431	18	2,035	20	422	18	529	33	72	38
1991.	979	31	1,533	23	3,145	19	1,180	18	636	31	35	32
1992.	172	20	215	23	243	22	1,541	10	749	14	62	21
1993.	375	18	271	19	681	20	1,258	11	516	26	133	36
1994.	646	19	2,114	22	999	35	1,084	11	663	33	126	32
1995.	474	19	1,014	20	1,266	27	1,407	11	575	14	139	31
1996.	564	17	1,393	18	1,166	18	1,395	9	386	12	44	22

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR. "OCEAN TO 3 MILES" INCLUDES TRIPS OFF OF FLORIDA'S WEST COAST FROM 0 TO 10 MILES FROM SHORE (STATE TERRITORIAL WATERS) WHILE "OCEAN 3 TO 200 MILES" INCLUDES TRIPS OFF FLORIDA'S WEST COAST BEYOND 10 MILES FROM SHORE. ESTIMATES DO NOT INCLUDE TEXAS DATA.

ESTIMATED WEIGHT (IN LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES AND FISHING AREA, 1981-1996 FOR THE ATLANTIC AND GULF COASTS.

YEAR	BLACK SEA BASS						BLUEFISH					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	552	21	448	21	1,228	21	19,875	14	27,832	20	48,160	10
1982.	512	22	8,465	40	3,186	20	14,104	24	21,692	12	47,628	16
1983.	339	26	434	19	4,315	18	23,565	10	24,680	21	44,466	21
1984.	269	16	755	23	2,738	14	17,918	20	18,983	11	31,431	11
1985.	508	11	845	21	3,893	18	21,375	9	11,520	16	20,285	23
1986.	1,963	16	5,538	37	6,536	23	25,631	12	31,360	8	36,752	13
1987.	501	21	1,465	17	1,807	10	25,008	7	26,211	9	25,985	10
1988.	580	14	880	14	4,228	15	16,791	9	17,063	9	15,539	8
1989.	819	8	1,751	12	3,630	9	13,670	7	13,033	7	14,281	9
1990.	469	10	726	16	2,710	9	15,043	7	8,780	7	7,090	8
1991.	751	10	955	11	3,769	12	16,889	7	9,593	7	7,484	8
1992.	404	9	1,007	12	2,332	8	10,273	6	8,699	8	5,809	9
1993.	562	18	2,188	41	3,143	18	9,979	6	5,816	6	4,846	10
1994.	377	12	453	13	3,251	12	8,128	7	5,172	8	2,692	14
1995.	608	15	476	20	5,956	14	7,268	8	3,989	10	3,517	11
1996.	481	16	560	18	5,759	15	5,365	11	3,919	9	5,804	17
YEAR	DOLPHIN						GAG					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	0	0	54	32	5,415	21	11	85	74	40	334	24
1982.	49	46	3,399	34	3,590	19	1	0	276	38	1,035	24
1983.	0	0	774	22	6,094	17	0	0	214	38	1,360	19
1984.	4	100	1,484	22	2,917	21	19	94	2,235	73	794	20
1985.	1	0	1,647	25	5,248	24	4	52	176	38	1,648	23
1986.	12	100	2,326	20	7,615	17	7	54	167	60	498	19
1987.	13	62	1,731	24	5,419	12	65	36	418	25	999	21
1988.	82	94	1,724	31	5,800	12	59	58	280	27	1,293	19
1989.	3	100	2,288	25	11,208	8	158	48	379	24	853	17
1990.	18	87	4,121	35	8,638	12	23	54	341	30	1,417	30
1991.	84	34	2,923	15	14,202	8	507	48	1,064	19	762	20
1992.	106	38	1,721	37	7,964	10	111	32	546	16	1,549	10
1993.	65	38	1,138	17	9,615	9	91	37	529	15	2,207	9
1994.	76	49	1,335	40	11,367	6	158	30	418	17	1,931	10
1995.	10	100	2,659	11	17,123	5	222	20	516	20	2,193	9
1996.	63	31	1,389	18	11,096	7	126	24	447	16	1,669	9

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR. "OCEAN TO 3 MILES" INCLUDES TRIPS OFF OF FLORIDA'S WEST COAST FROM 0 TO 10 MILES FROM SHORE (STATE TERRITORIAL WATERS) WHILE "OCEAN 3 TO 200 MILES" INCLUDES TRIPS OFF FLORIDA'S WEST COAST BEYOND 10 MILES FROM SHORE. ESTIMATES DO NOT INCLUDE TEXAS DATA.

ESTIMATED WEIGHT (IN LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES AND FISHING AREA, 1981-1996 FOR THE ATLANTIC AND GULF COASTS.

YEAR	GRAY SNAPPER						GRAY TRIGGERFISH					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	139	40	456	24	677	50	4	64	509	49	216	30
1982.	99	21	577	19	308	31	0	0	223	28	927	26
1983.	165	27	765	25	297	26	20	50	159	28	135	19
1984.	211	38	3,272	33	359	40	11	91	53	25	132	18
1985.	95	35	1,672	24	505	37	1	100	129	30	152	30
1986.	233	35	800	20	287	26	15	20	484	38	912	30
1987.	143	29	852	48	447	32	19	62	512	47	1,241	40
1988.	379	13	554	20	155	31	66	44	787	22	592	19
1989.	444	11	749	15	396	30	58	46	653	19	1,088	18
1990.	642	24	454	21	293	41	29	35	722	23	1,604	22
1991.	614	16	1,112	15	330	26	103	38	1,072	20	1,250	18
1992.	590	9	578	10	291	15	37	42	563	14	1,031	10
1993.	386	8	647	11	336	18	49	21	514	17	902	11
1994.	412	9	583	11	422	19	43	30	362	21	944	11
1995.	388	10	499	10	491	14	26	28	334	17	820	9
1996.	325	11	437	11	425	14	42	23	188	17	606	14
YEAR	GREATER AMBERJACK						KING MACKEREL					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	2	51	841	37	1,069	11	1,506	79	1,980	14	2,539	21
1982.	47	100	710	39	3,117	21	10	55	7,629	54	4,923	13
1983.	0	0	257	33	1,965	18	57	83	5,296	25	4,162	25
1984.	74	100	540	27	2,115	20	23	100	3,669	22	6,053	17
1985.	293	76	378	41	1,999	17	39	77	2,492	22	6,141	28
1986.	34	74	1,204	26	8,536	18	4	79	2,902	15	5,132	11
1987.	2	70	1,048	31	7,073	21	84	57	3,711	11	4,529	12
1988.	0	0	501	24	3,856	18	188	25	3,255	11	5,423	11
1989.	59	83	1,787	28	4,549	22	24	68	3,134	16	3,456	9
1990.	48	98	262	27	1,717	21	53	49	3,386	13	3,984	9
1991.	1	44	1,729	28	2,222	15	90	38	4,775	12	4,698	8
1992.	9	120	650	19	3,127	11	44	36	3,711	8	6,455	9
1993.	0	62	1,099	15	3,231	19	121	30	4,171	8	4,253	6
1994.	273	64	379	44	2,537	12	353	23	3,523	10	4,944	6
1995.	73	100	332	32	1,029	13	287	21	3,715	8	5,717	6
1996.	0	0	446	21	1,524	10	187	29	3,396	8	6,436	6

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR. "OCEAN TO 3 MILES" INCLUDES TRIPS OFF OF FLORIDA'S WEST COAST FROM 0 TO 10 MILES FROM SHORE (STATE TERRITORIAL WATERS) WHILE "OCEAN 3 TO 200 MILES" INCLUDES TRIPS OFF FLORIDA'S WEST COAST BEYOND 10 MILES FROM SHORE. ESTIMATES DO NOT INCLUDE TEXAS DATA.

ESTIMATED WEIGHT (IN LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES AND FISHING AREA, 1981-1996 FOR THE ATLANTIC AND GULF COASTS.

YEAR	RED DRUM						RED SNAPPER					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	1,702	16	1,800	20	387	55	1	100	807	31	3,573	29
1982.	2,866	17	4,163	25	297	44	0	0	1,015	26	1,341	24
1983.	3,745	17	3,008	19	1,255	23	0	0	188	25	4,139	20
1984.	3,258	14	5,517	25	720	20	0	0	167	24	1,713	20
1985.	3,821	14	3,506	25	311	28	39	77	190	35	3,564	19
1986.	3,718	15	3,065	13	371	20	3	60	641	31	1,895	20
1987.	3,602	12	1,571	11	1,033	27	17	54	370	36	1,394	31
1988.	3,646	9	1,002	16	194	33	46	40	255	24	1,823	22
1989.	4,460	9	1,969	15	226	22	34	49	202	47	1,847	21
1990.	2,911	14	1,964	16	535	23	4	101	281	31	890	16
1991.	4,445	10	2,141	13	60	37	25	79	343	30	1,267	14
1992.	6,190	5	2,434	6	348	15	32	49	531	26	2,668	11
1993.	6,987	6	2,210	9	727	15	60	71	344	15	3,873	7
1994.	6,589	5	1,919	8	463	23	38	31	396	17	3,500	9
1995.	10,109	5	2,860	6	697	21	5	83	329	17	2,609	9
1996.	9,099	5	3,063	8	880	19	3	100	254	23	2,462	9
YEAR	SAND SEATROUT						SCUP					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	846	29	3,130	53	97	32	3,970	36	1,220	26	624	32
1982.	193	26	785	26	271	28	1,737	17	1,883	21	1,585	25
1983.	87	44	2,138	25	261	31	2,158	18	1,475	22	2,619	28
1984.	162	61	2,630	31	240	26	1,540	23	600	25	276	31
1985.	642	30	3,861	25	592	34	5,699	14	271	22	126	21
1986.	985	15	1,853	12	208	21	4,879	13	5,208	15	1,522	33
1987.	431	20	1,323	14	260	17	1,456	14	2,967	23	1,774	26
1988.	978	15	489	16	94	25	2,442	13	1,369	18	456	36
1989.	938	27	504	17	13	30	3,547	9	1,744	18	266	26
1990.	497	15	866	18	472	39	2,472	11	1,371	11	297	21
1991.	1,715	18	853	15	141	36	5,776	7	1,834	12	479	19
1992.	1,017	15	490	12	136	27	3,149	8	1,176	17	87	30
1993.	781	10	819	16	490	29	1,942	7	498	17	758	34
1994.	1,663	8	877	12	344	19	1,721	17	564	15	343	22
1995.	1,077	10	769	14	155	22	464	22	468	16	382	24
1996.	832	12	768	12	289	23	1,528	10	508	17	203	31

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR. "OCEAN TO 3 MILES" INCLUDES TRIPS OFF OF FLORIDA'S WEST COAST FROM 0 TO 10 MILES FROM SHORE (STATE TERRITORIAL WATERS) WHILE "OCEAN 3 TO 200 MILES" INCLUDES TRIPS OFF FLORIDA'S WEST COAST BEYOND 10 MILES FROM SHORE. ESTIMATES DO NOT INCLUDE TEXAS DATA.

ESTIMATED WEIGHT (IN LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES AND FISHING AREA, 1981-1996 FOR THE ATLANTIC AND GULF COASTS.

YEAR	SHEEPSHEAD						SOUTHERN FLOUNDER					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	995	24	494	25	39	32	736	25	243	18	3	66
1982.	1,882	15	1,601	15	354	41	973	51	758	18	29	45
1983.	1,484	18	2,203	28	339	27	520	16	2,834	42	94	65
1984.	1,769	14	2,536	16	364	39	422	13	346	15	14	39
1985.	1,303	19	2,418	27	106	46	775	17	601	24	25	44
1986.	2,706	26	938	18	36	24	2,436	22	701	24	22	27
1987.	2,361	14	1,287	12	82	20	419	10	252	14	35	41
1988.	3,098	10	2,611	14	244	27	1,062	9	269	15	16	35
1989.	3,414	12	2,366	19	94	64	847	16	170	31	21	43
1990.	2,410	12	1,154	21	157	28	927	10	397	28	34	28
1991.	3,595	11	1,468	19	27	53	949	9	600	13	31	38
1992.	5,450	6	1,571	10	341	30	1,209	7	409	14	19	28
1993.	5,025	7	1,078	12	353	36	1,010	7	305	12	34	30
1994.	4,198	8	1,262	12	176	25	1,435	6	320	11	42	30
1995.	5,362	7	1,690	12	455	27	1,264	7	350	14	24	41
1996.	3,554	6	1,164	9	337	23	1,021	7	189	14	48	56
YEAR	SPANISH MACKEREL						SPOT					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	567	29	1,631	16	825	23	2,015	19	4,331	17	611	28
1982.	203	59	2,635	18	1,166	35	1,218	15	2,758	14	26	34
1983.	155	49	1,175	22	1,543	22	2,146	27	2,406	18	477	27
1984.	155	68	845	25	1,458	29	745	14	822	15	239	47
1985.	192	32	1,039	16	664	16	2,589	13	3,285	13	78	32
1986.	260	30	6,466	16	1,271	14	2,507	10	880	18	39	27
1987.	124	17	3,077	8	1,429	14	3,052	9	748	11	101	20
1988.	414	19	3,308	8	1,428	8	1,509	10	1,023	20	21	34
1989.	347	15	1,857	7	862	10	2,318	7	920	9	56	20
1990.	599	15	2,456	8	1,037	9	3,024	8	555	19	60	33
1991.	933	14	2,964	7	1,096	10	3,721	8	801	9	18	65
1992.	1,110	11	3,336	5	926	8	3,518	10	540	15	9	43
1993.	826	10	1,881	7	671	10	2,244	9	1,234	10	1	55
1994.	719	8	1,911	6	505	9	2,211	6	2,144	9	28	29
1995.	348	12	1,392	8	708	14	1,380	8	1,755	9	22	34
1996.	799	10	1,244	8	661	11	968	11	1,068	14	15	45

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR. "OCEAN TO 3 MILES" INCLUDES TRIPS OFF OF FLORIDA'S WEST COAST FROM 0 TO 10 MILES FROM SHORE (STATE TERRITORIAL WATERS) WHILE "OCEAN 3 TO 200 MILES" INCLUDES TRIPS OFF FLORIDA'S WEST COAST BEYOND 10 MILES FROM SHORE. ESTIMATES DO NOT INCLUDE TEXAS DATA.

ESTIMATED WEIGHT (IN LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES AND FISHING AREA, 1981-1996 FOR THE ATLANTIC AND GULF COASTS.

YEAR	SPOTTED SEATROUT						STRIPED BASS					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	2,730	18	2,558	22	549	25	1,177	37	532	39	245	43
1982.	4,391	12	7,559	13	1,038	19	297	29	1,773	62	453	32
1983.	3,610	16	5,576	15	3,318	24	360	26	2,056	72	282	50
1984.	2,947	39	5,223	20	1,600	44	410	41	874	53	0	0
1985.	3,328	10	4,822	16	385	28	514	44	335	43	0	0
1986.	10,210	29	7,260	7	1,618	13	220	56	810	32	98	70
1987.	6,256	7	6,536	7	1,467	13	178	30	529	21	156	56
1988.	7,868	5	5,286	9	1,782	15	417	22	710	26	195	35
1989.	7,644	8	5,868	15	1,057	17	365	26	313	25	64	40
1990.	3,786	8	2,274	10	304	22	1,200	17	864	17	232	49
1991.	8,899	7	5,820	7	576	16	1,764	13	1,559	14	383	25
1992.	7,706	5	2,937	7	678	17	2,185	13	1,641	18	241	23
1993.	7,268	5	2,450	7	345	20	2,718	9	2,188	19	760	18
1994.	7,395	4	3,520	6	263	16	4,160	10	2,296	10	357	20
1995.	8,554	5	3,821	5	194	24	6,001	8	4,403	13	861	12
1996.	7,067	6	3,303	7	175	24	6,004	7	6,959	8	1,656	16
YEAR	SUMMER FLOUNDER						TAUTOG					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	6,358	11	3,023	15	717	22	1,536	24	1,976	32	604	26
1982.	6,905	8	10,077	45	1,281	30	2,502	15	2,928	17	2,908	20
1983.	17,452	8	8,443	16	2,112	20	1,890	20	2,474	16	1,386	28
1984.	7,751	10	9,652	12	1,433	21	1,457	13	3,650	17	273	22
1985.	4,675	11	5,472	20	2,343	26	2,260	18	1,245	29	800	26
1986.	8,733	13	8,349	12	803	19	5,250	11	8,551	17	3,105	22
1987.	4,361	9	6,865	15	1,003	27	3,135	16	2,469	18	3,285	18
1988.	8,000	5	4,732	8	1,926	10	4,751	19	2,562	17	1,989	29
1989.	1,659	7	1,375	11	142	17	4,035	8	1,286	14	1,059	21
1990.	2,720	6	2,119	7	302	15	3,114	10	1,072	14	970	15
1991.	4,584	6	3,189	6	385	19	3,948	8	2,739	12	1,418	14
1992.	4,255	5	2,669	7	233	17	4,167	12	1,670	12	1,834	18
1993.	5,563	5	3,052	9	229	21	3,824	10	1,065	19	1,038	22
1994.	6,461	5	2,564	7	321	18	2,029	20	510	23	929	22
1995.	4,084	6	1,196	9	224	17	2,388	13	1,407	16	809	19
1996.	6,256	4	3,483	6	676	14	1,516	13	702	23	1,048	18

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR. "OCEAN TO 3 MILES" INCLUDES TRIPS OFF OF FLORIDA'S WEST COAST FROM 0 TO 10 MILES FROM SHORE (STATE TERRITORIAL WATERS) WHILE "OCEAN 3 TO 200 MILES" INCLUDES TRIPS OFF FLORIDA'S WEST COAST BEYOND 10 MILES FROM SHORE. ESTIMATES DO NOT INCLUDE TEXAS DATA.

ESTIMATED WEIGHT (IN LBS.) AND PSE (%) OF FISH HARVESTED BY MARINE RECREATIONAL ANGLERS, BY SPECIES AND FISHING AREA, 1981-1996 FOR THE ATLANTIC AND GULF COASTS.

YEAR	VERMILON SNAPPER						WEAKFISH					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	0	0	51	24	39	33	4,897	15	6,878	31	4,330	25
1982.	0	0	80	29	424	22	1,754	20	4,693	22	1,838	34
1983.	0	0	32	22	448	42	7,078	14	2,606	34	2,046	26
1984.	0	0	51	28	314	15	917	35	4,436	31	1,661	22
1985.	1	100	214	46	467	36	2,288	16	1,549	23	1,652	17
1986.	5	47	162	38	777	47	4,704	12	3,032	17	2,406	22
1987.	0	0	116	44	583	28	4,081	21	1,531	23	1,138	17
1988.	0	93	164	26	437	17	3,188	13	1,618	33	1,525	15
1989.	0	0	108	34	409	19	1,645	9	459	17	74	32
1990.	2	59	107	26	530	25	904	10	320	17	123	19
1991.	0	0	390	31	544	25	1,758	8	235	18	138	28
1992.	1	55	101	24	619	10	1,142	9	218	24	39	33
1993.	2	78	96	24	760	11	882	11	184	18	37	41
1994.	0	0	92	44	589	11	1,320	12	318	14	157	21
1995.	2	73	159	18	447	12	1,486	8	311	19	55	24
1996.	0	0	60	23	249	18	2,417	8	452	18	83	17
YEAR	WINTER FLOUNDER						ALL FISHES					
	INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES		INLAND		OCEAN 0 TO 3MILES		OCEAN 3 TO 200 MILES	
	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)	EST	PSE (%)
----- THOUSANDS OF POUNDS -----												
1981.	10,022	12	2,171	18	231	25	71,676	6	98,352	7	108,186	6
1982.	9,063	25	7,123	54	232	27	63,442	7	133,832	7	118,086	8
1983.	7,351	12	2,134	16	155	22	86,471	4	112,276	6	140,402	8
1984.	12,239	13	2,826	17	92	35	63,233	7	110,956	5	87,776	5
1985.	12,398	18	2,916	25	59	67	75,551	4	84,193	5	104,484	14
1986.	6,030	13	1,216	12	389	53	99,744	5	125,953	4	141,926	5
1987.	5,860	12	4,346	18	762	35	73,288	4	101,190	4	115,498	4
1988.	7,527	12	880	16	373	20	81,229	3	73,900	3	96,185	3
1989.	4,354	15	958	18	52	23	66,085	3	63,606	3	93,632	4
1990.	2,404	11	684	10	68	25	56,246	3	52,613	4	64,145	5
1991.	2,404	12	441	11	54	31	86,109	3	71,720	3	70,412	4
1992.	804	10	234	14	34	33	74,019	2	52,204	3	56,177	3
1993.	1,766	27	309	19	55	28	67,278	2	50,334	4	67,633	3
1994.	1,311	14	163	17	23	39	70,268	2	47,577	2	65,591	3
1995.	1,362	12	108	24	79	65	68,871	2	47,165	3	86,120	6
1996.	1,617	12	80	18	15	34	66,313	2	47,625	3	71,416	3

NOTE:--EST = ESTIMATE. PSE (%) = PROPORTIONAL STANDARD ERROR. "OCEAN TO 3 MILES" INCLUDES TRIPS OFF OF FLORIDA'S WEST COAST FROM 0 TO 10 MILES FROM SHORE (STATE TERRITORIAL WATERS) WHILE "OCEAN 3 TO 200 MILES" INCLUDES TRIPS OFF FLORIDA'S WEST COAST BEYOND 10 MILES FROM SHORE. ESTIMATES DO NOT INCLUDE TEXAS DATA.