



## METHYLPHENIDATE

(Trade Names: Ritalin- (IR, LA, and SR), Concerta, Metadate- (CD and ER),  
Methylin- (IR and ER) and Focalin- (IR and ER))

February 2011  
DEA/OD/ODE

### Introduction:

Methylphenidate (methyl-alpha-phenyl-2-piperidine-acetate hydrochloride) is a central nervous system (CNS) stimulant that has been marketed in the United States since the 1950s. For many years, Ritalin® (immediate release (IR) product), was the only brand-name product available. In recent years, other IR, extended release (ER), and long acting (LA) methylphenidate products have entered the market. These products are primarily prescribed to children for the treatment of attention deficit hyperactivity disorder (ADHD).

Domestic and worldwide use of methylphenidate has increased dramatically since 1990. According to the 2004 United Nations International Narcotic Control Board (INCB) report, the United States is the main consumer of methylphenidate accounting for about 70 percent of the global medical use of methylphenidate.

### Licit Use:

Methylphenidate is used almost exclusively for the treatment of ADHD. There is a considerable body of literature on the short-term efficacy of methylphenidate pharmacotherapy for the treatment of ADHD. However, attentional improvement is not diagnostic of ADHD. There is no diagnostic test that can confirm an ADHD diagnosis.

Recent data suggests that some children may continue to have significant ADHD-symptoms into adulthood. As a consequence, the prescription of methylphenidate for individuals 18 and older is the most rapidly growing market. Longer acting products, primarily Concerta®, have gained a significant share of the total methylphenidate market. The IMS Health National Prescription Audit *Plus*™ reported 14.8 million methylphenidate prescriptions dispensed in 2008 and 15.0 million dispensed in 2009. From January – September 2010, there were 11.4 million methylphenidate prescriptions dispensed.

### Chemistry/Pharmacology:

Methylphenidate is a CNS stimulant and produces a number of effects including appetite suppression, increased alertness and increases in blood pressure, heart rate, respiration, and body temperature. Almost complete absorption of IR methylphenidate occurs after oral administration with peak plasma levels in about 2 hours. It is extensively metabolized and about 80% of the dose is excreted in the urine as ritalinic acid. Only 20% of the administered oral dose is bioavailable due to extensive first-pass metabolism.

Biochemically, methylphenidate enhances the release and blocks the reuptake of dopamine (DA) and norepinephrine (NE) in mammalian brain. Pharmacologically methylphenidate is most closely related to cocaine. In human subjects, methylphenidate binds to the same receptor sites as cocaine in the brain and produces effects that are indistinguishable from cocaine.

### Illicit Use:

Like other potent stimulants, methylphenidate is abused for its “feel good” stimulant effects. The occasional abuser may use methylphenidate as a study aid to increase attention and stay awake. Others may use methylphenidate recreationally and combine it with alcohol or some other depressant to feel more alert

or less drunk. Serious methylphenidate abusers often snort or inject methylphenidate for its intense euphoric effects or to alleviate the severe depression and craving associated with a stimulant withdrawal syndrome.

Monitoring the Future (MTF) is a National Institute on Drug Abuse (NIDA) funded study conducted by the University of Michigan. In 2010, the MTF survey indicated that 1.5% of 8<sup>th</sup> grade students, 2.7% of 10<sup>th</sup> grade students and 2.7% of 12<sup>th</sup> grade students reported nonmedical use of Ritalin in the past year.

The National Survey on Drug Use and Health (NSDUH) is a database that measures drug use by people living in U.S. households. In 2009, the highest non-medical use of methylphenidate among youth was in the 18-25 year old age group, with 5.2% reporting lifetime non-medical use. An estimated 5.36 million people (2.1% of this population) misused methylphenidate in their lifetime, according to the 2009 NSDUH report.

The American Association of Poison Control Centers (AAPCC) report indicates that in 2009, there were a total of 9,215 methylphenidate exposures. Of this total, 6,592 were single exposures of methylphenidate. According to the Drug Abuse Warning Network (New DAWN ED), nonmedical use of methylphenidate accounted for an estimated 4,953 visits to the emergency department in 2009, more than twice the estimated 2,446 visits in 2004.

The National Forensic Laboratory Information System (NFLIS) is a DEA database that collects scientifically verified data on drug items and cases submitted to and analyzed by state and local forensic laboratories. The System to Retrieve Information from Drug Evidence (STRIDE) provides information on drug seizures reported to and analyzed by DEA laboratories. Of the substance exhibits submitted to federal, state and local forensic laboratories in 2009, there were 2,131 exhibits identified as methylphenidate. Of the exhibits submitted in 2010, 1,596 have been identified as methylphenidate, to date.

### User Population:

While a wide spectrum of the population has abused methylphenidate products, the primary abusers are individuals younger than 25 years of age; who often obtain methylphenidate from a friend or classmate and use this drug as a study aid or to party.

### Illicit Distribution:

Unlike other potent stimulants, there is no clandestine production of methylphenidate and diverted pharmaceutical products are the only source for abuse purposes. Methylphenidate is obtained from fraudulent prescriptions, doctor shopping, pharmacy theft and from friends or associates who have obtained the drug through a prescription.

### Control Status:

Methylphenidate is a Schedule II substance under the Controlled Substances Act.

Comments and additional information are welcomed by the Office of Diversion Control, Drug and Chemical Evaluation Section. Fax 202-353-1263, Telephone 202-307-7183, or Email ODE@usdoj.gov.