

**Surface and Bottom
Temperature Distributions from
the Northeast Fisheries Center
Spring and Fall
Bottom Trawl Survey Program,
1963-1987**

With addendum for 1988-1990

**by
Tamara Holzwarth
and David Mountain**

NOAA/National Marine Fisheries Service
Northeast Fisheries Science Center
Environmental Processes Division
Woods Hole, MA 02543-1097

February 1990
Revised October 1992

**Surface and Bottom
Temperature Distributions from
the Northeast Fisheries Center
Spring and Fall
Bottom Trawl Survey Program,
1963-1987**

With addendum for 1988-1990

by

**Tamara Holzwarth
and David Mountain**

**NOAA/National Marine Fisheries Service
Northeast Fisheries Science Center
Environmental Processes Division
Woods Hole, MA 02543-1097**

**February 1990
Revised October 1992**

Table of Contents

Abstract.....	1
Introduction	1
Data and methods	1
Results.....	4
Discussion	9
Acknowledgements	10
References.....	10

List of Figures

Figure 1. The region of the northeast continental shelf covered by the Northeast Fisheries Center bottom trawl survey.....	1
Figure 2. The surface (left) and bottom (right) temperature distribution for the spring of 1968	11
Figure 3. The surface (left) and bottom (right) temperature distribution for the spring of 1969	12
Figure 4. The surface (left) and bottom (right) temperature distribution for the spring of 1970	13
Figure 5. The surface (left) and bottom (right) temperature distribution for the spring of 1971	14
Figure 6. The surface (left) and bottom (right) temperature distribution for the spring of 1972	15
Figure 7. The surface (left) and bottom (right) temperature distribution for the spring of 1973	16
Figure 8. The surface (left) and bottom (right) temperature distribution for the spring of 1974	17
Figure 9. The surface (left) and bottom (right) temperature distribution for the spring of 1975	18
Figure 10. The surface (left) and bottom (right) temperature distribution for the spring of 1976	19
Figure 11. The surface (left) and bottom (right) temperature distribution for the spring of 1977	20
Figure 12. The surface (left) and bottom (right) temperature distribution for the spring of 1978	21
Figure 13. The surface (left) and bottom (right) temperature distribution for the spring of 1979	22
Figure 14. The surface (left) and bottom (right) temperature distribution for the spring of 1980	23
Figure 15. The surface (left) and bottom (right) temperature distribution for the spring of 1981	24
Figure 16. The surface (left) and bottom (right) temperature distribution for the spring of 1982	25
Figure 17. The surface (left) and bottom (right) temperature distribution for the spring of 1983	26

Figure 18. The surface (left) and bottom (right) temperature distribution for the spring of 1984	27
Figure 19. The surface (left) and bottom (right) temperature distribution for the spring of 1985	28
Figure 20. The surface (left) and bottom (right) temperature distribution for the spring of 1986	29
Figure 21. The surface (left) and bottom (right) temperature distribution for the spring of 1987	30
Figure 22. The surface (left) and bottom (right) temperature distribution for the fall of 1963	31
Figure 23. The surface (left) and bottom (right) temperature distribution for the fall of 1964	32
Figure 24. The surface (left) and bottom (right) temperature distribution for the fall of 1965	33
Figure 25. The surface (left) and bottom (right) temperature distribution for the fall of 1966	34
Figure 26. The surface (left) and bottom (right) temperature distribution for the fall of 1967	35
Figure 27. The surface (left) and bottom (right) temperature distribution for the fall of 1968	36
Figure 28. The surface (left) and bottom (right) temperature distribution for the fall of 1969	37
Figure 29. The surface (left) and bottom (right) temperature distribution for the fall of 1970	38
Figure 30. The surface (left) and bottom (right) temperature distribution for the fall of 1971	39
Figure 31. The surface (left) and bottom (right) temperature distribution for the fall of 1972	40
Figure 32. The surface (left) and bottom (right) temperature distribution for the fall of 1973	41
Figure 33. The surface (left) and bottom (right) temperature distribution for the fall of 1974	42
Figure 34. The surface (left) and bottom (right) temperature distribution for the fall of 1975	43
Figure 35. The surface (left) and bottom (right) temperature distribution for the fall of 1976	44
Figure 36. The surface (left) and bottom (right) temperature distribution for the fall of 1977	45
Figure 37. The surface (left) and bottom (right) temperature distribution for the fall of 1978	46
Figure 38. The surface (left) and bottom (right) temperature distribution for the fall of 1979	47
Figure 39. The surface (left) and bottom (right) temperature distribution for the fall of 1980	48
Figure 40. The surface (left) and bottom (right) temperature distribution for the fall of 1981	49

Figure 41. The surface (left) and bottom (right) temperature distribution for the fall of 1982	50
Figure 42. The surface (left) and bottom (right) temperature distribution for the fall of 1983	51
Figure 43. The surface (left) and bottom (right) temperature distribution for the fall of 1984	52
Figure 44. The surface (left) and bottom (right) temperature distribution for the fall of 1985	53
Figure 45. The surface (left) and bottom (right) temperature distribution for the fall of 1986	54
Figure 46. The surface (left) and bottom (right) temperature distribution for the fall of 1987.	55
Figure 47. Average surface temperature in the four regions of the shelf shown in Figure 1 for the spring (top) and the fall (bottom)	56
Figure 48. Average bottom temperature in the four regions of the continental shelf shown in Figure 1 for the spring (top) and the fall (bottom)	57
Figure 49. Average surface temperature anomaly in the spring for the four regions of the shelf shown in Figure 1	58
Figure 50. Average bottom temperature anomaly in the spring for the four regions of the shelf shown in Figure 1	59
Figure 51. Average surface temperature anomaly in the fall for the four regions of the shelf shown in Figure 1	60
Figure 52. Average bottom temperature anomaly in the fall for the four regions of the shelf shown in Figure 1	61
Figure 53. Comparison of the average spring and fall bottom temperatures (top) and temperature anomalies (bottom) in the Southern Middle Atlantic Bight calculated by Davis (1979) and in this report	62

List of Tables

Table 1. Inherent uncertainties in the three measurement techniques used in obtaining the temperature data.....	2
Table 2. Areal average surveys of surface and bottom temperature anomaly for the spring and fall bottom trawl surveys in the four regions of the northeast continental shelf shown in Figure 1	5

ABSTRACT

Temperature observations from the Northeast Fisheries Center spring and fall bottom-trawl survey program are summarized for the period 1963 to 1987. The distributions of surface and bottom temperature are presented for each survey. The average surface and bottom temperatures and temperature anomalies have been calculated for each survey in four regions of the northeast continental shelf: the Gulf of Maine, Georges Bank, northern Middle Atlantic Bight, and southern Middle Atlantic Bight. The temperature anomalies are determined relative to an established mean temperature field.

INTRODUCTION

The Northeast Fisheries Center has conducted bottom-trawl surveys each spring since 1968 and each fall since 1963 to determine the distribution and relative abundance of the groundfish stocks off the northeast coast of the United States. The trawls have been made on a stratified random grid with approximately 300 stations from Cape Hatteras to the Gulf of Maine, and from near the coast to the edge of the continental shelf (Grosslein 1969). On selected stations, temperature observations were made to record surface and bottom temperature.

Earlier presentations and summaries of the bottom temperature data from the bottom-trawl surveys are included in Davis (1978 and 1979). This report presents contoured distributions of the surface and bottom temperatures for all of the surveys from 1963 to 1987. Temperature anomalies relative to an established mean temperature field have been determined for all of the surface and bottom observations on each survey. The areal average temperatures and temperature anomalies for four regions of the shelf have been calculated and are presented in time series form.

DATA AND METHODS

Surface temperature measurements were made using a surface-sampling bucket with thermometer. In the early years of the survey

program, bottom-temperature measurements were made using a mechanical bathythermograph (MBT). Since 1971, the bottom measurements have been made using expendable bathythermograph probes (XBT). The accuracies of these different measurement techniques are listed in Table 1.

The temperature data used in this report were retrieved from the survey data base maintained on computer at the Northeast Fisheries Center in Woods Hole, Massachusetts. The date, position, bottom depth, and station number for the observations were also retrieved from the data base. Observations from stations without assigned station numbers were considered. When duplicate observations existed at the same location on a survey, only the first observation at the location was used.

Surface and bottom temperature distributions for each spring and fall survey were generated using the SURFACE III contouring software on a VAX 11/785 computer.

The shelf-wide surveys require five or six weeks to complete. Distributions of mean surface temperature presented by Mountain and Holzwarth (1989) suggest that changes of 2° to 4° C could be expected in water temperature, particularly at the surface, over the period of a survey. This means that the survey temperature distributions are not synoptic. Also, the timing of surveys varied by a few weeks between years, depending upon ship schedules, so that direct comparison between years may not be a reliable indication of

Table 1. Inherent uncertainties in the three measurement techniques used in obtaining the temperature data

Surface bucket with thermometer	$\pm 0.2^\circ \text{C}$
Mechanical bathythermograph (MBT)	$\pm 1.0^\circ \text{C}$
Expendable bathythermograph (XBT)	$\pm 0.2^\circ \text{C}$

indication of actual interannual variation in water temperature. In order to account for the different timing of observations within a survey and between years, a temperature anomaly is calculated for each temperature observation. The anomaly represents the difference between the observed temperature and the expected temperature at the location and on the day of the year that the observation was made.

The expected temperatures are derived through a method described by Mountain (1989). This method uses a series of mean annual temperature curves for about 160 standard station locations on the continental shelf. These curves were derived for both surface and bottom temperature from analysis of an eleven-year time series of hydrographic observations at the standard station locations. The expected temperature for a given location and day is determined by first identifying the closest standard stations. Then the expected temperature at each selected standard station is determined for the given day from its annual curve. The expected temperature at the given location is then determined by a weighted average of the temperatures at the nearby standard stations, with the weighting being inversely proportional to the square of the distance from the standard station to the given location.

Bottom temperature on the shelf can vary considerably with bottom depth. When estimating the expected value for a bottom temperature observation, the difference in depth between the observation site and a potential nearby standard station was determined. If the depth difference was greater than 25 m

and greater than one quarter of the water depth at the observation site, the standard station was judged to not represent the bottom conditions at the observation site and was not included in determining the expected bottom temperature. This depth selection process is somewhat arbitrary and the criteria used were selected after trials with a range of values. The depth filtering is especially important in the region of large gradient in bottom depth between the northern edge of Georges Bank and the southern Gulf of Maine.

To summarize the temperature and temperature anomaly data, the survey area is divided into four regions - Gulf of Maine, Georges Bank, northern Middle Atlantic Bight, and southern Middle Atlantic Bight (Figure 1). These are the same regions used by Davis (1978 and 1979). If the temperature observations within a region were not uniformly distributed, a simple average of them may not provide the best characterization of the temperature conditions in the region. Instead, an areal weighted average is desired. An areal average was calculated by first gridding the region into 0.25° longitude by 0.20° latitude boxes. For each box, the nearest survey stations were identified and the area of the box was divided among these stations in proportion to the inverse square of the distance from the station to the center of the box. This was done for all of the boxes in a region so that all of the area in the region was divided among the stations. For each station the areas assigned from the different boxes were summed to determine the total assigned area to the station. Then, a weighting factor was calculated for each station by dividing the total area assigned to the station by the total area of the region. These weighting factors indicate the proportion of the region each station represents. The areal average temperature or anomaly for the region was calculated by summing the products of the station weights and station temperature or anomaly values:

$$M = \sum a_i V_i \quad (1)$$




Figure 1. The region of the northeast continental shelf covered by the Northeast Fisheries Center bottom trawl survey. The boundaries of the four areas of the shelf for which average temperature and anomaly values are calculated are shown - Gulf of Maine, Georges Bank, northern Middle Atlantic Bight, and southern Middle Atlantic Bight.

where:

- M = the areal average value
 a_i = the weighting factor for the i^{th} station
 V_i = the temperature or temperature anomaly value for the i^{th} station

If, for any grid box in a region, no survey stations were found within approximately 60 km, an areal average for the region was not calculated. Instead, a simple average of the observations that were within the boundaries of the region was determined.

To establish confidence limits on the calculated average temperatures, the measurement error must be considered. The original temperature measurements have inherent uncertainties, as listed in Table 1. The measurement errors for the temperature observations are assumed to be normally distributed with a standard deviation equal to these uncertainties. The regional average temperatures are determined by averaging a number of observations and therefore the expected standard deviation associated with the average will decrease in proportion to the inverse square root of the number of stations. Since 40 or more stations are usually included in an average, this means that the standard deviations for the averages are generally less than 0.1°C . For the MBT data they are less than 0.2°C . By calculating average temperature over broad areas of the shelf so that many observations are included, the confidence limits on the average temperature values are relatively narrow.

In calculating the average temperature anomalies, an additional source of error must be considered. The anomaly for an individual observation is the difference between the observed temperature and the expected temperature at the same location for the same day of the year. The uncertainty in the expected temperature is determined from the standard deviations associated with annual curves for the standard stations used to calcu-

late the expected temperature (Mountain 1989). The resulting standard deviations for the expected temperatures are generally on the order of 1°C . The areal average temperature anomaly is, in essence, the difference of two means - the mean of the observed temperatures minus the mean of the expected temperatures. Therefore the uncertainty in the areal average anomaly is determined by:

$$\text{SDV1} = \sqrt{a_i^2 \sigma_i^2 + a_i^2 \alpha_i^2} \quad (2)$$

where

- a_i = the weighting factor for the survey station in the areal averaging
 σ_i = the standard deviation of the temperature observation for the i^{th} survey station
 α_i = the standard deviation of the expected temperature for the i^{th} survey station

This value indicates how well the calculated anomaly represents the true average temperature anomaly for the region as a whole. The values for SDV1 are generally on the order of 0.1° to 0.3°C .

Another question of interest is how well does the areal average value represent the anomaly one might find at any particular location within the region. This uncertainty is represented by the standard deviation of the individual anomalies within the region and is referred to in this report as SDV2.

RESULTS

The areal average temperatures and temperature anomalies have been calculated for the four regions in both the spring and the fall and for the surface and the bottom. The results are listed in Table 2. Cases where a simple average was determined are indicated in the table by an asterisk. The standard deviations SDV1 and SDV2 are also included in the table.

Table 2. Areal average surface and bottom temperature and temperature anomaly for the spring and fall bottom trawl surveys in the four regions of the northeast continental shelf shown in Figure 1: "#Obs", the number of observations included in each average; "Temp", the areal average temperature; "Anomaly", the areal average temperature anomaly; "SDV1", the standard deviation associated with the average temperature anomaly; "SDV2", the standard deviation of the individual anomalies from which the average anomaly was derived. An asterisk indicates that a true areal average could not be calculated due to poor station coverage and that the average values listed were derived from a simple average of the observations that were within the region. All of the temperature values are in °C.

Year	Spring - Gulf of Maine									
	Surface					Bottom				
	#Obs	Temp	Anomaly	SDV1	SDV2	#Obs	Temp	Anomaly	SDV1	SDV2
1968	74	3.9	-0.7	0.1	0.7	67	5.1	-0.5	0.2	1.0
1969	68	3.7	-0.5	0.2	1.0	59	5.4	-0.2	0.2	1.1
1970	93	3.7	-0.4	0.1	0.7	87	6.4	0.6	0.2	1.1
1971	89	3.9	-1.0	0.1	0.6	81	6.1	0.6	0.2	1.2
1972	97	4.6	-0.0	0.1	0.8	93	6.2	0.8	0.1	1.0
1973	73	6.0	-0.2	0.2	0.9	71	6.4	0.6	0.1	1.2
1974	71	5.7	0.2	0.2	1.0	47*	6.6	1.1	0.1	0.7
1975	68	5.2	-1.0	0.2	1.1	62	6.6	0.6	0.1	1.3
1976	102	6.2	1.0	0.1	0.7	98	7.0	1.3	0.1	1.0
1977	97	6.3	-0.1	0.1	1.2	93	5.5	-0.2	0.1	1.2
1978	99	5.8	-1.1	0.1	0.8	93	5.5	-0.2	0.1	0.9
1979	120	5.4	0.0	0.1	0.9	114	5.4	-0.2	0.1	0.9
1980	81	5.8	0.1	0.1	0.7	78	5.6	0.0	0.1	1.0
1981	89	7.2	0.1	0.1	0.6	82	5.5	-0.3	0.1	1.0
1982	81	5.7	-0.2	0.1	1.0	75	5.8	0.1	0.1	0.9
1983	82	5.3	0.1	0.1	1.0	78	5.6	-0.0	0.1	1.2
1984	76	4.4	-0.4	0.1	1.0	75	5.9	0.2	0.1	1.2
1985	28	4.4	0.1	0.2	1.1	25*	5.4	0.3	0.2	1.1
1986	39	6.0	1.0	0.2	0.7	37	7.2	1.7	0.2	0.9
1987	39	4.3	-1.0	0.2	1.1	38	5.6	0.0	0.2	1.2
	Fall - Gulf of Maine									
1963	89	8.6	-1.0	0.1	0.7	86	7.0	-0.7	0.2	1.2
1964	74	8.3	-1.6	0.2	0.9	73	5.8	-1.8	0.2	1.0
1965	75	10.5	-2.2	0.2	1.4	73	6.1	-1.8	0.2	1.6
1966	72	10.8	-1.5	0.1	1.2	65	5.9	-2.0	0.2	1.6
1967	79	8.6	-1.3	0.1	0.7	58	6.2	-1.6	0.2	1.1
1968	68	9.6	-0.5	0.2	1.2	60	7.6	-0.4	0.2	1.3
1969	79	9.8	-0.2	0.2	0.7	67	7.4	-0.5	0.2	1.1
1970	80	10.2	-0.1	0.2	0.7	77	7.6	-0.2	0.2	1.4
1971	88	11.6	1.0	0.1	1.0	83	8.4	0.5	0.2	1.7
1972	87	9.9	-0.7	0.2	0.8	78	8.2	0.3	0.1	1.3
1973	83	10.1	-0.4	0.1	0.9	73	8.5	0.7	0.1	1.4
1974	91	11.6	0.1	0.1	0.8	84	9.2	1.3	0.1	1.6
1975	99	11.7	0.5	0.1	0.8	96	8.2	0.4	0.1	1.5
1976	79	10.0	-0.4	0.1	0.7	70	9.1	1.2	0.1	1.2
1977	114	9.5	-0.0	0.1	0.8	103	7.9	0.1	0.1	1.3
1978	184	11.1	-0.3	0.1	0.7	171	7.4	-0.4	0.1	1.1
1979	177	10.9	0.4	0.1	0.6	165	8.1	0.4	0.1	1.4
1980	93	10.1	-0.8	0.1	0.8	84	7.8	-0.2	0.1	1.4
1981	104	10.6	-0.4	0.1	0.7	99	7.4	-0.6	0.1	1.2
1982	100	11.2	0.1	0.1	0.8	95	7.8	-0.2	0.1	1.6
1983	106	11.2	0.1	0.1	0.8	103	8.1	0.1	0.1	1.7
1984	57	12.4	0.9	0.2	0.8	45	8.9	0.8	0.2	1.6
1985	34	11.7	0.8	0.2	1.1	33	8.6	0.6	0.2	2.3
1986	40	11.3	-0.0	0.2	0.7	52	8.5	0.6	0.2	1.4
1987	38	11.4	-0.5	0.2	1.0	35	7.8	-0.1	0.2	2.2

Table 2. Continued

Year	Surface					Bottom				
	#Obs	Temp	Anomly	SDV1	SDV2	#Obs	Temp	Anomaly	SDV1	SDV2
1968	49	4.1	-0.8	0.2	0.5	36	4.0	-1.1	0.3	0.7
1969	59	5.1	0.5	0.2	0.8	45	4.9	0.2	0.3	0.7
1970	76	4.5	-0.6	0.2	0.8	62	4.6	-0.7	0.3	1.1
1971	64	3.8	-0.7	0.2	0.5	51	4.3	-0.6	0.3	1.0
1972	59	5.0	0.2	0.2	1.2	46	5.1	0.2	0.2	0.8
1973	59	5.6	0.0	0.2	0.7	47	6.4	1.0	0.2	1.4
1974	56	5.9	0.8	0.2	1.1	44	6.5	1.1	0.3	0.9
1975	51	5.5	-0.2	0.2	0.8	41	6.1	0.6	0.2	0.8
1976	60	6.0	1.1	0.2	0.6	51	6.0	0.9	0.2	1.1
1977	63	7.2	1.4	0.2	1.1	50	6.0	0.4	0.2	1.0
1978	61	4.7	-0.8	0.2	0.9	52	4.6	-0.9	0.2	1.0
1979	109	5.5	-0.0	0.2	0.7	97	5.5	-0.2	0.2	0.8
1980	59	6.4	0.5	0.2	1.1	51	6.5	0.7	0.2	1.0
1981	57	5.8	-0.3	0.2	0.7	43	5.7	-0.2	0.2	0.7
1982	58	5.2	-0.4	0.2	0.7	42	5.0	-0.3	0.2	1.3
1983	55	6.3	1.1	0.2	0.9	45	6.0	0.7	0.3	0.8
1984	54	5.6	0.8	0.2	0.8	43	6.0	1.0	0.2	0.9
1985	23	5.4	0.7	0.3	1.0	17	5.2	0.4	0.3	0.7
1986	23	6.1	1.1	0.3	0.5	20	6.4	1.4	0.3	1.2
1987	27	6.6	0.9	0.2	2.6	22	6.4	0.6	0.3	1.4

Fall - Georges Bank

1963	43	8.9	-1.4	0.2	1.1	30	8.7	-1.6	0.3	1.6
1964	52	10.7	-2.0	0.2	1.1	36	9.3	-2.3	0.3	1.0
1965	56	12.3	-1.3	0.2	1.8	40	11.0	-1.2	0.3	1.8
1966	54	11.4	-1.8	0.2	1.2	37	10.1	-2.0	0.3	1.6
1967	50	8.8	-1.7	0.2	1.0	36	8.6	-1.8	0.3	0.9
1968	52	12.7	-0.4	0.2	1.4	41	12.3	0.2	0.3	1.4
1969	55	12.3	-1.0	0.2	1.3	29	11.9	-0.6	0.3	1.3
1970	55	13.3	-0.5	0.2	1.2	40	11.3	-0.9	0.3	1.7
1971	55	15.0	1.0	0.2	1.5	49	12.3	-0.0	0.3	1.9
1972	55	13.0	-1.0	0.2	1.3	43	12.1	-0.4	0.2	1.5
1973	57	14.6	0.3	0.2	1.4	42	13.4	1.0	0.2	1.7
1974	56	15.1	0.5	0.2	1.4	47	13.1	0.5	0.2	1.4
1975	63	14.7	-0.1	0.2	2.0	50	11.9	-0.6	0.2	1.4
1976	50	13.6	0.1	0.2	1.0	43	13.4	1.1	0.2	1.3
1977	73	13.5	-0.2	0.2	1.3	61	13.2	0.8	0.2	2.1
1978	118	14.1	-0.2	0.2	1.3	105	11.6	-0.8	0.2	1.4
1979	105	14.3	0.5	0.1	1.0	91	13.2	0.9	0.2	1.5
1980	78	15.0	0.7	0.2	1.8	62	13.2	0.9	0.2	1.2
1981	74	12.8	-1.3	0.2	1.4	62	11.4	-1.1	0.2	1.5
1982	69	13.8	-0.2	0.2	1.7	61	11.5	-0.9	0.2	1.7
1983	89	15.1	0.7	0.2	1.2	77	11.8	-0.7	0.2	1.8
1984	32	14.9	0.0	0.3	1.0	26	13.1	0.6	0.3	2.0
1985	26	15.5	1.5	0.3	1.5	22	13.2	1.1	0.3	2.5
1986	20	13.8	-0.6	0.3	1.1	28	12.4	-0.0	0.3	1.1
1987	29	13.9	-1.0	0.2	0.8	26	11.3	-1.6	0.3	2.6

Table 2. Continued

Year	Surface					Bottom				
	#Obs	Temp	Anomaly	SDV1	SDV2	#Obs	Temp	Anomaly	SDV1	SDV2
1968	30	3.0	-1.4	0.3	0.9	20*	3.5	-1.2	0.4	2.5
1969	41	4.1	-0.3	0.3	1.2	28	5.4	-0.3	0.4	1.8
1970	43	5.8	-0.6	0.3	0.8	32*	4.9	-0.7	0.3	1.7
1971	47	4.0	-0.8	0.3	0.7	32*	5.9	0.6	0.3	2.3
1972	48	5.4	1.0	0.3	1.0	34	6.3	1.0	0.4	1.4
1973	50	6.5	1.9	0.3	1.8	37	7.4	2.0	0.4	1.8
1974	39	6.7	1.9	0.3	1.3	30	8.2	2.4	0.4	1.8
1975	28	5.8	1.0	0.4	0.9	23*	5.7	0.8	0.3	1.9
1976	44	6.2	1.7	0.3	1.0	34*	7.3	2.5	0.3	1.5
1977	39	6.1	0.4	0.3	1.3	27	5.2	-1.0	0.4	2.1
1978	55	4.4	-0.6	0.3	0.6	45	3.9	-1.5	0.4	1.4
1979	55	6.0	0.1	0.3	1.1	44	6.2	0.2	0.3	1.6
1980	93	6.3	0.8	0.2	1.0	81	6.3	0.3	0.3	1.1
1981	50	6.0	0.3	0.3	0.7	38	5.6	0.6	0.4	1.6
1982	17*	4.3	-1.0	0.4	0.5	15*	5.1	0.0	0.4	2.4
1983	34	6.0	1.2	0.3	0.9	23*	5.4	1.2	0.3	1.1
1984	41	5.0	0.5	0.3	0.9	31	5.7	-0.2	0.4	1.7
1985	14	5.5	1.2	0.4	1.4	13*	6.1	1.5	0.4	1.8
1986	15*	6.5	1.9	0.4	0.8	11*	6.1	1.7	0.5	1.3
1987	22	6.2	0.8	0.4	1.8	16*	5.6	-0.1	0.4	0.7

Fall - Middle Atlantic Bight North

1963	30*	10.3	-0.8	0.3	1.2	20*	11.4	0.2	0.4	1.2
1964	32	13.2	-1.9	0.3	0.9	20*	11.1	-2.2	0.4	1.3
1965	35	13.1	-1.3	0.3	1.5	23	10.3	-3.0	0.4	1.8
1966	34	12.4	-1.7	0.4	1.4	23*	9.6	-3.5	0.4	1.2
1967	46	14.3	-0.7	0.3	0.9	29	9.3	-3.7	0.4	1.2
1968	39	16.5	0.4	0.3	0.9	29	11.0	-1.9	0.4	1.8
1969	37	16.5	0.4	0.3	1.4	29*	12.3	-0.4	0.3	1.6
1970	43	16.9	1.0	0.3	2.0	30*	10.7	-2.5	0.3	1.4
1971	47	19.3	2.2	0.3	1.0	38	11.1	-1.5	0.4	2.2
1972	43	18.2	1.2	0.3	1.3	37	13.0	0.2	0.4	1.6
1973	43	17.9	0.6	0.3	1.0	30*	12.9	0.6	0.3	1.5
1974	40	17.9	0.1	0.3	1.3	28	12.5	-0.2	0.4	1.5
1975	36	16.0	0.0	0.3	1.1	28*	12.0	-0.8	0.3	1.3
1976	42	17.8	0.6	0.3	1.2	32	12.4	-0.3	0.4	1.4
1977	41	16.7	-0.1	0.3	1.2	31*	13.0	0.0	0.3	1.6
1978	73	16.6	-0.5	0.3	0.9	59	11.6	-0.7	0.3	1.7
1979	67	16.5	-0.2	0.3	1.2	56	11.4	-1.3	0.3	1.4
1980	32	18.4	1.7	0.3	1.6	27*	12.5	-0.4	0.3	1.9
1981	41	14.7	-2.1	0.3	1.5	33*	10.8	-1.7	0.3	1.2
1982	37	17.6	0.6	0.3	1.5	25	12.7	-0.2	0.4	1.6
1983	36	18.2	1.0	0.3	0.7	27	11.2	-1.3	0.4	1.4
1984	30	17.6	-0.1	0.4	1.4	21	12.0	-0.2	0.4	2.0
1985	13	17.5	1.1	0.5	1.7	9*	13.4	0.8	0.6	1.0
1986	24	18.1	0.8	0.4	1.3	22	12.4	-0.3	0.5	1.7
1987	18*	17.6	0.0	0.4	0.8	17*	10.8	-1.1	0.4	1.9

Spring - Middle Atlantic Bight South

Year	Surface					Bottom				
	#Obs	Temp	Anomaly	SDV1	SDV2	#Obs	Temp	Anomaly	SDV1	SDV2
1968	57	4.8	-1.0	0.3	1.2	44	5.5	-0.5	0.4	1.4
1969	51	4.9	-1.0	0.3	1.3	36	5.0	-1.0	0.4	1.4
1970	54	8.2	-1.4	0.3	1.2	37	6.9	-0.9	0.4	1.3
1971	51	6.4	-0.5	0.3	1.6	39	6.6	-0.1	0.4	2.4
1972	55	7.3	1.6	0.3	1.4	46	8.1	2.2	0.3	1.4
1973	62	6.9	0.7	0.3	1.6	46	7.4	1.2	0.3	1.3
1974	41	9.6	3.2	0.4	1.7	31	9.8	3.5	0.5	1.7
1975	41*	7.6	1.1	0.3	1.1	30*	7.4	1.2	0.3	1.0
1976	59	7.8	1.9	0.2	1.4	48	8.3	2.3	0.3	1.2
1977	58	7.0	0.4	0.2	1.7	50	6.2	-0.2	0.3	1.5
1978	56	6.2	-0.4	0.3	1.4	48	6.3	0.0	0.3	1.6
1979	55	7.2	0.3	0.3	1.5	39	6.7	0.1	0.4	1.3
1980	48	7.4	0.9	0.3	2.0	38	7.4	0.9	0.4	1.7
1981	52	7.0	0.2	0.3	1.3	41	6.8	0.5	0.4	1.2
1982	17*	6.6	-0.4	0.5	1.4	14*	6.6	-0.1	0.6	1.2
1983	47*	7.8	1.3	0.3	1.1	37*	7.7	1.1	0.3	1.4
1984	49	6.7	1.0	0.3	1.3	38	7.3	1.4	0.4	2.1
1985	16*	8.0	1.5	0.4	1.6	12*	6.9	0.9	0.5	1.1
1986	26*	7.4	1.3	0.4	1.3	22*	7.2	1.1	0.4	1.0
1987	31*	6.5	-0.3	0.3	1.7	35	6.2	-0.3	0.4	2.0

Fall - Middle Atlantic Bight South

1963	10*	10.5	-1.2	0.5	0.9	8*	10.3	-1.2	0.6	0.6
1964	9*	14.1	-2.0	0.6	1.2	5*	9.7	-2.6	0.7	1.3
1965	8*	14.1	-1.8	0.7	1.5	2*	10.6	-2.8	1.0	1.9
1966	12*	12.9	-2.4	0.6	1.5	3*	11.3	-1.8	0.8	2.4
1967	61	16.5	-0.5	0.2	0.9	38	12.9	-1.6	0.4	2.0
1968	62	19.3	1.2	0.2	1.0	51	12.8	-1.9	0.3	2.8
1969	49	18.7	0.4	0.3	0.7	41	15.3	1.1	0.3	2.2
1970	61	22.8	1.0	0.2	0.8	47	10.0	-3.3	0.3	2.4
1971	57	21.0	1.9	0.3	1.3	41	12.8	-1.6	0.4	3.9
1972	49	19.6	-0.2	0.3	1.1	37	15.1	1.1	0.3	2.5
1973	47	21.3	1.4	0.3	1.4	39	14.5	0.3	0.3	1.9
1974	50	20.8	0.6	0.3	1.2	40	14.6	0.6	0.3	1.8
1975	56	16.8	0.4	0.3	1.2	44	14.3	-0.2	0.3	1.7
1976	64	19.3	0.4	0.2	0.8	54	14.5	0.2	0.3	2.4
1977	58	19.7	-0.1	0.3	1.1	48	13.2	-1.1	0.3	1.7
1978	46	22.2	0.4	0.3	1.1	40	11.0	-2.0	0.3	1.7
1979	47	20.3	0.1	0.3	1.4	37	12.5	-1.4	0.3	2.4
1980	52*	21.3	1.5	0.3	0.8	40*	11.4	-1.8	0.3	1.8
1981	48	19.3	-1.0	0.3	1.1	35	13.8	0.2	0.4	2.5
1982	50	20.8	0.3	0.3	1.6	42	12.5	-1.1	0.4	2.0
1983	53	21.6	0.9	0.3	1.1	42	13.3	-0.4	0.3	1.7
1984	45	20.7	-0.5	0.3	1.8	37	11.8	-1.6	0.3	2.6
1985	26	21.7	1.6	0.4	1.7	22*	15.2	1.4	0.4	2.4
1986	30	21.0	-0.1	0.3	1.6	35	14.4	0.6	0.4	3.4
1987	25	22.7	1.2	0.4	1.0	21	11.5	-1.6	0.4	2.1

The surface and bottom temperature distributions for all the spring and fall surveys are presented in Figures 2 through 46. The time series of average temperature and of temperature anomaly for each region for spring and fall and for surface and bottom are presented in Figures 47 through 52. Since the standard deviations associated with the temperature and the anomaly values are relatively small (0.1° to 0.3° C) no error bars are included in these figures.

DISCUSSION

The results presented in this report include the period for which Davis (1978, 1979) calculated similar sets of areal average bottom temperatures and anomalies from the same survey data. Generally, the results compare well. For example, Figure 53 shows the results for the southern Middle Atlantic Bight from the two studies. The average temperatures are very close, particularly considering that Davis (1979) used 2° C increments in his areal averaging. The temperature-anomaly series also show similar trends. The differences that exist are not surprising since the anomalies were determined relative to different reference periods, 1967 to 1976 for Davis and 1977 to 1987 for this report, and since Davis had only a single point (the Nantucket Lightship) with which to compensate for different sampling dates over the whole shelf.

The time-series plots of the areal average temperature data (Figures 47 to 48) illustrate many of the characteristic features of the temperature patterns on the northeast continental shelf. In the spring, the surface and the bottom temperatures in the different regions are all very similar, although the southern Middle Atlantic Bight temperatures are about 1° C warmer than the other regions. The similarity in surface and bottom temperatures indicates that seasonal warming and thermocline formation have not begun when

the spring survey is conducted (mid-March to the end of April).

In the fall, the surface temperatures are considerably warmer than the bottom temperatures within the different regions. At the surface there is an increase in temperature from north (Gulf of Maine) to south (southern Middle Atlantic Bight). The bottom temperature in the Gulf of Maine stands out as being a few degrees colder than the other areas, which exhibit fairly similar average bottom temperatures. The colder bottom temperatures in the Gulf of Maine are due in large part to the Gulf being considerably deeper than the other three regions such that heat from the seasonal surface warming does not penetrate to the bottom.

The temperature anomaly time-series plots (Figures 49 to 52) illustrate characteristics of the interannual variability of temperatures on the continental shelf. The variability in the fall is generally larger than that in the spring. Within a region and for either season the surface and bottom temperatures exhibit a comparable degree of variability. The range of interannual temperature variation is somewhat larger in the Middle Atlantic Bight (3° to 4° C) than on Georges Bank or in the Gulf of Maine (2° to 3° C).

The different temperature anomaly time series appear autocorrelated, the positive values tend to be grouped together as do the negative values. This means that there were multiple-year periods when the shelf waters were generally warmer or cooler than average. As shown by Davis (1978 and 1979) the mid-1960s were quite cold, while the early to middle 1970s were warm. The temperature anomalies presented here (Figures 49 to 52) suggest that during the period of the late 1970s and early 1980s (1978 to 1982) the temperatures were generally intermediate between the two earlier extremes. In the mid-1980s (1983 to 1986) the temperatures were again warm, being comparable to the mid-1970s. This grouping of years into warm and cool periods is not precise and there is consid-

erable variability within and between the different time series on a year-to-year basis. The relatively small uncertainties associated with the temperature anomaly values (SDV1 in Table 1), however, suggest that the differences in temperature between individual years from the cool and the warm periods are significant.

Acknowledgements

James Manning provided valuable assistance in modifying the computer programs used to generate the contoured temperature figures in this report. Michael Pennington provided valuable advice on statistical procedures for determining the uncertainty in the data presented.

References

- Davis, C.W. 1978. Seasonal bottom-water temperature trends in the Gulf of Maine and on Georges Bank, 1967-1975. Seattle: NOAA/NMFS Scientific Publication Office. NOAA Tech. Rep., NMFS-SSRF-725. 17 p. Available from: National Technical Information Service, 5285 Port Royal Road, Springfield, VA, 22161.
- Davis, C.W. 1979. Bottom-water temperature trends in the Middle Atlantic Bight during spring and autumn, 1964-1976. Seattle, Washington: NOAA/NMFS Scientific Publications Office. NOAA Tech. Rep. NMFS-SSRF-738. 13 p. Available from: National Technical Information Service, 5285 Port Royal Road, Springfield, VA, 22161.
- Grosslein, M. D. 1969. Groundfish survey program of BCF Woods Hole. *Commer. Fish. Rev.* 31(8-9):22-30.
- Mountain, David G. 1989. TEMPEST a computer program for estimating temperature on the northeast continental shelf. Woods Hole, MA: NMFS Northeast Fisheries Center. Reference Document 89-02. Available from: Information Services Section, NMFS/Northeast Fisheries Center, Woods Hole, MA, 02543.
- Mountain, David G. and Tamara J. Holzwarth. 1990. Surface and bottom temperature distributions for the northeast continental shelf. Woods Hole, MA: NMFS/Northeast Fisheries Center. NOAA Tech. Mem. NMFS-F/NEC-73. Available from: Information Services Section, NMFS/Northeast Fisheries Center, Woods Hole, MA, 02543.




Figure 2. The surface (left) and bottom (right) temperature distribution for the spring of 1968. The dashed contour is the 200 m isobath.




Figure 3. The surface (left) and bottom (right) temperature distribution for the spring of 1969. The dashed contour is the 200 m isobath.




Figure 4. The surface (left) and bottom (right) temperature distribution for the spring of 1970. The dashed contour is the 200 m isobath.




Figure 5. The surface (left) and bottom (right) temperature distribution for the spring of 1971. The dashed contour is the 200 m isobath.




Figure 6. The surface (left) and bottom (right) temperature distribution for the spring of 1972. The dashed contour is the 200 m isobath.




Figure 7. The surface (left) and bottom (right) temperature distribution for the spring of 1973. The dashed contour is the 200 m isobath.




Figure 8. The surface (left) and bottom (right) temperature distribution for the spring of 1974. The dashed contour is the 200 m isobath.




Figure 9. The surface (left) and bottom (right) temperature distribution for the spring of 1975. The dashed contour is the 200 m isobath.




Figure 10. The surface (left) and bottom (right) temperature distribution for the spring of 1976. The dashed contour is the 200 m isobath.




Figure 11. The surface (left) and bottom (right) temperature distribution for the spring of 1977. The dashed contour is the 200 m isobath.




Figure 12. The surface (left) and bottom (right) temperature distribution for the spring of 1978. The dashed contour is the 200 m isobath.




Figure 13. The surface (left) and bottom (right) temperature distribution for the spring of 1979. The dashed contour is the 200 m isobath.




Figure 14. The surface (left) and bottom (right) temperature distribution for the spring of 1980. The dashed contour is the 200 m isobath.




Figure 15. The surface (left) and bottom (right) temperature distribution for the spring of 1981. The dashed contour is the 200 m isobath.




Figure 16. The surface (left) and bottom (right) temperature distribution for the spring of 1982. The dashed contour is the 200 m isobath.




Figure 17. The surface (left) and bottom (right) temperature distribution for the spring of 1983. The dashed contour is the 200 m isobath.




Figure 18. The surface (left) and bottom (right) temperature distribution for the spring of 1984. The dashed contour is the 200 m isobath.




Figure 19. The surface (left) and bottom (right) temperature distribution for the spring of 1985. The dashed contour is the 200 m isobath.




Figure 20. The surface (left) and bottom (right) temperature distribution for the spring of 1986. The dashed contour is the 200 m isobath.




Figure 21. The surface (left) and bottom (right) temperature distribution for the spring of 1987. The dashed contour is the 200 m isobath.




Figure 22. The surface (left) and bottom (right) temperature distribution for the fall of 1963. The dashed contour is the 200 m isobath.




Figure 23. The surface (left) and bottom (right) temperature distribution for the fall of 1964. The dashed contour is the 200 m isobath.




Figure 24. The surface (left) and bottom (right) temperature distribution for the fall of 1965. The dashed contour is the 200 m isobath.




Figure 25. The surface (left) and bottom (right) temperature distribution for the fall of 1966. The dashed contour is the 200 m isobath.




Figure 26. The surface (left) and bottom (right) temperature distribution for the fall of 1967. The dashed contour is the 200 m isobath.




Figure 27. The surface (left) and bottom (right) temperature distribution for the fall of 1968. The dashed contour is the 200 m isobath.




Figure 28. The surface (left) and bottom (right) temperature distribution for the fall of 1969. The dashed contour is the 200 m isobath.




Figure 29. The surface (left) and bottom (right) temperature distribution for the fall of 1970. The dashed contour is the 200 m isobath.




Figure 30. The surface (left) and bottom (right) temperature distribution for the fall of 1971. The dashed contour is the 200 m isobath.




Figure 31. The surface (left) and bottom (right) temperature distribution for the fall of 1972. The dashed contour is the 200 m isobath.




Figure 32. The surface (left) and bottom (right) temperature distribution for the fall of 1973. The dashed contour is the 200 m isobath.




Figure 33. The surface (left) and bottom (right) temperature distribution for the fall of 1974. The dashed contour is the 200 m isobath.




Figure 34. The surface (left) and bottom (right) temperature distribution for the fall of 1975. The dashed contour is the 200 m isobath.





Figure 35. The surface (left) and bottom (right) temperature distribution for the fall of 1976. The dashed contour is the 200 m isobath.






Figure 37. The surface (left) and bottom (right) temperature distribution for the fall of 1978. The dashed contour is the 200 m isobath.





Figure 38. The surface (left) and bottom (right) temperature distribution for the fall of 1979. The dashed contour is the 200 m isobath.






Figure 40. The surface (left) and bottom (right) temperature distribution for the fall of 1981. The dashed contour is the 200 m isobath.




Figure 41. The surface (left) and bottom (right) temperature distribution for the fall of 1982. The dashed contour is the 200 m isobath.




Figure 42. The surface (left) and bottom (right) temperature distribution for the fall of 1983. The dashed contour is the 200 m isobath.




Figure 43. The surface (left) and bottom (right) temperature distribution for the fall of 1984. The dashed contour is the 200 m isobath.




Figure 44. The surface (left) and bottom (right) temperature distribution for the fall of 1985. The dashed contour is the 200 m isobath.




Figure 45. The surface (left) and bottom (right) temperature distribution for the fall of 1986. The dashed contour is the 200 m isobath.




Figure 46. The surface (left) and bottom (right) temperature distribution for the fall of 1987. The dashed contour is the 200 m isobath.





Figure 47. Average surface temperature in the four regions of the shelf shown in Figure 1 for the spring (top) and the fall (bottom). The data are listed in Table 2.





Figure 48. Average bottom temperature in the four regions of the continental shelf shown in Figure 1 for the spring (top) and the fall (bottom). The data are listed in Table 2.




Figure 49. Average surface temperature anomaly in the spring for the four regions of the shelf shown in Figure 1. The data are listed in Table 2.




Figure 50. Average bottom temperature anomaly in the spring for the four regions of the shelf shown in Figure 1. The data are listed in Table 2.




Figure 51. Average surface temperature anomaly in the fall for the four regions of the shelf shown in Figure 1. The data are listed in Table 2.




Figure 52. Average bottom temperature anomaly in the fall for the four regions of the shelf shown in Figure 1. The data are listed in Table 2.






Figure 53. Comparison of the average spring and fall bottom temperatures (top) and temperature anomalies (bottom) in the southern Middle Atlantic Bight calculated by Davis (1979) and in this report.



UNITED STATES DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
NATIONAL MARINE FISHERIES SERVICE

Northeast Fisheries Center
Environmental Processes Division
Woods Hole, MA 02543

DATE : March 18, 1991

MEMORANDUM FOR : NEFC Investigators

THROUGH : David Mountain, Chief, FOI *DM*

FROM : Tamara Holzwarth-Davis, Technician, FOI *THD*

SUBJECT : 1988-1990 Addendum to NEFC Reference Document 90-03

NEFC Laboratory Reference Document 90-03 [Surface and Bottom Temperature Distributions from the Northeast Fisheries Center Spring and Fall Bottom Trawl Survey Program, 1963-1987 by Tamara Holzwarth and David Mountain] summarized the temperature data collected by the bottom trawl survey program for the period 1963-1987.

This addendum has been created in an effort to ensure complete and timely distribution of subsequent temperature data taken during the spring and fall bottom trawl survey program.

Enclosed are contoured distributions of the surface and bottom temperatures for the spring and fall surveys of 1988, 1989 and 1990. The areal average temperatures and temperature anomalies for these three years have been calculated according to the methods described in NEFC Ref. Doc. 90-03 and are presented in a table. Time series plots of all the temperature and temperature anomalies for the period 1963-1990 are also included.

Please note, no XBT temperature data was collected on Georges Bank during the 1990 fall survey.

Enclosures



Table 1. Areal average surface and bottom temperature anomaly for the spring and fall bottom trawl surveys in the four regions of the northeast continental shelf shown in Figure 1 NEFC Reference Document 90-03: "#Obs", the number of observations included in each average; "Temp", the areal average temperature; "Anomaly", the areal average temperature anomaly; "SDV1", the standard deviation associated with the average temperature anomaly; "SDV2", the standard deviation of the individual anomalies from which the average anomaly was derived. An asterisk indicates that a true areal average could not be calculated due to poor station coverage and that the average values listed were derived from a simple average of the observations that were within the region. All of the temperature values are in °C.

Spring - Gulf of Maine										
Surface						Bottom				
YEAR	#OBS	TEMP	ANOMALY	SDV1	SDV2	#OBS	TEMP	ANOMALY	SDV1	SDV2
1988	33	4.2	-0.3	0.2	0.9	31	6.0	0.9	0.2	0.9
1989	24*	4.3	-0.1	0.2	0.7	24*	5.5	-0.2	0.2	0.7
1990	24	4.5	0.2	0.3	0.7	24	5.2	0.1	0.3	0.7
Fall - Gulf of Maine										
1988	41	11.2	-0.8	0.2	0.7	41	7.7	-0.4	0.2	1.4
1989	43	11.7	-0.0	0.2	0.8	41	7.4	-0.5	0.2	1.5
1990	26*	13.0	1.1	0.2	0.9	25*	7.5	-0.2	0.2	1.0
Spring - Georges Bank										
1988	22	4.6	-0.0	0.3	0.7	27	4.5	-0.4	0.3	1.0
1989	28	4.5	0.0	0.3	0.6	22	5.0	0.0	0.4	1.1
1990	26	5.4	0.7	0.3	0.7	24	5.4	0.6	0.3	0.8
Fall - Georges Bank										
1988	19	13.9	-1.0	0.3	1.5	17	11.5	-1.0	0.4	1.6
1989	33	14.4	-0.1	0.2	1.1	31	12.0	-0.7	0.3	1.9
1990	0					0				

Table 1. Continued

Spring - Middle Atlantic Bight North

YEAR	#OBS	Surface				Bottom				
		TEMP	ANOMALY	SDV1	SDV2	#OBS	TEMP	ANOMALY	SDV1	SDV2
1988	19	4.6	0.1	0.4	1.0	23	6.8	1.3	0.5	1.4
1989	6*	5.8	1.5	0.7	1.6	4*	5.0	1.1	0.8	1.7
1990	23	6.4	2.0	0.4	0.7	20*	6.1	1.3	0.4	1.7

Fall - Middle Atlantic Bight North

1988	22	18.7	0.7	0.4	1.1	17	11.2	-1.1	0.5	2.0
1989	16	19.4	1.7	0.5	1.5	15*	12.6	0.1	0.4	2.2
1990	17*	19.0	0.5	0.4	0.7	14*	15.3	1.6	0.4	1.5

Spring - Middle Atlantic Bight South

1988	21	6.2	0.3	0.4	1.2	19	7.0	0.8	0.5	1.3
1989	15*	8.1	0.9	0.5	1.3	12*	8.1	1.4	0.6	0.6
1990	29	8.0	2.4	0.3	1.5	24	7.5	1.8	0.5	1.2

Fall - Middle Atlantic Bight South

1988	23	21.2	-0.2	0.4	1.3	21	10.6	-3.0	0.5	3.5
1989	20*	23.0	1.4	0.4	1.1	17*	12.0	-2.1	0.5	3.2
1990	36	22.3	0.8	0.3	1.0	31*	17.6	1.2	0.3	2.7




Figure 1. The surface (left) and bottom (right) temperature distribution for the spring of 1988. The dashed contour is the 200 m isobath.





Figure 2. The surface (left) and bottom (right) temperature distribution for the spring of 1989. The dashed contour is the 200 m isobath.





Figure 3. The surface (left) and bottom (right) temperature distribution for the spring of 1990. The dashed contour is the 200 m isobath.





Figure 4. The surface (left) and bottom (right) temperature distribution for the fall of 1988. The dashed contour is the 200 m isobath.




Figure 5. The surface (left) and bottom (right) temperature distribution for the fall of 1989. The dashed contour is the 200 m isobath.






Figure 6. The surface (left) and bottom (right) temperature distribution for the fall of 1990. The dashed contour is the 200 m isobath.

Spring Surface Temperature



Fall Surface Temperature





Figure 7. Average surface temperature in the four regions of the shelf shown in Figure 1 of NEFC Reference Document 90-03 for the spring (top) and the fall (bottom). Additional data (1988-1990) are listed in Table 1.

Spring Bottom Temperature



Fall Bottom Temperature




Figure 8. Average bottom temperature in the four regions of the shelf shown in Figure 1 of NEFC Reference Document 90-03 for the spring (top) and the fall (bottom). Additional data (1988-1990) are listed in Table 1.







Figure 9. Average surface temperature anomaly in the spring for the four regions of the shelf shown in Figure 1 NEFC Reference Document 90-03. The additional data (1989-1990) are listed in Table 1.







Figure 10. Average bottom temperature anomaly in the spring for the four regions of the shelf shown in Figure 1 NEFC Reference Document 90-03. The additional data (1989-1990) are listed in Table 1.







Figure 11. Average surface temperature anomaly in the fall for the four regions of the shelf shown in Figure 1 NEFC Reference Document 90-03. The additional data (1989-1990) are listed in Table 1.







Figure 12. Average bottom temperature anomaly in the fall for the four regions of the shelf shown in Figure 1 NEFC Reference Document 90-03. The additional data (1989-1990) are listed in Table 1.

Northeast Fisheries Center
Environmental Processes Division
Woods Hole, MA 02543

DATE : March 18, 1991

MEMORANDUM FOR : NEFC Investigators

THROUGH : David Mountain, Chief, FOI

FROM : Tamara Holzwarth-Davis, Technician, FOI

SUBJECT : 1988-1990 Addendum to NEFC Reference Document 90-03

NEFC Laboratory Reference Document 90-03 [Surface and Bottom Temperature Distributions from the Northeast Fisheries Center Spring and Fall Bottom Trawl Survey Program, 1963-1987 by Tamara Holzwarth and David Mountain] summarized the temperature data collected by the bottom trawl survey program for the period 1963-1987.

This addendum has been created in an effort to ensure complete and timely distribution of subsequent temperature data taken during the spring and fall bottom trawl survey program.

Enclosed are contoured distributions of the surface and bottom temperatures for the spring and fall surveys of 1988, 1989 and 1990. The areal average temperatures and temperature anomalies for these three years have been calculated according to the methods described in NEFC Ref. Doc. 90-03 and are presented in a table. Time series plots of all the temperature and temperature anomalies for the period 1963-1990 are also included.

Please note, no XBT temperature data was collected on Georges Bank during the 1990 fall survey.

Enclosures

Table 1. Areal average surface and bottom temperature anomaly for the spring and fall bottom trawl surveys in the four regions of the northeast continental shelf shown in Figure 1 NEFC Reference Document 90-03: "#Obs", the number of observations included in each average; "Temp", the areal average temperature; "Anomaly", the areal average temperature anomaly; "SDV1", the standard deviation associated with the average temperature anomaly; "SDV2", the standard deviation of the individual anomalies from which the average anomaly was derived. An asterisk indicates that a true areal average could not be calculated due to poor station coverage and that the average values listed were derived from a simple average of the observations that were within the region. All of the temperature values are in °C.

Spring - Gulf of Maine										
Surface						Bottom				
YEAR	#OBS	TEMP	ANOMALY	SDV1	SDV2	#OBS	TEMP	ANOMALY	SDV1	SDV2
1988	33	4.2	-0.3	0.2	0.9	31	6.0	0.9	0.2	0.9
1989	24*	4.3	-0.1	0.2	0.7	24*	5.5	-0.2	0.2	0.7
1990	24	4.5	0.2	0.3	0.7	24	5.2	0.1	0.3	0.7
Fall - Gulf of Maine										
1988	41	11.2	-0.8	0.2	0.7	41	7.7	-0.4	0.2	1.4
1989	43	11.7	-0.0	0.2	0.8	41	7.4	-0.5	0.2	1.5
1990	26*	13.0	1.1	0.2	0.9	25*	7.5	-0.2	0.2	1.0
Spring - Georges Bank										
1988	22	4.6	-0.0	0.3	0.7	27	4.5	-0.4	0.3	1.0
1989	28	4.5	0.0	0.3	0.6	22	5.0	0.0	0.4	1.1
1990	26	5.4	0.7	0.3	0.7	24	5.4	0.6	0.3	0.8
Fall - Georges Bank										
1988	19	13.9	-1.0	0.3	1.5	17	11.5	-1.0	0.4	1.6
1989	33	14.4	-0.1	0.2	1.1	31	12.0	-0.7	0.3	1.9
1990	0					0				

Table 1. Continued

Spring - Middle Atlantic Bight North

YEAR	#OBS	Surface				Bottom				
		TEMP	ANOMALY	SDV1	SDV2	#OBS	TEMP	ANOMALY	SDV1	SDV2
1988	19	4.6	0.1	0.4	1.0	23	6.8	1.3	0.5	1.4
1989	6*	5.8	1.5	0.7	1.6	4*	5.0	1.1	0.8	1.7
1990	23	6.4	2.0	0.4	0.7	20*	6.1	1.3	0.4	1.7

Fall - Middle Atlantic Bight North

1988	22	18.7	0.7	0.4	1.1	17	11.2	-1.1	0.5	2.0
1989	16	19.4	1.7	0.5	1.5	15*	12.6	0.1	0.4	2.2
1990	17*	19.0	0.5	0.4	0.7	14*	15.3	1.6	0.4	1.5

Spring - Middle Atlantic Bight South

1988	21	6.2	0.3	0.4	1.2	19	7.0	0.8	0.5	1.3
1989	15*	8.1	0.9	0.5	1.3	12*	8.1	1.4	0.6	0.6
1990	29	8.0	2.4	0.3	1.5	24	7.5	1.8	0.5	1.2

Fall - Middle Atlantic Bight South

1988	23	21.2	-0.2	0.4	1.3	21	10.6	-3.0	0.5	3.5
1989	20*	23.0	1.4	0.4	1.1	17*	12.0	-2.1	0.5	3.2
1990	36	22.3	0.8	0.3	1.0	31*	17.6	1.2	0.3	2.7




Figure 1. The surface (left) and bottom (right) temperature distribution for the spring of 1988. The dashed contour is the 200 m isobath.





Figure 2. The surface (left) and bottom (right) temperature distribution for the spring of 1989. The dashed contour is the 200 m isobath.





Figure 3. The surface (left) and bottom (right) temperature distribution for the spring of 1990. The dashed contour is the 200 m isobath.





Figure 4. The surface (left) and bottom (right) temperature distribution for the fall of 1988. The dashed contour is the 200 m isobath.





Figure 5. The surface (left) and bottom (right) temperature distribution for the fall of 1989. The dashed contour is the 200 m isobath.






Figure 6. The surface (left) and bottom (right) temperature distribution for the fall of 1990. The dashed contour is the 200 m isobath.

Spring Surface Temperature



Fall Surface Temperature





Figure 7. Average surface temperature in the four regions of the shelf shown in Figure 1 of NEFC Reference Document 90-03 for the spring (top) and the fall (bottom). Additional data (1988-1990) are listed in Table 1.

Spring Bottom Temperature



Fall Bottom Temperature




Figure 8. Average bottom temperature in the four regions of the shelf shown in Figure 1 of NEFC Reference Document 90-03 for the spring (top) and the fall (bottom). Additional data (1988-1990) are listed in Table 1.




Figure 9. Average surface temperature anomaly in the spring for the four regions of the shelf shown in Figure 1 NEFC Reference Document 90-03. The additional data (1989-1990) are listed in Table 1.







Figure 10. Average bottom temperature anomaly in the spring for the four regions of the shelf shown in Figure 1 NEFC Reference Document 90-03. The additional data (1989-1990) are listed in Table 1.







Figure 11. Average surface temperature anomaly in the fall for the four regions of the shelf shown in Figure 1 NEFC Reference Document 90-03. The additional data (1989-1990) are listed in Table 1.




Figure 12. Average bottom temperature anomaly in the fall for the four regions of the shelf shown in Figure 1 NEFC Reference Document 90-03. The additional data (1989-1990) are listed in Table 1.