




2013

Fourth National Report on Human Exposure to Environmental Chemicals

Updated Tables, September 2013



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

Table of Contents

General Information

Introduction	1
What's New and Different?	2
Chemicals in the Fourth Report: Updated Tables, September 2013	3

Tobacco Smoke

Cotinine	6
NNAL	8

Disinfection By-Products

Bromodichloromethane	10
Dibromochloromethane	11
Tribromomethane (Bromoform)	12
Trichloromethane (Chloroform)	13

Environmental Phenols

Benzophenone-3	14
Bisphenol A	16
4- <i>tert</i> -Octylphenol	18
Triclosan	20

Fungicides and Metabolites

<i>ortho</i> -Phenylphenol	22
Ethylene thiourea	24
Pentachlorophenol	26
Propylene thiourea	28

Herbicides and Metabolites

2,4-Dichlorophenoxyacetic acid	30
2,4,5-Trichlorophenoxyacetic acid	32

Sulfonyl Urea Herbicides

Urinary Bensulfuron-methyl	34
Urinary Chlorsulfuron	36
Urinary Ethametsulfuron-methyl	38
Urinary Foramsulfuron	40
Urinary Halosulfuron	42
Urinary Mesosulfuron-methyl	44
Urinary Metsulfuron-methyl	46
Urinary Nicosulfuron	48
Urinary Oxasulfuron	50
Urinary Primisulfuron-methyl	52
Urinary Prosulfuron	54
Urinary Rimsulfuron	56
Urinary Sulfometuron-methyl	58
Urinary Sulfosulfuron	60
Urinary Thifensulfuron-methyl	62

Urinary Triasulfuron	64
Urinary Triflusulfuron-methyl	66
Carbamate Pesticide Metabolites	
Carbofuranphenol	68
2-Isopropoxyphenol	70
Organochlorine Pesticides and Metabolites	
Aldrin	72
Dieldrin	74
Endrin	76
Heptachlor epoxide	78
<i>o,p'</i> -DDT	80
2,4,5-Trichlorophenol	82
2,4,6-Trichlorophenol	84
Other Pesticide Metabolites	
2,4-Dichlorophenol	86
2,5-Dichlorophenol	88
Organophosphorus Insecticides: Specific Metabolites	
Acephate	90
Dimethoate	92
Methamidophos	94
Omethoate	96
Organophosphorus Insecticides: Dialkyl Phosphate Metabolites	
Diethylphosphate (DEP)	98
Dimethylphosphate (DMP)	102
Diethylthiophosphate (DETP)	106
Dimethylthiophosphate (DMTP)	110
Diethyldithiophosphate (DEDTP)	114
Dimethyldithiophosphate (DMDTP)	118
Metals and Metalloids	
Antimony	122
Arsenic, Total	126
Arsenic (V) acid	128
Arsenobetaine	130
Arsenocholine	132
Arsenous (III) acid	134
Dimethylarsinic acid	136
Monomethylarsonic acid	138
Trimethylarsine oxide	140
Barium	142
Beryllium	146
Cadmium	150
Cesium	156
Cobalt	160
Lead	164
Mercury, Total	170

Mercury, Inorganic	171
Mercury, Total	172
Molybdenum	174
Platinum	178
Thallium	182
Tungsten	186
Uranium	190

Parabens

Butyl paraben	194
Ethyl paraben	196
Methyl paraben	198
<i>n</i> -Propyl paraben	200

Perchlorate and Other Anions

Nitrate	202
Perchlorate	204
Thiocyanate	206

Perfluorinated Compounds: Surfactants

Perfluorobutane sulfonic acid (PFBuS)	208
Perfluorodecanoic acid (PFDeA)	209
Perfluorododecanoic acid (PFDoA)	210
Perfluoroheptanoic acid (PFHpA)	211
Perfluorohexane sulfonic acid (PFHxS)	212
Perfluorononanoic acid (PFNA)	213
Perfluorooctanoic acid (PFOA)	214
Perfluorooctane sulfonic acid (PFOS)	215
Perfluorooctane sulfonamide (PFOSA)	216
2-(N-Ethyl-perfluorooctane sulfonamido) acetic acid (Et-PFOSA-AcOH)	217
2-(N-Methyl-perfluorooctane sulfonamido) acetic acid (Me-PFOSA-AcOH)	218
Perfluoroundecanoic acid (PFUA)	219

Phthalate Metabolites

Mono-benzyl phthalate (MBzP)	220
Mono-isobutyl phthalate (MiBP)	224
Mono- <i>n</i> -butyl phthalate (MnBP)	226
Mono-cyclohexyl phthalate (MCHP)	230
Mono-ethyl phthalate (MEP)	234
Mono-2-ethylhexyl phthalate (MEHP)	238
Mono-(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP)	242
Mono-(2-ethyl-5-oxohexyl) phthalate (MEOHP)	246
Mono-(2-ethyl-5-carboxypentyl) phthalate (MECPP)	250
Mono-(carboxynonyl) phthalate (MCNP)	252
Mono-isononyl phthalate (MiNP)	254
Mono-(carboxyoctyl) phthalate (MCOP)	258
Mono-methyl phthalate (MMP)	260
Mono-(3-carboxypropyl) phthalate (MCPP)	264
Mono- <i>n</i> -octyl phthalate (MOP)	268

Phytoestrogens and Metabolites

Urinary Daidzein	272
Urinary Enterodiol	276
Urinary Enterolactone	280
Urinary Equol	284
Urinary Genistein	288
Urinary O-Desmethylangolensin	292

Polycyclic Aromatic Hydrocarbon Metabolites

2-Hydroxyfluorene	296
3-Hydroxyfluorene	298
9-Hydroxyfluorene	300
1-Hydroxyphenanthrene	302
2-Hydroxyphenanthrene	304
3-Hydroxyphenanthrene	306
4-Hydroxyphenanthrene	308
1-Hydroxypyrene	310
1-Hydroxynaphthalene (1-Naphthol)	312
2-Hydroxynaphthalene (2-Naphthol)	314

Volatile Organic Compounds (VOCs)

1,1,1-Trichloroethane (Methyl chloroform)	316
1,1,2,2-Tetrachloroethane	317
1,1,2-Trichloroethane	318
1,1-Dichloroethane	319
1,1-Dichloroethene (Vinylidene chloride)	320
1,2-Dibromo-3-chloropropane (DBCP)	321
1,2-Dichlorobenzene	322
1,2-Dichloroethane (Ethylene dichloride)	323
<i>cis</i> -1,2-Dichloroethene	324
<i>trans</i> -1,2-Dichloroethene	325
1,2-Dichloropropane	326
1,3-Dichlorobenzene	327
1,4-Dichlorobenzene (Paradichlorobenzene)	328
2,5-Dimethylfuran	329
Benzene	330
Chlorobenzene	331
Dibromomethane	332
Dichloromethane (Methylene chloride)	333
Ethylbenzene	334
Hexachloroethane	335
Methyl- <i>tert</i> -butyl ether (MTBE)	336
Nitrobenzene	337
Styrene	338
Tetrachloroethene (Perchloroethylene)	339
Tetrachloromethane (Carbon tetrachloride)	340
Toluene	341
Trichloroethene (Trichloroethylene)	342

<i>m-p</i> -Xylene	343
<i>o</i> -Xylene	344
Special Analysis	
Special Analysis of Pooled Samples for Select Chemicals	345
Organochlorine Pesticides and Metabolites (Pooled Samples)	
Oxychlorthane	347
<i>trans</i> -Nonachlor	353
<i>p,p'</i> -DDT	359
<i>p,p'</i> -DDE	365
Hexachlorobenzene	371
<i>beta</i> -Hexachlorocyclohexane	377
<i>gamma</i> -Hexachlorocyclohexane	383
Mirex	389
Polybrominated Diphenyl Ethers and PBB 153 (Pooled Samples)	
2,2',4'-Tribromodiphenyl ether (BDE 17)	395
2,4,4'-Tribromodiphenyl ether (BDE 28)	398
2,2',4,4'-Tetrabromodiphenyl ether (BDE 47)	401
2,3',4,4'-Tetrabromodiphenyl ether (BDE 66)	404
2,2',3,4,4'-Pentabromodiphenyl ether (BDE 85)	407
2,2',4,4',5-Pentabromodiphenyl ether (BDE 99)	410
2,2',4,4',6-Pentabromodiphenyl ether (BDE 100)	413
2,2',4,4',5,6'-Hexabromodiphenyl ether (BDE 154)	418
2,2',3,4,4',5',6'-Heptabromodiphenyl ether (BDE 183)	421
2,2',3,3',4,4',5,5',6,6'-Decabromodiphenyl ether (BDE 209)	424
2,2',4,4',5,5'-Hexabromobiphenyl (PBB 153)	426
Polybrominated Dibenzo-<i>p</i>-dioxins (Pooled Samples)	
1,2,3,4,6,7,8-Heptabromodibenzo- <i>p</i> -dioxin (HpBDD)	429
1,2,3,4,7,8 and 1,2,3,6,7,8-Hexabromodibenzo- <i>p</i> -dioxin (HxBDD)	431
1,2,3,7,8,9-Hexabromodibenzo- <i>p</i> -dioxin (HxBDD)	433
1,2,3,7,8-Pentabromodibenzo- <i>p</i> -dioxin (PeBDD)	435
2,3,7,8-Tetrabromodibenzo- <i>p</i> -dioxin (TBDD)	437
Polybrominated Dibenzofurans (Pooled Samples)	
1,2,3,4,6,7,8-Heptabromodibenzofuran (HpBDF)	439
1,2,3,4,7,8-Hexabromodibenzofuran (HxBDF)	441
1,2,3,7,8-Pentabromodibenzofuran (PeBDF)	443
2,3,4,7,8-Pentabromodibenzofuran (PeBDF)	445
2,3,7,8-Tetrabromodibenzofuran (TBDF)	447
Polychlorinated Dibenzo-<i>p</i>-dioxins (Pooled Samples)	
1,2,3,4,6,7,8-Heptachlorodibenzo- <i>p</i> -dioxin (HpCDD)	449
1,2,3,4,7,8-Hexachlorodibenzo- <i>p</i> -dioxin (HxCDD)	453
1,2,3,6,7,8-Hexachlorodibenzo- <i>p</i> -dioxin (HxCDD)	457
1,2,3,7,8,9-Hexachlorodibenzo- <i>p</i> -dioxin (HxCDD)	461
1,2,3,4,6,7,8,9-Octachlorodibenzo- <i>p</i> -dioxin (OCDD)	465
1,2,3,7,8-Pentachlorodibenzo- <i>p</i> -dioxin (PeCDD)	469
2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin (TCDD)	473

Polychlorinated Dibenzofurans (Pooled Samples)

1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	477
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	481
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	485
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	489
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	493
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	497
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	501
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	505
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	509
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	513

Dioxin-like Polychlorinated Biphenyls: Coplanar PCBs (Pooled Samples)

3,3',4,4'-Tetrachlorobiphenyl (PCB 77)	517
3,4,4',5-Tetrachlorobiphenyl (PCB 81)	519
3,3',4,4',5-Pentachlorobiphenyl (PCB 126)	523
3,3',4,4',5,5'-Hexachlorobiphenyl (PCB 169)	527

Dioxin-like Polychlorinated Biphenyls: mono-ortho-substituted PCBs (Pooled Samples)

2,3,3',4,4'-Pentachlorobiphenyl (PCB 105)	531
2,3,3',4,4'-Pentachlorobiphenyl (PCB 114)	537
2,3',4,4',5-Pentachlorobiphenyl (PCB 118)	541
2',3,4,4',5-Pentachlorobiphenyl (PCB 123)	547
2,3,3',4,4',5-Hexachlorobiphenyl (PCB 156)	551
2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157)	557
2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167)	563
2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189)	569

Non-Dioxin -Like Polychlorinated Biphenyls (Pooled Samples)

2,4,4'-Trichlorobiphenyl (PCB 28)	575
2,2',3,5'-Tetrachloro biphenyl (PCB 44)	581
2,2',4,5'-Tetrachloro biphenyl (PCB 49)	587
2,2',5,5'-Tetrachlorobiphenyl (PCB 52)	593
2,3',4,4'-Tetrachlorobiphenyl (PCB 66)	599
2,4,4',5-Tetrachlorobiphenyl (PCB 74)	605
2,2',3,4,5'-Pentachlorobiphenyl (PCB 87)	611
2,2',4,4',5-Pentachlorobiphenyl (PCB 99)	617
2,2',4,5,5'-Pentachlorobiphenyl (PCB 101)	623
2,3,3',4',6-Pentachlorobiphenyl (PCB 110)	629
2,2',3,3',4,4'-Hexachlorobiphenyl (PCB 128)	635
2,2',3,4,4',5' and 2,3,3',4,4',6-Hexachlorobiphenyl (PCB 138 & 158)	641
2,2',3,4',5,5'-Hexachlorobiphenyl (PCB 146)	647
2,2',3,4',5',6-Hexachlorobiphenyl (PCB 149)	653
2,2',3,5,5',6-Hexachlorobiphenyl (PCB 151)	659
2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153)	665
2,2',3,3',4,4',5-Heptachlorobiphenyl (PCB 170)	671
2,2',3,3',4,4',5'-Heptachlorobiphenyl (PCB 172)	677
2,2',3,3',4,4',5',6'-Heptachlorobiphenyl (PCB 177)	683

2,2',3,3',5,5',6-Heptachlorobiphenyl (PCB 178)	689
2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180)	695
2,2',3,4,4',5,6-Heptachlorobiphenyl (PCB 183)	701
2,2',3,4',5,5',6-Heptachlorobiphenyl (PCB 187)	707
2,2',3,3',4,4',5,5'-Octachlorobiphenyl (PCB 194)	713
2,2',3,3',4,4',5,6-Octachlorobiphenyl (PCB 195)	719
2,2',3,3',4,4',5,6' and 2,2',3,4,4',5,5',6-Octachlorobiphenyl (PCB 196 & 203)	725
2,2',3,3',4,5,5',6-Octachlorobiphenyl (PCB 199)	731
2,2',3,3',4,4',5,5',6-Nonachlorobiphenyl (PCB 206)	737
2,2',3,3',4,4',5,5',6,6'-Decachlorobiphenyl (PCB 209)	743

General Information

Appendix C: References for Biomonitoring Analytical Methods	749
Appendix D. Limit of Detection Table	755

Fourth National Report on Human Exposure to Environmental Chemicals, Updated Tables, September 2013

The *Fourth National Report on Human Exposure to Environmental Chemicals, Updated Tables, September 2013* provides nationally representative biomonitoring data that has become available since the publication of the *Fourth National Report on Human Exposure to Environmental Chemicals, 2009*. The *Updated Tables, September 2013* includes all the updates previously provided in *Updated Tables, July 2010* through *Updated Tables, March 2013*. The *Updated Tables, September 2013* present data for 91 chemicals measured in serum pooled samples that have not been reported in any previous *Updated Tables*. Since publication of the *Fourth Report, 2009*, 201 chemicals have updated tables and 49 chemicals have been added, for a total of 250 chemicals presented in these *Updated Tables*.

A link to information about specific chemicals or the chemical group is provided at the bottom of the data table page, and it is referred to as a Biomonitoring Summary. For many chemicals, there is also a link provided to a brief overview about the chemical, referred to as a Fact Sheet. Two appendices are updated to include information pertinent to these chemicals, as well as the other chemicals included in the *Updated Tables* (Appendix C. References for Biomonitoring Methods; Appendix D. Limit of Detection Table).

The chemicals included in this update are listed in the table that follows.

The *Report* website is the best resource for the most recent updates of available data (<http://www.cdc.gov/exposurereport>). The *Report* website also contains details of the Data Sources and Data Analysis, Interpretation of *Report* Data, and Chemical and Toxicological Information that pertain to the *Updated Tables*.

What's New and Different?

The *Updated Tables* in this release include several chemicals that have results available from one or more of the three survey periods of 2005-2006, 2007-2008, and 2009-2010. New chemicals measured for the first time include one serum polybrominated diphenyl ether (PBDE), three serum polychlorinated biphenyls (PCBs), five polybrominated dioxins, and five polybrominated furans. Chemicals with updated data in this release are the serum dioxin-like and non-dioxin-like chemicals, serum PBDEs and PBB 153, serum organochlorine pesticides, urinary dialkylphosphate metabolites of organophosphorus pesticides, and urinary phytoestrogens and metabolites.

Beginning with the 2005-2006 survey period, measurement of a number of chemicals was made on pooled serum samples (see [Special Analysis of Pooled Samples for Select Chemicals](#)). Many of these chemicals were measured in individual samples prior to the 2005-2006 survey period: non-dioxin-like PCBs; dioxins, furans, and dioxin-like PCBs; PDBEs and PBB 153; and organochlorine pesticides. Serum polybrominated dioxins and furans were measured in pooled samples for the first time in the 2005-2006 survey period. Several organochlorine pesticides (e.g., aldrin, dieldrin, *o,p'* DDT, endrin, and heptachlor epoxide) were not measured in the pooled samples because they were largely undetected in previous NHANES survey periods.

NHANES made changes in oversampling of population subgroups for the 2007-2010 survey periods. Beginning in 2007, adolescents were no longer oversampled, and all Hispanics were oversampled, rather than just Mexican American persons. As a result of these changes, the numbers of participants in subsamples, particularly 12-19 year olds, may be smaller than in survey periods prior to NHANES 2007-2008. More information about in the NHANES 2007-2010 survey design are available from http://www.cdc.gov/nchs/data/nhanes/analyticnote_2007-2010.pdf.

CDC has learned that the analytical standard for Perfluorononanoic acid (PFNA) used for all CDC projects since the mid 2000's had a lower purity than previously described. As a result, a correction factor equal to 0.82 has been applied to PFNA for the NHANES periods 2007-2008, and 2009-2010. Consequently, the data tables in these *Updated Tables* for PFNA show results that are slightly different from the data tables presented in the *Fourth National Report on Human Exposure to Environmental Chemicals, 2009, Updated Tables, February 2013*, and in all of the *Updated Tables* since February 2011.

For the information in this section, "What's New and Different," contained in previous releases of *Updated Tables*, please see the Archives from *Updated Tables* at: <http://www.cdc.gov/exposurereport/>.

Chemicals in the *Fourth Report: Updated Tables, September 2013*

CDC's *Fourth National Report on Human Exposure to Environmental Chemicals: Updated Tables* provides exposure data on the following chemicals or classes of chemicals. The Updated Tables contain cumulative data from national samples collected beginning in 1999–2000 and as recently as 2009–2010. Not all chemicals were measured in each national sample. The data tables are available at <http://www.cdc.gov/exposurereport>. An asterisk (*) indicates the chemical has been added since publication of the *Fourth Report* in 2009.

Tobacco Smoke

Cotinine
NNAL*

Disinfection By Products

Bromodichloromethane
Dibromochloromethane
Tribromomethane (Bromoform)
Trichloromethane (Chloroform)

Environmental Phenols

Benzophenone-3
Bisphenol A
4-*tert*-Octylphenol
Triclosan

Fungicides and Metabolites

ortho-Phenylphenol
Ethylene thiourea*
Pentachlorophenol
Propylene thiourea*

Herbicides and Metabolites

2,4-Dichlorophenoxyacetic acid
2,4,5-Trichlorophenoxyacetic acid

Sulfonyl Urea Herbicides

Urinary Bensulfuron-methyl*
Urinary Chlorsulfuron*
Urinary Ethametsulfuron-methyl*
Urinary Foramsulfuron*
Urinary Halosulfuron*
Urinary Mesosulfuron-methyl*
Urinary Metsulfuron-methyl*
Urinary Nicosulfuron*
Urinary Oxasulfuron*
Urinary Primisulfuron-methyl*
Urinary Prosulfuron*
Urinary Rimsulfuron*
Urinary Sulfometuron-methyl*
Urinary Sulfosulfuron*
Urinary Thifensulfuron-methyl*
Urinary Triasulfuron*
Urinary Triflusulfuron-methyl*

Carbamate Pesticide Metabolites

Carbofuranphenol
2-Isopropoxyphenol

Organochlorine Pesticides and Metabolites

Aldrin
Dieldrin
Endrin
Heptachlor epoxide
o,p'-DDT
2,4,5-Trichlorophenol
2,4,6-Trichlorophenol

Other Pesticide Metabolites

2,4-Dichlorophenol*
2,5-Dichlorophenol*

Organophosphorus Insecticides: Specific Metabolites

Acephate*
Dimethoate*
Methamidophos*
Omethoate*

Organophosphorus Insecticides: Dialkyl Phosphate Metabolites

Diethylphosphate (DEP)
Dimethylphosphate (DMP)
Diethylthiophosphate (DETP)
Dimethylthiophosphate (DMTP)
Diethyldithiophosphate (DEDTP)
Dimethyldithiophosphate (DMDTP)

Metals and Metalloids

Antimony
Arsenic, Total
Arsenic (V) acid
Arsenobetaine
Arsenocholine
Arsenous (III) acid
Dimethylarsinic acid
Monomethylarsonic acid
Trimethylarsine oxide
Barium
Beryllium
Cadmium
Cesium
Cobalt
Lead
Mercury, Total
Mercury, Inorganic
Molybdenum
Platinum
Thallium
Tungsten
Uranium

Parabens

Butyl paraben*
Ethyl paraben*
Methyl paraben*
n-Propyl paraben*

Perchlorate and Other Anions

Nitrate*
Perchlorate
Thiocyanate*

Perfluorinated Compounds: Surfactants

Perfluorobutane sulfonic acid (PFBuS)
Perfluorodecanoic acid (PFDeA)
Perfluorododecanoic acid (PFDoA)
Perfluoroheptanoic acid (PFHpA)
Perfluorohexane sulfonic acid (PFHxS)
Perfluorononanoic acid (PFNA)
Perfluorooctanoic acid (PFOA)
Perfluorooctane sulfonic acid (PFOS)
Perfluorooctane sulfonamide (PFOSA)
2-(*N*-Ethyl-perfluorooctane sulfonamido) acetic acid (Et-PFOSA-AcOH)
2-(*N*-Methyl-perfluorooctane sulfonamido) acetic acid (Me-PFOSA-AcOH)
Perfluoroundecanoic acid (PFUA)

Chemicals in the *Fourth Report: Updated Tables, September 2013*

CDC's *Fourth National Report on Human Exposure to Environmental Chemicals: Updated Tables* provides exposure data on the following chemicals or classes of chemicals. The Updated Tables contain cumulative data from national samples collected beginning in 1999–2000 and as recently as 2009–2010. Not all chemicals were measured in each national sample. The data tables are available at <http://www.cdc.gov/exposurereport>. An asterisk (*) indicates the chemical has been added since publication of the *Fourth Report* in 2009.

Phthalate Metabolites

Mono-benzyl phthalate (MBzP)
 Mono-isobutyl phthalate (MiBP)
 Mono-n-butyl phthalate (MnBP)
 Mono-cyclohexyl phthalate (MCHP)
 Mono-ethyl phthalate (MEP)
 Mono-2-ethylhexyl phthalate (MEHP)
 Mono-(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP)
 Mono-(2-ethyl-5-oxohexyl) phthalate (MEOHP)
 Mono-(2-ethyl-5-carboxypentyl) phthalate (MECPP)
 Mono-(carboxynonyl) phthalate (MCNP)*
 Mono-isononyl phthalate (MiNP)
 Mono-(carboxyoctyl) phthalate (MCOP)*
 Mono-methyl phthalate (MMP)
 Mono-(3-carboxypropyl) phthalate (MCPP)
 Mono-n-octyl phthalate (MOP)

Phytoestrogens and Metabolites

Urinary Daidzein
 Urinary Enterodiol
 Urinary Enterolactone
 Urinary Equol
 Urinary Genistein
 Urinary O-Desmethylangolensin

Polycyclic Aromatic Hydrocarbon Metabolites

2-Hydroxyfluorene
 3-Hydroxyfluorene
 9-Hydroxyfluorene
 1-Hydroxyphenanthrene
 2-Hydroxyphenanthrene
 3-Hydroxyphenanthrene
 4-Hydroxyphenanthrene
 1-Hydroxypyrene
 1-Hydroxynaphthalene (1-Naphthol)
 2-Hydroxynaphthalene (2-Naphthol)

Volatile Organic Compounds

1,1,1-Trichloroethane (Methyl chloroform)
 1,1,2,2-Tetrachloroethane
 1,1,2-Trichloroethane
 1,1-Dichloroethane
 1,1-Dichloroethene (Vinylidene chloride)
 1,2-Dibromo-3-chloropropane (DBCP)
 1,2-Dichlorobenzene
 1,2-Dichloroethane (Ethylene dichloride)
cis-1,2-Dichloroethene
trans-1,2-Dichloroethene
 1,2-Dichloropropane
 1,3-Dichlorobenzene
 1,4-Dichlorobenzene (Paradichlorobenzene)
 2,5-Dimethylfuran
 Benzene
 Chlorobenzene
 Dibromomethane
 Dichloromethane (Methylene chloride)
 Ethylbenzene
 Hexachloroethane
 Methyl-*tert*-butyl ether (MTBE)
 Nitrobenzene
 Styrene
 Tetrachloroethene (Perchloroethylene)
 Tetrachloromethane (Carbon tetrachloride)
 Toluene
 Trichloroethene (Trichloroethylene)
m-*p*-Xylene
o-Xylene

Organochlorine Pesticides and Metabolites (Pooled Samples)

Oxychlorane
trans-Nonachlor
p,p'-DDT
p,p'-DDE
 Hexachlorobenzene
beta-Hexachlorocyclohexane
gamma-Hexachlorocyclohexane
 Mirex

Polybrominated Diphenyl Ethers and PBB 153 (Pooled Samples)

2,2',4'-Tribromodiphenyl ether (BDE 17)
 2,4,4'-Tribromodiphenyl ether (BDE 28)
 2,2',4,4'-Tetrabromodiphenyl ether (BDE 47)
 2,3',4,4'-Tetrabromodiphenyl ether (BDE 66)
 2,2',3,4,4'-Pentabromodiphenyl ether (BDE 85)
 2,2',4,4',5-Pentabromodiphenyl ether (BDE 99)
 2,2',4,4',6-Pentabromodiphenyl ether (BDE 100)
 2,2',4,4',5,5'-Hexabromodiphenyl ether (BDE 153)
 2,2',4,4',5,6'-Hexabromodiphenyl ether (BDE 154)
 2,2',3,4,4',5',6-Heptabromodiphenyl ether (BDE 183)
 2,2',3,3',4,4',5,5',6,6'-Decabromodiphenyl ether (BDE 209)*
 2,2',4,4',5,5'-Hexabromobiphenyl (PBB 153)

Polybrominated Dibenzo-*p*-dioxins (Pooled Samples)

1,2,3,4,6,7,8-Heptabromodibenzo-*p*-dioxin (HpBDD)*
 1,2,3,4,7,8 and 1,2,3,6,7,8-Hexabromodibenzo-*p*-dioxin (HxBDD)*
 1,2,3,7,8,9-Hexabromodibenzo-*p*-dioxin (HxBDD)*
 1,2,3,7,8-Pentabromodibenzo-*p*-dioxin (PeBDD)*
 2,3,7,8-Tetrabromodibenzo-*p*-dioxin (TBDD)*

Polybrominated Dibenzofurans (Pooled Samples)

1,2,3,4,6,7,8-Heptabromodibenzofuran (HpBDF)*
 1,2,3,4,7,8-Hexabromodibenzofuran (HxBDF)*
 1,2,3,7,8-Pentabromodibenzofuran (PeBDF)*
 2,3,4,7,8-Pentabromodibenzofuran (PeBDF)*
 2,3,7,8-Tetrabromodibenzofuran (TBDF)*

Polychlorinated Dibenzo-*p*-dioxins (Pooled Samples)

1,2,3,4,6,7,8-Heptachlorodibenzo-*p*-dioxin (HpCDD)
 1,2,3,4,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD)
 1,2,3,6,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD)
 1,2,3,7,8,9-Hexachlorodibenzo-*p*-dioxin (HxCDD)
 1,2,3,4,6,7,8,9-Octachlorodibenzo-*p*-dioxin (OCDD)
 1,2,3,7,8-Pentachlorodibenzo-*p*-dioxin (PeCDD)
 2,3,7,8-Tetrachlorodibenzo-*p*-dioxin (TCDD)

Polychlorinated Dibenzofurans (Pooled Samples)

1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)
 1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)
 1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)
 1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)
 1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)
 2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)
 1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)
 1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)
 2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)
 2,3,7,8-Tetrachlorodibenzofuran (TCDF)

Chemicals in the *Fourth Report: Updated Tables, September 2013*

CDC's *Fourth National Report on Human Exposure to Environmental Chemicals: Updated Tables* provides exposure data on the following chemicals or classes of chemicals. The Updated Tables contain cumulative data from national samples collected beginning in 1999–2000 and as recently as 2009–2010. Not all chemicals were measured in each national sample. The data tables are available at <http://www.cdc.gov/exposurereport>. An asterisk (*) indicates the chemical has been added since publication of the *Fourth Report* in 2009.

Dioxin like Polychlorinated Biphenyls: Coplanar PCBs (Pooled Samples)

3,3',4,4'-Tetrachlorobiphenyl (PCB 77)*
3,4,4',5-Tetrachlorobiphenyl (PCB 81)
3,3',4,4',5-Pentachlorobiphenyl (PCB 126)
3,3',4,4',5,5'-Hexachlorobiphenyl (PCB 169)

Dioxin like Polychlorinated Biphenyls: Mono-ortho-substituted PCBs (Pooled Samples)

2,3,3',4,4'-Pentachlorobiphenyl (PCB 105)
2,3,3',4,4'-Pentachlorobiphenyl (PCB 114)*
2,3',4,4',5-Pentachlorobiphenyl (PCB 118)
2',3,4,4',5-Pentachlorobiphenyl (PCB 123)*
2,3,3',4,4',5-Hexachlorobiphenyl (PCB 156)
2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157)
2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167)
2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189)

Non-Dioxin Like Polychlorinated Biphenyls (Pooled Samples)

2,4,4'-Trichlorobiphenyl (PCB 28)
2,2',3,5'-Tetrachloro biphenyl (PCB 44)
2,2',4,5'-Tetrachloro biphenyl (PCB 49)
2,2',5,5'-Tetrachlorobiphenyl (PCB 52)
2,3',4,4'-Tetrachlorobiphenyl (PCB 66)
2,4,4',5-Tetrachlorobiphenyl (PCB 74)
2,2',3,4,5'-Pentachlorobiphenyl (PCB 87)
2,2',4,4',5-Pentachlorobiphenyl (PCB 99)
2,2',4,5,5'-Pentachlorobiphenyl (PCB 101)
2,3,3',4',6-Pentachlorobiphenyl (PCB 110)
2,2',3,3',4,4'-Hexachlorobiphenyl (PCB 128)
2,2',3,4,4',5' and 2,3,3',4,4',6-Hexachlorobiphenyl (PCB 138 & 158)
2,2',3,4',5,5'-Hexachlorobiphenyl (PCB 146)
2,2',3,4',5',6-Hexachlorobiphenyl (PCB 149)
2,2',3,5,5',6-Hexachlorobiphenyl (PCB 151)
2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153)
2,2',3,3',4,4',5-Heptachlorobiphenyl (PCB 170)
2,2',3,3',4,5,5'-Heptachlorobiphenyl (PCB 172)
2,2',3,3',4,5',6'-Heptachlorobiphenyl (PCB 177)
2,2',3,3',5,5',6-Heptachlorobiphenyl (PCB 178)
2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180)
2,2',3,4,4',5',6-Heptachlorobiphenyl (PCB 183)
2,2',3,4',5,5',6-Heptachlorobiphenyl (PCB 187)
2,2',3,3',4,4',5,5'-Octachlorobiphenyl (PCB 194)
2,2',3,3',4,4',5,6-Octachlorobiphenyl (PCB 195)
2,2',3,3',4,4',5,6' and 2,2',3,4,4',5,5',6-Octachlorobiphenyl (PCB 196 & 203)
2,2',3,3',4,5,5',6-Octachlorobiphenyl (PCB 199)
2,2',3,3',4,4',5,5',6-Nonachlorobiphenyl (PCB 206)
2,2',3,3',4,4',5,5',6,6'-Decachlorobiphenyl (PCB 209)

Serum Cotinine

Metabolite of nicotine (component of tobacco smoke)

Geometric mean and selected percentiles of serum concentrations (in ng/mL) for the ***non-smoking U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	.060 (<LOD-.080)	.240 (.190-.302)	1.02 (.770-1.28)	1.96 (1.60-2.62)	5999
	01-02**	.062 (.050-.077)	< LOD	.160 (.120-.220)	.930 (.740-1.17)	2.20 (1.83-2.44)	6819
	03-04	.071 (.057-.089)	.050 (.040-.070)	.210 (.140-.310)	.990 (.740-1.30)	2.17 (1.81-2.54)	6320
	05-06	.054 (.047-.061)	.040 (.030-.040)	.120 (.100-.150)	.630 (.460-.870)	1.47 (1.15-1.92)	6347
	07-08	.057 (.048-.068)	.040 (.030-.040)	.130 (.100-.180)	.760 (.550-1.09)	1.81 (1.45-2.43)	6197
	09-10	.041 (.037-.046)	.030 (.020-.030)	.070 (.070-.090)	.450 (.350-.580)	1.29 (1.04-1.61)	6678
Age group							
3-11 years	99-00	.164 (.115-.234)	.110 (.066-.188)	.500 (.260-1.16)	1.88 (.997-3.44)	3.44 (1.42-4.79)	1174
	01-02**	.110 (.076-.160)	.070 (<LOD-.130)	.570 (.310-1.00)	2.23 (1.63-2.78)	3.23 (2.53-4.01)	1415
	03-04	.137 (.088-.213)	.120 (.060-.220)	.620 (.310-1.20)	2.04 (1.38-2.94)	3.35 (2.12-4.68)	1252
	05-06	.078 (.062-.097)	.050 (.040-.070)	.220 (.160-.350)	1.22 (.880-1.82)	2.42 (1.63-3.46)	1296
	07-08	.095 (.067-.136)	.060 (.040-.100)	.380 (.170-.840)	1.67 (1.10-2.54)	2.81 (2.26-3.54)	1337
	09-10	.060 (.047-.075)	.040 (.030-.050)	.160 (.110-.250)	.920 (.560-1.58)	2.15 (1.35-3.00)	1355
12-19 years	99-00	.163 (.142-.187)	.110 (.080-.163)	.540 (.428-.660)	1.66 (1.50-1.95)	2.62 (2.09-3.39)	1773
	01-02**	.086 (.059-.126)	.050 (<LOD-.110)	.350 (.190-.580)	1.53 (1.09-2.12)	3.12 (2.47-3.99)	1902
	03-04	.110 (.087-.139)	.080 (.060-.120)	.510 (.350-.670)	1.55 (1.21-1.93)	2.68 (1.96-4.02)	1783
	05-06	.074 (.060-.092)	.050 (.040-.060)	.230 (.150-.350)	1.16 (.860-1.69)	2.26 (1.69-2.72)	1714
	07-08	.081 (.061-.106)	.050 (.030-.070)	.350 (.170-.550)	1.25 (.930-1.99)	2.54 (2.04-2.94)	934
	09-10	.056 (.044-.072)	.030 (.020-.040)	.130 (.080-.230)	.980 (.510-1.80)	2.49 (1.31-3.65)	1042
20 years and older	99-00	*	.050 (<LOD-.061)	.167 (.140-.193)	.630 (.533-.820)	1.50 (1.28-1.66)	3052
	01-02**	.052 (<LOD-.063)	< LOD	.110 (.090-.150)	.630 (.470-.790)	1.42 (1.14-1.89)	3502
	03-04	.058 (.047-.071)	.040 (.030-.050)	.140 (.100-.200)	.630 (.480-.840)	1.54 (1.26-1.92)	3285
	05-06	.047 (.042-.053)	.030 (.030-.040)	.100 (.080-.120)	.440 (.310-.620)	1.14 (.870-1.41)	3337
	07-08	.049 (.043-.057)	.030 (.030-.040)	.100 (.080-.120)	.490 (.390-.640)	1.37 (.970-1.70)	3926
	09-10	.037 (.034-.040)	.020 (.020-.030)	.070 (.060-.070)	.310 (.240-.390)	.990 (.700-1.24)	4281
Gender							
Males	99-00	.124 (.106-.145)	.080 (.060-.110)	.308 (.220-.410)	1.20 (.950-1.49)	2.39 (1.66-3.22)	2789
	01-02**	.075 (.059-.094)	.050 (<LOD-.070)	.230 (.160-.320)	1.17 (.960-1.49)	2.44 (2.23-2.99)	3152
	03-04	.087 (.070-.108)	.060 (.040-.080)	.280 (.190-.360)	1.23 (.910-1.68)	2.63 (2.09-3.19)	2937
	05-06	.064 (.055-.074)	.040 (.040-.050)	.150 (.120-.190)	.720 (.550-1.07)	1.85 (1.42-2.24)	2922
	07-08	.068 (.056-.081)	.040 (.040-.050)	.170 (.120-.240)	1.07 (.690-1.39)	2.52 (1.84-2.92)	2948
	09-10	.046 (.042-.051)	.030 (.030-.030)	.080 (.070-.100)	.550 (.410-.700)	1.60 (1.19-2.38)	3181
Females	99-00	*	< LOD	.180 (.148-.230)	.850 (.600-1.14)	1.85 (1.33-2.45)	3210
	01-02**	.053 (<LOD-.066)	< LOD	.120 (.090-.180)	.710 (.540-.990)	1.77 (1.32-2.20)	3667
	03-04	.060 (.047-.077)	.040 (.030-.060)	.160 (.110-.260)	.860 (.580-1.15)	1.76 (1.32-2.22)	3383
	05-06	.047 (.040-.054)	.030 (.030-.040)	.100 (.070-.130)	.510 (.300-.830)	1.23 (1.04-1.52)	3425
	07-08	.050 (.042-.059)	.030 (.030-.040)	.110 (.080-.150)	.630 (.400-.900)	1.40 (1.09-1.87)	3249
	09-10	.037 (.033-.042)	.020 (.020-.030)	.070 (.050-.080)	.380 (.270-.570)	1.03 (.720-1.61)	3497

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.05, 0.05, 0.015, 0.015, 0.015, and 0.015 respectively. .

** In the 2001-2002 survey period, 83% of measurements had an LOD of 0.015 ng/mL, and 17% had an LOD of 0.05 ng/mL.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

***Non-smoking is defined as a serum cotinine concentration of 10 ng/mL or less.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cotinine_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Cotinine_FactSheet.html

Serum Cotinine

Metabolite of nicotine (component of tobacco smoke)

Geometric mean and selected percentiles of serum concentrations (in ng/mL) for the ***non-smoking U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	.140 (.110-.180)	.506 (.370-.726)	1.21 (.900-1.70)	2241
	01-02**	.060 (<LOD-.084)	< LOD	.160 (.080-.310)	.730 (.480-1.19)	2.12 (1.19-2.96)	1878
	03-04	.054 (.043-.068)	.030 (.020-.050)	.120 (.080-.180)	.690 (.430-1.00)	2.65 (1.87-3.57)	1707
	05-06	.047 (.038-.059)	.030 (.020-.040)	.100 (.070-.150)	.500 (.370-.730)	1.54 (.830-2.49)	1807
	07-08	.044 (.033-.057)	.030 (.020-.040)	.080 (.060-.120)	.480 (.290-.880)	1.59 (1.00-3.07)	1412
	09-10	.038 (.032-.045)	.020 (.020-.030)	.060 (.050-.080)	.360 (.230-.640)	1.35 (.710-2.25)	1615
Non-Hispanic blacks	99-00	.175 (.153-.201)	.131 (.111-.150)	.505 (.400-.625)	1.43 (1.21-1.75)	2.34 (1.84-3.50)	1333
	01-02**	.164 (.137-.197)	.130 (.110-.160)	.580 (.450-.770)	1.77 (1.55-2.05)	3.15 (2.50-4.30)	1602
	03-04	.144 (.104-.198)	.120 (.080-.180)	.520 (.350-.770)	1.54 (1.20-2.14)	2.77 (2.18-3.54)	1704
	05-06	.114 (.085-.153)	.080 (.060-.120)	.440 (.240-.690)	1.42 (.900-2.03)	2.45 (1.70-3.70)	1630
	07-08	.094 (.079-.112)	.060 (.050-.090)	.320 (.250-.390)	1.19 (.980-1.46)	2.37 (1.75-2.88)	1244
	09-10	.095 (.074-.122)	.070 (.050-.090)	.320 (.210-.540)	1.33 (.900-2.05)	2.85 (1.97-3.60)	1129
Non-Hispanic whites	99-00	*	.050 (<LOD-.073)	.216 (.154-.312)	.950 (.621-1.40)	1.92 (1.48-3.02)	1950
	01-02**	.052 (<LOD-.068)	< LOD	.120 (.090-.180)	.800 (.570-1.11)	1.88 (1.48-2.30)	2847
	03-04	.066 (.050-.087)	.040 (.030-.070)	.180 (.120-.300)	.920 (.620-1.32)	2.01 (1.70-2.49)	2500
	05-06	.049 (.042-.056)	.030 (.030-.040)	.100 (.080-.130)	.530 (.330-.800)	1.27 (.980-1.62)	2404
	07-08	.056 (.043-.073)	.040 (.030-.050)	.130 (.080-.240)	.750 (.440-1.27)	1.70 (1.24-2.54)	2485
	09-10	.037 (.033-.043)	.020 (.020-.030)	.070 (.050-.080)	.350 (.240-.560)	1.06 (.780-1.60)	2743

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.05, 0.05, 0.015, 0.015, 0.015, and 0.015 respectively.

** In the 2001-2002 survey period, 83% of measurements had an LOD of 0.015 ng/mL, and 17% had an LOD of 0.05 ng/mL.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result. .

***Non-smoking is defined as a serum cotinine concentration of 10 ng/mL or less

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cotinine_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Cotinine_FactSheet.html

Urinary NNAL (4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol)

Metabolite of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK)

Geometric mean and selected percentiles of urine concentrations (in pg/mL) for the ****non-smoking** U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	07-08	*	< LOD	2.80 (2.20-3.70)	11.1 (8.80-14.4)	24.5 (18.0-30.4)	5212
	09-10	*	< LOD	1.90 (1.60-2.40)	7.30 (6.40-8.50)	16.7 (13.9-19.6)	6067
Age group							
6-11 years	07-08	*	1.30 (.600-2.60)	8.50 (4.70-13.8)	31.4 (17.5-49.8)	60.6 (37.1-75.7)	875
	09-10	*	.600 (<LOD-1.00)	3.10 (2.20-5.00)	15.5 (10.4-22.8)	29.3 (18.1-59.4)	937
12-19 years	07-08	*	1.20 (.600-2.10)	5.00 (3.80-9.20)	20.0 (11.9-27.5)	39.1 (20.6-60.9)	843
	09-10	*	.900 (<LOD-1.50)	3.70 (2.70-5.20)	12.5 (6.90-25.5)	29.1 (13.9-51.4)	1004
20 years and older	07-08	*	< LOD	2.10 (1.70-2.50)	8.20 (6.90-9.80)	16.6 (13.9-21.5)	3494
	09-10	*	< LOD	1.60 (1.30-1.90)	6.10 (5.10-6.90)	13.1 (10.7-15.8)	4126
Gender							
Males	07-08	*	< LOD	3.80 (3.00-4.50)	12.6 (10.6-15.0)	27.5 (18.9-38.1)	2463
	09-10	*	< LOD	2.40 (1.90-3.00)	8.30 (7.20-9.90)	19.3 (15.2-22.5)	2903
Females	07-08	*	< LOD	2.10 (1.50-2.90)	9.70 (7.10-13.6)	21.6 (16.6-28.0)	2749
	09-10	*	< LOD	1.60 (1.30-1.90)	6.50 (4.70-8.80)	14.0 (10.7-19.0)	3164
Race/ethnicity							
Mexican Americans	07-08	*	< LOD	2.00 (1.20-3.10)	6.90 (4.90-9.00)	13.9 (10.8-17.3)	1206
	09-10	*	< LOD	1.60 (1.20-2.20)	4.90 (3.70-7.00)	11.1 (5.70-18.4)	1476
Non-Hispanic blacks	07-08	*	.800 (<LOD-1.90)	4.90 (3.60-6.30)	14.0 (11.6-17.2)	26.1 (18.5-33.2)	1057
	09-10	1.81 (1.30-2.54)	1.40 (<LOD-2.40)	5.90 (3.80-8.10)	18.9 (11.7-26.1)	34.0 (18.9-61.9)	1004
Non-Hispanic whites	07-08	*	< LOD	2.80 (1.90-4.10)	11.9 (8.20-17.7)	27.5 (19.6-35.4)	2047
	09-10	*	< LOD	1.80 (1.40-2.40)	6.80 (5.40-8.40)	15.2 (11.9-19.3)	2514

Limit of detection (LOD, see Data Analysis section) for Survey years 07-08 and 09-10 are 0.6 and 0.6 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**Non-smoking is defined as a serum cotinine concentration of 10 ng/mL or less.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NNAL_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/NNAL_FactSheet.html

NNAL (4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol) (creatinine corrected)

Metabolite of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK)

Geometric mean and selected percentiles of urine concentrations (in pg/mg of creatinine) for the ****non-smoking** U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	07-08	*	< LOD	2.79 (2.31-3.50)	10.3 (7.36-14.0)	20.5 (16.4-27.5)	5210
	09-10	*	< LOD	2.05 (1.88-2.28)	6.40 (5.71-7.41)	14.6 (11.5-17.3)	6066
Age group							
6-11 years	07-08	*	2.15 (1.38-3.22)	10.7 (5.95-17.9)	38.3 (21.2-57.1)	60.9 (46.4-72.1)	875
	09-10	*	1.19 (<LOD-1.52)	4.25 (2.88-6.37)	19.2 (11.0-28.8)	37.2 (22.9-53.0)	936
12-19 years	07-08	*	1.14 (.710-1.75)	4.40 (2.59-7.36)	14.2 (10.5-23.1)	29.2 (15.6-41.9)	841
	09-10	*	1.05 (<LOD-1.26)	3.08 (1.96-4.37)	10.0 (6.02-15.2)	20.7 (12.3-30.2)	1004
20 years and older	07-08	*	< LOD	2.34 (2.00-2.73)	6.82 (5.49-8.65)	15.1 (11.8-17.7)	3494
	09-10	*	< LOD	1.84 (1.71-2.00)	5.21 (4.57-6.07)	11.4 (8.42-14.1)	4126
Gender							
Males	07-08	*	< LOD	2.98 (2.39-3.71)	10.5 (7.57-12.9)	20.9 (15.7-29.0)	2462
	09-10	*	< LOD	2.00 (1.86-2.25)	6.61 (6.00-7.83)	15.1 (11.7-19.8)	2902
Females	07-08	*	< LOD	2.67 (2.14-3.37)	9.87 (6.79-14.9)	20.0 (15.7-27.9)	2748
	09-10	*	< LOD	2.09 (1.82-2.36)	6.15 (4.88-8.21)	12.7 (9.26-17.3)	3164
Race/ethnicity							
Mexican Americans	07-08	*	< LOD	2.16 (1.55-3.00)	5.91 (4.32-8.30)	13.3 (8.36-17.9)	1205
	09-10	*	< LOD	1.73 (1.36-2.11)	5.00 (3.72-6.84)	10.0 (6.79-14.7)	1476
Non-Hispanic blacks	07-08	*	.960 (<LOD-1.42)	3.70 (2.50-5.31)	11.5 (9.18-14.2)	20.8 (14.5-28.8)	1056
	09-10	1.40 (.982-2.01)	1.21 (<LOD-2.06)	3.96 (2.78-5.99)	12.2 (7.29-20.3)	23.4 (14.3-35.4)	1003
Non-Hispanic whites	07-08	*	< LOD	2.93 (2.22-4.00)	11.4 (7.22-17.1)	23.9 (17.1-32.0)	2047
	09-10	*	< LOD	1.96 (1.82-2.15)	6.09 (5.00-7.83)	14.7 (10.0-18.2)	2514

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**Non-smoking is defined as a serum cotinine concentration of 10 ng/mL or less.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NNAL_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/NNAL_FactSheet.html

Blood Bromodichloromethane

Geometric mean and selected percentiles of whole blood concentrations (in pg/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	2.21 (1.65-2.97)	2.30 (1.56-3.21)	4.63 (3.24-6.20)	8.45 (5.86-12.0)	12.0 (7.68-19.2)	785
	03-04	1.50 (1.20-1.86)	1.40 (1.10-1.90)	3.40 (2.60-4.20)	6.20 (5.30-7.00)	9.50 (7.00-12.0)	1322
	05-06	1.41 (1.09-1.83)	1.30 (.880-1.80)	3.00 (2.10-4.40)	6.30 (4.30-9.70)	10.0 (6.80-14.0)	3139
Age group							
12-19 years	05-06	1.23 (.954-1.58)	1.00 (.620-1.60)	2.80 (1.70-4.10)	5.50 (4.10-7.20)	8.20 (6.20-12.0)	932
20-59 years	01-02	2.21 (1.65-2.97)	2.30 (1.56-3.21)	4.63 (3.24-6.20)	8.45 (5.86-12.0)	12.0 (7.68-19.2)	785
	03-04	1.50 (1.20-1.86)	1.40 (1.10-1.90)	3.40 (2.60-4.20)	6.20 (5.30-7.00)	9.50 (7.00-12.0)	1322
	05-06	1.45 (1.11-1.89)	1.30 (.900-1.90)	3.10 (2.10-4.60)	6.40 (4.30-10.0)	11.0 (6.90-14.0)	1537
60 years and older	05-06	1.43 (.996-2.05)	1.40 (.850-2.00)	3.20 (1.60-5.90)	6.50 (3.20-15.0)	9.70 (5.00-18.0)	670
Gender							
Males	01-02	2.19 (1.60-3.00)	2.31 (1.63-3.21)	4.64 (3.21-6.08)	7.96 (5.74-15.3)	13.0 (6.93-20.5)	382
	03-04	1.48 (1.18-1.85)	1.40 (.940-2.00)	3.40 (2.60-4.30)	6.60 (5.40-7.20)	11.0 (7.20-14.0)	650
	05-06	1.39 (1.07-1.80)	1.20 (.830-1.70)	3.00 (2.00-4.30)	6.50 (4.30-10.0)	11.0 (6.80-16.0)	1489
Females	01-02	2.24 (1.66-3.01)	2.28 (1.49-3.24)	4.63 (3.09-7.01)	8.62 (5.26-12.9)	11.1 (7.68-25.0)	403
	03-04	1.51 (1.21-1.90)	1.50 (1.10-1.90)	3.30 (2.50-4.20)	6.10 (4.69-7.30)	7.80 (6.40-12.0)	672
	05-06	1.44 (1.10-1.88)	1.30 (.900-1.90)	3.10 (2.10-4.60)	6.20 (4.20-9.40)	9.40 (6.30-13.0)	1650
Race/ethnicity							
Mexican Americans	01-02	3.28 (2.29-4.68)	3.32 (2.19-4.70)	6.81 (3.71-10.4)	10.8 (8.24-14.7)	14.7 (11.1-20.5)	227
	03-04	1.65 (1.15-2.38)	1.60 (.820-2.80)	3.50 (2.60-4.90)	7.30 (4.50-10.0)	10.0 (7.30-11.0)	244
	05-06	1.95 (1.19-3.18)	1.90 (1.00-3.70)	4.40 (2.10-9.10)	9.10 (4.80-17.0)	14.0 (7.50-22.0)	771
Non-Hispanic blacks	01-02	2.32 (1.82-2.94)	2.50 (1.56-3.55)	4.57 (3.60-5.56)	8.69 (5.63-9.49)	10.0 (5.89-13.5)	130
	03-04	1.56 (1.15-2.13)	1.70 (1.10-2.20)	2.90 (2.15-3.80)	5.10 (3.80-6.60)	6.60 (4.90-13.0)	290
	05-06	1.74 (1.27-2.37)	1.70 (1.00-2.70)	3.80 (2.70-4.80)	6.40 (4.50-8.90)	8.70 (6.60-11.0)	817
Non-Hispanic whites	01-02	2.02 (1.42-2.87)	2.16 (1.36-3.09)	4.34 (2.92-6.01)	7.33 (4.72-15.3)	11.1 (6.01-26.1)	365
	03-04	1.42 (1.11-1.81)	1.30 (.850-1.90)	3.30 (2.30-4.40)	6.20 (5.20-7.20)	9.80 (6.70-13.0)	684
	05-06	1.29 (.989-1.67)	1.10 (.710-1.70)	2.70 (1.80-4.10)	5.80 (4.00-8.60)	9.40 (6.20-14.0)	1318

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.233, 0.62, and 0.62 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/THM-DBP_BiomonitoringSummary.html

Blood Dibromochloromethane (Chlorodibromomethane)

Geometric mean and selected percentiles of whole blood concentrations (in pg/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	.867 (.521-1.44)	.780 (.340-1.90)	2.61 (1.22-4.38)	5.46 (3.53-9.71)	8.96 (5.04-12.9)	781
	03-04	*	< LOD	1.30 (1.00-1.80)	3.60 (2.70-4.80)	7.20 (4.80-8.60)	1333
	05-06	*	< LOD	1.40 (.870-2.40)	4.30 (2.40-7.40)	7.80 (4.60-13.0)	3122
Age group							
12-19 years	05-06	*	< LOD	1.30 (.750-2.50)	4.10 (2.60-5.50)	6.50 (4.10-11.0)	925
20-59 years	01-02	.867 (.521-1.44)	.780 (.340-1.90)	2.61 (1.22-4.38)	5.46 (3.53-9.71)	8.96 (5.04-12.9)	781
	03-04	*	< LOD	1.30 (1.00-1.80)	3.60 (2.70-4.80)	7.20 (4.80-8.60)	1333
	05-06	*	< LOD	1.40 (.860-2.30)	4.10 (2.40-7.10)	7.80 (4.20-14.0)	1526
60 years and older	05-06	*	< LOD	1.60 (.780-4.00)	5.10 (1.30-13.0)	7.90 (2.20-20.0)	671
Gender							
Males	01-02	.850 (.481-1.50)	.730 (.300-2.25)	2.66 (.960-4.38)	4.77 (3.33-9.20)	8.06 (4.31-14.6)	371
	03-04	*	< LOD	1.30 (1.00-1.90)	3.70 (2.70-5.70)	7.20 (5.20-8.60)	657
	05-06	*	< LOD	1.40 (.830-2.30)	4.40 (2.30-7.80)	8.30 (4.30-14.0)	1487
Females	01-02	.884 (.550-1.42)	.820 (.340-1.69)	2.54 (1.37-4.31)	6.30 (3.28-10.1)	9.91 (5.02-13.0)	410
	03-04	*	< LOD	1.30 (1.00-1.80)	3.60 (2.50-4.80)	6.60 (3.80-9.20)	676
	05-06	*	< LOD	1.50 (.920-2.50)	4.20 (2.20-7.80)	7.20 (4.10-14.0)	1635
Race/ethnicity							
Mexican Americans	01-02	1.61 (.843-3.06)	1.49 (.670-3.99)	4.59 (1.93-8.89)	9.26 (5.21-12.1)	12.0 (9.63-16.1)	233
	03-04	1.20 (.963-1.50)	1.10 (.810-1.40)	2.30 (1.50-4.10)	5.20 (3.80-6.90)	7.70 (5.20-11.0)	256
	05-06	1.25 (.774-2.02)	.950 (<LOD-1.70)	2.50 (1.00-7.40)	7.40 (2.50-19.0)	13.0 (5.80-21.0)	759
Non-Hispanic blacks	01-02	1.03 (.505-2.09)	.930 (.530-2.03)	2.03 (.770-7.06)	4.22 (2.01-10.5)	8.09 (2.80-16.5)	128
	03-04	*	< LOD	.970 (.650-1.60)	1.90 (1.20-3.20)	3.20 (1.80-5.20)	288
	05-06	*	.620 (<LOD-.820)	1.30 (1.00-1.90)	3.40 (2.10-4.90)	5.50 (4.10-7.10)	822
Non-Hispanic whites	01-02	.736 (.413-1.31)	.640 (<LOD-1.93)	2.49 (.870-4.27)	4.57 (3.00-7.12)	6.98 (4.27-11.1)	357
	03-04	*	< LOD	1.30 (.950-1.80)	3.30 (2.50-4.60)	6.60 (3.70-9.20)	685
	05-06	*	< LOD	1.30 (.710-2.30)	4.10 (2.30-7.10)	7.70 (4.60-11.0)	1311

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.271, 0.62, and 0.62 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/THM-DBP_BiomonitoringSummary.html

Blood Tribromomethane (Bromoform)

Geometric mean and selected percentiles of whole blood concentrations (in pg/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	1.57 (1.07-2.31)	1.39 (.960-2.02)	2.78 (1.76-4.63)	6.05 (2.92-29.2)	15.5 (3.68-85.4)	774
	03-04	*	< LOD	1.80 (<LOD-2.80)	3.74 (2.30-7.10)	6.40 (3.60-14.0)	1310
	05-06	*	< LOD	1.20 (<LOD-1.60)	2.70 (2.00-4.00)	5.20 (3.10-7.60)	3016
Age group							
12-19 years	05-06	*	< LOD	1.20 (<LOD-1.60)	2.40 (1.70-4.40)	4.80 (2.80-7.10)	892
20-59 years	01-02	1.57 (1.07-2.31)	1.39 (.960-2.02)	2.78 (1.76-4.63)	6.05 (2.92-29.2)	15.5 (3.68-85.4)	774
	03-04	*	< LOD	1.80 (<LOD-2.80)	3.74 (2.30-7.10)	6.40 (3.60-14.0)	1310
	05-06	*	< LOD	1.20 (<LOD-1.60)	2.90 (1.90-4.20)	5.20 (3.10-8.80)	1478
60 years and older	05-06	*	< LOD	1.28 (<LOD-2.10)	2.60 (2.30-3.50)	4.70 (3.00-7.70)	646
Gender							
Males	01-02	1.49 (.944-2.34)	1.29 (.850-1.98)	2.65 (1.49-5.05)	6.12 (2.26-33.9)	14.9 (2.79-69.9)	374
	03-04	*	< LOD	1.90 (<LOD-2.87)	4.00 (2.40-6.80)	6.50 (4.00-13.0)	645
	05-06	*	< LOD	1.16 (<LOD-1.60)	2.70 (1.90-4.00)	5.00 (3.10-7.60)	1424
Females	01-02	1.67 (1.17-2.39)	1.46 (1.05-2.21)	2.86 (1.89-4.57)	5.69 (3.30-27.5)	22.2 (5.09-49.6)	400
	03-04	*	< LOD	1.72 (<LOD-2.65)	3.20 (1.93-7.70)	6.10 (3.10-31.0)	665
	05-06	*	< LOD	1.23 (1.00-1.60)	2.70 (1.90-4.20)	5.20 (3.10-8.80)	1592
Race/ethnicity							
Mexican Americans	01-02	2.34 (1.15-4.77)	1.66 (.990-3.38)	4.03 (1.42-36.5)	28.3 (4.39-49.2)	40.8 (31.5-57.9)	234
	03-04	*	1.60 (<LOD-3.10)	3.30 (<LOD-9.40)	7.60 (3.60-14.0)	11.0 (5.60-210)	242
	05-06	*	< LOD	1.60 (1.04-3.20)	4.00 (2.40-5.90)	5.90 (4.30-8.30)	721
Non-Hispanic blacks	01-02	1.51 (.857-2.67)	1.47 (.780-3.15)	2.58 (1.39-4.64)	4.34 (2.57-8.48)	6.27 (3.28-15.2)	121
	03-04	*	< LOD	1.60 (<LOD-2.30)	2.50 (1.80-3.20)	3.20 (2.40-6.10)	289
	05-06	*	< LOD	1.10 (<LOD-1.30)	2.00 (1.70-2.40)	3.40 (2.30-4.30)	796
Non-Hispanic whites	01-02	1.47 (.980-2.22)	1.29 (.840-2.06)	2.58 (1.51-4.91)	5.69 (2.84-21.2)	11.0 (3.30-69.9)	362
	03-04	*	< LOD	1.70 (<LOD-2.90)	3.50 (2.00-7.70)	5.90 (3.30-21.0)	680
	05-06	*	< LOD	1.20 (<LOD-1.60)	2.80 (2.00-3.90)	5.40 (3.10-9.90)	1271

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.596, 1.5, and 1.0 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/THM-DBP_BiomonitoringSummary.html

Blood Trichloromethane (Chloroform)

Geometric mean and selected percentiles of whole blood concentrations (in pg/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	16.6 (13.0-21.1)	16.1 (11.9-22.2)	31.7 (23.9-40.4)	55.5 (44.5-68.6)	72.1 (57.3-105)	744
	03-04	10.2 (8.56-12.2)	10.0 (8.50-13.0)	20.0 (17.0-24.0)	35.0 (29.0-40.0)	50.0 (37.0-65.0)	1222
	05-06	10.1 (9.09-11.3)	9.90 (8.90-11.0)	19.0 (17.0-21.0)	36.0 (31.0-41.0)	54.0 (46.0-64.0)	3111
Age group							
12-19 years	05-06	9.22 (8.20-10.4)	9.10 (8.10-10.0)	17.0 (14.0-20.0)	33.0 (27.0-40.0)	46.0 (36.0-64.0)	913
20-59 years	01-02	16.6 (13.0-21.1)	16.1 (11.9-22.2)	31.7 (23.9-40.4)	55.5 (44.5-68.6)	72.1 (57.3-105)	744
	03-04	10.2 (8.56-12.2)	10.0 (8.50-13.0)	20.0 (17.0-24.0)	35.0 (29.0-40.0)	50.0 (37.0-65.0)	1222
	05-06	10.5 (9.29-11.8)	10.0 (8.90-12.0)	20.0 (18.0-22.0)	37.0 (31.0-44.0)	55.0 (46.0-71.0)	1529
60 years and older	05-06	9.71 (8.25-11.4)	9.60 (8.20-11.0)	19.0 (14.0-22.0)	35.0 (27.0-40.0)	46.0 (39.0-59.0)	669
Gender							
Males	01-02	16.8 (12.0-23.5)	16.1 (11.0-24.8)	34.3 (22.4-48.7)	57.0 (39.9-76.4)	75.2 (54.5-156)	358
	03-04	10.1 (8.43-12.1)	10.0 (7.90-14.0)	20.0 (17.0-25.0)	36.8 (29.0-49.0)	53.0 (36.8-69.0)	599
	05-06	9.94 (8.80-11.2)	9.80 (8.50-11.0)	19.0 (17.0-22.0)	38.0 (33.0-44.0)	54.0 (46.0-67.0)	1483
Females	01-02	16.4 (13.4-20.1)	16.6 (12.0-21.5)	29.2 (24.0-36.5)	53.5 (38.4-68.9)	69.5 (53.3-104)	386
	03-04	10.4 (8.41-12.7)	10.0 (8.40-13.0)	20.0 (16.0-23.9)	33.0 (26.0-40.0)	46.0 (35.0-65.0)	623
	05-06	10.4 (9.24-11.6)	10.0 (9.10-11.0)	19.0 (17.0-21.0)	32.0 (29.0-39.0)	52.0 (41.0-67.0)	1628
Race/ethnicity							
Mexican Americans	01-02	17.0 (10.5-27.6)	14.5 (10.0-32.7)	35.1 (18.6-57.6)	60.7 (41.5-100)	93.0 (49.7-243)	223
	03-04	9.17 (7.45-11.3)	9.30 (7.60-11.0)	19.0 (15.0-24.0)	34.0 (24.0-44.4)	44.4 (30.0-59.0)	225
	05-06	10.3 (8.59-12.5)	10.0 (7.60-14.0)	20.0 (17.0-22.0)	36.0 (30.0-41.0)	47.0 (41.0-64.0)	740
Non-Hispanic blacks	01-02	19.1 (12.9-28.1)	20.9 (9.28-37.4)	38.4 (27.7-46.0)	55.9 (45.5-69.8)	68.9 (51.9-74.0)	116
	03-04	11.8 (9.54-14.6)	12.0 (8.90-15.0)	20.0 (15.0-30.0)	35.0 (28.0-59.0)	61.0 (34.0-100)	272
	05-06	11.9 (9.78-14.4)	12.0 (9.10-15.0)	22.0 (19.0-26.0)	39.0 (31.0-47.0)	55.0 (46.0-70.0)	822
Non-Hispanic whites	01-02	15.6 (12.0-20.2)	15.2 (11.2-20.5)	26.8 (20.4-39.5)	53.5 (35.3-72.1)	69.5 (53.5-105)	348
	03-04	9.84 (8.09-12.0)	10.0 (8.10-13.0)	20.0 (16.0-23.0)	33.8 (27.0-40.0)	47.0 (35.0-67.0)	630
	05-06	9.87 (8.79-11.1)	9.70 (8.80-11.0)	19.0 (16.0-21.0)	34.0 (29.0-43.0)	53.0 (45.0-67.0)	1318

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 2.37, 2.11, and 2.1 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/THM-DBP_BiomonitoringSummary.html

Urinary Benzophenone-3

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	22.9 (18.1-28.9)	18.1 (15.5-23.2)	94.0 (67.5-123)	370 (225-570)	1040 (698-1390)	2517
	05-06	19.4 (16.5-22.8)	14.1 (12.5-17.5)	68.8 (54.5-87.2)	433 (294-597)	1150 (936-1510)	2548
	07-08	18.3 (13.7-24.3)	14.3 (10.7-19.8)	64.1 (42.8-116)	365 (223-524)	801 (471-2080)	2604
	09-10	22.3 (18.0-27.7)	16.4 (12.7-20.5)	81.2 (55.0-126)	493 (320-742)	1750 (931-3120)	2749
Age group							
6-11 years	03-04	21.2 (16.4-27.3)	17.2 (14.9-25.9)	66.7 (38.7-102)	158 (106-246)	246 (154-618)	314
	05-06	21.2 (14.3-31.3)	19.4 (11.7-26.1)	58.8 (40.5-102)	259 (115-433)	852 (355-2200)	356
	07-08	24.1 (15.0-38.7)	17.2 (11.8-25.0)	79.9 (32.9-168)	392 (129-1610)	1410 (165-43600)	389
	09-10	21.7 (15.6-30.2)	14.6 (12.0-19.7)	52.7 (27.2-119)	470 (80.7-1900)	1570 (153-6440)	415
12-19 years	03-04	22.9 (18.0-29.3)	20.1 (16.1-25.1)	67.1 (45.2-93.8)	170 (137-240)	407 (183-717)	715
	05-06	21.2 (15.8-28.4)	14.9 (12.1-21.4)	64.3 (39.7-95.3)	227 (130-382)	633 (252-1260)	702
	07-08	21.5 (16.0-28.8)	15.0 (11.9-21.3)	59.3 (36.7-111)	214 (113-613)	613 (206-1350)	401
	09-10	23.3 (16.6-32.5)	16.6 (13.1-19.7)	62.3 (35.1-129)	423 (184-843)	988 (550-2670)	420
20 years and older	03-04	23.1 (18.0-29.6)	18.1 (14.7-23.3)	109 (72.1-140)	450 (315-733)	1220 (769-1750)	1488
	05-06	19.0 (16.5-21.8)	13.5 (12.0-16.3)	70.9 (55.2-87.2)	543 (390-638)	1200 (954-1690)	1490
	07-08	17.3 (13.0-23.0)	13.7 (10.0-19.5)	64.1 (41.5-117)	371 (240-516)	801 (516-1790)	1814
	09-10	22.3 (17.9-27.7)	16.5 (12.3-22.5)	86.7 (63.1-129)	505 (339-760)	1890 (1240-3170)	1914
Gender							
Males	03-04	16.8 (13.2-21.3)	13.7 (11.4-16.8)	55.3 (33.2-86.6)	178 (134-324)	567 (238-1350)	1229
	05-06	14.6 (10.8-19.7)	11.8 (8.80-15.9)	43.4 (31.0-68.8)	227 (103-552)	909 (479-1190)	1270
	07-08	11.9 (8.84-16.0)	9.40 (7.70-12.3)	37.6 (22.6-54.7)	172 (79.9-376)	471 (208-1410)	1294
	09-10	15.3 (12.7-18.6)	12.2 (10.0-15.2)	52.8 (33.9-72.6)	223 (129-332)	610 (318-1270)	1399
Females	03-04	30.7 (23.7-39.8)	26.0 (20.2-34.1)	137 (106-172)	596 (403-769)	1340 (776-1790)	1288
	05-06	25.5 (21.7-29.9)	17.9 (14.3-23.1)	129 (82.9-166)	638 (478-900)	1410 (945-2620)	1278
	07-08	27.6 (20.4-37.3)	25.6 (17.5-33.9)	117 (72.6-211)	535 (361-757)	1260 (643-2280)	1310
	09-10	32.0 (23.7-43.3)	23.0 (16.4-29.8)	148 (73.3-260)	1020 (540-1800)	3200 (1760-4400)	1350
Race/ethnicity							
Mexican Americans	03-04	16.5 (10.9-25.1)	11.9 (8.50-18.3)	45.5 (25.9-78.2)	178 (76.4-412)	412 (178-2180)	613
	05-06	14.3 (10.7-19.1)	10.2 (7.40-14.1)	45.6 (24.1-75.7)	214 (166-313)	635 (337-1090)	637
	07-08	12.1 (9.49-15.3)	9.90 (7.70-12.4)	32.6 (25.0-48.3)	142 (85.0-268)	409 (214-778)	531
	09-10	17.2 (13.7-21.6)	12.5 (10.5-14.5)	41.9 (30.2-57.4)	351 (248-546)	1130 (493-2140)	566
Non-Hispanic blacks	03-04	12.8 (9.38-17.4)	10.2 (7.40-14.4)	34.3 (22.8-50.6)	127 (90.8-176)	247 (143-499)	652
	05-06	12.4 (9.77-15.6)	9.40 (7.30-11.8)	33.0 (22.3-46.5)	208 (115-291)	556 (269-1060)	678
	07-08	8.55 (6.75-10.8)	7.40 (5.00-9.70)	25.5 (17.5-34.5)	117 (66.8-217)	422 (195-786)	597
	09-10	14.0 (10.9-18.1)	8.90 (7.30-10.5)	30.9 (21.9-50.5)	276 (143-437)	705 (320-3000)	516
Non-Hispanic whites	03-04	27.7 (20.3-37.8)	24.4 (16.8-32.0)	121 (83.6-162)	507 (316-769)	1340 (733-2070)	1092
	05-06	21.6 (18.5-25.2)	16.0 (13.5-19.0)	82.4 (64.2-114)	552 (392-754)	1370 (945-2050)	1038
	07-08	23.1 (15.0-35.5)	18.7 (11.0-33.7)	101 (47.4-206)	459 (241-749)	1160 (487-2620)	1077
	09-10	26.1 (19.0-35.9)	21.1 (14.6-29.7)	94.9 (59.8-186)	610 (325-1240)	2130 (934-3820)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.3, 0.4, 0.4, and 0.4 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Benzophenone-3_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Benzophenone-3_FactSheet.html

Urinary Benzophenone-3 (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	22.2 (17.6-28.0)	16.2 (12.7-21.6)	82.0 (58.7-108)	415 (283-577)	1080 (686-1600)	2514
	05-06	18.9 (15.8-22.6)	13.5 (11.2-16.8)	60.4 (47.6-81.4)	445 (346-553)	1210 (794-1970)	2548
	07-08	18.4 (13.9-24.4)	13.5 (10.2-17.8)	67.9 (45.2-117)	338 (215-584)	935 (525-1800)	2604
	09-10	23.3 (18.7-29.1)	16.1 (12.8-19.8)	79.1 (52.2-124)	579 (355-1040)	2340 (1380-3010)	2749
Age group							
6-11 years	03-04	25.8 (19.5-34.1)	22.4 (14.4-33.7)	84.6 (41.0-131)	171 (132-365)	427 (171-710)	314
	05-06	23.3 (15.9-34.1)	17.5 (11.6-28.3)	67.7 (39.7-105)	249 (106-631)	868 (264-2080)	356
	07-08	29.7 (19.0-46.6)	19.3 (12.4-32.3)	103 (51.1-159)	442 (133-1550)	1430 (187-29100)	389
	09-10	28.3 (20.4-39.4)	18.8 (14.7-21.1)	53.9 (33.2-132)	565 (104-3410)	1780 (253-8230)	415
12-19 years	03-04	17.2 (13.7-21.5)	12.9 (10.4-16.5)	43.6 (29.5-57.7)	136 (91.7-239)	350 (173-646)	713
	05-06	15.8 (11.6-21.6)	12.1 (9.16-17.0)	42.5 (28.6-58.7)	164 (101-438)	625 (225-956)	702
	07-08	16.7 (12.4-22.4)	12.1 (8.57-17.6)	55.7 (30.4-74.8)	211 (88.2-505)	505 (163-902)	401
	09-10	18.7 (13.4-26.1)	13.6 (10.1-19.7)	39.9 (25.5-91.6)	412 (89.8-750)	1240 (629-2540)	420
20 years and older	03-04	22.8 (17.8-29.1)	16.2 (12.7-21.9)	93.2 (66.0-130)	491 (361-700)	1330 (880-1880)	1487
	05-06	19.0 (16.3-22.2)	13.4 (11.2-16.2)	64.2 (49.3-87.0)	509 (406-628)	1480 (816-2200)	1490
	07-08	17.7 (13.3-23.6)	12.6 (9.93-17.5)	67.7 (44.1-127)	362 (249-584)	960 (581-1970)	1814
	09-10	23.6 (18.8-29.5)	16.3 (12.7-20.2)	83.6 (58.3-146)	602 (364-1230)	2470 (1350-3080)	1914
Gender							
Males	03-04	13.6 (10.8-17.1)	10.3 (8.36-12.9)	40.0 (24.9-62.5)	169 (93.3-316)	381 (229-685)	1228
	05-06	11.7 (8.74-15.8)	9.43 (6.72-12.5)	31.9 (19.7-49.4)	176 (77.1-406)	553 (317-948)	1270
	07-08	10.0 (7.37-13.7)	7.50 (5.74-10.0)	28.7 (18.5-47.3)	142 (65.8-225)	332 (181-960)	1294
	09-10	13.7 (11.4-16.6)	10.7 (8.53-12.7)	44.6 (29.7-57.4)	186 (123-287)	420 (287-1230)	1399
Females	03-04	35.5 (27.1-46.4)	28.2 (20.2-37.0)	144 (101-224)	686 (491-1130)	1850 (1220-2580)	1286
	05-06	30.0 (25.3-35.6)	20.9 (17.0-29.0)	129 (88.0-175)	707 (527-912)	1880 (1150-2770)	1278
	07-08	33.0 (25.0-43.7)	26.2 (19.1-41.1)	156 (90.2-235)	617 (385-935)	1500 (746-2710)	1310
	09-10	38.6 (28.1-53.0)	25.7 (17.5-35.2)	161 (76.3-378)	1580 (955-2410)	3250 (2540-4640)	1350
Race/ethnicity							
Mexican Americans	03-04	15.1 (9.44-24.0)	11.1 (6.95-16.0)	40.7 (18.3-85.8)	158 (87.4-362)	595 (118-1860)	612
	05-06	12.9 (10.2-16.2)	8.63 (6.16-12.5)	32.1 (23.1-51.8)	216 (135-408)	948 (441-1220)	637
	07-08	11.8 (8.87-15.6)	8.97 (7.00-12.4)	33.6 (26.0-43.2)	154 (93.4-249)	413 (217-932)	531
	09-10	17.1 (13.4-21.7)	12.2 (9.64-14.5)	40.4 (26.4-60.3)	288 (155-603)	1310 (441-2180)	566
Non-Hispanic blacks	03-04	8.78 (6.49-11.9)	6.80 (5.27-9.00)	19.7 (13.5-33.4)	79.8 (46.8-139)	185 (79.8-536)	651
	05-06	8.69 (7.02-10.8)	6.29 (5.52-7.78)	21.2 (14.5-34.5)	118 (74.9-245)	448 (208-876)	678
	07-08	6.65 (5.24-8.42)	5.09 (4.15-6.10)	17.8 (13.2-24.2)	86.6 (57.5-122)	297 (113-903)	597
	09-10	10.2 (8.12-12.7)	6.46 (5.40-7.67)	23.1 (15.5-38.2)	160 (79.9-396)	653 (305-1580)	516
Non-Hispanic whites	03-04	28.3 (20.6-38.8)	22.0 (14.6-32.7)	116 (73.5-175)	510 (380-760)	1330 (852-2410)	1091
	05-06	22.7 (19.4-26.7)	16.2 (13.0-19.7)	77.3 (56.5-106)	538 (406-679)	1500 (828-2430)	1038
	07-08	24.1 (15.6-37.2)	18.1 (11.0-30.0)	103 (53.6-198)	494 (227-960)	1170 (522-2710)	1077
	09-10	29.3 (20.7-41.5)	22.1 (14.6-33.1)	99.2 (54.3-241)	767 (412-1550)	2590 (1510-3250)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Benzophenone-3_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Benzophenone-3_FactSheet.html

Urinary Bisphenol A

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	2.64 (2.38-2.94)	2.80 (2.50-3.10)	5.50 (5.00-6.20)	10.6 (9.40-12.0)	16.0 (14.4-17.2)	2517
	05-06	1.90 (1.79-2.02)	2.00 (1.90-2.00)	3.70 (3.50-3.90)	7.00 (6.40-7.60)	11.5 (10.0-13.6)	2548
	07-08	2.08 (1.92-2.26)	2.10 (1.90-2.30)	4.10 (3.60-4.60)	7.70 (6.80-8.70)	13.0 (10.0-15.6)	2604
	09-10	1.83 (1.72-1.94)	1.90 (1.70-2.00)	3.50 (3.30-3.80)	6.60 (6.00-7.20)	9.60 (8.50-11.3)	2749
Age group							
6-11 years	03-04	3.55 (2.95-4.29)	3.80 (2.70-5.00)	6.90 (6.00-8.30)	12.6 (9.50-15.1)	16.0 (11.5-23.3)	314
	05-06	2.86 (2.52-3.24)	2.70 (2.30-2.90)	5.00 (4.40-5.80)	13.5 (9.30-16.8)	22.8 (13.6-34.6)	356
	07-08	2.48 (2.21-2.77)	2.40 (1.90-3.00)	4.50 (3.70-5.50)	7.60 (6.30-9.50)	13.4 (8.80-17.8)	389
	09-10	1.81 (1.55-2.10)	1.70 (1.50-2.00)	3.40 (3.00-4.00)	6.50 (4.30-9.70)	9.20 (6.50-15.1)	415
12-19 years	03-04	3.74 (3.31-4.22)	4.30 (3.60-4.60)	7.80 (6.50-9.00)	13.5 (11.8-15.2)	16.5 (15.2-20.9)	715
	05-06	2.42 (2.18-2.68)	2.40 (2.10-2.70)	4.30 (3.90-5.20)	8.40 (6.50-10.8)	11.9 (10.7-14.8)	702
	07-08	2.45 (2.14-2.80)	2.40 (2.10-2.60)	4.40 (3.60-5.50)	9.70 (7.30-11.9)	12.2 (9.70-19.0)	401
	09-10	2.11 (1.86-2.40)	2.20 (1.90-2.40)	3.80 (3.20-4.50)	6.90 (4.70-10.8)	11.1 (6.90-21.0)	420
20 years and older	03-04	2.41 (2.15-2.72)	2.60 (2.30-2.80)	5.10 (4.50-5.70)	9.50 (8.10-11.3)	15.2 (12.4-18.1)	1488
	05-06	1.75 (1.62-1.89)	1.80 (1.70-2.00)	3.40 (3.10-3.70)	6.40 (5.80-7.50)	10.7 (8.80-12.1)	1490
	07-08	1.99 (1.82-2.18)	2.00 (1.80-2.30)	3.90 (3.40-4.60)	7.40 (6.60-8.50)	13.2 (9.10-15.7)	1814
	09-10	1.79 (1.67-1.93)	1.80 (1.60-2.00)	3.50 (3.30-3.70)	6.50 (6.00-7.20)	9.60 (8.30-11.3)	1914
Gender							
Males	03-04	2.92 (2.63-3.24)	3.20 (2.70-3.60)	6.10 (5.40-6.60)	10.4 (9.50-11.6)	16.0 (12.7-17.6)	1229
	05-06	2.09 (1.92-2.28)	2.10 (2.00-2.30)	3.70 (3.40-4.20)	7.70 (6.80-9.30)	12.8 (10.1-15.8)	1270
	07-08	2.20 (2.01-2.41)	2.10 (1.90-2.40)	4.00 (3.60-4.70)	8.10 (6.70-9.70)	14.0 (8.70-20.8)	1294
	09-10	1.94 (1.82-2.07)	1.90 (1.70-2.00)	3.70 (3.40-4.10)	6.90 (5.90-7.90)	10.9 (8.40-13.5)	1399
Females	03-04	2.41 (2.11-2.75)	2.50 (2.20-2.80)	5.00 (4.20-6.20)	10.6 (8.70-12.5)	15.9 (13.5-20.1)	1288
	05-06	1.74 (1.55-1.95)	1.80 (1.60-2.00)	3.70 (3.10-4.00)	6.20 (5.80-7.60)	10.4 (7.70-14.7)	1278
	07-08	1.97 (1.80-2.16)	2.00 (1.80-2.20)	4.10 (3.60-4.60)	7.40 (6.80-8.20)	12.0 (9.60-14.1)	1310
	09-10	1.73 (1.60-1.87)	1.80 (1.60-2.00)	3.40 (3.20-3.70)	6.50 (5.60-7.10)	9.20 (8.00-11.2)	1350
Race/ethnicity							
Mexican Americans	03-04	2.58 (2.15-3.08)	2.60 (2.10-3.20)	5.20 (4.40-6.50)	9.90 (7.30-13.9)	15.4 (10.2-19.7)	613
	05-06	2.05 (1.75-2.40)	2.00 (1.70-2.40)	3.60 (3.00-4.20)	6.90 (5.40-10.1)	12.2 (8.00-15.6)	637
	07-08	2.09 (1.94-2.26)	2.00 (1.80-2.30)	3.80 (3.30-4.30)	7.10 (6.10-8.60)	11.7 (7.70-14.6)	531
	09-10	1.92 (1.69-2.19)	2.10 (1.80-2.30)	3.50 (3.10-4.10)	6.50 (5.20-7.50)	9.00 (6.90-16.2)	566
Non-Hispanic blacks	03-04	4.24 (3.73-4.82)	4.30 (3.80-5.10)	8.20 (7.10-9.80)	14.2 (11.7-16.9)	20.6 (14.9-25.2)	652
	05-06	2.50 (2.25-2.77)	2.70 (2.30-3.00)	4.60 (4.00-5.00)	8.00 (6.80-9.30)	11.3 (8.90-14.2)	678
	07-08	2.66 (2.38-2.97)	2.80 (2.60-3.10)	5.40 (4.40-6.40)	9.10 (7.50-10.4)	13.3 (10.0-15.1)	597
	09-10	2.51 (2.22-2.83)	2.60 (2.30-3.00)	4.40 (3.80-5.00)	7.30 (6.10-8.80)	10.3 (7.90-13.6)	516
Non-Hispanic whites	03-04	2.51 (2.26-2.79)	2.70 (2.50-3.00)	5.20 (4.70-5.80)	9.60 (8.30-10.9)	15.1 (12.6-16.7)	1092
	05-06	1.76 (1.62-1.91)	1.80 (1.60-2.00)	3.50 (3.20-3.80)	6.80 (5.90-7.60)	11.0 (8.80-13.7)	1038
	07-08	2.06 (1.87-2.27)	2.00 (1.80-2.30)	4.00 (3.40-4.70)	7.80 (6.70-8.90)	13.7 (9.80-16.1)	1077
	09-10	1.73 (1.60-1.87)	1.70 (1.60-1.90)	3.40 (3.10-3.70)	6.40 (5.40-7.60)	9.20 (8.20-11.2)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.4, 0.4, 0.4, and 0.4 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/BisphenolA_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/BisphenolA_FactSheet.html

Urinary Bisphenol A (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	2.58 (2.36-2.82)	2.50 (2.31-2.80)	4.29 (3.88-4.75)	7.67 (6.62-8.66)	11.2 (9.78-12.4)	2514
	05-06	1.86 (1.79-1.92)	1.71 (1.64-1.79)	3.01 (2.86-3.20)	5.73 (5.29-6.36)	9.70 (8.31-10.9)	2548
	07-08	2.10 (1.95-2.25)	1.95 (1.84-2.03)	3.45 (3.02-3.83)	6.09 (5.10-7.45)	10.0 (7.48-13.2)	2604
	09-10	1.91 (1.79-2.04)	1.76 (1.67-1.84)	3.06 (2.80-3.33)	5.33 (4.68-6.06)	8.03 (6.94-10.0)	2749
Age group							
6-11 years	03-04	4.32 (3.63-5.14)	4.29 (3.63-5.23)	7.14 (5.83-9.56)	12.2 (9.84-14.8)	15.7 (12.2-23.2)	314
	05-06	3.14 (2.79-3.54)	2.80 (2.55-3.06)	5.04 (4.46-5.71)	15.6 (8.15-22.4)	24.6 (19.9-48.5)	356
	07-08	3.05 (2.73-3.41)	2.69 (2.38-3.12)	5.06 (4.33-5.60)	11.9 (6.30-17.4)	20.8 (12.7-26.3)	389
	09-10	2.36 (2.09-2.65)	2.18 (2.00-2.53)	3.70 (3.14-4.29)	5.97 (4.57-9.15)	9.57 (6.52-13.7)	415
12-19 years	03-04	2.80 (2.52-3.11)	2.74 (2.35-3.22)	4.74 (4.21-5.09)	7.79 (6.41-8.87)	11.8 (8.05-14.2)	713
	05-06	1.80 (1.67-1.95)	1.65 (1.50-1.76)	2.73 (2.41-2.98)	5.71 (4.07-7.50)	8.52 (6.94-12.2)	702
	07-08	1.90 (1.72-2.10)	1.69 (1.50-2.00)	2.94 (2.38-3.60)	5.10 (4.17-6.82)	7.72 (5.32-13.9)	401
	09-10	1.70 (1.47-1.95)	1.60 (1.34-1.88)	2.63 (2.11-3.58)	4.47 (3.76-6.19)	6.71 (4.40-17.8)	420
20 years and older	03-04	2.39 (2.17-2.64)	2.36 (2.15-2.59)	3.93 (3.44-4.33)	6.64 (5.97-7.74)	10.0 (9.01-11.4)	1487
	05-06	1.75 (1.67-1.84)	1.64 (1.56-1.75)	2.84 (2.67-3.08)	5.38 (4.89-5.87)	8.54 (7.58-9.77)	1490
	07-08	2.04 (1.90-2.20)	1.92 (1.79-2.03)	3.36 (2.91-3.78)	6.02 (4.88-7.48)	9.32 (7.48-12.1)	1814
	09-10	1.90 (1.76-2.05)	1.76 (1.66-1.84)	3.04 (2.76-3.24)	5.33 (4.67-6.11)	8.00 (6.97-10.0)	1914
Gender							
Males	03-04	2.38 (2.15-2.63)	2.31 (2.08-2.70)	4.19 (3.81-4.64)	7.10 (6.41-8.28)	9.94 (9.06-11.7)	1228
	05-06	1.68 (1.57-1.80)	1.56 (1.45-1.64)	2.73 (2.43-3.14)	5.27 (4.64-6.04)	8.40 (6.81-11.4)	1270
	07-08	1.85 (1.71-2.01)	1.77 (1.57-1.91)	2.97 (2.59-3.33)	5.41 (4.47-7.51)	9.52 (6.29-13.4)	1294
	09-10	1.74 (1.62-1.87)	1.60 (1.48-1.72)	2.76 (2.53-2.96)	5.00 (4.16-5.86)	7.50 (6.22-9.00)	1399
Females	03-04	2.78 (2.50-3.08)	2.68 (2.40-2.94)	4.41 (3.81-5.15)	7.93 (6.48-10.2)	12.4 (9.29-18.2)	1286
	05-06	2.04 (1.95-2.14)	1.85 (1.76-2.02)	3.21 (2.98-3.39)	6.47 (5.51-7.50)	9.86 (8.62-11.8)	1278
	07-08	2.36 (2.17-2.57)	2.14 (1.98-2.33)	3.77 (3.43-4.31)	6.67 (5.41-7.83)	10.5 (7.45-15.2)	1310
	09-10	2.09 (1.92-2.27)	1.93 (1.80-2.15)	3.33 (2.94-3.84)	5.49 (4.87-6.74)	8.80 (6.47-12.5)	1350
Race/ethnicity							
Mexican Americans	03-04	2.34 (2.02-2.71)	2.38 (2.00-2.65)	3.85 (3.24-4.55)	7.09 (5.00-9.04)	10.9 (8.50-14.3)	612
	05-06	1.84 (1.66-2.04)	1.62 (1.45-1.84)	2.99 (2.56-3.60)	6.00 (4.51-8.00)	10.3 (6.84-14.9)	637
	07-08	2.04 (1.86-2.24)	1.88 (1.72-2.11)	3.33 (2.96-3.95)	5.68 (5.05-6.59)	8.90 (6.30-11.8)	531
	09-10	1.91 (1.71-2.13)	1.76 (1.67-1.87)	3.13 (2.63-3.57)	5.51 (4.72-6.97)	8.71 (5.86-14.4)	566
Non-Hispanic blacks	03-04	2.92 (2.58-3.32)	2.95 (2.51-3.27)	4.90 (4.07-6.13)	8.64 (7.53-9.63)	11.9 (10.2-13.3)	651
	05-06	1.76 (1.62-1.90)	1.68 (1.53-1.92)	2.82 (2.67-3.04)	4.56 (3.85-5.43)	6.81 (5.00-9.27)	678
	07-08	2.06 (1.91-2.23)	2.03 (1.83-2.25)	3.24 (2.86-3.78)	5.71 (5.15-6.54)	8.59 (6.54-10.9)	597
	09-10	1.82 (1.65-2.01)	1.70 (1.56-1.89)	2.90 (2.39-3.17)	4.94 (4.02-6.03)	6.74 (5.95-8.09)	516
Non-Hispanic whites	03-04	2.58 (2.37-2.81)	2.55 (2.32-2.80)	4.30 (3.93-4.67)	7.58 (6.32-8.87)	11.0 (9.34-12.4)	1091
	05-06	1.85 (1.75-1.96)	1.70 (1.60-1.82)	3.08 (2.78-3.31)	5.80 (5.13-6.72)	9.77 (7.87-11.5)	1038
	07-08	2.15 (1.97-2.35)	1.97 (1.84-2.11)	3.50 (3.04-3.95)	6.44 (5.04-8.00)	10.9 (7.47-14.5)	1077
	09-10	1.94 (1.79-2.10)	1.78 (1.67-1.92)	3.16 (2.80-3.59)	5.45 (4.67-6.94)	8.55 (6.73-10.0)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/BisphenolA_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/BisphenolA_FactSheet.html

Urinary 4-*tert*-Octylphenol

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	05-06	*	< LOD	< LOD	< LOD	.200 (<LOD-.300)	2548
	07-08	*	< LOD	< LOD	< LOD	.300 (.200-.300)	2604
	09-10	*	< LOD	< LOD	.200 (<LOD-.300)	.300 (.300-.400)	2749
Age group							
6-11 years	05-06	*	< LOD	< LOD	< LOD	.300 (.200-.300)	356
	07-08	*	< LOD	< LOD	.200 (<LOD-.300)	.300 (.200-.400)	389
	09-10	*	< LOD	< LOD	.300 (<LOD-.400)	.400 (.300-.600)	415
12-19 years	05-06	*	< LOD	< LOD	< LOD	.200 (<LOD-.400)	702
	07-08	*	< LOD	< LOD	< LOD	.300 (<LOD-.400)	401
	09-10	*	< LOD	< LOD	.300 (.200-.300)	.300 (.300-.400)	420
20 years and older	05-06	*	< LOD	< LOD	< LOD	.200 (<LOD-.300)	1490
	07-08	*	< LOD	< LOD	< LOD	.300 (.200-.400)	1814
	09-10	*	< LOD	< LOD	< LOD	.300 (.200-.400)	1914
Gender							
Males	05-06	*	< LOD	< LOD	< LOD	.200 (<LOD-.300)	1270
	07-08	*	< LOD	< LOD	< LOD	.300 (.200-.400)	1294
	09-10	*	< LOD	< LOD	.200 (<LOD-.300)	.300 (.300-.400)	1399
Females	05-06	*	< LOD	< LOD	< LOD	< LOD	1278
	07-08	*	< LOD	< LOD	< LOD	.300 (.200-.300)	1310
	09-10	*	< LOD	< LOD	.200 (<LOD-.300)	.300 (.200-.400)	1350
Race/ethnicity							
Mexican Americans	05-06	*	< LOD	< LOD	< LOD	.300 (<LOD-.800)	637
	07-08	*	< LOD	< LOD	< LOD	.300 (.200-.300)	531
	09-10	*	< LOD	< LOD	< LOD	.300 (<LOD-.500)	566
Non-Hispanic blacks	05-06	*	< LOD	< LOD	< LOD	.200 (<LOD-.400)	678
	07-08	*	< LOD	< LOD	.200 (<LOD-.300)	.300 (.200-.500)	597
	09-10	*	< LOD	< LOD	.300 (.200-.400)	.500 (.400-.600)	516
Non-Hispanic whites	05-06	*	< LOD	< LOD	< LOD	.200 (<LOD-.300)	1038
	07-08	*	< LOD	< LOD	< LOD	.200 (.200-.300)	1077
	09-10	*	< LOD	< LOD	.200 (<LOD-.300)	.300 (.300-.400)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06, 07-08, and 09-10 are 0.2, 0.2, and 0.2 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**The 2003-2004 data for this chemical have been removed, due to potential for contamination that may have occurred during sampling.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Octylphenol_BiomonitoringSummary.html

Urinary 4-*tert*-Octylphenol (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	05-06	*	< LOD	< LOD	< LOD	.560 (<LOD-.620)	2548
	07-08	*	< LOD	< LOD	< LOD	.670 (.560-.880)	2604
	09-10	*	< LOD	< LOD	.470 (<LOD-.500)	.640 (.580-.680)	2749
Age group							
6-11 years	05-06	*	< LOD	< LOD	< LOD	.410 (.350-.610)	356
	07-08	*	< LOD	< LOD	.440 (<LOD-.540)	.600 (.420-.930)	389
	09-10	*	< LOD	< LOD	.560 (<LOD-.670)	.740 (.640-.830)	415
12-19 years	05-06	*	< LOD	< LOD	< LOD	.450 (<LOD-.560)	702
	07-08	*	< LOD	< LOD	< LOD	.420 (<LOD-.540)	401
	09-10	*	< LOD	< LOD	.360 (.270-.440)	.450 (.340-.560)	420
20 years and older	05-06	*	< LOD	< LOD	< LOD	.580 (<LOD-.670)	1490
	07-08	*	< LOD	< LOD	< LOD	.700 (.580-.930)	1814
	09-10	*	< LOD	< LOD	< LOD	.640 (.580-.700)	1914
Gender							
Males	05-06	*	< LOD	< LOD	< LOD	.390 (<LOD-.440)	1270
	07-08	*	< LOD	< LOD	< LOD	.510 (.390-.600)	1294
	09-10	*	< LOD	< LOD	.380 (<LOD-.420)	.480 (.450-.610)	1399
Females	05-06	*	< LOD	< LOD	< LOD	< LOD	1278
	07-08	*	< LOD	< LOD	< LOD	.880 (.670-1.08)	1310
	09-10	*	< LOD	< LOD	.540 (<LOD-.580)	.700 (.640-.880)	1350
Race/ethnicity							
Mexican Americans	05-06	*	< LOD	< LOD	< LOD	.480 (<LOD-.740)	637
	07-08	*	< LOD	< LOD	< LOD	.520 (.440-.610)	531
	09-10	*	< LOD	< LOD	< LOD	.560 (<LOD-.750)	566
Non-Hispanic blacks	05-06	*	< LOD	< LOD	< LOD	.320 (<LOD-.400)	678
	07-08	*	< LOD	< LOD	.320 (<LOD-.410)	.460 (.340-.610)	597
	09-10	*	< LOD	< LOD	.300 (.250-.330)	.440 (.320-.560)	516
Non-Hispanic whites	05-06	*	< LOD	< LOD	< LOD	.610 (<LOD-.670)	1038
	07-08	*	< LOD	< LOD	< LOD	.700 (.560-.930)	1077
	09-10	*	< LOD	< LOD	.500 (<LOD-.560)	.670 (.580-.740)	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**The 2003-2004 data for this chemical have been removed, due to potential for contamination that may have occurred during sampling.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Octylphenol_BiomonitoringSummary.html

Urinary Triclosan

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	13.0 (11.6-14.6)	9.20 (7.90-10.9)	47.4 (38.2-58.4)	249 (188-304)	461 (383-522)	2517
	05-06	18.5 (16.1-21.3)	15.1 (11.8-18.5)	76.2 (57.9-97.6)	334 (279-402)	655 (573-739)	2548
	07-08	15.3 (13.5-17.4)	12.1 (10.2-13.8)	57.2 (46.1-65.9)	225 (176-288)	494 (371-615)	2604
	09-10	14.5 (12.6-16.6)	10.7 (8.80-12.6)	51.2 (39.4-67.7)	238 (200-284)	483 (398-569)	2749
Age group							
6-11 years	03-04	8.16 (6.20-10.8)	6.00 (4.00-8.50)	20.7 (14.3-31.6)	123 (36.4-163)	157 (113-380)	314
	05-06	12.8 (9.89-16.7)	10.3 (8.30-17.2)	35.4 (23.9-65.8)	97.6 (67.4-181)	246 (99.5-462)	356
	07-08	11.8 (7.57-18.2)	9.80 (6.70-13.9)	27.7 (14.8-52.2)	98.5 (40.5-364)	296 (67.4-826)	389
	09-10	10.9 (9.35-12.8)	9.90 (7.10-11.9)	28.3 (22.1-35.5)	95.5 (71.5-117)	200 (114-474)	415
12-19 years	03-04	14.5 (11.0-19.1)	10.3 (8.20-13.1)	39.0 (26.5-86.4)	304 (134-566)	655 (310-890)	715
	05-06	18.8 (14.9-23.8)	15.4 (11.0-21.0)	67.5 (45.3-100)	330 (174-461)	566 (389-707)	702
	07-08	18.2 (13.8-23.8)	13.8 (9.40-20.1)	63.2 (38.7-110)	296 (144-395)	401 (308-853)	401
	09-10	11.7 (9.89-13.8)	8.80 (7.30-10.9)	30.2 (24.3-40.6)	165 (56.7-289)	301 (220-431)	420
20 years and older	03-04	13.6 (12.0-15.3)	9.60 (8.20-11.5)	51.7 (39.6-65.7)	261 (198-317)	472 (406-522)	1488
	05-06	19.3 (16.4-22.6)	15.5 (11.8-19.4)	84.3 (61.0-114)	366 (289-462)	738 (583-864)	1490
	07-08	15.4 (13.7-17.3)	12.3 (10.1-14.4)	60.1 (48.5-69.0)	225 (185-286)	504 (378-573)	1814
	09-10	15.5 (12.9-18.5)	11.1 (8.60-14.2)	61.8 (41.8-86.0)	262 (214-327)	544 (415-621)	1914
Gender							
Males	03-04	16.2 (13.4-19.6)	11.7 (9.30-14.8)	84.9 (50.6-111)	317 (231-433)	574 (461-716)	1229
	05-06	21.3 (17.6-25.7)	17.6 (11.9-23.2)	103 (69.9-143)	446 (366-488)	738 (601-873)	1270
	07-08	15.2 (12.9-17.9)	12.3 (9.50-15.3)	60.6 (45.8-72.8)	236 (159-338)	467 (367-636)	1294
	09-10	14.8 (12.7-17.4)	10.9 (8.60-13.3)	55.1 (40.4-77.5)	243 (214-295)	455 (327-600)	1399
Females	03-04	10.6 (9.29-12.1)	7.60 (6.10-9.10)	33.2 (27.1-39.4)	144 (96.5-250)	380 (258-430)	1288
	05-06	16.2 (13.9-18.8)	12.6 (10.1-15.6)	58.7 (41.5-81.9)	226 (169-304)	513 (310-773)	1278
	07-08	15.5 (12.6-18.9)	12.0 (9.90-14.1)	52.1 (37.1-74.4)	210 (133-367)	504 (285-648)	1310
	09-10	14.2 (12.0-16.8)	10.5 (8.70-12.6)	50.0 (34.4-63.8)	235 (149-302)	488 (332-661)	1350
Race/ethnicity							
Mexican Americans	03-04	14.6 (10.6-20.1)	8.80 (5.40-17.5)	65.4 (32.8-127)	357 (225-456)	597 (372-992)	613
	05-06	26.7 (21.2-33.7)	18.7 (13.5-25.5)	196 (99.4-269)	668 (475-759)	866 (750-1180)	637
	07-08	17.1 (12.9-22.6)	11.8 (8.40-17.8)	67.4 (42.5-106)	358 (208-474)	556 (363-856)	531
	09-10	14.9 (12.2-18.3)	10.2 (8.20-12.8)	54.7 (33.3-86.5)	345 (260-494)	691 (443-1180)	566
Non-Hispanic blacks	03-04	14.4 (11.4-18.2)	11.1 (8.70-16.1)	37.6 (30.2-58.0)	203 (87.5-341)	450 (254-750)	652
	05-06	17.3 (13.3-22.4)	14.0 (10.4-19.0)	59.2 (37.7-98.3)	258 (138-460)	541 (273-1190)	678
	07-08	13.7 (11.7-16.1)	11.3 (8.80-13.9)	41.4 (28.9-49.6)	150 (93.5-265)	480 (190-757)	597
	09-10	12.8 (10.8-15.0)	9.30 (7.70-11.7)	34.3 (24.0-44.6)	168 (88.4-263)	451 (202-959)	516
Non-Hispanic whites	03-04	12.9 (11.2-14.9)	9.20 (7.40-11.0)	49.2 (37.8-63.4)	245 (163-334)	461 (383-527)	1092
	05-06	17.5 (14.9-20.6)	15.1 (10.9-19.0)	74.3 (54.1-90.3)	288 (231-366)	569 (462-693)	1038
	07-08	15.0 (12.6-17.7)	12.3 (9.80-14.5)	59.6 (41.9-73.4)	197 (147-266)	408 (296-537)	1077
	09-10	14.0 (11.8-16.7)	10.5 (8.30-12.9)	51.7 (33.1-79.2)	216 (174-266)	431 (301-565)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 2.3, 2.3, 2.3, and 2.3 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Triclosan_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Triclosan_FactSheet.html

Urinary Triclosan (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	12.7 (11.5-14.1)	9.48 (8.22-10.4)	43.9 (33.8-60.6)	212 (172-241)	368 (294-463)	2514
	05-06	18.0 (16.0-20.3)	13.0 (11.5-16.1)	73.2 (57.3-91.6)	304 (240-364)	532 (434-674)	2548
	07-08	15.5 (13.7-17.5)	12.4 (10.6-14.2)	50.4 (40.2-59.8)	233 (171-300)	443 (330-559)	2604
	09-10	15.1 (13.2-17.4)	10.9 (9.41-12.4)	49.4 (37.0-67.3)	256 (191-322)	454 (352-557)	2749
Age group							
6-11 years	03-04	9.93 (7.43-13.3)	7.55 (4.72-13.4)	25.1 (15.3-35.6)	116 (39.9-236)	236 (115-336)	314
	05-06	14.1 (10.8-18.5)	13.5 (7.97-18.5)	38.6 (26.0-54.1)	108 (62.4-169)	241 (108-598)	356
	07-08	14.5 (9.47-22.1)	13.0 (8.00-17.5)	33.1 (17.5-72.4)	132 (52.7-331)	331 (131-599)	389
	09-10	14.2 (12.4-16.4)	12.0 (8.84-15.0)	37.1 (27.0-47.3)	129 (89.1-155)	300 (148-469)	415
12-19 years	03-04	10.9 (8.32-14.2)	7.45 (5.48-10.7)	31.8 (21.9-61.1)	193 (90.7-318)	356 (169-580)	713
	05-06	14.0 (11.0-17.8)	11.1 (8.68-13.6)	51.0 (33.0-73.4)	193 (125-306)	385 (203-739)	702
	07-08	14.1 (11.0-18.1)	11.0 (7.87-14.4)	50.2 (27.1-78.8)	206 (95.3-293)	378 (238-571)	401
	09-10	9.38 (7.86-11.2)	7.71 (5.71-9.21)	23.2 (17.3-29.3)	153 (63.0-201)	247 (166-379)	420
20 years and older	03-04	13.4 (12.0-15.1)	10.0 (8.89-11.4)	50.0 (36.0-73.8)	224 (186-272)	385 (308-506)	1487
	05-06	19.3 (16.9-22.0)	13.7 (11.6-17.0)	86.1 (64.0-109)	343 (262-411)	581 (440-718)	1490
	07-08	15.8 (14.0-17.8)	12.6 (10.7-14.5)	52.3 (43.8-64.6)	244 (186-309)	484 (336-568)	1814
	09-10	16.4 (13.7-19.6)	11.5 (9.63-14.8)	57.1 (38.4-80.2)	279 (202-364)	504 (372-601)	1914
Gender							
Males	03-04	13.2 (11.3-15.6)	9.21 (6.86-12.1)	73.1 (45.8-85.9)	237 (175-294)	384 (294-506)	1228
	05-06	17.1 (14.1-20.7)	12.4 (10.0-18.5)	82.6 (58.2-109)	308 (232-368)	472 (355-721)	1270
	07-08	12.8 (10.9-15.0)	10.0 (8.12-13.0)	45.7 (32.1-57.3)	191 (132-241)	330 (253-456)	1294
	09-10	13.3 (11.3-15.7)	9.21 (7.40-11.7)	43.9 (33.5-67.4)	230 (178-276)	345 (276-429)	1399
Females	03-04	12.2 (10.6-14.2)	9.54 (8.45-10.4)	32.3 (26.2-46.6)	182 (138-217)	336 (225-480)	1286
	05-06	19.0 (16.1-22.6)	13.8 (11.7-16.7)	64.0 (47.6-89.8)	301 (209-434)	619 (418-898)	1278
	07-08	18.5 (15.6-22.0)	14.4 (12.4-16.8)	55.3 (38.5-77.0)	300 (182-435)	571 (359-729)	1310
	09-10	17.1 (14.6-20.1)	12.3 (10.7-14.8)	50.6 (37.0-71.3)	300 (187-422)	556 (422-668)	1350
Race/ethnicity							
Mexican Americans	03-04	13.3 (9.38-18.8)	9.18 (5.45-13.9)	66.7 (28.8-112)	292 (151-432)	453 (263-1150)	612
	05-06	24.1 (19.3-29.9)	17.6 (13.3-22.8)	154 (96.4-242)	440 (379-601)	736 (601-818)	637
	07-08	16.6 (12.2-22.6)	12.6 (8.51-16.5)	60.1 (39.6-80.3)	325 (182-475)	637 (368-828)	531
	09-10	14.8 (12.1-18.2)	9.38 (7.63-12.4)	45.4 (32.2-110)	352 (244-458)	578 (402-737)	566
Non-Hispanic blacks	03-04	9.94 (7.92-12.5)	7.74 (5.50-10.0)	30.2 (25.6-37.3)	132 (78.0-213)	260 (127-513)	651
	05-06	12.2 (9.47-15.6)	9.50 (7.64-11.8)	41.0 (23.3-77.4)	179 (106-243)	352 (203-674)	678
	07-08	10.6 (8.91-12.7)	8.34 (6.47-10.1)	28.6 (22.3-36.7)	120 (71.0-203)	266 (151-407)	597
	09-10	9.23 (7.62-11.2)	6.62 (5.63-8.18)	22.7 (16.9-29.1)	136 (63.6-240)	356 (180-689)	516
Non-Hispanic whites	03-04	13.3 (11.6-15.1)	9.82 (8.11-11.5)	47.0 (34.3-67.7)	213 (160-272)	358 (276-480)	1091
	05-06	18.4 (16.0-21.2)	13.5 (11.6-17.0)	73.4 (56.8-97.0)	282 (231-343)	472 (367-699)	1038
	07-08	15.6 (13.1-18.7)	13.0 (10.7-14.8)	50.4 (35.9-65.9)	222 (129-303)	418 (262-586)	1077
	09-10	15.8 (13.1-19.0)	11.4 (9.41-14.8)	52.0 (37.0-71.3)	242 (159-323)	410 (317-547)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Triclosan_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Triclosan_FactSheet.html

Urinary *ortho*-Phenylphenol

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	.100 (<LOD-.200)	.300 (.200-.400)	.500 (.400-.700)	.800 (.600-1.10)	2525
	05-06	*	< LOD	< LOD	.300 (.200-.400)	.500 (.400-.800)	2548
	07-08	*	< LOD	< LOD	.100 (.100-.200)	.300 (.200-.400)	2604
	09-10	*	< LOD	< LOD	.300 (.300-.400)	.500 (.500-.600)	2749
Age group							
6-11 years	03-04	.180 (.158-.206)	.200 (.200-.200)	.300 (.300-.400)	.600 (.500-.800)	.800 (.500-1.30)	314
	05-06	*	< LOD	.100 (<LOD-.200)	.500 (.300-1.10)	1.30 (.700-1.90)	356
	07-08	*	< LOD	< LOD	.300 (.200-.500)	.600 (.300-1.00)	389
	09-10	*	< LOD	.200 (<LOD-.300)	.400 (.300-.700)	.600 (.400-1.20)	415
12-19 years	03-04	.175 (.149-.206)	.200 (.100-.200)	.400 (.300-.500)	.600 (.500-.800)	.900 (.700-1.30)	722
	05-06	*	< LOD	< LOD	.300 (.200-.500)	.600 (.500-.800)	702
	07-08	*	< LOD	< LOD	.100 (.100-.200)	.300 (.200-.600)	401
	09-10	*	< LOD	.200 (<LOD-.300)	.400 (.300-.500)	.600 (.500-.900)	420
20 years and older	03-04	*	.100 (<LOD-.100)	.300 (.200-.400)	.500 (.400-.700)	.800 (.600-1.10)	1489
	05-06	*	< LOD	< LOD	.300 (.100-.400)	.500 (.400-.800)	1490
	07-08	*	< LOD	< LOD	.100 (<LOD-.200)	.300 (.200-.400)	1814
	09-10	*	< LOD	< LOD	.300 (.300-.400)	.500 (.400-.500)	1914
Gender							
Males	03-04	*	.100 (<LOD-.200)	.400 (.300-.400)	.600 (.500-.700)	.800 (.600-1.00)	1231
	05-06	*	< LOD	< LOD	.300 (.200-.400)	.500 (.400-.900)	1270
	07-08	*	< LOD	< LOD	.100 (<LOD-.200)	.300 (.200-.500)	1294
	09-10	*	< LOD	< LOD	.300 (.300-.400)	.500 (.500-.700)	1399
Females	03-04	*	.100 (<LOD-.200)	.300 (.200-.400)	.500 (.400-.700)	.800 (.600-1.20)	1294
	05-06	*	< LOD	< LOD	.300 (.200-.400)	.500 (.400-.800)	1278
	07-08	*	< LOD	< LOD	.100 (<LOD-.200)	.300 (.200-.500)	1310
	09-10	*	< LOD	< LOD	.300 (.300-.400)	.500 (.400-.500)	1350
Race/ethnicity							
Mexican Americans	03-04	.191 (.150-.245)	.200 (.100-.300)	.400 (.300-.600)	.700 (.500-1.10)	1.10 (.600-1.60)	617
	05-06	*	< LOD	< LOD	.400 (.100-.900)	.800 (.400-3.20)	637
	07-08	*	< LOD	< LOD	.100 (<LOD-.300)	.400 (.200-1.10)	531
	09-10	*	< LOD	< LOD	.300 (.200-.400)	.400 (.300-.900)	566
Non-Hispanic blacks	03-04	*	.100 (<LOD-.200)	.400 (.300-.400)	.600 (.500-.800)	1.00 (.600-1.40)	636
	05-06	*	< LOD	< LOD	.400 (.300-.500)	.500 (.400-.700)	678
	07-08	*	< LOD	< LOD	.200 (<LOD-.300)	.400 (.200-.600)	597
	09-10	*	< LOD	.200 (<LOD-.300)	.400 (.300-.600)	.700 (.500-.800)	516
Non-Hispanic whites	03-04	*	.100 (<LOD-.100)	.300 (.200-.400)	.500 (.400-.700)	.800 (.600-1.00)	1077
	05-06	*	< LOD	< LOD	.300 (.100-.400)	.500 (.300-1.10)	1038
	07-08	*	< LOD	< LOD	.100 (<LOD-.200)	.300 (.200-.500)	1077
	09-10	*	< LOD	< LOD	.300 (.300-.400)	.500 (.400-.600)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.1, 0.1, 0.1, and 0.2 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Orthophenylphenol_BiomonitoringSummary.html

Urinary *ortho*-Phenylphenol (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	.150 (<LOD-.170)	.320 (.270-.380)	.670 (.500-.800)	.950 (.750-1.31)	2522
	05-06	*	< LOD	< LOD	.370 (.310-.410)	.600 (.470-.750)	2548
	07-08	*	< LOD	< LOD	.290 (.250-.330)	.470 (.400-.520)	2604
	09-10	*	< LOD	< LOD	.560 (.480-.640)	.820 (.700-.910)	2749
Age group							
6-11 years	03-04	.219 (.189-.255)	.230 (.170-.280)	.400 (.330-.540)	.760 (.530-.860)	.880 (.760-1.90)	314
	05-06	*	< LOD	.180 (<LOD-.240)	.590 (.430-.930)	1.94 (.710-2.94)	356
	07-08	*	< LOD	< LOD	.390 (.240-.660)	.660 (.390-1.31)	389
	09-10	*	< LOD	.410 (<LOD-.480)	.820 (.560-1.06)	1.22 (.850-1.56)	415
12-19 years	03-04	.131 (.110-.156)	.140 (.100-.160)	.270 (.230-.340)	.490 (.400-.690)	.790 (.550-1.30)	720
	05-06	*	< LOD	< LOD	.280 (.220-.320)	.410 (.330-.710)	702
	07-08	*	< LOD	< LOD	.190 (.170-.250)	.330 (.240-.470)	401
	09-10	*	< LOD	.250 (<LOD-.300)	.420 (.300-.550)	.550 (.390-1.20)	420
20 years and older	03-04	*	.140 (<LOD-.170)	.320 (.260-.370)	.670 (.500-.800)	1.00 (.740-1.38)	1488
	05-06	*	< LOD	< LOD	.370 (.300-.410)	.540 (.430-.710)	1490
	07-08	*	< LOD	< LOD	.300 (<LOD-.330)	.470 (.390-.520)	1814
	09-10	*	< LOD	< LOD	.560 (.480-.640)	.820 (.700-.880)	1914
Gender							
Males	03-04	*	.130 (<LOD-.150)	.290 (.240-.340)	.570 (.430-.740)	.880 (.720-1.06)	1230
	05-06	*	< LOD	< LOD	.280 (.200-.350)	.470 (.350-.670)	1270
	07-08	*	< LOD	< LOD	.220 (<LOD-.250)	.350 (.300-.440)	1294
	09-10	*	< LOD	< LOD	.440 (.360-.540)	.670 (.510-.850)	1399
Females	03-04	*	.170 (<LOD-.200)	.350 (.280-.440)	.700 (.500-.940)	1.07 (.730-1.57)	1292
	05-06	*	< LOD	< LOD	.430 (.370-.500)	.700 (.500-1.03)	1278
	07-08	*	< LOD	< LOD	.350 (<LOD-.440)	.500 (.440-.580)	1310
	09-10	*	< LOD	< LOD	.640 (.570-.700)	.930 (.780-1.17)	1350
Race/ethnicity							
Mexican Americans	03-04	.174 (.132-.229)	.170 (.120-.240)	.440 (.280-.590)	.770 (.560-1.09)	1.11 (.770-1.72)	616
	05-06	*	< LOD	< LOD	.370 (.210-.670)	.740 (.390-1.71)	637
	07-08	*	< LOD	< LOD	.250 (<LOD-.290)	.430 (.320-.910)	531
	09-10	*	< LOD	< LOD	.420 (.370-.560)	.680 (.560-.880)	566
Non-Hispanic blacks	03-04	*	.110 (<LOD-.130)	.250 (.200-.310)	.530 (.380-.650)	.770 (.600-1.02)	635
	05-06	*	< LOD	< LOD	.290 (.220-.370)	.440 (.380-.540)	678
	07-08	*	< LOD	< LOD	.210 (<LOD-.250)	.330 (.250-.640)	597
	09-10	*	< LOD	.220 (<LOD-.240)	.370 (.310-.450)	.570 (.440-.860)	516
Non-Hispanic whites	03-04	*	.150 (<LOD-.180)	.330 (.260-.390)	.690 (.470-.830)	1.00 (.740-1.38)	1076
	05-06	*	< LOD	< LOD	.390 (.320-.430)	.610 (.470-.880)	1038
	07-08	*	< LOD	< LOD	.300 (<LOD-.370)	.470 (.390-.580)	1077
	09-10	*	< LOD	< LOD	.610 (.520-.700)	.870 (.700-.940)	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Orthophenylphenol_BiomonitoringSummary.html

Urinary Ethylene thiourea

Metabolite of Several Dithiocarbamate Fungicides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	.310 (<LOD-.560)	2169
	05-06	*	< LOD	< LOD	.600 (<LOD-1.08)	1.34 (.900-1.75)	2585
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	280
	05-06	*	< LOD	< LOD	.830 (<LOD-1.53)	1.55 (.960-3.11)	342
12-19 years	03-04	*	< LOD	< LOD	< LOD	.600 (<LOD-.920)	622
	05-06	*	< LOD	< LOD	.580 (<LOD-1.11)	1.34 (.660-2.23)	698
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1267
	05-06	*	< LOD	< LOD	.590 (<LOD-1.09)	1.32 (.880-1.66)	1545
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	.360 (<LOD-.530)	1056
	05-06	*	< LOD	< LOD	.940 (.530-1.19)	1.51 (1.08-2.09)	1219
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1113
	05-06	*	< LOD	< LOD	.370 (<LOD-.890)	.950 (.500-1.58)	1366
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	495
	05-06	*	< LOD	< LOD	.440 (<LOD-.960)	1.12 (.650-1.76)	686
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	.480 (<LOD-1.06)	607
	05-06	*	< LOD	< LOD	1.37 (.790-2.05)	2.22 (1.37-3.77)	695
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	.330 (<LOD-.560)	925
	05-06	*	< LOD	< LOD	.570 (<LOD-1.02)	1.18 (.940-1.42)	998

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.1 and 0.24 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ETUPTU_BiomonitoringSummary.html

Urinary Ethylene thiourea (creatinine corrected)

Metabolite of Several Dithiocarbamate Fungicides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	.434 (<LOD-.538)	2167
	05-06	*	< LOD	< LOD	.792 (<LOD-1.06)	1.32 (1.06-1.58)	2585
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	280
	05-06	*	< LOD	< LOD	1.04 (<LOD-1.36)	2.36 (1.16-3.12)	342
12-19 years	03-04	*	< LOD	< LOD	< LOD	.423 (<LOD-.700)	621
	05-06	*	< LOD	< LOD	.582 (<LOD-.893)	.967 (.680-1.58)	698
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1266
	05-06	*	< LOD	< LOD	.792 (<LOD-1.06)	1.32 (1.06-1.49)	1545
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	.350 (<LOD-.467)	1055
	05-06	*	< LOD	< LOD	.783 (.634-1.04)	1.28 (1.05-1.45)	1219
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1112
	05-06	*	< LOD	< LOD	.827 (<LOD-1.21)	1.33 (1.00-2.01)	1366
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	494
	05-06	*	< LOD	< LOD	.748 (<LOD-1.17)	1.33 (.837-1.89)	686
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	.368 (<LOD-.619)	607
	05-06	*	< LOD	< LOD	1.02 (.649-1.74)	1.84 (1.04-2.50)	695
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	.434 (<LOD-.583)	924
	05-06	*	< LOD	< LOD	.773 (<LOD-1.04)	1.25 (.968-1.48)	998

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ETUPTU_BiomonitoringSummary.html

Urinary Pentachlorophenol

Also a Metabolite of Several Organochlorine Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	1.12 (.570-1.58)	2.57 (2.08-2.99)	3.63 (2.98-4.61)	2354
Age group							
6-11 years	03-04	*	< LOD	1.57 (.970-2.25)	3.23 (2.12-5.67)	5.67 (2.94-6.38)	290
12-19 years	03-04	*	< LOD	1.36 (.760-1.99)	2.88 (2.08-3.53)	3.80 (3.06-6.38)	674
20 years and older	03-04	*	< LOD	1.00 (<LOD-1.47)	2.41 (1.92-2.84)	3.40 (2.95-3.81)	1390
Gender							
Males	03-04	*	< LOD	1.32 (.720-2.05)	2.79 (2.46-3.40)	4.58 (3.50-5.49)	1147
Females	03-04	*	< LOD	.880 (<LOD-1.41)	2.12 (1.71-2.74)	3.20 (2.44-3.84)	1207
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	1.73 (.640-3.38)	2.44 (1.73-3.84)	550
Non-Hispanic blacks	03-04	*	< LOD	1.46 (1.11-1.93)	2.79 (2.22-3.65)	4.06 (3.20-4.77)	610
Non-Hispanic whites	03-04	*	< LOD	1.17 (.580-1.74)	2.66 (2.06-3.23)	3.69 (2.99-5.17)	1039

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 0.5.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Pentachlorophenol_BiomonitoringSummary.html

Urinary Pentachlorophenol (creatinine corrected)

Also a Metabolite of Several Organochlorine Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	1.22 (1.01-1.52)	2.30 (1.84-2.77)	3.44 (2.69-3.96)	2352
Age group							
6-11 years	03-04	*	< LOD	1.75 (1.38-2.69)	3.72 (2.50-4.96)	4.96 (3.59-10.6)	290
12-19 years	03-04	*	< LOD	1.11 (.800-1.28)	1.67 (1.31-2.65)	2.76 (1.64-3.89)	673
20 years and older	03-04	*	< LOD	1.19 (<LOD-1.50)	2.19 (1.86-2.64)	3.24 (2.54-3.96)	1389
Gender							
Males	03-04	*	< LOD	1.10 (.825-1.38)	1.93 (1.62-2.65)	3.23 (2.06-4.94)	1146
Females	03-04	*	< LOD	1.37 (<LOD-1.74)	2.50 (2.13-2.98)	3.50 (2.79-4.07)	1206
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	1.54 (1.01-2.69)	2.33 (1.40-4.62)	549
Non-Hispanic blacks	03-04	*	< LOD	.919 (.679-1.25)	1.88 (1.42-2.77)	2.94 (2.21-3.76)	610
Non-Hispanic whites	03-04	*	< LOD	1.35 (1.08-1.64)	2.42 (1.94-3.18)	3.54 (2.77-4.78)	1038

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Pentachlorophenol_BiomonitoringSummary.html

Urinary Propylene thiourea

Metabolite of Propineb

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2456
	05-06	*	< LOD	< LOD	< LOD	< LOD	2628
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	305
	05-06	*	< LOD	< LOD	< LOD	< LOD	349
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	702
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1449
	05-06	*	< LOD	< LOD	< LOD	< LOD	1561
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1209
	05-06	*	< LOD	< LOD	< LOD	< LOD	1246
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1247
	05-06	*	< LOD	< LOD	< LOD	< LOD	1382
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	598
	05-06	*	< LOD	< LOD	< LOD	< LOD	693
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	637
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1062
	05-06	*	< LOD	< LOD	< LOD	< LOD	1006

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.1 and 0.37 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ETUPTU_BiomonitoringSummary.html

Urinary Propylene thiourea (creatinine corrected)

Metabolite of Propineb

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2454
	05-06	*	< LOD	< LOD	< LOD	< LOD	2628
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	305
	05-06	*	< LOD	< LOD	< LOD	< LOD	349
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	701
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1448
	05-06	*	< LOD	< LOD	< LOD	< LOD	1561
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1208
	05-06	*	< LOD	< LOD	< LOD	< LOD	1246
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1246
	05-06	*	< LOD	< LOD	< LOD	< LOD	1382
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	597
	05-06	*	< LOD	< LOD	< LOD	< LOD	693
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	637
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1061
	05-06	*	< LOD	< LOD	< LOD	< LOD	1006

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ETUPTU_BiomonitoringSummary.html

Urinary 2,4-Dichlorophenoxyacetic acid

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1977
	01-02	*	< LOD	.230 (<LOD-.320)	.690 (.560-.910)	1.27 (1.02-1.37)	2413
	03-04	.245 (.210-.285)	.240 (.180-.320)	.580 (.490-.670)	.990 (.890-1.31)	1.63 (1.31-2.37)	1960
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	1.30 (<LOD-2.40)	477
	01-02	*	< LOD	.310 (.210-.400)	.740 (.550-1.13)	1.55 (1.00-2.21)	546
	03-04	.266 (.214-.332)	.290 (.200-.390)	.670 (.440-.920)	1.03 (.890-1.40)	1.88 (1.01-2.54)	309
12-19 years	99-00	*	< LOD	< LOD	< LOD	1.10 (<LOD-1.60)	677
	01-02	*	< LOD	.250 (<LOD-.420)	.690 (.440-1.16)	1.24 (.690-1.66)	797
	03-04	.256 (.212-.310)	.260 (.180-.380)	.580 (.470-.710)	1.04 (.890-1.31)	1.66 (1.20-2.97)	714
20-59 years	99-00	*	< LOD	< LOD	< LOD	< LOD	823
	01-02	*	< LOD	.210 (<LOD-.310)	.690 (.540-.910)	1.27 (.930-1.49)	1070
	03-04	.239 (.205-.279)	.220 (.170-.300)	.570 (.480-.640)	.980 (.840-1.35)	1.55 (1.25-2.50)	937
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	1.10 (<LOD-1.80)	962
	01-02	*	< LOD	.330 (.230-.490)	.930 (.680-1.22)	1.51 (1.27-2.08)	1135
	03-04	.277 (.240-.319)	.290 (.210-.380)	.620 (.540-.740)	1.16 (.910-1.40)	1.98 (1.41-2.54)	943
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1015
	01-02	*	< LOD	< LOD	.490 (.370-.660)	.890 (.670-1.22)	1278
	03-04	.217 (.180-.261)	.190 (.120-.280)	.480 (.400-.640)	.960 (.800-1.22)	1.44 (1.10-2.29)	1017
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	695
	01-02	*	< LOD	.260 (<LOD-.350)	.730 (.610-.910)	1.20 (.960-1.43)	659
	03-04	.320 (.263-.389)	.350 (.260-.450)	.730 (.610-.840)	1.42 (1.01-1.52)	1.78 (1.22-4.47)	495
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.70)	520
	01-02	*	< LOD	< LOD	.610 (.420-.890)	1.07 (.810-1.48)	668
	03-04	*	.190 (<LOD-.280)	.480 (.370-.600)	.890 (.740-1.10)	1.25 (.910-2.21)	579
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	589
	01-02	*	< LOD	.250 (<LOD-.410)	.760 (.560-1.10)	1.32 (1.05-2.03)	892
	03-04	.252 (.209-.304)	.240 (.170-.370)	.590 (.470-.720)	1.10 (.860-1.47)	1.68 (1.26-2.73)	755

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 0.89, 0.2, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/2,4-DichlorophenoxyaceticAcid_BiomonitoringSummary.html

Urinary 2,4-Dichlorophenoxyacetic acid (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1977
	01-02	*	< LOD	.378 (<LOD-.414)	.667 (.609-.778)	1.08 (.926-1.26)	2412
	03-04	.230 (.193-.274)	.243 (.198-.279)	.469 (.414-.557)	.962 (.797-1.20)	1.58 (1.24-2.34)	1958
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	1.32 (<LOD-2.24)	477
	01-02	*	< LOD	.485 (.378-.679)	1.13 (.825-1.35)	1.41 (1.27-1.73)	546
	03-04	.323 (.249-.421)	.320 (.250-.440)	.744 (.500-1.06)	1.30 (.990-2.55)	2.55 (1.23-5.16)	309
12-19 years	99-00	*	< LOD	< LOD	< LOD	.593 (<LOD-1.05)	677
	01-02	*	< LOD	.275 (<LOD-.376)	.483 (.328-.662)	.662 (.517-.918)	796
	03-04	.193 (.160-.232)	.205 (.157-.250)	.419 (.328-.460)	.709 (.540-.925)	1.23 (.837-2.35)	713
20-59 years	99-00	*	< LOD	< LOD	< LOD	< LOD	823
	01-02	*	< LOD	.378 (<LOD-.412)	.667 (.593-.778)	1.08 (.806-1.29)	1070
	03-04	.227 (.188-.274)	.242 (.196-.278)	.452 (.397-.545)	.923 (.708-1.20)	1.48 (1.14-2.43)	936
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	.667 (<LOD-1.16)	962
	01-02	*	< LOD	.336 (.268-.412)	.640 (.564-.789)	1.14 (.893-1.39)	1135
	03-04	.222 (.183-.268)	.234 (.193-.273)	.442 (.392-.531)	.913 (.718-1.11)	1.48 (.969-2.79)	942
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1015
	01-02	*	< LOD	< LOD	.700 (.609-.778)	1.08 (.809-1.26)	1277
	03-04	.239 (.199-.287)	.254 (.200-.297)	.489 (.412-.615)	1.00 (.737-1.26)	1.64 (1.23-2.43)	1016
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	695
	01-02	*	< LOD	.350 (<LOD-.386)	.720 (.562-.853)	1.13 (.778-1.56)	659
	03-04	.287 (.223-.369)	.318 (.194-.461)	.582 (.460-.762)	1.07 (.771-1.36)	1.48 (1.08-3.19)	494
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	.593 (<LOD-1.19)	520
	01-02	*	< LOD	< LOD	.438 (.337-.568)	.778 (.552-.975)	667
	03-04	*	.130 (<LOD-.169)	.294 (.250-.331)	.531 (.423-.692)	.938 (.612-1.34)	579
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	589
	01-02	*	< LOD	.412 (<LOD-.467)	.737 (.616-.894)	1.17 (.988-1.40)	892
	03-04	.250 (.201-.311)	.259 (.212-.304)	.500 (.415-.631)	1.03 (.842-1.34)	1.90 (1.30-2.55)	754

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/2,4-DichlorophenoxyaceticAcid_BiomonitoringSummary.html

Urinary 2,4,5-Trichlorophenoxyacetic acid

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1814
	01-02	*	< LOD	< LOD	< LOD	< LOD	2538
	03-04	*	< LOD	< LOD	< LOD	.150 (<LOD-.210)	1912
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	< LOD	430
	01-02	*	< LOD	< LOD	< LOD	< LOD	580
	03-04	*	< LOD	< LOD	< LOD	.110 (<LOD-.160)	303
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	618
	01-02	*	< LOD	< LOD	< LOD	< LOD	831
	03-04	*	< LOD	< LOD	< LOD	.140 (<LOD-.190)	701
20-59 years	99-00	*	< LOD	< LOD	< LOD	< LOD	766
	01-02	*	< LOD	< LOD	< LOD	< LOD	1127
	03-04	*	< LOD	< LOD	< LOD	.160 (<LOD-.250)	908
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	891
	01-02	*	< LOD	< LOD	< LOD	< LOD	1192
	03-04	*	< LOD	< LOD	< LOD	.140 (<LOD-.220)	920
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	923
	01-02	*	< LOD	< LOD	< LOD	< LOD	1346
	03-04	*	< LOD	< LOD	< LOD	.150 (<LOD-.210)	992
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	652
	01-02	*	< LOD	< LOD	< LOD	< LOD	679
	03-04	*	< LOD	< LOD	.120 (<LOD-.340)	.200 (<LOD-.390)	494
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	483
	01-02	*	< LOD	< LOD	< LOD	< LOD	701
	03-04	*	< LOD	< LOD	< LOD	< LOD	572
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	531
	01-02	*	< LOD	< LOD	< LOD	< LOD	957
	03-04	*	< LOD	< LOD	< LOD	.160 (<LOD-.230)	716

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 1.1, 0.1, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/2,4,5-TrichlorophenoxyaceticAcid_BiomonitoringSummary.html

Urinary 2,4,5-Trichlorophenoxyacetic acid (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1814
	01-02	*	< LOD	< LOD	< LOD	< LOD	2537
	03-04	*	< LOD	< LOD	< LOD	.333 (<LOD-.412)	1910
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	< LOD	430
	01-02	*	< LOD	< LOD	< LOD	< LOD	580
	03-04	*	< LOD	< LOD	< LOD	.318 (<LOD-.368)	303
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	618
	01-02	*	< LOD	< LOD	< LOD	< LOD	830
	03-04	*	< LOD	< LOD	< LOD	.226 (<LOD-.269)	700
20-59 years	99-00	*	< LOD	< LOD	< LOD	< LOD	766
	01-02	*	< LOD	< LOD	< LOD	< LOD	1127
	03-04	*	< LOD	< LOD	< LOD	.350 (<LOD-.438)	907
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	891
	01-02	*	< LOD	< LOD	< LOD	< LOD	1192
	03-04	*	< LOD	< LOD	< LOD	.250 (<LOD-.350)	919
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	923
	01-02	*	< LOD	< LOD	< LOD	< LOD	1345
	03-04	*	< LOD	< LOD	< LOD	.368 (<LOD-.500)	991
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	652
	01-02	*	< LOD	< LOD	< LOD	< LOD	679
	03-04	*	< LOD	< LOD	.193 (<LOD-.333)	.318 (<LOD-.467)	493
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	483
	01-02	*	< LOD	< LOD	< LOD	< LOD	700
	03-04	*	< LOD	< LOD	< LOD	< LOD	572
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	531
	01-02	*	< LOD	< LOD	< LOD	< LOD	957
	03-04	*	< LOD	< LOD	< LOD	.350 (<LOD-.436)	715

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/2,4,5-TrichlorophenoxyaceticAcid_BiomonitoringSummary.html

Urinary Bensulfuron-methyl

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2355
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2494
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	286
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	374
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	671
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	381
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1398
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1739
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1150
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1240
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1205
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1254
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	567
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	477
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	617
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	558
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1018
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1053

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.05, 0.05, and 0.05 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Bensulfuron-methyl (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2353
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2492
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	286
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	374
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	670
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	379
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1397
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1739
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1149
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1239
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1204
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1253
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	566
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	476
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	617
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	557
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1017
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1053

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Chlorsulfuron

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2100
	05-06	*	< LOD	< LOD	< LOD	< LOD	2419
	07-08	*	< LOD	< LOD	< LOD	< LOD	2566
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	259
	05-06	*	< LOD	< LOD	< LOD	< LOD	314
	07-08	*	< LOD	< LOD	< LOD	< LOD	380
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	601
	05-06	*	< LOD	< LOD	< LOD	< LOD	653
	07-08	*	< LOD	< LOD	< LOD	< LOD	386
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1240
	05-06	*	< LOD	< LOD	< LOD	< LOD	1452
	07-08	*	< LOD	< LOD	< LOD	< LOD	1800
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1031
	05-06	*	< LOD	< LOD	< LOD	< LOD	1151
	07-08	*	< LOD	< LOD	< LOD	< LOD	1285
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1069
	05-06	*	< LOD	< LOD	< LOD	< LOD	1268
	07-08	*	< LOD	< LOD	< LOD	< LOD	1281
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	506
	05-06	*	< LOD	< LOD	< LOD	< LOD	633
	07-08	*	< LOD	< LOD	< LOD	< LOD	491
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	568
	05-06	*	< LOD	< LOD	< LOD	< LOD	658
	07-08	*	< LOD	< LOD	< LOD	< LOD	570
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	894
	05-06	*	< LOD	< LOD	< LOD	< LOD	940
	07-08	*	< LOD	< LOD	< LOD	< LOD	1081

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.06, 0.06, and 0.06 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Chlorsulfuron (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2098
	05-06	*	< LOD	< LOD	< LOD	< LOD	2419
	07-08	*	< LOD	< LOD	< LOD	< LOD	2564
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	259
	05-06	*	< LOD	< LOD	< LOD	< LOD	314
	07-08	*	< LOD	< LOD	< LOD	< LOD	380
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	600
	05-06	*	< LOD	< LOD	< LOD	< LOD	653
	07-08	*	< LOD	< LOD	< LOD	< LOD	384
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1239
	05-06	*	< LOD	< LOD	< LOD	< LOD	1452
	07-08	*	< LOD	< LOD	< LOD	< LOD	1800
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1030
	05-06	*	< LOD	< LOD	< LOD	< LOD	1151
	07-08	*	< LOD	< LOD	< LOD	< LOD	1284
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1068
	05-06	*	< LOD	< LOD	< LOD	< LOD	1268
	07-08	*	< LOD	< LOD	< LOD	< LOD	1280
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	505
	05-06	*	< LOD	< LOD	< LOD	< LOD	633
	07-08	*	< LOD	< LOD	< LOD	< LOD	490
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	568
	05-06	*	< LOD	< LOD	< LOD	< LOD	658
	07-08	*	< LOD	< LOD	< LOD	< LOD	569
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	893
	05-06	*	< LOD	< LOD	< LOD	< LOD	940
	07-08	*	< LOD	< LOD	< LOD	< LOD	1081

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Ethametsulfuron-methyl

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2321
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2458
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	294
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	671
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	381
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1356
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1711
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1132
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1218
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1189
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1240
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	590
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	468
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	608
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	534
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	974
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1052

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04, 05-06, and 07-08 are 0.1, 0.1, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Ethametsulfuron-methyl (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2319
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2456
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	294
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	670
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	379
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1355
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1711
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1131
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1217
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1188
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1239
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	589
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	467
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	608
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	533
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	973
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1052

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Foramsulfuron

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2255
	05-06	*	< LOD	< LOD	< LOD	< LOD	2534
	07-08	*	< LOD	< LOD	< LOD	< LOD	2423
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	283
	05-06	*	< LOD	< LOD	< LOD	< LOD	334
	07-08	*	< LOD	< LOD	< LOD	< LOD	364
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	653
	05-06	*	< LOD	< LOD	< LOD	< LOD	688
	07-08	*	< LOD	< LOD	< LOD	< LOD	364
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1319
	05-06	*	< LOD	< LOD	< LOD	< LOD	1512
	07-08	*	< LOD	< LOD	< LOD	< LOD	1695
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1097
	05-06	*	< LOD	< LOD	< LOD	< LOD	1203
	07-08	*	< LOD	< LOD	< LOD	< LOD	1205
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1158
	05-06	*	< LOD	< LOD	< LOD	< LOD	1331
	07-08	*	< LOD	< LOD	< LOD	< LOD	1218
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	587
	05-06	*	< LOD	< LOD	< LOD	< LOD	632
	07-08	*	< LOD	< LOD	< LOD	< LOD	487
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	584
	05-06	*	< LOD	< LOD	< LOD	< LOD	711
	07-08	*	< LOD	< LOD	< LOD	< LOD	523
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	935
	05-06	*	< LOD	< LOD	< LOD	< LOD	990
	07-08	*	< LOD	< LOD	< LOD	< LOD	1004

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.05, 0.05, and 0.05 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Foramsulfuron (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2253
	05-06	*	< LOD	< LOD	< LOD	< LOD	2534
	07-08	*	< LOD	< LOD	< LOD	< LOD	2421
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	283
	05-06	*	< LOD	< LOD	< LOD	< LOD	334
	07-08	*	< LOD	< LOD	< LOD	< LOD	364
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	652
	05-06	*	< LOD	< LOD	< LOD	< LOD	688
	07-08	*	< LOD	< LOD	< LOD	< LOD	362
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1318
	05-06	*	< LOD	< LOD	< LOD	< LOD	1512
	07-08	*	< LOD	< LOD	< LOD	< LOD	1695
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1096
	05-06	*	< LOD	< LOD	< LOD	< LOD	1203
	07-08	*	< LOD	< LOD	< LOD	< LOD	1204
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1157
	05-06	*	< LOD	< LOD	< LOD	< LOD	1331
	07-08	*	< LOD	< LOD	< LOD	< LOD	1217
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	586
	05-06	*	< LOD	< LOD	< LOD	< LOD	632
	07-08	*	< LOD	< LOD	< LOD	< LOD	486
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	584
	05-06	*	< LOD	< LOD	< LOD	< LOD	711
	07-08	*	< LOD	< LOD	< LOD	< LOD	522
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	934
	05-06	*	< LOD	< LOD	< LOD	< LOD	990
	07-08	*	< LOD	< LOD	< LOD	< LOD	1004

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Halosulfuron

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2356
	05-06	*	< LOD	< LOD	< LOD	< LOD	2485
	07-08	*	< LOD	< LOD	< LOD	< LOD	2530
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	291
	05-06	*	< LOD	< LOD	< LOD	< LOD	331
	07-08	*	< LOD	< LOD	< LOD	< LOD	377
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	672
	05-06	*	< LOD	< LOD	< LOD	< LOD	679
	07-08	*	< LOD	< LOD	< LOD	< LOD	385
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1393
	05-06	*	< LOD	< LOD	< LOD	< LOD	1475
	07-08	*	< LOD	< LOD	< LOD	< LOD	1768
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1148
	05-06	*	< LOD	< LOD	< LOD	< LOD	1184
	07-08	*	< LOD	< LOD	< LOD	< LOD	1261
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1208
	05-06	*	< LOD	< LOD	< LOD	< LOD	1301
	07-08	*	< LOD	< LOD	< LOD	< LOD	1269
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	592
	05-06	*	< LOD	< LOD	< LOD	< LOD	650
	07-08	*	< LOD	< LOD	< LOD	< LOD	485
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	609
	05-06	*	< LOD	< LOD	< LOD	< LOD	694
	07-08	*	< LOD	< LOD	< LOD	< LOD	565
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1005
	05-06	*	< LOD	< LOD	< LOD	< LOD	950
	07-08	*	< LOD	< LOD	< LOD	< LOD	1068

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.1, 0.1, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Halosulfuron (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2354
	05-06	*	< LOD	< LOD	< LOD	< LOD	2485
	07-08	*	< LOD	< LOD	< LOD	< LOD	2528
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	291
	05-06	*	< LOD	< LOD	< LOD	< LOD	331
	07-08	*	< LOD	< LOD	< LOD	< LOD	377
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	671
	05-06	*	< LOD	< LOD	< LOD	< LOD	679
	07-08	*	< LOD	< LOD	< LOD	< LOD	383
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1392
	05-06	*	< LOD	< LOD	< LOD	< LOD	1475
	07-08	*	< LOD	< LOD	< LOD	< LOD	1768
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1147
	05-06	*	< LOD	< LOD	< LOD	< LOD	1184
	07-08	*	< LOD	< LOD	< LOD	< LOD	1260
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1207
	05-06	*	< LOD	< LOD	< LOD	< LOD	1301
	07-08	*	< LOD	< LOD	< LOD	< LOD	1268
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	591
	05-06	*	< LOD	< LOD	< LOD	< LOD	650
	07-08	*	< LOD	< LOD	< LOD	< LOD	484
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	609
	05-06	*	< LOD	< LOD	< LOD	< LOD	694
	07-08	*	< LOD	< LOD	< LOD	< LOD	564
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1004
	05-06	*	< LOD	< LOD	< LOD	< LOD	950
	07-08	*	< LOD	< LOD	< LOD	< LOD	1068

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Mesosulfuron-methyl

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2427
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2494
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	302
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	372
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	694
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	371
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1431
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1751
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1189
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1245
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1238
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1249
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	596
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	465
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	634
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	554
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1039
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1059

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.06, 0.06, and 0.06 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Mesosulfuron-methyl (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2425
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2492
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	302
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	372
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	693
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	369
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1430
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1751
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1188
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1244
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1237
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1248
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	595
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	464
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	634
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	553
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1038
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1059

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Metsulfuron-methyl

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2360
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2494
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	297
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	373
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	675
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	374
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1388
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1747
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1152
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1246
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1208
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1248
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	595
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	470
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	596
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	538
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1013
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1066

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.05, 0.05, and 0.05 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Metsulfuron-methyl (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2358
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2492
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	297
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	373
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	674
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	372
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1387
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1747
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1151
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1245
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1207
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1247
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	594
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	469
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	596
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	537
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1012
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1066

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Nicosulfuron

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2322
	05-06	*	< LOD	< LOD	< LOD	< LOD	2570
	07-08	*	< LOD	< LOD	< LOD	< LOD	2422
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	296
	05-06	*	< LOD	< LOD	< LOD	< LOD	341
	07-08	*	< LOD	< LOD	< LOD	< LOD	363
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	666
	05-06	*	< LOD	< LOD	< LOD	< LOD	699
	07-08	*	< LOD	< LOD	< LOD	< LOD	364
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1360
	05-06	*	< LOD	< LOD	< LOD	< LOD	1530
	07-08	*	< LOD	< LOD	< LOD	< LOD	1695
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1131
	05-06	*	< LOD	< LOD	< LOD	< LOD	1222
	07-08	*	< LOD	< LOD	< LOD	< LOD	1210
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1191
	05-06	*	< LOD	< LOD	< LOD	< LOD	1348
	07-08	*	< LOD	< LOD	< LOD	< LOD	1212
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	589
	05-06	*	< LOD	< LOD	< LOD	< LOD	656
	07-08	*	< LOD	< LOD	< LOD	< LOD	486
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	601
	05-06	*	< LOD	< LOD	< LOD	< LOD	719
	07-08	*	< LOD	< LOD	< LOD	< LOD	524
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	980
	05-06	*	< LOD	< LOD	< LOD	< LOD	993
	07-08	*	< LOD	< LOD	< LOD	< LOD	1007

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.1, 0.1, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Nicosulfuron (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2320
	05-06	*	< LOD	< LOD	< LOD	< LOD	2570
	07-08	*	< LOD	< LOD	< LOD	< LOD	2420
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	296
	05-06	*	< LOD	< LOD	< LOD	< LOD	341
	07-08	*	< LOD	< LOD	< LOD	< LOD	363
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	665
	05-06	*	< LOD	< LOD	< LOD	< LOD	699
	07-08	*	< LOD	< LOD	< LOD	< LOD	362
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1359
	05-06	*	< LOD	< LOD	< LOD	< LOD	1530
	07-08	*	< LOD	< LOD	< LOD	< LOD	1695
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1130
	05-06	*	< LOD	< LOD	< LOD	< LOD	1222
	07-08	*	< LOD	< LOD	< LOD	< LOD	1209
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1190
	05-06	*	< LOD	< LOD	< LOD	< LOD	1348
	07-08	*	< LOD	< LOD	< LOD	< LOD	1211
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	588
	05-06	*	< LOD	< LOD	< LOD	< LOD	656
	07-08	*	< LOD	< LOD	< LOD	< LOD	485
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	601
	05-06	*	< LOD	< LOD	< LOD	< LOD	719
	07-08	*	< LOD	< LOD	< LOD	< LOD	523
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	979
	05-06	*	< LOD	< LOD	< LOD	< LOD	993
	07-08	*	< LOD	< LOD	< LOD	< LOD	1007

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Oxasulfuron

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2355
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2540
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	290
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	379
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	672
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	381
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1393
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1780
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1150
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1269
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1205
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1271
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	579
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	487
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	616
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	567
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1018
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1076

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.06, 0.06, and 0.06 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Oxasulfuron (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2353
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2538
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	290
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	379
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	671
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	379
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1392
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1780
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1149
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1268
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1204
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1270
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	578
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	486
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	616
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	566
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1017
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1076

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Primisulfuron-methyl

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2250
	05-06	*	< LOD	< LOD	< LOD	< LOD	2455
	07-08	*	< LOD	< LOD	< LOD	< LOD	2422
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	285
	05-06	*	< LOD	< LOD	< LOD	< LOD	327
	07-08	*	< LOD	< LOD	< LOD	< LOD	356
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	627
	05-06	*	< LOD	< LOD	< LOD	< LOD	665
	07-08	*	< LOD	< LOD	< LOD	< LOD	376
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1338
	05-06	*	< LOD	< LOD	< LOD	< LOD	1463
	07-08	*	< LOD	< LOD	< LOD	< LOD	1690
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1107
	05-06	*	< LOD	< LOD	< LOD	< LOD	1160
	07-08	*	< LOD	< LOD	< LOD	< LOD	1207
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1143
	05-06	*	< LOD	< LOD	< LOD	< LOD	1295
	07-08	*	< LOD	< LOD	< LOD	< LOD	1215
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	574
	05-06	*	< LOD	< LOD	< LOD	< LOD	667
	07-08	*	< LOD	< LOD	< LOD	< LOD	456
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	556
	05-06	*	< LOD	< LOD	< LOD	< LOD	661
	07-08	*	< LOD	< LOD	< LOD	< LOD	549
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	975
	05-06	*	< LOD	< LOD	< LOD	< LOD	938
	07-08	*	< LOD	< LOD	< LOD	< LOD	1030

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.07, 0.07, and 0.07 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Primisulfuron-methyl (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2249
	05-06	*	< LOD	< LOD	< LOD	< LOD	2455
	07-08	*	< LOD	< LOD	< LOD	< LOD	2420
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	285
	05-06	*	< LOD	< LOD	< LOD	< LOD	327
	07-08	*	< LOD	< LOD	< LOD	< LOD	356
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	626
	05-06	*	< LOD	< LOD	< LOD	< LOD	665
	07-08	*	< LOD	< LOD	< LOD	< LOD	374
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1338
	05-06	*	< LOD	< LOD	< LOD	< LOD	1463
	07-08	*	< LOD	< LOD	< LOD	< LOD	1690
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1107
	05-06	*	< LOD	< LOD	< LOD	< LOD	1160
	07-08	*	< LOD	< LOD	< LOD	< LOD	1206
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1142
	05-06	*	< LOD	< LOD	< LOD	< LOD	1295
	07-08	*	< LOD	< LOD	< LOD	< LOD	1214
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	573
	05-06	*	< LOD	< LOD	< LOD	< LOD	667
	07-08	*	< LOD	< LOD	< LOD	< LOD	455
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	556
	05-06	*	< LOD	< LOD	< LOD	< LOD	661
	07-08	*	< LOD	< LOD	< LOD	< LOD	548
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	975
	05-06	*	< LOD	< LOD	< LOD	< LOD	938
	07-08	*	< LOD	< LOD	< LOD	< LOD	1030

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Prosulfuron

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2247
	05-06	*	< LOD	< LOD	< LOD	< LOD	2461
	07-08	*	< LOD	< LOD	< LOD	< LOD	2566
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	284
	05-06	*	< LOD	< LOD	< LOD	< LOD	326
	07-08	*	< LOD	< LOD	< LOD	< LOD	380
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	640
	05-06	*	< LOD	< LOD	< LOD	< LOD	676
	07-08	*	< LOD	< LOD	< LOD	< LOD	386
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1323
	05-06	*	< LOD	< LOD	< LOD	< LOD	1459
	07-08	*	< LOD	< LOD	< LOD	< LOD	1800
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1100
	05-06	*	< LOD	< LOD	< LOD	< LOD	1165
	07-08	*	< LOD	< LOD	< LOD	< LOD	1285
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1147
	05-06	*	< LOD	< LOD	< LOD	< LOD	1296
	07-08	*	< LOD	< LOD	< LOD	< LOD	1281
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	589
	05-06	*	< LOD	< LOD	< LOD	< LOD	673
	07-08	*	< LOD	< LOD	< LOD	< LOD	491
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	565
	05-06	*	< LOD	< LOD	< LOD	< LOD	677
	07-08	*	< LOD	< LOD	< LOD	< LOD	570
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	940
	05-06	*	< LOD	< LOD	< LOD	< LOD	924
	07-08	*	< LOD	< LOD	< LOD	< LOD	1081

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.05, 0.05, and 0.05 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Prosulfuron (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2245
	05-06	*	< LOD	< LOD	< LOD	< LOD	2461
	07-08	*	< LOD	< LOD	< LOD	< LOD	2564
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	284
	05-06	*	< LOD	< LOD	< LOD	< LOD	326
	07-08	*	< LOD	< LOD	< LOD	< LOD	380
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	639
	05-06	*	< LOD	< LOD	< LOD	< LOD	676
	07-08	*	< LOD	< LOD	< LOD	< LOD	384
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1322
	05-06	*	< LOD	< LOD	< LOD	< LOD	1459
	07-08	*	< LOD	< LOD	< LOD	< LOD	1800
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1099
	05-06	*	< LOD	< LOD	< LOD	< LOD	1165
	07-08	*	< LOD	< LOD	< LOD	< LOD	1284
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1146
	05-06	*	< LOD	< LOD	< LOD	< LOD	1296
	07-08	*	< LOD	< LOD	< LOD	< LOD	1280
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	588
	05-06	*	< LOD	< LOD	< LOD	< LOD	673
	07-08	*	< LOD	< LOD	< LOD	< LOD	490
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	565
	05-06	*	< LOD	< LOD	< LOD	< LOD	677
	07-08	*	< LOD	< LOD	< LOD	< LOD	569
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	939
	05-06	*	< LOD	< LOD	< LOD	< LOD	924
	07-08	*	< LOD	< LOD	< LOD	< LOD	1081

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Rimsulfuron

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2284
	05-06	*	< LOD	< LOD	< LOD	< LOD	2593
	07-08	*	< LOD	< LOD	< LOD	< LOD	2530
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	279
	05-06	*	< LOD	< LOD	< LOD	< LOD	346
	07-08	*	< LOD	< LOD	< LOD	< LOD	376
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	654
	05-06	*	< LOD	< LOD	< LOD	< LOD	706
	07-08	*	< LOD	< LOD	< LOD	< LOD	383
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1351
	05-06	*	< LOD	< LOD	< LOD	< LOD	1541
	07-08	*	< LOD	< LOD	< LOD	< LOD	1771
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1110
	05-06	*	< LOD	< LOD	< LOD	< LOD	1224
	07-08	*	< LOD	< LOD	< LOD	< LOD	1271
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1174
	05-06	*	< LOD	< LOD	< LOD	< LOD	1369
	07-08	*	< LOD	< LOD	< LOD	< LOD	1259
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	561
	05-06	*	< LOD	< LOD	< LOD	< LOD	675
	07-08	*	< LOD	< LOD	< LOD	< LOD	463
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	592
	05-06	*	< LOD	< LOD	< LOD	< LOD	715
	07-08	*	< LOD	< LOD	< LOD	< LOD	568
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	985
	05-06	*	< LOD	< LOD	< LOD	< LOD	998
	07-08	*	< LOD	< LOD	< LOD	< LOD	1080

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.05, 0.05, and 0.05 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Rimsulfuron (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2282
	05-06	*	< LOD	< LOD	< LOD	< LOD	2593
	07-08	*	< LOD	< LOD	< LOD	< LOD	2528
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	279
	05-06	*	< LOD	< LOD	< LOD	< LOD	346
	07-08	*	< LOD	< LOD	< LOD	< LOD	376
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	653
	05-06	*	< LOD	< LOD	< LOD	< LOD	706
	07-08	*	< LOD	< LOD	< LOD	< LOD	381
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1350
	05-06	*	< LOD	< LOD	< LOD	< LOD	1541
	07-08	*	< LOD	< LOD	< LOD	< LOD	1771
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1109
	05-06	*	< LOD	< LOD	< LOD	< LOD	1224
	07-08	*	< LOD	< LOD	< LOD	< LOD	1270
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1173
	05-06	*	< LOD	< LOD	< LOD	< LOD	1369
	07-08	*	< LOD	< LOD	< LOD	< LOD	1258
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	560
	05-06	*	< LOD	< LOD	< LOD	< LOD	675
	07-08	*	< LOD	< LOD	< LOD	< LOD	462
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	592
	05-06	*	< LOD	< LOD	< LOD	< LOD	715
	07-08	*	< LOD	< LOD	< LOD	< LOD	567
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	984
	05-06	*	< LOD	< LOD	< LOD	< LOD	998
	07-08	*	< LOD	< LOD	< LOD	< LOD	1080

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Sulfometuron-methyl

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2067
	05-06	*	< LOD	< LOD	< LOD	< LOD	2598
	07-08	*	< LOD	< LOD	< LOD	< LOD	2458
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	252
	05-06	*	< LOD	< LOD	< LOD	< LOD	345
	07-08	*	< LOD	< LOD	< LOD	< LOD	352
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	587
	05-06	*	< LOD	< LOD	< LOD	< LOD	710
	07-08	*	< LOD	< LOD	< LOD	< LOD	373
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1228
	05-06	*	< LOD	< LOD	< LOD	< LOD	1543
	07-08	*	< LOD	< LOD	< LOD	< LOD	1733
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1007
	05-06	*	< LOD	< LOD	< LOD	< LOD	1233
	07-08	*	< LOD	< LOD	< LOD	< LOD	1226
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1060
	05-06	*	< LOD	< LOD	< LOD	< LOD	1365
	07-08	*	< LOD	< LOD	< LOD	< LOD	1232
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	441
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	569
	05-06	*	< LOD	< LOD	< LOD	< LOD	698
	07-08	*	< LOD	< LOD	< LOD	< LOD	557
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	916
	05-06	*	< LOD	< LOD	< LOD	< LOD	996
	07-08	*	< LOD	< LOD	< LOD	< LOD	1045

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.05, 0.05, and 0.05 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Sulfometuron-methyl (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2065
	05-06	*	< LOD	< LOD	< LOD	< LOD	2598
	07-08	*	< LOD	< LOD	< LOD	< LOD	2456
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	252
	05-06	*	< LOD	< LOD	< LOD	< LOD	345
	07-08	*	< LOD	< LOD	< LOD	< LOD	352
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	586
	05-06	*	< LOD	< LOD	< LOD	< LOD	710
	07-08	*	< LOD	< LOD	< LOD	< LOD	371
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1227
	05-06	*	< LOD	< LOD	< LOD	< LOD	1543
	07-08	*	< LOD	< LOD	< LOD	< LOD	1733
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1006
	05-06	*	< LOD	< LOD	< LOD	< LOD	1233
	07-08	*	< LOD	< LOD	< LOD	< LOD	1225
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1059
	05-06	*	< LOD	< LOD	< LOD	< LOD	1365
	07-08	*	< LOD	< LOD	< LOD	< LOD	1231
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	440
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	453
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	569
	05-06	*	< LOD	< LOD	< LOD	< LOD	698
	07-08	*	< LOD	< LOD	< LOD	< LOD	556
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	915
	05-06	*	< LOD	< LOD	< LOD	< LOD	996
	07-08	*	< LOD	< LOD	< LOD	< LOD	1045

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Sulfosulfuron

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2391
	05-06	*	< LOD	< LOD	< LOD	< LOD	2593
	07-08	*	< LOD	< LOD	< LOD	< LOD	2530
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	294
	05-06	*	< LOD	< LOD	< LOD	< LOD	344
	07-08	*	< LOD	< LOD	< LOD	< LOD	377
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	681
	05-06	*	< LOD	< LOD	< LOD	< LOD	708
	07-08	*	< LOD	< LOD	< LOD	< LOD	385
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1416
	05-06	*	< LOD	< LOD	< LOD	< LOD	1541
	07-08	*	< LOD	< LOD	< LOD	< LOD	1768
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1170
	05-06	*	< LOD	< LOD	< LOD	< LOD	1230
	07-08	*	< LOD	< LOD	< LOD	< LOD	1261
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1221
	05-06	*	< LOD	< LOD	< LOD	< LOD	1363
	07-08	*	< LOD	< LOD	< LOD	< LOD	1269
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	594
	05-06	*	< LOD	< LOD	< LOD	< LOD	673
	07-08	*	< LOD	< LOD	< LOD	< LOD	485
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	617
	05-06	*	< LOD	< LOD	< LOD	< LOD	719
	07-08	*	< LOD	< LOD	< LOD	< LOD	565
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1026
	05-06	*	< LOD	< LOD	< LOD	< LOD	997
	07-08	*	< LOD	< LOD	< LOD	< LOD	1068

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.1, 0.1, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Sulfosulfuron (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2389
	05-06	*	< LOD	< LOD	< LOD	< LOD	2593
	07-08	*	< LOD	< LOD	< LOD	< LOD	2528
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	294
	05-06	*	< LOD	< LOD	< LOD	< LOD	344
	07-08	*	< LOD	< LOD	< LOD	< LOD	377
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	680
	05-06	*	< LOD	< LOD	< LOD	< LOD	708
	07-08	*	< LOD	< LOD	< LOD	< LOD	383
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1415
	05-06	*	< LOD	< LOD	< LOD	< LOD	1541
	07-08	*	< LOD	< LOD	< LOD	< LOD	1768
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1169
	05-06	*	< LOD	< LOD	< LOD	< LOD	1230
	07-08	*	< LOD	< LOD	< LOD	< LOD	1260
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1220
	05-06	*	< LOD	< LOD	< LOD	< LOD	1363
	07-08	*	< LOD	< LOD	< LOD	< LOD	1268
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	593
	05-06	*	< LOD	< LOD	< LOD	< LOD	673
	07-08	*	< LOD	< LOD	< LOD	< LOD	484
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	617
	05-06	*	< LOD	< LOD	< LOD	< LOD	719
	07-08	*	< LOD	< LOD	< LOD	< LOD	564
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1025
	05-06	*	< LOD	< LOD	< LOD	< LOD	997
	07-08	*	< LOD	< LOD	< LOD	< LOD	1068

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Thifensulfuron-methyl

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2355
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2495
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	291
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	370
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	677
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	375
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1387
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1750
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1145
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1252
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1210
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1243
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	566
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	479
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	616
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	542
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1022
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1056

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.08, 0.08, and 0.08 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Thifensulfuron-methyl (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2353
	05-06	*	< LOD	< LOD	< LOD	< LOD	2634
	07-08	*	< LOD	< LOD	< LOD	< LOD	2493
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	291
	05-06	*	< LOD	< LOD	< LOD	< LOD	350
	07-08	*	< LOD	< LOD	< LOD	< LOD	370
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	676
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
	07-08	*	< LOD	< LOD	< LOD	< LOD	373
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1386
	05-06	*	< LOD	< LOD	< LOD	< LOD	1566
	07-08	*	< LOD	< LOD	< LOD	< LOD	1750
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1144
	05-06	*	< LOD	< LOD	< LOD	< LOD	1247
	07-08	*	< LOD	< LOD	< LOD	< LOD	1251
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1209
	05-06	*	< LOD	< LOD	< LOD	< LOD	1387
	07-08	*	< LOD	< LOD	< LOD	< LOD	1242
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	565
	05-06	*	< LOD	< LOD	< LOD	< LOD	697
	07-08	*	< LOD	< LOD	< LOD	< LOD	478
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	616
	05-06	*	< LOD	< LOD	< LOD	< LOD	720
	07-08	*	< LOD	< LOD	< LOD	< LOD	541
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1021
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007
	07-08	*	< LOD	< LOD	< LOD	< LOD	1056

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Triasulfuron

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2250
	05-06	*	< LOD	< LOD	< LOD	< LOD	2557
	07-08	*	< LOD	< LOD	< LOD	< LOD	2530
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	278
	05-06	*	< LOD	< LOD	< LOD	< LOD	341
	07-08	*	< LOD	< LOD	< LOD	< LOD	378
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	643
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	385
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1329
	05-06	*	< LOD	< LOD	< LOD	< LOD	1521
	07-08	*	< LOD	< LOD	< LOD	< LOD	1767
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1101
	05-06	*	< LOD	< LOD	< LOD	< LOD	1210
	07-08	*	< LOD	< LOD	< LOD	< LOD	1265
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1149
	05-06	*	< LOD	< LOD	< LOD	< LOD	1347
	07-08	*	< LOD	< LOD	< LOD	< LOD	1265
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	549
	05-06	*	< LOD	< LOD	< LOD	< LOD	677
	07-08	*	< LOD	< LOD	< LOD	< LOD	488
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	580
	05-06	*	< LOD	< LOD	< LOD	< LOD	713
	07-08	*	< LOD	< LOD	< LOD	< LOD	547
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	973
	05-06	*	< LOD	< LOD	< LOD	< LOD	968
	07-08	*	< LOD	< LOD	< LOD	< LOD	1073

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.07, 0.07, and 0.07 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Triasulfuron (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2248
	05-06	*	< LOD	< LOD	< LOD	< LOD	2557
	07-08	*	< LOD	< LOD	< LOD	< LOD	2528
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	278
	05-06	*	< LOD	< LOD	< LOD	< LOD	341
	07-08	*	< LOD	< LOD	< LOD	< LOD	378
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	642
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	383
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1328
	05-06	*	< LOD	< LOD	< LOD	< LOD	1521
	07-08	*	< LOD	< LOD	< LOD	< LOD	1767
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1100
	05-06	*	< LOD	< LOD	< LOD	< LOD	1210
	07-08	*	< LOD	< LOD	< LOD	< LOD	1264
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1148
	05-06	*	< LOD	< LOD	< LOD	< LOD	1347
	07-08	*	< LOD	< LOD	< LOD	< LOD	1264
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	548
	05-06	*	< LOD	< LOD	< LOD	< LOD	677
	07-08	*	< LOD	< LOD	< LOD	< LOD	487
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	580
	05-06	*	< LOD	< LOD	< LOD	< LOD	713
	07-08	*	< LOD	< LOD	< LOD	< LOD	546
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	972
	05-06	*	< LOD	< LOD	< LOD	< LOD	968
	07-08	*	< LOD	< LOD	< LOD	< LOD	1073

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Triflurosulfuron-methyl

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2427
	05-06	*	< LOD	< LOD	< LOD	< LOD	2562
	07-08	*	< LOD	< LOD	< LOD	< LOD	2494
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	302
	05-06	*	< LOD	< LOD	< LOD	< LOD	340
	07-08	*	< LOD	< LOD	< LOD	< LOD	374
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	694
	05-06	*	< LOD	< LOD	< LOD	< LOD	702
	07-08	*	< LOD	< LOD	< LOD	< LOD	381
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1431
	05-06	*	< LOD	< LOD	< LOD	< LOD	1520
	07-08	*	< LOD	< LOD	< LOD	< LOD	1739
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1189
	05-06	*	< LOD	< LOD	< LOD	< LOD	1216
	07-08	*	< LOD	< LOD	< LOD	< LOD	1240
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1238
	05-06	*	< LOD	< LOD	< LOD	< LOD	1346
	07-08	*	< LOD	< LOD	< LOD	< LOD	1254
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	596
	05-06	*	< LOD	< LOD	< LOD	< LOD	693
	07-08	*	< LOD	< LOD	< LOD	< LOD	477
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	634
	05-06	*	< LOD	< LOD	< LOD	< LOD	699
	07-08	*	< LOD	< LOD	< LOD	< LOD	558
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1039
	05-06	*	< LOD	< LOD	< LOD	< LOD	972
	07-08	*	< LOD	< LOD	< LOD	< LOD	1053

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 0.05, 0.05, and 0.05 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Triflurosulfuron-methyl (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2425
	05-06	*	< LOD	< LOD	< LOD	< LOD	2562
	07-08	*	< LOD	< LOD	< LOD	< LOD	2492
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	302
	05-06	*	< LOD	< LOD	< LOD	< LOD	340
	07-08	*	< LOD	< LOD	< LOD	< LOD	374
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	693
	05-06	*	< LOD	< LOD	< LOD	< LOD	702
	07-08	*	< LOD	< LOD	< LOD	< LOD	379
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1430
	05-06	*	< LOD	< LOD	< LOD	< LOD	1520
	07-08	*	< LOD	< LOD	< LOD	< LOD	1739
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1188
	05-06	*	< LOD	< LOD	< LOD	< LOD	1216
	07-08	*	< LOD	< LOD	< LOD	< LOD	1239
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1237
	05-06	*	< LOD	< LOD	< LOD	< LOD	1346
	07-08	*	< LOD	< LOD	< LOD	< LOD	1253
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	595
	05-06	*	< LOD	< LOD	< LOD	< LOD	693
	07-08	*	< LOD	< LOD	< LOD	< LOD	476
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	634
	05-06	*	< LOD	< LOD	< LOD	< LOD	699
	07-08	*	< LOD	< LOD	< LOD	< LOD	557
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1038
	05-06	*	< LOD	< LOD	< LOD	< LOD	972
	07-08	*	< LOD	< LOD	< LOD	< LOD	1053

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/SulfonylureaHerbicides_BiomonitoringSummary.html

Urinary Carbofuranphenol

Metabolite of Benfuracarb, Carbofuran, Carbosulfan, and Furathiocarb

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	.770 (<LOD-1.30)	1994
	01-02	*	< LOD	< LOD	< LOD	< LOD	2530
	03-04	*	< LOD	< LOD	< LOD	< LOD	1879
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	.450 (<LOD-2.20)	482
	01-02	*	< LOD	< LOD	< LOD	< LOD	578
	03-04	*	< LOD	< LOD	< LOD	< LOD	293
12-19 years	99-00	*	< LOD	< LOD	< LOD	.570 (<LOD-1.20)	681
	01-02	*	< LOD	< LOD	< LOD	< LOD	827
	03-04	*	< LOD	< LOD	< LOD	< LOD	684
20-59 years	99-00	*	< LOD	< LOD	< LOD	.840 (<LOD-1.50)	831
	01-02	*	< LOD	< LOD	< LOD	< LOD	1125
	03-04	*	< LOD	< LOD	< LOD	< LOD	902
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	.740 (<LOD-1.50)	973
	01-02	*	< LOD	< LOD	< LOD	< LOD	1190
	03-04	*	< LOD	< LOD	< LOD	< LOD	900
Females	99-00	*	< LOD	< LOD	< LOD	.840 (<LOD-1.50)	1021
	01-02	*	< LOD	< LOD	< LOD	< LOD	1340
	03-04	*	< LOD	< LOD	< LOD	< LOD	979
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	.590 (<LOD-2.00)	1.90 (<LOD-5.10)	696
	01-02	*	< LOD	< LOD	< LOD	< LOD	680
	03-04	*	< LOD	< LOD	< LOD	< LOD	464
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	.550 (<LOD-1.60)	521
	01-02	*	< LOD	< LOD	< LOD	< LOD	696
	03-04	*	< LOD	< LOD	< LOD	< LOD	544
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	.740 (<LOD-1.50)	603
	01-02	*	< LOD	< LOD	< LOD	< LOD	953
	03-04	*	< LOD	< LOD	< LOD	< LOD	741

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 0.4, 0.4, and 0.4 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Carbofuran_BiomonitoringSummary.html

Urinary Carbofuranphenol (creatinine corrected)

Metabolite of Benfuracarb, Carbofuran, Carbosulfan, and Furathiocarb

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	.778 (<LOD-1.00)	1994
	01-02	*	< LOD	< LOD	< LOD	< LOD	2529
	03-04	*	< LOD	< LOD	< LOD	< LOD	1877
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	.988 (<LOD-2.80)	482
	01-02	*	< LOD	< LOD	< LOD	< LOD	578
	03-04	*	< LOD	< LOD	< LOD	< LOD	293
12-19 years	99-00	*	< LOD	< LOD	< LOD	.482 (<LOD-.853)	681
	01-02	*	< LOD	< LOD	< LOD	< LOD	826
	03-04	*	< LOD	< LOD	< LOD	< LOD	683
20-59 years	99-00	*	< LOD	< LOD	< LOD	.875 (<LOD-1.06)	831
	01-02	*	< LOD	< LOD	< LOD	< LOD	1125
	03-04	*	< LOD	< LOD	< LOD	< LOD	901
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	.667 (<LOD-1.08)	973
	01-02	*	< LOD	< LOD	< LOD	< LOD	1190
	03-04	*	< LOD	< LOD	< LOD	< LOD	899
Females	99-00	*	< LOD	< LOD	< LOD	.881 (<LOD-1.13)	1021
	01-02	*	< LOD	< LOD	< LOD	< LOD	1339
	03-04	*	< LOD	< LOD	< LOD	< LOD	978
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	.778 (<LOD-1.94)	1.83 (<LOD-4.16)	696
	01-02	*	< LOD	< LOD	< LOD	< LOD	680
	03-04	*	< LOD	< LOD	< LOD	< LOD	463
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	.700 (<LOD-1.08)	521
	01-02	*	< LOD	< LOD	< LOD	< LOD	695
	03-04	*	< LOD	< LOD	< LOD	< LOD	544
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	.737 (<LOD-.929)	603
	01-02	*	< LOD	< LOD	< LOD	< LOD	953
	03-04	*	< LOD	< LOD	< LOD	< LOD	740

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Carbofuran_BiomonitoringSummary.html

Urinary 2-Isopropoxyphenol

Metabolite of Propoxur

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1917
	01-02	*	< LOD	< LOD	< LOD	< LOD	2503
	03-04	*	< LOD	< LOD	< LOD	< LOD	1881
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	< LOD	456
	01-02	*	< LOD	< LOD	< LOD	< LOD	574
	03-04	*	< LOD	< LOD	< LOD	< LOD	295
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	655
	01-02	*	< LOD	< LOD	< LOD	< LOD	820
	03-04	*	< LOD	< LOD	< LOD	< LOD	686
20-59 years	99-00	*	< LOD	< LOD	< LOD	< LOD	806
	01-02	*	< LOD	< LOD	< LOD	< LOD	1109
	03-04	*	< LOD	< LOD	< LOD	< LOD	900
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	936
	01-02	*	< LOD	< LOD	< LOD	< LOD	1178
	03-04	*	< LOD	< LOD	< LOD	< LOD	899
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	981
	01-02	*	< LOD	< LOD	< LOD	< LOD	1325
	03-04	*	< LOD	< LOD	< LOD	< LOD	982
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	664
	01-02	*	< LOD	< LOD	< LOD	< LOD	677
	03-04	*	< LOD	< LOD	< LOD	< LOD	484
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	500
	01-02	*	< LOD	< LOD	< LOD	< LOD	696
	03-04	*	< LOD	< LOD	< LOD	< LOD	535
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	585
	01-02	*	< LOD	< LOD	< LOD	< LOD	931
	03-04	*	< LOD	< LOD	< LOD	< LOD	736

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 1.1, 0.4, and 0.4 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Propoxur_BiomonitoringSummary.html

Urinary 2-Isopropoxyphenol (creatinine corrected)

Metabolite of Propoxur

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1917
	01-02	*	< LOD	< LOD	< LOD	< LOD	2502
	03-04	*	< LOD	< LOD	< LOD	< LOD	1879
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	< LOD	456
	01-02	*	< LOD	< LOD	< LOD	< LOD	574
	03-04	*	< LOD	< LOD	< LOD	< LOD	295
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	655
	01-02	*	< LOD	< LOD	< LOD	< LOD	819
	03-04	*	< LOD	< LOD	< LOD	< LOD	685
20-59 years	99-00	*	< LOD	< LOD	< LOD	< LOD	806
	01-02	*	< LOD	< LOD	< LOD	< LOD	1109
	03-04	*	< LOD	< LOD	< LOD	< LOD	899
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	936
	01-02	*	< LOD	< LOD	< LOD	< LOD	1178
	03-04	*	< LOD	< LOD	< LOD	< LOD	898
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	981
	01-02	*	< LOD	< LOD	< LOD	< LOD	1324
	03-04	*	< LOD	< LOD	< LOD	< LOD	981
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	664
	01-02	*	< LOD	< LOD	< LOD	< LOD	677
	03-04	*	< LOD	< LOD	< LOD	< LOD	483
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	500
	01-02	*	< LOD	< LOD	< LOD	< LOD	695
	03-04	*	< LOD	< LOD	< LOD	< LOD	535
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	585
	01-02	*	< LOD	< LOD	< LOD	< LOD	931
	03-04	*	< LOD	< LOD	< LOD	< LOD	735

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Propoxur_BiomonitoringSummary.html

Serum Aldrin (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2275
	03-04	*	< LOD	< LOD	< LOD	< LOD	1946
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	< LOD	< LOD	< LOD	< LOD	588
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1519
	03-04	*	< LOD	< LOD	< LOD	< LOD	1358
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1057
	03-04	*	< LOD	< LOD	< LOD	< LOD	946
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1218
	03-04	*	< LOD	< LOD	< LOD	< LOD	1000
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	559
	03-04	*	< LOD	< LOD	< LOD	< LOD	456
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	512
	03-04	*	< LOD	< LOD	< LOD	< LOD	485
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1045
	03-04	*	< LOD	< LOD	< LOD	< LOD	881

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 5.94 and 7.8.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/AldrinDieldrin_BiomonitoringSummary.html

Serum Aldrin (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2275
	03-04	*	< LOD	< LOD	< LOD	< LOD	1946
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	< LOD	< LOD	< LOD	< LOD	588
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1519
	03-04	*	< LOD	< LOD	< LOD	< LOD	1358
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1057
	03-04	*	< LOD	< LOD	< LOD	< LOD	946
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1218
	03-04	*	< LOD	< LOD	< LOD	< LOD	1000
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	559
	03-04	*	< LOD	< LOD	< LOD	< LOD	456
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	512
	03-04	*	< LOD	< LOD	< LOD	< LOD	485
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1045
	03-04	*	< LOD	< LOD	< LOD	< LOD	881

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/AldrinDieldrin_BiomonitoringSummary.html

Serum Dieldrin (lipid adjusted)

Also a Metabolite of Aldrin

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	15.3 (14.5-17.4)	20.3 (18.7-22.4)	2159
	03-04	*	< LOD	9.00 (8.40-9.90)	14.4 (12.1-16.4)	19.0 (15.8-24.2)	1952
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	716
	03-04	*	< LOD	< LOD	< LOD	9.10 (<LOD-16.4)	587
20 years and older	01-02	*	< LOD	10.5 (<LOD-11.6)	16.6 (15.1-18.2)	21.3 (19.1-24.0)	1443
	03-04	*	< LOD	9.50 (8.80-10.4)	14.9 (12.8-17.0)	19.5 (16.0-25.7)	1365
Gender							
Males	01-02	*	< LOD	< LOD	15.7 (14.4-18.7)	20.3 (18.6-24.0)	1007
	03-04	*	< LOD	9.30 (8.40-10.8)	15.1 (13.1-19.1)	21.9 (14.9-38.5)	954
Females	01-02	*	< LOD	< LOD	15.3 (13.4-17.2)	19.8 (18.0-21.6)	1152
	03-04	*	< LOD	8.70 (7.80-9.50)	12.8 (11.2-15.4)	16.9 (13.9-22.4)	998
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	11.7 (<LOD-15.1)	15.4 (12.7-19.1)	539
	03-04	*	< LOD	< LOD	10.8 (9.00-14.1)	14.0 (10.6-24.1)	456
Non-Hispanic blacks	01-02	*	< LOD	< LOD	15.0 (11.8-19.1)	20.6 (15.8-25.2)	484
	03-04	*	< LOD	8.80 (<LOD-10.1)	13.0 (10.5-15.8)	15.9 (13.3-21.5)	487
Non-Hispanic whites	01-02	*	< LOD	< LOD	15.6 (14.8-17.8)	21.1 (18.9-23.6)	980
	03-04	*	< LOD	9.30 (8.60-10.2)	14.9 (12.5-17.5)	19.7 (15.6-33.4)	885

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 10.5 and 7.8.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/AldrinDieldrin_BiomonitoringSummary.html

Serum Dieldrin (whole weight)

Metabolite of Aldrin

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	.110 (.100-.120)	.150 (.130-.170)	2159
	03-04	*	< LOD	.059 (.054-.064)	.098 (.083-.112)	.138 (.109-.177)	1952
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	716
	03-04	*	< LOD	< LOD	< LOD	.048 (<LOD-.093)	587
20 years and older	01-02	*	< LOD	.070 (<LOD-.080)	.120 (.110-.130)	.160 (.140-.180)	1443
	03-04	*	< LOD	.062 (.056-.069)	.102 (.088-.117)	.139 (.112-.182)	1365
Gender							
Males	01-02	*	< LOD	< LOD	.120 (.100-.130)	.160 (.130-.190)	1007
	03-04	*	< LOD	.064 (.054-.075)	.109 (.084-.138)	.147 (.108-.242)	954
Females	01-02	*	< LOD	< LOD	.100 (.090-.110)	.140 (.120-.160)	1152
	03-04	*	< LOD	.055 (.049-.060)	.089 (.077-.100)	.110 (.096-.147)	998
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	.090 (<LOD-.100)	.120 (.090-.140)	539
	03-04	*	< LOD	< LOD	.077 (.062-.116)	.113 (.073-.158)	456
Non-Hispanic blacks	01-02	*	< LOD	< LOD	.090 (.070-.110)	.130 (.090-.190)	484
	03-04	*	< LOD	.053 (<LOD-.058)	.080 (.063-.101)	.103 (.080-.149)	487
Non-Hispanic whites	01-02	*	< LOD	< LOD	.110 (.100-.130)	.150 (.130-.180)	980
	03-04	*	< LOD	.062 (.054-.070)	.103 (.086-.124)	.139 (.109-.202)	885

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/AldrinDieldrin_BiomonitoringSummary.html

Serum Endrin (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	5.10 (<LOD-5.40)	2187
	03-04	*	< LOD	< LOD	< LOD	< LOD	1825
Age group							
12-19 years	01-02	*	< LOD	< LOD	5.20 (<LOD-5.50)	5.60 (5.40-5.70)	730
	03-04	*	< LOD	< LOD	< LOD	< LOD	539
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1457
	03-04	*	< LOD	< LOD	< LOD	< LOD	1286
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	5.20 (<LOD-5.40)	1022
	03-04	*	< LOD	< LOD	< LOD	< LOD	885
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1165
	03-04	*	< LOD	< LOD	< LOD	< LOD	940
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	5.30 (<LOD-6.50)	547
	03-04	*	< LOD	< LOD	< LOD	< LOD	433
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	5.40 (<LOD-6.30)	487
	03-04	*	< LOD	< LOD	< LOD	< LOD	446
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	5.10 (<LOD-5.40)	1000
	03-04	*	< LOD	< LOD	< LOD	< LOD	831

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 5.09 and 7.8.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Endrin_BiomonitoringSummary.html

Serum Endrin (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	.020 (<LOD-.020)	2187
	03-04	*	< LOD	< LOD	< LOD	< LOD	1825
Age group							
12-19 years	01-02	*	< LOD	< LOD	.020 (<LOD-.020)	.020 (.020-.020)	730
	03-04	*	< LOD	< LOD	< LOD	< LOD	539
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1457
	03-04	*	< LOD	< LOD	< LOD	< LOD	1286
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	.020 (<LOD-.020)	1022
	03-04	*	< LOD	< LOD	< LOD	< LOD	885
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1165
	03-04	*	< LOD	< LOD	< LOD	< LOD	940
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	.020 (<LOD-.020)	547
	03-04	*	< LOD	< LOD	< LOD	< LOD	433
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	.020 (<LOD-.020)	487
	03-04	*	< LOD	< LOD	< LOD	< LOD	446
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	.020 (<LOD-.020)	1000
	03-04	*	< LOD	< LOD	< LOD	< LOD	831

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Endrin_BiomonitoringSummary.html

Serum Heptachlor epoxide (lipid adjusted)

Metabolite of Heptachlor

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	15.4 (<LOD-19.8)	24.0 (15.1-38.8)	1589
	01-02	*	< LOD	< LOD	14.8 (13.0-17.8)	21.8 (18.2-27.3)	2259
	03-04	*	< LOD	< LOD	13.4 (11.1-15.9)	18.9 (15.9-23.0)	1963
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	638
	01-02	*	< LOD	< LOD	< LOD	< LOD	741
	03-04	*	< LOD	< LOD	< LOD	< LOD	592
20 years and older	99-00	*	< LOD	< LOD	17.8 (<LOD-23.9)	27.1 (16.8-46.1)	951
	01-02	*	< LOD	< LOD	15.7 (13.7-18.8)	23.1 (19.1-29.1)	1518
	03-04	*	< LOD	8.20 (<LOD-9.50)	14.5 (11.6-17.6)	20.6 (16.8-24.0)	1371
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	19.6 (<LOD-27.2)	760
	01-02	*	< LOD	< LOD	13.9 (12.0-17.8)	20.8 (15.9-25.3)	1047
	03-04	*	< LOD	< LOD	13.4 (11.2-16.4)	18.6 (16.6-21.3)	956
Females	99-00	*	< LOD	< LOD	18.2 (<LOD-25.2)	27.7 (16.0-54.3)	829
	01-02	*	< LOD	< LOD	15.6 (13.3-18.2)	23.2 (18.9-29.8)	1212
	03-04	*	< LOD	< LOD	13.5 (11.0-16.8)	19.6 (14.8-24.5)	1007
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	15.3 (<LOD-25.6)	22.2 (<LOD-62.3)	598
	01-02	*	< LOD	< LOD	13.2 (<LOD-16.3)	16.8 (13.8-23.1)	553
	03-04	*	< LOD	< LOD	10.6 (8.10-13.5)	13.7 (10.9-16.5)	460
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	19.5 (<LOD-32.4)	336
	01-02	*	< LOD	< LOD	14.6 (11.7-19.0)	21.5 (18.2-27.3)	503
	03-04	*	< LOD	< LOD	13.5 (10.1-16.8)	18.6 (12.7-25.2)	490
Non-Hispanic whites	99-00	*	< LOD	< LOD	16.5 (<LOD-21.8)	26.4 (<LOD-54.3)	539
	01-02	*	< LOD	< LOD	15.3 (13.0-19.6)	22.8 (18.9-29.8)	1041
	03-04	*	< LOD	7.90 (<LOD-9.40)	14.0 (11.2-17.6)	20.4 (15.8-24.5)	888

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 14.6, 10.5, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum Heptachlor epoxide (whole weight)

Metabolite of Heptachlor

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	.110 (<LOD-.140)	.180 (.110-.220)	1589
	01-02	*	< LOD	< LOD	.100 (.090-.120)	.150 (.130-.180)	2259
	03-04	*	< LOD	< LOD	.094 (.076-.108)	.128 (.108-.161)	1963
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	638
	01-02	*	< LOD	< LOD	< LOD	< LOD	741
	03-04	*	< LOD	< LOD	< LOD	< LOD	592
20 years and older	99-00	*	< LOD	< LOD	.130 (<LOD-.170)	.190 (.130-.270)	951
	01-02	*	< LOD	< LOD	.110 (.090-.140)	.170 (.140-.190)	1518
	03-04	*	< LOD	.057 (<LOD-.063)	.101 (.082-.116)	.135 (.113-.173)	1371
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	.150 (<LOD-.180)	760
	01-02	*	< LOD	< LOD	.100 (.090-.120)	.150 (.110-.180)	1047
	03-04	*	< LOD	< LOD	.094 (.077-.111)	.126 (.106-.149)	956
Females	99-00	*	< LOD	< LOD	.120 (<LOD-.190)	.200 (.120-.310)	829
	01-02	*	< LOD	< LOD	.100 (.090-.130)	.170 (.130-.200)	1212
	03-04	*	< LOD	< LOD	.096 (.071-.108)	.133 (.107-.173)	1007
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	.100 (<LOD-.170)	.170 (<LOD-.380)	598
	01-02	*	< LOD	< LOD	.090 (<LOD-.110)	.120 (.100-.240)	553
	03-04	*	< LOD	< LOD	.069 (.053-.097)	.098 (.074-.113)	460
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	.110 (<LOD-.200)	336
	01-02	*	< LOD	< LOD	.090 (.070-.110)	.130 (.100-.180)	503
	03-04	*	< LOD	< LOD	.087 (.067-.104)	.110 (.090-.157)	490
Non-Hispanic whites	99-00	*	< LOD	< LOD	.120 (<LOD-.170)	.180 (<LOD-.310)	539
	01-02	*	< LOD	< LOD	.100 (.090-.130)	.170 (.130-.190)	1041
	03-04	*	< LOD	.055 (<LOD-.063)	.101 (.077-.117)	.135 (.111-.180)	888

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum *o,p'*-Dichlorodiphenyltrichloroethane (DDT) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1669
	01-02	*	< LOD	< LOD	< LOD	< LOD	2279
	03-04	*	< LOD	< LOD	< LOD	< LOD	1946
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	< LOD	< LOD	< LOD	< LOD	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1002
	01-02	*	< LOD	< LOD	< LOD	< LOD	1523
	03-04	*	< LOD	< LOD	< LOD	< LOD	1358
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	796
	01-02	*	< LOD	< LOD	< LOD	< LOD	1059
	03-04	*	< LOD	< LOD	< LOD	< LOD	949
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	873
	01-02	*	< LOD	< LOD	< LOD	< LOD	1220
	03-04	*	< LOD	< LOD	< LOD	< LOD	997
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	632
	01-02	*	< LOD	< LOD	< LOD	< LOD	565
	03-04	*	< LOD	< LOD	< LOD	< LOD	458
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	354
	01-02	*	< LOD	< LOD	< LOD	< LOD	507
	03-04	*	< LOD	< LOD	< LOD	< LOD	486
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	560
	01-02	*	< LOD	< LOD	< LOD	< LOD	1045
	03-04	*	< LOD	< LOD	< LOD	< LOD	880

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 20.7, 17.4, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *o,p'*-Dichlorodiphenyltrichloroethane (DDT) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1669
	01-02	*	< LOD	< LOD	< LOD	< LOD	2279
	03-04	*	< LOD	< LOD	< LOD	< LOD	1946
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	< LOD	< LOD	< LOD	< LOD	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1002
	01-02	*	< LOD	< LOD	< LOD	< LOD	1523
	03-04	*	< LOD	< LOD	< LOD	< LOD	1358
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	796
	01-02	*	< LOD	< LOD	< LOD	< LOD	1059
	03-04	*	< LOD	< LOD	< LOD	< LOD	949
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	873
	01-02	*	< LOD	< LOD	< LOD	< LOD	1220
	03-04	*	< LOD	< LOD	< LOD	< LOD	997
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	632
	01-02	*	< LOD	< LOD	< LOD	< LOD	565
	03-04	*	< LOD	< LOD	< LOD	< LOD	458
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	354
	01-02	*	< LOD	< LOD	< LOD	< LOD	507
	03-04	*	< LOD	< LOD	< LOD	< LOD	486
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	560
	01-02	*	< LOD	< LOD	< LOD	< LOD	1045
	03-04	*	< LOD	< LOD	< LOD	< LOD	880

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Urinary 2,4,5-Trichlorophenol

Metabolite of Several Organochlorine Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	.100 (.100-.100)	.200 (.200-.300)	.400 (.300-.400)	2525
	05-06	*	< LOD	.100 (.100-.200)	.300 (.200-.300)	.400 (.300-.500)	2548
	07-08	*	< LOD	.100 (<LOD-.100)	.200 (.200-.200)	.300 (.200-.300)	2604
	09-10	*	< LOD	< LOD	.200 (.200-.200)	.300 (.200-.300)	2749
Age group							
6-11 years	03-04	*	< LOD	.100 (.100-.200)	.200 (.200-.300)	.300 (.200-.500)	314
	05-06	*	< LOD	.100 (.100-.200)	.300 (.200-.400)	.400 (.300-.500)	356
	07-08	*	< LOD	.100 (<LOD-.100)	.200 (.100-.300)	.300 (.200-.500)	389
	09-10	*	< LOD	< LOD	.200 (.100-.200)	.200 (.200-.300)	415
12-19 years	03-04	*	< LOD	.100 (.100-.200)	.200 (.200-.300)	.300 (.200-.500)	722
	05-06	*	< LOD	.100 (.100-.200)	.300 (.200-.300)	.400 (.300-.500)	702
	07-08	*	< LOD	.100 (<LOD-.100)	.200 (.100-.200)	.200 (.200-.500)	401
	09-10	*	< LOD	.100 (<LOD-.100)	.200 (.200-.300)	.300 (.200-.500)	420
20 years and older	03-04	*	< LOD	.100 (.100-.100)	.300 (.200-.300)	.400 (.300-.500)	1489
	05-06	*	< LOD	.100 (.100-.200)	.300 (.200-.300)	.400 (.300-.500)	1490
	07-08	*	< LOD	.100 (<LOD-.100)	.200 (.200-.300)	.300 (.200-.400)	1814
	09-10	*	< LOD	< LOD	.200 (.200-.200)	.300 (.200-.300)	1914
Gender							
Males	03-04	*	< LOD	.100 (.100-.100)	.200 (.200-.300)	.400 (.300-.400)	1231
	05-06	*	< LOD	.100 (.100-.200)	.200 (.200-.300)	.400 (.300-.500)	1270
	07-08	*	< LOD	.100 (<LOD-.100)	.200 (.200-.200)	.300 (.200-.300)	1294
	09-10	*	< LOD	< LOD	.200 (.200-.200)	.300 (.200-.300)	1399
Females	03-04	*	< LOD	.100 (.100-.200)	.200 (.200-.300)	.400 (.300-.400)	1294
	05-06	*	< LOD	.100 (.100-.200)	.300 (.200-.400)	.500 (.300-.500)	1278
	07-08	*	< LOD	.100 (<LOD-.100)	.200 (.200-.300)	.300 (.200-.400)	1310
	09-10	*	< LOD	.100 (<LOD-.100)	.200 (.200-.200)	.300 (.200-.300)	1350
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	.100 (<LOD-.200)	.200 (.200-.300)	.300 (.200-.400)	617
	05-06	*	< LOD	.100 (<LOD-.200)	.300 (.200-.300)	.400 (.300-.500)	637
	07-08	*	< LOD	< LOD	.200 (.100-.200)	.200 (.200-.300)	531
	09-10	*	< LOD	< LOD	.100 (<LOD-.200)	.200 (.100-.300)	566
Non-Hispanic blacks	03-04	*	< LOD	.200 (.100-.200)	.300 (.200-.500)	.400 (.300-.700)	636
	05-06	*	< LOD	.200 (.100-.200)	.300 (.200-.400)	.500 (.300-.500)	678
	07-08	*	< LOD	.100 (.100-.200)	.200 (.200-.300)	.400 (.300-.500)	597
	09-10	*	< LOD	.100 (<LOD-.100)	.200 (.200-.200)	.200 (.200-.300)	516
Non-Hispanic whites	03-04	*	< LOD	.100 (.100-.100)	.200 (.200-.300)	.400 (.300-.400)	1077
	05-06	*	< LOD	.100 (.100-.200)	.300 (.200-.300)	.400 (.300-.600)	1038
	07-08	*	< LOD	.100 (<LOD-.100)	.200 (.200-.300)	.300 (.200-.400)	1077
	09-10	*	< LOD	< LOD	.200 (.200-.200)	.300 (.200-.300)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.1, 0.1, 0.1, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Trichlorophenols_BiomonitoringSummary.html

Urinary 2,4,5-Trichlorophenol (creatinine corrected)

Metabolite of Several Organochlorine Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	.170 (.160-.180)	.280 (.260-.310)	.370 (.330-.420)	2522
	05-06	*	< LOD	.160 (.150-.180)	.290 (.260-.320)	.410 (.360-.450)	2548
	07-08	*	< LOD	.150 (<LOD-.160)	.280 (.230-.320)	.390 (.330-.470)	2604
	09-10	*	< LOD	< LOD	.260 (.240-.280)	.350 (.320-.390)	2749
Age group							
6-11 years	03-04	*	< LOD	.180 (.150-.230)	.290 (.250-.320)	.370 (.310-.540)	314
	05-06	*	< LOD	.180 (.140-.200)	.310 (.210-.450)	.450 (.320-.610)	356
	07-08	*	< LOD	.180 (<LOD-.190)	.270 (.210-.390)	.430 (.270-.580)	389
	09-10	*	< LOD	< LOD	.320 (.240-.350)	.390 (.320-.470)	415
12-19 years	03-04	*	< LOD	.120 (.100-.140)	.200 (.170-.220)	.240 (.220-.280)	720
	05-06	*	< LOD	.120 (.110-.130)	.210 (.180-.240)	.290 (.240-.330)	702
	07-08	*	< LOD	.100 (<LOD-.120)	.170 (.150-.210)	.250 (.170-.310)	401
	09-10	*	< LOD	.110 (<LOD-.130)	.190 (.150-.250)	.280 (.190-.380)	420
20 years and older	03-04	*	< LOD	.180 (.160-.180)	.290 (.270-.320)	.390 (.350-.470)	1488
	05-06	*	< LOD	.170 (.150-.190)	.300 (.260-.330)	.410 (.370-.470)	1490
	07-08	*	< LOD	.150 (<LOD-.180)	.290 (.230-.350)	.410 (.340-.500)	1814
	09-10	*	< LOD	< LOD	.270 (.240-.280)	.350 (.320-.410)	1914
Gender							
Males	03-04	*	< LOD	.130 (.110-.150)	.230 (.190-.260)	.320 (.270-.350)	1230
	05-06	*	< LOD	.130 (.120-.140)	.220 (.190-.240)	.310 (.260-.360)	1270
	07-08	*	< LOD	.110 (<LOD-.120)	.190 (.180-.230)	.300 (.230-.340)	1294
	09-10	*	< LOD	< LOD	.210 (.190-.230)	.280 (.240-.330)	1399
Females	03-04	*	< LOD	.200 (.180-.210)	.320 (.290-.350)	.440 (.350-.510)	1292
	05-06	*	< LOD	.210 (.180-.230)	.350 (.300-.410)	.470 (.410-.550)	1278
	07-08	*	< LOD	.190 (<LOD-.230)	.330 (.280-.410)	.470 (.370-.580)	1310
	09-10	*	< LOD	.180 (<LOD-.190)	.300 (.280-.330)	.420 (.330-.470)	1350
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	.140 (<LOD-.150)	.240 (.200-.280)	.330 (.280-.460)	616
	05-06	*	< LOD	.140 (<LOD-.160)	.240 (.190-.320)	.350 (.290-.380)	637
	07-08	*	< LOD	< LOD	.210 (.180-.250)	.270 (.240-.320)	531
	09-10	*	< LOD	< LOD	.190 (<LOD-.230)	.290 (.220-.330)	566
Non-Hispanic blacks	03-04	*	< LOD	.120 (.100-.150)	.230 (.170-.290)	.310 (.230-.390)	635
	05-06	*	< LOD	.110 (.100-.140)	.210 (.170-.260)	.320 (.260-.360)	678
	07-08	*	< LOD	.120 (.100-.140)	.200 (.170-.250)	.290 (.230-.420)	597
	09-10	*	< LOD	.090 (<LOD-.110)	.160 (.130-.200)	.220 (.160-.380)	516
Non-Hispanic whites	03-04	*	< LOD	.180 (.160-.190)	.290 (.260-.320)	.370 (.340-.440)	1076
	05-06	*	< LOD	.180 (.160-.190)	.300 (.270-.350)	.410 (.360-.500)	1038
	07-08	*	< LOD	.160 (<LOD-.190)	.300 (.250-.360)	.440 (.330-.510)	1077
	09-10	*	< LOD	< LOD	.280 (.250-.320)	.370 (.330-.410)	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Trichlorophenols_BiomonitoringSummary.html

Urinary 2,4,6-Trichlorophenol

Metabolite of Several Organochlorine Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	.500 (<LOD-.600)	1.00 (.800-1.20)	1.40 (1.20-1.80)	2525
	05-06	*	< LOD	.600 (<LOD-.700)	1.00 (.800-1.20)	1.40 (1.20-1.80)	2548
	07-08	*	< LOD	< LOD	.800 (.700-.900)	1.20 (1.00-1.30)	2604
	09-10	*	< LOD	.500 (<LOD-.600)	.800 (.700-.900)	1.10 (1.00-1.40)	2749
Age group							
6-11 years	03-04	*	< LOD	.600 (.500-.700)	1.10 (.800-1.40)	1.90 (1.10-3.10)	314
	05-06	*	< LOD	.700 (.600-.900)	1.30 (1.00-2.30)	2.70 (1.30-5.40)	356
	07-08	*	< LOD	.600 (<LOD-.700)	1.10 (.900-1.40)	1.60 (1.30-2.10)	389
	09-10	*	< LOD	.500 (<LOD-.600)	.900 (.700-1.20)	1.30 (.900-2.20)	415
12-19 years	03-04	*	< LOD	.600 (.500-.800)	1.20 (.900-1.70)	1.80 (1.50-2.10)	722
	05-06	*	< LOD	.600 (<LOD-.800)	1.00 (.800-1.30)	1.30 (1.20-1.70)	702
	07-08	*	< LOD	.600 (<LOD-.700)	.800 (.700-1.10)	1.10 (.800-1.70)	401
	09-10	*	< LOD	.600 (<LOD-.700)	.900 (.700-1.30)	1.30 (.900-1.90)	420
20 years and older	03-04	*	< LOD	.500 (<LOD-.600)	1.00 (.800-1.10)	1.30 (1.10-1.70)	1489
	05-06	*	< LOD	.600 (<LOD-.700)	1.00 (.800-1.20)	1.30 (1.20-1.80)	1490
	07-08	*	< LOD	< LOD	.800 (.700-.900)	1.10 (.900-1.30)	1814
	09-10	*	< LOD	< LOD	.800 (.700-.900)	1.10 (1.00-1.20)	1914
Gender							
Males	03-04	*	< LOD	.600 (<LOD-.600)	1.00 (.800-1.10)	1.30 (1.10-1.80)	1231
	05-06	*	< LOD	.600 (<LOD-.800)	1.10 (.900-1.30)	1.60 (1.20-2.00)	1270
	07-08	*	< LOD	.500 (<LOD-.600)	.900 (.700-1.00)	1.20 (1.10-1.40)	1294
	09-10	*	< LOD	< LOD	.800 (.700-.900)	1.10 (1.00-1.20)	1399
Females	03-04	*	< LOD	.500 (<LOD-.600)	1.10 (.900-1.20)	1.40 (1.10-2.00)	1294
	05-06	*	< LOD	.500 (<LOD-.600)	.900 (.800-1.20)	1.30 (1.10-1.70)	1278
	07-08	*	< LOD	< LOD	.800 (.700-.900)	1.10 (.900-1.40)	1310
	09-10	*	< LOD	.500 (<LOD-.600)	.800 (.700-1.00)	1.10 (.900-1.60)	1350
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	.700 (.600-.800)	1.20 (1.10-1.60)	1.80 (1.30-2.00)	617
	05-06	*	< LOD	.600 (.500-.700)	1.00 (.800-1.20)	1.30 (1.20-1.70)	637
	07-08	*	< LOD	< LOD	.700 (.700-.900)	1.00 (.900-1.20)	531
	09-10	*	< LOD	.500 (<LOD-.600)	.900 (.700-1.10)	1.10 (.900-1.30)	566
Non-Hispanic blacks	03-04	*	< LOD	.900 (.700-1.00)	1.40 (1.10-1.90)	2.00 (1.50-2.70)	636
	05-06	*	< LOD	.800 (.700-1.10)	1.50 (1.20-1.90)	2.20 (1.60-3.30)	678
	07-08	*	< LOD	.600 (.500-.600)	1.00 (.900-1.10)	1.30 (1.10-1.60)	597
	09-10	*	< LOD	.700 (.600-.800)	1.10 (.900-1.30)	1.50 (1.20-2.00)	516
Non-Hispanic whites	03-04	*	< LOD	< LOD	.800 (.700-1.00)	1.20 (1.00-1.50)	1077
	05-06	*	< LOD	.500 (<LOD-.700)	.900 (.700-1.30)	1.30 (1.10-1.80)	1038
	07-08	*	< LOD	< LOD	.800 (.700-.900)	1.20 (1.00-1.40)	1077
	09-10	*	< LOD	< LOD	.700 (.600-.900)	1.00 (.800-1.30)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.5, 0.5, 0.5, and 0.5 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Trichlorophenols_BiomonitoringSummary.html

Urinary 2,4,6-Trichlorophenol (creatinine corrected)

Metabolite of Several Organochlorine Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	.710 (<LOD-.780)	1.25 (1.17-1.35)	1.75 (1.59-2.06)	2522
	05-06	*	< LOD	.720 (<LOD-.760)	1.27 (1.17-1.38)	1.75 (1.59-1.94)	2548
	07-08	*	< LOD	< LOD	1.25 (1.06-1.42)	1.75 (1.52-2.19)	2604
	09-10	*	< LOD	.710 (<LOD-.760)	1.21 (1.13-1.35)	1.67 (1.55-1.84)	2749
Age group							
6-11 years	03-04	*	< LOD	.920 (.850-1.13)	1.59 (1.22-1.91)	2.11 (1.46-4.55)	314
	05-06	*	< LOD	.880 (.740-1.06)	1.59 (1.21-2.06)	2.50 (1.61-5.20)	356
	07-08	*	< LOD	.900 (<LOD-.930)	1.46 (1.14-1.67)	2.33 (1.52-2.92)	389
	09-10	*	< LOD	.850 (<LOD-1.03)	1.59 (1.17-1.74)	1.85 (1.67-2.33)	415
12-19 years	03-04	*	< LOD	.580 (.510-.660)	.970 (.830-1.10)	1.21 (1.09-1.49)	720
	05-06	*	< LOD	.550 (<LOD-.630)	.970 (.690-1.17)	1.40 (1.08-1.59)	702
	07-08	*	< LOD	.550 (<LOD-.610)	.830 (.730-1.03)	1.30 (.930-1.48)	401
	09-10	*	< LOD	.590 (<LOD-.700)	.970 (.690-1.23)	1.23 (.920-1.66)	420
20 years and older	03-04	*	< LOD	.710 (<LOD-.770)	1.25 (1.17-1.35)	1.75 (1.59-2.00)	1488
	05-06	*	< LOD	.730 (<LOD-.780)	1.30 (1.17-1.40)	1.75 (1.57-2.06)	1490
	07-08	*	< LOD	< LOD	1.30 (1.06-1.46)	1.84 (1.52-2.33)	1814
	09-10	*	< LOD	< LOD	1.21 (1.13-1.35)	1.67 (1.52-1.94)	1914
Gender							
Males	03-04	*	< LOD	.560 (<LOD-.600)	.920 (.820-1.10)	1.30 (1.17-1.46)	1230
	05-06	*	< LOD	.600 (<LOD-.650)	1.00 (.850-1.13)	1.43 (1.25-1.59)	1270
	07-08	*	< LOD	.530 (<LOD-.600)	.930 (.830-1.06)	1.46 (1.18-1.59)	1294
	09-10	*	< LOD	< LOD	.980 (.850-1.12)	1.33 (1.13-1.59)	1399
Females	03-04	*	< LOD	.900 (<LOD-.960)	1.59 (1.35-1.75)	2.19 (1.75-2.63)	1292
	05-06	*	< LOD	.850 (<LOD-.970)	1.46 (1.30-1.67)	2.06 (1.67-2.50)	1278
	07-08	*	< LOD	< LOD	1.46 (1.23-1.75)	2.19 (1.67-2.50)	1310
	09-10	*	< LOD	.850 (<LOD-.920)	1.46 (1.30-1.59)	2.06 (1.62-2.33)	1350
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	.760 (.600-.900)	1.15 (.970-1.38)	1.59 (1.18-2.42)	616
	05-06	*	< LOD	.650 (.570-.700)	1.03 (.860-1.19)	1.46 (1.11-1.94)	637
	07-08	*	< LOD	< LOD	1.05 (.910-1.17)	1.35 (1.13-1.57)	531
	09-10	*	< LOD	.690 (<LOD-.760)	1.00 (.900-1.33)	1.67 (1.13-2.18)	566
Non-Hispanic blacks	03-04	*	< LOD	.600 (.560-.640)	.950 (.800-1.10)	1.34 (1.06-1.67)	635
	05-06	*	< LOD	.630 (.540-.740)	1.03 (.830-1.49)	1.59 (1.13-2.07)	678
	07-08	*	< LOD	.520 (.470-.620)	1.00 (.830-1.13)	1.40 (1.13-1.52)	597
	09-10	*	< LOD	.560 (.490-.610)	.830 (.750-.950)	1.17 (.920-1.52)	516
Non-Hispanic whites	03-04	*	< LOD	< LOD	1.30 (1.17-1.46)	1.75 (1.59-2.11)	1076
	05-06	*	< LOD	.760 (<LOD-.830)	1.35 (1.25-1.50)	1.79 (1.60-2.06)	1038
	07-08	*	< LOD	< LOD	1.35 (1.13-1.52)	1.84 (1.52-2.50)	1077
	09-10	*	< LOD	< LOD	1.30 (1.18-1.46)	1.73 (1.59-2.06)	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Trichlorophenols_BiomonitoringSummary.html

Urinary Acephate

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2491
	05-06	*	< LOD	< LOD	< LOD	< LOD	2546
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	309
	05-06	*	< LOD	< LOD	< LOD	< LOD	343
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	715
	05-06	*	< LOD	< LOD	< LOD	< LOD	696
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1467
	05-06	*	< LOD	< LOD	< LOD	< LOD	1507
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1221
	05-06	*	< LOD	< LOD	< LOD	< LOD	1205
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1270
	05-06	*	< LOD	< LOD	< LOD	< LOD	1341
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	608
	05-06	*	< LOD	< LOD	< LOD	< LOD	666
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	647
	05-06	*	< LOD	< LOD	< LOD	< LOD	698
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1077
	05-06	*	< LOD	< LOD	< LOD	< LOD	979

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.1 and 0.25 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Acephate_BiomonitoringSummary.html

Urinary Acephate (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2489
	05-06	*	< LOD	< LOD	< LOD	< LOD	2546
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	309
	05-06	*	< LOD	< LOD	< LOD	< LOD	343
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	714
	05-06	*	< LOD	< LOD	< LOD	< LOD	696
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1466
	05-06	*	< LOD	< LOD	< LOD	< LOD	1507
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1220
	05-06	*	< LOD	< LOD	< LOD	< LOD	1205
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1269
	05-06	*	< LOD	< LOD	< LOD	< LOD	1341
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	607
	05-06	*	< LOD	< LOD	< LOD	< LOD	666
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	647
	05-06	*	< LOD	< LOD	< LOD	< LOD	698
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1076
	05-06	*	< LOD	< LOD	< LOD	< LOD	979

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Acephate_BiomonitoringSummary.html

Urinary Dimethoate

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2455
	05-06	*	< LOD	< LOD	< LOD	< LOD	2628
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	306
	05-06	*	< LOD	< LOD	< LOD	< LOD	349
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	708
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1441
	05-06	*	< LOD	< LOD	< LOD	< LOD	1561
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1197
	05-06	*	< LOD	< LOD	< LOD	< LOD	1245
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1258
	05-06	*	< LOD	< LOD	< LOD	< LOD	1383
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	605
	05-06	*	< LOD	< LOD	< LOD	< LOD	693
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	629
	05-06	*	< LOD	< LOD	< LOD	< LOD	719
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1068
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.1 and 0.25 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DimethoateOmethoate_BiomonitoringSummary.html

Urinary Dimethoate (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2453
	05-06	*	< LOD	< LOD	< LOD	< LOD	2628
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	306
	05-06	*	< LOD	< LOD	< LOD	< LOD	349
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	707
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1440
	05-06	*	< LOD	< LOD	< LOD	< LOD	1561
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1196
	05-06	*	< LOD	< LOD	< LOD	< LOD	1245
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1257
	05-06	*	< LOD	< LOD	< LOD	< LOD	1383
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	604
	05-06	*	< LOD	< LOD	< LOD	< LOD	693
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	629
	05-06	*	< LOD	< LOD	< LOD	< LOD	719
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1067
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DimethoateOmethoate_BiomonitoringSummary.html

Urinary Methamidophos

Also a metabolite of Acephate

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2379
	05-06	*	< LOD	< LOD	< LOD	< LOD	2591
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	295
	05-06	*	< LOD	< LOD	< LOD	< LOD	346
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	683
	05-06	*	< LOD	< LOD	< LOD	< LOD	710
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1401
	05-06	*	< LOD	< LOD	< LOD	< LOD	1535
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1171
	05-06	*	< LOD	< LOD	< LOD	< LOD	1232
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1208
	05-06	*	< LOD	< LOD	< LOD	< LOD	1359
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	564
	05-06	*	< LOD	< LOD	< LOD	< LOD	688
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	631
	05-06	*	< LOD	< LOD	< LOD	< LOD	717
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1034
	05-06	*	< LOD	< LOD	< LOD	< LOD	979

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.1 and 0.37 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Methamidophos_BiomonitoringSummary.html

Urinary Methamidophos (creatinine corrected)

Also a metabolite of Acephate

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2377
	05-06	*	< LOD	< LOD	< LOD	< LOD	2591
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	295
	05-06	*	< LOD	< LOD	< LOD	< LOD	346
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	682
	05-06	*	< LOD	< LOD	< LOD	< LOD	710
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1400
	05-06	*	< LOD	< LOD	< LOD	< LOD	1535
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1170
	05-06	*	< LOD	< LOD	< LOD	< LOD	1232
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1207
	05-06	*	< LOD	< LOD	< LOD	< LOD	1359
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	563
	05-06	*	< LOD	< LOD	< LOD	< LOD	688
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	631
	05-06	*	< LOD	< LOD	< LOD	< LOD	717
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1033
	05-06	*	< LOD	< LOD	< LOD	< LOD	979

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Methamidophos_BiomonitoringSummary.html

Urinary Omethoate

Metabolite of Dimethoate

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2455
	05-06	*	< LOD	< LOD	< LOD	< LOD	2628
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	303
	05-06	*	< LOD	< LOD	< LOD	< LOD	349
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	707
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1445
	05-06	*	< LOD	< LOD	< LOD	< LOD	1561
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1207
	05-06	*	< LOD	< LOD	< LOD	< LOD	1245
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1248
	05-06	*	< LOD	< LOD	< LOD	< LOD	1383
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	606
	05-06	*	< LOD	< LOD	< LOD	< LOD	693
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	637
	05-06	*	< LOD	< LOD	< LOD	< LOD	719
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1055
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.1 and 0.14 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DimethoateOmethoate_BiomonitoringSummary.html

Urinary Omethoate (creatinine corrected)

Metabolite of Dimethoate

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2453
	05-06	*	< LOD	< LOD	< LOD	< LOD	2628
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	303
	05-06	*	< LOD	< LOD	< LOD	< LOD	349
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	706
	05-06	*	< LOD	< LOD	< LOD	< LOD	718
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1444
	05-06	*	< LOD	< LOD	< LOD	< LOD	1561
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1206
	05-06	*	< LOD	< LOD	< LOD	< LOD	1245
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1247
	05-06	*	< LOD	< LOD	< LOD	< LOD	1383
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	605
	05-06	*	< LOD	< LOD	< LOD	< LOD	693
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	637
	05-06	*	< LOD	< LOD	< LOD	< LOD	719
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1054
	05-06	*	< LOD	< LOD	< LOD	< LOD	1007

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DimethoateOmethoate_BiomonitoringSummary.html

Urinary Diethylphosphate (DEP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	1.03 (.670-1.58)	1.20 (.750-1.70)	3.20 (2.30-4.80)	7.60 (5.00-12.0)	13.0 (7.70-23.0)	1949
	01-02	*	< LOD	2.90 (2.50-3.26)	6.68 (5.78-7.52)	10.6 (9.14-12.0)	3018
	03-04	*	< LOD	4.56 (3.57-5.83)	10.2 (9.00-11.5)	15.5 (13.5-16.8)	2453
	05-06	*	< LOD	1.46 (.780-2.17)	7.01 (5.67-8.94)	13.3 (10.2-16.0)	2635
	07-08	*	< LOD	2.48 (1.44-3.33)	8.74 (7.51-10.3)	15.3 (13.1-17.8)	2591
Age group							
6-11 years	99-00	1.32 (.757-2.29)	1.50 (.860-2.60)	4.50 (2.10-7.90)	11.0 (4.80-24.0)	16.0 (8.50-36.0)	471
	01-02	*	.290 (<LOD-1.04)	3.45 (2.36-4.47)	9.56 (6.33-18.0)	20.0 (9.44-38.2)	576
	03-04	*	< LOD	5.13 (2.48-7.94)	10.9 (8.08-15.9)	16.1 (10.9-18.5)	308
	05-06	*	< LOD	1.45 (<LOD-4.51)	8.21 (4.85-13.4)	13.5 (8.36-20.4)	350
	07-08	*	< LOD	4.31 (.420-6.49)	12.9 (9.14-17.7)	20.2 (11.9-37.5)	385
12-19 years	99-00	1.21 (.758-1.94)	1.40 (.970-2.10)	3.70 (2.40-5.50)	8.00 (4.70-19.0)	20.0 (8.00-27.0)	664
	01-02	*	< LOD	2.86 (1.96-3.95)	7.58 (5.71-9.15)	11.0 (9.35-12.4)	822
	03-04	*	.530 (<LOD-2.32)	5.80 (4.34-7.67)	14.8 (9.12-19.8)	20.8 (14.8-32.7)	701
	05-06	*	< LOD	.460 (<LOD-3.69)	9.47 (4.67-14.1)	14.4 (10.2-22.4)	718
	07-08	*	< LOD	3.81 (<LOD-5.42)	10.7 (7.76-14.2)	23.3 (11.3-35.6)	391
20-59 years	99-00	.955 (.623-1.47)	1.10 (.700-1.60)	3.00 (1.80-4.80)	7.30 (4.70-11.0)	11.0 (6.80-22.0)	814
	01-02	*	< LOD	2.71 (2.34-3.12)	5.79 (5.05-7.21)	10.4 (7.43-12.3)	1122
	03-04	*	< LOD	4.37 (3.02-5.81)	9.74 (8.35-11.3)	14.2 (11.5-16.2)	922
	05-06	*	< LOD	1.13 (<LOD-2.17)	6.15 (4.51-8.24)	11.5 (8.23-15.5)	1092
	07-08	*	< LOD	1.55 (.470-2.48)	7.54 (6.10-8.98)	14.0 (9.62-15.7)	1178
60 years and older	01-02	*	.440 (<LOD-1.63)	3.30 (2.63-4.51)	7.06 (5.17-8.69)	9.41 (8.31-10.6)	498
	03-04	*	1.80 (<LOD-2.98)	4.73 (3.98-5.91)	10.0 (8.09-11.3)	13.6 (10.6-16.7)	522
	05-06	*	< LOD	2.69 (1.71-3.95)	8.33 (6.60-12.0)	16.3 (12.0-18.7)	475
	07-08	*	< LOD	3.55 (2.80-5.31)	9.86 (7.53-14.2)	17.1 (12.6-22.1)	637
Gender							
Males	99-00	1.11 (.717-1.73)	1.20 (.810-1.70)	3.80 (2.50-5.00)	8.00 (5.00-19.0)	19.0 (7.20-30.0)	952
	01-02	*	< LOD	3.21 (2.72-3.56)	6.99 (5.83-7.71)	11.5 (9.36-12.1)	1420
	03-04	*	< LOD	5.01 (3.90-6.50)	11.0 (10.1-12.7)	16.2 (14.3-18.8)	1198
	05-06	*	< LOD	.770 (<LOD-1.71)	6.82 (5.14-9.54)	13.4 (9.36-16.1)	1246
	07-08	*	< LOD	1.95 (<LOD-3.34)	9.39 (7.90-11.8)	17.0 (13.7-21.5)	1295
Females	99-00	.954 (.599-1.52)	1.20 (.620-1.70)	2.90 (1.90-4.80)	7.50 (4.60-11.0)	11.0 (7.40-16.0)	997
	01-02	*	.290 (<LOD-.790)	2.63 (2.22-3.03)	6.21 (4.59-8.64)	10.2 (7.52-14.7)	1598
	03-04	*	.110 (<LOD-2.29)	4.29 (3.21-5.80)	9.38 (7.95-10.7)	13.3 (11.2-15.9)	1255
	05-06	*	< LOD	2.15 (1.27-2.76)	7.23 (5.79-9.32)	12.9 (9.64-16.0)	1389
	07-08	*	< LOD	2.76 (1.69-3.62)	7.76 (6.23-10.7)	14.5 (10.7-18.8)	1296

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.2, 0.2, 0.1, 0.37, and 0.37 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethylphosphate (DEP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Mexican Americans	99-00	1.22 (.740-2.01)	1.20 (.840-1.80)	4.10 (2.20-7.00)	11.0 (6.40-19.0)	18.0 (12.0-27.0)	672
	01-02	*	.600 (<LOD-1.49)	3.07 (2.22-3.83)	6.26 (5.00-7.82)	11.2 (7.82-12.3)	763
	03-04	*	1.18 (<LOD-2.97)	5.81 (3.99-7.59)	11.0 (8.48-16.1)	16.9 (12.1-24.0)	581
	05-06	*	< LOD	< LOD	6.30 (1.57-11.6)	14.9 (5.72-33.5)	695
	07-08	*	< LOD	1.63 (<LOD-3.88)	8.36 (6.44-9.61)	14.8 (8.85-25.1)	501
Non-Hispanic blacks	99-00	1.56 (1.13-2.14)	1.60 (1.40-1.80)	4.30 (2.90-5.80)	10.0 (5.60-18.0)	18.0 (8.00-27.0)	509
	01-02	*	1.03 (<LOD-2.33)	4.81 (3.48-6.52)	9.98 (7.39-13.3)	15.1 (9.81-23.4)	756
	03-04	*	1.04 (<LOD-3.28)	6.29 (5.05-8.44)	11.9 (9.92-15.5)	16.2 (14.1-22.5)	628
	05-06	*	< LOD	2.10 (<LOD-4.57)	8.87 (6.52-12.0)	14.3 (11.2-19.1)	723
	07-08	*	< LOD	3.92 (2.67-5.19)	11.3 (9.44-14.0)	19.5 (14.1-22.0)	574
Non-Hispanic whites	99-00	.981 (.579-1.66)	1.10 (.490-2.10)	3.30 (2.20-4.90)	7.70 (4.70-14.0)	14.0 (7.60-25.0)	595
	01-02	*	< LOD	2.63 (2.21-3.13)	5.79 (5.05-7.35)	9.90 (8.38-11.7)	1253
	03-04	*	< LOD	4.25 (3.12-5.77)	9.74 (8.37-11.3)	14.4 (12.4-16.0)	1055
	05-06	*	< LOD	1.50 (.900-2.27)	6.82 (5.08-9.74)	13.3 (8.37-16.9)	1007
	07-08	*	< LOD	2.14 (1.27-3.03)	7.57 (6.57-8.97)	14.5 (11.4-16.5)	1086

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.2, 0.2, 0.1, 0.37, and 0.37 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethylphosphate (DEP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.924 (.608-1.41)	.920 (.570-1.40)	2.73 (1.68-4.60)	7.94 (4.40-12.2)	12.2 (8.00-19.6)	1949
	01-02	*	< LOD	2.56 (2.17-2.85)	5.36 (4.75-6.06)	8.72 (7.19-10.3)	3017
	03-04	*	< LOD	4.64 (3.82-5.79)	9.18 (7.85-10.8)	13.1 (10.5-16.1)	2450
	05-06	*	< LOD	2.23 (1.49-3.09)	7.72 (6.26-9.36)	13.5 (11.0-19.0)	2634
	07-08	*	< LOD	3.20 (2.37-4.12)	8.86 (7.52-10.7)	15.7 (11.9-19.8)	2589
Age group							
6-11 years	99-00	1.43 (.870-2.34)	1.47 (1.02-2.41)	3.94 (2.20-8.57)	10.5 (4.55-20.8)	16.6 (10.5-32.7)	471
	01-02	*	.886 (<LOD-1.76)	4.02 (2.87-5.25)	8.85 (6.88-15.6)	18.4 (9.40-28.8)	576
	03-04	*	< LOD	6.10 (3.79-9.03)	11.9 (9.54-15.2)	16.1 (11.9-28.3)	308
	05-06	*	< LOD	3.09 (<LOD-6.77)	9.50 (5.67-17.6)	20.7 (9.50-26.5)	350
	07-08	*	< LOD	4.67 (2.01-6.96)	17.2 (13.1-20.3)	28.2 (19.3-34.8)	385
12-19 years	99-00	.818 (.533-1.26)	.790 (.560-1.25)	2.35 (1.37-3.75)	5.44 (2.82-14.4)	12.4 (4.66-34.2)	664
	01-02	*	< LOD	2.05 (1.54-2.67)	4.40 (3.40-5.28)	7.28 (5.28-9.75)	821
	03-04	*	.438 (<LOD-2.05)	4.47 (3.90-5.43)	10.1 (7.10-13.5)	14.7 (9.82-26.4)	699
	05-06	*	< LOD	1.27 (<LOD-2.81)	6.74 (4.24-9.87)	11.7 (6.98-16.6)	718
	07-08	*	< LOD	2.86 (<LOD-4.53)	8.56 (5.70-17.0)	17.0 (8.05-23.0)	389
20-59 years	99-00	.883 (.574-1.36)	.860 (.500-1.35)	2.66 (1.54-4.95)	7.37 (4.32-12.1)	12.1 (8.00-17.5)	814
	01-02	*	< LOD	2.28 (2.01-2.56)	4.75 (3.92-5.83)	7.37 (5.93-9.72)	1122
	03-04	*	< LOD	4.29 (2.98-5.71)	8.35 (6.09-11.3)	11.9 (9.02-14.7)	921
	05-06	*	< LOD	1.92 (<LOD-2.95)	6.64 (5.22-9.25)	12.3 (9.25-19.4)	1091
	07-08	*	< LOD	2.38 (1.81-3.27)	7.45 (6.11-9.43)	11.4 (9.22-15.8)	1178
60+ years	01-02	*	1.00 (<LOD-1.88)	3.29 (2.74-4.14)	6.52 (4.92-8.21)	9.82 (7.05-14.4)	498
	03-04	*	2.39 (<LOD-3.35)	5.65 (4.36-7.50)	9.44 (8.38-10.5)	13.3 (10.5-18.2)	522
	05-06	*	< LOD	3.75 (2.30-5.14)	9.88 (7.72-12.6)	19.3 (12.6-26.6)	475
	07-08	*	< LOD	5.28 (4.03-6.23)	11.8 (9.36-14.4)	17.7 (13.2-21.9)	637
Gender							
Males	99-00	.855 (.566-1.29)	.820 (.510-1.34)	2.61 (1.76-4.03)	7.69 (4.41-12.1)	12.2 (6.94-23.8)	952
	01-02	*	< LOD	2.20 (1.86-2.64)	4.49 (3.92-5.31)	7.50 (5.79-9.10)	1420
	03-04	*	< LOD	4.20 (3.28-5.30)	8.35 (7.10-10.1)	12.0 (9.33-14.7)	1197
	05-06	*	< LOD	1.16 (<LOD-2.25)	6.24 (4.58-7.63)	10.0 (7.60-17.8)	1246
	07-08	*	< LOD	2.17 (<LOD-3.23)	8.14 (6.35-10.7)	13.6 (9.79-18.6)	1294
Females	99-00	.996 (.620-1.60)	.960 (.540-1.62)	2.81 (1.45-5.85)	8.00 (4.00-13.0)	12.1 (6.67-19.6)	997
	01-02	*	.778 (<LOD-1.31)	2.81 (2.44-3.33)	6.49 (4.95-7.76)	9.61 (7.12-13.6)	1597
	03-04	*	.636 (<LOD-2.84)	5.06 (4.03-6.86)	9.85 (8.25-11.8)	13.8 (10.9-20.3)	1253
	05-06	*	< LOD	3.50 (2.43-4.11)	9.36 (6.91-11.8)	16.5 (12.4-22.3)	1388
	07-08	*	< LOD	4.09 (2.80-4.91)	9.46 (8.43-11.5)	16.5 (13.1-20.9)	1295

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethylphosphate (DEP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	1.09 (.633-1.89)	1.05 (.650-1.98)	3.78 (2.11-6.46)	9.84 (5.66-15.7)	15.7 (8.61-29.0)	672
	01-02	*	.891 (<LOD-1.41)	2.39 (1.77-3.14)	5.17 (4.10-6.53)	8.04 (6.40-10.9)	763
	03-04	*	1.07 (<LOD-2.79)	4.94 (3.30-7.01)	9.94 (7.89-13.3)	16.6 (10.1-22.4)	580
	05-06	*	< LOD	< LOD	6.26 (3.71-10.1)	14.9 (5.75-37.3)	695
	07-08	*	< LOD	3.19 (<LOD-4.27)	8.56 (6.00-10.3)	13.3 (9.77-15.9)	500
Non-Hispanic blacks	99-00	1.07 (.773-1.47)	1.18 (.830-1.54)	2.61 (1.89-3.47)	5.98 (3.94-9.56)	11.9 (5.98-22.2)	509
	01-02	*	.797 (<LOD-1.62)	2.98 (2.57-3.61)	6.91 (5.17-8.69)	9.58 (7.82-12.1)	755
	03-04	*	.741 (<LOD-2.21)	4.25 (3.46-5.36)	8.30 (6.57-10.7)	11.7 (10.5-13.2)	627
	05-06	*	< LOD	1.92 (<LOD-3.58)	7.08 (5.16-9.35)	11.4 (8.91-12.3)	722
	07-08	*	< LOD	3.33 (2.33-4.15)	8.72 (7.27-10.2)	16.7 (11.5-19.4)	573
Non-Hispanic whites	99-00	.932 (.549-1.58)	.900 (.430-1.68)	2.87 (1.51-5.88)	8.57 (4.40-14.4)	13.0 (8.21-23.8)	595
	01-02	*	< LOD	2.52 (2.08-2.95)	5.26 (4.57-5.93)	8.26 (6.61-10.8)	1253
	03-04	*	< LOD	4.65 (3.73-5.87)	9.12 (7.79-11.2)	12.7 (10.1-15.7)	1054
	05-06	*	< LOD	2.50 (1.74-3.30)	7.72 (6.24-9.25)	13.9 (10.0-19.5)	1007
	07-08	*	< LOD	3.10 (2.19-4.18)	8.44 (6.84-9.74)	13.6 (10.9-17.5)	1086

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethylphosphate (DMP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	.740 (<LOD-1.40)	2.90 (2.10-4.00)	7.90 (6.20-8.90)	14.0 (10.0-19.0)	1949
	01-02	*	< LOD	3.32 (2.90-3.74)	8.40 (7.11-9.23)	13.4 (11.4-14.9)	3017
	03-04	*	< LOD	4.15 (3.39-4.94)	9.54 (7.46-11.9)	15.1 (12.8-18.2)	2494
	05-06	*	< LOD	5.21 (4.15-6.56)	17.2 (13.8-20.5)	29.6 (24.3-35.2)	2635
	07-08	*	< LOD	7.91 (6.64-9.07)	20.8 (18.2-24.4)	35.6 (30.3-39.4)	2593
Age group							
6-11 years	99-00	1.58 (1.15-2.18)	1.10 (.580-2.20)	4.40 (2.80-6.80)	10.0 (7.80-21.0)	22.0 (15.0-33.0)	471
	01-02	*	.970 (<LOD-2.00)	5.04 (3.31-7.66)	12.2 (9.10-15.1)	18.3 (12.6-41.7)	576
	03-04	*	< LOD	4.53 (3.34-5.96)	11.0 (5.62-17.9)	16.2 (7.46-28.3)	310
	05-06	*	< LOD	9.92 (3.24-20.0)	27.1 (16.5-46.6)	46.6 (27.1-63.0)	350
	07-08	*	< LOD	12.5 (7.22-16.4)	29.7 (20.3-39.1)	43.3 (29.7-62.7)	385
12-19 years	99-00	*	.670 (<LOD-1.80)	3.80 (2.50-4.90)	9.90 (6.20-18.0)	22.0 (13.0-29.0)	664
	01-02	*	.670 (<LOD-1.31)	4.27 (3.41-5.35)	9.27 (7.80-12.3)	14.7 (11.8-21.3)	822
	03-04	*	1.20 (<LOD-2.27)	4.61 (3.47-6.72)	10.9 (7.90-15.0)	20.9 (12.5-26.8)	717
	05-06	*	< LOD	7.32 (1.35-12.4)	22.6 (16.1-36.1)	44.0 (27.6-54.0)	718
	07-08	*	< LOD	10.0 (7.22-14.5)	25.5 (16.8-35.6)	36.2 (25.5-49.6)	391
20-59 years	99-00	*	.680 (<LOD-1.30)	2.70 (1.80-3.70)	6.60 (5.70-8.10)	9.70 (8.80-14.0)	814
	01-02	*	< LOD	2.95 (2.35-3.41)	6.95 (5.80-8.82)	11.5 (9.66-13.7)	1121
	03-04	*	< LOD	3.75 (2.84-4.88)	8.52 (6.86-10.7)	14.1 (10.8-17.5)	938
	05-06	*	< LOD	4.09 (2.45-6.03)	13.8 (9.82-19.8)	24.6 (19.8-32.8)	1092
	07-08	*	< LOD	6.07 (4.26-7.69)	18.0 (13.6-22.6)	30.3 (23.2-37.9)	1180
60 years and older	01-02	*	.700 (<LOD-1.50)	3.68 (2.91-4.56)	8.93 (6.79-10.7)	14.4 (9.60-19.1)	498
	03-04	*	2.04 (<LOD-2.63)	4.60 (3.83-5.15)	11.3 (7.68-13.4)	16.3 (12.4-19.4)	529
	05-06	*	< LOD	6.75 (4.75-9.32)	16.8 (11.7-22.1)	26.3 (18.9-38.8)	475
	07-08	*	< LOD	10.8 (8.64-13.0)	24.9 (19.2-32.4)	40.5 (32.5-47.4)	637
Gender							
Males	99-00	*	.670 (<LOD-1.30)	2.90 (2.20-4.00)	7.90 (6.00-9.30)	18.0 (10.0-24.0)	952
	01-02	*	< LOD	3.49 (2.90-4.15)	8.40 (6.95-10.3)	12.8 (11.3-14.7)	1420
	03-04	*	< LOD	4.03 (3.30-4.91)	8.49 (6.82-11.9)	15.1 (11.3-20.0)	1221
	05-06	*	< LOD	4.10 (2.37-6.40)	15.9 (12.4-20.9)	28.1 (22.3-36.7)	1246
	07-08	*	< LOD	7.75 (6.50-9.63)	20.8 (17.6-26.6)	36.1 (27.0-47.4)	1295
Females	99-00	*	.790 (<LOD-1.60)	2.90 (2.00-4.20)	7.80 (5.70-9.00)	11.0 (9.00-18.0)	997
	01-02	*	< LOD	3.06 (2.63-3.63)	8.38 (6.56-9.63)	13.7 (10.9-17.6)	1597
	03-04	*	1.08 (<LOD-1.92)	4.29 (3.31-5.28)	10.4 (7.64-12.9)	14.8 (13.3-18.0)	1273
	05-06	*	< LOD	6.40 (5.04-7.21)	18.7 (14.6-20.8)	30.9 (24.6-41.1)	1389
	07-08	*	< LOD	7.91 (6.18-9.36)	20.6 (16.5-25.5)	34.2 (26.6-41.5)	1298

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.58, 0.5, 0.5, 0.47, and 0.47 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethylphosphate (DMP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Mexican Americans	99-00	*	1.10 (<LOD-1.80)	3.80 (2.70-5.30)	9.60 (6.00-16.0)	16.0 (8.90-31.0)	672
	01-02	*	.710 (<LOD-1.52)	3.34 (2.45-4.42)	9.13 (7.29-10.7)	14.4 (11.0-20.6)	763
	03-04	*	< LOD	4.37 (3.12-6.86)	10.3 (7.85-17.4)	22.3 (9.61-31.6)	610
	05-06	*	< LOD	2.65 (<LOD-6.47)	18.9 (13.3-27.9)	39.0 (30.0-47.2)	695
	07-08	*	< LOD	6.85 (3.69-8.82)	20.9 (14.1-26.4)	35.6 (26.3-49.4)	501
Non-Hispanic blacks	99-00	1.42 (1.16-1.74)	1.00 (.650-1.50)	3.60 (2.50-5.50)	8.90 (6.90-15.0)	21.0 (14.0-24.0)	509
	01-02	*	.890 (<LOD-2.20)	4.71 (3.68-6.62)	10.7 (8.54-13.9)	19.4 (14.7-22.8)	755
	03-04	*	< LOD	5.28 (4.18-6.37)	10.8 (8.33-12.5)	15.8 (12.2-23.4)	631
	05-06	*	< LOD	9.16 (2.74-15.1)	22.7 (18.2-33.7)	42.5 (27.3-61.9)	723
	07-08	*	< LOD	8.67 (6.43-11.1)	23.7 (18.3-28.6)	38.0 (27.7-48.1)	574
Non-Hispanic whites	99-00	*	< LOD	2.90 (1.80-4.20)	7.90 (5.90-9.00)	11.0 (9.00-18.0)	595
	01-02	*	< LOD	3.13 (2.49-3.58)	7.79 (6.67-9.22)	12.5 (10.3-14.7)	1253
	03-04	*	< LOD	3.95 (3.23-4.91)	8.83 (6.56-12.5)	14.8 (11.9-20.0)	1061
	05-06	*	< LOD	5.04 (3.60-6.75)	15.3 (11.8-20.5)	27.8 (20.8-36.7)	1007
	07-08	*	< LOD	7.46 (6.27-9.03)	20.7 (17.5-25.5)	35.6 (28.7-39.1)	1088

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.58, 0.5, 0.5, 0.47, and 0.47 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethylphosphate (DMP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	.810 (<LOD-1.15)	2.93 (2.11-3.92)	8.50 (6.96-10.4)	16.1 (13.3-17.6)	1949
	01-02	*	< LOD	3.16 (2.69-3.59)	8.00 (7.12-9.28)	12.9 (11.5-15.0)	3016
	03-04	*	< LOD	4.05 (3.50-5.15)	9.96 (8.78-11.2)	15.0 (13.0-18.1)	2491
	05-06	*	< LOD	5.97 (4.75-7.43)	18.0 (15.9-20.5)	27.9 (24.0-32.7)	2634
	07-08	*	< LOD	9.10 (8.08-10.2)	20.0 (18.0-24.2)	33.6 (28.8-40.2)	2591
Age group							
6-11 years	99-00	1.71 (1.29-2.27)	1.38 (.890-2.38)	4.48 (2.88-7.89)	16.7 (8.21-21.2)	22.1 (19.2-30.1)	471
	01-02	*	1.93 (<LOD-2.97)	5.99 (4.32-8.28)	12.9 (9.34-18.5)	20.6 (13.3-34.8)	576
	03-04	*	< LOD	6.87 (3.80-9.00)	13.9 (9.68-19.3)	19.6 (13.9-25.9)	310
	05-06	*	< LOD	12.4 (6.71-21.9)	41.2 (23.4-62.2)	71.5 (37.9-107)	350
	07-08	*	< LOD	15.6 (10.8-21.6)	35.3 (27.6-39.2)	54.3 (36.5-67.1)	385
12-19 years	99-00	*	.590 (<LOD-.950)	2.28 (1.70-2.80)	7.78 (4.16-14.4)	16.0 (8.70-35.3)	664
	01-02	*	.913 (<LOD-1.27)	3.29 (2.76-3.78)	6.29 (5.51-7.30)	9.83 (7.94-14.2)	821
	03-04	*	1.00 (<LOD-1.68)	4.16 (3.37-5.07)	8.81 (7.78-10.2)	12.7 (10.3-17.7)	715
	05-06	*	< LOD	5.89 (2.00-10.1)	18.1 (13.9-24.0)	28.9 (19.9-37.9)	718
	07-08	*	< LOD	9.10 (5.62-10.9)	19.9 (15.5-22.5)	26.1 (19.8-27.2)	389
20-59 years	99-00	*	.760 (<LOD-1.12)	2.88 (1.89-3.99)	8.11 (5.89-10.3)	14.6 (10.4-16.8)	814
	01-02	*	< LOD	2.55 (2.05-3.03)	6.92 (5.85-8.00)	11.5 (9.38-13.6)	1121
	03-04	*	< LOD	3.77 (2.72-4.55)	8.78 (6.68-10.5)	13.5 (11.2-15.6)	937
	05-06	*	< LOD	4.71 (3.22-6.50)	16.3 (13.3-18.7)	24.0 (19.9-27.6)	1091
	07-08	*	< LOD	7.32 (6.02-8.62)	17.2 (13.0-20.0)	28.2 (19.3-40.7)	1180
60+ years	01-02	*	1.46 (<LOD-2.00)	4.31 (3.44-5.39)	10.6 (7.92-12.0)	14.3 (11.8-18.5)	498
	03-04	*	2.19 (<LOD-2.69)	5.88 (4.65-7.51)	12.3 (9.79-14.8)	17.4 (13.0-24.2)	529
	05-06	*	< LOD	9.16 (6.36-11.1)	19.7 (13.7-26.4)	30.5 (22.0-46.2)	475
	07-08	*	< LOD	12.9 (10.5-15.3)	29.9 (20.9-38.5)	45.5 (30.5-80.3)	637
Gender							
Males	99-00	*	.620 (<LOD-.940)	2.38 (1.86-3.18)	7.58 (4.64-11.6)	15.2 (9.74-19.5)	952
	01-02	*	< LOD	2.77 (2.32-3.19)	7.06 (5.44-8.80)	11.2 (9.70-12.7)	1420
	03-04	*	< LOD	3.27 (2.55-3.94)	8.58 (6.79-10.4)	12.5 (10.5-15.4)	1220
	05-06	*	< LOD	4.15 (2.16-5.95)	14.7 (11.1-18.0)	21.9 (18.9-26.9)	1246
	07-08	*	< LOD	7.08 (5.54-8.54)	17.5 (15.1-20.4)	28.0 (21.2-35.4)	1294
Females	99-00	*	1.00 (<LOD-1.71)	3.63 (2.30-5.19)	9.12 (7.82-11.7)	16.4 (11.7-19.7)	997
	01-02	*	< LOD	3.50 (2.86-4.49)	9.09 (7.83-10.8)	15.0 (12.5-17.8)	1596
	03-04	*	1.67 (<LOD-2.25)	5.28 (4.05-6.49)	11.2 (9.65-13.8)	18.1 (14.8-20.6)	1271
	05-06	*	< LOD	8.22 (6.99-9.20)	22.0 (17.7-24.9)	37.9 (27.0-47.2)	1388
	07-08	*	< LOD	10.7 (9.10-12.4)	24.3 (19.5-28.2)	38.6 (27.2-54.6)	1297

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethylphosphate (DMP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	1.06 (<LOD-1.55)	3.89 (2.54-5.45)	9.41 (7.69-11.5)	16.7 (11.7-23.6)	672
	01-02	*	.947 (<LOD-1.35)	3.18 (2.54-3.80)	8.44 (6.24-10.9)	14.3 (11.4-16.2)	763
	03-04	*	< LOD	4.16 (2.57-6.30)	12.3 (7.50-16.5)	18.8 (15.3-21.5)	609
	05-06	*	< LOD	4.27 (<LOD-8.50)	19.1 (14.9-22.7)	29.9 (23.8-39.0)	695
	07-08	*	< LOD	7.47 (5.57-9.46)	19.9 (16.8-24.7)	29.8 (24.7-33.4)	500
Non-Hispanic blacks	99-00	.973 (.780-1.21)	.690 (.530-1.06)	2.67 (1.89-3.77)	7.07 (5.09-11.4)	14.0 (10.6-19.1)	509
	01-02	*	.851 (<LOD-1.34)	3.36 (2.68-4.22)	7.63 (6.35-9.18)	13.2 (10.1-14.9)	754
	03-04	*	< LOD	3.70 (2.84-4.48)	7.52 (6.59-8.89)	12.7 (8.89-19.3)	630
	05-06	*	< LOD	5.89 (2.56-8.89)	18.7 (13.1-26.4)	29.9 (24.0-38.7)	722
	07-08	*	< LOD	6.75 (4.80-8.65)	20.1 (15.5-25.1)	30.7 (24.2-37.7)	573
Non-Hispanic whites	99-00	*	< LOD	3.15 (1.97-4.32)	8.73 (5.89-13.3)	15.8 (10.0-21.2)	595
	01-02	*	< LOD	3.12 (2.50-3.76)	8.30 (6.91-10.0)	13.6 (11.2-16.9)	1253
	03-04	*	< LOD	4.05 (3.49-5.64)	10.2 (8.83-11.8)	14.8 (12.1-19.2)	1060
	05-06	*	< LOD	5.97 (4.53-8.08)	17.7 (15.2-20.6)	26.3 (23.3-29.3)	1007
	07-08	*	< LOD	9.33 (7.92-10.8)	20.0 (17.5-24.2)	35.1 (27.2-43.9)	1088

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethylthiophosphate (DETP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	.500 (<LOD-.690)	.760 (.620-1.10)	1.40 (1.10-1.80)	2.20 (1.70-2.80)	1949
	01-02	.484 (.381-.615)	.630 (.450-.930)	1.53 (1.33-1.80)	2.57 (2.28-3.01)	3.94 (3.26-4.94)	3016
	03-04	*	< LOD	.830 (.710-.960)	1.83 (1.51-2.30)	3.00 (2.39-3.78)	2422
	05-06	*	< LOD	.620 (<LOD-.710)	1.43 (1.17-1.84)	2.55 (1.94-3.37)	2634
	07-08	*	< LOD	1.00 (.880-1.16)	2.18 (1.75-2.75)	4.35 (2.95-6.01)	2590
Age group							
6-11 years	99-00	*	.600 (<LOD-.810)	.910 (.720-1.30)	1.70 (1.20-3.20)	3.20 (1.70-7.30)	471
	01-02	.453 (.350-.585)	.550 (.350-.850)	1.58 (1.33-2.04)	2.75 (2.22-3.38)	4.08 (2.95-5.16)	575
	03-04	*	< LOD	.820 (.580-.970)	1.45 (1.05-2.16)	2.18 (1.45-4.13)	296
	05-06	*	< LOD	.700 (<LOD-.950)	1.38 (1.03-2.05)	2.15 (1.37-4.06)	350
	07-08	*	< LOD	1.18 (.990-1.42)	2.93 (2.06-5.58)	6.44 (2.87-13.2)	385
12-19 years	99-00	*	.210 (<LOD-.710)	.780 (.600-1.20)	1.50 (1.20-2.30)	2.30 (1.60-4.30)	664
	01-02	.505 (.388-.657)	.690 (.440-.960)	1.61 (1.32-1.94)	2.57 (2.23-3.39)	4.08 (2.73-5.86)	822
	03-04	*	.260 (<LOD-.400)	.930 (.750-1.13)	2.14 (1.75-2.89)	3.27 (2.69-4.83)	690
	05-06	*	< LOD	.700 (<LOD-.920)	1.42 (1.07-1.94)	2.46 (1.47-4.42)	717
	07-08	*	< LOD	1.04 (.880-1.45)	2.20 (1.58-3.11)	4.02 (2.16-7.07)	390
20-59 years	99-00	*	.490 (<LOD-.670)	.740 (.600-.930)	1.30 (.990-1.80)	2.00 (1.50-2.60)	814
	01-02	.449 (.340-.592)	.540 (.380-.880)	1.45 (1.19-1.79)	2.46 (2.11-3.17)	3.83 (2.96-5.34)	1122
	03-04	*	< LOD	.800 (.650-.960)	1.76 (1.37-2.32)	2.65 (2.31-3.89)	919
	05-06	*	< LOD	.600 (<LOD-.680)	1.42 (1.12-1.77)	2.45 (1.82-3.31)	1092
	07-08	*	< LOD	.930 (.780-1.22)	2.16 (1.68-2.76)	4.20 (2.76-5.92)	1179
60 years and older	01-02	.662 (.571-.768)	1.00 (.740-1.14)	1.77 (1.50-2.10)	2.81 (2.44-3.46)	4.45 (2.93-8.31)	497
	03-04	*	.310 (<LOD-.520)	.890 (.740-1.05)	1.99 (1.52-2.67)	3.72 (2.19-6.81)	517
	05-06	*	< LOD	.660 (<LOD-.840)	1.62 (1.25-2.31)	3.20 (2.04-4.18)	475
	07-08	*	< LOD	.980 (.860-1.16)	2.04 (1.60-3.01)	4.25 (2.76-5.40)	636
Gender							
Males	99-00	*	.510 (<LOD-.700)	.790 (.680-1.10)	1.50 (1.20-2.20)	2.70 (1.90-4.10)	952
	01-02	.495 (.393-.623)	.660 (.470-.900)	1.53 (1.37-1.82)	2.68 (2.28-3.34)	3.96 (3.07-5.85)	1419
	03-04	*	.250 (<LOD-.380)	.890 (.780-1.04)	2.20 (1.65-2.41)	3.15 (2.42-4.29)	1174
	05-06	*	< LOD	.590 (<LOD-.720)	1.48 (1.15-1.97)	2.61 (1.89-3.58)	1246
	07-08	*	< LOD	1.03 (.890-1.25)	2.49 (1.82-3.29)	4.79 (3.44-5.92)	1293
Females	99-00	*	< LOD	.720 (.570-1.00)	1.30 (.910-1.60)	1.70 (1.30-3.20)	997
	01-02	.474 (.360-.624)	.620 (.410-.960)	1.51 (1.22-1.86)	2.49 (2.15-3.04)	3.94 (2.82-5.27)	1597
	03-04	*	< LOD	.770 (.600-.930)	1.59 (1.24-2.08)	2.67 (2.07-3.81)	1248
	05-06	*	< LOD	.630 (.580-.710)	1.43 (1.14-1.77)	2.42 (1.78-3.52)	1388
	07-08	*	< LOD	.960 (.840-1.14)	2.00 (1.60-2.50)	4.01 (2.41-6.59)	1297

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.09, 0.1, 0.2, 0.56, and 0.56 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethylthiophosphate (DETP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	.570 (<LOD-.780)	.840 (.740-.980)	1.50 (1.20-1.90)	2.20 (2.00-2.90)	672
	01-02	.549 (.403-.749)	.710 (.470-.960)	1.40 (1.04-1.97)	2.63 (2.06-3.14)	3.98 (2.75-5.21)	763
	03-04	*	.250 (<LOD-.380)	.970 (.720-1.52)	2.35 (1.69-3.83)	4.19 (2.38-8.80)	584
	05-06	*	< LOD	.700 (<LOD-.900)	1.42 (.980-2.48)	2.72 (1.42-4.48)	695
	07-08	*	< LOD	.970 (.830-1.15)	1.98 (1.60-2.65)	3.46 (2.78-4.46)	500
Non-Hispanic blacks	99-00	.343 (.201-.584)	.570 (<LOD-.750)	.820 (.690-1.20)	1.80 (1.30-3.20)	3.60 (2.00-4.80)	509
	01-02	.755 (.589-.966)	1.17 (.740-1.46)	1.85 (1.73-1.99)	3.54 (2.95-3.87)	5.27 (3.89-6.74)	755
	03-04	.466 (.384-.564)	.430 (.210-.710)	1.08 (.930-1.29)	2.32 (1.74-2.70)	3.25 (2.55-4.83)	605
	05-06	*	< LOD	.850 (.640-1.18)	1.74 (1.43-2.27)	2.96 (2.09-4.20)	722
	07-08	*	< LOD	1.23 (.890-1.40)	2.85 (2.05-4.12)	4.81 (4.00-6.59)	574
Non-Hispanic whites	99-00	*	.160 (<LOD-.700)	.740 (.580-1.10)	1.30 (.960-1.90)	1.90 (1.50-2.80)	595
	01-02	.464 (.344-.626)	.600 (.410-.950)	1.53 (1.27-1.83)	2.46 (2.15-3.17)	3.75 (2.82-5.54)	1252
	03-04	*	< LOD	.750 (.610-.890)	1.75 (1.33-2.20)	2.68 (2.20-3.78)	1044
	05-06	*	< LOD	.600 (<LOD-.690)	1.42 (1.04-1.99)	2.56 (1.82-3.70)	1007
	07-08	*	< LOD	.910 (.820-1.04)	1.80 (1.56-2.46)	3.52 (2.20-5.61)	1087

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.09, 0.1, 0.2, 0.56, and 0.56 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethylthiophosphate (DETP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	.250 (<LOD-.480)	.710 (.460-1.07)	1.72 (1.17-2.32)	2.64 (2.08-3.06)	1949
	01-02	.496 (.388-.636)	.569 (.403-.789)	1.42 (1.16-1.79)	3.11 (2.58-3.82)	4.88 (3.78-6.65)	3015
	03-04	*	< LOD	.740 (.667-.839)	1.59 (1.32-2.05)	2.87 (2.22-3.77)	2420
	05-06	*	< LOD	.956 (<LOD-1.09)	1.88 (1.67-2.21)	2.89 (2.35-3.37)	2633
	07-08	*	< LOD	1.24 (1.15-1.37)	2.46 (1.98-3.11)	3.92 (3.05-5.12)	2588
Age group							
6-11 years	99-00	*	.470 (<LOD-.870)	1.08 (.800-1.32)	1.75 (1.44-2.36)	2.45 (2.04-5.32)	471
	01-02	.591 (.471-.742)	.636 (.397-1.05)	1.63 (1.31-1.94)	3.22 (2.72-4.16)	5.70 (3.84-6.80)	575
	03-04	*	< LOD	.869 (.592-1.09)	1.57 (1.08-2.67)	2.67 (1.57-4.05)	296
	05-06	*	< LOD	1.23 (<LOD-1.33)	1.95 (1.60-2.86)	3.97 (1.95-5.88)	350
	07-08	*	< LOD	1.89 (1.32-2.09)	3.74 (2.46-5.55)	6.54 (3.49-16.0)	385
12-19 years	99-00	*	.180 (<LOD-.400)	.510 (.320-.820)	1.07 (.720-1.61)	1.97 (1.07-3.92)	664
	01-02	.393 (.300-.515)	.534 (.313-.735)	1.23 (.978-1.53)	2.19 (1.61-3.07)	3.14 (2.25-3.97)	821
	03-04	*	.300 (<LOD-.354)	.645 (.559-.734)	1.49 (1.16-1.60)	1.97 (1.57-2.43)	689
	05-06	*	< LOD	.710 (<LOD-.889)	1.35 (1.00-2.03)	2.21 (1.35-3.33)	717
	07-08	*	< LOD	.902 (.692-1.15)	1.78 (1.15-3.32)	3.32 (1.68-5.48)	388
20-59 years	99-00	*	.250 (<LOD-.460)	.680 (.440-1.08)	1.79 (1.08-2.39)	2.75 (2.02-3.22)	814
	01-02	.447 (.335-.597)	.494 (.316-.742)	1.32 (.986-1.71)	2.87 (2.08-3.95)	4.69 (3.20-7.81)	1122
	03-04	*	< LOD	.700 (.550-.875)	1.47 (1.11-2.23)	2.82 (2.02-3.81)	918
	05-06	*	< LOD	.913 (<LOD-1.13)	1.84 (1.65-2.11)	2.86 (2.25-3.33)	1091
	07-08	*	< LOD	1.24 (1.09-1.40)	2.27 (1.79-3.07)	3.51 (2.59-4.78)	1179
60+ years	01-02	.821 (.708-.952)	.971 (.849-1.10)	2.25 (1.81-2.66)	3.97 (3.33-5.06)	5.51 (4.79-8.07)	497
	03-04	*	.491 (<LOD-.560)	.933 (.814-1.17)	2.14 (1.46-3.00)	3.92 (2.28-8.15)	517
	05-06	*	< LOD	1.11 (<LOD-1.38)	2.50 (1.67-3.11)	3.80 (2.86-8.71)	475
	07-08	*	< LOD	1.32 (1.20-1.58)	2.91 (2.25-3.38)	4.49 (3.05-9.80)	636
Gender							
Males	99-00	*	.270 (<LOD-.470)	.670 (.520-.840)	1.34 (1.08-2.17)	2.67 (1.67-3.23)	952
	01-02	.410 (.319-.526)	.517 (.342-.721)	1.22 (1.02-1.41)	2.46 (1.71-3.18)	3.67 (2.70-4.71)	1419
	03-04	*	.276 (<LOD-.326)	.667 (.560-.804)	1.49 (1.15-1.91)	2.87 (1.74-3.72)	1173
	05-06	*	< LOD	.773 (<LOD-.907)	1.42 (1.21-2.00)	2.35 (1.90-3.27)	1246
	07-08	*	< LOD	1.02 (.943-1.09)	1.94 (1.60-2.39)	3.22 (2.50-4.42)	1292
Females	99-00	*	< LOD	.790 (.380-1.50)	1.89 (1.07-2.52)	2.52 (1.89-3.75)	997
	01-02	.597 (.457-.779)	.647 (.423-.961)	1.77 (1.34-2.50)	3.82 (2.93-5.14)	6.57 (4.34-9.28)	1596
	03-04	*	< LOD	.821 (.700-.946)	1.76 (1.34-2.29)	3.02 (2.25-4.33)	1247
	05-06	*	< LOD	1.17 (1.03-1.38)	2.21 (1.83-2.50)	3.18 (2.50-4.47)	1387
	07-08	*	< LOD	1.47 (1.28-1.72)	2.86 (2.29-3.51)	4.48 (3.30-5.73)	1296

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethylthiophosphate (DETP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	.330 (<LOD-.790)	.830 (.550-1.20)	1.69 (1.20-2.43)	2.71 (1.75-3.78)	672
	01-02	.520 (.390-.693)	.570 (.416-.824)	1.30 (1.05-1.74)	2.58 (1.82-3.72)	3.82 (2.77-6.43)	763
	03-04	*	.329 (<LOD-.414)	.824 (.555-1.22)	2.03 (1.26-3.63)	3.74 (2.47-6.18)	584
	05-06	*	< LOD	.907 (<LOD-1.10)	1.74 (1.35-2.07)	2.86 (2.02-4.00)	695
	07-08	*	< LOD	1.20 (.989-1.32)	2.20 (1.83-2.86)	3.32 (2.59-5.11)	499
Non-Hispanic blacks	99-00	.234 (.136-.403)	.310 (<LOD-.580)	.720 (.510-.850)	1.39 (1.03-2.10)	2.91 (1.49-4.24)	509
	01-02	.556 (.457-.676)	.726 (.565-.938)	1.50 (1.34-1.70)	2.80 (2.35-3.30)	4.00 (3.29-4.99)	754
	03-04	.321 (.270-.381)	.313 (.231-.407)	.699 (.561-.831)	1.45 (1.09-1.75)	2.64 (1.71-3.64)	604
	05-06	*	< LOD	.720 (.606-.824)	1.48 (1.24-1.73)	2.45 (1.76-3.19)	721
	07-08	*	< LOD	1.07 (.880-1.21)	2.40 (1.85-2.89)	4.12 (3.19-5.67)	573
Non-Hispanic whites	99-00	*	.230 (<LOD-.550)	.710 (.390-1.22)	1.88 (1.05-2.58)	2.64 (2.08-3.07)	595
	01-02	.506 (.373-.687)	.579 (.368-.850)	1.51 (1.18-2.00)	3.38 (2.64-4.35)	5.83 (3.95-7.81)	1252
	03-04	*	< LOD	.735 (.636-.848)	1.52 (1.20-2.07)	2.69 (2.07-3.97)	1043
	05-06	*	< LOD	1.00 (<LOD-1.21)	2.00 (1.73-2.27)	3.08 (2.50-3.74)	1007
	07-08	*	< LOD	1.24 (1.07-1.39)	2.28 (1.71-3.30)	3.49 (2.56-4.60)	1087

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethylthiophosphate (DMTP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	1.82 (1.36-2.44)	2.70 (1.40-4.10)	11.0 (8.40-16.0)	38.0 (25.0-41.0)	48.0 (38.0-62.0)	1948
	01-02	*	.580 (<LOD-1.45)	4.23 (3.10-5.88)	17.5 (13.0-22.7)	32.9 (27.2-45.3)	3016
	03-04	2.25 (1.98-2.56)	2.03 (1.73-2.37)	6.36 (5.11-7.34)	19.6 (16.2-21.5)	35.3 (27.4-45.3)	2494
	05-06	1.96 (1.74-2.20)	1.71 (1.43-1.94)	5.34 (4.49-6.08)	14.2 (10.5-20.8)	28.2 (22.1-36.0)	2631
	07-08	2.28 (1.97-2.64)	2.10 (1.77-2.44)	5.90 (4.94-7.01)	18.3 (13.5-23.0)	36.8 (27.8-45.7)	2591
Age group							
6-11 years	99-00	2.72 (1.93-3.85)	4.20 (2.50-7.20)	20.0 (13.0-29.0)	40.0 (38.0-52.0)	62.0 (40.0-92.0)	471
	01-02	*	1.46 (.600-2.69)	8.33 (5.75-14.0)	28.4 (19.7-41.4)	45.7 (28.5-74.5)	575
	03-04	2.79 (2.25-3.45)	2.67 (1.81-3.88)	6.95 (5.64-8.58)	19.4 (10.5-27.4)	30.9 (19.4-76.5)	310
	05-06	2.70 (2.25-3.26)	2.38 (1.76-3.33)	8.21 (4.96-11.8)	21.4 (12.7-40.7)	40.7 (18.3-90.4)	349
	07-08	3.43 (2.88-4.09)	3.50 (2.69-4.10)	9.90 (7.14-13.9)	32.0 (19.1-45.7)	52.5 (36.7-67.4)	385
12-19 years	99-00	2.53 (1.64-3.92)	3.70 (1.70-6.80)	16.0 (11.0-31.0)	38.0 (33.0-58.0)	69.0 (38.0-260)	664
	01-02	*	1.04 (<LOD-2.12)	4.83 (3.35-6.48)	20.8 (12.2-27.9)	34.9 (23.6-54.7)	822
	03-04	2.21 (1.81-2.70)	1.83 (1.46-2.16)	5.91 (4.04-8.78)	18.7 (12.2-33.9)	47.1 (22.2-80.8)	717
	05-06	2.51 (2.00-3.16)	2.53 (1.90-3.02)	6.44 (4.91-8.77)	21.7 (13.1-28.2)	38.5 (25.3-79.2)	717
	07-08	2.10 (1.68-2.61)	1.82 (1.53-2.31)	5.18 (3.55-7.48)	13.6 (8.16-27.9)	31.5 (12.4-54.5)	391
20-59 years	99-00	1.59 (1.17-2.16)	2.30 (.830-3.80)	9.10 (7.10-13.0)	38.0 (19.0-39.0)	39.0 (38.0-58.0)	813
	01-02	*	< LOD	3.32 (2.29-4.96)	13.6 (9.50-20.0)	30.0 (20.5-45.3)	1121
	03-04	1.98 (1.71-2.30)	1.78 (1.48-2.18)	5.11 (4.31-6.53)	16.7 (12.1-20.8)	28.5 (24.1-40.0)	938
	05-06	1.72 (1.50-1.97)	1.37 (1.20-1.73)	4.25 (3.68-5.22)	12.4 (8.37-19.6)	26.6 (16.4-37.9)	1092
	07-08	2.03 (1.68-2.46)	1.90 (1.46-2.39)	5.18 (4.24-6.09)	13.3 (10.4-19.2)	30.6 (21.4-44.6)	1179
60 years and older	01-02	*	.860 (<LOD-2.16)	5.85 (3.33-9.56)	19.6 (13.0-27.9)	32.9 (22.7-51.0)	498
	03-04	3.18 (2.47-4.08)	2.64 (2.16-3.81)	9.63 (6.32-15.5)	25.0 (15.5-45.3)	45.3 (20.9-119)	529
	05-06	2.18 (1.81-2.63)	2.03 (1.46-2.68)	6.38 (4.62-8.23)	14.0 (9.72-20.9)	25.2 (18.2-37.6)	473
	07-08	2.91 (2.37-3.58)	2.57 (1.99-3.18)	9.19 (5.54-13.0)	26.2 (19.4-32.5)	42.7 (30.5-54.1)	636
Gender							
Males	99-00	2.10 (1.48-2.98)	3.50 (2.20-4.80)	14.0 (8.00-24.0)	38.0 (21.0-49.0)	42.0 (38.0-53.0)	952
	01-02	*	.690 (<LOD-1.52)	4.44 (3.32-6.48)	18.3 (12.7-25.6)	31.1 (25.0-43.3)	1420
	03-04	2.30 (1.95-2.70)	2.15 (1.70-2.53)	6.42 (4.90-7.75)	18.8 (13.3-22.5)	30.0 (23.3-45.3)	1221
	05-06	1.88 (1.64-2.14)	1.61 (1.28-1.94)	5.38 (3.96-6.38)	13.1 (9.59-20.9)	28.7 (19.5-41.7)	1243
	07-08	2.33 (2.00-2.71)	2.15 (1.85-2.50)	5.98 (4.78-7.26)	16.7 (11.8-24.7)	31.5 (21.2-48.9)	1293
Females	99-00	1.59 (1.23-2.06)	2.00 (.690-3.60)	9.70 (7.30-14.0)	38.0 (26.0-39.0)	52.0 (38.0-110)	996
	01-02	*	.430 (<LOD-1.45)	3.81 (2.68-5.85)	16.0 (10.9-22.7)	35.2 (25.0-46.1)	1596
	03-04	2.21 (1.92-2.55)	1.95 (1.68-2.29)	6.27 (4.93-7.39)	20.4 (16.4-24.0)	36.7 (27.7-47.8)	1273
	05-06	2.04 (1.79-2.33)	1.75 (1.46-2.13)	5.23 (4.49-6.10)	15.6 (10.9-20.8)	28.1 (20.8-37.9)	1388
	07-08	2.24 (1.88-2.66)	1.99 (1.57-2.52)	5.85 (4.92-7.31)	18.7 (13.4-26.0)	37.7 (27.0-50.1)	1298

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.18, 0.4, 0.5, 0.55, and 0.55 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethylthiophosphate (DMTP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	1.79 (1.05-3.05)	2.00 (.530-4.40)	11.0 (6.70-17.0)	39.0 (32.0-62.0)	140 (46.0-230)	671
	01-02	*	< LOD	3.93 (3.05-5.06)	15.1 (11.1-19.1)	35.6 (22.7-46.0)	763
	03-04	2.41 (1.88-3.09)	2.54 (1.41-3.98)	6.43 (4.90-8.02)	17.2 (12.0-22.3)	28.9 (19.2-39.8)	610
	05-06	2.37 (2.07-2.71)	2.26 (1.95-2.68)	5.65 (4.86-6.44)	16.4 (12.9-21.4)	28.3 (23.3-41.4)	693
	07-08	2.10 (1.83-2.42)	1.98 (1.63-2.31)	5.18 (4.17-6.61)	13.6 (11.5-20.0)	31.0 (20.0-43.5)	501
Non-Hispanic blacks	99-00	2.13 (1.57-2.88)	3.60 (2.10-4.70)	12.0 (8.80-18.0)	38.0 (20.0-41.0)	41.0 (37.0-110)	509
	01-02	1.54 (1.15-2.07)	1.18 (.580-1.97)	5.21 (3.11-8.25)	20.0 (13.4-27.2)	39.1 (25.9-62.8)	755
	03-04	2.13 (1.83-2.48)	1.85 (1.65-2.20)	5.44 (3.87-8.28)	20.4 (15.8-24.9)	31.1 (24.9-51.9)	631
	05-06	1.94 (1.63-2.31)	1.63 (1.23-2.02)	5.21 (3.93-7.18)	16.8 (13.3-21.6)	28.4 (21.5-40.6)	723
	07-08	2.12 (1.89-2.38)	1.97 (1.64-2.26)	5.63 (4.15-6.94)	20.8 (12.9-23.9)	33.5 (26.3-47.6)	573
Non-Hispanic whites	99-00	1.77 (1.23-2.53)	2.70 (.830-4.40)	11.0 (7.50-17.0)	38.0 (17.0-53.0)	48.0 (38.0-69.0)	595
	01-02	*	.540 (<LOD-1.71)	4.21 (2.77-6.48)	18.8 (10.9-25.4)	33.1 (26.6-46.5)	1252
	03-04	2.31 (2.00-2.66)	2.07 (1.73-2.41)	6.51 (5.12-7.85)	19.8 (15.4-23.3)	36.7 (27.4-47.8)	1061
	05-06	1.94 (1.67-2.26)	1.76 (1.40-1.99)	5.33 (4.32-6.19)	13.6 (9.80-22.1)	28.9 (20.2-42.8)	1005
	07-08	2.22 (1.83-2.69)	1.96 (1.57-2.48)	5.84 (4.70-7.68)	17.1 (12.4-24.3)	37.3 (25.9-48.7)	1088

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.18, 0.4, 0.5, 0.55, and 0.55 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethylthiophosphate (DMTP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	1.64 (1.22-2.20)	2.12 (1.22-3.35)	9.57 (6.59-15.8)	32.0 (23.9-41.1)	51.0 (39.0-70.1)	1948
	01-02	*	.933 (<LOD-1.40)	4.08 (2.65-6.06)	14.6 (11.5-20.5)	28.9 (24.7-39.4)	3015
	03-04	2.21 (1.92-2.55)	1.98 (1.69-2.38)	5.96 (5.07-7.16)	19.1 (14.8-22.4)	32.6 (25.8-41.9)	2491
	05-06	1.99 (1.76-2.26)	1.75 (1.56-1.90)	5.00 (4.16-6.24)	16.8 (11.8-22.1)	30.0 (24.1-38.3)	2630
	07-08	2.34 (2.05-2.66)	2.00 (1.77-2.31)	5.91 (4.88-7.09)	17.5 (13.9-23.2)	33.7 (25.7-45.9)	2589
Age group							
6-11 years	99-00	2.95 (2.25-3.86)	5.32 (3.75-6.33)	19.1 (12.2-28.0)	47.0 (32.1-60.3)	66.1 (50.9-95.0)	471
	01-02	*	2.16 (1.32-3.12)	10.6 (7.84-13.6)	28.7 (18.8-45.0)	48.1 (33.4-71.1)	575
	03-04	3.40 (2.70-4.28)	3.41 (2.40-4.17)	7.91 (6.43-12.2)	25.2 (16.7-34.2)	36.1 (25.4-67.7)	310
	05-06	3.35 (2.67-4.21)	2.78 (2.03-3.66)	9.26 (5.67-15.1)	33.4 (16.8-54.3)	71.5 (28.7-161)	349
	07-08	4.35 (3.78-5.01)	4.26 (3.36-4.75)	14.1 (9.35-16.6)	32.8 (23.8-45.4)	51.3 (41.1-77.9)	385
12-19 years	99-00	1.71 (1.07-2.75)	2.14 (.890-4.83)	13.5 (6.46-22.6)	36.0 (25.6-51.4)	61.5 (41.7-109)	664
	01-02	*	.933 (<LOD-1.57)	3.56 (2.38-5.57)	12.2 (8.96-16.0)	22.5 (13.2-34.7)	821
	03-04	1.66 (1.37-2.03)	1.52 (1.19-1.82)	4.38 (3.29-5.66)	13.3 (9.94-20.5)	26.5 (15.5-36.0)	715
	05-06	1.89 (1.54-2.33)	1.81 (1.38-2.33)	4.55 (3.47-6.17)	14.3 (8.75-27.5)	32.9 (14.3-51.4)	717
	07-08	1.62 (1.27-2.06)	1.46 (1.19-1.74)	4.14 (3.12-5.52)	11.1 (5.52-20.5)	20.5 (9.61-32.2)	389
20-59 years	99-00	1.47 (1.07-2.02)	1.90 (.870-3.11)	8.09 (5.19-14.6)	27.0 (19.8-37.6)	47.5 (34.2-70.1)	813
	01-02	*	< LOD	3.16 (1.99-4.62)	11.9 (7.79-17.2)	25.2 (15.9-37.0)	1121
	03-04	1.88 (1.61-2.19)	1.67 (1.45-1.94)	4.88 (4.20-5.68)	13.9 (10.3-19.7)	30.3 (19.7-38.2)	937
	05-06	1.72 (1.52-1.96)	1.46 (1.28-1.65)	4.18 (3.14-4.99)	13.2 (9.30-21.9)	25.4 (21.4-30.3)	1091
	07-08	2.02 (1.73-2.36)	1.79 (1.52-2.05)	4.53 (3.90-5.90)	13.2 (9.47-17.2)	26.7 (17.1-41.8)	1179
60+ years	01-02	*	1.39 (<LOD-2.40)	6.38 (3.68-9.97)	24.0 (14.9-35.2)	45.5 (26.5-69.7)	498
	03-04	3.86 (2.99-4.97)	3.18 (2.46-4.84)	11.2 (7.10-18.6)	29.2 (18.6-47.9)	48.2 (24.9-113)	529
	05-06	2.61 (2.12-3.20)	2.40 (1.75-2.75)	7.66 (5.71-11.3)	19.2 (12.1-31.1)	35.3 (20.5-56.4)	473
	07-08	3.56 (2.79-4.55)	3.05 (2.31-4.13)	10.6 (6.86-14.8)	31.8 (17.7-50.2)	54.2 (34.1-87.5)	636
	Gender						
Males	99-00	1.61 (1.11-2.34)	2.39 (1.27-3.51)	9.27 (6.00-16.9)	28.9 (19.0-40.4)	41.1 (34.9-52.9)	952
	01-02	*	.778 (<LOD-1.14)	3.82 (2.65-4.98)	13.6 (10.0-17.8)	25.2 (17.2-37.6)	1420
	03-04	1.88 (1.58-2.23)	1.76 (1.51-2.12)	4.87 (3.87-6.31)	14.4 (9.18-20.4)	25.7 (17.7-36.9)	1220
	05-06	1.59 (1.37-1.86)	1.33 (1.14-1.57)	4.06 (2.85-5.27)	13.1 (8.56-20.5)	26.0 (17.2-34.2)	1243
	07-08	1.93 (1.68-2.21)	1.62 (1.42-1.94)	4.36 (3.77-5.50)	14.2 (10.4-17.7)	25.9 (18.2-39.7)	1292
Females	99-00	1.66 (1.26-2.18)	2.01 (.870-3.33)	10.0 (6.67-16.2)	34.9 (26.2-47.1)	70.1 (39.0-118)	996
	01-02	*	1.08 (<LOD-1.65)	4.44 (2.55-7.02)	17.4 (11.9-24.8)	38.0 (27.4-51.6)	1595
	03-04	2.60 (2.23-3.02)	2.24 (1.83-2.63)	7.11 (5.54-10.2)	22.8 (18.9-28.7)	41.6 (29.2-58.9)	1271
	05-06	2.47 (2.16-2.83)	2.19 (1.83-2.44)	6.07 (4.74-8.34)	20.7 (13.8-25.6)	36.7 (26.1-46.2)	1387
	07-08	2.81 (2.40-3.29)	2.50 (2.07-2.93)	7.40 (5.83-8.83)	21.9 (16.7-27.4)	41.2 (27.4-60.8)	1297

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethylthiophosphate (DMTP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	1.60 (.899-2.86)	1.83 (.680-4.23)	10.4 (5.95-16.9)	37.0 (23.1-63.1)	112 (40.5-190)	671
	01-02	*	< LOD	3.55 (2.65-4.68)	13.8 (9.88-22.9)	30.4 (22.9-47.2)	763
	03-04	2.20 (1.65-2.94)	2.24 (1.70-3.00)	5.95 (4.14-8.68)	15.7 (10.7-20.1)	24.5 (18.9-36.9)	609
	05-06	2.24 (1.96-2.55)	2.14 (1.89-2.40)	5.19 (4.33-6.18)	14.7 (10.9-20.0)	28.4 (21.5-36.3)	693
	07-08	2.14 (1.92-2.38)	1.88 (1.61-2.22)	4.84 (4.08-6.48)	13.2 (10.0-18.6)	26.2 (18.6-34.1)	500
Non-Hispanic blacks	99-00	1.45 (1.03-2.06)	1.75 (1.17-3.06)	8.48 (4.36-13.4)	25.5 (15.4-39.3)	54.4 (25.5-97.6)	509
	01-02	1.13 (.889-1.44)	1.00 (.693-1.23)	3.62 (2.29-5.41)	12.4 (9.17-17.5)	23.3 (17.4-43.1)	754
	03-04	1.46 (1.28-1.67)	1.22 (1.06-1.37)	3.90 (2.77-5.34)	12.7 (8.09-18.7)	26.0 (14.6-38.7)	630
	05-06	1.40 (1.14-1.71)	1.12 (.938-1.37)	3.58 (2.63-5.06)	12.0 (8.44-19.3)	25.6 (17.4-31.2)	722
	07-08	1.61 (1.45-1.80)	1.38 (1.22-1.61)	4.26 (3.41-5.20)	14.7 (10.7-18.8)	27.5 (20.9-37.4)	572
Non-Hispanic whites	99-00	1.68 (1.16-2.43)	2.22 (.870-3.51)	9.40 (5.58-17.0)	33.3 (20.6-49.4)	52.9 (39.0-71.1)	595
	01-02	*	.966 (<LOD-1.61)	4.22 (2.41-7.00)	15.9 (11.6-24.1)	33.1 (24.8-45.4)	1252
	03-04	2.40 (2.05-2.81)	2.12 (1.75-2.53)	6.27 (5.10-8.10)	20.0 (15.3-25.7)	36.2 (25.7-52.7)	1060
	05-06	2.09 (1.80-2.42)	1.78 (1.56-2.06)	5.14 (4.02-6.91)	16.8 (11.3-23.4)	30.0 (23.4-38.3)	1005
	07-08	2.40 (1.99-2.90)	2.04 (1.67-2.57)	6.00 (4.64-7.49)	16.9 (13.0-25.3)	37.6 (23.1-54.2)	1088

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethyldithiophosphate (DEDTP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	.090 (<LOD-.140)	.210 (.140-.290)	.470 (.380-.640)	.870 (.640-1.10)	1949
	01-02	*	< LOD	< LOD	.610 (.410-.780)	.890 (.710-1.32)	3013
	03-04	*	< LOD	< LOD	< LOD	.290 (.160-.510)	2494
	05-06	*	< LOD	< LOD	< LOD	< LOD	2420
	07-08	*	< LOD	< LOD	< LOD	< LOD	2585
Age group							
6-11 years	99-00	*	.090 (<LOD-.160)	.190 (.130-.280)	.430 (.300-.650)	.850 (.470-1.00)	471
	01-02	*	< LOD	< LOD	.630 (.380-.870)	.940 (.690-1.42)	576
	03-04	*	< LOD	< LOD	< LOD	.540 (<LOD-.650)	310
	05-06	*	< LOD	< LOD	< LOD	< LOD	317
	07-08	*	< LOD	< LOD	< LOD	< LOD	383
12-19 years	99-00	*	.080 (<LOD-.180)	.260 (.120-.350)	.640 (.420-.840)	.930 (.720-1.30)	664
	01-02	*	< LOD	< LOD	.560 (.330-.730)	.820 (.610-.990)	822
	03-04	*	< LOD	< LOD	.150 (<LOD-.350)	.450 (.370-.570)	717
	05-06	*	< LOD	< LOD	< LOD	< LOD	660
	07-08	*	< LOD	< LOD	< LOD	< LOD	390
20-59 years	99-00	*	.090 (<LOD-.130)	.210 (.130-.290)	.450 (.360-.640)	.900 (.610-1.10)	814
	01-02	*	< LOD	< LOD	.620 (.430-.760)	.830 (.700-1.32)	1118
	03-04	*	< LOD	< LOD	< LOD	.220 (<LOD-.580)	938
	05-06	*	< LOD	< LOD	< LOD	< LOD	1006
	07-08	*	< LOD	< LOD	< LOD	< LOD	1176
60 years and older	01-02	*	< LOD	< LOD	.630 (.360-1.08)	1.11 (.610-1.41)	497
	03-04	*	< LOD	< LOD	< LOD	.150 (<LOD-.550)	529
	05-06	*	< LOD	< LOD	< LOD	< LOD	437
	07-08	*	< LOD	< LOD	< LOD	< LOD	636
Gender							
Males	99-00	*	.090 (<LOD-.150)	.220 (.140-.310)	.490 (.380-.680)	.870 (.680-1.10)	952
	01-02	*	< LOD	< LOD	.600 (.380-.740)	.780 (.700-1.03)	1420
	03-04	*	< LOD	< LOD	< LOD	.390 (.160-.540)	1221
	05-06	*	< LOD	< LOD	< LOD	< LOD	1142
	07-08	*	< LOD	< LOD	< LOD	< LOD	1290
Females	99-00	*	.090 (<LOD-.130)	.190 (.110-.310)	.460 (.320-.840)	.870 (.440-1.40)	997
	01-02	*	< LOD	< LOD	.660 (.460-.880)	1.03 (.700-1.47)	1593
	03-04	*	< LOD	< LOD	< LOD	.230 (<LOD-.600)	1273
	05-06	*	< LOD	< LOD	< LOD	< LOD	1278
	07-08	*	< LOD	< LOD	< LOD	< LOD	1295

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.05, 0.1, 0.1, 0.39, and 0.39 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethyldithiophosphate (DEDTP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.130 (.099-.171)	.100 (.050-.200)	.310 (.230-.390)	.650 (.530-.860)	1.10 (.860-1.60)	672
	01-02	*	< LOD	.200 (<LOD-.520)	.720 (.410-1.12)	1.12 (.700-1.58)	763
	03-04	*	< LOD	< LOD	.110 (<LOD-.680)	.680 (.180-1.15)	610
	05-06	*	< LOD	< LOD	< LOD	< LOD	665
	07-08	*	< LOD	< LOD	< LOD	< LOD	495
Non-Hispanic blacks	99-00	.117 (.084-.162)	.090 (<LOD-.230)	.270 (.140-.410)	.560 (.400-.830)	.870 (.650-1.20)	509
	01-02	*	< LOD	< LOD	.620 (.440-.800)	.810 (.670-.930)	754
	03-04	*	< LOD	< LOD	< LOD	.250 (.110-.540)	631
	05-06	*	< LOD	< LOD	< LOD	< LOD	660
	07-08	*	< LOD	< LOD	< LOD	< LOD	574
Non-Hispanic whites	99-00	*	.080 (<LOD-.160)	.190 (.120-.290)	.450 (.310-.720)	.870 (.510-1.30)	595
	01-02	*	< LOD	< LOD	.610 (.350-.800)	.910 (.660-1.41)	1252
	03-04	*	< LOD	< LOD	< LOD	.240 (.120-.540)	1061
	05-06	*	< LOD	< LOD	< LOD	< LOD	917
	07-08	*	< LOD	< LOD	< LOD	< LOD	1087

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.05, 0.1, 0.1, 0.39, and 0.39 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethyldithiophosphate (DEDTP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	.070 (<LOD-.110)	.200 (.140-.290)	.550 (.390-.700)	.860 (.670-1.14)	1949
	01-02	*	< LOD	< LOD	.604 (.389-.783)	1.08 (.768-1.52)	3012
	03-04	*	< LOD	< LOD	< LOD	.412 (.350-.507)	2491
	05-06	*	< LOD	< LOD	< LOD	< LOD	2419
	07-08	*	< LOD	< LOD	< LOD	< LOD	2583
Age group							
6-11 years	99-00	*	.100 (<LOD-.140)	.190 (.150-.270)	.570 (.410-.760)	1.03 (.570-1.58)	471
	01-02	*	< LOD	< LOD	.778 (.606-1.12)	1.36 (1.02-1.86)	576
	03-04	*	< LOD	< LOD	< LOD	.475 (<LOD-.967)	310
	05-06	*	< LOD	< LOD	< LOD	< LOD	317
	07-08	*	< LOD	< LOD	< LOD	< LOD	383
12-19 years	99-00	*	.050 (<LOD-.080)	.170 (.100-.220)	.440 (.230-.730)	.730 (.380-1.09)	664
	01-02	*	< LOD	< LOD	.359 (.250-.537)	.667 (.375-.987)	821
	03-04	*	< LOD	< LOD	.226 (<LOD-.259)	.333 (.241-.600)	715
	05-06	*	< LOD	< LOD	< LOD	< LOD	660
	07-08	*	< LOD	< LOD	< LOD	< LOD	388
20-59 years	99-00	*	.080 (<LOD-.110)	.210 (.140-.310)	.550 (.360-.730)	.860 (.650-1.20)	814
	01-02	*	< LOD	< LOD	.583 (.379-.740)	1.03 (.699-1.60)	1118
	03-04	*	< LOD	< LOD	< LOD	.400 (<LOD-.537)	937
	05-06	*	< LOD	< LOD	< LOD	< LOD	1005
	07-08	*	< LOD	< LOD	< LOD	< LOD	1176
60+ years	01-02	*	< LOD	< LOD	.833 (.368-1.28)	1.44 (.853-2.61)	497
	03-04	*	< LOD	< LOD	< LOD	.438 (<LOD-.700)	529
	05-06	*	< LOD	< LOD	< LOD	< LOD	437
	07-08	*	< LOD	< LOD	< LOD	< LOD	636
Gender							
Males	99-00	*	.070 (<LOD-.110)	.190 (.140-.230)	.410 (.340-.500)	.720 (.520-.940)	952
	01-02	*	< LOD	< LOD	.389 (.309-.658)	.768 (.637-1.01)	1420
	03-04	*	< LOD	< LOD	< LOD	.339 (.269-.412)	1220
	05-06	*	< LOD	< LOD	< LOD	< LOD	1142
	07-08	*	< LOD	< LOD	< LOD	< LOD	1289
Females	99-00	*	.090 (<LOD-.120)	.220 (.140-.360)	.670 (.410-.870)	.890 (.660-1.62)	997
	01-02	*	< LOD	< LOD	.733 (.527-1.03)	1.27 (.868-2.37)	1592
	03-04	*	< LOD	< LOD	< LOD	.500 (<LOD-.604)	1271
	05-06	*	< LOD	< LOD	< LOD	< LOD	1277
	07-08	*	< LOD	< LOD	< LOD	< LOD	1294

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Diethyldithiophosphate (DEDTP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.116 (.084-.161)	.090 (.060-.170)	.300 (.190-.410)	.810 (.570-.990)	1.19 (.860-2.66)	672
	01-02	*	< LOD	.292 (<LOD-.444)	.845 (.487-1.24)	1.29 (.901-1.67)	763
	03-04	*	< LOD	< LOD	.318 (<LOD-.467)	.579 (.404-.940)	609
	05-06	*	< LOD	< LOD	< LOD	< LOD	665
	07-08	*	< LOD	< LOD	< LOD	< LOD	494
Non-Hispanic blacks	99-00	.080 (.057-.111)	.070 (<LOD-.110)	.170 (.110-.280)	.450 (.300-.580)	.700 (.500-1.02)	509
	01-02	*	< LOD	< LOD	.456 (.362-.593)	.769 (.541-1.07)	753
	03-04	*	< LOD	< LOD	< LOD	.269 (.196-.368)	630
	05-06	*	< LOD	< LOD	< LOD	< LOD	659
	07-08	*	< LOD	< LOD	< LOD	< LOD	573
Non-Hispanic whites	99-00	*	.070 (<LOD-.120)	.200 (.140-.310)	.560 (.380-.730)	.880 (.600-1.38)	595
	01-02	*	< LOD	< LOD	.583 (.368-.868)	1.14 (.682-1.70)	1252
	03-04	*	< LOD	< LOD	< LOD	.389 (.333-.523)	1060
	05-06	*	< LOD	< LOD	< LOD	< LOD	917
	07-08	*	< LOD	< LOD	< LOD	< LOD	1087

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethyldithiophosphate (DMDTP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	2.30 (1.30-3.90)	13.0 (5.00-17.0)	19.0 (17.0-38.0)	1949
	01-02	*	< LOD	.920 (.330-1.30)	2.61 (2.04-3.45)	5.96 (4.41-7.90)	3015
	03-04	*	< LOD	.700 (.570-.920)	2.54 (1.64-3.87)	5.74 (3.63-9.85)	2458
	05-06	*	< LOD	< LOD	1.96 (1.41-2.59)	3.87 (2.77-6.09)	2635
	07-08	*	< LOD	< LOD	2.37 (1.93-2.93)	5.60 (4.62-6.79)	2571
Age group							
6-11 years	99-00	.691 (.425-1.12)	.740 (.080-1.80)	4.30 (2.40-8.60)	17.0 (6.90-37.0)	32.0 (17.0-44.0)	471
	01-02	*	< LOD	1.30 (.750-2.11)	3.53 (2.20-4.50)	7.33 (4.32-9.74)	575
	03-04	*	< LOD	.900 (.620-1.14)	2.94 (1.14-5.48)	5.48 (2.94-8.53)	306
	05-06	*	< LOD	< LOD	1.84 (.540-3.35)	3.69 (2.67-9.95)	350
	07-08	*	< LOD	.950 (<LOD-1.48)	3.80 (2.45-5.70)	6.66 (5.11-9.64)	379
12-19 years	99-00	*	< LOD	2.30 (1.40-4.50)	13.0 (5.40-20.0)	20.0 (17.0-38.0)	664
	01-02	*	< LOD	.830 (.400-1.14)	2.52 (1.85-3.07)	4.63 (3.59-5.83)	821
	03-04	*	< LOD	.580 (.350-.770)	1.46 (1.12-1.99)	2.67 (1.82-4.49)	699
	05-06	*	< LOD	< LOD	2.78 (1.51-3.76)	5.98 (3.03-11.5)	718
	07-08	*	< LOD	< LOD	1.45 (.810-2.03)	2.45 (1.65-3.93)	384
20-59 years	99-00	*	< LOD	2.10 (.840-3.60)	11.0 (4.00-17.0)	17.0 (7.70-50.0)	814
	01-02	*	< LOD	.840 (<LOD-1.31)	2.32 (1.70-3.40)	4.90 (2.90-9.52)	1122
	03-04	*	< LOD	.610 (.350-.800)	2.00 (1.36-3.63)	5.07 (3.62-8.62)	925
	05-06	*	< LOD	< LOD	1.71 (.880-2.36)	3.60 (2.40-5.18)	1092
	07-08	*	< LOD	< LOD	1.84 (1.34-2.32)	4.27 (2.84-6.79)	1172
60 years and older	01-02	*	< LOD	1.02 (.540-1.37)	3.57 (2.58-5.96)	8.16 (5.04-16.1)	497
	03-04	*	< LOD	1.39 (.720-2.34)	4.09 (2.35-13.4)	13.4 (3.36-33.3)	528
	05-06	*	< LOD	< LOD	2.13 (1.67-3.34)	5.12 (2.62-8.17)	475
	07-08	*	< LOD	1.03 (<LOD-1.99)	5.47 (4.31-6.33)	8.89 (6.33-12.2)	636
Gender							
Males	99-00	*	.110 (<LOD-.610)	2.30 (1.20-4.90)	16.0 (5.70-17.0)	19.0 (17.0-38.0)	952
	01-02	*	< LOD	.890 (.370-1.28)	2.59 (1.88-3.41)	5.96 (4.47-7.86)	1420
	03-04	*	< LOD	.680 (.360-1.01)	2.26 (1.14-4.45)	5.30 (3.20-10.2)	1209
	05-06	*	< LOD	< LOD	1.96 (.980-3.04)	4.33 (2.42-7.65)	1246
	07-08	*	< LOD	< LOD	2.38 (1.76-3.70)	5.70 (3.80-10.0)	1278
Females	99-00	*	< LOD	2.20 (1.10-3.90)	11.0 (4.20-17.0)	20.0 (13.0-40.0)	997
	01-02	*	< LOD	.960 (.250-1.37)	2.61 (1.96-3.68)	5.72 (3.79-9.87)	1595
	03-04	*	< LOD	.730 (.590-.970)	2.83 (1.99-3.96)	6.15 (3.91-11.3)	1249
	05-06	*	< LOD	< LOD	1.96 (1.49-2.59)	3.75 (2.90-4.60)	1389
	07-08	*	< LOD	< LOD	2.34 (1.79-3.12)	5.54 (4.06-6.69)	1293

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.08, 0.1, 0.1, 0.51, and 0.51 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethyldithiophosphate (DMDTP)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	*	.250 (<LOD-.870)	1.90 (1.10-3.00)	5.80 (4.10-9.70)	12.0 (5.90-28.0)	672
	01-02	*	< LOD	1.08 (.780-1.38)	2.80 (2.19-3.54)	4.93 (3.73-7.42)	763
	03-04	*	< LOD	.730 (.360-1.21)	2.11 (1.38-3.15)	5.26 (2.31-6.94)	610
	05-06	*	< LOD	< LOD	2.04 (1.62-2.96)	5.12 (2.98-10.7)	695
	07-08	*	< LOD	< LOD	1.63 (1.18-2.12)	3.88 (2.12-6.07)	491
Non-Hispanic blacks	99-00	*	.330 (<LOD-1.20)	3.20 (1.40-7.00)	14.0 (5.70-30.0)	19.0 (17.0-39.0)	509
	01-02	*	< LOD	.640 (<LOD-1.41)	1.96 (1.36-4.15)	4.42 (2.68-8.38)	755
	03-04	*	< LOD	.590 (.310-.730)	1.77 (1.05-3.33)	4.77 (2.75-7.80)	604
	05-06	*	< LOD	< LOD	2.15 (1.66-2.82)	5.45 (2.82-8.39)	723
	07-08	*	< LOD	< LOD	2.04 (1.54-2.65)	4.27 (2.66-7.79)	568
Non-Hispanic whites	99-00	*	< LOD	2.00 (.800-4.00)	13.0 (3.90-20.0)	20.0 (16.0-40.0)	595
	01-02	*	< LOD	1.01 (.180-1.43)	2.74 (1.94-4.18)	6.72 (4.88-8.54)	1252
	03-04	*	< LOD	.740 (.540-1.04)	2.66 (1.55-4.56)	5.89 (3.63-11.6)	1056
	05-06	*	< LOD	< LOD	1.93 (1.23-2.65)	3.60 (2.71-5.51)	1007
	07-08	*	< LOD	< LOD	2.32 (1.75-3.28)	5.60 (4.46-6.88)	1087

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, and 07-08 are 0.08, 0.1, 0.1, 0.51, and 0.51 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethyldithiophosphate (DMDTP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	1.88 (.970-3.86)	10.1 (5.31-18.3)	21.7 (12.8-33.7)	1949
	01-02	*	< LOD	.717 (.389-1.12)	2.80 (2.21-3.96)	6.84 (4.96-8.09)	3014
	03-04	*	< LOD	.572 (.411-.772)	2.62 (1.72-4.12)	6.40 (3.85-9.75)	2455
	05-06	*	< LOD	< LOD	2.18 (1.64-3.00)	4.62 (3.23-7.12)	2634
	07-08	*	< LOD	< LOD	2.63 (2.17-3.08)	6.12 (4.15-8.24)	2569
Age group							
6-11 years	99-00	.748 (.474-1.18)	.790 (.190-1.60)	4.07 (2.31-7.18)	16.2 (8.22-27.0)	30.8 (20.2-38.9)	471
	01-02	*	< LOD	1.36 (.805-2.31)	4.10 (2.67-6.24)	6.98 (4.40-12.8)	575
	03-04	*	< LOD	.962 (.584-1.57)	3.39 (2.29-6.15)	7.12 (4.48-7.55)	306
	05-06	*	< LOD	< LOD	2.57 (1.39-4.14)	4.50 (3.17-9.35)	350
	07-08	*	< LOD	1.50 (<LOD-1.95)	4.35 (3.05-6.59)	8.01 (4.75-12.9)	379
12-19 years	99-00	*	< LOD	1.52 (.620-3.47)	9.48 (4.04-16.8)	21.5 (9.48-42.3)	664
	01-02	*	< LOD	.543 (.306-.773)	2.02 (1.49-2.40)	3.13 (2.51-4.67)	820
	03-04	*	< LOD	.361 (.241-.584)	1.02 (.628-1.57)	2.46 (1.33-3.39)	697
	05-06	*	< LOD	< LOD	2.05 (1.16-3.56)	4.27 (2.30-8.66)	718
	07-08	*	< LOD	< LOD	1.16 (.782-1.85)	1.94 (1.39-3.79)	382
20-59 years	99-00	*	< LOD	1.71 (.850-3.56)	8.50 (4.00-19.1)	20.5 (8.57-40.7)	814
	01-02	*	< LOD	.605 (<LOD-1.05)	2.56 (1.64-4.03)	6.33 (3.96-8.17)	1122
	03-04	*	< LOD	.451 (.340-.583)	2.17 (1.10-3.64)	5.71 (2.47-10.1)	924
	05-06	*	< LOD	< LOD	1.90 (1.44-2.57)	3.69 (2.37-7.07)	1091
	07-08	*	< LOD	< LOD	1.99 (1.80-2.32)	4.15 (2.77-6.77)	1172
60+ years	01-02	*	< LOD	1.10 (.631-1.86)	4.89 (3.36-7.82)	13.4 (6.98-19.8)	497
	03-04	*	< LOD	1.55 (.765-2.48)	6.52 (2.73-13.3)	13.3 (5.44-23.0)	528
	05-06	*	< LOD	< LOD	3.22 (2.25-4.54)	6.53 (3.70-10.4)	475
	07-08	*	< LOD	1.72 (<LOD-2.58)	5.92 (3.29-8.44)	10.3 (7.25-18.4)	636
Gender							
Males	99-00	*	.150 (<LOD-.370)	1.79 (.840-3.97)	11.0 (4.62-17.4)	18.1 (7.51-44.7)	952
	01-02	*	< LOD	.628 (.333-.850)	2.39 (1.75-3.13)	5.37 (4.23-7.29)	1420
	03-04	*	< LOD	.467 (.280-.743)	2.03 (.990-3.56)	5.38 (2.29-8.25)	1208
	05-06	*	< LOD	< LOD	1.80 (1.14-3.17)	4.14 (2.18-7.15)	1246
	07-08	*	< LOD	< LOD	1.89 (1.44-2.71)	4.92 (2.77-8.44)	1277
Females	99-00	*	< LOD	2.06 (.940-4.00)	9.30 (4.96-25.5)	27.0 (9.66-47.5)	997
	01-02	*	< LOD	.831 (.412-1.50)	3.35 (2.50-4.96)	7.81 (5.29-11.9)	1594
	03-04	*	< LOD	.680 (.538-.887)	3.59 (2.41-5.18)	7.12 (4.13-13.1)	1247
	05-06	*	< LOD	< LOD	2.40 (2.00-3.18)	5.25 (3.18-7.60)	1388
	07-08	*	< LOD	< LOD	3.05 (2.44-4.00)	7.18 (4.22-11.1)	1292

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Dimethyldithiophosphate (DMDTP) (creatinine corrected)

Metabolite of Several Organophosphorus Insecticides

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Mexican Americans	99-00	*	.270 (<LOD-.660)	1.35 (.860-2.53)	6.55 (3.83-11.8)	16.7 (6.25-38.8)	672
	01-02	*	< LOD	.833 (.589-1.15)	2.72 (2.05-3.32)	5.16 (3.69-7.83)	763
	03-04	*	< LOD	.651 (.318-1.09)	2.04 (1.10-3.74)	4.50 (2.17-5.88)	609
	05-06	*	< LOD	< LOD	2.40 (1.29-4.00)	5.14 (3.45-10.5)	695
	07-08	*	< LOD	< LOD	1.97 (1.50-2.77)	4.30 (2.77-6.92)	490
Non-Hispanic blacks	99-00	*	.250 (<LOD-.700)	2.40 (.690-5.44)	9.41 (4.81-17.8)	17.9 (11.5-40.7)	509
	01-02	*	< LOD	.403 (<LOD-.847)	1.69 (.793-3.50)	3.50 (2.19-5.91)	754
	03-04	*	< LOD	.343 (.242-.477)	1.07 (.727-1.84)	3.21 (1.45-6.07)	603
	05-06	*	< LOD	< LOD	1.80 (.990-3.89)	4.95 (2.93-5.59)	722
	07-08	*	< LOD	< LOD	1.79 (1.38-2.31)	4.27 (2.44-6.16)	567
Non-Hispanic whites	99-00	*	< LOD	1.77 (.780-4.02)	11.4 (4.07-21.5)	21.5 (11.4-34.8)	595
	01-02	*	< LOD	.787 (.389-1.37)	3.42 (2.48-5.24)	7.75 (5.80-9.47)	1252
	03-04	*	< LOD	.631 (.412-.897)	3.06 (1.75-5.84)	6.74 (4.42-12.0)	1055
	05-06	*	< LOD	< LOD	2.16 (1.64-3.00)	4.14 (2.91-7.59)	1007
	07-08	*	< LOD	< LOD	2.74 (2.12-3.58)	5.92 (4.00-9.71)	1087

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OP-DPM_BiomonitoringSummary.html

Urinary Antimony

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.132 (.120-.145)	.130 (.120-.150)	.220 (.200-.230)	.330 (.300-.350)	.430 (.390-.470)	2276
	01-02	.134 (.126-.142)	.130 (.130-.140)	.190 (.180-.200)	.270 (.250-.310)	.350 (.320-.400)	2690
	03-04	*	.080 (<LOD-.090)	.130 (.120-.150)	.200 (.190-.220)	.280 (.250-.320)	2558
	05-06	.073 (.066-.081)	.070 (.070-.080)	.120 (.110-.140)	.220 (.180-.250)	.300 (.270-.360)	2576
	07-08	.061 (.057-.066)	.060 (.060-.060)	.100 (.090-.110)	.170 (.140-.200)	.240 (.220-.260)	2627
	09-10	.056 (.053-.059)	.050 (.050-.060)	.090 (.090-.100)	.170 (.140-.180)	.230 (.200-.280)	2847
Age group							
6-11 years	99-00	.176 (.154-.200)	.190 (.160-.210)	.260 (.230-.280)	.350 (.300-.400)	.440 (.320-.600)	316
	01-02	.146 (.134-.160)	.150 (.130-.160)	.200 (.180-.210)	.270 (.240-.330)	.340 (.280-.440)	368
	03-04	.099 (.087-.114)	.100 (.070-.120)	.160 (.120-.200)	.240 (.190-.310)	.310 (.230-.330)	290
	05-06	.075 (.063-.088)	.080 (.060-.090)	.110 (.090-.130)	.190 (.120-.260)	.240 (.170-.340)	355
	07-08	.068 (.061-.077)	.070 (.060-.080)	.110 (.090-.130)	.170 (.150-.210)	.230 (.180-.280)	394
	09-10	.069 (.061-.079)	.070 (.060-.080)	.120 (.100-.150)	.220 (.150-.260)	.260 (.230-.350)	378
12-19 years	99-00	.158 (.141-.178)	.170 (.150-.180)	.240 (.210-.270)	.350 (.290-.420)	.460 (.350-.510)	663
	01-02	.169 (.156-.184)	.160 (.150-.180)	.240 (.220-.260)	.350 (.320-.410)	.460 (.400-.500)	762
	03-04	.105 (.095-.115)	.100 (.090-.120)	.150 (.140-.160)	.230 (.200-.270)	.290 (.250-.370)	725
	05-06	.092 (.083-.101)	.090 (.080-.100)	.140 (.130-.160)	.240 (.200-.270)	.280 (.250-.320)	701
	07-08	.079 (.069-.091)	.080 (.070-.090)	.130 (.110-.140)	.210 (.150-.230)	.230 (.210-.340)	376
	09-10	.063 (.056-.071)	.060 (.050-.070)	.100 (.090-.120)	.180 (.150-.210)	.270 (.180-.370)	451
20 years and older	99-00	.123 (.112-.137)	.120 (.110-.130)	.200 (.180-.220)	.310 (.290-.350)	.430 (.390-.470)	1297
	01-02	.128 (.119-.136)	.130 (.120-.130)	.180 (.170-.190)	.250 (.220-.300)	.330 (.280-.390)	1560
	03-04	*	.070 (<LOD-.080)	.120 (.100-.140)	.190 (.170-.210)	.270 (.220-.320)	1543
	05-06	.070 (.064-.078)	.070 (.060-.080)	.120 (.110-.140)	.220 (.180-.270)	.320 (.260-.420)	1520
	07-08	.058 (.054-.062)	.060 (.050-.060)	.090 (.090-.100)	.160 (.130-.190)	.240 (.210-.270)	1857
	09-10	.054 (.051-.057)	.050 (.050-.050)	.090 (.080-.090)	.150 (.140-.180)	.220 (.190-.270)	2018
Gender							
Males	99-00	.143 (.131-.157)	.150 (.130-.160)	.240 (.220-.260)	.350 (.330-.390)	.470 (.390-.570)	1132
	01-02	.145 (.136-.154)	.140 (.130-.150)	.200 (.190-.210)	.310 (.280-.330)	.390 (.350-.440)	1335
	03-04	.095 (.088-.103)	.090 (.080-.100)	.140 (.130-.160)	.220 (.200-.250)	.320 (.270-.350)	1281
	05-06	.085 (.076-.095)	.080 (.080-.090)	.140 (.120-.160)	.250 (.210-.290)	.350 (.260-.460)	1271
	07-08	.068 (.062-.076)	.070 (.060-.070)	.110 (.100-.120)	.210 (.170-.230)	.280 (.230-.340)	1327
	09-10	.060 (.055-.065)	.060 (.050-.070)	.100 (.090-.110)	.170 (.150-.200)	.250 (.200-.290)	1397
Females	99-00	.122 (.109-.137)	.120 (.110-.140)	.200 (.180-.220)	.300 (.280-.340)	.400 (.350-.460)	1144
	01-02	.125 (.117-.133)	.120 (.120-.130)	.180 (.160-.190)	.240 (.220-.280)	.320 (.260-.360)	1355
	03-04	*	< LOD	.120 (.090-.140)	.180 (.150-.220)	.230 (.190-.330)	1277
	05-06	.063 (.057-.071)	.060 (.050-.070)	.100 (.090-.120)	.180 (.150-.230)	.270 (.200-.330)	1305
	07-08	.055 (.052-.058)	.050 (.050-.060)	.090 (.080-.100)	.130 (.120-.150)	.200 (.170-.230)	1300
	09-10	.052 (.049-.056)	.050 (.040-.050)	.090 (.080-.090)	.150 (.130-.170)	.220 (.190-.270)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.04, 0.04, 0.07, 0.032, 0.032, and 0.032 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Antimony_BiomonitoringSummary.html

Urinary Antimony

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.132 (.108-.161)	.140 (.120-.170)	.210 (.180-.240)	.300 (.260-.390)	.430 (.330-.560)	787
	01-02	.142 (.130-.154)	.130 (.130-.150)	.200 (.170-.230)	.260 (.240-.320)	.360 (.300-.400)	683
	03-04	.093 (.079-.110)	.090 (<LOD-.120)	.140 (.120-.160)	.190 (.160-.260)	.270 (.210-.330)	618
	05-06	.093 (.082-.105)	.090 (.080-.100)	.150 (.140-.170)	.250 (.210-.340)	.470 (.270-.850)	652
	07-08	.069 (.060-.079)	.070 (.060-.080)	.110 (.100-.120)	.190 (.150-.250)	.270 (.220-.390)	515
	09-10	.063 (.060-.067)	.060 (.060-.070)	.110 (.090-.120)	.170 (.150-.200)	.250 (.200-.270)	613
Non-Hispanic blacks	99-00	.175 (.148-.207)	.180 (.150-.200)	.260 (.230-.300)	.400 (.310-.490)	.490 (.410-.710)	554
	01-02	.180 (.164-.197)	.170 (.160-.190)	.250 (.220-.280)	.360 (.320-.410)	.460 (.370-.530)	667
	03-04	.108 (.098-.119)	.110 (.100-.120)	.160 (.150-.190)	.230 (.200-.280)	.310 (.250-.360)	723
	05-06	.088 (.077-.100)	.090 (.080-.100)	.140 (.130-.170)	.210 (.190-.250)	.280 (.240-.320)	692
	07-08	.085 (.079-.092)	.080 (.080-.090)	.130 (.120-.140)	.210 (.180-.250)	.290 (.250-.370)	589
	09-10	.073 (.065-.081)	.070 (.060-.080)	.120 (.110-.140)	.190 (.160-.250)	.280 (.220-.350)	544
Non-Hispanic whites	99-00	.128 (.115-.144)	.130 (.110-.140)	.210 (.190-.230)	.330 (.280-.350)	.400 (.360-.500)	768
	01-02	.126 (.117-.135)	.130 (.120-.130)	.180 (.170-.190)	.250 (.230-.300)	.340 (.310-.390)	1132
	03-04	*	.070 (<LOD-.080)	.130 (.110-.140)	.190 (.170-.210)	.280 (.230-.320)	1074
	05-06	.069 (.062-.077)	.070 (.060-.080)	.110 (.100-.130)	.210 (.170-.260)	.300 (.240-.380)	1041
	07-08	.057 (.052-.063)	.060 (.050-.060)	.090 (.080-.110)	.150 (.130-.200)	.230 (.190-.260)	1095
	09-10	.053 (.050-.057)	.050 (.040-.050)	.090 (.080-.090)	.160 (.130-.190)	.230 (.190-.280)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.04, 0.04, 0.07, 0.032, 0.032, and 0.032 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Antimony_BiomonitoringSummary.html

Urinary Antimony (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.124 (.108-.143)	.119 (.102-.143)	.185 (.164-.214)	.276 (.233-.333)	.385 (.333-.430)	2276
	01-02	.126 (.119-.134)	.120 (.115-.126)	.173 (.162-.188)	.267 (.242-.300)	.364 (.320-.414)	2689
	03-04	*	.080 (<LOD-.086)	.135 (.119-.143)	.208 (.192-.230)	.277 (.250-.294)	2558
	05-06	.072 (.068-.077)	.070 (.060-.070)	.100 (.100-.110)	.160 (.150-.190)	.230 (.190-.290)	2576
	07-08	.064 (.060-.068)	.060 (.060-.060)	.090 (.080-.100)	.140 (.140-.160)	.200 (.170-.230)	2627
	09-10	.060 (.056-.064)	.060 (.050-.060)	.090 (.080-.090)	.140 (.120-.160)	.200 (.180-.230)	2847
Age group							
6-11 years	99-00	.191 (.147-.248)	.185 (.156-.220)	.250 (.200-.417)	.447 (.271-.741)	.741 (.333-1.30)	316
	01-02	.178 (.159-.200)	.173 (.150-.193)	.228 (.200-.272)	.338 (.265-.480)	.471 (.313-.727)	368
	03-04	.116 (.103-.130)	.118 (.098-.136)	.167 (.146-.187)	.256 (.194-.317)	.333 (.250-.500)	290
	05-06	.092 (.081-.104)	.090 (.080-.110)	.130 (.110-.150)	.180 (.150-.210)	.220 (.180-.270)	355
	07-08	.089 (.079-.100)	.090 (.070-.100)	.120 (.110-.140)	.200 (.150-.240)	.300 (.200-.370)	394
	09-10	.094 (.084-.106)	.090 (.080-.100)	.140 (.120-.160)	.200 (.170-.250)	.280 (.220-.320)	378
12-19 years	99-00	.121 (.104-.140)	.120 (.095-.146)	.176 (.146-.207)	.259 (.206-.310)	.310 (.228-.421)	663
	01-02	.121 (.112-.131)	.115 (.106-.127)	.160 (.138-.186)	.224 (.199-.245)	.266 (.244-.310)	762
	03-04	.075 (.068-.082)	.068 (.061-.077)	.100 (.092-.113)	.156 (.126-.173)	.193 (.172-.255)	725
	05-06	.070 (.065-.076)	.070 (.060-.080)	.100 (.090-.110)	.140 (.120-.150)	.170 (.150-.250)	701
	07-08	.062 (.054-.070)	.060 (.050-.070)	.090 (.070-.100)	.120 (.100-.160)	.160 (.110-.240)	376
	09-10	.059 (.053-.066)	.060 (.050-.060)	.090 (.070-.100)	.130 (.110-.170)	.180 (.150-.220)	451
20 years and older	99-00	.118 (.104-.135)	.111 (.097-.135)	.175 (.149-.209)	.263 (.227-.320)	.352 (.320-.391)	1297
	01-02	.122 (.115-.129)	.115 (.108-.121)	.167 (.153-.181)	.265 (.241-.300)	.364 (.318-.405)	1559
	03-04	*	.079 (<LOD-.087)	.135 (.116-.145)	.209 (.195-.233)	.278 (.250-.294)	1543
	05-06	.070 (.066-.075)	.060 (.060-.070)	.100 (.090-.110)	.170 (.150-.190)	.250 (.190-.300)	1520
	07-08	.062 (.058-.066)	.060 (.050-.060)	.090 (.080-.100)	.140 (.130-.160)	.200 (.160-.240)	1857
	09-10	.057 (.053-.061)	.050 (.050-.060)	.080 (.080-.090)	.130 (.120-.140)	.190 (.160-.220)	2018
Gender							
Males	99-00	.112 (.099-.127)	.109 (.095-.127)	.164 (.146-.181)	.226 (.204-.268)	.320 (.235-.391)	1132
	01-02	.114 (.107-.123)	.108 (.103-.115)	.153 (.138-.171)	.228 (.205-.250)	.333 (.281-.438)	1334
	03-04	.080 (.076-.084)	.075 (.069-.081)	.122 (.111-.132)	.192 (.173-.209)	.253 (.230-.278)	1281
	05-06	.070 (.064-.077)	.060 (.060-.070)	.100 (.090-.120)	.160 (.130-.220)	.250 (.170-.310)	1271
	07-08	.061 (.057-.066)	.060 (.050-.060)	.090 (.080-.100)	.140 (.130-.160)	.210 (.160-.260)	1327
	09-10	.055 (.050-.060)	.050 (.050-.060)	.080 (.070-.100)	.130 (.120-.150)	.190 (.160-.210)	1397
Females	99-00	.137 (.117-.161)	.131 (.108-.164)	.213 (.176-.247)	.320 (.263-.417)	.429 (.357-.485)	1144
	01-02	.139 (.131-.148)	.132 (.124-.140)	.196 (.178-.211)	.295 (.267-.317)	.371 (.333-.444)	1355
	03-04	*	< LOD	.143 (.125-.161)	.225 (.188-.261)	.288 (.250-.333)	1277
	05-06	.074 (.070-.078)	.070 (.070-.070)	.110 (.100-.110)	.170 (.150-.190)	.220 (.180-.300)	1305
	07-08	.067 (.062-.071)	.060 (.060-.070)	.100 (.090-.100)	.140 (.130-.160)	.200 (.160-.230)	1300
	09-10	.064 (.060-.069)	.060 (.060-.070)	.090 (.090-.100)	.150 (.130-.170)	.220 (.180-.260)	1450

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Antimony_BiomonitoringSummary.html

Urinary Antimony (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.120 (.107-.135)	.114 (.105-.129)	.167 (.148-.203)	.250 (.209-.315)	.333 (.280-.357)	787
	01-02	.138 (.128-.149)	.130 (.117-.143)	.182 (.159-.203)	.269 (.229-.308)	.338 (.308-.429)	682
	03-04	.086 (.076-.098)	.082 (<LOD-.092)	.129 (.107-.151)	.189 (.154-.238)	.238 (.185-.321)	618
	05-06	.087 (.076-.099)	.080 (.070-.080)	.120 (.110-.130)	.190 (.150-.310)	.370 (.200-.800)	652
	07-08	.069 (.059-.081)	.060 (.050-.080)	.100 (.080-.120)	.160 (.130-.180)	.200 (.160-.360)	515
	09-10	.066 (.063-.071)	.060 (.060-.060)	.100 (.080-.110)	.160 (.130-.190)	.240 (.190-.280)	613
Non-Hispanic blacks	99-00	.114 (.099-.133)	.112 (.098-.130)	.163 (.144-.183)	.236 (.195-.338)	.343 (.255-.425)	554
	01-02	.123 (.113-.134)	.115 (.106-.127)	.163 (.150-.181)	.233 (.208-.267)	.300 (.248-.373)	667
	03-04	.078 (.071-.085)	.074 (.069-.082)	.109 (.096-.124)	.170 (.148-.192)	.222 (.179-.257)	723
	05-06	.064 (.058-.071)	.060 (.050-.070)	.090 (.080-.090)	.130 (.120-.150)	.190 (.150-.220)	692
	07-08	.062 (.059-.066)	.060 (.050-.070)	.090 (.080-.090)	.140 (.120-.160)	.180 (.160-.220)	589
	09-10	.058 (.053-.063)	.060 (.050-.060)	.080 (.070-.090)	.130 (.110-.160)	.170 (.150-.190)	544
Non-Hispanic whites	99-00	.129 (.109-.152)	.125 (.102-.152)	.195 (.167-.225)	.298 (.239-.352)	.400 (.333-.444)	768
	01-02	.127 (.117-.138)	.120 (.113-.130)	.176 (.159-.198)	.280 (.241-.317)	.380 (.318-.471)	1132
	03-04	*	.081 (<LOD-.089)	.139 (.124-.147)	.217 (.200-.238)	.286 (.253-.333)	1074
	05-06	.072 (.068-.077)	.070 (.060-.070)	.110 (.100-.110)	.170 (.150-.190)	.230 (.190-.280)	1041
	07-08	.064 (.060-.069)	.060 (.050-.070)	.090 (.080-.100)	.140 (.140-.160)	.210 (.170-.230)	1095
	09-10	.060 (.055-.065)	.060 (.050-.060)	.090 (.080-.100)	.140 (.120-.170)	.200 (.170-.250)	1225

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Antimony_BiomonitoringSummary.html

Urinary Total Arsenic

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	8.30 (7.19-9.57)	7.70 (6.90-8.90)	16.0 (14.1-18.7)	37.4 (31.6-43.5)	65.4 (48.7-83.3)	2557
	05-06	9.29 (8.05-10.7)	8.65 (7.48-9.99)	17.1 (14.9-20.6)	41.1 (33.3-49.7)	66.7 (53.7-87.0)	2576
	07-08	8.10 (7.44-8.83)	7.49 (6.90-8.12)	14.9 (13.2-17.0)	33.3 (29.8-38.7)	50.8 (42.3-65.1)	2605
	09-10	9.28 (8.47-10.2)	8.15 (7.20-8.98)	18.0 (15.3-20.8)	44.6 (39.0-55.1)	85.6 (64.7-114)	2860
Age group							
6-11 years	03-04	7.08 (5.66-8.84)	6.80 (5.90-7.70)	10.9 (8.90-14.2)	24.6 (13.8-61.8)	46.9 (17.5-178)	290
	05-06	7.19 (5.81-8.90)	6.96 (5.32-8.88)	11.5 (9.19-16.0)	19.6 (13.1-51.5)	34.1 (19.6-58.5)	355
	07-08	6.85 (5.98-7.83)	6.40 (5.74-7.23)	10.8 (9.75-12.3)	22.5 (16.9-34.7)	41.0 (21.1-52.8)	390
	09-10	6.63 (5.74-7.66)	5.94 (5.14-7.19)	10.8 (9.37-12.5)	26.0 (18.2-33.4)	37.7 (27.8-65.6)	378
12-19 years	03-04	8.55 (7.34-9.97)	8.10 (6.80-9.40)	15.2 (12.2-17.8)	30.5 (23.1-40.4)	46.1 (32.9-62.5)	725
	05-06	8.19 (6.87-9.77)	7.92 (6.37-9.50)	14.0 (11.6-18.1)	28.2 (22.9-32.9)	41.9 (32.7-48.0)	701
	07-08	7.09 (6.17-8.14)	6.87 (5.88-7.86)	11.4 (9.41-13.7)	20.4 (16.1-26.6)	38.2 (21.6-53.3)	373
	09-10	6.45 (5.58-7.47)	6.11 (5.26-6.89)	10.8 (8.59-13.7)	25.9 (16.2-32.9)	38.8 (27.8-55.1)	454
20 years and older	03-04	8.41 (7.25-9.77)	7.90 (7.00-9.10)	17.0 (15.0-19.7)	40.5 (34.9-46.2)	66.2 (51.2-93.1)	1542
	05-06	9.76 (8.43-11.3)	9.12 (7.85-10.4)	18.9 (15.8-22.9)	44.2 (35.2-56.1)	71.4 (57.7-98.3)	1520
	07-08	8.43 (7.70-9.22)	7.94 (7.09-8.67)	16.2 (14.5-18.6)	35.2 (30.4-42.3)	59.0 (44.2-75.6)	1842
	09-10	10.2 (9.14-11.3)	8.75 (7.95-9.81)	20.4 (17.2-24.1)	52.1 (42.4-66.1)	93.1 (74.2-127)	2028
Gender							
Males	03-04	9.50 (8.34-10.8)	8.90 (7.70-9.80)	17.6 (15.2-20.1)	41.6 (32.5-52.8)	65.8 (48.7-95.4)	1281
	05-06	10.1 (8.61-11.8)	8.95 (8.05-10.0)	18.3 (15.5-22.9)	40.8 (31.0-52.6)	63.7 (46.4-78.7)	1271
	07-08	9.25 (8.28-10.3)	8.50 (7.37-9.53)	17.0 (14.6-19.4)	36.0 (32.1-44.2)	62.5 (44.3-84.6)	1318
	09-10	10.1 (9.06-11.3)	8.80 (7.80-9.75)	20.4 (16.1-23.9)	47.4 (42.1-64.1)	89.1 (71.6-114)	1401
Females	03-04	7.30 (6.02-8.84)	6.90 (5.90-8.30)	15.0 (11.3-19.5)	33.4 (26.5-41.7)	60.5 (40.8-77.1)	1276
	05-06	8.60 (7.38-10.0)	8.18 (6.64-9.97)	15.9 (13.7-19.9)	41.5 (32.2-53.7)	72.6 (54.8-122)	1305
	07-08	7.14 (6.51-7.82)	6.54 (6.09-7.14)	12.7 (11.6-14.4)	30.1 (26.0-34.0)	49.1 (40.1-57.5)	1287
	09-10	8.55 (7.44-9.83)	7.63 (6.45-8.62)	15.8 (13.1-19.9)	41.5 (31.7-55.5)	81.5 (54.3-132)	1459
Race/ethnicity							
Mexican Americans	03-04	9.29 (8.12-10.6)	9.20 (8.10-10.3)	16.2 (13.5-19.9)	34.4 (24.0-60.5)	68.2 (41.3-111)	618
	05-06	9.55 (8.54-10.7)	9.11 (7.99-10.3)	15.6 (14.0-17.1)	29.2 (21.4-56.8)	67.6 (41.7-81.4)	652
	07-08	8.98 (8.13-9.92)	8.84 (7.80-9.48)	15.4 (12.3-19.7)	35.2 (25.0-46.0)	53.0 (44.2-77.4)	510
	09-10	8.47 (7.30-9.84)	7.96 (6.87-9.08)	13.9 (11.3-17.7)	34.7 (25.0-53.2)	60.9 (49.3-78.7)	613
Non-Hispanic blacks	03-04	11.6 (9.50-14.1)	10.4 (7.90-11.8)	21.5 (14.9-34.4)	43.5 (36.2-61.8)	78.0 (43.6-141)	722
	05-06	11.0 (8.60-14.0)	9.55 (6.99-13.3)	21.9 (14.9-28.9)	44.9 (31.1-71.4)	82.3 (49.2-164)	692
	07-08	10.5 (9.40-11.7)	9.21 (8.22-10.4)	18.4 (16.1-21.5)	42.4 (32.9-52.8)	65.6 (45.5-112)	585
	09-10	10.9 (9.46-12.5)	9.26 (7.70-11.4)	21.7 (17.6-24.2)	49.1 (32.2-81.7)	84.8 (51.3-174)	546
Non-Hispanic whites	03-04	7.12 (6.13-8.27)	7.00 (6.10-7.90)	13.7 (11.3-15.8)	29.0 (22.6-35.9)	53.1 (38.4-65.6)	1074
	05-06	8.66 (7.20-10.4)	8.05 (6.52-9.66)	16.3 (13.4-20.6)	40.8 (29.4-50.2)	58.5 (46.0-88.0)	1041
	07-08	6.98 (6.31-7.71)	6.46 (5.93-7.29)	12.4 (11.3-14.3)	28.3 (21.6-32.6)	42.1 (32.3-50.0)	1088
	09-10	8.18 (7.46-8.96)	7.24 (6.46-8.21)	14.8 (12.7-17.4)	38.9 (31.7-44.4)	66.3 (49.3-88.7)	1224

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.74, 0.74, 0.74, and 0.74 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Total Arsenic (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in $\mu\text{g/g}$ of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	8.24 (7.07-9.59)	7.04 (5.93-8.51)	14.1 (11.6-17.2)	30.4 (26.0-38.7)	50.4 (40.3-64.5)	2557
	05-06	9.15 (7.93-10.6)	7.70 (6.55-8.98)	15.2 (11.7-19.4)	35.1 (26.5-44.7)	62.8 (44.7-85.0)	2576
	07-08	8.46 (7.78-9.21)	7.06 (6.56-7.74)	13.8 (12.5-15.2)	28.9 (24.2-36.7)	49.0 (38.8-70.5)	2605
	09-10	9.90 (9.06-10.8)	7.90 (6.98-8.97)	17.6 (15.4-20.2)	45.2 (36.6-53.3)	80.8 (60.5-94.4)	2860
Age group							
6-11 years	03-04	8.25 (6.58-10.3)	7.18 (5.93-9.45)	11.7 (9.10-16.3)	22.2 (12.0-69.5)	40.1 (14.7-188)	290
	05-06	8.88 (7.05-11.2)	7.87 (6.19-9.42)	11.8 (9.32-18.9)	24.5 (13.0-62.8)	45.4 (22.9-80.9)	355
	07-08	8.87 (8.06-9.77)	7.53 (6.73-7.88)	13.4 (10.2-16.0)	26.7 (19.9-34.0)	37.2 (28.6-71.4)	390
	09-10	8.97 (7.93-10.2)	7.42 (6.70-8.32)	11.0 (9.90-14.0)	33.5 (17.0-54.6)	60.8 (36.6-84.9)	378
12-19 years	03-04	6.11 (5.23-7.13)	5.06 (4.47-6.04)	9.66 (7.44-11.2)	17.8 (12.0-26.0)	27.8 (20.7-35.9)	725
	05-06	6.30 (5.56-7.14)	5.19 (4.80-6.19)	9.62 (8.12-11.1)	19.4 (13.9-25.8)	28.0 (21.9-33.2)	701
	07-08	5.50 (4.91-6.16)	4.96 (4.25-5.40)	7.69 (6.09-9.31)	16.8 (10.8-21.1)	22.5 (16.8-29.1)	373
	09-10	6.06 (5.34-6.87)	4.95 (4.39-5.81)	9.18 (7.00-11.0)	19.2 (14.5-21.3)	28.4 (20.8-35.7)	454
20 years and older	03-04	8.64 (7.38-10.1)	7.47 (6.20-9.01)	15.4 (12.7-18.8)	33.8 (27.3-41.2)	53.9 (45.4-64.5)	1542
	05-06	9.75 (8.46-11.2)	8.22 (6.98-9.75)	17.0 (12.8-21.3)	41.0 (29.6-52.5)	68.4 (52.8-89.7)	1520
	07-08	9.00 (8.20-9.88)	7.55 (6.79-8.53)	14.9 (13.1-17.1)	32.5 (25.8-41.0)	59.4 (41.0-86.2)	1842
	09-10	10.8 (9.71-12.0)	8.73 (7.69-9.71)	20.1 (16.5-24.2)	50.8 (40.5-59.7)	87.3 (70.0-105)	2028
Gender							
Males	03-04	8.00 (6.81-9.40)	6.75 (5.66-8.35)	13.7 (11.0-18.0)	28.7 (25.1-36.4)	45.6 (35.3-62.1)	1281
	05-06	8.26 (7.09-9.63)	7.16 (5.87-8.54)	12.9 (10.2-18.0)	28.8 (22.9-36.3)	46.1 (35.1-66.5)	1271
	07-08	8.30 (7.49-9.18)	6.79 (6.39-7.61)	13.2 (11.8-15.2)	28.9 (23.5-38.5)	47.7 (35.7-68.1)	1318
	09-10	9.21 (8.55-9.93)	7.22 (6.43-8.67)	17.0 (14.9-18.6)	41.8 (35.4-47.9)	66.9 (52.0-81.5)	1401
Females	03-04	8.47 (7.12-10.1)	7.33 (6.10-8.75)	14.4 (11.7-17.7)	32.3 (24.2-46.6)	58.4 (42.8-75.0)	1276
	05-06	10.1 (8.72-11.7)	8.29 (7.23-9.87)	17.4 (13.0-21.4)	43.8 (29.2-61.5)	74.1 (55.0-96.2)	1305
	07-08	8.63 (7.91-9.41)	7.13 (6.53-8.33)	14.1 (12.3-16.9)	27.9 (24.1-37.0)	51.4 (39.7-83.3)	1287
	09-10	10.6 (9.36-12.0)	8.38 (7.40-9.41)	18.6 (14.9-23.9)	50.8 (35.1-72.9)	87.8 (66.8-109)	1459
Race/ethnicity							
Mexican Americans	03-04	8.61 (7.33-10.1)	7.76 (6.30-9.44)	12.6 (10.2-15.9)	24.0 (17.7-34.8)	42.4 (24.8-62.4)	618
	05-06	8.98 (7.89-10.2)	7.48 (6.72-8.87)	12.8 (11.0-16.3)	28.1 (21.1-35.5)	49.1 (31.0-108)	652
	07-08	8.88 (7.71-10.2)	7.38 (6.51-8.63)	14.1 (10.6-17.8)	28.6 (22.5-35.3)	48.2 (31.1-75.7)	510
	09-10	8.88 (7.87-10.0)	7.62 (6.69-8.45)	13.2 (10.7-17.2)	31.4 (23.7-38.5)	49.8 (37.5-68.9)	613
Non-Hispanic blacks	03-04	8.31 (6.99-9.88)	6.88 (5.66-8.41)	13.8 (11.5-17.0)	27.6 (17.9-56.0)	54.3 (27.5-120)	722
	05-06	7.96 (6.40-9.92)	6.48 (5.21-8.30)	13.4 (10.0-18.6)	32.5 (18.7-66.5)	71.4 (35.6-98.8)	692
	07-08	7.72 (6.98-8.54)	6.60 (6.01-7.56)	13.4 (11.5-15.4)	25.7 (22.1-30.0)	42.7 (31.4-60.3)	585
	09-10	8.67 (7.54-9.98)	7.26 (5.96-9.02)	15.6 (12.4-21.3)	38.7 (31.5-48.7)	63.9 (43.1-102)	546
Non-Hispanic whites	03-04	7.50 (6.25-9.01)	6.32 (5.28-7.96)	12.5 (9.86-17.1)	26.8 (21.8-32.0)	40.0 (31.3-53.9)	1074
	05-06	9.01 (7.57-10.7)	7.68 (6.18-9.56)	14.3 (11.1-20.8)	31.9 (24.1-46.1)	59.4 (37.9-96.2)	1041
	07-08	7.82 (7.05-8.68)	6.72 (6.00-7.54)	12.5 (11.1-13.9)	24.6 (19.6-32.3)	43.1 (29.1-64.0)	1088
	09-10	9.14 (8.35-10.0)	7.10 (6.43-8.06)	15.8 (13.7-18.4)	40.9 (30.4-50.8)	66.4 (50.9-90.0)	1224

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Arsenic (V) acid

Geometric mean and selected percentiles of urine concentrations (in µg As/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	1.10 (<LOD-1.50)	2568
	05-06	*	< LOD	< LOD	< LOD	1.06 (<LOD-1.43)	2588
	07-08	*	< LOD	< LOD	< LOD	< LOD	2576
	09-10	*	< LOD	< LOD	< LOD	< LOD	2852
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	1.10 (<LOD-1.30)	292
	05-06	*	< LOD	< LOD	< LOD	< LOD	354
	07-08	*	< LOD	< LOD	< LOD	< LOD	390
	09-10	*	< LOD	< LOD	< LOD	< LOD	379
12-19 years	03-04	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.60)	728
	05-06	*	< LOD	< LOD	< LOD	1.00 (<LOD-1.30)	703
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
	09-10	*	< LOD	< LOD	< LOD	< LOD	453
20 years and older	03-04	*	< LOD	< LOD	< LOD	1.10 (<LOD-1.50)	1548
	05-06	*	< LOD	< LOD	< LOD	1.09 (<LOD-1.71)	1531
	07-08	*	< LOD	< LOD	< LOD	< LOD	1820
	09-10	*	< LOD	< LOD	< LOD	< LOD	2020
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.50)	1284
	05-06	*	< LOD	< LOD	< LOD	1.14 (<LOD-1.71)	1276
	07-08	*	< LOD	< LOD	< LOD	< LOD	1289
	09-10	*	< LOD	< LOD	< LOD	< LOD	1396
Females	03-04	*	< LOD	< LOD	< LOD	1.10 (<LOD-1.30)	1284
	05-06	*	< LOD	< LOD	< LOD	1.01 (<LOD-1.22)	1312
	07-08	*	< LOD	< LOD	< LOD	< LOD	1287
	09-10	*	< LOD	< LOD	< LOD	< LOD	1456
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.60)	621
	05-06	*	< LOD	< LOD	< LOD	< LOD	651
	07-08	*	< LOD	< LOD	< LOD	< LOD	513
	09-10	*	< LOD	< LOD	< LOD	< LOD	618
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.80)	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	586
	09-10	*	< LOD	< LOD	< LOD	< LOD	543
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	1.10 (<LOD-1.50)	1078
	05-06	*	< LOD	< LOD	< LOD	1.10 (<LOD-1.74)	1050
	07-08	*	< LOD	< LOD	< LOD	< LOD	1063
	09-10	*	< LOD	< LOD	< LOD	< LOD	1210

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 1.0, 1.0, 1.0, and 1.0 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Arsenic (V) acid (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg As/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	3.04 (<LOD-3.50)	2568
	05-06	*	< LOD	< LOD	< LOD	3.23 (<LOD-3.55)	2588
	07-08	*	< LOD	< LOD	< LOD	< LOD	2576
	09-10	*	< LOD	< LOD	< LOD	< LOD	2851
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	2.80 (<LOD-4.00)	292
	05-06	*	< LOD	< LOD	< LOD	< LOD	354
	07-08	*	< LOD	< LOD	< LOD	< LOD	390
	09-10	*	< LOD	< LOD	< LOD	< LOD	378
12-19 years	03-04	*	< LOD	< LOD	< LOD	1.75 (<LOD-2.41)	728
	05-06	*	< LOD	< LOD	< LOD	1.69 (<LOD-2.73)	703
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
	09-10	*	< LOD	< LOD	< LOD	< LOD	453
20 years and older	03-04	*	< LOD	< LOD	< LOD	3.18 (<LOD-3.70)	1548
	05-06	*	< LOD	< LOD	< LOD	3.38 (<LOD-3.94)	1531
	07-08	*	< LOD	< LOD	< LOD	< LOD	1820
	09-10	*	< LOD	< LOD	< LOD	< LOD	2020
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	2.61 (<LOD-3.18)	1284
	05-06	*	< LOD	< LOD	< LOD	2.14 (<LOD-2.73)	1276
	07-08	*	< LOD	< LOD	< LOD	< LOD	1289
	09-10	*	< LOD	< LOD	< LOD	< LOD	1395
Females	03-04	*	< LOD	< LOD	< LOD	3.33 (<LOD-3.89)	1284
	05-06	*	< LOD	< LOD	< LOD	3.55 (<LOD-4.44)	1312
	07-08	*	< LOD	< LOD	< LOD	< LOD	1287
	09-10	*	< LOD	< LOD	< LOD	< LOD	1456
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	2.69 (<LOD-3.50)	621
	05-06	*	< LOD	< LOD	< LOD	< LOD	651
	07-08	*	< LOD	< LOD	< LOD	< LOD	513
	09-10	*	< LOD	< LOD	< LOD	< LOD	618
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	1.75 (<LOD-2.19)	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	586
	09-10	*	< LOD	< LOD	< LOD	< LOD	542
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	3.33 (<LOD-3.95)	1078
	05-06	*	< LOD	< LOD	< LOD	3.38 (<LOD-3.80)	1050
	07-08	*	< LOD	< LOD	< LOD	< LOD	1063
	09-10	*	< LOD	< LOD	< LOD	< LOD	1210

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Arsenobetaine

Geometric mean and selected percentiles of urine concentrations (in µg As/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	1.55 (1.31-1.83)	1.00 (.800-1.40)	5.20 (4.00-6.50)	16.8 (12.7-22.3)	35.0 (27.6-44.6)	2568
	05-06	1.86 (1.43-2.41)	1.53 (.980-2.26)	6.73 (4.80-8.91)	22.6 (16.5-30.3)	40.6 (30.7-59.4)	2588
	07-08	*	.670 (.510-.910)	4.26 (3.53-4.88)	16.6 (12.5-20.2)	29.5 (22.8-37.1)	2576
	09-10	1.59 (1.38-1.83)	.940 (.720-1.26)	6.18 (4.67-7.87)	23.5 (19.8-30.1)	50.4 (35.7-63.2)	2870
Age group							
6-11 years	03-04	*	< LOD	1.80 (.800-4.00)	8.80 (3.90-29.9)	29.9 (6.20-190)	292
	05-06	*	< LOD	2.22 (.620-5.35)	7.14 (2.34-29.9)	18.9 (5.46-45.0)	354
	07-08	*	< LOD	1.24 (.780-2.42)	6.71 (3.37-8.38)	13.0 (7.22-25.2)	390
	09-10	*	< LOD	1.20 (.590-3.12)	7.98 (5.20-15.7)	19.4 (9.25-47.9)	380
12-19 years	03-04	*	.600 (.400-.800)	3.20 (2.00-4.70)	13.9 (7.20-25.1)	31.8 (17.2-35.8)	728
	05-06	*	.550 (<LOD-1.14)	3.76 (1.95-7.16)	12.4 (9.15-20.4)	25.6 (16.4-32.4)	703
	07-08	*	< LOD	1.72 (.780-2.62)	4.88 (3.49-9.17)	10.4 (6.81-23.6)	366
	09-10	*	< LOD	2.07 (1.26-3.60)	11.0 (5.35-16.7)	17.8 (12.8-24.0)	455
20 years and older	03-04	1.74 (1.48-2.05)	1.30 (1.00-1.60)	6.10 (4.90-7.10)	18.5 (14.0-23.5)	35.5 (26.8-50.5)	1548
	05-06	2.20 (1.71-2.84)	1.89 (1.41-2.64)	8.14 (5.86-10.4)	27.6 (18.7-35.7)	43.9 (36.0-67.9)	1531
	07-08	1.46 (1.28-1.67)	.960 (.750-1.22)	5.08 (4.30-6.20)	19.1 (14.3-23.7)	31.7 (23.9-46.1)	1820
	09-10	1.92 (1.63-2.26)	1.33 (.960-1.91)	7.60 (6.06-9.90)	28.2 (21.9-34.9)	59.5 (45.7-76.7)	2035
Gender							
Males	03-04	1.66 (1.43-1.93)	1.20 (.900-1.50)	5.80 (4.40-7.10)	18.6 (13.9-23.7)	35.0 (26.8-40.5)	1284
	05-06	1.83 (1.40-2.40)	1.44 (.920-2.04)	6.63 (5.06-8.66)	23.2 (12.8-33.4)	37.1 (28.5-51.3)	1276
	07-08	*	.910 (.660-1.26)	5.07 (4.01-6.81)	19.1 (13.8-22.6)	30.3 (22.8-46.1)	1289
	09-10	1.71 (1.48-1.97)	.990 (.780-1.33)	7.14 (5.38-9.86)	26.8 (21.5-31.8)	57.4 (36.4-64.6)	1402
Females	03-04	1.45 (1.17-1.80)	.900 (.700-1.40)	4.70 (3.40-6.20)	15.6 (11.1-25.3)	32.7 (21.1-51.3)	1284
	05-06	1.88 (1.42-2.50)	1.64 (.950-2.51)	6.81 (4.03-9.88)	21.8 (17.3-30.1)	43.3 (33.8-67.9)	1312
	07-08	*	.530 (.410-.650)	3.52 (2.94-4.35)	14.0 (10.9-17.5)	29.5 (19.4-34.0)	1287
	09-10	1.49 (1.20-1.84)	.890 (.590-1.35)	5.54 (3.56-7.74)	21.2 (14.7-32.7)	47.1 (32.5-76.4)	1468
Race/ethnicity							
Mexican Americans	03-04	1.19 (.871-1.62)	.800 (.500-1.30)	3.20 (1.80-5.20)	10.2 (6.70-21.4)	31.4 (16.3-39.1)	621
	05-06	1.44 (1.23-1.69)	1.04 (.770-1.35)	3.90 (3.03-5.06)	16.8 (11.1-32.0)	41.4 (23.4-57.9)	651
	07-08	*	.560 (<LOD-.910)	3.27 (2.00-5.65)	16.9 (9.73-20.8)	29.6 (19.3-40.5)	513
	09-10	*	.640 (.440-.800)	3.58 (1.80-5.89)	15.5 (7.83-24.6)	31.9 (18.7-44.7)	617
Non-Hispanic blacks	03-04	2.29 (1.60-3.28)	2.00 (1.20-3.50)	7.70 (5.00-12.0)	23.7 (13.2-38.7)	45.6 (25.1-94.0)	725
	05-06	2.55 (1.64-3.99)	2.42 (1.15-4.58)	8.74 (6.01-14.4)	23.8 (15.9-48.9)	59.6 (30.5-121)	695
	07-08	1.58 (1.31-1.90)	1.11 (.750-1.60)	5.88 (4.12-8.13)	21.9 (16.4-24.9)	35.9 (25.4-47.7)	586
	09-10	1.99 (1.60-2.47)	1.79 (1.05-2.41)	7.61 (5.34-10.0)	25.5 (15.4-37.6)	51.2 (31.0-90.9)	546
Non-Hispanic whites	03-04	1.37 (1.11-1.68)	.800 (.700-1.20)	4.30 (2.50-6.30)	13.3 (9.70-21.4)	29.3 (21.4-35.5)	1078
	05-06	1.74 (1.27-2.40)	1.41 (.750-2.27)	6.40 (3.93-9.04)	22.4 (14.1-30.1)	38.4 (27.8-66.8)	1050
	07-08	*	.530 (<LOD-.780)	3.63 (3.06-4.38)	12.6 (9.08-17.1)	24.3 (17.6-30.3)	1063
	09-10	1.43 (1.24-1.65)	.800 (.610-1.18)	5.40 (3.77-7.27)	20.6 (17.7-24.5)	41.3 (30.6-59.5)	1226

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.4, 0.4, 0.4, and 0.4 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Arsenobetaine (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg As/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	1.54 (1.30-1.82)	1.16 (.959-1.43)	5.00 (3.62-6.91)	16.2 (12.5-20.3)	29.4 (24.0-36.4)	2568
	05-06	1.82 (1.40-2.37)	1.64 (1.10-2.41)	6.64 (4.48-8.79)	20.3 (14.5-28.1)	37.0 (27.5-58.3)	2588
	07-08	*	1.00 (.850-1.18)	4.07 (3.36-4.98)	14.7 (11.1-17.8)	27.7 (18.9-45.4)	2576
	09-10	1.69 (1.45-1.97)	1.23 (.980-1.47)	6.48 (4.90-8.00)	23.6 (18.5-30.7)	49.6 (36.6-61.3)	2869
Age group							
6-11 years	03-04	*	< LOD	2.00 (1.15-4.83)	12.2 (4.13-39.7)	29.6 (6.80-153)	292
	05-06	*	< LOD	2.88 (1.04-6.25)	8.77 (2.88-40.5)	23.6 (6.86-74.8)	354
	07-08	*	< LOD	2.11 (1.27-3.79)	8.98 (4.76-14.8)	16.4 (8.98-23.8)	390
	09-10	*	< LOD	1.70 (1.21-3.02)	9.52 (5.08-20.7)	24.9 (9.81-50.7)	379
12-19 years	03-04	*	.531 (.400-.638)	2.14 (1.39-3.51)	9.29 (4.29-14.7)	17.3 (10.4-28.7)	728
	05-06	*	.620 (<LOD-.800)	2.82 (1.58-4.77)	10.5 (7.03-13.7)	15.4 (11.8-22.3)	703
	07-08	*	< LOD	1.28 (.650-1.72)	5.17 (2.31-6.53)	9.88 (5.72-16.0)	366
	09-10	*	< LOD	2.25 (1.17-3.76)	8.69 (5.03-12.8)	14.0 (9.82-15.5)	455
20 years and older	03-04	1.79 (1.51-2.12)	1.47 (1.15-1.88)	5.91 (4.32-7.72)	17.2 (13.4-21.8)	30.1 (26.1-36.4)	1548
	05-06	2.19 (1.70-2.82)	2.13 (1.40-2.83)	7.79 (5.38-10.2)	24.4 (16.6-32.9)	43.1 (30.5-71.4)	1531
	07-08	1.55 (1.36-1.78)	1.29 (1.13-1.52)	5.27 (3.91-6.40)	16.6 (12.2-21.6)	33.4 (21.0-55.9)	1820
	09-10	2.03 (1.70-2.42)	1.52 (1.18-2.16)	7.65 (6.04-10.4)	27.7 (22.1-35.9)	57.0 (39.7-71.9)	2035
Gender							
Males	03-04	1.40 (1.18-1.67)	1.11 (.909-1.28)	4.78 (3.61-6.70)	14.4 (11.1-18.5)	26.5 (18.6-29.9)	1284
	05-06	1.50 (1.15-1.96)	1.25 (.800-1.93)	5.11 (3.41-7.54)	15.3 (11.2-20.8)	30.5 (19.6-44.0)	1276
	07-08	*	1.04 (.870-1.25)	4.11 (3.16-5.30)	14.9 (10.1-17.9)	27.0 (17.8-49.2)	1289
	09-10	1.56 (1.35-1.80)	1.04 (.880-1.34)	6.20 (4.34-7.79)	23.4 (18.7-27.6)	39.2 (32.0-54.8)	1401
Females	03-04	1.68 (1.37-2.05)	1.25 (.938-1.67)	5.58 (3.50-7.43)	17.2 (12.3-24.5)	32.9 (25.6-46.3)	1284
	05-06	2.20 (1.65-2.93)	2.04 (1.30-2.92)	7.86 (5.38-10.8)	26.4 (16.6-35.6)	46.2 (30.3-74.3)	1312
	07-08	*	.980 (.800-1.18)	4.06 (3.35-5.29)	14.6 (11.1-17.8)	29.6 (18.2-50.0)	1287
	09-10	1.83 (1.48-2.26)	1.36 (1.04-1.75)	6.65 (4.68-9.21)	24.1 (16.5-37.3)	57.0 (36.3-71.3)	1468
Race/ethnicity							
Mexican Americans	03-04	1.10 (.786-1.55)	.877 (.612-1.40)	2.93 (1.78-5.21)	8.88 (5.50-15.4)	19.0 (9.64-29.4)	621
	05-06	1.35 (1.13-1.62)	.970 (.810-1.19)	4.21 (2.79-5.68)	16.6 (10.5-21.3)	36.9 (18.4-69.5)	651
	07-08	*	.780 (<LOD-1.21)	3.04 (1.99-5.09)	13.7 (9.09-16.7)	28.9 (16.2-46.7)	513
	09-10	*	.790 (.670-.940)	3.64 (2.28-4.41)	13.5 (7.78-19.1)	26.8 (16.5-42.0)	617
Non-Hispanic blacks	03-04	1.65 (1.19-2.30)	1.53 (.901-2.45)	5.81 (4.25-7.82)	13.6 (9.76-27.9)	32.9 (13.4-82.1)	725
	05-06	1.84 (1.20-2.83)	1.71 (.910-2.95)	6.07 (3.83-9.25)	19.4 (10.4-45.2)	47.5 (25.3-62.0)	695
	07-08	1.16 (.955-1.41)	.830 (.620-1.12)	4.44 (3.23-6.58)	14.9 (10.5-17.6)	25.1 (17.4-40.0)	586
	09-10	1.59 (1.29-1.96)	1.30 (.850-2.04)	5.92 (4.20-8.40)	19.5 (13.5-27.1)	37.9 (26.1-51.7)	545
Non-Hispanic whites	03-04	1.44 (1.15-1.80)	1.05 (.833-1.36)	4.47 (2.73-6.83)	14.3 (10.9-18.6)	26.5 (18.6-32.0)	1078
	05-06	1.81 (1.32-2.47)	1.65 (.970-2.55)	6.65 (3.95-9.34)	19.7 (12.6-30.5)	36.6 (25.1-71.4)	1050
	07-08	*	.970 (<LOD-1.18)	3.56 (2.93-4.53)	11.6 (8.98-16.6)	23.8 (15.5-41.7)	1063
	09-10	1.60 (1.36-1.87)	1.15 (.900-1.43)	5.85 (4.33-7.65)	22.8 (15.6-29.4)	38.5 (32.0-57.0)	1226

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Arsenocholine

Geometric mean and selected percentiles of urine concentrations (in µg As/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2568
	05-06	*	< LOD	< LOD	< LOD	< LOD	2588
	07-08	*	< LOD	< LOD	< LOD	< LOD	2576
	09-10	*	< LOD	< LOD	< LOD	< LOD	2871
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	292
	05-06	*	< LOD	< LOD	< LOD	< LOD	354
	07-08	*	< LOD	< LOD	< LOD	< LOD	390
	09-10	*	< LOD	< LOD	< LOD	< LOD	380
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	728
	05-06	*	< LOD	< LOD	< LOD	< LOD	703
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
	09-10	*	< LOD	< LOD	< LOD	< LOD	456
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1548
	05-06	*	< LOD	< LOD	< LOD	< LOD	1531
	07-08	*	< LOD	< LOD	< LOD	< LOD	1820
	09-10	*	< LOD	< LOD	< LOD	< LOD	2035
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1276
	07-08	*	< LOD	< LOD	< LOD	< LOD	1289
	09-10	*	< LOD	< LOD	< LOD	< LOD	1403
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1312
	07-08	*	< LOD	< LOD	< LOD	< LOD	1287
	09-10	*	< LOD	< LOD	< LOD	< LOD	1468
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	621
	05-06	*	< LOD	< LOD	< LOD	< LOD	651
	07-08	*	< LOD	< LOD	< LOD	< LOD	513
	09-10	*	< LOD	< LOD	< LOD	< LOD	618
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	586
	09-10	*	< LOD	< LOD	< LOD	< LOD	546
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1078
	05-06	*	< LOD	< LOD	< LOD	< LOD	1050
	07-08	*	< LOD	< LOD	< LOD	< LOD	1063
	09-10	*	< LOD	< LOD	< LOD	< LOD	1226

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.6, 0.6, 0.6, and 0.6 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Arsenocholine (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in $\mu\text{g As/g}$ of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2568
	05-06	*	< LOD	< LOD	< LOD	< LOD	2588
	07-08	*	< LOD	< LOD	< LOD	< LOD	2576
	09-10	*	< LOD	< LOD	< LOD	< LOD	2870
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	292
	05-06	*	< LOD	< LOD	< LOD	< LOD	354
	07-08	*	< LOD	< LOD	< LOD	< LOD	390
	09-10	*	< LOD	< LOD	< LOD	< LOD	379
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	728
	05-06	*	< LOD	< LOD	< LOD	< LOD	703
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
	09-10	*	< LOD	< LOD	< LOD	< LOD	456
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1548
	05-06	*	< LOD	< LOD	< LOD	< LOD	1531
	07-08	*	< LOD	< LOD	< LOD	< LOD	1820
	09-10	*	< LOD	< LOD	< LOD	< LOD	2035
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1276
	07-08	*	< LOD	< LOD	< LOD	< LOD	1289
	09-10	*	< LOD	< LOD	< LOD	< LOD	1402
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1312
	07-08	*	< LOD	< LOD	< LOD	< LOD	1287
	09-10	*	< LOD	< LOD	< LOD	< LOD	1468
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	621
	05-06	*	< LOD	< LOD	< LOD	< LOD	651
	07-08	*	< LOD	< LOD	< LOD	< LOD	513
	09-10	*	< LOD	< LOD	< LOD	< LOD	618
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	586
	09-10	*	< LOD	< LOD	< LOD	< LOD	545
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1078
	05-06	*	< LOD	< LOD	< LOD	< LOD	1050
	07-08	*	< LOD	< LOD	< LOD	< LOD	1063
	09-10	*	< LOD	< LOD	< LOD	< LOD	1226

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Arsenous (III) acid

Geometric mean and selected percentiles of urine concentrations (in µg As/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2568
	05-06	*	< LOD	< LOD	< LOD	< LOD	2588
	07-08	*	< LOD	< LOD	< LOD	< LOD	2576
	09-10	*	< LOD	< LOD	< LOD	< LOD	2871
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	292
	05-06	*	< LOD	< LOD	< LOD	< LOD	354
	07-08	*	< LOD	< LOD	< LOD	< LOD	390
	09-10	*	< LOD	< LOD	< LOD	< LOD	380
12-19 years	03-04	*	< LOD	< LOD	< LOD	1.40 (<LOD-1.70)	728
	05-06	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.40)	703
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
	09-10	*	< LOD	< LOD	< LOD	< LOD	456
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1548
	05-06	*	< LOD	< LOD	< LOD	< LOD	1531
	07-08	*	< LOD	< LOD	< LOD	< LOD	1820
	09-10	*	< LOD	< LOD	< LOD	< LOD	2035
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	1.26 (<LOD-1.73)	1276
	07-08	*	< LOD	< LOD	< LOD	< LOD	1289
	09-10	*	< LOD	< LOD	< LOD	1.24 (<LOD-1.50)	1403
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1312
	07-08	*	< LOD	< LOD	< LOD	< LOD	1287
	09-10	*	< LOD	< LOD	< LOD	< LOD	1468
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	2.00 (<LOD-3.00)	621
	05-06	*	< LOD	< LOD	< LOD	< LOD	651
	07-08	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.76)	513
	09-10	*	< LOD	< LOD	< LOD	< LOD	618
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.80)	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	586
	09-10	*	< LOD	< LOD	< LOD	< LOD	546
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1078
	05-06	*	< LOD	< LOD	< LOD	< LOD	1050
	07-08	*	< LOD	< LOD	< LOD	< LOD	1063
	09-10	*	< LOD	< LOD	< LOD	< LOD	1226

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 1.2, 1.2, 1.2, and 1.2 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Arsenous (III) acid (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg As/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2568
	05-06	*	< LOD	< LOD	< LOD	< LOD	2588
	07-08	*	< LOD	< LOD	< LOD	< LOD	2576
	09-10	*	< LOD	< LOD	< LOD	< LOD	2870
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	292
	05-06	*	< LOD	< LOD	< LOD	< LOD	354
	07-08	*	< LOD	< LOD	< LOD	< LOD	390
	09-10	*	< LOD	< LOD	< LOD	< LOD	379
12-19 years	03-04	*	< LOD	< LOD	< LOD	1.95 (<LOD-2.76)	728
	05-06	*	< LOD	< LOD	< LOD	2.02 (<LOD-3.04)	703
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
	09-10	*	< LOD	< LOD	< LOD	< LOD	456
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1548
	05-06	*	< LOD	< LOD	< LOD	< LOD	1531
	07-08	*	< LOD	< LOD	< LOD	< LOD	1820
	09-10	*	< LOD	< LOD	< LOD	< LOD	2035
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	2.43 (<LOD-3.15)	1276
	07-08	*	< LOD	< LOD	< LOD	< LOD	1289
	09-10	*	< LOD	< LOD	< LOD	2.93 (<LOD-3.54)	1402
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1312
	07-08	*	< LOD	< LOD	< LOD	< LOD	1287
	09-10	*	< LOD	< LOD	< LOD	< LOD	1468
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	3.08 (<LOD-4.44)	621
	05-06	*	< LOD	< LOD	< LOD	< LOD	651
	07-08	*	< LOD	< LOD	< LOD	3.54 (<LOD-4.72)	513
	09-10	*	< LOD	< LOD	< LOD	< LOD	618
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	2.00 (<LOD-2.29)	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	586
	09-10	*	< LOD	< LOD	< LOD	< LOD	545
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1078
	05-06	*	< LOD	< LOD	< LOD	< LOD	1050
	07-08	*	< LOD	< LOD	< LOD	< LOD	1063
	09-10	*	< LOD	< LOD	< LOD	< LOD	1226

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Dimethylarsinic acid

Metabolite of Arsenic

Geometric mean and selected percentiles of urine concentrations (in µg As/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	3.71 (3.33-4.14)	3.90 (3.00-4.00)	6.00 (5.00-7.00)	11.0 (9.20-12.0)	16.0 (13.0-17.8)	2568
	05-06	3.95 (3.59-4.35)	3.90 (3.46-4.31)	6.17 (5.53-7.04)	10.3 (9.00-12.2)	15.0 (12.2-17.8)	2588
	07-08	3.67 (3.44-3.91)	3.58 (3.44-3.84)	5.96 (5.44-6.57)	9.78 (8.72-11.2)	13.3 (11.8-15.3)	2576
	09-10	3.67 (3.43-3.93)	3.50 (3.25-3.76)	6.16 (5.59-6.73)	11.5 (9.35-13.7)	17.4 (14.0-21.0)	2871
Age group							
6-11 years	03-04	3.73 (3.12-4.45)	4.00 (3.00-4.00)	6.00 (5.00-7.00)	9.00 (7.00-12.0)	12.0 (8.00-22.0)	292
	05-06	3.96 (3.49-4.49)	3.94 (3.20-4.71)	5.94 (5.01-7.10)	10.5 (7.10-12.1)	13.0 (10.7-15.3)	354
	07-08	3.86 (3.45-4.32)	3.76 (3.47-4.01)	6.10 (5.29-6.99)	10.4 (7.57-13.9)	15.2 (9.32-39.2)	390
	09-10	3.53 (3.17-3.92)	3.27 (2.81-3.87)	5.76 (5.07-6.53)	10.7 (7.84-11.9)	12.8 (11.0-14.4)	380
12-19 years	03-04	3.85 (3.34-4.42)	4.00 (3.00-4.00)	6.00 (5.00-7.10)	9.30 (7.70-12.0)	13.0 (10.0-16.0)	728
	05-06	3.97 (3.46-4.55)	3.97 (3.31-4.64)	5.74 (5.11-7.10)	9.43 (7.63-11.6)	12.0 (10.7-13.9)	703
	07-08	3.60 (3.16-4.09)	3.66 (3.24-4.26)	5.53 (4.75-6.72)	8.90 (7.33-9.95)	10.7 (9.08-13.3)	366
	09-10	3.08 (2.77-3.43)	2.89 (2.64-3.29)	5.05 (4.44-5.77)	7.89 (6.87-10.1)	11.3 (8.75-13.8)	456
20 years and older	03-04	3.69 (3.31-4.11)	3.70 (3.00-4.00)	6.00 (5.00-7.00)	11.0 (10.0-12.0)	16.0 (13.0-19.0)	1548
	05-06	3.95 (3.58-4.36)	3.87 (3.46-4.32)	6.30 (5.59-7.28)	10.3 (9.25-13.3)	16.2 (12.5-19.4)	1531
	07-08	3.66 (3.43-3.91)	3.56 (3.41-3.80)	5.97 (5.44-6.63)	9.89 (8.72-11.4)	13.8 (11.8-15.5)	1820
	09-10	3.79 (3.51-4.09)	3.60 (3.33-3.87)	6.40 (5.71-7.03)	12.1 (9.52-16.0)	18.2 (15.8-23.6)	2035
Gender							
Males	03-04	4.12 (3.60-4.71)	4.00 (3.70-4.30)	6.00 (5.60-7.70)	11.0 (9.00-15.0)	17.0 (12.1-22.0)	1284
	05-06	4.17 (3.79-4.59)	4.04 (3.68-4.43)	6.32 (5.41-7.61)	10.5 (8.68-13.3)	14.8 (11.8-19.2)	1276
	07-08	4.05 (3.73-4.40)	3.96 (3.63-4.27)	6.53 (5.77-7.15)	10.3 (9.31-12.0)	14.6 (11.6-20.2)	1289
	09-10	3.89 (3.50-4.32)	3.71 (3.35-4.11)	6.33 (5.80-7.02)	11.6 (8.80-17.3)	19.2 (13.0-24.7)	1403
Females	03-04	3.37 (3.00-3.78)	3.00 (3.00-4.00)	5.50 (4.80-6.20)	10.0 (8.00-11.0)	14.0 (11.0-17.7)	1284
	05-06	3.75 (3.37-4.18)	3.70 (3.28-4.21)	6.03 (5.46-6.99)	9.95 (8.81-12.2)	15.5 (11.6-19.0)	1312
	07-08	3.34 (3.13-3.57)	3.37 (3.15-3.53)	5.48 (5.00-6.05)	8.99 (7.97-10.4)	11.9 (10.7-13.9)	1287
	09-10	3.48 (3.22-3.76)	3.28 (2.95-3.63)	5.85 (5.21-6.69)	11.5 (9.42-13.3)	16.1 (13.8-17.7)	1468
Race/ethnicity							
Mexican Americans	03-04	4.72 (4.27-5.22)	4.80 (4.00-5.00)	7.00 (6.00-9.00)	12.0 (10.0-16.0)	17.0 (12.0-25.0)	621
	05-06	4.43 (4.15-4.72)	4.66 (4.26-5.13)	6.93 (6.38-7.58)	9.51 (8.70-11.1)	13.1 (10.3-15.1)	651
	07-08	4.36 (4.05-4.69)	4.62 (4.07-5.07)	7.14 (6.23-7.76)	9.67 (8.39-10.8)	11.7 (9.96-17.6)	513
	09-10	3.90 (3.45-4.40)	3.98 (3.18-4.63)	6.53 (5.61-7.15)	10.7 (8.95-13.2)	16.2 (13.1-19.2)	618
Non-Hispanic blacks	03-04	4.27 (3.71-4.92)	4.00 (3.50-5.00)	7.00 (6.00-8.00)	11.6 (9.00-15.0)	16.0 (14.0-18.7)	725
	05-06	4.28 (3.68-4.98)	4.02 (3.37-4.82)	6.42 (5.31-7.36)	10.9 (8.22-13.1)	15.2 (11.5-27.1)	695
	07-08	4.19 (3.92-4.49)	3.98 (3.63-4.27)	6.69 (5.80-7.43)	11.2 (10.1-12.5)	14.7 (12.1-19.2)	586
	09-10	4.09 (3.63-4.61)	3.93 (3.31-4.36)	7.13 (5.99-8.63)	12.3 (10.1-17.0)	18.2 (13.7-27.8)	546
Non-Hispanic whites	03-04	3.27 (2.95-3.62)	3.00 (3.00-3.80)	5.00 (4.60-6.00)	9.00 (7.00-10.0)	12.0 (9.50-15.0)	1078
	05-06	3.69 (3.28-4.15)	3.60 (3.19-4.08)	5.66 (5.10-6.70)	9.62 (8.11-11.8)	13.9 (10.6-17.8)	1050
	07-08	3.23 (3.01-3.48)	3.34 (3.04-3.53)	5.14 (4.77-5.59)	8.27 (7.48-9.07)	10.5 (9.25-13.2)	1063
	09-10	3.21 (3.07-3.37)	3.12 (2.91-3.29)	5.32 (5.00-5.66)	8.65 (7.23-10.1)	12.0 (10.1-14.4)	1226

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 1.7, 1.7, 1.7, and 1.7 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Monomethylarsonic acid

Metabolite of Arsenic

Geometric mean and selected percentiles of urine concentrations (in µg As/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	1.20 (1.00-1.30)	1.90 (1.60-2.10)	2.40 (2.00-2.80)	2567
	05-06	*	< LOD	1.19 (1.05-1.34)	1.72 (1.51-1.94)	2.12 (1.79-2.70)	2588
	07-08	*	< LOD	1.11 (1.00-1.20)	1.61 (1.51-1.69)	2.09 (1.91-2.26)	2576
	09-10	*	< LOD	1.02 (.920-1.10)	1.64 (1.41-1.83)	2.01 (1.82-2.28)	2871
Age group							
6-11 years	03-04	*	< LOD	1.00 (<LOD-1.40)	1.80 (1.30-2.60)	2.30 (1.70-2.90)	292
	05-06	*	< LOD	1.03 (<LOD-1.37)	1.54 (1.37-2.12)	2.12 (1.51-2.73)	354
	07-08	*	< LOD	1.05 (<LOD-1.28)	1.60 (1.32-1.83)	2.04 (1.64-2.86)	390
	09-10	*	< LOD	< LOD	1.38 (1.00-1.84)	1.81 (1.26-2.13)	380
12-19 years	03-04	*	< LOD	1.50 (1.10-1.80)	2.20 (1.70-3.00)	2.90 (2.20-3.60)	728
	05-06	*	< LOD	1.31 (1.09-1.46)	1.99 (1.62-2.23)	2.41 (2.04-2.89)	703
	07-08	*	< LOD	1.26 (1.11-1.35)	1.83 (1.54-2.22)	2.31 (1.87-2.90)	366
	09-10	*	< LOD	1.10 (.910-1.25)	1.62 (1.33-1.92)	2.03 (1.74-2.30)	456
20 years and older	03-04	*	< LOD	1.20 (1.00-1.30)	1.80 (1.50-2.10)	2.30 (2.00-2.60)	1547
	05-06	*	< LOD	1.20 (1.05-1.33)	1.70 (1.48-1.90)	2.06 (1.74-2.75)	1531
	07-08	*	< LOD	1.08 (.970-1.17)	1.58 (1.44-1.71)	1.98 (1.82-2.21)	1820
	09-10	*	< LOD	1.02 (.930-1.11)	1.65 (1.42-1.86)	2.03 (1.81-2.35)	2035
Gender							
Males	03-04	*	< LOD	1.30 (1.10-1.60)	2.00 (1.80-2.40)	2.60 (2.10-3.00)	1283
	05-06	*	< LOD	1.31 (1.12-1.49)	1.79 (1.57-2.09)	2.23 (1.85-2.86)	1276
	07-08	*	< LOD	1.24 (1.13-1.35)	1.78 (1.63-1.94)	2.33 (1.97-2.71)	1289
	09-10	*	< LOD	1.07 (.970-1.23)	1.77 (1.47-2.01)	2.19 (1.83-2.68)	1403
Females	03-04	*	< LOD	1.00 (<LOD-1.20)	1.60 (1.30-1.90)	2.10 (1.70-2.60)	1284
	05-06	*	< LOD	1.08 (.920-1.24)	1.57 (1.36-1.90)	2.04 (1.67-2.50)	1312
	07-08	*	< LOD	.970 (<LOD-1.06)	1.42 (1.30-1.57)	1.81 (1.68-1.90)	1287
	09-10	*	< LOD	.950 (<LOD-1.06)	1.52 (1.27-1.74)	1.89 (1.66-2.12)	1468
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	1.50 (1.20-1.90)	2.20 (1.70-2.80)	2.80 (2.00-4.40)	621
	05-06	*	< LOD	1.19 (1.10-1.32)	1.73 (1.51-1.97)	2.04 (1.67-2.51)	651
	07-08	*	< LOD	1.22 (1.02-1.35)	1.69 (1.43-2.12)	2.32 (1.75-2.85)	513
	09-10	*	< LOD	1.02 (<LOD-1.37)	1.65 (1.38-1.90)	2.02 (1.65-2.33)	618
Non-Hispanic blacks	03-04	*	< LOD	1.10 (<LOD-1.30)	1.80 (1.40-2.00)	2.20 (1.70-2.70)	725
	05-06	*	< LOD	1.02 (.950-1.07)	1.35 (1.29-1.44)	1.70 (1.50-1.88)	695
	07-08	*	< LOD	1.21 (1.09-1.33)	1.79 (1.61-2.05)	2.32 (2.15-2.63)	586
	09-10	*	< LOD	1.02 (<LOD-1.18)	1.51 (1.36-1.84)	1.88 (1.73-2.24)	546
Non-Hispanic whites	03-04	*	< LOD	1.10 (.900-1.30)	1.80 (1.40-2.00)	2.10 (1.80-2.50)	1077
	05-06	*	< LOD	1.22 (1.02-1.39)	1.75 (1.42-2.12)	2.14 (1.72-2.93)	1050
	07-08	*	< LOD	1.04 (.930-1.15)	1.50 (1.37-1.63)	1.91 (1.69-2.09)	1063
	09-10	*	< LOD	.940 (<LOD-1.04)	1.45 (1.24-1.76)	1.90 (1.65-2.12)	1226

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.9, 0.9, 0.9, and 0.9 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Monomethylarsonic acid (creatinine corrected)

Metabolite of Arsenic

Geometric mean and selected percentiles of urine concentrations (in µg As/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	1.33 (1.18-1.54)	2.22 (1.82-2.57)	2.86 (2.40-3.53)	2567
	05-06	*	< LOD	1.28 (1.14-1.52)	2.37 (1.94-2.78)	3.13 (2.67-3.76)	2588
	07-08	*	< LOD	1.31 (1.23-1.42)	2.24 (2.00-2.56)	3.19 (2.78-3.58)	2576
	09-10	*	< LOD	1.36 (1.25-1.45)	2.16 (2.00-2.29)	2.91 (2.67-3.15)	2870
Age group							
6-11 years	03-04	*	< LOD	1.63 (<LOD-1.81)	2.31 (1.88-2.50)	2.52 (2.31-3.07)	292
	05-06	*	< LOD	1.44 (<LOD-1.83)	2.37 (1.65-3.20)	3.13 (2.29-4.92)	354
	07-08	*	< LOD	1.69 (<LOD-2.06)	2.46 (2.09-2.91)	3.20 (2.56-4.32)	390
	09-10	*	< LOD	< LOD	2.37 (2.07-2.78)	3.16 (2.55-3.76)	379
12-19 years	03-04	*	< LOD	1.10 (.853-1.23)	1.53 (1.30-1.85)	2.07 (1.71-2.22)	728
	05-06	*	< LOD	1.05 (.930-1.14)	1.56 (1.49-1.73)	1.97 (1.78-2.13)	703
	07-08	*	< LOD	.970 (.880-1.07)	1.45 (1.19-1.73)	1.97 (1.49-2.34)	366
	09-10	*	< LOD	1.21 (1.02-1.45)	1.98 (1.60-2.56)	2.56 (1.94-4.57)	456
20 years and older	03-04	*	< LOD	1.36 (1.18-1.58)	2.28 (1.82-2.79)	3.00 (2.43-3.53)	1547
	05-06	*	< LOD	1.31 (1.13-1.58)	2.46 (2.00-2.91)	3.20 (2.78-3.81)	1531
	07-08	*	< LOD	1.32 (1.23-1.45)	2.35 (2.02-2.67)	3.38 (2.78-3.83)	1820
	09-10	*	< LOD	1.36 (1.25-1.45)	2.18 (2.00-2.29)	2.91 (2.78-3.20)	2035
Gender							
Males	03-04	*	< LOD	1.20 (1.05-1.36)	1.88 (1.53-2.34)	2.50 (2.07-3.45)	1283
	05-06	*	< LOD	1.11 (.950-1.28)	1.78 (1.47-2.29)	2.46 (1.88-3.32)	1276
	07-08	*	< LOD	1.18 (1.07-1.26)	2.06 (1.85-2.34)	2.71 (2.46-3.37)	1289
	09-10	*	< LOD	1.16 (1.07-1.25)	1.79 (1.64-1.99)	2.29 (2.08-2.78)	1402
Females	03-04	*	< LOD	1.50 (<LOD-1.77)	2.40 (1.96-2.86)	3.00 (2.61-3.53)	1284
	05-06	*	< LOD	1.58 (1.36-1.76)	2.78 (2.29-3.20)	3.56 (3.05-4.35)	1312
	07-08	*	< LOD	1.45 (<LOD-1.60)	2.41 (1.94-2.91)	3.56 (2.78-4.27)	1287
	09-10	*	< LOD	1.56 (<LOD-1.73)	2.46 (2.29-2.62)	3.37 (2.91-3.56)	1468
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	1.46 (1.11-1.93)	2.30 (1.84-3.00)	3.16 (2.40-3.85)	621
	05-06	*	< LOD	1.19 (1.08-1.35)	1.92 (1.60-2.13)	2.53 (2.06-2.91)	651
	07-08	*	< LOD	1.28 (1.10-1.60)	2.27 (1.64-2.71)	3.05 (2.35-3.76)	513
	09-10	*	< LOD	1.31 (<LOD-1.49)	2.00 (1.73-2.56)	3.05 (2.29-3.56)	618
Non-Hispanic blacks	03-04	*	< LOD	.816 (<LOD-.985)	1.37 (1.14-1.61)	1.88 (1.46-2.17)	725
	05-06	*	< LOD	.790 (.720-.880)	1.11 (.980-1.34)	1.57 (1.23-1.98)	695
	07-08	*	< LOD	.920 (.840-1.06)	1.45 (1.33-1.73)	2.09 (1.73-2.56)	586
	09-10	*	< LOD	.970 (<LOD-1.08)	1.73 (1.39-1.94)	2.29 (1.83-2.91)	545
Non-Hispanic whites	03-04	*	< LOD	1.33 (1.15-1.62)	2.28 (1.73-2.86)	2.86 (2.35-3.75)	1077
	05-06	*	< LOD	1.35 (1.16-1.64)	2.50 (2.00-3.05)	3.20 (2.78-3.81)	1050
	07-08	*	< LOD	1.35 (1.23-1.52)	2.37 (2.04-2.78)	3.37 (2.78-4.00)	1063
	09-10	*	< LOD	1.36 (<LOD-1.52)	2.21 (1.99-2.37)	2.91 (2.67-3.20)	1226

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Trimethylarsine oxide

Geometric mean and selected percentiles of urine concentrations (in µg As/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2568
	05-06	*	< LOD	< LOD	< LOD	< LOD	2588
	07-08	*	< LOD	< LOD	< LOD	< LOD	2576
	09-10	*	< LOD	< LOD	< LOD	< LOD	2871
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	292
	05-06	*	< LOD	< LOD	< LOD	< LOD	354
	07-08	*	< LOD	< LOD	< LOD	< LOD	390
	09-10	*	< LOD	< LOD	< LOD	< LOD	380
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	728
	05-06	*	< LOD	< LOD	< LOD	< LOD	703
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
	09-10	*	< LOD	< LOD	< LOD	< LOD	456
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1548
	05-06	*	< LOD	< LOD	< LOD	< LOD	1531
	07-08	*	< LOD	< LOD	< LOD	< LOD	1820
	09-10	*	< LOD	< LOD	< LOD	< LOD	2035
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1276
	07-08	*	< LOD	< LOD	< LOD	< LOD	1289
	09-10	*	< LOD	< LOD	< LOD	< LOD	1403
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1312
	07-08	*	< LOD	< LOD	< LOD	< LOD	1287
	09-10	*	< LOD	< LOD	< LOD	< LOD	1468
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	621
	05-06	*	< LOD	< LOD	< LOD	< LOD	651
	07-08	*	< LOD	< LOD	< LOD	< LOD	513
	09-10	*	< LOD	< LOD	< LOD	< LOD	618
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	586
	09-10	*	< LOD	< LOD	< LOD	< LOD	546
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1078
	05-06	*	< LOD	< LOD	< LOD	< LOD	1050
	07-08	*	< LOD	< LOD	< LOD	< LOD	1063
	09-10	*	< LOD	< LOD	< LOD	< LOD	1226

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 1.0, 1.0, 1.0, and 1.0 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Trimethylarsine oxide (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg As/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2568
	05-06	*	< LOD	< LOD	< LOD	< LOD	2588
	07-08	*	< LOD	< LOD	< LOD	< LOD	2576
	09-10	*	< LOD	< LOD	< LOD	< LOD	2870
Age group							
6-11 years	03-04	*	< LOD	< LOD	< LOD	< LOD	292
	05-06	*	< LOD	< LOD	< LOD	< LOD	354
	07-08	*	< LOD	< LOD	< LOD	< LOD	390
	09-10	*	< LOD	< LOD	< LOD	< LOD	379
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	728
	05-06	*	< LOD	< LOD	< LOD	< LOD	703
	07-08	*	< LOD	< LOD	< LOD	< LOD	366
	09-10	*	< LOD	< LOD	< LOD	< LOD	456
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1548
	05-06	*	< LOD	< LOD	< LOD	< LOD	1531
	07-08	*	< LOD	< LOD	< LOD	< LOD	1820
	09-10	*	< LOD	< LOD	< LOD	< LOD	2035
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1276
	07-08	*	< LOD	< LOD	< LOD	< LOD	1289
	09-10	*	< LOD	< LOD	< LOD	< LOD	1402
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
	05-06	*	< LOD	< LOD	< LOD	< LOD	1312
	07-08	*	< LOD	< LOD	< LOD	< LOD	1287
	09-10	*	< LOD	< LOD	< LOD	< LOD	1468
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	621
	05-06	*	< LOD	< LOD	< LOD	< LOD	651
	07-08	*	< LOD	< LOD	< LOD	< LOD	513
	09-10	*	< LOD	< LOD	< LOD	< LOD	618
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	695
	07-08	*	< LOD	< LOD	< LOD	< LOD	586
	09-10	*	< LOD	< LOD	< LOD	< LOD	545
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	1078
	05-06	*	< LOD	< LOD	< LOD	< LOD	1050
	07-08	*	< LOD	< LOD	< LOD	< LOD	1063
	09-10	*	< LOD	< LOD	< LOD	< LOD	1226

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Arsenic_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html

Urinary Barium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	1.50 (1.35-1.66)	1.60 (1.50-1.90)	3.10 (2.70-3.40)	5.40 (4.60-6.10)	6.90 (6.20-8.40)	2180
	01-02	1.52 (1.41-1.65)	1.63 (1.50-1.76)	3.12 (2.77-3.51)	5.24 (4.73-5.84)	7.48 (6.54-8.12)	2690
	03-04	1.49 (1.36-1.64)	1.51 (1.35-1.72)	2.91 (2.64-3.28)	5.36 (4.86-5.71)	7.54 (6.93-8.63)	2558
	05-06	1.52 (1.39-1.66)	1.58 (1.44-1.70)	2.99 (2.63-3.41)	5.34 (4.82-5.83)	7.10 (6.46-8.18)	2576
	07-08	1.56 (1.45-1.69)	1.64 (1.49-1.80)	3.01 (2.72-3.34)	4.93 (4.39-5.69)	7.04 (6.07-8.20)	2627
	09-10	1.47 (1.34-1.60)	1.48 (1.35-1.62)	2.81 (2.52-3.06)	4.66 (4.34-5.35)	6.78 (5.86-7.76)	2848
Age group							
6-11 years	99-00	2.15 (1.70-2.72)	2.20 (1.90-2.50)	4.00 (2.60-6.10)	6.40 (5.20-8.30)	8.30 (5.00-76.2)	297
	01-02	1.80 (1.44-2.26)	2.09 (1.74-2.49)	3.63 (2.86-4.39)	5.37 (4.26-7.38)	6.88 (5.37-8.49)	368
	03-04	2.21 (1.81-2.71)	2.50 (1.81-2.75)	4.76 (3.80-5.65)	8.63 (5.59-11.8)	11.8 (6.87-14.8)	290
	05-06	1.73 (1.53-1.97)	1.87 (1.61-2.03)	3.55 (2.41-4.30)	6.21 (4.30-7.32)	7.72 (6.26-8.92)	355
	07-08	1.75 (1.50-2.04)	1.86 (1.58-2.23)	3.41 (2.85-3.79)	5.57 (4.49-6.59)	7.32 (5.71-9.18)	394
	09-10	1.49 (1.28-1.74)	1.50 (1.22-1.68)	2.72 (2.19-3.13)	4.63 (4.02-5.37)	6.59 (5.24-7.72)	378
12-19 years	99-00	1.97 (1.78-2.19)	2.00 (1.70-2.30)	3.50 (3.10-4.00)	5.90 (4.80-7.00)	9.70 (5.90-13.1)	621
	01-02	2.03 (1.76-2.34)	2.27 (1.96-2.53)	4.11 (3.48-4.72)	6.73 (5.55-7.87)	9.02 (7.25-11.4)	762
	03-04	2.16 (1.93-2.41)	2.35 (2.06-2.63)	4.11 (3.48-4.71)	7.18 (6.00-8.29)	9.63 (8.15-11.4)	725
	05-06	2.16 (1.92-2.42)	2.33 (1.99-2.56)	3.80 (3.60-4.21)	6.35 (5.48-7.49)	8.24 (6.06-10.5)	701
	07-08	2.18 (1.99-2.39)	2.31 (2.07-2.59)	3.62 (3.34-4.10)	6.19 (4.75-7.51)	8.20 (6.22-11.1)	376
	09-10	1.79 (1.52-2.09)	1.90 (1.46-2.45)	3.25 (2.82-4.14)	5.43 (4.45-6.07)	6.87 (5.73-8.16)	451
20 years and older	99-00	1.36 (1.24-1.51)	1.50 (1.30-1.70)	2.80 (2.60-3.20)	5.10 (4.30-5.50)	6.40 (5.70-8.40)	1262
	01-02	1.43 (1.32-1.54)	1.50 (1.39-1.65)	2.85 (2.55-3.27)	4.86 (4.53-5.47)	7.14 (6.08-8.12)	1560
	03-04	1.34 (1.20-1.50)	1.39 (1.19-1.56)	2.54 (2.21-2.91)	4.61 (3.99-5.30)	6.61 (5.57-7.43)	1543
	05-06	1.42 (1.28-1.57)	1.43 (1.28-1.63)	2.75 (2.47-3.11)	5.12 (4.45-5.81)	6.83 (6.14-7.90)	1520
	07-08	1.46 (1.35-1.58)	1.51 (1.39-1.67)	2.79 (2.54-3.15)	4.61 (4.19-5.42)	6.80 (5.50-8.37)	1857
	09-10	1.42 (1.30-1.55)	1.43 (1.31-1.58)	2.67 (2.36-3.02)	4.58 (4.22-5.14)	6.80 (5.70-8.63)	2019
Gender							
Males	99-00	1.70 (1.54-1.88)	1.90 (1.80-2.00)	3.20 (3.00-3.60)	5.50 (4.20-6.70)	7.50 (5.90-9.70)	1083
	01-02	1.64 (1.47-1.82)	1.80 (1.65-1.98)	3.15 (2.76-3.73)	5.52 (4.82-6.35)	7.87 (6.49-9.32)	1335
	03-04	1.62 (1.47-1.78)	1.69 (1.49-1.85)	3.09 (2.81-3.54)	5.65 (5.14-6.16)	8.56 (6.71-9.67)	1281
	05-06	1.66 (1.48-1.85)	1.71 (1.50-1.94)	3.21 (2.76-3.79)	5.48 (4.68-6.22)	7.10 (6.18-8.84)	1271
	07-08	1.62 (1.48-1.77)	1.80 (1.57-1.93)	3.14 (2.79-3.42)	5.00 (4.31-5.70)	6.84 (5.83-7.72)	1327
	09-10	1.53 (1.35-1.74)	1.57 (1.36-1.82)	3.01 (2.53-3.36)	4.83 (4.25-5.77)	6.81 (5.77-8.76)	1398
Females	99-00	1.33 (1.15-1.53)	1.50 (1.20-1.60)	2.80 (2.30-3.30)	5.20 (4.20-5.90)	6.80 (5.60-10.4)	1097
	01-02	1.43 (1.30-1.56)	1.44 (1.29-1.63)	3.11 (2.74-3.43)	4.93 (4.44-5.88)	7.15 (6.32-7.86)	1355
	03-04	1.38 (1.24-1.54)	1.39 (1.18-1.57)	2.71 (2.41-3.15)	4.95 (4.29-5.51)	6.87 (5.65-8.10)	1277
	05-06	1.40 (1.27-1.54)	1.42 (1.30-1.63)	2.70 (2.39-3.19)	5.21 (4.65-5.71)	7.09 (6.24-7.94)	1305
	07-08	1.51 (1.36-1.68)	1.54 (1.38-1.70)	2.84 (2.52-3.37)	4.87 (4.23-6.13)	7.18 (5.94-8.83)	1300
	09-10	1.40 (1.30-1.51)	1.38 (1.31-1.47)	2.61 (2.41-2.90)	4.58 (4.33-4.95)	6.59 (5.37-7.86)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.12, 0.12, 0.31, 0.12, 0.12, and 0.12 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Barium_BiomonitoringSummary.html

Urinary Barium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	1.35 (1.25-1.46)	1.40 (1.20-1.50)	2.60 (2.30-2.90)	4.50 (4.10-5.30)	6.30 (5.50-6.80)	692
	01-02	1.21 (1.06-1.37)	1.25 (1.11-1.46)	2.56 (2.04-2.91)	4.35 (3.65-5.49)	6.43 (5.21-8.22)	683
	03-04	1.40 (1.15-1.70)	1.45 (1.22-1.73)	2.62 (1.78-3.38)	4.07 (2.95-6.37)	6.01 (4.01-7.88)	618
	05-06	1.33 (1.24-1.42)	1.36 (1.22-1.50)	2.63 (2.35-2.89)	4.47 (3.80-5.23)	6.21 (5.23-7.00)	652
	07-08	1.33 (1.16-1.53)	1.37 (1.19-1.48)	2.60 (2.29-2.84)	4.21 (3.53-4.90)	5.79 (4.90-8.00)	515
	09-10	1.24 (1.07-1.45)	1.18 (1.05-1.40)	2.25 (1.88-2.71)	4.15 (3.55-5.40)	6.35 (5.00-8.75)	613
Non-Hispanic blacks	99-00	1.34 (1.12-1.62)	1.40 (1.20-1.60)	2.50 (2.30-2.90)	5.10 (3.70-6.60)	7.40 (5.40-13.9)	540
	01-02	1.30 (1.14-1.48)	1.42 (1.22-1.62)	2.61 (2.31-2.82)	4.30 (3.70-5.18)	5.99 (4.87-7.26)	667
	03-04	1.27 (1.17-1.39)	1.38 (1.26-1.48)	2.34 (2.05-2.59)	3.77 (3.35-4.36)	5.86 (4.76-7.45)	723
	05-06	1.25 (1.15-1.37)	1.31 (1.21-1.41)	2.40 (2.22-2.61)	4.24 (3.69-4.83)	6.19 (5.23-6.96)	692
	07-08	1.17 (1.07-1.28)	1.23 (1.12-1.37)	2.14 (2.01-2.34)	3.70 (3.27-4.00)	5.00 (3.93-5.90)	589
	09-10	1.22 (1.08-1.38)	1.22 (1.07-1.38)	2.13 (1.91-2.51)	3.76 (3.09-4.88)	5.50 (4.06-7.32)	544
Non-Hispanic whites	99-00	1.56 (1.36-1.80)	1.80 (1.60-2.00)	3.30 (2.80-3.70)	5.50 (4.50-6.30)	7.50 (6.20-8.80)	765
	01-02	1.61 (1.46-1.77)	1.68 (1.54-1.85)	3.31 (2.87-3.74)	5.66 (4.94-6.30)	7.73 (6.61-8.49)	1132
	03-04	1.56 (1.37-1.78)	1.61 (1.28-1.92)	3.12 (2.75-3.70)	5.57 (5.04-6.43)	8.08 (6.87-9.53)	1074
	05-06	1.59 (1.42-1.78)	1.66 (1.50-1.86)	3.08 (2.63-3.79)	5.50 (4.90-6.08)	7.10 (6.46-8.24)	1041
	07-08	1.71 (1.55-1.88)	1.82 (1.62-1.97)	3.36 (2.91-3.71)	5.38 (4.51-6.30)	7.18 (6.22-9.05)	1095
	09-10	1.56 (1.42-1.73)	1.60 (1.40-1.76)	3.00 (2.64-3.36)	4.89 (4.48-5.84)	7.16 (6.12-9.31)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.12, 0.12, 0.31, 0.12, 0.12, and 0.12 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Barium_BiomonitoringSummary.html

Urinary Barium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	1.21 (1.10-1.33)	1.18 (1.05-1.38)	2.39 (2.10-2.59)	4.00 (3.33-4.80)	5.31 (4.80-6.51)	692
	01-02	1.18 (1.03-1.34)	1.16 (1.00-1.38)	2.33 (1.90-2.61)	3.68 (3.29-4.10)	4.96 (4.24-6.80)	682
	03-04	1.29 (1.08-1.55)	1.28 (1.03-1.53)	2.25 (1.73-2.97)	3.99 (2.79-5.03)	4.99 (4.24-6.56)	618
	05-06	1.25 (1.17-1.33)	1.21 (1.09-1.33)	2.27 (1.84-2.61)	3.66 (3.07-4.35)	5.11 (4.00-5.95)	652
	07-08	1.33 (1.16-1.53)	1.28 (1.14-1.49)	2.40 (1.87-2.90)	4.23 (3.45-4.87)	6.18 (4.75-7.27)	515
	09-10	1.30 (1.18-1.44)	1.26 (1.14-1.41)	2.33 (1.98-2.79)	3.55 (3.11-4.42)	5.50 (4.27-6.58)	613
Non-Hispanic blacks	99-00	.881 (.703-1.11)	.905 (.710-1.06)	1.64 (1.36-2.00)	3.27 (2.26-4.76)	4.84 (3.57-10.8)	540
	01-02	.891 (.777-1.02)	.921 (.754-1.11)	1.64 (1.44-2.03)	2.86 (2.48-3.37)	4.02 (3.52-4.68)	667
	03-04	.915 (.832-1.01)	.963 (.880-1.04)	1.51 (1.39-1.75)	2.62 (2.29-3.04)	3.76 (3.22-4.72)	723
	05-06	.909 (.845-.979)	.880 (.820-.970)	1.78 (1.56-1.97)	2.99 (2.63-3.14)	3.91 (3.39-4.73)	692
	07-08	.858 (.807-.913)	.850 (.790-.920)	1.53 (1.34-1.61)	2.51 (2.01-3.03)	3.43 (2.83-5.05)	589
	09-10	.965 (.853-1.09)	.930 (.860-1.05)	1.63 (1.40-1.85)	2.92 (2.33-3.34)	3.71 (3.24-4.69)	544
Non-Hispanic whites	99-00	1.56 (1.38-1.77)	1.55 (1.36-1.74)	2.72 (2.27-3.24)	5.00 (3.81-6.02)	6.60 (5.52-10.2)	765
	01-02	1.62 (1.49-1.76)	1.66 (1.49-1.82)	3.04 (2.76-3.32)	4.96 (4.55-5.41)	6.74 (5.57-7.64)	1132
	03-04	1.64 (1.49-1.82)	1.60 (1.40-1.76)	2.88 (2.56-3.26)	5.38 (4.67-6.28)	7.57 (6.69-9.27)	1074
	05-06	1.65 (1.52-1.80)	1.61 (1.46-1.83)	2.95 (2.70-3.26)	4.82 (4.32-5.65)	7.21 (5.69-8.51)	1041
	07-08	1.91 (1.74-2.10)	1.85 (1.65-2.09)	3.22 (2.84-3.76)	5.42 (4.55-6.91)	7.81 (6.33-9.45)	1095
	09-10	1.75 (1.59-1.92)	1.70 (1.58-1.86)	2.99 (2.67-3.33)	5.00 (4.14-5.69)	6.63 (5.69-8.50)	1225

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Barium_BiomonitoringSummary.html

Urinary Beryllium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	2465
	01-02	*	< LOD	< LOD	< LOD	< LOD	2690
	03-04	*	< LOD	< LOD	< LOD	< LOD	2558
	05-06	*	< LOD	< LOD	< LOD	< LOD	2576
	07-08	*	< LOD	< LOD	< LOD	< LOD	2627
	09-10	*	< LOD	< LOD	< LOD	< LOD	2848
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	< LOD	340
	01-02	*	< LOD	< LOD	< LOD	< LOD	368
	03-04	*	< LOD	< LOD	< LOD	< LOD	290
	05-06	*	< LOD	< LOD	< LOD	< LOD	355
	07-08	*	< LOD	< LOD	< LOD	< LOD	394
	09-10	*	< LOD	< LOD	< LOD	< LOD	378
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	719
	01-02	*	< LOD	< LOD	< LOD	.140 (<LOD-.170)	762
	03-04	*	< LOD	< LOD	< LOD	< LOD	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	701
	07-08	*	< LOD	< LOD	< LOD	< LOD	376
	09-10	*	< LOD	< LOD	< LOD	< LOD	451
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1406
	01-02	*	< LOD	< LOD	< LOD	< LOD	1560
	03-04	*	< LOD	< LOD	< LOD	< LOD	1543
	05-06	*	< LOD	< LOD	< LOD	< LOD	1520
	07-08	*	< LOD	< LOD	< LOD	< LOD	1857
	09-10	*	< LOD	< LOD	< LOD	< LOD	2019
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	1227
	01-02	*	< LOD	< LOD	< LOD	.130 (<LOD-.150)	1335
	03-04	*	< LOD	< LOD	< LOD	< LOD	1281
	05-06	*	< LOD	< LOD	< LOD	< LOD	1271
	07-08	*	< LOD	< LOD	< LOD	< LOD	1327
	09-10	*	< LOD	< LOD	< LOD	< LOD	1398
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1238
	01-02	*	< LOD	< LOD	< LOD	< LOD	1355
	03-04	*	< LOD	< LOD	< LOD	< LOD	1277
	05-06	*	< LOD	< LOD	< LOD	< LOD	1305
	07-08	*	< LOD	< LOD	< LOD	< LOD	1300
	09-10	*	< LOD	< LOD	< LOD	< LOD	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.13, 0.13, 0.13, 0.072, 0.072, and 0.072 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Beryllium_BiomonitoringSummary.html

Urinary Beryllium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	884
	01-02	*	< LOD	< LOD	< LOD	.130 (<LOD-.160)	683
	03-04	*	< LOD	< LOD	< LOD	< LOD	618
	05-06	*	< LOD	< LOD	< LOD	< LOD	652
	07-08	*	< LOD	< LOD	< LOD	< LOD	515
	09-10	*	< LOD	< LOD	< LOD	< LOD	613
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	568
	01-02	*	< LOD	< LOD	< LOD	< LOD	667
	03-04	*	< LOD	< LOD	< LOD	< LOD	723
	05-06	*	< LOD	< LOD	< LOD	< LOD	692
	07-08	*	< LOD	< LOD	< LOD	< LOD	589
	09-10	*	< LOD	< LOD	< LOD	< LOD	544
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	822
	01-02	*	< LOD	< LOD	< LOD	< LOD	1132
	03-04	*	< LOD	< LOD	< LOD	< LOD	1074
	05-06	*	< LOD	< LOD	< LOD	< LOD	1041
	07-08	*	< LOD	< LOD	< LOD	< LOD	1095
	09-10	*	< LOD	< LOD	< LOD	< LOD	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.13, 0.13, 0.13, 0.13, 0.072, 0.072, and 0.072 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Beryllium_BiomonitoringSummary.html

Urinary Beryllium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	2465
	01-02	*	< LOD	< LOD	< LOD	< LOD	2689
	03-04	*	< LOD	< LOD	< LOD	< LOD	2558
	05-06	*	< LOD	< LOD	< LOD	< LOD	2576
	07-08	*	< LOD	< LOD	< LOD	< LOD	2627
	09-10	*	< LOD	< LOD	< LOD	< LOD	2848
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	< LOD	340
	01-02	*	< LOD	< LOD	< LOD	< LOD	368
	03-04	*	< LOD	< LOD	< LOD	< LOD	290
	05-06	*	< LOD	< LOD	< LOD	< LOD	355
	07-08	*	< LOD	< LOD	< LOD	< LOD	394
	09-10	*	< LOD	< LOD	< LOD	< LOD	378
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	719
	01-02	*	< LOD	< LOD	< LOD	.231 (<LOD-.273)	762
	03-04	*	< LOD	< LOD	< LOD	< LOD	725
	05-06	*	< LOD	< LOD	< LOD	< LOD	701
	07-08	*	< LOD	< LOD	< LOD	< LOD	376
	09-10	*	< LOD	< LOD	< LOD	< LOD	451
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1406
	01-02	*	< LOD	< LOD	< LOD	< LOD	1559
	03-04	*	< LOD	< LOD	< LOD	< LOD	1543
	05-06	*	< LOD	< LOD	< LOD	< LOD	1520
	07-08	*	< LOD	< LOD	< LOD	< LOD	1857
	09-10	*	< LOD	< LOD	< LOD	< LOD	2019
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	1227
	01-02	*	< LOD	< LOD	< LOD	.281 (<LOD-.333)	1334
	03-04	*	< LOD	< LOD	< LOD	< LOD	1281
	05-06	*	< LOD	< LOD	< LOD	< LOD	1271
	07-08	*	< LOD	< LOD	< LOD	< LOD	1327
	09-10	*	< LOD	< LOD	< LOD	< LOD	1398
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1238
	01-02	*	< LOD	< LOD	< LOD	< LOD	1355
	03-04	*	< LOD	< LOD	< LOD	< LOD	1277
	05-06	*	< LOD	< LOD	< LOD	< LOD	1305
	07-08	*	< LOD	< LOD	< LOD	< LOD	1300
	09-10	*	< LOD	< LOD	< LOD	< LOD	1450

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Beryllium_BiomonitoringSummary.html

Urinary Beryllium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	884
	01-02	*	< LOD	< LOD	< LOD	.346 (<LOD-.391)	682
	03-04	*	< LOD	< LOD	< LOD	< LOD	618
	05-06	*	< LOD	< LOD	< LOD	< LOD	652
	07-08	*	< LOD	< LOD	< LOD	< LOD	515
	09-10	*	< LOD	< LOD	< LOD	< LOD	613
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	568
	01-02	*	< LOD	< LOD	< LOD	< LOD	667
	03-04	*	< LOD	< LOD	< LOD	< LOD	723
	05-06	*	< LOD	< LOD	< LOD	< LOD	692
	07-08	*	< LOD	< LOD	< LOD	< LOD	589
	09-10	*	< LOD	< LOD	< LOD	< LOD	544
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	822
	01-02	*	< LOD	< LOD	< LOD	< LOD	1132
	03-04	*	< LOD	< LOD	< LOD	< LOD	1074
	05-06	*	< LOD	< LOD	< LOD	< LOD	1041
	07-08	*	< LOD	< LOD	< LOD	< LOD	1095
	09-10	*	< LOD	< LOD	< LOD	< LOD	1225

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Beryllium_BiomonitoringSummary.html

Blood Cadmium

Geometric mean and selected percentiles of blood concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.412 (.378-.449)	.400 (.300-.400)	.600 (.500-.700)	1.00 (1.00-1.10)	1.40 (1.30-1.40)	7970
	01-02	*	.300 (.300-.400)	.500 (.500-.600)	1.00 (.900-1.10)	1.40 (1.20-1.70)	8945
	03-04	.304 (.289-.320)	.300 (.300-.300)	.500 (.500-.600)	1.10 (1.00-1.20)	1.60 (1.50-1.60)	8372
	05-06	.310 (.294-.327)	.270 (.250-.280)	.510 (.460-.560)	1.02 (.910-1.13)	1.53 (1.34-1.75)	8407
	07-08	.315 (.300-.331)	.270 (.260-.280)	.500 (.460-.560)	1.00 (.900-1.13)	1.52 (1.30-1.77)	8266
	09-10	.302 (.293-.311)	.260 (.250-.270)	.480 (.460-.510)	.960 (.880-1.01)	1.40 (1.29-1.53)	8793
Age group							
1-5 years	99-00	*	< LOD	.300 (<LOD-.400)	.400 (.300-.400)	.400 (.300-.700)	723
	01-02	*	< LOD	< LOD	< LOD	.300 (.300-.300)	898
	03-04	*	< LOD	< LOD	.200 (.200-.300)	.200 (.200-.400)	910
	05-06	*	< LOD	< LOD	< LOD	.230 (.210-.250)	968
	07-08	*	< LOD	< LOD	.210 (<LOD-.230)	.240 (.220-.260)	817
	09-10	*	< LOD	< LOD	.200 (<LOD-.210)	.220 (.200-.240)	836
6-11 years	99-00	*	< LOD	.300 (<LOD-.400)	.400 (.300-.500)	.500 (.400-.500)	905
	01-02	*	< LOD	< LOD	.300 (.300-.300)	.400 (.400-.400)	1044
	03-04	*	< LOD	.200 (<LOD-.200)	.300 (.200-.300)	.300 (.300-.300)	856
	05-06	*	< LOD	< LOD	.220 (.200-.240)	.260 (.230-.280)	934
	07-08	*	< LOD	< LOD	.230 (.210-.240)	.260 (.240-.280)	1011
	09-10	*	< LOD	< LOD	.210 (.200-.240)	.240 (.220-.280)	1009
12-19 years	99-00	.333 (.304-.366)	.300 (.300-.400)	.400 (.400-.500)	.800 (.600-1.00)	1.10 (.900-1.20)	2135
	01-02	*	< LOD	.300 (.300-.400)	.500 (.400-.700)	.900 (.700-1.20)	2231
	03-04	*	.200 (<LOD-.200)	.300 (.300-.300)	.600 (.500-.700)	.900 (.800-1.10)	2081
	05-06	*	< LOD	.250 (.240-.270)	.520 (.410-.680)	.960 (.820-1.08)	1996
	07-08	*	< LOD	.260 (.240-.270)	.520 (.400-.670)	.900 (.730-1.19)	1074
	09-10	*	< LOD	.240 (.220-.260)	.340 (.310-.450)	.620 (.460-.850)	1183
20 years and older	99-00	.468 (.426-.513)	.400 (.400-.500)	.700 (.600-.800)	1.10 (1.00-1.20)	1.50 (1.40-1.60)	4207
	01-02	.425 (.400-.452)	.400 (.400-.400)	.600 (.600-.700)	1.10 (1.00-1.30)	1.60 (1.40-1.90)	4772
	03-04	.378 (.359-.398)	.400 (.300-.400)	.600 (.600-.700)	1.20 (1.20-1.30)	1.80 (1.60-1.90)	4525
	05-06	.373 (.352-.395)	.330 (.310-.350)	.610 (.570-.660)	1.17 (1.06-1.26)	1.72 (1.53-1.95)	4509
	07-08	.376 (.354-.399)	.330 (.310-.350)	.600 (.550-.670)	1.16 (1.02-1.30)	1.70 (1.50-1.96)	5364
	09-10	.358 (.347-.370)	.320 (.310-.330)	.580 (.550-.600)	1.10 (1.00-1.18)	1.55 (1.43-1.72)	5765
Gender							
Males	99-00	.403 (.368-.441)	.400 (.300-.400)	.600 (.500-.700)	1.00 (.900-1.10)	1.40 (1.30-1.50)	3913
	01-02	*	.300 (<LOD-.300)	.500 (.500-.600)	1.00 (.900-1.10)	1.50 (1.20-1.80)	4339
	03-04	.283 (.266-.300)	.300 (.200-.300)	.500 (.500-.500)	1.10 (1.00-1.20)	1.60 (1.50-1.60)	4131
	05-06	*	.240 (.220-.260)	.470 (.420-.530)	1.02 (.910-1.12)	1.53 (1.27-1.86)	4092
	07-08	.299 (.283-.317)	.250 (.230-.260)	.470 (.420-.540)	1.05 (.930-1.19)	1.60 (1.30-1.90)	4147
	09-10	.279 (.268-.289)	.230 (.220-.240)	.410 (.390-.460)	.920 (.830-1.00)	1.45 (1.30-1.58)	4366
Females	99-00	.421 (.386-.460)	.400 (.400-.400)	.600 (.500-.700)	1.00 (1.00-1.10)	1.30 (1.20-1.40)	4057
	01-02	.382 (.362-.403)	.400 (.300-.400)	.600 (.500-.600)	1.00 (.900-1.10)	1.40 (1.20-1.70)	4606
	03-04	.326 (.309-.344)	.300 (.300-.300)	.600 (.500-.600)	1.10 (1.00-1.20)	1.60 (1.50-1.70)	4241
	05-06	.329 (.311-.349)	.290 (.280-.310)	.530 (.480-.580)	1.02 (.870-1.18)	1.54 (1.33-1.79)	4315
	07-08	.331 (.316-.348)	.290 (.280-.310)	.530 (.480-.570)	.980 (.860-1.10)	1.43 (1.29-1.63)	4119
	09-10	.326 (.314-.339)	.290 (.280-.310)	.520 (.490-.550)	.960 (.880-1.10)	1.40 (1.20-1.53)	4427

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.3, 0.3, 0.14, 0.2, 0.2, and 0.2 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cadmium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Cadmium_FactSheet.html

Blood Cadmium

Geometric mean and selected percentiles of blood concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.395 (.367-.424)	.400 (.400-.400)	.500 (.500-.600)	.800 (.700-1.00)	1.20 (.900-1.30)	2742
	01-02	*	< LOD	.400 (.400-.500)	.600 (.500-.800)	1.00 (.700-1.30)	2268
	03-04	.235 (.216-.255)	.200 (.200-.300)	.400 (.300-.400)	.600 (.500-.800)	1.00 (.800-1.50)	2085
	05-06	*	.220 (.200-.240)	.350 (.300-.400)	.580 (.510-.680)	.820 (.710-1.00)	2236
	07-08	*	.220 (.210-.230)	.350 (.320-.370)	.570 (.510-.660)	.840 (.690-1.02)	1712
	09-10	*	.220 (.200-.230)	.330 (.300-.370)	.580 (.500-.670)	.800 (.690-.960)	1966
	Non-Hispanic blacks	99-00	.393 (.361-.427)	.400 (.300-.400)	.600 (.500-.600)	1.00 (.900-1.20)	1.40 (1.20-1.60)
01-02		*	.300 (<LOD-.300)	.500 (.500-.600)	1.00 (.900-1.10)	1.40 (1.20-1.70)	2219
03-04		.304 (.275-.337)	.300 (.300-.300)	.500 (.400-.600)	1.00 (.900-1.20)	1.50 (1.30-1.70)	2292
05-06		.307 (.290-.326)	.260 (.250-.280)	.490 (.440-.570)	1.03 (.880-1.21)	1.50 (1.23-1.79)	2193
07-08		.333 (.316-.352)	.280 (.270-.300)	.550 (.480-.620)	1.20 (1.01-1.36)	1.81 (1.45-2.13)	1746
09-10		.328 (.310-.347)	.280 (.260-.290)	.550 (.490-.640)	1.10 (1.00-1.20)	1.52 (1.38-1.73)	1593
Non-Hispanic whites		99-00	.420 (.376-.470)	.400 (.300-.500)	.600 (.500-.700)	1.10 (1.00-1.20)	1.40 (1.30-1.50)
	01-02	*	.300 (.300-.400)	.600 (.500-.600)	1.00 (.900-1.20)	1.50 (1.30-1.80)	3806
	03-04	.313 (.296-.331)	.300 (.300-.300)	.600 (.500-.600)	1.10 (1.00-1.20)	1.60 (1.50-1.70)	3478
	05-06	.321 (.300-.343)	.270 (.250-.300)	.540 (.470-.610)	1.08 (.930-1.23)	1.64 (1.40-1.94)	3310
	07-08	.321 (.303-.341)	.270 (.260-.290)	.520 (.470-.580)	1.05 (.920-1.20)	1.55 (1.30-1.80)	3461
	09-10	.307 (.297-.318)	.270 (.260-.280)	.490 (.460-.540)	.990 (.900-1.10)	1.50 (1.36-1.60)	3760

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.3, 0.3, 0.14, 0.2, 0.2, and 0.2 respectively. < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cadmium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Cadmium_FactSheet.html

Urinary Cadmium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.193 (.169-.220)	.232 (.214-.249)	.475 (.436-.519)	.858 (.763-.980)	1.20 (1.06-1.34)	2257
	01-02	.210 (.189-.235)	.230 (.207-.255)	.458 (.423-.482)	.839 (.753-.919)	1.20 (1.07-1.28)	2690
	03-04	.211 (.196-.226)	.210 (.200-.230)	.450 (.400-.500)	.800 (.730-.880)	1.15 (.980-1.26)	2543
	05-06	.191 (.170-.216)	.200 (.170-.220)	.400 (.360-.460)	.780 (.700-.860)	1.05 (.960-1.17)	2576
	07-08	.185 (.172-.198)	.180 (.170-.200)	.380 (.360-.400)	.700 (.650-.770)	1.00 (.920-1.12)	2627
	09-10	.179 (.169-.190)	.180 (.170-.200)	.370 (.340-.400)	.690 (.630-.770)	1.03 (.910-1.13)	2848
Age group							
6-11 years	99-00	*	.078 (.061-.101)	.141 (.115-.173)	.219 (.178-.233)	.279 (.211-.507)	310
	01-02	.061 (<LOD-.081)	.077 (.067-.092)	.140 (.112-.160)	.219 (.184-.262)	.282 (.260-.326)	368
	03-04	.077 (.065-.090)	.080 (.060-.090)	.120 (.100-.160)	.180 (.160-.310)	.310 (.170-.610)	287
	05-06	.066 (.056-.078)	.060 (.050-.080)	.110 (.090-.130)	.180 (.130-.240)	.240 (.160-.290)	355
	07-08	.064 (.058-.071)	.060 (.050-.070)	.110 (.090-.130)	.180 (.140-.210)	.230 (.180-.310)	394
	09-10	.057 (.053-.061)	.060 (.050-.060)	.090 (.080-.100)	.130 (.120-.160)	.170 (.140-.230)	378
12-19 years	99-00	.092 (.067-.126)	.128 (.107-.148)	.203 (.183-.232)	.329 (.272-.372)	.426 (.366-.596)	648
	01-02	.109 (.087-.136)	.135 (.114-.157)	.210 (.189-.247)	.327 (.289-.366)	.452 (.366-.480)	762
	03-04	.121 (.109-.134)	.130 (.110-.150)	.200 (.170-.210)	.300 (.260-.360)	.390 (.330-.490)	724
	05-06	.099 (.090-.109)	.110 (.100-.120)	.170 (.150-.190)	.240 (.210-.280)	.310 (.250-.430)	701
	07-08	.089 (.080-.100)	.080 (.070-.110)	.160 (.140-.170)	.260 (.200-.300)	.330 (.280-.410)	376
	09-10	.081 (.076-.086)	.080 (.070-.080)	.140 (.120-.150)	.220 (.180-.250)	.280 (.240-.300)	451
20 years and older	99-00	.281 (.253-.313)	.306 (.261-.339)	.551 (.510-.623)	.980 (.836-1.13)	1.32 (1.13-1.57)	1299
	01-02	.273 (.249-.299)	.280 (.261-.308)	.545 (.493-.607)	.972 (.855-1.06)	1.28 (1.20-1.43)	1560
	03-04	.260 (.238-.284)	.270 (.240-.300)	.530 (.470-.580)	.890 (.800-.990)	1.25 (1.09-1.46)	1532
	05-06	.241 (.213-.272)	.250 (.220-.290)	.490 (.440-.560)	.860 (.790-.930)	1.21 (1.06-1.32)	1520
	07-08	.232 (.215-.251)	.240 (.210-.260)	.450 (.410-.490)	.790 (.730-.870)	1.13 (.990-1.44)	1857
	09-10	.229 (.213-.245)	.230 (.210-.260)	.430 (.400-.470)	.790 (.700-.910)	1.13 (1.03-1.28)	2019
Gender							
Males	99-00	.199 (.165-.241)	.227 (.193-.263)	.462 (.381-.539)	.892 (.748-1.15)	1.41 (.980-1.83)	1121
	01-02	.201 (.177-.229)	.223 (.191-.257)	.445 (.393-.481)	.875 (.741-1.03)	1.22 (1.12-1.38)	1335
	03-04	.206 (.190-.222)	.210 (.190-.230)	.440 (.390-.490)	.790 (.700-.870)	1.01 (.890-1.25)	1277
	05-06	.195 (.176-.217)	.210 (.190-.230)	.400 (.360-.440)	.800 (.730-.890)	1.17 (1.01-1.30)	1271
	07-08	.179 (.162-.197)	.180 (.160-.200)	.360 (.320-.400)	.670 (.540-.780)	.950 (.800-1.16)	1327
	09-10	.170 (.157-.185)	.180 (.150-.200)	.350 (.310-.380)	.650 (.560-.700)	.950 (.770-1.14)	1398
Females	99-00	.187 (.153-.229)	.239 (.220-.255)	.492 (.456-.540)	.818 (.705-.980)	1.10 (1.01-1.19)	1136
	01-02	.219 (.192-.251)	.234 (.202-.265)	.466 (.433-.519)	.817 (.733-.886)	1.17 (.918-1.36)	1355
	03-04	.216 (.195-.238)	.210 (.200-.240)	.450 (.400-.530)	.820 (.700-.960)	1.20 (1.02-1.37)	1266
	05-06	.188 (.160-.221)	.190 (.160-.220)	.400 (.350-.480)	.750 (.640-.860)	.980 (.830-1.20)	1305
	07-08	.191 (.177-.207)	.190 (.170-.200)	.400 (.370-.430)	.740 (.660-.840)	1.09 (.940-1.38)	1300
	09-10	.188 (.172-.206)	.200 (.170-.210)	.400 (.360-.440)	.740 (.620-.880)	1.07 (.910-1.31)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.06, 0.06, 0.06, 0.042, 0.042, and 0.042 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**All results are corrected for molybdenum oxide interference in the ICP-MS method.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cadmium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Cadmium_FactSheet.html

Urinary Cadmium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	.191 (.157-.233)	.202 (.167-.221)	.447 (.351-.551)	.813 (.686-.977)	1.12 (.886-1.38)	780
	01-02	.160 (.135-.189)	.181 (.171-.198)	.322 (.285-.362)	.559 (.430-.733)	.766 (.633-1.15)	683
	03-04	.175 (.151-.203)	.170 (.150-.210)	.350 (.290-.430)	.680 (.520-.820)	1.04 (.820-1.20)	614
	05-06	.173 (.152-.196)	.180 (.160-.200)	.340 (.300-.380)	.560 (.500-.630)	.780 (.660-.900)	652
	07-08	.159 (.139-.181)	.170 (.140-.200)	.320 (.290-.370)	.570 (.510-.620)	.710 (.630-.830)	515
	09-10	.153 (.136-.173)	.150 (.130-.170)	.300 (.240-.360)	.530 (.440-.680)	.790 (.610-.840)	613
	Non-Hispanic blacks	99-00	.283 (.208-.387)	.316 (.243-.412)	.633 (.498-.806)	1.22 (.892-1.38)	1.48 (1.30-1.72)
01-02		.277 (.229-.336)	.302 (.257-.354)	.589 (.476-.713)	1.04 (.843-1.38)	1.51 (1.28-1.74)	667
03-04		.265 (.237-.295)	.270 (.220-.320)	.550 (.440-.640)	.960 (.810-1.17)	1.52 (1.06-1.82)	717
05-06		.236 (.218-.254)	.240 (.210-.260)	.480 (.420-.530)	.830 (.670-.930)	1.04 (.870-1.26)	692
07-08		.246 (.218-.277)	.260 (.220-.300)	.460 (.420-.530)	.840 (.690-.980)	1.40 (.900-1.85)	589
09-10		.234 (.211-.260)	.240 (.220-.270)	.450 (.410-.540)	.910 (.750-1.06)	1.42 (.990-1.74)	544
Non-Hispanic whites		99-00	.175 (.148-.206)	.220 (.194-.246)	.455 (.388-.510)	.848 (.714-1.01)	1.17 (.963-1.47)
	01-02	.204 (.179-.231)	.221 (.191-.255)	.445 (.394-.479)	.817 (.717-.875)	1.17 (.989-1.24)	1132
	03-04	.209 (.192-.226)	.200 (.190-.220)	.440 (.390-.500)	.790 (.700-.860)	1.13 (.940-1.26)	1070
	05-06	.185 (.159-.216)	.200 (.160-.230)	.400 (.330-.480)	.780 (.670-.920)	1.05 (.940-1.25)	1041
	07-08	.177 (.161-.195)	.170 (.150-.190)	.370 (.320-.400)	.690 (.620-.780)	1.00 (.880-1.13)	1095
	09-10	.171 (.158-.186)	.180 (.150-.200)	.350 (.320-.390)	.680 (.590-.780)	1.05 (.880-1.18)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.06, 0.06, 0.06, 0.042, 0.042, and 0.042 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result

**All results are corrected for molybdenum oxide interference in the ICP-MS method.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cadmium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Cadmium_FactSheet.html

Urinary Cadmium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.181 (.157-.209)	.219 (.199-.238)	.423 (.391-.446)	.712 (.645-.757)	.941 (.826-1.07)	2257
	01-02	.199 (.181-.218)	.212 (.194-.232)	.404 (.377-.440)	.690 (.630-.754)	.919 (.813-.998)	2689
	03-04	.210 (.201-.219)	.208 (.189-.226)	.412 (.381-.438)	.678 (.650-.716)	.940 (.833-1.04)	2543
	05-06	.189 (.169-.210)	.180 (.160-.200)	.370 (.310-.430)	.650 (.590-.720)	.910 (.770-1.08)	2576
	07-08	.193 (.177-.210)	.190 (.180-.210)	.370 (.330-.410)	.660 (.580-.740)	.960 (.840-1.06)	2627
	09-10	.191 (.184-.199)	.180 (.170-.190)	.370 (.340-.400)	.680 (.630-.730)	.910 (.840-1.00)	2848
Age group							
6-11 years	99-00	*	.085 (.063-.107)	.148 (.123-.182)	.210 (.171-.316)	.316 (.184-.607)	310
	01-02	.075 (<LOD-.094)	.100 (.083-.112)	.166 (.136-.192)	.233 (.206-.281)	.318 (.221-.440)	368
	03-04	.090 (.078-.104)	.091 (.075-.104)	.126 (.111-.156)	.200 (.147-.350)	.308 (.178-.415)	287
	05-06	.081 (.072-.092)	.080 (.070-.090)	.130 (.110-.140)	.170 (.150-.190)	.200 (.180-.240)	355
	07-08	.084 (.076-.092)	.080 (.080-.090)	.120 (.110-.140)	.170 (.150-.240)	.260 (.180-.430)	394
	09-10	.077 (.070-.085)	.080 (.070-.090)	.110 (.100-.130)	.160 (.130-.220)	.230 (.140-.280)	378
12-19 years	99-00	.071 (.051-.098)	.093 (.084-.106)	.147 (.130-.163)	.215 (.204-.240)	.283 (.222-.404)	648
	01-02	.078 (.067-.091)	.091 (.085-.101)	.137 (.123-.143)	.191 (.175-.234)	.280 (.234-.321)	762
	03-04	.086 (.077-.096)	.084 (.074-.097)	.122 (.113-.135)	.176 (.154-.198)	.234 (.187-.274)	724
	05-06	.076 (.071-.081)	.080 (.070-.090)	.120 (.110-.130)	.150 (.140-.180)	.210 (.160-.240)	701
	07-08	.070 (.062-.079)	.070 (.070-.080)	.110 (.100-.110)	.150 (.140-.160)	.180 (.160-.210)	376
	09-10	.076 (.068-.084)	.080 (.070-.090)	.120 (.100-.130)	.160 (.140-.180)	.190 (.170-.270)	451
20 years and older	99-00	.267 (.247-.289)	.288 (.261-.304)	.484 (.433-.545)	.769 (.727-.818)	1.07 (.927-1.17)	1299
	01-02	.261 (.236-.289)	.273 (.247-.303)	.481 (.426-.518)	.779 (.691-.850)	.979 (.874-1.12)	1559
	03-04	.268 (.255-.281)	.270 (.247-.292)	.490 (.444-.538)	.767 (.688-.830)	1.02 (.909-1.14)	1532
	05-06	.240 (.216-.267)	.230 (.200-.260)	.440 (.400-.500)	.730 (.660-.820)	1.02 (.850-1.18)	1520
	07-08	.247 (.227-.270)	.240 (.220-.270)	.430 (.390-.470)	.740 (.670-.840)	1.05 (.930-1.16)	1857
	09-10	.242 (.232-.253)	.230 (.220-.250)	.440 (.420-.460)	.760 (.700-.820)	1.01 (.910-1.14)	2019
Gender							
Males	99-00	.154 (.131-.182)	.176 (.158-.191)	.329 (.293-.382)	.617 (.537-.700)	.789 (.696-.929)	1121
	01-02	.159 (.143-.177)	.168 (.157-.182)	.335 (.304-.364)	.536 (.491-.653)	.757 (.690-.856)	1334
	03-04	.173 (.161-.187)	.162 (.143-.185)	.325 (.300-.352)	.591 (.560-.631)	.740 (.678-.795)	1277
	05-06	.160 (.145-.177)	.150 (.130-.160)	.300 (.260-.340)	.620 (.560-.670)	.820 (.710-.990)	1271
	07-08	.160 (.146-.175)	.160 (.140-.180)	.290 (.260-.330)	.520 (.440-.650)	.740 (.600-.960)	1327
	09-10	.155 (.149-.161)	.140 (.140-.160)	.280 (.250-.300)	.540 (.510-.600)	.800 (.670-.900)	1398
Females	99-00	.211 (.170-.261)	.267 (.239-.308)	.473 (.423-.551)	.783 (.690-.917)	1.09 (.813-1.38)	1136
	01-02	.245 (.216-.278)	.263 (.228-.297)	.479 (.414-.541)	.792 (.687-.884)	.985 (.876-1.16)	1355
	03-04	.252 (.238-.266)	.253 (.227-.288)	.487 (.438-.533)	.802 (.716-.906)	1.06 (.940-1.21)	1266
	05-06	.220 (.193-.252)	.220 (.180-.250)	.420 (.370-.470)	.690 (.590-.850)	.990 (.780-1.31)	1305
	07-08	.231 (.210-.254)	.230 (.200-.270)	.440 (.390-.480)	.770 (.700-.860)	1.08 (.970-1.16)	1300
	09-10	.233 (.218-.248)	.240 (.220-.260)	.450 (.420-.490)	.770 (.680-.860)	1.09 (.910-1.19)	1450

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**All results are corrected for molybdenum oxide interference in the ICP-MS method.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cadmium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Cadmium_FactSheet.html

Urinary Cadmium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	.175 (.137-.223)	.181 (.144-.225)	.331 (.266-.418)	.622 (.441-.828)	.856 (.674-1.13)	780
	01-02	.156 (.136-.178)	.170 (.150-.184)	.282 (.263-.340)	.501 (.388-.614)	.693 (.507-.839)	682
	03-04	.163 (.147-.181)	.159 (.140-.183)	.296 (.256-.311)	.531 (.418-.667)	.718 (.562-.950)	614
	05-06	.162 (.146-.181)	.150 (.140-.170)	.280 (.220-.340)	.480 (.430-.510)	.570 (.520-.640)	652
	07-08	.159 (.141-.178)	.160 (.130-.180)	.270 (.220-.330)	.480 (.370-.620)	.720 (.520-.950)	515
	09-10	.161 (.149-.173)	.150 (.130-.170)	.280 (.240-.320)	.510 (.410-.630)	.720 (.540-.890)	613
	Non-Hispanic blacks	99-00	.183 (.140-.240)	.202 (.168-.241)	.414 (.343-.472)	.663 (.516-.827)	.873 (.722-.962)
01-02		.190 (.156-.232)	.196 (.174-.225)	.387 (.336-.449)	.686 (.559-.850)	.917 (.725-1.08)	667
03-04		.190 (.173-.210)	.185 (.168-.207)	.338 (.288-.431)	.700 (.500-.818)	.865 (.708-1.10)	717
05-06		.171 (.159-.183)	.160 (.140-.180)	.320 (.290-.370)	.550 (.500-.580)	.700 (.610-.730)	692
07-08		.180 (.164-.197)	.170 (.160-.190)	.330 (.290-.390)	.600 (.470-.700)	.770 (.690-.900)	589
09-10		.186 (.170-.203)	.180 (.160-.200)	.360 (.310-.430)	.680 (.570-.760)	.860 (.740-1.04)	544
Non-Hispanic whites		99-00	.175 (.146-.209)	.219 (.191-.250)	.432 (.387-.470)	.729 (.666-.783)	1.00 (.826-1.16)
	01-02	.205 (.184-.229)	.224 (.208-.242)	.421 (.382-.470)	.719 (.668-.784)	.931 (.806-1.05)	1132
	03-04	.220 (.207-.235)	.221 (.197-.253)	.434 (.398-.476)	.687 (.647-.767)	1.00 (.830-1.08)	1070
	05-06	.193 (.169-.221)	.180 (.160-.220)	.390 (.310-.480)	.680 (.630-.750)	.930 (.800-1.07)	1041
	07-08	.199 (.178-.222)	.200 (.180-.230)	.380 (.340-.440)	.700 (.630-.810)	1.03 (.880-1.13)	1095
	09-10	.192 (.183-.201)	.180 (.170-.200)	.390 (.340-.420)	.700 (.630-.760)	.910 (.840-1.02)	1225

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**All results are corrected for molybdenum oxide interference in the ICP-MS method.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cadmium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Cadmium_FactSheet.html

Urinary Cesium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	4.35 (4.00-4.74)	4.90 (4.40-5.40)	7.10 (6.60-7.70)	9.60 (8.80-10.3)	11.5 (10.2-13.0)	2464
	01-02	4.81 (4.40-5.26)	5.49 (5.12-5.90)	7.91 (7.47-8.39)	10.4 (9.56-11.4)	12.6 (11.1-13.8)	2690
	03-04	4.67 (4.39-4.97)	5.14 (4.84-5.49)	7.68 (7.20-8.21)	10.6 (9.55-11.4)	12.7 (11.5-13.9)	2558
	05-06	4.69 (4.41-4.98)	5.19 (4.91-5.43)	7.68 (7.21-8.09)	10.1 (9.65-10.9)	12.6 (12.0-13.4)	2576
	07-08	4.42 (4.25-4.60)	4.90 (4.64-5.12)	7.02 (6.79-7.38)	9.46 (9.04-9.86)	11.3 (10.5-11.8)	2627
	09-10	4.06 (3.87-4.26)	4.47 (4.22-4.66)	6.54 (6.35-6.75)	8.98 (8.25-9.68)	11.0 (10.1-11.6)	2848
Age group							
6-11 years	99-00	4.87 (4.08-5.81)	5.70 (4.60-6.70)	7.30 (6.70-8.00)	9.00 (7.90-10.1)	9.70 (9.00-10.8)	340
	01-02	4.87 (4.08-5.82)	5.64 (4.69-6.56)	7.96 (6.77-8.84)	9.88 (8.64-10.6)	11.1 (10.2-12.4)	368
	03-04	5.21 (4.74-5.71)	5.50 (4.76-6.37)	8.08 (7.10-8.83)	11.5 (8.86-12.9)	12.9 (10.8-13.6)	290
	05-06	4.74 (4.31-5.21)	5.38 (4.67-5.60)	7.00 (6.38-7.50)	9.30 (8.60-9.95)	11.2 (9.62-12.9)	355
	07-08	4.58 (4.22-4.96)	5.10 (4.50-5.44)	6.91 (6.45-7.47)	8.25 (7.77-8.93)	9.88 (8.47-10.9)	394
	09-10	4.21 (3.84-4.62)	4.58 (4.01-5.12)	6.68 (6.25-6.98)	8.28 (7.47-9.11)	9.56 (8.68-11.9)	378
12-19 years	99-00	4.55 (4.09-5.05)	5.20 (4.40-5.60)	6.90 (6.10-7.80)	8.80 (8.10-9.50)	10.7 (8.90-12.5)	718
	01-02	5.22 (4.57-5.95)	5.62 (5.16-6.12)	7.55 (7.13-8.04)	9.77 (9.12-11.1)	12.0 (10.0-15.0)	762
	03-04	5.04 (4.59-5.54)	5.70 (5.16-6.07)	7.53 (6.91-8.37)	9.71 (8.80-10.4)	11.6 (9.92-13.2)	725
	05-06	4.76 (4.24-5.34)	5.27 (4.80-5.74)	7.50 (6.71-8.04)	9.60 (8.52-10.5)	10.6 (9.68-11.8)	701
	07-08	4.58 (4.10-5.13)	5.14 (4.30-5.76)	6.93 (6.33-7.37)	9.20 (7.67-9.91)	10.6 (8.91-15.1)	376
	09-10	3.70 (3.39-4.02)	4.11 (3.54-4.70)	6.17 (5.72-6.47)	8.20 (7.16-8.79)	9.52 (8.66-10.6)	451
20 years and older	99-00	4.26 (3.94-4.62)	4.80 (4.40-5.30)	7.10 (6.50-7.60)	9.80 (8.90-10.7)	11.7 (10.2-13.4)	1406
	01-02	4.74 (4.32-5.20)	5.43 (5.05-5.87)	7.97 (7.43-8.52)	10.6 (9.73-11.5)	12.9 (11.2-14.2)	1560
	03-04	4.56 (4.23-4.90)	5.03 (4.60-5.42)	7.66 (7.01-8.34)	10.7 (9.40-11.5)	12.9 (11.5-14.9)	1543
	05-06	4.67 (4.38-4.99)	5.14 (4.84-5.44)	7.77 (7.32-8.30)	10.4 (9.72-11.5)	13.2 (12.4-13.9)	1520
	07-08	4.38 (4.20-4.58)	4.79 (4.59-4.99)	7.12 (6.81-7.52)	9.69 (9.23-10.2)	11.4 (10.6-12.0)	1857
	09-10	4.10 (3.88-4.34)	4.49 (4.21-4.71)	6.57 (6.33-6.84)	9.26 (8.37-10.1)	11.1 (10.3-11.9)	2019
Gender							
Males	99-00	4.84 (4.35-5.38)	5.50 (4.60-6.00)	7.50 (7.00-8.20)	9.70 (8.80-11.3)	11.7 (10.3-13.0)	1226
	01-02	5.34 (4.89-5.84)	6.13 (5.61-6.64)	8.27 (7.84-9.08)	10.8 (10.1-12.1)	12.8 (11.3-15.0)	1335
	03-04	5.03 (4.73-5.36)	5.59 (5.17-6.00)	7.98 (7.31-8.63)	11.0 (9.53-11.8)	12.9 (11.5-16.1)	1281
	05-06	5.13 (4.78-5.51)	5.37 (5.03-5.71)	7.92 (7.61-8.41)	10.9 (9.96-11.6)	13.2 (12.4-13.9)	1271
	07-08	4.77 (4.59-4.96)	5.26 (4.98-5.43)	7.18 (6.93-7.54)	9.68 (8.93-10.6)	11.6 (10.6-12.6)	1327
	09-10	4.27 (4.00-4.55)	4.70 (4.46-5.00)	6.57 (6.39-6.81)	8.67 (7.91-9.57)	10.7 (9.66-11.3)	1398
Females	99-00	3.95 (3.63-4.29)	4.50 (4.20-4.90)	6.70 (6.20-7.30)	9.10 (8.30-10.0)	11.2 (9.90-12.9)	1238
	01-02	4.36 (3.95-4.81)	4.87 (4.45-5.25)	7.33 (6.71-8.01)	9.77 (9.07-11.0)	12.4 (10.4-13.8)	1355
	03-04	4.35 (4.03-4.70)	4.79 (4.25-5.26)	7.30 (6.87-7.81)	10.2 (9.40-11.0)	12.1 (11.2-13.6)	1277
	05-06	4.30 (3.98-4.65)	4.89 (4.58-5.25)	7.43 (6.99-7.83)	9.79 (9.30-10.5)	12.1 (10.6-13.1)	1305
	07-08	4.12 (3.88-4.37)	4.40 (4.17-4.74)	6.84 (6.32-7.36)	9.31 (8.69-9.76)	10.5 (9.93-11.7)	1300
	09-10	3.88 (3.63-4.14)	4.13 (3.83-4.55)	6.46 (5.95-6.90)	9.29 (8.62-10.0)	11.2 (10.3-12.1)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.14, 0.14, 0.2, 0.066, 0.066, and 0.066 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cesium_BiomonitoringSummary.html

Urinary Cesium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	4.32 (3.82-4.89)	4.80 (4.30-5.20)	6.70 (6.40-7.20)	9.10 (8.00-9.90)	11.1 (9.60-12.7)	884
	01-02	4.63 (4.10-5.24)	5.29 (4.59-5.89)	7.08 (6.42-7.99)	9.13 (7.86-11.3)	11.3 (8.81-14.9)	683
	03-04	4.94 (4.64-5.27)	5.62 (5.01-6.09)	7.86 (7.15-8.46)	10.3 (8.99-11.3)	11.9 (10.7-14.7)	618
	05-06	4.64 (4.41-4.88)	5.18 (4.91-5.55)	7.29 (6.74-7.73)	9.52 (8.76-9.98)	11.5 (9.88-11.9)	652
	07-08	4.50 (4.06-4.99)	4.93 (4.40-5.56)	6.98 (6.38-7.49)	9.12 (8.25-9.82)	10.8 (9.45-12.1)	515
	09-10	4.20 (3.95-4.48)	4.47 (4.16-4.87)	6.49 (6.00-7.01)	8.64 (7.81-9.40)	10.2 (9.15-11.4)	613
Non-Hispanic blacks	99-00	4.94 (4.33-5.64)	5.40 (4.80-6.40)	7.50 (6.90-8.40)	9.80 (8.80-10.8)	11.6 (9.80-13.1)	568
	01-02	4.93 (4.70-5.17)	5.33 (5.05-5.64)	7.36 (6.97-7.59)	9.44 (8.71-9.68)	10.7 (10.1-12.3)	667
	03-04	4.71 (4.47-4.97)	5.12 (4.71-5.49)	7.17 (6.72-7.60)	9.13 (8.52-9.99)	10.7 (9.99-11.4)	723
	05-06	4.78 (4.42-5.18)	5.26 (5.03-5.44)	7.20 (6.63-7.66)	9.14 (8.26-10.6)	11.0 (9.63-13.7)	692
	07-08	4.64 (4.29-5.01)	4.90 (4.48-5.31)	6.90 (6.35-7.42)	9.39 (8.29-10.3)	11.3 (9.59-12.6)	589
	09-10	4.07 (3.82-4.35)	4.68 (4.29-4.84)	6.56 (6.19-7.01)	8.98 (8.12-9.31)	9.97 (9.27-11.3)	544
Non-Hispanic whites	99-00	4.25 (3.83-4.72)	4.80 (4.20-5.50)	7.20 (6.60-7.90)	9.70 (8.90-10.7)	11.8 (10.3-13.3)	821
	01-02	4.77 (4.27-5.32)	5.49 (4.99-6.05)	7.98 (7.45-8.61)	10.4 (9.54-11.4)	12.6 (11.0-13.8)	1132
	03-04	4.56 (4.22-4.94)	5.02 (4.59-5.42)	7.60 (7.00-8.23)	10.7 (9.26-11.8)	12.9 (11.5-14.7)	1074
	05-06	4.64 (4.25-5.05)	5.19 (4.76-5.49)	7.77 (7.18-8.46)	10.3 (9.62-11.5)	13.1 (12.1-13.9)	1041
	07-08	4.29 (4.07-4.53)	4.76 (4.48-5.08)	7.02 (6.67-7.49)	9.50 (8.91-9.93)	11.3 (10.3-12.2)	1095
	09-10	3.96 (3.73-4.20)	4.38 (4.10-4.64)	6.43 (6.15-6.63)	8.62 (7.86-9.29)	10.5 (9.35-11.5)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.14, 0.14, 0.2, 0.066, 0.066, and 0.066 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cesium_BiomonitoringSummary.html

Urinary Cesium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	4.10 (3.96-4.25)	4.13 (3.97-4.27)	5.41 (5.21-5.70)	7.14 (6.83-7.50)	8.64 (8.00-9.30)	2464
	01-02	4.54 (4.30-4.79)	4.44 (4.20-4.64)	6.06 (5.66-6.47)	8.18 (7.62-8.95)	10.2 (8.84-11.7)	2689
	03-04	4.64 (4.42-4.87)	4.42 (4.21-4.74)	6.11 (5.76-6.48)	8.51 (7.99-9.15)	10.6 (9.75-11.0)	2558
	05-06	4.62 (4.37-4.88)	4.51 (4.20-4.76)	6.19 (5.90-6.55)	8.84 (8.14-9.24)	10.8 (9.71-12.1)	2576
	07-08	4.62 (4.42-4.82)	4.58 (4.36-4.79)	6.23 (5.96-6.61)	8.67 (7.79-9.52)	10.6 (9.78-11.2)	2627
	09-10	4.33 (4.17-4.49)	4.28 (4.12-4.45)	6.00 (5.70-6.23)	8.06 (7.76-8.36)	9.75 (9.05-10.4)	2848
Age group							
6-11 years	99-00	5.34 (5.03-5.67)	5.42 (5.03-6.04)	6.63 (6.18-7.13)	8.23 (7.13-9.41)	9.90 (7.88-10.1)	340
	01-02	5.95 (5.48-6.46)	5.91 (5.43-6.53)	7.77 (7.00-8.28)	9.27 (8.35-11.9)	11.9 (9.38-12.3)	368
	03-04	6.07 (5.63-6.53)	6.02 (5.45-6.49)	8.30 (7.16-8.99)	10.8 (9.15-11.8)	11.9 (10.3-15.8)	290
	05-06	5.85 (5.41-6.34)	5.74 (5.21-6.41)	7.65 (6.67-9.11)	10.1 (9.38-10.8)	11.6 (10.1-14.9)	355
	07-08	5.96 (5.53-6.43)	6.12 (5.79-6.55)	7.88 (7.33-8.53)	10.7 (8.91-12.1)	12.5 (10.9-14.1)	394
	09-10	5.74 (5.30-6.22)	5.58 (5.15-6.12)	7.16 (6.71-7.87)	9.19 (8.13-10.4)	12.2 (8.69-14.8)	378
12-19 years	99-00	3.43 (3.29-3.58)	3.54 (3.29-3.68)	4.35 (4.17-4.56)	5.31 (4.97-5.79)	6.67 (5.33-8.09)	718
	01-02	3.73 (3.41-4.08)	3.55 (3.36-3.74)	4.74 (4.40-5.14)	6.10 (5.35-7.63)	8.08 (6.44-9.82)	762
	03-04	3.60 (3.37-3.85)	3.51 (3.33-3.72)	4.53 (4.24-4.87)	6.08 (5.14-6.94)	7.27 (6.13-9.07)	725
	05-06	3.66 (3.40-3.93)	3.59 (3.40-3.83)	4.76 (4.30-5.20)	5.93 (5.62-6.62)	7.62 (6.03-8.07)	701
	07-08	3.58 (3.31-3.88)	3.70 (3.22-3.96)	4.98 (4.60-5.17)	6.36 (5.38-7.28)	7.93 (6.06-9.76)	376
	09-10	3.46 (3.27-3.65)	3.48 (3.23-3.74)	4.39 (4.16-4.70)	5.54 (5.01-6.32)	6.56 (5.95-7.35)	451
20 years and older	99-00	4.08 (3.88-4.29)	4.06 (3.85-4.29)	5.39 (5.04-5.85)	7.17 (6.84-7.58)	8.61 (7.91-9.30)	1406
	01-02	4.54 (4.30-4.78)	4.43 (4.20-4.59)	5.94 (5.64-6.40)	8.15 (7.46-8.97)	10.2 (8.74-11.7)	1559
	03-04	4.68 (4.46-4.91)	4.47 (4.27-4.80)	6.11 (5.83-6.43)	8.47 (7.76-9.17)	10.5 (9.68-11.2)	1543
	05-06	4.67 (4.42-4.93)	4.54 (4.25-4.78)	6.24 (5.93-6.66)	8.92 (8.29-9.31)	11.2 (9.71-12.7)	1520
	07-08	4.67 (4.46-4.88)	4.58 (4.34-4.82)	6.19 (5.93-6.61)	8.65 (7.79-9.52)	10.6 (9.68-11.3)	1857
	09-10	4.34 (4.17-4.53)	4.28 (4.10-4.49)	6.07 (5.70-6.33)	8.20 (7.81-8.65)	9.80 (9.05-11.1)	2019
Gender							
Males	99-00	3.78 (3.65-3.91)	3.78 (3.61-3.96)	4.96 (4.72-5.20)	6.50 (6.18-6.70)	7.71 (7.01-8.64)	1226
	01-02	4.22 (3.96-4.51)	4.10 (3.87-4.41)	5.60 (5.27-6.03)	7.67 (6.90-8.48)	9.46 (8.22-11.5)	1334
	03-04	4.24 (3.99-4.50)	4.12 (3.89-4.37)	5.66 (5.19-6.06)	7.66 (6.90-8.40)	9.00 (8.36-10.3)	1281
	05-06	4.21 (3.94-4.50)	4.11 (3.72-4.46)	5.75 (5.14-6.11)	7.68 (6.91-8.78)	9.33 (8.84-9.84)	1271
	07-08	4.27 (4.04-4.51)	4.19 (4.02-4.42)	5.81 (5.46-6.29)	7.84 (7.20-8.65)	9.61 (8.65-10.2)	1327
	09-10	3.89 (3.70-4.09)	3.85 (3.60-4.06)	5.29 (4.89-5.69)	7.02 (6.76-7.42)	8.38 (7.87-9.18)	1398
Females	99-00	4.43 (4.20-4.68)	4.45 (4.14-4.77)	5.92 (5.36-6.47)	7.70 (7.16-8.07)	9.41 (8.00-10.4)	1238
	01-02	4.86 (4.58-5.16)	4.72 (4.50-5.06)	6.54 (5.93-7.00)	8.50 (7.84-9.79)	10.3 (8.95-12.2)	1355
	03-04	5.05 (4.77-5.35)	4.77 (4.44-5.14)	6.58 (6.14-7.22)	9.43 (8.56-10.5)	11.3 (10.7-12.3)	1277
	05-06	5.05 (4.78-5.33)	4.86 (4.57-5.17)	6.72 (6.31-7.23)	9.65 (9.23-10.3)	12.6 (11.0-13.3)	1305
	07-08	4.98 (4.76-5.20)	4.90 (4.69-5.20)	6.72 (6.20-7.06)	9.50 (8.37-10.5)	11.3 (10.7-12.1)	1300
	09-10	4.79 (4.63-4.97)	4.69 (4.50-4.93)	6.57 (6.34-6.85)	8.88 (8.36-9.33)	10.5 (9.56-12.0)	1450

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cesium_BiomonitoringSummary.html

Urinary Cesium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	3.99 (3.73-4.25)	3.95 (3.65-4.17)	5.09 (4.79-5.39)	6.65 (6.08-7.10)	7.98 (7.20-8.95)	884
	01-02	4.51 (4.00-5.08)	4.51 (3.82-4.95)	5.91 (5.31-6.64)	7.77 (6.60-10.0)	10.0 (7.60-20.5)	682
	03-04	4.58 (4.16-5.05)	4.51 (4.10-4.92)	5.74 (5.42-6.09)	7.53 (6.59-8.91)	9.44 (8.24-10.6)	618
	05-06	4.36 (4.10-4.64)	4.40 (3.97-4.73)	5.69 (5.24-6.22)	7.23 (6.69-7.80)	8.14 (7.65-9.33)	652
	07-08	4.50 (4.22-4.79)	4.39 (3.96-4.86)	6.07 (5.57-6.59)	7.63 (7.30-7.89)	9.06 (7.89-10.5)	515
	09-10	4.41 (4.17-4.65)	4.38 (4.04-4.59)	5.70 (5.34-6.07)	7.68 (7.17-8.24)	8.89 (8.50-9.52)	613
Non-Hispanic blacks	99-00	3.21 (2.90-3.56)	3.26 (3.05-3.44)	4.30 (4.00-4.55)	5.50 (5.00-5.98)	6.33 (5.91-7.04)	568
	01-02	3.38 (3.19-3.57)	3.35 (3.05-3.60)	4.41 (4.15-4.78)	5.87 (5.63-6.29)	6.75 (6.41-7.03)	667
	03-04	3.38 (3.21-3.56)	3.30 (3.08-3.50)	4.31 (4.02-4.62)	5.79 (5.12-6.47)	6.98 (6.38-7.18)	723
	05-06	3.47 (3.34-3.60)	3.40 (3.21-3.60)	4.47 (4.25-4.68)	5.94 (5.52-6.31)	6.95 (6.20-7.51)	692
	07-08	3.39 (3.21-3.58)	3.33 (3.18-3.54)	4.50 (4.21-4.78)	5.84 (5.41-6.18)	6.87 (5.94-7.58)	589
	09-10	3.23 (3.05-3.42)	3.20 (2.99-3.46)	4.31 (4.12-4.59)	5.88 (5.30-6.28)	6.69 (6.32-7.89)	544
Non-Hispanic whites	99-00	4.26 (4.07-4.47)	4.28 (4.05-4.50)	5.66 (5.26-6.05)	7.27 (6.84-7.83)	8.75 (7.93-9.38)	821
	01-02	4.81 (4.55-5.08)	4.63 (4.42-4.96)	6.33 (5.91-6.68)	8.46 (7.84-9.39)	10.3 (9.04-11.8)	1132
	03-04	4.81 (4.52-5.12)	4.56 (4.31-4.98)	6.28 (5.95-6.71)	8.63 (7.99-9.28)	10.6 (9.43-11.0)	1074
	05-06	4.82 (4.50-5.17)	4.68 (4.32-5.04)	6.41 (6.00-6.81)	9.18 (8.67-9.53)	11.0 (9.71-12.7)	1041
	07-08	4.81 (4.54-5.09)	4.73 (4.42-5.12)	6.43 (6.01-6.80)	9.24 (8.20-9.82)	11.0 (10.2-11.7)	1095
	09-10	4.42 (4.23-4.63)	4.36 (4.15-4.54)	6.16 (5.82-6.51)	8.13 (7.68-8.65)	9.83 (8.79-11.4)	1225

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cesium_BiomonitoringSummary.html

Urinary Cobalt

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.375 (.336-.419)	.410 (.370-.450)	.630 (.570-.680)	.950 (.890-1.03)	1.32 (1.16-1.48)	2465
	01-02	.379 (.355-.404)	.410 (.380-.430)	.610 (.570-.660)	.940 (.870-1.00)	1.28 (1.15-1.44)	2690
	03-04	.316 (.291-.343)	.330 (.300-.350)	.520 (.490-.550)	.820 (.750-.890)	1.16 (1.08-1.26)	2558
	05-06	.374 (.357-.392)	.390 (.370-.410)	.600 (.560-.630)	.880 (.840-.950)	1.33 (1.17-1.47)	2576
	07-08	.369 (.351-.389)	.380 (.360-.390)	.580 (.550-.600)	.920 (.840-1.01)	1.32 (1.10-1.43)	2627
	09-10	.369 (.341-.398)	.380 (.350-.400)	.600 (.560-.650)	.960 (.870-1.06)	1.40 (1.20-1.58)	2848
Age group							
6-11 years	99-00	.499 (.427-.583)	.530 (.450-.640)	.750 (.610-.900)	1.03 (.880-1.12)	1.22 (1.03-1.50)	340
	01-02	.452 (.377-.543)	.520 (.430-.610)	.710 (.670-.810)	1.07 (.940-1.21)	1.32 (1.17-1.53)	368
	03-04	.454 (.393-.523)	.460 (.410-.520)	.750 (.590-.900)	1.24 (.910-1.47)	1.68 (1.26-1.81)	290
	05-06	.424 (.383-.468)	.450 (.410-.490)	.660 (.580-.710)	.870 (.830-1.04)	1.16 (.910-1.39)	355
	07-08	.433 (.367-.509)	.450 (.360-.550)	.700 (.560-.810)	.960 (.790-1.28)	1.40 (.880-1.82)	394
	09-10	.424 (.379-.474)	.440 (.400-.490)	.670 (.610-.740)	.960 (.850-1.10)	1.14 (1.01-1.69)	378
12-19 years	99-00	.519 (.463-.581)	.520 (.490-.570)	.820 (.670-.890)	1.17 (1.01-1.47)	1.52 (1.26-2.56)	719
	01-02	.515 (.469-.564)	.520 (.480-.570)	.750 (.690-.840)	1.24 (1.07-1.32)	1.59 (1.37-1.99)	762
	03-04	.461 (.428-.496)	.480 (.450-.520)	.740 (.650-.800)	1.03 (.940-1.23)	1.60 (1.14-1.92)	725
	05-06	.510 (.451-.577)	.500 (.460-.570)	.820 (.690-.980)	1.30 (1.08-1.40)	1.74 (1.39-2.08)	701
	07-08	.529 (.478-.585)	.510 (.450-.570)	.820 (.690-.960)	1.36 (1.09-1.53)	1.70 (1.40-1.78)	376
	09-10	.465 (.406-.533)	.460 (.430-.530)	.830 (.710-.920)	1.21 (1.05-1.51)	1.56 (1.19-2.11)	451
20 years and older	99-00	.343 (.305-.386)	.370 (.330-.420)	.570 (.520-.640)	.880 (.810-.980)	1.28 (1.07-1.39)	1406
	01-02	.352 (.333-.373)	.380 (.350-.410)	.560 (.520-.590)	.860 (.800-.930)	1.15 (1.04-1.42)	1560
	03-04	.285 (.259-.313)	.300 (.270-.330)	.460 (.410-.510)	.690 (.660-.730)	1.06 (.890-1.14)	1543
	05-06	.351 (.332-.371)	.370 (.350-.390)	.550 (.520-.600)	.810 (.760-.880)	1.30 (1.05-1.47)	1520
	07-08	.343 (.326-.362)	.350 (.340-.370)	.530 (.500-.560)	.830 (.750-.920)	1.20 (1.00-1.38)	1857
	09-10	.351 (.322-.382)	.350 (.330-.380)	.570 (.520-.630)	.890 (.800-1.02)	1.35 (1.12-1.58)	2019
Gender							
Males	99-00	.371 (.331-.416)	.410 (.370-.450)	.580 (.540-.640)	.820 (.740-.900)	1.01 (.900-1.12)	1227
	01-02	.367 (.338-.399)	.390 (.360-.420)	.550 (.520-.600)	.790 (.740-.850)	1.05 (.960-1.14)	1335
	03-04	.294 (.270-.319)	.320 (.290-.340)	.480 (.440-.500)	.670 (.620-.710)	.870 (.790-.920)	1281
	05-06	.369 (.354-.383)	.380 (.370-.400)	.540 (.510-.570)	.760 (.690-.850)	1.06 (.910-1.27)	1271
	07-08	.350 (.330-.371)	.370 (.350-.390)	.530 (.500-.560)	.750 (.690-.820)	1.00 (.860-1.14)	1327
	09-10	.336 (.307-.368)	.360 (.330-.380)	.530 (.480-.570)	.750 (.690-.850)	.960 (.850-1.12)	1398
Females	99-00	.379 (.333-.431)	.410 (.340-.460)	.680 (.590-.790)	1.17 (.930-1.36)	1.50 (1.28-2.05)	1238
	01-02	.390 (.364-.417)	.430 (.400-.450)	.670 (.620-.700)	1.06 (.980-1.16)	1.46 (1.22-1.81)	1355
	03-04	.339 (.308-.372)	.340 (.310-.370)	.580 (.540-.610)	1.04 (.900-1.13)	1.47 (1.33-1.73)	1277
	05-06	.379 (.351-.409)	.410 (.370-.440)	.660 (.620-.700)	1.04 (.900-1.21)	1.54 (1.30-1.93)	1305
	07-08	.389 (.367-.411)	.390 (.360-.410)	.650 (.590-.690)	1.08 (.980-1.26)	1.43 (1.34-1.62)	1300
	09-10	.403 (.364-.447)	.400 (.360-.440)	.700 (.640-.770)	1.19 (1.05-1.36)	1.68 (1.51-2.11)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.07, 0.07, 0.08, 0.041, 0.041, and 0.041 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cobalt_BiomonitoringSummary.html

Urinary Cobalt

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.418 (.348-.502)	.470 (.370-.530)	.670 (.630-.770)	1.05 (.950-1.19)	1.47 (1.25-1.67)	884
	01-02	.398 (.373-.424)	.430 (.410-.450)	.650 (.600-.710)	.950 (.850-1.03)	1.20 (1.06-1.48)	683
	03-04	.374 (.350-.398)	.350 (.340-.390)	.580 (.530-.620)	1.09 (.920-1.16)	1.33 (1.16-1.73)	618
	05-06	.437 (.404-.473)	.450 (.420-.500)	.680 (.620-.720)	1.11 (.940-1.30)	1.58 (1.33-1.79)	652
	07-08	.396 (.349-.450)	.400 (.330-.480)	.610 (.560-.710)	.980 (.870-1.08)	1.26 (1.05-1.69)	515
	09-10	.378 (.351-.407)	.380 (.360-.410)	.590 (.560-.640)	.920 (.860-1.04)	1.24 (1.04-1.58)	613
Non-Hispanic blacks	99-00	.434 (.405-.465)	.430 (.390-.470)	.680 (.620-.760)	1.17 (1.04-1.26)	1.45 (1.23-2.04)	568
	01-02	.435 (.388-.487)	.420 (.380-.460)	.650 (.540-.810)	1.16 (.850-1.64)	1.75 (1.32-2.22)	667
	03-04	.380 (.348-.414)	.380 (.360-.410)	.600 (.540-.680)	1.01 (.890-1.09)	1.28 (1.01-2.03)	723
	05-06	.412 (.374-.453)	.410 (.360-.440)	.600 (.560-.670)	.930 (.830-1.11)	1.47 (1.12-2.00)	692
	07-08	.403 (.367-.443)	.390 (.360-.420)	.620 (.560-.740)	1.06 (.880-1.31)	1.57 (1.22-1.86)	589
	09-10	.368 (.335-.403)	.380 (.340-.400)	.590 (.520-.650)	.990 (.870-1.15)	1.47 (1.20-1.58)	544
Non-Hispanic whites	99-00	.369 (.316-.431)	.410 (.350-.460)	.630 (.550-.700)	.930 (.830-1.08)	1.29 (1.02-1.65)	822
	01-02	.359 (.327-.394)	.390 (.360-.430)	.590 (.520-.660)	.870 (.800-.950)	1.16 (1.04-1.32)	1132
	03-04	.301 (.270-.334)	.310 (.280-.340)	.500 (.460-.540)	.760 (.690-.850)	1.09 (.950-1.26)	1074
	05-06	.361 (.340-.384)	.380 (.360-.410)	.580 (.530-.630)	.860 (.800-.940)	1.30 (1.06-1.50)	1041
	07-08	.358 (.338-.379)	.370 (.350-.390)	.560 (.520-.600)	.900 (.780-1.00)	1.32 (1.02-1.46)	1095
	09-10	.369 (.335-.408)	.370 (.350-.400)	.600 (.540-.660)	.970 (.850-1.10)	1.40 (1.17-1.64)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.07, 0.07, 0.08, 0.041, 0.041, and 0.041 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cobalt_BiomonitoringSummary.html

Urinary Cobalt (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.353 (.319-.391)	.328 (.302-.365)	.515 (.457-.581)	.821 (.679-.963)	1.16 (.938-1.50)	2465
	01-02	.358 (.333-.384)	.335 (.313-.360)	.523 (.487-.562)	.844 (.750-.955)	1.16 (1.00-1.28)	2689
	03-04	.314 (.303-.325)	.290 (.278-.306)	.455 (.434-.481)	.737 (.703-.781)	1.02 (.911-1.10)	2558
	05-06	.368 (.353-.384)	.340 (.320-.360)	.520 (.490-.550)	.810 (.750-.920)	1.17 (.990-1.27)	2576
	07-08	.385 (.366-.406)	.360 (.350-.380)	.560 (.520-.590)	.870 (.760-.990)	1.22 (1.01-1.46)	2627
	09-10	.393 (.372-.415)	.370 (.350-.390)	.560 (.530-.600)	.890 (.830-.940)	1.29 (1.08-1.49)	2848
Age group							
6-11 years	99-00	.547 (.467-.640)	.554 (.449-.647)	.774 (.626-.938)	1.00 (.833-1.49)	1.25 (.895-1.50)	340
	01-02	.552 (.508-.599)	.548 (.503-.609)	.756 (.660-.829)	1.00 (.900-1.27)	1.30 (1.03-1.73)	368
	03-04	.529 (.471-.593)	.500 (.463-.543)	.689 (.634-.750)	1.04 (.760-1.29)	1.29 (1.04-1.36)	290
	05-06	.523 (.494-.554)	.520 (.480-.570)	.720 (.660-.760)	.900 (.840-1.04)	1.27 (.980-1.45)	355
	07-08	.564 (.501-.635)	.580 (.500-.630)	.760 (.660-.910)	1.12 (.830-1.46)	1.39 (1.06-1.81)	394
	09-10	.578 (.538-.620)	.570 (.530-.620)	.760 (.690-.840)	1.06 (.920-1.21)	1.26 (1.06-1.58)	378
12-19 years	99-00	.391 (.353-.433)	.378 (.329-.407)	.537 (.469-.595)	.824 (.638-1.17)	1.44 (.821-3.54)	719
	01-02	.368 (.343-.396)	.352 (.327-.372)	.534 (.471-.611)	.851 (.673-.949)	1.06 (.932-1.24)	762
	03-04	.329 (.304-.355)	.316 (.277-.348)	.495 (.442-.561)	.738 (.630-.847)	.952 (.792-1.09)	725
	05-06	.392 (.360-.427)	.360 (.330-.400)	.530 (.480-.620)	.810 (.720-.920)	1.02 (.870-1.48)	701
	07-08	.414 (.382-.448)	.390 (.340-.450)	.610 (.520-.700)	.880 (.750-1.13)	1.17 (.890-1.35)	376
	09-10	.435 (.398-.476)	.410 (.370-.470)	.610 (.550-.700)	.960 (.790-1.07)	1.30 (1.00-1.53)	451
20 years and older	99-00	.328 (.297-.362)	.306 (.280-.328)	.471 (.428-.522)	.727 (.632-.905)	1.12 (.861-1.36)	1406
	01-02	.337 (.313-.363)	.313 (.294-.337)	.475 (.435-.513)	.792 (.704-.955)	1.15 (.963-1.33)	1559
	03-04	.293 (.282-.304)	.271 (.257-.286)	.400 (.380-.429)	.691 (.616-.744)	.976 (.829-1.10)	1543
	05-06	.350 (.334-.367)	.320 (.300-.340)	.490 (.450-.520)	.780 (.690-.940)	1.17 (.970-1.31)	1520
	07-08	.366 (.348-.384)	.340 (.330-.360)	.510 (.480-.540)	.830 (.720-.960)	1.19 (.980-1.47)	1857
	09-10	.371 (.350-.394)	.340 (.320-.360)	.520 (.490-.550)	.840 (.750-.930)	1.29 (1.02-1.52)	2019
Gender							
Males	99-00	.290 (.259-.324)	.279 (.248-.301)	.402 (.365-.449)	.608 (.534-.728)	.838 (.667-1.10)	1227
	01-02	.290 (.272-.310)	.278 (.256-.297)	.392 (.361-.425)	.644 (.574-.707)	.848 (.786-.929)	1334
	03-04	.247 (.237-.259)	.234 (.215-.250)	.333 (.313-.352)	.513 (.476-.585)	.700 (.630-.753)	1281
	05-06	.302 (.289-.317)	.290 (.270-.300)	.410 (.370-.440)	.610 (.580-.640)	.770 (.710-.850)	1271
	07-08	.313 (.298-.329)	.300 (.280-.310)	.420 (.400-.440)	.630 (.580-.680)	.830 (.680-1.02)	1327
	09-10	.306 (.290-.323)	.290 (.280-.300)	.420 (.390-.450)	.630 (.570-.690)	.810 (.720-.920)	1398
Females	99-00	.426 (.378-.479)	.407 (.362-.457)	.606 (.550-.694)	.960 (.781-1.29)	1.50 (1.11-1.83)	1238
	01-02	.435 (.404-.468)	.408 (.382-.438)	.635 (.560-.708)	1.00 (.879-1.19)	1.29 (1.12-1.60)	1355
	03-04	.393 (.378-.409)	.361 (.342-.381)	.554 (.513-.615)	.937 (.850-1.00)	1.29 (1.10-1.33)	1277
	05-06	.445 (.423-.468)	.420 (.400-.440)	.630 (.590-.700)	.990 (.900-1.17)	1.34 (1.18-1.60)	1305
	07-08	.470 (.442-.499)	.450 (.430-.480)	.690 (.630-.750)	1.09 (.930-1.27)	1.46 (1.26-1.69)	1300
	09-10	.498 (.465-.534)	.460 (.450-.480)	.690 (.650-.720)	1.09 (.990-1.33)	1.64 (1.34-2.15)	1450

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cobalt_BiomonitoringSummary.html

Urinary Cobalt (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.386 (.339-.439)	.376 (.333-.419)	.598 (.500-.669)	.898 (.826-1.00)	1.23 (1.11-1.35)	884
	01-02	.388 (.361-.417)	.361 (.333-.394)	.591 (.500-.662)	.872 (.777-.990)	1.10 (.990-1.27)	682
	03-04	.346 (.326-.368)	.327 (.296-.349)	.542 (.487-.594)	.850 (.753-.963)	1.14 (.963-1.35)	618
	05-06	.411 (.388-.436)	.380 (.360-.400)	.580 (.520-.650)	.900 (.820-1.02)	1.19 (1.05-1.35)	652
	07-08	.396 (.348-.450)	.370 (.330-.410)	.560 (.470-.690)	.900 (.770-1.12)	1.17 (.990-1.54)	515
	09-10	.396 (.367-.427)	.380 (.340-.420)	.590 (.530-.650)	.930 (.820-1.03)	1.22 (1.04-1.30)	613
Non-Hispanic blacks	99-00	.282 (.275-.289)	.257 (.243-.279)	.417 (.378-.462)	.723 (.600-.785)	.975 (.757-1.60)	568
	01-02	.298 (.275-.323)	.268 (.251-.296)	.444 (.393-.505)	.728 (.582-.917)	1.03 (.740-1.55)	667
	03-04	.273 (.248-.300)	.259 (.239-.281)	.388 (.344-.461)	.700 (.563-.842)	.964 (.733-1.15)	723
	05-06	.299 (.281-.318)	.280 (.240-.300)	.420 (.380-.460)	.700 (.570-.780)	.950 (.790-1.17)	692
	07-08	.295 (.277-.314)	.270 (.260-.290)	.420 (.390-.450)	.710 (.570-.860)	1.07 (.850-1.28)	589
	09-10	.291 (.271-.314)	.280 (.240-.310)	.420 (.380-.480)	.700 (.610-.750)	.840 (.750-1.06)	544
Non-Hispanic whites	99-00	.369 (.324-.421)	.352 (.313-.387)	.533 (.452-.611)	.861 (.667-1.16)	1.25 (.895-1.57)	822
	01-02	.362 (.331-.396)	.343 (.313-.368)	.523 (.479-.562)	.830 (.736-.983)	1.16 (.983-1.33)	1132
	03-04	.317 (.301-.334)	.291 (.274-.309)	.457 (.425-.488)	.738 (.683-.804)	1.00 (.857-1.13)	1074
	05-06	.376 (.357-.395)	.350 (.330-.370)	.520 (.490-.580)	.810 (.710-.950)	1.18 (.950-1.43)	1041
	07-08	.401 (.380-.423)	.380 (.360-.400)	.590 (.530-.630)	.890 (.760-1.06)	1.27 (.980-1.55)	1095
	09-10	.413 (.388-.440)	.390 (.360-.410)	.580 (.530-.620)	.910 (.830-.980)	1.33 (1.08-1.58)	1225

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Cobalt_BiomonitoringSummary.html

Blood Lead

Geometric mean and selected percentiles of blood concentrations (in µg/dL) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	1.83 (1.75-1.91)	1.80 (1.70-1.90)	2.80 (2.60-2.90)	4.20 (3.90-4.60)	5.80 (5.10-6.60)	2742
	01-02	1.46 (1.34-1.60)	1.50 (1.30-1.60)	2.30 (2.10-2.60)	3.60 (3.40-4.20)	5.40 (4.40-6.70)	2268
	03-04	1.55 (1.43-1.69)	1.50 (1.40-1.60)	2.30 (2.10-2.50)	3.50 (2.90-4.20)	4.90 (3.90-6.40)	2085
	05-06	1.29 (1.21-1.38)	1.26 (1.15-1.36)	2.00 (1.87-2.20)	3.16 (2.79-3.58)	4.22 (3.47-5.36)	2236
	07-08	1.25 (1.15-1.36)	1.20 (1.10-1.31)	1.88 (1.70-2.06)	2.81 (2.60-3.20)	3.92 (3.20-5.00)	1712
	09-10	1.14 (1.03-1.28)	1.04 (.940-1.18)	1.76 (1.52-1.99)	2.93 (2.63-3.30)	3.92 (3.60-4.69)	1966
Non-Hispanic blacks	99-00	1.87 (1.75-2.00)	1.80 (1.70-2.00)	2.80 (2.60-3.00)	4.30 (4.00-4.60)	5.70 (5.20-6.10)	1842
	01-02	1.65 (1.52-1.80)	1.60 (1.40-1.70)	2.60 (2.30-2.90)	4.20 (3.80-4.70)	5.80 (5.30-6.50)	2219
	03-04	1.69 (1.52-1.89)	1.60 (1.40-1.80)	2.60 (2.20-3.00)	4.10 (3.50-4.70)	5.30 (4.60-6.60)	2293
	05-06	1.39 (1.26-1.53)	1.31 (1.19-1.45)	2.16 (1.95-2.42)	3.48 (3.19-3.80)	4.65 (4.21-5.14)	2193
	07-08	1.39 (1.30-1.48)	1.30 (1.20-1.42)	2.10 (2.00-2.20)	3.22 (3.08-3.50)	4.50 (4.00-4.80)	1746
	09-10	1.24 (1.18-1.30)	1.19 (1.12-1.25)	1.87 (1.76-1.99)	2.90 (2.68-3.18)	3.86 (3.57-4.29)	1593
Non-Hispanic whites	99-00	1.62 (1.55-1.69)	1.60 (1.50-1.70)	2.40 (2.30-2.50)	3.60 (3.40-3.90)	5.00 (4.40-5.70)	2716
	01-02	1.43 (1.37-1.48)	1.40 (1.30-1.50)	2.20 (2.10-2.20)	3.20 (3.10-3.40)	4.20 (3.90-4.50)	3806
	03-04	1.37 (1.32-1.43)	1.30 (1.30-1.40)	2.10 (2.00-2.10)	3.00 (2.80-3.20)	3.90 (3.60-4.30)	3478
	05-06	1.28 (1.19-1.37)	1.27 (1.17-1.38)	1.97 (1.86-2.14)	2.99 (2.73-3.25)	3.82 (3.41-4.20)	3310
	07-08	1.24 (1.16-1.33)	1.20 (1.10-1.30)	1.86 (1.72-2.00)	2.70 (2.54-2.90)	3.50 (3.20-3.89)	3461
	09-10	1.10 (1.04-1.16)	1.07 (1.00-1.15)	1.67 (1.59-1.76)	2.49 (2.35-2.63)	3.14 (2.99-3.36)	3760

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.3, 0.3, 0.28, 0.25, 0.25, and 0.25 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Lead_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Lead_FactSheet.html

Urinary Lead

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.766 (.708-.828)	.800 (.800-.900)	1.40 (1.30-1.50)	2.20 (2.00-2.30)	2.90 (2.60-3.30)	2465
	01-02	.677 (.637-.718)	.700 (.700-.800)	1.20 (1.20-1.30)	2.00 (1.90-2.20)	2.70 (2.50-2.80)	2690
	03-04	.636 (.595-.680)	.640 (.580-.690)	1.04 (.960-1.12)	1.73 (1.52-1.86)	2.29 (2.03-2.62)	2558
	05-06	.554 (.523-.587)	.570 (.540-.600)	.990 (.910-1.05)	1.58 (1.44-1.73)	2.14 (1.94-2.45)	2576
	07-08	.493 (.467-.520)	.500 (.470-.530)	.850 (.790-.910)	1.38 (1.23-1.58)	1.97 (1.75-2.17)	2627
	09-10	.458 (.441-.476)	.470 (.450-.480)	.790 (.750-.830)	1.24 (1.11-1.38)	1.65 (1.46-1.84)	2848
Age group							
6-11 years	99-00	1.07 (.955-1.20)	1.10 (.900-1.30)	1.50 (1.40-1.70)	2.40 (1.80-3.10)	3.40 (2.40-5.00)	340
	01-02	.753 (.661-.857)	.800 (.600-.900)	1.20 (1.10-1.40)	2.10 (1.60-2.40)	2.60 (2.10-3.70)	368
	03-04	.795 (.671-.941)	.790 (.640-.900)	1.35 (.970-1.86)	2.27 (1.62-4.09)	3.33 (2.23-4.41)	290
	05-06	.508 (.447-.579)	.510 (.440-.560)	.740 (.620-.980)	1.33 (1.00-1.86)	1.90 (1.33-2.60)	355
	07-08	.494 (.411-.593)	.520 (.410-.590)	.790 (.660-.900)	1.23 (.980-1.73)	1.90 (1.51-2.18)	394
	09-10	.443 (.399-.493)	.460 (.430-.490)	.710 (.630-.820)	1.09 (.880-1.33)	1.40 (1.06-1.82)	378
12-19 years	99-00	.659 (.579-.749)	.700 (.600-.800)	1.10 (.900-1.30)	1.80 (1.40-2.20)	2.20 (1.90-2.80)	719
	01-02	.564 (.526-.605)	.600 (.500-.600)	1.00 (.800-1.10)	1.60 (1.40-1.70)	2.00 (1.80-2.40)	762
	03-04	.604 (.553-.660)	.630 (.570-.680)	.920 (.810-1.02)	1.32 (1.14-1.80)	1.86 (1.44-2.29)	725
	05-06	.472 (.421-.530)	.490 (.450-.540)	.780 (.710-.830)	1.19 (1.10-1.45)	1.65 (1.34-1.80)	701
	07-08	.386 (.346-.431)	.380 (.350-.410)	.640 (.560-.720)	1.03 (.870-1.24)	1.38 (1.09-1.91)	376
	09-10	.320 (.283-.361)	.320 (.270-.370)	.530 (.460-.630)	.830 (.710-1.10)	1.16 (.860-1.37)	451
20 years and older	99-00	.752 (.691-.818)	.800 (.700-.900)	1.40 (1.30-1.50)	2.20 (2.00-2.40)	2.90 (2.60-3.30)	1406
	01-02	.688 (.641-.738)	.700 (.700-.800)	1.20 (1.20-1.30)	2.00 (1.90-2.30)	2.80 (2.50-2.90)	1560
	03-04	.625 (.579-.674)	.620 (.560-.700)	1.04 (.960-1.11)	1.70 (1.52-1.80)	2.21 (2.04-2.49)	1543
	05-06	.574 (.539-.612)	.600 (.560-.640)	1.02 (.970-1.13)	1.65 (1.53-1.81)	2.21 (1.99-2.57)	1520
	07-08	.512 (.485-.540)	.530 (.490-.560)	.900 (.820-.950)	1.49 (1.29-1.64)	2.01 (1.78-2.33)	1857
	09-10	.486 (.465-.507)	.490 (.470-.520)	.830 (.780-.880)	1.32 (1.16-1.46)	1.71 (1.52-2.03)	2019
Gender							
Males	99-00	.923 (.822-1.04)	.900 (.900-1.00)	1.60 (1.40-1.80)	2.50 (2.20-2.90)	3.40 (2.90-3.80)	1227
	01-02	.808 (.757-.862)	.800 (.800-.900)	1.40 (1.30-1.50)	2.50 (2.20-2.70)	3.20 (2.90-3.50)	1335
	03-04	.731 (.680-.785)	.730 (.680-.800)	1.17 (1.07-1.27)	2.03 (1.78-2.22)	2.66 (2.33-2.91)	1281
	05-06	.672 (.638-.707)	.690 (.620-.760)	1.16 (1.04-1.28)	1.78 (1.60-2.00)	2.45 (2.00-2.97)	1271
	07-08	.560 (.518-.606)	.570 (.530-.620)	.930 (.840-1.02)	1.63 (1.36-1.97)	2.30 (1.97-3.07)	1327
	09-10	.527 (.491-.566)	.540 (.480-.580)	.880 (.820-.960)	1.41 (1.25-1.58)	1.83 (1.62-2.21)	1398
Females	99-00	.642 (.589-.701)	.700 (.600-.800)	1.20 (1.10-1.30)	1.90 (1.60-2.20)	2.40 (2.10-3.00)	1238
	01-02	.573 (.535-.613)	.600 (.600-.600)	1.10 (1.00-1.10)	1.60 (1.50-1.80)	2.20 (1.90-2.40)	1355
	03-04	.558 (.506-.616)	.540 (.480-.620)	.920 (.820-1.04)	1.49 (1.24-1.75)	1.82 (1.59-2.30)	1277
	05-06	.461 (.425-.499)	.460 (.430-.510)	.810 (.730-.880)	1.27 (1.17-1.46)	1.86 (1.54-2.17)	1305
	07-08	.436 (.412-.461)	.430 (.400-.480)	.760 (.690-.830)	1.22 (1.05-1.34)	1.67 (1.44-1.81)	1300
	09-10	.400 (.378-.424)	.420 (.390-.440)	.690 (.660-.740)	1.05 (.990-1.13)	1.42 (1.26-1.59)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.1, 0.1, 0.33, 0.1, 0.1, and 0.1 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Lead_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Lead_FactSheet.html

Urinary Lead

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	1.02 (.915-1.13)	1.10 (.900-1.20)	1.80 (1.60-1.90)	2.90 (2.50-3.40)	4.30 (3.10-5.40)	884
	01-02	.833 (.745-.931)	.900 (.700-1.00)	1.50 (1.20-1.70)	2.50 (2.00-2.90)	3.30 (2.70-3.80)	683
	03-04	.815 (.710-.935)	.840 (.700-.990)	1.31 (1.18-1.59)	2.19 (1.86-2.50)	2.66 (2.13-3.97)	618
	05-06	.729 (.653-.815)	.770 (.680-.860)	1.32 (1.15-1.59)	2.22 (1.81-2.64)	3.08 (2.50-4.03)	652
	07-08	.607 (.528-.698)	.610 (.530-.690)	.990 (.890-1.21)	1.76 (1.37-2.17)	2.29 (1.82-2.80)	515
	09-10	.546 (.483-.618)	.520 (.460-.580)	.930 (.800-1.09)	1.66 (1.34-1.89)	2.16 (1.80-2.87)	613
Non-Hispanic blacks	99-00	1.11 (1.00-1.23)	1.10 (1.00-1.20)	1.90 (1.50-2.10)	3.00 (2.40-3.50)	4.20 (3.30-5.70)	568
	01-02	.940 (.833-1.06)	.900 (.800-1.00)	1.60 (1.30-1.80)	2.70 (2.10-3.40)	3.70 (2.90-4.80)	667
	03-04	.848 (.729-.986)	.850 (.710-1.00)	1.40 (1.10-1.72)	2.14 (1.78-2.64)	2.82 (2.31-3.89)	723
	05-06	.666 (.604-.734)	.660 (.590-.760)	1.09 (.980-1.18)	1.62 (1.42-1.85)	2.24 (1.65-2.90)	692
	07-08	.618 (.558-.685)	.620 (.560-.690)	1.01 (.920-1.13)	1.56 (1.38-1.90)	2.06 (1.88-2.60)	589
	09-10	.560 (.522-.601)	.550 (.490-.600)	.910 (.830-1.01)	1.53 (1.26-1.71)	1.96 (1.68-2.80)	544
Non-Hispanic whites	99-00	.695 (.625-.773)	.700 (.700-.900)	1.30 (1.10-1.40)	2.00 (1.80-2.40)	2.70 (2.30-3.10)	822
	01-02	.610 (.572-.651)	.700 (.600-.700)	1.10 (1.10-1.20)	1.90 (1.70-2.00)	2.40 (2.30-2.60)	1132
	03-04	.591 (.556-.628)	.590 (.540-.650)	.960 (.910-.990)	1.52 (1.40-1.75)	2.14 (1.78-2.51)	1074
	05-06	.520 (.477-.566)	.540 (.490-.580)	.950 (.820-1.05)	1.53 (1.30-1.76)	2.07 (1.78-2.45)	1041
	07-08	.452 (.422-.485)	.460 (.430-.490)	.780 (.700-.880)	1.23 (1.03-1.50)	1.79 (1.58-1.97)	1095
	09-10	.431 (.406-.458)	.450 (.410-.480)	.750 (.690-.820)	1.11 (1.05-1.24)	1.51 (1.39-1.67)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.1, 0.1, 0.33, 0.1, 0.1, and 0.1 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Lead_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Lead_FactSheet.html

Urinary Lead (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.721 (.700-.742)	.701 (.677-.725)	1.11 (1.05-1.15)	1.70 (1.62-1.85)	2.38 (2.22-2.79)	2465
	01-02	.639 (.603-.677)	.635 (.588-.676)	1.03 (.963-1.08)	1.52 (1.43-1.61)	2.03 (1.89-2.22)	2689
	03-04	.632 (.603-.662)	.622 (.594-.655)	.979 (.920-1.03)	1.49 (1.33-1.64)	1.97 (1.73-2.26)	2558
	05-06	.546 (.520-.573)	.530 (.510-.560)	.860 (.810-.900)	1.27 (1.15-1.37)	1.71 (1.50-1.89)	2576
	07-08	.514 (.482-.548)	.500 (.460-.530)	.800 (.720-.880)	1.30 (1.23-1.42)	1.85 (1.73-1.96)	2627
	09-10	.488 (.466-.512)	.470 (.440-.500)	.760 (.710-.810)	1.16 (1.08-1.28)	1.53 (1.41-1.62)	2848
Age group							
6-11 years	99-00	1.17 (.975-1.41)	1.06 (.918-1.22)	1.55 (1.22-1.97)	2.71 (1.67-4.66)	4.66 (1.97-18.0)	340
	01-02	.918 (.841-1.00)	.870 (.800-.933)	1.27 (1.12-1.43)	2.33 (1.59-3.64)	3.64 (1.89-5.56)	368
	03-04	.926 (.812-1.06)	.914 (.781-1.03)	1.45 (1.17-1.72)	2.14 (1.62-3.47)	3.47 (2.19-5.31)	290
	05-06	.628 (.563-.701)	.590 (.530-.680)	.870 (.770-.940)	1.29 (1.03-1.82)	1.96 (1.32-2.42)	355
	07-08	.643 (.543-.763)	.630 (.530-.730)	1.02 (.770-1.24)	1.50 (1.24-2.02)	2.04 (1.70-2.58)	394
	09-10	.604 (.551-.662)	.580 (.520-.650)	.870 (.780-1.00)	1.32 (1.09-1.51)	1.60 (1.38-1.75)	378
12-19 years	99-00	.496 (.460-.535)	.469 (.408-.508)	.709 (.655-.828)	1.11 (.981-1.28)	1.65 (1.15-2.79)	719
	01-02	.404 (.380-.428)	.375 (.342-.400)	.603 (.541-.702)	.990 (.882-1.18)	1.41 (1.07-1.63)	762
	03-04	.432 (.404-.461)	.404 (.383-.436)	.623 (.551-.730)	.938 (.828-1.06)	1.23 (1.09-1.35)	725
	05-06	.363 (.333-.395)	.340 (.310-.360)	.510 (.460-.600)	.800 (.690-.930)	1.07 (.940-1.23)	701
	07-08	.302 (.270-.337)	.290 (.250-.340)	.430 (.400-.490)	.670 (.550-.790)	.900 (.700-1.09)	376
	09-10	.299 (.273-.328)	.290 (.260-.320)	.420 (.370-.490)	.620 (.550-.760)	.880 (.740-1.01)	451
20 years and older	99-00	.720 (.683-.758)	.712 (.667-.739)	1.10 (1.02-1.18)	1.69 (1.53-1.87)	2.31 (2.15-2.62)	1406
	01-02	.658 (.617-.703)	.652 (.608-.702)	1.05 (.992-1.11)	1.51 (1.40-1.61)	2.00 (1.85-2.19)	1559
	03-04	.641 (.606-.679)	.633 (.605-.670)	.988 (.917-1.04)	1.47 (1.28-1.63)	1.94 (1.72-2.12)	1543
	05-06	.573 (.548-.600)	.570 (.530-.600)	.890 (.850-.960)	1.32 (1.22-1.41)	1.77 (1.53-1.94)	1520
	07-08	.545 (.513-.579)	.530 (.490-.570)	.840 (.760-.910)	1.36 (1.27-1.49)	1.92 (1.78-2.08)	1857
	09-10	.514 (.489-.539)	.500 (.460-.530)	.790 (.730-.840)	1.22 (1.10-1.35)	1.57 (1.46-1.71)	2019
Gender							
Males	99-00	.720 (.679-.763)	.693 (.645-.734)	1.10 (.992-1.22)	1.68 (1.50-2.09)	2.43 (2.15-3.03)	1227
	01-02	.639 (.607-.673)	.638 (.586-.686)	1.01 (.957-1.08)	1.55 (1.41-1.61)	2.06 (1.88-2.43)	1334
	03-04	.615 (.588-.644)	.593 (.561-.639)	.914 (.862-.977)	1.44 (1.25-1.53)	2.00 (1.71-2.28)	1281
	05-06	.551 (.522-.582)	.530 (.510-.580)	.830 (.770-.910)	1.25 (1.13-1.39)	1.77 (1.42-2.20)	1271
	07-08	.501 (.471-.534)	.490 (.450-.530)	.750 (.700-.810)	1.29 (1.15-1.49)	1.88 (1.71-1.98)	1327
	09-10	.481 (.458-.505)	.450 (.430-.490)	.740 (.690-.800)	1.15 (1.04-1.29)	1.56 (1.37-1.70)	1398
Females	99-00	.722 (.681-.765)	.707 (.667-.746)	1.11 (1.05-1.18)	1.74 (1.50-2.02)	2.38 (2.03-2.88)	1238
	01-02	.639 (.594-.688)	.625 (.571-.682)	1.03 (.946-1.11)	1.50 (1.39-1.61)	1.98 (1.85-2.15)	1355
	03-04	.648 (.601-.698)	.649 (.604-.718)	1.03 (.938-1.10)	1.56 (1.34-1.73)	1.96 (1.72-2.20)	1277
	05-06	.541 (.507-.577)	.530 (.500-.580)	.880 (.820-.940)	1.28 (1.12-1.41)	1.64 (1.38-1.91)	1305
	07-08	.527 (.489-.568)	.500 (.470-.550)	.840 (.740-.960)	1.34 (1.23-1.47)	1.79 (1.56-2.08)	1300
	09-10	.495 (.466-.526)	.480 (.450-.520)	.780 (.710-.830)	1.20 (1.08-1.32)	1.52 (1.39-1.62)	1450

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Lead_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Lead_FactSheet.html

Urinary Lead (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	.940 (.876-1.01)	.887 (.796-1.03)	1.43 (1.37-1.58)	2.38 (2.08-2.77)	3.46 (2.78-4.18)	884
	01-02	.810 (.731-.898)	.774 (.702-.893)	1.29 (1.09-1.44)	2.05 (1.75-2.50)	2.78 (2.56-3.33)	682
	03-04	.755 (.681-.838)	.708 (.612-.851)	1.18 (1.09-1.31)	1.86 (1.50-2.26)	2.31 (1.98-2.92)	618
	05-06	.686 (.638-.737)	.680 (.620-.740)	1.00 (.930-1.13)	1.63 (1.39-1.88)	2.20 (1.77-3.20)	652
	07-08	.607 (.514-.717)	.590 (.500-.690)	.970 (.770-1.19)	1.56 (1.25-2.07)	2.20 (1.78-2.46)	515
	09-10	.573 (.526-.623)	.540 (.470-.610)	.950 (.780-1.06)	1.59 (1.41-1.82)	2.02 (1.82-2.40)	613
Non-Hispanic blacks	99-00	.722 (.659-.790)	.671 (.583-.753)	1.11 (.988-1.20)	2.00 (1.56-2.51)	2.83 (2.20-3.88)	568
	01-02	.644 (.559-.742)	.608 (.510-.710)	.962 (.853-1.20)	1.79 (1.36-2.33)	2.75 (2.04-3.98)	667
	03-04	.609 (.529-.701)	.569 (.492-.698)	.900 (.793-1.03)	1.48 (1.11-1.97)	2.24 (1.65-2.88)	723
	05-06	.483 (.459-.508)	.470 (.440-.490)	.740 (.670-.820)	1.18 (1.06-1.29)	1.60 (1.37-1.85)	692
	07-08	.452 (.414-.492)	.440 (.400-.490)	.710 (.610-.780)	1.13 (.930-1.33)	1.57 (1.21-1.88)	589
	09-10	.444 (.417-.473)	.420 (.390-.470)	.680 (.600-.740)	1.07 (.940-1.18)	1.51 (1.18-1.65)	544
Non-Hispanic whites	99-00	.696 (.668-.725)	.677 (.645-.718)	1.07 (.997-1.14)	1.66 (1.50-1.83)	2.31 (1.94-2.82)	822
	01-02	.615 (.579-.654)	.621 (.571-.667)	1.00 (.933-1.07)	1.46 (1.37-1.52)	1.88 (1.62-2.03)	1132
	03-04	.623 (.592-.655)	.618 (.587-.657)	.971 (.914-1.03)	1.44 (1.25-1.61)	1.85 (1.64-2.10)	1074
	05-06	.541 (.500-.585)	.530 (.490-.580)	.850 (.790-.920)	1.24 (1.12-1.37)	1.62 (1.37-1.94)	1041
	07-08	.506 (.466-.550)	.490 (.460-.540)	.790 (.700-.880)	1.27 (1.15-1.42)	1.81 (1.59-1.96)	1095
	09-10	.482 (.448-.518)	.460 (.430-.500)	.740 (.690-.820)	1.15 (1.01-1.30)	1.46 (1.32-1.59)	1225

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Lead_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Lead_FactSheet.html

Total Blood Mercury

Geometric mean and selected percentiles of blood concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	.797 (.703-.903)	.800 (.700-.900)	1.70 (1.50-1.90)	3.30 (2.90-3.90)	4.90 (4.30-5.50)	8373
	05-06	.863 (.787-.946)	.830 (.760-.920)	1.66 (1.48-1.93)	3.20 (2.87-3.54)	4.64 (4.17-5.25)	8407
	07-08	.769 (.689-.859)	.740 (.660-.830)	1.48 (1.29-1.69)	2.95 (2.46-3.59)	4.64 (3.74-5.79)	8266
	09-10	.863 (.792-.941)	.790 (.730-.880)	1.68 (1.49-1.91)	3.43 (3.07-3.84)	5.13 (4.57-5.67)	8793
Age group							
1-5 years	03-04	.326 (.285-.372)	.300 (.300-.300)	.500 (.500-.700)	1.00 (.800-1.60)	1.80 (1.30-2.50)	911
	05-06	*	< LOD	.500 (.470-.550)	.940 (.820-1.24)	1.43 (1.25-1.59)	968
	07-08	*	< LOD	.440 (.380-.540)	.830 (.620-1.12)	1.32 (.960-2.40)	817
	09-10	*	< LOD	.490 (.430-.590)	.890 (.740-1.08)	1.30 (1.08-1.52)	836
6-11 years	03-04	.419 (.363-.484)	.400 (.400-.500)	.700 (.700-.900)	1.30 (1.00-1.60)	1.90 (1.40-3.50)	856
	05-06	*	.410 (.330-.460)	.740 (.630-1.00)	1.43 (1.21-1.87)	2.34 (1.53-3.42)	934
	07-08	*	.380 (.340-.440)	.700 (.600-.790)	1.21 (.970-1.36)	1.56 (1.34-1.80)	1011
	09-10	*	.360 (<LOD-.400)	.670 (.590-.770)	1.22 (1.05-1.45)	1.88 (1.43-2.61)	1009
12-19 years	03-04	.490 (.418-.574)	.500 (.400-.600)	1.00 (.800-1.20)	1.80 (1.40-2.30)	2.60 (2.10-3.30)	2081
	05-06	.513 (.461-.570)	.460 (.390-.530)	.850 (.740-1.04)	1.66 (1.31-1.98)	2.41 (2.12-2.90)	1996
	07-08	.469 (.426-.516)	.440 (.390-.490)	.800 (.670-.970)	1.55 (1.30-1.72)	2.05 (1.77-2.34)	1074
	09-10	.534 (.473-.602)	.450 (.370-.540)	.910 (.770-1.11)	2.04 (1.53-2.55)	3.01 (2.53-3.63)	1183
20 years and older	03-04	.979 (.860-1.12)	1.00 (.800-1.10)	2.00 (1.70-2.30)	3.80 (3.20-4.40)	5.40 (4.60-6.70)	4525
	05-06	1.06 (.967-1.15)	1.03 (.930-1.15)	1.98 (1.73-2.22)	3.64 (3.33-4.01)	5.31 (4.82-5.67)	4509
	07-08	.944 (.833-1.07)	.890 (.780-1.03)	1.73 (1.47-2.09)	3.41 (2.82-4.17)	5.32 (4.32-6.72)	5364
	09-10	1.04 (.956-1.14)	.970 (.870-1.08)	2.00 (1.80-2.20)	3.96 (3.55-4.27)	5.75 (5.14-6.50)	5765
Gender							
Males	03-04	.814 (.714-.927)	.800 (.700-.900)	1.80 (1.50-2.00)	3.70 (3.20-4.30)	5.40 (4.60-6.50)	4132
	05-06	.864 (.783-.954)	.810 (.720-.940)	1.69 (1.48-2.01)	3.30 (2.86-3.73)	4.83 (4.08-5.45)	4092
	07-08	.809 (.709-.923)	.760 (.670-.850)	1.56 (1.31-1.81)	3.21 (2.72-4.06)	5.16 (4.12-6.97)	4147
	09-10	.883 (.810-.962)	.790 (.730-.870)	1.75 (1.54-2.02)	3.84 (3.35-4.26)	5.65 (5.13-6.34)	4366
Females	03-04	.781 (.689-.886)	.800 (.700-.900)	1.60 (1.40-1.80)	3.00 (2.50-3.50)	4.40 (3.60-5.30)	4241
	05-06	.864 (.791-.943)	.850 (.770-.920)	1.63 (1.44-1.89)	3.09 (2.75-3.46)	4.51 (4.01-5.28)	4315
	07-08	.748 (.677-.827)	.720 (.660-.810)	1.42 (1.24-1.60)	2.70 (2.27-3.27)	3.93 (3.17-5.16)	4119
	09-10	.845 (.772-.924)	.800 (.720-.880)	1.61 (1.43-1.81)	3.13 (2.76-3.48)	4.43 (4.04-5.11)	4427
Race/ethnicity							
Mexican Americans	03-04	.563 (.472-.672)	.600 (.500-.700)	1.00 (.800-1.30)	1.90 (1.60-2.40)	3.00 (2.20-3.80)	2085
	05-06	.597 (.524-.679)	.580 (.490-.670)	1.04 (.870-1.24)	1.70 (1.40-2.12)	2.58 (1.96-3.31)	2236
	07-08	.594 (.536-.658)	.580 (.520-.670)	1.03 (.900-1.17)	1.73 (1.49-2.04)	2.48 (2.10-2.91)	1712
	09-10	.613 (.571-.659)	.580 (.540-.630)	1.01 (.890-1.15)	1.63 (1.47-1.90)	2.45 (2.03-2.93)	1966
Non-Hispanic blacks	03-04	.877 (.753-1.02)	.900 (.800-1.00)	1.60 (1.40-1.80)	3.00 (2.30-4.00)	4.40 (3.30-6.00)	2293
	05-06	.823 (.697-.972)	.800 (.670-.940)	1.50 (1.21-1.92)	2.72 (2.14-3.59)	4.09 (3.22-5.16)	2193
	07-08	.766 (.711-.825)	.780 (.710-.830)	1.32 (1.23-1.42)	2.25 (1.99-2.58)	3.42 (2.74-3.90)	1746
	09-10	.928 (.805-1.07)	.900 (.800-1.02)	1.67 (1.38-1.96)	2.93 (2.20-4.21)	4.56 (3.34-6.69)	1593
Non-Hispanic whites	03-04	.776 (.655-.919)	.800 (.700-.900)	1.70 (1.40-2.00)	3.20 (2.60-3.90)	4.70 (4.00-5.60)	3478
	05-06	.891 (.801-.992)	.870 (.770-1.00)	1.74 (1.50-2.10)	3.37 (2.88-3.76)	4.76 (4.18-5.37)	3310
	07-08	.743 (.651-.847)	.720 (.620-.820)	1.43 (1.18-1.70)	2.79 (2.33-3.41)	4.18 (3.57-4.83)	3461
	09-10	.856 (.766-.957)	.790 (.690-.920)	1.70 (1.46-1.98)	3.43 (2.94-3.94)	4.92 (4.30-5.65)	3760

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.2, 0.33, 0.33, and 0.33 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mercury_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Mercury_FactSheet.html

Inorganic Blood Mercury

Geometric mean and selected percentiles of blood concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	.600 (.500-.600)	.700 (.700-.700)	8147
	05-06	*	< LOD	< LOD	.540 (.500-.580)	.660 (.620-.710)	8371
	07-08	*	< LOD	.350 (<LOD-.370)	.520 (.500-.540)	.650 (.620-.690)	8162
	09-10	*	< LOD	< LOD	.390 (.360-.430)	.510 (.480-.570)	8733
Age group							
1-5 years	03-04	*	< LOD	< LOD	< LOD	.500 (<LOD-.600)	792
	05-06	*	< LOD	< LOD	.430 (<LOD-.470)	.510 (.430-.670)	948
	07-08	*	< LOD	< LOD	.350 (<LOD-.450)	.500 (.410-.550)	726
	09-10	*	< LOD	< LOD	< LOD	.360 (<LOD-.460)	789
6-11 years	03-04	*	< LOD	< LOD	< LOD	.600 (.500-.600)	842
	05-06	*	< LOD	< LOD	.450 (<LOD-.520)	.560 (.470-.640)	932
	07-08	*	< LOD	< LOD	.380 (.350-.410)	.470 (.420-.520)	1010
	09-10	*	< LOD	< LOD	< LOD	.380 (.350-.440)	1006
12-19 years	03-04	*	< LOD	< LOD	.500 (<LOD-.500)	.600 (.500-.600)	2060
	05-06	*	< LOD	< LOD	.430 (.410-.460)	.540 (.480-.590)	1984
	07-08	*	< LOD	< LOD	.370 (<LOD-.400)	.480 (.410-.530)	1069
	09-10	*	< LOD	< LOD	< LOD	.420 (.350-.500)	1184
20 years and older	03-04	*	< LOD	< LOD	.600 (.500-.600)	.700 (.700-.800)	4453
	05-06	*	< LOD	< LOD	.570 (.530-.610)	.690 (.650-.750)	4507
	07-08	*	< LOD	.380 (.360-.390)	.550 (.530-.570)	.700 (.660-.730)	5357
	09-10	*	< LOD	< LOD	.420 (.390-.450)	.540 (.490-.600)	5754
Gender							
Males	03-04	*	< LOD	< LOD	.500 (.500-.600)	.600 (.600-.700)	4015
	05-06	*	< LOD	< LOD	.480 (.450-.520)	.600 (.550-.640)	4076
	07-08	*	< LOD	< LOD	.500 (.470-.520)	.600 (.570-.650)	4093
	09-10	*	< LOD	< LOD	.370 (<LOD-.420)	.500 (.440-.560)	4336
Females	03-04	*	< LOD	< LOD	.600 (.500-.600)	.700 (.700-.800)	4132
	05-06	*	< LOD	< LOD	.580 (.550-.640)	.700 (.670-.780)	4295
	07-08	*	< LOD	.380 (.360-.390)	.550 (.520-.570)	.700 (.670-.740)	4069
	09-10	*	< LOD	< LOD	.410 (.380-.440)	.530 (.490-.600)	4397
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	.500 (.500-.600)	.700 (.600-.800)	2007
	05-06	*	< LOD	< LOD	.530 (.470-.580)	.670 (.560-.830)	2224
	07-08	*	< LOD	< LOD	.430 (.400-.480)	.560 (.520-.610)	1685
	09-10	*	< LOD	< LOD	< LOD	.470 (.390-.530)	1947
Non-Hispanic blacks	03-04	*	< LOD	< LOD	.600 (.500-.600)	.700 (.600-.800)	2240
	05-06	*	< LOD	< LOD	.530 (.470-.600)	.670 (.600-.760)	2183
	07-08	*	< LOD	< LOD	.490 (.450-.530)	.610 (.560-.650)	1729
	09-10	*	< LOD	< LOD	.370 (.350-.390)	.480 (.410-.530)	1580
Non-Hispanic whites	03-04	*	< LOD	< LOD	.600 (.500-.600)	.700 (.600-.700)	3406
	05-06	*	< LOD	< LOD	.540 (.500-.580)	.650 (.610-.710)	3298
	07-08	*	< LOD	.360 (<LOD-.390)	.530 (.500-.550)	.660 (.620-.700)	3421
	09-10	*	< LOD	< LOD	.410 (.370-.450)	.520 (.480-.590)	3739

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.42, 0.4, 0.35, and 0.35 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mercury_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Mercury_FactSheet.html

Urinary Mercury

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	.447 (.406-.492)	.420 (.360-.480)	1.00 (.870-1.14)	2.08 (1.78-2.42)	3.19 (2.76-3.55)	2538
	05-06	.468 (.426-.514)	.460 (.410-.510)	1.03 (.900-1.12)	2.11 (1.88-2.36)	2.94 (2.58-3.26)	2578
	07-08	.443 (.408-.482)	.440 (.400-.470)	.880 (.760-1.00)	1.74 (1.62-1.96)	2.66 (2.29-3.08)	2634
	09-10	*	.400 (.360-.450)	.850 (.770-.910)	1.53 (1.30-1.81)	2.42 (2.07-2.72)	2865
Age group							
6-11 years	03-04	.254 (.213-.304)	.200 (.160-.250)	.440 (.330-.580)	1.16 (.610-1.61)	1.96 (1.13-2.97)	287
	05-06	.333 (.267-.416)	.320 (.250-.390)	.650 (.470-.840)	1.32 (.930-1.88)	2.18 (1.28-3.40)	355
	07-08	.301 (.260-.347)	.290 (.230-.340)	.520 (.430-.620)	1.03 (.770-1.23)	1.87 (1.03-3.48)	398
	09-10	*	.260 (.220-.320)	.510 (.430-.620)	1.03 (.730-1.31)	1.58 (1.18-1.88)	379
12-19 years	03-04	.358 (.313-.408)	.330 (.290-.370)	.700 (.530-.840)	1.60 (1.14-2.52)	2.93 (1.88-3.66)	722
	05-06	.372 (.286-.486)	.350 (.270-.470)	.740 (.580-.920)	1.61 (.970-2.81)	2.59 (1.40-4.45)	703
	07-08	.364 (.326-.406)	.380 (.320-.450)	.590 (.550-.650)	1.24 (.830-1.71)	1.82 (1.41-2.29)	375
	09-10	*	.290 (.230-.360)	.530 (.470-.630)	1.09 (.890-1.31)	1.73 (1.28-2.31)	455
20 years and older	03-04	.495 (.442-.555)	.480 (.410-.570)	1.12 (.930-1.29)	2.20 (1.85-2.65)	3.33 (2.76-3.88)	1529
	05-06	.505 (.468-.545)	.510 (.460-.560)	1.11 (1.04-1.16)	2.23 (1.97-2.50)	3.11 (2.64-3.37)	1520
	07-08	.477 (.435-.523)	.470 (.430-.520)	.970 (.850-1.10)	1.89 (1.69-2.20)	2.82 (2.33-3.56)	1861
	09-10	*	.450 (.390-.510)	.890 (.810-1.00)	1.66 (1.40-2.01)	2.53 (2.21-2.84)	2031
Gender							
Males	03-04	.433 (.405-.463)	.400 (.350-.460)	.940 (.840-1.05)	1.88 (1.63-2.18)	2.68 (2.34-3.05)	1266
	05-06	.464 (.411-.523)	.450 (.400-.520)	.980 (.860-1.11)	2.03 (1.57-2.48)	3.00 (2.48-3.37)	1270
	07-08	.457 (.417-.501)	.460 (.400-.520)	.880 (.780-1.01)	1.68 (1.53-1.77)	2.40 (2.11-2.76)	1326
	09-10	*	.410 (.340-.480)	.860 (.750-.950)	1.46 (1.29-1.66)	2.21 (1.93-2.53)	1404
Females	03-04	.460 (.396-.534)	.430 (.330-.530)	1.07 (.870-1.28)	2.26 (1.77-2.90)	3.54 (2.76-4.31)	1272
	05-06	.472 (.424-.525)	.470 (.390-.550)	1.07 (.900-1.19)	2.14 (1.84-2.50)	2.89 (2.60-3.38)	1308
	07-08	.431 (.388-.478)	.430 (.380-.460)	.870 (.710-1.05)	1.88 (1.55-2.38)	2.92 (2.27-4.17)	1308
	09-10	*	.390 (.360-.450)	.840 (.730-.940)	1.61 (1.29-2.03)	2.61 (2.16-3.12)	1461
Race/ethnicity							
Mexican Americans	03-04	.416 (.340-.509)	.360 (.280-.430)	.960 (.700-1.23)	2.19 (1.39-3.24)	3.16 (1.99-6.30)	619
	05-06	.451 (.369-.551)	.420 (.310-.560)	1.01 (.780-1.25)	2.22 (1.48-2.64)	3.00 (2.27-4.01)	651
	07-08	.409 (.349-.480)	.370 (.330-.450)	.780 (.700-.950)	1.82 (1.26-1.97)	2.55 (1.87-3.08)	514
	09-10	*	.350 (.280-.430)	.670 (.520-.890)	1.53 (1.06-1.84)	2.29 (1.81-2.76)	615
Non-Hispanic blacks	03-04	.476 (.413-.549)	.430 (.360-.530)	.890 (.770-1.00)	1.96 (1.60-2.31)	3.09 (2.03-4.89)	713
	05-06	.453 (.384-.533)	.450 (.380-.550)	.890 (.710-1.13)	1.78 (1.34-2.29)	2.57 (2.21-3.15)	691
	07-08	.478 (.411-.556)	.460 (.380-.540)	.910 (.770-1.06)	1.85 (1.42-2.41)	2.76 (1.97-4.19)	589
	09-10	*	.410 (.340-.490)	.840 (.650-1.08)	1.66 (1.34-1.95)	2.64 (1.88-3.30)	546
Non-Hispanic whites	03-04	.441 (.382-.509)	.420 (.330-.520)	1.01 (.840-1.23)	2.08 (1.67-2.46)	3.24 (2.67-3.60)	1066
	05-06	.459 (.409-.513)	.440 (.400-.510)	1.00 (.860-1.12)	2.07 (1.77-2.40)	2.81 (2.47-3.37)	1044
	07-08	.431 (.378-.493)	.430 (.380-.480)	.880 (.700-1.07)	1.71 (1.50-2.18)	2.70 (2.18-3.59)	1100
	09-10	*	.390 (.330-.470)	.850 (.750-.950)	1.52 (1.26-2.01)	2.42 (1.93-2.85)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.14, 0.11, 0.08, and 0.08 respectively.
 * Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mercury_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Mercury_FactSheet.html

Urinary Mercury (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	.443 (.404-.486)	.447 (.392-.498)	.909 (.785-1.00)	1.65 (1.40-1.86)	2.35 (1.88-2.85)	2537
	05-06	.460 (.414-.511)	.450 (.410-.510)	.870 (.790-1.00)	1.63 (1.44-1.75)	2.26 (2.12-2.50)	2578
	07-08	.462 (.425-.502)	.450 (.400-.490)	.820 (.750-.960)	1.57 (1.38-1.73)	2.32 (2.00-2.89)	2634
	09-10	*	.409 (.367-.459)	.793 (.691-.893)	1.43 (1.24-1.67)	2.09 (1.79-2.39)	2865
Age group							
6-11 years	03-04	.297 (.246-.358)	.276 (.208-.347)	.485 (.391-.630)	1.25 (.667-1.79)	1.79 (1.11-2.61)	286
	05-06	.411 (.323-.524)	.390 (.290-.500)	.710 (.510-.960)	1.30 (.990-2.12)	2.55 (1.38-3.50)	355
	07-08	.393 (.351-.440)	.350 (.300-.440)	.630 (.540-.770)	1.15 (.860-1.50)	1.68 (1.18-2.99)	398
	09-10	*	.357 (.306-.406)	.632 (.500-.750)	1.04 (.863-1.26)	1.62 (1.19-1.98)	379
12-19 years	03-04	.255 (.225-.289)	.217 (.196-.275)	.464 (.376-.535)	1.06 (.714-1.39)	1.67 (1.13-2.03)	722
	05-06	.286 (.230-.356)	.260 (.200-.320)	.500 (.380-.660)	1.09 (.660-1.70)	1.76 (1.11-2.67)	703
	07-08	.284 (.251-.320)	.280 (.230-.300)	.500 (.400-.550)	.890 (.620-1.08)	1.18 (.980-1.36)	375
	09-10	*	.226 (.202-.287)	.481 (.429-.553)	.917 (.736-1.18)	1.41 (1.12-1.62)	455
20 years and older	03-04	.508 (.455-.566)	.525 (.447-.616)	1.00 (.875-1.09)	1.76 (1.46-2.11)	2.54 (2.04-3.00)	1529
	05-06	.503 (.461-.549)	.510 (.470-.550)	.940 (.850-1.07)	1.69 (1.50-1.86)	2.31 (2.12-2.54)	1520
	07-08	.507 (.463-.555)	.500 (.450-.550)	.940 (.810-1.02)	1.69 (1.51-2.01)	2.56 (2.09-3.17)	1861
	09-10	*	.454 (.395-.517)	.861 (.731-.988)	1.51 (1.29-1.85)	2.15 (1.88-2.57)	2031
Gender							
Males	03-04	.365 (.333-.400)	.362 (.309-.417)	.696 (.620-.784)	1.31 (1.18-1.44)	1.87 (1.51-2.30)	1266
	05-06	.380 (.336-.431)	.390 (.330-.440)	.740 (.600-.890)	1.27 (1.09-1.47)	1.73 (1.62-1.85)	1270
	07-08	.408 (.374-.445)	.390 (.350-.450)	.730 (.650-.810)	1.22 (1.11-1.36)	1.69 (1.54-2.11)	1326
	09-10	*	.337 (.298-.391)	.675 (.585-.802)	1.19 (1.06-1.29)	1.50 (1.33-1.78)	1404
Females	03-04	.532 (.472-.599)	.545 (.455-.652)	1.06 (.969-1.21)	1.88 (1.64-2.30)	2.77 (2.12-3.56)	1271
	05-06	.552 (.494-.617)	.540 (.490-.620)	1.09 (.850-1.27)	1.96 (1.72-2.14)	2.78 (2.35-3.17)	1308
	07-08	.520 (.469-.576)	.490 (.460-.540)	.960 (.820-1.11)	1.92 (1.58-2.24)	2.83 (2.24-3.50)	1308
	09-10	*	.475 (.423-.552)	.890 (.771-1.07)	1.81 (1.43-2.09)	2.57 (2.09-2.94)	1461
Race/ethnicity							
Mexican Americans	03-04	.384 (.307-.480)	.365 (.280-.455)	.768 (.619-.990)	1.62 (1.23-2.16)	2.32 (1.78-4.01)	618
	05-06	.425 (.337-.536)	.400 (.310-.490)	.840 (.560-1.29)	1.82 (1.30-2.47)	2.63 (2.22-3.20)	651
	07-08	.409 (.350-.479)	.380 (.310-.480)	.790 (.690-.850)	1.55 (1.08-1.98)	2.03 (1.55-2.70)	514
	09-10	*	.333 (.272-.400)	.660 (.494-.861)	1.29 (1.02-1.54)	1.95 (1.52-2.89)	615
Non-Hispanic blacks	03-04	.343 (.301-.391)	.306 (.265-.368)	.587 (.522-.687)	1.28 (.964-1.63)	2.13 (1.41-2.87)	713
	05-06	.328 (.285-.378)	.320 (.270-.370)	.610 (.470-.780)	1.15 (.930-1.40)	1.64 (1.29-1.96)	691
	07-08	.350 (.303-.404)	.330 (.280-.380)	.590 (.490-.690)	1.10 (.840-1.46)	1.85 (1.13-2.77)	589
	09-10	*	.317 (.259-.393)	.582 (.500-.659)	1.05 (.900-1.30)	1.55 (1.18-1.96)	546
Non-Hispanic whites	03-04	.463 (.400-.537)	.476 (.385-.588)	.970 (.800-1.07)	1.67 (1.32-2.11)	2.40 (1.88-2.90)	1066
	05-06	.475 (.426-.531)	.490 (.440-.540)	.890 (.820-1.02)	1.61 (1.42-1.75)	2.23 (1.98-2.50)	1044
	07-08	.481 (.423-.546)	.480 (.390-.540)	.890 (.750-1.03)	1.58 (1.34-2.02)	2.49 (1.89-3.18)	1100
	09-10	*	.434 (.370-.500)	.833 (.689-1.04)	1.50 (1.26-1.87)	2.12 (1.80-2.64)	1225

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mercury_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Mercury_FactSheet.html

Urinary Molybdenum

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	45.9 (40.1-52.6)	50.7 (44.6-58.4)	84.9 (78.7-92.3)	135 (125-146)	180 (154-216)	2257
	01-02	45.0 (42.1-48.0)	52.4 (48.9-55.5)	83.4 (79.1-88.7)	124 (117-130)	165 (145-176)	2690
	03-04	39.7 (37.7-41.7)	44.5 (41.6-46.7)	78.5 (74.9-82.3)	111 (105-118)	138 (133-146)	2558
	05-06	45.0 (42.4-47.8)	51.0 (47.2-54.7)	81.5 (77.4-85.8)	127 (117-132)	158 (144-174)	2576
	07-08	45.2 (43.1-47.3)	49.7 (47.4-52.0)	86.2 (82.3-89.3)	136 (128-142)	163 (154-171)	2627
	09-10	42.7 (40.8-44.7)	45.2 (43.6-47.3)	79.9 (75.9-83.0)	121 (112-130)	160 (147-180)	2848
Age group							
6-11 years	99-00	78.2 (61.0-100)	84.8 (67.7-105)	126 (106-147)	178 (147-259)	267 (159-840)	310
	01-02	63.3 (53.4-75.0)	69.2 (63.0-77.6)	109 (94.5-124)	169 (138-197)	197 (161-291)	368
	03-04	62.2 (56.7-68.3)	71.3 (55.7-84.1)	108 (92.7-122)	138 (127-152)	181 (138-235)	290
	05-06	65.6 (56.6-76.0)	73.5 (62.8-85.5)	108 (98.6-115)	154 (130-176)	181 (154-205)	355
	07-08	69.3 (60.8-79.0)	72.8 (62.1-83.9)	123 (99.9-148)	169 (154-217)	235 (169-282)	394
	09-10	65.0 (57.8-73.2)	69.7 (61.1-84.2)	123 (103-138)	180 (146-205)	218 (180-263)	378
12-19 years	99-00	54.3 (47.6-62.0)	60.6 (52.2-70.3)	93.3 (79.9-109)	146 (112-171)	188 (146-216)	648
	01-02	60.6 (55.5-66.2)	65.7 (58.7-73.1)	97.1 (91.8-108)	145 (129-159)	179 (155-227)	762
	03-04	52.5 (49.0-56.3)	59.6 (55.5-65.1)	87.3 (84.5-91.4)	118 (105-125)	143 (130-156)	725
	05-06	59.1 (53.7-65.1)	63.8 (57.9-69.4)	96.6 (86.3-103)	136 (127-147)	173 (148-202)	701
	07-08	64.1 (58.6-70.2)	68.6 (63.7-80.2)	107 (97.2-118)	149 (125-167)	174 (151-196)	376
	09-10	52.4 (47.5-57.7)	58.5 (51.4-65.6)	96.2 (85.0-104)	148 (120-170)	178 (151-201)	451
20 years and older	99-00	41.7 (36.7-47.4)	46.6 (40.5-52.5)	76.7 (73.4-82.2)	126 (114-134)	168 (143-206)	1299
	01-02	41.1 (38.3-44.1)	47.6 (43.7-51.2)	79.1 (71.9-83.6)	114 (103-124)	150 (130-166)	1560
	03-04	35.9 (34.0-38.0)	40.3 (37.6-42.1)	71.5 (67.3-75.2)	105 (98.6-111)	133 (119-144)	1543
	05-06	41.3 (38.7-44.0)	46.0 (41.7-49.6)	75.7 (72.1-80.3)	116 (107-127)	153 (135-171)	1520
	07-08	40.8 (38.7-43.0)	44.5 (42.2-47.8)	77.7 (74.1-81.2)	123 (114-135)	152 (145-164)	1857
	09-10	39.6 (37.5-41.8)	42.0 (39.8-43.9)	73.5 (69.0-76.8)	107 (100-113)	144 (130-163)	2019
Gender							
Males	99-00	52.7 (45.7-60.7)	57.5 (48.5-68.4)	93.2 (83.8-106)	150 (128-187)	215 (161-278)	1121
	01-02	51.0 (46.6-55.7)	56.9 (52.0-62.6)	88.5 (81.6-96.5)	130 (120-141)	169 (145-194)	1335
	03-04	45.5 (43.3-47.8)	51.0 (48.1-55.4)	85.8 (82.8-90.7)	119 (112-130)	148 (136-163)	1281
	05-06	50.9 (48.2-53.9)	56.0 (51.0-60.3)	85.7 (80.3-92.8)	130 (122-136)	161 (145-177)	1271
	07-08	50.6 (48.1-53.2)	56.7 (52.4-60.6)	93.6 (88.7-98.7)	148 (137-150)	171 (159-178)	1327
	09-10	46.2 (43.5-49.1)	50.2 (46.4-52.7)	81.8 (77.0-85.9)	127 (113-144)	169 (147-197)	1398
Females	99-00	40.4 (34.8-46.8)	45.6 (40.4-52.0)	77.3 (71.0-85.7)	119 (105-138)	154 (132-180)	1136
	01-02	39.9 (37.2-42.9)	45.8 (42.8-49.3)	78.4 (72.6-82.9)	115 (104-128)	158 (130-177)	1355
	03-04	34.9 (32.2-37.7)	37.9 (33.5-41.6)	67.3 (64.6-72.2)	101 (97.3-108)	127 (114-139)	1277
	05-06	40.0 (36.4-43.8)	45.6 (40.0-51.0)	75.6 (71.2-82.0)	118 (107-128)	154 (136-179)	1305
	07-08	40.5 (38.1-43.0)	43.5 (39.0-48.0)	77.0 (72.0-84.5)	123 (114-134)	152 (142-165)	1300
	09-10	39.6 (36.9-42.4)	41.4 (39.0-43.8)	76.8 (70.7-82.0)	114 (106-127)	151 (135-170)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.8, 0.8, 1.5, 0.92, 0.92, and 0.92 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Molybdenum_BiomonitoringSummary.html

Urinary Molybdenum

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	47.0 (42.1-52.4)	53.2 (49.2-59.0)	80.6 (73.7-91.7)	121 (103-139)	152 (120-217)	780
	01-02	49.3 (46.5-52.3)	55.7 (50.4-61.0)	86.4 (80.8-94.1)	133 (113-155)	177 (142-207)	683
	03-04	47.0 (43.1-51.1)	54.4 (48.2-59.9)	82.9 (73.2-91.0)	121 (106-143)	152 (141-169)	618
	05-06	55.1 (51.7-58.7)	58.4 (54.1-63.6)	98.3 (91.5-105)	140 (134-150)	177 (153-200)	652
	07-08	53.3 (47.6-59.6)	57.3 (49.6-64.3)	90.4 (82.5-97.2)	140 (125-156)	176 (154-191)	515
	09-10	49.4 (45.9-53.2)	51.1 (46.4-55.4)	81.4 (76.2-85.9)	134 (115-149)	180 (148-231)	613
Non-Hispanic blacks	99-00	57.7 (51.0-65.2)	62.2 (55.0-71.5)	97.8 (85.0-110)	153 (126-188)	206 (150-274)	546
	01-02	53.2 (49.9-56.7)	60.3 (55.1-63.8)	90.0 (81.0-101)	132 (121-147)	166 (147-170)	667
	03-04	46.0 (41.7-50.8)	46.2 (40.9-55.2)	82.3 (73.3-91.3)	117 (109-129)	156 (135-175)	723
	05-06	48.1 (44.7-51.8)	52.0 (48.3-55.7)	80.0 (71.6-89.4)	125 (112-133)	160 (139-189)	692
	07-08	51.9 (45.6-59.1)	56.9 (47.5-65.8)	90.3 (83.0-102)	137 (118-158)	168 (147-180)	589
	09-10	45.8 (41.4-50.8)	52.1 (47.9-55.6)	85.6 (77.7-94.0)	132 (107-166)	184 (152-216)	544
Non-Hispanic whites	99-00	44.5 (37.0-53.4)	48.5 (41.1-59.8)	85.0 (76.7-96.5)	135 (119-154)	187 (146-223)	760
	01-02	42.2 (38.5-46.2)	48.9 (44.2-53.2)	80.7 (71.9-85.8)	117 (108-129)	152 (134-180)	1132
	03-04	37.1 (34.7-39.6)	41.3 (38.1-44.5)	75.2 (69.2-79.6)	107 (99.3-115)	130 (119-142)	1074
	05-06	42.8 (39.7-46.1)	48.6 (42.3-53.4)	79.0 (73.5-84.0)	123 (108-132)	156 (136-178)	1041
	07-08	41.7 (39.2-44.5)	46.5 (42.9-49.8)	81.2 (76.6-86.6)	131 (119-142)	153 (149-164)	1095
	09-10	40.6 (38.1-43.1)	42.6 (39.6-45.3)	75.9 (70.6-81.0)	114 (104-123)	148 (134-166)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.8, 0.8, 1.5, 0.92, 0.92, and 0.92 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Molybdenum_BiomonitoringSummary.html

Urinary Molybdenum (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	43.2 (40.0-46.6)	41.6 (38.5-45.2)	63.5 (59.3-68.8)	108 (97.3-115)	144 (125-171)	2257
	01-02	42.5 (39.9-45.2)	42.2 (40.1-45.2)	62.0 (58.4-66.4)	98.8 (90.1-109)	130 (120-149)	2689
	03-04	39.4 (37.6-41.3)	39.2 (37.9-40.7)	58.3 (55.6-61.5)	89.0 (80.5-99.0)	120 (107-135)	2558
	05-06	44.3 (42.4-46.3)	43.9 (41.8-45.7)	65.9 (61.8-68.6)	97.3 (92.5-103)	132 (125-143)	2576
	07-08	47.1 (45.8-48.4)	46.3 (44.8-47.7)	72.4 (68.7-74.3)	104 (96.7-111)	137 (127-152)	2627
	09-10	45.5 (43.2-47.8)	44.4 (41.9-47.4)	67.2 (64.7-69.4)	105 (95.4-121)	143 (129-157)	2848
Age group							
6-11 years	99-00	85.9 (73.7-100)	79.3 (71.6-88.4)	122 (107-133)	173 (130-243)	214 (154-1040)	310
	01-02	77.2 (73.1-81.5)	77.6 (71.8-84.5)	109 (99.4-120)	159 (129-170)	185 (165-219)	368
	03-04	72.5 (65.2-80.7)	73.5 (65.1-79.9)	101 (84.9-117)	132 (107-158)	160 (129-257)	290
	05-06	81.0 (71.9-91.3)	78.6 (72.1-89.0)	112 (93.7-134)	161 (133-199)	201 (160-261)	355
	07-08	90.4 (81.8-99.8)	88.2 (79.2-101)	133 (113-147)	206 (159-259)	274 (224-354)	394
	09-10	88.6 (81.9-95.9)	89.0 (79.2-95.4)	131 (123-148)	174 (154-191)	195 (178-216)	378
12-19 years	99-00	41.9 (39.3-44.6)	40.5 (37.7-44.4)	57.3 (51.5-62.5)	85.4 (67.4-107)	112 (78.4-185)	648
	01-02	43.4 (40.8-46.1)	44.1 (40.8-47.2)	60.8 (57.6-63.7)	85.5 (79.7-93.8)	106 (94.8-118)	762
	03-04	37.5 (35.4-39.8)	38.9 (36.9-41.8)	53.2 (50.3-56.1)	71.9 (64.6-76.9)	81.0 (74.3-102)	725
	05-06	45.5 (42.5-48.7)	45.7 (41.3-49.2)	64.6 (58.5-71.2)	91.4 (75.3-102)	109 (95.0-131)	701
	07-08	50.1 (47.2-53.2)	50.1 (44.2-53.4)	71.6 (64.0-79.6)	95.8 (85.1-107)	129 (99.5-138)	376
	09-10	49.0 (45.3-53.0)	50.7 (44.6-56.2)	67.6 (63.0-73.1)	93.6 (85.8-101)	126 (96.4-134)	451
20 years and older	99-00	39.6 (36.9-42.6)	38.5 (36.1-41.0)	56.4 (53.5-60.7)	92.5 (83.1-100)	122 (116-147)	1299
	01-02	39.3 (36.8-42.0)	39.6 (36.4-42.1)	57.2 (52.9-61.0)	86.7 (75.2-96.8)	123 (109-139)	1559
	03-04	36.9 (35.0-38.9)	37.0 (35.7-38.4)	53.5 (50.0-56.9)	79.8 (75.9-87.5)	118 (101-134)	1543
	05-06	41.2 (39.3-43.1)	40.5 (38.8-42.7)	59.1 (55.6-62.8)	87.9 (79.7-96.9)	119 (103-132)	1520
	07-08	43.5 (42.1-44.9)	42.9 (41.3-44.7)	64.9 (61.7-69.9)	94.7 (90.8-97.1)	122 (110-132)	1857
	09-10	41.9 (40.0-43.9)	41.2 (39.4-43.0)	61.9 (59.0-64.0)	90.4 (84.3-98.5)	127 (115-141)	2019
Gender							
Males	99-00	40.8 (37.5-44.3)	38.5 (37.2-40.4)	62.4 (55.9-68.4)	101 (83.9-118)	131 (112-179)	1121
	01-02	40.3 (37.1-43.8)	40.2 (36.3-43.3)	60.5 (54.8-66.3)	91.3 (83.4-106)	123 (107-155)	1334
	03-04	38.3 (36.1-40.7)	37.8 (36.1-39.3)	56.3 (53.3-59.3)	85.7 (77.2-96.7)	118 (100-139)	1281
	05-06	41.8 (39.5-44.2)	41.7 (39.2-44.1)	62.1 (58.1-67.4)	94.6 (84.2-101)	126 (112-132)	1271
	07-08	45.3 (43.8-46.8)	44.6 (42.9-46.7)	72.3 (65.4-74.8)	105 (96.0-118)	135 (126-152)	1327
	09-10	42.1 (39.6-44.9)	41.3 (38.7-44.2)	63.4 (59.4-67.4)	101 (88.3-121)	139 (123-151)	1398
Females	99-00	45.5 (41.5-50.0)	44.1 (39.5-48.8)	64.4 (59.5-70.5)	112 (95.2-121)	152 (122-181)	1136
	01-02	44.6 (42.2-47.1)	45.1 (42.2-46.9)	63.6 (59.5-69.4)	107 (92.5-119)	136 (117-169)	1355
	03-04	40.5 (38.1-43.0)	41.1 (38.7-43.7)	61.4 (56.8-65.1)	90.1 (82.2-103)	122 (115-142)	1277
	05-06	46.9 (44.7-49.2)	46.5 (43.9-48.8)	67.2 (63.3-71.7)	103 (93.6-110)	157 (126-173)	1305
	07-08	48.9 (47.2-50.7)	48.3 (45.7-51.0)	72.6 (69.2-74.4)	103 (93.6-116)	139 (113-177)	1300
	09-10	48.9 (46.4-51.5)	48.8 (45.8-51.5)	70.1 (67.2-73.9)	109 (97.1-125)	148 (137-162)	1450

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Molybdenum_BiomonitoringSummary.html

Urinary Molybdenum (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	42.9 (40.6-45.4)	43.2 (40.9-45.6)	61.6 (57.2-65.5)	89.5 (80.0-103)	115 (93.7-137)	780
	01-02	48.1 (44.3-52.2)	48.4 (44.8-52.3)	71.7 (66.4-76.0)	103 (90.0-120)	129 (109-155)	682
	03-04	43.5 (40.5-46.8)	43.5 (41.0-46.2)	62.8 (56.8-67.2)	85.9 (79.5-97.0)	112 (97.0-133)	618
	05-06	51.8 (49.9-53.8)	52.1 (49.7-54.7)	73.5 (69.5-77.4)	106 (99.5-111)	133 (112-151)	652
	07-08	53.3 (50.2-56.5)	51.9 (47.1-55.3)	77.9 (70.6-82.1)	113 (103-129)	151 (132-185)	515
	09-10	51.8 (48.9-54.9)	51.4 (48.3-54.8)	74.0 (68.7-78.8)	109 (92.7-125)	148 (125-173)	613
Non-Hispanic blacks	99-00	37.2 (33.4-41.6)	37.1 (33.0-41.2)	55.9 (49.6-63.3)	88.2 (69.1-112)	119 (88.3-141)	546
	01-02	36.5 (34.1-39.0)	37.5 (35.1-38.9)	57.1 (49.7-62.4)	78.3 (71.5-92.0)	109 (81.1-127)	667
	03-04	33.1 (30.5-35.9)	31.7 (30.1-34.7)	47.2 (43.7-52.1)	72.9 (64.6-78.4)	90.5 (78.4-118)	723
	05-06	34.9 (33.1-36.8)	35.4 (32.7-38.1)	53.5 (49.7-55.8)	81.1 (75.4-89.4)	102 (89.4-119)	692
	07-08	38.0 (34.8-41.4)	38.7 (34.2-43.0)	57.0 (52.7-60.2)	82.0 (73.2-92.4)	105 (92.0-118)	589
	09-10	36.3 (32.9-40.1)	37.2 (35.1-39.5)	54.7 (50.3-60.4)	85.2 (68.6-110)	121 (97.1-155)	544
Non-Hispanic whites	99-00	44.5 (40.2-49.2)	42.1 (38.8-47.3)	65.3 (58.9-71.3)	116 (101-126)	172 (131-195)	760
	01-02	42.5 (39.3-46.0)	41.9 (39.3-45.6)	61.2 (57.1-67.2)	104 (88.7-120)	138 (120-163)	1132
	03-04	39.1 (37.2-41.1)	39.3 (37.7-40.8)	58.1 (54.6-61.4)	87.4 (78.9-96.7)	118 (106-134)	1074
	05-06	44.5 (42.4-46.6)	43.9 (41.7-45.7)	65.0 (59.9-68.5)	97.3 (90.7-105)	132 (119-152)	1041
	07-08	46.7 (45.0-48.6)	46.1 (43.6-48.4)	72.4 (65.8-75.1)	103 (95.7-113)	135 (120-152)	1095
	09-10	45.3 (42.4-48.5)	44.2 (41.5-48.4)	66.6 (63.1-69.3)	101 (88.4-124)	141 (121-159)	1225

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Molybdenum_BiomonitoringSummary.html

Urinary Platinum

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	2465
	01-02	*	< LOD	< LOD	< LOD	< LOD	2690
	03-04	*	< LOD	< LOD	< LOD	< LOD	2558
	05-06	*	< LOD	< LOD	.009 (<LOD-.012)	.014 (.009-.025)	2576
	07-08	*	< LOD	< LOD	< LOD	< LOD	2627
	09-10	*	< LOD	< LOD	.009 (<LOD-.015)	.016 (.009-.049)	2847
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	< LOD	340
	01-02	*	< LOD	< LOD	< LOD	< LOD	368
	03-04	*	< LOD	< LOD	< LOD	< LOD	290
	05-06	*	< LOD	< LOD	.010 (<LOD-.011)	.014 (.010-.020)	355
	07-08	*	< LOD	< LOD	< LOD	.009 (<LOD-.011)	394
	09-10	*	< LOD	< LOD	.010 (<LOD-.014)	.015 (.009-.036)	378
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	719
	01-02	*	< LOD	< LOD	< LOD	< LOD	762
	03-04	*	< LOD	< LOD	< LOD	< LOD	725
	05-06	*	< LOD	< LOD	< LOD	.012 (.009-.019)	701
	07-08	*	< LOD	< LOD	< LOD	.009 (<LOD-.017)	376
	09-10	*	< LOD	< LOD	.010 (<LOD-.019)	.018 (<LOD-.078)	451
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1406
	01-02	*	< LOD	< LOD	< LOD	< LOD	1560
	03-04	*	< LOD	< LOD	< LOD	< LOD	1543
	05-06	*	< LOD	< LOD	.009 (<LOD-.014)	.014 (.009-.032)	1520
	07-08	*	< LOD	< LOD	< LOD	< LOD	1857
	09-10	*	< LOD	< LOD	.009 (<LOD-.016)	.017 (.009-.049)	2018
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	1227
	01-02	*	< LOD	< LOD	< LOD	< LOD	1335
	03-04	*	< LOD	< LOD	< LOD	< LOD	1281
	05-06	*	< LOD	< LOD	.009 (<LOD-.012)	.012 (.009-.020)	1271
	07-08	*	< LOD	< LOD	< LOD	< LOD	1327
	09-10	*	< LOD	< LOD	.009 (<LOD-.015)	.016 (<LOD-.047)	1397
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1238
	01-02	*	< LOD	< LOD	< LOD	< LOD	1355
	03-04	*	< LOD	< LOD	< LOD	< LOD	1277
	05-06	*	< LOD	< LOD	.010 (<LOD-.014)	.015 (.010-.033)	1305
	07-08	*	< LOD	< LOD	< LOD	.009 (<LOD-.009)	1300
	09-10	*	< LOD	< LOD	.010 (<LOD-.015)	.017 (.009-.051)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.04, 0.04, 0.07, 0.009, 0.009, and 0.009 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Platinum_BiomonitoringSummary.html

Urinary Platinum

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	884
	01-02	*	< LOD	< LOD	< LOD	< LOD	683
	03-04	*	< LOD	< LOD	< LOD	< LOD	618
	05-06	*	< LOD	< LOD	.010 (<LOD-.014)	.014 (.009-.023)	652
	07-08	*	< LOD	< LOD	< LOD	.009 (<LOD-.010)	515
	09-10	*	< LOD	< LOD	< LOD	.010 (.009-.012)	613
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	568
	01-02	*	< LOD	< LOD	< LOD	< LOD	667
	03-04	*	< LOD	< LOD	< LOD	< LOD	723
	05-06	*	< LOD	< LOD	< LOD	.011 (.009-.016)	692
	07-08	*	< LOD	< LOD	< LOD	.009 (<LOD-.013)	589
	09-10	*	< LOD	< LOD	.012 (<LOD-.032)	.024 (.009-.103)	544
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	822
	01-02	*	< LOD	< LOD	< LOD	< LOD	1132
	03-04	*	< LOD	< LOD	< LOD	< LOD	1074
	05-06	*	< LOD	< LOD	.009 (<LOD-.012)	.013 (.009-.026)	1041
	07-08	*	< LOD	< LOD	< LOD	.009 (<LOD-.009)	1095
	09-10	*	< LOD	< LOD	.009 (<LOD-.019)	.018 (<LOD-.073)	1224

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.04, 0.04, 0.07, 0.009, 0.009, and 0.009 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Platinum_BiomonitoringSummary.html

Urinary Platinum (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	2465
	01-02	*	< LOD	< LOD	< LOD	< LOD	2689
	03-04	*	< LOD	< LOD	< LOD	< LOD	2558
	05-06	*	< LOD	< LOD	.021 (<LOD-.023)	.029 (.026-.035)	2576
	07-08	*	< LOD	< LOD	< LOD	< LOD	2627
	09-10	*	< LOD	< LOD	.023 (<LOD-.027)	.033 (.026-.046)	2847
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	< LOD	340
	01-02	*	< LOD	< LOD	< LOD	< LOD	368
	03-04	*	< LOD	< LOD	< LOD	< LOD	290
	05-06	*	< LOD	< LOD	.020 (<LOD-.027)	.030 (.021-.050)	355
	07-08	*	< LOD	< LOD	< LOD	.026 (<LOD-.032)	394
	09-10	*	< LOD	< LOD	.026 (<LOD-.030)	.043 (.027-.050)	378
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	719
	01-02	*	< LOD	< LOD	< LOD	< LOD	762
	03-04	*	< LOD	< LOD	< LOD	< LOD	725
	05-06	*	< LOD	< LOD	< LOD	.017 (.012-.026)	701
	07-08	*	< LOD	< LOD	< LOD	.017 (<LOD-.025)	376
	09-10	*	< LOD	< LOD	.021 (<LOD-.029)	.032 (<LOD-.050)	451
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1406
	01-02	*	< LOD	< LOD	< LOD	< LOD	1559
	03-04	*	< LOD	< LOD	< LOD	< LOD	1543
	05-06	*	< LOD	< LOD	.022 (<LOD-.025)	.030 (.026-.035)	1520
	07-08	*	< LOD	< LOD	< LOD	< LOD	1857
	09-10	*	< LOD	< LOD	.023 (<LOD-.030)	.035 (.024-.050)	2018
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	1227
	01-02	*	< LOD	< LOD	< LOD	< LOD	1334
	03-04	*	< LOD	< LOD	< LOD	< LOD	1281
	05-06	*	< LOD	< LOD	.014 (<LOD-.016)	.021 (.016-.025)	1271
	07-08	*	< LOD	< LOD	< LOD	< LOD	1327
	09-10	*	< LOD	< LOD	.019 (<LOD-.025)	.027 (<LOD-.036)	1397
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1238
	01-02	*	< LOD	< LOD	< LOD	< LOD	1355
	03-04	*	< LOD	< LOD	< LOD	< LOD	1277
	05-06	*	< LOD	< LOD	.026 (<LOD-.030)	.033 (.029-.050)	1305
	07-08	*	< LOD	< LOD	< LOD	.033 (<LOD-.043)	1300
	09-10	*	< LOD	< LOD	.027 (<LOD-.032)	.038 (.030-.054)	1450

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Platinum_BiomonitoringSummary.html

Urinary Platinum (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	884
	01-02	*	< LOD	< LOD	< LOD	< LOD	682
	03-04	*	< LOD	< LOD	< LOD	< LOD	618
	05-06	*	< LOD	< LOD	.016 (<LOD-.019)	.023 (.017-.032)	652
	07-08	*	< LOD	< LOD	< LOD	.025 (<LOD-.033)	515
	09-10	*	< LOD	< LOD	< LOD	.025 (.021-.029)	613
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	568
	01-02	*	< LOD	< LOD	< LOD	< LOD	667
	03-04	*	< LOD	< LOD	< LOD	< LOD	723
	05-06	*	< LOD	< LOD	< LOD	.018 (.014-.022)	692
	07-08	*	< LOD	< LOD	< LOD	.017 (<LOD-.021)	589
	09-10	*	< LOD	< LOD	.017 (<LOD-.024)	.027 (.019-.050)	544
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	822
	01-02	*	< LOD	< LOD	< LOD	< LOD	1132
	03-04	*	< LOD	< LOD	< LOD	< LOD	1074
	05-06	*	< LOD	< LOD	.023 (<LOD-.025)	.030 (.027-.035)	1041
	07-08	*	< LOD	< LOD	< LOD	.030 (<LOD-.038)	1095
	09-10	*	< LOD	< LOD	.025 (<LOD-.032)	.036 (<LOD-.058)	1224

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Platinum_BiomonitoringSummary.html

Urinary Thallium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.176 (.162-.192)	.200 (.180-.220)	.290 (.270-.330)	.400 (.370-.420)	.450 (.430-.480)	2413
	01-02	.165 (.154-.177)	.190 (.180-.200)	.280 (.260-.290)	.370 (.350-.390)	.440 (.410-.470)	2653
	03-04	.155 (.145-.165)	.170 (.160-.180)	.270 (.250-.290)	.370 (.340-.400)	.440 (.410-.490)	2558
	05-06	.158 (.151-.165)	.180 (.170-.190)	.270 (.260-.280)	.360 (.350-.390)	.430 (.410-.460)	2576
	07-08	.146 (.139-.153)	.160 (.150-.170)	.250 (.230-.260)	.330 (.320-.360)	.400 (.390-.420)	2627
	09-10	.144 (.137-.152)	.160 (.150-.170)	.240 (.230-.250)	.340 (.310-.360)	.410 (.380-.440)	2848
Age group							
6-11 years	99-00	.201 (.167-.243)	.210 (.150-.280)	.310 (.250-.350)	.410 (.330-.450)	.450 (.350-.590)	336
	01-02	.172 (.147-.202)	.200 (.160-.220)	.290 (.230-.330)	.350 (.340-.370)	.390 (.360-.430)	362
	03-04	.191 (.170-.215)	.190 (.170-.230)	.300 (.250-.370)	.430 (.360-.500)	.510 (.430-.690)	290
	05-06	.174 (.158-.192)	.190 (.170-.210)	.280 (.250-.300)	.380 (.320-.430)	.430 (.400-.480)	355
	07-08	.166 (.150-.185)	.170 (.150-.200)	.250 (.230-.270)	.350 (.310-.410)	.420 (.360-.470)	394
	09-10	.161 (.147-.176)	.170 (.150-.200)	.280 (.230-.310)	.360 (.330-.410)	.440 (.400-.460)	378
12-19 years	99-00	.202 (.181-.225)	.220 (.200-.240)	.300 (.270-.340)	.410 (.390-.430)	.470 (.430-.510)	697
	01-02	.200 (.182-.220)	.220 (.190-.250)	.310 (.290-.320)	.370 (.350-.420)	.470 (.400-.500)	746
	03-04	.201 (.185-.218)	.220 (.210-.240)	.310 (.290-.320)	.410 (.360-.470)	.500 (.420-.560)	725
	05-06	.182 (.165-.200)	.210 (.190-.230)	.290 (.270-.310)	.370 (.350-.400)	.430 (.390-.480)	701
	07-08	.172 (.155-.190)	.180 (.160-.210)	.270 (.230-.300)	.330 (.310-.350)	.390 (.340-.420)	376
	09-10	.150 (.137-.163)	.160 (.150-.180)	.250 (.240-.270)	.340 (.290-.380)	.400 (.350-.430)	451
20 years and older	99-00	.170 (.157-.183)	.190 (.180-.210)	.290 (.260-.320)	.400 (.370-.420)	.450 (.420-.480)	1380
	01-02	.159 (.147-.173)	.190 (.170-.200)	.270 (.250-.290)	.380 (.350-.400)	.440 (.410-.490)	1545
	03-04	.145 (.134-.156)	.160 (.150-.170)	.250 (.240-.270)	.360 (.330-.390)	.420 (.390-.460)	1543
	05-06	.152 (.144-.161)	.170 (.160-.180)	.260 (.240-.270)	.360 (.340-.390)	.440 (.400-.470)	1520
	07-08	.140 (.133-.148)	.150 (.140-.160)	.240 (.230-.250)	.330 (.310-.360)	.400 (.380-.440)	1857
	09-10	.142 (.133-.150)	.160 (.150-.170)	.240 (.220-.250)	.330 (.310-.360)	.410 (.370-.440)	2019
Gender							
Males	99-00	.197 (.179-.217)	.220 (.200-.240)	.320 (.280-.350)	.400 (.370-.440)	.450 (.420-.520)	1200
	01-02	.184 (.173-.196)	.210 (.200-.230)	.290 (.280-.300)	.380 (.360-.400)	.430 (.400-.470)	1313
	03-04	.167 (.156-.178)	.190 (.180-.200)	.280 (.260-.300)	.370 (.340-.400)	.430 (.400-.480)	1281
	05-06	.171 (.163-.179)	.190 (.180-.200)	.270 (.260-.280)	.370 (.360-.390)	.430 (.410-.470)	1271
	07-08	.154 (.148-.161)	.170 (.160-.180)	.250 (.240-.270)	.340 (.320-.360)	.390 (.380-.440)	1327
	09-10	.152 (.140-.164)	.170 (.150-.180)	.250 (.240-.270)	.330 (.310-.360)	.410 (.360-.440)	1398
Females	99-00	.159 (.145-.175)	.180 (.150-.200)	.270 (.250-.300)	.390 (.350-.420)	.460 (.410-.490)	1213
	01-02	.149 (.137-.163)	.160 (.150-.180)	.260 (.230-.290)	.370 (.330-.400)	.440 (.400-.500)	1340
	03-04	.144 (.133-.156)	.160 (.140-.170)	.250 (.230-.280)	.370 (.330-.410)	.450 (.410-.510)	1277
	05-06	.146 (.138-.155)	.160 (.150-.180)	.260 (.240-.270)	.360 (.350-.380)	.440 (.400-.460)	1305
	07-08	.139 (.129-.148)	.150 (.130-.160)	.240 (.220-.260)	.330 (.300-.390)	.410 (.380-.430)	1300
	09-10	.137 (.129-.146)	.150 (.140-.160)	.230 (.210-.250)	.340 (.310-.360)	.410 (.380-.450)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.02, 0.02, 0.02, 0.015, 0.015, and 0.015 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Thallium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Thallium_FactSheet.html

Urinary Thallium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.172 (.150-.196)	.200 (.160-.230)	.270 (.250-.300)	.370 (.320-.420)	.450 (.370-.520)	861
	01-02	.160 (.148-.173)	.180 (.160-.200)	.260 (.240-.270)	.340 (.310-.360)	.400 (.350-.440)	675
	03-04	.171 (.160-.183)	.200 (.170-.220)	.280 (.260-.310)	.360 (.340-.420)	.450 (.390-.480)	618
	05-06	.158 (.149-.167)	.180 (.170-.190)	.250 (.240-.270)	.330 (.300-.360)	.400 (.360-.440)	652
	07-08	.151 (.140-.164)	.170 (.150-.180)	.250 (.220-.270)	.330 (.310-.360)	.380 (.350-.390)	515
	09-10	.147 (.139-.155)	.160 (.150-.180)	.240 (.210-.260)	.310 (.290-.320)	.390 (.320-.430)	613
Non-Hispanic blacks	99-00	.217 (.197-.239)	.230 (.220-.260)	.350 (.300-.390)	.450 (.400-.520)	.550 (.460-.630)	561
	01-02	.202 (.187-.218)	.220 (.200-.230)	.300 (.270-.340)	.410 (.380-.440)	.520 (.440-.590)	657
	03-04	.185 (.167-.206)	.190 (.170-.220)	.290 (.250-.330)	.410 (.330-.490)	.490 (.410-.640)	723
	05-06	.188 (.169-.210)	.210 (.180-.230)	.300 (.270-.320)	.390 (.360-.440)	.500 (.430-.530)	692
	07-08	.171 (.160-.182)	.180 (.170-.190)	.270 (.250-.290)	.350 (.330-.390)	.430 (.390-.460)	589
	09-10	.162 (.148-.177)	.180 (.160-.200)	.270 (.250-.280)	.350 (.330-.400)	.450 (.390-.510)	544
Non-Hispanic whites	99-00	.170 (.153-.188)	.200 (.170-.220)	.290 (.260-.330)	.400 (.360-.420)	.450 (.420-.480)	801
	01-02	.159 (.147-.172)	.180 (.170-.200)	.270 (.250-.290)	.360 (.330-.390)	.430 (.390-.460)	1114
	03-04	.146 (.135-.158)	.160 (.150-.170)	.260 (.240-.280)	.360 (.330-.380)	.410 (.380-.460)	1074
	05-06	.150 (.140-.160)	.170 (.160-.180)	.260 (.240-.270)	.350 (.320-.380)	.420 (.390-.450)	1041
	07-08	.138 (.130-.147)	.150 (.130-.160)	.240 (.220-.260)	.320 (.300-.360)	.390 (.350-.440)	1095
	09-10	.139 (.130-.149)	.150 (.140-.170)	.240 (.210-.250)	.330 (.300-.360)	.400 (.360-.430)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.02, 0.02, 0.02, 0.015, 0.015, and 0.015 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Thallium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Thallium_FactSheet.html

Urinary Thallium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.166 (.159-.173)	.168 (.162-.176)	.224 (.217-.233)	.297 (.273-.319)	.366 (.338-.387)	2413
	01-02	.156 (.151-.162)	.156 (.148-.164)	.215 (.208-.222)	.287 (.278-.300)	.349 (.337-.365)	2652
	03-04	.154 (.149-.158)	.153 (.146-.160)	.214 (.203-.222)	.286 (.274-.304)	.350 (.328-.369)	2558
	05-06	.155 (.149-.162)	.150 (.140-.160)	.210 (.200-.220)	.300 (.290-.320)	.370 (.350-.390)	2576
	07-08	.152 (.147-.158)	.150 (.140-.160)	.210 (.200-.220)	.290 (.280-.310)	.370 (.350-.380)	2627
	09-10	.153 (.145-.162)	.150 (.140-.160)	.220 (.200-.230)	.300 (.280-.320)	.370 (.350-.400)	2848
Age group							
6-11 years	99-00	.221 (.197-.248)	.222 (.196-.236)	.297 (.229-.356)	.375 (.318-.469)	.424 (.356-.600)	336
	01-02	.211 (.198-.226)	.207 (.198-.221)	.286 (.260-.321)	.370 (.333-.402)	.412 (.389-.456)	362
	03-04	.223 (.208-.238)	.216 (.198-.229)	.306 (.280-.346)	.412 (.346-.458)	.458 (.400-.532)	290
	05-06	.215 (.200-.231)	.210 (.190-.250)	.310 (.270-.320)	.360 (.320-.410)	.430 (.360-.510)	355
	07-08	.217 (.201-.234)	.220 (.210-.240)	.310 (.270-.340)	.410 (.370-.440)	.470 (.410-.540)	394
	09-10	.219 (.199-.241)	.220 (.200-.260)	.300 (.280-.340)	.400 (.350-.440)	.500 (.390-.600)	378
12-19 years	99-00	.153 (.146-.160)	.154 (.146-.162)	.205 (.191-.219)	.258 (.231-.278)	.321 (.265-.364)	697
	01-02	.143 (.137-.150)	.145 (.135-.152)	.196 (.184-.207)	.272 (.250-.289)	.312 (.299-.333)	746
	03-04	.143 (.135-.152)	.146 (.131-.155)	.194 (.179-.208)	.254 (.234-.280)	.304 (.271-.327)	725
	05-06	.140 (.133-.147)	.140 (.130-.150)	.190 (.170-.200)	.240 (.220-.270)	.290 (.250-.310)	701
	07-08	.134 (.124-.145)	.140 (.120-.150)	.190 (.180-.220)	.260 (.220-.300)	.300 (.250-.370)	376
	09-10	.140 (.127-.155)	.140 (.120-.160)	.190 (.170-.220)	.260 (.220-.310)	.310 (.260-.370)	451
20 years and older	99-00	.162 (.153-.171)	.167 (.155-.176)	.218 (.207-.230)	.286 (.271-.300)	.364 (.325-.389)	1380
	01-02	.153 (.147-.159)	.153 (.144-.161)	.210 (.200-.217)	.278 (.263-.293)	.343 (.313-.362)	1544
	03-04	.148 (.144-.153)	.149 (.141-.156)	.206 (.192-.215)	.273 (.258-.289)	.333 (.306-.353)	1543
	05-06	.152 (.145-.159)	.150 (.140-.160)	.210 (.200-.210)	.290 (.270-.310)	.370 (.350-.400)	1520
	07-08	.149 (.143-.156)	.150 (.140-.160)	.210 (.200-.220)	.280 (.270-.300)	.350 (.330-.380)	1857
	09-10	.150 (.142-.157)	.150 (.140-.160)	.210 (.200-.220)	.290 (.270-.320)	.370 (.330-.400)	2019
Gender							
Males	99-00	.154 (.147-.161)	.156 (.149-.164)	.202 (.192-.214)	.269 (.254-.297)	.338 (.300-.364)	1200
	01-02	.146 (.140-.153)	.148 (.142-.157)	.192 (.184-.204)	.260 (.246-.278)	.307 (.291-.342)	1312
	03-04	.140 (.135-.146)	.142 (.134-.149)	.188 (.180-.198)	.264 (.235-.286)	.317 (.287-.350)	1281
	05-06	.140 (.134-.147)	.140 (.130-.140)	.190 (.180-.200)	.270 (.240-.300)	.320 (.300-.340)	1271
	07-08	.138 (.131-.145)	.130 (.130-.140)	.190 (.180-.210)	.270 (.250-.290)	.330 (.300-.360)	1327
	09-10	.138 (.129-.148)	.130 (.130-.150)	.190 (.170-.210)	.270 (.240-.300)	.330 (.290-.360)	1398
Females	99-00	.178 (.167-.189)	.182 (.169-.197)	.244 (.226-.259)	.317 (.281-.366)	.380 (.333-.462)	1213
	01-02	.167 (.158-.176)	.167 (.153-.180)	.233 (.217-.250)	.313 (.282-.348)	.378 (.348-.402)	1340
	03-04	.167 (.162-.173)	.166 (.157-.177)	.235 (.222-.243)	.313 (.286-.333)	.368 (.340-.412)	1277
	05-06	.171 (.162-.181)	.170 (.160-.180)	.230 (.220-.250)	.330 (.300-.360)	.430 (.370-.490)	1305
	07-08	.167 (.163-.173)	.170 (.160-.180)	.230 (.230-.240)	.320 (.300-.340)	.390 (.360-.430)	1300
	09-10	.170 (.161-.178)	.170 (.160-.180)	.230 (.220-.250)	.340 (.310-.370)	.420 (.370-.470)	1450

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Thallium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Thallium_FactSheet.html

Urinary Thallium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.158 (.147-.170)	.160 (.148-.176)	.213 (.200-.237)	.282 (.266-.304)	.343 (.306-.389)	861
	01-02	.156 (.145-.169)	.155 (.145-.167)	.204 (.191-.221)	.286 (.250-.317)	.361 (.301-.424)	674
	03-04	.159 (.148-.170)	.157 (.143-.172)	.211 (.187-.241)	.293 (.273-.324)	.369 (.326-.422)	618
	05-06	.149 (.139-.158)	.150 (.140-.160)	.200 (.180-.210)	.270 (.240-.290)	.320 (.280-.370)	652
	07-08	.151 (.143-.160)	.160 (.130-.170)	.200 (.190-.220)	.270 (.250-.280)	.320 (.290-.370)	515
	09-10	.154 (.141-.167)	.150 (.140-.170)	.210 (.180-.230)	.280 (.250-.330)	.360 (.310-.430)	613
Non-Hispanic blacks	99-00	.142 (.133-.152)	.140 (.129-.151)	.200 (.184-.214)	.278 (.244-.307)	.383 (.286-.462)	561
	01-02	.138 (.128-.150)	.136 (.125-.146)	.194 (.170-.212)	.256 (.238-.278)	.328 (.271-.387)	657
	03-04	.133 (.122-.145)	.128 (.119-.143)	.185 (.171-.200)	.255 (.237-.269)	.323 (.267-.377)	723
	05-06	.137 (.128-.145)	.130 (.130-.140)	.190 (.180-.200)	.240 (.220-.290)	.320 (.270-.370)	692
	07-08	.125 (.118-.132)	.130 (.120-.130)	.170 (.160-.180)	.230 (.210-.260)	.280 (.250-.330)	589
	09-10	.128 (.120-.137)	.120 (.110-.130)	.180 (.160-.210)	.250 (.230-.280)	.310 (.270-.350)	544
Non-Hispanic whites	99-00	.169 (.160-.179)	.173 (.167-.181)	.227 (.215-.240)	.300 (.272-.329)	.364 (.333-.377)	801
	01-02	.161 (.155-.167)	.161 (.153-.171)	.222 (.214-.231)	.292 (.278-.304)	.348 (.330-.383)	1114
	03-04	.154 (.148-.160)	.153 (.143-.162)	.214 (.200-.223)	.283 (.271-.304)	.333 (.313-.363)	1074
	05-06	.156 (.148-.164)	.150 (.140-.160)	.210 (.200-.230)	.300 (.280-.320)	.360 (.330-.400)	1041
	07-08	.154 (.146-.164)	.160 (.140-.170)	.220 (.210-.230)	.300 (.280-.320)	.370 (.340-.390)	1095
	09-10	.155 (.145-.166)	.150 (.140-.170)	.220 (.200-.240)	.310 (.280-.330)	.380 (.340-.410)	1225

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Thallium_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Thallium_FactSheet.html

Urinary Tungsten

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.093 (.087-.100)	.090 (.080-.090)	.180 (.160-.190)	.320 (.280-.370)	.500 (.430-.550)	2338
	01-02	.082 (.073-.092)	.070 (.060-.090)	.160 (.140-.180)	.300 (.260-.350)	.460 (.370-.560)	2652
	03-04	.071 (.064-.078)	.070 (.060-.080)	.130 (.120-.140)	.270 (.230-.300)	.400 (.330-.480)	2558
	05-06	.085 (.079-.092)	.090 (.080-.090)	.170 (.150-.180)	.310 (.280-.340)	.470 (.400-.520)	2576
	07-08	.099 (.093-.105)	.100 (.090-.110)	.190 (.180-.200)	.360 (.320-.380)	.500 (.440-.570)	2627
	09-10	.081 (.075-.087)	.080 (.070-.090)	.160 (.150-.170)	.310 (.270-.330)	.460 (.370-.540)	2847
Age group							
6-11 years	99-00	.158 (.123-.204)	.160 (.120-.220)	.270 (.220-.350)	.510 (.380-.560)	.620 (.510-.950)	320
	01-02	.137 (.110-.170)	.140 (.110-.170)	.260 (.200-.350)	.460 (.360-.690)	.770 (.510-1.53)	363
	03-04	.130 (.111-.151)	.140 (.120-.150)	.240 (.190-.290)	.410 (.290-.500)	.500 (.370-.630)	290
	05-06	.142 (.121-.166)	.140 (.120-.170)	.230 (.210-.290)	.430 (.330-.610)	.640 (.440-.850)	355
	07-08	.160 (.138-.186)	.150 (.130-.170)	.310 (.240-.390)	.510 (.450-.580)	.810 (.560-1.15)	394
	09-10	.151 (.126-.180)	.150 (.110-.200)	.310 (.230-.400)	.560 (.400-.770)	.770 (.560-1.04)	378
12-19 years	99-00	.113 (.097-.132)	.110 (.090-.130)	.210 (.180-.250)	.360 (.310-.440)	.530 (.380-.800)	679
	01-02	.113 (.095-.135)	.110 (.090-.130)	.210 (.180-.260)	.400 (.310-.520)	.570 (.430-.790)	744
	03-04	.105 (.090-.122)	.100 (.090-.120)	.190 (.160-.230)	.350 (.290-.460)	.530 (.350-1.00)	725
	05-06	.131 (.108-.158)	.130 (.100-.160)	.250 (.200-.310)	.430 (.360-.530)	.620 (.410-1.04)	701
	07-08	.150 (.136-.167)	.160 (.140-.180)	.250 (.220-.290)	.470 (.350-.660)	.660 (.430-1.05)	376
	09-10	.101 (.089-.115)	.100 (.080-.120)	.210 (.190-.240)	.360 (.300-.460)	.540 (.440-.650)	451
20 years and older	99-00	.084 (.078-.091)	.080 (.070-.090)	.160 (.130-.180)	.280 (.260-.320)	.450 (.360-.520)	1339
	01-02	.073 (.065-.082)	.060 (.050-.070)	.140 (.110-.160)	.260 (.210-.310)	.380 (.310-.490)	1545
	03-04	.062 (.056-.068)	.060 (.050-.070)	.110 (.100-.120)	.210 (.180-.250)	.360 (.270-.430)	1543
	05-06	.075 (.070-.081)	.080 (.070-.080)	.140 (.130-.160)	.250 (.220-.310)	.410 (.340-.470)	1520
	07-08	.088 (.081-.095)	.090 (.080-.100)	.170 (.150-.190)	.310 (.280-.340)	.450 (.390-.500)	1857
	09-10	.073 (.068-.079)	.080 (.070-.080)	.150 (.130-.160)	.260 (.230-.300)	.370 (.330-.460)	2018
Gender							
Males	99-00	.107 (.096-.120)	.100 (.090-.120)	.210 (.190-.230)	.390 (.310-.470)	.530 (.470-.650)	1160
	01-02	.088 (.074-.105)	.080 (.060-.100)	.170 (.140-.220)	.330 (.260-.390)	.490 (.380-.580)	1307
	03-04	.081 (.071-.093)	.080 (.070-.090)	.140 (.130-.170)	.300 (.250-.340)	.430 (.340-.560)	1281
	05-06	.095 (.090-.101)	.090 (.080-.100)	.180 (.160-.190)	.340 (.310-.380)	.480 (.420-.610)	1271
	07-08	.109 (.102-.116)	.110 (.100-.130)	.210 (.190-.240)	.380 (.350-.450)	.550 (.450-.660)	1327
	09-10	.090 (.082-.098)	.090 (.080-.100)	.170 (.150-.190)	.320 (.290-.350)	.500 (.410-.550)	1398
Females	99-00	.082 (.077-.087)	.070 (.060-.080)	.150 (.130-.160)	.270 (.240-.300)	.400 (.320-.470)	1178
	01-02	.076 (.069-.084)	.060 (.060-.080)	.150 (.120-.170)	.280 (.230-.330)	.430 (.340-.560)	1345
	03-04	.062 (.056-.069)	.060 (.050-.070)	.110 (.100-.120)	.220 (.190-.250)	.370 (.270-.460)	1277
	05-06	.077 (.069-.085)	.080 (.070-.090)	.160 (.150-.170)	.280 (.250-.320)	.440 (.350-.510)	1305
	07-08	.090 (.083-.098)	.090 (.080-.100)	.170 (.160-.180)	.300 (.270-.350)	.460 (.420-.550)	1300
	09-10	.074 (.067-.081)	.070 (.070-.080)	.150 (.130-.170)	.280 (.240-.330)	.400 (.320-.600)	1449

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.04, 0.04, 0.04, 0.021, 0.021, and 0.021 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Tungsten_BiomonitoringSummary.html

Urinary Tungsten

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.113 (.095-.133)	.110 (.090-.130)	.200 (.160-.250)	.400 (.300-.520)	.550 (.420-.830)	790
	01-02	.101 (.093-.109)	.100 (.090-.110)	.190 (.170-.210)	.370 (.310-.430)	.570 (.450-.670)	680
	03-04	.086 (.073-.100)	.080 (.070-.100)	.170 (.120-.220)	.360 (.230-.620)	.640 (.410-.800)	618
	05-06	.102 (.088-.118)	.100 (.080-.110)	.200 (.170-.220)	.450 (.310-.650)	.730 (.510-1.04)	652
	07-08	.106 (.084-.135)	.110 (.080-.150)	.210 (.170-.270)	.400 (.250-.670)	.580 (.400-1.12)	515
	09-10	.099 (.089-.109)	.090 (.090-.100)	.200 (.170-.220)	.380 (.340-.470)	.540 (.480-.580)	613
Non-Hispanic blacks	99-00	.113 (.101-.126)	.100 (.090-.130)	.210 (.180-.250)	.370 (.290-.460)	.560 (.420-.810)	562
	01-02	.096 (.080-.116)	.090 (.070-.120)	.160 (.130-.250)	.310 (.270-.400)	.460 (.400-.590)	649
	03-04	.092 (.082-.104)	.090 (.080-.110)	.160 (.150-.180)	.300 (.250-.330)	.470 (.340-.550)	723
	05-06	.101 (.088-.116)	.110 (.090-.120)	.190 (.160-.220)	.320 (.280-.370)	.450 (.340-.650)	692
	07-08	.120 (.099-.146)	.120 (.090-.160)	.240 (.200-.290)	.390 (.340-.440)	.540 (.460-.690)	589
	09-10	.093 (.086-.100)	.100 (.090-.110)	.180 (.160-.200)	.290 (.270-.340)	.400 (.320-.590)	543
Non-Hispanic whites	99-00	.092 (.084-.100)	.080 (.070-.100)	.180 (.160-.200)	.320 (.270-.380)	.470 (.380-.550)	802
	01-02	.076 (.066-.088)	.060 (.050-.080)	.150 (.120-.180)	.290 (.230-.360)	.430 (.330-.620)	1117
	03-04	.065 (.058-.073)	.060 (.060-.070)	.120 (.100-.130)	.230 (.190-.290)	.380 (.320-.410)	1074
	05-06	.082 (.075-.089)	.080 (.080-.090)	.160 (.140-.180)	.300 (.250-.340)	.430 (.380-.480)	1041
	07-08	.093 (.087-.101)	.100 (.090-.110)	.180 (.160-.190)	.330 (.290-.370)	.470 (.410-.570)	1095
	09-10	.077 (.070-.083)	.080 (.070-.090)	.150 (.140-.170)	.280 (.250-.310)	.400 (.330-.540)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.04, 0.04, 0.04, 0.021, 0.021, and 0.021 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Tungsten_BiomonitoringSummary.html

Urinary Tungsten (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.087 (.080-.095)	.080 (.075-.086)	.146 (.136-.158)	.270 (.206-.333)	.383 (.302-.459)	2338
	01-02	.078 (.069-.087)	.074 (.064-.084)	.138 (.122-.154)	.255 (.216-.300)	.359 (.315-.436)	2651
	03-04	.070 (.063-.078)	.065 (.059-.074)	.117 (.107-.133)	.215 (.179-.253)	.333 (.255-.439)	2558
	05-06	.084 (.080-.089)	.080 (.080-.080)	.140 (.130-.150)	.250 (.210-.290)	.350 (.290-.440)	2576
	07-08	.103 (.096-.111)	.100 (.090-.110)	.170 (.160-.190)	.320 (.290-.360)	.470 (.410-.540)	2627
	09-10	.086 (.080-.093)	.080 (.070-.090)	.150 (.130-.160)	.270 (.230-.310)	.400 (.340-.470)	2847
Age group							
6-11 years	99-00	.174 (.150-.201)	.169 (.136-.198)	.293 (.216-.333)	.439 (.331-.667)	.667 (.452-.880)	320
	01-02	.168 (.144-.197)	.158 (.139-.190)	.275 (.231-.326)	.412 (.333-.554)	.634 (.436-1.28)	363
	03-04	.151 (.131-.174)	.144 (.119-.167)	.250 (.205-.283)	.333 (.278-.484)	.484 (.333-.739)	290
	05-06	.175 (.152-.202)	.160 (.140-.180)	.270 (.220-.320)	.560 (.330-.790)	.790 (.470-1.18)	355
	07-08	.209 (.185-.235)	.200 (.180-.240)	.340 (.290-.370)	.540 (.450-.660)	.780 (.620-.910)	394
	09-10	.205 (.177-.239)	.200 (.160-.230)	.330 (.290-.440)	.620 (.450-.850)	.930 (.690-1.08)	378
12-19 years	99-00	.084 (.078-.091)	.079 (.074-.084)	.138 (.124-.158)	.231 (.180-.287)	.339 (.237-.465)	679
	01-02	.081 (.071-.092)	.081 (.072-.091)	.148 (.122-.167)	.250 (.208-.301)	.359 (.272-.431)	744
	03-04	.075 (.065-.086)	.071 (.061-.082)	.122 (.098-.148)	.197 (.167-.308)	.379 (.197-.582)	725
	05-06	.101 (.086-.117)	.100 (.080-.110)	.170 (.140-.200)	.300 (.200-.370)	.370 (.300-.770)	701
	07-08	.118 (.104-.132)	.120 (.110-.130)	.180 (.160-.210)	.340 (.230-.550)	.530 (.310-.810)	376
	09-10	.094 (.085-.106)	.090 (.080-.110)	.150 (.120-.180)	.270 (.210-.290)	.370 (.280-.500)	451
20 years and older	99-00	.080 (.072-.089)	.075 (.067-.082)	.130 (.116-.146)	.218 (.179-.301)	.347 (.245-.426)	1339
	01-02	.070 (.063-.079)	.067 (.058-.075)	.119 (.099-.139)	.216 (.176-.267)	.333 (.253-.431)	1544
	03-04	.063 (.057-.071)	.059 (.053-.065)	.105 (.094-.117)	.181 (.155-.215)	.279 (.217-.370)	1543
	05-06	.075 (.071-.079)	.070 (.070-.080)	.120 (.110-.130)	.210 (.200-.220)	.290 (.240-.360)	1520
	07-08	.093 (.087-.101)	.090 (.080-.100)	.160 (.140-.170)	.280 (.250-.310)	.420 (.350-.490)	1857
	09-10	.078 (.072-.084)	.070 (.070-.080)	.130 (.120-.140)	.220 (.190-.260)	.340 (.270-.400)	2018
Gender							
Males	99-00	.083 (.074-.094)	.073 (.063-.086)	.146 (.126-.165)	.279 (.198-.386)	.439 (.329-.605)	1160
	01-02	.071 (.060-.083)	.065 (.056-.077)	.125 (.098-.153)	.255 (.203-.306)	.364 (.300-.431)	1306
	03-04	.068 (.059-.079)	.062 (.054-.071)	.111 (.098-.133)	.216 (.170-.284)	.341 (.240-.500)	1281
	05-06	.078 (.073-.083)	.070 (.070-.080)	.130 (.120-.150)	.240 (.200-.300)	.360 (.300-.440)	1271
	07-08	.097 (.090-.106)	.090 (.090-.110)	.170 (.150-.180)	.310 (.280-.360)	.440 (.410-.480)	1327
	09-10	.082 (.074-.089)	.070 (.070-.080)	.140 (.120-.150)	.250 (.220-.310)	.440 (.320-.500)	1398
Females	99-00	.091 (.085-.098)	.084 (.080-.091)	.145 (.136-.158)	.265 (.200-.301)	.339 (.300-.381)	1178
	01-02	.085 (.077-.094)	.083 (.075-.091)	.143 (.130-.161)	.258 (.216-.317)	.353 (.317-.538)	1345
	03-04	.072 (.065-.079)	.069 (.063-.078)	.121 (.108-.138)	.211 (.176-.237)	.333 (.261-.439)	1277
	05-06	.090 (.085-.096)	.080 (.080-.090)	.150 (.140-.160)	.250 (.210-.290)	.350 (.260-.530)	1305
	07-08	.109 (.100-.118)	.100 (.090-.120)	.180 (.160-.210)	.330 (.290-.360)	.520 (.400-.590)	1300
	09-10	.091 (.083-.100)	.090 (.080-.090)	.150 (.130-.170)	.270 (.210-.340)	.380 (.330-.450)	1449

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Tungsten_BiomonitoringSummary.html

Urinary Tungsten (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	.106 (.093-.120)	.100 (.086-.116)	.184 (.152-.214)	.329 (.267-.392)	.497 (.354-.727)	790
	01-02	.098 (.090-.108)	.089 (.081-.100)	.164 (.143-.187)	.294 (.258-.375)	.555 (.410-.797)	679
	03-04	.079 (.065-.096)	.073 (.054-.093)	.136 (.103-.197)	.300 (.233-.426)	.482 (.344-.823)	618
	05-06	.096 (.083-.111)	.090 (.080-.090)	.160 (.150-.180)	.330 (.250-.420)	.670 (.360-.890)	652
	07-08	.106 (.083-.137)	.110 (.080-.140)	.200 (.160-.270)	.400 (.260-.710)	.610 (.310-1.12)	515
	09-10	.103 (.096-.111)	.100 (.090-.110)	.180 (.160-.200)	.330 (.290-.390)	.480 (.390-.540)	613
Non-Hispanic blacks	99-00	.073 (.064-.083)	.071 (.061-.081)	.124 (.109-.154)	.201 (.188-.222)	.360 (.217-.465)	562
	01-02	.066 (.056-.077)	.060 (.049-.079)	.109 (.090-.125)	.199 (.153-.285)	.340 (.250-.414)	649
	03-04	.066 (.059-.074)	.067 (.055-.075)	.105 (.095-.120)	.186 (.150-.224)	.317 (.214-.358)	723
	05-06	.073 (.065-.082)	.070 (.060-.080)	.120 (.110-.140)	.210 (.190-.260)	.320 (.260-.390)	692
	07-08	.088 (.076-.102)	.090 (.070-.100)	.150 (.140-.160)	.240 (.220-.300)	.380 (.270-.630)	589
	09-10	.074 (.067-.080)	.070 (.060-.080)	.120 (.110-.140)	.210 (.180-.250)	.310 (.220-.410)	543
Non-Hispanic whites	99-00	.091 (.083-.100)	.082 (.077-.088)	.150 (.136-.169)	.279 (.200-.354)	.385 (.302-.462)	802
	01-02	.078 (.068-.088)	.073 (.061-.085)	.139 (.121-.157)	.253 (.209-.308)	.353 (.286-.453)	1117
	03-04	.069 (.060-.078)	.063 (.057-.071)	.116 (.104-.133)	.199 (.167-.237)	.299 (.222-.439)	1074
	05-06	.085 (.080-.090)	.080 (.080-.090)	.140 (.130-.160)	.240 (.210-.280)	.340 (.280-.420)	1041
	07-08	.104 (.096-.114)	.100 (.090-.110)	.180 (.160-.200)	.310 (.280-.360)	.460 (.390-.530)	1095
	09-10	.086 (.078-.094)	.080 (.070-.090)	.140 (.130-.160)	.250 (.200-.310)	.370 (.330-.400)	1225

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Tungsten_BiomonitoringSummary.html

Urinary Uranium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.008 (.007-.009)	.007 (.006-.008)	.013 (.010-.017)	.027 (.021-.038)	.046 (.037-.056)	2464
	01-02	.009 (.007-.010)	.008 (.007-.009)	.014 (.012-.018)	.030 (.023-.039)	.046 (.034-.062)	2690
	03-04	.008 (.007-.008)	.007 (.006-.007)	.011 (.010-.013)	.021 (.017-.026)	.031 (.026-.037)	2557
	05-06	.006 (.005-.006)	.005 (.005-.006)	.010 (.009-.012)	.019 (.016-.022)	.033 (.023-.041)	2576
	07-08	.007 (.006-.008)	.007 (.005-.008)	.013 (.011-.015)	.024 (.018-.033)	.039 (.026-.057)	2627
	09-10	.007 (.006-.008)	.007 (.006-.008)	.013 (.011-.015)	.022 (.019-.030)	.036 (.029-.044)	2848
Age group							
6-11 years	99-00	.009 (.007-.011)	.007 (.006-.009)	.013 (.009-.022)	.032 (.019-.048)	.048 (.033-.066)	340
	01-02	.008 (.007-.010)	.008 (.006-.010)	.014 (.010-.020)	.026 (.020-.036)	.040 (.025-.049)	368
	03-04	.008 (.007-.009)	.007 (.006-.009)	.012 (.009-.016)	.020 (.016-.026)	.028 (.020-.039)	289
	05-06	.006 (.005-.007)	.005 (.004-.007)	.010 (.008-.011)	.015 (.012-.031)	.031 (.013-.051)	355
	07-08	.007 (.006-.008)	.006 (.005-.008)	.012 (.009-.016)	.021 (.016-.027)	.030 (.022-.039)	394
	09-10	.006 (.005-.008)	.006 (.005-.008)	.012 (.009-.014)	.018 (.013-.034)	.033 (.016-.052)	378
12-19 years	99-00	.009 (.008-.011)	.009 (.008-.010)	.015 (.012-.018)	.026 (.020-.043)	.044 (.028-.072)	719
	01-02	.010 (.008-.012)	.010 (.008-.012)	.017 (.013-.023)	.030 (.022-.042)	.042 (.027-.088)	762
	03-04	.010 (.009-.011)	.009 (.008-.010)	.015 (.012-.018)	.028 (.023-.036)	.038 (.036-.053)	725
	05-06	.007 (.006-.008)	.007 (.006-.008)	.013 (.011-.015)	.023 (.018-.032)	.034 (.027-.045)	701
	07-08	.009 (.007-.011)	.008 (.007-.011)	.016 (.014-.020)	.029 (.022-.056)	.056 (.027-.156)	376
	09-10	.008 (.007-.010)	.008 (.006-.010)	.014 (.011-.018)	.023 (.020-.038)	.040 (.024-.059)	451
20 years and older	99-00	.008 (.006-.009)	.007 (.005-.008)	.013 (.010-.017)	.027 (.021-.040)	.046 (.036-.056)	1405
	01-02	.009 (.007-.010)	.008 (.007-.009)	.014 (.012-.017)	.031 (.022-.040)	.046 (.034-.065)	1560
	03-04	*	.006 (.005-.007)	.011 (.009-.012)	.019 (.016-.026)	.029 (.024-.038)	1543
	05-06	.006 (.005-.006)	.005 (.005-.006)	.010 (.008-.012)	.019 (.015-.022)	.032 (.022-.041)	1520
	07-08	.007 (.005-.008)	.006 (.005-.008)	.013 (.010-.015)	.024 (.017-.035)	.039 (.026-.052)	1857
	09-10	.007 (.006-.008)	.007 (.006-.008)	.013 (.011-.015)	.023 (.019-.030)	.036 (.029-.044)	2019
Gender							
Males	99-00	.009 (.008-.011)	.008 (.007-.010)	.015 (.012-.021)	.036 (.024-.046)	.053 (.040-.067)	1227
	01-02	.009 (.008-.011)	.009 (.007-.010)	.015 (.013-.021)	.033 (.024-.045)	.047 (.035-.065)	1335
	03-04	.008 (.007-.009)	.007 (.006-.008)	.013 (.011-.016)	.023 (.019-.027)	.031 (.027-.035)	1280
	05-06	.006 (.006-.007)	.006 (.005-.006)	.011 (.009-.012)	.019 (.015-.022)	.030 (.021-.043)	1271
	07-08	.007 (.006-.009)	.007 (.006-.009)	.014 (.013-.016)	.026 (.021-.037)	.046 (.030-.056)	1327
	09-10	.007 (.006-.009)	.007 (.006-.009)	.014 (.012-.017)	.024 (.020-.034)	.040 (.032-.046)	1398
Females	99-00	.007 (.006-.008)	.006 (.005-.007)	.012 (.009-.015)	.023 (.016-.033)	.036 (.026-.050)	1237
	01-02	.008 (.007-.010)	.008 (.006-.009)	.014 (.011-.017)	.027 (.019-.037)	.041 (.029-.063)	1355
	03-04	*	.006 (.005-.007)	.010 (.009-.011)	.018 (.013-.027)	.031 (.022-.039)	1277
	05-06	.005 (.005-.006)	.005 (.004-.006)	.010 (.008-.011)	.019 (.016-.023)	.034 (.025-.040)	1305
	07-08	.006 (.005-.008)	.006 (.005-.007)	.011 (.009-.015)	.024 (.016-.033)	.035 (.022-.067)	1300
	09-10	.007 (.006-.008)	.007 (.006-.008)	.012 (.011-.014)	.022 (.017-.029)	.033 (.023-.045)	1450

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.004, 0.004, 0.005, 0.002, 0.002, and 0.002 respectively.
 * Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Uranium_BiomonitoringSummary.html

Urinary Uranium

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.017 (.012-.023)	.016 (.011-.021)	.033 (.020-.054)	.060 (.040-.127)	.114 (.054-.279)	883
	01-02	.013 (.010-.016)	.012 (.009-.016)	.022 (.017-.030)	.040 (.031-.054)	.055 (.046-.069)	683
	03-04	.014 (.011-.017)	.013 (.009-.018)	.024 (.017-.034)	.041 (.028-.073)	.064 (.039-.158)	618
	05-06	.008 (.007-.009)	.009 (.007-.010)	.014 (.013-.017)	.025 (.021-.033)	.042 (.028-.051)	652
	07-08	.009 (.008-.011)	.009 (.008-.010)	.017 (.014-.022)	.032 (.026-.039)	.047 (.032-.073)	515
	09-10	.009 (.008-.011)	.009 (.008-.011)	.018 (.015-.021)	.034 (.027-.040)	.046 (.039-.055)	613
Non-Hispanic blacks	99-00	.009 (.007-.011)	.008 (.006-.010)	.014 (.010-.020)	.028 (.018-.049)	.052 (.030-.067)	568
	01-02	.008 (.007-.009)	.008 (.007-.009)	.012 (.011-.015)	.021 (.017-.027)	.030 (.023-.037)	667
	03-04	.008 (.008-.009)	.007 (.007-.008)	.012 (.011-.013)	.021 (.017-.027)	.031 (.023-.045)	722
	05-06	.006 (.005-.007)	.006 (.005-.007)	.010 (.009-.011)	.016 (.014-.020)	.023 (.018-.031)	692
	07-08	.007 (.006-.009)	.007 (.006-.008)	.013 (.010-.016)	.024 (.016-.038)	.038 (.025-.055)	589
	09-10	.008 (.007-.010)	.008 (.007-.010)	.014 (.012-.016)	.027 (.017-.044)	.044 (.027-.074)	544
Non-Hispanic whites	99-00	.007 (.006-.009)	.007 (.006-.007)	.012 (.009-.016)	.023 (.017-.037)	.043 (.027-.051)	822
	01-02	.008 (.007-.009)	.007 (.006-.009)	.013 (.011-.016)	.026 (.019-.035)	.037 (.029-.050)	1132
	03-04	*	.006 (.005-.007)	.010 (.009-.012)	.018 (.015-.023)	.027 (.020-.036)	1074
	05-06	.005 (.005-.006)	.005 (.004-.006)	.010 (.008-.012)	.018 (.014-.022)	.033 (.021-.043)	1041
	07-08	.006 (.005-.008)	.006 (.005-.008)	.013 (.009-.016)	.023 (.015-.039)	.038 (.022-.086)	1095
	09-10	.006 (.005-.007)	.006 (.005-.007)	.012 (.011-.014)	.021 (.017-.027)	.032 (.023-.041)	1225

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.004, 0.004, 0.005, 0.002, 0.002, and 0.002 respectively.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Uranium_BiomonitoringSummary.html

Urinary Uranium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.007 (.006-.009)	.007 (.006-.009)	.013 (.010-.016)	.024 (.019-.030)	.034 (.027-.053)	2464
	01-02	.008 (.007-.010)	.007 (.006-.009)	.014 (.011-.018)	.026 (.020-.034)	.040 (.028-.058)	2689
	03-04	.008 (.007-.008)	.007 (.006-.008)	.012 (.010-.014)	.021 (.017-.025)	.029 (.023-.039)	2557
	05-06	.006 (.005-.006)	.005 (.005-.006)	.009 (.008-.010)	.017 (.014-.020)	.026 (.020-.039)	2576
	07-08	.007 (.006-.009)	.006 (.005-.008)	.012 (.009-.016)	.024 (.016-.038)	.038 (.025-.065)	2627
	09-10	.007 (.006-.008)	.007 (.006-.008)	.012 (.010-.013)	.020 (.017-.024)	.029 (.024-.035)	2848
Age group							
6-11 years	99-00	.009 (.007-.012)	.008 (.006-.011)	.015 (.010-.024)	.030 (.016-.044)	.037 (.030-.077)	340
	01-02	.010 (.008-.011)	.010 (.008-.012)	.015 (.013-.019)	.027 (.018-.032)	.033 (.027-.048)	368
	03-04	.009 (.008-.010)	.008 (.007-.010)	.013 (.011-.017)	.024 (.016-.039)	.033 (.022-.050)	289
	05-06	.007 (.006-.008)	.006 (.005-.008)	.010 (.008-.014)	.018 (.013-.035)	.034 (.018-.048)	355
	07-08	.009 (.007-.011)	.008 (.007-.010)	.014 (.010-.022)	.026 (.016-.053)	.042 (.023-.065)	394
	09-10	.009 (.007-.010)	.008 (.007-.010)	.014 (.011-.016)	.020 (.015-.030)	.029 (.022-.033)	378
12-19 years	99-00	.007 (.006-.008)	.006 (.005-.008)	.010 (.009-.014)	.020 (.014-.030)	.030 (.019-.074)	719
	01-02	.007 (.006-.008)	.007 (.006-.008)	.012 (.009-.016)	.020 (.015-.026)	.026 (.020-.042)	762
	03-04	.007 (.006-.008)	.006 (.005-.007)	.010 (.008-.013)	.019 (.015-.027)	.034 (.022-.041)	725
	05-06	.006 (.005-.007)	.005 (.005-.006)	.009 (.007-.012)	.017 (.013-.022)	.023 (.019-.026)	701
	07-08	.007 (.006-.009)	.006 (.005-.008)	.012 (.009-.016)	.025 (.013-.050)	.033 (.022-.077)	376
	09-10	.007 (.006-.009)	.007 (.006-.008)	.011 (.010-.013)	.017 (.014-.021)	.025 (.017-.038)	451
20 years and older	99-00	.007 (.006-.009)	.007 (.006-.009)	.013 (.010-.016)	.024 (.019-.029)	.034 (.025-.051)	1405
	01-02	.008 (.007-.010)	.007 (.006-.009)	.014 (.011-.019)	.027 (.020-.039)	.043 (.030-.063)	1559
	03-04	*	.007 (.006-.008)	.012 (.010-.014)	.020 (.017-.024)	.028 (.022-.038)	1543
	05-06	.006 (.005-.006)	.005 (.005-.006)	.009 (.008-.010)	.016 (.014-.019)	.026 (.020-.039)	1520
	07-08	.007 (.006-.008)	.006 (.005-.008)	.012 (.009-.016)	.024 (.016-.036)	.037 (.025-.065)	1857
	09-10	.007 (.006-.008)	.007 (.006-.008)	.012 (.010-.013)	.020 (.017-.025)	.029 (.024-.036)	2019
Gender							
Males	99-00	.007 (.006-.009)	.006 (.005-.008)	.011 (.009-.015)	.021 (.017-.028)	.035 (.024-.056)	1227
	01-02	.007 (.006-.008)	.007 (.006-.008)	.012 (.010-.015)	.022 (.018-.028)	.033 (.025-.047)	1334
	03-04	.007 (.006-.008)	.006 (.006-.007)	.010 (.009-.012)	.019 (.015-.024)	.026 (.019-.039)	1280
	05-06	.005 (.005-.005)	.005 (.004-.005)	.008 (.007-.009)	.014 (.013-.016)	.021 (.016-.031)	1271
	07-08	.007 (.005-.008)	.006 (.005-.007)	.012 (.009-.015)	.022 (.016-.031)	.032 (.024-.056)	1327
	09-10	.007 (.006-.008)	.006 (.006-.007)	.011 (.010-.012)	.019 (.015-.023)	.028 (.022-.033)	1398
Females	99-00	.008 (.007-.010)	.007 (.006-.010)	.013 (.010-.017)	.025 (.019-.033)	.034 (.027-.054)	1237
	01-02	.009 (.008-.011)	.009 (.007-.011)	.016 (.012-.021)	.029 (.021-.042)	.045 (.031-.067)	1355
	03-04	*	.008 (.007-.009)	.013 (.011-.016)	.022 (.018-.028)	.031 (.025-.041)	1277
	05-06	.006 (.006-.007)	.006 (.005-.006)	.010 (.009-.011)	.019 (.015-.024)	.035 (.022-.041)	1305
	07-08	.008 (.006-.009)	.007 (.006-.008)	.013 (.010-.018)	.026 (.016-.043)	.042 (.024-.083)	1300
	09-10	.008 (.007-.009)	.008 (.006-.009)	.013 (.011-.015)	.021 (.017-.027)	.031 (.024-.041)	1450

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Uranium_BiomonitoringSummary.html

Urinary Uranium (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	.015 (.011-.022)	.015 (.011-.020)	.029 (.016-.058)	.059 (.027-.146)	.100 (.042-.270)	883
	01-02	.012 (.010-.016)	.012 (.009-.016)	.021 (.015-.028)	.033 (.024-.053)	.050 (.034-.080)	682
	03-04	.013 (.010-.016)	.013 (.009-.017)	.022 (.016-.029)	.035 (.026-.051)	.051 (.034-.061)	618
	05-06	.008 (.007-.009)	.007 (.006-.008)	.013 (.010-.015)	.022 (.016-.031)	.035 (.025-.060)	652
	07-08	.009 (.008-.012)	.009 (.007-.011)	.017 (.014-.021)	.031 (.021-.048)	.042 (.027-.097)	515
	09-10	.010 (.009-.011)	.010 (.008-.011)	.016 (.013-.018)	.026 (.022-.030)	.036 (.030-.044)	613
Non-Hispanic blacks	99-00	.006 (.004-.007)	.005 (.004-.006)	.008 (.006-.013)	.017 (.011-.029)	.028 (.018-.048)	568
	01-02	.005 (.005-.006)	.005 (.005-.006)	.008 (.007-.010)	.013 (.011-.014)	.017 (.014-.029)	667
	03-04	.006 (.005-.006)	.005 (.005-.006)	.009 (.008-.009)	.013 (.012-.015)	.018 (.014-.024)	722
	05-06	.004 (.004-.005)	.004 (.003-.005)	.006 (.006-.007)	.011 (.009-.015)	.017 (.012-.021)	692
	07-08	.005 (.005-.006)	.005 (.004-.006)	.009 (.007-.011)	.014 (.011-.017)	.018 (.014-.041)	589
	09-10	.006 (.005-.008)	.006 (.005-.007)	.010 (.008-.012)	.018 (.013-.021)	.023 (.018-.044)	544
Non-Hispanic whites	99-00	.007 (.006-.009)	.007 (.006-.009)	.012 (.010-.015)	.021 (.017-.027)	.030 (.024-.050)	822
	01-02	.008 (.007-.009)	.007 (.006-.009)	.013 (.011-.016)	.025 (.018-.032)	.034 (.025-.051)	1132
	03-04	*	.007 (.006-.008)	.011 (.010-.013)	.019 (.015-.024)	.027 (.020-.040)	1074
	05-06	.006 (.005-.006)	.005 (.005-.006)	.009 (.008-.010)	.016 (.013-.020)	.024 (.019-.039)	1041
	07-08	.007 (.006-.009)	.006 (.005-.008)	.012 (.009-.019)	.025 (.015-.043)	.039 (.024-.081)	1095
	09-10	.007 (.006-.008)	.007 (.006-.008)	.012 (.010-.013)	.018 (.015-.024)	.028 (.021-.035)	1225

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Uranium_BiomonitoringSummary.html

Urinary Butyl paraben

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	05-06	*	< LOD	1.30 (.800-1.80)	6.60 (4.80-10.7)	19.6 (16.4-26.7)	2548
	07-08	*	< LOD	1.10 (.900-1.50)	9.00 (6.20-11.4)	20.9 (15.2-30.1)	2604
	09-10	*	< LOD	.600 (.500-.800)	5.10 (3.40-8.10)	17.1 (10.5-29.4)	2749
Age group							
6-11 years	05-06	*	< LOD	.400 (.200-.600)	1.90 (.800-5.80)	7.50 (1.70-18.5)	356
	07-08	*	< LOD	.300 (.200-.500)	1.30 (.900-1.70)	3.80 (1.00-11.4)	389
	09-10	*	< LOD	.300 (<LOD-.400)	.900 (.500-1.70)	2.20 (1.30-5.10)	415
12-19 years	05-06	*	.200 (<LOD-.400)	.900 (.600-1.80)	9.60 (4.10-16.1)	24.6 (14.5-33.4)	702
	07-08	*	< LOD	.800 (.600-1.40)	6.20 (2.50-11.5)	15.8 (6.20-33.2)	401
	09-10	*	< LOD	.500 (.400-.900)	5.70 (1.40-15.7)	18.6 (5.70-39.0)	420
20 years and older	05-06	*	< LOD	1.70 (1.00-2.20)	6.80 (4.80-11.7)	20.5 (15.8-28.3)	1490
	07-08	*	< LOD	1.50 (1.00-1.90)	9.80 (6.70-13.4)	24.6 (16.5-32.0)	1814
	09-10	*	< LOD	.700 (.500-1.00)	6.00 (3.70-10.0)	18.7 (11.9-35.1)	1914
Gender							
Males	05-06	*	< LOD	.300 (<LOD-.300)	1.10 (.600-2.00)	3.20 (1.90-5.50)	1270
	07-08	*	< LOD	.200 (.200-.300)	1.00 (.700-1.50)	3.60 (2.00-7.20)	1294
	09-10	*	< LOD	< LOD	.800 (.500-1.20)	2.70 (1.50-12.0)	1399
Females	05-06	.904 (.760-1.07)	.500 (.400-.700)	3.70 (2.90-4.80)	17.4 (11.7-20.8)	34.9 (28.2-41.1)	1278
	07-08	.926 (.825-1.04)	.500 (.400-.700)	3.90 (3.00-4.70)	17.9 (12.1-20.5)	34.8 (25.2-52.1)	1310
	09-10	*	.300 (.200-.400)	1.80 (1.30-2.30)	11.3 (6.60-18.8)	31.8 (17.7-49.4)	1350
Race/ethnicity							
Mexican Americans	05-06	*	.200 (<LOD-.300)	1.10 (.700-2.30)	11.3 (6.40-15.6)	27.3 (16.5-35.9)	637
	07-08	*	< LOD	.900 (.500-1.30)	6.80 (3.50-13.2)	22.7 (13.2-30.7)	531
	09-10	*	< LOD	.900 (.400-1.30)	8.30 (4.50-13.8)	18.8 (13.8-35.1)	566
Non-Hispanic blacks	05-06	*	.300 (.200-.400)	2.10 (1.00-3.50)	9.10 (4.40-25.0)	31.8 (10.1-82.6)	678
	07-08	*	.200 (<LOD-.300)	1.10 (.700-1.70)	7.10 (4.20-9.90)	20.0 (8.90-31.3)	597
	09-10	*	< LOD	.500 (.300-.900)	3.10 (1.40-7.30)	9.00 (5.60-16.7)	516
Non-Hispanic whites	05-06	*	< LOD	1.00 (.700-1.80)	5.80 (4.00-9.00)	17.7 (11.7-25.9)	1038
	07-08	*	< LOD	1.20 (.800-1.60)	9.10 (5.00-13.3)	19.6 (12.1-32.0)	1077
	09-10	*	< LOD	.700 (.500-.900)	4.90 (2.70-9.20)	17.7 (9.20-34.5)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06, 07-08, and 09-10 are 0.2, 0.2, and 0.2 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Parabens_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Parabens_FactSheet.html

Urinary Butyl paraben (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	05-06	*	< LOD	1.16 (.910-1.62)	7.82 (6.26-11.2)	21.2 (16.8-27.3)	2548
	07-08	*	< LOD	1.16 (.880-1.59)	9.83 (6.51-13.5)	23.1 (18.1-30.4)	2604
	09-10	*	< LOD	.710 (.610-.930)	5.58 (3.55-10.0)	18.0 (10.2-27.6)	2749
Age group							
6-11 years	05-06	*	< LOD	.440 (.370-.660)	1.65 (1.00-5.29)	7.92 (1.64-30.5)	356
	07-08	*	< LOD	.540 (.410-.720)	1.24 (.920-2.80)	5.91 (.930-76.0)	389
	09-10	*	< LOD	.530 (<LOD-.630)	1.36 (.740-1.89)	2.74 (1.56-6.58)	415
12-19 years	05-06	*	.190 (<LOD-.250)	.690 (.490-1.20)	6.67 (3.60-11.2)	18.7 (9.52-28.9)	702
	07-08	*	< LOD	.670 (.450-.930)	3.70 (1.80-7.41)	9.64 (6.00-18.7)	401
	09-10	*	< LOD	.550 (.420-.790)	3.75 (1.21-11.9)	13.8 (3.75-23.7)	420
20 years and older	05-06	*	< LOD	1.49 (1.06-1.96)	9.07 (6.55-14.8)	25.5 (17.9-30.4)	1490
	07-08	*	< LOD	1.52 (1.04-2.21)	12.4 (8.35-15.1)	27.5 (21.2-36.8)	1814
	09-10	*	< LOD	.830 (.640-1.19)	7.06 (3.79-11.9)	22.0 (13.1-31.2)	1914
Gender							
Males	05-06	*	< LOD	.290 (<LOD-.350)	.830 (.570-1.16)	2.28 (1.43-2.91)	1270
	07-08	*	< LOD	.290 (.260-.340)	.830 (.620-1.36)	2.40 (1.56-4.85)	1294
	09-10	*	< LOD	< LOD	.700 (.560-1.09)	2.50 (1.18-5.87)	1399
Females	05-06	1.06 (.914-1.24)	.750 (.600-.860)	4.44 (3.01-6.26)	19.8 (13.5-25.7)	34.7 (26.9-42.7)	1278
	07-08	1.11 (.984-1.25)	.670 (.550-.920)	4.44 (3.02-6.12)	20.6 (14.6-25.9)	39.4 (28.1-77.7)	1310
	09-10	*	.450 (.420-.520)	2.17 (1.67-3.10)	13.5 (8.75-22.5)	28.2 (18.4-45.7)	1350
Race/ethnicity							
Mexican Americans	05-06	*	.240 (<LOD-.290)	1.20 (.760-1.68)	9.15 (5.36-15.2)	20.3 (15.8-29.2)	637
	07-08	*	< LOD	.810 (.610-1.07)	8.57 (3.28-12.0)	23.0 (12.7-29.5)	531
	09-10	*	< LOD	.930 (.570-1.76)	8.01 (4.77-10.5)	17.0 (10.5-27.1)	566
Non-Hispanic blacks	05-06	*	.220 (.180-.270)	1.46 (.750-2.64)	8.18 (3.65-17.9)	21.0 (10.9-51.3)	678
	07-08	*	.220 (<LOD-.280)	.900 (.570-1.27)	5.21 (3.13-6.18)	12.1 (6.14-21.4)	597
	09-10	*	< LOD	.390 (.270-.590)	2.63 (.890-4.71)	7.00 (3.26-11.9)	516
Non-Hispanic whites	05-06	*	< LOD	1.12 (.850-1.65)	6.93 (5.21-9.91)	20.9 (15.4-27.3)	1038
	07-08	*	< LOD	1.18 (.810-2.09)	10.7 (6.32-14.8)	26.8 (15.1-40.4)	1077
	09-10	*	< LOD	.780 (.620-1.13)	6.34 (3.44-12.6)	20.8 (9.12-33.2)	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Parabens_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Parabens_FactSheet.html

Urinary Ethyl paraben

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	05-06	*	< LOD	4.80 (3.40-6.00)	26.2 (19.4-31.9)	57.2 (40.3-83.4)	2548
	07-08	*	< LOD	5.40 (3.80-6.90)	29.3 (21.9-42.4)	72.1 (54.6-91.8)	2604
	09-10	*	1.10 (<LOD-1.20)	6.60 (4.90-8.30)	37.6 (24.4-51.3)	82.6 (60.3-119)	2749
Age group							
6-11 years	05-06	*	< LOD	< LOD	2.60 (1.70-4.70)	9.90 (2.80-30.7)	356
	07-08	*	< LOD	1.70 (1.10-2.60)	4.60 (2.30-36.7)	35.6 (2.60-282)	389
	09-10	*	< LOD	1.50 (1.20-1.80)	3.70 (2.50-5.30)	11.5 (4.50-18.4)	415
12-19 years	05-06	*	< LOD	2.30 (1.30-3.20)	11.3 (7.90-24.1)	38.0 (17.7-122)	702
	07-08	*	< LOD	3.00 (2.30-4.90)	23.0 (7.30-60.3)	60.3 (12.5-159)	401
	09-10	*	< LOD	3.50 (2.10-5.90)	33.8 (19.4-49.6)	83.2 (34.5-284)	420
20 years and older	05-06	*	< LOD	5.90 (4.60-8.10)	30.4 (22.3-38.6)	69.5 (48.1-89.5)	1490
	07-08	*	1.10 (<LOD-1.40)	6.20 (4.60-9.20)	34.7 (22.8-50.4)	77.4 (51.4-120)	1814
	09-10	*	1.30 (1.10-1.50)	8.40 (6.00-10.6)	45.0 (29.1-60.3)	91.4 (62.6-129)	1914
Gender							
Males	05-06	*	< LOD	1.90 (1.20-3.00)	8.90 (5.80-13.7)	25.2 (15.7-29.2)	1270
	07-08	*	< LOD	2.40 (1.50-3.00)	11.2 (6.60-15.3)	28.3 (15.3-42.8)	1294
	09-10	*	< LOD	2.20 (1.90-2.70)	10.4 (7.40-19.1)	36.4 (19.1-61.3)	1399
Females	05-06	*	1.30 (<LOD-2.20)	10.0 (7.50-13.2)	42.7 (32.3-62.8)	98.7 (74.7-147)	1278
	07-08	*	1.80 (1.40-2.50)	10.1 (6.90-16.5)	62.0 (38.2-80.8)	123 (78.5-259)	1310
	09-10	3.84 (3.35-4.41)	2.00 (1.60-2.40)	15.4 (12.5-18.4)	62.4 (45.7-78.7)	138 (92.2-211)	1350
Race/ethnicity							
Mexican Americans	05-06	*	< LOD	3.70 (1.90-6.10)	24.0 (11.6-37.7)	55.6 (40.3-120)	637
	07-08	*	< LOD	4.10 (2.90-5.90)	23.6 (12.8-30.1)	48.3 (31.0-65.2)	531
	09-10	*	< LOD	4.30 (2.60-7.70)	31.0 (21.5-46.3)	65.5 (46.3-92.2)	566
Non-Hispanic blacks	05-06	*	1.00 (<LOD-1.50)	5.70 (3.70-9.80)	31.3 (20.4-52.1)	92.3 (41.5-166)	678
	07-08	*	1.00 (<LOD-1.50)	7.90 (3.90-15.8)	39.9 (24.0-63.4)	97.2 (58.3-256)	597
	09-10	3.62 (3.11-4.22)	1.90 (1.60-2.40)	11.1 (8.10-17.5)	58.6 (33.8-93.7)	166 (66.0-297)	516
Non-Hispanic whites	05-06	*	< LOD	4.80 (3.10-6.10)	24.0 (15.5-32.3)	54.7 (36.4-86.8)	1038
	07-08	*	1.00 (<LOD-1.30)	5.20 (3.50-7.10)	27.2 (18.0-40.8)	67.1 (42.8-98.1)	1077
	09-10	*	1.10 (<LOD-1.20)	6.50 (4.50-8.50)	37.5 (22.7-52.7)	78.7 (49.7-138)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06, 07-08, and 09-10 are 1.0, 1.0, and 1.0 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Parabens_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Parabens_FactSheet.html

Urinary Ethyl paraben (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	05-06	*	< LOD	4.61 (3.39-6.30)	27.1 (19.1-34.0)	66.5 (46.5-84.2)	2548
	07-08	*	< LOD	5.56 (4.31-7.56)	32.7 (22.5-50.2)	80.0 (53.2-117)	2604
	09-10	*	1.65 (<LOD-1.82)	6.11 (4.90-7.96)	36.6 (28.0-47.2)	94.6 (67.3-127)	2749
Age group							
6-11 years	05-06	*	< LOD	< LOD	4.18 (1.91-10.2)	13.5 (4.18-29.0)	356
	07-08	*	< LOD	2.54 (1.82-3.30)	6.13 (3.26-34.3)	34.3 (3.30-207)	389
	09-10	*	< LOD	2.37 (1.97-3.09)	5.00 (3.74-6.67)	10.1 (6.11-27.3)	415
12-19 years	05-06	*	< LOD	1.94 (1.29-2.65)	10.9 (3.57-20.2)	32.0 (14.1-57.8)	702
	07-08	*	< LOD	2.93 (1.93-3.97)	12.9 (4.60-37.9)	37.9 (8.79-110)	401
	09-10	*	< LOD	3.11 (2.22-5.94)	26.2 (13.0-33.9)	52.4 (28.8-177)	420
20 years and older	05-06	*	< LOD	6.36 (4.63-7.92)	32.4 (23.7-46.0)	76.8 (57.3-97.9)	1490
	07-08	*	1.58 (<LOD-1.97)	6.97 (4.94-10.9)	38.5 (24.8-61.9)	94.1 (53.6-146)	1814
	09-10	*	1.82 (1.61-2.13)	8.08 (5.92-10.7)	43.7 (33.1-56.0)	106 (78.1-137)	1914
Gender							
Males	05-06	*	< LOD	1.82 (1.48-2.15)	6.38 (3.55-9.95)	15.5 (10.2-22.3)	1270
	07-08	*	< LOD	2.14 (1.67-3.03)	8.61 (5.54-12.4)	21.7 (11.7-32.7)	1294
	09-10	*	< LOD	2.45 (2.22-2.92)	8.63 (6.14-13.7)	28.8 (14.4-43.2)	1399
Females	05-06	*	2.09 (<LOD-2.60)	12.7 (8.45-16.2)	57.3 (40.7-75.2)	107 (82.9-141)	1278
	07-08	*	2.63 (2.09-3.55)	13.9 (9.09-22.5)	66.1 (40.9-95.1)	130 (78.8-203)	1310
	09-10	4.63 (4.13-5.19)	2.90 (2.47-3.43)	16.8 (13.7-22.0)	71.9 (53.4-98.8)	173 (121-227)	1350
Race/ethnicity							
Mexican Americans	05-06	*	< LOD	3.49 (2.09-5.23)	23.6 (11.8-33.2)	68.5 (32.1-107)	637
	07-08	*	< LOD	4.44 (2.86-5.81)	22.5 (15.9-27.2)	47.7 (26.7-74.2)	531
	09-10	*	< LOD	4.40 (3.09-6.31)	30.0 (16.0-36.9)	59.1 (36.9-87.2)	566
Non-Hispanic blacks	05-06	*	.970 (<LOD-1.14)	4.48 (2.90-6.07)	22.0 (14.1-36.8)	47.5 (33.7-106)	678
	07-08	*	1.18 (<LOD-1.47)	6.37 (3.20-10.9)	38.8 (16.5-53.1)	90.7 (42.6-172)	597
	09-10	2.62 (2.14-3.21)	1.57 (1.20-2.05)	7.35 (5.15-9.97)	44.9 (22.0-75.0)	98.8 (60.0-162)	516
Non-Hispanic whites	05-06	*	< LOD	4.73 (3.38-6.58)	27.5 (16.2-38.4)	70.6 (46.0-91.2)	1038
	07-08	*	1.48 (<LOD-1.82)	5.71 (3.74-8.97)	32.3 (20.0-53.6)	75.7 (40.5-121)	1077
	09-10	*	1.79 (<LOD-2.03)	6.63 (4.84-9.18)	37.7 (27.7-53.4)	114 (62.8-164)	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Parabens_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Parabens_FactSheet.html

Urinary Methyl paraben

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	05-06	56.4 (46.9-67.9)	63.5 (47.9-74.5)	216 (168-282)	560 (432-777)	974 (827-1110)	2548
	07-08	60.1 (49.6-72.8)	59.8 (47.5-73.3)	204 (171-253)	544 (444-675)	902 (746-1100)	2604
	09-10	57.1 (51.0-63.9)	55.1 (46.7-62.6)	213 (179-259)	621 (546-716)	1010 (919-1170)	2749
Age group							
6-11 years	05-06	33.5 (25.6-43.8)	25.0 (19.4-35.3)	86.7 (61.5-153)	289 (238-868)	1560 (389-2090)	356
	07-08	35.2 (25.0-49.6)	30.1 (19.6-38.5)	94.0 (46.6-214)	340 (218-527)	651 (430-995)	389
	09-10	33.9 (27.4-41.9)	26.5 (20.4-33.3)	79.4 (59.0-126)	415 (163-849)	873 (477-1230)	415
12-19 years	05-06	53.8 (43.9-66.0)	53.5 (39.4-65.7)	192 (163-264)	491 (380-720)	901 (571-1210)	702
	07-08	64.9 (50.3-83.8)	66.3 (48.0-96.7)	221 (180-276)	521 (424-789)	971 (574-1390)	401
	09-10	53.9 (40.9-70.9)	56.5 (39.4-75.6)	176 (122-291)	716 (399-934)	984 (805-1210)	420
20 years and older	05-06	60.3 (48.7-74.7)	70.0 (57.1-85.3)	225 (177-322)	602 (439-814)	974 (827-1110)	1490
	07-08	63.0 (51.8-76.7)	64.0 (51.4-79.8)	210 (172-267)	558 (447-678)	916 (743-1190)	1814
	09-10	61.0 (53.5-69.6)	59.1 (50.6-72.6)	239 (188-276)	625 (566-711)	1010 (909-1270)	1914
Gender							
Males	05-06	29.8 (24.8-35.8)	23.7 (19.5-29.4)	97.6 (78.1-121)	299 (229-355)	491 (385-743)	1270
	07-08	33.6 (27.3-41.3)	27.4 (20.8-39.9)	93.5 (71.0-137)	312 (260-407)	598 (460-734)	1294
	09-10	31.7 (27.6-36.4)	25.3 (20.6-30.6)	94.0 (76.8-122)	366 (275-482)	727 (569-919)	1399
Females	05-06	104 (80.8-135)	137 (93.0-168)	356 (279-427)	842 (660-974)	1110 (956-1330)	1278
	07-08	105 (90.2-123)	116 (95.2-139)	315 (248-424)	683 (558-901)	1140 (917-1420)	1310
	09-10	100 (87.9-114)	106 (90.9-125)	349 (271-405)	812 (696-931)	1230 (1010-1550)	1350
Race/ethnicity							
Mexican Americans	05-06	78.2 (61.0-100)	86.3 (64.7-121)	301 (233-349)	742 (474-896)	1100 (896-1170)	637
	07-08	80.5 (66.0-98.2)	87.0 (63.4-117)	283 (222-348)	630 (533-791)	1030 (773-1240)	531
	09-10	82.9 (68.6-100)	85.5 (73.0-99.8)	293 (231-385)	821 (536-1120)	1580 (855-2360)	566
Non-Hispanic blacks	05-06	174 (133-229)	216 (150-306)	616 (465-789)	1180 (1010-1480)	1690 (1210-2880)	678
	07-08	152 (135-171)	159 (135-194)	483 (421-552)	1080 (938-1350)	1720 (1210-2700)	597
	09-10	191 (171-213)	223 (170-288)	678 (533-849)	1470 (1220-2010)	2340 (1650-2750)	516
Non-Hispanic whites	05-06	43.8 (36.7-52.3)	44.3 (36.7-58.2)	165 (132-209)	412 (320-520)	806 (485-974)	1038
	07-08	48.9 (39.9-59.9)	47.6 (37.8-62.0)	158 (129-198)	430 (301-557)	671 (524-921)	1077
	09-10	43.5 (39.4-48.1)	41.3 (33.5-47.7)	150 (127-176)	463 (361-557)	757 (619-888)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06, 07-08, and 09-10 are 1.0, 1.0, and 1.0 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Parabens_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Parabens_FactSheet.html

Urinary Methyl paraben (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	05-06	55.0 (46.8-64.6)	58.8 (47.0-73.4)	221 (170-264)	527 (426-685)	902 (753-995)	2548
	07-08	60.6 (49.8-73.8)	61.0 (45.3-83.2)	228 (188-288)	538 (439-645)	820 (694-1000)	2604
	09-10	59.6 (53.5-66.3)	57.7 (52.3-68.5)	232 (195-262)	544 (468-632)	887 (734-1020)	2749
Age group							
6-11 years	05-06	36.8 (27.9-48.7)	26.9 (21.0-36.6)	108 (55.9-164)	259 (195-753)	1540 (313-2510)	356
	07-08	43.4 (31.3-60.2)	33.1 (21.8-55.0)	118 (62.3-197)	424 (218-668)	817 (558-1020)	389
	09-10	44.2 (35.7-54.6)	32.3 (26.5-42.9)	110 (67.6-158)	483 (258-722)	849 (535-1080)	415
12-19 years	05-06	40.1 (33.7-47.8)	41.7 (31.5-56.3)	160 (129-178)	362 (280-428)	549 (434-656)	702
	07-08	50.5 (40.1-63.7)	52.8 (36.6-78.1)	167 (111-245)	336 (292-395)	475 (376-623)	401
	09-10	43.3 (33.5-55.9)	43.5 (27.5-68.6)	168 (123-221)	399 (292-558)	632 (425-890)	420
20 years and older	05-06	60.5 (50.5-72.5)	69.6 (51.5-90.9)	251 (189-320)	602 (462-729)	930 (788-1010)	1490
	07-08	64.7 (52.9-79.2)	69.7 (51.1-92.5)	256 (209-316)	576 (450-710)	865 (697-1130)	1814
	09-10	64.6 (57.0-73.3)	71.2 (56.6-84.7)	251 (217-287)	566 (495-673)	924 (753-1060)	1914
Gender							
Males	05-06	23.9 (20.6-27.8)	21.1 (17.4-25.3)	72.1 (58.0-89.5)	209 (160-260)	368 (262-498)	1270
	07-08	28.3 (22.8-35.0)	23.5 (17.2-33.2)	80.7 (56.9-111)	257 (181-318)	444 (386-558)	1294
	09-10	28.4 (25.1-32.2)	23.3 (19.3-26.9)	77.7 (59.6-105)	260 (212-370)	516 (412-633)	1399
Females	05-06	123 (99.3-152)	147 (111-196)	377 (319-445)	788 (676-910)	1050 (937-1290)	1278
	07-08	126 (109-145)	160 (134-182)	374 (325-422)	734 (576-879)	1040 (835-1260)	1310
	09-10	121 (107-137)	146 (125-162)	359 (307-426)	745 (606-890)	1080 (924-1390)	1350
Race/ethnicity							
Mexican Americans	05-06	70.3 (57.1-86.7)	85.8 (67.2-106)	249 (191-303)	515 (431-656)	870 (656-1080)	637
	07-08	78.5 (63.6-97.0)	96.2 (59.4-127)	293 (245-370)	627 (541-728)	831 (729-963)	531
	09-10	82.3 (71.1-95.2)	95.2 (76.0-113)	284 (237-337)	623 (459-899)	1130 (657-2120)	566
Non-Hispanic blacks	05-06	122 (93.0-161)	161 (98.2-225)	377 (334-462)	792 (581-1100)	1220 (916-1590)	678
	07-08	118 (105-133)	134 (111-155)	356 (311-403)	757 (573-882)	1030 (847-1290)	597
	09-10	138 (118-163)	165 (127-203)	403 (356-498)	928 (729-1120)	1350 (1080-1680)	516
Non-Hispanic whites	05-06	46.1 (39.0-54.4)	45.6 (34.5-59.7)	170 (142-234)	447 (339-706)	814 (640-995)	1038
	07-08	51.0 (40.5-64.2)	46.7 (35.8-66.3)	191 (154-230)	481 (383-614)	747 (574-966)	1077
	09-10	48.9 (44.5-53.7)	45.9 (39.5-51.9)	184 (154-215)	468 (388-560)	734 (631-924)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Parabens_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Parabens_FactSheet.html

Urinary *n*-Propyl paraben

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	05-06	7.91 (6.41-9.77)	8.70 (5.90-12.4)	48.9 (36.6-64.8)	160 (129-200)	299 (237-354)	2548
	07-08	7.59 (6.22-9.26)	7.60 (5.70-10.6)	44.1 (34.1-55.6)	144 (115-196)	268 (231-290)	2604
	09-10	6.97 (5.96-8.16)	6.40 (5.10-8.00)	44.3 (38.6-50.3)	153 (133-183)	297 (233-334)	2749
Age group							
6-11 years	05-06	3.41 (2.43-4.77)	2.50 (1.60-4.30)	11.1 (7.10-21.0)	47.5 (30.0-91.9)	125 (55.8-383)	356
	07-08	3.61 (2.39-5.44)	3.50 (1.90-6.00)	13.7 (7.60-22.8)	49.1 (21.7-99.6)	99.6 (80.4-120)	389
	09-10	3.28 (2.58-4.17)	2.70 (2.00-4.00)	10.2 (7.00-14.0)	42.9 (28.2-61.6)	114 (56.8-239)	415
12-19 years	05-06	8.16 (5.70-11.7)	8.40 (5.00-13.9)	46.4 (24.1-68.5)	165 (96.7-222)	310 (201-361)	702
	07-08	8.85 (6.73-11.6)	7.90 (5.30-12.0)	49.2 (31.1-62.2)	193 (100-234)	271 (195-314)	401
	09-10	7.01 (5.13-9.58)	7.90 (4.40-11.0)	39.7 (24.8-57.8)	153 (78.1-299)	367 (136-694)	420
20 years and older	05-06	8.67 (6.83-11.0)	11.2 (7.70-14.4)	55.6 (41.2-75.6)	172 (130-233)	302 (237-367)	1490
	07-08	8.04 (6.56-9.84)	8.40 (6.10-11.9)	48.8 (38.3-61.1)	152 (118-202)	272 (232-312)	1814
	09-10	7.58 (6.28-9.15)	7.50 (5.30-9.90)	48.9 (42.6-58.0)	166 (139-199)	298 (251-334)	1914
Gender							
Males	05-06	2.96 (2.33-3.77)	2.30 (1.60-3.10)	13.8 (9.00-18.0)	51.7 (39.7-70.9)	125 (77.2-185)	1270
	07-08	3.02 (2.47-3.71)	2.30 (1.70-3.30)	12.6 (8.40-17.4)	56.4 (38.9-89.5)	140 (98.1-209)	1294
	09-10	2.77 (2.26-3.40)	2.00 (1.40-2.60)	11.4 (7.70-15.4)	57.3 (44.6-67.8)	134 (86.1-184)	1399
Females	05-06	20.4 (16.0-25.9)	29.1 (21.6-37.5)	93.0 (75.7-129)	254 (193-318)	357 (318-395)	1278
	07-08	18.4 (15.7-21.5)	23.3 (20.5-26.1)	76.3 (62.9-100)	210 (164-269)	312 (271-363)	1310
	09-10	16.9 (14.6-19.4)	20.2 (17.0-25.9)	83.3 (70.0-92.4)	227 (171-295)	361 (297-502)	1350
Race/ethnicity							
Mexican Americans	05-06	10.6 (8.08-13.8)	12.1 (7.00-17.5)	65.9 (43.5-93.3)	247 (174-293)	389 (293-453)	637
	07-08	7.82 (5.98-10.2)	7.00 (4.20-12.9)	52.1 (36.9-64.1)	146 (105-201)	217 (161-401)	531
	09-10	8.94 (7.25-11.0)	8.70 (5.70-13.3)	58.9 (45.7-68.5)	182 (117-299)	451 (269-911)	566
Non-Hispanic blacks	05-06	26.8 (21.5-33.3)	34.7 (28.3-44.2)	125 (89.5-165)	331 (236-411)	531 (367-730)	678
	07-08	21.3 (18.4-24.7)	28.6 (21.8-34.8)	85.4 (70.8-96.9)	222 (176-282)	410 (277-506)	597
	09-10	26.1 (21.8-31.3)	29.9 (23.4-39.5)	129 (87.4-167)	322 (229-503)	572 (369-654)	516
Non-Hispanic whites	05-06	6.21 (5.02-7.68)	6.00 (4.60-9.10)	35.7 (28.4-49.6)	130 (92.1-157)	229 (156-318)	1038
	07-08	6.49 (5.05-8.33)	6.10 (4.50-9.20)	36.8 (26.9-47.5)	129 (91.3-196)	251 (190-290)	1077
	09-10	5.34 (4.55-6.27)	4.40 (3.60-5.70)	35.6 (26.1-44.3)	128 (92.4-141)	235 (170-300)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06, 07-08, and 09-10 are 0.2, 0.2, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Parabens_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Parabens_FactSheet.html

Urinary *n*-Propyl paraben (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	05-06	7.71 (6.40-9.30)	8.27 (5.87-11.3)	52.9 (41.6-66.1)	161 (140-178)	263 (226-288)	2548
	07-08	7.65 (6.22-9.40)	8.00 (5.34-11.6)	50.0 (39.5-59.8)	138 (115-168)	235 (199-293)	2604
	09-10	7.28 (6.30-8.41)	6.67 (5.35-8.50)	45.1 (37.3-53.5)	155 (128-187)	253 (223-300)	2749
Age group							
6-11 years	05-06	3.75 (2.67-5.26)	2.70 (1.89-4.43)	11.5 (8.07-16.0)	52.1 (22.9-100)	121 (55.5-352)	356
	07-08	4.45 (3.03-6.53)	3.50 (1.79-6.85)	16.5 (8.82-28.8)	59.1 (28.6-94.4)	98.4 (65.0-139)	389
	09-10	4.27 (3.34-5.47)	3.33 (2.41-4.62)	12.0 (8.52-16.6)	46.1 (27.5-108)	143 (72.5-243)	415
12-19 years	05-06	6.08 (4.36-8.49)	5.44 (3.03-10.5)	33.0 (20.7-48.7)	116 (81.9-161)	175 (140-239)	702
	07-08	6.88 (5.30-8.94)	5.68 (3.51-9.90)	33.2 (26.7-46.5)	103 (87.2-112)	132 (105-235)	401
	09-10	5.64 (4.17-7.62)	5.19 (3.53-9.03)	32.9 (20.4-42.4)	107 (57.6-217)	218 (101-397)	420
20 years and older	05-06	8.70 (7.08-10.7)	10.8 (7.39-15.2)	63.2 (49.7-80.5)	178 (152-198)	270 (236-293)	1490
	07-08	8.26 (6.70-10.2)	9.91 (6.15-12.6)	57.9 (47.3-69.1)	161 (127-199)	273 (212-312)	1814
	09-10	8.03 (6.74-9.56)	8.27 (6.10-10.8)	55.5 (43.5-66.0)	164 (136-200)	270 (232-314)	1914
Gender							
Males	05-06	2.38 (1.92-2.95)	1.84 (1.42-2.46)	9.52 (6.79-12.8)	40.5 (31.1-51.5)	90.8 (56.8-125)	1270
	07-08	2.55 (2.06-3.15)	1.74 (1.40-2.53)	10.7 (5.99-15.4)	44.7 (29.0-66.1)	112 (77.4-144)	1294
	09-10	2.48 (2.05-3.00)	1.71 (1.35-2.25)	8.85 (6.12-14.2)	40.3 (33.8-57.6)	105 (74.5-143)	1399
Females	05-06	23.9 (19.9-28.8)	34.9 (26.9-46.6)	114 (101-132)	235 (193-263)	306 (277-342)	1278
	07-08	22.0 (18.7-25.8)	30.4 (25.2-36.3)	93.3 (80.8-110)	199 (161-245)	302 (271-370)	1310
	09-10	20.4 (17.8-23.3)	25.7 (21.8-30.7)	97.1 (81.8-115)	223 (191-248)	334 (295-389)	1350
Race/ethnicity							
Mexican Americans	05-06	9.49 (7.29-12.4)	11.0 (6.69-16.4)	55.2 (44.9-79.6)	186 (147-271)	329 (194-441)	637
	07-08	7.63 (5.72-10.2)	6.60 (4.10-13.4)	47.9 (39.5-66.7)	162 (123-187)	273 (187-304)	531
	09-10	8.87 (7.42-10.6)	7.97 (5.87-12.8)	54.7 (42.3-73.8)	184 (140-269)	325 (221-575)	566
Non-Hispanic blacks	05-06	18.8 (15.2-23.3)	23.6 (19.0-30.8)	83.7 (61.8-105)	217 (169-255)	318 (242-377)	678
	07-08	16.6 (14.1-19.5)	22.9 (18.0-27.2)	64.2 (51.4-73.3)	141 (117-172)	254 (169-311)	597
	09-10	18.9 (15.2-23.5)	21.1 (18.2-27.7)	80.9 (59.5-109)	200 (151-249)	301 (223-376)	516
Non-Hispanic whites	05-06	6.53 (5.32-8.02)	6.45 (3.96-9.11)	47.4 (32.6-65.3)	146 (119-184)	241 (184-278)	1038
	07-08	6.77 (5.14-8.93)	5.81 (3.94-10.5)	45.4 (32.2-61.4)	132 (105-172)	215 (169-295)	1077
	09-10	6.00 (5.19-6.94)	5.12 (4.05-6.25)	37.8 (32.1-49.2)	148 (115-183)	242 (215-272)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Parabens_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Parabens_FactSheet.html

Urinary Nitrate

Geometric mean and selected percentiles of urine concentrations (in mg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50 th	75 th	90 th	95 th	
Total	01-02	48.2 (46.2-50.3)	49.0 (46.0-52.0)	78.0 (73.0-83.0)	100 (100-130)	140 (130-150)	1617
	05-06	42.7 (39.6-46.1)	47.8 (44.4-51.2)	74.6 (69.8-79.4)	108 (101-114)	133 (125-144)	7697
	07-08	46.3 (44.6-48.1)	50.3 (48.4-52.0)	76.0 (72.7-79.1)	110 (104-116)	138 (132-146)	7629
Age group							
6-11 years	01-02	62.2 (53.8-71.8)	68.0 (58.0-79.0)	94.0 (84.0-100)	130 (100-160)	150 (120-380)	374
	05-06	51.2 (47.4-55.4)	54.7 (51.9-58.2)	79.2 (72.8-87.2)	113 (101-128)	141 (120-158)	1054
	07-08	55.2 (51.7-58.9)	60.2 (56.1-64.3)	84.5 (80.1-92.5)	117 (107-135)	149 (132-189)	1143
12-19 years	01-02	57.4 (53.5-61.6)	66.0 (60.0-69.0)	91.0 (86.0-95.0)	120 (100-130)	150 (130-160)	827
	05-06	52.5 (48.5-56.8)	57.5 (52.3-62.6)	84.2 (79.9-88.4)	119 (111-124)	144 (129-153)	2106
	07-08	55.5 (51.5-59.7)	56.8 (51.9-61.5)	84.1 (76.2-94.5)	119 (107-133)	144 (133-162)	1135
20 years and older	01-02	45.4 (43.3-47.5)	49.0 (46.0-52.0)	78.0 (73.0-83.0)	100 (100-130)	140 (130-150)	1617
	05-06	40.5 (37.4-43.9)	45.0 (41.3-48.3)	71.7 (67.1-77.4)	105 (98.0-113)	129 (122-142)	4537
	07-08	44.2 (42.5-45.9)	48.1 (46.0-49.7)	73.2 (70.1-76.5)	107 (101-113)	135 (128-146)	5351
Gender							
Males	01-02	57.5 (54.6-60.6)	63.0 (59.0-67.0)	89.0 (83.0-94.0)	130 (100-140)	150 (140-170)	1335
	05-06	48.4 (44.6-52.6)	52.7 (48.3-57.8)	79.4 (72.8-86.5)	110 (103-121)	136 (123-152)	3765
	07-08	51.9 (49.9-54.1)	56.1 (53.8-58.0)	79.5 (75.7-83.5)	112 (105-119)	137 (131-149)	3839
Females	01-02	40.7 (38.4-43.2)	43.0 (41.0-48.0)	72.0 (68.0-76.0)	100 (98.0-120)	130 (120-150)	1483
	05-06	37.9 (35.1-40.8)	42.0 (38.2-46.0)	69.2 (65.5-73.4)	104 (96.6-110)	130 (124-140)	3932
	07-08	41.4 (39.3-43.7)	43.9 (41.5-46.3)	71.3 (67.5-74.8)	108 (99.8-115)	138 (132-149)	3790
Race/ethnicity							
Mexican Americans	01-02	53.2 (48.7-58.2)	59.0 (52.0-66.0)	84.0 (79.0-91.0)	120 (100-150)	160 (130-180)	707
	05-06	47.8 (44.7-51.2)	52.4 (49.9-56.2)	77.9 (75.1-83.1)	113 (104-120)	148 (133-156)	1972
	07-08	48.7 (45.0-52.6)	51.9 (47.4-56.7)	75.6 (69.7-81.3)	111 (100-122)	148 (127-164)	1505
Non-Hispanic blacks	01-02	53.8 (47.8-60.5)	58.0 (51.0-64.0)	84.0 (77.0-93.0)	120 (100-130)	140 (130-170)	680
	05-06	45.9 (42.1-50.0)	50.4 (46.6-54.9)	75.0 (68.8-80.8)	101 (95.3-110)	127 (114-148)	2078
	07-08	47.5 (45.0-50.3)	50.3 (48.6-52.5)	74.7 (71.9-77.3)	105 (97.1-116)	134 (125-150)	1707
Non-Hispanic whites	01-02	46.3 (44.1-48.6)	51.0 (47.0-53.0)	81.0 (78.0-85.0)	120 (100-130)	140 (130-150)	1228
	05-06	41.2 (37.6-45.2)	46.2 (41.3-50.6)	73.3 (67.3-80.1)	107 (98.2-115)	129 (122-142)	3056
	07-08	45.0 (42.7-47.5)	49.2 (46.1-52.6)	75.5 (70.5-80.0)	108 (101-116)	134 (128-140)	3190

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 05-06, and 07-08 are 0.7, 0.7, and 0.7 respectively.

Urinary Nitrate (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in mg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50 th	75 th	90 th	95 th	
Total	01-02	49.8 (47.7-51.9)	46.9 (44.2-49.6)	63.8 (61.2-67.7)	90.9 (84.3-98.8)	120 (111-128)	1616
	05-06	42.6 (40.2-45.1)	42.4 (40.1-44.7)	59.7 (55.8-64.1)	85.5 (81.3-91.3)	113 (106-118)	7697
	07-08	47.7 (45.9-49.7)	46.0 (44.0-48.3)	66.5 (62.4-70.5)	98.0 (92.3-102)	127 (119-135)	7628
Age group							
6-11 years	01-02	72.0 (66.1-78.4)	66.0 (62.6-70.4)	87.0 (80.2-97.7)	129 (96.5-144)	144 (130-235)	374
	05-06	60.8 (57.4-64.5)	57.3 (53.6-60.6)	76.5 (70.9-82.1)	109 (95.0-123)	134 (121-164)	1054
	07-08	70.2 (65.7-74.9)	65.9 (62.4-69.5)	89.0 (83.2-96.5)	128 (112-152)	173 (140-216)	1143
12-19 years	01-02	44.8 (43.4-46.2)	43.8 (42.6-45.0)	56.2 (52.2-59.7)	73.2 (65.2-85.1)	93.4 (79.0-104)	826
	05-06	39.8 (37.8-41.9)	38.1 (36.3-40.3)	51.8 (47.7-56.0)	70.6 (63.1-78.8)	88.9 (79.4-103)	2106
	07-08	43.4 (41.3-45.5)	40.5 (38.6-43.5)	56.0 (52.7-59.2)	76.6 (69.0-86.2)	98.0 (85.4-121)	1134
20 years and older	01-02	48.3 (45.9-50.9)	46.9 (44.2-49.6)	63.8 (61.2-67.7)	90.9 (84.3-98.8)	120 (111-128)	1616
	05-06	41.4 (38.9-43.9)	41.0 (38.8-43.7)	58.6 (54.8-63.1)	85.3 (80.2-91.0)	111 (105-116)	4537
	07-08	46.5 (44.6-48.4)	44.7 (42.5-47.0)	65.0 (60.7-69.2)	96.1 (90.0-102)	125 (117-132)	5351
Gender							
Males	01-02	47.6 (44.7-50.7)	46.1 (43.4-48.7)	61.3 (58.0-64.5)	86.7 (77.1-97.3)	114 (97.3-125)	1335
	05-06	40.1 (37.5-42.9)	39.5 (36.7-42.8)	55.3 (51.6-59.5)	77.2 (70.7-83.0)	95.9 (89.6-102)	3765
	07-08	44.6 (42.8-46.6)	42.8 (40.7-45.1)	60.6 (58.1-64.4)	85.6 (81.1-91.3)	111 (101-121)	3839
Females	01-02	51.9 (49.9-54.1)	51.2 (48.4-53.0)	69.1 (66.7-71.2)	100 (91.7-111)	129 (118-140)	1481
	05-06	45.1 (42.8-47.6)	45.0 (42.4-47.4)	64.4 (60.0-69.6)	96.8 (87.8-105)	128 (117-134)	3932
	07-08	51.0 (48.9-53.2)	50.0 (47.5-52.7)	72.3 (67.8-76.8)	107 (101-116)	146 (129-163)	3789
Race/ethnicity							
Mexican Americans	01-02	50.9 (45.7-56.8)	48.1 (44.9-51.4)	67.4 (60.3-77.6)	97.3 (85.7-117)	135 (100-161)	707
	05-06	44.6 (42.6-46.7)	44.0 (42.2-45.3)	60.4 (58.1-62.9)	89.9 (82.0-95.4)	120 (111-128)	1972
	07-08	48.7 (45.1-52.7)	47.0 (43.5-50.5)	64.8 (59.7-69.9)	93.5 (88.9-101)	128 (108-156)	1505
Non-Hispanic blacks	01-02	38.7 (36.3-41.3)	38.0 (34.6-41.3)	53.5 (50.3-57.4)	70.3 (64.9-79.3)	91.7 (78.9-100)	679
	05-06	32.9 (30.9-35.0)	31.9 (29.8-34.1)	45.4 (41.5-49.6)	64.0 (60.1-68.0)	81.2 (75.3-89.6)	2078
	07-08	35.9 (34.2-37.7)	34.8 (33.3-36.6)	49.0 (44.8-53.6)	69.1 (63.5-77.4)	87.8 (78.1-97.4)	1706
Non-Hispanic whites	01-02	51.4 (49.4-53.4)	49.1 (46.9-51.5)	66.9 (64.2-69.4)	95.2 (87.7-100)	124 (115-132)	1227
	05-06	43.7 (40.8-46.8)	43.9 (40.8-46.8)	61.4 (56.5-66.2)	85.5 (80.6-91.9)	110 (102-116)	3056
	07-08	49.0 (46.7-51.4)	47.4 (44.9-50.5)	68.2 (63.3-73.1)	98.3 (91.2-105)	126 (118-135)	3190

Urinary Perchlorate

Geometric mean and selected percentiles of urine concentrations (in $\mu\text{g/L}$) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	3.72 (3.46-4.01)	3.70 (3.40-3.90)	6.20 (5.60-6.90)	10.0 (8.90-12.0)	14.0 (12.0-18.0)	1618
	03-04	3.39 (3.08-3.73)	3.50 (3.00-4.00)	5.80 (5.20-6.70)	10.0 (8.80-12.0)	14.0 (13.0-16.0)	2522
	05-06	3.47 (3.20-3.76)	3.62 (3.35-3.98)	6.10 (5.67-6.52)	9.76 (9.19-10.5)	13.4 (12.4-14.6)	7697
	07-08	3.88 (3.60-4.18)	3.96 (3.59-4.30)	7.10 (6.50-7.74)	12.0 (11.2-13.0)	17.3 (16.1-18.8)	7629
Age group							
6-11 years	01-02	5.19 (4.44-6.05)	5.50 (4.60-6.70)	8.50 (7.20-10.0)	13.0 (9.80-20.0)	20.0 (13.0-24.0)	374
	03-04	4.54 (3.85-5.36)	4.80 (4.20-5.50)	8.30 (6.00-10.0)	14.0 (9.20-17.0)	17.0 (12.0-30.0)	314
	05-06	4.64 (4.15-5.18)	4.81 (4.26-5.38)	7.67 (6.71-8.60)	11.6 (10.0-12.7)	14.1 (12.4-16.3)	1054
	07-08	4.88 (4.31-5.53)	4.86 (4.22-5.93)	9.11 (7.84-10.3)	14.4 (12.1-16.9)	18.9 (16.1-23.2)	1143
12-19 years	01-02	4.00 (3.62-4.42)	4.60 (4.00-5.00)	7.10 (6.60-7.70)	10.0 (9.70-12.0)	14.0 (12.0-18.0)	828
	03-04	3.81 (3.35-4.33)	4.00 (3.40-4.60)	6.70 (5.80-7.40)	10.0 (8.20-13.0)	14.0 (10.0-19.0)	721
	05-06	4.14 (3.77-4.53)	4.25 (3.94-4.65)	7.00 (6.41-7.69)	10.9 (9.71-12.4)	15.6 (12.4-17.9)	2106
	07-08	4.50 (4.06-4.97)	4.73 (4.39-5.05)	7.93 (6.89-8.89)	13.0 (11.1-15.4)	17.7 (15.8-20.2)	1135
20 years and older	01-02	3.53 (3.24-3.84)	3.70 (3.40-3.90)	6.20 (5.60-6.90)	10.0 (8.90-12.0)	14.0 (12.0-18.0)	1618
	03-04	3.21 (2.89-3.56)	3.40 (2.80-3.80)	5.50 (4.90-6.40)	9.50 (8.30-10.0)	13.0 (12.0-15.0)	1487
	05-06	3.27 (3.01-3.55)	3.42 (3.14-3.72)	5.75 (5.33-6.21)	9.38 (8.73-9.86)	13.1 (12.2-14.5)	4537
	07-08	3.70 (3.43-3.99)	3.69 (3.42-4.09)	6.75 (6.09-7.38)	11.6 (10.8-12.7)	16.8 (15.4-18.7)	5351
Gender							
Males	01-02	4.41 (4.13-4.70)	4.60 (4.40-4.80)	7.40 (6.70-8.30)	12.0 (10.0-13.0)	15.0 (12.0-20.0)	1335
	03-04	3.95 (3.56-4.37)	4.10 (3.60-4.60)	6.70 (5.90-7.90)	12.0 (9.70-13.0)	15.0 (14.0-18.0)	1229
	05-06	3.93 (3.64-4.25)	4.07 (3.77-4.38)	6.50 (6.11-6.92)	10.4 (9.65-11.6)	14.7 (13.2-16.3)	3765
	07-08	4.40 (4.13-4.70)	4.42 (4.12-4.78)	7.90 (7.16-8.46)	13.3 (12.0-14.4)	18.6 (17.0-19.7)	3839
Females	01-02	3.17 (2.89-3.48)	3.30 (2.80-3.60)	5.70 (5.20-6.30)	9.70 (8.60-12.0)	14.0 (12.0-18.0)	1485
	03-04	2.93 (2.62-3.28)	3.00 (2.60-3.40)	5.10 (4.60-5.80)	8.60 (7.20-10.0)	12.0 (9.20-16.0)	1293
	05-06	3.08 (2.80-3.38)	3.24 (2.96-3.51)	5.55 (5.14-6.07)	9.20 (8.22-10.3)	12.4 (11.3-14.2)	3932
	07-08	3.42 (3.11-3.76)	3.44 (3.10-3.79)	6.41 (5.95-7.01)	10.8 (10.0-12.0)	16.0 (14.1-17.9)	3790
Race/ethnicity							
Mexican Americans	01-02	4.23 (3.65-4.90)	4.60 (3.90-5.20)	7.40 (6.10-8.80)	13.0 (9.90-14.0)	15.0 (13.0-19.0)	708
	03-04	3.96 (3.63-4.32)	4.20 (3.70-4.60)	6.50 (5.60-7.90)	12.0 (9.50-13.0)	16.0 (13.0-18.0)	617
	05-06	4.02 (3.77-4.28)	4.16 (3.84-4.46)	6.93 (6.62-7.26)	11.3 (10.4-12.0)	15.7 (14.6-16.8)	1972
	07-08	4.20 (3.95-4.46)	4.38 (4.08-4.65)	7.32 (6.73-7.95)	12.1 (10.9-14.0)	16.7 (15.0-17.7)	1505
Non-Hispanic blacks	01-02	3.69 (3.22-4.23)	3.90 (3.30-4.30)	6.20 (5.30-7.30)	9.70 (8.20-13.0)	16.0 (12.0-21.0)	681
	03-04	3.38 (3.05-3.74)	3.40 (3.00-3.70)	5.70 (4.80-6.60)	9.00 (7.90-12.0)	14.0 (9.80-18.0)	652
	05-06	3.48 (3.19-3.80)	3.54 (3.24-3.94)	5.95 (5.33-6.65)	9.99 (9.02-11.0)	14.4 (12.8-15.7)	2078
	07-08	3.82 (3.37-4.33)	3.82 (3.37-4.48)	6.96 (6.18-7.97)	12.0 (10.6-13.8)	17.2 (14.3-20.6)	1707
Non-Hispanic whites	01-02	3.70 (3.35-4.08)	3.90 (3.60-4.30)	6.60 (6.00-7.40)	10.0 (9.20-12.0)	15.0 (12.0-19.0)	1228
	03-04	3.43 (3.04-3.86)	3.50 (2.90-4.20)	5.90 (5.10-7.10)	9.90 (8.50-12.0)	14.0 (12.0-16.0)	1092
	05-06	3.42 (3.08-3.80)	3.59 (3.23-4.04)	6.05 (5.52-6.55)	9.62 (8.90-10.5)	13.2 (11.9-14.8)	3056
	07-08	3.84 (3.52-4.18)	3.93 (3.48-4.36)	7.07 (6.37-7.77)	11.7 (10.8-12.7)	16.8 (15.3-18.4)	3190

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06, and 07-08 are 0.05, 0.05, 0.05, and 0.05 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Perchlorate_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Perchlorate_FactSheet.html

Urinary Perchlorate (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	3.74 (3.51-3.99)	3.70 (3.40-3.90)	6.20 (5.60-6.90)	10.0 (8.90-12.0)	14.0 (12.0-18.0)	1618
	03-04	3.31 (3.04-3.59)	3.18 (2.90-3.45)	5.19 (4.80-5.55)	8.86 (7.69-10.0)	12.4 (11.2-14.5)	2519
	05-06	3.46 (3.27-3.66)	3.37 (3.17-3.59)	5.42 (5.08-5.86)	8.56 (8.19-9.17)	11.6 (10.4-13.4)	7697
	07-08	4.00 (3.69-4.33)	3.89 (3.59-4.26)	6.59 (6.10-7.13)	10.9 (10.1-11.8)	15.7 (14.6-16.8)	7628
Age group							
6-11 years	01-02	6.01 (5.49-6.57)	5.50 (4.60-6.70)	8.50 (7.20-10.0)	13.0 (9.80-20.0)	20.0 (13.0-24.0)	374
	03-04	5.52 (4.86-6.28)	5.34 (4.92-5.78)	7.64 (6.67-10.5)	13.5 (9.66-21.3)	21.3 (12.0-46.9)	314
	05-06	5.51 (5.06-6.00)	5.66 (5.08-6.22)	8.39 (7.61-9.25)	12.3 (10.3-13.6)	15.4 (14.3-17.0)	1054
	07-08	6.21 (5.52-6.99)	6.34 (5.56-7.10)	9.78 (8.63-11.3)	14.7 (13.3-16.0)	20.6 (16.4-25.0)	1143
12-19 years	01-02	3.10 (2.77-3.46)	4.60 (4.00-5.00)	7.10 (6.60-7.70)	10.0 (9.70-12.0)	14.0 (12.0-18.0)	828
	03-04	2.85 (2.59-3.14)	2.83 (2.62-3.17)	4.39 (3.88-4.67)	6.63 (5.56-7.82)	8.62 (7.24-11.4)	719
	05-06	3.14 (2.89-3.41)	3.17 (2.87-3.40)	4.86 (4.32-5.41)	7.30 (6.40-8.43)	9.54 (8.46-11.0)	2106
	07-08	3.52 (3.27-3.78)	3.43 (3.07-3.80)	5.62 (5.13-6.11)	8.84 (7.70-9.73)	12.6 (10.4-15.8)	1134
20 years and older	01-02	3.64 (3.37-3.93)	3.70 (3.40-3.90)	6.20 (5.60-6.90)	10.0 (8.90-12.0)	14.0 (12.0-18.0)	1618
	03-04	3.18 (2.90-3.49)	3.02 (2.74-3.33)	4.94 (4.53-5.35)	8.64 (7.18-10.0)	11.7 (10.3-15.2)	1486
	05-06	3.34 (3.15-3.53)	3.24 (3.08-3.45)	5.12 (4.82-5.45)	8.24 (7.67-8.64)	11.4 (10.0-13.0)	4537
	07-08	3.88 (3.57-4.22)	3.77 (3.49-4.14)	6.35 (5.85-6.91)	10.5 (9.64-11.5)	15.4 (13.6-16.9)	5351
Gender							
Males	01-02	3.57 (3.36-3.79)	4.60 (4.40-4.80)	7.40 (6.70-8.30)	12.0 (10.0-13.0)	15.0 (12.0-20.0)	1335
	03-04	3.23 (2.96-3.52)	3.09 (2.83-3.33)	5.00 (4.60-5.55)	8.00 (6.86-10.2)	12.5 (10.2-15.3)	1228
	05-06	3.25 (3.08-3.44)	3.16 (2.99-3.36)	4.98 (4.71-5.37)	8.07 (7.60-8.58)	11.0 (10.0-12.9)	3765
	07-08	3.78 (3.47-4.12)	3.64 (3.28-4.03)	6.29 (5.72-6.85)	10.3 (9.28-11.3)	14.4 (13.1-15.5)	3839
Females	01-02	3.92 (3.57-4.30)	3.30 (2.80-3.60)	5.70 (5.20-6.30)	9.70 (8.60-12.0)	14.0 (12.0-18.0)	1485
	03-04	3.38 (3.02-3.78)	3.28 (2.90-3.61)	5.24 (4.73-6.00)	9.40 (7.24-11.4)	12.4 (10.7-15.2)	1291
	05-06	3.67 (3.43-3.93)	3.58 (3.31-3.91)	5.86 (5.39-6.19)	9.21 (8.35-9.77)	12.1 (10.5-14.5)	3932
	07-08	4.22 (3.90-4.56)	4.16 (3.81-4.55)	6.89 (6.40-7.54)	11.5 (10.4-12.6)	16.6 (14.5-18.6)	3789
Race/ethnicity							
Mexican Americans	01-02	3.96 (3.39-4.63)	4.60 (3.90-5.20)	7.40 (6.10-8.80)	13.0 (9.90-14.0)	15.0 (13.0-19.0)	708
	03-04	3.59 (3.32-3.88)	3.45 (3.10-3.68)	5.69 (4.82-6.45)	9.77 (8.00-11.2)	13.5 (11.2-20.0)	616
	05-06	3.75 (3.56-3.95)	3.60 (3.42-3.81)	5.97 (5.53-6.48)	9.83 (9.19-10.4)	12.7 (11.3-14.5)	1972
	07-08	4.20 (3.96-4.46)	4.17 (3.81-4.58)	6.95 (6.33-7.73)	11.8 (10.4-12.3)	15.7 (14.4-17.6)	1505
Non-Hispanic blacks	01-02	2.66 (2.35-3.01)	3.90 (3.30-4.30)	6.20 (5.30-7.30)	9.70 (8.20-13.0)	16.0 (12.0-21.0)	681
	03-04	2.33 (2.11-2.57)	2.24 (2.02-2.47)	3.60 (3.22-4.13)	6.54 (4.91-8.47)	9.02 (6.84-11.8)	651
	05-06	2.50 (2.32-2.69)	2.40 (2.24-2.61)	3.97 (3.69-4.26)	6.71 (6.30-7.13)	9.41 (8.53-10.4)	2078
	07-08	2.89 (2.60-3.21)	2.76 (2.47-3.17)	4.92 (4.36-5.54)	8.63 (7.43-9.78)	11.8 (10.6-13.6)	1706
Non-Hispanic whites	01-02	3.95 (3.64-4.29)	3.90 (3.60-4.30)	6.60 (6.00-7.40)	10.0 (9.20-12.0)	15.0 (12.0-19.0)	1228
	03-04	3.53 (3.16-3.94)	3.35 (3.01-3.68)	5.29 (4.81-5.96)	9.23 (7.61-11.0)	12.8 (11.0-16.5)	1091
	05-06	3.63 (3.36-3.91)	3.51 (3.20-3.85)	5.60 (5.10-6.10)	8.58 (7.98-9.42)	11.8 (10.0-14.3)	3056
	07-08	4.17 (3.83-4.54)	4.06 (3.71-4.52)	6.63 (6.18-7.23)	10.8 (9.86-11.8)	15.5 (14.0-16.8)	3190

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Perchlorate_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Perchlorate_FactSheet.html

Urinary Thiocyanate

Geometric mean and selected percentiles of urine concentrations (in mg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50 th	75 th	90 th	95 th	
Total	01-02	1.52 (1.44-1.61)	1.50 (1.40-1.60)	3.60 (3.00-4.10)	7.90 (6.90-9.00)	12.0 (10.0-13.0)	1617
	05-06	1.23 (1.12-1.35)	1.18 (1.08-1.31)	2.55 (2.31-2.83)	6.11 (5.10-6.85)	9.04 (8.14-9.93)	7697
	07-08	1.35 (1.26-1.45)	1.32 (1.23-1.42)	2.77 (2.47-3.13)	6.36 (5.39-7.33)	9.54 (8.58-11.1)	7629
Age group							
6-11 years	01-02	1.35 (1.20-1.51)	1.50 (1.30-1.70)	2.50 (1.90-3.00)	3.70 (3.30-4.10)	4.90 (3.90-5.60)	374
	05-06	.980 (.892-1.08)	1.03 (.916-1.15)	1.72 (1.55-1.92)	2.65 (2.31-3.09)	3.45 (2.96-4.21)	1054
	07-08	1.10 (.989-1.22)	1.12 (1.02-1.24)	1.88 (1.68-2.07)	2.92 (2.48-3.52)	4.08 (3.35-5.30)	1143
12-19 years	01-02	1.59 (1.43-1.77)	1.50 (1.40-1.70)	2.90 (2.50-3.30)	5.30 (4.30-7.40)	7.70 (5.90-10.0)	826
	05-06	1.24 (1.11-1.39)	1.29 (1.15-1.47)	2.27 (1.99-2.62)	3.84 (3.18-4.65)	6.07 (4.76-6.95)	2106
	07-08	1.29 (1.15-1.46)	1.35 (1.14-1.53)	2.39 (2.06-2.69)	3.93 (3.36-4.57)	6.14 (4.55-7.27)	1135
20 years and older	01-02	1.54 (1.43-1.65)	1.50 (1.40-1.60)	3.60 (3.00-4.10)	7.90 (6.90-9.00)	12.0 (10.0-13.0)	1617
	05-06	1.26 (1.14-1.40)	1.19 (1.08-1.34)	2.82 (2.50-3.15)	6.87 (6.06-7.58)	9.79 (8.78-10.7)	4537
	07-08	1.39 (1.29-1.50)	1.34 (1.25-1.46)	3.11 (2.68-3.57)	7.21 (6.05-8.31)	10.8 (8.99-12.0)	5351
Gender							
Males	01-02	1.86 (1.67-2.07)	1.80 (1.60-2.10)	4.00 (3.40-4.90)	8.40 (7.00-10.0)	12.0 (10.0-13.0)	1335
	05-06	1.49 (1.36-1.64)	1.44 (1.27-1.62)	3.10 (2.82-3.56)	6.90 (6.13-7.64)	9.93 (8.99-10.7)	3765
	07-08	1.61 (1.50-1.72)	1.60 (1.50-1.71)	3.32 (2.96-3.77)	7.27 (6.01-8.22)	10.9 (9.04-12.3)	3839
Females	01-02	1.26 (1.16-1.36)	1.30 (1.00-1.40)	2.50 (2.40-2.80)	6.00 (5.20-6.70)	8.70 (7.20-10.0)	1482
	05-06	1.02 (.920-1.13)	.995 (.886-1.13)	2.07 (1.86-2.25)	4.61 (3.81-5.98)	8.04 (7.26-8.78)	3932
	07-08	1.14 (1.04-1.24)	1.09 (.991-1.19)	2.31 (2.06-2.62)	5.51 (4.58-6.52)	8.59 (7.36-9.85)	3790
Race/ethnicity							
Mexican Americans	01-02	1.14 (.977-1.33)	1.20 (.940-1.40)	2.20 (1.80-2.60)	3.90 (3.00-4.90)	5.70 (4.20-6.60)	707
	05-06	.823 (.744-.912)	.844 (.763-.937)	1.51 (1.36-1.71)	2.70 (2.26-3.22)	4.48 (3.49-5.40)	1972
	07-08	.993 (.926-1.06)	1.04 (.987-1.11)	1.88 (1.75-2.02)	3.30 (2.98-3.52)	4.56 (3.92-5.43)	1505
Non-Hispanic blacks	01-02	1.91 (1.72-2.11)	2.00 (1.80-2.20)	3.90 (3.30-4.90)	7.60 (6.20-10.0)	10.0 (7.00-15.0)	679
	05-06	1.50 (1.32-1.72)	1.51 (1.36-1.68)	3.13 (2.56-3.81)	6.94 (5.66-8.09)	10.3 (8.01-12.4)	2078
	07-08	1.73 (1.61-1.86)	1.73 (1.61-1.86)	3.78 (3.53-4.23)	8.29 (7.25-9.07)	12.8 (11.2-14.5)	1707
Non-Hispanic whites	01-02	1.56 (1.45-1.68)	1.50 (1.40-1.70)	3.40 (2.80-3.80)	7.20 (6.30-8.50)	12.0 (8.90-13.0)	1228
	05-06	1.30 (1.18-1.44)	1.25 (1.12-1.40)	2.74 (2.46-3.01)	6.53 (5.57-7.31)	9.43 (8.58-10.4)	3056
	07-08	1.42 (1.29-1.56)	1.41 (1.28-1.53)	2.98 (2.52-3.44)	6.67 (5.24-8.14)	9.71 (8.32-11.9)	3190

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 05-06, and 07-08 are 0.02, 0.02, and 0.02 respectively.

Urinary Thiocyanate (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in mg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50 th	75 th	90 th	95 th	
Total	01-02	1.53 (1.43-1.64)	1.47 (1.37-1.61)	3.35 (2.95-3.74)	7.34 (6.67-7.81)	10.3 (8.81-11.3)	1616
	05-06	1.23 (1.13-1.33)	1.15 (1.05-1.27)	2.44 (2.18-2.68)	5.67 (5.16-6.28)	8.23 (7.71-8.98)	7697
	07-08	1.39 (1.28-1.51)	1.33 (1.22-1.43)	2.72 (2.42-3.04)	6.11 (5.25-7.30)	9.59 (8.03-11.5)	7628
Age group							
6-11 years	01-02	1.56 (1.35-1.80)	1.55 (1.35-1.82)	2.67 (2.33-3.13)	4.46 (3.79-5.20)	5.33 (4.58-6.16)	374
	05-06	1.16 (1.06-1.28)	1.18 (1.06-1.34)	2.03 (1.81-2.32)	3.15 (2.83-3.43)	4.25 (3.49-4.58)	1054
	07-08	1.40 (1.27-1.54)	1.42 (1.24-1.59)	2.28 (2.02-2.53)	3.74 (3.29-4.26)	5.23 (4.39-6.65)	1143
12-19 years	01-02	1.23 (1.12-1.35)	1.14 (1.08-1.24)	2.12 (1.78-2.45)	3.88 (3.28-4.91)	5.75 (3.93-8.13)	825
	05-06	.941 (.867-1.02)	.937 (.855-1.03)	1.69 (1.51-1.86)	2.95 (2.43-3.64)	4.15 (3.64-5.00)	2106
	07-08	1.01 (.910-1.12)	1.04 (.906-1.17)	1.75 (1.60-1.92)	2.94 (2.49-3.54)	4.69 (3.64-5.44)	1134
20 years and older	01-02	1.59 (1.47-1.71)	1.47 (1.37-1.61)	3.35 (2.95-3.74)	7.34 (6.67-7.81)	10.3 (8.81-11.3)	1616
	05-06	1.29 (1.17-1.42)	1.20 (1.08-1.33)	2.68 (2.37-3.10)	6.43 (5.88-7.07)	8.99 (8.23-10.2)	4537
	07-08	1.46 (1.34-1.59)	1.37 (1.27-1.47)	3.00 (2.69-3.45)	7.02 (5.82-8.26)	10.5 (8.63-12.9)	5351
Gender							
Males	01-02	1.51 (1.37-1.67)	1.42 (1.31-1.59)	3.33 (3.00-3.67)	6.38 (5.81-6.92)	8.51 (7.41-9.72)	1335
	05-06	1.24 (1.12-1.37)	1.18 (1.05-1.32)	2.61 (2.26-3.01)	5.67 (5.00-6.43)	8.17 (7.15-9.13)	3765
	07-08	1.38 (1.28-1.50)	1.32 (1.23-1.42)	2.77 (2.42-3.08)	6.13 (5.47-6.97)	9.27 (7.82-11.1)	3839
Females	01-02	1.55 (1.43-1.69)	1.45 (1.33-1.58)	2.79 (2.42-3.25)	7.07 (5.86-7.69)	10.0 (8.18-11.1)	1480
	05-06	1.22 (1.12-1.33)	1.13 (1.04-1.24)	2.27 (2.00-2.54)	5.62 (4.91-6.37)	8.40 (7.20-10.5)	3932
	07-08	1.40 (1.27-1.55)	1.34 (1.19-1.45)	2.69 (2.40-3.03)	6.07 (4.89-7.80)	9.88 (7.93-12.1)	3789
Race/ethnicity							
Mexican Americans	01-02	1.07 (.898-1.27)	1.06 (.819-1.29)	1.97 (1.51-2.50)	3.62 (2.86-4.52)	5.10 (3.52-7.33)	707
	05-06	.768 (.685-.861)	.773 (.684-.881)	1.37 (1.20-1.59)	2.44 (2.19-2.85)	3.66 (3.28-4.27)	1972
	07-08	.995 (.923-1.07)	1.03 (.918-1.14)	1.88 (1.71-2.05)	3.27 (2.77-3.76)	4.59 (3.86-5.59)	1505
Non-Hispanic blacks	01-02	1.37 (1.18-1.59)	1.37 (1.18-1.61)	2.80 (2.21-3.43)	5.67 (4.17-6.74)	7.64 (5.91-11.3)	678
	05-06	1.08 (.972-1.20)	1.05 (.972-1.16)	2.15 (1.84-2.57)	4.77 (3.93-5.61)	6.91 (5.84-7.80)	2078
	07-08	1.31 (1.21-1.41)	1.25 (1.18-1.36)	2.81 (2.55-3.24)	6.19 (5.29-6.77)	8.74 (7.44-9.59)	1706
Non-Hispanic whites	01-02	1.67 (1.55-1.80)	1.55 (1.42-1.73)	3.29 (3.00-3.73)	7.09 (6.38-7.95)	10.3 (8.78-11.1)	1227
	05-06	1.38 (1.25-1.52)	1.30 (1.14-1.44)	2.78 (2.48-3.12)	6.31 (5.67-7.08)	9.00 (8.13-10.5)	3056
	07-08	1.55 (1.40-1.70)	1.45 (1.32-1.58)	2.96 (2.56-3.53)	7.03 (5.71-8.56)	10.9 (8.70-13.1)	3190

Serum Perfluorobutane sulfonic acid (PFBuS)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	2094
	05-06	*	< LOD	< LOD	< LOD	.100 (<LOD-.200)	2120
	07-08	*	< LOD	< LOD	< LOD	< LOD	2100
	09-10	*	< LOD	< LOD	< LOD	< LOD	2233
Age group 12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	640
	05-06	*	< LOD	< LOD	< LOD	.100 (<LOD-.100)	640
	07-08	*	< LOD	< LOD	< LOD	< LOD	357
	09-10	*	< LOD	< LOD	< LOD	< LOD	364
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1454
	05-06	*	< LOD	< LOD	< LOD	.100 (<LOD-.200)	1480
	07-08	*	< LOD	< LOD	< LOD	< LOD	1743
	09-10	*	< LOD	< LOD	< LOD	< LOD	1869
Gender Males	03-04	*	< LOD	< LOD	< LOD	< LOD	1053
	05-06	*	< LOD	< LOD	< LOD	.100 (<LOD-.200)	1048
	07-08	*	< LOD	< LOD	< LOD	< LOD	1059
	09-10	*	< LOD	< LOD	< LOD	< LOD	1075
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1041
	05-06	*	< LOD	< LOD	< LOD	.100 (<LOD-.200)	1072
	07-08	*	< LOD	< LOD	< LOD	< LOD	1041
	09-10	*	< LOD	< LOD	< LOD	< LOD	1158
Race/ethnicity Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	485
	05-06	*	< LOD	< LOD	< LOD	.100 (<LOD-.200)	499
	07-08	*	< LOD	< LOD	< LOD	< LOD	391
	09-10	*	< LOD	< LOD	< LOD	< LOD	461
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	538
	05-06	*	< LOD	< LOD	< LOD	.100 (<LOD-.100)	544
	07-08	*	< LOD	< LOD	< LOD	< LOD	419
	09-10	*	< LOD	< LOD	< LOD	< LOD	391
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	962
	05-06	*	< LOD	< LOD	< LOD	.100 (<LOD-.400)	935
	07-08	*	< LOD	< LOD	< LOD	< LOD	931
	09-10	*	< LOD	< LOD	< LOD	< LOD	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.4, 0.1, 0.1, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum Perfluorodecanoic acid (PFDeA)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	.200 (.200-.200)	.200 (.200-.300)	.400 (.300-.700)	.600 (.400-1.10)	1562
	03-04	*	< LOD	.300 (<LOD-.500)	.600 (.400-1.10)	.900 (.500-1.80)	2094
	05-06	.355 (.297-.423)	.300 (.300-.400)	.500 (.400-.700)	.900 (.600-1.60)	1.50 (.900-2.60)	2120
	07-08	.286 (.264-.309)	.300 (.300-.300)	.400 (.400-.500)	.700 (.600-.700)	.900 (.800-1.00)	2100
	09-10	.279 (.258-.303)	.300 (.300-.300)	.400 (.400-.500)	.700 (.600-.800)	.900 (.800-1.10)	2233
Age group 12-19 years	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.300 (.200-.400)	.400 (.300-.600)	543
	03-04	*	< LOD	< LOD	.500 (<LOD-1.00)	.800 (.300-1.20)	640
	05-06	.295 (.258-.338)	.300 (.200-.300)	.500 (.400-.500)	.600 (.500-.800)	.800 (.600-1.60)	640
	07-08	.231 (.214-.248)	.200 (.200-.300)	.300 (.300-.400)	.500 (.400-.500)	.600 (.500-.700)	357
	09-10	.220 (.198-.245)	.200 (.200-.200)	.300 (.300-.300)	.400 (.400-.600)	.600 (.400-.800)	364
20 years and older	99-00	*	.200 (.200-.200)	.300 (.200-.300)	.400 (.300-.800)	.600 (.400-1.50)	1019
	03-04	*	< LOD	.400 (<LOD-.500)	.700 (.400-1.00)	.900 (.500-1.80)	1454
	05-06	.364 (.303-.438)	.300 (.300-.400)	.500 (.400-.700)	1.00 (.600-1.70)	1.50 (.900-2.60)	1480
	07-08	.295 (.271-.321)	.300 (.300-.300)	.400 (.400-.500)	.700 (.600-.800)	.900 (.800-1.10)	1743
	09-10	.289 (.265-.314)	.300 (.300-.300)	.400 (.400-.500)	.700 (.600-.800)	.900 (.800-1.20)	1869
Gender Males	99-00	*	.200 (.200-.200)	.300 (.200-.300)	.400 (.300-.700)	.500 (.300-1.90)	743
	03-04	*	< LOD	.400 (<LOD-.500)	.800 (.400-1.40)	1.10 (.600-2.10)	1053
Females	05-06	.381 (.318-.456)	.400 (.300-.400)	.600 (.400-.800)	1.00 (.600-2.20)	1.70 (1.00-2.60)	1048
	07-08	.306 (.283-.331)	.300 (.300-.300)	.400 (.400-.500)	.700 (.600-.800)	.900 (.800-1.20)	1059
	09-10	.289 (.264-.318)	.300 (.300-.300)	.400 (.400-.500)	.600 (.500-.800)	.800 (.600-1.20)	1075
	99-00	*	.200 (.200-.200)	.200 (.200-.300)	.400 (.300-.800)	.600 (.400-1.40)	819
	03-04	*	< LOD	.300 (<LOD-.400)	.500 (.400-.800)	.800 (.500-1.20)	1041
Race/ethnicity Mexican Americans	05-06	.331 (.277-.396)	.300 (.300-.400)	.500 (.400-.600)	.900 (.600-1.40)	1.30 (.800-2.30)	1072
	07-08	.267 (.245-.292)	.300 (.200-.300)	.400 (.400-.500)	.600 (.500-.800)	.800 (.700-1.10)	1041
	09-10	.270 (.248-.295)	.300 (.200-.300)	.400 (.400-.500)	.700 (.600-.800)	1.00 (.800-1.10)	1158
	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.300 (.200-.300)	.300 (.200-1.50)	584
	03-04	*	< LOD	< LOD	.500 (.400-.500)	.600 (.500-.800)	485
Non-Hispanic blacks	05-06	.283 (.245-.327)	.300 (.200-.300)	.400 (.300-.500)	.600 (.500-1.00)	1.00 (.500-2.40)	499
	07-08	.253 (.222-.289)	.200 (.200-.300)	.400 (.300-.500)	.600 (.400-.700)	.600 (.500-1.40)	391
	09-10	.242 (.215-.272)	.200 (.200-.300)	.400 (.300-.400)	.500 (.500-.600)	.700 (.600-.800)	461
	99-00	*	.200 (.200-.200)	.400 (.200-.700)	.800 (.500-1.20)	1.10 (.600-2.30)	303
	03-04	*	< LOD	.400 (<LOD-.700)	.800 (.400-1.50)	1.00 (.500-3.10)	538
Non-Hispanic whites	05-06	.405 (.309-.531)	.400 (.300-.500)	.600 (.400-.900)	1.20 (.600-2.90)	2.30 (1.10-3.70)	544
	07-08	.331 (.298-.368)	.300 (.300-.400)	.500 (.400-.600)	.800 (.600-.900)	1.00 (.800-1.50)	419
	09-10	.336 (.298-.379)	.300 (.300-.300)	.500 (.400-.600)	.800 (.600-1.00)	1.10 (.800-2.60)	391
	99-00	*	.200 (.200-.200)	.300 (.200-.300)	.400 (.300-.600)	.500 (.400-.800)	519
	03-04	*	< LOD	.300 (<LOD-.500)	.600 (.400-1.00)	.900 (.500-1.80)	962
05-06	.350 (.294-.417)	.300 (.300-.400)	.500 (.400-.700)	.900 (.600-1.50)	1.40 (.800-2.30)	935	
07-08	.276 (.254-.301)	.300 (.200-.300)	.400 (.400-.500)	.600 (.500-.700)	.900 (.700-.900)	931	
09-10	.270 (.240-.304)	.300 (.200-.300)	.400 (.300-.500)	.600 (.500-.800)	.800 (.600-1.10)	1031	

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.2, 0.3, 0.2, 0.2, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum Perfluorododecanoic acid (PFDoA)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	1562
	03-04	*	< LOD	< LOD	< LOD	< LOD	2094
	05-06	*	< LOD	< LOD	< LOD	< LOD	2120
	07-08	*	< LOD	< LOD	< LOD	< LOD	2100
	09-10	*	< LOD	< LOD	< LOD	< LOD	2233
Age group 12-19 years	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	543
	03-04	*	< LOD	< LOD	< LOD	< LOD	640
	05-06	*	< LOD	< LOD	< LOD	< LOD	640
	07-08	*	< LOD	< LOD	< LOD	< LOD	357
	09-10	*	< LOD	< LOD	< LOD	< LOD	364
20 years and older	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	1019
	03-04	*	< LOD	< LOD	< LOD	< LOD	1454
	05-06	*	< LOD	< LOD	< LOD	< LOD	1480
	07-08	*	< LOD	< LOD	< LOD	< LOD	1743
	09-10	*	< LOD	< LOD	< LOD	.100 (<LOD-.100)	1869
Gender Males	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	743
	03-04	*	< LOD	< LOD	< LOD	< LOD	1053
	05-06	*	< LOD	< LOD	< LOD	< LOD	1048
	07-08	*	< LOD	< LOD	< LOD	< LOD	1059
	09-10	*	< LOD	< LOD	< LOD	< LOD	1075
Females	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	819
	03-04	*	< LOD	< LOD	< LOD	< LOD	1041
	05-06	*	< LOD	< LOD	< LOD	< LOD	1072
	07-08	*	< LOD	< LOD	< LOD	< LOD	1041
	09-10	*	< LOD	< LOD	< LOD	.100 (<LOD-.100)	1158
Race/ethnicity Mexican Americans	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	584
	03-04	*	< LOD	< LOD	< LOD	< LOD	485
	05-06	*	< LOD	< LOD	< LOD	< LOD	499
	07-08	*	< LOD	< LOD	< LOD	< LOD	391
	09-10	*	< LOD	< LOD	< LOD	< LOD	461
Non-Hispanic blacks	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	303
	03-04	*	< LOD	< LOD	< LOD	< LOD	538
	05-06	*	< LOD	< LOD	< LOD	< LOD	544
	07-08	*	< LOD	< LOD	< LOD	< LOD	419
	09-10	*	< LOD	< LOD	< LOD	.100 (<LOD-.400)	391
Non-Hispanic whites	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	519
	03-04	*	< LOD	< LOD	< LOD	< LOD	962
	05-06	*	< LOD	< LOD	< LOD	< LOD	935
	07-08	*	< LOD	< LOD	< LOD	< LOD	931
	09-10	*	< LOD	< LOD	< LOD	< LOD	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.2, 1.0, 0.2, 0.2, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum Perfluoroheptanoic acid (PFHpA)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	.400 (.400-.400)	.400 (.400-.400)	.500 (.400-.600)	.700 (.500-1.00)	1562
	03-04	*	< LOD	< LOD	< LOD	.400 (<LOD-.500)	2094
	05-06	*	< LOD	< LOD	< LOD	.700 (<LOD-1.70)	2120
	07-08	*	< LOD	< LOD	< LOD	.500 (.400-.600)	2100
	09-10	*	< LOD	< LOD	.200 (.100-.200)	.200 (.200-.300)	2233
Age group 12-19 years	99-00	*	.400 (.400-.400)	.400 (.400-.500)	.800 (.600-1.10)	1.20 (.800-1.80)	543
	03-04	*	< LOD	< LOD	.400 (<LOD-.600)	.600 (.500-.900)	640
	05-06	*	< LOD	< LOD	.700 (<LOD-1.10)	1.10 (.600-2.70)	640
	07-08	*	< LOD	< LOD	.500 (<LOD-.600)	.600 (.500-.700)	357
	09-10	*	< LOD	.100 (.100-.200)	.200 (.200-.400)	.400 (.200-.500)	364
20 years and older	99-00	*	.400 (.400-.400)	.400 (.400-.400)	.400 (.400-.500)	.600 (.400-.900)	1019
	03-04	*	< LOD	< LOD	< LOD	< LOD	1454
	05-06	*	< LOD	< LOD	< LOD	.600 (<LOD-2.10)	1480
	07-08	*	< LOD	< LOD	< LOD	.500 (<LOD-.500)	1743
	09-10	*	< LOD	< LOD	.100 (.100-.200)	.200 (.200-.200)	1869
Gender Males	99-00	*	.400 (.400-.400)	.400 (.400-.400)	.500 (.400-.600)	.700 (.500-1.00)	743
	03-04	*	< LOD	< LOD	< LOD	.300 (<LOD-.400)	1053
Females	05-06	*	< LOD	< LOD	.400 (<LOD-1.00)	.800 (<LOD-2.10)	1048
	07-08	*	< LOD	< LOD	< LOD	.600 (.400-.700)	1059
	09-10	*	< LOD	< LOD	.200 (.100-.200)	.200 (.200-.300)	1075
	99-00	*	.400 (.400-.400)	.400 (.400-.400)	.500 (.400-.700)	.700 (.400-1.20)	819
	03-04	*	< LOD	< LOD	< LOD	.400 (.300-.600)	1041
Race/ethnicity Mexican Americans	05-06	*	< LOD	< LOD	< LOD	.600 (<LOD-1.30)	1072
	07-08	*	< LOD	< LOD	< LOD	.500 (.400-.500)	1041
	09-10	*	< LOD	< LOD	.200 (.100-.200)	.200 (.200-.300)	1158
	99-00	*	.400 (.400-.400)	.400 (.400-.400)	.500 (.400-.900)	.900 (.500-1.30)	584
	03-04	*	< LOD	< LOD	< LOD	.500 (<LOD-.900)	485
Non-Hispanic blacks	05-06	*	< LOD	< LOD	< LOD	.500 (<LOD-1.50)	499
	07-08	*	< LOD	< LOD	< LOD	.500 (<LOD-.500)	391
	09-10	*	< LOD	< LOD	.200 (.100-.200)	.200 (.200-.300)	461
	99-00	*	.400 (.400-.400)	.400 (.400-.400)	.400 (.400-.500)	.600 (.400-1.20)	303
	03-04	*	< LOD	< LOD	< LOD	< LOD	538
Non-Hispanic whites	05-06	*	< LOD	< LOD	< LOD	.500 (<LOD-1.80)	544
	07-08	*	< LOD	< LOD	< LOD	.600 (.400-.700)	419
	09-10	*	< LOD	< LOD	.100 (.100-.200)	.200 (.200-.200)	391
	99-00	*	.400 (.400-.400)	.400 (.400-.400)	.400 (.400-.600)	.600 (.500-.900)	519
	03-04	*	< LOD	< LOD	< LOD	.300 (<LOD-.500)	962
Non-Hispanic whites	05-06	*	< LOD	< LOD	< LOD	.800 (<LOD-2.10)	935
	07-08	*	< LOD	< LOD	< LOD	.500 (.400-.700)	931
	09-10	*	< LOD	< LOD	.200 (.100-.200)	.200 (.200-.300)	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.4, 0.3, 0.4, 0.4, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum Perfluorohexane sulfonic acid (PFHxS)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	2.13 (1.91-2.38)	2.10 (1.80-2.30)	3.40 (3.10-4.00)	5.80 (5.20-6.90)	8.70 (7.00-10.0)	1562
	03-04	1.93 (1.73-2.16)	1.90 (1.70-2.10)	3.30 (2.80-3.90)	5.90 (4.80-7.20)	8.30 (7.10-9.70)	2094
	05-06	1.67 (1.42-1.98)	1.80 (1.50-2.10)	3.20 (2.60-4.00)	5.40 (4.40-7.10)	8.30 (5.80-11.9)	2120
	07-08	1.95 (1.76-2.17)	2.00 (1.80-2.10)	3.50 (3.10-4.00)	5.90 (4.90-8.30)	9.80 (6.10-15.2)	2100
	09-10	1.66 (1.51-1.82)	1.70 (1.50-1.90)	3.00 (2.60-3.30)	4.80 (4.40-5.30)	6.90 (5.90-7.60)	2233
Age group 12-19 years	99-00	2.69 (2.14-3.39)	2.50 (2.00-3.70)	5.00 (3.60-6.50)	8.00 (5.80-12.9)	12.9 (6.90-16.4)	543
	03-04	2.44 (2.05-2.90)	2.40 (1.80-3.20)	4.90 (4.00-6.30)	9.50 (6.80-12.5)	13.3 (9.90-19.6)	640
	05-06	2.09 (1.74-2.52)	2.40 (2.00-2.90)	4.30 (3.70-5.40)	9.50 (6.40-13.1)	14.1 (9.80-17.8)	640
	07-08	2.40 (2.09-2.75)	2.30 (1.80-2.70)	4.30 (3.70-5.70)	9.90 (6.50-12.5)	15.9 (11.1-21.5)	357
	09-10	2.03 (1.77-2.33)	1.90 (1.60-2.10)	3.80 (3.20-4.20)	8.10 (6.30-9.70)	12.3 (8.60-19.0)	364
20 years and older	99-00	2.05 (1.85-2.28)	2.00 (1.80-2.30)	3.30 (3.00-3.70)	5.40 (4.90-5.90)	8.20 (6.70-9.50)	1019
	03-04	1.86 (1.67-2.08)	1.80 (1.60-2.10)	3.00 (2.60-3.60)	5.50 (4.50-6.70)	7.60 (6.30-9.40)	1454
	05-06	1.62 (1.37-1.92)	1.70 (1.50-2.10)	3.10 (2.50-3.80)	5.20 (4.20-6.50)	7.20 (5.10-11.6)	1480
	07-08	1.90 (1.69-2.13)	1.90 (1.70-2.10)	3.30 (2.90-3.90)	5.80 (4.50-8.00)	9.00 (5.60-15.0)	1743
	09-10	1.61 (1.45-1.79)	1.70 (1.50-1.90)	2.90 (2.50-3.30)	4.50 (4.00-5.20)	6.00 (5.40-7.10)	1869
Gender Males	99-00	2.61 (2.27-3.01)	2.60 (2.20-3.00)	4.10 (3.50-4.70)	6.60 (5.30-9.40)	10.5 (6.90-15.0)	743
	03-04	2.17 (1.87-2.51)	2.10 (1.80-2.40)	3.40 (2.80-4.50)	6.10 (4.60-8.10)	8.50 (6.50-10.5)	1053
	05-06	2.08 (1.76-2.45)	2.20 (1.80-2.70)	3.70 (3.20-4.60)	6.40 (4.80-9.30)	9.50 (6.70-13.7)	1048
	07-08	2.63 (2.36-2.94)	2.60 (2.40-2.80)	4.20 (3.80-4.90)	7.30 (5.50-11.0)	11.4 (7.10-18.1)	1059
	09-10	2.21 (1.99-2.46)	2.20 (1.90-2.50)	3.40 (3.00-4.00)	5.50 (5.00-6.40)	7.80 (6.50-9.90)	1075
Females	99-00	1.79 (1.56-2.06)	1.70 (1.50-2.10)	2.80 (2.30-3.50)	5.10 (4.10-6.70)	8.00 (5.70-9.00)	819
	03-04	1.72 (1.56-1.91)	1.60 (1.40-1.80)	2.90 (2.50-3.50)	5.80 (4.60-7.10)	8.20 (6.70-10.0)	1041
	05-06	1.37 (1.15-1.63)	1.50 (1.20-1.70)	2.80 (2.20-3.30)	4.60 (3.90-5.80)	6.60 (5.60-9.30)	1072
	07-08	1.46 (1.30-1.64)	1.40 (1.30-1.50)	2.60 (2.20-3.10)	4.70 (3.80-6.10)	7.50 (5.00-12.5)	1041
	09-10	1.26 (1.14-1.40)	1.30 (1.20-1.40)	2.20 (2.00-2.60)	4.00 (3.50-4.50)	5.40 (4.70-6.50)	1158
Race/ethnicity Mexican Americans	99-00	1.47 (1.13-1.92)	1.50 (1.00-2.00)	2.80 (1.80-3.80)	4.50 (3.30-6.90)	6.70 (4.60-9.50)	584
	03-04	1.42 (1.17-1.72)	1.50 (1.20-1.70)	2.30 (1.90-2.90)	4.30 (3.10-5.10)	5.50 (4.00-8.90)	485
	05-06	1.25 (.884-1.76)	1.60 (1.20-2.10)	2.80 (2.50-3.40)	4.30 (3.60-5.20)	6.00 (4.60-7.20)	499
	07-08	1.65 (1.46-1.86)	1.70 (1.40-1.90)	3.00 (2.70-3.40)	5.00 (3.70-6.10)	6.70 (5.20-10.7)	391
	09-10	1.01 (.835-1.22)	1.10 (.800-1.30)	1.80 (1.50-2.20)	3.10 (2.50-3.70)	4.00 (3.20-5.30)	461
Non-Hispanic blacks	99-00	2.14 (1.58-2.89)	2.10 (1.40-3.00)	3.40 (2.70-5.00)	6.70 (3.90-10.7)	10.7 (6.80-14.3)	303
	03-04	1.92 (1.62-2.26)	1.90 (1.60-2.20)	3.50 (2.80-4.60)	6.00 (5.00-7.10)	8.30 (6.30-12.0)	538
	05-06	1.53 (1.13-2.07)	1.70 (1.40-2.20)	3.30 (2.50-4.10)	5.40 (3.90-7.50)	7.50 (5.30-11.0)	544
	07-08	1.98 (1.55-2.52)	2.10 (1.80-2.50)	3.90 (2.80-5.40)	7.90 (4.20-12.5)	11.1 (5.90-17.2)	419
	09-10	1.44 (1.26-1.64)	1.40 (1.20-1.70)	2.60 (2.10-3.00)	4.40 (3.80-5.20)	6.50 (4.60-8.20)	391
Non-Hispanic whites	99-00	2.30 (2.07-2.56)	2.30 (1.90-2.60)	3.60 (3.20-4.20)	5.90 (5.30-7.00)	8.40 (7.00-9.70)	519
	03-04	2.01 (1.77-2.27)	1.90 (1.70-2.20)	3.30 (2.80-4.10)	6.10 (4.70-7.80)	8.20 (6.90-10.1)	962
	05-06	1.79 (1.48-2.17)	1.80 (1.50-2.20)	3.30 (2.70-4.30)	5.60 (4.60-8.20)	9.40 (6.10-13.1)	935
	07-08	2.13 (1.89-2.41)	2.10 (1.80-2.40)	3.70 (3.20-4.20)	6.40 (5.30-9.00)	10.5 (6.50-15.6)	931
	09-10	1.93 (1.77-2.10)	2.00 (1.80-2.20)	3.30 (2.90-3.70)	5.30 (4.80-5.60)	7.30 (6.20-8.80)	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.1, 0.3, 0.1, 0.1, and 0.1 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum Perfluorononanoic acid (PFNA)

Geometric mean and selected percentiles of serum concentrations (in $\mu\text{g/L}$) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.557 (.463-.671)	.600 (.500-.700)	.900 (.700-1.00)	1.30 (1.00-1.90)	1.80 (1.30-2.40)	1562
	03-04	.966 (.816-1.14)	1.00 (.900-1.10)	1.50 (1.20-1.80)	2.30 (1.60-4.30)	3.20 (1.80-7.70)	2094
	05-06	1.09 (.909-1.29)	1.10 (.900-1.20)	1.70 (1.30-2.10)	2.60 (1.80-4.20)	3.60 (2.50-5.70)	2120
	07-08	1.22 (1.12-1.32)	1.23 (1.15-1.31)	1.72 (1.56-1.89)	2.62 (2.30-2.87)	3.28 (2.95-3.77)	2100
	09-10	1.26 (1.11-1.44)	1.23 (1.07-1.31)	1.80 (1.48-2.13)	2.71 (1.97-4.26)	3.77 (2.30-7.22)	2233
Age group 12-19 years	99-00	.476 (.410-.553)	.500 (.400-.600)	.700 (.600-.800)	.900 (.800-1.10)	1.20 (.900-1.40)	543
	03-04	.852 (.697-1.04)	.800 (.700-1.00)	1.20 (1.00-1.60)	1.90 (1.20-3.70)	2.80 (1.30-6.30)	640
	05-06	.929 (.782-1.10)	.900 (.800-1.10)	1.40 (1.10-1.80)	2.00 (1.60-2.70)	2.70 (2.10-3.40)	640
	07-08	1.16 (1.04-1.30)	1.15 (.980-1.31)	1.56 (1.31-1.97)	2.05 (1.64-2.71)	2.54 (2.13-3.28)	357
	09-10	1.10 (.950-1.27)	1.07 (.900-1.23)	1.48 (1.31-1.80)	2.21 (1.48-2.95)	2.62 (1.89-4.84)	364
20 years and older	99-00	.573 (.472-.695)	.600 (.500-.700)	.900 (.700-1.20)	1.40 (1.00-1.90)	1.90 (1.40-2.40)	1019
	03-04	.984 (.835-1.16)	1.00 (.900-1.10)	1.50 (1.20-1.80)	2.40 (1.60-4.40)	3.40 (1.80-8.40)	1454
	05-06	1.11 (.929-1.33)	1.10 (1.00-1.20)	1.70 (1.30-2.20)	2.60 (1.80-4.40)	3.90 (2.50-5.80)	1480
	07-08	1.23 (1.13-1.33)	1.23 (1.15-1.31)	1.72 (1.64-1.89)	2.62 (2.38-2.95)	3.36 (3.03-3.77)	1743
	09-10	1.29 (1.13-1.47)	1.23 (1.15-1.39)	1.80 (1.56-2.21)	2.71 (1.97-4.67)	3.94 (2.38-8.36)	1869
Gender Males	99-00	.608 (.515-.717)	.600 (.500-.700)	.900 (.800-1.10)	1.30 (1.10-1.80)	1.70 (1.30-2.40)	743
	03-04	1.09 (.912-1.30)	1.10 (.900-1.20)	1.60 (1.40-1.90)	2.40 (1.70-5.00)	4.00 (1.80-8.70)	1053
	05-06	1.19 (.993-1.43)	1.20 (1.00-1.30)	1.80 (1.40-2.50)	2.90 (1.80-4.60)	3.90 (2.50-6.70)	1048
	07-08	1.36 (1.27-1.47)	1.31 (1.23-1.39)	1.89 (1.72-2.05)	2.87 (2.54-3.12)	3.53 (3.12-4.18)	1059
	09-10	1.37 (1.17-1.59)	1.31 (1.15-1.48)	1.80 (1.56-2.30)	2.79 (1.89-4.84)	4.10 (2.21-12.1)	1075
Females	99-00	.517 (.419-.638)	.500 (.400-.600)	.800 (.600-1.00)	1.30 (.900-2.00)	1.80 (1.30-2.60)	819
	03-04	.861 (.721-1.03)	.900 (.800-1.00)	1.30 (1.00-1.70)	2.20 (1.40-3.40)	3.00 (1.70-6.10)	1041
	05-06	.992 (.833-1.18)	.900 (.800-1.10)	1.50 (1.20-1.90)	2.40 (1.70-3.50)	3.10 (2.10-5.70)	1072
	07-08	1.09 (.985-1.21)	1.07 (.980-1.23)	1.56 (1.48-1.72)	2.21 (1.97-2.54)	2.79 (2.54-3.53)	1041
	09-10	1.17 (1.05-1.31)	1.15 (.980-1.23)	1.72 (1.39-2.05)	2.62 (1.97-3.77)	3.53 (2.30-6.15)	1158
Race/ethnicity Mexican Americans	99-00	.349 (.303-.403)	.400 (.300-.400)	.500 (.400-.700)	.700 (.600-.900)	.900 (.700-1.30)	584
	03-04	.689 (.586-.809)	.700 (.600-.900)	1.10 (.900-1.40)	1.60 (1.30-2.00)	2.00 (1.60-2.80)	485
	05-06	.821 (.710-.951)	.800 (.700-.900)	1.20 (1.10-1.40)	1.60 (1.30-2.30)	2.30 (1.50-3.20)	499
	07-08	1.10 (.967-1.25)	1.15 (.980-1.31)	1.56 (1.39-1.80)	2.13 (1.80-2.71)	2.71 (2.13-3.28)	391
	09-10	.993 (.906-1.09)	.980 (.900-1.07)	1.39 (1.31-1.56)	2.05 (1.80-2.38)	2.46 (2.21-3.03)	461
Non-Hispanic blacks	99-00	.765 (.607-.965)	.700 (.600-1.00)	1.30 (.800-1.80)	1.90 (1.50-2.50)	2.50 (1.90-2.90)	303
	03-04	1.14 (.834-1.54)	1.10 (.900-1.40)	1.70 (1.20-2.90)	3.20 (1.50-6.50)	4.70 (2.10-9.30)	538
	05-06	1.17 (.916-1.50)	1.20 (.900-1.40)	1.60 (1.30-2.50)	2.60 (1.70-5.60)	4.40 (2.40-8.20)	544
	07-08	1.35 (1.23-1.48)	1.39 (1.23-1.48)	1.89 (1.72-2.13)	2.79 (2.30-3.36)	3.36 (2.79-4.67)	419
	09-10	1.42 (1.29-1.56)	1.39 (1.23-1.56)	1.97 (1.80-2.30)	2.79 (2.38-3.36)	3.36 (2.62-4.84)	391
Non-Hispanic whites	99-00	.571 (.471-.691)	.600 (.500-.700)	.900 (.700-1.10)	1.30 (1.00-1.90)	1.70 (1.30-2.30)	519
	03-04	.963 (.826-1.12)	.900 (.900-1.10)	1.50 (1.20-1.70)	2.30 (1.60-3.60)	3.00 (1.80-6.20)	962
	05-06	1.10 (.914-1.31)	1.10 (.900-1.20)	1.70 (1.30-2.10)	2.60 (1.80-4.20)	3.50 (2.40-5.70)	935
	07-08	1.22 (1.10-1.34)	1.23 (1.15-1.31)	1.72 (1.48-1.89)	2.54 (2.13-3.03)	3.20 (2.79-3.85)	931
	09-10	1.28 (1.08-1.51)	1.23 (1.07-1.39)	1.80 (1.48-2.21)	2.62 (1.80-5.08)	3.85 (2.05-10.8)	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.1, 0.1, 0.1, 0.082, and 0.082 respectively.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.82 for survey periods 2007-2008 and 2009-2010 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum Perfluorooctanoic acid (PFOA)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles (95% confidence interval)				Sample size
		(95% conf. interval)	50th	75th	90th	95th	
Total	99-00	5.21 (4.72-5.74)	5.20 (4.80-5.90)	6.90 (6.30-7.80)	9.40 (8.20-11.1)	11.9 (10.9-13.5)	1562
	03-04	3.95 (3.65-4.27)	4.10 (3.80-4.40)	5.80 (5.30-6.40)	7.80 (6.70-9.60)	9.80 (7.40-14.1)	2094
	05-06	3.92 (3.48-4.42)	4.20 (3.80-4.50)	6.20 (5.40-7.20)	9.00 (7.40-11.2)	11.3 (8.80-14.5)	2120
	07-08	4.12 (4.01-4.24)	4.30 (4.20-4.50)	5.90 (5.70-6.20)	7.90 (7.50-8.30)	9.60 (8.90-10.1)	2100
	09-10	3.07 (2.81-3.36)	3.20 (2.90-3.50)	4.60 (4.10-5.10)	6.00 (5.30-7.20)	7.50 (6.20-9.70)	2233
Age group 12-19 years	99-00	5.46 (4.98-5.99)	5.60 (4.90-6.10)	6.90 (6.20-7.70)	9.40 (7.70-11.0)	11.6 (10.2-12.5)	543
	03-04	3.89 (3.47-4.35)	4.00 (3.50-4.50)	5.40 (4.60-6.10)	7.00 (5.60-9.20)	8.60 (5.90-12.6)	640
	05-06	3.59 (3.26-3.96)	3.80 (3.30-4.20)	5.40 (4.60-5.90)	6.90 (6.30-7.90)	8.40 (7.30-10.1)	640
	07-08	3.91 (3.71-4.12)	4.00 (3.70-4.40)	5.00 (4.70-5.50)	6.10 (5.70-6.70)	7.30 (6.20-8.00)	357
	09-10	2.74 (2.46-3.05)	2.90 (2.60-3.10)	3.80 (3.30-4.20)	4.80 (4.10-5.30)	5.00 (4.40-7.20)	364
20 years and older	99-00	5.16 (4.66-5.72)	5.20 (4.70-5.70)	7.00 (6.30-7.80)	9.40 (8.20-11.1)	12.0 (10.8-14.0)	1019
	03-04	3.96 (3.67-4.27)	4.10 (3.90-4.40)	5.90 (5.40-6.50)	7.80 (6.80-9.60)	9.90 (7.60-14.2)	1454
	05-06	3.97 (3.51-4.49)	4.20 (3.80-4.60)	6.40 (5.40-7.50)	9.30 (7.40-11.9)	11.6 (9.00-14.8)	1480
	07-08	4.15 (4.02-4.30)	4.30 (4.20-4.60)	6.10 (5.80-6.50)	8.10 (7.70-8.70)	9.80 (9.00-10.4)	1743
	09-10	3.12 (2.84-3.43)	3.30 (3.00-3.60)	4.70 (4.20-5.30)	6.40 (5.40-7.50)	7.70 (6.30-10.2)	1869
Gender Males	99-00	5.71 (5.17-6.31)	6.00 (5.40-6.50)	7.70 (6.80-8.70)	10.6 (9.00-11.8)	12.1 (11.2-13.1)	743
	03-04	4.47 (4.07-4.91)	4.60 (4.30-5.00)	6.30 (5.70-7.20)	8.40 (6.80-12.5)	10.7 (7.40-17.5)	1053
	05-06	4.69 (4.23-5.20)	4.90 (4.40-5.40)	7.20 (6.10-8.00)	9.90 (8.00-12.2)	12.2 (9.60-15.2)	1048
	07-08	4.80 (4.55-5.06)	4.90 (4.60-5.20)	6.70 (6.30-7.10)	8.70 (8.10-9.30)	10.1 (9.50-11.1)	1059
	09-10	3.53 (3.22-3.87)	3.70 (3.40-4.00)	4.90 (4.50-5.40)	6.80 (5.40-8.00)	7.90 (6.40-10.2)	1075
Females	99-00	4.80 (4.32-5.34)	4.70 (4.40-5.20)	6.30 (5.60-7.10)	8.30 (7.50-9.90)	11.3 (9.20-14.4)	819
	03-04	3.50 (3.21-3.82)	3.60 (3.30-3.90)	5.20 (4.70-5.80)	7.10 (6.30-8.20)	8.60 (7.40-10.6)	1041
	05-06	3.31 (2.89-3.79)	3.50 (3.10-4.00)	5.20 (4.40-6.00)	7.90 (6.10-9.70)	10.1 (7.50-14.2)	1072
	07-08	3.55 (3.38-3.73)	3.70 (3.50-3.90)	5.20 (4.80-5.60)	7.00 (6.50-7.50)	8.30 (7.20-9.90)	1041
	09-10	2.69 (2.45-2.96)	2.70 (2.50-3.00)	4.10 (3.60-4.70)	5.60 (5.10-6.50)	6.90 (5.80-8.40)	1158
Race/ethnicity Mexican Americans	99-00	3.89 (3.58-4.22)	4.20 (3.80-4.60)	5.80 (5.20-6.30)	7.70 (6.40-8.00)	8.20 (7.70-8.90)	584
	03-04	3.11 (2.84-3.40)	3.30 (3.10-3.70)	4.50 (4.20-5.20)	6.70 (5.70-7.30)	7.60 (6.70-10.5)	485
	05-06	2.62 (2.33-2.95)	2.80 (2.50-3.30)	4.30 (3.80-4.70)	5.80 (5.30-6.70)	7.40 (5.90-8.10)	499
	07-08	3.54 (3.35-3.75)	3.80 (3.50-4.00)	5.20 (4.90-5.60)	6.60 (6.20-7.10)	7.60 (6.80-9.00)	391
	09-10	2.26 (2.00-2.54)	2.40 (2.10-2.60)	3.60 (3.10-3.80)	4.60 (4.00-5.30)	5.40 (4.50-6.60)	461
Non-Hispanic blacks	99-00	4.78 (4.10-5.57)	4.80 (3.80-5.90)	6.40 (5.90-7.50)	9.00 (7.40-11.5)	11.5 (9.20-14.0)	303
	03-04	3.37 (2.99-3.79)	3.70 (3.20-4.20)	5.20 (4.40-6.30)	7.70 (5.30-11.6)	9.60 (6.50-13.9)	538
	05-06	3.27 (2.61-4.08)	3.70 (3.00-4.20)	5.50 (4.40-6.80)	8.10 (6.00-11.3)	10.4 (7.80-12.3)	544
	07-08	3.86 (3.57-4.16)	4.00 (3.50-4.30)	5.90 (5.20-6.50)	7.80 (7.10-8.70)	9.20 (8.50-10.1)	419
	09-10	2.74 (2.47-3.04)	2.80 (2.60-3.00)	4.00 (3.70-4.40)	5.50 (5.00-6.20)	6.70 (5.60-9.40)	391
Non-Hispanic whites	99-00	5.61 (5.06-6.23)	5.60 (4.90-6.20)	7.30 (6.50-8.20)	10.3 (8.40-12.1)	13.0 (11.0-14.9)	519
	03-04	4.18 (3.85-4.53)	4.30 (3.90-4.70)	6.00 (5.50-6.70)	7.90 (7.20-9.20)	9.90 (7.60-13.3)	962
	05-06	4.27 (3.80-4.81)	4.40 (4.00-5.00)	6.60 (5.60-7.80)	9.60 (7.40-12.2)	11.6 (8.80-14.8)	935
	07-08	4.38 (4.20-4.56)	4.60 (4.30-4.70)	6.10 (5.80-6.60)	8.20 (7.80-8.80)	9.90 (9.30-10.6)	931
	09-10	3.36 (3.06-3.69)	3.50 (3.20-3.90)	4.80 (4.40-5.40)	6.60 (5.40-7.80)	7.80 (6.20-10.5)	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.1, 0.1, 0.1, 0.1, and 0.1 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum Perfluorooctane sulfonic acid (PFOS)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	30.4 (27.1-33.9)	30.2 (27.8-33.9)	43.7 (37.5-47.3)	57.0 (50.2-71.7)	75.7 (58.1-97.5)	1562
	03-04	20.7 (19.2-22.3)	21.2 (19.8-22.4)	30.0 (27.5-33.0)	41.3 (35.6-50.0)	54.6 (44.0-66.5)	2094
	05-06	17.1 (16.0-18.2)	17.5 (16.8-18.6)	27.2 (24.9-29.6)	39.4 (34.9-43.1)	47.5 (42.7-56.8)	2120
	07-08	13.2 (12.2-14.2)	13.6 (12.8-14.7)	21.0 (18.9-23.3)	32.6 (29.4-36.3)	40.5 (35.4-47.4)	2100
	09-10	9.32 (8.13-10.7)	9.70 (8.50-10.8)	14.8 (12.9-17.3)	23.7 (18.3-30.2)	32.0 (22.6-48.5)	2233
Age group 12-19 years	99-00	29.1 (26.2-32.4)	29.5 (26.9-34.2)	39.0 (35.9-45.0)	52.7 (46.0-56.2)	57.4 (52.7-66.5)	543
	03-04	19.3 (17.5-21.4)	19.9 (17.8-22.0)	27.1 (23.7-30.2)	36.5 (28.6-45.6)	42.6 (35.1-52.1)	640
	05-06	15.0 (14.3-15.7)	14.9 (13.6-16.6)	22.7 (19.7-24.9)	30.6 (27.8-34.1)	38.5 (33.0-44.6)	640
	07-08	11.3 (10.3-12.3)	11.3 (10.3-13.0)	15.9 (15.1-17.7)	21.7 (17.7-28.2)	28.0 (22.0-32.2)	357
	09-10	6.84 (5.81-8.06)	6.90 (6.00-8.40)	10.7 (8.90-12.5)	14.4 (12.4-18.1)	18.1 (13.5-26.0)	364
20 years and older	99-00	30.6 (27.1-34.4)	30.3 (27.9-33.9)	44.4 (37.9-48.0)	58.0 (50.1-75.7)	78.0 (59.9-107)	1019
	03-04	20.9 (19.3-22.5)	21.4 (19.8-22.8)	30.4 (28.1-33.0)	42.7 (35.7-53.3)	57.8 (45.7-69.4)	1454
	05-06	17.4 (16.2-18.7)	18.0 (17.1-19.4)	27.8 (25.3-30.8)	40.2 (35.6-44.5)	49.6 (42.8-60.7)	1480
	07-08	13.5 (12.4-14.6)	14.0 (13.0-15.5)	21.7 (19.5-24.6)	33.9 (29.9-39.0)	42.8 (37.3-50.3)	1743
	09-10	9.72 (8.45-11.2)	10.1 (8.90-11.2)	15.7 (13.4-18.4)	25.3 (19.3-32.9)	34.1 (23.4-52.7)	1869
Gender Males	99-00	33.4 (29.6-37.6)	34.9 (31.1-37.9)	46.3 (41.2-51.6)	58.4 (50.2-78.3)	78.3 (58.0-108)	743
	03-04	23.2 (21.1-25.6)	23.9 (22.4-25.5)	32.2 (28.8-35.9)	45.3 (35.5-62.7)	62.7 (43.8-81.8)	1053
Females	05-06	20.5 (19.4-21.8)	21.3 (20.0-22.5)	31.4 (27.9-33.7)	43.3 (38.7-49.7)	54.3 (43.5-80.7)	1048
	07-08	16.3 (15.0-17.7)	17.0 (15.7-17.8)	23.9 (21.6-26.9)	36.4 (33.5-41.1)	45.3 (40.4-53.1)	1059
	09-10	11.5 (9.93-13.3)	11.8 (10.6-12.9)	16.8 (14.3-20.4)	25.7 (19.4-39.7)	37.4 (22.5-72.3)	1075
	99-00	28.0 (24.6-31.8)	27.8 (24.5-30.2)	39.0 (32.7-46.0)	55.4 (46.3-70.2)	75.7 (56.1-98.4)	819
	03-04	18.4 (17.0-20.0)	18.2 (16.9-19.8)	27.4 (23.8-30.2)	39.8 (34.4-42.6)	46.6 (42.3-61.5)	1041
Race/ethnicity Mexican Americans	05-06	14.4 (13.3-15.4)	14.6 (13.5-15.9)	23.3 (21.1-25.3)	34.2 (30.6-37.7)	42.8 (38.0-46.5)	1072
	07-08	10.7 (9.70-11.7)	10.7 (9.80-11.7)	17.2 (15.4-19.1)	28.7 (21.7-32.5)	33.6 (29.9-41.6)	1041
	09-10	7.65 (6.73-8.71)	7.80 (6.70-9.00)	12.0 (10.8-14.4)	21.1 (16.4-26.9)	28.8 (22.3-34.1)	1158
	99-00	22.7 (19.8-25.9)	23.7 (20.8-27.2)	33.0 (27.9-39.7)	41.9 (36.7-54.5)	53.6 (41.3-72.0)	584
	03-04	14.7 (13.0-16.6)	15.9 (13.4-17.9)	21.2 (18.7-23.5)	28.1 (24.1-35.0)	35.5 (28.9-38.5)	485
Non-Hispanic blacks	05-06	11.2 (10.3-12.2)	11.6 (9.90-13.3)	17.2 (15.4-19.3)	24.9 (22.0-30.3)	31.9 (26.4-40.6)	499
	07-08	10.6 (9.47-11.9)	10.8 (9.70-12.2)	17.1 (14.9-19.2)	26.3 (20.7-30.4)	31.5 (25.7-36.1)	391
	09-10	6.23 (5.28-7.36)	6.10 (5.10-8.10)	11.2 (8.50-14.1)	16.8 (14.9-18.8)	20.4 (17.5-22.7)	461
	99-00	32.9 (26.0-41.6)	32.0 (23.9-45.8)	50.6 (37.4-62.2)	69.3 (60.9-75.9)	81.5 (68.0-129)	303
	03-04	21.6 (19.1-24.4)	22.1 (19.6-24.9)	32.3 (28.1-36.2)	43.8 (37.2-57.3)	57.7 (43.8-78.4)	538
Non-Hispanic whites	05-06	18.4 (15.6-21.8)	19.0 (16.7-22.3)	27.6 (24.3-35.5)	45.9 (34.4-58.1)	57.9 (45.0-84.4)	544
	07-08	15.0 (12.6-17.8)	15.2 (12.9-17.7)	25.8 (21.0-33.3)	42.7 (31.5-57.3)	57.3 (43.4-79.2)	419
	09-10	9.11 (7.93-10.5)	9.50 (8.20-11.2)	16.0 (13.4-17.4)	23.0 (20.5-26.1)	28.7 (25.7-39.7)	391
	99-00	32.3 (29.5-35.3)	32.8 (29.5-36.0)	45.1 (40.4-47.4)	56.3 (51.7-70.0)	78.0 (58.0-98.4)	519
	03-04	21.4 (19.9-23.1)	22.0 (20.5-23.0)	30.2 (27.7-33.3)	41.7 (35.7-49.6)	56.3 (44.0-70.0)	962
05-06	18.1 (17.1-19.1)	18.6 (17.3-19.8)	28.3 (26.1-30.8)	39.7 (35.6-43.3)	46.6 (42.7-54.3)	935	
07-08	13.7 (12.7-14.8)	14.3 (13.2-15.5)	21.1 (18.7-23.7)	32.9 (29.7-36.0)	40.4 (36.1-44.9)	931	
09-10	10.2 (8.69-11.9)	10.4 (9.20-11.5)	15.6 (13.0-18.7)	24.7 (17.5-35.1)	33.1 (20.7-56.7)	1031	

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.2, 0.4, 0.2, 0.2, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum Perfluorooctane sulfonamide (PFOSA)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.361 (.303-.431)	.400 (.300-.400)	.700 (.500-.800)	1.00 (.900-1.40)	1.40 (1.00-2.00)	1562
	03-04	*	< LOD	< LOD	.300 (.200-.300)	.300 (.300-.400)	2094
	05-06	*	< LOD	.100 (<LOD-.200)	.200 (.200-.300)	.300 (.200-.500)	2120
	07-08	*	< LOD	< LOD	< LOD	< LOD	2100
	09-10	*	< LOD	< LOD	< LOD	< LOD	2233
Age group 12-19 years	99-00	.432 (.372-.501)	.400 (.400-.500)	.700 (.700-.900)	1.10 (.900-1.40)	1.60 (1.10-2.90)	543
	03-04	*	< LOD	< LOD	.300 (.200-.300)	.300 (.200-.500)	640
	05-06	*	< LOD	< LOD	.200 (.100-.300)	.300 (.200-.500)	640
	07-08	*	< LOD	< LOD	< LOD	< LOD	357
	09-10	*	< LOD	< LOD	< LOD	< LOD	364
20 years and older	99-00	.351 (.291-.422)	.300 (.300-.400)	.600 (.500-.800)	1.00 (.800-1.40)	1.40 (1.00-2.00)	1019
	03-04	*	< LOD	< LOD	.300 (.200-.300)	.300 (.300-.400)	1454
	05-06	*	< LOD	.100 (<LOD-.200)	.200 (.200-.300)	.300 (.200-.500)	1480
	07-08	*	< LOD	< LOD	< LOD	< LOD	1743
	09-10	*	< LOD	< LOD	< LOD	< LOD	1869
Gender Males	99-00	.359 (.301-.428)	.400 (.300-.400)	.600 (.500-.700)	.900 (.700-1.30)	1.30 (.900-1.90)	743
	03-04	*	< LOD	< LOD	.300 (.200-.300)	.300 (.200-.500)	1053
	05-06	*	< LOD	.100 (.100-.200)	.200 (.200-.300)	.300 (.200-.600)	1048
	07-08	*	< LOD	< LOD	< LOD	< LOD	1059
	09-10	*	< LOD	< LOD	< LOD	< LOD	1075
Females	99-00	.364 (.302-.438)	.300 (.300-.500)	.700 (.500-.900)	1.10 (.900-1.40)	1.50 (1.20-2.00)	819
	03-04	*	< LOD	< LOD	.300 (.200-.300)	.300 (.300-.500)	1041
	05-06	*	< LOD	.100 (<LOD-.100)	.200 (.200-.300)	.300 (.200-.400)	1072
	07-08	*	< LOD	< LOD	< LOD	< LOD	1041
	09-10	*	< LOD	< LOD	< LOD	< LOD	1158
Race/ethnicity Mexican Americans	99-00	.294 (.255-.340)	.300 (.200-.400)	.500 (.400-.600)	.800 (.700-1.00)	1.10 (.800-1.60)	584
	03-04	*	< LOD	< LOD	< LOD	.200 (<LOD-.300)	485
	05-06	*	< LOD	< LOD	.200 (<LOD-.300)	.200 (<LOD-.500)	499
	07-08	*	< LOD	< LOD	< LOD	< LOD	391
	09-10	*	< LOD	< LOD	< LOD	< LOD	461
Non-Hispanic blacks	99-00	.352 (.292-.425)	.300 (.300-.400)	.600 (.500-.800)	1.00 (.700-1.70)	1.70 (.900-2.60)	303
	03-04	*	< LOD	< LOD	.300 (.200-.300)	.300 (.200-.500)	538
	05-06	*	< LOD	< LOD	.200 (.100-.200)	.200 (.200-.400)	544
	07-08	*	< LOD	< LOD	< LOD	< LOD	419
	09-10	*	< LOD	< LOD	< LOD	< LOD	391
Non-Hispanic whites	99-00	.390 (.315-.483)	.400 (.300-.500)	.700 (.600-.900)	1.10 (.900-1.40)	1.50 (1.10-2.00)	519
	03-04	*	< LOD	< LOD	.300 (.200-.300)	.300 (.300-.500)	962
	05-06	*	< LOD	.100 (.100-.200)	.200 (.200-.300)	.300 (.200-.500)	935
	07-08	*	< LOD	< LOD	< LOD	< LOD	931
	09-10	*	< LOD	< LOD	< LOD	< LOD	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.05, 0.2, 0.1, 0.1, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum 2-(N-Ethyl-perfluorooctane sulfonamido) acetic acid (Et-PFOA-AcOH)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.654 (.611-.701)	.600 (.500-.700)	1.10 (1.00-1.20)	1.60 (1.50-1.80)	2.30 (1.90-2.70)	1562
	03-04	*	< LOD	< LOD	< LOD	< LOD	2094
	05-06	*	< LOD	< LOD	< LOD	.300 (<LOD-.300)	2120
	07-08	*	< LOD	< LOD	< LOD	< LOD	2100
	09-10	*	< LOD	< LOD	< LOD	.100 (<LOD-.100)	2233
Age group 12-19 years	99-00	.827 (.765-.894)	.800 (.700-.800)	1.20 (1.10-1.30)	1.70 (1.50-2.00)	2.40 (1.90-3.30)	543
	03-04	*	< LOD	< LOD	< LOD	< LOD	640
	05-06	*	< LOD	< LOD	< LOD	.200 (<LOD-.400)	640
	07-08	*	< LOD	< LOD	< LOD	< LOD	357
	09-10	*	< LOD	< LOD	< LOD	< LOD	364
20 years and older	99-00	.628 (.586-.674)	.600 (.500-.700)	1.00 (.900-1.10)	1.60 (1.50-1.80)	2.30 (1.80-2.70)	1019
	03-04	*	< LOD	< LOD	< LOD	< LOD	1454
	05-06	*	< LOD	< LOD	< LOD	.300 (<LOD-.300)	1480
	07-08	*	< LOD	< LOD	< LOD	< LOD	1743
	09-10	*	< LOD	< LOD	< LOD	.100 (<LOD-.100)	1869
Gender Males	99-00	.632 (.586-.681)	.600 (.500-.700)	1.00 (.900-1.10)	1.50 (1.30-1.80)	1.90 (1.60-2.90)	743
	03-04	*	< LOD	< LOD	< LOD	< LOD	1053
	05-06	*	< LOD	< LOD	< LOD	.300 (.200-.300)	1048
	07-08	*	< LOD	< LOD	< LOD	< LOD	1059
	09-10	*	< LOD	< LOD	< LOD	.100 (<LOD-.100)	1075
Females	99-00	.675 (.616-.740)	.700 (.500-.700)	1.10 (1.00-1.20)	1.70 (1.50-2.20)	2.50 (2.00-2.80)	819
	03-04	*	< LOD	< LOD	< LOD	< LOD	1041
	05-06	*	< LOD	< LOD	< LOD	.300 (<LOD-.300)	1072
	07-08	*	< LOD	< LOD	< LOD	< LOD	1041
	09-10	*	< LOD	< LOD	< LOD	.100 (<LOD-.100)	1158
Race/ethnicity Mexican Americans	99-00	.591 (.498-.702)	.600 (.400-.700)	1.00 (.800-1.20)	1.70 (1.30-2.00)	2.50 (1.90-3.00)	584
	03-04	*	< LOD	< LOD	< LOD	< LOD	485
	05-06	*	< LOD	< LOD	< LOD	< LOD	499
	07-08	*	< LOD	< LOD	< LOD	< LOD	391
	09-10	*	< LOD	< LOD	< LOD	< LOD	461
Non-Hispanic blacks	99-00	.570 (.495-.655)	.500 (.500-.600)	.900 (.700-1.00)	1.50 (1.20-1.90)	2.20 (1.40-3.20)	303
	03-04	*	< LOD	< LOD	< LOD	.400 (<LOD-.500)	538
	05-06	*	< LOD	< LOD	< LOD	< LOD	544
	07-08	*	< LOD	< LOD	< LOD	< LOD	419
	09-10	*	< LOD	< LOD	< LOD	< LOD	391
Non-Hispanic whites	99-00	.673 (.625-.725)	.700 (.600-.700)	1.10 (1.00-1.20)	1.70 (1.50-1.90)	2.40 (1.90-2.80)	519
	03-04	*	< LOD	< LOD	< LOD	< LOD	962
	05-06	*	< LOD	< LOD	< LOD	.300 (.200-.400)	935
	07-08	*	< LOD	< LOD	< LOD	< LOD	931
	09-10	*	< LOD	< LOD	< LOD	.100 (<LOD-.200)	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.2, 0.4, 0.2, 0.2, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum 2-(N-Methyl-perfluorooctane sulfonamido) acetic acid (Me-PFOSA-AcOH)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.852 (.746-.974)	.870 (.780-.960)	1.40 (1.13-1.66)	2.18 (1.83-2.62)	2.79 (2.44-3.23)	1562
	03-04	*	< LOD	.610 (<LOD-.610)	.870 (.780-.960)	1.13 (.960-1.31)	2094
	05-06	.410 (.372-.451)	.440 (.350-.440)	.700 (.610-.780)	1.22 (1.13-1.22)	1.57 (1.48-1.74)	2120
	07-08	.301 (.276-.329)	.260 (.260-.350)	.520 (.440-.610)	.960 (.780-1.05)	1.31 (1.22-1.40)	2100
	09-10	.189 (.176-.204)	.170 (.170-.170)	.350 (.350-.440)	.700 (.610-.700)	.960 (.870-1.05)	2233
Age group							
12-19 years	99-00	1.17 (1.03-1.34)	1.22 (1.05-1.31)	1.83 (1.31-2.18)	2.70 (2.18-3.05)	3.23 (2.79-4.10)	543
	03-04	*	< LOD	.610 (<LOD-.700)	.960 (.780-1.13)	1.31 (1.05-1.57)	640
	05-06	.432 (.380-.490)	.440 (.350-.520)	.780 (.610-.870)	1.13 (.960-1.31)	1.48 (1.13-1.66)	640
	07-08	.338 (.302-.379)	.350 (.260-.440)	.610 (.520-.610)	.960 (.780-1.48)	1.57 (.960-2.01)	357
	09-10	.225 (.193-.263)	.260 (.170-.260)	.440 (.350-.520)	.700 (.520-.870)	.870 (.610-1.66)	364
20 years and older	99-00	.806 (.701-.927)	.780 (.700-.960)	1.31 (1.13-1.57)	2.09 (1.74-2.44)	2.70 (2.27-3.14)	1019
	03-04	*	< LOD	.610 (<LOD-.610)	.870 (.780-.960)	1.05 (.960-1.40)	1454
	05-06	.406 (.369-.448)	.440 (.350-.440)	.700 (.610-.780)	1.22 (1.13-1.31)	1.66 (1.48-1.74)	1480
	07-08	.296 (.269-.326)	.260 (.260-.350)	.520 (.440-.610)	.960 (.780-1.05)	1.22 (1.13-1.40)	1743
	09-10	.185 (.170-.200)	.170 (.170-.170)	.350 (.260-.350)	.700 (.610-.780)	.960 (.870-1.05)	1869
Gender							
Males	99-00	.891 (.764-1.04)	.870 (.780-1.05)	1.40 (1.13-1.74)	2.27 (2.01-2.79)	2.79 (2.44-3.66)	743
	03-04	*	< LOD	.610 (<LOD-.610)	.960 (.780-1.05)	1.13 (.960-1.48)	1053
	05-06	.435 (.400-.472)	.440 (.350-.440)	.780 (.700-.870)	1.22 (1.13-1.31)	1.66 (1.40-1.74)	1048
	07-08	.305 (.279-.333)	.260 (.260-.350)	.520 (.440-.610)	.870 (.780-.960)	1.31 (1.13-1.40)	1059
	09-10	.192 (.179-.205)	.170 (.170-.170)	.350 (.350-.440)	.700 (.610-.780)	.960 (.780-1.13)	1075
Females	99-00	.820 (.724-.928)	.780 (.700-.960)	1.31 (1.13-1.66)	2.01 (1.74-2.35)	2.62 (2.27-3.23)	819
	03-04	*	< LOD	< LOD	.870 (.780-.960)	.960 (.870-1.57)	1041
	05-06	.387 (.342-.438)	.350 (.350-.440)	.700 (.610-.780)	1.13 (1.05-1.31)	1.57 (1.40-1.83)	1072
	07-08	.298 (.266-.334)	.260 (.260-.350)	.520 (.440-.610)	.960 (.780-1.13)	1.31 (1.13-1.66)	1041
	09-10	.187 (.170-.206)	.170 (.170-.170)	.350 (.260-.350)	.610 (.520-.780)	.960 (.780-1.05)	1158
Race/ethnicity							
Mexican Americans	99-00	.675 (.552-.825)	.700 (.520-.780)	1.13 (.870-1.31)	1.83 (1.40-2.35)	2.35 (1.74-3.92)	584
	03-04	*	< LOD	< LOD	.610 (<LOD-.780)	.780 (<LOD-1.13)	485
	05-06	.292 (.245-.348)	.260 (.260-.350)	.440 (.350-.700)	.780 (.610-1.05)	1.05 (.870-1.22)	499
	07-08	*	< LOD	.350 (.260-.440)	.520 (.440-.780)	.780 (.520-1.13)	391
	09-10	.133 (.113-.157)	.090 (.090-.170)	.260 (.170-.350)	.520 (.440-.610)	.700 (.520-.960)	461
Non-Hispanic blacks	99-00	.991 (.857-1.15)	1.05 (.780-1.22)	1.57 (1.31-1.83)	2.35 (1.92-2.79)	3.14 (2.27-4.27)	303
	03-04	*	< LOD	.700 (<LOD-.780)	.960 (.780-1.22)	1.31 (.960-1.57)	538
	05-06	.418 (.361-.486)	.440 (.350-.520)	.700 (.610-.870)	1.22 (1.05-1.48)	1.66 (1.48-2.01)	544
	07-08	.329 (.302-.357)	.350 (.260-.350)	.520 (.520-.700)	1.05 (.780-1.22)	1.40 (1.22-1.66)	419
	09-10	.184 (.158-.215)	.170 (.170-.170)	.350 (.260-.440)	.610 (.440-.780)	.870 (.610-1.13)	391
Non-Hispanic whites	99-00	.889 (.766-1.03)	.870 (.780-1.05)	1.40 (1.22-1.74)	2.18 (1.92-2.70)	2.79 (2.44-3.66)	519
	03-04	*	< LOD	.610 (<LOD-.700)	.870 (.780-.960)	1.13 (.960-1.40)	962
	05-06	.428 (.380-.483)	.440 (.350-.520)	.780 (.700-.870)	1.22 (1.13-1.31)	1.66 (1.40-1.83)	935
	07-08	.332 (.296-.372)	.350 (.260-.350)	.610 (.520-.700)	.960 (.780-1.22)	1.31 (1.13-1.57)	931
	09-10	.206 (.191-.222)	.170 (.170-.260)	.350 (.350-.440)	.700 (.610-.780)	.960 (.870-1.13)	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.2, 0.5, 0.2, 0.2, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.872 for survey periods 1999-2000 and 2003-2010 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Serum Perfluoroundecanoic acid (PFUA)

Geometric mean and selected percentiles of serum concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.300 (.200-.400)	.400 (.300-.700)	1562
	03-04	*	< LOD	< LOD	< LOD	.600 (<LOD-1.30)	2094
	05-06	*	< LOD	.300 (.200-.400)	.500 (.400-.800)	.700 (.600-1.20)	2120
	07-08	*	< LOD	.200 (<LOD-.300)	.400 (.300-.500)	.600 (.500-.800)	2100
	09-10	.172 (.151-.196)	.200 (.100-.200)	.300 (.200-.400)	.600 (.400-.800)	.900 (.600-1.10)	2233
Age group 12-19 years	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.300)	543
	03-04	*	< LOD	< LOD	< LOD	.300 (<LOD-1.10)	640
	05-06	*	< LOD	< LOD	.300 (.200-.500)	.500 (.300-.900)	640
	07-08	*	< LOD	< LOD	.200 (<LOD-.200)	.300 (.200-.300)	357
	09-10	*	.100 (<LOD-.100)	.200 (.100-.200)	.300 (.200-.600)	.400 (.200-.900)	364
20 years and older	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.300 (.200-.500)	.500 (.300-.700)	1019
	03-04	*	< LOD	< LOD	.300 (<LOD-.600)	.600 (<LOD-1.40)	1454
	05-06	*	< LOD	.300 (.200-.400)	.500 (.400-.800)	.800 (.600-1.10)	1480
	07-08	*	< LOD	.200 (.200-.300)	.400 (.300-.600)	.600 (.500-.900)	1743
	09-10	.182 (.160-.207)	.200 (.200-.200)	.300 (.300-.400)	.600 (.400-.800)	.900 (.700-1.20)	1869
Gender Males	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.300 (.200-.500)	.500 (.300-.600)	743
	03-04	*	< LOD	< LOD	.400 (<LOD-1.00)	.700 (<LOD-2.30)	1053
	05-06	*	< LOD	.300 (.200-.400)	.600 (.400-.700)	.800 (.600-1.10)	1048
	07-08	*	< LOD	.200 (<LOD-.300)	.400 (.300-.500)	.700 (.500-1.00)	1059
	09-10	.170 (.147-.196)	.200 (.100-.200)	.300 (.200-.400)	.500 (.400-.900)	.800 (.500-1.20)	1075
Females	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.300 (.200-.400)	.400 (.200-.800)	819
	03-04	*	< LOD	< LOD	< LOD	.400 (<LOD-.700)	1041
	05-06	*	< LOD	.300 (<LOD-.400)	.500 (.400-.700)	.700 (.500-1.10)	1072
	07-08	*	< LOD	.200 (<LOD-.200)	.400 (.300-.500)	.600 (.400-.700)	1041
	09-10	.175 (.153-.199)	.200 (.100-.200)	.300 (.200-.400)	.600 (.400-.800)	.900 (.700-1.20)	1158
Race/ethnicity Mexican Americans	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.200)	584
	03-04	*	< LOD	< LOD	< LOD	< LOD	485
	05-06	*	< LOD	< LOD	.300 (<LOD-.700)	.500 (.300-1.00)	499
	07-08	*	< LOD	< LOD	.300 (<LOD-.400)	.300 (<LOD-1.40)	391
	09-10	*	.100 (.100-.100)	.200 (.200-.200)	.300 (.300-.400)	.400 (.300-.600)	461
Non-Hispanic blacks	99-00	*	.200 (.200-.200)	.300 (.200-.500)	.600 (.300-1.00)	.900 (.600-1.80)	303
	03-04	*	< LOD	< LOD	.600 (<LOD-1.40)	.900 (.300-2.90)	538
	05-06	*	< LOD	.300 (<LOD-.500)	.600 (.400-1.50)	1.20 (.600-2.70)	544
	07-08	*	< LOD	.300 (.200-.400)	.500 (.400-.700)	.800 (.600-1.20)	419
	09-10	.237 (.199-.283)	.200 (.200-.200)	.400 (.300-.500)	.800 (.500-1.10)	1.30 (.800-1.80)	391
Non-Hispanic whites	99-00	*	.200 (.200-.200)	.200 (.200-.200)	.200 (.200-.300)	.300 (.200-.800)	519
	03-04	*	< LOD	< LOD	< LOD	.500 (<LOD-.900)	962
	05-06	*	< LOD	.300 (.200-.400)	.500 (.400-.700)	.600 (.400-1.50)	935
	07-08	*	< LOD	.200 (<LOD-.200)	.300 (.300-.500)	.500 (.400-.600)	931
	09-10	.162 (.136-.193)	.200 (.100-.200)	.300 (.200-.400)	.500 (.300-.800)	.800 (.500-1.00)	1031

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 03-04, 05-06, 07-08, and 09-10 are 0.2, 0.3, 0.2, 0.2, and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PFCs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PFCs_FactSheet.html

Urinary Mono-benzyl phthalate (MBzP)

Metabolite of Benzylbutyl phthalate (BzBP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	11.0 (9.88-12.3)	12.2 (11.0-13.6)	25.5 (23.5-28.6)	48.3 (39.8-59.3)	74.4 (68.1-83.4)	2541
	01-02	10.8 (10.0-11.7)	11.4 (10.7-12.5)	27.4 (24.8-29.7)	58.2 (51.3-63.5)	87.6 (73.7-102)	2782
	03-04	9.89 (9.12-10.7)	10.3 (9.22-11.8)	23.3 (21.7-25.2)	47.9 (41.3-54.4)	72.6 (61.4-90.2)	2605
	05-06	8.32 (7.28-9.51)	8.93 (7.78-10.4)	20.7 (18.2-23.7)	44.6 (37.6-52.8)	67.1 (57.4-77.7)	2548
	07-08	7.24 (6.47-8.09)	8.42 (7.20-9.50)	18.3 (16.6-20.4)	37.3 (31.3-43.3)	58.8 (48.7-69.1)	2604
	09-10	6.46 (5.76-7.24)	6.70 (5.97-7.49)	15.4 (13.4-17.1)	31.0 (26.8-36.1)	48.3 (43.1-55.4)	2749
Age group							
6-11 years	99-00	28.4 (23.7-34.0)	29.0 (24.3-35.0)	59.0 (40.2-70.6)	92.8 (70.6-154)	154 (77.8-287)	328
	01-02	24.1 (21.0-27.6)	26.6 (19.2-31.2)	49.3 (44.4-66.8)	120 (83.7-137)	169 (132-237)	393
	03-04	24.4 (20.3-29.2)	25.2 (21.7-28.7)	45.9 (36.3-66.5)	105 (79.1-154)	184 (105-263)	342
	05-06	21.3 (19.2-23.7)	23.5 (19.4-26.6)	46.1 (37.1-56.9)	78.6 (64.3-104)	116 (88.4-137)	356
	07-08	15.4 (12.3-19.3)	17.6 (13.8-20.8)	33.8 (27.0-47.5)	69.9 (51.3-104)	131 (69.9-221)	389
	09-10	11.6 (9.51-14.1)	12.6 (10.4-14.5)	28.4 (20.2-37.8)	63.9 (47.4-76.8)	87.8 (65.4-212)	415
12-19 years	99-00	18.4 (15.7-21.6)	20.4 (16.1-25.1)	36.9 (31.5-42.1)	63.7 (48.4-82.6)	90.2 (67.5-122)	752
	01-02	16.7 (14.3-19.6)	17.9 (15.3-22.3)	40.0 (34.1-45.3)	81.1 (66.1-95.5)	122 (96.3-143)	742
	03-04	15.9 (14.0-18.1)	17.9 (16.1-19.6)	35.9 (28.7-46.5)	64.7 (51.2-86.3)	110 (71.9-136)	729
	05-06	14.2 (11.9-16.9)	14.8 (11.5-18.3)	31.8 (24.2-41.0)	62.6 (44.0-85.0)	94.2 (67.7-134)	702
	07-08	11.6 (9.56-14.2)	14.6 (11.5-16.6)	27.2 (22.0-31.5)	46.7 (35.4-58.3)	63.9 (54.3-102)	401
	09-10	10.6 (8.75-12.9)	11.9 (10.1-15.4)	24.8 (21.9-29.2)	43.1 (36.0-45.8)	56.1 (44.7-74.3)	420
20 years and older	99-00	8.96 (7.83-10.3)	9.94 (8.71-11.2)	21.0 (18.1-23.8)	37.4 (31.6-45.0)	62.1 (39.4-85.5)	1461
	01-02	9.16 (8.40-9.98)	10.0 (9.22-10.7)	22.9 (20.5-24.1)	47.1 (38.7-54.9)	71.8 (59.6-87.1)	1647
	03-04	8.22 (7.42-9.11)	8.71 (7.78-9.72)	19.4 (17.0-21.2)	39.0 (33.9-44.3)	57.2 (47.9-68.3)	1534
	05-06	6.87 (6.00-7.86)	7.34 (6.41-8.71)	17.1 (14.5-19.7)	34.0 (28.3-43.4)	56.9 (45.9-66.7)	1490
	07-08	6.18 (5.40-7.08)	7.13 (6.12-8.06)	15.5 (13.4-17.6)	30.5 (25.2-37.7)	49.5 (40.3-63.9)	1814
	09-10	5.61 (4.97-6.34)	5.95 (5.17-6.61)	12.7 (10.9-14.8)	25.6 (21.5-29.4)	39.6 (34.2-47.7)	1914
Gender							
Males	99-00	11.6 (10.1-13.4)	12.7 (10.9-14.4)	25.5 (22.7-29.0)	50.0 (43.1-62.8)	77.8 (69.3-93.4)	1215
	01-02	11.2 (9.78-12.9)	11.5 (10.5-13.3)	26.6 (23.8-31.0)	56.4 (45.7-70.1)	87.5 (63.5-132)	1371
	03-04	10.5 (9.50-11.6)	10.9 (9.50-12.6)	23.3 (21.2-25.6)	47.2 (38.6-57.3)	72.6 (56.6-94.8)	1250
	05-06	9.47 (8.20-10.9)	10.2 (8.78-12.2)	23.3 (19.3-25.9)	51.6 (39.8-61.6)	71.0 (61.6-83.2)	1270
	07-08	7.80 (6.96-8.74)	8.64 (7.49-10.0)	17.6 (15.8-20.4)	33.8 (28.4-41.3)	49.3 (41.5-62.1)	1294
	09-10	6.93 (6.07-7.92)	6.98 (6.21-7.80)	15.6 (13.3-17.6)	33.0 (27.4-37.8)	49.0 (40.4-62.8)	1399
Females	99-00	10.5 (9.17-12.0)	11.6 (10.2-13.8)	25.8 (22.2-29.8)	45.9 (38.7-59.3)	73.9 (60.6-83.4)	1326
	01-02	10.5 (9.40-11.7)	11.1 (9.94-12.9)	27.4 (23.3-31.5)	58.6 (49.2-66.0)	88.0 (73.2-103)	1411
	03-04	9.33 (8.53-10.2)	9.58 (8.71-11.2)	24.0 (21.2-26.1)	48.3 (42.3-52.1)	72.5 (60.8-86.3)	1355
	05-06	7.34 (6.30-8.55)	7.70 (6.55-9.36)	18.4 (15.6-22.2)	39.5 (33.0-46.3)	62.1 (48.0-73.7)	1278
	07-08	6.73 (5.77-7.85)	7.99 (6.41-9.36)	18.8 (15.4-22.5)	40.1 (32.2-50.9)	64.4 (51.1-78.8)	1310
	09-10	6.04 (5.38-6.77)	6.36 (5.18-7.47)	15.3 (12.9-16.9)	29.3 (24.5-36.9)	47.7 (38.0-60.4)	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.576, 0.216, 0.072, 0.216, 0.216, and 0.216 respectively.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.72 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/BzBP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-benzyl phthalate (MBzP)

Metabolite of Benzylbutyl phthalate (BzBP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	10.0 (8.69-11.6)	11.3 (9.65-12.2)	23.8 (19.8-26.0)	48.6 (40.0-60.5)	71.1 (58.0-108)	814
	01-02	9.49 (7.75-11.6)	10.7 (7.78-13.3)	21.2 (18.9-27.4)	50.7 (38.2-61.5)	68.2 (50.6-116)	677
	03-04	10.7 (9.52-12.0)	11.2 (9.72-13.2)	23.3 (19.9-25.5)	51.4 (35.0-66.5)	71.9 (62.3-104)	652
	05-06	8.95 (7.12-11.3)	10.2 (8.57-11.9)	20.1 (14.3-31.0)	43.1 (33.9-49.8)	59.8 (43.6-77.3)	637
	07-08	7.87 (6.39-9.68)	9.43 (6.91-11.5)	18.7 (14.1-25.3)	38.0 (30.2-47.5)	60.4 (40.5-73.9)	531
	09-10	7.58 (6.42-8.96)	7.39 (6.33-9.00)	17.1 (14.3-20.3)	33.8 (26.5-43.5)	54.5 (34.4-79.7)	566
Non-Hispanic blacks	99-00	16.6 (14.9-18.4)	16.6 (14.8-18.4)	35.5 (31.7-40.0)	68.2 (57.6-93.4)	99.1 (76.6-174)	603
	01-02	17.1 (15.1-19.4)	17.4 (14.3-20.2)	36.4 (29.9-45.3)	72.6 (62.2-91.3)	103 (91.2-129)	703
	03-04	13.6 (11.8-15.7)	14.4 (11.1-17.7)	31.5 (28.0-35.3)	58.2 (50.4-76.0)	86.2 (71.8-124)	699
	05-06	11.9 (9.03-15.7)	12.7 (10.1-17.1)	26.1 (21.0-35.0)	60.6 (47.2-74.6)	83.2 (74.6-102)	678
	07-08	9.37 (8.28-10.6)	9.72 (8.35-11.7)	21.4 (18.4-24.8)	46.9 (34.3-61.6)	71.3 (63.7-89.6)	597
	09-10	8.98 (7.67-10.5)	9.14 (7.76-10.5)	21.0 (16.4-24.7)	38.9 (30.3-52.6)	67.4 (51.3-80.7)	516
Non-Hispanic whites	99-00	10.3 (9.12-11.6)	11.6 (10.3-13.4)	24.5 (22.0-27.6)	42.3 (36.9-53.4)	73.9 (53.4-83.4)	912
	01-02	10.1 (9.12-11.1)	10.5 (9.65-11.2)	25.6 (23.2-28.4)	55.2 (47.6-65.0)	87.5 (67.1-112)	1216
	03-04	9.25 (8.30-10.3)	9.50 (8.42-11.5)	22.0 (20.0-25.3)	45.9 (38.6-52.1)	65.9 (54.9-87.9)	1088
	05-06	7.67 (6.61-8.90)	8.06 (6.98-10.1)	19.8 (17.1-23.5)	43.4 (32.4-52.5)	65.8 (52.5-76.8)	1038
	07-08	7.10 (6.33-7.97)	8.14 (7.06-9.29)	17.9 (16.5-20.3)	37.7 (31.0-44.2)	57.0 (46.0-68.6)	1077
	09-10	6.07 (5.24-7.03)	6.31 (5.54-7.23)	14.7 (12.4-16.7)	29.5 (23.4-36.9)	44.6 (34.9-64.1)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.576, 0.216, 0.072, 0.216, 0.216, and 0.216 respectively.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.72 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/BzBP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-benzyl phthalate (MBzP) (creatinine corrected)

Metabolite of Benzylbutyl phthalate (BzBP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	10.0 (9.33-10.8)	9.60 (9.22-10.3)	18.1 (16.8-19.6)	36.1 (29.9-42.3)	55.7 (50.1-62.1)	2541
	01-02	10.2 (9.50-10.9)	9.71 (9.12-10.4)	19.1 (17.8-20.9)	39.5 (35.3-42.1)	65.1 (53.6-73.2)	2782
	03-04	9.29 (8.76-9.86)	9.04 (8.07-9.82)	17.7 (16.0-19.0)	33.1 (29.5-37.1)	50.4 (44.9-57.5)	2605
	05-06	8.11 (7.28-9.04)	8.24 (7.37-8.98)	15.3 (13.9-17.4)	30.2 (26.4-35.1)	47.4 (42.7-53.3)	2548
	07-08	7.29 (6.71-7.93)	7.20 (6.57-8.17)	14.3 (13.1-15.4)	28.3 (24.4-31.8)	42.6 (36.3-49.1)	2604
	09-10	6.74 (6.06-7.50)	6.40 (5.63-7.30)	12.4 (11.3-13.8)	24.0 (20.4-26.6)	34.1 (30.5-41.4)	2749
Age group							
6-11 years	99-00	28.8 (24.2-34.3)	27.8 (21.7-36.9)	52.9 (40.8-71.4)	75.1 (64.4-102)	102 (71.8-124)	328
	01-02	27.4 (24.8-30.3)	26.9 (24.3-30.4)	48.9 (40.2-57.9)	96.7 (83.3-127)	140 (86.9-219)	393
	03-04	25.8 (21.9-30.3)	23.2 (20.1-29.1)	43.7 (36.1-57.2)	97.8 (61.6-154)	165 (71.8-286)	342
	05-06	23.5 (21.1-26.2)	25.5 (20.5-29.5)	47.1 (35.8-57.8)	77.6 (62.8-94.3)	104 (85.8-130)	356
	07-08	19.0 (16.0-22.5)	20.7 (15.6-27.0)	35.4 (32.3-40.0)	67.2 (51.9-91.5)	118 (74.3-159)	389
	09-10	15.1 (12.7-18.0)	14.6 (12.0-17.2)	29.3 (25.9-37.6)	66.5 (45.3-84.1)	92.2 (67.1-163)	415
12-19 years	99-00	12.4 (11.1-14.0)	12.3 (10.5-14.6)	20.4 (17.5-25.1)	35.8 (27.9-49.9)	50.4 (35.7-59.0)	752
	01-02	12.9 (11.3-14.8)	13.0 (11.3-15.0)	24.4 (21.1-27.8)	48.7 (40.2-61.9)	72.0 (58.1-88.6)	742
	03-04	11.9 (11.1-12.8)	12.2 (11.1-13.1)	21.6 (18.0-25.6)	37.5 (29.3-44.5)	56.0 (39.5-82.6)	729
	05-06	10.6 (9.14-12.2)	10.5 (8.33-13.0)	18.5 (15.9-23.7)	35.8 (28.1-46.0)	48.8 (41.3-65.3)	702
	07-08	9.06 (7.76-10.6)	9.39 (8.12-11.1)	16.7 (13.8-21.0)	32.1 (22.0-45.9)	48.4 (32.3-55.6)	401
	09-10	8.54 (7.36-9.90)	8.73 (7.44-10.8)	14.9 (13.2-17.5)	24.9 (21.6-28.9)	32.5 (25.7-42.7)	420
20 years and older	99-00	8.47 (7.73-9.27)	8.68 (8.01-9.26)	14.5 (13.2-16.8)	24.9 (22.0-29.4)	41.1 (29.7-53.2)	1461
	01-02	8.66 (8.08-9.28)	8.47 (8.00-9.00)	15.6 (14.3-17.1)	30.5 (25.8-34.7)	46.8 (39.0-56.4)	1647
	03-04	7.90 (7.39-8.45)	7.58 (6.98-8.37)	14.2 (13.1-15.4)	26.3 (25.0-28.2)	40.2 (33.7-44.9)	1534
	05-06	6.89 (6.20-7.66)	7.10 (6.60-7.71)	13.1 (12.1-14.2)	22.4 (19.9-25.5)	34.6 (28.9-43.1)	1490
	07-08	6.35 (5.74-7.03)	6.41 (5.65-7.15)	12.3 (11.1-13.6)	21.8 (19.1-25.7)	32.6 (28.7-40.5)	1814
	09-10	5.94 (5.31-6.66)	5.67 (5.03-6.53)	10.8 (9.18-12.5)	19.2 (16.6-23.5)	29.1 (24.2-33.4)	1914
Gender							
Males	99-00	9.14 (8.51-9.82)	8.96 (8.32-9.39)	17.1 (15.5-18.8)	32.0 (25.5-41.0)	52.9 (34.9-71.8)	1215
	01-02	9.13 (8.18-10.2)	8.55 (7.85-9.53)	17.4 (15.1-19.0)	35.3 (31.5-40.4)	57.8 (43.8-74.7)	1371
	03-04	8.25 (7.60-8.95)	7.99 (7.01-9.03)	15.2 (14.0-17.0)	28.4 (24.6-35.8)	45.3 (37.4-59.8)	1250
	05-06	7.61 (6.65-8.70)	7.57 (6.65-8.91)	14.3 (13.0-16.6)	30.0 (25.5-34.5)	43.7 (36.2-48.2)	1270
	07-08	6.57 (5.91-7.31)	6.66 (5.68-7.46)	12.8 (11.3-14.8)	24.4 (21.2-29.5)	36.3 (30.6-45.5)	1294
	09-10	6.21 (5.56-6.94)	5.88 (5.13-6.74)	11.6 (10.1-12.9)	21.3 (18.8-24.1)	32.2 (27.6-35.6)	1399
Females	99-00	11.0 (9.96-12.1)	10.6 (9.60-11.5)	18.6 (17.3-21.1)	40.6 (33.7-43.6)	57.9 (43.3-84.2)	1326
	01-02	11.3 (10.2-12.4)	10.8 (9.97-11.9)	21.2 (18.6-24.5)	42.1 (35.9-50.0)	69.0 (50.0-83.4)	1411
	03-04	10.4 (9.65-11.2)	9.97 (9.20-10.9)	20.1 (17.8-22.6)	37.4 (31.2-43.4)	52.9 (45.8-65.0)	1355
	05-06	8.63 (7.84-9.51)	8.67 (8.12-9.26)	15.9 (14.8-18.3)	32.0 (26.3-41.4)	52.3 (45.2-62.8)	1278
	07-08	8.07 (7.29-8.93)	8.37 (7.20-9.16)	15.5 (13.9-18.0)	31.5 (27.5-34.2)	48.3 (40.2-54.9)	1310
	09-10	7.29 (6.47-8.21)	6.90 (5.90-8.04)	13.6 (11.9-15.3)	25.6 (21.9-30.4)	38.8 (30.7-47.3)	1350

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.72 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/BzBP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-benzyl phthalate (MBzP) (creatinine corrected)

Metabolite of Benzylbutyl phthalate (BzBP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
Race/ethnicity			50th	75th	90th	95th	
Mexican Americans	99-00	9.10 (8.20-10.1)	8.60 (7.87-9.53)	17.4 (15.5-19.1)	33.5 (30.2-38.7)	49.1 (40.6-67.5)	814
	01-02	8.95 (7.69-10.4)	8.55 (7.16-10.7)	17.1 (14.1-20.9)	33.6 (26.3-44.4)	51.5 (36.8-86.5)	677
	03-04	9.66 (8.41-11.1)	9.00 (8.25-9.66)	17.3 (15.1-20.0)	38.2 (27.6-46.7)	56.0 (41.1-89.8)	652
	05-06	8.05 (6.69-9.69)	8.38 (7.48-8.91)	16.0 (12.6-20.2)	31.0 (22.4-40.6)	45.7 (31.3-54.2)	637
	07-08	7.68 (6.13-9.62)	8.06 (6.42-10.3)	15.5 (12.5-19.2)	29.1 (21.9-36.9)	42.9 (33.5-51.8)	531
	09-10	7.53 (6.52-8.69)	7.29 (6.30-8.18)	13.3 (11.6-16.3)	32.3 (23.3-43.6)	52.2 (33.0-66.1)	566
Non-Hispanic blacks	99-00	10.7 (9.75-11.7)	9.83 (8.76-10.9)	19.4 (16.2-22.9)	40.5 (28.4-54.7)	62.5 (46.4-71.9)	603
	01-02	12.0 (10.5-13.7)	11.3 (9.84-13.9)	24.1 (19.1-27.4)	43.3 (38.7-50.0)	77.7 (54.4-83.4)	703
	03-04	9.65 (8.46-11.0)	9.81 (8.32-11.2)	17.4 (15.8-20.7)	36.0 (29.6-41.6)	53.8 (42.5-65.2)	699
	05-06	8.38 (6.44-10.9)	8.24 (6.72-10.6)	16.1 (12.8-21.5)	34.1 (25.8-43.9)	52.3 (39.9-62.9)	678
	07-08	7.28 (6.51-8.14)	7.07 (5.97-8.28)	14.7 (12.2-16.0)	25.8 (22.6-31.5)	42.3 (32.6-50.3)	597
	09-10	6.50 (5.49-7.70)	6.50 (5.38-7.45)	11.7 (9.80-14.9)	23.9 (17.0-31.8)	34.4 (25.8-48.9)	516
Non-Hispanic whites	99-00	10.0 (9.16-11.0)	9.66 (9.25-10.7)	18.2 (16.6-19.8)	38.3 (27.9-46.7)	56.1 (48.5-65.0)	912
	01-02	9.97 (9.24-10.8)	9.33 (8.69-10.3)	18.5 (17.1-20.1)	38.2 (33.5-42.1)	64.2 (49.7-78.5)	1216
	03-04	9.14 (8.48-9.84)	8.97 (7.77-9.93)	17.6 (15.6-19.2)	30.2 (28.1-35.2)	47.2 (41.4-57.3)	1088
	05-06	8.08 (7.17-9.10)	8.35 (7.31-9.49)	15.2 (13.7-17.4)	29.9 (25.1-36.5)	47.4 (40.0-57.1)	1038
	07-08	7.42 (6.90-7.97)	7.38 (6.65-8.23)	14.0 (12.9-15.4)	28.3 (24.2-32.3)	44.5 (36.3-52.4)	1077
	09-10	6.83 (6.08-7.67)	6.37 (5.58-7.34)	12.6 (11.1-14.5)	24.0 (19.5-27.6)	33.4 (27.9-44.6)	1206

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.72 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/BzBP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-*n*-butyl phthalate (MnBP)

Metabolite of Dibutyl phthalate (DBP) and Benzylbutyl phthalate (BzBP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.*

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	24.6 (22.1-27.4)	26.0 (23.6-29.2)	51.6 (44.5-60.3)	98.6 (90.2-114)	150 (121-169)	2541
	01-02	18.9 (17.4-20.6)	20.4 (19.2-21.8)	40.4 (36.5-44.2)	73.6 (65.3-85.6)	108 (94.1-122)	2782
	03-04	21.1 (19.8-22.5)	23.2 (21.2-24.8)	42.7 (38.5-47.2)	80.7 (70.2-93.9)	122 (104-137)	2605
	05-06	19.6 (18.2-21.3)	20.6 (18.6-23.0)	40.4 (36.4-44.9)	71.5 (66.5-79.5)	107 (96.3-124)	2548
	07-08	18.9 (17.2-20.7)	20.0 (18.1-21.9)	39.1 (34.5-43.4)	72.4 (64.5-77.7)	110 (94.5-120)	2604
	09-10	14.6 (13.2-16.2)	15.9 (14.3-17.3)	30.5 (27.2-33.3)	54.4 (49.7-59.1)	75.9 (67.5-86.8)	2749
Age group							
6-11 years	99-00	41.4 (35.6-48.0)	40.0 (36.2-49.2)	75.5 (59.1-92.8)	124 (98.4-159)	166 (127-279)	328
	01-02	31.1 (26.6-36.5)	32.4 (25.6-37.1)	62.1 (51.3-76.9)	107 (84.3-136)	159 (110-290)	393
	03-04	36.3 (30.7-42.9)	36.7 (28.5-47.5)	71.3 (56.9-87.2)	137 (107-162)	191 (150-243)	342
	05-06	30.5 (27.5-33.8)	32.3 (27.8-35.0)	57.4 (46.8-65.4)	93.4 (75.6-130)	136 (108-181)	356
	07-08	26.9 (22.5-32.1)	28.7 (21.3-38.0)	53.6 (45.6-63.6)	85.6 (71.4-102)	119 (91.6-150)	389
	09-10	21.7 (19.0-24.8)	23.3 (20.8-26.2)	44.9 (39.1-49.7)	83.8 (59.6-121)	124 (92.3-173)	415
12-19 years	99-00	36.0 (30.8-42.1)	36.3 (30.6-44.9)	68.6 (55.9-79.7)	119 (90.2-159)	165 (121-227)	752
	01-02	25.1 (21.6-29.2)	26.7 (22.0-32.7)	52.6 (48.4-60.4)	92.4 (72.7-121)	148 (106-185)	742
	03-04	26.7 (24.1-29.5)	28.2 (25.2-33.1)	52.0 (45.3-60.9)	97.5 (74.7-116)	134 (110-158)	729
	05-06	27.5 (24.3-31.3)	29.0 (23.7-36.6)	57.2 (48.7-64.9)	88.6 (75.1-105)	125 (96.2-154)	702
	07-08	26.2 (22.5-30.6)	27.3 (22.1-32.8)	51.7 (46.2-63.8)	111 (77.7-132)	149 (103-188)	401
	09-10	18.9 (16.1-22.2)	22.3 (19.4-24.4)	37.2 (31.3-43.4)	61.7 (52.8-73.2)	81.4 (68.9-86.3)	420
20 years and older	99-00	21.6 (19.0-24.5)	23.1 (19.7-26.1)	46.3 (36.9-53.6)	95.0 (78.7-111)	143 (117-161)	1461
	01-02	17.0 (15.4-18.8)	19.1 (17.1-20.4)	35.1 (31.6-40.2)	64.8 (57.3-79.7)	95.4 (84.6-113)	1647
	03-04	19.0 (17.7-20.5)	20.7 (18.9-22.9)	38.4 (35.7-42.7)	74.7 (64.1-82.6)	108 (90.7-127)	1534
	05-06	17.7 (16.0-19.6)	18.4 (16.6-20.6)	36.7 (32.5-41.4)	65.1 (55.4-74.3)	101 (84.3-121)	1490
	07-08	17.2 (15.6-19.1)	18.6 (16.7-20.3)	34.5 (31.5-39.1)	63.8 (56.4-72.5)	98.3 (83.0-115)	1814
	09-10	13.5 (12.0-15.1)	14.5 (13.0-16.0)	28.1 (24.1-32.0)	50.0 (44.1-56.1)	68.9 (60.6-77.3)	1914
Gender							
Males	99-00	22.0 (20.1-24.1)	23.2 (20.4-26.3)	43.1 (36.6-49.5)	84.4 (71.3-96.2)	116 (97.8-132)	1215
	01-02	17.7 (16.0-19.6)	19.3 (17.3-21.0)	34.5 (30.3-40.6)	62.1 (54.1-75.5)	95.2 (75.5-117)	1371
	03-04	20.0 (18.1-22.0)	21.1 (19.0-24.0)	39.4 (35.5-43.4)	65.5 (59.0-73.0)	95.9 (79.8-111)	1250
	05-06	19.8 (18.4-21.4)	20.7 (19.0-23.0)	38.5 (34.6-42.0)	63.8 (56.8-72.0)	96.2 (79.0-113)	1270
	07-08	18.4 (16.8-20.3)	19.0 (17.3-21.3)	36.0 (32.5-40.4)	60.9 (52.8-68.8)	88.3 (72.6-110)	1294
	09-10	14.5 (13.0-16.3)	15.4 (13.7-16.9)	28.4 (25.9-31.2)	51.7 (44.3-54.8)	68.9 (58.9-78.0)	1399
Females	99-00	27.3 (23.6-31.5)	30.0 (25.9-33.3)	59.7 (51.6-69.6)	120 (98.3-145)	167 (143-223)	1326
	01-02	20.2 (18.2-22.4)	21.7 (19.7-24.3)	46.7 (43.1-51.1)	85.0 (72.7-92.5)	121 (106-136)	1411
	03-04	22.2 (21.2-23.3)	24.4 (23.4-25.8)	47.3 (42.7-53.0)	95.9 (79.9-114)	137 (122-156)	1355
	05-06	19.5 (17.2-22.0)	20.4 (17.1-24.7)	42.3 (36.7-52.0)	83.2 (71.0-90.2)	124 (101-137)	1278
	07-08	19.3 (17.2-21.7)	20.8 (18.1-23.2)	42.5 (35.6-50.6)	83.7 (68.6-109)	132 (109-151)	1310
	09-10	14.7 (13.1-16.5)	16.4 (13.9-19.3)	32.7 (29.9-37.1)	57.7 (52.7-63.9)	86.2 (68.9-113)	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.9, 1.1, 0.4, 0.6, 0.6, and 0.4 respectively.

*In the 1999-2000 survey period, concentrations of mono-isobutyl and mono-*n*-butyl phthalates were measured together and expressed as the combined value, monobutyl phthalate (MBP)

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DBP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-*n*-butyl phthalate (MnBP)

Metabolite of Dibutyl phthalate (DBP) and Benzylbutyl phthalate (BzBP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.*

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	23.4 (21.8-25.1)	26.3 (23.9-28.1)	48.1 (41.2-56.7)	92.2 (78.9-101)	117 (104-131)	814
	01-02	20.1 (16.6-24.5)	23.1 (18.0-26.5)	42.1 (34.0-51.5)	77.1 (62.9-92.5)	112 (84.6-143)	677
	03-04	24.1 (19.8-29.5)	26.9 (20.2-32.0)	47.4 (38.0-58.5)	85.6 (61.7-117)	127 (99.8-165)	652
	05-06	22.4 (18.6-27.0)	23.5 (19.0-29.0)	45.4 (37.1-55.4)	90.2 (71.1-119)	132 (106-174)	637
	07-08	22.2 (18.5-26.7)	22.4 (17.8-27.1)	43.8 (36.2-50.4)	76.4 (59.8-106)	145 (85.6-223)	531
	09-10	17.2 (15.3-19.4)	17.3 (14.3-20.9)	34.3 (31.4-38.8)	56.1 (47.7-67.9)	80.0 (65.1-107)	566
Non-Hispanic blacks	99-00	37.0 (31.9-42.9)	38.7 (33.4-44.5)	78.2 (58.7-91.8)	118 (108-143)	167 (143-197)	603
	01-02	29.6 (26.6-33.1)	31.5 (28.7-34.1)	58.3 (51.2-63.4)	93.2 (79.5-121)	138 (110-184)	703
	03-04	30.1 (28.4-31.9)	31.5 (29.7-34.1)	64.1 (58.3-67.1)	106 (94.8-119)	144 (115-168)	699
	05-06	27.1 (23.1-31.8)	28.5 (25.0-31.2)	53.1 (39.9-65.8)	105 (77.3-135)	143 (118-183)	678
	07-08	25.0 (21.9-28.4)	25.8 (22.0-30.4)	52.4 (43.9-59.5)	91.2 (75.6-106)	114 (96.8-148)	597
	09-10	21.9 (19.2-25.0)	23.2 (20.0-26.3)	42.4 (37.1-47.5)	69.1 (59.4-84.2)	103 (74.8-134)	516
Non-Hispanic whites	99-00	21.8 (19.3-24.6)	23.2 (19.5-27.5)	46.3 (37.5-53.3)	90.2 (74.7-106)	142 (111-161)	912
	01-02	17.6 (16.0-19.3)	19.2 (17.0-21.0)	36.6 (32.4-42.6)	69.2 (59.2-87.6)	107 (89.8-123)	1216
	03-04	18.9 (17.6-20.3)	20.7 (18.9-22.8)	38.4 (35.5-42.7)	71.3 (60.1-80.0)	101 (90.7-124)	1088
	05-06	17.8 (16.2-19.5)	18.6 (16.9-20.7)	37.8 (33.1-42.1)	65.4 (55.4-73.9)	95.9 (79.5-113)	1038
	07-08	17.8 (16.2-19.5)	18.9 (17.2-20.9)	36.7 (32.3-41.4)	68.0 (58.9-77.2)	110 (86.5-120)	1077
	09-10	13.0 (11.6-14.5)	14.1 (12.7-15.8)	26.6 (23.4-31.0)	48.8 (42.2-54.3)	67.5 (57.6-76.5)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.9, 1.1, 0.4, 0.6, 0.6, and 0.4 respectively.

*In the 1999-2000 survey period, concentrations of mono-isobutyl and mono-*n*-butyl phthalates were measured together and expressed as the combined value, monobutyl phthalate (MBP)

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DBP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-*n*-butyl phthalate (MnBP) (creatinine corrected)

Metabolite of Dibutyl phthalate (DBP) and Benzylbutyl phthalate (BzBP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.*

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	22.4 (20.6-24.4)	21.9 (19.8-24.3)	38.9 (35.0-41.8)	68.3 (60.3-78.3)	97.7 (81.4-131)	2541
	01-02	17.8 (16.7-19.0)	17.4 (16.3-18.4)	30.3 (28.1-32.3)	52.4 (47.4-61.0)	81.3 (71.0-92.5)	2782
	03-04	19.8 (18.5-21.2)	19.3 (17.7-20.8)	33.9 (30.9-38.5)	59.0 (52.9-68.1)	91.6 (74.1-115)	2605
	05-06	19.2 (18.0-20.4)	18.3 (16.8-19.8)	30.8 (29.0-33.0)	50.8 (47.8-56.6)	77.8 (67.2-94.8)	2548
	07-08	19.0 (17.7-20.5)	18.8 (17.2-20.0)	31.4 (28.7-34.1)	52.0 (46.4-59.5)	77.1 (64.2-88.6)	2604
	09-10	15.3 (14.0-16.6)	14.4 (13.4-15.8)	23.6 (21.7-25.2)	38.7 (34.1-45.8)	57.4 (49.8-69.9)	2749
Age group							
6-11 years	99-00	41.9 (37.4-47.1)	39.1 (34.3-49.0)	65.9 (56.7-80.0)	108 (71.2-179)	159 (102-263)	328
	01-02	35.4 (31.7-39.6)	35.1 (29.3-38.9)	55.4 (50.1-62.3)	84.0 (69.0-113)	147 (93.8-235)	393
	03-04	38.4 (33.8-43.6)	39.0 (34.7-42.6)	59.3 (52.9-68.1)	104 (83.6-119)	137 (108-198)	342
	05-06	33.6 (30.8-36.6)	33.9 (30.6-37.4)	50.6 (45.0-58.3)	78.5 (66.8-111)	128 (73.4-172)	356
	07-08	33.1 (27.6-39.7)	34.2 (27.5-41.3)	51.9 (42.6-63.8)	81.8 (61.3-101)	109 (83.1-159)	389
	09-10	28.3 (25.0-32.0)	26.7 (24.0-31.1)	47.8 (39.4-53.9)	78.6 (59.8-101)	130 (98.9-156)	415
12-19 years	99-00	24.3 (21.2-27.8)	23.7 (20.6-27.4)	37.6 (31.6-43.8)	63.3 (52.4-76.4)	88.1 (61.5-142)	752
	01-02	19.4 (17.3-21.7)	20.3 (17.5-22.3)	34.9 (30.5-37.9)	53.4 (45.2-73.9)	89.7 (60.3-106)	742
	03-04	20.0 (18.7-21.3)	19.8 (18.2-21.7)	30.7 (27.9-34.8)	52.7 (43.4-65.4)	74.4 (56.0-90.9)	729
	05-06	20.5 (19.0-22.2)	21.2 (18.8-22.9)	31.9 (29.6-33.8)	49.3 (42.1-56.4)	70.2 (52.0-108)	702
	07-08	20.4 (18.4-22.6)	21.9 (18.7-25.7)	33.8 (30.0-38.5)	50.7 (42.4-64.0)	72.4 (50.7-84.5)	401
	09-10	15.2 (13.8-16.8)	15.3 (13.8-18.4)	22.5 (21.1-24.8)	31.8 (28.2-34.6)	42.3 (33.8-51.2)	420
20 years and older	99-00	20.4 (18.6-22.4)	19.5 (18.1-21.4)	34.9 (30.3-40.0)	62.4 (53.4-72.1)	91.0 (70.4-135)	1461
	01-02	16.1 (15.0-17.3)	15.5 (14.2-16.5)	26.3 (24.2-28.6)	44.2 (38.7-51.1)	71.6 (61.2-85.6)	1647
	03-04	18.3 (17.0-19.6)	17.7 (16.6-19.2)	31.0 (27.4-34.0)	53.3 (46.5-64.2)	83.8 (65.3-114)	1534
	05-06	17.8 (16.4-19.2)	16.8 (15.6-18.4)	28.1 (25.3-30.9)	46.1 (41.5-53.0)	74.3 (55.4-94.8)	1490
	07-08	17.7 (16.3-19.2)	17.2 (15.7-18.8)	28.2 (25.8-30.5)	46.7 (42.0-53.3)	71.8 (57.8-89.5)	1814
	09-10	14.3 (13.0-15.7)	13.5 (12.5-14.8)	21.6 (20.0-23.8)	34.5 (30.1-40.2)	50.9 (42.2-67.7)	1914
Gender							
Males	99-00	17.3 (16.1-18.6)	17.0 (15.5-18.8)	28.6 (25.8-32.1)	49.3 (42.6-53.5)	64.7 (57.3-71.5)	1215
	01-02	14.4 (13.5-15.4)	13.7 (12.9-14.9)	22.9 (20.8-24.6)	39.9 (35.6-44.0)	60.0 (50.5-76.2)	1371
	03-04	15.7 (14.5-16.9)	14.8 (13.6-16.0)	25.4 (23.6-28.1)	41.4 (37.0-47.5)	59.4 (50.3-81.5)	1250
	05-06	15.9 (15.1-16.8)	15.4 (14.2-16.4)	24.9 (23.1-27.0)	38.6 (35.7-42.3)	51.2 (46.1-63.5)	1270
	07-08	15.5 (14.4-16.8)	14.7 (13.4-16.3)	24.9 (22.6-27.6)	40.8 (35.2-44.9)	55.2 (46.1-65.2)	1294
	09-10	13.0 (12.0-14.2)	12.8 (11.7-13.6)	19.9 (18.1-21.2)	30.3 (28.4-33.2)	45.3 (39.7-53.8)	1399
Females	99-00	28.6 (25.3-32.3)	28.8 (25.5-30.5)	50.6 (41.9-56.3)	84.3 (69.2-106)	134 (93.6-155)	1326
	01-02	21.7 (19.6-23.9)	21.6 (19.7-23.6)	35.8 (33.0-38.7)	64.9 (58.9-70.2)	91.5 (81.4-103)	1411
	03-04	24.8 (22.9-26.8)	24.1 (21.6-26.8)	42.9 (39.2-48.9)	74.9 (64.2-86.6)	117 (83.6-139)	1355
	05-06	22.9 (20.9-25.1)	21.5 (19.4-24.6)	37.1 (33.3-41.9)	67.2 (56.8-74.4)	99.6 (77.8-128)	1278
	07-08	23.1 (21.0-25.5)	22.8 (20.9-25.2)	38.0 (32.7-43.1)	62.7 (52.1-78.5)	89.8 (76.2-126)	1310
	09-10	17.8 (16.0-19.7)	17.6 (15.1-19.4)	27.1 (24.4-30.3)	47.8 (37.9-53.8)	70.6 (53.8-91.5)	1350

*In the 1999-2000 survey period, concentrations of mono-isobutyl and mono-*n*-butyl phthalates were measured together and expressed as the combined value, monobutyl phthalate (MBP)

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DBP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-*n*-butyl phthalate (MnBP) (creatinine corrected)

Metabolite of Dibutyl phthalate (DBP) and Benzylbutyl phthalate (BzBP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.*

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	21.2 (19.3-23.3)	20.0 (18.2-22.9)	40.1 (32.6-44.3)	63.6 (57.5-70.1)	82.9 (73.9-100)	814
	01-02	19.0 (16.2-22.2)	19.2 (16.3-21.9)	33.7 (28.3-39.6)	61.0 (43.9-84.0)	86.7 (60.6-128)	677
	03-04	21.8 (17.7-26.8)	20.4 (17.4-24.0)	37.6 (29.6-50.1)	75.3 (55.6-92.0)	96.3 (76.4-164)	652
	05-06	20.1 (17.8-22.8)	19.1 (16.4-23.0)	34.3 (29.2-42.6)	64.8 (54.6-75.9)	98.1 (70.2-154)	637
	07-08	21.7 (17.7-26.5)	20.9 (16.4-27.1)	36.9 (29.1-46.7)	59.5 (47.1-88.9)	102 (63.4-161)	531
	09-10	17.1 (15.5-18.8)	15.0 (14.0-16.3)	27.4 (23.6-31.6)	51.0 (42.9-69.6)	88.1 (59.8-100)	566
Non-Hispanic blacks	99-00	23.9 (21.3-26.8)	25.0 (20.7-28.1)	42.2 (35.9-49.6)	70.0 (61.1-83.9)	96.2 (83.9-105)	603
	01-02	20.8 (18.8-23.1)	20.2 (19.2-22.8)	34.5 (30.9-36.8)	62.8 (50.6-74.6)	85.6 (72.1-99.0)	703
	03-04	21.4 (19.5-23.4)	22.0 (19.6-24.8)	35.4 (31.6-42.7)	65.7 (54.0-75.3)	94.0 (71.1-128)	699
	05-06	19.1 (16.4-22.1)	18.5 (16.2-20.7)	30.1 (24.9-39.5)	54.6 (42.0-81.3)	84.7 (51.3-125)	678
	07-08	19.4 (17.3-21.7)	19.2 (16.8-22.0)	30.5 (27.0-36.5)	50.7 (42.8-60.4)	78.3 (59.3-125)	597
	09-10	15.9 (14.5-17.4)	15.3 (13.5-18.6)	24.9 (22.6-27.2)	37.6 (33.8-45.4)	51.2 (44.7-70.6)	516
Non-Hispanic whites	99-00	21.3 (19.1-23.8)	20.5 (18.6-23.2)	36.4 (31.5-41.0)	67.1 (56.7-78.4)	97.7 (73.5-142)	912
	01-02	17.4 (16.2-18.6)	16.5 (15.3-17.8)	29.0 (26.6-32.2)	51.1 (46.0-60.0)	81.4 (68.1-99.0)	1216
	03-04	18.6 (17.3-20.1)	17.9 (16.7-19.8)	31.6 (27.7-37.8)	53.3 (48.2-61.5)	81.5 (64.2-108)	1088
	05-06	18.7 (17.5-19.9)	17.9 (16.3-19.5)	29.7 (27.5-32.0)	48.7 (42.2-55.4)	74.3 (58.3-94.5)	1038
	07-08	18.5 (17.3-19.9)	18.1 (16.6-19.6)	30.4 (27.3-33.3)	50.9 (43.6-61.0)	75.8 (62.6-88.0)	1077
	09-10	14.6 (13.5-15.7)	13.9 (13.2-15.0)	22.1 (20.3-24.2)	35.3 (30.9-40.6)	52.5 (45.8-63.0)	1206

*In the 1999-2000 survey period, concentrations of mono-isobutyl and mono-*n*-butyl phthalates were measured together and expressed as the combined value, monobutyl phthalate (MBP)

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DBP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-cyclohexyl phthalate (MCHP)

Metabolite of Dicyclohexyl phthalate (DCHP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	2.21 (<LOD-3.42)	2541
	01-02	*	< LOD	< LOD	.804 (.603-1.01)	1.01 (.804-1.21)	2782
	03-04	*	< LOD	< LOD	< LOD	.603 (.402-.804)	2605
	05-06	*	< LOD	< LOD	< LOD	< LOD	2548
	07-08	*	< LOD	< LOD	< LOD	< LOD	2604
	09-10	*	< LOD	< LOD	< LOD	< LOD	2749
Age group							
6-11 years	99-00	*	< LOD	< LOD	2.01 (<LOD-2.21)	3.42 (2.01-7.64)	328
	01-02	*	< LOD	< LOD	.804 (.603-1.01)	1.21 (1.01-1.41)	393
	03-04	*	< LOD	< LOD	.603 (<LOD-.804)	1.01 (.603-1.01)	342
	05-06	*	< LOD	< LOD	< LOD	< LOD	356
	07-08	*	< LOD	< LOD	< LOD	< LOD	389
	09-10	*	< LOD	< LOD	< LOD	.960 (.640-1.57)	415
12-19 years	99-00	*	< LOD	< LOD	2.01 (<LOD-3.02)	3.42 (2.01-5.03)	752
	01-02	*	< LOD	< LOD	.804 (.603-1.01)	1.01 (.804-1.21)	742
	03-04	*	< LOD	< LOD	.402 (<LOD-.603)	.804 (.603-1.21)	729
	05-06	*	< LOD	< LOD	< LOD	< LOD	702
	07-08	*	< LOD	< LOD	< LOD	< LOD	401
	09-10	*	< LOD	< LOD	< LOD	< LOD	420
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1461
	01-02	*	< LOD	< LOD	.804 (<LOD-1.01)	1.01 (.804-1.21)	1647
	03-04	*	< LOD	< LOD	< LOD	.603 (.402-.603)	1534
	05-06	*	< LOD	< LOD	< LOD	< LOD	1490
	07-08	*	< LOD	< LOD	< LOD	< LOD	1814
	09-10	*	< LOD	< LOD	< LOD	< LOD	1914
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	2.21 (<LOD-4.02)	1215
	01-02	*	< LOD	< LOD	.804 (.603-1.01)	1.01 (.804-1.21)	1371
	03-04	*	< LOD	< LOD	< LOD	.603 (.603-.804)	1250
	05-06	*	< LOD	< LOD	< LOD	< LOD	1270
	07-08	*	< LOD	< LOD	< LOD	< LOD	1294
	09-10	*	< LOD	< LOD	< LOD	< LOD	1399
Females	99-00	*	< LOD	< LOD	< LOD	2.21 (<LOD-3.82)	1326
	01-02	*	< LOD	< LOD	.804 (<LOD-.804)	1.01 (.804-1.01)	1411
	03-04	*	< LOD	< LOD	< LOD	.603 (.402-1.01)	1355
	05-06	*	< LOD	< LOD	< LOD	< LOD	1278
	07-08	*	< LOD	< LOD	< LOD	< LOD	1310
	09-10	*	< LOD	< LOD	< LOD	< LOD	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.81, 0.603, 0.402, 0.603, 0.603, and 0.402 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 2.01 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DCHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-cyclohexyl phthalate (MCHP)

Metabolite of Dicyclohexyl phthalate (DCHP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

Race/ethnicity	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	814
	01-02	*	< LOD	< LOD	.804 (<LOD-1.01)	1.01 (.603-1.41)	677
	03-04	*	< LOD	< LOD	< LOD	.603 (.402-1.01)	652
	05-06	*	< LOD	< LOD	< LOD	< LOD	637
	07-08	*	< LOD	< LOD	< LOD	< LOD	531
	09-10	*	< LOD	< LOD	< LOD	.440 (<LOD-.640)	566
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	2.21 (1.81-2.41)	603
	01-02	*	< LOD	< LOD	.804 (.603-1.01)	1.01 (.804-1.41)	703
	03-04	*	< LOD	< LOD	.603 (.402-.603)	.804 (.603-1.01)	699
	05-06	*	< LOD	< LOD	< LOD	< LOD	678
	07-08	*	< LOD	< LOD	< LOD	< LOD	597
	09-10	*	< LOD	< LOD	< LOD	.480 (<LOD-.620)	516
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	2.01 (<LOD-3.42)	912
	01-02	*	< LOD	< LOD	.804 (<LOD-.804)	1.01 (.804-1.21)	1216
	03-04	*	< LOD	< LOD	< LOD	.603 (.402-.804)	1088
	05-06	*	< LOD	< LOD	< LOD	< LOD	1038
	07-08	*	< LOD	< LOD	< LOD	< LOD	1077
	09-10	*	< LOD	< LOD	< LOD	< LOD	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.81, 0.603, 0.402, 0.603, 0.603, and 0.402 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 2.01 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DCHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-cyclohexyl phthalate (MCHP) (creatinine corrected)

Metabolite of Dicyclohexyl phthalate (DCHP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	6.03 (<LOD-6.69)	2541
	01-02	*	< LOD	< LOD	1.23 (1.07-1.39)	1.83 (1.55-2.01)	2782
	03-04	*	< LOD	< LOD	< LOD	.905 (.804-1.01)	2605
	05-06	*	< LOD	< LOD	< LOD	< LOD	2548
	07-08	*	< LOD	< LOD	< LOD	< LOD	2604
	09-10	*	< LOD	< LOD	< LOD	< LOD	2749
Age group							
6-11 years	99-00	*	< LOD	< LOD	3.10 (<LOD-4.70)	5.67 (3.10-12.9)	328
	01-02	*	< LOD	< LOD	1.39 (1.03-1.49)	1.89 (1.39-2.35)	393
	03-04	*	< LOD	< LOD	.744 (<LOD-1.07)	1.07 (.704-1.67)	342
	05-06	*	< LOD	< LOD	< LOD	< LOD	356
	07-08	*	< LOD	< LOD	< LOD	< LOD	389
	09-10	*	< LOD	< LOD	< LOD	1.80 (1.34-2.55)	415
12-19 years	99-00	*	< LOD	< LOD	2.45 (<LOD-3.10)	3.36 (2.73-3.66)	752
	01-02	*	< LOD	< LOD	.945 (.764-1.33)	1.55 (1.07-2.37)	742
	03-04	*	< LOD	< LOD	.442 (<LOD-.543)	.764 (.482-1.25)	729
	05-06	*	< LOD	< LOD	< LOD	< LOD	702
	07-08	*	< LOD	< LOD	< LOD	< LOD	401
	09-10	*	< LOD	< LOD	< LOD	< LOD	420
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1461
	01-02	*	< LOD	< LOD	1.27 (<LOD-1.39)	1.83 (1.55-2.11)	1647
	03-04	*	< LOD	< LOD	< LOD	.905 (.804-1.01)	1534
	05-06	*	< LOD	< LOD	< LOD	< LOD	1490
	07-08	*	< LOD	< LOD	< LOD	< LOD	1814
	09-10	*	< LOD	< LOD	< LOD	< LOD	1914
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	4.30 (<LOD-6.35)	1215
	01-02	*	< LOD	< LOD	1.03 (.844-1.33)	1.77 (1.35-2.13)	1371
	03-04	*	< LOD	< LOD	< LOD	.663 (.583-1.01)	1250
	05-06	*	< LOD	< LOD	< LOD	< LOD	1270
	07-08	*	< LOD	< LOD	< LOD	< LOD	1294
	09-10	*	< LOD	< LOD	< LOD	< LOD	1399
Females	99-00	*	< LOD	< LOD	< LOD	6.69 (<LOD-7.10)	1326
	01-02	*	< LOD	< LOD	1.35 (<LOD-1.55)	1.83 (1.61-2.21)	1411
	03-04	*	< LOD	< LOD	< LOD	1.01 (.844-1.19)	1355
	05-06	*	< LOD	< LOD	< LOD	< LOD	1278
	07-08	*	< LOD	< LOD	< LOD	< LOD	1310
	09-10	*	< LOD	< LOD	< LOD	< LOD	1350

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 2.01 for survey periods 1999-2008 compared with results previously released

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DCHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-cyclohexyl phthalate (MCHP) (creatinine corrected)

Metabolite of Dicyclohexyl phthalate (DCHP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	814
	01-02	*	< LOD	< LOD	1.19 (<LOD-1.39)	1.91 (1.59-2.23)	677
	03-04	*	< LOD	< LOD	< LOD	.784 (.623-1.13)	652
	05-06	*	< LOD	< LOD	< LOD	< LOD	637
	07-08	*	< LOD	< LOD	< LOD	< LOD	531
	09-10	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.56)	566
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	2.87 (2.25-3.50)	603
	01-02	*	< LOD	< LOD	.824 (.724-.985)	1.19 (1.01-1.43)	703
	03-04	*	< LOD	< LOD	.503 (.342-.623)	.663 (.523-.945)	699
	05-06	*	< LOD	< LOD	< LOD	< LOD	678
	07-08	*	< LOD	< LOD	< LOD	< LOD	597
	09-10	*	< LOD	< LOD	< LOD	.880 (<LOD-1.08)	516
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	6.35 (<LOD-7.10)	912
	01-02	*	< LOD	< LOD	1.27 (<LOD-1.49)	1.83 (1.55-2.13)	1216
	03-04	*	< LOD	< LOD	< LOD	.965 (.844-1.07)	1088
	05-06	*	< LOD	< LOD	< LOD	< LOD	1038
	07-08	*	< LOD	< LOD	< LOD	< LOD	1077
	09-10	*	< LOD	< LOD	< LOD	< LOD	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 2.01 for survey periods 1999-2008 compared with results previously released

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DCHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-ethyl phthalate (MEP)

Metabolite of Diethyl phthalate (DEP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	118 (103-135)	108 (90.0-132)	300 (244-355)	830 (664-977)	1870 (1420-2490)	2536
	01-02	117 (105-131)	112 (93.0-128)	307 (274-348)	813 (688-947)	1650 (1230-2130)	2782
	03-04	127 (112-145)	115 (99.9-137)	332 (302-366)	909 (769-1150)	1780 (1420-2180)	2605
	05-06	108 (95.5-123)	102 (83.4-121)	284 (257-323)	750 (707-825)	1410 (1140-1830)	2548
	07-08	90.4 (80.3-102)	81.1 (70.8-93.5)	239 (205-272)	605 (493-713)	1180 (821-1760)	2604
	09-10	64.4 (58.3-71.2)	54.9 (50.5-62.7)	163 (145-187)	485 (374-604)	988 (805-1180)	2749
Age group							
6-11 years	99-00	60.3 (49.4-73.6)	49.8 (41.0-61.8)	130 (85.2-164)	249 (192-481)	499 (250-703)	328
	01-02	56.2 (47.0-67.2)	47.5 (40.9-61.1)	121 (93.7-143)	297 (208-420)	533 (377-717)	393
	03-04	62.9 (54.3-72.9)	53.9 (46.4-67.9)	130 (112-159)	344 (195-377)	546 (358-949)	342
	05-06	55.7 (45.7-67.9)	47.9 (38.3-61.6)	111 (73.9-147)	289 (173-368)	522 (326-719)	356
	07-08	49.0 (42.4-56.6)	45.2 (38.5-53.3)	108 (80.5-135)	193 (155-227)	296 (214-359)	389
	09-10	35.2 (31.2-39.8)	33.0 (29.8-37.8)	68.7 (54.3-80.9)	151 (114-207)	288 (205-530)	415
12-19 years	99-00	139 (105-183)	127 (92.8-169)	372 (276-540)	994 (694-1420)	2150 (1020-2920)	752
	01-02	130 (105-161)	121 (97.9-150)	316 (256-429)	835 (649-974)	1360 (969-2020)	742
	03-04	148 (123-178)	146 (109-194)	367 (285-459)	828 (642-1030)	1520 (900-2180)	729
	05-06	126 (108-146)	114 (96.4-134)	315 (256-390)	765 (636-1060)	1360 (1030-1680)	702
	07-08	98.6 (79.1-123)	97.4 (68.3-130)	214 (173-330)	477 (356-705)	821 (511-1270)	401
	09-10	64.0 (50.1-81.8)	53.9 (39.0-75.4)	151 (101-228)	354 (260-497)	662 (371-1190)	420
20 years and older	99-00	125 (108-145)	119 (92.6-146)	318 (257-389)	882 (666-1100)	2300 (1470-3060)	1456
	01-02	126 (113-141)	120 (101-140)	329 (291-374)	888 (702-1100)	1800 (1420-2420)	1647
	03-04	135 (116-157)	124 (104-145)	351 (311-415)	1050 (807-1370)	1970 (1480-2510)	1534
	05-06	114 (99.0-132)	111 (87.1-131)	299 (268-346)	782 (724-901)	1480 (1140-2080)	1490
	07-08	95.3 (85.4-106)	84.7 (75.6-97.8)	264 (235-299)	673 (539-823)	1470 (913-1920)	1814
	09-10	69.0 (61.8-76.9)	60.5 (54.0-68.7)	183 (159-207)	557 (445-712)	1110 (875-1350)	1914
Gender							
Males	99-00	118 (98.5-142)	102 (78.3-130)	345 (246-429)	948 (672-1500)	2310 (1400-3010)	1214
	01-02	120 (103-139)	113 (91.8-131)	331 (276-398)	957 (703-1390)	2040 (1390-2900)	1371
	03-04	130 (114-148)	113 (97.8-133)	354 (296-432)	1000 (801-1370)	1920 (1460-2300)	1250
	05-06	107 (92.1-124)	95.9 (75.8-120)	279 (227-340)	757 (651-855)	1470 (923-1930)	1270
	07-08	92.8 (80.1-107)	81.0 (70.4-93.3)	245 (194-298)	659 (481-867)	1450 (851-1900)	1294
	09-10	61.0 (54.2-68.6)	52.8 (45.4-58.3)	157 (126-202)	436 (342-586)	965 (768-1170)	1399
Females	99-00	118 (102-136)	115 (91.3-139)	280 (231-335)	652 (581-813)	1470 (903-2560)	1322
	01-02	115 (101-131)	110 (91.5-128)	282 (255-329)	692 (580-867)	1230 (982-1650)	1411
	03-04	125 (106-148)	120 (90.9-145)	316 (258-372)	830 (639-1080)	1710 (1190-2260)	1355
	05-06	110 (92.8-130)	109 (86.8-130)	291 (252-346)	734 (671-871)	1310 (1140-1900)	1278
	07-08	88.2 (77.8-99.9)	82.4 (67.6-101)	233 (202-264)	553 (462-703)	1050 (713-1680)	1310
	09-10	67.8 (60.3-76.4)	59.6 (52.4-73.7)	164 (148-192)	548 (392-675)	988 (777-1340)	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.792, 0.594, 0.264, 0.528, 0.462, and 0.396 respectively.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.66 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-ethyl phthalate (MEP)

Metabolite of Diethyl phthalate (DEP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	119 (103-138)	115 (96.5-138)	291 (257-357)	842 (561-996)	1130 (966-1410)	813
	01-02	149 (129-173)	145 (125-174)	349 (293-435)	984 (690-1390)	1730 (1010-2940)	677
	03-04	176 (158-196)	164 (140-202)	394 (345-454)	1080 (873-1440)	2010 (1470-2840)	652
	05-06	163 (130-203)	161 (124-186)	384 (310-466)	1030 (682-1830)	1930 (1170-3330)	637
	07-08	122 (105-141)	111 (82.0-144)	314 (270-372)	850 (604-971)	1180 (971-1520)	531
	09-10	82.2 (70.0-96.5)	72.6 (59.6-84.0)	199 (142-254)	586 (415-869)	1300 (818-1760)	566
Non-Hispanic blacks	99-00	213 (182-249)	202 (169-231)	521 (419-626)	1250 (932-1500)	2380 (1410-3060)	603
	01-02	233 (214-253)	235 (191-269)	563 (468-718)	1430 (1070-1630)	2340 (1860-3350)	703
	03-04	236 (205-272)	202 (167-273)	626 (507-760)	1650 (1220-1970)	2880 (1840-3900)	699
	05-06	219 (189-254)	199 (163-253)	522 (440-700)	1450 (1070-1900)	2610 (1900-3600)	678
	07-08	168 (145-195)	165 (132-202)	415 (343-493)	994 (711-1260)	1920 (1090-3370)	597
	09-10	156 (130-188)	137 (113-159)	405 (276-543)	1190 (841-1570)	1930 (1430-3380)	516
Non-Hispanic whites	99-00	101 (87.6-116)	88.4 (71.0-104)	242 (189-318)	651 (527-882)	1630 (1050-2560)	908
	01-02	104 (92.8-117)	96.8 (78.8-117)	273 (241-297)	672 (598-813)	1530 (1030-1800)	1216
	03-04	110 (95.4-128)	97.8 (82.4-115)	276 (242-315)	801 (630-975)	1480 (1050-2170)	1088
	05-06	91.3 (80.3-104)	84.2 (69.4-111)	232 (196-278)	600 (500-715)	1140 (757-1550)	1038
	07-08	78.9 (70.4-88.5)	68.9 (59.8-80.1)	199 (175-233)	506 (429-650)	880 (670-1500)	1077
	09-10	52.1 (48.1-56.5)	45.1 (40.0-51.6)	127 (106-140)	309 (259-394)	768 (468-988)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.792, 0.594, 0.264, 0.528, 0.462, and 0.396 respectively.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.66 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-ethyl phthalate (MEP) (creatinine corrected)

Metabolite of Diethyl phthalate (DEP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	107 (98.0-118)	93.0 (85.4-104)	237 (203-278)	598 (497-781)	1290 (1100-1530)	2536
	01-02	110 (99.3-122)	97.4 (87.9-107)	256 (218-287)	640 (532-778)	1220 (959-1450)	2782
	03-04	120 (107-133)	101 (90.1-117)	298 (254-332)	733 (587-853)	1350 (1080-1680)	2605
	05-06	106 (95.3-117)	92.3 (80.9-104)	242 (211-286)	625 (528-711)	1140 (942-1310)	2548
	07-08	91.1 (82.6-101)	75.3 (68.6-83.5)	204 (185-231)	547 (433-680)	987 (821-1190)	2604
	09-10	67.2 (60.5-74.7)	56.1 (48.9-66.8)	150 (128-172)	433 (373-524)	850 (709-999)	2749
Age group							
6-11 years	99-00	61.1 (51.4-72.7)	52.5 (43.3-72.4)	109 (83.6-137)	225 (145-365)	412 (264-517)	328
	01-02	64.0 (54.4-75.2)	53.6 (43.7-69.4)	117 (88.8-148)	338 (191-529)	556 (338-874)	393
	03-04	66.6 (57.5-77.1)	57.4 (43.9-74.7)	119 (95.6-134)	310 (205-394)	474 (310-877)	342
	05-06	61.3 (49.6-75.8)	50.1 (42.7-61.0)	107 (76.2-145)	279 (167-441)	512 (366-1210)	356
	07-08	60.4 (53.9-67.7)	52.8 (42.4-64.2)	108 (96.3-134)	205 (166-237)	278 (228-386)	389
	09-10	45.9 (41.0-51.3)	39.0 (34.7-41.8)	73.0 (63.2-87.2)	168 (125-234)	296 (182-590)	415
12-19 years	99-00	93.7 (78.6-112)	80.7 (61.4-103)	240 (181-326)	580 (446-834)	1160 (661-1320)	752
	01-02	100 (83.0-121)	92.4 (73.1-119)	218 (164-270)	533 (389-729)	878 (573-1220)	742
	03-04	111 (93.1-132)	99.1 (81.2-121)	240 (193-320)	522 (389-714)	969 (651-1470)	729
	05-06	93.6 (80.5-109)	85.3 (69.0-104)	201 (174-238)	500 (389-635)	813 (590-1110)	702
	07-08	76.7 (61.8-95.1)	72.8 (53.3-91.3)	161 (108-218)	334 (213-491)	547 (334-866)	401
	09-10	51.4 (40.8-64.9)	46.6 (31.2-68.8)	108 (84.2-137)	236 (150-393)	419 (268-524)	420
20 years and older	99-00	118 (107-131)	102 (89.5-117)	257 (222-298)	669 (530-965)	1430 (1180-2210)	1456
	01-02	119 (108-132)	105 (96.3-120)	277 (240-321)	702 (583-871)	1400 (1010-1840)	1647
	03-04	130 (115-147)	117 (96.8-133)	338 (279-417)	815 (652-1060)	1510 (1170-2110)	1534
	05-06	115 (103-128)	100 (87.6-117)	278 (233-309)	700 (574-803)	1220 (972-1530)	1490
	07-08	97.9 (89.6-107)	78.7 (73.2-87.7)	236 (208-266)	624 (518-759)	1100 (875-1380)	1814
	09-10	73.0 (65.1-81.9)	62.5 (52.4-72.8)	173 (149-195)	523 (407-662)	989 (810-1120)	1914
Gender							
Males	99-00	92.8 (82.0-105)	79.0 (70.8-88.2)	214 (165-274)	661 (457-978)	1290 (965-1920)	1214
	01-02	97.6 (85.8-111)	83.2 (72.6-96.8)	232 (186-280)	729 (554-974)	1400 (980-2000)	1371
	03-04	102 (91.9-113)	83.7 (71.0-99.0)	258 (225-307)	704 (562-896)	1320 (1060-1590)	1250
	05-06	85.7 (75.4-97.4)	73.0 (61.0-90.2)	204 (172-231)	504 (368-673)	898 (607-1240)	1270
	07-08	78.1 (68.5-89.1)	65.5 (57.0-73.8)	181 (162-216)	490 (390-625)	936 (662-1190)	1294
	09-10	54.7 (48.3-62.0)	43.3 (38.2-54.2)	125 (97.7-153)	314 (262-436)	792 (570-935)	1399
Females	99-00	123 (109-139)	104 (93.7-118)	249 (203-326)	543 (460-770)	1270 (770-2250)	1322
	01-02	123 (110-139)	113 (97.4-124)	269 (235-312)	568 (470-725)	944 (784-1330)	1411
	03-04	139 (119-163)	120 (97.8-143)	335 (250-418)	741 (587-923)	1490 (951-2200)	1355
	05-06	129 (116-144)	106 (97.4-121)	293 (247-333)	759 (629-920)	1370 (1150-1730)	1278
	07-08	106 (95.5-117)	88.5 (77.6-98.6)	227 (188-278)	610 (491-727)	1040 (759-1520)	1310
	09-10	81.9 (72.7-92.2)	69.9 (57.2-80.9)	178 (151-214)	523 (407-661)	987 (716-1220)	1350

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.66 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-ethyl phthalate (MEP) (creatinine corrected)

Metabolite of Diethyl phthalate (DEP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	108 (93.9-125)	102 (89.8-115)	252 (207-312)	538 (444-643)	875 (643-1270)	813
	01-02	141 (120-165)	131 (108-160)	304 (261-378)	714 (568-1090)	1280 (928-1740)	677
	03-04	159 (141-179)	154 (133-178)	352 (275-439)	815 (631-1060)	1500 (1190-2070)	652
	05-06	146 (123-173)	125 (107-157)	324 (235-425)	767 (600-1020)	1410 (926-2510)	637
	07-08	119 (100-140)	105 (81.9-142)	260 (224-306)	616 (502-750)	1230 (628-2160)	531
	09-10	81.6 (70.9-93.8)	64.9 (56.3-72.7)	170 (131-254)	523 (324-792)	945 (685-1510)	566
Non-Hispanic blacks	99-00	137 (121-156)	130 (109-151)	292 (257-333)	682 (503-1120)	1270 (810-1710)	603
	01-02	163 (149-179)	150 (122-178)	367 (316-408)	818 (634-979)	1380 (1020-1850)	703
	03-04	167 (148-189)	140 (124-162)	409 (314-522)	1010 (793-1300)	1710 (1420-2310)	699
	05-06	154 (131-181)	134 (113-160)	360 (311-412)	919 (590-1290)	1700 (1070-2030)	678
	07-08	131 (115-148)	121 (105-132)	280 (224-334)	614 (489-843)	1100 (836-1780)	597
	09-10	113 (89.0-143)	92.7 (77.5-121)	236 (180-338)	722 (443-1210)	1300 (890-2240)	516
Non-Hispanic whites	99-00	98.3 (88.9-109)	84.5 (72.9-93.7)	206 (158-255)	565 (432-916)	1290 (975-1810)	908
	01-02	103 (93.5-114)	89.6 (82.1-98.9)	223 (190-265)	607 (470-765)	1050 (874-1440)	1216
	03-04	109 (95.3-125)	88.9 (79.2-105)	254 (223-320)	652 (548-876)	1270 (922-1830)	1088
	05-06	96.1 (86.5-107)	84.5 (71.2-97.6)	211 (189-242)	532 (450-686)	993 (828-1240)	1038
	07-08	82.4 (74.7-90.9)	66.4 (58.5-73.8)	185 (158-219)	523 (397-640)	871 (708-1040)	1077
	09-10	58.6 (53.2-64.4)	49.1 (41.4-56.8)	125 (110-146)	366 (278-437)	706 (515-965)	1206

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 0.66 for survey periods 1999-2008 compared with results previously released.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-2-ethylhexyl phthalate (MEHP)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	3.43 (3.19-3.69)	3.20 (3.00-3.60)	7.60 (6.80-8.40)	14.9 (13.5-17.4)	23.8 (19.2-28.6)	2541
	01-02	4.27 (3.80-4.79)	4.20 (3.70-4.90)	9.80 (8.40-11.6)	23.0 (19.1-27.9)	39.2 (31.8-50.0)	2782
	03-04	2.34 (2.10-2.62)	1.90 (1.70-2.40)	5.30 (4.50-6.60)	15.1 (11.4-20.6)	31.0 (21.4-42.0)	2605
	05-06	3.04 (2.78-3.32)	2.50 (2.10-2.80)	6.30 (5.70-7.10)	17.7 (14.0-22.5)	39.7 (28.6-52.1)	2548
	07-08	2.64 (2.29-3.05)	2.20 (1.80-2.50)	5.40 (4.30-6.90)	14.1 (11.2-20.2)	27.6 (19.8-39.8)	2604
	09-10	1.59 (1.41-1.79)	1.51 (1.33-1.71)	3.52 (2.99-4.00)	7.54 (5.96-9.54)	14.1 (9.91-21.1)	2749
Age group							
6-11 years	99-00	5.12 (4.42-5.92)	4.90 (3.70-6.40)	11.1 (8.30-13.6)	19.0 (13.8-36.1)	35.3 (15.6-130)	328
	01-02	4.41 (3.90-5.00)	4.40 (4.10-5.30)	9.30 (7.90-11.7)	19.7 (14.6-25.9)	31.4 (21.8-47.9)	393
	03-04	2.84 (2.10-3.84)	2.70 (1.80-4.10)	6.40 (4.40-9.60)	13.9 (7.80-27.6)	27.6 (11.3-64.7)	342
	05-06	3.10 (2.78-3.47)	3.00 (2.60-3.30)	6.20 (5.10-7.10)	14.1 (9.40-19.3)	19.7 (14.7-36.6)	356
	07-08	2.39 (2.05-2.80)	2.20 (1.70-2.90)	4.50 (3.70-6.20)	8.70 (6.40-13.9)	15.1 (10.6-24.1)	389
	09-10	1.64 (1.45-1.85)	1.71 (1.26-2.02)	3.50 (3.09-3.94)	5.95 (4.56-7.56)	8.92 (6.94-12.9)	415
12-19 years	99-00	3.75 (3.24-4.35)	3.70 (2.90-4.60)	8.10 (6.40-9.40)	15.3 (11.4-20.5)	22.8 (19.1-29.2)	752
	01-02	4.57 (3.96-5.27)	4.50 (3.70-5.10)	11.0 (9.50-14.4)	23.0 (17.7-32.7)	42.5 (25.9-57.5)	742
	03-04	2.77 (2.25-3.41)	2.50 (2.00-3.00)	6.40 (4.50-8.60)	18.6 (10.2-35.6)	40.6 (20.7-58.4)	729
	05-06	3.72 (3.04-4.56)	3.20 (2.40-4.10)	8.80 (6.20-13.3)	22.6 (13.8-43.4)	48.7 (23.1-62.9)	702
	07-08	2.99 (2.39-3.75)	2.30 (1.80-2.70)	6.00 (4.40-9.90)	21.1 (11.8-32.8)	37.6 (24.8-74.1)	401
	09-10	1.82 (1.52-2.16)	1.66 (1.42-1.94)	3.98 (3.35-4.73)	9.53 (6.27-14.0)	17.6 (9.54-27.4)	420
20 years and older	99-00	3.21 (2.94-3.51)	3.00 (2.70-3.40)	7.30 (6.40-8.00)	14.5 (12.1-17.0)	22.7 (17.5-27.0)	1461
	01-02	4.20 (3.63-4.86)	4.10 (3.50-5.00)	9.50 (8.10-11.9)	23.5 (18.0-29.8)	39.5 (30.3-57.1)	1647
	03-04	2.23 (2.03-2.44)	1.70 (1.50-2.00)	5.10 (4.50-6.00)	15.1 (10.9-19.7)	29.5 (20.4-40.0)	1534
	05-06	2.94 (2.68-3.21)	2.30 (1.90-2.70)	6.20 (5.60-6.70)	17.7 (13.6-24.5)	41.5 (28.6-54.1)	1490
	07-08	2.62 (2.27-3.02)	2.10 (1.80-2.50)	5.40 (4.20-7.10)	14.4 (11.2-20.2)	27.3 (19.2-40.6)	1814
	09-10	1.55 (1.36-1.78)	1.44 (1.25-1.68)	3.40 (2.88-4.03)	7.39 (5.94-9.58)	14.6 (9.91-21.1)	1914
Gender							
Males	99-00	3.68 (3.31-4.10)	3.40 (2.90-3.90)	8.00 (7.40-8.80)	16.0 (14.0-19.0)	25.3 (19.5-36.7)	1215
	01-02	4.31 (3.84-4.83)	4.30 (3.70-5.10)	9.70 (8.30-11.2)	23.0 (16.9-29.8)	37.9 (29.9-48.4)	1371
	03-04	2.56 (2.26-2.90)	2.20 (1.70-2.60)	6.00 (4.60-7.70)	17.2 (11.3-26.3)	33.3 (24.9-55.5)	1250
	05-06	3.40 (3.01-3.85)	2.80 (2.40-3.30)	7.00 (5.70-9.00)	22.6 (15.0-35.8)	49.8 (35.2-67.4)	1270
	07-08	2.77 (2.35-3.27)	2.30 (1.90-2.80)	5.50 (4.20-7.60)	14.4 (11.2-20.5)	28.9 (19.2-40.0)	1294
	09-10	1.83 (1.63-2.05)	1.77 (1.63-1.91)	3.97 (3.49-4.51)	8.63 (6.94-11.9)	18.0 (12.6-29.0)	1399
Females	99-00	3.21 (2.91-3.54)	3.10 (2.80-3.50)	7.10 (5.90-8.50)	13.6 (12.1-17.2)	21.9 (15.6-28.5)	1326
	01-02	4.23 (3.67-4.86)	4.10 (3.50-5.00)	9.80 (8.40-12.2)	23.0 (19.5-28.4)	43.5 (31.4-53.7)	1411
	03-04	2.15 (1.92-2.42)	1.80 (1.50-2.10)	4.90 (4.10-5.70)	13.2 (10.0-18.1)	27.8 (17.5-40.7)	1355
	05-06	2.72 (2.49-2.98)	2.10 (1.90-2.40)	6.00 (5.20-6.80)	13.9 (11.7-17.5)	30.8 (21.9-36.2)	1278
	07-08	2.52 (2.18-2.92)	2.00 (1.70-2.40)	5.10 (4.20-6.30)	14.1 (9.50-21.4)	26.4 (19.2-42.1)	1310
	09-10	1.39 (1.21-1.60)	1.30 (1.12-1.53)	3.00 (2.51-3.62)	6.48 (5.45-8.09)	10.3 (8.33-14.9)	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.2, 1.0, 0.9, 1.2, 1.1, and 0.5 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-2-ethylhexyl phthalate (MEHP)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	3.49 (3.16-3.85)	3.50 (3.10-3.90)	7.00 (5.70-8.60)	13.3 (10.7-18.7)	23.9 (17.4-27.3)	814
	01-02	4.32 (3.75-4.98)	4.70 (3.80-5.70)	10.1 (8.50-11.4)	19.6 (16.6-23.0)	28.5 (24.2-39.9)	677
	03-04	2.35 (1.87-2.96)	2.20 (1.50-3.00)	5.10 (4.30-6.60)	11.2 (7.50-16.5)	18.5 (11.6-38.2)	652
	05-06	2.99 (2.50-3.57)	2.30 (1.70-3.30)	5.70 (4.70-7.60)	18.4 (12.1-30.6)	36.4 (26.2-63.8)	637
	07-08	2.89 (2.38-3.50)	2.60 (2.00-3.10)	5.30 (4.40-8.20)	16.9 (10.3-27.3)	30.2 (22.9-34.3)	531
	09-10	2.08 (1.84-2.36)	2.10 (1.86-2.40)	4.44 (3.73-5.63)	9.67 (6.89-16.0)	17.9 (13.1-24.3)	566
Non-Hispanic blacks	99-00	4.82 (3.92-5.93)	5.20 (4.10-5.80)	9.50 (7.60-11.4)	19.5 (12.9-26.5)	29.5 (18.6-60.3)	603
	01-02	6.60 (5.57-7.82)	6.70 (5.40-8.10)	15.4 (13.0-18.7)	32.9 (26.5-41.4)	52.6 (41.0-84.0)	703
	03-04	3.61 (3.07-4.23)	3.50 (3.00-4.00)	8.50 (7.10-11.4)	22.9 (16.5-28.6)	35.2 (29.3-49.1)	699
	05-06	4.09 (3.51-4.75)	3.70 (3.10-4.10)	9.10 (6.90-11.9)	22.5 (17.4-40.7)	59.3 (34.5-86.7)	678
	07-08	3.30 (2.98-3.64)	3.20 (2.70-3.60)	7.20 (6.30-8.90)	15.3 (13.3-20.2)	25.5 (19.2-38.1)	597
	09-10	2.08 (1.79-2.41)	2.05 (1.70-2.42)	4.79 (3.80-5.95)	10.1 (7.50-14.4)	16.4 (9.60-38.4)	516
Non-Hispanic whites	99-00	3.16 (2.89-3.46)	2.80 (2.50-3.10)	7.40 (6.30-8.40)	14.5 (12.2-17.4)	22.4 (16.9-28.5)	912
	01-02	3.85 (3.37-4.40)	3.70 (3.10-4.40)	8.70 (7.80-9.90)	20.9 (17.3-25.9)	37.9 (29.9-49.5)	1216
	03-04	2.14 (1.92-2.39)	1.70 (1.40-1.90)	4.80 (4.00-5.80)	13.6 (9.50-20.0)	31.0 (18.1-48.9)	1088
	05-06	2.83 (2.59-3.10)	2.20 (1.80-2.60)	5.90 (5.30-6.90)	17.0 (13.3-21.6)	36.3 (26.6-51.0)	1038
	07-08	2.44 (2.07-2.88)	2.00 (1.70-2.30)	4.90 (3.70-6.20)	13.1 (8.70-20.1)	25.0 (15.8-42.1)	1077
	09-10	1.41 (1.22-1.62)	1.30 (1.12-1.57)	3.04 (2.53-3.60)	6.12 (5.23-7.65)	11.8 (7.63-21.1)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.2, 1.0, 0.9, 1.2, 1.1, and 0.5 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-2-ethylhexyl phthalate (MEHP) (creatinine corrected)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	3.12 (2.95-3.31)	3.08 (2.82-3.27)	5.88 (5.38-6.25)	10.8 (9.62-12.5)	18.9 (15.0-21.8)	2541
	01-02	4.00 (3.58-4.48)	3.90 (3.44-4.47)	7.94 (7.22-9.02)	18.0 (15.3-21.5)	32.8 (25.2-42.9)	2782
	03-04	2.20 (2.01-2.41)	1.89 (1.68-2.19)	4.31 (3.84-4.74)	10.8 (8.72-13.8)	25.4 (16.7-34.7)	2605
	05-06	2.96 (2.72-3.23)	2.61 (2.37-2.86)	5.69 (5.00-6.47)	13.7 (11.4-17.8)	30.1 (21.2-35.2)	2548
	07-08	2.66 (2.37-2.99)	2.36 (2.11-2.67)	5.15 (4.35-6.00)	11.8 (8.89-15.6)	21.9 (14.6-33.4)	2604
	09-10	1.66 (1.45-1.90)	1.52 (1.36-1.77)	3.09 (2.69-3.60)	6.28 (4.98-8.48)	11.2 (8.33-19.1)	2749
Age group							
6-11 years	99-00	5.19 (4.55-5.93)	5.37 (4.52-5.95)	9.11 (8.06-11.4)	21.6 (11.5-41.9)	41.9 (13.5-86.2)	328
	01-02	5.03 (4.47-5.65)	5.38 (4.51-6.21)	9.90 (7.87-11.5)	21.1 (13.8-28.8)	31.4 (24.3-40.7)	393
	03-04	3.00 (2.30-3.93)	2.80 (1.93-4.09)	5.86 (4.69-7.70)	14.3 (8.54-24.4)	28.7 (14.1-45.3)	342
	05-06	3.42 (3.08-3.79)	3.26 (2.63-3.92)	6.18 (5.40-6.85)	11.3 (8.96-17.4)	20.7 (11.3-31.8)	356
	07-08	2.95 (2.49-3.49)	2.80 (2.17-3.33)	5.42 (3.95-6.51)	10.6 (7.47-14.0)	15.6 (10.6-23.7)	389
	09-10	2.13 (1.90-2.40)	2.08 (1.88-2.33)	3.69 (3.13-4.20)	5.83 (4.80-7.95)	8.89 (5.88-20.1)	415
12-19 years	99-00	2.53 (2.14-2.99)	2.35 (2.05-2.76)	5.83 (4.38-6.29)	9.66 (7.41-11.5)	12.1 (10.5-17.3)	752
	01-02	3.53 (3.09-4.03)	3.67 (2.89-4.48)	7.47 (6.51-8.67)	15.2 (11.7-21.9)	25.2 (17.7-32.8)	742
	03-04	2.07 (1.74-2.48)	1.88 (1.60-2.23)	4.25 (3.19-5.62)	11.6 (6.83-23.2)	24.8 (11.6-37.9)	729
	05-06	2.77 (2.27-3.38)	2.43 (2.03-2.87)	5.24 (4.06-7.75)	15.2 (9.86-23.2)	27.1 (16.0-43.7)	702
	07-08	2.33 (1.90-2.86)	2.00 (1.67-2.57)	4.33 (3.55-6.23)	12.6 (8.31-16.3)	21.9 (12.0-45.7)	401
	09-10	1.46 (1.20-1.77)	1.33 (1.09-1.61)	2.85 (2.24-3.63)	7.47 (4.31-11.6)	13.2 (8.11-20.9)	420
20 years and older	99-00	3.03 (2.83-3.25)	2.98 (2.73-3.23)	5.55 (4.90-6.06)	10.0 (8.60-12.9)	17.5 (13.8-22.1)	1461
	01-02	3.97 (3.49-4.52)	3.82 (3.26-4.38)	7.79 (7.00-9.00)	18.3 (15.3-21.8)	34.5 (23.1-47.9)	1647
	03-04	2.14 (1.98-2.31)	1.84 (1.63-2.08)	4.14 (3.78-4.40)	10.5 (8.38-12.9)	25.6 (15.9-36.3)	1534
	05-06	2.94 (2.69-3.23)	2.60 (2.36-2.83)	5.67 (4.77-6.52)	13.8 (11.3-18.1)	33.1 (21.9-47.4)	1490
	07-08	2.69 (2.39-3.02)	2.36 (2.14-2.66)	5.20 (4.38-6.04)	11.8 (8.94-16.6)	22.1 (13.5-37.1)	1814
	09-10	1.65 (1.43-1.90)	1.51 (1.34-1.75)	3.04 (2.63-3.60)	6.24 (4.98-8.61)	11.1 (8.03-19.4)	1914
Gender							
Males	99-00	2.89 (2.60-3.22)	2.76 (2.52-2.96)	5.58 (4.71-6.08)	10.3 (9.35-12.4)	21.6 (14.1-27.7)	1215
	01-02	3.50 (3.08-3.99)	3.33 (2.83-3.90)	7.00 (6.49-7.77)	16.2 (12.8-20.9)	31.6 (20.5-49.4)	1371
	03-04	2.01 (1.82-2.21)	1.71 (1.46-1.89)	4.14 (3.49-4.81)	10.4 (7.68-16.2)	23.3 (15.1-41.1)	1250
	05-06	2.73 (2.43-3.07)	2.30 (2.12-2.61)	5.08 (4.29-6.14)	14.3 (11.2-20.5)	31.0 (21.5-50.9)	1270
	07-08	2.33 (2.03-2.68)	2.00 (1.71-2.34)	4.36 (3.71-5.41)	11.6 (8.33-14.2)	20.2 (14.6-26.3)	1294
	09-10	1.64 (1.45-1.85)	1.52 (1.39-1.74)	3.06 (2.70-3.60)	6.53 (5.18-9.49)	13.0 (8.65-22.2)	1399
Females	99-00	3.36 (3.11-3.63)	3.33 (2.91-3.80)	6.15 (5.55-6.77)	11.1 (9.11-14.0)	17.3 (12.4-24.6)	1326
	01-02	4.54 (4.02-5.13)	4.47 (3.85-5.14)	9.28 (7.94-10.3)	20.3 (16.6-24.4)	34.7 (27.1-42.0)	1411
	03-04	2.40 (2.15-2.69)	2.16 (1.84-2.40)	4.40 (3.97-4.89)	10.9 (8.27-16.0)	27.0 (17.5-34.6)	1355
	05-06	3.20 (2.89-3.55)	2.89 (2.58-3.17)	6.07 (5.00-7.00)	13.3 (10.4-16.3)	28.2 (18.2-37.4)	1278
	07-08	3.02 (2.70-3.38)	2.76 (2.36-3.02)	5.57 (4.90-6.50)	12.1 (8.64-18.2)	24.7 (14.8-44.4)	1310
	09-10	1.68 (1.44-1.96)	1.53 (1.35-1.89)	3.10 (2.68-3.69)	5.75 (4.58-8.03)	10.5 (7.36-17.5)	1350

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-2-ethylhexyl phthalate (MEHP) (creatinine corrected)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	3.16 (2.72-3.68)	3.15 (2.52-3.81)	5.88 (4.86-7.24)	11.6 (9.63-13.1)	15.7 (12.6-23.1)	814
	01-02	4.07 (3.60-4.61)	4.18 (3.82-4.90)	7.80 (6.64-9.49)	16.4 (13.6-18.9)	24.9 (19.8-28.7)	677
	03-04	2.12 (1.74-2.59)	1.94 (1.50-2.42)	4.06 (3.29-4.93)	9.38 (5.72-15.4)	16.8 (9.86-38.6)	652
	05-06	2.69 (2.36-3.07)	2.41 (2.04-2.73)	4.82 (4.09-6.05)	14.3 (10.0-16.9)	27.2 (16.3-40.1)	637
	07-08	2.82 (2.38-3.34)	2.45 (2.08-2.89)	5.00 (4.14-6.86)	12.3 (9.58-20.7)	29.0 (17.0-50.3)	531
	09-10	2.07 (1.83-2.34)	1.89 (1.69-2.31)	4.06 (3.50-4.98)	8.38 (6.08-11.3)	13.9 (9.70-20.0)	566
	Non-Hispanic blacks	99-00	3.11 (2.59-3.73)	3.13 (2.50-3.61)	5.84 (4.43-7.32)	10.2 (8.05-15.6)	18.4 (11.6-35.2)
01-02		4.63 (3.96-5.42)	4.59 (3.97-5.02)	9.93 (7.95-12.4)	21.2 (16.0-33.2)	39.9 (27.7-48.1)	703
03-04		2.56 (2.24-2.92)	2.28 (2.02-2.78)	5.17 (4.48-6.83)	13.2 (10.5-16.2)	27.5 (18.4-36.0)	699
05-06		2.87 (2.45-3.38)	2.41 (2.09-2.78)	5.72 (4.40-7.29)	15.0 (12.0-20.6)	54.4 (18.4-84.0)	678
07-08		2.56 (2.26-2.90)	2.40 (2.17-2.73)	4.77 (4.07-5.76)	11.4 (8.75-15.4)	18.0 (16.1-26.3)	597
09-10		1.51 (1.22-1.86)	1.48 (1.14-1.93)	2.87 (2.26-3.69)	5.73 (3.69-10.7)	10.7 (5.61-19.1)	516
Non-Hispanic whites		99-00	3.09 (2.84-3.36)	3.08 (2.73-3.47)	5.87 (5.11-6.67)	10.6 (8.95-13.5)	20.0 (14.0-24.6)
	01-02	3.81 (3.34-4.35)	3.67 (3.11-4.33)	7.78 (6.74-9.35)	17.0 (14.1-21.8)	32.8 (21.5-46.9)	1216
	03-04	2.12 (1.91-2.35)	1.82 (1.60-2.13)	4.11 (3.49-4.42)	10.7 (7.42-15.1)	27.0 (15.1-37.4)	1088
	05-06	2.98 (2.77-3.21)	2.66 (2.43-2.93)	5.73 (5.00-6.47)	13.4 (11.3-17.8)	27.7 (19.5-37.4)	1038
	07-08	2.55 (2.20-2.94)	2.26 (1.97-2.67)	4.90 (3.99-5.92)	11.0 (7.80-14.8)	20.5 (11.9-30.2)	1077
	09-10	1.58 (1.36-1.84)	1.49 (1.27-1.76)	2.94 (2.50-3.49)	5.64 (4.53-7.96)	10.3 (6.85-18.3)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	20.0 (17.8-22.5)	20.1 (17.8-22.4)	43.6 (38.0-49.7)	92.3 (77.0-108)	192 (131-256)	2782
	03-04	21.7 (19.3-24.4)	21.2 (18.7-24.1)	49.1 (40.5-56.9)	121 (91.3-164)	266 (165-383)	2605
	05-06	25.5 (23.0-28.2)	23.8 (21.5-26.8)	55.1 (50.2-61.0)	153 (132-180)	306 (240-421)	2548
	07-08	22.1 (18.7-26.0)	20.7 (17.6-23.7)	48.2 (39.8-61.8)	123 (83.7-181)	238 (171-336)	2604
	09-10	12.9 (11.3-14.7)	12.9 (11.4-14.8)	25.8 (22.1-30.3)	53.2 (41.7-75.2)	103 (74.2-149)	2749
Age group							
6-11 years	01-02	33.6 (29.7-37.9)	32.9 (26.9-39.1)	66.9 (49.7-74.0)	127 (103-148)	216 (137-280)	393
	03-04	36.9 (28.4-47.9)	36.5 (26.5-47.0)	77.4 (49.1-103)	164 (79.9-350)	318 (164-400)	342
	05-06	34.9 (30.8-39.6)	35.7 (31.3-40.7)	68.9 (56.5-76.0)	140 (101-169)	206 (133-401)	356
	07-08	28.6 (23.4-34.8)	27.0 (20.1-36.4)	56.1 (45.4-67.8)	113 (76.5-218)	242 (109-351)	389
	09-10	15.0 (13.2-17.1)	17.0 (14.1-19.8)	28.9 (24.4-37.4)	49.3 (41.2-70.2)	75.1 (55.2-117)	415
12-19 years	01-02	24.9 (21.3-29.1)	25.3 (22.9-31.3)	50.6 (40.7-64.5)	107 (78.5-148)	216 (117-330)	742
	03-04	28.3 (23.0-34.8)	29.8 (25.9-33.9)	56.9 (45.4-73.7)	157 (84.1-299)	317 (176-553)	729
	05-06	34.8 (28.0-43.3)	32.5 (27.1-42.2)	79.5 (66.9-103)	213 (131-384)	424 (232-836)	702
	07-08	29.8 (22.5-39.3)	26.6 (20.0-35.4)	66.7 (43.7-96.6)	224 (101-417)	417 (209-615)	401
	09-10	15.3 (12.4-18.8)	14.9 (12.0-18.0)	28.8 (22.7-41.0)	70.2 (41.0-110)	117 (61.0-215)	420
20 years and older	01-02	18.1 (15.7-20.9)	17.8 (14.7-20.7)	39.8 (32.7-48.0)	86.2 (65.7-107)	175 (110-279)	1647
	03-04	19.5 (17.7-21.5)	18.4 (16.6-21.0)	41.9 (36.9-51.2)	107 (88.2-136)	225 (148-384)	1534
	05-06	23.4 (21.1-25.9)	21.4 (19.5-23.7)	48.6 (43.7-55.1)	148 (121-172)	306 (238-421)	1490
	07-08	20.5 (17.4-24.1)	19.6 (17.1-22.2)	46.3 (37.2-59.9)	110 (75.0-169)	214 (157-303)	1814
	09-10	12.4 (10.7-14.3)	12.5 (10.8-14.3)	24.5 (21.1-29.3)	52.5 (40.1-72.6)	104 (72.6-151)	1914
Gender							
Males	01-02	22.0 (19.5-24.7)	21.2 (19.4-24.2)	48.0 (41.4-54.4)	94.2 (80.8-110)	212 (130-256)	1371
	03-04	24.1 (20.9-27.9)	22.9 (19.2-27.9)	51.0 (40.5-59.8)	133 (94.8-220)	317 (162-470)	1250
	05-06	29.6 (26.0-33.8)	27.2 (22.9-31.0)	63.1 (54.3-71.9)	180 (143-263)	494 (285-626)	1270
	07-08	23.2 (19.4-27.8)	20.8 (18.1-24.5)	46.0 (39.2-57.3)	132 (85.4-190)	277 (162-349)	1294
	09-10	15.2 (13.1-17.6)	14.6 (12.9-16.4)	29.5 (24.6-36.2)	71.0 (50.3-93.5)	128 (90.6-181)	1399
Females	01-02	18.3 (15.7-21.4)	18.2 (14.9-22.1)	39.8 (34.3-46.0)	86.0 (69.4-115)	170 (119-273)	1411
	03-04	19.7 (17.4-22.2)	19.4 (16.7-22.8)	46.4 (37.5-54.4)	103 (84.1-148)	214 (140-318)	1355
	05-06	22.0 (19.2-25.2)	21.4 (19.5-23.4)	47.1 (42.6-54.3)	135 (113-156)	232 (186-300)	1278
	07-08	21.0 (17.8-24.8)	19.9 (16.8-23.3)	51.0 (39.9-64.6)	121 (76.9-183)	223 (171-336)	1310
	09-10	11.0 (9.58-12.8)	11.6 (9.73-13.5)	22.5 (19.2-27.2)	42.8 (33.2-59.1)	82.2 (53.1-116)	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.0, 0.3, 0.7, 0.7, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	01-02	18.5 (16.2-21.1)	19.1 (16.3-21.6)	36.3 (31.6-44.0)	79.9 (66.4-93.9)	123 (100-161)	677
	03-04	18.9 (15.4-23.4)	19.8 (17.6-22.3)	37.5 (30.0-45.6)	72.2 (52.4-115)	116 (71.6-327)	652
	05-06	23.0 (18.0-29.3)	19.9 (15.7-23.9)	47.8 (34.8-65.3)	136 (84.6-223)	244 (157-520)	637
	07-08	22.7 (18.5-27.7)	19.8 (17.5-23.8)	43.6 (33.5-66.6)	104 (82.8-157)	238 (158-282)	531
	09-10	15.3 (12.9-18.2)	15.8 (13.3-18.3)	31.5 (26.6-39.2)	64.5 (46.4-94.9)	108 (70.7-153)	566
Non-Hispanic blacks	01-02	29.8 (26.1-34.1)	30.9 (27.2-34.3)	61.9 (52.6-69.4)	126 (108-157)	276 (157-339)	703
	03-04	30.8 (26.8-35.5)	29.1 (25.3-32.3)	65.6 (53.7-76.3)	154 (113-178)	275 (174-401)	699
	05-06	34.8 (30.3-39.9)	30.2 (27.6-33.2)	73.8 (61.0-96.7)	206 (156-275)	395 (274-547)	678
	07-08	25.7 (23.1-28.6)	25.8 (22.4-28.9)	55.9 (47.1-64.7)	121 (99.1-150)	184 (137-255)	597
	09-10	15.5 (12.8-18.8)	15.7 (13.4-17.7)	30.4 (24.4-35.7)	54.5 (37.7-84.1)	94.6 (51.5-229)	516
Non-Hispanic whites	01-02	19.1 (16.7-21.9)	19.2 (16.9-21.4)	41.7 (35.3-50.7)	91.1 (75.6-110)	212 (130-275)	1216
	03-04	20.8 (18.6-23.3)	19.7 (17.2-22.5)	47.5 (39.4-56.1)	120 (91.3-165)	270 (155-403)	1088
	05-06	24.3 (21.9-26.9)	23.0 (21.1-26.0)	54.6 (48.4-61.0)	148 (121-172)	302 (221-421)	1038
	07-08	21.3 (17.6-25.8)	20.2 (16.5-24.1)	46.7 (36.1-65.2)	123 (75.1-203)	277 (161-373)	1077
	09-10	12.2 (10.5-14.1)	12.3 (10.5-14.2)	24.4 (20.7-29.1)	51.7 (37.7-77.1)	104 (72.6-151)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.0, 0.3, 0.7, 0.7, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP) (creatinine corrected)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	18.8 (17.0-20.7)	16.6 (14.9-18.5)	32.2 (27.8-37.1)	71.1 (58.7-88.3)	143 (101-200)	2782
	03-04	20.4 (18.7-22.3)	17.7 (16.3-19.6)	35.8 (30.5-43.3)	93.5 (74.0-128)	182 (134-262)	2605
	05-06	24.8 (22.4-27.5)	21.4 (19.1-23.4)	46.1 (40.0-52.1)	117 (97.8-148)	235 (197-272)	2548
	07-08	22.2 (19.4-25.5)	19.3 (17.4-22.1)	40.5 (34.1-50.3)	99.3 (69.7-134)	179 (135-252)	2604
	09-10	13.5 (11.6-15.6)	11.9 (10.5-13.9)	22.3 (18.6-26.4)	44.0 (35.3-61.0)	86.3 (59.2-131)	2749
Age group							
6-11 years	01-02	38.2 (34.3-42.6)	34.3 (29.9-38.9)	60.6 (51.9-76.4)	107 (96.3-147)	211 (122-313)	393
	03-04	39.0 (31.1-48.9)	36.6 (25.3-49.3)	65.6 (49.8-91.3)	129 (77.1-253)	211 (123-708)	342
	05-06	38.5 (34.2-43.3)	37.0 (32.9-40.3)	65.5 (55.4-71.9)	115 (85.1-165)	213 (119-333)	356
	07-08	35.2 (29.1-42.5)	31.4 (25.6-37.8)	53.5 (44.8-72.3)	139 (79.9-230)	258 (141-303)	389
	09-10	19.6 (17.3-22.2)	18.6 (16.5-21.5)	34.2 (27.3-40.4)	55.4 (44.0-64.8)	72.1 (56.3-140)	415
12-19 years	01-02	19.2 (17.0-21.8)	17.8 (15.6-20.0)	34.9 (29.2-42.7)	73.4 (58.4-80.7)	102 (86.6-160)	742
	03-04	21.2 (18.1-24.7)	18.6 (16.9-21.7)	38.7 (29.7-53.4)	103 (62.7-209)	212 (100-358)	729
	05-06	26.0 (21.2-31.7)	23.7 (18.8-28.4)	49.4 (37.4-74.0)	131 (79.0-228)	278 (132-375)	702
	07-08	23.2 (18.1-29.6)	20.0 (14.6-23.9)	46.1 (31.5-66.3)	148 (66.6-234)	234 (146-373)	401
	09-10	12.3 (9.73-15.5)	10.5 (8.95-13.3)	20.9 (14.2-31.0)	45.3 (28.9-91.9)	110 (44.6-200)	420
20 years and older	01-02	17.1 (15.2-19.3)	15.0 (13.3-16.7)	27.7 (23.2-34.0)	63.7 (48.3-86.9)	137 (84.4-203)	1647
	03-04	18.8 (17.5-20.2)	16.3 (15.4-17.5)	31.6 (28.1-35.3)	83.8 (67.2-106)	171 (129-246)	1534
	05-06	23.4 (21.1-26.0)	19.4 (17.6-21.8)	42.5 (37.6-49.2)	115 (93.1-153)	235 (184-298)	1490
	07-08	21.0 (18.5-24.0)	18.5 (16.3-20.2)	38.1 (30.9-49.2)	94.3 (62.6-126)	164 (120-235)	1814
	09-10	13.1 (11.3-15.2)	11.5 (10.1-13.4)	20.9 (17.7-25.1)	40.6 (32.4-64.9)	86.3 (59.2-131)	1914
Gender							
Males	01-02	17.9 (16.2-19.7)	15.4 (13.8-17.9)	32.2 (27.8-36.8)	73.4 (55.3-91.8)	137 (97.7-224)	1371
	03-04	18.9 (17.1-20.9)	17.1 (15.2-18.6)	32.7 (26.6-41.6)	93.4 (68.8-123)	193 (108-291)	1250
	05-06	23.8 (21.0-26.9)	20.2 (17.5-23.3)	44.8 (38.5-54.1)	129 (92.5-166)	251 (202-352)	1270
	07-08	19.6 (16.8-22.7)	16.8 (14.6-20.0)	34.8 (28.7-41.8)	90.3 (65.2-119)	164 (111-258)	1294
	09-10	13.6 (11.7-15.9)	11.6 (10.5-13.1)	22.9 (17.8-27.4)	52.8 (39.1-74.3)	103 (73.3-173)	1399
Females	01-02	19.7 (17.3-22.4)	17.6 (15.4-19.5)	32.1 (26.8-38.6)	70.5 (57.8-93.7)	156 (93.7-201)	1411
	03-04	21.9 (19.7-24.5)	18.7 (16.8-20.9)	39.3 (33.8-46.9)	94.3 (72.8-136)	171 (146-261)	1355
	05-06	25.9 (23.2-28.8)	22.5 (19.6-25.5)	47.3 (40.5-52.3)	108 (88.8-131)	202 (157-278)	1278
	07-08	25.2 (22.1-28.6)	22.1 (19.6-24.4)	47.4 (38.5-56.1)	112 (84.6-150)	190 (162-268)	1310
	09-10	13.3 (11.4-15.6)	12.4 (10.4-14.6)	21.9 (18.3-26.7)	38.8 (30.9-52.2)	74.0 (46.9-115)	1350

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP) (creatinine corrected)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	01-02	17.4 (15.9-19.1)	15.7 (14.4-17.5)	30.6 (26.0-34.7)	65.9 (50.6-83.9)	103 (75.5-128)	677
	03-04	17.1 (14.3-20.4)	15.4 (13.2-17.7)	29.3 (23.8-36.8)	57.3 (45.7-97.6)	105 (70.1-195)	652
	05-06	20.7 (17.2-25.0)	17.5 (15.3-21.3)	37.3 (30.9-45.7)	99.9 (67.6-159)	181 (110-357)	637
	07-08	22.1 (17.7-27.7)	18.8 (15.2-23.4)	38.6 (28.3-54.9)	91.4 (61.0-170)	197 (141-282)	531
	09-10	15.2 (12.9-17.8)	14.6 (12.8-16.2)	27.0 (21.3-33.3)	60.0 (42.3-77.7)	94.8 (66.4-142)	566
Non-Hispanic blacks	01-02	20.9 (18.8-23.3)	19.7 (17.5-21.8)	38.3 (32.1-46.0)	93.5 (69.2-123)	164 (130-183)	703
	03-04	21.9 (20.1-23.8)	19.5 (17.3-22.6)	40.1 (35.8-45.3)	102 (75.5-122)	164 (133-269)	699
	05-06	24.5 (21.2-28.2)	18.9 (17.1-22.8)	46.1 (37.9-59.7)	128 (100-158)	308 (158-399)	678
	07-08	20.0 (18.1-22.0)	18.3 (16.5-19.2)	36.9 (27.9-46.7)	85.5 (65.8-103)	136 (107-227)	597
	09-10	11.2 (8.71-14.5)	10.5 (8.26-13.5)	18.9 (13.6-25.3)	32.9 (22.3-58.4)	47.2 (33.6-153)	516
Non-Hispanic whites	01-02	18.9 (17.0-21.0)	16.3 (14.8-18.4)	32.1 (27.3-37.3)	70.8 (56.9-93.7)	177 (98.0-242)	1216
	03-04	20.5 (18.5-22.8)	17.8 (16.2-19.7)	35.3 (29.7-44.9)	96.2 (75.8-136)	211 (136-283)	1088
	05-06	25.6 (23.1-28.2)	22.4 (19.6-24.5)	47.6 (40.7-54.1)	119 (98.2-148)	231 (181-297)	1038
	07-08	22.2 (18.7-26.3)	19.5 (16.7-23.0)	40.4 (32.1-54.8)	99.1 (68.1-145)	179 (129-258)	1077
	09-10	13.7 (11.7-16.1)	11.9 (10.4-13.9)	22.4 (18.0-27.6)	43.0 (34.2-61.0)	84.1 (54.7-142)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-oxohexyl) phthalate (MEOHP)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	13.5 (12.0-15.0)	14.0 (12.5-15.1)	29.6 (25.2-34.0)	59.9 (50.4-70.9)	120 (87.2-156)	2782
	03-04	14.5 (13.0-16.1)	14.4 (12.4-16.7)	31.4 (27.4-36.6)	76.7 (59.4-102)	157 (106-232)	2605
	05-06	16.2 (14.6-18.0)	15.1 (13.5-17.1)	35.5 (32.1-40.3)	94.8 (78.5-112)	183 (147-250)	2548
	07-08	12.2 (10.3-14.3)	11.3 (10.0-13.3)	27.1 (21.8-33.8)	64.2 (47.3-93.6)	130 (93.6-177)	2604
	09-10	8.02 (7.11-9.06)	8.02 (7.27-9.04)	16.1 (13.8-19.0)	32.5 (25.2-41.8)	55.7 (41.8-80.9)	2749
Age group							
6-11 years	01-02	23.3 (20.9-26.1)	22.9 (18.5-28.1)	46.5 (38.1-52.0)	81.6 (64.7-109)	142 (93.9-178)	393
	03-04	25.1 (19.6-32.3)	25.8 (19.3-31.4)	51.1 (32.1-76.5)	97.9 (58.8-197)	197 (97.6-261)	342
	05-06	23.0 (20.3-26.1)	24.5 (19.9-27.6)	44.3 (38.1-52.3)	83.8 (63.6-115)	126 (77.3-253)	356
	07-08	16.9 (13.9-20.6)	16.6 (12.4-22.6)	34.5 (26.2-40.9)	64.4 (46.9-129)	137 (63.8-179)	389
	09-10	9.78 (8.72-11.0)	11.1 (8.87-12.7)	20.0 (17.1-21.6)	35.4 (24.8-41.4)	48.4 (36.0-74.0)	415
12-19 years	01-02	17.5 (15.1-20.3)	18.6 (16.2-20.7)	35.0 (27.7-42.1)	70.7 (52.2-104)	118 (74.0-174)	742
	03-04	19.5 (16.0-23.7)	20.3 (18.4-23.5)	37.8 (32.6-44.6)	110 (54.6-168)	212 (103-326)	729
	05-06	23.0 (18.7-28.4)	22.1 (18.0-26.2)	50.7 (42.7-62.2)	134 (82.3-240)	263 (134-511)	702
	07-08	16.9 (12.8-22.3)	15.9 (11.9-19.3)	38.2 (24.0-55.5)	121 (58.1-258)	258 (120-354)	401
	09-10	10.0 (8.32-12.1)	9.82 (7.96-12.3)	19.0 (15.9-23.5)	40.0 (26.7-61.5)	68.4 (32.6-154)	420
20 years and older	01-02	12.0 (10.5-13.9)	12.3 (10.4-14.1)	26.0 (21.6-32.1)	52.3 (41.8-68.3)	116 (74.9-160)	1647
	03-04	12.9 (11.8-14.1)	12.4 (10.9-14.5)	27.0 (25.0-30.9)	68.9 (55.0-86.5)	139 (92.7-216)	1534
	05-06	14.7 (13.2-16.4)	13.4 (12.2-15.1)	31.9 (28.2-36.2)	91.6 (74.6-104)	182 (138-247)	1490
	07-08	11.1 (9.48-13.1)	10.7 (9.30-12.3)	25.5 (20.2-31.4)	59.8 (42.3-88.9)	108 (80.0-155)	1814
	09-10	7.59 (6.64-8.68)	7.55 (6.69-8.58)	15.3 (12.6-18.6)	30.6 (24.1-41.5)	54.9 (41.8-81.0)	1914
Gender							
Males	01-02	14.5 (13.0-16.2)	14.6 (13.1-16.2)	31.6 (25.6-34.7)	60.4 (52.3-71.4)	129 (84.4-167)	1371
	03-04	15.6 (13.6-17.9)	14.7 (12.7-18.1)	31.8 (27.2-39.5)	83.8 (59.4-134)	185 (96.2-277)	1250
	05-06	18.3 (16.0-20.9)	16.3 (14.7-19.7)	39.3 (33.4-48.1)	104 (80.7-140)	258 (180-337)	1270
	07-08	12.5 (10.5-15.0)	11.3 (9.80-13.4)	26.1 (21.8-32.2)	61.8 (46.8-98.1)	139 (83.7-189)	1294
	09-10	9.14 (8.01-10.4)	8.76 (7.87-9.88)	18.2 (14.8-21.0)	39.3 (27.5-50.9)	69.6 (50.0-109)	1399
Females	01-02	12.5 (10.8-14.6)	13.1 (11.2-15.0)	28.1 (23.7-33.5)	57.5 (45.8-72.7)	115 (81.8-147)	1411
	03-04	13.4 (11.9-15.1)	13.7 (11.4-16.4)	29.5 (26.1-36.6)	68.6 (53.7-88.1)	143 (88.2-210)	1355
	05-06	14.4 (12.6-16.5)	13.8 (12.5-15.7)	32.5 (29.3-36.4)	81.7 (68.6-104)	159 (114-182)	1278
	07-08	11.8 (10.0-14.0)	11.7 (9.80-13.5)	27.9 (21.3-37.5)	64.2 (43.9-93.6)	122 (92.0-191)	1310
	09-10	7.09 (6.17-8.14)	7.35 (6.13-8.67)	15.2 (12.1-17.5)	27.1 (21.6-36.9)	48.2 (31.3-67.1)	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.1, 0.5, 0.7, 0.6, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-oxohexyl) phthalate (MEOHP)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	01-02	13.1 (11.6-14.9)	13.4 (11.6-15.0)	25.5 (21.6-30.8)	56.6 (40.6-70.3)	77.3 (70.5-101)	677
	03-04	12.8 (10.5-15.5)	13.6 (11.4-15.6)	25.3 (20.4-29.9)	46.6 (32.3-70.8)	76.0 (51.6-153)	652
	05-06	14.8 (11.7-18.8)	12.8 (10.5-16.1)	30.9 (22.6-42.9)	79.1 (51.7-131)	152 (92.9-276)	637
	07-08	12.6 (10.5-15.1)	11.4 (10.4-12.9)	25.1 (18.7-37.2)	53.4 (43.2-91.7)	118 (88.4-147)	531
	09-10	9.57 (8.10-11.3)	9.65 (8.17-11.3)	19.2 (16.3-23.3)	39.0 (30.9-54.7)	64.6 (43.0-95.3)	566
Non-Hispanic blacks	01-02	19.6 (17.1-22.5)	20.1 (17.9-22.4)	39.0 (34.8-44.2)	80.5 (71.4-97.4)	153 (102-228)	703
	03-04	20.2 (17.7-23.0)	20.1 (17.0-22.5)	40.0 (33.9-46.9)	92.6 (68.8-130)	173 (104-247)	699
	05-06	21.8 (18.9-25.2)	18.8 (16.8-21.0)	46.0 (38.0-59.1)	130 (104-168)	243 (159-304)	678
	07-08	14.2 (12.7-15.9)	14.0 (12.7-16.3)	30.5 (25.8-35.9)	64.2 (52.1-76.1)	110 (71.9-136)	597
	09-10	9.57 (8.17-11.2)	9.64 (8.15-11.1)	19.8 (16.5-22.1)	31.4 (25.5-44.0)	50.9 (31.4-129)	516
Non-Hispanic whites	01-02	12.8 (11.2-14.6)	13.2 (11.6-14.6)	28.5 (23.6-34.0)	58.6 (48.8-70.9)	126 (83.7-172)	1216
	03-04	13.8 (12.4-15.4)	13.4 (11.3-16.3)	31.0 (27.0-36.3)	77.6 (59.4-102)	161 (98.7-241)	1088
	05-06	15.5 (13.9-17.3)	15.0 (13.3-16.5)	35.3 (30.1-40.8)	92.6 (74.6-111)	182 (134-247)	1038
	07-08	11.7 (9.66-14.2)	11.0 (9.20-13.6)	26.9 (20.2-35.1)	64.2 (43.4-108)	137 (90.0-197)	1077
	09-10	7.59 (6.60-8.73)	7.55 (6.81-8.63)	15.7 (12.9-18.6)	29.1 (22.9-46.5)	55.7 (41.5-83.5)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.1, 0.5, 0.7, 0.6, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-oxohexyl) phthalate (MEOHP) (creatinine corrected)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	12.6 (11.5-13.9)	11.2 (10.2-12.3)	21.3 (18.3-23.8)	45.2 (37.1-58.1)	87.0 (68.0-124)	2782
	03-04	13.6 (12.4-14.8)	12.1 (11.0-12.9)	24.3 (20.9-27.8)	63.0 (47.8-75.8)	118 (94.1-153)	2605
	05-06	15.8 (14.2-17.5)	13.5 (12.4-14.7)	28.9 (26.0-33.8)	77.7 (62.4-91.1)	144 (118-172)	2548
	07-08	12.3 (10.7-14.0)	11.0 (9.72-12.2)	22.3 (18.4-27.9)	52.9 (37.9-74.6)	107 (74.8-136)	2604
	09-10	8.37 (7.31-9.59)	7.44 (6.71-8.42)	13.5 (11.5-15.7)	25.9 (21.4-34.6)	47.7 (34.6-67.8)	2749
Age group							
6-11 years	01-02	26.6 (24.0-29.4)	22.8 (20.3-25.0)	43.3 (33.6-47.1)	74.7 (69.0-91.9)	131 (83.0-183)	393
	03-04	26.6 (21.4-33.0)	25.3 (17.8-32.4)	43.6 (34.2-63.2)	77.1 (63.0-118)	121 (76.3-435)	342
	05-06	25.4 (22.5-28.6)	24.4 (21.4-26.4)	42.7 (36.0-46.9)	70.4 (52.8-117)	136 (77.2-195)	356
	07-08	20.8 (17.1-25.4)	18.5 (15.6-21.5)	35.0 (26.4-46.0)	84.1 (47.2-137)	145 (84.8-187)	389
	09-10	12.7 (11.4-14.2)	12.6 (11.5-14.1)	20.8 (16.5-25.1)	33.3 (29.5-44.1)	45.5 (33.7-74.5)	415
12-19 years	01-02	13.5 (12.0-15.2)	12.0 (10.8-14.3)	23.4 (20.0-28.5)	48.4 (39.2-54.9)	70.5 (55.0-97.2)	742
	03-04	14.6 (12.6-16.9)	12.7 (11.6-14.4)	25.5 (20.7-33.8)	67.9 (42.3-143)	153 (61.8-209)	729
	05-06	17.2 (14.1-20.9)	15.3 (12.5-18.6)	32.5 (25.7-41.3)	84.2 (49.1-147)	163 (93.1-250)	702
	07-08	13.2 (10.3-16.7)	11.1 (9.08-13.5)	25.9 (18.9-35.6)	86.3 (39.4-128)	132 (84.3-203)	401
	09-10	8.06 (6.57-9.88)	7.05 (6.16-8.25)	13.3 (9.53-18.3)	24.8 (17.8-47.9)	55.2 (23.6-110)	420
20 years and older	01-02	11.4 (10.2-12.8)	10.1 (8.89-11.4)	17.5 (15.2-21.8)	38.4 (30.5-52.5)	84.3 (53.3-128)	1647
	03-04	12.4 (11.5-13.3)	11.0 (10.0-12.0)	20.9 (18.6-22.8)	53.9 (40.7-70.2)	109 (88.6-130)	1534
	05-06	14.8 (13.3-16.4)	12.4 (11.2-13.5)	26.7 (23.8-31.3)	76.3 (61.0-89.9)	144 (106-170)	1490
	07-08	11.4 (10.0-13.0)	10.0 (9.15-11.2)	20.8 (16.7-25.6)	47.3 (33.2-65.3)	87.4 (62.7-136)	1814
	09-10	8.04 (6.99-9.24)	7.19 (6.36-8.00)	12.7 (10.7-14.7)	24.3 (19.7-34.6)	47.4 (33.7-67.3)	1914
Gender							
Males	01-02	11.8 (10.7-13.0)	10.2 (8.93-11.7)	21.2 (18.5-23.3)	46.1 (35.3-58.7)	84.2 (69.6-104)	1371
	03-04	12.3 (11.1-13.5)	11.1 (10.0-12.0)	21.6 (17.6-26.9)	59.1 (45.4-72.0)	120 (72.0-162)	1250
	05-06	14.7 (12.9-16.7)	12.6 (10.9-14.3)	27.1 (23.3-35.6)	79.5 (59.6-96.9)	147 (120-190)	1270
	07-08	10.5 (9.08-12.2)	9.25 (8.08-10.6)	18.9 (15.9-23.1)	46.7 (32.6-62.4)	86.4 (56.5-144)	1294
	09-10	8.19 (7.15-9.38)	7.07 (6.59-7.55)	13.4 (10.8-15.8)	30.2 (21.7-41.2)	52.1 (41.2-82.3)	1399
Females	01-02	13.5 (11.9-15.2)	12.0 (10.8-13.7)	21.5 (18.0-25.6)	44.8 (36.8-61.6)	92.3 (61.0-139)	1411
	03-04	14.9 (13.4-16.7)	12.7 (11.4-14.2)	26.6 (21.8-30.6)	65.6 (48.0-90.1)	118 (97.0-157)	1355
	05-06	16.9 (15.1-19.0)	14.7 (13.2-16.5)	30.3 (26.6-34.7)	76.4 (57.6-97.2)	137 (106-170)	1278
	07-08	14.2 (12.5-16.0)	12.5 (11.5-13.8)	25.9 (21.2-31.0)	61.1 (40.9-84.1)	114 (84.3-142)	1310
	09-10	8.55 (7.37-9.93)	8.06 (6.76-9.33)	13.6 (11.6-16.2)	24.1 (20.0-32.7)	43.2 (27.7-64.5)	1350

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-oxohexyl) phthalate (MEOHP) (creatinine corrected)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	01-02	12.4 (11.4-13.5)	11.0 (10.5-12.3)	20.9 (18.5-24.4)	44.6 (33.4-56.2)	65.9 (53.1-83.1)	677
	03-04	11.5 (9.81-13.6)	10.7 (9.04-12.3)	18.8 (15.6-24.6)	39.1 (31.8-53.9)	63.0 (47.2-121)	652
	05-06	13.3 (11.1-16.1)	11.4 (9.58-13.6)	24.0 (19.4-30.4)	61.2 (45.6-85.9)	102 (69.9-200)	637
	07-08	12.3 (10.0-15.0)	10.6 (8.57-14.1)	21.7 (16.1-28.7)	53.1 (32.7-85.2)	113 (79.7-149)	531
	09-10	9.50 (8.16-11.1)	8.72 (8.13-9.61)	16.6 (12.9-20.5)	35.4 (24.4-46.6)	56.7 (40.9-82.3)	566
Non-Hispanic blacks	01-02	13.8 (12.3-15.4)	13.1 (12.0-14.2)	23.9 (20.0-29.3)	58.3 (45.3-79.7)	101 (81.3-124)	703
	03-04	14.3 (13.1-15.6)	13.3 (11.3-15.5)	24.8 (21.7-27.7)	61.2 (46.8-76.6)	105 (79.7-152)	699
	05-06	15.3 (13.2-17.8)	11.6 (9.87-14.6)	29.1 (24.7-38.5)	77.1 (61.9-113)	172 (99.8-251)	678
	07-08	11.0 (9.94-12.2)	9.90 (9.16-11.1)	20.6 (16.6-23.5)	42.2 (33.8-58.6)	75.8 (57.2-124)	597
	09-10	6.92 (5.60-8.57)	6.49 (5.12-8.24)	11.4 (9.09-14.1)	19.2 (14.9-27.3)	28.1 (19.8-61.1)	516
Non-Hispanic whites	01-02	12.7 (11.4-14.0)	11.1 (9.90-12.3)	20.8 (18.0-23.9)	45.7 (35.9-64.9)	96.0 (68.5-161)	1216
	03-04	13.7 (12.2-15.3)	12.0 (10.5-12.9)	24.9 (20.7-28.6)	69.5 (51.4-95.3)	124 (90.3-182)	1088
	05-06	16.3 (14.8-18.0)	14.0 (12.9-15.7)	30.8 (27.3-34.8)	79.3 (66.0-93.8)	139 (117-163)	1038
	07-08	12.2 (10.3-14.5)	11.0 (9.38-13.1)	22.3 (17.4-30.4)	51.4 (35.7-77.7)	107 (74.6-139)	1077
	09-10	8.53 (7.35-9.90)	7.62 (6.76-8.54)	13.7 (11.2-16.2)	24.9 (20.3-35.7)	47.8 (32.8-76.4)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-carboxypentyl) phthalate (MECPP)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	34.7 (31.0-38.9)	33.0 (29.1-37.4)	71.8 (61.7-84.8)	168 (133-240)	339 (235-506)	2605
	05-06	38.6 (34.7-42.9)	35.6 (31.1-40.3)	79.7 (70.9-92.9)	211 (180-246)	386 (311-484)	2548
	07-08	33.3 (28.7-38.6)	31.2 (27.1-36.0)	69.5 (55.8-86.3)	153 (120-216)	308 (220-397)	2604
	09-10	20.7 (18.5-23.3)	20.4 (18.1-23.5)	39.9 (34.9-45.5)	78.4 (67.2-95.5)	127 (99.5-199)	2749
Age group							
6-11 years	03-04	58.2 (44.7-75.6)	51.6 (39.2-67.6)	112 (71.4-182)	314 (124-524)	391 (238-781)	342
	05-06	57.4 (50.2-65.7)	53.5 (42.7-67.2)	94.7 (83.2-108)	200 (154-247)	297 (196-548)	356
	07-08	46.6 (39.2-55.3)	44.1 (34.9-54.7)	92.7 (64.7-112)	156 (117-278)	357 (176-441)	389
	09-10	27.7 (24.7-31.2)	29.4 (25.4-33.2)	48.5 (39.0-60.2)	87.1 (64.4-102)	118 (87.1-202)	415
12-19 years	03-04	44.6 (36.8-54.0)	42.7 (38.4-47.6)	86.5 (67.3-108)	220 (120-397)	448 (235-808)	729
	05-06	52.9 (43.0-65.2)	46.7 (39.4-61.6)	114 (85.6-173)	314 (195-515)	560 (324-1180)	702
	07-08	44.3 (35.2-55.9)	38.9 (28.9-49.3)	97.3 (64.3-127)	247 (140-456)	476 (231-977)	401
	09-10	26.2 (22.4-30.6)	25.7 (21.2-30.0)	50.1 (38.3-57.7)	90.3 (58.2-162)	147 (83.9-349)	420
20 years and older	03-04	31.3 (28.6-34.4)	29.2 (26.2-33.0)	63.5 (56.5-73.9)	157 (130-187)	312 (199-457)	1534
	05-06	35.1 (31.5-39.0)	31.3 (28.1-35.7)	72.7 (65.4-82.4)	193 (163-237)	377 (285-460)	1490
	07-08	30.7 (26.4-35.8)	29.2 (24.3-34.4)	63.2 (51.6-80.0)	145 (109-206)	286 (182-378)	1814
	09-10	19.4 (17.0-22.0)	18.8 (15.7-22.3)	37.7 (32.8-44.0)	76.9 (63.1-92.8)	126 (96.1-197)	1914
Gender							
Males	03-04	37.9 (33.1-43.5)	34.7 (30.0-39.5)	73.7 (60.8-91.9)	187 (133-300)	388 (222-660)	1250
	05-06	43.6 (38.1-49.8)	39.7 (31.9-46.2)	87.0 (76.8-103)	260 (188-308)	460 (347-670)	1270
	07-08	34.4 (29.3-40.5)	31.5 (27.3-35.7)	65.1 (52.1-87.4)	161 (120-217)	321 (213-422)	1294
	09-10	23.4 (20.7-26.4)	23.0 (20.2-25.6)	44.9 (40.1-51.0)	87.1 (70.3-110)	162 (107-288)	1399
Females	03-04	31.9 (28.1-36.2)	31.3 (27.5-35.8)	69.3 (58.9-81.9)	154 (128-199)	312 (182-441)	1355
	05-06	34.3 (30.4-38.6)	31.7 (28.0-36.6)	71.0 (63.0-85.3)	192 (156-217)	309 (251-386)	1278
	07-08	32.3 (27.8-37.5)	31.0 (26.4-38.3)	73.0 (56.9-90.2)	147 (112-220)	297 (185-420)	1310
	09-10	18.5 (16.2-21.0)	18.8 (15.8-22.4)	34.7 (29.3-41.5)	72.9 (56.3-93.2)	111 (90.3-146)	1350
Race/ethnicity							
Mexican Americans	03-04	31.9 (27.1-37.6)	31.5 (26.8-37.4)	57.4 (45.9-71.8)	116 (86.0-162)	175 (133-355)	652
	05-06	39.4 (30.3-51.2)	34.2 (24.1-47.1)	74.6 (54.0-111)	220 (124-338)	394 (222-673)	637
	07-08	36.7 (30.6-44.0)	31.3 (27.0-35.7)	71.9 (54.9-91.7)	162 (123-241)	321 (209-477)	531
	09-10	26.2 (22.5-30.4)	25.8 (22.3-29.8)	51.0 (41.3-57.4)	92.6 (74.5-122)	160 (113-245)	566
Non-Hispanic blacks	03-04	42.6 (37.0-49.2)	38.3 (33.8-46.9)	82.5 (68.7-103)	191 (146-246)	339 (244-468)	699
	05-06	46.6 (41.3-52.5)	40.3 (35.6-46.1)	96.3 (76.6-132)	256 (208-347)	455 (328-528)	678
	07-08	35.0 (31.1-39.4)	35.6 (30.7-38.8)	71.6 (59.1-84.5)	151 (113-192)	235 (184-338)	597
	09-10	21.9 (18.6-25.7)	22.3 (18.2-25.6)	40.5 (35.7-47.1)	77.0 (61.1-97.1)	127 (76.0-268)	516
Non-Hispanic whites	03-04	33.8 (30.1-37.9)	32.1 (27.6-37.5)	72.4 (62.0-87.7)	167 (133-240)	354 (220-560)	1088
	05-06	37.0 (33.4-41.0)	35.4 (30.4-40.9)	79.7 (70.0-93.3)	203 (174-237)	380 (284-484)	1038
	07-08	32.0 (26.7-38.3)	30.4 (24.5-38.0)	67.6 (52.5-90.9)	145 (105-244)	316 (197-476)	1077
	09-10	19.6 (17.2-22.2)	19.3 (16.4-23.1)	39.3 (33.0-45.3)	75.0 (61.9-95.7)	120 (95.5-197)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, 07-08 and 09-10 are 0.3, 0.6, 0.5, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(2-ethyl-5-carboxypentyl) phthalate (MECPP) (creatinine corrected)

Metabolite of Di-2-ethylhexyl phthalate (DEHP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	32.6 (29.6-36.0)	27.0 (24.3-30.6)	54.6 (48.0-63.5)	139 (109-186)	251 (192-356)	2605
	05-06	37.6 (33.7-42.0)	32.2 (29.5-37.0)	67.5 (58.2-80.8)	168 (139-209)	290 (261-328)	2548
	07-08	33.6 (29.7-38.0)	29.1 (25.8-32.2)	58.7 (49.6-68.8)	138 (109-164)	233 (178-319)	2604
	09-10	21.6 (19.0-24.7)	19.2 (16.8-22.1)	33.7 (30.0-40.0)	69.4 (54.0-84.3)	121 (85.1-184)	2749
Age group							
6-11 years	03-04	61.5 (49.0-77.2)	52.2 (41.6-73.8)	104 (74.2-140)	210 (111-500)	372 (192-988)	342
	05-06	63.2 (55.6-71.9)	54.2 (48.1-63.6)	92.8 (82.6-111)	160 (124-247)	312 (172-480)	356
	07-08	57.4 (49.2-66.9)	49.6 (42.7-61.5)	94.2 (79.5-122)	185 (138-294)	376 (188-404)	389
	09-10	36.1 (32.4-40.3)	33.6 (32.3-36.8)	55.4 (43.9-66.9)	88.7 (69.2-113)	121 (90.6-224)	415
12-19 years	03-04	33.4 (28.7-38.7)	27.1 (23.9-32.0)	55.0 (43.8-83.8)	168 (92.5-289)	294 (159-387)	729
	05-06	39.5 (32.5-47.8)	33.9 (27.9-41.0)	72.7 (56.9-96.7)	197 (106-327)	385 (216-531)	702
	07-08	34.5 (28.1-42.3)	28.0 (23.2-35.0)	65.9 (47.5-85.9)	159 (90.1-233)	246 (159-489)	401
	09-10	21.1 (17.4-25.5)	18.2 (15.6-20.7)	33.2 (25.5-43.3)	64.9 (42.7-158)	158 (58.4-246)	420
20 years and older	03-04	30.1 (27.7-32.7)	25.1 (22.9-27.6)	49.1 (44.1-55.2)	126 (101-154)	237 (191-315)	1534
	05-06	35.2 (31.2-39.6)	29.8 (26.1-33.3)	60.8 (54.2-74.0)	167 (131-206)	279 (247-322)	1490
	07-08	31.5 (27.8-35.8)	27.8 (24.1-30.8)	53.1 (45.0-65.4)	128 (101-155)	214 (162-319)	1814
	09-10	20.5 (17.9-23.5)	18.2 (15.9-20.5)	31.7 (27.4-36.8)	65.0 (52.1-80.6)	118 (81.3-173)	1914
Gender							
Males	03-04	29.8 (26.8-33.1)	23.5 (21.4-27.1)	50.7 (42.2-61.7)	132 (98.0-191)	248 (159-422)	1250
	05-06	35.0 (30.7-39.9)	29.0 (25.7-32.1)	69.3 (54.2-82.9)	172 (141-210)	301 (249-376)	1270
	07-08	29.0 (25.3-33.2)	25.1 (21.7-28.8)	47.6 (40.8-57.4)	120 (91.7-157)	210 (157-331)	1294
	09-10	21.0 (18.4-23.9)	18.4 (16.1-20.4)	33.1 (29.2-38.5)	70.2 (53.4-94.6)	125 (92.8-210)	1399
Females	03-04	35.5 (31.6-40.0)	30.6 (26.4-35.5)	58.3 (48.8-71.8)	144 (108-192)	251 (192-349)	1355
	05-06	40.3 (36.2-44.9)	36.7 (31.6-40.0)	66.1 (58.2-83.7)	168 (127-206)	279 (240-341)	1278
	07-08	38.7 (34.5-43.3)	32.8 (30.0-36.4)	67.8 (58.5-80.0)	147 (117-176)	266 (176-379)	1310
	09-10	22.3 (19.3-25.7)	20.5 (17.2-23.9)	34.9 (30.0-41.7)	67.0 (53.5-80.8)	104 (80.8-170)	1350
Race/ethnicity							
Mexican Americans	03-04	28.8 (25.4-32.6)	24.7 (22.4-26.3)	46.7 (39.0-56.3)	94.7 (73.2-137)	152 (118-238)	652
	05-06	35.5 (28.7-43.7)	29.8 (25.5-34.9)	61.5 (48.4-86.2)	165 (105-201)	278 (181-501)	637
	07-08	35.8 (29.1-44.0)	30.6 (23.2-38.6)	64.1 (47.2-84.8)	151 (92.3-240)	286 (198-402)	531
	09-10	26.0 (22.5-30.0)	23.8 (21.3-27.0)	40.1 (35.0-52.3)	88.8 (69.5-104)	148 (100-203)	566
Non-Hispanic blacks	03-04	30.3 (27.7-33.2)	27.0 (23.2-30.7)	51.1 (41.6-64.0)	135 (100-161)	212 (173-252)	699
	05-06	32.7 (28.9-37.1)	27.1 (22.4-31.5)	62.4 (53.1-76.0)	166 (119-260)	370 (231-429)	678
	07-08	27.2 (24.3-30.5)	23.5 (21.9-26.1)	49.1 (37.7-58.6)	103 (85.9-132)	178 (140-259)	597
	09-10	15.9 (12.6-19.9)	14.5 (11.5-18.8)	26.3 (19.5-32.9)	48.0 (32.6-66.4)	70.7 (46.1-209)	516
Non-Hispanic whites	03-04	33.4 (29.5-37.7)	27.0 (23.5-31.6)	56.8 (48.6-69.4)	145 (109-198)	294 (193-385)	1088
	05-06	39.0 (34.9-43.4)	34.3 (30.3-39.1)	69.5 (59.5-83.7)	182 (143-214)	284 (237-324)	1038
	07-08	33.4 (28.4-39.3)	29.2 (25.1-34.6)	57.7 (46.6-72.4)	138 (101-166)	233 (164-338)	1077
	09-10	22.0 (19.1-25.3)	19.4 (16.8-22.9)	34.0 (29.3-42.5)	67.6 (52.5-84.3)	113 (81.3-188)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DEHP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(carboxynonyl) phthalate (MCNP)

Metabolite of Di-isodecyl phthalate (DiDP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	05-06	2.73 (2.50-2.98)	2.70 (2.40-3.00)	5.30 (4.80-5.90)	10.2 (8.80-11.9)	17.5 (14.0-21.4)	2548
	07-08	2.42 (2.17-2.71)	2.40 (2.20-2.70)	4.70 (4.20-5.30)	9.20 (8.00-10.6)	16.1 (12.3-21.0)	2604
	09-10	2.88 (2.58-3.22)	2.91 (2.51-3.26)	5.52 (4.83-6.30)	10.6 (9.14-12.6)	16.8 (13.9-22.1)	2749
Age group							
6-11 years	05-06	4.54 (3.94-5.24)	4.70 (4.00-5.60)	8.10 (6.80-9.60)	14.6 (10.8-19.1)	22.8 (16.7-35.3)	356
	07-08	3.15 (2.65-3.74)	3.40 (2.90-3.80)	5.40 (4.60-6.80)	9.10 (7.40-12.3)	14.9 (9.10-24.8)	389
	09-10	3.45 (2.95-4.03)	3.59 (3.04-4.34)	6.39 (5.39-7.41)	9.65 (8.75-11.2)	14.5 (10.4-23.2)	415
12-19 years	05-06	3.18 (2.74-3.68)	3.30 (2.90-3.80)	6.00 (4.70-7.20)	10.3 (7.60-13.7)	16.5 (11.5-21.5)	702
	07-08	2.86 (2.59-3.16)	2.70 (2.40-2.90)	5.10 (4.30-5.70)	10.3 (8.90-13.0)	16.8 (10.7-32.4)	401
	09-10	3.02 (2.61-3.51)	3.05 (2.41-3.65)	5.11 (4.49-6.43)	11.0 (7.24-14.6)	16.9 (10.9-32.0)	420
20 years and older	05-06	2.51 (2.27-2.78)	2.40 (2.20-2.70)	4.90 (4.40-5.40)	9.50 (7.90-11.9)	17.0 (13.1-21.5)	1490
	07-08	2.29 (2.04-2.58)	2.30 (2.00-2.50)	4.50 (4.00-5.10)	9.10 (7.60-10.7)	16.1 (12.2-21.4)	1814
	09-10	2.81 (2.49-3.16)	2.80 (2.44-3.09)	5.52 (4.69-6.45)	10.7 (8.95-13.5)	17.3 (13.5-23.7)	1914
Gender							
Males	05-06	3.16 (2.82-3.54)	3.00 (2.70-3.50)	5.90 (5.10-6.90)	11.0 (8.80-14.0)	19.5 (14.0-28.0)	1270
	07-08	2.75 (2.50-3.03)	2.60 (2.40-2.90)	5.10 (4.50-5.70)	9.50 (8.00-12.2)	19.4 (12.4-29.1)	1294
	09-10	3.14 (2.80-3.53)	3.11 (2.72-3.61)	5.75 (5.13-6.56)	10.4 (9.14-11.5)	15.1 (12.5-20.5)	1399
Females	05-06	2.37 (2.13-2.65)	2.30 (2.00-2.70)	4.80 (4.30-5.30)	9.30 (7.60-11.5)	14.7 (13.1-18.2)	1278
	07-08	2.14 (1.84-2.49)	2.20 (1.80-2.60)	4.40 (3.80-4.90)	9.00 (7.30-10.3)	14.1 (10.8-18.5)	1310
	09-10	2.66 (2.32-3.04)	2.60 (2.21-2.98)	5.21 (4.43-6.17)	11.6 (8.88-14.8)	17.5 (14.6-23.4)	1350
Race/ethnicity							
Mexican Americans	05-06	2.72 (2.40-3.08)	2.70 (2.40-3.20)	4.90 (4.30-5.70)	9.00 (7.00-12.4)	14.4 (9.00-26.8)	637
	07-08	2.37 (2.15-2.62)	2.30 (2.10-2.60)	4.30 (3.70-5.10)	8.40 (6.50-10.5)	12.3 (8.40-16.2)	531
	09-10	2.35 (2.15-2.57)	2.51 (2.18-2.69)	4.35 (3.83-4.83)	7.72 (6.56-9.53)	11.0 (8.09-14.1)	566
Non-Hispanic blacks	05-06	3.18 (2.75-3.67)	3.20 (2.70-3.80)	5.90 (4.90-7.30)	13.0 (9.50-14.6)	19.2 (14.6-30.5)	678
	07-08	2.89 (2.64-3.17)	2.90 (2.50-3.10)	5.60 (4.80-6.20)	11.9 (9.10-15.4)	21.7 (17.9-28.9)	597
	09-10	3.51 (2.92-4.22)	3.64 (2.97-4.24)	6.94 (5.75-8.47)	13.2 (10.2-17.5)	20.8 (13.5-41.7)	516
Non-Hispanic whites	05-06	2.67 (2.39-2.98)	2.60 (2.30-2.90)	5.30 (4.70-6.10)	10.1 (8.20-12.6)	17.6 (12.6-24.3)	1038
	07-08	2.42 (2.09-2.81)	2.40 (2.10-2.70)	4.70 (4.10-5.60)	9.10 (7.60-11.6)	14.9 (10.5-29.1)	1077
	09-10	2.98 (2.61-3.40)	2.96 (2.53-3.48)	5.59 (4.78-6.57)	10.7 (8.83-14.3)	16.6 (12.6-23.7)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06, 07-08, and 09-10 are 0.6, 0.5, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DiDP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(carboxynonyl) phthalate (MCNP) (creatinine corrected)

Metabolite of Di-isodecyl phthalate (DiDP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	05-06	2.66 (2.43-2.91)	2.48 (2.29-2.71)	4.48 (4.05-4.85)	8.67 (7.17-9.71)	13.2 (10.8-17.5)	2548
	07-08	2.44 (2.25-2.65)	2.21 (2.00-2.36)	4.00 (3.68-4.44)	8.31 (6.96-9.04)	12.8 (11.2-15.1)	2604
	09-10	3.01 (2.74-3.31)	2.79 (2.48-3.10)	5.04 (4.47-5.66)	9.76 (8.41-11.2)	15.5 (12.8-18.9)	2749
Age group							
6-11 years	05-06	5.00 (4.45-5.63)	5.06 (4.18-5.76)	7.48 (6.70-8.84)	16.1 (12.5-18.9)	23.5 (16.1-31.7)	356
	07-08	3.88 (3.34-4.50)	3.82 (3.17-4.18)	6.07 (5.14-7.00)	11.2 (9.01-13.6)	18.2 (12.6-27.0)	389
	09-10	4.49 (4.01-5.03)	4.33 (3.95-4.82)	7.30 (6.41-8.54)	11.2 (9.74-13.6)	14.9 (12.7-23.1)	415
12-19 years	05-06	2.37 (2.04-2.75)	2.18 (1.91-2.52)	3.64 (2.94-4.55)	6.26 (5.15-8.70)	10.7 (7.57-14.1)	702
	07-08	2.23 (2.07-2.40)	1.96 (1.83-2.08)	3.68 (3.08-3.83)	6.39 (4.88-9.01)	10.0 (7.26-13.5)	401
	09-10	2.43 (2.10-2.81)	2.17 (1.79-2.54)	3.94 (3.23-4.87)	7.67 (5.32-11.2)	12.8 (8.43-14.4)	420
20 years and older	05-06	2.52 (2.29-2.78)	2.37 (2.14-2.57)	4.05 (3.77-4.51)	7.90 (6.46-9.40)	12.0 (9.71-16.7)	1490
	07-08	2.36 (2.15-2.58)	2.14 (1.91-2.32)	3.88 (3.44-4.29)	8.21 (6.67-9.00)	12.3 (11.0-15.3)	1814
	09-10	2.97 (2.69-3.29)	2.72 (2.43-3.03)	4.93 (4.32-5.62)	9.68 (7.90-11.6)	15.6 (11.9-22.5)	1914
Gender							
Males	05-06	2.53 (2.28-2.82)	2.36 (2.14-2.56)	4.38 (3.74-5.04)	7.41 (6.47-8.88)	13.0 (9.40-19.6)	1270
	07-08	2.32 (2.11-2.55)	2.12 (1.90-2.32)	3.88 (3.50-4.29)	7.13 (5.78-9.15)	11.4 (9.68-15.5)	1294
	09-10	2.82 (2.55-3.12)	2.56 (2.28-2.96)	4.73 (4.19-5.49)	8.68 (7.21-10.2)	13.4 (11.2-15.4)	1399
Females	05-06	2.79 (2.47-3.15)	2.65 (2.34-2.95)	4.51 (4.03-5.15)	9.41 (6.82-11.6)	13.5 (10.7-19.7)	1278
	07-08	2.57 (2.33-2.83)	2.27 (2.03-2.57)	4.13 (3.78-4.82)	8.71 (7.41-9.50)	13.5 (12.0-16.0)	1310
	09-10	3.20 (2.88-3.57)	2.98 (2.72-3.28)	5.40 (4.70-6.23)	10.6 (8.39-13.8)	16.9 (12.5-27.3)	1350
Race/ethnicity							
Mexican Americans	05-06	2.45 (2.13-2.81)	2.32 (2.14-2.60)	4.04 (3.64-4.58)	6.96 (5.73-8.14)	11.0 (7.89-19.1)	637
	07-08	2.32 (2.10-2.56)	2.10 (1.90-2.34)	4.12 (3.54-4.68)	6.89 (5.73-7.67)	9.75 (7.69-11.7)	531
	09-10	2.33 (2.08-2.62)	2.12 (1.80-2.66)	3.80 (3.27-4.58)	7.02 (5.44-9.66)	11.3 (8.09-13.0)	566
Non-Hispanic blacks	05-06	2.24 (1.96-2.55)	1.97 (1.78-2.37)	3.60 (3.25-4.24)	7.82 (6.38-10.4)	12.1 (10.1-17.4)	678
	07-08	2.25 (2.05-2.47)	2.00 (1.88-2.17)	3.88 (3.50-4.27)	8.18 (6.25-10.4)	14.9 (9.31-20.4)	597
	09-10	2.54 (2.24-2.88)	2.46 (2.02-2.96)	4.52 (4.20-5.13)	8.51 (6.35-10.9)	11.3 (9.81-16.0)	516
Non-Hispanic whites	05-06	2.81 (2.51-3.14)	2.57 (2.33-2.85)	4.62 (4.10-5.19)	9.26 (7.00-10.8)	13.5 (10.3-23.5)	1038
	07-08	2.53 (2.23-2.86)	2.26 (1.98-2.57)	4.11 (3.66-4.70)	8.51 (6.59-10.2)	13.3 (10.7-17.6)	1077
	09-10	3.35 (2.98-3.77)	3.06 (2.76-3.36)	5.51 (4.74-6.35)	10.4 (8.43-13.5)	15.8 (12.3-24.8)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DiDP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-isononyl phthalate (MiNP)

Metabolite of Di-isononyl phthalate (DiNP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	5.39 (<LOD-21.3)	2541
	01-02	*	< LOD	< LOD	< LOD	< LOD	2782
	03-04	*	< LOD	< LOD	< LOD	1.54 (<LOD-2.00)	2605
	05-06	*	< LOD	< LOD	1.69 (<LOD-2.00)	3.54 (2.31-6.16)	2548
	07-08	*	< LOD	< LOD	1.54 (<LOD-1.69)	3.08 (2.16-4.16)	2604
	09-10	*	< LOD	1.80 (1.49-2.13)	5.62 (4.48-7.28)	14.6 (10.7-18.3)	2749
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	9.55 (<LOD-34.7)	328
	01-02	*	< LOD	< LOD	< LOD	< LOD	393
	03-04	*	< LOD	< LOD	1.54 (<LOD-2.31)	2.62 (1.85-4.77)	342
	05-06	*	< LOD	< LOD	1.69 (<LOD-2.00)	2.16 (2.00-2.62)	356
	07-08	*	< LOD	< LOD	1.39 (<LOD-2.00)	2.16 (1.54-3.85)	389
	09-10	*	< LOD	1.72 (1.09-2.94)	5.30 (3.08-8.98)	13.1 (5.33-21.5)	415
12-19 years	99-00	*	< LOD	< LOD	< LOD	3.54 (<LOD-31.3)	752
	01-02	*	< LOD	< LOD	< LOD	1.39 (<LOD-1.69)	742
	03-04	*	< LOD	< LOD	< LOD	2.16 (<LOD-2.77)	729
	05-06	*	< LOD	< LOD	2.16 (<LOD-4.31)	5.39 (2.00-12.2)	702
	07-08	*	< LOD	< LOD	2.46 (1.39-5.39)	5.54 (2.46-8.01)	401
	09-10	*	< LOD	2.20 (1.42-3.96)	10.7 (4.87-17.1)	20.6 (14.8-36.1)	420
20 years and older	99-00	*	< LOD	< LOD	< LOD	4.77 (<LOD-20.3)	1461
	01-02	*	< LOD	< LOD	< LOD	< LOD	1647
	03-04	*	< LOD	< LOD	< LOD	< LOD	1534
	05-06	*	< LOD	< LOD	1.54 (<LOD-2.00)	3.54 (2.16-6.47)	1490
	07-08	*	< LOD	< LOD	1.39 (<LOD-1.69)	2.77 (1.85-3.85)	1814
	09-10	*	< LOD	1.76 (1.43-2.11)	5.44 (4.31-6.64)	12.9 (8.32-17.9)	1914
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	7.55 (<LOD-29.1)	1215
	01-02	*	< LOD	< LOD	< LOD	< LOD	1371
	03-04	*	< LOD	< LOD	< LOD	1.54 (<LOD-2.77)	1250
	05-06	*	< LOD	< LOD	2.00 (1.39-2.93)	4.00 (3.23-6.31)	1270
	07-08	*	< LOD	< LOD	1.54 (<LOD-2.16)	3.23 (2.31-4.93)	1294
	09-10	*	< LOD	2.06 (1.74-2.68)	7.19 (5.62-10.5)	17.1 (11.7-27.6)	1399
Females	99-00	*	< LOD	< LOD	< LOD	3.85 (<LOD-10.5)	1326
	01-02	*	< LOD	< LOD	< LOD	1.39 (<LOD-1.69)	1411
	03-04	*	< LOD	< LOD	< LOD	< LOD	1355
	05-06	*	< LOD	< LOD	1.39 (<LOD-1.54)	2.62 (1.85-4.31)	1278
	07-08	*	< LOD	< LOD	1.39 (<LOD-1.69)	2.77 (2.00-3.70)	1310
	09-10	*	< LOD	1.52 (1.12-1.91)	4.70 (3.80-5.47)	12.3 (6.61-16.5)	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.23, 1.23, 1.54, 1.23, 1.23, and 0.77 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 1.54 for survey periods 1999-2008 compared with results previously reported.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DiNP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-isononyl phthalate (MiNP)

Metabolite of Di-isononyl phthalate (DiNP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

Race/ethnicity	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	2.31 (<LOD-4.31)	814
	01-02	*	< LOD	< LOD	< LOD	1.54 (<LOD-2.16)	677
	03-04	*	< LOD	< LOD	< LOD	< LOD	652
	05-06	*	< LOD	< LOD	1.54 (<LOD-2.00)	4.31 (2.00-8.62)	637
	07-08	*	< LOD	< LOD	1.39 (<LOD-2.93)	3.08 (2.00-3.85)	531
	09-10	*	< LOD	1.52 (1.19-1.91)	5.11 (3.77-7.67)	12.9 (6.27-22.2)	566
	Non-Hispanic blacks	99-00	*	< LOD	< LOD	3.54 (<LOD-21.3)	10.5 (<LOD-46.5)
01-02		*	< LOD	< LOD	< LOD	1.54 (<LOD-2.62)	703
03-04		*	< LOD	< LOD	< LOD	2.00 (1.54-3.08)	699
05-06		*	< LOD	< LOD	2.00 (1.54-2.77)	5.70 (3.54-8.62)	678
07-08		*	< LOD	< LOD	1.69 (<LOD-2.31)	3.70 (2.16-5.70)	597
09-10		*	< LOD	2.14 (1.79-2.56)	6.05 (4.48-10.5)	17.1 (8.32-21.9)	516
Non-Hispanic whites		99-00	*	< LOD	< LOD	< LOD	5.39 (<LOD-24.6)
	01-02	*	< LOD	< LOD	< LOD	< LOD	1216
	03-04	*	< LOD	< LOD	< LOD	< LOD	1088
	05-06	*	< LOD	< LOD	1.54 (<LOD-2.16)	3.23 (1.85-6.47)	1038
	07-08	*	< LOD	< LOD	1.39 (<LOD-2.00)	2.77 (1.69-4.93)	1077
	09-10	*	< LOD	1.83 (1.40-2.53)	5.67 (4.59-8.07)	14.7 (10.3-21.5)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.23, 1.23, 1.54, 1.23, 1.23, and 0.77 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 1.54 for survey periods 1999-2008 compared with results previously reported

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DiNP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-isononyl phthalate (MiNP) (creatinine corrected)

Metabolite of Di-isononyl phthalate (DiNP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	6.61 (<LOD-12.8)	2541
	01-02	*	< LOD	< LOD	< LOD	< LOD	2782
	03-04	*	< LOD	< LOD	< LOD	4.50 (<LOD-4.90)	2605
	05-06	*	< LOD	< LOD	3.20 (<LOD-3.82)	5.16 (3.99-5.67)	2548
	07-08	*	< LOD	< LOD	3.35 (<LOD-3.96)	5.05 (4.15-6.22)	2604
	09-10	*	< LOD	2.16 (1.91-2.41)	5.40 (4.21-7.00)	11.8 (9.59-14.2)	2749
Age group							
6-11 years	99-00	*	< LOD	< LOD	< LOD	9.24 (<LOD-21.9)	328
	01-02	*	< LOD	< LOD	< LOD	< LOD	393
	03-04	*	< LOD	< LOD	3.71 (<LOD-4.82)	5.04 (3.71-7.19)	342
	05-06	*	< LOD	< LOD	2.37 (<LOD-2.93)	3.03 (2.54-5.16)	356
	07-08	*	< LOD	< LOD	3.35 (<LOD-5.33)	5.33 (3.11-7.26)	389
	09-10	*	< LOD	2.53 (1.93-3.09)	6.61 (3.77-9.73)	10.7 (6.70-21.7)	415
12-19 years	99-00	*	< LOD	< LOD	< LOD	3.08 (<LOD-11.8)	752
	01-02	*	< LOD	< LOD	< LOD	3.19 (<LOD-5.13)	742
	03-04	*	< LOD	< LOD	< LOD	3.03 (<LOD-3.99)	729
	05-06	*	< LOD	< LOD	2.66 (<LOD-3.59)	3.99 (2.97-5.28)	702
	07-08	*	< LOD	< LOD	2.56 (2.10-3.23)	3.68 (2.90-6.84)	401
	09-10	*	< LOD	2.14 (1.40-3.40)	7.58 (4.06-11.2)	13.2 (10.2-23.4)	420
20 years and older	99-00	*	< LOD	< LOD	< LOD	7.11 (<LOD-12.4)	1461
	01-02	*	< LOD	< LOD	< LOD	< LOD	1647
	03-04	*	< LOD	< LOD	< LOD	< LOD	1534
	05-06	*	< LOD	< LOD	3.37 (<LOD-4.19)	5.17 (4.17-6.11)	1490
	07-08	*	< LOD	< LOD	3.48 (<LOD-4.15)	5.45 (4.15-6.39)	1814
	09-10	*	< LOD	2.12 (1.86-2.35)	5.00 (3.91-6.67)	11.7 (8.29-14.5)	1914
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	6.53 (<LOD-15.7)	1215
	01-02	*	< LOD	< LOD	< LOD	< LOD	1371
	03-04	*	< LOD	< LOD	< LOD	3.56 (<LOD-4.54)	1250
	05-06	*	< LOD	< LOD	2.51 (1.96-3.51)	3.99 (2.79-5.61)	1270
	07-08	*	< LOD	< LOD	2.42 (<LOD-2.95)	3.96 (3.23-5.05)	1294
	09-10	*	< LOD	1.91 (1.59-2.37)	5.54 (4.14-7.89)	12.9 (9.90-18.8)	1399
Females	99-00	*	< LOD	< LOD	< LOD	6.61 (<LOD-11.8)	1326
	01-02	*	< LOD	< LOD	< LOD	4.20 (<LOD-4.40)	1411
	03-04	*	< LOD	< LOD	< LOD	< LOD	1355
	05-06	*	< LOD	< LOD	3.67 (<LOD-4.39)	5.37 (4.88-6.27)	1278
	07-08	*	< LOD	< LOD	4.08 (<LOD-4.84)	5.81 (4.36-7.26)	1310
	09-10	*	< LOD	2.35 (2.01-2.49)	5.40 (4.13-7.00)	10.8 (7.58-14.9)	1350

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 1.54 for survey periods 1999-2008 compared with results previously reported

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DiNP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-isononyl phthalate (MiNP) (creatinine corrected)

Metabolite of Di-isononyl phthalate (DiNP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	5.44 (<LOD-7.70)	814
	01-02	*	< LOD	< LOD	< LOD	3.70 (<LOD-4.20)	677
	03-04	*	< LOD	< LOD	< LOD	< LOD	652
	05-06	*	< LOD	< LOD	2.71 (<LOD-3.73)	4.88 (3.23-6.27)	637
	07-08	*	< LOD	< LOD	2.81 (<LOD-3.23)	3.96 (3.23-5.13)	531
	09-10	*	< LOD	1.74 (1.42-2.35)	4.35 (3.10-9.92)	11.5 (8.42-19.1)	566
	Non-Hispanic blacks	99-00	*	< LOD	< LOD	3.13 (<LOD-8.18)	6.61 (<LOD-22.1)
01-02		*	< LOD	< LOD	< LOD	2.71 (<LOD-3.30)	703
03-04		*	< LOD	< LOD	< LOD	3.37 (2.57-3.85)	699
05-06		*	< LOD	< LOD	2.22 (1.83-2.59)	4.39 (2.83-6.75)	678
07-08		*	< LOD	< LOD	2.49 (<LOD-3.35)	4.19 (3.11-4.62)	597
09-10		*	< LOD	1.51 (1.25-1.80)	4.00 (2.90-5.61)	8.97 (5.15-15.1)	516
Non-Hispanic whites		99-00	*	< LOD	< LOD	< LOD	8.39 (<LOD-15.4)
	01-02	*	< LOD	< LOD	< LOD	< LOD	1216
	03-04	*	< LOD	< LOD	< LOD	< LOD	1088
	05-06	*	< LOD	< LOD	3.37 (<LOD-3.99)	4.88 (3.99-5.41)	1038
	07-08	*	< LOD	< LOD	3.60 (<LOD-4.15)	5.45 (3.96-6.96)	1077
	09-10	*	< LOD	2.41 (2.08-2.70)	6.00 (4.23-8.05)	12.9 (10.4-16.6)	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 1.54 for survey periods 1999-2008 compared with results previously reported

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DiNP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(carboxyoctyl) phthalate (MCOP)

Metabolite of Di-isononyl phthalate (DiNP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	05-06	5.39 (4.68-6.22)	5.10 (4.40-6.00)	10.9 (9.10-13.1)	25.5 (19.1-35.4)	54.4 (32.5-85.2)	2548
	07-08	6.79 (5.83-7.92)	6.40 (5.60-7.40)	14.9 (12.1-18.2)	34.6 (25.2-43.8)	63.0 (41.8-80.0)	2604
	09-10	12.6 (10.4-15.2)	11.8 (9.26-14.7)	33.1 (25.8-40.2)	83.3 (61.2-113)	160 (117-191)	2749
Age group							
6-11 years	05-06	8.52 (7.19-10.1)	8.90 (7.60-9.90)	15.0 (11.3-19.3)	26.4 (19.9-37.3)	40.3 (26.4-61.6)	356
	07-08	10.8 (9.25-12.6)	10.7 (9.10-13.1)	19.8 (15.8-24.5)	33.8 (26.1-50.3)	59.9 (33.5-79.5)	389
	09-10	15.0 (10.8-20.7)	14.2 (9.15-20.7)	34.7 (18.6-56.5)	74.6 (41.6-135)	120 (74.6-340)	415
12-19 years	05-06	6.61 (5.42-8.06)	6.30 (5.20-7.30)	11.9 (9.60-16.0)	28.4 (18.2-58.7)	63.2 (28.4-106)	702
	07-08	9.25 (7.33-11.7)	8.10 (6.80-10.3)	20.8 (15.4-28.7)	60.5 (29.6-79.4)	79.4 (49.7-166)	401
	09-10	16.1 (12.3-20.9)	14.5 (11.6-17.0)	39.8 (22.5-61.2)	96.8 (66.7-191)	209 (135-263)	420
20 years and older	05-06	4.96 (4.27-5.76)	4.50 (3.90-5.20)	9.70 (8.20-12.3)	25.1 (18.2-35.7)	54.4 (30.8-88.5)	1490
	07-08	6.15 (5.20-7.29)	5.80 (5.00-6.60)	13.2 (10.5-16.8)	32.0 (23.0-41.8)	61.2 (38.9-80.0)	1814
	09-10	11.9 (9.89-14.3)	11.0 (8.58-13.7)	32.1 (25.8-39.0)	83.7 (56.0-114)	158 (109-188)	1914
Gender							
Males	05-06	6.24 (5.25-7.42)	5.70 (4.70-6.90)	12.1 (10.2-15.1)	30.8 (19.0-58.8)	66.4 (33.8-122)	1270
	07-08	7.14 (6.10-8.36)	6.60 (5.70-7.70)	15.6 (12.4-19.3)	35.7 (24.7-43.2)	61.2 (40.1-89.1)	1294
	09-10	14.0 (11.6-16.8)	13.4 (10.5-15.6)	36.0 (27.5-45.9)	92.9 (66.1-134)	168 (111-217)	1399
Females	05-06	4.69 (4.14-5.30)	4.50 (4.00-5.20)	9.30 (8.20-11.1)	21.5 (17.5-27.0)	41.7 (28.1-61.7)	1278
	07-08	6.48 (5.49-7.65)	6.20 (5.40-7.40)	14.3 (12.0-17.0)	32.6 (24.8-47.0)	64.8 (47.0-73.3)	1310
	09-10	11.4 (9.17-14.1)	10.6 (8.05-13.7)	29.6 (20.8-38.4)	70.4 (50.1-113)	146 (85.2-188)	1350
Race/ethnicity							
Mexican Americans	05-06	5.77 (4.84-6.87)	5.20 (4.50-5.70)	9.90 (8.50-12.0)	23.6 (14.9-49.7)	54.4 (23.0-185)	637
	07-08	6.83 (6.23-7.49)	6.20 (5.60-6.80)	12.9 (11.3-14.9)	29.1 (25.2-33.8)	52.0 (36.8-61.2)	531
	09-10	9.96 (7.99-12.4)	8.59 (6.21-12.1)	20.3 (15.8-32.6)	59.3 (44.0-69.5)	86.2 (63.2-136)	566
Non-Hispanic blacks	05-06	6.08 (5.60-6.61)	6.00 (5.10-6.70)	11.4 (10.4-13.0)	24.5 (20.3-28.1)	50.0 (32.0-70.3)	678
	07-08	7.55 (6.21-9.17)	7.00 (6.00-8.00)	16.6 (12.0-22.9)	40.3 (25.5-51.9)	60.4 (41.8-97.8)	597
	09-10	12.6 (9.85-16.1)	11.0 (7.35-16.2)	34.5 (26.5-42.1)	82.4 (63.8-109)	137 (83.3-188)	516
Non-Hispanic whites	05-06	5.25 (4.40-6.27)	5.00 (4.10-6.20)	11.1 (8.60-13.6)	25.5 (17.7-43.5)	56.1 (28.1-88.5)	1038
	07-08	6.81 (5.48-8.46)	6.40 (5.20-7.90)	15.1 (11.0-20.8)	35.7 (23.2-62.0)	64.7 (39.4-89.0)	1077
	09-10	13.4 (10.7-16.8)	12.9 (10.1-16.1)	34.8 (27.5-43.4)	88.9 (64.4-136)	168 (113-220)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06, 07-08, and 09-10 are 0.7, 0.7, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DiNP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(carboxyoctyl) phthalate (MCOP) (creatinine corrected)

Metabolite of Di-isononyl phthalate (DiNP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	05-06	5.26 (4.54-6.10)	4.53 (3.95-5.11)	9.25 (7.64-11.2)	24.0 (17.1-30.8)	40.2 (30.3-55.2)	2548
	07-08	6.85 (6.05-7.75)	5.94 (5.19-6.67)	12.6 (10.8-14.4)	30.6 (23.4-38.5)	50.2 (40.0-60.0)	2604
	09-10	13.1 (11.1-15.5)	11.6 (9.69-13.8)	29.3 (23.7-36.2)	72.3 (55.0-93.8)	119 (89.1-147)	2749
Age group							
6-11 years	05-06	9.38 (8.08-10.9)	8.48 (7.66-10.1)	14.8 (12.2-18.7)	30.3 (20.0-38.8)	40.1 (28.2-57.3)	356
	07-08	13.3 (11.4-15.5)	11.6 (9.90-14.2)	23.1 (16.2-32.5)	44.2 (27.0-64.1)	60.3 (44.2-110)	389
	09-10	19.5 (14.7-25.9)	18.6 (11.7-26.9)	35.9 (25.7-56.9)	87.8 (43.7-132)	139 (80.1-312)	415
12-19 years	05-06	4.93 (4.10-5.93)	4.32 (3.61-4.89)	7.44 (5.93-11.0)	19.4 (12.4-31.8)	34.7 (24.1-58.3)	702
	07-08	7.20 (5.94-8.73)	6.28 (4.79-8.08)	12.6 (10.0-20.7)	32.7 (22.6-56.9)	56.9 (31.4-103)	401
	09-10	12.9 (10.5-15.8)	11.5 (8.38-15.1)	26.3 (19.2-37.9)	74.8 (43.5-120)	125 (101-167)	420
20 years and older	05-06	4.97 (4.26-5.81)	4.17 (3.68-4.77)	8.58 (7.23-10.6)	23.4 (15.5-31.8)	41.1 (29.2-55.9)	1490
	07-08	6.32 (5.49-7.27)	5.31 (4.77-6.05)	11.8 (9.27-13.9)	27.5 (19.5-36.8)	47.7 (34.4-60.9)	1814
	09-10	12.6 (10.7-14.8)	10.9 (9.42-12.7)	28.5 (23.3-35.9)	69.5 (52.5-89.4)	113 (81.9-153)	1914
Gender							
Males	05-06	5.01 (4.21-5.97)	4.29 (3.75-5.03)	8.44 (7.17-10.9)	21.6 (15.1-35.4)	44.6 (25.7-85.9)	1270
	07-08	6.01 (5.29-6.84)	5.14 (4.54-6.26)	11.5 (9.32-13.6)	24.5 (19.2-31.0)	43.2 (31.2-54.9)	1294
	09-10	12.5 (10.6-14.7)	11.4 (9.29-13.9)	28.6 (23.5-36.6)	67.2 (52.0-85.9)	106 (82.2-138)	1399
Females	05-06	5.51 (4.75-6.39)	4.73 (4.23-5.32)	9.55 (8.26-12.2)	24.7 (17.3-30.3)	39.3 (30.3-45.0)	1278
	07-08	7.76 (6.84-8.81)	6.52 (5.95-7.39)	13.7 (11.9-16.5)	35.3 (26.6-44.2)	57.7 (43.5-74.4)	1310
	09-10	13.7 (11.5-16.4)	11.8 (9.80-14.4)	29.5 (22.8-37.9)	75.3 (54.2-110)	123 (93.8-197)	1350
Race/ethnicity							
Mexican Americans	05-06	5.19 (4.38-6.14)	4.50 (3.64-5.29)	8.04 (7.13-10.1)	19.5 (12.3-37.7)	43.6 (20.9-93.4)	637
	07-08	6.66 (5.97-7.44)	5.29 (4.69-6.20)	11.6 (10.4-13.0)	25.7 (20.9-29.8)	38.3 (30.8-58.3)	531
	09-10	9.89 (7.75-12.6)	8.53 (6.98-10.0)	20.9 (12.8-28.3)	51.6 (34.2-72.3)	89.4 (59.3-125)	566
Non-Hispanic blacks	05-06	4.28 (3.86-4.74)	3.67 (3.26-4.09)	7.60 (6.80-8.81)	17.1 (13.2-21.5)	31.8 (22.5-43.7)	678
	07-08	5.87 (4.82-7.13)	5.37 (4.16-6.50)	11.1 (8.68-15.2)	28.1 (17.7-37.2)	40.2 (32.0-57.8)	597
	09-10	9.11 (7.52-11.0)	7.62 (5.84-10.8)	19.8 (15.5-26.5)	47.6 (36.9-57.7)	77.8 (57.0-104)	516
Non-Hispanic whites	05-06	5.53 (4.61-6.63)	4.81 (4.13-5.56)	10.0 (7.63-13.3)	25.2 (16.3-34.5)	39.3 (26.8-59.7)	1038
	07-08	7.11 (5.91-8.56)	6.27 (5.09-7.50)	13.0 (10.4-16.3)	31.1 (21.3-43.2)	54.2 (40.2-64.1)	1077
	09-10	15.1 (12.6-18.1)	13.5 (11.3-16.4)	33.6 (27.0-43.5)	79.2 (60.0-106)	126 (97.9-186)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DiNP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-methyl phthalate (MMP)

Metabolite of Dimethyl phthalate (DMP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	1.15 (.985-1.34)	1.50 (1.30-1.80)	3.30 (2.90-3.70)	6.00 (5.10-7.50)	9.80 (8.00-12.5)	2782
	03-04	*	1.30 (<LOD-1.70)	3.90 (3.00-4.90)	9.70 (7.70-12.2)	16.3 (12.0-20.4)	2605
	05-06	*	< LOD	2.50 (1.90-3.40)	7.40 (6.10-8.60)	12.4 (10.8-14.7)	2548
	07-08	*	< LOD	2.20 (1.80-2.70)	6.50 (4.70-8.30)	11.3 (8.40-16.3)	2604
	09-10	*	.940 (.670-1.21)	2.89 (2.51-3.28)	6.72 (5.76-8.16)	12.0 (9.29-15.3)	2749
Age group							
6-11 years	01-02	1.45 (1.13-1.87)	1.90 (1.40-2.80)	4.00 (3.40-4.90)	7.20 (6.00-8.00)	11.8 (7.60-20.8)	393
	03-04	2.10 (1.67-2.63)	1.70 (1.20-2.40)	4.30 (2.80-6.40)	9.50 (6.40-18.8)	18.8 (9.40-33.4)	342
	05-06	*	< LOD	4.00 (3.30-4.80)	10.6 (7.40-13.4)	15.4 (10.6-30.6)	356
	07-08	*	1.20 (<LOD-1.60)	3.70 (3.00-5.30)	8.00 (6.50-8.70)	12.1 (8.30-35.8)	389
	09-10	2.13 (1.83-2.47)	2.36 (1.72-2.92)	4.83 (3.85-5.61)	10.8 (8.59-16.7)	22.3 (11.4-25.8)	415
12-19 years	01-02	1.59 (1.28-1.96)	2.10 (1.80-2.50)	3.90 (3.30-4.70)	8.50 (5.30-10.5)	13.0 (9.60-17.8)	742
	03-04	*	1.50 (1.00-2.20)	4.40 (3.00-7.00)	13.3 (7.40-20.9)	21.8 (17.8-31.5)	729
	05-06	*	< LOD	3.70 (2.70-4.60)	8.60 (7.00-9.50)	12.4 (9.30-20.7)	702
	07-08	*	< LOD	2.30 (1.70-3.20)	6.70 (4.20-8.90)	12.0 (7.90-17.4)	401
	09-10	1.45 (1.12-1.89)	1.37 (.870-1.98)	3.33 (2.76-4.53)	7.68 (5.27-11.7)	12.0 (7.33-19.6)	420
20 years and older	01-02	1.06 (.904-1.25)	1.40 (1.10-1.60)	3.10 (2.50-3.50)	5.60 (4.60-7.10)	9.20 (7.40-12.3)	1647
	03-04	*	1.20 (<LOD-1.50)	3.80 (2.80-4.90)	9.30 (7.10-11.8)	14.3 (10.8-19.9)	1534
	05-06	*	< LOD	2.20 (1.60-2.80)	6.90 (5.30-8.40)	11.9 (9.50-14.7)	1490
	07-08	*	< LOD	2.00 (1.60-2.50)	6.00 (4.20-8.30)	11.1 (8.10-17.3)	1814
	09-10	*	.750 (<LOD-1.02)	2.59 (2.19-2.90)	6.30 (5.22-7.60)	10.5 (8.49-14.2)	1914
Gender							
Males	01-02	1.17 (.962-1.43)	1.50 (1.30-1.90)	3.30 (2.90-4.00)	5.90 (4.80-7.90)	9.20 (7.10-13.5)	1371
	03-04	*	1.30 (1.00-1.80)	4.00 (3.20-5.20)	10.0 (7.30-13.5)	17.7 (11.8-20.4)	1250
	05-06	*	< LOD	2.40 (1.80-3.50)	7.40 (5.70-9.20)	13.4 (11.0-15.7)	1270
	07-08	*	< LOD	2.40 (1.80-3.20)	6.90 (4.70-8.70)	10.3 (8.20-17.2)	1294
	09-10	*	1.04 (.730-1.40)	2.92 (2.60-3.47)	6.53 (5.38-8.66)	11.4 (8.86-13.9)	1399
Females	01-02	1.13 (.973-1.31)	1.40 (1.10-1.70)	3.30 (2.80-3.70)	6.40 (5.00-7.80)	10.3 (8.20-16.2)	1411
	03-04	*	1.20 (<LOD-1.60)	3.60 (2.80-4.70)	9.50 (7.70-12.2)	15.3 (11.7-22.9)	1355
	05-06	*	< LOD	2.60 (1.80-3.40)	7.40 (5.10-9.30)	12.0 (9.30-16.0)	1278
	07-08	*	< LOD	2.10 (1.70-2.50)	6.20 (4.30-8.20)	12.0 (7.90-16.1)	1310
	09-10	*	.840 (.600-1.09)	2.81 (2.30-3.25)	6.81 (5.49-8.27)	13.5 (8.91-16.8)	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.2, 1.0, 1.1, 1.1, and 0.5 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DMP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-methyl phthalate (MMP)

Metabolite of Dimethyl phthalate (DMP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	01-02	1.21 (1.02-1.45)	1.60 (1.30-1.70)	3.30 (2.70-4.00)	5.60 (4.70-7.30)	8.60 (6.40-15.2)	677
	03-04	*	1.40 (1.10-1.80)	4.00 (2.50-6.30)	9.60 (7.30-13.4)	15.0 (11.4-18.8)	652
	05-06	*	< LOD	3.10 (2.20-4.30)	7.30 (5.90-9.40)	11.3 (8.50-18.0)	637
	07-08	*	< LOD	2.50 (1.30-3.70)	6.40 (4.20-11.1)	12.1 (6.80-16.3)	531
	09-10	1.36 (1.22-1.51)	1.26 (.970-1.53)	3.67 (2.97-4.16)	8.30 (6.51-9.24)	12.0 (9.36-18.2)	566
Non-Hispanic blacks	01-02	1.64 (1.37-1.98)	2.10 (1.70-2.70)	4.40 (3.60-5.10)	8.30 (6.20-10.1)	11.0 (9.50-13.4)	703
	03-04	2.16 (1.64-2.84)	1.70 (1.00-2.80)	5.10 (3.70-7.00)	11.4 (7.30-17.3)	17.8 (10.2-39.4)	699
	05-06	*	< LOD	3.70 (2.40-4.70)	8.20 (6.10-10.6)	13.0 (10.6-15.8)	678
	07-08	*	< LOD	2.40 (1.50-3.40)	6.10 (4.10-8.80)	10.6 (6.70-16.4)	597
	09-10	1.71 (1.32-2.21)	1.89 (1.33-2.66)	4.11 (3.38-5.12)	7.87 (5.90-12.5)	14.2 (8.47-17.0)	516
Non-Hispanic whites	01-02	1.08 (.906-1.29)	1.40 (1.10-1.70)	3.20 (2.50-3.60)	5.60 (4.70-6.70)	9.70 (7.10-14.0)	1216
	03-04	*	1.20 (<LOD-1.60)	3.60 (2.80-4.80)	9.70 (7.50-12.2)	16.5 (12.0-20.4)	1088
	05-06	*	< LOD	2.30 (1.70-2.80)	7.40 (5.50-8.80)	12.5 (10.4-15.0)	1038
	07-08	*	< LOD	2.10 (1.70-2.60)	6.80 (4.30-8.50)	11.2 (8.00-18.7)	1077
	09-10	*	.690 (<LOD-.930)	2.37 (1.95-2.85)	6.17 (5.23-8.04)	12.3 (8.97-15.9)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.2, 1.0, 1.1, 1.1, and 0.5 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DMP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-methyl phthalate (MMP) (creatinine corrected)

Metabolite of Dimethyl phthalate (DMP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	1.08 (.936-1.25)	1.33 (1.13-1.55)	2.62 (2.36-2.97)	5.00 (3.97-6.02)	8.00 (6.07-11.0)	2782
	03-04	*	1.53 (<LOD-1.79)	3.45 (2.76-4.37)	7.95 (5.83-11.8)	13.5 (11.3-18.1)	2605
	05-06	*	< LOD	2.85 (2.52-3.25)	6.53 (5.49-7.67)	12.2 (9.74-14.3)	2548
	07-08	*	< LOD	2.79 (2.31-3.25)	5.68 (4.77-7.36)	10.8 (8.00-13.6)	2604
	09-10	*	1.18 (1.03-1.38)	2.56 (2.29-2.92)	5.55 (4.81-6.54)	10.6 (7.70-13.1)	2749
Age group							
6-11 years	01-02	1.65 (1.28-2.14)	2.32 (1.72-2.86)	3.97 (3.27-4.71)	7.20 (5.99-9.41)	13.2 (7.60-22.5)	393
	03-04	2.22 (1.75-2.81)	1.88 (1.49-2.60)	4.96 (2.81-6.49)	9.67 (6.15-14.6)	16.2 (12.1-34.5)	342
	05-06	*	< LOD	4.59 (3.09-6.35)	11.0 (8.07-15.0)	18.6 (10.4-21.9)	356
	07-08	*	2.05 (<LOD-2.33)	4.56 (3.85-5.36)	9.06 (6.63-11.4)	13.6 (10.4-36.0)	389
	09-10	2.77 (2.44-3.15)	2.85 (2.46-3.29)	5.24 (4.34-6.18)	14.7 (9.87-16.6)	20.5 (14.7-27.7)	415
12-19 years	01-02	1.23 (1.01-1.48)	1.51 (1.32-1.82)	2.84 (2.52-3.33)	5.36 (3.68-6.39)	7.27 (5.64-11.4)	742
	03-04	*	1.34 (1.12-1.73)	3.19 (2.26-4.43)	7.84 (4.95-12.1)	13.3 (9.49-19.1)	729
	05-06	*	< LOD	2.80 (2.15-3.55)	5.65 (4.33-6.52)	12.4 (6.45-18.9)	702
	07-08	*	< LOD	2.00 (1.63-2.63)	4.32 (3.10-6.05)	7.25 (4.77-10.8)	401
	09-10	1.17 (.907-1.51)	1.18 (.930-1.49)	2.24 (1.81-2.62)	4.17 (3.09-6.47)	7.33 (4.56-15.9)	420
20 years and older	01-02	1.00 (.868-1.16)	1.21 (1.05-1.40)	2.44 (2.14-2.68)	4.53 (3.49-6.02)	7.72 (5.52-11.4)	1647
	03-04	*	1.52 (<LOD-1.79)	3.39 (2.69-4.31)	7.69 (5.40-11.8)	13.3 (10.1-18.9)	1534
	05-06	*	< LOD	2.76 (2.44-3.12)	6.09 (5.00-7.75)	11.2 (9.17-13.4)	1490
	07-08	*	< LOD	2.75 (2.26-3.25)	5.58 (4.33-7.59)	10.4 (7.09-14.8)	1814
	09-10	*	1.09 (<LOD-1.23)	2.34 (2.02-2.69)	5.16 (4.25-6.39)	9.53 (7.07-12.1)	1914
Gender							
Males	01-02	.954 (.794-1.15)	1.17 (1.02-1.40)	2.37 (2.03-2.75)	4.18 (3.45-5.64)	6.42 (4.94-9.59)	1371
	03-04	*	1.35 (1.15-1.52)	2.95 (2.26-4.09)	7.11 (5.00-10.2)	11.8 (9.05-13.3)	1250
	05-06	*	< LOD	2.19 (1.89-2.50)	5.03 (4.09-6.45)	10.2 (7.75-13.4)	1270
	07-08	*	< LOD	2.33 (1.92-2.88)	4.88 (3.85-7.25)	8.56 (6.25-13.8)	1294
	09-10	*	1.02 (.850-1.21)	2.49 (2.00-2.86)	5.23 (4.25-6.84)	9.51 (7.20-12.4)	1399
Females	01-02	1.21 (1.06-1.38)	1.45 (1.23-1.82)	2.87 (2.58-3.06)	5.56 (4.55-7.14)	10.0 (7.20-15.3)	1411
	03-04	*	1.83 (<LOD-2.10)	3.87 (3.04-4.96)	9.41 (6.41-13.6)	16.5 (12.5-23.5)	1355
	05-06	*	< LOD	3.47 (3.00-4.11)	7.67 (6.00-9.75)	13.2 (9.75-18.7)	1278
	07-08	*	< LOD	3.12 (2.72-3.71)	6.50 (5.20-8.67)	11.7 (8.56-14.8)	1310
	09-10	*	1.35 (1.16-1.57)	2.76 (2.34-3.18)	5.77 (4.94-7.22)	11.3 (8.29-15.5)	1350

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DMP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-methyl phthalate (MMP) (creatinine corrected)

Metabolite of Dimethyl phthalate (DMP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	01-02	1.14 (.975-1.34)	1.48 (1.30-1.63)	2.50 (2.20-2.94)	4.19 (3.77-5.76)	8.47 (5.58-12.5)	677
	03-04	*	1.36 (1.15-1.71)	3.56 (2.45-4.90)	7.53 (5.62-9.64)	12.5 (9.44-16.9)	652
	05-06	*	< LOD	2.86 (2.07-3.67)	6.48 (4.21-8.24)	11.5 (6.94-21.7)	637
	07-08	*	< LOD	2.59 (2.06-3.39)	5.65 (3.60-8.13)	9.40 (6.06-13.9)	531
	09-10	1.35 (1.19-1.53)	1.37 (1.13-1.67)	3.13 (2.61-3.57)	6.18 (5.33-8.40)	10.7 (8.33-15.2)	566
Non-Hispanic blacks	01-02	1.15 (.949-1.40)	1.39 (1.29-1.67)	2.70 (2.36-2.95)	4.87 (4.21-5.93)	8.09 (5.86-11.0)	703
	03-04	1.53 (1.21-1.94)	1.40 (1.07-1.75)	3.14 (2.26-4.37)	7.50 (4.29-14.6)	13.7 (7.70-20.7)	699
	05-06	*	< LOD	2.59 (1.87-3.28)	5.20 (4.53-6.29)	8.50 (6.79-10.3)	678
	07-08	*	< LOD	2.10 (1.69-2.44)	4.22 (2.94-5.74)	7.36 (4.88-8.79)	597
	09-10	1.24 (.983-1.56)	1.35 (1.08-1.58)	2.69 (2.22-3.23)	4.94 (4.03-6.54)	8.18 (5.74-15.5)	516
Non-Hispanic whites	01-02	1.07 (.914-1.26)	1.30 (1.05-1.58)	2.62 (2.32-2.99)	5.22 (3.68-6.82)	8.26 (6.07-12.6)	1216
	03-04	*	1.56 (<LOD-1.89)	3.50 (2.71-4.67)	8.10 (6.00-11.9)	13.6 (11.3-18.9)	1088
	05-06	*	< LOD	2.98 (2.60-3.39)	7.19 (5.58-9.51)	13.4 (10.4-16.6)	1038
	07-08	*	< LOD	2.89 (2.26-3.67)	6.01 (4.63-8.57)	11.7 (7.47-15.6)	1077
	09-10	*	1.13 (<LOD-1.29)	2.48 (2.06-2.80)	5.38 (4.44-6.76)	10.9 (7.68-13.8)	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DMP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(3-carboxypropyl) phthalate (MCPP)

Metabolite of Di-n-octyl phthalate (DOP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	2.75 (2.49-3.04)	3.10 (2.80-3.30)	5.70 (5.10-6.40)	10.0 (8.90-11.3)	14.6 (12.7-17.5)	2782
	03-04	2.91 (2.79-3.04)	3.10 (3.00-3.30)	5.70 (5.20-6.10)	10.2 (9.20-11.8)	15.3 (13.8-16.2)	2605
	05-06	2.05 (1.87-2.26)	2.10 (1.90-2.40)	4.40 (4.00-4.80)	8.90 (7.60-10.1)	13.1 (10.6-17.4)	2548
	07-08	2.76 (2.52-3.04)	2.70 (2.50-3.00)	5.80 (5.20-6.50)	11.1 (9.70-13.1)	18.1 (15.4-22.3)	2604
	09-10	3.02 (2.58-3.53)	3.10 (2.57-3.65)	6.44 (5.44-7.70)	13.4 (11.4-16.5)	22.3 (19.5-26.2)	2749
Age group							
6-11 years	01-02	6.11 (5.46-6.84)	6.70 (5.40-7.50)	11.8 (10.2-13.3)	20.1 (17.8-23.1)	24.7 (22.2-31.6)	393
	03-04	6.86 (5.80-8.11)	7.10 (6.00-8.90)	12.7 (10.4-16.4)	22.0 (17.2-27.7)	29.2 (22.3-36.6)	342
	05-06	4.82 (4.09-5.68)	4.90 (4.30-5.90)	8.50 (6.70-10.4)	13.1 (10.3-17.9)	17.9 (12.6-34.5)	356
	07-08	6.01 (4.80-7.53)	6.40 (5.10-8.00)	11.3 (8.40-15.9)	20.3 (13.2-27.3)	23.9 (19.1-34.1)	389
	09-10	4.54 (3.74-5.52)	4.86 (3.71-6.49)	9.08 (7.14-11.2)	14.4 (10.8-20.3)	20.3 (14.3-51.9)	415
12-19 years	01-02	3.71 (3.18-4.33)	4.00 (3.40-4.70)	7.10 (6.10-8.10)	11.5 (9.50-12.7)	14.1 (11.8-19.0)	742
	03-04	3.72 (3.33-4.15)	4.10 (3.50-4.60)	7.20 (5.80-8.40)	11.0 (9.50-13.6)	15.4 (13.4-17.9)	729
	05-06	2.96 (2.52-3.46)	3.10 (2.50-3.60)	5.90 (4.80-7.30)	11.8 (9.10-14.5)	17.0 (12.7-21.5)	702
	07-08	3.81 (3.15-4.61)	4.00 (3.20-4.70)	8.20 (6.90-9.40)	13.3 (11.1-17.1)	24.1 (13.8-25.6)	401
	09-10	3.65 (2.88-4.64)	3.74 (2.90-4.74)	7.16 (5.48-9.32)	15.9 (10.8-25.8)	32.3 (20.9-54.3)	420
20 years and older	01-02	2.37 (2.11-2.66)	2.60 (2.20-3.00)	4.80 (4.30-5.40)	8.10 (7.20-9.50)	12.0 (10.1-14.2)	1647
	03-04	2.53 (2.41-2.66)	2.80 (2.60-2.90)	4.90 (4.50-5.20)	8.10 (7.50-8.90)	13.2 (10.8-14.3)	1534
	05-06	1.76 (1.60-1.93)	1.70 (1.60-2.00)	3.60 (3.20-4.10)	7.10 (6.10-8.40)	11.3 (9.60-13.6)	1490
	07-08	2.41 (2.18-2.67)	2.50 (2.20-2.60)	5.00 (4.60-5.50)	9.60 (8.00-11.1)	15.6 (12.9-19.9)	1814
	09-10	2.80 (2.40-3.28)	2.88 (2.42-3.39)	6.00 (5.04-7.15)	13.1 (10.8-16.4)	21.8 (18.7-27.9)	1914
Gender							
Males	01-02	2.89 (2.64-3.17)	3.10 (2.80-3.40)	5.70 (5.00-6.80)	9.90 (8.70-12.0)	14.2 (12.4-18.1)	1371
	03-04	3.25 (3.01-3.52)	3.30 (3.00-3.60)	6.00 (5.60-6.80)	11.8 (10.1-13.6)	16.0 (14.2-20.0)	1250
	05-06	2.32 (2.11-2.54)	2.40 (2.00-2.70)	4.70 (4.30-5.20)	9.60 (7.90-11.8)	14.5 (11.5-22.3)	1270
	07-08	3.02 (2.70-3.39)	2.90 (2.60-3.30)	5.80 (5.10-6.60)	12.3 (10.1-15.3)	21.3 (16.7-28.5)	1294
	09-10	3.37 (2.86-3.96)	3.42 (2.91-4.04)	7.09 (6.04-8.32)	15.5 (12.9-18.7)	23.6 (19.9-33.0)	1399
Females	01-02	2.62 (2.29-2.99)	3.00 (2.50-3.30)	5.70 (5.00-6.30)	10.0 (8.50-11.5)	14.7 (11.2-20.3)	1411
	03-04	2.63 (2.44-2.83)	3.00 (2.60-3.20)	5.20 (4.70-5.80)	9.20 (8.20-10.3)	13.4 (11.0-16.2)	1355
	05-06	1.83 (1.59-2.10)	1.80 (1.60-2.10)	4.00 (3.30-4.60)	8.00 (6.60-9.50)	12.1 (9.50-14.7)	1278
	07-08	2.54 (2.30-2.79)	2.60 (2.30-3.00)	5.90 (5.00-6.80)	10.5 (9.10-12.3)	15.4 (13.3-21.5)	1310
	09-10	2.72 (2.29-3.22)	2.79 (2.25-3.40)	5.78 (4.67-7.20)	12.6 (9.86-15.9)	20.7 (15.9-29.7)	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.4, 0.2, 0.2, 0.2, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DOP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(3-carboxypropyl) phthalate (MCPP)

Metabolite of Di-n-octyl phthalate (DOP) and of several high molecular weight phthalates

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	01-02	2.67 (2.26-3.16)	3.00 (2.30-3.50)	5.30 (4.40-5.90)	9.20 (7.30-12.4)	13.6 (10.4-18.7)	677
	03-04	3.08 (2.86-3.32)	3.10 (2.80-3.40)	5.50 (4.80-6.10)	9.90 (7.90-12.6)	13.7 (11.5-19.8)	652
	05-06	2.24 (1.71-2.94)	2.20 (1.60-3.10)	4.50 (3.30-6.50)	9.40 (6.50-13.1)	14.1 (9.90-25.4)	637
	07-08	2.66 (2.37-3.00)	2.70 (2.40-3.20)	5.00 (4.30-5.70)	8.80 (7.60-11.2)	14.5 (11.5-17.8)	531
	09-10	2.31 (1.95-2.74)	2.32 (1.92-2.93)	5.11 (4.44-5.58)	8.95 (7.39-11.3)	14.8 (11.1-23.2)	566
Non-Hispanic blacks	01-02	3.09 (2.81-3.40)	3.30 (2.90-3.60)	6.30 (5.50-6.50)	10.9 (9.10-13.0)	15.1 (13.5-22.4)	703
	03-04	3.30 (3.01-3.62)	3.20 (2.90-3.60)	6.10 (5.50-6.90)	11.6 (9.30-13.4)	20.1 (13.6-25.5)	699
	05-06	2.28 (2.03-2.57)	2.40 (2.10-2.80)	4.50 (3.80-5.60)	8.70 (7.10-10.3)	13.3 (10.4-16.5)	678
	07-08	2.94 (2.50-3.44)	2.90 (2.50-3.40)	5.80 (4.90-6.80)	12.2 (8.50-17.1)	21.1 (13.7-42.4)	597
	09-10	3.38 (2.66-4.31)	3.35 (2.49-4.56)	6.92 (5.86-8.97)	13.6 (10.2-21.0)	22.9 (17.9-28.7)	516
Non-Hispanic whites	01-02	2.72 (2.40-3.08)	3.00 (2.60-3.30)	5.80 (4.80-6.80)	10.3 (8.90-11.9)	15.8 (12.6-19.5)	1216
	03-04	2.87 (2.73-3.02)	3.10 (2.90-3.30)	5.60 (5.00-6.00)	10.1 (9.10-12.0)	15.2 (13.4-16.1)	1088
	05-06	1.97 (1.81-2.15)	2.00 (1.80-2.20)	4.20 (3.80-4.60)	8.70 (6.80-10.3)	13.1 (10.0-17.5)	1038
	07-08	2.84 (2.51-3.21)	2.90 (2.50-3.40)	6.10 (5.30-7.00)	11.4 (9.70-14.0)	18.1 (14.0-24.1)	1077
	09-10	3.17 (2.65-3.79)	3.16 (2.60-3.87)	6.86 (5.72-8.32)	14.5 (12.6-17.8)	23.6 (18.9-35.7)	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.4, 0.2, 0.2, 0.2, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DOP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(3-carboxypropyl) phthalate (MCPP) (creatinine corrected)

Metabolite of Di-n-octyl phthalate (DOP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	2.58 (2.35-2.83)	2.47 (2.25-2.76)	4.08 (3.87-4.48)	7.24 (6.58-8.00)	11.4 (10.0-12.5)	2782
	03-04	2.74 (2.56-2.93)	2.60 (2.42-2.79)	4.39 (3.94-4.86)	7.70 (6.65-8.60)	10.7 (9.53-11.9)	2605
	05-06	2.00 (1.82-2.20)	1.86 (1.67-2.06)	3.55 (3.14-4.00)	6.80 (5.77-7.95)	10.3 (8.68-12.7)	2548
	07-08	2.79 (2.63-2.96)	2.53 (2.37-2.70)	4.76 (4.33-5.16)	9.70 (8.52-10.9)	14.8 (12.7-17.0)	2604
	09-10	3.15 (2.77-3.59)	2.94 (2.55-3.38)	5.71 (4.88-6.72)	11.8 (10.5-13.5)	18.8 (16.0-21.9)	2749
Age group							
6-11 years	01-02	6.96 (6.29-7.70)	7.08 (5.83-7.87)	11.2 (9.15-14.0)	20.7 (15.5-22.3)	26.4 (20.7-27.0)	393
	03-04	7.25 (6.43-8.18)	7.06 (5.93-7.59)	11.2 (9.38-12.4)	19.1 (14.4-25.6)	26.9 (19.3-29.0)	342
	05-06	5.31 (4.57-6.17)	5.06 (4.57-5.79)	8.78 (7.50-10.0)	13.7 (10.2-21.3)	21.3 (13.2-36.7)	356
	07-08	7.41 (5.87-9.34)	6.81 (5.50-9.05)	11.6 (9.43-14.1)	20.2 (12.7-39.2)	30.4 (16.5-68.9)	389
	09-10	5.92 (5.09-6.88)	5.94 (5.09-6.92)	9.47 (8.38-10.3)	15.5 (10.7-23.7)	22.7 (13.6-47.2)	415
12-19 years	01-02	2.86 (2.52-3.25)	2.94 (2.50-3.36)	4.59 (4.03-5.23)	6.69 (6.30-7.25)	9.55 (8.03-10.7)	742
	03-04	2.78 (2.54-3.05)	2.75 (2.53-2.99)	4.55 (4.05-5.15)	6.61 (6.15-7.20)	8.24 (7.00-10.2)	729
	05-06	2.20 (1.90-2.55)	2.07 (1.76-2.46)	3.67 (2.99-4.39)	6.84 (5.35-8.67)	9.67 (7.12-11.4)	702
	07-08	2.96 (2.59-3.38)	2.95 (2.51-3.62)	5.04 (4.33-6.00)	8.18 (6.75-10.4)	10.9 (9.38-15.1)	401
	09-10	2.94 (2.40-3.60)	2.67 (2.26-3.28)	4.68 (4.04-5.56)	10.8 (7.30-17.8)	21.4 (15.6-29.9)	420
20 years and older	01-02	2.24 (2.03-2.47)	2.20 (2.00-2.40)	3.51 (3.11-3.95)	5.39 (4.82-6.18)	7.71 (6.71-9.28)	1647
	03-04	2.43 (2.27-2.61)	2.38 (2.19-2.56)	3.70 (3.33-4.08)	6.09 (5.38-7.06)	8.79 (7.84-9.56)	1534
	05-06	1.76 (1.60-1.94)	1.67 (1.47-1.86)	2.96 (2.60-3.38)	5.55 (4.61-6.46)	8.46 (7.00-10.3)	1490
	07-08	2.48 (2.33-2.64)	2.26 (2.14-2.36)	4.00 (3.68-4.39)	8.03 (7.11-9.09)	12.7 (11.1-14.8)	1814
	09-10	2.97 (2.60-3.39)	2.75 (2.38-3.22)	5.25 (4.45-6.31)	11.3 (9.63-13.2)	18.1 (15.4-21.6)	1914
Gender							
Males	01-02	2.35 (2.17-2.56)	2.20 (2.02-2.42)	3.76 (3.45-4.20)	7.17 (6.27-8.06)	11.5 (9.28-15.1)	1371
	03-04	2.55 (2.39-2.72)	2.36 (2.22-2.47)	4.14 (3.52-4.86)	7.84 (6.80-8.79)	11.6 (10.2-13.5)	1250
	05-06	1.86 (1.70-2.03)	1.70 (1.53-1.88)	3.33 (2.92-3.70)	6.70 (5.59-8.33)	10.7 (8.72-14.2)	1270
	07-08	2.54 (2.34-2.76)	2.32 (2.10-2.50)	4.17 (3.80-4.81)	8.91 (7.37-11.2)	14.7 (12.9-17.5)	1294
	09-10	3.02 (2.61-3.49)	2.82 (2.43-3.36)	5.51 (4.70-6.36)	10.9 (9.84-12.1)	18.5 (13.7-23.7)	1399
Females	01-02	2.81 (2.48-3.18)	2.76 (2.46-3.02)	4.39 (3.97-4.92)	7.66 (6.17-8.69)	11.0 (8.62-15.9)	1411
	03-04	2.93 (2.68-3.20)	2.89 (2.58-3.23)	4.53 (4.09-5.00)	7.58 (6.44-8.57)	10.0 (8.60-11.6)	1355
	05-06	2.15 (1.88-2.45)	2.00 (1.72-2.44)	3.77 (3.15-4.47)	6.90 (5.62-8.43)	10.0 (8.14-12.5)	1278
	07-08	3.04 (2.84-3.25)	2.80 (2.50-3.11)	5.08 (4.64-5.52)	10.3 (8.67-11.3)	15.0 (11.8-17.8)	1310
	09-10	3.28 (2.88-3.74)	3.00 (2.58-3.46)	5.87 (4.96-7.05)	13.0 (10.3-14.8)	19.1 (15.4-29.5)	1350

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DOP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-(3-carboxypropyl) phthalate (MCPP) (creatinine corrected)

Metabolite of Di-n-octyl phthalate (DOP) and of several high molecular weight phthalates

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	01-02	2.52 (2.21-2.88)	2.37 (2.07-2.73)	4.29 (3.73-5.00)	7.36 (5.58-10.3)	11.4 (8.36-14.5)	677
	03-04	2.78 (2.56-3.02)	2.63 (2.30-2.96)	4.32 (3.66-4.96)	8.14 (6.72-10.3)	12.8 (9.37-17.1)	652
	05-06	2.02 (1.62-2.51)	1.79 (1.42-2.53)	3.85 (2.74-5.22)	7.00 (5.40-8.82)	10.2 (7.64-16.4)	637
	07-08	2.60 (2.26-2.99)	2.45 (2.05-3.10)	4.31 (3.49-5.22)	8.31 (6.67-9.76)	12.9 (9.50-15.5)	531
	09-10	2.29 (1.92-2.74)	2.08 (1.76-2.60)	4.29 (3.61-5.05)	9.02 (7.16-11.4)	13.7 (10.9-17.8)	566
Non-Hispanic blacks	01-02	2.17 (2.02-2.33)	2.07 (1.88-2.28)	3.68 (3.25-4.07)	6.73 (5.46-7.70)	10.0 (8.27-13.1)	703
	03-04	2.34 (2.13-2.58)	2.17 (1.99-2.45)	3.96 (3.33-4.43)	6.93 (6.03-8.80)	13.4 (9.32-15.9)	699
	05-06	1.61 (1.42-1.81)	1.58 (1.38-1.77)	2.76 (2.27-3.37)	5.63 (4.33-7.59)	8.68 (6.86-10.4)	678
	07-08	2.28 (1.96-2.65)	2.11 (1.72-2.35)	3.91 (3.33-4.56)	7.97 (5.58-12.8)	16.4 (8.21-57.0)	597
	09-10	2.45 (2.02-2.97)	2.26 (1.81-2.84)	4.14 (3.35-5.19)	8.02 (6.24-11.3)	14.7 (9.92-19.7)	516
Non-Hispanic whites	01-02	2.69 (2.42-3.00)	2.57 (2.25-2.94)	4.19 (3.87-4.79)	7.63 (6.58-8.45)	11.8 (10.0-14.7)	1216
	03-04	2.83 (2.60-3.09)	2.65 (2.43-2.96)	4.55 (3.90-5.16)	7.79 (6.59-8.79)	10.6 (9.43-11.3)	1088
	05-06	2.08 (1.92-2.24)	1.95 (1.76-2.12)	3.60 (3.26-4.00)	6.47 (5.56-7.93)	10.3 (7.95-13.1)	1038
	07-08	2.96 (2.72-3.23)	2.70 (2.50-2.93)	5.06 (4.35-5.77)	10.3 (8.61-11.5)	15.1 (12.3-19.3)	1077
	09-10	3.57 (3.12-4.07)	3.28 (2.81-3.73)	6.33 (5.25-7.39)	13.2 (11.1-14.9)	21.6 (16.6-26.2)	1206

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DOP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-*n*-octyl phthalate (MOP)

Metabolite of Di-*n*-octyl phthalate (DOP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	2.69 (2.02-3.36)	4.87 (3.86-5.71)	2541
	01-02	*	< LOD	< LOD	< LOD	< LOD	2782
	03-04	*	< LOD	< LOD	< LOD	< LOD	2605
	05-06	*	< LOD	< LOD	< LOD	< LOD	2548
	07-08	*	< LOD	< LOD	< LOD	< LOD	2604
	09-10	*	< LOD	< LOD	< LOD	< LOD	2749
Age group							
6-11 years	99-00	*	< LOD	< LOD	3.70 (<LOD-5.88)	5.38 (2.86-8.40)	328
	01-02	*	< LOD	< LOD	< LOD	< LOD	393
	03-04	*	< LOD	< LOD	< LOD	< LOD	342
	05-06	*	< LOD	< LOD	< LOD	< LOD	356
	07-08	*	< LOD	< LOD	< LOD	< LOD	389
	09-10	*	< LOD	< LOD	< LOD	< LOD	415
12-19 years	99-00	*	< LOD	< LOD	2.86 (<LOD-4.20)	4.70 (3.36-7.06)	752
	01-02	*	< LOD	< LOD	< LOD	< LOD	742
	03-04	*	< LOD	< LOD	< LOD	< LOD	729
	05-06	*	< LOD	< LOD	< LOD	< LOD	702
	07-08	*	< LOD	< LOD	< LOD	< LOD	401
	09-10	*	< LOD	< LOD	< LOD	< LOD	420
20 years and older	99-00	*	< LOD	< LOD	2.52 (1.85-3.19)	4.87 (3.53-5.88)	1461
	01-02	*	< LOD	< LOD	< LOD	< LOD	1647
	03-04	*	< LOD	< LOD	< LOD	< LOD	1534
	05-06	*	< LOD	< LOD	< LOD	< LOD	1490
	07-08	*	< LOD	< LOD	< LOD	< LOD	1814
	09-10	*	< LOD	< LOD	< LOD	< LOD	1914
Gender							
Males	99-00	*	< LOD	< LOD	2.69 (1.85-3.70)	4.70 (3.36-5.88)	1215
	01-02	*	< LOD	< LOD	< LOD	< LOD	1371
	03-04	*	< LOD	< LOD	< LOD	< LOD	1250
	05-06	*	< LOD	< LOD	< LOD	< LOD	1270
	07-08	*	< LOD	< LOD	< LOD	< LOD	1294
	09-10	*	< LOD	< LOD	< LOD	< LOD	1399
Females	99-00	*	< LOD	< LOD	2.52 (2.02-3.53)	5.21 (3.70-6.38)	1326
	01-02	*	< LOD	< LOD	< LOD	< LOD	1411
	03-04	*	< LOD	< LOD	< LOD	< LOD	1355
	05-06	*	< LOD	< LOD	< LOD	< LOD	1278
	07-08	*	< LOD	< LOD	< LOD	< LOD	1310
	09-10	*	< LOD	< LOD	< LOD	< LOD	1350

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.51, 1.68, 1.68, 1.85, 1.85, and 0.84 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 1.68 for survey periods 1999-2008 compared with results previously reported.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DOP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-*n*-octyl phthalate (MOP)

Metabolite of Di-*n*-octyl phthalate (DOP)

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.**

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	1.85 (<LOD-2.35)	2.69 (2.35-4.37)	814
	01-02	*	< LOD	< LOD	< LOD	< LOD	677
	03-04	*	< LOD	< LOD	< LOD	< LOD	652
	05-06	*	< LOD	< LOD	< LOD	< LOD	637
	07-08	*	< LOD	< LOD	< LOD	< LOD	531
	09-10	*	< LOD	< LOD	< LOD	< LOD	566
Non-Hispanic blacks	99-00	*	< LOD	< LOD	3.19 (<LOD-5.04)	5.04 (3.70-6.89)	603
	01-02	*	< LOD	< LOD	< LOD	< LOD	703
	03-04	*	< LOD	< LOD	< LOD	< LOD	699
	05-06	*	< LOD	< LOD	< LOD	< LOD	678
	07-08	*	< LOD	< LOD	< LOD	< LOD	597
	09-10	*	< LOD	< LOD	< LOD	< LOD	516
Non-Hispanic whites	99-00	*	< LOD	< LOD	2.69 (2.02-3.53)	5.04 (3.86-5.88)	912
	01-02	*	< LOD	< LOD	< LOD	< LOD	1216
	03-04	*	< LOD	< LOD	< LOD	< LOD	1088
	05-06	*	< LOD	< LOD	< LOD	< LOD	1038
	07-08	*	< LOD	< LOD	< LOD	< LOD	1077
	09-10	*	< LOD	< LOD	< LOD	< LOD	1206

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 1.51, 1.68, 1.68, 1.85, 1.85, and 0.84 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 1.68 for survey periods 1999-2008 compared with results previously reported.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DOP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-*n*-octyl phthalate (MOP) (creatinine corrected)

Metabolite of Di-*n*-octyl phthalate (DOP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	4.03 (3.48-4.38)	5.93 (4.96-7.21)	2541
	01-02	*	< LOD	< LOD	< LOD	< LOD	2782
	03-04	*	< LOD	< LOD	< LOD	< LOD	2605
	05-06	*	< LOD	< LOD	< LOD	< LOD	2548
	07-08	*	< LOD	< LOD	< LOD	< LOD	2604
	09-10	*	< LOD	< LOD	< LOD	< LOD	2749
Age group							
6-11 years	99-00	*	< LOD	< LOD	3.73 (<LOD-6.30)	6.30 (3.31-17.2)	328
	01-02	*	< LOD	< LOD	< LOD	< LOD	393
	03-04	*	< LOD	< LOD	< LOD	< LOD	342
	05-06	*	< LOD	< LOD	< LOD	< LOD	356
	07-08	*	< LOD	< LOD	< LOD	< LOD	389
	09-10	*	< LOD	< LOD	< LOD	< LOD	415
12-19 years	99-00	*	< LOD	< LOD	2.50 (<LOD-2.87)	3.16 (2.59-5.59)	752
	01-02	*	< LOD	< LOD	< LOD	< LOD	742
	03-04	*	< LOD	< LOD	< LOD	< LOD	729
	05-06	*	< LOD	< LOD	< LOD	< LOD	702
	07-08	*	< LOD	< LOD	< LOD	< LOD	401
	09-10	*	< LOD	< LOD	< LOD	< LOD	420
20 years and older	99-00	*	< LOD	< LOD	4.37 (3.48-4.89)	5.93 (5.04-7.76)	1461
	01-02	*	< LOD	< LOD	< LOD	< LOD	1647
	03-04	*	< LOD	< LOD	< LOD	< LOD	1534
	05-06	*	< LOD	< LOD	< LOD	< LOD	1490
	07-08	*	< LOD	< LOD	< LOD	< LOD	1814
	09-10	*	< LOD	< LOD	< LOD	< LOD	1914
Gender							
Males	99-00	*	< LOD	< LOD	3.06 (2.59-3.48)	4.30 (3.26-5.80)	1215
	01-02	*	< LOD	< LOD	< LOD	< LOD	1371
	03-04	*	< LOD	< LOD	< LOD	< LOD	1250
	05-06	*	< LOD	< LOD	< LOD	< LOD	1270
	07-08	*	< LOD	< LOD	< LOD	< LOD	1294
	09-10	*	< LOD	< LOD	< LOD	< LOD	1399
Females	99-00	*	< LOD	< LOD	5.04 (4.20-5.96)	7.21 (5.59-10.5)	1326
	01-02	*	< LOD	< LOD	< LOD	< LOD	1411
	03-04	*	< LOD	< LOD	< LOD	< LOD	1355
	05-06	*	< LOD	< LOD	< LOD	< LOD	1278
	07-08	*	< LOD	< LOD	< LOD	< LOD	1310
	09-10	*	< LOD	< LOD	< LOD	< LOD	1350

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 1.68 for survey periods 1999-2008 compared with results previously reported.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DOP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Mono-*n*-octyl phthalate (MOP) (creatinine corrected)

Metabolite of Di-*n*-octyl phthalate (DOP)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.**

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	*	< LOD	< LOD	3.06 (<LOD-4.70)	5.59 (4.37-6.72)	814
	01-02	*	< LOD	< LOD	< LOD	< LOD	677
	03-04	*	< LOD	< LOD	< LOD	< LOD	652
	05-06	*	< LOD	< LOD	< LOD	< LOD	637
	07-08	*	< LOD	< LOD	< LOD	< LOD	531
	09-10	*	< LOD	< LOD	< LOD	< LOD	566
Non-Hispanic blacks	99-00	*	< LOD	< LOD	2.28 (<LOD-3.26)	3.73 (2.52-5.49)	603
	01-02	*	< LOD	< LOD	< LOD	< LOD	703
	03-04	*	< LOD	< LOD	< LOD	< LOD	699
	05-06	*	< LOD	< LOD	< LOD	< LOD	678
	07-08	*	< LOD	< LOD	< LOD	< LOD	597
	09-10	*	< LOD	< LOD	< LOD	< LOD	516
Non-Hispanic whites	99-00	*	< LOD	< LOD	4.38 (3.60-5.17)	6.12 (5.29-8.40)	912
	01-02	*	< LOD	< LOD	< LOD	< LOD	1216
	03-04	*	< LOD	< LOD	< LOD	< LOD	1088
	05-06	*	< LOD	< LOD	< LOD	< LOD	1038
	07-08	*	< LOD	< LOD	< LOD	< LOD	1077
	09-10	*	< LOD	< LOD	< LOD	< LOD	1206

< LOD means less than the limit of detection for the urine levels not corrected for creatinine.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

**To correct for inadequate purity in the analytical standards, all results and LODs have been adjusted by a factor of 1.68 for survey periods 1999-2008 compared with results previously reported.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DOP_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/Phthalates_FactSheet.html

Urinary Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	75.1 (61.9-91.1)	69.8 (57.8-82.6)	229 (184-298)	538 (471-702)	1320 (1020-1540)	2553
	01-02	51.7 (46.6-57.5)	52.3 (48.9-57.4)	192 (151-226)	577 (447-725)	1250 (863-1640)	2794
	03-04	66.7 (60.4-73.7)	62.0 (54.1-69.7)	195 (171-219)	590 (500-675)	1070 (893-1330)	2594
	05-06	66.5 (59.7-74.1)	58.8 (51.0-67.7)	204 (187-226)	648 (485-745)	1180 (1000-1350)	2528
	07-08	67.1 (60.2-74.8)	57.6 (49.1-66.0)	203 (182-222)	747 (551-994)	1740 (1210-2240)	2424
	09-10	69.3 (61.6-77.9)	57.2 (48.8-64.3)	228 (190-262)	753 (671-880)	1850 (1380-2480)	2792
Age group							
6-11 years	99-00	90.5 (75.1-109)	101 (70.3-138)	257 (172-430)	510 (437-840)	1130 (657-1740)	330
	01-02	84.9 (71.6-101)	72.7 (56.3-97.0)	261 (155-385)	605 (437-989)	1060 (645-1500)	396
	03-04	84.9 (71.6-101)	66.8 (55.4-92.8)	229 (151-314)	574 (371-654)	818 (625-1060)	341
	05-06	94.1 (74.3-119)	74.0 (56.0-95.9)	287 (149-471)	1010 (557-1600)	1870 (1040-3830)	351
	07-08	105 (80.2-137)	90.6 (61.4-123)	277 (183-431)	1370 (491-2190)	2190 (1340-3500)	360
	09-10	109 (88.1-135)	98.5 (75.4-119)	461 (262-605)	1000 (730-1520)	1900 (1120-2800)	371
12-19 years	99-00	123 (91.4-166)	124 (85.6-168)	326 (227-454)	833 (445-1490)	1460 (861-2410)	753
	01-02	69.3 (52.6-91.3)	70.2 (52.5-87.5)	255 (185-344)	774 (573-984)	1360 (922-1950)	744
	03-04	89.0 (75.2-105)	78.0 (59.0-104)	248 (197-332)	808 (500-968)	1200 (900-1790)	729
	05-06	94.4 (76.0-117)	77.1 (59.7-103)	286 (200-416)	836 (530-1170)	1390 (929-2220)	693
	07-08	79.7 (64.4-98.6)	76.0 (52.2-107)	256 (189-300)	828 (396-1100)	1220 (874-2070)	375
	09-10	94.2 (74.1-120)	82.9 (60.8-116)	295 (201-395)	1020 (690-1210)	1390 (1150-2500)	447
20 years and older	99-00	67.6 (55.4-82.4)	60.9 (49.3-74.1)	215 (167-239)	518 (459-573)	1320 (978-1540)	1470
	01-02	46.4 (41.4-52.0)	49.1 (40.8-53.4)	176 (133-216)	520 (396-703)	1210 (771-1900)	1654
	03-04	61.9 (55.2-69.4)	57.9 (48.7-68.5)	180 (150-214)	554 (416-695)	1110 (857-1360)	1524
	05-06	60.5 (53.4-68.7)	54.0 (46.8-63.8)	197 (169-209)	512 (427-664)	1100 (795-1270)	1484
	07-08	62.2 (55.0-70.4)	51.7 (44.3-60.3)	189 (169-211)	686 (514-939)	1640 (1060-2500)	1689
	09-10	63.0 (54.7-72.5)	49.3 (43.0-59.8)	203 (168-250)	701 (572-856)	1980 (1220-2780)	1974
Gender							
Males	99-00	88.9 (71.4-111)	80.6 (66.6-112)	262 (198-355)	587 (501-989)	1540 (989-2080)	1220
	01-02	49.8 (42.8-57.9)	50.8 (46.0-55.0)	190 (137-240)	498 (386-694)	920 (717-1380)	1375
	03-04	73.8 (63.4-85.9)	65.2 (56.1-76.4)	214 (156-311)	709 (535-900)	1200 (969-1380)	1244
	05-06	73.9 (65.3-83.6)	66.4 (53.8-78.5)	210 (180-241)	694 (471-823)	1180 (979-1400)	1252
	07-08	72.7 (62.2-85.0)	63.4 (49.3-76.5)	210 (174-259)	706 (520-1180)	2010 (1210-3520)	1198
	09-10	68.8 (59.0-80.3)	55.0 (46.0-65.0)	207 (168-257)	674 (559-840)	1430 (1160-2370)	1377
Females	99-00	64.1 (52.9-77.6)	57.8 (45.0-73.2)	199 (150-244)	476 (389-722)	1220 (566-1700)	1333
	01-02	53.6 (48.1-59.8)	55.7 (49.8-62.7)	199 (149-234)	642 (511-816)	1470 (1170-1980)	1419
	03-04	60.7 (53.6-68.8)	57.4 (49.1-67.6)	175 (159-201)	466 (381-622)	884 (654-1380)	1350
	05-06	60.1 (52.5-68.9)	52.5 (45.0-61.1)	200 (166-231)	587 (420-717)	1140 (811-1440)	1276
	07-08	62.2 (54.4-71.0)	52.2 (44.4-58.7)	191 (169-217)	752 (476-905)	1570 (994-2180)	1226
	09-10	69.8 (60.7-80.3)	60.2 (46.1-71.1)	249 (200-309)	840 (694-973)	2140 (1210-2910)	1415

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.5, 1.6, 0.3, 0.4, 0.4, and 0.4 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	78.9 (59.8-104)	66.2 (48.4-87.1)	254 (170-402)	806 (534-1020)	1360 (968-2780)	816
	01-02	39.2 (28.5-54.0)	39.9 (28.8-59.9)	169 (100-291)	515 (388-669)	896 (613-1480)	679
	03-04	57.4 (50.2-65.7)	45.5 (35.2-53.3)	178 (130-236)	686 (487-934)	1390 (934-1650)	653
	05-06	60.6 (53.3-68.9)	54.1 (44.3-65.2)	193 (155-239)	671 (424-966)	1350 (966-1550)	634
	07-08	60.5 (46.6-78.6)	45.8 (34.4-67.3)	206 (143-313)	815 (510-1010)	1410 (994-2010)	520
	09-10	60.6 (48.8-75.2)	48.8 (38.6-74.3)	219 (142-283)	622 (495-853)	1190 (884-1570)	604
Non-Hispanic blacks	99-00	91.9 (71.9-118)	103 (81.6-133)	286 (243-377)	553 (459-824)	1190 (640-1900)	607
	01-02	66.1 (48.2-90.7)	72.9 (52.8-97.3)	255 (182-393)	757 (448-1400)	1410 (757-2480)	706
	03-04	75.0 (56.1-100)	66.0 (50.7-86.6)	241 (151-340)	622 (408-875)	1190 (660-2300)	699
	05-06	81.3 (71.6-92.4)	76.5 (64.7-97.1)	254 (199-331)	676 (463-874)	1210 (760-1790)	662
	07-08	84.0 (63.3-111)	69.8 (43.4-102)	338 (248-419)	1080 (735-1500)	2180 (1230-2610)	562
	09-10	77.2 (64.6-92.1)	69.9 (58.1-82.6)	229 (192-281)	711 (534-1080)	1350 (894-3480)	534
Non-Hispanic whites	99-00	74.4 (61.5-89.9)	66.9 (56.2-78.2)	216 (157-298)	512 (438-745)	1360 (989-1710)	917
	01-02	48.6 (43.8-54.0)	49.8 (42.8-54.2)	171 (137-204)	504 (389-658)	1140 (774-1620)	1222
	03-04	65.8 (58.6-74.0)	62.0 (52.8-71.9)	191 (165-215)	572 (416-722)	1070 (823-1330)	1079
	05-06	62.0 (54.4-70.7)	54.2 (46.9-64.1)	200 (164-216)	581 (430-717)	1040 (811-1310)	1039
	07-08	63.6 (56.8-71.2)	56.7 (46.2-65.0)	180 (158-212)	603 (455-832)	1430 (915-2170)	993
	09-10	68.3 (58.2-80.2)	55.0 (45.5-66.8)	219 (168-284)	726 (584-887)	1740 (1150-2510)	1186

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.5, 1.6, 0.3, 0.4, 0.4, and 0.4 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Daidzein (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	68.5 (55.9-83.9)	65.1 (52.8-80.8)	204 (156-249)	560 (471-629)	944 (836-1150)	2553
	01-02	48.5 (43.7-54.0)	48.3 (43.1-56.5)	166 (140-196)	500 (391-608)	957 (801-1180)	2794
	03-04	62.5 (58.3-67.0)	56.1 (50.0-63.5)	164 (144-187)	502 (429-591)	1060 (825-1150)	2594
	05-06	64.8 (58.7-71.6)	52.4 (47.4-59.7)	184 (162-209)	564 (461-679)	1070 (896-1310)	2528
	07-08	67.5 (60.5-75.2)	55.6 (47.4-64.0)	197 (178-217)	672 (538-897)	1400 (1070-2250)	2424
	09-10	74.2 (67.0-82.2)	55.9 (51.6-61.7)	219 (184-272)	790 (688-1090)	1920 (1550-2390)	2791
Age group							
6-11 years	99-00	92.6 (76.3-112)	93.1 (71.2-114)	251 (157-324)	552 (363-838)	1070 (781-2150)	330
	01-02	96.4 (79.0-118)	85.9 (60.3-127)	275 (159-395)	655 (452-733)	938 (683-1140)	396
	03-04	90.4 (77.2-106)	72.8 (58.1-99.5)	201 (152-240)	454 (329-638)	702 (526-1200)	341
	05-06	104 (83.1-130)	73.8 (59.5-96.7)	305 (175-453)	1040 (686-1830)	2430 (1190-3630)	351
	07-08	128 (101-164)	106 (80.0-141)	307 (197-429)	1420 (543-2630)	2630 (1380-3670)	360
	09-10	149 (124-179)	128 (91.0-162)	500 (346-601)	1230 (814-1530)	2150 (1450-3070)	370
12-19 years	99-00	83.1 (58.4-118)	85.6 (53.8-126)	207 (138-386)	628 (295-1060)	1000 (628-2380)	753
	01-02	53.4 (40.8-70.0)	50.9 (38.3-77.6)	181 (142-248)	549 (385-718)	1030 (622-1360)	744
	03-04	66.6 (55.7-79.6)	56.4 (43.9-74.6)	179 (151-217)	517 (370-753)	877 (617-1240)	729
	05-06	70.0 (56.9-86.2)	55.1 (42.9-74.3)	203 (149-263)	545 (338-845)	914 (659-1750)	693
	07-08	61.3 (51.5-72.9)	61.9 (39.9-75.0)	191 (144-217)	470 (295-645)	821 (544-1110)	375
	09-10	88.6 (71.3-110)	76.5 (53.6-107)	249 (177-312)	689 (522-969)	1410 (823-2740)	447
20 years and older	99-00	63.8 (51.5-79.1)	59.3 (48.7-75.0)	194 (151-234)	554 (471-624)	908 (783-1180)	1470
	01-02	43.9 (39.4-48.9)	43.8 (37.4-53.3)	153 (125-186)	428 (348-590)	946 (728-1220)	1654
	03-04	59.2 (54.3-64.5)	51.7 (46.4-62.6)	155 (131-185)	501 (404-658)	1090 (808-1240)	1524
	05-06	60.7 (54.3-67.8)	50.6 (43.1-59.5)	171 (151-199)	483 (434-609)	1030 (847-1200)	1484
	07-08	63.9 (56.0-72.9)	49.2 (42.6-57.7)	187 (164-216)	675 (514-904)	1340 (962-2780)	1689
	09-10	67.1 (59.4-75.7)	50.3 (44.2-55.5)	194 (168-232)	775 (613-1120)	1980 (1550-2640)	1974
Gender							
Males	99-00	69.7 (54.7-88.8)	70.2 (51.3-84.3)	198 (147-276)	623 (494-836)	1050 (884-1290)	1220
	01-02	40.5 (34.8-47.1)	41.7 (34.5-47.9)	139 (107-179)	348 (260-553)	788 (585-947)	1375
	03-04	57.7 (49.4-67.3)	48.9 (41.6-58.2)	160 (126-205)	467 (368-600)	919 (665-1200)	1244
	05-06	59.5 (53.3-66.3)	51.0 (43.2-57.6)	164 (134-203)	467 (403-612)	927 (781-1100)	1252
	07-08	61.3 (52.5-71.7)	48.7 (41.5-63.2)	178 (141-205)	585 (412-845)	1290 (907-2810)	1198
	09-10	62.8 (54.8-72.1)	49.9 (42.6-57.3)	174 (145-196)	613 (497-790)	1740 (1170-2570)	1376
Females	99-00	67.4 (54.8-82.9)	62.7 (51.1-80.9)	207 (152-250)	509 (356-624)	850 (610-1410)	1333
	01-02	57.6 (50.8-65.2)	59.5 (48.1-73.4)	191 (159-227)	615 (536-722)	1180 (924-1430)	1419
	03-04	67.4 (60.8-74.9)	65.1 (54.6-73.4)	165 (143-191)	564 (409-699)	1090 (757-1370)	1350
	05-06	70.4 (61.3-80.8)	57.1 (45.1-71.7)	194 (169-216)	609 (463-887)	1200 (957-1750)	1276
	07-08	73.8 (64.9-84.0)	63.0 (52.1-77.2)	223 (190-259)	763 (577-1000)	1410 (1070-1990)	1226
	09-10	87.2 (77.9-97.7)	64.1 (55.6-76.7)	292 (240-336)	972 (800-1230)	2280 (1560-2880)	1415

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Daidzein (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	72.5 (59.1-88.9)	64.2 (47.1-91.1)	243 (176-310)	677 (468-1080)	1380 (753-2690)	816
	01-02	36.9 (27.8-49.0)	37.1 (24.8-57.9)	146 (101-225)	441 (307-596)	722 (565-1160)	679
	03-04	51.8 (45.0-59.6)	47.7 (35.9-58.3)	159 (117-209)	502 (354-686)	1130 (779-1390)	653
	05-06	54.7 (45.6-65.7)	47.2 (35.7-61.1)	174 (136-219)	558 (380-617)	925 (642-1140)	634
	07-08	59.5 (44.1-80.1)	41.1 (27.4-76.1)	200 (150-288)	715 (502-1060)	1410 (1090-1860)	520
	09-10	63.7 (52.5-77.1)	49.2 (37.9-65.4)	213 (167-273)	614 (487-861)	1380 (865-1940)	604
Non-Hispanic blacks	99-00	59.1 (46.5-75.1)	67.6 (52.1-87.1)	172 (134-207)	381 (316-533)	802 (562-1010)	607
	01-02	46.4 (33.7-63.8)	49.8 (35.5-69.7)	169 (104-249)	504 (263-773)	939 (542-1530)	706
	03-04	53.1 (42.5-66.4)	44.7 (28.6-72.1)	147 (121-191)	415 (275-538)	727 (538-1040)	699
	05-06	57.0 (50.3-64.5)	48.5 (40.3-59.7)	157 (124-221)	467 (370-654)	913 (603-1290)	662
	07-08	65.0 (48.7-86.6)	53.0 (36.0-80.6)	240 (171-292)	728 (520-937)	1330 (800-2670)	562
	09-10	62.1 (53.0-72.8)	55.7 (42.7-65.0)	179 (157-223)	576 (457-690)	956 (710-1840)	533
Non-Hispanic whites	99-00	72.8 (60.3-88.0)	67.5 (55.3-81.5)	207 (160-249)	560 (442-659)	908 (742-1350)	917
	01-02	48.1 (43.4-53.3)	47.0 (42.0-56.0)	163 (137-191)	463 (375-627)	957 (805-1220)	1222
	03-04	64.9 (59.8-70.3)	58.2 (50.4-66.3)	165 (138-204)	516 (385-667)	1090 (786-1250)	1079
	05-06	65.2 (57.9-73.3)	52.7 (46.5-60.7)	184 (155-211)	564 (443-737)	1070 (894-1390)	1039
	07-08	66.2 (58.4-75.1)	57.7 (46.0-65.5)	187 (160-216)	604 (474-763)	1210 (907-1670)	993
	09-10	76.6 (67.1-87.4)	56.7 (51.2-64.1)	225 (173-306)	809 (617-1180)	1920 (1410-2470)	1186

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Enterodiol

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	26.6 (21.9-32.3)	34.0 (29.4-38.7)	78.8 (62.6-95.5)	165 (135-215)	266 (215-335)	2527
	01-02	35.7 (32.5-39.3)	39.4 (36.3-43.4)	89.2 (80.7-96.9)	181 (162-205)	253 (223-308)	2794
	03-04	39.5 (36.1-43.3)	44.9 (40.6-50.3)	106 (97.2-115)	242 (212-264)	367 (311-423)	2594
	05-06	37.7 (33.2-42.9)	42.4 (39.4-47.2)	106 (95.5-120)	244 (226-266)	379 (305-440)	2528
	07-08	36.8 (32.0-42.4)	43.5 (37.8-49.2)	109 (92.1-124)	238 (199-294)	429 (279-723)	2424
	09-10	39.5 (36.3-43.1)	44.5 (41.0-48.5)	107 (98.4-120)	264 (226-288)	545 (446-651)	2792
Age group							
6-11 years	99-00	26.5 (17.1-41.0)	29.4 (21.2-44.2)	78.7 (44.2-109)	215 (91.1-279)	276 (131-458)	327
	01-02	33.6 (29.8-37.8)	35.5 (29.4-43.7)	78.1 (63.8-87.0)	152 (113-171)	203 (167-327)	396
	03-04	42.0 (34.5-51.1)	42.1 (34.4-49.2)	91.4 (74.4-110)	199 (126-289)	312 (217-525)	341
	05-06	33.2 (26.1-42.3)	38.5 (34.7-43.2)	70.4 (55.7-91.7)	129 (99.1-162)	181 (129-366)	351
	07-08	38.5 (30.1-49.3)	42.9 (31.3-55.3)	80.4 (62.7-118)	181 (119-318)	318 (186-439)	360
	09-10	38.6 (33.9-44.0)	39.9 (32.5-44.2)	88.8 (67.7-106)	195 (131-265)	277 (210-380)	371
12-19 years	99-00	29.8 (23.8-37.2)	34.0 (27.4-42.0)	84.4 (59.1-101)	166 (112-234)	253 (182-337)	744
	01-02	35.3 (30.5-40.9)	37.7 (34.8-43.4)	84.2 (72.1-96.9)	165 (128-206)	238 (169-343)	744
	03-04	45.1 (39.4-51.6)	46.3 (38.7-56.9)	99.5 (84.9-109)	191 (161-269)	325 (275-415)	729
	05-06	39.1 (34.0-45.0)	41.1 (36.8-46.5)	92.0 (80.2-109)	193 (151-238)	265 (231-303)	693
	07-08	37.3 (29.5-47.2)	45.8 (32.2-54.1)	97.9 (73.4-126)	199 (151-253)	293 (212-483)	375
	09-10	37.5 (30.4-46.2)	38.3 (32.5-50.7)	98.0 (72.3-119)	184 (122-286)	315 (177-820)	447
20 years and older	99-00	26.1 (21.8-31.3)	34.3 (29.8-38.7)	78.3 (63.5-94.8)	160 (132-196)	263 (189-335)	1456
	01-02	36.1 (31.8-41.0)	40.4 (36.0-45.7)	91.2 (79.3-105)	190 (161-220)	256 (224-312)	1654
	03-04	38.4 (34.2-43.0)	44.9 (40.1-51.9)	112 (98.8-125)	255 (215-274)	398 (320-448)	1524
	05-06	38.1 (32.7-44.4)	43.8 (39.0-49.5)	116 (103-130)	266 (242-281)	413 (326-479)	1484
	07-08	36.5 (31.2-42.8)	43.5 (37.6-49.3)	116 (97.5-130)	247 (194-326)	473 (300-757)	1689
	09-10	40.0 (36.2-44.1)	47.6 (42.2-53.0)	114 (101-130)	277 (230-366)	596 (460-746)	1974
Gender							
Males	99-00	25.3 (19.5-32.7)	33.0 (28.0-38.0)	72.6 (54.7-94.3)	149 (109-219)	258 (169-286)	1206
	01-02	35.2 (31.8-39.1)	40.5 (36.8-44.8)	90.6 (82.3-103)	184 (158-198)	263 (223-338)	1375
	03-04	39.7 (36.2-43.6)	45.1 (40.5-51.1)	102 (92.9-115)	231 (186-266)	361 (268-460)	1244
	05-06	40.4 (32.8-49.8)	46.2 (38.9-52.3)	107 (90.6-127)	264 (227-289)	413 (287-496)	1252
	07-08	33.1 (28.6-38.2)	39.8 (34.7-43.3)	91.8 (78.6-116)	216 (180-242)	378 (238-738)	1198
	09-10	37.4 (30.9-45.3)	41.7 (35.5-48.4)	94.7 (78.9-111)	227 (173-312)	476 (366-664)	1377
Females	99-00	27.9 (23.4-33.3)	36.0 (29.9-40.3)	84.4 (71.8-97.9)	177 (146-219)	280 (219-375)	1321
	01-02	36.2 (32.2-40.7)	38.5 (35.3-43.4)	87.0 (75.6-98.5)	175 (152-212)	248 (220-283)	1419
	03-04	39.3 (33.8-45.5)	44.9 (38.3-53.0)	110 (97.3-124)	250 (203-294)	398 (309-464)	1350
	05-06	35.4 (30.3-41.2)	41.1 (35.6-45.4)	106 (94.2-123)	225 (186-267)	352 (268-477)	1276
	07-08	40.7 (33.0-50.2)	48.4 (39.3-58.1)	123 (101-145)	258 (206-336)	479 (311-757)	1226
	09-10	41.7 (37.6-46.3)	48.9 (41.6-56.7)	123 (111-140)	277 (243-304)	553 (448-778)	1415

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.8, 1.5, 0.3, 0.04, 0.04, and 0.04 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Enterodiol

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	21.7 (19.5-24.1)	28.0 (24.7-34.7)	70.4 (60.8-78.8)	143 (117-169)	213 (169-256)	791
	01-02	30.5 (25.7-36.3)	34.0 (29.0-39.2)	75.9 (58.8-89.7)	159 (119-202)	244 (192-298)	679
	03-04	33.1 (26.4-41.6)	38.7 (31.2-47.6)	95.2 (78.1-115)	198 (125-307)	307 (186-480)	653
	05-06	38.1 (32.2-45.1)	40.6 (34.6-47.6)	102 (86.8-123)	262 (177-359)	383 (289-511)	634
	07-08	28.2 (24.8-32.0)	29.7 (26.7-33.4)	70.8 (59.9-83.8)	173 (139-208)	305 (207-456)	520
	09-10	27.9 (24.5-31.7)	31.1 (28.8-36.7)	75.5 (66.6-85.5)	157 (120-192)	255 (174-408)	604
Non-Hispanic blacks	99-00	25.8 (21.7-30.7)	31.2 (24.4-35.9)	66.0 (50.2-86.7)	157 (122-193)	260 (185-336)	608
	01-02	35.0 (28.9-42.3)	38.7 (33.2-49.2)	83.7 (70.0-103)	169 (132-191)	225 (175-339)	706
	03-04	40.4 (34.9-46.7)	47.1 (40.4-53.7)	100 (81.3-123)	212 (157-254)	293 (232-417)	699
	05-06	34.7 (30.1-40.1)	38.4 (32.6-44.7)	85.1 (70.9-109)	190 (137-242)	269 (245-364)	662
	07-08	34.4 (26.9-44.0)	38.9 (28.9-48.5)	95.7 (78.5-126)	202 (147-252)	327 (232-514)	562
	09-10	32.4 (27.3-38.4)	35.7 (30.4-42.3)	98.5 (76.2-126)	211 (178-245)	307 (221-546)	534
Non-Hispanic whites	99-00	29.2 (24.0-35.4)	37.6 (31.3-43.8)	85.8 (68.3-99.4)	171 (138-228)	270 (187-375)	915
	01-02	35.6 (31.8-40.0)	40.4 (36.1-44.7)	89.6 (78.5-101)	175 (153-198)	254 (214-337)	1222
	03-04	39.8 (35.5-44.7)	45.8 (38.9-54.9)	109 (95.8-126)	255 (206-267)	365 (294-424)	1079
	05-06	37.3 (31.2-44.6)	42.6 (37.1-50.1)	108 (94.0-127)	243 (219-267)	371 (285-456)	1039
	07-08	39.1 (31.5-48.5)	46.8 (39.3-55.5)	116 (91.8-140)	257 (188-369)	444 (269-839)	993
	09-10	44.1 (39.0-49.9)	51.7 (44.2-56.7)	114 (100-129)	277 (236-311)	596 (414-778)	1186

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.8, 1.5, 0.3, 0.04, 0.04, and 0.04 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Enterodiol (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	24.2 (20.3-28.9)	29.9 (25.0-34.7)	70.5 (59.6-82.1)	146 (124-177)	240 (199-320)	2527
	01-02	33.5 (30.7-36.7)	37.6 (33.0-42.1)	79.2 (72.6-87.3)	149 (136-164)	224 (197-250)	2794
	03-04	37.0 (33.6-40.7)	41.7 (37.9-45.4)	95.9 (85.6-106)	194 (171-213)	294 (247-368)	2594
	05-06	36.8 (32.7-41.4)	41.0 (35.9-46.3)	103 (94.3-114)	204 (174-219)	323 (280-424)	2528
	07-08	37.0 (32.3-42.3)	42.1 (36.8-48.1)	103 (87.1-117)	224 (164-328)	395 (292-634)	2424
	09-10	42.3 (38.9-46.1)	47.1 (44.9-49.8)	108 (101-117)	281 (234-333)	522 (415-602)	2791
Age group							
6-11 years	99-00	27.0 (18.6-39.3)	33.7 (21.8-43.7)	62.7 (43.0-108)	168 (81.6-290)	290 (150-411)	327
	01-02	38.1 (32.5-44.7)	39.3 (30.8-53.1)	78.3 (66.6-103)	188 (147-244)	304 (242-389)	396
	03-04	44.7 (37.4-53.5)	46.6 (38.5-58.3)	93.3 (73.9-125)	177 (128-205)	215 (181-413)	341
	05-06	36.6 (28.9-46.4)	37.8 (32.3-46.5)	75.6 (60.9-105)	156 (109-219)	223 (165-326)	351
	07-08	47.2 (37.5-59.4)	48.0 (38.0-65.2)	86.3 (67.9-178)	230 (181-297)	335 (230-854)	360
	09-10	52.6 (45.0-61.4)	54.3 (39.8-64.5)	104 (79.5-139)	235 (145-335)	336 (239-413)	370
12-19 years	99-00	20.1 (16.7-24.2)	24.1 (19.6-30.1)	55.1 (42.6-71.7)	99.5 (91.5-116)	158 (121-177)	744
	01-02	27.2 (23.3-31.8)	28.7 (24.9-34.9)	62.7 (54.9-71.0)	104 (84.2-139)	152 (111-225)	744
	03-04	33.8 (30.3-37.7)	34.8 (33.2-38.8)	67.6 (64.1-73.9)	130 (107-164)	221 (138-324)	729
	05-06	29.0 (25.2-33.3)	31.4 (27.8-36.1)	68.8 (57.7-80.7)	131 (113-160)	167 (160-216)	693
	07-08	28.7 (22.3-36.9)	33.8 (25.1-43.3)	73.4 (52.9-99.3)	131 (100-207)	207 (128-280)	375
	09-10	35.3 (29.5-42.1)	37.8 (29.7-44.1)	81.7 (66.9-101)	170 (111-303)	306 (170-763)	447
20 years and older	99-00	24.7 (20.6-29.5)	30.6 (26.0-34.8)	73.0 (62.5-84.5)	157 (129-184)	242 (199-344)	1456
	01-02	34.2 (30.3-38.5)	39.6 (32.5-45.3)	84.1 (73.8-93.3)	152 (136-169)	224 (193-250)	1654
	03-04	36.7 (32.5-41.4)	42.0 (37.1-47.8)	103 (91.8-111)	205 (173-239)	327 (256-431)	1524
	05-06	38.2 (33.1-44.0)	43.1 (36.4-49.4)	111 (96.0-128)	216 (187-242)	358 (291-475)	1484
	07-08	37.5 (32.2-43.7)	42.5 (36.8-49.9)	109 (90.7-134)	242 (167-372)	447 (328-718)	1689
	09-10	42.6 (39.0-46.4)	48.3 (46.0-51.1)	113 (103-128)	303 (252-355)	579 (461-747)	1974
Gender							
Males	99-00	19.8 (15.4-25.4)	25.4 (20.0-31.4)	55.1 (46.4-64.2)	122 (94.6-168)	199 (145-282)	1206
	01-02	28.7 (26.0-31.7)	31.4 (27.0-36.8)	71.3 (64.2-78.0)	136 (114-160)	212 (179-259)	1375
	03-04	31.1 (27.5-35.0)	34.8 (32.3-38.5)	78.0 (68.8-92.7)	165 (128-211)	249 (187-361)	1244
	05-06	32.5 (26.7-39.5)	34.2 (29.5-41.1)	90.9 (75.2-102)	192 (147-235)	311 (218-490)	1252
	07-08	27.9 (23.9-32.6)	33.0 (28.5-38.6)	79.3 (72.4-87.1)	164 (122-234)	259 (172-499)	1198
	09-10	34.2 (27.9-42.0)	39.0 (33.0-45.6)	80.9 (69.7-103)	197 (150-253)	345 (235-494)	1376
Females	99-00	29.3 (25.0-34.4)	35.4 (29.5-41.6)	85.2 (74.5-92.3)	165 (141-201)	321 (226-370)	1321
	01-02	38.9 (34.9-43.3)	43.9 (39.3-48.7)	88.3 (74.7-102)	157 (139-174)	235 (180-304)	1419
	03-04	43.6 (38.3-49.7)	49.0 (43.4-57.7)	109 (98.8-123)	215 (178-270)	377 (278-443)	1350
	05-06	41.4 (36.1-47.4)	48.2 (41.8-55.7)	116 (98.0-131)	210 (183-231)	376 (235-470)	1276
	07-08	48.4 (40.6-57.6)	53.2 (44.2-62.9)	128 (105-154)	300 (215-388)	529 (372-759)	1226
	09-10	52.1 (46.7-58.1)	59.4 (49.4-66.2)	139 (121-156)	384 (311-450)	694 (526-938)	1415

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Enterodiol (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	19.6 (17.3-22.2)	23.5 (19.7-28.7)	60.6 (49.8-77.7)	134 (114-154)	193 (154-227)	791
	01-02	28.7 (24.5-33.7)	31.0 (27.0-37.3)	64.1 (55.0-77.5)	133 (94.4-183)	223 (143-337)	679
	03-04	29.9 (23.8-37.4)	34.0 (26.5-42.8)	74.1 (62.0-101)	176 (118-234)	241 (189-364)	653
	05-06	34.4 (28.3-41.8)	35.3 (30.7-40.0)	84.6 (71.2-110)	200 (145-271)	315 (221-515)	634
	07-08	27.7 (24.5-31.3)	29.5 (23.9-33.8)	75.9 (61.3-88.2)	169 (124-198)	240 (178-387)	520
	09-10	29.3 (25.8-33.3)	33.6 (29.5-38.0)	72.3 (61.4-82.7)	157 (116-193)	265 (193-351)	604
Non-Hispanic blacks	99-00	16.6 (13.9-19.7)	18.8 (14.4-22.9)	47.3 (37.7-55.8)	113 (85.9-145)	169 (131-272)	608
	01-02	24.5 (19.7-30.6)	27.0 (22.6-33.3)	57.5 (48.2-73.2)	117 (92.2-143)	157 (118-246)	706
	03-04	28.6 (24.5-33.4)	30.4 (26.1-36.1)	68.3 (53.4-87.5)	126 (96.3-191)	203 (153-227)	699
	05-06	24.3 (20.5-28.9)	25.3 (22.1-30.4)	68.6 (50.3-86.0)	125 (110-147)	190 (139-245)	662
	07-08	26.6 (20.8-34.0)	30.0 (23.6-38.0)	62.9 (51.9-80.8)	139 (102-191)	257 (156-459)	562
	09-10	26.1 (22.1-30.8)	29.7 (25.5-35.9)	73.9 (56.7-88.3)	153 (115-191)	252 (191-339)	533
Non-Hispanic whites	99-00	28.6 (24.3-33.6)	34.4 (28.9-39.3)	75.8 (65.3-87.2)	163 (130-197)	252 (203-363)	915
	01-02	35.3 (31.6-39.4)	40.9 (34.3-46.8)	83.3 (74.7-90.3)	152 (136-173)	225 (189-275)	1222
	03-04	39.3 (34.4-44.8)	45.5 (38.4-53.0)	105 (90.2-116)	202 (168-234)	299 (244-374)	1079
	05-06	39.2 (33.3-46.1)	45.0 (39.4-50.2)	111 (96.2-124)	210 (175-235)	331 (277-439)	1039
	07-08	40.7 (32.7-50.6)	46.4 (38.6-56.7)	109 (86.3-138)	240 (154-384)	424 (257-802)	993
	09-10	49.5 (43.6-56.1)	53.5 (48.3-59.0)	123 (106-140)	325 (252-384)	572 (414-766)	1186

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Enterolactone

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	239 (200-286)	315 (245-381)	726 (595-879)	1970 (1440-2370)	2800 (2500-3140)	2548
	01-02	259 (233-287)	350 (314-389)	807 (739-873)	1590 (1440-1820)	2720 (1870-3430)	2794
	03-04	298 (265-334)	395 (331-464)	900 (819-969)	1790 (1560-2040)	2620 (2360-2880)	2594
	05-06	283 (245-327)	385 (338-460)	935 (824-1060)	1850 (1710-2040)	2880 (2480-3160)	2528
	07-08	242 (211-277)	311 (263-371)	876 (741-1010)	1740 (1480-2050)	2910 (2170-3320)	2424
	09-10	215 (193-240)	267 (229-306)	734 (659-853)	1760 (1490-2090)	2690 (2220-3160)	2792
Age group							
6-11 years	99-00	308 (219-432)	356 (243-474)	721 (520-1320)	1730 (973-2840)	2840 (1700-3590)	331
	01-02	288 (245-339)	329 (271-412)	680 (566-794)	1380 (929-1620)	2200 (1420-2550)	396
	03-04	384 (287-513)	414 (299-567)	926 (589-1190)	1660 (1140-2280)	2360 (1700-3440)	341
	05-06	300 (242-373)	399 (280-546)	851 (667-984)	1390 (1000-1770)	1910 (1390-2460)	351
	07-08	255 (203-320)	314 (247-418)	706 (547-882)	1340 (897-1610)	1730 (1150-2600)	360
	09-10	255 (206-316)	275 (209-346)	666 (552-749)	1480 (1180-1720)	1870 (1500-2870)	371
12-19 years	99-00	250 (191-327)	317 (242-410)	672 (454-888)	1760 (973-2480)	2920 (1950-4330)	746
	01-02	267 (231-308)	321 (255-399)	729 (617-856)	1480 (1230-1800)	2180 (1560-3440)	744
	03-04	314 (267-369)	400 (333-468)	866 (736-1050)	1690 (1410-2080)	2620 (2000-2890)	729
	05-06	292 (245-348)	393 (350-492)	877 (681-1010)	1620 (1470-1800)	2360 (1780-2960)	693
	07-08	250 (203-309)	281 (217-444)	832 (645-1000)	1450 (1100-2030)	2390 (1450-3320)	375
	09-10	194 (154-244)	229 (181-318)	632 (479-735)	1340 (812-2040)	2190 (1340-2960)	447
20 years and older	99-00	230 (193-274)	310 (242-375)	734 (599-888)	2000 (1490-2390)	2790 (2510-3540)	1471
	01-02	254 (223-289)	357 (314-397)	835 (760-914)	1660 (1460-1890)	2840 (1890-3610)	1654
	03-04	286 (253-324)	394 (311-465)	900 (824-960)	1820 (1560-2060)	2630 (2350-3100)	1524
	05-06	280 (238-330)	379 (332-460)	958 (843-1110)	1970 (1760-2310)	3130 (2540-3570)	1484
	07-08	239 (205-280)	311 (258-374)	923 (746-1090)	1800 (1480-2370)	3000 (2210-4030)	1689
	09-10	215 (189-243)	272 (228-320)	768 (666-930)	1850 (1590-2140)	2770 (2350-3690)	1974
Gender							
Males	99-00	254 (212-304)	351 (266-418)	778 (579-1050)	2000 (1580-2400)	2730 (2430-3350)	1219
	01-02	262 (233-295)	340 (314-387)	873 (769-957)	1810 (1490-2470)	3050 (1990-4070)	1375
	03-04	314 (280-351)	425 (376-477)	938 (840-1060)	1760 (1540-2050)	2620 (2060-3230)	1244
	05-06	301 (247-368)	421 (341-515)	1000 (854-1150)	2020 (1730-2330)	3130 (2660-3340)	1252
	07-08	245 (215-278)	321 (261-389)	882 (714-1030)	1620 (1420-2050)	2700 (2000-3360)	1198
	09-10	210 (173-256)	254 (207-324)	718 (583-870)	1670 (1300-2110)	2680 (1850-4060)	1377
Females	99-00	226 (180-284)	287 (236-339)	684 (560-799)	1890 (1200-2460)	2830 (2100-4330)	1329
	01-02	255 (226-288)	357 (298-397)	759 (680-840)	1450 (1190-1700)	2200 (1710-2950)	1419
	03-04	283 (233-343)	371 (278-465)	859 (706-984)	1810 (1440-2170)	2630 (2210-3440)	1350
	05-06	267 (220-324)	359 (318-445)	874 (716-1030)	1770 (1440-2020)	2660 (2040-3270)	1276
	07-08	240 (196-293)	301 (234-394)	875 (747-1030)	1870 (1510-2200)	2920 (2160-3510)	1226
	09-10	220 (196-247)	281 (237-325)	749 (662-868)	1850 (1520-2220)	2690 (2350-3200)	1415

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.6, 1.9, 0.3, 0.1, 0.1, and 0.1 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Enterolactone

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	212 (169-265)	281 (230-335)	631 (539-732)	1650 (950-2210)	2690 (2380-3350)	813
	01-02	275 (221-342)	347 (312-395)	778 (671-913)	1520 (1090-1920)	2340 (1610-2990)	679
	03-04	275 (239-316)	376 (316-435)	849 (744-958)	1560 (1320-1860)	2240 (1860-3100)	653
	05-06	359 (311-415)	420 (364-515)	914 (750-1080)	1710 (1340-2060)	2660 (1950-3180)	634
	07-08	227 (182-283)	320 (240-385)	747 (634-876)	1520 (1300-1740)	2070 (1650-2840)	520
	09-10	198 (166-237)	265 (215-303)	642 (524-752)	1290 (981-1640)	1930 (1380-2280)	604
Non-Hispanic blacks	99-00	262 (196-349)	363 (293-440)	759 (629-925)	1730 (1000-2420)	2500 (1870-3280)	605
	01-02	278 (226-342)	418 (341-479)	769 (686-853)	1450 (1160-1840)	2000 (1540-2420)	706
	03-04	328 (285-378)	437 (362-526)	942 (768-1110)	1580 (1360-1820)	2280 (1880-2640)	699
	05-06	276 (237-322)	411 (346-483)	924 (827-1050)	1660 (1430-2040)	2620 (2220-3080)	662
	07-08	252 (185-343)	362 (272-452)	902 (700-1120)	1820 (1320-2100)	2350 (1920-3160)	562
	09-10	187 (140-250)	267 (180-346)	777 (543-945)	1410 (1150-1810)	1920 (1580-2340)	534
Non-Hispanic whites	99-00	247 (196-311)	317 (240-403)	752 (616-955)	2040 (1600-2450)	3000 (2460-3880)	917
	01-02	267 (235-303)	357 (307-397)	834 (750-923)	1630 (1420-1890)	2780 (1820-3740)	1222
	03-04	299 (254-352)	396 (299-488)	900 (789-996)	1810 (1510-2140)	2680 (2310-3230)	1079
	05-06	274 (228-329)	373 (318-460)	923 (745-1080)	1800 (1610-2020)	2910 (2360-3220)	1039
	07-08	265 (218-322)	328 (266-409)	926 (733-1130)	1790 (1420-2580)	3030 (2050-4310)	993
	09-10	236 (210-265)	293 (244-334)	765 (660-899)	1900 (1590-2310)	2990 (2210-4300)	1186

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.6, 1.9, 0.3, 0.1, 0.1, and 0.1 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Enterolactone (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	218 (184-260)	284 (247-336)	733 (613-869)	1580 (1290-1830)	2250 (1860-2830)	2548
	01-02	243 (220-268)	324 (293-360)	756 (668-858)	1430 (1250-1580)	2120 (1720-2450)	2794
	03-04	279 (245-317)	371 (324-430)	810 (697-911)	1590 (1350-1910)	2400 (2000-2890)	2594
	05-06	276 (237-321)	382 (329-448)	856 (775-975)	1640 (1480-1840)	2650 (2240-3380)	2528
	07-08	243 (211-281)	345 (292-402)	826 (733-946)	1650 (1390-2050)	2730 (2080-3510)	2424
	09-10	230 (206-258)	301 (268-342)	779 (695-882)	1800 (1540-2090)	2820 (2400-3420)	2791
Age group							
6-11 years	99-00	315 (238-416)	384 (266-435)	704 (495-1110)	1580 (1110-2010)	2100 (1580-3040)	331
	01-02	327 (274-391)	349 (263-478)	738 (603-994)	1420 (1140-1800)	2020 (1420-2940)	396
	03-04	409 (310-540)	470 (354-632)	904 (669-1150)	1480 (1090-2420)	2300 (1480-2690)	341
	05-06	331 (268-408)	459 (325-528)	778 (637-1010)	1480 (1130-1750)	1820 (1430-2510)	351
	07-08	313 (255-384)	402 (309-484)	766 (725-824)	1310 (1020-1590)	1650 (1310-2220)	360
	09-10	346 (283-424)	416 (323-471)	753 (607-904)	1440 (1100-2000)	2290 (1440-3340)	370
12-19 years	99-00	169 (133-214)	210 (172-264)	486 (371-615)	1150 (742-1540)	1850 (1310-2350)	746
	01-02	206 (178-239)	255 (223-293)	619 (466-753)	1110 (869-1480)	1500 (1140-2080)	744
	03-04	235 (202-273)	285 (255-307)	631 (533-744)	1230 (993-1350)	1510 (1240-2230)	729
	05-06	216 (184-254)	297 (255-336)	627 (514-701)	1010 (853-1170)	1410 (1150-1700)	693
	07-08	192 (149-248)	273 (197-373)	540 (477-676)	936 (708-1220)	1620 (936-3420)	375
	09-10	182 (147-226)	228 (176-295)	514 (362-809)	1140 (817-1530)	1740 (1140-3160)	447
20 years and older	99-00	217 (181-261)	288 (249-350)	785 (653-923)	1640 (1330-1890)	2310 (1890-3110)	1471
	01-02	240 (213-271)	331 (300-375)	786 (672-915)	1460 (1300-1630)	2180 (1890-2470)	1654
	03-04	274 (236-318)	377 (316-448)	823 (697-937)	1700 (1430-1980)	2480 (2040-3100)	1524
	05-06	281 (236-335)	399 (326-491)	942 (831-1040)	1730 (1570-2200)	2930 (2350-3530)	1484
	07-08	246 (210-287)	350 (291-418)	906 (784-1040)	1810 (1420-2360)	2820 (2320-3570)	1689
	09-10	229 (203-257)	300 (271-346)	828 (727-939)	1920 (1640-2350)	3070 (2530-3760)	1974
Gender							
Males	99-00	199 (170-234)	263 (228-309)	664 (490-828)	1380 (1200-1710)	2030 (1780-2480)	1219
	01-02	213 (191-238)	287 (255-316)	682 (601-764)	1350 (1090-1580)	1980 (1570-2550)	1375
	03-04	245 (215-280)	330 (288-372)	729 (676-808)	1330 (1160-1580)	2040 (1590-2380)	1244
	05-06	243 (196-300)	336 (279-418)	800 (707-897)	1500 (1270-1680)	2200 (1890-2480)	1252
	07-08	206 (176-241)	303 (251-358)	715 (594-859)	1410 (1080-1850)	2140 (1560-3060)	1198
	09-10	192 (157-235)	261 (210-310)	650 (543-766)	1440 (1170-1670)	2300 (1540-3220)	1376
Females	99-00	238 (191-297)	303 (260-379)	819 (662-968)	1720 (1390-2010)	2550 (1940-3390)	1329
	01-02	274 (241-312)	356 (313-407)	833 (710-1000)	1490 (1320-1670)	2150 (1890-2410)	1419
	03-04	314 (257-385)	414 (327-510)	866 (704-1090)	1840 (1460-2280)	2840 (2220-3910)	1350
	05-06	313 (261-375)	419 (337-529)	954 (809-1060)	1840 (1550-2320)	3100 (2350-4580)	1276
	07-08	285 (234-346)	396 (327-469)	966 (843-1060)	1910 (1560-2410)	3040 (2730-3740)	1226
	09-10	274 (244-309)	362 (305-416)	907 (789-1100)	2200 (1730-2810)	3570 (2800-4200)	1415

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Enterolactone (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	194 (165-228)	254 (225-282)	605 (519-695)	1340 (969-1740)	2100 (1620-2790)	813
	01-02	259 (213-314)	362 (296-409)	730 (586-905)	1240 (1010-1490)	1630 (1240-2540)	679
	03-04	248 (217-282)	350 (310-403)	724 (657-826)	1320 (1020-1560)	1820 (1460-2170)	653
	05-06	324 (267-393)	379 (315-471)	773 (679-953)	1560 (1290-1680)	2060 (1630-2560)	634
	07-08	223 (185-268)	312 (267-379)	733 (642-871)	1330 (1110-1580)	1860 (1620-2160)	520
	09-10	208 (174-249)	284 (214-322)	624 (505-728)	1270 (1090-1460)	1870 (1460-2130)	604
Non-Hispanic blacks	99-00	168 (125-226)	214 (173-274)	546 (411-732)	1130 (874-1450)	1590 (1120-2560)	605
	01-02	195 (155-245)	302 (257-331)	562 (470-629)	985 (821-1130)	1490 (1080-1730)	706
	03-04	233 (197-274)	303 (276-347)	621 (512-707)	1050 (917-1350)	1550 (1170-2220)	699
	05-06	193 (167-223)	286 (252-331)	630 (556-729)	1120 (967-1230)	1490 (1220-1650)	662
	07-08	195 (144-264)	291 (191-408)	659 (564-741)	1220 (993-1360)	1480 (1350-2060)	562
	09-10	150 (113-200)	256 (183-328)	539 (457-659)	927 (757-1160)	1310 (1000-1720)	533
Non-Hispanic whites	99-00	241 (194-300)	323 (279-388)	828 (674-997)	1780 (1390-2020)	2490 (1930-3340)	917
	01-02	264 (232-301)	339 (299-393)	833 (719-944)	1530 (1340-1880)	2410 (1940-2800)	1222
	03-04	294 (246-353)	399 (326-488)	836 (707-980)	1660 (1330-2040)	2500 (1990-3350)	1079
	05-06	288 (238-347)	403 (326-500)	926 (809-1050)	1700 (1500-2200)	2930 (2230-3710)	1039
	07-08	276 (225-337)	386 (318-467)	911 (736-1100)	1870 (1370-2730)	2850 (2200-3640)	993
	09-10	264 (233-299)	329 (290-385)	895 (779-1030)	2140 (1640-2780)	3420 (2740-3990)	1186

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Equol

Metabolite of Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	8.37 (7.21-9.72)	8.02 (6.27-9.90)	17.2 (15.2-19.8)	35.0 (28.9-41.6)	53.7 (40.1-74.2)	2182
	01-02	9.17 (7.76-10.8)	9.00 (7.40-10.5)	19.6 (16.6-23.5)	42.1 (34.3-51.5)	73.5 (53.6-89.2)	2794
	03-04	8.02 (7.07-9.10)	8.00 (6.80-9.40)	18.2 (15.2-20.8)	36.6 (31.3-40.0)	64.9 (45.4-97.9)	2590
	05-06	8.40 (7.63-9.24)	8.61 (7.89-9.17)	18.1 (16.4-20.4)	37.6 (33.9-41.8)	64.5 (54.8-85.3)	2527
	07-08	7.29 (6.21-8.57)	7.18 (6.18-8.44)	15.3 (12.5-18.5)	30.7 (24.5-43.2)	65.9 (50.7-131)	2424
	09-10	7.82 (6.91-8.85)	7.39 (6.59-8.34)	17.1 (14.9-18.9)	38.0 (33.9-42.4)	68.6 (52.4-95.6)	2792
Age group							
6-11 years	99-00	10.5 (7.65-14.3)	11.7 (5.43-18.6)	24.9 (17.5-29.5)	34.4 (29.5-53.3)	56.1 (30.1-149)	272
	01-02	12.2 (10.2-14.6)	13.7 (11.1-16.2)	26.2 (17.9-37.9)	50.4 (35.0-84.3)	85.4 (50.4-159)	396
	03-04	12.4 (9.71-15.8)	12.1 (8.40-17.8)	24.4 (20.1-31.2)	46.7 (33.5-82.3)	85.8 (44.9-118)	341
	05-06	13.3 (11.3-15.7)	13.5 (12.1-16.8)	26.5 (20.8-34.1)	45.6 (39.3-69.6)	70.3 (44.7-144)	351
	07-08	10.9 (8.38-14.3)	11.3 (8.60-13.2)	22.0 (15.8-27.5)	55.9 (24.2-77.8)	73.3 (59.5-115)	360
	09-10	13.4 (9.85-18.4)	12.7 (9.01-16.7)	28.5 (22.6-41.8)	68.0 (45.3-179)	242 (68.0-598)	371
12-19 years	99-00	10.9 (8.64-13.8)	10.8 (8.52-13.4)	22.3 (16.0-34.9)	42.9 (34.1-71.3)	71.6 (48.1-210)	657
	01-02	10.2 (8.50-12.1)	10.5 (8.20-12.5)	20.5 (16.8-24.8)	43.1 (30.1-56.1)	70.0 (46.3-99.2)	744
	03-04	10.6 (8.97-12.4)	10.5 (8.70-12.2)	22.6 (19.3-26.4)	39.9 (34.8-45.1)	61.9 (45.1-113)	729
	05-06	11.6 (10.3-13.0)	11.4 (9.25-12.3)	23.2 (20.3-28.8)	46.3 (37.2-64.0)	81.8 (50.1-146)	693
	07-08	8.47 (7.19-9.98)	7.91 (6.79-9.30)	16.0 (12.8-19.0)	32.7 (24.5-43.2)	65.6 (33.6-148)	375
	09-10	11.4 (8.90-14.6)	9.60 (7.57-14.1)	24.6 (19.6-29.6)	48.0 (35.9-112)	220 (69.4-701)	447
20 years and older	99-00	7.79 (6.79-8.94)	7.43 (5.71-8.85)	16.0 (13.6-18.1)	33.1 (24.4-39.7)	52.2 (36.3-93.9)	1253
	01-02	8.70 (7.29-10.4)	8.00 (6.20-10.2)	18.6 (15.0-22.3)	41.3 (34.1-47.4)	73.5 (53.9-89.0)	1654
	03-04	7.28 (6.37-8.33)	7.20 (6.00-8.70)	16.0 (14.2-19.2)	33.9 (28.2-38.9)	63.0 (41.3-121)	1520
	05-06	7.58 (6.82-8.42)	7.57 (7.01-8.30)	16.6 (14.9-18.2)	33.6 (29.1-42.6)	58.1 (50.6-89.0)	1483
	07-08	6.82 (5.70-8.17)	6.75 (5.70-7.98)	14.3 (11.6-18.3)	28.9 (23.6-41.9)	65.8 (47.7-135)	1689
	09-10	6.97 (6.07-8.00)	6.72 (5.78-7.75)	15.0 (13.1-17.1)	33.2 (28.4-38.3)	57.7 (44.6-75.4)	1974
Gender							
Males	99-00	9.15 (7.37-11.4)	8.44 (6.36-11.2)	19.0 (15.9-24.0)	35.6 (29.2-54.8)	71.3 (39.7-166)	1042
	01-02	9.41 (7.99-11.1)	9.20 (7.70-10.8)	20.1 (16.7-26.1)	43.1 (32.4-53.1)	61.7 (51.8-81.5)	1375
	03-04	8.56 (7.54-9.72)	8.70 (7.20-10.2)	19.0 (15.9-21.6)	38.0 (31.9-44.2)	72.6 (45.4-100)	1240
	05-06	9.13 (7.99-10.4)	9.42 (8.16-10.8)	20.0 (16.7-23.8)	39.4 (33.8-45.3)	63.1 (52.0-88.1)	1252
	07-08	7.58 (6.40-8.97)	7.74 (6.62-8.80)	15.5 (12.5-18.2)	32.7 (24.5-41.9)	65.6 (47.7-133)	1198
	09-10	8.20 (7.14-9.42)	8.18 (6.86-9.67)	18.4 (15.2-21.3)	39.3 (31.4-48.0)	70.7 (53.4-112)	1377
Females	99-00	7.70 (6.79-8.75)	7.57 (5.79-9.04)	15.6 (12.7-18.9)	33.6 (26.7-37.7)	48.2 (37.1-62.9)	1140
	01-02	8.94 (7.38-10.8)	8.60 (6.70-10.7)	19.0 (15.6-22.9)	41.6 (33.3-51.5)	79.8 (56.6-122)	1419
	03-04	7.55 (6.44-8.84)	7.30 (6.20-9.00)	16.9 (14.7-19.8)	33.8 (28.3-40.0)	60.3 (42.5-116)	1350
	05-06	7.75 (6.85-8.78)	7.57 (6.87-8.37)	16.7 (15.0-18.0)	35.6 (30.9-43.1)	69.2 (51.7-125)	1275
	07-08	7.03 (5.84-8.47)	6.75 (5.85-7.85)	15.1 (12.3-19.0)	30.0 (24.4-48.3)	76.3 (49.5-163)	1226
	09-10	7.47 (6.33-8.81)	6.75 (5.70-7.85)	15.4 (13.1-18.5)	36.4 (29.4-45.3)	65.2 (46.4-130)	1415

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 3.0, 3.3, 0.3, 0.06, 0.06, and 0.06 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Equol

Metabolite of Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	5.24 (4.77-5.76)	4.51 (3.65-5.18)	9.48 (7.96-10.3)	18.5 (14.5-22.6)	30.9 (21.6-48.4)	726
	01-02	7.22 (6.04-8.62)	6.50 (4.40-9.10)	14.2 (11.3-20.1)	31.4 (21.1-41.0)	42.4 (38.1-60.2)	679
	03-04	6.08 (5.08-7.28)	5.70 (4.70-6.80)	12.5 (9.80-15.4)	26.1 (18.1-39.7)	43.6 (34.3-88.2)	653
	05-06	6.04 (5.31-6.87)	6.13 (5.28-7.35)	13.3 (11.2-14.7)	25.2 (19.6-33.4)	47.6 (33.6-67.1)	634
	07-08	5.52 (5.02-6.07)	5.15 (4.71-5.86)	11.4 (10.1-14.0)	23.2 (20.2-30.3)	39.9 (30.7-60.9)	520
	09-10	5.07 (4.51-5.71)	5.13 (4.45-5.74)	11.1 (8.72-13.0)	20.2 (17.1-29.7)	46.3 (29.7-75.4)	604
	Non-Hispanic blacks	99-00	6.73 (5.20-8.71)	6.24 (3.86-10.0)	15.1 (12.7-17.6)	27.6 (19.4-35.6)	36.4 (28.9-49.4)
01-02		7.15 (6.06-8.43)	6.10 (4.70-7.60)	14.7 (12.0-18.7)	30.9 (22.8-41.9)	45.7 (36.1-90.5)	706
03-04		7.35 (6.16-8.79)	7.90 (6.00-9.60)	16.5 (14.2-18.3)	32.0 (25.1-37.5)	47.0 (37.8-88.2)	696
05-06		6.96 (5.89-8.22)	6.37 (5.44-7.95)	15.1 (11.7-20.2)	30.1 (23.8-39.4)	48.8 (38.7-64.9)	662
07-08		6.73 (5.28-8.59)	6.69 (5.17-8.19)	15.2 (11.9-18.5)	31.3 (21.7-50.7)	72.0 (36.1-402)	562
09-10		5.59 (4.98-6.27)	5.98 (5.37-6.48)	12.2 (10.8-14.2)	25.0 (21.6-31.0)	43.3 (32.4-53.7)	534
Non-Hispanic whites		99-00	9.26 (7.80-11.0)	8.98 (6.73-11.9)	19.0 (16.1-22.8)	36.2 (30.1-45.4)	56.1 (42.1-89.4)
	01-02	9.91 (7.95-12.4)	10.0 (7.30-12.6)	22.0 (17.4-27.4)	44.4 (35.1-57.5)	74.4 (55.1-107)	1222
	03-04	8.52 (7.26-10.0)	8.80 (7.10-10.3)	19.3 (15.4-23.0)	37.7 (31.7-44.2)	73.1 (46.7-120)	1078
	05-06	9.05 (8.08-10.1)	9.49 (8.47-10.4)	19.4 (16.6-23.2)	39.2 (33.9-45.1)	69.2 (52.2-130)	1038
	07-08	8.01 (6.45-9.93)	8.19 (6.59-9.49)	16.1 (12.5-21.7)	32.7 (24.1-55.9)	79.5 (47.7-152)	993
	09-10	9.45 (8.04-11.1)	9.15 (7.87-10.2)	19.3 (17.5-22.4)	41.3 (35.4-48.7)	75.4 (52.4-189)	1186

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 3.0, 3.3, 0.3, 0.06, 0.06, and 0.06 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Equol (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	7.70 (6.82-8.70)	7.96 (6.87-9.35)	16.2 (13.2-18.6)	30.6 (26.9-35.1)	50.3 (41.8-67.3)	2182
	01-02	8.60 (7.26-10.2)	7.98 (6.62-9.76)	17.6 (14.8-21.8)	37.8 (30.3-46.3)	62.6 (50.0-85.0)	2794
	03-04	7.52 (6.83-8.29)	7.29 (6.56-8.18)	14.6 (13.3-15.7)	27.8 (24.7-31.9)	50.6 (39.9-75.0)	2590
	05-06	8.18 (7.33-9.14)	7.92 (7.01-8.76)	15.9 (14.2-17.4)	32.7 (29.9-35.1)	69.1 (48.4-105)	2527
	07-08	7.33 (6.34-8.48)	6.91 (6.13-7.86)	14.4 (12.2-16.7)	28.9 (23.5-37.4)	60.9 (40.5-93.1)	2424
	09-10	8.38 (7.52-9.33)	7.70 (6.80-8.69)	16.6 (15.3-18.1)	36.3 (30.5-44.5)	73.4 (54.2-122)	2791
Age group							
6-11 years	99-00	10.3 (7.83-13.5)	11.4 (7.46-16.3)	22.6 (14.7-27.2)	32.7 (25.5-46.0)	47.8 (32.7-150)	272
	01-02	13.9 (11.2-17.2)	14.0 (10.6-17.0)	28.8 (19.6-39.8)	54.4 (34.9-99.8)	88.2 (56.8-186)	396
	03-04	13.2 (10.9-15.9)	13.2 (9.19-17.5)	24.2 (20.6-28.0)	41.6 (35.2-64.1)	93.5 (45.4-117)	341
	05-06	14.7 (12.6-17.1)	15.3 (13.8-17.4)	26.2 (21.2-34.4)	48.0 (40.6-70.3)	88.9 (55.4-267)	351
	07-08	13.4 (10.8-16.6)	13.1 (10.6-15.2)	22.6 (16.9-32.8)	45.2 (28.8-84.4)	72.8 (45.8-199)	360
	09-10	18.3 (13.5-24.7)	15.7 (11.5-19.3)	42.0 (28.1-54.1)	84.1 (63.5-183)	232 (98.4-498)	370
12-19 years	99-00	7.61 (6.17-9.39)	8.02 (6.72-9.29)	14.0 (11.4-20.4)	27.5 (20.8-38.3)	47.4 (27.5-149)	657
	01-02	7.83 (6.68-9.17)	7.77 (6.29-9.13)	17.4 (14.9-19.2)	31.9 (25.3-41.7)	54.3 (33.0-76.7)	744
	03-04	7.91 (6.59-9.49)	8.11 (6.80-9.09)	14.5 (11.8-17.5)	30.1 (21.0-38.2)	40.6 (32.6-66.9)	729
	05-06	8.61 (7.54-9.83)	8.03 (7.00-9.21)	16.3 (13.4-19.0)	31.8 (25.3-37.6)	62.0 (37.0-129)	693
	07-08	6.51 (5.69-7.46)	5.83 (4.78-6.71)	12.6 (10.5-16.4)	28.7 (20.8-35.3)	50.4 (30.7-72.7)	375
	09-10	10.7 (8.50-13.5)	8.95 (7.40-10.3)	20.6 (15.9-23.3)	46.3 (31.3-101)	215 (58.9-919)	447
20 years and older	99-00	7.45 (6.60-8.41)	7.63 (6.35-9.22)	15.3 (12.7-17.7)	30.8 (25.0-37.2)	53.2 (41.2-71.8)	1253
	01-02	8.23 (6.93-9.79)	7.52 (5.97-9.56)	16.6 (14.0-20.7)	36.4 (29.3-42.9)	58.7 (47.9-85.0)	1654
	03-04	6.98 (6.30-7.73)	6.76 (6.05-7.65)	13.6 (12.5-14.8)	24.6 (21.9-29.4)	42.8 (35.7-81.6)	1520
	05-06	7.60 (6.72-8.59)	7.43 (6.63-8.21)	14.1 (12.4-16.4)	30.1 (23.9-34.9)	65.9 (42.7-123)	1483
	07-08	7.00 (5.90-8.31)	6.64 (5.73-7.57)	13.2 (11.2-16.2)	26.4 (21.6-38.4)	57.7 (38.4-97.0)	1689
	09-10	7.43 (6.61-8.34)	6.97 (6.09-8.08)	15.0 (13.5-16.7)	29.2 (25.3-34.5)	54.7 (41.4-75.7)	1974
Gender							
Males	99-00	7.01 (5.93-8.29)	7.32 (5.63-8.78)	13.8 (11.7-17.6)	29.3 (21.4-41.8)	54.1 (35.8-81.0)	1042
	01-02	7.66 (6.39-9.18)	7.43 (5.81-9.16)	16.2 (14.0-19.4)	32.9 (27.9-40.4)	54.3 (37.8-67.1)	1375
	03-04	6.71 (6.02-7.47)	6.63 (5.78-7.29)	13.3 (11.9-15.5)	25.7 (23.1-30.3)	51.1 (37.1-76.7)	1240
	05-06	7.35 (6.40-8.43)	7.03 (5.90-8.67)	14.9 (12.0-18.3)	31.4 (26.8-34.0)	57.6 (42.7-83.5)	1252
	07-08	6.39 (5.52-7.41)	6.10 (5.28-6.85)	12.2 (10.6-14.3)	25.7 (20.4-33.0)	48.4 (35.3-77.1)	1198
	09-10	7.49 (6.54-8.59)	6.90 (5.98-8.20)	15.7 (14.1-17.3)	32.4 (25.3-44.1)	73.4 (44.2-130)	1376
Females	99-00	8.41 (7.33-9.66)	8.71 (7.33-10.1)	17.7 (15.1-20.0)	31.6 (27.5-37.2)	46.3 (41.0-56.5)	1140
	01-02	9.60 (7.99-11.5)	8.66 (6.97-10.6)	19.2 (15.7-23.7)	41.7 (32.3-55.9)	85.0 (61.7-115)	1419
	03-04	8.38 (7.39-9.51)	8.29 (7.03-9.28)	15.1 (14.0-16.6)	30.6 (25.0-32.5)	45.4 (38.7-102)	1350
	05-06	9.07 (8.04-10.2)	8.49 (7.90-9.22)	16.7 (15.3-18.1)	34.3 (29.9-40.7)	85.1 (48.4-144)	1275
	07-08	8.35 (6.98-9.99)	7.97 (6.83-9.31)	16.0 (13.4-20.0)	32.8 (23.7-46.1)	67.5 (41.6-167)	1226
	09-10	9.34 (7.98-10.9)	8.44 (7.24-9.76)	17.4 (15.1-20.4)	39.6 (30.7-54.2)	72.9 (51.6-158)	1415

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Equol (creatinine corrected)

Metabolite of Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	4.89 (4.36-5.47)	4.73 (3.90-5.27)	8.85 (8.18-9.91)	22.3 (16.5-26.8)	37.1 (25.3-57.6)	726
	01-02	6.79 (5.82-7.92)	6.81 (5.55-8.04)	14.9 (11.7-17.8)	29.1 (22.9-36.0)	41.3 (31.4-47.1)	679
	03-04	5.48 (4.60-6.54)	4.72 (3.95-5.78)	10.1 (7.76-13.4)	24.4 (17.2-30.9)	35.6 (27.6-102)	653
	05-06	5.45 (4.84-6.14)	5.00 (4.37-5.85)	9.94 (8.80-12.4)	22.3 (19.3-30.0)	39.4 (30.0-61.9)	634
	07-08	5.42 (4.82-6.09)	4.87 (4.28-5.66)	10.9 (9.21-12.2)	20.9 (17.0-26.9)	34.9 (26.9-60.9)	520
	09-10	5.33 (4.60-6.18)	5.00 (4.17-6.16)	9.49 (8.54-11.2)	19.6 (15.1-35.6)	48.5 (30.3-126)	604
Non-Hispanic blacks	99-00	4.36 (3.41-5.57)	4.57 (2.94-6.23)	10.2 (7.96-12.0)	17.1 (14.8-19.8)	26.0 (19.6-32.0)	514
	01-02	5.01 (4.26-5.89)	4.48 (3.80-5.42)	11.0 (8.80-13.0)	22.7 (16.5-29.3)	35.4 (24.8-46.7)	706
	03-04	5.23 (4.44-6.16)	5.28 (4.67-6.10)	10.3 (8.61-12.3)	20.2 (17.1-23.8)	29.4 (23.9-42.3)	696
	05-06	4.88 (4.11-5.78)	4.72 (3.55-6.00)	9.40 (7.54-12.0)	21.5 (16.8-24.4)	31.6 (24.4-51.4)	662
	07-08	5.21 (4.11-6.61)	4.73 (3.77-6.22)	9.81 (7.77-12.7)	19.3 (14.7-28.8)	50.6 (24.2-314)	562
	09-10	4.50 (4.03-5.01)	4.57 (3.80-5.29)	9.76 (8.00-11.3)	18.4 (15.6-22.3)	26.8 (22.8-29.7)	533
Non-Hispanic whites	99-00	9.13 (7.85-10.6)	9.51 (7.63-11.2)	18.0 (15.2-21.6)	35.4 (29.5-41.3)	56.5 (46.0-73.0)	758
	01-02	9.81 (7.93-12.1)	9.17 (7.06-12.2)	19.8 (15.4-25.6)	41.3 (30.7-55.9)	66.4 (56.8-85.0)	1222
	03-04	8.40 (7.50-9.41)	8.17 (7.13-9.19)	15.6 (14.1-17.1)	30.6 (25.0-32.8)	56.8 (38.0-107)	1078
	05-06	9.51 (8.48-10.7)	9.17 (8.06-10.5)	17.1 (15.0-19.4)	34.5 (31.0-40.6)	77.7 (48.4-129)	1038
	07-08	8.33 (6.76-10.3)	7.94 (6.62-9.61)	15.7 (12.5-20.4)	30.7 (23.4-46.1)	65.2 (38.9-144)	993
	09-10	10.6 (9.17-12.2)	9.69 (8.74-10.8)	20.5 (17.8-22.5)	44.2 (33.4-60.8)	84.1 (60.3-187)	1186

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Genistein

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	24.4 (19.7-30.3)	27.0 (22.5-32.8)	93.6 (75.8-118)	284 (244-331)	563 (413-709)	2557
	01-02	33.0 (30.1-36.2)	29.0 (26.8-31.8)	92.5 (77.9-109)	306 (240-372)	619 (523-719)	2794
	03-04	31.1 (29.0-33.3)	26.2 (23.9-29.7)	87.8 (78.8-102)	286 (239-313)	528 (402-610)	2594
	05-06	28.7 (25.8-32.0)	26.0 (20.9-29.7)	81.0 (73.3-92.4)	261 (232-308)	523 (422-591)	2528
	07-08	31.3 (27.3-35.8)	25.7 (22.8-28.9)	89.7 (73.4-109)	309 (227-411)	665 (527-999)	2424
	09-10	32.5 (29.3-36.2)	28.2 (23.2-33.1)	102 (88.8-120)	356 (295-428)	691 (530-1060)	2792
Age group							
6-11 years	99-00	27.6 (21.1-36.1)	31.9 (18.1-42.6)	104 (67.6-151)	220 (151-315)	376 (272-725)	331
	01-02	39.2 (33.4-46.0)	31.7 (25.8-39.6)	94.1 (62.3-158)	258 (190-426)	502 (258-830)	396
	03-04	33.6 (27.8-40.6)	29.3 (23.1-37.8)	79.4 (56.7-111)	193 (146-274)	351 (207-376)	341
	05-06	38.6 (29.8-49.9)	34.6 (25.3-42.5)	109 (76.6-181)	367 (252-568)	629 (367-2390)	351
	07-08	43.7 (32.6-58.5)	34.3 (22.1-56.4)	139 (82.5-194)	505 (251-763)	840 (505-2200)	360
	09-10	46.1 (37.8-56.3)	38.4 (30.7-46.9)	167 (96.9-253)	467 (280-616)	958 (528-1210)	371
12-19 years	99-00	43.7 (34.2-55.7)	45.4 (34.3-60.5)	138 (93.7-179)	319 (245-464)	547 (321-777)	754
	01-02	34.1 (27.2-42.8)	29.0 (26.1-32.9)	90.6 (71.5-113)	278 (216-363)	470 (360-687)	744
	03-04	34.7 (29.3-41.0)	29.0 (23.4-37.6)	111 (69.3-141)	304 (206-376)	530 (358-671)	729
	05-06	34.1 (28.8-40.5)	26.9 (20.5-35.6)	90.5 (76.1-115)	307 (209-394)	486 (372-982)	693
	07-08	35.2 (28.5-43.5)	28.4 (23.7-34.5)	111 (63.7-138)	279 (160-483)	538 (320-954)	375
	09-10	39.2 (29.5-52.0)	34.4 (23.5-47.1)	141 (77.0-188)	373 (242-520)	620 (445-1160)	447
20 years and older	99-00	21.9 (17.6-27.2)	24.0 (21.7-28.4)	86.2 (67.5-108)	293 (235-343)	566 (412-744)	1472
	01-02	32.1 (28.8-35.8)	28.8 (25.4-33.4)	93.4 (77.3-110)	312 (235-389)	627 (537-790)	1654
	03-04	30.2 (27.8-32.8)	25.9 (22.3-29.8)	87.8 (77.9-103)	296 (239-321)	557 (412-653)	1524
	05-06	27.0 (23.9-30.6)	25.0 (19.8-28.8)	77.4 (66.7-89.5)	244 (190-308)	518 (413-591)	1484
	07-08	29.6 (25.4-34.5)	24.7 (21.3-28.4)	84.1 (68.0-98.0)	278 (207-422)	671 (512-1150)	1689
	09-10	30.5 (26.7-34.8)	26.3 (20.3-31.7)	97.5 (77.3-115)	327 (281-425)	691 (523-1110)	1974
Gender							
Males	99-00	29.8 (22.2-40.0)	31.9 (26.3-37.2)	108 (79.1-151)	335 (257-440)	709 (437-981)	1222
	01-02	32.2 (27.9-37.2)	29.5 (25.4-33.7)	91.2 (73.4-103)	239 (190-331)	474 (335-719)	1375
	03-04	33.7 (29.6-38.4)	27.3 (23.4-33.0)	92.6 (68.6-124)	310 (228-351)	562 (443-653)	1244
	05-06	32.1 (28.6-36.0)	29.1 (22.7-33.8)	90.2 (77.5-101)	269 (216-340)	494 (394-591)	1252
	07-08	33.3 (27.7-40.0)	26.2 (22.3-31.2)	98.0 (71.9-119)	316 (220-477)	723 (501-1510)	1198
	09-10	33.5 (29.3-38.2)	28.8 (23.6-35.5)	89.8 (74.7-114)	351 (286-425)	623 (476-967)	1377
Females	99-00	20.3 (17.0-24.2)	23.1 (20.1-26.3)	84.7 (59.6-105)	242 (203-288)	446 (339-619)	1335
	01-02	33.7 (30.9-36.8)	28.7 (26.0-32.3)	97.0 (79.9-118)	387 (253-500)	666 (598-807)	1419
	03-04	28.7 (25.5-32.4)	25.0 (22.4-28.4)	85.0 (71.2-106)	256 (218-308)	467 (356-620)	1350
	05-06	25.9 (22.3-29.9)	22.4 (17.9-27.4)	74.5 (61.0-87.0)	246 (195-329)	526 (398-657)	1276
	07-08	29.5 (25.1-34.6)	24.6 (22.2-28.4)	83.5 (63.7-109)	296 (194-432)	661 (448-964)	1226
	09-10	31.7 (27.6-36.4)	27.0 (21.8-32.1)	115 (101-129)	371 (290-489)	826 (523-1530)	1415

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.3, 0.8, 0.3, 1.0, 0.2, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Genistein

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	31.1 (25.1-38.5)	30.0 (25.1-37.3)	117 (82.9-179)	328 (248-494)	573 (419-1180)	819
	01-02	28.3 (22.0-36.4)	25.6 (19.6-32.5)	74.8 (48.8-111)	225 (174-314)	424 (323-523)	679
	03-04	31.1 (27.5-35.2)	25.2 (19.1-32.6)	83.2 (63.6-120)	319 (252-537)	653 (537-851)	653
	05-06	27.6 (24.4-31.2)	26.6 (21.5-30.2)	77.0 (63.3-87.3)	255 (187-319)	412 (320-568)	634
	07-08	30.5 (24.3-38.4)	24.0 (18.9-30.2)	86.8 (68.0-123)	318 (210-414)	603 (414-1130)	520
	09-10	28.7 (23.7-34.7)	24.2 (18.4-32.3)	84.8 (57.8-120)	257 (197-327)	484 (334-623)	604
Non-Hispanic blacks	99-00	26.7 (19.2-37.0)	32.9 (24.4-41.5)	103 (84.8-137)	257 (213-367)	495 (329-926)	608
	01-02	37.6 (27.4-51.6)	35.3 (23.6-49.5)	95.5 (71.1-142)	378 (192-530)	598 (375-1120)	706
	03-04	32.3 (24.0-43.4)	27.4 (18.8-44.9)	84.8 (60.5-113)	279 (163-412)	514 (323-852)	699
	05-06	31.0 (26.7-36.1)	27.9 (23.0-34.9)	92.3 (69.3-106)	265 (187-355)	458 (371-643)	662
	07-08	36.2 (29.3-44.7)	27.3 (22.6-36.8)	124 (84.3-157)	395 (310-530)	912 (567-1490)	562
	09-10	33.5 (29.0-38.6)	27.8 (23.2-36.4)	99.6 (79.3-120)	321 (218-392)	625 (410-1010)	534
Non-Hispanic whites	99-00	23.6 (19.1-29.3)	25.6 (21.7-32.0)	91.4 (68.0-122)	288 (227-353)	566 (395-734)	917
	01-02	30.9 (27.8-34.4)	27.6 (24.6-30.8)	89.7 (71.1-105)	278 (226-365)	626 (485-755)	1222
	03-04	30.8 (28.2-33.6)	26.2 (22.6-31.0)	93.6 (78.8-112)	279 (225-313)	504 (376-610)	1079
	05-06	26.9 (23.6-30.6)	23.8 (19.0-29.1)	77.2 (66.2-87.9)	244 (198-316)	518 (391-609)	1039
	07-08	29.3 (25.2-34.0)	25.2 (22.1-28.5)	84.1 (65.8-102)	235 (182-346)	620 (446-840)	993
	09-10	31.9 (27.4-37.2)	28.3 (21.3-34.6)	102 (75.4-125)	334 (280-450)	691 (501-1060)	1186

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.3, 0.8, 0.3, 1.0, 0.2, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Genistein (creatinine corrected)

Metabolite of Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	22.3 (17.7-28.1)	23.8 (18.8-28.9)	84.7 (67.2-105)	222 (182-279)	381 (334-497)	2557
	01-02	30.9 (28.5-33.6)	25.9 (23.4-29.2)	83.3 (72.2-96.3)	256 (211-296)	427 (375-490)	2794
	03-04	29.1 (27.3-31.0)	24.5 (21.3-27.4)	77.7 (67.3-90.9)	231 (203-279)	510 (388-619)	2594
	05-06	28.0 (25.3-31.0)	23.2 (20.3-26.4)	72.0 (63.1-85.2)	242 (197-280)	577 (406-734)	2528
	07-08	31.4 (27.3-36.1)	23.7 (21.0-27.7)	85.2 (70.3-99.5)	264 (214-375)	751 (464-1130)	2424
	09-10	34.9 (31.6-38.4)	25.4 (22.8-30.0)	100 (83.3-124)	367 (313-424)	863 (612-1090)	2791
Age group							
6-11 years	99-00	28.3 (21.1-37.9)	27.8 (15.8-41.3)	94.3 (60.5-145)	209 (148-317)	490 (279-895)	331
	01-02	44.5 (37.0-53.5)	37.9 (29.7-49.0)	112 (76.5-146)	252 (173-371)	504 (252-713)	396
	03-04	35.8 (29.7-43.0)	35.4 (24.9-43.2)	78.6 (60.9-119)	172 (130-243)	297 (168-618)	341
	05-06	42.5 (33.1-54.6)	34.1 (23.7-41.9)	138 (89.5-195)	414 (256-850)	1040 (429-1590)	351
	07-08	53.5 (41.5-69.0)	44.2 (33.0-63.9)	149 (83.2-224)	451 (236-937)	940 (411-1800)	360
	09-10	62.8 (52.5-75.1)	47.8 (36.7-63.0)	200 (144-249)	493 (405-644)	875 (551-1140)	370
12-19 years	99-00	29.4 (22.3-38.8)	32.0 (23.8-41.6)	83.2 (64.1-104)	184 (130-295)	336 (184-816)	754
	01-02	26.3 (21.3-32.5)	21.0 (17.8-26.5)	66.2 (47.9-91.5)	200 (149-298)	321 (261-435)	744
	03-04	25.9 (21.8-30.9)	21.8 (17.0-29.1)	65.6 (51.0-83.1)	201 (145-313)	366 (297-455)	729
	05-06	25.3 (22.1-29.0)	19.6 (16.4-24.0)	76.5 (51.4-91.6)	164 (130-249)	368 (213-520)	693
	07-08	27.1 (23.1-31.7)	23.7 (17.8-29.3)	77.7 (56.2-95.3)	159 (125-264)	324 (214-568)	375
	09-10	36.8 (28.5-47.7)	27.6 (20.4-40.1)	102 (67.5-174)	371 (234-455)	848 (424-1100)	447
20 years and older	99-00	20.6 (16.3-26.2)	21.6 (17.7-26.2)	83.1 (64.9-107)	234 (190-287)	381 (325-562)	1472
	01-02	30.4 (27.6-33.4)	24.8 (21.9-30.0)	83.1 (68.5-99.0)	269 (208-328)	435 (374-518)	1654
	03-04	28.9 (26.7-31.3)	24.0 (20.1-27.4)	78.5 (66.1-92.9)	253 (209-302)	542 (399-673)	1524
	05-06	27.1 (24.3-30.3)	23.0 (19.7-26.8)	70.3 (61.2-81.3)	233 (184-275)	575 (349-762)	1484
	07-08	30.4 (25.8-35.8)	22.8 (19.9-25.9)	81.2 (63.2-99.5)	263 (205-423)	842 (464-1310)	1689
	09-10	32.5 (28.9-36.5)	23.1 (20.5-28.2)	91.9 (72.8-116)	340 (295-409)	863 (590-1150)	1974
Gender							
Males	99-00	23.3 (16.8-32.3)	23.8 (17.5-32.2)	86.2 (64.7-115)	236 (178-330)	523 (323-889)	1222
	01-02	26.2 (23.1-29.8)	22.0 (19.4-26.0)	67.6 (57.4-78.1)	186 (144-237)	350 (278-418)	1375
	03-04	26.4 (22.8-30.5)	21.5 (17.3-26.2)	70.0 (51.1-90.5)	203 (159-235)	415 (346-600)	1244
	05-06	25.8 (23.4-28.4)	22.8 (19.3-26.4)	65.0 (54.6-82.2)	207 (154-261)	431 (305-628)	1252
	07-08	28.1 (23.4-33.7)	20.9 (17.6-24.8)	76.1 (58.9-92.9)	249 (174-369)	634 (370-1130)	1198
	09-10	30.6 (27.3-34.3)	23.5 (20.7-27.5)	78.7 (63.0-101)	313 (241-370)	631 (453-1020)	1376
Females	99-00	21.3 (17.5-26.0)	23.2 (17.5-29.3)	83.1 (57.2-106)	211 (154-283)	357 (283-398)	1335
	01-02	36.2 (32.8-39.9)	29.6 (25.2-34.3)	107 (88.4-129)	321 (269-355)	547 (427-729)	1419
	03-04	31.9 (28.7-35.5)	27.1 (23.6-31.6)	87.1 (71.1-102)	278 (209-324)	548 (363-763)	1350
	05-06	30.3 (25.8-35.5)	23.5 (19.8-29.1)	81.3 (67.4-104)	271 (208-406)	658 (444-894)	1276
	07-08	35.0 (30.1-40.8)	27.7 (23.0-34.7)	95.6 (80.3-107)	290 (220-506)	842 (506-1350)	1226
	09-10	39.6 (35.2-44.5)	29.4 (23.9-35.5)	131 (101-151)	430 (340-542)	944 (670-1540)	1415

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary Genistein (creatinine corrected)

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Race/ethnicity							
Mexican Americans	99-00	28.4 (23.3-34.7)	27.9 (22.5-35.0)	109 (91.5-137)	257 (209-380)	562 (257-981)	819
	01-02	26.6 (21.6-32.9)	21.0 (16.1-28.9)	61.6 (50.6-76.8)	205 (147-270)	372 (271-479)	679
	03-04	28.0 (24.8-31.8)	23.5 (18.5-28.1)	69.2 (52.3-92.8)	254 (187-390)	608 (417-764)	653
	05-06	24.9 (22.1-28.0)	21.7 (17.8-23.8)	72.4 (56.2-90.2)	202 (140-274)	338 (275-648)	634
	07-08	30.0 (23.2-38.8)	22.5 (16.5-30.5)	88.3 (55.4-127)	287 (210-373)	583 (385-921)	520
	09-10	30.1 (25.4-35.7)	23.2 (19.0-30.5)	79.2 (60.6-108)	262 (192-353)	601 (371-818)	604
Non-Hispanic blacks	99-00	17.1 (12.4-23.7)	19.5 (15.7-26.1)	59.5 (43.1-93.7)	179 (132-245)	299 (222-446)	608
	01-02	26.4 (19.3-36.1)	22.7 (16.4-33.6)	69.4 (42.2-115)	217 (139-317)	384 (217-747)	706
	03-04	22.8 (18.1-28.8)	19.6 (14.3-27.1)	55.0 (39.6-78.1)	182 (107-240)	311 (210-514)	699
	05-06	21.7 (18.8-25.1)	18.6 (14.9-21.6)	55.3 (42.6-78.7)	157 (124-235)	351 (235-543)	662
	07-08	28.0 (22.5-34.9)	20.8 (15.9-25.9)	91.2 (63.7-112)	252 (184-413)	668 (310-1530)	562
	09-10	26.9 (23.7-30.6)	19.5 (17.6-23.1)	78.0 (58.1-107)	245 (158-341)	505 (341-879)	533
Non-Hispanic whites	99-00	23.2 (18.5-29.0)	24.9 (19.0-31.7)	86.1 (68.4-105)	232 (178-295)	381 (325-523)	917
	01-02	30.6 (28.2-33.2)	25.4 (22.7-29.5)	82.0 (68.3-96.3)	248 (207-320)	427 (365-518)	1222
	03-04	30.4 (27.9-33.0)	26.2 (21.5-31.4)	79.6 (69.1-99.6)	238 (195-321)	534 (352-688)	1079
	05-06	28.2 (25.1-31.8)	23.5 (19.8-29.0)	71.7 (58.9-89.6)	242 (192-305)	615 (423-802)	1039
	07-08	30.5 (25.8-36.0)	23.7 (20.2-28.3)	80.1 (63.4-96.6)	256 (185-369)	651 (412-1010)	993
	09-10	35.8 (31.2-41.0)	25.6 (22.5-32.7)	107 (78.5-140)	398 (304-448)	881 (576-1300)	1186

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary O-Desmethylangolensin

Metabolite of Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	4.39 (3.37-5.73)	4.98 (3.65-6.77)	22.7 (18.7-30.1)	100 (74.8-141)	222 (182-250)	2271
	01-02	4.08 (3.53-4.73)	3.30 (2.70-4.20)	19.8 (16.7-24.6)	96.0 (70.1-135)	260 (161-437)	2794
	03-04	4.91 (4.34-5.55)	4.60 (4.00-5.20)	23.6 (18.6-27.2)	95.9 (70.7-122)	230 (185-342)	2581
	05-06	4.71 (4.17-5.31)	3.80 (3.10-4.30)	22.5 (19.7-26.6)	101 (86.4-124)	266 (200-385)	2528
	07-08	4.65 (4.02-5.39)	3.40 (2.90-4.20)	23.0 (17.9-29.8)	132 (98.2-181)	328 (236-501)	2424
	09-10	4.73 (4.08-5.48)	3.60 (2.90-4.30)	21.8 (16.8-28.4)	125 (99.2-156)	394 (237-615)	2792
Age group							
6-11 years	99-00	5.60 (3.85-8.15)	7.52 (3.43-15.2)	36.2 (20.3-45.0)	78.7 (43.4-191)	176 (74.8-264)	287
	01-02	6.19 (4.51-8.49)	5.90 (3.80-9.30)	26.9 (15.7-52.1)	122 (61.5-215)	281 (161-466)	396
	03-04	6.33 (4.30-9.30)	6.10 (3.50-10.6)	24.0 (17.0-33.8)	70.8 (51.4-122)	138 (80.9-256)	341
	05-06	7.20 (5.20-9.96)	6.20 (4.00-11.0)	37.8 (24.2-55.2)	145 (83.9-295)	386 (201-516)	351
	07-08	7.50 (5.43-10.4)	5.40 (3.90-9.70)	34.5 (16.5-52.6)	133 (79.9-275)	490 (133-3020)	360
	09-10	7.85 (6.20-9.94)	7.00 (4.30-11.7)	34.9 (27.7-49.6)	153 (89.1-199)	263 (167-467)	371
12-19 years	99-00	6.04 (3.76-9.70)	7.60 (5.13-13.5)	36.6 (22.0-57.3)	107 (63.4-165)	194 (107-238)	667
	01-02	5.92 (4.46-7.87)	5.20 (3.70-7.60)	33.6 (18.0-56.8)	125 (91.2-172)	299 (172-435)	744
	03-04	6.37 (4.95-8.18)	5.30 (3.80-8.10)	33.9 (18.2-50.5)	110 (82.2-198)	257 (175-415)	729
	05-06	8.50 (6.48-11.2)	8.10 (5.50-11.6)	41.3 (33.6-49.6)	105 (80.2-213)	385 (139-829)	693
	07-08	4.74 (3.00-7.48)	3.80 (2.20-7.20)	20.6 (11.2-34.6)	105 (29.5-357)	328 (78.1-638)	375
	09-10	6.40 (4.93-8.31)	5.30 (4.00-7.70)	26.8 (16.6-47.8)	117 (77.8-165)	377 (156-965)	447
20 years and older	99-00	4.05 (3.12-5.26)	4.46 (3.31-5.64)	19.8 (16.0-26.5)	101 (80.8-150)	228 (179-259)	1317
	01-02	3.65 (3.08-4.32)	2.80 (2.30-3.70)	17.0 (13.9-22.4)	81.5 (63.0-128)	260 (135-526)	1654
	03-04	4.56 (4.02-5.17)	4.40 (3.60-4.90)	22.0 (17.8-25.6)	94.0 (67.6-131)	230 (187-398)	1511
	05-06	4.09 (3.51-4.77)	3.20 (2.60-4.00)	17.9 (14.9-22.3)	91.2 (72.3-124)	248 (182-302)	1484
	07-08	4.41 (3.72-5.22)	3.20 (2.40-4.20)	23.0 (11.2-30.7)	133 (93.3-187)	320 (206-589)	1689
	09-10	4.28 (3.59-5.11)	3.00 (2.40-3.90)	19.3 (14.9-27.0)	127 (95.2-159)	438 (225-681)	1974
Gender							
Males	99-00	4.97 (3.71-6.66)	5.62 (4.12-8.73)	29.1 (19.8-42.9)	121 (74.1-190)	235 (177-332)	1087
	01-02	3.81 (3.08-4.71)	3.30 (2.60-4.50)	17.4 (13.2-24.6)	82.8 (58.4-116)	194 (123-324)	1375
	03-04	4.90 (3.93-6.12)	4.60 (3.40-5.50)	23.0 (16.6-30.1)	92.9 (70.7-129)	222 (159-332)	1240
	05-06	4.93 (4.13-5.89)	4.00 (3.40-5.50)	22.3 (16.7-28.1)	90.8 (63.5-137)	235 (171-311)	1252
	07-08	4.49 (3.67-5.49)	3.10 (2.50-4.10)	22.1 (14.5-33.7)	117 (81.4-194)	293 (195-589)	1198
	09-10	4.39 (3.46-5.59)	3.30 (2.40-4.40)	18.1 (14.5-24.3)	99.2 (65.2-142)	293 (153-615)	1377
Females	99-00	3.92 (2.97-5.16)	4.22 (3.18-5.51)	19.4 (14.1-26.1)	83.8 (61.1-114)	192 (123-250)	1184
	01-02	4.36 (3.64-5.23)	3.40 (2.50-4.60)	21.3 (16.8-29.2)	107 (70.6-199)	394 (230-746)	1419
	03-04	4.91 (4.26-5.66)	4.60 (3.90-5.50)	23.9 (18.1-27.4)	99.3 (63.5-155)	283 (160-433)	1341
	05-06	4.50 (3.90-5.20)	3.30 (2.70-4.30)	22.8 (17.9-28.3)	121 (83.2-168)	284 (182-499)	1276
	07-08	4.82 (4.03-5.75)	3.90 (2.90-5.30)	24.6 (18.1-33.9)	137 (99.3-195)	362 (236-468)	1226
	09-10	5.09 (4.19-6.18)	3.90 (2.70-5.00)	28.5 (18.4-42.8)	152 (110-230)	494 (309-766)	1415

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.2, 0.4, 0.2, 0.2, 0.2, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary O-Desmethylangolensin

Metabolite of Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/L) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	2.41 (1.55-3.73)	2.14 (1.31-3.37)	21.0 (10.6-30.5)	97.6 (59.7-140)	191 (122-320)	721
	01-02	2.44 (1.51-3.94)	1.40 (.500-3.40)	13.1 (5.80-27.6)	66.4 (33.5-102)	152 (75.8-265)	679
	03-04	2.54 (1.86-3.48)	1.90 (1.00-3.40)	14.4 (9.00-18.1)	62.3 (40.3-99.5)	146 (99.5-254)	652
	05-06	3.04 (2.29-4.05)	2.10 (1.40-2.90)	14.5 (8.20-24.7)	62.8 (39.1-104)	158 (87.0-246)	634
	07-08	3.16 (2.40-4.15)	2.50 (1.50-3.30)	14.7 (10.5-20.7)	84.9 (52.1-152)	211 (159-340)	520
	09-10	3.44 (2.78-4.27)	2.70 (2.10-3.50)	15.2 (10.6-21.2)	93.1 (63.5-114)	216 (127-294)	604
Non-Hispanic blacks	99-00	5.74 (4.55-7.24)	8.71 (5.82-10.9)	33.5 (22.1-52.4)	108 (78.3-156)	192 (149-255)	538
	01-02	5.35 (4.00-7.14)	5.30 (2.90-7.30)	32.7 (22.0-52.4)	128 (75.9-216)	308 (150-436)	706
	03-04	5.49 (4.05-7.46)	4.20 (3.30-6.10)	26.1 (13.4-49.0)	117 (86.2-166)	221 (177-354)	698
	05-06	5.83 (4.49-7.58)	4.90 (3.90-6.30)	32.6 (21.2-46.7)	110 (72.4-189)	297 (162-398)	662
	07-08	6.63 (4.50-9.75)	5.80 (3.90-9.10)	40.8 (23.6-63.3)	183 (119-230)	347 (199-822)	562
	09-10	5.03 (3.75-6.74)	4.30 (2.90-6.00)	27.0 (16.7-35.9)	120 (80.5-168)	293 (168-464)	534
Non-Hispanic whites	99-00	4.50 (3.26-6.22)	4.99 (3.43-7.10)	22.5 (17.1-34.4)	103 (72.0-152)	228 (177-259)	826
	01-02	4.13 (3.43-4.96)	3.40 (2.60-4.40)	17.9 (15.5-23.8)	98.7 (67.4-153)	260 (153-526)	1222
	03-04	5.27 (4.63-5.99)	5.30 (4.40-6.00)	24.2 (18.8-28.5)	99.5 (67.6-149)	246 (183-409)	1070
	05-06	4.72 (4.09-5.45)	3.80 (3.20-4.70)	21.9 (17.5-27.6)	97.7 (72.3-142)	246 (181-363)	1039
	07-08	4.67 (3.88-5.62)	3.50 (2.80-4.60)	21.3 (16.9-27.8)	113 (79.9-193)	357 (193-628)	993
	09-10	4.95 (4.20-5.84)	3.70 (2.90-4.50)	20.4 (16.0-30.1)	139 (97.5-156)	471 (206-766)	1186

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, 03-04, 05-06, 07-08, and 09-10 are 0.2, 0.4, 0.2, 0.2, 0.2, and 0.2 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary O-Desmethylangolensin (creatinine corrected)

Metabolite of Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	4.03 (2.97-5.45)	4.44 (3.10-6.35)	21.8 (15.3-31.6)	90.4 (62.9-122)	167 (140-218)	2271
	01-02	3.83 (3.32-4.43)	3.24 (2.55-4.12)	18.9 (16.1-23.2)	85.2 (65.1-117)	281 (149-412)	2794
	03-04	4.58 (4.20-5.01)	4.09 (3.53-4.59)	19.6 (17.4-22.6)	92.9 (76.5-120)	201 (183-261)	2581
	05-06	4.59 (4.04-5.21)	3.70 (3.12-4.42)	20.7 (18.4-22.8)	95.6 (78.0-129)	215 (180-325)	2528
	07-08	4.68 (4.04-5.41)	3.51 (3.06-4.02)	22.5 (16.7-30.5)	125 (94.8-164)	350 (199-523)	2424
	09-10	5.07 (4.38-5.87)	3.78 (3.06-4.61)	23.6 (19.3-29.7)	138 (101-185)	433 (285-547)	2791
Age group							
6-11 years	99-00	6.00 (4.04-8.91)	7.15 (3.94-16.6)	28.8 (14.3-45.0)	83.3 (45.0-167)	179 (88.2-262)	287
	01-02	7.03 (5.05-9.77)	6.49 (4.24-11.6)	29.9 (16.3-54.1)	101 (75.3-199)	305 (134-464)	396
	03-04	6.73 (4.55-9.97)	5.97 (4.09-8.28)	27.7 (15.6-42.3)	86.1 (45.2-135)	149 (84.3-189)	341
	05-06	7.93 (5.73-11.0)	7.15 (3.90-11.3)	40.9 (27.5-55.9)	164 (85.8-280)	346 (215-497)	351
	07-08	9.20 (6.70-12.6)	6.81 (5.05-10.7)	31.1 (20.1-58.9)	164 (100-296)	465 (175-2310)	360
	09-10	10.7 (8.43-13.6)	9.58 (5.71-13.9)	43.0 (31.3-63.3)	201 (111-260)	330 (226-478)	370
12-19 years	99-00	4.13 (2.33-7.35)	5.71 (2.82-11.5)	26.0 (14.7-44.4)	71.4 (40.8-122)	122 (75.2-262)	667
	01-02	4.57 (3.44-6.07)	3.88 (2.82-5.32)	26.0 (18.0-35.9)	95.0 (61.4-129)	259 (129-331)	744
	03-04	4.76 (3.71-6.11)	4.26 (3.13-5.97)	22.8 (14.3-39.2)	86.4 (49.0-149)	185 (101-261)	729
	05-06	6.30 (4.89-8.13)	5.69 (3.44-8.40)	28.0 (22.5-37.1)	89.0 (58.5-138)	258 (103-686)	693
	07-08	3.64 (2.28-5.83)	2.99 (1.91-5.69)	18.3 (7.73-24.0)	75.3 (21.4-200)	199 (69.4-664)	375
	09-10	6.02 (4.74-7.65)	5.71 (3.30-7.53)	23.6 (16.0-36.8)	109 (59.2-189)	287 (132-558)	447
20 years and older	99-00	3.82 (2.84-5.13)	3.90 (2.69-5.73)	20.2 (12.9-29.2)	96.5 (61.8-133)	172 (140-252)	1317
	01-02	3.46 (2.95-4.04)	2.74 (2.03-3.64)	16.8 (13.7-20.3)	78.9 (58.2-124)	281 (143-467)	1654
	03-04	4.35 (3.96-4.78)	3.75 (3.23-4.35)	18.6 (15.7-22.6)	95.2 (76.5-131)	228 (192-284)	1511
	05-06	4.10 (3.51-4.79)	3.35 (2.57-4.30)	17.4 (14.0-20.6)	89.3 (64.5-140)	197 (160-332)	1484
	07-08	4.52 (3.82-5.36)	3.13 (2.55-3.92)	22.7 (16.1-32.4)	130 (89.1-183)	353 (189-641)	1689
	09-10	4.56 (3.83-5.42)	3.29 (2.42-4.07)	21.2 (16.2-26.8)	139 (95.3-212)	437 (246-634)	1974
Gender							
Males	99-00	3.95 (2.79-5.58)	4.50 (2.88-6.50)	24.5 (13.8-40.5)	96.5 (62.4-122)	210 (125-265)	1087
	01-02	3.10 (2.48-3.86)	2.87 (2.00-3.96)	15.0 (10.4-19.7)	60.7 (41.5-90.2)	154 (96.4-301)	1375
	03-04	3.83 (3.10-4.73)	3.33 (2.41-4.35)	15.6 (13.9-18.9)	76.2 (51.1-99.5)	193 (124-255)	1240
	05-06	3.97 (3.33-4.72)	3.49 (2.73-4.33)	17.7 (14.1-21.4)	68.3 (55.8-85.8)	165 (140-216)	1252
	07-08	3.79 (3.10-4.63)	2.79 (2.33-3.49)	18.2 (12.4-25.7)	86.7 (59.2-153)	264 (139-432)	1198
	09-10	4.02 (3.10-5.19)	3.19 (2.13-4.58)	17.3 (12.3-21.7)	86.2 (59.8-137)	317 (140-485)	1376
Females	99-00	4.10 (3.00-5.61)	4.17 (2.91-6.35)	20.4 (14.4-27.2)	86.0 (49.1-145)	155 (100-205)	1184
	01-02	4.68 (3.87-5.68)	3.74 (2.77-4.88)	26.8 (18.7-34.0)	111 (70.3-205)	399 (175-739)	1419
	03-04	5.45 (4.71-6.30)	4.82 (4.09-5.92)	24.3 (19.6-28.9)	106 (78.5-163)	255 (193-343)	1341
	05-06	5.27 (4.56-6.09)	4.00 (3.07-5.49)	24.1 (19.8-31.1)	140 (96.8-166)	325 (197-613)	1276
	07-08	5.72 (4.75-6.88)	4.23 (3.33-5.71)	29.0 (20.4-43.2)	164 (105-213)	394 (212-743)	1226
	09-10	6.35 (5.34-7.57)	4.42 (3.46-5.79)	34.1 (25.4-43.5)	199 (139-248)	504 (350-761)	1415

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary O-Desmethylangolensin (creatinine corrected)

Metabolite of Daidzein

Geometric mean and selected percentiles of urine concentrations (in µg/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

Race/ethnicity	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Mexican Americans	99-00	2.19 (1.49-3.24)	1.88 (1.15-3.10)	16.6 (11.3-25.8)	71.2 (52.3-113)	136 (90.5-251)	721
	01-02	2.30 (1.48-3.57)	1.45 (.769-3.37)	11.8 (4.91-23.6)	46.7 (31.4-87.8)	108 (54.7-218)	679
	03-04	2.30 (1.68-3.15)	1.65 (.845-3.06)	13.2 (6.63-21.2)	51.6 (36.9-75.4)	125 (84.1-165)	652
	05-06	2.75 (1.92-3.93)	1.69 (1.04-2.99)	12.8 (7.27-24.9)	53.0 (33.3-93.2)	135 (72.6-204)	634
	07-08	3.10 (2.26-4.27)	2.22 (1.42-3.32)	16.0 (8.84-23.8)	82.0 (45.8-145)	215 (98.9-467)	520
	09-10	3.62 (2.92-4.50)	2.71 (1.82-4.92)	13.5 (10.8-19.6)	76.5 (54.2-128)	226 (113-461)	604
Non-Hispanic blacks	99-00	3.65 (2.90-4.60)	5.22 (3.59-6.75)	23.8 (17.6-32.1)	67.1 (52.1-79.7)	116 (81.9-239)	538
	01-02	3.75 (2.76-5.10)	3.57 (2.14-5.48)	22.8 (14.5-30.4)	78.5 (46.5-175)	218 (98.5-339)	706
	03-04	3.89 (3.03-4.99)	3.12 (1.96-4.96)	16.9 (11.3-26.7)	80.0 (58.6-104)	159 (120-255)	698
	05-06	4.09 (3.13-5.34)	4.22 (2.49-5.71)	20.1 (13.5-24.5)	69.9 (44.8-151)	225 (90.2-369)	662
	07-08	5.13 (3.47-7.57)	4.22 (2.84-7.71)	27.0 (15.6-44.5)	133 (81.9-182)	308 (164-485)	562
	09-10	4.06 (3.11-5.28)	3.61 (2.41-5.07)	22.4 (12.9-30.2)	95.1 (59.2-142)	208 (133-418)	533
Non-Hispanic whites	99-00	4.42 (3.12-6.27)	4.68 (3.05-7.23)	22.3 (15.3-37.9)	102 (57.6-148)	177 (140-215)	826
	01-02	4.08 (3.41-4.89)	3.51 (2.45-4.57)	21.2 (16.3-27.4)	96.4 (69.8-126)	300 (153-464)	1222
	03-04	5.18 (4.70-5.70)	4.58 (3.89-5.39)	21.6 (17.4-27.7)	103 (77.7-149)	225 (185-271)	1070
	05-06	4.96 (4.33-5.69)	3.97 (3.26-4.67)	21.5 (17.3-26.2)	101 (74.5-142)	227 (183-332)	1039
	07-08	4.86 (4.00-5.91)	3.85 (3.06-4.57)	22.2 (15.1-32.4)	124 (84.2-180)	355 (175-728)	993
	09-10	5.55 (4.77-6.47)	3.88 (3.19-4.87)	23.8 (19.2-32.5)	146 (97.6-248)	461 (330-739)	1186

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phytoestrogens_BiomonitoringSummary.html

Urinary 2-Hydroxyfluorene

Metabolite of Fluorene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	304 (262-354)	280 (242-319)	679 (561-815)	1850 (1430-2190)	2670 (2230-3130)	2521
	05-06	316 (297-338)	272 (243-296)	661 (605-743)	2060 (1600-2420)	3150 (2670-3730)	2431
	07-08	303 (271-338)	259 (231-293)	618 (489-817)	1630 (1340-2020)	2510 (2100-2890)	2581
Age group							
6-11 years	03-04	209 (183-239)	228 (189-259)	341 (295-411)	576 (423-728)	763 (613-827)	338
	05-06	201 (184-221)	195 (170-223)	352 (306-402)	497 (436-618)	709 (536-833)	343
	07-08	233 (205-264)	239 (194-280)	400 (326-454)	620 (484-714)	851 (695-1000)	382
12-19 years	03-04	281 (245-321)	292 (259-323)	502 (445-601)	1000 (753-1320)	1480 (1320-1930)	707
	05-06	283 (258-311)	272 (247-300)	482 (425-540)	1080 (843-1250)	1750 (1100-2820)	670
	07-08	287 (243-339)	255 (217-293)	541 (434-631)	963 (774-1350)	1510 (1050-2120)	395
20 years and older	03-04	323 (272-383)	290 (247-334)	832 (653-1020)	2090 (1680-2460)	2920 (2410-3650)	1476
	05-06	340 (311-371)	284 (252-317)	813 (705-917)	2410 (1880-2820)	3460 (2820-4160)	1418
	07-08	314 (276-357)	267 (231-306)	699 (518-1020)	1790 (1470-2220)	2700 (2190-3270)	1804
Gender							
Males	03-04	385 (333-446)	338 (300-415)	900 (739-1150)	2120 (1770-2450)	2930 (2490-3390)	1213
	05-06	376 (345-410)	332 (302-364)	823 (658-1030)	2160 (1660-2570)	3120 (2570-3560)	1213
	07-08	366 (334-402)	306 (280-348)	756 (613-983)	1740 (1430-2130)	2650 (2130-3270)	1280
Females	03-04	243 (204-290)	226 (184-269)	490 (405-617)	1400 (907-1930)	2310 (1740-3070)	1308
	05-06	268 (234-307)	227 (199-270)	528 (445-638)	1910 (1020-2830)	3180 (2180-4590)	1218
	07-08	252 (219-289)	216 (192-236)	505 (387-629)	1470 (1050-1960)	2330 (1770-2840)	1301
Race/ethnicity							
Mexican Americans	03-04	248 (215-286)	235 (203-293)	502 (412-580)	1100 (771-1330)	1640 (1250-2240)	629
	05-06	233 (200-271)	216 (180-247)	413 (334-488)	901 (704-1080)	1360 (1060-1950)	615
	07-08	248 (226-273)	240 (215-261)	418 (355-496)	801 (643-979)	1300 (933-1680)	523
Non-Hispanic blacks	03-04	432 (361-516)	381 (317-462)	856 (702-1270)	2200 (1830-2610)	2960 (2350-4460)	684
	05-06	404 (346-471)	330 (307-377)	805 (626-1090)	2190 (1610-2930)	3160 (2530-4330)	657
	07-08	387 (333-450)	325 (281-403)	879 (724-1040)	1940 (1500-2550)	3170 (2290-5120)	590
Non-Hispanic whites	03-04	308 (255-372)	280 (233-332)	728 (554-898)	1940 (1540-2390)	2920 (2400-3500)	1045
	05-06	319 (290-350)	270 (237-315)	705 (626-829)	2180 (1700-2540)	3390 (2670-4250)	972
	07-08	307 (261-362)	257 (221-308)	629 (452-995)	1720 (1260-2190)	2580 (2070-3020)	1070

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 5.0, 5.0, and 5.0 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Fluorene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 2-Hydroxyfluorene (creatinine corrected)

Metabolite of Fluorene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	286 (256-320)	221 (200-249)	495 (429-629)	1510 (1150-1800)	2070 (1850-2390)	2521
	05-06	308 (288-330)	227 (209-245)	567 (509-658)	1730 (1540-1850)	2450 (2020-2930)	2431
	07-08	304 (272-339)	234 (211-260)	526 (403-711)	1500 (1140-1780)	2150 (1850-2500)	2581
Age group							
6-11 years	03-04	221 (199-246)	217 (188-238)	305 (270-356)	462 (370-535)	695 (465-807)	338
	05-06	224 (207-243)	219 (197-254)	320 (288-365)	451 (384-523)	527 (450-720)	343
	07-08	287 (255-324)	262 (240-292)	407 (355-503)	721 (555-849)	864 (757-956)	382
12-19 years	03-04	213 (189-240)	189 (172-211)	310 (275-377)	649 (512-778)	937 (762-1320)	707
	05-06	214 (201-227)	189 (178-197)	286 (266-320)	607 (516-769)	1190 (803-1430)	670
	07-08	223 (200-249)	204 (176-236)	351 (270-419)	592 (504-683)	900 (683-1080)	395
20 years and older	03-04	310 (274-350)	233 (203-269)	659 (494-835)	1730 (1420-1940)	2310 (1930-2590)	1476
	05-06	339 (313-367)	241 (216-262)	754 (629-891)	1870 (1750-2110)	2830 (2250-3200)	1418
	07-08	321 (282-365)	234 (210-269)	633 (438-873)	1650 (1360-2060)	2390 (2090-2690)	1804
Gender							
Males	03-04	302 (271-337)	242 (218-272)	635 (494-758)	1510 (1200-1700)	1940 (1700-2310)	1213
	05-06	302 (278-328)	233 (212-259)	610 (510-744)	1480 (1240-1680)	1920 (1680-2390)	1213
	07-08	308 (283-336)	239 (218-261)	615 (464-764)	1480 (1200-1640)	2120 (1780-2390)	1280
Females	03-04	271 (235-314)	208 (183-244)	415 (352-521)	1530 (991-1890)	2220 (1730-2590)	1308
	05-06	314 (284-347)	217 (198-247)	526 (437-622)	1900 (1680-2240)	2930 (2310-3280)	1218
	07-08	300 (257-350)	225 (201-265)	461 (347-683)	1510 (929-2100)	2250 (1630-2770)	1301
Race/ethnicity							
Mexican Americans	03-04	224 (197-255)	198 (178-227)	370 (306-432)	745 (562-1030)	1200 (745-1730)	629
	05-06	210 (190-232)	179 (155-205)	294 (250-368)	624 (439-834)	1000 (720-1190)	615
	07-08	244 (216-275)	214 (187-238)	364 (301-438)	718 (582-824)	958 (775-1480)	523
Non-Hispanic blacks	03-04	308 (259-365)	249 (210-305)	609 (430-803)	1430 (939-1870)	1880 (1430-2500)	684
	05-06	284 (250-322)	222 (202-245)	502 (350-714)	1230 (941-1760)	1880 (1510-2440)	657
	07-08	299 (254-351)	237 (198-287)	602 (468-768)	1370 (1050-1910)	2220 (1570-2990)	590
Non-Hispanic whites	03-04	304 (266-349)	226 (199-261)	581 (449-796)	1630 (1410-1940)	2310 (1940-2540)	1045
	05-06	335 (307-366)	242 (216-270)	687 (561-806)	1840 (1650-2060)	2800 (2070-3260)	972
	07-08	319 (274-372)	238 (205-290)	579 (393-841)	1620 (1180-2090)	2340 (1940-2640)	1070

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Fluorene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 3-Hydroxyfluorene

Metabolite of Fluorene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	134 (115-155)	111 (96.0-126)	253 (207-349)	959 (666-1300)	1620 (1390-1900)	2745
	03-04	126 (108-148)	103 (90.4-118)	302 (231-404)	1090 (934-1270)	1740 (1400-2070)	2502
	05-06	124 (115-134)	99.1 (88.5-110)	304 (268-365)	1060 (830-1210)	1780 (1500-1910)	2398
	07-08	116 (103-132)	92.9 (85.7-100)	258 (203-341)	821 (681-1050)	1380 (1170-1630)	2581
Age group							
6-11 years	01-02	106 (94.8-119)	106 (91.0-124)	174 (145-200)	287 (235-358)	385 (306-455)	387
	03-04	89.2 (77.3-103)	92.3 (79.7-103)	151 (113-172)	238 (184-331)	343 (241-423)	336
	05-06	81.7 (73.6-90.7)	80.6 (68.3-98.9)	140 (122-174)	235 (189-272)	287 (242-345)	336
	07-08	98.0 (85.8-112)	94.2 (84.3-106)	178 (149-198)	302 (239-332)	458 (247-752)	382
12-19 years	01-02	129 (103-161)	113 (97.0-137)	222 (177-266)	542 (325-1070)	1210 (680-2130)	733
	03-04	116 (100-134)	114 (94.7-133)	214 (181-269)	476 (317-748)	924 (609-1410)	701
	05-06	119 (107-133)	112 (94.4-126)	212 (181-229)	588 (436-659)	986 (609-1800)	661
	07-08	114 (95.5-136)	93.7 (77.4-114)	218 (168-258)	494 (341-740)	830 (725-1020)	400
20 years and older	01-02	138 (119-161)	111 (94.0-129)	311 (228-429)	1130 (823-1400)	1850 (1470-2080)	1625
	03-04	134 (112-160)	105 (86.5-125)	408 (300-537)	1240 (1050-1480)	1910 (1620-2290)	1465
	05-06	131 (118-145)	101 (86.2-117)	394 (308-448)	1220 (968-1510)	1900 (1620-2050)	1401
	07-08	119 (103-138)	91.4 (83.5-100)	302 (205-449)	992 (745-1200)	1550 (1230-1820)	1799
Gender							
Males	01-02	163 (137-194)	134 (115-155)	352 (242-481)	1110 (721-1500)	1850 (1390-2190)	1346
	03-04	165 (140-194)	133 (108-162)	458 (313-681)	1270 (1030-1530)	1920 (1580-2120)	1205
	05-06	155 (138-174)	127 (118-136)	420 (304-510)	1130 (887-1330)	1810 (1440-2050)	1196
	07-08	146 (131-162)	115 (103-132)	330 (262-450)	944 (748-1190)	1530 (1160-1940)	1284
Females	01-02	111 (95.0-130)	94.0 (83.0-108)	204 (169-250)	796 (498-1040)	1390 (1250-1670)	1399
	03-04	98.2 (81.2-119)	80.5 (66.7-101)	195 (163-272)	842 (530-1200)	1590 (1180-1800)	1297
	05-06	99.9 (87.0-115)	78.2 (68.3-88.3)	225 (176-278)	961 (558-1390)	1720 (1300-2020)	1202
	07-08	93.9 (81.2-109)	77.2 (70.4-83.8)	192 (134-276)	688 (457-1090)	1280 (1030-1590)	1297
Race/ethnicity							
Mexican Americans	01-02	108 (87.1-134)	99.0 (81.0-119)	190 (144-255)	476 (302-641)	718 (476-1140)	662
	03-04	93.4 (79.6-110)	83.4 (67.0-96.9)	190 (150-252)	520 (388-726)	1030 (688-1320)	622
	05-06	82.5 (71.0-95.8)	71.8 (60.2-83.9)	143 (116-173)	435 (265-716)	763 (565-1020)	596
	07-08	90.4 (79.6-103)	81.2 (74.5-92.4)	160 (136-192)	363 (259-513)	674 (454-1180)	529
Non-Hispanic blacks	01-02	203 (169-244)	162 (135-188)	449 (308-830)	1420 (1130-1780)	2350 (1520-3000)	692
	03-04	195 (157-242)	160 (124-187)	482 (303-827)	1460 (1070-1800)	2110 (1580-2830)	683
	05-06	170 (143-202)	134 (121-149)	345 (246-558)	1300 (929-1760)	1880 (1550-2370)	652
	07-08	164 (142-190)	133 (112-157)	393 (303-494)	1290 (963-1420)	1770 (1330-2660)	587
Non-Hispanic whites	01-02	130 (108-157)	108 (93.0-127)	246 (202-352)	948 (621-1320)	1620 (1320-1990)	1207
	03-04	129 (106-155)	105 (87.6-124)	313 (237-464)	1200 (1010-1410)	1870 (1430-2220)	1035
	05-06	123 (111-137)	96.8 (85.4-112)	343 (275-420)	1060 (760-1260)	1870 (1390-2050)	966
	07-08	118 (98.0-141)	91.6 (81.6-103)	274 (195-403)	857 (646-1190)	1370 (1050-1740)	1066

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06 and 07-08 are 2.0, 5.0, 5.0 and 5.0 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Fluorene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 3-Hydroxyfluorene (creatinine corrected)

Metabolite of Fluorene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	125 (108-144)	94.4 (83.3-106)	220 (174-303)	754 (622-873)	1060 (909-1290)	2745
	03-04	119 (106-134)	86.6 (76.8-97.3)	256 (201-337)	856 (668-1060)	1330 (1130-1590)	2502
	05-06	121 (111-131)	84.5 (77.5-93.4)	245 (203-297)	930 (806-1020)	1280 (1150-1400)	2398
	07-08	117 (104-132)	84.6 (76.8-98.5)	226 (177-310)	789 (601-1000)	1230 (1040-1410)	2581
Age group							
6-11 years	01-02	119 (103-137)	110 (96.7-121)	153 (135-197)	265 (206-377)	382 (254-631)	387
	03-04	94.6 (84.3-106)	92.4 (83.1-98.5)	130 (115-154)	202 (168-282)	311 (197-435)	336
	05-06	90.7 (82.6-99.6)	90.3 (80.1-99.7)	130 (119-142)	210 (167-242)	290 (237-353)	336
	07-08	120 (105-138)	108 (92.3-127)	186 (150-215)	330 (231-401)	419 (304-918)	382
12-19 years	01-02	99.3 (81.6-121)	81.9 (74.2-92.6)	144 (109-197)	390 (214-700)	711 (372-1380)	733
	03-04	88.1 (78.0-99.6)	79.0 (72.5-83.2)	137 (114-170)	319 (225-440)	586 (356-880)	701
	05-06	90.1 (84.3-96.3)	73.8 (70.0-81.7)	133 (112-153)	381 (323-426)	618 (437-819)	661
	07-08	88.5 (78.9-99.2)	74.8 (65.2-84.6)	141 (118-185)	317 (242-449)	515 (338-656)	400
20 years and older	01-02	131 (113-151)	94.4 (81.0-108)	279 (195-367)	862 (727-923)	1210 (1010-1350)	1625
	03-04	128 (113-146)	88.2 (76.8-104)	364 (242-484)	1020 (828-1240)	1500 (1290-1710)	1465
	05-06	131 (119-144)	85.9 (77.1-98.5)	334 (251-476)	1020 (931-1180)	1380 (1220-1550)	1401
	07-08	122 (106-142)	83.9 (76.0-99.9)	276 (186-389)	926 (710-1110)	1320 (1150-1500)	1799
Gender							
Males	01-02	132 (111-156)	100 (88.5-117)	260 (194-361)	745 (560-914)	1130 (862-1380)	1346
	03-04	129 (114-147)	95.8 (82.5-111)	338 (226-456)	833 (672-1010)	1220 (980-1510)	1205
	05-06	125 (112-139)	92.4 (84.3-103)	272 (215-404)	784 (660-936)	1100 (874-1310)	1196
	07-08	123 (111-136)	92.3 (80.8-105)	282 (227-354)	791 (675-969)	1110 (989-1330)	1284
Females	01-02	119 (104-136)	88.9 (78.6-103)	179 (156-227)	777 (604-888)	1030 (923-1270)	1399
	03-04	110 (93.4-129)	81.7 (69.5-91.5)	183 (132-285)	920 (488-1270)	1460 (1130-1750)	1297
	05-06	117 (105-130)	77.6 (71.9-90.2)	203 (160-290)	1040 (836-1240)	1400 (1220-1780)	1202
	07-08	112 (95.7-131)	82.1 (72.7-96.8)	180 (132-265)	789 (450-1140)	1240 (1050-1500)	1297
Race/ethnicity							
Mexican Americans	01-02	101 (84.2-121)	84.0 (74.8-97.2)	159 (124-214)	364 (227-598)	604 (357-1230)	662
	03-04	84.8 (73.1-98.4)	68.8 (60.0-80.2)	170 (119-213)	353 (319-435)	668 (353-1050)	622
	05-06	74.0 (66.0-83.1)	59.9 (52.5-69.7)	112 (88.0-136)	282 (217-410)	585 (320-763)	596
	07-08	88.1 (75.3-103)	72.7 (62.2-84.1)	141 (107-184)	366 (232-536)	568 (397-722)	529
Non-Hispanic blacks	01-02	143 (118-173)	110 (90.7-133)	316 (186-525)	849 (622-1230)	1240 (882-1430)	692
	03-04	139 (113-169)	106 (93.1-118)	328 (211-488)	961 (580-1310)	1310 (961-1770)	683
	05-06	119 (103-138)	87.4 (78.4-97.0)	214 (154-388)	711 (523-1000)	1100 (838-1380)	652
	07-08	128 (109-150)	94.1 (82.2-110)	309 (204-433)	829 (715-981)	1310 (981-1640)	587
Non-Hispanic whites	01-02	128 (108-152)	94.7 (83.4-109)	223 (171-323)	816 (642-914)	1150 (910-1410)	1207
	03-04	127 (110-146)	89.8 (79.1-104)	311 (212-418)	980 (768-1180)	1460 (1270-1700)	1035
	05-06	130 (117-144)	91.1 (81.0-104)	290 (229-426)	973 (836-1180)	1380 (1130-1570)	966
	07-08	122 (103-145)	89.3 (76.2-108)	244 (166-377)	895 (579-1110)	1250 (1080-1470)	1066

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Fluorene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 9-Hydroxyfluorene

Metabolite of Fluorene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	267 (234-305)	269 (233-313)	541 (463-620)	929 (839-1060)	1390 (1130-1600)	2504
	05-06	308 (289-328)	300 (271-325)	609 (563-652)	1270 (1090-1420)	1820 (1560-2110)	2422
	07-08	337 (310-367)	327 (295-367)	677 (582-769)	1300 (1060-1490)	1960 (1670-2420)	2608
Age group							
6-11 years	03-04	209 (184-238)	216 (173-240)	331 (267-458)	594 (446-809)	853 (553-1500)	333
	05-06	184 (162-209)	190 (153-210)	324 (261-397)	657 (448-783)	823 (671-1220)	342
	07-08	242 (208-282)	229 (196-280)	449 (326-580)	791 (565-1040)	1090 (787-1610)	389
12-19 years	03-04	253 (216-297)	271 (215-324)	506 (413-560)	894 (773-988)	1210 (929-1510)	698
	05-06	266 (245-290)	278 (253-314)	528 (454-574)	842 (738-1020)	1290 (1070-1510)	668
	07-08	308 (255-371)	295 (241-374)	583 (459-736)	1050 (829-1290)	1400 (1080-1870)	401
20 years and older	03-04	277 (240-320)	279 (237-335)	583 (476-656)	979 (842-1140)	1490 (1140-1770)	1473
	05-06	335 (309-363)	323 (292-349)	653 (593-729)	1360 (1180-1610)	1950 (1650-2530)	1412
	07-08	355 (323-390)	352 (311-384)	716 (629-835)	1410 (1170-1670)	2160 (1770-2780)	1818
Gender							
Males	03-04	330 (291-374)	330 (285-372)	632 (583-709)	1090 (898-1370)	1720 (1360-2250)	1208
	05-06	355 (325-387)	334 (296-387)	693 (608-783)	1310 (1150-1620)	2000 (1670-2560)	1208
	07-08	401 (365-441)	394 (356-442)	778 (661-897)	1420 (1260-1580)	2120 (1770-2820)	1294
Females	03-04	218 (188-253)	226 (191-253)	439 (357-521)	833 (697-973)	1100 (955-1310)	1296
	05-06	269 (242-299)	257 (230-297)	503 (426-580)	1090 (845-1400)	1610 (1400-1920)	1214
	07-08	286 (260-313)	271 (246-299)	557 (474-671)	1110 (914-1400)	1830 (1400-2400)	1314
Race/ethnicity							
Mexican Americans	03-04	229 (200-261)	232 (197-274)	435 (362-492)	773 (561-1010)	1030 (849-1170)	614
	05-06	232 (204-264)	221 (186-262)	407 (361-435)	738 (596-840)	1030 (788-1470)	614
	07-08	267 (246-288)	259 (236-285)	476 (430-515)	859 (732-1100)	1470 (1070-1980)	532
Non-Hispanic blacks	03-04	380 (311-465)	357 (291-475)	709 (559-825)	1260 (1080-1600)	1880 (1560-2330)	679
	05-06	416 (360-481)	406 (366-450)	716 (593-943)	1430 (1010-2250)	2310 (1590-3410)	655
	07-08	426 (376-482)	423 (356-477)	817 (693-978)	1490 (1200-1950)	2260 (1870-3040)	597
Non-Hispanic whites	03-04	266 (226-313)	266 (228-324)	558 (453-659)	939 (824-1130)	1450 (1090-1770)	1048
	05-06	306 (279-335)	296 (257-331)	613 (548-669)	1290 (1150-1450)	1730 (1530-2140)	966
	07-08	345 (302-394)	338 (286-398)	700 (564-849)	1380 (1030-1690)	2040 (1570-2650)	1080

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 5.0, 5.0, and 5.0 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Fluorene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 9-Hydroxyfluorene (creatinine corrected)

Metabolite of Fluorene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	252 (230-276)	233 (205-266)	412 (371-461)	729 (609-905)	1100 (918-1390)	2504
	05-06	301 (281-322)	270 (252-290)	511 (460-575)	1040 (908-1230)	1510 (1360-1750)	2422
	07-08	339 (309-372)	303 (277-333)	562 (493-645)	1110 (908-1300)	1660 (1430-1960)	2608
Age group							
6-11 years	03-04	223 (200-247)	208 (176-264)	343 (299-389)	548 (433-689)	866 (533-958)	333
	05-06	205 (180-234)	195 (176-209)	309 (272-371)	590 (402-668)	866 (620-1300)	342
	07-08	298 (256-346)	280 (249-331)	463 (399-576)	817 (576-1130)	1130 (862-1230)	389
12-19 years	03-04	192 (172-215)	182 (161-200)	323 (286-360)	484 (412-696)	738 (571-1110)	698
	05-06	201 (183-220)	194 (173-223)	325 (283-343)	511 (397-666)	848 (521-941)	668
	07-08	239 (204-280)	235 (190-283)	381 (312-462)	596 (479-722)	768 (660-1090)	401
20 years and older	03-04	267 (242-294)	243 (219-280)	446 (392-490)	788 (641-994)	1280 (991-1560)	1473
	05-06	336 (314-358)	304 (284-323)	573 (506-650)	1200 (1000-1360)	1720 (1490-1900)	1412
	07-08	363 (330-399)	314 (293-351)	618 (533-738)	1200 (1030-1450)	1770 (1570-2040)	1818
Gender							
Males	03-04	260 (238-283)	236 (209-269)	429 (382-480)	739 (591-952)	1300 (916-1800)	1208
	05-06	286 (266-308)	258 (235-291)	498 (422-576)	993 (858-1180)	1410 (1240-1730)	1208
	07-08	338 (308-372)	303 (269-338)	575 (516-626)	1150 (987-1300)	1690 (1420-1940)	1294
Females	03-04	244 (217-275)	227 (198-269)	396 (342-467)	717 (569-905)	1020 (905-1280)	1296
	05-06	316 (287-348)	278 (253-303)	532 (466-623)	1100 (840-1370)	1610 (1360-1840)	1214
	07-08	340 (306-377)	303 (274-335)	555 (451-683)	1030 (852-1330)	1620 (1330-2040)	1314
Race/ethnicity							
Mexican Americans	03-04	209 (188-232)	195 (181-209)	335 (293-379)	541 (447-722)	793 (569-1440)	614
	05-06	208 (194-224)	201 (180-220)	304 (270-342)	502 (425-584)	663 (527-919)	614
	07-08	260 (236-285)	244 (226-259)	388 (348-419)	654 (527-858)	1160 (775-1640)	532
Non-Hispanic blacks	03-04	272 (226-326)	258 (208-307)	463 (349-568)	777 (628-920)	1040 (863-1300)	679
	05-06	293 (258-332)	261 (230-297)	449 (377-554)	851 (635-1080)	1480 (925-2070)	655
	07-08	331 (289-379)	293 (257-333)	542 (489-620)	990 (805-1250)	1550 (1160-2470)	597
Non-Hispanic whites	03-04	263 (235-295)	244 (211-287)	439 (387-490)	773 (607-1020)	1290 (967-1700)	1048
	05-06	323 (296-353)	293 (266-322)	570 (496-650)	1180 (962-1360)	1630 (1370-1840)	966
	07-08	359 (316-407)	314 (280-366)	605 (497-750)	1170 (935-1420)	1740 (1460-2040)	1080

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Fluorene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 1-Hydroxyphenanthrene

Metabolite of Phenanthrene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	140 (125-158)	141 (130-154)	266 (229-312)	476 (426-539)	684 (581-763)	2741
	03-04	156 (140-173)	166 (150-179)	287 (263-321)	464 (423-508)	625 (538-745)	2496
	05-06	152 (143-161)	151 (138-163)	275 (257-303)	493 (428-549)	669 (580-852)	2449
	07-08	139 (130-149)	140 (127-154)	252 (228-272)	442 (402-481)	683 (577-760)	2608
Age group							
6-11 years	01-02	119 (104-137)	121 (98.0-142)	228 (182-249)	357 (265-501)	501 (367-661)	387
	03-04	138 (124-154)	131 (111-161)	222 (200-275)	397 (304-500)	615 (384-916)	328
	05-06	111 (101-123)	110 (96.9-121)	176 (150-240)	320 (251-375)	381 (329-540)	345
	07-08	121 (105-139)	120 (96.5-154)	200 (174-235)	334 (247-401)	405 (278-557)	389
12-19 years	01-02	133 (110-162)	131 (110-153)	238 (198-303)	431 (376-546)	579 (439-820)	733
	03-04	158 (141-178)	171 (138-193)	300 (262-333)	470 (437-537)	654 (563-745)	692
	05-06	144 (132-157)	154 (130-175)	260 (236-286)	425 (383-496)	666 (496-845)	677
	07-08	136 (122-152)	144 (128-164)	219 (199-283)	374 (315-415)	503 (392-607)	401
20 years and older	01-02	145 (127-164)	145 (133-164)	274 (237-329)	499 (440-565)	713 (595-819)	1621
	03-04	157 (141-176)	167 (152-182)	296 (264-337)	469 (430-513)	625 (534-783)	1476
	05-06	159 (147-172)	156 (141-172)	287 (265-328)	516 (438-615)	744 (580-996)	1427
	07-08	142 (132-153)	141 (125-158)	258 (241-280)	470 (417-531)	733 (622-856)	1818
Gender							
Males	01-02	150 (133-169)	145 (132-164)	284 (235-348)	501 (424-593)	713 (575-845)	1344
	03-04	176 (161-194)	182 (167-198)	321 (283-363)	492 (459-536)	662 (541-929)	1196
	05-06	169 (156-183)	166 (153-180)	289 (265-334)	519 (438-593)	791 (562-1080)	1222
	07-08	153 (140-167)	152 (137-171)	268 (251-292)	448 (406-488)	722 (566-813)	1294
Females	01-02	132 (115-152)	137 (121-151)	254 (221-297)	464 (390-520)	654 (538-769)	1397
	03-04	138 (122-156)	144 (126-164)	263 (240-294)	430 (404-469)	589 (489-680)	1300
	05-06	137 (123-152)	136 (117-153)	260 (234-297)	460 (391-553)	648 (540-809)	1227
	07-08	128 (120-136)	128 (117-139)	230 (205-258)	432 (373-500)	663 (527-740)	1314
Race/ethnicity							
Mexican Americans	01-02	117 (90.8-152)	116 (88.0-147)	214 (157-306)	369 (254-583)	549 (342-847)	664
	03-04	138 (120-158)	152 (129-175)	246 (213-291)	413 (311-492)	518 (352-896)	611
	05-06	125 (104-151)	129 (107-151)	223 (174-276)	376 (298-460)	523 (399-617)	621
	07-08	118 (108-129)	108 (103-119)	194 (172-216)	341 (271-437)	491 (417-738)	532
Non-Hispanic blacks	01-02	150 (127-179)	145 (126-170)	287 (243-327)	493 (423-629)	713 (568-975)	690
	03-04	182 (152-217)	178 (149-212)	346 (274-418)	552 (459-645)	797 (606-1050)	679
	05-06	162 (148-177)	155 (141-173)	307 (259-349)	517 (408-695)	753 (545-1110)	658
	07-08	142 (129-158)	141 (128-163)	264 (234-302)	448 (401-511)	613 (522-722)	597
Non-Hispanic whites	01-02	144 (125-166)	144 (131-162)	276 (241-330)	489 (436-552)	661 (550-793)	1204
	03-04	159 (140-180)	168 (151-186)	296 (265-330)	475 (421-539)	641 (528-843)	1046
	05-06	156 (143-170)	155 (137-173)	279 (251-322)	500 (414-606)	743 (570-1050)	982
	07-08	146 (133-159)	149 (132-174)	261 (235-292)	453 (405-503)	725 (590-836)	1080

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06 and 07-08 are 3.5, 5.0, 5.0 and 5.0 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phenanthrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 1-Hydroxyphenanthrene (creatinine corrected)

Metabolite of Phenanthrene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	132 (118-147)	125 (113-141)	210 (191-231)	344 (310-385)	464 (404-539)	2741
	03-04	146 (138-155)	141 (133-150)	222 (205-244)	352 (330-384)	487 (416-546)	2496
	05-06	147 (138-157)	140 (130-148)	229 (211-247)	372 (339-406)	502 (446-600)	2449
	07-08	140 (130-151)	137 (124-151)	221 (207-236)	345 (319-372)	457 (407-522)	2608
Age group							
6-11 years	01-02	133 (116-153)	126 (110-147)	188 (165-225)	344 (245-437)	467 (364-598)	387
	03-04	145 (129-163)	138 (121-155)	205 (169-240)	315 (244-456)	477 (313-803)	328
	05-06	124 (114-134)	115 (103-133)	191 (162-210)	275 (230-340)	368 (321-396)	345
	07-08	149 (132-169)	139 (127-158)	221 (191-264)	334 (275-410)	424 (332-796)	389
12-19 years	01-02	103 (87.9-121)	97.3 (81.4-117)	158 (131-192)	240 (198-322)	354 (235-531)	733
	03-04	119 (108-130)	115 (101-131)	176 (157-197)	275 (230-331)	345 (269-496)	692
	05-06	108 (97.5-119)	108 (95.9-116)	163 (148-180)	268 (199-330)	372 (262-609)	677
	07-08	106 (94.9-117)	106 (90.0-116)	158 (147-171)	243 (197-303)	324 (250-407)	401
20 years and older	01-02	137 (122-153)	132 (117-145)	223 (200-243)	351 (319-395)	476 (421-541)	1621
	03-04	151 (142-161)	147 (134-157)	237 (212-254)	365 (335-397)	500 (422-580)	1476
	05-06	158 (147-170)	151 (137-163)	244 (224-265)	397 (362-420)	554 (478-666)	1427
	07-08	145 (135-157)	142 (127-156)	229 (214-251)	354 (336-383)	495 (420-560)	1818
Gender							
Males	01-02	122 (108-137)	116 (104-131)	194 (173-218)	321 (262-395)	455 (348-642)	1344
	03-04	138 (131-146)	130 (117-143)	201 (187-227)	335 (301-391)	437 (392-550)	1196
	05-06	135 (126-145)	125 (117-132)	208 (187-229)	341 (300-385)	498 (391-634)	1222
	07-08	129 (117-141)	126 (112-141)	207 (191-223)	327 (290-344)	410 (354-487)	1294
Females	01-02	142 (125-160)	136 (118-156)	226 (199-256)	355 (323-399)	473 (421-541)	1397
	03-04	154 (142-168)	150 (138-160)	241 (212-269)	364 (323-425)	505 (418-635)	1300
	05-06	160 (149-172)	152 (136-166)	251 (229-275)	398 (367-428)	509 (435-638)	1227
	07-08	152 (140-165)	148 (133-162)	231 (213-262)	373 (329-411)	478 (409-577)	1314
Race/ethnicity							
Mexican Americans	01-02	110 (88.6-136)	101 (85.5-131)	162 (138-203)	275 (187-452)	400 (268-755)	664
	03-04	124 (111-140)	122 (107-136)	183 (159-242)	304 (246-392)	392 (318-437)	611
	05-06	112 (99.2-126)	109 (87.1-129)	170 (145-201)	259 (228-307)	353 (297-414)	621
	07-08	115 (102-129)	107 (96.7-118)	159 (143-183)	287 (218-344)	424 (324-520)	532
Non-Hispanic blacks	01-02	106 (88.7-127)	102 (92.4-111)	172 (141-203)	274 (224-356)	384 (308-598)	690
	03-04	129 (112-148)	126 (108-143)	199 (173-242)	337 (272-400)	428 (374-522)	679
	05-06	114 (106-121)	107 (98.6-114)	179 (163-189)	301 (229-370)	449 (321-771)	658
	07-08	111 (98.8-124)	109 (90.0-126)	181 (153-201)	293 (257-338)	385 (338-446)	597
Non-Hispanic whites	01-02	142 (126-161)	137 (119-153)	228 (206-247)	363 (321-406)	476 (411-552)	1204
	03-04	157 (146-168)	150 (140-159)	241 (216-256)	372 (335-424)	515 (426-658)	1046
	05-06	164 (152-176)	154 (141-165)	247 (228-268)	398 (366-428)	554 (466-739)	982
	07-08	152 (139-166)	152 (135-167)	236 (217-258)	354 (330-383)	464 (407-551)	1080

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phenanthrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 2-Hydroxyphenanthrene

Metabolite of Phenanthrene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	54.0 (46.0-63.5)	58.0 (50.0-68.0)	117 (102-140)	240 (201-271)	332 (299-377)	2742
	03-04	59.3 (52.5-66.9)	62.2 (55.4-69.7)	117 (105-129)	206 (180-235)	291 (252-336)	2512
	05-06	65.3 (60.9-70.0)	63.3 (58.5-68.6)	126 (111-139)	234 (203-254)	314 (282-383)	2410
	07-08	63.6 (58.8-68.7)	62.9 (58.0-68.7)	117 (104-130)	204 (184-223)	300 (256-367)	2572
Age group							
6-11 years	01-02	40.5 (34.3-47.7)	46.0 (37.0-57.0)	87.0 (71.0-101)	170 (133-207)	257 (195-320)	387
	03-04	45.3 (40.0-51.3)	48.1 (41.9-56.8)	72.7 (63.2-89.4)	135 (104-176)	225 (136-299)	337
	05-06	42.1 (38.4-46.1)	40.1 (36.4-47.2)	70.4 (58.1-83.2)	110 (89.2-132)	152 (112-213)	341
	07-08	48.1 (41.8-55.4)	46.4 (41.7-54.6)	80.3 (68.2-95.4)	144 (98.8-184)	184 (122-229)	382
12-19 years	01-02	49.5 (37.3-65.7)	52.0 (41.0-68.0)	108 (93.0-122)	210 (145-270)	281 (214-524)	733
	03-04	58.0 (50.7-66.2)	62.5 (54.5-71.6)	104 (88.8-132)	198 (166-215)	258 (199-318)	707
	05-06	60.4 (55.4-65.8)	62.4 (56.3-67.1)	114 (93.8-133)	175 (156-195)	250 (198-291)	664
	07-08	59.4 (52.9-66.7)	60.5 (51.5-67.0)	97.4 (81.2-121)	160 (133-204)	223 (175-264)	398
20 years and older	01-02	56.8 (48.1-66.9)	60.0 (52.0-73.0)	126 (104-152)	249 (207-292)	342 (308-398)	1622
	03-04	61.5 (53.7-70.4)	64.9 (57.2-73.5)	125 (111-138)	216 (187-250)	310 (259-389)	1468
	05-06	69.6 (63.6-76.2)	67.3 (61.5-75.3)	137 (118-154)	251 (215-280)	348 (298-446)	1405
	07-08	66.2 (60.8-72.1)	65.9 (60.4-71.2)	123 (109-136)	218 (195-251)	334 (272-412)	1792
Gender							
Males	01-02	62.1 (53.3-72.5)	67.0 (58.0-80.0)	136 (109-161)	274 (245-303)	367 (329-414)	1345
	03-04	72.3 (64.4-81.3)	71.6 (65.8-83.3)	138 (123-150)	240 (200-279)	337 (279-428)	1211
	05-06	76.5 (70.3-83.3)	72.9 (67.4-81.3)	141 (122-156)	254 (219-296)	383 (289-555)	1204
	07-08	75.8 (70.3-81.8)	73.5 (67.6-81.4)	136 (126-149)	229 (199-256)	354 (276-408)	1277
Females	01-02	47.4 (39.1-57.5)	50.0 (41.0-60.0)	105 (88.0-129)	200 (171-240)	294 (236-357)	1397
	03-04	49.0 (42.4-56.6)	52.5 (45.3-58.7)	95.4 (84.1-110)	166 (146-202)	247 (202-291)	1301
	05-06	56.0 (50.2-62.5)	54.4 (49.5-61.0)	107 (94.7-125)	206 (173-245)	275 (248-343)	1206
	07-08	53.7 (48.9-59.0)	53.0 (48.9-59.2)	97.0 (87.0-111)	173 (146-207)	280 (206-327)	1295
Race/ethnicity							
Mexican Americans	01-02	46.8 (32.7-66.8)	51.0 (36.0-73.0)	97.0 (72.0-140)	191 (122-332)	303 (187-652)	665
	03-04	53.9 (46.7-62.1)	60.1 (50.8-68.6)	104 (89.5-115)	164 (144-187)	200 (187-219)	627
	05-06	54.6 (45.0-66.3)	56.2 (47.1-67.7)	96.2 (84.4-108)	164 (120-202)	213 (175-294)	612
	07-08	58.2 (53.9-62.7)	59.3 (50.8-63.0)	98.6 (92.5-113)	167 (141-193)	252 (192-395)	528
Non-Hispanic blacks	01-02	71.1 (58.0-87.1)	74.0 (66.0-90.0)	152 (124-182)	262 (217-311)	374 (284-560)	690
	03-04	81.2 (68.1-96.7)	73.4 (61.8-101)	158 (125-199)	280 (219-374)	390 (289-574)	678
	05-06	77.3 (69.0-86.6)	75.3 (65.8-83.5)	141 (114-172)	267 (202-340)	363 (275-546)	641
	07-08	72.8 (65.2-81.3)	70.7 (62.6-80.2)	127 (115-149)	251 (206-284)	367 (284-455)	597
Non-Hispanic whites	01-02	53.1 (43.6-64.6)	57.0 (48.0-68.0)	117 (99.0-144)	242 (199-286)	333 (303-385)	1204
	03-04	58.2 (50.1-67.7)	60.7 (52.5-71.0)	115 (97.9-133)	209 (172-255)	299 (251-351)	1044
	05-06	65.2 (59.4-71.5)	63.7 (57.0-70.4)	130 (109-144)	244 (206-261)	343 (282-439)	974
	07-08	63.3 (56.6-70.8)	64.0 (56.4-71.1)	121 (102-135)	203 (174-234)	298 (238-410)	1050

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06 and 07-08 are 3.2, 5.0, 5.0 and 5.0 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phenanthrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 2-Hydroxyphenanthrene (creatinine corrected)

Metabolite of Phenanthrene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	50.6 (43.3-59.2)	52.5 (46.4-59.6)	91.4 (78.7-108)	164 (138-200)	233 (206-275)	2742
	03-04	55.4 (51.1-60.0)	52.3 (48.8-55.8)	85.8 (79.4-97.0)	150 (128-174)	212 (180-252)	2512
	05-06	63.1 (58.9-67.5)	57.0 (53.3-61.9)	101 (92.8-110)	178 (157-194)	245 (209-295)	2410
	07-08	64.0 (58.9-69.4)	60.3 (55.3-66.5)	98.2 (87.5-108)	163 (151-179)	229 (199-248)	2572
Age group							
6-11 years	01-02	45.3 (38.8-52.8)	50.0 (40.0-57.8)	85.9 (69.6-100)	144 (105-217)	234 (138-397)	387
	03-04	47.9 (43.6-52.7)	46.6 (41.0-51.4)	66.3 (60.3-72.8)	106 (81.0-125)	183 (108-295)	337
	05-06	45.9 (42.0-50.1)	43.6 (38.4-48.7)	65.8 (56.1-85.2)	101 (85.4-110)	116 (101-135)	341
	07-08	59.7 (51.2-69.4)	56.5 (49.4-65.2)	85.1 (73.1-107)	134 (105-187)	187 (121-437)	382
12-19 years	01-02	38.4 (29.6-49.7)	38.5 (30.6-48.7)	64.9 (52.4-84.0)	117 (86.7-167)	173 (128-305)	733
	03-04	43.6 (39.3-48.5)	40.8 (37.7-45.0)	70.6 (55.8-84.7)	106 (88.2-123)	127 (115-176)	707
	05-06	44.5 (39.9-49.7)	42.7 (38.9-47.1)	66.1 (58.0-73.7)	111 (80.0-129)	146 (115-232)	664
	07-08	46.1 (42.0-50.5)	47.0 (39.7-53.0)	69.7 (62.0-78.0)	100 (83.0-113)	127 (105-188)	398
20 years and older	01-02	53.7 (46.2-62.4)	55.6 (49.4-62.7)	94.6 (84.1-112)	173 (145-201)	241 (209-287)	1622
	03-04	58.7 (53.6-64.2)	54.8 (50.7-60.9)	91.6 (83.5-105)	164 (139-182)	227 (183-291)	1468
	05-06	69.1 (64.1-74.6)	63.6 (58.4-67.8)	111 (102-123)	191 (168-216)	267 (232-316)	1405
	07-08	67.8 (62.4-73.8)	64.1 (57.6-71.6)	105 (95.5-119)	177 (160-194)	239 (211-255)	1792
Gender							
Males	01-02	50.4 (43.3-58.5)	52.5 (47.2-59.2)	91.4 (78.6-103)	176 (140-208)	245 (206-349)	1345
	03-04	56.6 (52.5-61.0)	52.3 (48.1-57.1)	84.7 (79.2-96.4)	157 (132-180)	227 (184-284)	1211
	05-06	61.2 (56.7-65.9)	55.7 (50.0-59.3)	102 (89.3-111)	161 (137-194)	244 (190-310)	1204
	07-08	64.0 (59.3-69.1)	59.8 (55.3-64.1)	99.7 (90.5-108)	172 (152-191)	229 (211-243)	1277
Females	01-02	50.9 (42.7-60.7)	52.8 (45.5-61.0)	91.6 (75.0-115)	158 (133-183)	214 (200-245)	1397
	03-04	54.3 (48.7-60.5)	52.3 (47.0-57.2)	88.1 (76.4-101)	146 (118-179)	202 (175-246)	1301
	05-06	65.0 (60.0-70.3)	60.5 (55.1-65.3)	101 (91.6-113)	184 (165-201)	256 (224-312)	1206
	07-08	63.9 (57.8-70.7)	61.5 (54.0-69.3)	95.8 (83.7-115)	159 (140-189)	229 (186-256)	1295
Race/ethnicity							
Mexican Americans	01-02	43.7 (31.9-59.8)	44.0 (34.3-58.8)	80.8 (59.8-115)	146 (103-245)	245 (144-394)	665
	03-04	48.8 (42.5-55.9)	50.1 (42.4-58.8)	79.5 (68.2-86.1)	124 (96.2-149)	155 (127-206)	627
	05-06	48.7 (43.0-55.2)	46.0 (39.4-56.2)	75.1 (62.9-85.8)	111 (94.5-138)	146 (121-178)	612
	07-08	56.7 (51.1-62.8)	55.0 (47.7-62.2)	83.7 (76.2-90.4)	133 (107-180)	198 (145-233)	528
Non-Hispanic blacks	01-02	50.2 (40.5-62.2)	49.7 (44.3-56.0)	87.7 (72.6-103)	174 (127-224)	257 (178-459)	690
	03-04	57.6 (49.8-66.7)	53.1 (45.7-63.5)	90.3 (77.9-117)	152 (128-186)	218 (173-348)	678
	05-06	54.3 (49.7-59.3)	47.7 (45.4-51.3)	84.8 (71.3-97.8)	150 (116-192)	211 (167-298)	641
	07-08	56.6 (50.4-63.6)	51.2 (46.6-55.9)	95.5 (77.7-107)	154 (136-178)	212 (188-232)	597
Non-Hispanic whites	01-02	52.5 (43.8-62.9)	54.8 (47.7-61.9)	92.7 (80.5-113)	170 (140-204)	233 (202-293)	1204
	03-04	57.1 (51.7-63.0)	52.9 (49.1-57.2)	88.5 (79.2-102)	159 (128-187)	237 (180-295)	1044
	05-06	67.9 (62.6-73.7)	60.9 (56.5-66.5)	111 (101-123)	191 (162-224)	274 (231-323)	974
	07-08	65.9 (59.4-73.3)	62.8 (55.1-72.0)	100 (87.2-119)	166 (147-190)	229 (197-252)	1050

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phenanthrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 3-Hydroxyphenanthrene

Metabolite of Phenanthrene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	105 (92.5-118)	105 (92.0-118)	200 (179-225)	401 (336-480)	649 (542-747)	2741
	03-04	115 (104-128)	118 (107-129)	219 (201-237)	424 (351-487)	647 (509-788)	2426
	05-06	103 (96.1-109)	99.8 (90.3-107)	200 (185-210)	399 (365-429)	593 (463-746)	2448
	07-08	97.6 (91.6-104)	97.8 (90.9-106)	178 (164-195)	319 (283-353)	490 (426-557)	2601
Age group							
6-11 years	01-02	105 (91.1-122)	109 (82.0-138)	195 (160-231)	298 (245-346)	412 (319-545)	387
	03-04	116 (99.9-134)	107 (93.0-135)	205 (170-223)	333 (241-435)	472 (322-881)	325
	05-06	89.6 (79.6-101)	84.0 (72.6-103)	154 (123-176)	245 (193-316)	316 (245-569)	345
	07-08	100 (85.9-117)	102 (83.4-126)	175 (144-183)	251 (194-328)	348 (239-862)	389
12-19 years	01-02	104 (87.3-125)	107 (90.0-122)	201 (165-231)	329 (255-445)	459 (331-631)	733
	03-04	120 (103-141)	127 (107-149)	221 (195-262)	366 (327-424)	564 (391-730)	677
	05-06	105 (96.5-113)	104 (97.3-116)	193 (166-217)	323 (275-383)	453 (377-604)	676
	07-08	103 (92.9-114)	103 (89.5-116)	173 (146-195)	293 (246-352)	371 (323-436)	401
20 years and older	01-02	105 (91.7-119)	105 (89.0-118)	201 (180-231)	433 (366-515)	683 (597-806)	1621
	03-04	114 (103-128)	118 (105-130)	221 (203-243)	450 (353-505)	696 (511-885)	1424
	05-06	104 (95.3-113)	101 (89.6-113)	206 (188-223)	419 (384-466)	657 (481-832)	1427
	07-08	96.5 (89.3-104)	96.9 (87.3-106)	183 (163-201)	331 (285-393)	535 (432-624)	1811
Gender							
Males	01-02	122 (107-138)	117 (106-132)	224 (188-278)	474 (372-597)	734 (597-1010)	1344
	03-04	137 (122-154)	135 (119-157)	251 (221-287)	487 (398-579)	754 (606-913)	1167
	05-06	125 (114-137)	119 (107-136)	227 (204-261)	427 (395-491)	768 (520-951)	1222
	07-08	117 (109-127)	115 (103-130)	209 (191-229)	352 (322-390)	557 (462-674)	1290
Females	01-02	90.8 (77.7-106)	93.0 (80.0-108)	184 (155-207)	328 (291-399)	518 (434-649)	1397
	03-04	97.7 (86.5-110)	100 (86.6-114)	193 (170-210)	351 (287-445)	531 (447-647)	1259
	05-06	84.6 (75.6-94.5)	82.4 (73.1-92.5)	159 (143-176)	363 (285-404)	480 (421-577)	1226
	07-08	81.9 (76.8-87.3)	81.7 (74.7-91.0)	148 (134-173)	272 (237-326)	439 (355-501)	1311
Race/ethnicity							
Mexican Americans	01-02	83.5 (64.1-109)	84.0 (64.0-111)	144 (117-198)	259 (193-430)	454 (253-1140)	664
	03-04	92.7 (81.3-106)	92.8 (84.1-111)	164 (133-204)	317 (236-374)	436 (338-501)	576
	05-06	77.0 (64.1-92.5)	74.8 (60.0-90.8)	138 (110-162)	260 (191-312)	366 (299-471)	620
	07-08	79.1 (71.4-87.7)	75.3 (66.4-85.2)	133 (118-154)	250 (176-330)	414 (302-511)	529
Non-Hispanic blacks	01-02	145 (122-172)	135 (120-163)	281 (228-346)	516 (414-684)	957 (609-1410)	690
	03-04	166 (141-195)	159 (137-193)	313 (249-399)	542 (458-670)	798 (579-1200)	686
	05-06	138 (120-159)	125 (113-147)	260 (211-329)	466 (365-671)	716 (491-1100)	658
	07-08	129 (115-145)	129 (112-148)	237 (205-266)	427 (349-514)	657 (514-839)	596
Non-Hispanic whites	01-02	104 (90.8-120)	106 (92.0-120)	199 (177-224)	401 (331-495)	649 (518-747)	1204
	03-04	114 (100-129)	117 (104-128)	220 (201-237)	440 (332-509)	702 (505-915)	1010
	05-06	101 (93.6-110)	101 (89.3-112)	200 (181-213)	402 (368-434)	657 (449-829)	982
	07-08	97.0 (87.7-107)	97.9 (87.3-111)	179 (155-202)	315 (270-355)	468 (371-590)	1077

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06 and 07-08 are 3.6, 5.0, 5.0 and 5.0 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phenanthrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 3-Hydroxyphenanthrene (creatinine corrected)

Metabolite of Phenanthrene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	98.0 (87.4-110)	86.5 (78.8-97.1)	158 (142-174)	299 (252-342)	428 (365-621)	2741
	03-04	108 (101-115)	99.5 (91.9-106)	172 (158-189)	321 (273-363)	497 (406-566)	2426
	05-06	99.5 (93.6-106)	88.1 (83.8-93.5)	164 (151-175)	326 (273-364)	443 (402-525)	2448
	07-08	98.1 (91.3-106)	91.3 (81.8-100)	159 (141-174)	273 (246-301)	380 (333-442)	2601
Age group							
6-11 years	01-02	118 (103-135)	114 (94.9-131)	173 (150-201)	289 (238-368)	410 (308-544)	387
	03-04	124 (110-138)	119 (110-129)	165 (145-202)	283 (231-402)	484 (269-710)	325
	05-06	99.6 (89.4-111)	93.5 (85.7-107)	146 (118-165)	232 (171-292)	292 (232-352)	345
	07-08	123 (107-142)	120 (105-143)	183 (159-200)	230 (201-382)	382 (222-948)	389
12-19 years	01-02	81.1 (70.1-93.8)	78.0 (66.7-89.4)	116 (96.5-144)	205 (153-275)	283 (198-511)	733
	03-04	90.3 (81.5-100)	84.9 (76.1-93.9)	139 (116-160)	212 (193-231)	295 (225-429)	677
	05-06	78.2 (70.9-86.3)	72.5 (66.7-82.1)	111 (98.8-121)	183 (141-262)	291 (211-465)	676
	07-08	80.0 (73.6-86.9)	74.4 (67.0-87.5)	117 (110-126)	188 (151-223)	243 (213-284)	401
20 years and older	01-02	98.9 (88.3-111)	85.3 (78.3-96.6)	164 (147-183)	320 (273-356)	488 (379-774)	1621
	03-04	109 (102-117)	99.1 (91.7-106)	180 (161-198)	343 (282-385)	523 (417-608)	1424
	05-06	103 (96.3-111)	90.3 (84.4-97.8)	177 (161-200)	350 (319-387)	506 (426-538)	1427
	07-08	98.8 (91.2-107)	90.3 (80.2-100)	163 (140-184)	290 (262-321)	388 (338-464)	1811
Gender							
Males	01-02	98.6 (87.7-111)	87.1 (79.5-98.2)	159 (138-188)	310 (245-368)	505 (356-761)	1344
	03-04	107 (98.2-116)	99.2 (87.4-109)	169 (154-184)	305 (261-343)	491 (383-592)	1167
	05-06	100 (92.6-109)	89.2 (81.5-97.0)	166 (147-184)	326 (260-365)	430 (364-553)	1222
	07-08	98.9 (91.2-107)	91.3 (81.3-101)	166 (146-184)	288 (263-310)	382 (321-449)	1290
Females	01-02	97.5 (84.9-112)	86.4 (76.9-100)	158 (139-178)	292 (245-333)	410 (352-583)	1397
	03-04	109 (99.6-119)	100 (88.9-110)	176 (155-202)	328 (265-398)	497 (380-594)	1259
	05-06	98.7 (91.6-106)	87.3 (80.4-95.1)	160 (144-177)	332 (284-375)	446 (393-518)	1226
	07-08	97.4 (89.6-106)	90.9 (79.4-102)	150 (131-181)	262 (223-305)	377 (299-446)	1311
Race/ethnicity							
Mexican Americans	01-02	78.2 (62.4-98.0)	70.8 (56.2-87.4)	130 (92.9-171)	224 (173-321)	333 (224-656)	664
	03-04	83.8 (75.1-93.6)	79.5 (72.9-87.0)	137 (111-165)	224 (188-274)	305 (247-406)	576
	05-06	68.8 (61.0-77.7)	61.3 (51.5-73.2)	96.9 (82.9-122)	175 (159-206)	291 (200-387)	620
	07-08	77.1 (68.5-86.7)	71.6 (63.1-78.8)	113 (99.6-133)	200 (160-227)	275 (230-334)	529
Non-Hispanic blacks	01-02	102 (85.8-122)	91.9 (79.8-108)	160 (133-202)	296 (234-475)	673 (347-1260)	690
	03-04	118 (103-135)	106 (90.5-126)	186 (156-219)	357 (257-439)	505 (384-772)	686
	05-06	97.0 (86.6-109)	85.7 (77.3-94.1)	141 (125-172)	271 (217-366)	443 (283-623)	658
	07-08	101 (88.7-114)	91.3 (80.9-104)	176 (143-208)	297 (238-360)	411 (344-512)	596
Non-Hispanic whites	01-02	103 (91.5-117)	89.4 (80.2-102)	167 (148-193)	324 (272-368)	447 (365-738)	1204
	03-04	112 (104-121)	103 (96.9-108)	183 (160-200)	346 (278-402)	544 (417-656)	1010
	05-06	106 (98.8-114)	93.6 (86.4-102)	178 (160-206)	348 (317-387)	506 (416-551)	982
	07-08	101 (91.1-112)	96.5 (83.5-107)	163 (137-190)	280 (243-317)	378 (316-447)	1077

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phenanthrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 4-Hydroxyphenanthrene

Metabolite of Phenanthrene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.*

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	25.1 (22.6-27.9)	25.9 (22.5-29.6)	53.7 (47.4-59.7)	96.8 (85.0-116)	152 (118-176)	2443
	05-06	29.3 (26.7-32.2)	29.2 (25.1-31.8)	56.9 (50.1-66.1)	108 (93.1-124)	157 (134-176)	2131
Age group							
6-11 years	03-04	25.2 (22.3-28.4)	26.3 (22.8-30.0)	45.7 (38.6-53.3)	95.5 (61.7-137)	138 (95.1-202)	321
	05-06	20.4 (17.7-23.4)	20.2 (17.6-23.8)	35.1 (27.4-41.6)	59.6 (43.3-93.0)	96.1 (62.2-155)	307
12-19 years	03-04	23.7 (20.3-27.6)	25.8 (20.5-31.4)	49.9 (45.2-53.0)	77.4 (67.2-95.1)	123 (94.6-142)	683
	05-06	26.6 (23.0-30.8)	27.6 (22.4-31.2)	50.9 (43.5-56.8)	79.4 (71.1-89.9)	108 (89.9-140)	568
20 years and older	03-04	25.3 (22.4-28.6)	25.8 (22.4-29.5)	54.8 (48.6-61.9)	101 (87.1-120)	163 (117-197)	1439
	05-06	31.1 (28.1-34.3)	30.7 (26.4-33.7)	62.9 (52.6-75.1)	119 (97.1-140)	165 (138-188)	1256
Gender							
Males	03-04	30.0 (26.5-34.0)	29.6 (25.5-35.1)	60.4 (53.4-68.7)	110 (90.6-130)	169 (118-211)	1180
	05-06	31.9 (28.6-35.5)	30.7 (25.0-36.8)	60.3 (49.7-72.7)	111 (92.3-134)	153 (119-226)	1064
Females	03-04	21.1 (18.7-23.8)	21.3 (18.3-25.3)	46.5 (41.6-53.3)	88.4 (73.5-103)	136 (106-157)	1263
	05-06	27.1 (23.4-31.4)	26.6 (22.7-31.1)	54.0 (41.9-66.9)	107 (82.6-140)	165 (120-189)	1067
Race/ethnicity							
Mexican Americans	03-04	23.5 (20.9-26.3)	24.4 (21.4-29.4)	46.1 (40.1-53.0)	80.5 (60.9-94.1)	99.7 (80.2-154)	607
	05-06	24.5 (20.3-29.6)	24.8 (20.2-31.3)	43.7 (37.2-54.5)	81.2 (62.8-94.4)	112 (82.6-134)	466
Non-Hispanic blacks	03-04	35.4 (29.6-42.3)	37.1 (29.8-46.1)	71.4 (60.3-83.3)	123 (104-145)	167 (133-215)	657
	05-06	36.8 (31.8-42.5)	33.6 (29.7-42.0)	66.0 (56.0-80.5)	130 (98.0-168)	190 (133-276)	612
Non-Hispanic whites	03-04	24.4 (21.5-27.7)	25.2 (21.8-29.1)	51.5 (45.1-58.5)	97.9 (77.4-127)	163 (113-202)	1021
	05-06	28.6 (25.2-32.4)	28.6 (24.5-31.9)	55.5 (47.3-69.3)	108 (90.9-136)	163 (134-188)	890

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 5.0 and 5.0 respectively.

*This chemical was not measured after NHANES 2005-2006 because of potential interferences that would affect the quality of the results.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phenanthrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 4-Hydroxyphenanthrene (creatinine corrected)

Metabolite of Phenanthrene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.*

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	23.3 (21.7-25.0)	22.7 (20.5-24.9)	42.0 (38.6-44.6)	74.3 (66.3-85.4)	114 (102-130)	2443
	05-06	28.9 (26.6-31.4)	26.9 (24.1-29.4)	48.4 (43.9-53.7)	88.3 (77.5-102)	127 (106-145)	2131
Age group							
6-11 years	03-04	26.3 (23.7-29.2)	24.8 (21.6-28.6)	43.3 (35.2-48.8)	72.9 (56.3-113)	126 (79.3-173)	321
	05-06	22.9 (19.8-26.5)	21.7 (20.0-24.8)	36.2 (27.8-43.7)	50.0 (42.7-79.3)	79.3 (50.0-98.3)	307
12-19 years	03-04	17.7 (15.9-19.8)	16.7 (15.7-18.4)	31.4 (26.0-37.1)	47.9 (40.7-54.1)	61.7 (51.7-86.0)	683
	05-06	20.1 (18.2-22.2)	19.0 (16.5-22.9)	30.0 (28.3-33.0)	48.7 (41.0-63.5)	79.4 (53.8-98.2)	568
20 years and older	03-04	24.0 (22.0-26.2)	23.3 (21.1-25.4)	43.5 (39.4-48.1)	79.0 (66.8-100)	121 (104-138)	1439
	05-06	31.4 (29.0-34.1)	29.4 (26.9-32.4)	54.6 (48.0-61.6)	102 (86.4-109)	139 (115-166)	1256
Gender							
Males	03-04	23.4 (21.6-25.4)	22.7 (20.3-24.9)	40.9 (35.9-45.7)	72.7 (61.5-86.8)	112 (100-126)	1180
	05-06	26.1 (23.8-28.7)	24.1 (22.2-28.2)	44.3 (40.0-51.1)	74.3 (61.6-87.7)	102 (87.3-120)	1064
Females	03-04	23.2 (21.2-25.5)	22.8 (19.7-26.0)	42.5 (38.3-46.2)	75.1 (63.4-94.0)	119 (93.1-139)	1263
	05-06	31.8 (28.7-35.3)	28.9 (25.8-31.8)	54.2 (46.0-63.2)	103 (85.4-127)	145 (119-174)	1067
Race/ethnicity							
Mexican Americans	03-04	21.2 (18.8-23.8)	20.7 (19.1-22.9)	35.9 (29.8-40.7)	59.0 (43.3-87.5)	83.4 (60.9-139)	607
	05-06	22.5 (20.1-25.3)	21.6 (18.8-27.1)	34.7 (31.4-41.5)	54.5 (48.2-61.6)	81.3 (59.8-86.1)	466
Non-Hispanic blacks	03-04	25.0 (21.6-28.9)	25.3 (23.1-28.8)	42.0 (37.4-51.4)	75.0 (61.6-87.9)	102 (79.2-137)	657
	05-06	25.8 (23.0-29.0)	24.5 (22.0-26.8)	39.4 (36.2-44.7)	68.5 (59.8-88.6)	103 (80.0-146)	612
Non-Hispanic whites	03-04	23.7 (21.5-26.2)	22.4 (19.4-25.4)	43.6 (38.8-47.3)	82.2 (68.5-101)	126 (103-143)	1021
	05-06	30.5 (27.3-34.1)	28.5 (25.0-31.6)	53.0 (45.5-61.5)	102 (83.0-115)	138 (112-174)	890

*This chemical was not measured after NHANES 2005-2006 because of potential interferences that would affect the quality of the results.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Phenanthrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 1-Hydroxypyrene

Metabolite of Pyrene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	89.2 (79.8-99.7)	91.3 (83.5-98.8)	189 (168-208)	389 (345-459)	569 (493-676)	2515
	05-06	98.3 (91.0-106)	94.0 (85.5-102)	206 (181-227)	440 (373-521)	673 (585-829)	2415
	07-08	118 (108-129)	118 (107-133)	233 (208-263)	439 (392-487)	675 (543-844)	2581
Age group							
6-11 years	03-04	112 (96.9-130)	119 (99.0-143)	193 (163-229)	351 (233-484)	514 (336-680)	333
	05-06	103 (91.3-117)	99.8 (91.7-105)	170 (139-206)	342 (222-413)	469 (353-786)	342
	07-08	146 (127-168)	149 (131-171)	231 (203-288)	395 (293-557)	569 (395-1060)	387
12-19 years	03-04	119 (103-137)	115 (98.0-145)	244 (213-274)	506 (359-608)	705 (636-788)	705
	05-06	118 (104-134)	108 (93.2-132)	218 (179-266)	409 (332-557)	837 (481-992)	667
	07-08	149 (130-170)	145 (118-165)	282 (228-343)	452 (399-515)	614 (476-843)	401
20 years and older	03-04	82.8 (73.0-93.8)	83.8 (75.6-92.4)	177 (155-203)	387 (337-437)	553 (483-644)	1477
	05-06	94.9 (86.3-104)	89.1 (79.8-100)	208 (173-245)	476 (368-563)	677 (563-913)	1406
	07-08	111 (100-123)	114 (95.7-129)	226 (199-258)	446 (370-506)	719 (525-904)	1793
Gender							
Males	03-04	108 (96.0-122)	111 (98.7-121)	227 (197-265)	459 (387-518)	644 (526-811)	1214
	05-06	114 (104-125)	108 (98.4-119)	235 (196-279)	465 (386-585)	793 (634-980)	1208
	07-08	134 (120-150)	131 (113-155)	260 (229-293)	474 (409-560)	748 (554-956)	1279
Females	03-04	74.0 (64.3-85.1)	75.3 (66.3-83.9)	158 (143-173)	334 (258-407)	502 (389-604)	1301
	05-06	84.8 (74.5-96.5)	80.0 (69.8-89.9)	173 (148-213)	421 (313-521)	617 (534-677)	1207
	07-08	105 (94.7-116)	111 (92.6-123)	213 (194-227)	387 (342-444)	580 (486-769)	1302
Race/ethnicity							
Mexican Americans	03-04	89.4 (78.6-102)	92.4 (79.8-113)	173 (153-191)	331 (281-415)	495 (404-548)	623
	05-06	95.6 (81.0-113)	92.3 (80.1-112)	179 (149-220)	332 (261-455)	490 (367-756)	611
	07-08	119 (110-129)	117 (104-132)	196 (170-230)	380 (289-494)	593 (485-755)	531
Non-Hispanic blacks	03-04	128 (105-155)	126 (108-153)	296 (226-355)	553 (400-669)	699 (564-935)	681
	05-06	116 (98.5-137)	108 (91.0-124)	230 (185-332)	517 (401-639)	743 (559-1060)	653
	07-08	136 (122-153)	142 (122-160)	282 (250-320)	548 (480-610)	752 (617-871)	592
Non-Hispanic whites	03-04	84.8 (73.5-97.9)	85.4 (75.6-97.3)	182 (156-214)	386 (336-462)	566 (470-749)	1050
	05-06	94.9 (85.4-106)	91.3 (80.3-104)	203 (170-235)	399 (351-538)	677 (563-899)	968
	07-08	116 (102-131)	117 (102-136)	235 (198-287)	436 (364-500)	719 (473-981)	1059

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 5.0, 5.0, and 5.0 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Pyrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 1-Hydroxypyrene (creatinine corrected)

Metabolite of Pyrene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	83.4 (77.4-90.0)	79.9 (72.6-86.9)	149 (133-167)	279 (236-343)	424 (352-474)	2515
	05-06	95.1 (89.1-102)	85.5 (81.0-91.1)	173 (161-184)	338 (292-371)	492 (427-550)	2415
	07-08	119 (110-129)	113 (103-127)	200 (180-224)	364 (321-411)	527 (467-587)	2581
Age group							
6-11 years	03-04	119 (102-138)	112 (98.2-137)	185 (156-225)	336 (227-440)	475 (336-566)	333
	05-06	115 (102-128)	105 (94.7-116)	181 (145-217)	285 (242-385)	407 (308-517)	342
	07-08	180 (156-206)	171 (145-197)	255 (233-316)	457 (329-593)	604 (457-1320)	387
12-19 years	03-04	89.4 (77.7-103)	81.6 (74.3-93.0)	146 (123-187)	269 (196-364)	364 (251-611)	705
	05-06	88.7 (78.7-99.9)	80.9 (73.8-92.3)	146 (115-176)	250 (199-358)	400 (252-682)	667
	07-08	116 (105-127)	112 (98.3-127)	178 (154-202)	301 (245-349)	405 (321-558)	401
20 years and older	03-04	79.1 (73.2-85.4)	73.5 (66.7-82.3)	142 (127-160)	278 (236-331)	424 (349-472)	1477
	05-06	94.1 (87.1-102)	82.5 (75.8-89.3)	177 (163-194)	352 (294-397)	502 (444-558)	1406
	07-08	114 (105-125)	106 (95.3-124)	194 (173-217)	364 (317-415)	510 (456-591)	1793
Gender							
Males	03-04	84.8 (77.3-93.1)	82.8 (72.7-91.0)	163 (145-175)	290 (241-352)	416 (353-513)	1214
	05-06	91.6 (84.3-99.6)	84.6 (76.9-95.5)	169 (152-182)	308 (268-364)	492 (426-589)	1208
	07-08	113 (102-125)	109 (94.3-124)	193 (176-214)	379 (308-440)	549 (456-629)	1279
Females	03-04	82.1 (73.5-91.7)	78.3 (69.2-86.8)	137 (119-165)	274 (215-350)	440 (313-479)	1301
	05-06	98.7 (91.7-106)	86.8 (78.5-92.8)	175 (156-195)	345 (295-390)	482 (395-558)	1207
	07-08	125 (114-137)	119 (104-135)	203 (179-238)	348 (302-405)	503 (425-619)	1302
Race/ethnicity							
Mexican Americans	03-04	81.2 (73.9-89.2)	80.9 (70.4-91.7)	146 (122-167)	247 (221-301)	372 (321-460)	623
	05-06	84.9 (76.4-94.2)	78.4 (67.7-91.2)	132 (118-145)	237 (203-287)	374 (279-482)	611
	07-08	116 (107-125)	105 (96.5-116)	173 (156-188)	296 (250-367)	501 (395-629)	531
Non-Hispanic blacks	03-04	91.0 (77.7-107)	92.4 (76.6-106)	173 (145-211)	315 (227-433)	451 (301-680)	681
	05-06	81.2 (70.7-93.2)	74.9 (65.6-84.8)	140 (113-173)	295 (206-403)	477 (324-643)	653
	07-08	106 (93.1-121)	104 (93.4-123)	201 (162-249)	352 (302-464)	537 (384-662)	592
Non-Hispanic whites	03-04	83.3 (75.4-92.0)	77.9 (69.1-86.8)	148 (127-174)	287 (241-360)	438 (360-506)	1050
	05-06	99.3 (91.2-108)	90.6 (82.2-99.5)	188 (169-200)	358 (305-395)	495 (427-555)	968
	07-08	120 (108-134)	113 (97.7-135)	206 (178-237)	379 (300-438)	510 (438-617)	1059

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Pyrene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 1-Hydroxynaphthalene (1-Naphthol)

Metabolite of Naphthalene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	2680 (2360-3050)	2260 (1980-2650)	7660 (6270-9420)	18500 (15500-20900)	26100 (23000-33600)	2595
	05-06	2640 (2390-2920)	1940 (1690-2240)	8450 (7050-9980)	29900 (23200-37100)	48100 (41300-65600)	2385
	07-08	2580 (2190-3030)	1980 (1750-2280)	6870 (5000-8760)	17300 (13000-22200)	29200 (22200-41100)	2495
Age group							
6-11 years	03-04	1540 (1360-1750)	1330 (1110-1680)	2790 (2180-3520)	5770 (4110-8200)	10500 (5870-17000)	340
	05-06	1190 (1010-1400)	1030 (775-1280)	2150 (1690-2730)	8100 (3870-10700)	11000 (7290-44100)	336
	07-08	1540 (1280-1850)	1500 (1220-1680)	3090 (2170-3800)	5940 (4160-9290)	9360 (4680-14300)	370
12-19 years	03-04	1960 (1680-2280)	1670 (1500-1970)	4190 (3100-5400)	10500 (8430-14300)	20900 (13500-26100)	727
	05-06	1710 (1490-1960)	1360 (1110-1700)	3760 (3100-4370)	14100 (9090-23500)	25800 (12400-46200)	655
	07-08	1830 (1490-2260)	1510 (1190-2010)	4030 (2550-5750)	10000 (6700-13500)	14800 (12000-21300)	386
20 years and older	03-04	3020 (2580-3520)	2650 (2200-3200)	9420 (7320-11500)	20500 (16500-23400)	29400 (23400-37400)	1528
	05-06	3100 (2740-3510)	2330 (2030-2820)	10500 (8180-13300)	34500 (26700-43400)	55100 (42200-80400)	1394
	07-08	2880 (2370-3490)	2260 (1830-2800)	8200 (6010-10700)	19600 (15000-25500)	33000 (23600-54900)	1739
Gender							
Males	03-04	3170 (2820-3560)	2840 (2370-3390)	8790 (7130-11000)	19600 (16100-22200)	25800 (22500-30600)	1243
	05-06	2970 (2620-3350)	2370 (1980-2970)	8760 (7140-10700)	28900 (21700-33000)	44100 (36000-52300)	1197
	07-08	2940 (2580-3360)	2370 (2050-2930)	7750 (5870-9830)	15900 (13800-21300)	28200 (21400-36800)	1239
Females	03-04	2290 (1940-2720)	1840 (1500-2310)	6380 (4450-8810)	17900 (14000-21500)	28500 (21500-37400)	1352
	05-06	2360 (2010-2770)	1590 (1420-1820)	7350 (5570-10200)	33600 (21600-43900)	56800 (41600-101E3)	1188
	07-08	2270 (1850-2790)	1580 (1320-1970)	5310 (4030-7760)	17700 (11900-24800)	32000 (21300-78900)	1256
Race/ethnicity							
Mexican Americans	03-04	1950 (1720-2190)	1650 (1370-2040)	4380 (3520-5300)	11600 (9040-15500)	20100 (14900-22500)	651
	05-06	1540 (1290-1830)	1290 (1090-1550)	3710 (2680-4430)	11800 (8100-15600)	22200 (15200-34000)	607
	07-08	1610 (1380-1890)	1480 (1230-1660)	3380 (2590-4040)	8750 (6120-11900)	16100 (8190-29400)	505
Non-Hispanic blacks	03-04	3340 (2800-3990)	2650 (2190-3460)	7840 (6520-10800)	20800 (17000-28600)	34700 (25400-48900)	695
	05-06	3420 (2830-4140)	2670 (2200-3430)	11900 (9120-14600)	29900 (24400-41000)	56400 (42800-71600)	641
	07-08	2880 (2480-3350)	2410 (2120-2910)	8150 (5310-10700)	19300 (13700-27100)	31600 (21300-42400)	574
Non-Hispanic whites	03-04	2800 (2410-3250)	2310 (1930-2840)	8630 (6820-11000)	19900 (15900-22600)	27600 (23400-35100)	1084
	05-06	2720 (2300-3210)	1980 (1560-2430)	8660 (6730-11300)	32500 (22600-42200)	51300 (41000-85500)	952
	07-08	2790 (2230-3500)	2080 (1740-2690)	7520 (4970-10400)	18700 (12600-26900)	32200 (22500-62000)	1022

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04, 05-06, and 07-08 are 46.7, 47.9, and 44.7 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Naphthalene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 1-Hydroxynaphthalene (1-Naphthol) (creatinine corrected)

Metabolite of Naphthalene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	2520 (2280-2790)	2100 (1870-2400)	6560 (5430-8180)	15100 (13500-17400)	21800 (18200-25400)	2595
	05-06	2560 (2290-2860)	1790 (1570-2060)	7570 (5890-9400)	24700 (20400-31000)	39600 (34100-53000)	2385
	07-08	2590 (2190-3060)	1980 (1660-2400)	6250 (4690-8220)	16100 (13600-18800)	25300 (20900-32800)	2495
Age group							
6-11 years	03-04	1630 (1430-1860)	1360 (1180-1740)	2770 (2260-3200)	5690 (3990-8950)	10200 (5690-16400)	340
	05-06	1300 (1090-1550)	1090 (938-1330)	2210 (1990-2810)	6150 (3330-10300)	10300 (6820-28000)	336
	07-08	1870 (1560-2240)	1670 (1350-2150)	3310 (2610-4650)	7360 (5130-8830)	9990 (8060-11300)	370
12-19 years	03-04	1470 (1260-1710)	1200 (1020-1390)	2840 (2490-4080)	7560 (5470-9120)	10800 (9120-13400)	727
	05-06	1290 (1140-1450)	995 (854-1100)	2700 (2270-3350)	8760 (5830-10800)	12500 (9370-19600)	655
	07-08	1390 (1180-1640)	1130 (833-1580)	3210 (2020-3630)	6950 (4790-8410)	9930 (7840-11900)	386
20 years and older	03-04	2900 (2590-3250)	2520 (2120-3060)	8210 (6450-9810)	16900 (14400-19800)	24200 (20000-27200)	1528
	05-06	3080 (2720-3500)	2320 (1910-2720)	10100 (7670-13900)	29800 (23900-34100)	46400 (39100-57300)	1394
	07-08	2960 (2430-3600)	2360 (1880-2770)	7420 (5750-10200)	18500 (14900-22600)	27900 (22300-45000)	1739
Gender							
Males	03-04	2490 (2250-2750)	2140 (1860-2590)	6560 (5500-8010)	13500 (12300-14400)	18300 (16600-20700)	1243
	05-06	2380 (2110-2680)	1890 (1590-2210)	6450 (4940-9210)	19200 (16100-22400)	31700 (22100-38400)	1197
	07-08	2480 (2210-2770)	2070 (1740-2390)	6170 (5070-7520)	13800 (11900-16000)	20700 (16700-24000)	1239
Females	03-04	2560 (2220-2950)	2040 (1730-2440)	6680 (5000-8850)	17500 (13500-21700)	24700 (19800-30600)	1352
	05-06	2750 (2370-3190)	1750 (1460-2060)	8050 (6080-11800)	31500 (24700-39500)	54100 (40800-64500)	1188
	07-08	2700 (2130-3430)	1940 (1580-2500)	6310 (4290-9330)	18900 (13800-23700)	32800 (22300-104E3)	1256
Race/ethnicity							
Mexican Americans	03-04	1750 (1590-1930)	1500 (1260-1690)	3240 (2900-4250)	8550 (7150-12000)	17500 (13500-21400)	651
	05-06	1370 (1150-1640)	1080 (940-1380)	2830 (2130-3820)	9500 (5950-12600)	18800 (12000-21600)	607
	07-08	1590 (1290-1950)	1290 (1030-1620)	3100 (2240-5050)	7580 (5420-10600)	11800 (7860-26100)	505
Non-Hispanic blacks	03-04	2370 (2020-2780)	2000 (1560-2420)	5900 (4440-7070)	13800 (8930-18700)	21100 (17100-27800)	695
	05-06	2400 (1990-2890)	1980 (1520-2510)	7170 (5280-8850)	20600 (15400-29200)	39200 (29200-54300)	641
	07-08	2220 (1870-2640)	1870 (1470-2410)	6050 (4340-7940)	13600 (10600-17000)	21300 (14100-31800)	574
Non-Hispanic whites	03-04	2770 (2460-3120)	2350 (1940-2900)	8030 (6330-9260)	16600 (14400-18500)	23600 (20000-26600)	1084
	05-06	2840 (2400-3360)	1990 (1600-2540)	8650 (5940-12600)	28200 (20800-35000)	43100 (38400-59000)	952
	07-08	2900 (2330-3610)	2160 (1710-2780)	7300 (5000-10200)	18400 (14800-22300)	27300 (22300-50800)	1022

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Naphthalene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 2-Hydroxynaphthalene (2-Naphthol)

Metabolite of Naphthalene

Geometric mean and selected percentiles of urine concentrations (in ng/L) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	2470 (2110-2890)	2280 (1930-2670)	5680 (4580-6830)	14700 (12800-19500)	26000 (22500-29700)	2748
	03-04	3180 (2760-3670)	2960 (2500-3500)	7500 (6190-9690)	17300 (14500-22100)	25800 (22600-27700)	2575
	05-06	3630 (3350-3920)	3440 (3110-3800)	8550 (7840-9420)	20200 (17000-24100)	33100 (28600-38200)	2446
	07-08	3830 (3370-4360)	3660 (3240-4120)	8460 (6670-10300)	17300 (14800-21900)	26700 (23100-30800)	2530
Age group							
6-11 years	01-02	1690 (1560-1840)	1700 (1400-1950)	3010 (2580-3470)	5410 (3890-6700)	7720 (6300-9540)	387
	03-04	2110 (1800-2470)	2140 (1790-2590)	3570 (3140-4380)	5580 (5050-6380)	9710 (5230-20000)	339
	05-06	2160 (1940-2410)	2090 (1950-2330)	3960 (3090-4460)	5640 (5000-9680)	10600 (7200-13300)	345
	07-08	2700 (2380-3060)	2590 (2140-2950)	4660 (4050-5230)	8720 (5910-10700)	11500 (9160-14800)	375
12-19 years	01-02	2220 (1700-2900)	2150 (1740-2530)	4390 (3150-6110)	11000 (6990-20400)	22500 (13900-28400)	735
	03-04	3040 (2680-3460)	2910 (2510-3500)	5290 (4670-6590)	12400 (8530-16800)	17600 (15100-25900)	721
	05-06	3460 (3030-3950)	3290 (2860-3780)	6750 (5870-7490)	14500 (10300-20800)	23900 (14700-33100)	677
	07-08	4290 (3480-5280)	4340 (3730-5310)	8740 (6570-10800)	17100 (13000-22500)	24000 (19600-29700)	387
20 years and older	01-02	2620 (2220-3100)	2440 (1940-2950)	6380 (5110-8110)	17600 (14000-21100)	28100 (23300-33700)	1626
	03-04	3360 (2860-3960)	3180 (2590-3960)	8770 (7210-11300)	19200 (16000-23200)	26600 (23400-28800)	1515
	05-06	3880 (3510-4290)	3770 (3250-4330)	9580 (8770-10600)	23000 (18600-28800)	37000 (29200-39500)	1424
	07-08	3910 (3410-4500)	3680 (3200-4320)	9170 (6820-11000)	18500 (15100-23600)	28600 (23800-33500)	1768
Gender							
Males	01-02	2750 (2360-3210)	2510 (2090-2970)	6060 (4820-7810)	16900 (11900-23000)	28100 (20800-35600)	1349
	03-04	3520 (3090-4010)	3370 (2910-3880)	8520 (6960-10500)	18400 (15300-22600)	26100 (22200-27900)	1233
	05-06	4020 (3650-4440)	3850 (3250-4340)	9490 (7840-11300)	19700 (16700-23100)	32900 (25500-38200)	1221
	07-08	4250 (3770-4780)	4200 (3740-4520)	9300 (7350-10800)	17500 (14500-22700)	26000 (20900-32600)	1256
Females	01-02	2220 (1860-2660)	2060 (1650-2480)	5240 (3890-6440)	13900 (12300-17700)	25300 (19700-28300)	1399
	03-04	2890 (2420-3450)	2600 (2140-3160)	6800 (5140-9210)	16600 (12700-21300)	23900 (22100-27600)	1342
	05-06	3280 (2830-3800)	3180 (2640-3600)	7830 (6770-9000)	21300 (14500-29200)	35000 (26700-39600)	1225
	07-08	3470 (2990-4030)	3180 (2650-3740)	7630 (5970-10000)	17100 (13600-21900)	26700 (21600-30800)	1274
Race/ethnicity							
Mexican Americans	01-02	2700 (2360-3080)	2710 (2350-3260)	5140 (4360-6150)	9640 (8150-10600)	14300 (10400-18700)	665
	03-04	3130 (2710-3630)	3110 (2570-3850)	6740 (5600-8280)	14500 (10400-15600)	18000 (15500-20500)	648
	05-06	4520 (4120-4960)	4790 (4130-5430)	8960 (7840-9810)	17300 (12400-23300)	28700 (20000-32600)	621
	07-08	4690 (4070-5410)	4500 (3930-5580)	8600 (6770-11300)	16400 (12600-20900)	24000 (19100-29700)	515
Non-Hispanic blacks	01-02	3970 (3470-4540)	3460 (3100-4020)	9290 (6820-13200)	22800 (16000-29100)	33000 (25900-38700)	692
	03-04	4690 (3830-5750)	4290 (3250-5540)	10300 (7670-13100)	21200 (17300-26100)	30100 (22300-40600)	690
	05-06	4990 (4410-5650)	4630 (3930-5620)	11600 (9370-14300)	22600 (20000-26900)	31900 (27600-38500)	657
	07-08	4900 (4420-5440)	4810 (4330-5190)	10100 (8050-12300)	22100 (17200-26300)	28600 (25000-36600)	587
Non-Hispanic whites	01-02	2190 (1760-2720)	1910 (1610-2420)	4970 (3710-6780)	14100 (10700-20200)	25900 (20700-30000)	1207
	03-04	3080 (2580-3670)	2660 (2200-3530)	7640 (5890-10400)	18000 (14600-22900)	26100 (22900-27900)	1072
	05-06	3250 (2900-3640)	2980 (2710-3330)	7900 (6690-9360)	19300 (15800-24700)	34000 (25500-38300)	980
	07-08	3600 (2980-4350)	3340 (2620-4170)	8040 (5730-10700)	17100 (11800-23800)	26400 (20400-35600)	1043

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, 05-06 and 07-08 are 2.4, 31.1, 13.2 and 42.0 respectively.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Naphthalene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Urinary 2-Hydroxynaphthalene (2-Naphthol) (creatinine corrected)

Metabolite of Naphthalene

Geometric mean and selected percentiles of urine concentrations (in ng/g of creatinine) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	2310 (1980-2680)	1940 (1670-2300)	4730 (3820-5860)	11500 (9980-13100)	16700 (14100-19200)	2748
	03-04	2990 (2670-3340)	2560 (2200-2970)	6340 (5010-7870)	14100 (11800-16300)	19900 (16000-23800)	2575
	05-06	3520 (3290-3770)	3080 (2670-3400)	7240 (6450-8270)	17000 (14800-18800)	23900 (21900-27200)	2446
	07-08	3860 (3440-4330)	3440 (3040-3910)	7100 (6010-8630)	15300 (12900-18300)	20700 (18800-23900)	2530
Age group							
6-11 years	01-02	1890 (1740-2070)	1830 (1720-1940)	3110 (2510-3470)	5040 (4380-5540)	6490 (5270-12400)	387
	03-04	2240 (1890-2650)	2040 (1660-2440)	3540 (2840-3910)	5950 (4140-8510)	8720 (7560-12000)	339
	05-06	2400 (2100-2750)	2270 (2000-2580)	3680 (3150-4430)	6100 (4840-7040)	7820 (6310-13800)	345
	07-08	3330 (2980-3730)	3380 (2900-3710)	5230 (4320-6780)	8000 (6790-9920)	10500 (8000-14500)	375
12-19 years	01-02	1720 (1350-2190)	1510 (1340-1830)	2750 (2080-4070)	7180 (4070-10900)	11100 (7860-17400)	735
	03-04	2280 (2060-2520)	2030 (1770-2350)	3780 (3200-4470)	7840 (6130-9110)	10500 (9100-12500)	721
	05-06	2590 (2370-2840)	2310 (2030-2620)	4580 (3990-5020)	8470 (6980-10500)	11900 (10100-15200)	677
	07-08	3300 (2740-3960)	3130 (2430-4080)	5730 (4880-6790)	9850 (7560-12000)	12500 (9910-18600)	387
20 years and older	01-02	2480 (2130-2880)	2080 (1680-2600)	5630 (4460-6940)	12400 (10900-13900)	17700 (15200-20600)	1626
	03-04	3230 (2870-3640)	2910 (2360-3410)	7540 (6200-9100)	15600 (13100-17900)	21800 (17200-24000)	1515
	05-06	3870 (3590-4170)	3400 (3080-3750)	9200 (7930-10500)	19000 (17400-21200)	27200 (22300-29700)	1424
	07-08	4020 (3550-4550)	3470 (3070-4040)	7750 (6170-9830)	17500 (13900-19800)	22700 (20200-25500)	1768
Gender							
Males	01-02	2230 (1900-2610)	1860 (1630-2100)	4790 (3510-6120)	11400 (8950-14300)	15800 (13100-19200)	1349
	03-04	2770 (2500-3060)	2360 (2130-2810)	6300 (5080-7360)	12800 (11300-14600)	16400 (14600-18300)	1233
	05-06	3230 (3000-3480)	2830 (2530-3140)	6750 (5620-8100)	13800 (12300-16200)	18900 (17400-21200)	1221
	07-08	3580 (3240-3950)	3130 (2700-3570)	6790 (5790-8100)	14400 (12200-17200)	19600 (18300-20900)	1256
Females	01-02	2380 (2050-2770)	1990 (1660-2560)	4730 (3790-6000)	11500 (10300-12800)	17400 (13700-21500)	1399
	03-04	3210 (2760-3730)	2680 (2200-3260)	6430 (4740-8580)	16000 (11500-19900)	22500 (18100-25000)	1342
	05-06	3830 (3390-4330)	3210 (2720-3760)	8080 (6430-9740)	20800 (17000-24700)	29000 (24900-34200)	1225
	07-08	4150 (3570-4830)	3590 (3190-4180)	7400 (5880-9630)	16800 (12900-20000)	22700 (20000-25500)	1274
Race/ethnicity							
Mexican Americans	01-02	2520 (2230-2850)	2350 (2010-2950)	4650 (3970-5210)	7320 (6050-8090)	12000 (7820-15300)	665
	03-04	2830 (2440-3280)	2570 (2200-3230)	5000 (4100-5940)	10300 (6790-13000)	13300 (11000-15500)	648
	05-06	4050 (3770-4350)	4030 (3650-4360)	7050 (6300-8050)	12600 (9530-15100)	17400 (13400-20600)	621
	07-08	4580 (3970-5280)	4370 (3550-5020)	7590 (6300-9380)	14200 (11600-16400)	19100 (16000-24700)	515
Non-Hispanic blacks	01-02	2790 (2390-3270)	2410 (2040-2770)	5980 (4840-6980)	11600 (9090-15300)	17400 (12200-23400)	692
	03-04	3320 (2700-4090)	3060 (2310-3860)	7050 (5140-9410)	12800 (10200-16500)	19000 (15700-23400)	690
	05-06	3510 (3160-3890)	3100 (2590-3530)	6710 (5330-8040)	12300 (10300-16100)	19700 (16100-25000)	657
	07-08	3800 (3470-4150)	3450 (3080-3900)	6730 (5560-8410)	13100 (11300-14900)	19200 (14800-26500)	587
Non-Hispanic whites	01-02	2160 (1780-2620)	1740 (1440-2180)	4340 (3320-6090)	11900 (9930-14300)	16900 (13800-20600)	1207
	03-04	3040 (2660-3460)	2480 (2110-3020)	7170 (5100-8950)	15500 (13300-17600)	21800 (17100-24000)	1072
	05-06	3400 (3080-3760)	2730 (2390-3260)	7450 (6100-9650)	17900 (15300-20700)	24900 (21200-29000)	980
	07-08	3750 (3180-4430)	3280 (2600-4080)	6910 (5530-9510)	16800 (12800-19800)	22200 (19400-25500)	1043

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Naphthalene_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PAHs_FactSheet.html

Blood 1,1,1-Trichloroethane (Methyl chloroform)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1345
	05-06	*	< LOD	< LOD	< LOD	< LOD	3085
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	911
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1345
	05-06	*	< LOD	< LOD	< LOD	< LOD	1524
60 years and older	05-06	*	< LOD	< LOD	< LOD	.014 (<LOD-.021)	650
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	660
	05-06	*	< LOD	< LOD	< LOD	< LOD	1464
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	685
	05-06	*	< LOD	< LOD	< LOD	< LOD	1621
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	762
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	289
	05-06	*	< LOD	< LOD	< LOD	< LOD	791
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	688
	05-06	*	< LOD	< LOD	< LOD	< LOD	1303

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.048 and 0.01 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood 1,1,2,2-Tetrachloroethane

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1235
	05-06	*	< LOD	< LOD	< LOD	< LOD	3131
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	924
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1235
	05-06	*	< LOD	< LOD	< LOD	< LOD	1542
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	665
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	613
	05-06	*	< LOD	< LOD	< LOD	< LOD	1484
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	622
	05-06	*	< LOD	< LOD	< LOD	< LOD	1647
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	250
	05-06	*	< LOD	< LOD	< LOD	< LOD	753
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	282
	05-06	*	< LOD	< LOD	< LOD	< LOD	823
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	606
	05-06	*	< LOD	< LOD	< LOD	< LOD	1323

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.01 and 0.01 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood 1,1,2-Trichloroethane

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1354
	05-06	*	< LOD	< LOD	< LOD	< LOD	3165
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	936
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1354
	05-06	*	< LOD	< LOD	< LOD	< LOD	1556
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	673
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	664
	05-06	*	< LOD	< LOD	< LOD	< LOD	1499
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	690
	05-06	*	< LOD	< LOD	< LOD	< LOD	1666
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	777
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	297
	05-06	*	< LOD	< LOD	< LOD	< LOD	824
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	686
	05-06	*	< LOD	< LOD	< LOD	< LOD	1331

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.01 and 0.01 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood 1,1-Dichloroethane

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1367
	05-06	*	< LOD	< LOD	< LOD	< LOD	3193
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	941
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1367
	05-06	*	< LOD	< LOD	< LOD	< LOD	1569
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	683
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	670
	05-06	*	< LOD	< LOD	< LOD	< LOD	1510
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	697
	05-06	*	< LOD	< LOD	< LOD	< LOD	1683
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	778
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	300
	05-06	*	< LOD	< LOD	< LOD	< LOD	832
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	695
	05-06	*	< LOD	< LOD	< LOD	< LOD	1347

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.01 and 0.01 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood 1,1 - Dichloroethene (Vinylidene chloride)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1367
	05-06	*	< LOD	< LOD	< LOD	< LOD	3163
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	931
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1367
	05-06	*	< LOD	< LOD	< LOD	< LOD	1555
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	677
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	670
	05-06	*	< LOD	< LOD	< LOD	< LOD	1502
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	697
	05-06	*	< LOD	< LOD	< LOD	< LOD	1661
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	774
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	300
	05-06	*	< LOD	< LOD	< LOD	< LOD	823
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	695
	05-06	*	< LOD	< LOD	< LOD	< LOD	1332

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.009 and 0.009 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood 1,2-Dibromo-3-chloropropane (DBCP)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1170
	05-06	*	< LOD	< LOD	< LOD	< LOD	2998
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	875
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1170
	05-06	*	< LOD	< LOD	< LOD	< LOD	1485
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	638
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	568
	05-06	*	< LOD	< LOD	< LOD	< LOD	1409
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	602
	05-06	*	< LOD	< LOD	< LOD	< LOD	1589
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	234
	05-06	*	< LOD	< LOD	< LOD	< LOD	752
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	239
	05-06	*	< LOD	< LOD	< LOD	< LOD	737
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	603
	05-06	*	< LOD	< LOD	< LOD	< LOD	1287

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.1 and 0.1 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DBCP_BiomonitoringSummary.html

Blood 1,2-Dichlorobenzene (*o*-Dichlorobenzene)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1327
	05-06	*	< LOD	< LOD	< LOD	< LOD	3083
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	921
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1327
	05-06	*	< LOD	< LOD	< LOD	< LOD	1523
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	639
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	647
	05-06	*	< LOD	< LOD	< LOD	< LOD	1457
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	680
	05-06	*	< LOD	< LOD	< LOD	< LOD	1626
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	250
	05-06	*	< LOD	< LOD	< LOD	< LOD	766
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	291
	05-06	*	< LOD	< LOD	< LOD	< LOD	801
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	682
	05-06	*	< LOD	< LOD	< LOD	< LOD	1288

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.1 and 0.1 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Chlorobenzenes_BiomonitoringSummary.html

Blood 1,2-Dichloroethane (Ethylene dichloride)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1346
	05-06	*	< LOD	< LOD	< LOD	< LOD	3082
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	906
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1346
	05-06	*	< LOD	< LOD	< LOD	< LOD	1520
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	656
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	661
	05-06	*	< LOD	< LOD	< LOD	< LOD	1452
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	685
	05-06	*	< LOD	< LOD	< LOD	< LOD	1630
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	763
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	289
	05-06	*	< LOD	< LOD	< LOD	< LOD	795
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	689
	05-06	*	< LOD	< LOD	< LOD	< LOD	1297

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.01 and 0.01 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood *cis*-1,2-Dichloroethene

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1366
	05-06	*	< LOD	< LOD	< LOD	< LOD	3140
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	924
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1366
	05-06	*	< LOD	< LOD	< LOD	< LOD	1539
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	677
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	669
	05-06	*	< LOD	< LOD	< LOD	< LOD	1481
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	697
	05-06	*	< LOD	< LOD	< LOD	< LOD	1659
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	757
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	300
	05-06	*	< LOD	< LOD	< LOD	< LOD	822
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	694
	05-06	*	< LOD	< LOD	< LOD	< LOD	1329

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.01 and 0.01 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood *trans*-1,2-Dichloroethene

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1367
	05-06	*	< LOD	< LOD	< LOD	< LOD	3193
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	941
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1367
	05-06	*	< LOD	< LOD	< LOD	< LOD	1569
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	683
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	670
	05-06	*	< LOD	< LOD	< LOD	< LOD	1511
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	697
	05-06	*	< LOD	< LOD	< LOD	< LOD	1682
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	779
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	300
	05-06	*	< LOD	< LOD	< LOD	< LOD	832
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	695
	05-06	*	< LOD	< LOD	< LOD	< LOD	1346

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.01 and 0.01 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood 1,2-Dichloropropane

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1364
	05-06	*	< LOD	< LOD	< LOD	< LOD	3120
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	918
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1364
	05-06	*	< LOD	< LOD	< LOD	< LOD	1536
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	666
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	667
	05-06	*	< LOD	< LOD	< LOD	< LOD	1478
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	697
	05-06	*	< LOD	< LOD	< LOD	< LOD	1642
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	755
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	300
	05-06	*	< LOD	< LOD	< LOD	< LOD	811
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	692
	05-06	*	< LOD	< LOD	< LOD	< LOD	1322

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.008 and 0.008 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood 1,3-Dichlorobenzene (*m*-Dichlorobenzene)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1334
	05-06	*	< LOD	< LOD	< LOD	< LOD	3148
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	928
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1334
	05-06	*	< LOD	< LOD	< LOD	< LOD	1545
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	675
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	659
	05-06	*	< LOD	< LOD	< LOD	< LOD	1491
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	675
	05-06	*	< LOD	< LOD	< LOD	< LOD	1657
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	266
	05-06	*	< LOD	< LOD	< LOD	< LOD	774
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	279
	05-06	*	< LOD	< LOD	< LOD	< LOD	809
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	686
	05-06	*	< LOD	< LOD	< LOD	< LOD	1330

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.05 and 0.05 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Chlorobenzenes_BiomonitoringSummary.html

Blood 1,4-Dichlorobenzene (Paradichlorobenzene)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	.300 (.190-.550)	1.19 (.610-2.90)	4.10 (1.40-7.60)	807
	03-04	*	< LOD	.320 (.250-.400)	1.10 (.710-1.60)	3.30 (1.70-5.10)	1322
	05-06	*	.085 (.085-.085)	.218 (.180-.260)	.917 (.643-1.31)	3.59 (1.80-5.50)	3126
Age group							
12-19 years	05-06	*	.085 (.085-.085)	.220 (.160-.280)	.960 (.560-2.00)	4.27 (2.37-6.10)	926
20-59 years	01-02	*	< LOD	.300 (.190-.550)	1.19 (.610-2.90)	4.10 (1.40-7.60)	807
	03-04	*	< LOD	.320 (.250-.400)	1.10 (.710-1.60)	3.30 (1.70-5.10)	1322
	05-06	*	.085 (.085-.085)	.220 (.170-.260)	.910 (.540-1.50)	3.77 (1.50-7.33)	1537
60 years and older	05-06	*	.085 (.085-.085)	.207 (.158-.313)	.877 (.525-1.80)	3.06 (1.50-4.20)	663
Gender							
Males	01-02	*	< LOD	.270 (.180-.530)	.900 (.570-2.90)	3.60 (1.10-7.20)	390
	03-04	*	< LOD	.320 (.250-.380)	.770 (.580-1.20)	1.90 (1.20-4.00)	651
	05-06	*	.085 (.085-.085)	.230 (.188-.290)	.950 (.603-1.50)	3.50 (1.50-7.20)	1479
Females	01-02	*	< LOD	.360 (.190-.620)	1.20 (.620-3.90)	4.10 (1.68-8.30)	417
	03-04	*	< LOD	.350 (.220-.450)	1.40 (.810-2.10)	4.10 (2.20-5.90)	671
	05-06	*	.085 (.085-.085)	.180 (.140-.240)	.865 (.522-1.70)	3.59 (2.10-5.18)	1647
Race/ethnicity							
Mexican Americans	01-02	.331 (.246-.446)	.180 (<LOD-.260)	.790 (.400-1.30)	6.00 (1.30-16.0)	16.0 (6.20-33.0)	217
	03-04	.381 (.256-.566)	.210 (.140-.400)	.730 (.370-2.90)	6.20 (2.90-9.30)	10.0 (6.30-19.0)	262
	05-06	.412 (.309-.549)	.210 (.150-.290)	1.20 (.680-2.97)	8.80 (3.80-16.0)	16.0 (8.10-33.0)	772
Non-Hispanic blacks	01-02	.558 (.376-.827)	.360 (.260-.560)	1.80 (.710-3.90)	6.30 (1.40-29.0)	15.0 (3.60-51.0)	136
	03-04	.423 (.292-.613)	.340 (.200-.480)	.980 (.550-1.50)	4.10 (1.60-9.20)	11.0 (2.50-19.0)	297
	05-06	.381 (.272-.532)	.230 (.160-.330)	.860 (.500-1.70)	5.24 (2.00-11.5)	12.7 (8.41-21.0)	805
Non-Hispanic whites	01-02	*	< LOD	.200 (<LOD-.380)	.570 (.300-1.20)	1.19 (.570-3.70)	396
	03-04	*	< LOD	.200 (.160-.260)	.490 (.370-.720)	.940 (.690-2.00)	658
	05-06	*	.085 (.085-.085)	.134 (.085-.180)	.438 (.260-.670)	1.16 (.520-2.30)	1319

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.12, 0.12, and 0.011 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Chlorobenzenes_BiomonitoringSummary.html

Blood 2,5-Dimethylfuran

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	.016 (<LOD-.032)	.100 (.083-.110)	.140 (.120-.180)	1213
	05-06	*	< LOD	< LOD	.100 (.069-.120)	.160 (.130-.180)	3160
Age group							
12-19 years	05-06	*	< LOD	< LOD	.013 (<LOD-.029)	.049 (.020-.069)	930
20-59 years	03-04	*	< LOD	.016 (<LOD-.032)	.100 (.083-.110)	.140 (.120-.180)	1213
	05-06	*	< LOD	.013 (<LOD-.035)	.120 (.097-.144)	.170 (.140-.210)	1553
60 years and older	05-06	*	< LOD	< LOD	.069 (.027-.100)	.140 (.106-.160)	677
Gender							
Males	03-04	*	< LOD	.019 (<LOD-.034)	.094 (.071-.110)	.130 (.110-.190)	598
	05-06	*	< LOD	< LOD	.110 (.079-.130)	.160 (.130-.190)	1490
Females	03-04	*	< LOD	.012 (<LOD-.048)	.110 (.071-.120)	.150 (.120-.210)	615
	05-06	*	< LOD	< LOD	.094 (.062-.120)	.158 (.120-.180)	1670
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	.012 (<LOD-.020)	.038 (.013-.054)	237
	05-06	*	< LOD	< LOD	.015 (<LOD-.029)	.033 (.020-.055)	778
Non-Hispanic blacks	03-04	*	< LOD	.042 (.019-.063)	.110 (.079-.170)	.170 (.099-.220)	258
	05-06	*	< LOD	.012 (<LOD-.030)	.085 (.058-.110)	.130 (.095-.180)	825
Non-Hispanic whites	03-04	*	< LOD	.021 (<LOD-.043)	.110 (.087-.120)	.150 (.120-.200)	624
	05-06	*	< LOD	< LOD	.118 (.091-.140)	.170 (.140-.230)	1325

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.012 and 0.011 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/2,5-Dimethylfuran_BiomonitoringSummary.html

Blood Benzene

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	.030 (<LOD-.050)	.100 (.060-.140)	.190 (.140-.290)	.320 (.190-.480)	837
	03-04	*	.027 (.025-.031)	.064 (.050-.084)	.170 (.150-.190)	.260 (.210-.320)	1345
	05-06	*	.026 (<LOD-.030)	.056 (.047-.072)	.220 (.157-.260)	.310 (.264-.370)	3091
Age group							
12-19 years	05-06	*	.024 (<LOD-.028)	.039 (.031-.056)	.089 (.066-.110)	.120 (.092-.180)	912
20-59 years	01-02	*	.030 (<LOD-.050)	.100 (.060-.140)	.190 (.140-.290)	.320 (.190-.480)	837
	03-04	*	.027 (.025-.031)	.064 (.050-.084)	.170 (.150-.190)	.260 (.210-.320)	1345
	05-06	*	.028 (.025-.032)	.072 (.053-.100)	.240 (.200-.290)	.340 (.300-.390)	1513
60 years and older	05-06	*	< LOD	.042 (.036-.048)	.180 (.100-.230)	.270 (.200-.340)	666
Gender							
Males	01-02	*	.030 (<LOD-.050)	.110 (.060-.160)	.230 (.150-.370)	.370 (.210-.510)	403
	03-04	.039 (.035-.043)	.030 (.027-.035)	.069 (.053-.084)	.160 (.140-.180)	.240 (.190-.320)	654
	05-06	*	.028 (.025-.032)	.066 (.051-.083)	.230 (.165-.280)	.330 (.260-.440)	1459
Females	01-02	*	.030 (<LOD-.050)	.100 (.060-.130)	.180 (.120-.240)	.250 (.170-.330)	434
	03-04	*	.025 (<LOD-.029)	.057 (.040-.090)	.180 (.150-.220)	.290 (.220-.420)	691
	05-06	*	.024 (<LOD-.028)	.049 (.040-.067)	.200 (.150-.230)	.290 (.250-.340)	1632
Race/ethnicity							
Mexican Americans	01-02	*	.030 (<LOD-.070)	.070 (.030-.170)	.140 (.060-.360)	.230 (.130-.370)	227
	03-04	*	.027 (<LOD-.035)	.041 (.034-.057)	.077 (.058-.110)	.130 (.084-.320)	254
	05-06	*	< LOD	.036 (.031-.045)	.060 (.048-.086)	.096 (.072-.140)	744
Non-Hispanic blacks	01-02	*	< LOD	.060 (<LOD-.160)	.180 (.090-.300)	.250 (.160-.480)	137
	03-04	.043 (.033-.058)	.029 (<LOD-.054)	.092 (.055-.140)	.210 (.140-.290)	.320 (.240-.460)	302
	05-06	*	.026 (<LOD-.034)	.077 (.051-.110)	.180 (.150-.221)	.280 (.220-.360)	812
Non-Hispanic whites	01-02	*	.030 (<LOD-.060)	.110 (.070-.160)	.210 (.140-.340)	.330 (.190-.510)	411
	03-04	*	.028 (.025-.033)	.068 (.053-.088)	.180 (.150-.200)	.280 (.210-.330)	687
	05-06	*	.027 (<LOD-.031)	.069 (.049-.091)	.240 (.190-.280)	.330 (.270-.390)	1309

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.024, 0.024, and 0.024 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Benzene_BiomonitoringSummary.html

Blood Chlorobenzene (Monochlorobenzene)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1366
	05-06	*	< LOD	< LOD	< LOD	< LOD	3072
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	910
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1366
	05-06	*	< LOD	< LOD	< LOD	< LOD	1495
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	667
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	669
	05-06	*	< LOD	< LOD	< LOD	< LOD	1453
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	697
	05-06	*	< LOD	< LOD	< LOD	< LOD	1619
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	740
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	300
	05-06	*	< LOD	< LOD	< LOD	< LOD	817
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	694
	05-06	*	< LOD	< LOD	< LOD	< LOD	1287

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.011 and 0.011 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Chlorobenzenes_BiomonitoringSummary.html

Blood Dibromomethane

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1355
	05-06	*	< LOD	< LOD	< LOD	< LOD	3152
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	929
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1355
	05-06	*	< LOD	< LOD	< LOD	< LOD	1554
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	669
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	666
	05-06	*	< LOD	< LOD	< LOD	< LOD	1494
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	689
	05-06	*	< LOD	< LOD	< LOD	< LOD	1658
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	778
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	292
	05-06	*	< LOD	< LOD	< LOD	< LOD	816
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	693
	05-06	*	< LOD	< LOD	< LOD	< LOD	1324

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.03 and 0.03 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood Dichloromethane (Methylene chloride)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1165
	05-06	*	< LOD	< LOD	< LOD	< LOD	2852
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	837
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1165
	05-06	*	< LOD	< LOD	< LOD	< LOD	1400
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	615
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	568
	05-06	*	< LOD	< LOD	< LOD	< LOD	1338
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	597
	05-06	*	< LOD	< LOD	< LOD	< LOD	1514
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	225
	05-06	*	< LOD	< LOD	< LOD	< LOD	731
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	245
	05-06	*	< LOD	< LOD	< LOD	< LOD	729
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	607
	05-06	*	< LOD	< LOD	< LOD	< LOD	1186

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.07 and 0.25 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/HalogenatedSolvents_BiomonitoringSummary.html

Blood Ethylbenzene

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	.034 (.029-.039)	.030 (.030-.040)	.050 (.050-.070)	.090 (.070-.120)	.140 (.090-.180)	879
	03-04	.035 (.033-.037)	.032 (.030-.036)	.053 (.048-.057)	.083 (.077-.088)	.110 (.098-.120)	1299
	05-06	.038 (.036-.041)	.035 (.033-.037)	.059 (.052-.067)	.100 (.093-.114)	.140 (.122-.160)	3119
Age group							
12-19 years	05-06	.032 (.030-.034)	.031 (.029-.033)	.046 (.042-.049)	.070 (.058-.085)	.096 (.075-.120)	909
20-59 years	01-02	.034 (.029-.039)	.030 (.030-.040)	.050 (.050-.070)	.090 (.070-.120)	.140 (.090-.180)	879
	03-04	.035 (.033-.037)	.032 (.030-.036)	.053 (.048-.057)	.083 (.077-.088)	.110 (.098-.120)	1299
	05-06	.040 (.037-.043)	.037 (.034-.039)	.062 (.054-.073)	.110 (.098-.126)	.150 (.130-.180)	1539
	05-06	.037 (.033-.040)	.033 (.030-.036)	.056 (.046-.068)	.098 (.090-.110)	.130 (.110-.160)	671
60 years and older							
Gender							
Males	01-02	.035 (.029-.041)	.030 (<LOD-.040)	.060 (.040-.070)	.100 (.070-.150)	.150 (.090-.220)	419
	03-04	.037 (.034-.041)	.036 (.031-.040)	.057 (.051-.065)	.086 (.078-.099)	.110 (.094-.130)	625
	05-06	.042 (.039-.044)	.038 (.036-.041)	.067 (.057-.074)	.114 (.099-.131)	.160 (.136-.180)	1474
Females	01-02	.033 (.028-.038)	.030 (.030-.040)	.050 (.040-.060)	.080 (.060-.120)	.130 (.070-.180)	460
	03-04	.032 (.030-.034)	.030 (.027-.033)	.048 (.041-.053)	.081 (.071-.089)	.100 (.091-.120)	674
	05-06	.035 (.032-.038)	.032 (.029-.035)	.052 (.047-.061)	.093 (.081-.103)	.120 (.106-.150)	1645
Race/ethnicity							
Mexican Americans	01-02	*	.030 (<LOD-.040)	.040 (.040-.060)	.070 (.050-.110)	.120 (.070-.210)	220
	03-04	.031 (.027-.036)	.029 (<LOD-.035)	.044 (.035-.048)	.064 (.051-.074)	.091 (.066-.170)	253
	05-06	.032 (.028-.036)	.030 (.026-.036)	.047 (.039-.057)	.072 (.051-.093)	.086 (.073-.120)	752
Non-Hispanic blacks	01-02	.032 (.027-.038)	.030 (<LOD-.040)	.050 (.030-.070)	.080 (.060-.130)	.130 (.070-.240)	159
	03-04	.032 (.027-.038)	.030 (<LOD-.037)	.050 (.037-.066)	.079 (.065-.100)	.110 (.084-.130)	281
	05-06	.037 (.035-.040)	.034 (.030-.038)	.057 (.052-.064)	.095 (.085-.100)	.120 (.104-.140)	809
Non-Hispanic whites	01-02	.035 (.029-.043)	.030 (<LOD-.040)	.060 (.050-.070)	.090 (.070-.140)	.150 (.090-.190)	432
	03-04	.036 (.034-.038)	.034 (.030-.038)	.055 (.050-.061)	.087 (.081-.092)	.110 (.098-.130)	669
	05-06	.039 (.037-.042)	.036 (.033-.038)	.061 (.053-.072)	.110 (.095-.120)	.150 (.125-.180)	1326

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.024, 0.024, and 0.024 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Ethylbenzene_BiomonitoringSummary.html

Blood Hexachloroethane

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1366
	05-06	*	< LOD	< LOD	< LOD	< LOD	3155
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	930
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1366
	05-06	*	< LOD	< LOD	< LOD	< LOD	1547
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	678
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	669
	05-06	*	< LOD	< LOD	< LOD	< LOD	1495
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	697
	05-06	*	< LOD	< LOD	< LOD	< LOD	1660
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	267
	05-06	*	< LOD	< LOD	< LOD	< LOD	758
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	300
	05-06	*	< LOD	< LOD	< LOD	< LOD	830
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	694
	05-06	*	< LOD	< LOD	< LOD	< LOD	1335

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.011 and 0.011 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Hexachloroethane_BiomonitoringSummary.html

Blood Methyl-*tert*-butyl ether (MTBE)

Geometric mean and selected percentiles of whole blood concentrations (in pg/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	16.4 (4.77-56.7)	27.7 (7.29-64.9)	73.8 (35.5-127)	132 (64.0-278)	188 (109-339)	672
	03-04	11.0 (5.98-20.1)	10.0 (4.60-25.1)	45.0 (16.0-98.0)	110 (70.0-180)	170 (110-340)	1307
	05-06	6.16 (2.94-12.9)	4.60 (1.80-15.0)	20.0 (5.20-73.0)	75.0 (22.0-140)	110 (49.0-250)	3080
Age group							
12-19 years	05-06	5.85 (2.61-13.1)	4.20 (1.40-18.0)	18.0 (4.60-75.0)	70.0 (18.0-140)	110 (62.0-180)	911
20-59 years	01-02	16.4 (4.77-56.7)	27.7 (7.29-64.9)	73.8 (35.5-127)	132 (64.0-278)	188 (109-339)	672
	03-04	11.0 (5.98-20.1)	10.0 (4.60-25.1)	45.0 (16.0-98.0)	110 (70.0-180)	170 (110-340)	1307
	05-06	6.26 (3.05-12.9)	4.80 (2.00-14.0)	20.0 (5.50-70.0)	71.0 (21.0-150)	110 (46.0-250)	1512
60 years and older	05-06	6.00 (2.67-13.5)	3.80 (1.60-14.0)	22.0 (4.00-94.0)	86.0 (19.0-180)	120 (55.0-290)	657
Gender							
Males	01-02	16.9 (4.96-57.7)	27.9 (6.82-64.6)	75.0 (35.5-131)	132 (54.9-307)	167 (109-417)	334
	03-04	12.2 (6.29-23.6)	11.0 (4.80-29.0)	55.0 (18.0-110)	130 (79.0-200)	200 (110-470)	641
	05-06	6.24 (3.07-12.7)	4.60 (1.90-15.0)	20.0 (5.90-69.0)	73.0 (25.0-120)	110 (52.0-230)	1462
Females	01-02	16.0 (4.12-61.8)	26.6 (5.93-74.6)	72.7 (32.6-132)	142 (73.5-255)	194 (92.3-336)	338
	03-04	9.88 (5.62-17.4)	8.90 (4.30-23.0)	38.0 (14.0-85.0)	94.0 (58.0-160)	140 (90.0-250)	666
	05-06	6.08 (2.81-13.1)	4.50 (1.70-14.0)	19.0 (4.90-79.0)	76.0 (20.0-160)	110 (48.0-250)	1618
Race/ethnicity							
Mexican Americans	01-02	23.3 (4.96-110)	33.4 (2.92-187)	91.3 (26.1-255)	225 (80.6-339)	273 (182-358)	166
	03-04	11.6 (5.35-25.3)	12.0 (3.80-29.0)	32.0 (14.0-80.0)	80.0 (38.0-190)	160 (74.0-220)	245
	05-06	7.61 (2.99-19.3)	5.40 (2.00-33.0)	30.0 (4.60-90.0)	67.0 (26.0-190)	100 (45.0-330)	735
Non-Hispanic blacks	01-02	14.9 (6.11-36.4)	26.4 (3.11-55.6)	52.6 (30.0-86.8)	87.4 (38.6-155)	120 (70.0-155)	119
	03-04	9.63 (4.83-19.2)	10.0 (3.50-28.0)	32.8 (11.0-85.0)	77.0 (36.0-160)	140 (61.0-210)	285
	05-06	9.14 (3.16-26.5)	5.70 (1.80-50.0)	48.0 (5.00-150)	120 (46.0-280)	180 (88.0-460)	820
Non-Hispanic whites	01-02	16.0 (4.13-62.4)	27.9 (4.71-74.6)	72.7 (33.3-132)	132 (59.6-249)	165 (92.8-366)	333
	03-04	11.5 (5.51-23.8)	11.0 (4.00-33.5)	59.0 (14.0-120)	120 (73.0-230)	180 (110-430)	673
	05-06	5.55 (2.47-12.5)	4.00 (<LOD-15.0)	16.0 (4.30-75.0)	70.0 (15.0-150)	110 (40.0-250)	1291

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.232, 2.0, and 1.4 respectively. < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/MTBE_BiomonitoringSummary.html

Blood Nitrobenzene

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1066
	05-06	*	< LOD	< LOD	< LOD	< LOD	3144
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	926
20-59 years	03-04	*	< LOD	< LOD	< LOD	< LOD	1066
	05-06	*	< LOD	< LOD	< LOD	< LOD	1549
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	669
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	529
	05-06	*	< LOD	< LOD	< LOD	< LOD	1491
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	537
	05-06	*	< LOD	< LOD	< LOD	< LOD	1653
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	206
	05-06	*	< LOD	< LOD	< LOD	< LOD	773
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	205
	05-06	*	< LOD	< LOD	< LOD	< LOD	816
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	564
	05-06	*	< LOD	< LOD	< LOD	< LOD	1322

Limit of detection (LOD, see Data Analysis section) for Survey years 03-04 and 05-06 are 0.3 and 0.3 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Nitrobenzene_BiomonitoringSummary.html

Blood Styrene

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	.080 (.050-.100)	.130 (.090-.200)	.200 (.140-.260)	950
	03-04	*	< LOD	.050 (.044-.061)	.089 (.081-.097)	.120 (.110-.130)	1245
	05-06	*	< LOD	.047 (.042-.056)	.099 (.086-.113)	.135 (.120-.160)	2808
Age group							
12-19 years	05-06	*	< LOD	.039 (.031-.049)	.058 (.050-.071)	.072 (.059-.089)	827
20-59 years	01-02	*	< LOD	.080 (.050-.100)	.130 (.090-.200)	.200 (.140-.260)	950
	03-04	*	< LOD	.050 (.044-.061)	.089 (.081-.097)	.120 (.110-.130)	1245
	05-06	*	< LOD	.051 (.044-.065)	.110 (.094-.120)	.150 (.120-.180)	1380
60 years and older	05-06	*	< LOD	.041 (.033-.055)	.083 (.072-.110)	.130 (.100-.150)	601
Gender							
Males	01-02	*	< LOD	.080 (.050-.110)	.140 (.100-.230)	.220 (.140-.340)	445
	03-04	*	< LOD	.056 (.045-.068)	.089 (.081-.100)	.120 (.095-.150)	608
	05-06	*	< LOD	.052 (.043-.067)	.110 (.096-.120)	.142 (.120-.170)	1326
Females	01-02	*	< LOD	.070 (.050-.100)	.120 (.090-.180)	.180 (.120-.260)	505
	03-04	*	< LOD	.048 (.040-.060)	.090 (.074-.100)	.110 (.097-.140)	637
	05-06	*	< LOD	.045 (.038-.052)	.087 (.077-.100)	.129 (.107-.150)	1482
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	.050 (.030-.070)	.090 (.070-.110)	.120 (.090-.170)	225
	03-04	*	< LOD	< LOD	.049 (.033-.066)	.062 (.048-.100)	241
	05-06	*	< LOD	.035 (<LOD-.047)	.060 (.042-.092)	.092 (.064-.110)	603
Non-Hispanic blacks	01-02	*	.040 (<LOD-.080)	.100 (.040-.180)	.180 (.110-.350)	.300 (.170-1.00)	192
	03-04	*	< LOD	.060 (.047-.077)	.100 (.077-.130)	.130 (.110-.160)	264
	05-06	*	< LOD	.056 (.047-.064)	.100 (.079-.120)	.145 (.115-.180)	781
Non-Hispanic whites	01-02	*	.030 (<LOD-.050)	.080 (.050-.110)	.130 (.090-.210)	.200 (.130-.260)	462
	03-04	*	< LOD	.056 (.046-.067)	.096 (.083-.110)	.130 (.110-.140)	646
	05-06	*	< LOD	.049 (.043-.058)	.103 (.087-.120)	.140 (.120-.174)	1217

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.03, 0.03, and 0.03 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Styrene_BiomonitoringSummary.html

Blood Tetrachloroethene (Perchloroethylene)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	.050 (<LOD-.060)	.100 (.070-.150)	.190 (.130-.260)	978
	03-04	*	< LOD	< LOD	.076 (.060-.097)	.140 (.091-.300)	1317
	05-06	*	< LOD	< LOD	.070 (.056-.088)	.126 (.098-.170)	2940
Age group							
12-19 years	05-06	*	< LOD	< LOD	.049 (<LOD-.062)	.079 (.057-.106)	865
20-59 years	01-02	*	< LOD	.050 (<LOD-.060)	.100 (.070-.150)	.190 (.130-.260)	978
	03-04	*	< LOD	< LOD	.076 (.060-.097)	.140 (.091-.300)	1317
	05-06	*	< LOD	< LOD	.074 (.056-.100)	.130 (.098-.199)	1458
60 years and older	05-06	*	< LOD	< LOD	.071 (.054-.092)	.113 (.092-.190)	617
Gender							
Males	01-02	*	< LOD	.050 (<LOD-.060)	.110 (.070-.170)	.210 (.170-.340)	457
	03-04	*	< LOD	< LOD	.082 (.060-.140)	.230 (.097-.410)	639
	05-06	*	< LOD	< LOD	.085 (.062-.126)	.172 (.113-.300)	1404
Females	01-02	*	< LOD	.050 (<LOD-.060)	.100 (.070-.140)	.150 (.100-.220)	521
	03-04	*	< LOD	< LOD	.069 (.050-.091)	.120 (.085-.180)	678
	05-06	*	< LOD	< LOD	.056 (.049-.071)	.090 (.070-.130)	1536
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	.060 (<LOD-.070)	.070 (.060-.230)	226
	03-04	*	< LOD	< LOD	.049 (<LOD-.097)	.100 (.054-.180)	248
	05-06	*	< LOD	< LOD	.069 (.056-.074)	.130 (.073-.284)	713
Non-Hispanic blacks	01-02	*	< LOD	< LOD	.070 (.050-.110)	.110 (.060-.190)	195
	03-04	*	< LOD	< LOD	.086 (.050-.220)	.220 (.082-.360)	284
	05-06	*	< LOD	< LOD	.074 (.057-.112)	.120 (.069-.280)	776
Non-Hispanic whites	01-02	*	< LOD	.050 (<LOD-.070)	.110 (.090-.170)	.210 (.150-.260)	487
	03-04	*	< LOD	< LOD	.072 (.060-.091)	.140 (.085-.330)	686
	05-06	*	< LOD	< LOD	.070 (.053-.090)	.130 (.092-.180)	1229

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.048, 0.048, and 0.048 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/HalogenatedSolvents_BiomonitoringSummary.html

Blood Tetrachloromethane (Carbon tetrachloride)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	.010 (<LOD-.010)	.010 (<LOD-.140)	.020 (.010-.050)	742
	03-04	*	< LOD	< LOD	< LOD	< LOD	1362
	05-06	*	< LOD	< LOD	< LOD	< LOD	2979
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	882
20-59 years	01-02	*	< LOD	.010 (<LOD-.010)	.010 (<LOD-.140)	.020 (.010-.050)	742
	03-04	*	< LOD	< LOD	< LOD	< LOD	1362
	05-06	*	< LOD	< LOD	< LOD	< LOD	1476
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	621
Gender							
Males	01-02	*	< LOD	.010 (<LOD-.010)	.010 (<LOD-.140)	.010 (.010-.040)	364
	03-04	*	< LOD	< LOD	< LOD	< LOD	667
	05-06	*	< LOD	< LOD	< LOD	< LOD	1419
Females	01-02	*	< LOD	.010 (<LOD-.010)	.010 (<LOD-.090)	.040 (.010-.080)	378
	03-04	*	< LOD	< LOD	< LOD	< LOD	695
	05-06	*	< LOD	< LOD	< LOD	< LOD	1560
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	.010 (<LOD-.750)	.020 (.010-.050)	193
	03-04	*	< LOD	< LOD	< LOD	< LOD	266
	05-06	*	< LOD	< LOD	< LOD	< LOD	740
Non-Hispanic blacks	01-02	*	< LOD	.010 (<LOD-.010)	.010 (<LOD-.020)	.010 (<LOD-.020)	132
	03-04	*	< LOD	< LOD	< LOD	< LOD	299
	05-06	*	< LOD	< LOD	< LOD	< LOD	779
Non-Hispanic whites	01-02	*	< LOD	.010 (<LOD-.010)	.010 (<LOD-.140)	.030 (.010-.080)	366
	03-04	*	< LOD	< LOD	< LOD	< LOD	692
	05-06	*	< LOD	< LOD	< LOD	< LOD	1237

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.01, 0.005, and 0.005 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/OtherHalogenatedSolvents_BiomonitoringSummary.html

Blood Toluene

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	.156 (.122-.198)	.160 (.120-.220)	.340 (.260-.430)	.670 (.480-.950)	1.06 (.700-1.43)	954
	03-04	.114 (.100-.129)	.096 (.087-.110)	.220 (.180-.260)	.430 (.380-.550)	.680 (.560-.880)	1336
	05-06	.137 (.123-.152)	.120 (.110-.130)	.230 (.200-.262)	.550 (.481-.640)	.814 (.702-.937)	3050
Age group							
12-19 years	05-06	.110 (.098-.122)	.100 (.094-.120)	.170 (.150-.180)	.280 (.240-.310)	.400 (.300-.610)	907
20-59 years	01-02	.156 (.122-.198)	.160 (.120-.220)	.340 (.260-.430)	.670 (.480-.950)	1.06 (.700-1.43)	954
	03-04	.114 (.100-.129)	.096 (.087-.110)	.220 (.180-.260)	.430 (.380-.550)	.680 (.560-.880)	1336
	05-06	.147 (.133-.163)	.120 (.110-.137)	.260 (.210-.340)	.594 (.505-.720)	.900 (.730-1.10)	1505
60 years and older	05-06	.124 (.100-.154)	.114 (.097-.138)	.190 (.167-.230)	.520 (.370-.600)	.720 (.600-.814)	638
Gender							
Males	01-02	.165 (.130-.209)	.170 (.120-.230)	.360 (.260-.520)	.780 (.580-1.06)	1.22 (.850-1.43)	450
	03-04	.128 (.112-.148)	.110 (.096-.130)	.250 (.190-.310)	.500 (.380-.660)	.730 (.590-1.10)	647
	05-06	.152 (.139-.166)	.130 (.120-.140)	.280 (.240-.330)	.640 (.550-.720)	.920 (.790-1.10)	1441
Females	01-02	.147 (.114-.190)	.150 (.110-.220)	.320 (.240-.390)	.550 (.400-.740)	.810 (.530-1.63)	504
	03-04	.101 (.086-.118)	.085 (.070-.100)	.190 (.150-.230)	.410 (.340-.500)	.580 (.480-.750)	689
	05-06	.124 (.107-.144)	.110 (.097-.130)	.190 (.170-.230)	.470 (.380-.550)	.690 (.550-.880)	1609
Race/ethnicity							
Mexican Americans	01-02	.136 (.106-.176)	.140 (.080-.210)	.270 (.210-.340)	.550 (.400-.980)	.990 (.500-1.30)	219
	03-04	.084 (.074-.096)	.076 (.064-.091)	.120 (.100-.170)	.280 (.170-.410)	.400 (.310-.620)	253
	05-06	.110 (.097-.125)	.110 (.096-.120)	.160 (.140-.170)	.240 (.220-.290)	.340 (.290-.460)	737
Non-Hispanic blacks	01-02	.137 (.089-.210)	.150 (.070-.200)	.310 (.200-.460)	.690 (.390-1.19)	1.15 (.660-1.69)	194
	03-04	.105 (.077-.144)	.095 (.070-.130)	.200 (.130-.330)	.440 (.290-.620)	.620 (.480-.710)	297
	05-06	.139 (.120-.161)	.120 (.099-.140)	.230 (.190-.290)	.450 (.370-.550)	.670 (.550-.830)	796
Non-Hispanic whites	01-02	.165 (.125-.217)	.170 (.120-.240)	.350 (.270-.450)	.710 (.450-1.12)	1.14 (.710-1.63)	467
	03-04	.123 (.110-.139)	.100 (.092-.120)	.240 (.210-.280)	.500 (.400-.590)	.750 (.590-.940)	685
	05-06	.144 (.125-.166)	.130 (.110-.140)	.260 (.210-.330)	.600 (.510-.710)	.880 (.760-.990)	1291

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.025, 0.025, and 0.025 respectively.
 < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Toluene_BiomonitoringSummary.html

Blood Trichloroethene (Trichloroethylene)

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	922
	03-04	*	< LOD	< LOD	< LOD	< LOD	1228
	05-06	*	< LOD	< LOD	< LOD	< LOD	3178
Age group							
12-19 years	05-06	*	< LOD	< LOD	< LOD	< LOD	937
20-59 years	01-02	*	< LOD	< LOD	< LOD	< LOD	922
	03-04	*	< LOD	< LOD	< LOD	< LOD	1228
	05-06	*	< LOD	< LOD	< LOD	< LOD	1562
60 years and older	05-06	*	< LOD	< LOD	< LOD	< LOD	679
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	434
	03-04	*	< LOD	< LOD	< LOD	< LOD	604
	05-06	*	< LOD	< LOD	< LOD	< LOD	1504
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	488
	03-04	*	< LOD	< LOD	< LOD	< LOD	624
	05-06	*	< LOD	< LOD	< LOD	< LOD	1674
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	228
	03-04	*	< LOD	< LOD	< LOD	< LOD	224
	05-06	*	< LOD	< LOD	< LOD	< LOD	773
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	191
	03-04	*	< LOD	< LOD	< LOD	< LOD	266
	05-06	*	< LOD	< LOD	< LOD	< LOD	828
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	441
	03-04	*	< LOD	< LOD	< LOD	< LOD	644
	05-06	*	< LOD	< LOD	< LOD	< LOD	1341

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.012, 0.012, and 0.012 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/HalogenatedSolvents_BiomonitoringSummary.html

Blood *m*- and *p*-Xylene

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	.156 (.124-.198)	.150 (.110-.200)	.280 (.190-.430)	.500 (.370-.690)	.670 (.500-.890)	962
	03-04	.136 (.123-.150)	.130 (.120-.150)	.200 (.190-.210)	.280 (.260-.300)	.340 (.310-.400)	1346
	05-06	.132 (.122-.143)	.120 (.110-.130)	.190 (.170-.210)	.301 (.270-.330)	.410 (.360-.450)	3133
Age group							
12-19 years	05-06	.118 (.106-.130)	.120 (.110-.120)	.160 (.150-.170)	.220 (.190-.270)	.330 (.240-.380)	923
20-59 years	01-02	.156 (.124-.198)	.150 (.110-.200)	.280 (.190-.430)	.500 (.370-.690)	.670 (.500-.890)	962
	03-04	.136 (.123-.150)	.130 (.120-.150)	.200 (.190-.210)	.280 (.260-.300)	.340 (.310-.400)	1346
	05-06	.137 (.127-.147)	.130 (.120-.140)	.200 (.180-.220)	.320 (.280-.350)	.430 (.360-.480)	1542
60 years and older	05-06	.126 (.112-.142)	.120 (.100-.130)	.184 (.150-.220)	.303 (.240-.360)	.400 (.335-.463)	668
Gender							
Males	01-02	.155 (.121-.200)	.140 (.110-.190)	.280 (.180-.440)	.510 (.350-.700)	.700 (.500-.890)	455
	03-04	.149 (.134-.166)	.140 (.130-.170)	.220 (.200-.230)	.290 (.270-.320)	.380 (.300-.490)	654
	05-06	.145 (.135-.157)	.130 (.130-.147)	.220 (.190-.240)	.340 (.310-.370)	.460 (.430-.490)	1480
Females	01-02	.157 (.124-.199)	.150 (.110-.220)	.270 (.190-.410)	.480 (.360-.650)	.650 (.460-.890)	507
	03-04	.124 (.112-.138)	.130 (.110-.140)	.180 (.160-.200)	.270 (.240-.300)	.340 (.290-.390)	692
	05-06	.120 (.109-.132)	.110 (.100-.123)	.170 (.151-.197)	.260 (.220-.320)	.340 (.300-.420)	1653
Race/ethnicity							
Mexican Americans	01-02	.134 (.097-.184)	.130 (.090-.180)	.240 (.150-.370)	.400 (.290-.540)	.540 (.380-1.10)	223
	03-04	.132 (.109-.160)	.120 (.110-.150)	.180 (.140-.230)	.250 (.220-.320)	.360 (.260-.490)	257
	05-06	.108 (.093-.125)	.110 (.098-.120)	.150 (.130-.170)	.220 (.170-.290)	.270 (.220-.370)	755
Non-Hispanic blacks	01-02	.147 (.107-.202)	.140 (.090-.220)	.280 (.160-.460)	.470 (.330-.850)	.590 (.420-1.08)	198
	03-04	.117 (.094-.146)	.120 (.099-.140)	.170 (.140-.220)	.270 (.210-.330)	.330 (.260-.450)	297
	05-06	.131 (.120-.144)	.120 (.110-.140)	.190 (.170-.210)	.280 (.250-.320)	.350 (.320-.392)	823
Non-Hispanic whites	01-02	.163 (.124-.214)	.160 (.110-.220)	.300 (.190-.480)	.520 (.370-.740)	.700 (.510-.960)	468
	03-04	.141 (.128-.155)	.140 (.130-.150)	.210 (.190-.220)	.290 (.270-.310)	.350 (.330-.400)	690
	05-06	.136 (.125-.147)	.130 (.120-.140)	.200 (.180-.220)	.324 (.280-.350)	.430 (.368-.480)	1323

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.034, 0.034, and 0.034 respectively. < LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Xylenes_BiomonitoringSummary.html

Blood *o*-Xylene

Geometric mean and selected percentiles of whole blood concentrations (in ng/mL) for the U.S. population from the National Health and Nutrition Examination Survey.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	.070 (.050-.080)	.100 (.080-.130)	.130 (.110-.180)	981
	03-04	*	< LOD	.051 (<LOD-.057)	.072 (.066-.079)	.090 (.081-.097)	1365
	05-06	.037 (.034-.040)	.035 (.032-.039)	.053 (.047-.061)	.083 (.072-.094)	.110 (.094-.120)	3153
Age group							
12-19 years	05-06	.032 (.030-.034)	.031 (.029-.034)	.044 (.040-.049)	.062 (.053-.075)	.088 (.072-.100)	929
20-59 years	01-02	*	< LOD	.070 (.050-.080)	.100 (.080-.130)	.130 (.110-.180)	981
	03-04	*	< LOD	.051 (<LOD-.057)	.072 (.066-.079)	.090 (.081-.097)	1365
	05-06	.038 (.034-.041)	.036 (.032-.039)	.055 (.048-.062)	.083 (.073-.094)	.110 (.093-.120)	1547
60 years and older	05-06	.038 (.033-.044)	.036 (.032-.042)	.054 (.046-.065)	.092 (.065-.115)	.115 (.094-.150)	677
Gender							
Males	01-02	*	< LOD	.070 (.050-.090)	.110 (.080-.140)	.150 (.110-.180)	465
	03-04	*	< LOD	.057 (.050-.060)	.074 (.068-.084)	.096 (.084-.110)	667
	05-06	.040 (.036-.045)	.038 (.033-.041)	.058 (.050-.070)	.092 (.082-.100)	.120 (.100-.140)	1487
Females	01-02	*	< LOD	.060 (.050-.080)	.100 (.070-.130)	.120 (.100-.180)	516
	03-04	*	< LOD	< LOD	.067 (.059-.076)	.085 (.074-.095)	698
	05-06	.034 (.032-.037)	.034 (.031-.036)	.049 (.044-.054)	.072 (.060-.086)	.094 (.078-.110)	1666
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	.060 (<LOD-.080)	.090 (.080-.110)	.120 (.090-.270)	227
	03-04	*	< LOD	.052 (<LOD-.067)	.074 (.067-.080)	.100 (.075-.130)	265
	05-06	.030 (.027-.033)	.029 (.026-.031)	.040 (.036-.045)	.057 (.050-.069)	.079 (.062-.097)	777
Non-Hispanic blacks	01-02	*	< LOD	.050 (<LOD-.070)	.100 (.080-.110)	.110 (.100-.160)	197
	03-04	*	< LOD	< LOD	.065 (.055-.078)	.084 (.068-.100)	301
	05-06	.035 (.033-.038)	.034 (.031-.038)	.049 (.045-.054)	.073 (.064-.077)	.088 (.078-.100)	824
Non-Hispanic whites	01-02	*	< LOD	.070 (.050-.090)	.110 (.080-.140)	.140 (.110-.180)	483
	03-04	*	< LOD	.052 (<LOD-.058)	.073 (.066-.082)	.090 (.081-.100)	694
	05-06	.039 (.035-.043)	.037 (.033-.041)	.056 (.049-.064)	.087 (.074-.100)	.110 (.096-.130)	1319

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02, 03-04, and 05-06 are 0.049, 0.049, and 0.024 respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Xylenes_BiomonitoringSummary.html

Special Analysis of Pooled Samples for Select Chemicals

Statement of need

In the National Biomonitoring Program, the decision to pool samples before making chemical measurements has been driven by at least two factors:

- The need to improve the sensitivity of the measurement; that is, the chemical concentrations are so low that a larger sample volume is necessary to achieve lower limits of detection and greater likelihood of detectable results; and
- The need to reduce the number of samples analyzed, based on weighing the costs of the analysis against a low frequency of detectable results.

Pooled Samples in the *Updated Tables, September 2013*

This release of the *Updated Tables* includes NHANES 2005-2006 and 2007-2008 results for chemicals that were measured in samples created from pools of serum samples. The historical data tables for these chemicals with results for NHANES survey periods prior to 2005-2006 are included and show results for individual sample measurements. Each historical data table is presented before its corresponding pooled sample data table. A few chemicals (e.g., PCBs 77, 114 and 123; the brominated flame retardant BDE 209) and the chemical groups brominated dibenzo-*p*-dioxins and furans were not measured prior to NHANES 2005-2006; therefore, only the pooled data tables are presented. In addition, several chemicals were not measured after NHANES 2003-2004, so there are no pooled data tables for these (e.g., five organochlorine pesticides).

Overview of Sample Pooling

Beginning with NHANES 2005-2006, the CDC is using a weighted pooled-sample design to measure serum concentrations of dioxins, furans, polychlorinated biphenyls (PCBs), organochlorine pesticides and metabolites, and the brominated flame retardants (polybrominated diphenyl ethers and the polybrominated diphenyl, PBB 153). In previous NHANES survey periods, measurements were made on individual serum samples from the U.S. population using stratified multistage selection.

Although NHANES sampling weights can be incorporated to make pooled sample estimates representative of the non-institutionalized U.S. population, there are statistical estimation challenges. Measurements of these chemicals in individuals tend to have a log-normal distribution with central tendency best estimated using a geometric mean. However, the measured value for a pooled sample is comparable to an arithmetic average of measurements in individuals. Consequently, the pooled sample result is expected to be higher than the geometric mean of multiple individual results. Another challenge is that direct calculation of the design effects required for accurate standard error and confidence interval estimation is not possible because samples are pooled across the design cells of the original survey. For these reasons, data tables showing the pooled sample results present only weighted arithmetic means and unadjusted standard errors for each category.

Interpreting the Data

The following cautions and suggestions apply to the pooled sample results and data tables:

a) Individual sample data, as explained above, tend to be log-normally distributed with central tendency best estimated using a geometric mean. Pooled data are comparable to an arithmetic mean of individual sample results and therefore, pooled data results are expected to be higher than the geometric mean of individual sample results. Therefore, the weighted arithmetic means from pooled samples are expected to be higher than the geometric means provided in the historical (companion) data tables or in other publications based on data provided on individual results.

b) The standard errors are unadjusted and therefore, do not reflect the design effects of the survey. In many cases, the standard errors are based on very few pooled sample measurements and cannot be expected to accurately reflect the true imprecision of the weighted arithmetic mean estimates. Therefore, when the unadjusted standard error was more than 30% of the weighted arithmetic mean, then a value is not reported because it is not reliable.

Additional information about sample pooling is available from the following sources:

Geometric mean estimation from pooled samples (2007)

<http://www.ncbi.nlm.nih.gov/pubmed/17618673>

Characterizing populations of individuals using pooled samples (2010)

<http://www.ncbi.nlm.nih.gov/pubmed/19002216>

Use of pooled samples from the National Health and Nutrition Examination Survey (2012)

<http://www.ncbi.nlm.nih.gov/pubmed/22492247>

Serum Oxychlordane (lipid adjusted)

Metabolite of Chlordane

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	20.8 (17.8-23.0)	34.4 (30.5-38.6)	44.8 (40.2-49.6)	1661
	01-02	11.4 (<LOD-12.5)	11.1 (<LOD-12.5)	21.7 (19.3-24.4)	36.4 (31.5-41.4)	49.7 (42.0-61.2)	2249
	03-04	9.37 (8.69-10.1)	10.3 (9.20-11.0)	18.0 (16.8-20.1)	29.0 (26.8-32.1)	37.7 (34.8-43.8)	1978
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	663
	01-02	*	< LOD	< LOD	< LOD	11.5 (<LOD-12.6)	752
	03-04	*	< LOD	< LOD	9.20 (<LOD-11.5)	11.5 (8.10-18.9)	595
20 years and older	99-00	*	< LOD	23.3 (21.0-25.9)	37.7 (32.3-43.5)	47.7 (43.1-50.8)	998
	01-02	12.9 (11.7-14.3)	13.3 (11.4-14.9)	23.9 (21.2-26.7)	38.5 (33.4-45.9)	53.1 (44.1-65.9)	1497
	03-04	10.6 (9.82-11.5)	11.4 (10.6-12.4)	19.9 (17.9-21.5)	31.3 (28.8-33.2)	39.2 (36.5-44.8)	1383
Gender							
Males	99-00	*	< LOD	18.1 (16.1-19.6)	31.3 (25.9-38.2)	42.4 (35.3-49.6)	793
	01-02	11.1 (<LOD-12.6)	11.1 (<LOD-12.6)	20.6 (16.6-24.9)	33.1 (27.5-43.8)	48.1 (40.2-56.9)	1049
	03-04	9.10 (8.20-10.1)	9.90 (8.30-11.2)	17.1 (15.6-18.4)	27.6 (25.3-32.2)	36.0 (32.7-39.2)	963
Females	99-00	*	< LOD	22.3 (20.1-25.9)	36.9 (31.5-40.3)	46.2 (39.1-51.8)	868
	01-02	11.7 (10.7-12.7)	11.0 (<LOD-12.9)	23.1 (20.7-25.0)	37.5 (34.5-42.1)	52.8 (42.7-70.0)	1200
	03-04	9.63 (8.89-10.4)	10.6 (9.10-11.3)	20.1 (17.4-21.7)	30.3 (27.5-32.7)	41.9 (36.3-45.5)	1015
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	16.3 (<LOD-19.9)	28.9 (18.8-42.0)	39.9 (26.8-61.0)	628
	01-02	*	< LOD	13.9 (11.0-18.4)	27.2 (21.0-33.1)	37.9 (29.9-42.0)	557
	03-04	*	< LOD	12.8 (10.1-15.8)	22.9 (15.8-31.4)	31.4 (22.4-51.6)	462
Non-Hispanic blacks	99-00	*	< LOD	18.7 (<LOD-32.2)	39.9 (26.5-47.3)	48.6 (43.5-65.5)	350
	01-02	11.7 (<LOD-13.6)	< LOD	22.8 (17.2-28.3)	41.4 (30.6-53.7)	56.5 (41.8-73.5)	501
	03-04	8.74 (<LOD-10.2)	8.70 (<LOD-10.6)	18.9 (15.9-21.5)	35.1 (25.4-40.2)	44.2 (37.7-56.8)	493
Non-Hispanic whites	99-00	*	< LOD	21.8 (18.6-24.6)	34.2 (28.9-40.9)	44.0 (37.2-49.8)	559
	01-02	12.1 (11.0-13.3)	11.8 (10.5-13.9)	23.0 (20.1-25.7)	37.5 (31.6-45.1)	52.2 (41.0-67.4)	1031
	03-04	10.2 (9.36-11.1)	11.2 (10.0-12.1)	19.7 (17.2-21.7)	30.3 (26.8-33.6)	37.7 (34.3-45.5)	898

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 14.5, 10.5, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum Oxychlordane (lipid adjusted)

Metabolite of Chlordane

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	4.80	1.1	9
		07-08	3.52	0.52	6
	20-39	05-06	6.50	1.06	12
		07-08	6.93	0.75	15
	40-59	05-06	14.9	1.4	12
		07-08	12.1	0.5	16
60+	05-06	25.4	2.5	15	
	07-08	26.1	1.4	23	
Non-Hispanic White Female	12-19	05-06	2.98	0.3	10
		07-08	3.66	0.74	7
	20-39	05-06	5.64	0.56	16
		07-08	5.79	0.51	13
	40-59	05-06	13.7	1	13
		07-08	11.8	0.8	17
60+	05-06	28.9	2.6	17	
	07-08	31.8	2.3	21	
Non-Hispanic Black Male	12-19	05-06	4.08	0.49	13
		07-08	2.80	0.75	6
	20-39	05-06	4.53	0.9	6
		07-08	5.72	0.28	6
	40-59	05-06	12.0	1	5
		07-08	12.1	1.1	6
60+	05-06	32.7	5.3	5	
	07-08	26.6	2.5	8	
Non-Hispanic Black Female	12-19	05-06	2.92	0.31	14
		07-08	*	*	5
	20-39	05-06	5.56	0.68	7
		07-08	4.45	0.79	8
	40-59	05-06	14.5	0.8	7
		07-08	14.3	1	8
60+	05-06	52.9	11.7	5	
	07-08	38.1	1.7	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum Oxychlordane (lipid adjusted)

Metabolite of Chlordane

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	2.54	0.33	11
		07-08	2.76	0.45	6
	20-39	05-06	5.84	1.22	9
		07-08	4.93	0.38	9
	40-59	05-06	11.0	1.4	4
		07-08	12.0	0.7	6
60+	05-06	18.6	2.4	4	
	07-08	21.9	2.2	5	
Mexican American Female	12-19	05-06	2.39	0.25	16
		07-08	*	*	5
	20-39	05-06	4.39	0.5	9
		07-08	4.17	0.55	8
	40-59	05-06	13.7	1.8	6
		07-08	13.4	2.2	6
	60+	05-06	27.4	3.8	3
		07-08	28.2	4.3	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum Oxychlordane (whole weight)

Metabolite of Chlordane

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	.140 (.120-.150)	.260 (.200-.290)	.310 (.290-.340)	1661
	01-02	.070 (<LOD-.077)	.070 (<LOD-.080)	.140 (.130-.160)	.250 (.220-.300)	.350 (.290-.440)	2249
	03-04	.057 (.053-.062)	.063 (.058-.068)	.119 (.106-.133)	.204 (.189-.213)	.269 (.246-.291)	1978
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	663
	01-02	*	< LOD	< LOD	< LOD	.060 (<LOD-.070)	752
	03-04	*	< LOD	< LOD	.047 (<LOD-.063)	.066 (.048-.092)	595
20 years and older	99-00	*	< LOD	.150 (.140-.180)	.280 (.230-.300)	.330 (.300-.400)	998
	01-02	.082 (.074-.091)	.080 (.070-.090)	.160 (.140-.180)	.270 (.230-.320)	.370 (.310-.450)	1497
	03-04	.067 (.061-.073)	.073 (.066-.079)	.130 (.115-.146)	.210 (.203-.227)	.286 (.258-.320)	1383
Gender							
Males	99-00	*	< LOD	.120 (.100-.140)	.220 (.180-.280)	.300 (.260-.340)	793
	01-02	.069 (<LOD-.079)	.070 (<LOD-.080)	.130 (.120-.160)	.230 (.190-.300)	.320 (.250-.430)	1049
	03-04	.056 (.050-.063)	.063 (.055-.076)	.115 (.100-.126)	.189 (.168-.207)	.258 (.216-.302)	963
Females	99-00	*	< LOD	.140 (.130-.170)	.270 (.200-.310)	.320 (.290-.400)	868
	01-02	.071 (.065-.077)	.070 (<LOD-.080)	.150 (.130-.160)	.260 (.230-.310)	.370 (.280-.510)	1200
	03-04	.058 (.053-.064)	.063 (.057-.068)	.126 (.104-.146)	.208 (.199-.231)	.286 (.245-.331)	1015
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	.100 (<LOD-.130)	.210 (.130-.320)	.290 (.190-.410)	628
	01-02	*	< LOD	.100 (.070-.130)	.200 (.150-.240)	.280 (.210-.360)	557
	03-04	*	< LOD	.083 (.066-.104)	.149 (.108-.230)	.230 (.148-.373)	462
Non-Hispanic blacks	99-00	*	< LOD	.110 (<LOD-.170)	.240 (.170-.290)	.320 (.240-.430)	350
	01-02	.066 (<LOD-.077)	< LOD	.130 (.090-.170)	.260 (.180-.350)	.350 (.240-.560)	501
	03-04	.049 (<LOD-.057)	.050 (<LOD-.062)	.112 (.087-.136)	.225 (.165-.287)	.315 (.253-.348)	493
Non-Hispanic whites	99-00	*	< LOD	.140 (.120-.170)	.270 (.200-.300)	.320 (.280-.380)	559
	01-02	.075 (.068-.083)	.080 (.070-.090)	.150 (.130-.170)	.250 (.220-.310)	.370 (.280-.450)	1031
	03-04	.063 (.058-.070)	.070 (.063-.077)	.128 (.110-.148)	.207 (.190-.223)	.271 (.242-.315)	898

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum Oxychlordane (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.026	0.006	9
		07-08	.018	0.003	6
	20-39	05-06	.049	0.01	12
		07-08	.047	0.006	15
	40-59	05-06	.113	0.013	12
		07-08	.089	0.005	16
60+	05-06	.171	0.015	15	
	07-08	.167	0.009	23	
Non-Hispanic White Female	12-19	05-06	.016	0.002	10
		07-08	.017	0.004	7
	20-39	05-06	.037	0.004	16
		07-08	.035	0.003	13
	40-59	05-06	.101	0.01	13
		07-08	.079	0.006	17
60+	05-06	.210	0.017	17	
	07-08	.214	0.017	21	
Non-Hispanic Black Male	12-19	05-06	.020	0.002	13
		07-08	.013	0.003	6
	20-39	05-06	.027	0.005	6
		07-08	.032	0.002	6
	40-59	05-06	.076	0.006	5
		07-08	.077	0.007	6
60+	05-06	.193	0.028	5	
	07-08	.162	0.017	8	
Non-Hispanic Black Female	12-19	05-06	.014	0.001	14
		07-08	*	*	5
	20-39	05-06	.032	0.004	7
		07-08	.022	0.004	8
	40-59	05-06	.093	0.006	7
		07-08	.086	0.008	8
60+	05-06	.347	0.067	5	
	07-08	.229	0.012	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum Oxychlordane (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.014	0.002	11
		07-08	.014	0.002	6
	20-39	05-06	.041	0.007	9
		07-08	.034	0.004	9
	40-59	05-06	.089	0.014	4
		07-08	.087	0.009	6
60+	05-06	.137	0.025	4	
	07-08	.156	0.011	5	
Mexican American Female	12-19	05-06	.013	0.001	16
		07-08	*	*	5
	20-39	05-06	.029	0.003	9
		07-08	.024	0.004	8
	40-59	05-06	.100	0.011	6
		07-08	.090	0.014	6
	60+	05-06	.208	0.035	3
		07-08	.193	0.03	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum *trans*-Nonachlor (lipid adjusted)

Metabolite of Chlordane

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	18.3 (16.7-20.0)	17.9 (16.1-20.1)	31.9 (28.9-36.0)	55.1 (48.4-62.6)	79.4 (67.6-88.1)	1933
	01-02	17.0 (15.2-18.9)	17.9 (15.5-20.5)	33.7 (30.2-37.2)	56.3 (49.6-66.0)	78.2 (64.0-113)	2286
	03-04	14.7 (13.1-16.5)	14.8 (13.5-17.0)	30.2 (26.7-32.5)	49.0 (42.6-54.7)	68.3 (58.6-82.3)	1955
Age group							
12-19 years	99-00	*	< LOD	< LOD	18.8 (<LOD-20.6)	25.2 (19.1-28.4)	664
	01-02	*	< LOD	< LOD	13.4 (11.8-16.4)	19.2 (15.2-23.5)	758
	03-04	*	< LOD	8.70 (<LOD-12.5)	16.1 (10.7-23.7)	22.6 (16.1-34.6)	589
20 years and older	99-00	20.8 (19.0-22.8)	20.7 (18.0-23.5)	35.4 (30.9-40.3)	59.9 (51.8-67.6)	82.7 (74.9-89.6)	1269
	01-02	19.8 (17.6-22.3)	20.9 (19.0-23.1)	36.6 (32.8-41.1)	60.6 (52.5-69.9)	84.9 (66.0-123)	1528
	03-04	16.9 (15.1-18.9)	17.3 (14.6-20.0)	31.8 (28.9-35.3)	51.4 (45.9-58.6)	74.7 (59.8-90.0)	1366
Gender							
Males	99-00	17.7 (16.5-19.1)	17.2 (14.9-20.1)	30.2 (27.7-34.2)	51.1 (47.3-58.6)	78.2 (60.2-88.1)	922
	01-02	17.0 (14.8-19.5)	18.3 (14.8-21.1)	34.4 (28.3-39.3)	54.8 (45.0-68.9)	78.2 (59.7-113)	1062
	03-04	14.8 (12.7-17.3)	14.6 (12.2-18.0)	30.8 (26.7-35.3)	51.0 (42.0-59.4)	68.6 (56.0-93.8)	955
Females	99-00	18.8 (16.7-21.1)	18.4 (16.1-22.2)	32.9 (29.0-38.3)	59.0 (48.4-67.6)	80.8 (71.5-95.5)	1011
	01-02	17.0 (15.4-18.7)	17.6 (15.0-20.3)	32.8 (30.4-36.7)	56.9 (51.9-65.5)	78.1 (65.5-111)	1224
	03-04	14.5 (13.1-16.1)	15.0 (13.8-16.3)	28.2 (25.3-32.8)	48.1 (41.4-52.7)	68.3 (56.8-79.9)	1000
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	25.1 (22.7-29.5)	40.7 (35.1-51.8)	56.3 (45.8-77.2)	650
	01-02	11.9 (<LOD-14.6)	10.6 (<LOD-14.5)	26.0 (19.3-30.4)	47.9 (36.3-57.2)	59.8 (49.3-74.1)	558
	03-04	10.2 (7.86-13.2)	9.10 (<LOD-11.1)	20.7 (11.1-34.7)	39.5 (25.9-65.5)	62.2 (36.0-93.4)	457
Non-Hispanic blacks	99-00	20.3 (17.0-24.1)	17.5 (15.4-23.5)	35.7 (28.9-45.5)	77.0 (60.8-90.7)	107 (84.0-143)	404
	01-02	18.8 (15.4-22.9)	19.2 (14.7-22.0)	36.8 (28.3-50.5)	73.6 (50.8-110)	112 (68.7-160)	514
	03-04	14.4 (12.2-17.0)	13.8 (11.2-16.3)	30.8 (26.5-36.1)	59.9 (47.7-77.7)	86.6 (56.8-129)	486
Non-Hispanic whites	99-00	19.1 (17.2-21.1)	19.0 (16.9-22.2)	32.8 (28.0-37.6)	52.5 (44.9-64.4)	74.0 (62.3-86.7)	722
	01-02	17.5 (15.6-19.7)	19.0 (16.3-21.1)	34.0 (29.7-38.1)	55.5 (45.9-69.4)	78.7 (59.1-126)	1052
	03-04	15.8 (13.7-18.2)	16.0 (13.8-19.3)	30.8 (26.4-35.0)	48.8 (42.1-55.7)	67.6 (57.5-87.3)	889

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 14.5, 10.5, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum *trans*-Nonachlor (lipid adjusted)

Metabolite of Chlordane

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	6.06	1.08	6
	20-39	05-06	11.3	2	12
		07-08	11.1	1.5	15
	40-59	05-06	26.9	3.9	12
		07-08	21.5	1.2	16
60+	05-06	49.4	7.9	15	
	07-08	50.6	4.2	23	
Non-Hispanic White Female	12-19	05-06	4.94	0.52	10
		07-08	6.70	0.89	7
	20-39	05-06	9.70	1.44	16
		07-08	8.95	0.97	13
	40-59	05-06	21.8	2.3	13
		07-08	16.4	1.5	17
60+	05-06	50.2	6.1	17	
	07-08	51.7	6.1	21	
Non-Hispanic Black Male	12-19	05-06	6.50	0.67	13
		07-08	4.71	0.65	6
	20-39	05-06	8.24	0.86	6
		07-08	8.87	0.97	6
	40-59	05-06	22.0	2.6	5
		07-08	21.7	2.1	6
60+	05-06	66.4	13.2	5	
	07-08	54.4	4.9	8	
Non-Hispanic Black Female	12-19	05-06	4.44	0.54	14
		07-08	*	*	5
	20-39	05-06	9.32	1.22	7
		07-08	6.42	0.6	8
	40-59	05-06	24.1	1.5	7
		07-08	20.4	1.7	8
60+	05-06	89.0	15.4	5	
	07-08	68.0	3.9	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.77 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum *trans*-Nonachlor (lipid adjusted)

Metabolite of Chlordane

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	3.71	0.49	11
		07-08	3.91	0.81	6
	20-39	05-06	7.58	0.75	9
		07-08	8.43	0.97	9
	40-59	05-06	18.8	2.4	4
		07-08	24.5	2.9	6
60+	05-06	30.7	6.3	4	
	07-08	40.3	4.5	5	
Mexican American Female	12-19	05-06	3.53	0.23	16
		07-08	*	*	5
	20-39	05-06	6.40	0.75	9
		07-08	7.55	0.9	8
	40-59	05-06	18.5	1.7	6
		07-08	22.6	4.4	6
60+	05-06	46.4	9.8	3	
	07-08	46.9	8	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.77 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum *trans*-Nonachlor (whole weight)

Metabolite of Chlordane

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.109 (.099-.119)	.110 (.090-.120)	.210 (.190-.240)	.370 (.330-.420)	.550 (.470-.630)	1933
	01-02	.104 (.093-.116)	.110 (.100-.120)	.220 (.190-.250)	.390 (.330-.480)	.590 (.430-.800)	2286
	03-04	.089 (.080-.100)	.094 (.084-.108)	.191 (.171-.211)	.324 (.290-.371)	.470 (.410-.558)	1955
Age group							
12-19 years	99-00	*	< LOD	< LOD	.090 (<LOD-.110)	.120 (.100-.130)	664
	01-02	*	< LOD	< LOD	.070 (.060-.080)	.090 (.080-.130)	758
	03-04	*	< LOD	.041 (<LOD-.060)	.081 (.054-.117)	.109 (.081-.161)	589
20 years and older	99-00	.128 (.116-.141)	.130 (.110-.150)	.230 (.210-.260)	.400 (.360-.460)	.580 (.490-.690)	1269
	01-02	.125 (.111-.141)	.130 (.120-.150)	.240 (.210-.280)	.420 (.350-.540)	.640 (.470-.840)	1528
	03-04	.106 (.095-.119)	.112 (.096-.127)	.210 (.186-.237)	.355 (.301-.405)	.520 (.430-.594)	1366
Gender							
Males	99-00	.106 (.098-.114)	.100 (.090-.120)	.210 (.180-.220)	.350 (.310-.400)	.520 (.400-.630)	922
	01-02	.105 (.091-.122)	.110 (.090-.130)	.220 (.190-.260)	.380 (.310-.500)	.580 (.390-.830)	1062
	03-04	.092 (.079-.108)	.098 (.085-.126)	.202 (.177-.232)	.343 (.285-.395)	.458 (.395-.594)	955
Females	99-00	.111 (.099-.125)	.110 (.090-.130)	.220 (.190-.250)	.390 (.310-.460)	.580 (.460-.690)	1011
	01-02	.103 (.093-.113)	.110 (.090-.120)	.220 (.180-.240)	.400 (.340-.450)	.590 (.430-.830)	1224
	03-04	.087 (.078-.097)	.092 (.079-.098)	.183 (.161-.210)	.317 (.286-.367)	.470 (.409-.565)	1000
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	.170 (.120-.210)	.310 (.240-.340)	.390 (.320-.520)	650
	01-02	.071 (<LOD-.091)	.060 (<LOD-.090)	.160 (.120-.210)	.330 (.270-.390)	.470 (.360-.590)	558
	03-04	.062 (.047-.082)	.055 (<LOD-.069)	.135 (.069-.237)	.288 (.158-.559)	.414 (.279-.651)	457
Non-Hispanic blacks	99-00	.112 (.093-.134)	.100 (.080-.130)	.220 (.180-.300)	.490 (.340-.600)	.760 (.510-.960)	404
	01-02	.106 (.085-.131)	.110 (.080-.130)	.220 (.160-.310)	.490 (.320-.680)	.680 (.410-1.20)	514
	03-04	.080 (.068-.096)	.078 (.061-.093)	.186 (.145-.242)	.417 (.272-.565)	.573 (.497-.684)	486
Non-Hispanic whites	99-00	.116 (.104-.129)	.120 (.100-.140)	.220 (.190-.240)	.370 (.300-.440)	.510 (.440-.630)	722
	01-02	.108 (.096-.122)	.120 (.100-.130)	.220 (.190-.250)	.390 (.310-.490)	.600 (.400-.930)	1052
	03-04	.098 (.085-.113)	.103 (.090-.124)	.205 (.173-.234)	.327 (.288-.390)	.461 (.397-.576)	889

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum *trans*-Nonachlor(whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	.030	0.005	6
	20-39	05-06	.086	0.019	12
		07-08	.076	0.011	15
	40-59	05-06	.206	0.034	12
		07-08	.159	0.012	16
60+	05-06	.331	0.049	15	
	07-08	.325	0.028	23	
Non-Hispanic White Female	12-19	05-06	.026	0.003	10
		07-08	.031	0.004	7
	20-39	05-06	.063	0.01	16
		07-08	.054	0.007	13
	40-59	05-06	.161	0.019	13
		07-08	.108	0.009	17
60+	05-06	.362	0.04	17	
	07-08	.352	0.045	21	
Non-Hispanic Black Male	12-19	05-06	.032	0.003	13
		07-08	.022	0.003	6
	20-39	05-06	.049	0.006	6
		07-08	.050	0.006	6
	40-59	05-06	.140	0.014	5
		07-08	.138	0.013	6
60+	05-06	.390	0.071	5	
	07-08	.331	0.032	8	
Non-Hispanic Black Female	12-19	05-06	.022	0.003	14
		07-08	*	*	5
	20-39	05-06	.054	0.007	7
		07-08	.032	0.003	8
	40-59	05-06	.154	0.009	7
		07-08	.124	0.012	8
60+	05-06	.588	0.088	5	
	07-08	.409	0.027	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum *trans*-Nonachlor(whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.020	0.002	11
		07-08	.020	0.004	6
	20-39	05-06	.054	0.005	9
		07-08	.058	0.008	9
	40-59	05-06	.153	0.022	4
		07-08	.176	0.024	6
60+	05-06	.228	0.06	4	
	07-08	.289	0.028	5	
Mexican American Female	12-19	05-06	.019	0.001	16
		07-08	*	*	5
	20-39	05-06	.042	0.005	9
		07-08	.044	0.006	8
	40-59	05-06	.135	0.011	6
		07-08	.151	0.029	6
	60+	05-06	.360	0.095	3
		07-08	.323	0.058	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/ChlordaneHeptachlor_FactSheet.html

Serum *p,p'*-Dichlorodiphenyltrichloroethane (DDT) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	28.0 (21.9-34.0)	1679
	01-02	*	< LOD	< LOD	< LOD	26.6 (22.5-36.0)	2305
	03-04	*	< LOD	< LOD	11.9 (10.0-15.1)	19.5 (15.1-27.5)	1965
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	677
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	< LOD	< LOD	< LOD	9.10 (<LOD-12.2)	595
20 years and older	99-00	*	< LOD	< LOD	< LOD	30.5 (23.0-37.3)	1002
	01-02	*	< LOD	< LOD	< LOD	28.1 (23.8-39.0)	1549
	03-04	*	< LOD	< LOD	12.9 (10.3-16.5)	20.7 (15.9-28.7)	1370
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	25.1 (<LOD-39.3)	799
	01-02	*	< LOD	< LOD	< LOD	21.6 (<LOD-25.8)	1073
	03-04	*	< LOD	< LOD	10.6 (9.10-13.7)	15.2 (11.8-26.9)	959
Females	99-00	*	< LOD	< LOD	< LOD	29.4 (23.0-35.8)	880
	01-02	*	< LOD	< LOD	18.3 (<LOD-21.9)	36.6 (25.5-54.3)	1232
	03-04	*	< LOD	< LOD	14.0 (10.8-17.5)	21.0 (18.0-27.8)	1006
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	61.3 (27.0-155)	155 (59.3-590)	635
	01-02	*	< LOD	< LOD	83.1 (33.3-236)	293 (104-541)	566
	03-04	*	< LOD	8.90 (<LOD-12.9)	24.0 (18.6-33.3)	48.6 (31.1-71.1)	461
Non-Hispanic blacks	99-00	*	< LOD	< LOD	22.3 (<LOD-31.5)	31.5 (23.2-65.0)	356
	01-02	*	< LOD	< LOD	23.2 (<LOD-40.9)	40.9 (21.2-95.8)	514
	03-04	*	< LOD	9.00 (<LOD-10.4)	17.5 (14.8-23.2)	30.7 (19.0-53.4)	490
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	564
	01-02	*	< LOD	< LOD	< LOD	17.9 (<LOD-20.7)	1061
	03-04	*	< LOD	< LOD	9.70 (8.50-11.2)	12.9 (10.7-16.6)	890

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 20.7, 17.4, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-DDT (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	2.23	0.15	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	2.05	0.37	15
	40-59	05-06	3.79	0.26	12
		07-08	2.81	0.25	16
60+	05-06	5.68	0.54	15	
	07-08	4.85	0.47	23	
Non-Hispanic White Female	12-19	05-06	2.27	0.45	10
		07-08	*	*	7
	20-39	05-06	2.72	0.22	16
		07-08	*	*	13
	40-59	05-06	3.81	0.41	13
		07-08	2.90	0.23	17
60+	05-06	6.21	0.41	17	
	07-08	6.37	0.91	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	1.80	0.37	6
	40-59	05-06	*	*	5
		07-08	4.68	0.69	6
60+	05-06	14.9	3.2	5	
	07-08	6.57	0.83	8	
Non-Hispanic Black Female	12-19	05-06	2.59	0.34	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	7.36	1.11	7
		07-08	5.86	0.51	8
60+	05-06	*	*	5	
	07-08	19.3	3	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-DDT (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	18.6	4.2	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	7.14	1.69	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	11.8	3.1	6
	60+	05-06	17.3	4.8	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-Dichlorodiphenyltrichloroethane (DDT) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	.170 (.130-.220)	1679
	01-02	*	< LOD	< LOD	< LOD	.180 (.160-.220)	2305
	03-04	*	< LOD	< LOD	.078 (.065-.097)	.128 (.098-.167)	1965
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	677
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	< LOD	< LOD	< LOD	.048 (<LOD-.069)	595
20 years and older	99-00	*	< LOD	< LOD	< LOD	.190 (.150-.230)	1002
	01-02	*	< LOD	< LOD	< LOD	.200 (.170-.260)	1549
	03-04	*	< LOD	< LOD	.084 (.068-.106)	.142 (.105-.189)	1370
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	.150 (<LOD-.240)	799
	01-02	*	< LOD	< LOD	< LOD	.150 (<LOD-.180)	1073
	03-04	*	< LOD	< LOD	.071 (.059-.095)	.108 (.078-.180)	959
Females	99-00	*	< LOD	< LOD	< LOD	.190 (.150-.230)	880
	01-02	*	< LOD	< LOD	.130 (<LOD-.150)	.240 (.180-.400)	1232
	03-04	*	< LOD	< LOD	.087 (.071-.106)	.146 (.106-.207)	1006
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	.400 (.190-1.00)	1.00 (.330-4.26)	635
	01-02	*	< LOD	< LOD	.530 (.250-1.34)	1.62 (.570-4.01)	566
	03-04	*	< LOD	.063 (<LOD-.079)	.146 (.114-.203)	.313 (.189-.627)	461
Non-Hispanic blacks	99-00	*	< LOD	< LOD	.120 (<LOD-.170)	.180 (.140-.420)	356
	01-02	*	< LOD	< LOD	.130 (<LOD-.290)	.250 (.120-.530)	514
	03-04	*	< LOD	.051 (<LOD-.061)	.112 (.080-.143)	.201 (.132-.343)	490
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	564
	01-02	*	< LOD	< LOD	< LOD	.130 (<LOD-.140)	1061
	03-04	*	< LOD	< LOD	.064 (.054-.075)	.086 (.074-.107)	890

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-DDT (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.012	0.001	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	.014	0.003	15
	40-59	05-06	.028	0.002	12
		07-08	.021	0.002	16
60+	05-06	.038	0.003	15	
	07-08	.031	0.003	23	
Non-Hispanic White Female	12-19	05-06	.012	0.002	10
		07-08	*	*	7
	20-39	05-06	.018	0.002	16
		07-08	*	*	13
	40-59	05-06	.028	0.003	13
		07-08	.019	0.002	17
60+	05-06	.045	0.003	17	
	07-08	.043	0.006	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	.010	0.002	6
	40-59	05-06	*	*	5
		07-08	.030	0.004	6
60+	05-06	.088	0.019	5	
	07-08	.040	0.005	8	
Non-Hispanic Black Female	12-19	05-06	.013	0.002	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	.047	0.007	7
		07-08	.035	0.003	8
60+	05-06	*	*	5	
	07-08	.115	0.017	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-DDT (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	.129	0.029	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	.053	0.015	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	.081	0.022	6
	60+	05-06	.138	0.05	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-Dichlorodiphenyldichloroethene (DDE) (lipid adjusted)

Metabolite of Dichlorodiphenyltrichloroethane

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	260 (226-298)	226 (184-278)	537 (476-631)	1150 (976-1350)	1830 (1410-2300)	1964
	01-02	295 (267-327)	251 (228-278)	598 (521-699)	1410 (1210-1500)	2320 (1830-2780)	2298
	03-04	238 (195-292)	203 (163-275)	509 (376-655)	1170 (836-1570)	1860 (1400-2380)	1956
Age group							
12-19 years	99-00	118 (102-135)	108 (97.7-119)	185 (141-237)	339 (243-479)	528 (339-812)	686
	01-02	124 (106-146)	113 (100-140)	213 (172-253)	319 (282-389)	456 (343-722)	758
	03-04	105 (84.7-129)	93.6 (81.0-114)	167 (123-240)	341 (211-586)	522 (313-1430)	588
20 years and older	99-00	297 (256-344)	269 (213-323)	608 (530-693)	1260 (1030-1550)	2020 (1520-2730)	1278
	01-02	338 (303-376)	285 (249-337)	695 (595-798)	1480 (1310-1700)	2550 (1980-3080)	1540
	03-04	268 (217-332)	233 (175-314)	557 (420-734)	1270 (877-1800)	1990 (1500-2470)	1368
Gender							
Males	99-00	249 (220-283)	223 (182-262)	494 (380-578)	1010 (789-1130)	1430 (1080-2160)	937
	01-02	285 (252-323)	248 (222-285)	520 (441-627)	1160 (937-1360)	1900 (1580-2490)	1069
	03-04	235 (193-288)	200 (164-262)	466 (331-653)	1000 (763-1400)	1610 (1210-2320)	955
Females	99-00	270 (226-322)	234 (184-302)	601 (492-711)	1350 (1040-1720)	2210 (1570-2810)	1027
	01-02	305 (273-341)	256 (219-297)	708 (567-844)	1480 (1410-1710)	2670 (1940-3300)	1229
	03-04	241 (193-301)	207 (161-281)	539 (386-735)	1250 (813-1900)	2010 (1500-2450)	1001
Race/ethnicity							
Mexican Americans	99-00	674 (574-792)	624 (545-701)	1350 (1090-1660)	3090 (2040-4950)	4950 (3070-9350)	657
	01-02	652 (569-747)	561 (455-690)	1400 (1050-1950)	4110 (2520-6550)	7080 (3080-15600)	566
	03-04	444 (362-545)	373 (283-522)	875 (608-1170)	2150 (1520-2470)	3290 (2380-9240)	457
Non-Hispanic blacks	99-00	295 (241-362)	251 (199-313)	668 (492-874)	1850 (1040-2220)	2300 (1560-5680)	416
	01-02	324 (262-400)	248 (223-296)	762 (583-999)	1620 (1180-2980)	3260 (1270-6900)	515
	03-04	262 (233-295)	216 (173-267)	589 (453-747)	1620 (1130-2310)	2860 (1880-3440)	487
Non-Hispanic whites	99-00	217 (189-249)	194 (162-238)	438 (355-507)	825 (647-1010)	1160 (1010-1350)	732
	01-02	253 (226-284)	225 (203-254)	463 (402-558)	1150 (878-1340)	1640 (1410-1940)	1053
	03-04	208 (165-263)	177 (148-238)	417 (302-564)	907 (574-1480)	1490 (909-2300)	888

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 18.6, 8.3, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-DDE (lipid adjusted)

Metabolite of Dichlorodiphenyltrichloroethane

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	113	20	9
		07-08	86.7	7.7	6
	20-39	05-06	*	*	12
		07-08	113	11	15
	40-59	05-06	220	19	12
		07-08	176	14	16
60+	05-06	696	53	15	
	07-08	502	46	23	
Non-Hispanic White Female	12-19	05-06	83.6	6.7	10
		07-08	70.2	14.9	7
	20-39	05-06	102	7	16
		07-08	91.8	6.3	13
	40-59	05-06	276	36	13
		07-08	179	12	17
60+	05-06	736	71	17	
	07-08	521	46	21	
Non-Hispanic Black Male	12-19	05-06	141	21	13
		07-08	91.5	8.5	6
	20-39	05-06	236	59	6
		07-08	123	12	6
	40-59	05-06	454	70	5
		07-08	333	20	6
60+	05-06	1910	448	5	
	07-08	854	88	8	
Non-Hispanic Black Female	12-19	05-06	81.8	4.8	14
		07-08	63.6	8.4	5
	20-39	05-06	*	*	7
		07-08	105	10	8
	40-59	05-06	691	130	7
		07-08	495	82	8
60+	05-06	2110	398	5	
	07-08	2390	354	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 4.02 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-DDE (lipid adjusted)

Metabolite of Dichlorodiphenyltrichloroethane

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	265	39	6
	20-39	05-06	1730	472	9
		07-08	1090	233	9
	40-59	05-06	*	*	4
		07-08	1010	149	6
	60+	05-06	*	*	4
		07-08	1160	223	5
Mexican American Female	12-19	05-06	464	118	16
		07-08	164	34	5
	20-39	05-06	895	239	9
		07-08	*	*	8
	40-59	05-06	1320	121	6
		07-08	1080	133	6
	60+	05-06	1550	153	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 4.02 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-Dichlorodiphenyldichloroethene (DDE) (whole weight)

Metabolite of Dichlorodiphenyltrichloroethane

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	1.54 (1.33-1.79)	1.31 (1.09-1.66)	3.50 (2.97-4.27)	7.49 (6.14-9.25)	11.6 (9.25-14.8)	1964
	01-02	1.81 (1.64-2.01)	1.57 (1.37-1.72)	3.97 (3.43-4.59)	8.81 (7.85-10.1)	15.4 (12.9-17.6)	2298
	03-04	1.45 (1.18-1.79)	1.26 (1.00-1.58)	3.16 (2.40-4.21)	7.07 (5.55-9.80)	12.1 (8.37-16.0)	1956
Age group							
12-19 years	99-00	.561 (.488-.646)	.520 (.430-.600)	.870 (.680-1.18)	1.52 (1.13-2.25)	2.32 (1.76-3.56)	686
	01-02	.623 (.534-.726)	.590 (.500-.730)	1.00 (.820-1.22)	1.65 (1.39-2.07)	2.30 (1.91-3.14)	758
	03-04	.516 (.419-.635)	.456 (.385-.557)	.796 (.611-1.19)	1.69 (.994-2.69)	2.51 (1.56-6.71)	588
20 years and older	99-00	1.83 (1.56-2.14)	1.61 (1.26-2.07)	4.17 (3.48-4.66)	8.12 (6.37-10.6)	12.3 (9.87-16.7)	1278
	01-02	2.14 (1.91-2.39)	1.77 (1.61-2.05)	4.59 (4.10-5.26)	9.75 (8.34-11.5)	16.8 (13.7-19.1)	1540
	03-04	1.69 (1.36-2.10)	1.46 (1.12-1.96)	3.68 (2.66-4.96)	7.91 (6.01-11.0)	12.8 (9.25-16.8)	1368
Gender							
Males	99-00	1.49 (1.30-1.70)	1.25 (1.10-1.44)	3.02 (2.57-3.80)	6.43 (5.40-8.00)	9.63 (6.63-15.6)	937
	01-02	1.77 (1.57-2.01)	1.59 (1.36-1.76)	3.40 (3.03-4.10)	7.48 (6.43-8.75)	13.1 (9.66-17.6)	1069
	03-04	1.46 (1.18-1.80)	1.30 (1.04-1.58)	2.80 (2.18-4.13)	6.71 (5.51-8.54)	9.93 (7.51-15.4)	955
Females	99-00	1.59 (1.32-1.92)	1.38 (1.03-1.99)	4.05 (3.15-4.79)	8.18 (6.36-11.5)	13.2 (9.81-18.5)	1027
	01-02	1.85 (1.66-2.06)	1.49 (1.32-1.75)	4.57 (3.81-5.47)	10.2 (9.01-11.9)	16.8 (13.4-19.7)	1229
	03-04	1.45 (1.16-1.82)	1.25 (.965-1.66)	3.55 (2.43-4.59)	7.87 (5.41-12.6)	13.7 (8.50-17.3)	1001
Race/ethnicity							
Mexican Americans	99-00	3.92 (3.40-4.51)	3.52 (3.17-3.91)	8.22 (7.26-10.4)	22.0 (12.2-32.2)	32.2 (19.7-48.1)	657
	01-02	3.92 (3.37-4.57)	3.53 (2.68-4.34)	9.34 (7.31-12.5)	26.6 (17.9-38.3)	40.9 (26.8-90.5)	566
	03-04	2.69 (2.18-3.32)	2.24 (1.70-3.24)	5.78 (4.54-7.21)	13.0 (9.53-15.6)	22.9 (15.3-43.4)	457
Non-Hispanic blacks	99-00	1.63 (1.31-2.02)	1.37 (1.11-1.66)	3.84 (3.01-5.69)	11.2 (6.57-13.3)	14.6 (8.88-35.2)	416
	01-02	1.82 (1.46-2.28)	1.38 (1.22-1.72)	4.39 (3.52-6.06)	10.6 (7.24-17.6)	19.4 (8.51-49.3)	515
	03-04	1.47 (1.30-1.65)	1.20 (.963-1.51)	3.76 (2.85-4.75)	9.23 (7.19-14.9)	16.8 (14.7-20.6)	487
Non-Hispanic whites	99-00	1.32 (1.14-1.53)	1.13 (1.01-1.35)	2.88 (2.34-3.36)	5.75 (4.62-6.53)	8.04 (6.32-9.81)	732
	01-02	1.57 (1.39-1.76)	1.41 (1.27-1.58)	3.11 (2.56-3.68)	7.00 (6.02-8.34)	11.3 (8.60-13.7)	1053
	03-04	1.29 (1.01-1.64)	1.12 (.890-1.49)	2.63 (1.84-3.90)	6.36 (3.90-8.71)	9.71 (6.01-15.0)	888

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-DDE (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.592	0.071	9
		07-08	.429	0.031	6
	20-39	05-06	*	*	12
		07-08	.745	0.073	15
	40-59	05-06	1.64	0.13	12
		07-08	1.27	0.09	16
	60+	05-06	4.72	0.32	15
		07-08	3.21	0.29	23
Non-Hispanic White Female	12-19	05-06	.444	0.034	10
		07-08	.334	0.079	7
	20-39	05-06	.657	0.046	16
		07-08	.546	0.037	13
	40-59	05-06	2.02	0.27	13
		07-08	1.18	0.08	17
	60+	05-06	5.38	0.51	17
		07-08	3.50	0.31	21
Non-Hispanic Black Male	12-19	05-06	.705	0.099	13
		07-08	.434	0.044	6
	20-39	05-06	1.43	0.39	6
		07-08	.694	0.076	6
	40-59	05-06	2.86	0.39	5
		07-08	2.14	0.15	6
	60+	05-06	11.4	2.7	5
		07-08	5.21	0.59	8
Non-Hispanic Black Female	12-19	05-06	.406	0.025	14
		07-08	.300	0.037	5
	20-39	05-06	*	*	7
		07-08	.524	0.05	8
	40-59	05-06	4.51	0.94	7
		07-08	2.97	0.49	8
	60+	05-06	14.0	2.5	5
		07-08	14.4	2.2	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum *p,p'*-DDE (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	1.39	0.18	6
	20-39	05-06	12.8	3.7	9
		07-08	7.50	1.68	9
	40-59	05-06	*	*	4
		07-08	7.34	1.34	6
60+	05-06	*	*	4	
	07-08	8.68	2.08	5	
Mexican American Female	12-19	05-06	2.52	0.65	16
		07-08	.803	0.172	5
	20-39	05-06	5.94	1.49	9
		07-08	*	*	8
	40-59	05-06	9.64	0.85	6
		07-08	7.28	0.97	6
	60+	05-06	12.1	2.3	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DDT_BiomonitoringSummary.html

Serum Hexachlorobenzene (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1702
	01-02	*	< LOD	< LOD	< LOD	< LOD	2277
	03-04	15.2 (14.5-15.9)	14.9 (14.2-15.7)	19.0 (18.1-20.3)	24.4 (22.6-26.4)	28.9 (25.6-32.8)	1961
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	591
	01-02	*	< LOD	< LOD	< LOD	< LOD	747
	03-04	13.3 (12.5-14.1)	13.4 (11.5-14.9)	16.7 (15.5-18.4)	20.7 (19.7-22.7)	25.3 (22.7-30.2)	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1111
	01-02	*	< LOD	< LOD	< LOD	< LOD	1530
	03-04	15.5 (14.7-16.2)	15.1 (14.5-15.9)	19.4 (18.3-20.9)	24.8 (22.7-26.6)	29.0 (25.6-33.6)	1373
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	807
	01-02	*	< LOD	< LOD	< LOD	< LOD	1058
	03-04	14.5 (13.8-15.3)	14.2 (13.4-15.0)	18.1 (17.0-19.3)	23.6 (21.0-25.6)	26.9 (25.2-31.3)	957
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	895
	01-02	*	< LOD	< LOD	< LOD	< LOD	1219
	03-04	15.8 (15.0-16.6)	15.7 (15.1-16.4)	20.0 (18.7-21.4)	24.9 (23.0-28.6)	29.8 (26.5-33.7)	1004
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	583
	01-02	*	< LOD	< LOD	< LOD	< LOD	554
	03-04	16.2 (14.9-17.7)	15.3 (14.7-16.4)	20.4 (18.3-22.5)	27.6 (24.7-29.4)	33.7 (27.6-44.6)	460
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	350
	01-02	*	< LOD	< LOD	< LOD	< LOD	511
	03-04	14.5 (13.9-15.0)	14.1 (13.7-15.1)	18.3 (16.6-19.9)	22.3 (20.9-24.0)	26.6 (23.9-30.9)	488
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	636
	01-02	*	< LOD	< LOD	< LOD	< LOD	1052
	03-04	15.1 (14.4-16.0)	15.0 (14.2-15.9)	19.2 (17.9-20.8)	24.3 (22.3-26.4)	28.2 (24.9-32.8)	888

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 118.0, 31.4, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Hexachlorobenzene_BiomonitoringSummary.html

Serum Hexachlorobenzene (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	7.62	1.17	9
		07-08	12.4	0.8	6
	20-39	05-06	7.34	0.41	12
		07-08	9.45	0.46	15
	40-59	05-06	8.85	0.38	12
		07-08	10.1	0.3	16
60+	05-06	11.1	0.5	15	
	07-08	11.8	0.5	23	
Non-Hispanic White Female	12-19	05-06	5.44	0.49	10
		07-08	7.06	0.29	7
	20-39	05-06	7.11	0.33	16
		07-08	7.94	0.23	13
	40-59	05-06	9.25	0.44	13
		07-08	10.8	0.7	17
60+	05-06	13.9	0.7	17	
	07-08	15.1	0.7	21	
Non-Hispanic Black Male	12-19	05-06	6.42	0.37	13
		07-08	9.67	0.79	6
	20-39	05-06	5.77	0.38	6
		07-08	9.11	0.58	6
	40-59	05-06	7.19	0.45	5
		07-08	9.58	0.43	6
60+	05-06	10.9	1.6	5	
	07-08	10.3	0.6	8	
Non-Hispanic Black Female	12-19	05-06	4.41	0.27	14
		07-08	7.20	0.57	5
	20-39	05-06	5.90	0.64	7
		07-08	7.04	0.3	8
	40-59	05-06	9.19	0.42	7
		07-08	9.90	0.58	8
60+	05-06	18.1	1.9	5	
	07-08	15.1	1.2	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 4.17 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorobenzene_BiomonitoringSummary.html

Serum Hexachlorobenzene (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	6.25	0.43	11
		07-08	9.49	0.7	6
	20-39	05-06	9.61	1.74	9
		07-08	9.47	0.57	9
	40-59	05-06	14.0	1.9	4
		07-08	*	*	6
60+	05-06	10.5	1.3	4	
	07-08	12.1	0.8	5	
Mexican American Female	12-19	05-06	5.93	0.55	16
		07-08	6.81	0.44	5
	20-39	05-06	9.78	0.81	9
		07-08	10.4	0.7	8
	40-59	05-06	15.9	1.4	6
		07-08	20.6	2.6	6
	60+	05-06	19.6	2.6	3
		07-08	20.7	1.9	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 4.17 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorobenzene_BiomonitoringSummary.html

Serum Hexachlorobenzene (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1702
	01-02	*	< LOD	< LOD	< LOD	< LOD	2277
	03-04	.092 (.088-.097)	.090 (.086-.095)	.120 (.113-.126)	.157 (.145-.167)	.186 (.169-.212)	1961
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	591
	01-02	*	< LOD	< LOD	< LOD	< LOD	747
	03-04	.065 (.062-.069)	.064 (.060-.069)	.079 (.072-.086)	.102 (.091-.111)	.123 (.111-.141)	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1111
	01-02	*	< LOD	< LOD	< LOD	< LOD	1530
	03-04	.097 (.092-.102)	.095 (.090-.099)	.123 (.118-.130)	.160 (.147-.175)	.191 (.174-.223)	1373
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	807
	01-02	*	< LOD	< LOD	< LOD	< LOD	1058
	03-04	.090 (.085-.095)	.087 (.082-.094)	.115 (.107-.122)	.147 (.135-.163)	.179 (.159-.203)	957
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	895
	01-02	*	< LOD	< LOD	< LOD	< LOD	1219
	03-04	.095 (.089-.100)	.092 (.088-.099)	.123 (.118-.132)	.163 (.148-.176)	.190 (.176-.226)	1004
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	583
	01-02	*	< LOD	< LOD	< LOD	< LOD	554
	03-04	.098 (.089-.109)	.090 (.081-.107)	.125 (.114-.152)	.171 (.157-.203)	.225 (.178-.258)	460
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	350
	01-02	*	< LOD	< LOD	< LOD	< LOD	511
	03-04	.081 (.077-.085)	.078 (.073-.083)	.104 (.095-.118)	.140 (.127-.155)	.167 (.145-.196)	488
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	636
	01-02	*	< LOD	< LOD	< LOD	< LOD	1052
	03-04	.094 (.088-.099)	.092 (.086-.097)	.121 (.114-.129)	.156 (.143-.173)	.182 (.163-.218)	888

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Hexachlorobenzene_BiomonitoringSummary.html

Serum Hexachlorobenzene (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.043	0.008	9
		07-08	.062	0.003	6
	20-39	05-06	.054	0.004	12
		07-08	.062	0.003	15
	40-59	05-06	.067	0.003	12
		07-08	.074	0.003	16
60+	05-06	.076	0.002	15	
	07-08	.075	0.003	23	
Non-Hispanic White Female	12-19	05-06	.029	0.003	10
		07-08	.033	0.001	7
	20-39	05-06	.046	0.002	16
		07-08	.047	0.002	13
	40-59	05-06	.069	0.005	13
		07-08	.071	0.005	17
60+	05-06	.102	0.005	17	
	07-08	.102	0.004	21	
Non-Hispanic Black Male	12-19	05-06	.032	0.001	13
		07-08	.046	0.004	6
	20-39	05-06	.034	0.003	6
		07-08	.051	0.003	6
	40-59	05-06	.046	0.003	5
		07-08	.061	0.003	6
60+	05-06	.064	0.008	5	
	07-08	.062	0.003	8	
Non-Hispanic Black Female	12-19	05-06	.022	0.001	14
		07-08	.034	0.003	5
	20-39	05-06	.034	0.004	7
		07-08	.035	0.001	8
	40-59	05-06	.059	0.003	7
		07-08	.059	0.004	8
60+	05-06	.121	0.012	5	
	07-08	.090	0.006	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorobenzene_BiomonitoringSummary.html

Serum Hexachlorobenzene (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.034	0.002	11
		07-08	.050	0.003	6
	20-39	05-06	.066	0.01	9
		07-08	.065	0.006	9
	40-59	05-06	.113	0.011	4
		07-08	*	*	6
60+	05-06	.076	0.011	4	
	07-08	.089	0.011	5	
Mexican American Female	12-19	05-06	.032	0.003	16
		07-08	.033	0.002	5
	20-39	05-06	.066	0.008	9
		07-08	.060	0.004	8
	40-59	05-06	.117	0.01	6
		07-08	.138	0.016	6
	60+	05-06	.148	0.02	3
		07-08	.141	0.013	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorobenzene_BiomonitoringSummary.html

Serum *beta*-Hexachlorocyclohexane (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	9.68 (<LOD-10.9)	< LOD	19.1 (16.0-21.6)	42.0 (33.4-50.6)	68.9 (50.6-89.5)	1893
	01-02	*	< LOD	16.0 (14.2-17.3)	36.7 (30.4-45.7)	67.2 (50.3-85.7)	2291
	03-04	*	< LOD	14.1 (12.1-16.5)	32.1 (27.1-37.8)	56.5 (43.7-69.4)	1959
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	11.7 (<LOD-16.1)	653
	01-02	*	< LOD	< LOD	< LOD	13.1 (9.70-19.0)	758
	03-04	*	< LOD	< LOD	< LOD	8.80 (<LOD-14.7)	589
20 years and older	99-00	10.9 (9.61-12.4)	< LOD	21.1 (18.9-24.0)	46.0 (35.9-56.8)	73.4 (52.7-96.0)	1240
	01-02	9.87 (9.04-10.8)	7.80 (6.90-8.90)	17.4 (16.0-20.5)	39.6 (33.2-52.0)	71.7 (53.7-96.2)	1533
	03-04	7.89 (<LOD-9.09)	< LOD	16.3 (13.6-18.4)	35.2 (29.6-42.5)	62.2 (48.2-87.6)	1370
Gender							
Males	99-00	*	< LOD	14.6 (10.8-19.1)	29.8 (23.3-38.7)	44.9 (32.8-68.9)	901
	01-02	*	< LOD	12.9 (11.1-15.7)	27.5 (24.0-34.4)	45.7 (35.2-55.6)	1067
	03-04	*	< LOD	10.4 (8.70-12.5)	21.6 (17.7-26.4)	35.3 (26.1-49.9)	952
Females	99-00	11.1 (9.56-12.8)	< LOD	22.0 (19.1-27.6)	51.3 (42.2-67.6)	81.8 (64.4-111)	992
	01-02	10.2 (9.46-11.0)	7.70 (6.90-8.50)	18.5 (16.2-22.7)	47.5 (37.6-62.0)	84.7 (62.0-111)	1224
	03-04	8.43 (<LOD-9.45)	< LOD	18.6 (16.9-21.5)	41.8 (33.9-51.4)	70.3 (62.2-98.4)	1007
Race/ethnicity							
Mexican Americans	99-00	16.7 (13.7-20.2)	15.5 (11.6-20.4)	37.7 (29.6-47.3)	97.9 (62.6-135)	142 (99.8-199)	632
	01-02	13.1 (11.6-14.7)	11.8 (9.60-13.4)	25.8 (21.1-32.7)	69.4 (50.8-87.6)	134 (85.7-166)	563
	03-04	10.5 (8.66-12.8)	10.0 (<LOD-12.7)	23.8 (17.5-29.9)	50.1 (30.0-70.3)	70.3 (42.5-123)	460
Non-Hispanic blacks	99-00	*	< LOD	15.4 (12.0-23.1)	37.9 (30.2-42.2)	49.9 (40.9-81.1)	403
	01-02	*	< LOD	12.0 (8.20-16.2)	36.2 (18.4-73.4)	71.2 (31.9-178)	513
	03-04	*	< LOD	9.70 (8.30-11.9)	27.1 (21.1-36.3)	48.2 (34.8-54.3)	487
Non-Hispanic whites	99-00	*	< LOD	17.5 (14.2-20.5)	34.7 (25.2-46.1)	51.6 (40.0-70.8)	702
	01-02	*	< LOD	14.4 (11.8-16.8)	31.9 (26.6-37.8)	52.0 (37.7-69.5)	1051
	03-04	*	< LOD	12.8 (10.9-14.7)	27.6 (22.1-32.2)	40.8 (32.3-56.5)	887

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 9.36, 6.76, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *beta*-Hexachlorocyclohexane (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	4.44	0.23	12
		07-08	3.34	0.35	16
	60+	05-06	14.3	2.1	15
		07-08	11.3	2.6	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	2.79	0.55	16
		07-08	*	*	13
	40-59	05-06	7.32	0.81	13
		07-08	3.90	1	17
	60+	05-06	19.6	1.3	17
		07-08	17.3	1.7	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	2.00	0.39	6
		07-08	*	*	6
	40-59	05-06	5.64	1.62	5
		07-08	3.62	0.43	6
	60+	05-06	15.0	3.5	5
		07-08	7.79	1.2	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	9.50	1.59	7
		07-08	6.79	1.13	8
	60+	05-06	56.1	12.3	5
		07-08	34.4	5.5	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *beta*-Hexachlorocyclohexane (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	10.7	2.2	9
		07-08	5.59	1.57	9
	40-59	05-06	*	*	4
		07-08	16.6	5	6
60+	05-06	20.5	1.6	4	
	07-08	21.9	5.7	5	
Mexican American Female	12-19	05-06	3.50	0.67	16
		07-08	*	*	5
	20-39	05-06	12.0	2.2	9
		07-08	5.65	1.61	8
	40-59	05-06	25.3	3.5	6
		07-08	16.8	2.1	6
	60+	05-06	36.8	5.5	3
		07-08	50.2	8.6	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *beta*-Hexachlorocyclohexane (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	.058 (<LOD-.066)	< LOD	.120 (.100-.150)	.290 (.220-.360)	.450 (.360-.560)	1893
	01-02	*	< LOD	.100 (.090-.120)	.250 (.210-.310)	.460 (.340-.600)	2291
	03-04	*	< LOD	.092 (.081-.103)	.216 (.173-.254)	.372 (.294-.442)	1959
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	.050 (<LOD-.070)	653
	01-02	*	< LOD	< LOD	< LOD	.080 (.050-.100)	758
	03-04	*	< LOD	< LOD	< LOD	.048 (<LOD-.064)	589
20 years and older	99-00	.067 (.059-.077)	< LOD	.140 (.120-.160)	.330 (.240-.410)	.480 (.410-.620)	1240
	01-02	.062 (.057-.069)	.050 (.050-.060)	.120 (.100-.140)	.270 (.230-.350)	.480 (.370-.690)	1533
	03-04	.050 (<LOD-.058)	< LOD	.103 (.089-.125)	.234 (.191-.290)	.412 (.308-.587)	1370
Gender							
Males	99-00	*	< LOD	.090 (.080-.120)	.210 (.160-.250)	.290 (.220-.470)	901
	01-02	*	< LOD	.080 (.070-.100)	.200 (.150-.260)	.310 (.250-.400)	1067
	03-04	*	< LOD	.072 (.056-.089)	.144 (.118-.174)	.222 (.170-.305)	952
Females	99-00	.065 (.056-.077)	< LOD	.150 (.120-.190)	.380 (.300-.450)	.560 (.420-.680)	992
	01-02	.062 (.057-.067)	.050 (.040-.050)	.130 (.110-.150)	.320 (.260-.400)	.570 (.450-.710)	1224
	03-04	.051 (<LOD-.057)	< LOD	.118 (.103-.130)	.290 (.244-.319)	.442 (.382-.661)	1007
Race/ethnicity							
Mexican Americans	99-00	.098 (.080-.119)	.090 (.070-.110)	.240 (.200-.310)	.580 (.390-.840)	.910 (.580-1.37)	632
	01-02	.078 (.068-.091)	.070 (.050-.080)	.160 (.130-.210)	.470 (.330-.700)	1.01 (.620-1.32)	563
	03-04	.064 (.051-.080)	.057 (<LOD-.086)	.139 (.124-.190)	.331 (.191-.501)	.521 (.297-.814)	460
Non-Hispanic blacks	99-00	*	< LOD	.100 (.070-.140)	.250 (.190-.290)	.360 (.280-.460)	403
	01-02	*	< LOD	.070 (.050-.110)	.220 (.120-.410)	.410 (.190-1.05)	513
	03-04	*	< LOD	.065 (.047-.083)	.167 (.131-.214)	.281 (.200-.404)	487
Non-Hispanic whites	99-00	*	< LOD	.120 (.100-.140)	.250 (.170-.340)	.390 (.280-.510)	702
	01-02	*	< LOD	.090 (.080-.110)	.210 (.180-.260)	.350 (.250-.480)	1051
	03-04	*	< LOD	.083 (.073-.096)	.175 (.146-.220)	.287 (.221-.372)	887

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *beta*-Hexachlorocyclohexane (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	.033	0.002	12
		07-08	.025	0.003	16
	60+	05-06	.095	0.012	15
		07-08	.071	0.015	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	.018	0.004	16
		07-08	*	*	13
	40-59	05-06	.055	0.007	13
		07-08	.026	0.007	17
	60+	05-06	.142	0.008	17
		07-08	.117	0.012	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.012	0.002	6
		07-08	*	*	6
	40-59	05-06	.036	0.011	5
		07-08	.023	0.003	6
	60+	05-06	.087	0.018	5
		07-08	.047	0.007	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	.062	0.011	7
		07-08	.041	0.007	8
	60+	05-06	.369	0.07	5
		07-08	.204	0.029	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *beta*-Hexachlorocyclohexane (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.075	0.015	9
		07-08	.041	0.013	9
	40-59	05-06	*	*	4
		07-08	.120	0.038	6
60+	05-06	.150	0.02	4	
	07-08	.171	0.058	5	
Mexican American Female	12-19	05-06	.019	0.003	16
		07-08	*	*	5
	20-39	05-06	.082	0.016	9
		07-08	.033	0.009	8
	40-59	05-06	.186	0.025	6
		07-08	.114	0.015	6
	60+	05-06	.275	0.037	3
		07-08	.344	0.06	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *gamma*-Hexachlorocyclohexane (Lindane) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1799
	01-02	*	< LOD	< LOD	< LOD	< LOD	2280
	03-04	*	< LOD	< LOD	< LOD	< LOD	1960
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	660
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	< LOD	< LOD	< LOD	593
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1139
	01-02	*	< LOD	< LOD	< LOD	< LOD	1522
	03-04	*	< LOD	< LOD	< LOD	< LOD	1367
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	863
	01-02	*	< LOD	< LOD	< LOD	< LOD	1060
	03-04	*	< LOD	< LOD	< LOD	< LOD	952
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	936
	01-02	*	< LOD	< LOD	< LOD	< LOD	1220
	03-04	*	< LOD	< LOD	< LOD	< LOD	1008
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	631
	01-02	*	< LOD	< LOD	< LOD	< LOD	563
	03-04	*	< LOD	< LOD	< LOD	< LOD	461
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	380
	01-02	*	< LOD	< LOD	< LOD	< LOD	509
	03-04	*	< LOD	< LOD	< LOD	< LOD	490
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	646
	01-02	*	< LOD	< LOD	< LOD	< LOD	1045
	03-04	*	< LOD	< LOD	< LOD	< LOD	884

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 14.5, 10.5, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *gamma*-Hexachlorocyclohexane (Lindane) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
60+	05-06	*	*	15	
	07-08	*	*	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
60+	05-06	*	*	17	
	07-08	*	*	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
60+	05-06	*	*	5	
	07-08	*	*	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
60+	05-06	*	*	5	
	07-08	*	*	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *gamma*-Hexachlorocyclohexane (Lindane) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *gamma*-Hexachlorocyclohexane (Lindane) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1799
	01-02	*	< LOD	< LOD	< LOD	< LOD	2280
	03-04	*	< LOD	< LOD	< LOD	< LOD	1960
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	660
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	< LOD	< LOD	< LOD	593
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1139
	01-02	*	< LOD	< LOD	< LOD	< LOD	1522
	03-04	*	< LOD	< LOD	< LOD	< LOD	1367
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	863
	01-02	*	< LOD	< LOD	< LOD	< LOD	1060
	03-04	*	< LOD	< LOD	< LOD	< LOD	952
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	936
	01-02	*	< LOD	< LOD	< LOD	< LOD	1220
	03-04	*	< LOD	< LOD	< LOD	< LOD	1008
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	631
	01-02	*	< LOD	< LOD	< LOD	< LOD	563
	03-04	*	< LOD	< LOD	< LOD	< LOD	461
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	380
	01-02	*	< LOD	< LOD	< LOD	< LOD	509
	03-04	*	< LOD	< LOD	< LOD	< LOD	490
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	646
	01-02	*	< LOD	< LOD	< LOD	< LOD	1045
	03-04	*	< LOD	< LOD	< LOD	< LOD	884

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *gamma*-Hexachlorocyclohexane (Lindane) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum *gamma*-Hexachlorocyclohexane (Lindane) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/Hexachlorocyclohexane_BiomonitoringSummary.html

Serum Mirex (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1853
	01-02	*	< LOD	< LOD	15.8 (<LOD-73.7)	57.1 (13.2-230)	2257
	03-04	*	< LOD	< LOD	8.40 (<LOD-13.0)	13.2 (7.90-29.6)	1951
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	659
	01-02	*	< LOD	< LOD	< LOD	< LOD	728
	03-04	*	< LOD	< LOD	< LOD	< LOD	592
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1194
	01-02	*	< LOD	< LOD	19.6 (<LOD-108)	71.0 (14.6-305)	1529
	03-04	*	< LOD	< LOD	9.10 (<LOD-15.6)	15.4 (8.10-37.1)	1359
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	887
	01-02	*	< LOD	< LOD	16.1 (<LOD-65.6)	50.8 (12.3-225)	1052
	03-04	*	< LOD	< LOD	9.70 (<LOD-15.4)	15.5 (9.70-24.4)	949
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	966
	01-02	*	< LOD	< LOD	15.0 (<LOD-108)	63.0 (12.0-374)	1205
	03-04	*	< LOD	< LOD	< LOD	11.6 (<LOD-31.3)	1002
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	617
	01-02	*	< LOD	< LOD	< LOD	< LOD	548
	03-04	*	< LOD	< LOD	< LOD	< LOD	459
Non-Hispanic blacks	99-00	*	< LOD	< LOD	15.5 (<LOD-42.2)	39.5 (<LOD-115)	398
	01-02	*	< LOD	13.7 (<LOD-47.3)	51.3 (15.4-230)	153 (30.5-425)	500
	03-04	*	< LOD	< LOD	18.1 (8.70-40.8)	40.3 (15.5-82.7)	484
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	688
	01-02	*	< LOD	< LOD	15.1 (<LOD-104)	66.7 (12.5-291)	1049
	03-04	*	< LOD	< LOD	< LOD	11.6 (<LOD-23.4)	884

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 14.6, 10.5, and 7.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mirex_BiomonitoringSummary.html

Serum Mirex (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	4.25	0.31	16
60+	05-06	5.32	0.61	15	
	07-08	6.36	1.34	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	2.42	0.14	13
		07-08	2.05	0.28	17
60+	05-06	3.51	0.24	17	
	07-08	3.90	0.39	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	2.68	0.59	6
		07-08	*	*	6
	40-59	05-06	5.90	0.49	5
		07-08	*	*	6
60+	05-06	*	*	5	
	07-08	13.9	2.1	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	1.62	0.32	7
		07-08	*	*	8
	40-59	05-06	5.92	0.65	7
		07-08	5.42	1.21	8
60+	05-06	10.3	2.7	5	
	07-08	*	*	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mirex_BiomonitoringSummary.html

Serum Mirex (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	2.66	0.74	4
		07-08	*	*	6
60+	05-06	2.89	0.78	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	1.84	0.34	6
		07-08	*	*	6
	60+	05-06	2.84	0.37	3
		07-08	2.59	0.49	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.46 and 1.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mirex_BiomonitoringSummary.html

Serum Mirex (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1853
	01-02	*	< LOD	< LOD	.100 (<LOD-.470)	.410 (.080-1.73)	2257
	03-04	*	< LOD	< LOD	.054 (<LOD-.084)	.093 (.052-.170)	1951
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	659
	01-02	*	< LOD	< LOD	< LOD	< LOD	728
	03-04	*	< LOD	< LOD	< LOD	< LOD	592
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1194
	01-02	*	< LOD	< LOD	.140 (<LOD-.690)	.470 (.090-1.92)	1529
	03-04	*	< LOD	< LOD	.059 (<LOD-.102)	.106 (.053-.215)	1359
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	887
	01-02	*	< LOD	< LOD	.110 (<LOD-.470)	.370 (.090-1.37)	1052
	03-04	*	< LOD	< LOD	.064 (<LOD-.106)	.108 (.062-.170)	949
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	966
	01-02	*	< LOD	< LOD	.090 (<LOD-.510)	.430 (.070-1.79)	1205
	03-04	*	< LOD	< LOD	< LOD	.077 (<LOD-.170)	1002
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	617
	01-02	*	< LOD	< LOD	< LOD	< LOD	548
	03-04	*	< LOD	< LOD	< LOD	< LOD	459
Non-Hispanic blacks	99-00	*	< LOD	< LOD	.090 (<LOD-.220)	.220 (<LOD-.450)	398
	01-02	*	< LOD	.090 (<LOD-.240)	.310 (.090-1.41)	1.08 (.170-3.02)	500
	03-04	*	< LOD	< LOD	.112 (.055-.268)	.256 (.089-.635)	484
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	688
	01-02	*	< LOD	< LOD	.100 (<LOD-.610)	.450 (.080-1.79)	1049
	03-04	*	< LOD	< LOD	< LOD	.079 (<LOD-.174)	884

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mirex_BiomonitoringSummary.html

Serum Mirex (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	.031	0.003	16
60+	05-06	.036	0.004	15	
	07-08	.040	0.008	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	.018	0.002	13
		07-08	.014	0.002	17
60+	05-06	.026	0.002	17	
	07-08	.026	0.003	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.016	0.004	6
		07-08	*	*	6
	40-59	05-06	.038	0.003	5
		07-08	*	*	6
60+	05-06	*	*	5	
	07-08	.084	0.012	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.009	0.002	7
		07-08	*	*	8
	40-59	05-06	.038	0.004	7
		07-08	.032	0.008	8
60+	05-06	.067	0.016	5	
	07-08	*	*	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mirex_BiomonitoringSummary.html

Serum Mirex (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	.022	0.007	4
		07-08	*	*	6
	60+	05-06	.022	0.008	4
		07-08	*	*	5
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.014	0.003	6
		07-08	*	*	6
	60+	05-06	.022	0.005	3
		07-08	.018	0.003	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/Mirex_BiomonitoringSummary.html

Serum 2,2',4-Tribromodiphenyl ether (BDE 17) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1992
Age group							
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	607
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1385
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	< LOD	964
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1028
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	482
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	< LOD	491
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	892

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 1.0.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4-Tribromodiphenyl ether (BDE 17) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4-Tribromodiphenyl ether (BDE 17) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,4,4'-Tribromodiphenyl ether (BDE 28) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	1.19 (1.03-1.37)	1.10 (1.00-1.30)	2.20 (1.90-2.60)	4.80 (3.30-6.70)	8.00 (5.40-11.3)	1987
Age group							
12-19 years	03-04	1.30 (1.15-1.46)	1.20 (1.10-1.50)	2.40 (2.00-2.60)	4.00 (2.90-5.60)	6.10 (4.00-9.40)	598
20 years and older	03-04	1.17 (1.01-1.37)	1.10 (.900-1.30)	2.20 (1.90-2.70)	5.00 (3.30-7.20)	8.20 (6.00-10.9)	1389
Gender							
Males	03-04	1.21 (1.05-1.39)	1.10 (1.00-1.30)	2.30 (2.00-2.70)	5.10 (3.50-7.30)	8.20 (6.00-11.3)	964
Females	03-04	1.17 (.990-1.38)	1.10 (.900-1.40)	2.10 (1.80-2.60)	4.60 (3.00-6.90)	7.80 (4.70-11.8)	1023
Race/ethnicity							
Mexican Americans	03-04	1.43 (1.29-1.59)	1.40 (1.20-1.50)	2.40 (2.00-2.90)	4.70 (3.70-5.50)	7.30 (5.60-8.30)	488
Non-Hispanic blacks	03-04	1.21 (1.03-1.42)	1.10 (.900-1.30)	2.20 (1.80-2.50)	5.30 (3.50-6.70)	8.40 (5.40-12.3)	470
Non-Hispanic whites	03-04	1.17 (.968-1.42)	1.10 (.900-1.30)	2.20 (1.80-2.70)	4.90 (3.00-8.00)	8.00 (4.50-13.6)	905

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 0.8.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,4,4'-Tribromodphenyl ether (BDE 28) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	1.82	0.44	9
		07-08	1.38	0.19	6
	20-39	05-06	2.16	0.45	12
		07-08	1.37	0.16	15
	40-59	05-06	1.23	0.12	12
		07-08	1.97	0.43	16
	60+	05-06	3.07	0.34	15
		07-08	2.39	0.29	23
Non-Hispanic White Female	12-19	05-06	2.37	0.49	10
		07-08	1.31	0.12	7
	20-39	05-06	1.70	0.32	16
		07-08	1.67	0.23	13
	40-59	05-06	1.44	0.2	13
		07-08	1.37	0.15	17
	60+	05-06	2.59	0.55	17
		07-08	2.51	0.38	21
Non-Hispanic Black Male	12-19	05-06	1.79	0.2	13
		07-08	1.21	0.1	6
	20-39	05-06	1.38	0.19	6
		07-08	1.25	0.14	6
	40-59	05-06	1.47	0.35	5
		07-08	1.33	0.17	6
	60+	05-06	1.91	0.5	5
		07-08	1.94	0.3	8
Non-Hispanic Black Female	12-19	05-06	2.29	0.39	14
		07-08	*	*	5
	20-39	05-06	1.93	0.37	7
		07-08	1.61	0.21	8
	40-59	05-06	*	*	7
		07-08	1.29	0.11	8
	60+	05-06	*	*	5
		07-08	2.33	0.61	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,4,4'-Tribromodphenyl ether (BDE 28) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	2.04	0.21	11
		07-08	2.06	0.38	6
	20-39	05-06	1.99	0.21	9
		07-08	1.52	0.21	9
	40-59	05-06	2.06	0.13	4
		07-08	2.15	0.28	6
60+	05-06	2.38	0.14	4	
	07-08	2.87	0.84	5	
Mexican American Female	12-19	05-06	2.08	0.19	16
		07-08	1.49	0.24	5
	20-39	05-06	1.51	0.14	9
		07-08	1.84	0.42	8
	40-59	05-06	1.57	0.29	6
		07-08	1.81	0.31	6
	60+	05-06	3.29	0.62	3
		07-08	2.34	0.33	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4'-Tetrabromodiphenyl ether (BDE 47) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	20.5 (17.6-23.9)	19.2 (15.7-22.3)	41.1 (35.6-49.2)	85.1 (66.8-127)	163 (108-240)	2016
Age group							
12-19 years	03-04	28.2 (24.6-32.3)	27.2 (22.1-33.6)	53.6 (44.9-63.6)	104 (82.4-145)	174 (115-211)	615
20 years and older	03-04	19.5 (16.5-23.1)	18.0 (14.6-21.6)	39.1 (32.8-47.0)	83.3 (63.0-127)	163 (102-240)	1401
Gender							
Males	03-04	21.4 (18.1-25.3)	19.2 (15.8-24.0)	45.2 (37.3-54.9)	94.3 (66.8-148)	168 (112-382)	981
Females	03-04	19.6 (16.4-23.5)	19.1 (14.1-23.2)	38.4 (31.8-46.4)	79.5 (60.7-121)	155 (102-239)	1035
Race/ethnicity							
Mexican Americans	03-04	25.5 (23.0-28.1)	23.6 (21.2-25.5)	47.1 (38.2-56.5)	87.2 (72.0-105)	151 (105-195)	478
Non-Hispanic blacks	03-04	24.3 (20.9-28.2)	21.4 (18.2-25.6)	47.5 (40.7-53.2)	116 (81.8-149)	242 (136-481)	499
Non-Hispanic whites	03-04	19.5 (16.1-23.7)	17.4 (14.4-22.2)	40.2 (33.1-51.9)	85.1 (60.3-142)	163 (90.2-283)	912

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 4.2.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4'-Tetrabromodiphenyl ether (BDE 47) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	59.0	16.8	9
		07-08	32.9	4.8	6
	20-39	05-06	*	*	12
		07-08	29.1	4.2	15
	40-59	05-06	21.2	2.6	12
		07-08	38.7	9.7	16
	60+	05-06	67.3	12.4	15
		07-08	40.1	6.5	23
Non-Hispanic White Female	12-19	05-06	62.5	14.4	10
		07-08	31.2	3.2	7
	20-39	05-06	*	*	16
		07-08	32.2	6.6	13
	40-59	05-06	22.5	4.6	13
		07-08	20.5	3.1	17
	60+	05-06	*	*	17
		07-08	40.4	7.5	21
Non-Hispanic Black Male	12-19	05-06	51.3	4.9	13
		07-08	31.0	4.9	6
	20-39	05-06	42.8	8.7	6
		07-08	28.1	3.3	6
	40-59	05-06	29.6	6.9	5
		07-08	30.5	5.3	6
	60+	05-06	65.8	18.9	5
		07-08	31.8	4.1	8
Non-Hispanic Black Female	12-19	05-06	74.9	13.9	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	32.2	4.1	8
	40-59	05-06	*	*	7
		07-08	20.9	2.4	8
	60+	05-06	41.4	8.5	5
		07-08	46.4	12.5	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4'-Tetrabromodiphenyl ether (BDE 47) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	46.8	5	11
		07-08	53.1	9.7	6
	20-39	05-06	41.2	4.2	9
		07-08	33.9	6	9
	40-59	05-06	36.4	3.7	4
		07-08	37.4	8.6	6
60+	05-06	36.8	2.9	4	
	07-08	41.5	6.6	5	
Mexican American Female	12-19	05-06	50.7	5.4	16
		07-08	34.2	6	5
	20-39	05-06	30.4	3.6	9
		07-08	31.2	8.1	8
	40-59	05-06	25.4	4	6
		07-08	23.0	3	6
	60+	05-06	52.9	12.1	3
		07-08	37.8	6.9	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,3',4,4'-Tetrabromodiphenyl ether (BDE 66) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	1.30 (1.00-2.10)	1999
Age group							
12-19 years	03-04	*	< LOD	< LOD	< LOD	1.30 (<LOD-1.90)	606
20 years and older	03-04	*	< LOD	< LOD	< LOD	1.30 (1.00-2.20)	1393
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	1.40 (1.00-2.60)	970
Females	03-04	*	< LOD	< LOD	< LOD	1.10 (<LOD-2.20)	1029
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	1.20 (<LOD-1.60)	461
Non-Hispanic blacks	03-04	*	< LOD	< LOD	1.00 (<LOD-2.00)	2.40 (1.40-5.10)	496
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	1.20 (<LOD-2.50)	914

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 1.0.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,3',4,4'-Tetrabromodiphenyl ether (BDE 66) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,3',4,4'-Tetrabromodiphenyl ether (BDE 66) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',3,4,4'-Pentabromodiphenyl ether (BDE 85) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	4.10 (2.80-6.30)	2000
Age group							
12-19 years	03-04	*	< LOD	< LOD	2.80 (<LOD-3.50)	4.00 (3.50-6.30)	610
20 years and older	03-04	*	< LOD	< LOD	< LOD	4.10 (2.50-6.70)	1390
Gender							
Males	03-04	*	< LOD	< LOD	2.40 (<LOD-3.30)	4.80 (3.10-8.40)	967
Females	03-04	*	< LOD	< LOD	< LOD	3.60 (<LOD-5.20)	1033
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	3.70 (2.80-4.80)	484
Non-Hispanic blacks	03-04	*	< LOD	< LOD	3.50 (2.50-4.40)	6.90 (3.10-18.0)	493
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	3.90 (<LOD-7.50)	895

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 2.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',3,4,4'-Pentabromodiphenyl ether (BDE 85) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	.648	0.088	6
	20-39	05-06	*	*	12
		07-08	.618	0.09	15
	40-59	05-06	*	*	12
		07-08	1.06	0.29	16
60+	05-06	1.58	0.32	15	
	07-08	.971	0.197	23	
Non-Hispanic White Female	12-19	05-06	1.50	0.41	10
		07-08	.807	0.166	7
	20-39	05-06	*	*	16
		07-08	.804	0.207	13
	40-59	05-06	*	*	13
		07-08	*	*	17
60+	05-06	*	*	17	
	07-08	.802	0.172	21	
Non-Hispanic Black Male	12-19	05-06	1.14	0.11	13
		07-08	.763	0.149	6
	20-39	05-06	1.08	0.2	6
		07-08	.727	0.104	6
	40-59	05-06	*	*	5
		07-08	.829	0.141	6
60+	05-06	*	*	5	
	07-08	.848	0.199	8	
Non-Hispanic Black Female	12-19	05-06	1.88	0.41	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	.905	0.109	8
	40-59	05-06	*	*	7
		07-08	*	*	8
60+	05-06	1.11	0.32	5	
	07-08	1.16	0.33	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',3,4,4'-Pentabromodiphenyl ether (BDE 85) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.899	0.078	11
		07-08	1.04	0.3	6
	20-39	05-06	.825	0.081	9
		07-08	.790	0.125	9
	40-59	05-06	.900	0.144	4
		07-08	.804	0.214	6
60+	05-06	.683	0.167	4	
	07-08	.967	0.239	5	
Mexican American Female	12-19	05-06	1.00	0.1	16
		07-08	.766	0.147	5
	20-39	05-06	.662	0.098	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	1.10	0.33	3
		07-08	1.00	0.25	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5-Pentabromodiphenyl ether (BDE 99) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	*	< LOD	9.20 (7.50-11.1)	21.7 (17.0-29.1)	42.2 (33.3-54.8)	1985
Age group							
12-19 years	03-04	6.88 (6.14-7.72)	5.70 (<LOD-7.70)	12.9 (11.4-15.7)	27.9 (19.6-37.9)	45.2 (35.9-56.8)	602
20 years and older	03-04	*	< LOD	8.50 (7.10-10.5)	20.6 (15.5-28.8)	41.6 (30.8-57.3)	1383
Gender							
Males	03-04	5.28 (<LOD-6.14)	< LOD	10.0 (8.40-11.6)	24.5 (18.0-37.3)	45.5 (33.8-57.3)	964
Females	03-04	*	< LOD	8.70 (6.60-10.6)	18.3 (14.7-28.8)	41.2 (22.9-60.3)	1021
Race/ethnicity							
Mexican Americans	03-04	5.90 (5.45-6.39)	5.50 (<LOD-5.80)	10.8 (9.30-12.6)	20.0 (17.0-23.5)	30.8 (24.5-41.7)	478
Non-Hispanic blacks	03-04	6.22 (5.42-7.12)	5.00 (<LOD-5.70)	11.5 (9.60-13.1)	30.2 (21.5-42.2)	74.7 (30.2-155)	479
Non-Hispanic whites	03-04	*	< LOD	8.90 (6.80-11.3)	21.7 (15.3-34.0)	43.6 (30.7-71.4)	903

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 5.0.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5-Pentabromodiphenyl ether (BDE 99) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	6.80	1.22	6
	20-39	05-06	*	*	12
		07-08	5.38	0.78	15
	40-59	05-06	4.45	0.74	12
		07-08	*	*	16
	60+	05-06	15.4	4.3	15
		07-08	9.80	2.13	23
Non-Hispanic White Female	12-19	05-06	17.5	4.8	10
		07-08	8.10	2.09	7
	20-39	05-06	*	*	16
		07-08	7.72	2.02	13
	40-59	05-06	4.16	0.79	13
		07-08	4.40	0.82	17
	60+	05-06	*	*	17
		07-08	8.00	1.79	21
Non-Hispanic Black Male	12-19	05-06	12.6	1.4	13
		07-08	7.39	1.79	6
	20-39	05-06	*	*	6
		07-08	6.25	0.82	6
	40-59	05-06	5.66	1.1	5
		07-08	7.22	1.5	6
	60+	05-06	*	*	5
		07-08	8.27	1.74	8
Non-Hispanic Black Female	12-19	05-06	23.3	4.8	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	7.19	1.14	8
	40-59	05-06	*	*	7
		07-08	4.30	0.6	8
	60+	05-06	11.2	3.1	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.2 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5-Pentabromodiphenyl ether (BDE 99) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	9.27	0.91	11
		07-08	10.7	2.5	6
	20-39	05-06	8.47	1	9
		07-08	7.77	1.66	9
	40-59	05-06	9.14	0.97	4
		07-08	6.94	1.71	6
60+	05-06	7.52	1.13	4	
	07-08	8.39	1.47	5	
Mexican American Female	12-19	05-06	10.7	1.3	16
		07-08	7.04	0.95	5
	20-39	05-06	6.51	0.65	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	4.47	0.66	6
	60+	05-06	13.0	3.7	3
		07-08	9.24	1.82	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 1.2 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',6-Pentabromodiphenyl ether (BDE 100) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	3.93 (3.42-4.51)	3.60 (3.10-4.10)	7.80 (6.80-9.00)	18.4 (15.4-22.0)	36.5 (24.6-54.0)	2040
Age group							
12-19 years	03-04	5.17 (4.46-6.00)	4.90 (3.80-6.10)	9.50 (7.90-12.8)	19.3 (14.4-26.2)	34.3 (25.2-45.0)	622
20 years and older	03-04	3.77 (3.24-4.38)	3.30 (2.90-4.00)	7.40 (6.20-9.00)	18.3 (15.0-22.2)	36.6 (23.2-59.2)	1418
Gender							
Males	03-04	4.16 (3.55-4.86)	3.80 (3.10-4.50)	8.50 (7.50-9.90)	18.5 (14.9-25.1)	44.1 (21.9-61.5)	994
Females	03-04	3.72 (3.15-4.40)	3.30 (2.80-4.10)	7.10 (6.00-8.00)	18.4 (14.4-23.1)	33.3 (23.3-46.0)	1046
Race/ethnicity							
Mexican Americans	03-04	4.58 (4.03-5.22)	4.30 (3.80-5.20)	8.10 (6.80-9.10)	14.9 (11.4-20.3)	26.7 (20.3-36.2)	488
Non-Hispanic blacks	03-04	4.72 (4.01-5.55)	4.30 (3.30-5.20)	9.50 (7.80-10.9)	24.1 (16.6-34.3)	41.3 (26.0-79.2)	503
Non-Hispanic whites	03-04	3.78 (3.17-4.51)	3.30 (2.90-4.10)	7.70 (6.20-9.70)	18.6 (14.5-23.3)	40.5 (22.6-59.2)	921

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 1.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',6-Pentabromodiphenyl ether (BDE 100) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	5.68	0.92	6
	20-39	05-06	*	*	12
		07-08	7.65	1.18	15
	40-59	05-06	4.07	0.61	12
		07-08	9.00	2.08	16
	60+	05-06	14.2	2.6	15
		07-08	7.76	1.41	23
Non-Hispanic White Female	12-19	05-06	10.8	1.9	10
		07-08	5.98	0.69	7
	20-39	05-06	9.89	2.83	16
		07-08	6.11	1.05	13
	40-59	05-06	4.06	0.66	13
		07-08	4.12	0.55	17
	60+	05-06	*	*	17
		07-08	8.12	1.66	21
Non-Hispanic Black Male	12-19	05-06	10.8	1	13
		07-08	6.45	0.92	6
	20-39	05-06	9.11	2.32	6
		07-08	7.18	1.05	6
	40-59	05-06	7.19	1.91	5
		07-08	7.41	0.94	6
	60+	05-06	12.1	2.6	5
		07-08	6.56	0.9	8
Non-Hispanic Black Female	12-19	05-06	14.2	2.8	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	7.39	1.17	8
	40-59	05-06	*	*	7
		07-08	4.37	0.48	8
	60+	05-06	8.03	2.35	5
		07-08	8.54	2.26	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',6-Pentabromodiphenyl ether (BDE 100) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	8.93	1.09	11
		07-08	11.0	2.6	6
	20-39	05-06	8.90	1.44	9
		07-08	7.71	1.43	9
	40-59	05-06	7.89	1.25	4
		07-08	8.15	1.99	6
60+	05-06	6.74	0.97	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	9.45	1.14	16
		07-08	6.40	1.09	5
	20-39	05-06	5.36	0.72	9
		07-08	7.45	1.76	8
	40-59	05-06	5.98	1.15	6
		07-08	4.87	0.76	6
	60+	05-06	10.2	2.3	3
		07-08	7.86	1.53	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5,5'-Hexabromodiphenyl ether (BDE 153) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	5.69 (5.11-6.34)	4.80 (4.20-5.30)	11.3 (9.90-12.8)	32.6 (27.9-40.3)	65.7 (54.9-88.4)	2039
Age group							
12-19 years	03-04	8.05 (6.68-9.70)	7.50 (6.10-9.90)	15.3 (11.7-19.4)	31.0 (22.9-44.2)	52.9 (35.9-68.5)	621
20 years and older	03-04	5.41 (4.83-6.05)	4.40 (3.80-5.10)	10.6 (9.00-12.3)	32.8 (26.5-45.9)	73.3 (58.2-90.4)	1418
Gender							
Males	03-04	6.85 (5.99-7.84)	5.50 (4.80-6.70)	12.7 (11.2-16.5)	49.3 (31.0-62.9)	88.4 (63.4-115)	994
Females	03-04	4.78 (4.20-5.43)	4.10 (3.40-5.00)	9.70 (7.40-11.9)	26.0 (20.2-31.6)	54.5 (34.6-62.9)	1045
Race/ethnicity							
Mexican Americans	03-04	5.11 (4.32-6.06)	4.80 (4.20-5.40)	8.70 (6.50-11.0)	17.2 (13.0-25.3)	34.0 (18.3-55.1)	487
Non-Hispanic blacks	03-04	6.05 (5.35-6.83)	5.50 (4.80-6.20)	12.9 (9.90-17.5)	30.0 (21.0-44.2)	53.1 (36.8-63.3)	503
Non-Hispanic whites	03-04	5.85 (5.03-6.81)	4.90 (3.90-5.70)	11.9 (9.90-15.2)	39.0 (28.5-54.9)	75.9 (58.0-93.2)	921

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 2.2.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5,5'-Hexabromodiphenyl ether (BDE 153) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	18.0	4.4	9
		07-08	13.8	2.3	6
	20-39	05-06	15.4	3.7	12
		07-08	20.9	3.1	15
	40-59	05-06	7.53	1.04	12
		07-08	16.0	3.1	16
60+	05-06	19.7	4.7	15	
	07-08	14.7	3.8	23	
Non-Hispanic White Female	12-19	05-06	12.8	2.2	10
		07-08	8.84	0.94	7
	20-39	05-06	11.1	1.9	16
		07-08	11.8	1.7	13
	40-59	05-06	10.4	2.3	13
		07-08	9.89	1.77	17
	60+	05-06	*	*	17
		07-08	13.2	2.5	21
Non-Hispanic Black Male	12-19	05-06	16.5	1.5	13
		07-08	20.7	3.9	6
	20-39	05-06	10.3	1.6	6
		07-08	13.4	2.8	6
	40-59	05-06	13.4	3.6	5
		07-08	10.2	1.1	6
	60+	05-06	14.7	1.6	5
		07-08	10.6	1.8	8
Non-Hispanic Black Female	12-19	05-06	12.1	1.5	14
		07-08	11.1	2	5
	20-39	05-06	9.55	1.69	7
		07-08	11.0	1.7	8
	40-59	05-06	7.41	2.22	7
		07-08	6.28	1.03	8
	60+	05-06	*	*	5
		07-08	7.91	2.1	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5,5'-Hexabromodiphenyl ether (BDE 153) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	11.0	1	11
		07-08	17.2	5	6
	20-39	05-06	11.9	2.6	9
		07-08	10.7	2.1	9
	40-59	05-06	*	*	4
		07-08	11.9	2.4	6
60+	05-06	8.00	1.44	4	
	07-08	17.6	4.5	5	
Mexican American Female	12-19	05-06	10.5	1.4	16
		07-08	7.14	0.72	5
	20-39	05-06	6.73	1.42	9
		07-08	11.1	2.3	8
	40-59	05-06	9.24	2.16	6
		07-08	6.55	1.61	6
	60+	05-06	11.0	1.8	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5,6'-Hexabromodiphenyl ether (BDE 154) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	.900 (.800-1.10)	2.10 (1.70-2.70)	4.20 (2.80-5.40)	2014
Age group							
12-19 years	03-04	*	< LOD	1.20 (1.00-1.40)	2.70 (2.00-3.00)	4.00 (3.00-4.80)	614
20 years and older	03-04	*	< LOD	.900 (.800-1.10)	2.00 (1.60-2.60)	4.20 (2.70-5.70)	1400
Gender							
Males	03-04	*	< LOD	1.00 (.800-1.20)	2.30 (1.80-3.00)	4.30 (3.20-6.50)	976
Females	03-04	*	< LOD	.900 (.800-1.00)	1.80 (1.40-2.80)	4.20 (2.50-5.70)	1038
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	1.00 (.900-1.10)	1.80 (1.40-2.30)	3.90 (2.30-4.50)	477
Non-Hispanic blacks	03-04	*	< LOD	1.20 (.900-1.40)	2.70 (2.30-4.40)	5.30 (3.10-8.70)	498
Non-Hispanic whites	03-04	*	< LOD	.900 (.800-1.10)	2.00 (1.50-2.80)	4.20 (2.50-6.70)	913

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 0.8.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5,6'-Hexabromodiphenyl ether (BDE 154) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]	
Non-Hispanic White Male	12-19	05-06	*	*	9	
		07-08	*	*	6	
	20-39	05-06	.802	0.237	12	
		07-08	.686	0.089	15	
	40-59	05-06	*	*	12	
		07-08	.974	0.279	16	
	60+	05-06	1.36	0.28	15	
		07-08	.833	0.152	23	
Non-Hispanic White Female	12-19	05-06	1.23	0.3	10	
		07-08	*	*	7	
	20-39	05-06	*	*	16	
		07-08	*	*	13	
	40-59	05-06	*	*	13	
		07-08	*	*	17	
	60+	05-06	*	*	17	
		07-08	.753	0.141	21	
	Non-Hispanic Black Male	12-19	05-06	1.20	0.16	13
			07-08	.712	0.108	6
20-39		05-06	*	*	6	
		07-08	.637	0.083	6	
40-59		05-06	.623	0.16	5	
		07-08	.733	0.099	6	
60+		05-06	1.40	0.38	5	
		07-08	.693	0.103	8	
Non-Hispanic Black Female	12-19	05-06	1.73	0.39	14	
		07-08	*	*	5	
	20-39	05-06	*	*	7	
		07-08	.704	0.088	8	
	40-59	05-06	*	*	7	
		07-08	*	*	8	
	60+	05-06	*	*	5	
		07-08	*	*	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5,6'-Hexabromodiphenyl ether (BDE 154) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.902	0.091	11
		07-08	.973	0.224	6
	20-39	05-06	.827	0.103	9
		07-08	.789	0.148	9
	40-59	05-06	.870	0.104	4
		07-08	.732	0.19	6
60+	05-06	.657	0.135	4	
	07-08	.797	0.177	5	
Mexican American Female	12-19	05-06	.965	0.106	16
		07-08	.649	0.13	5
	20-39	05-06	.609	0.061	9
		07-08	.671	0.177	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	1.09	0.25	3
		07-08	.898	0.201	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',3,4,4',5',6-Heptabromodiphenyl ether (BDE 183) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	03-04	*	< LOD	< LOD	< LOD	< LOD	1993
Age group							
12-19 years	03-04	*	< LOD	< LOD	< LOD	< LOD	604
20 years and older	03-04	*	< LOD	< LOD	< LOD	< LOD	1389
Gender							
Males	03-04	*	< LOD	< LOD	< LOD	1.70 (<LOD-2.60)	962
Females	03-04	*	< LOD	< LOD	< LOD	< LOD	1031
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	< LOD	< LOD	< LOD	484
Non-Hispanic blacks	03-04	*	< LOD	< LOD	< LOD	1.80 (<LOD-2.70)	482
Non-Hispanic whites	03-04	*	< LOD	< LOD	< LOD	< LOD	901

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 1.7.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',3,4,4',5',6-Heptabromodiphenyl ether (BDE 183) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	.611	0.077	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	.693	0.188	5
		07-08	.733	0.087	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',3,4,4',5',6-Heptabromodiphenyl ether (BDE 183) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',3,3',4,4',5,5',6,6'-Decabromodiphenyl ether (BDE 209) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]	
Non-Hispanic White Male	12-19	05-06	*	*	9	
		07-08	*	*	6	
	20-39	05-06	*	*	12	
		07-08	*	*	15	
	40-59	05-06	*	*	12	
		07-08	*	*	16	
	60+	05-06	*	*	15	
		07-08	*	*	23	
	Non-Hispanic White Female	12-19	05-06	*	*	10
			07-08	*	*	7
20-39		05-06	*	*	16	
		07-08	*	*	13	
40-59		05-06	*	*	13	
		07-08	*	*	17	
60+		05-06	*	*	17	
		07-08	*	*	21	
Non-Hispanic Black Male		12-19	05-06	*	*	13
			07-08	*	*	6
	20-39	05-06	*	*	6	
		07-08	*	*	6	
	40-59	05-06	*	*	5	
		07-08	*	*	6	
	60+	05-06	*	*	5	
		07-08	*	*	8	
	Non-Hispanic Black Female	12-19	05-06	*	*	14
			07-08	*	*	5
20-39		05-06	*	*	7	
		07-08	*	*	8	
40-59		05-06	*	*	7	
		07-08	*	*	8	
60+		05-06	*	*	5	
		07-08	*	*	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 6.0 and 5.8.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',3,3',4,4',5,5',6,6'-Decabromodiphenyl ether (BDE 209) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]	
Mexican American Male	12-19	05-06	*	*	11	
		07-08	*	*	6	
	20-39	05-06	*	*	9	
		07-08	*	*	9	
	40-59	05-06	*	*	4	
		07-08	*	*	6	
	60+	05-06	*	*	4	
		07-08	*	*	5	
	Mexican American Female	12-19	05-06	*	*	16
			07-08	*	*	5
20-39		05-06	*	*	9	
		07-08	*	*	8	
40-59		05-06	*	*	6	
		07-08	*	*	6	
60+		05-06	*	*	3	
		07-08	*	*	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 6.0 and 5.8.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5,5'-Hexabromobiphenyl (PBB 153) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	2.29 (1.82-2.87)	2.20 (2.00-2.60)	4.40 (3.40-6.30)	12.8 (6.60-25.5)	27.2 (11.7-60.9)	2032
Age group							
12-19 years	03-04	*	< LOD	1.10 (.900-1.60)	2.70 (2.00-3.00)	4.10 (2.90-4.70)	616
20 years and older	03-04	2.72 (2.14-3.47)	2.50 (2.20-2.80)	4.90 (3.60-7.90)	13.6 (7.20-34.6)	34.6 (12.8-66.8)	1416
Gender							
Males	03-04	2.76 (2.21-3.45)	2.70 (2.30-3.20)	5.40 (3.90-8.50)	15.8 (9.30-27.8)	35.4 (13.5-70.3)	987
Females	03-04	1.92 (1.50-2.46)	2.00 (1.60-2.30)	3.60 (2.80-4.90)	9.70 (4.60-27.5)	23.9 (7.40-56.6)	1045
Race/ethnicity							
Mexican Americans	03-04	1.11 (.917-1.33)	1.10 (<LOD-1.30)	2.60 (2.00-3.00)	6.00 (3.50-8.60)	10.0 (6.50-15.2)	484
Non-Hispanic blacks	03-04	2.35 (1.64-3.37)	2.10 (1.70-2.80)	4.80 (3.00-12.1)	13.9 (7.00-38.2)	29.7 (12.1-70.2)	503
Non-Hispanic whites	03-04	2.66 (2.06-3.42)	2.50 (2.20-2.80)	4.90 (3.40-7.80)	13.5 (6.60-36.4)	34.6 (11.6-70.3)	917

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 0.8.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5,5'-Hexabromobiphenyl (PBB 153) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	.824	0.161	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	8.92	1.44	16
	60+	05-06	8.95	2.02	15
		07-08	6.84	0.74	23
Non-Hispanic White Female	12-19	05-06	.826	0.18	10
		07-08	*	*	7
	20-39	05-06	2.63	0.45	16
		07-08	1.81	0.34	13
	40-59	05-06	*	*	13
		07-08	4.63	1.07	17
	60+	05-06	8.83	1.74	17
		07-08	6.74	1.65	21
Non-Hispanic Black Male	12-19	05-06	1.13	0.18	13
		07-08	.689	0.111	6
	20-39	05-06	*	*	6
		07-08	2.10	0.56	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	8.56	1.13	5
		07-08	7.25	1.17	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	2.38	0.55	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	6.31	1.13	8
	60+	05-06	*	*	5
		07-08	8.01	1.65	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 2,2',4,4',5,5'-Hexabromobiphenyl (PBB 153) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.601	0.074	11
		07-08	*	*	6
	20-39	05-06	1.69	0.44	9
		07-08	.962	0.178	9
	40-59	05-06	4.29	1.26	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	2.12	0.4	6
		07-08	1.64	0.35	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.6 and 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/PBDEs_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/PBDEs_FactSheet.html

Serum 1,2,3,4,6,7,8-Heptabromodibenzo-*p*-dioxin (HpBDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from
the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	11.8	1.7	9
	20-39	05-06	22.5	4.6	12
	40-59	05-06	15.3	2.4	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	29.1	8.7	9
	40-59	05-06	*	*	4
	60+	05-06	19.6	4.9	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 11.1.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,4,6,7,8-Heptabromodibenzo-*p*-dioxin (HpBDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools ‡
Non-Hispanic White Male	12-19	05-06	66.6	11.3	9
	20-39	05-06	177	44	12
	40-59	05-06	116	19	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	216	69	9
	40-59	05-06	*	*	4
	60+	05-06	148	48	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,4,7,8-Hexabromodibenzo-*p*-dioxin / 1,2,3,6,7,8-Hexbromodibenzo-*p*-dioxin (HxBDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 19.9.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,4,7,8-Hexabromodibenzo-*p*-dioxin / 1,2,3,6,7,8-Hexbromodibenzo-*p*-dioxin (HxBDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,7,8,9-Hexabromodibenzo-*p*-dioxin (HxBDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 11.1.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,7,8,9-Hexabromodibenzo-*p*-dioxin (HxBDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,7,8-Pentabromodibenzo-*p*-dioxin (PeBDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 8.8.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,7,8-Pentabromodibenzo-*p*-dioxin (PeBDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools ‡
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 2,3,7,8-Tetrabromodibenzo-*p*-dioxin (TBDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 6.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 2,3,7,8-Tetrabromodibenzo-p-dioxin (TBDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and
Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools ‡
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,4,6,7,8-Heptabromodibenzofuran (HpBDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from
the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	19.3	5.1	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	19.6	4.6	14
	20-39	05-06	18.9	2.4	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 15.5.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,4,6,7,8-Heptabromodibenzofuran (HpBDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	98.2	26.5	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	96.0	21.5	14
	20-39	05-06	108	11	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,4,7,8-Hexabromodibenzofuran (HxBDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 11.1.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,4,7,8-Hexabromodibenzofuran (HxBDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,7,8-Pentabromodibenzofuran (PeBDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 8.8.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,7,8-Pentabromodibenzofuran (PeBDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools ‡
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 2,3,4,7,8-Pentabromodibenzofuran (PeBDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 8.8.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 2,3,4,7,8-Pentabromodibenzofuran (PeBDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools ‡
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 2,3,7,8,-Tetrabromodibenzofuran (TBDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 11.1.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 2,3,7,8,-Tetrabromodibenzofuran (TBDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled concentrations[†]
(in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and
Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects .

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Serum 1,2,3,4,6,7,8-Heptachlorodibenzo-*p*-dioxin (HpCDD) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	58.2 (<LOD-63.6)	86.0 (75.5-96.7)	112 (101-131)	1894
	01-02	39.0 (33.7-45.0)	40.2 (34.9-46.9)	68.7 (56.7-82.2)	115 (88.2-138)	147 (126-181)	1220
	03-04	25.3 (23.4-27.3)	24.9 (22.8-26.9)	42.5 (38.8-48.1)	70.4 (62.7-80.1)	91.3 (73.5-117)	1874
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	63.6 (<LOD-75.6)	657
	01-02	†	†	†	†	†	†
	03-04	16.7 (15.1-18.4)	16.4 (15.1-18.3)	23.6 (21.5-25.8)	33.4 (28.6-36.8)	46.7 (34.5-78.1)	586
20 years and older	99-00	*	< LOD	62.0 (57.1-66.7)	92.9 (81.2-101)	120 (102-139)	1237
	01-02	39.0 (33.7-45.0)	40.2 (34.9-46.9)	68.7 (56.7-82.2)	115 (88.2-138)	147 (126-181)	1220
	03-04	26.8 (24.6-29.2)	27.3 (24.6-29.0)	45.6 (41.3-53.2)	73.7 (64.1-88.6)	95.0 (76.1-126)	1288
Gender							
Males	99-00	*	< LOD	< LOD	73.6 (69.0-80.8)	94.7 (83.1-103)	910
	01-02	36.6 (31.7-42.3)	39.0 (33.3-42.6)	62.1 (49.7-75.0)	102 (75.8-132)	138 (103-169)	553
	03-04	24.2 (21.7-27.0)	23.2 (21.1-25.6)	40.6 (35.3-46.9)	64.2 (58.8-73.7)	85.0 (65.8-113)	920
Females	99-00	*	< LOD	62.7 (<LOD-69.1)	102 (86.0-118)	131 (111-164)	984
	01-02	41.2 (34.9-48.7)	43.6 (35.3-52.4)	76.0 (59.5-90.1)	125 (93.4-150)	158 (130-191)	667
	03-04	26.3 (24.4-28.3)	26.8 (24.3-28.3)	44.4 (41.1-50.2)	76.1 (65.3-89.1)	95.7 (80.7-128)	954
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	61.4 (<LOD-69.0)	97.7 (82.8-111)	132 (108-159)	621
	01-02	39.6 (35.7-43.9)	39.7 (33.6-47.4)	64.0 (55.8-74.7)	107 (82.4-128)	149 (111-171)	262
	03-04	25.8 (22.6-29.4)	26.1 (20.9-30.9)	41.9 (36.7-44.7)	61.0 (49.7-71.9)	80.1 (65.0-89.1)	424
Non-Hispanic blacks	99-00	*	< LOD	58.1 (<LOD-71.1)	95.0 (75.1-110)	125 (102-183)	408
	01-02	43.7 (35.4-54.0)	42.8 (32.2-59.8)	80.6 (60.9-106)	134 (101-166)	167 (130-230)	218
	03-04	25.8 (22.6-29.4)	23.7 (20.7-27.1)	41.2 (32.6-56.4)	69.2 (54.6-115)	115 (67.1-164)	454
Non-Hispanic whites	99-00	*	< LOD	59.0 (<LOD-64.8)	84.9 (72.0-97.0)	106 (96.7-122)	709
	01-02	39.3 (33.0-46.8)	40.5 (34.0-50.1)	71.0 (56.3-87.5)	117 (87.1-147)	147 (125-186)	657
	03-04	25.0 (22.6-27.7)	24.6 (22.3-27.4)	42.6 (39.4-48.5)	73.5 (60.4-86.7)	93.7 (71.6-127)	875

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 55.9, 10.3, and 13.0, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8-Heptachlorodibenzo-*p*-dioxin (HpCDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	19.2	1.6	9
	20-39	05-06	19.4	1.4	12
	40-59	05-06	36.1	4.6	12
	60+	05-06	47.8	4.3	15
Non-Hispanic White Female	12-19	05-06	15.0	1	10
	20-39	05-06	20.5	2.1	16
	40-59	05-06	32.2	2.9	13
	60+	05-06	49.7	3	17
Non-Hispanic Black Male	12-19	05-06	17.2	0.7	13
	20-39	05-06	16.1	1	6
	40-59	05-06	23.4	1.6	5
	60+	05-06	44.9	5.7	5
Non-Hispanic Black Female	12-19	05-06	15.9	1.7	14
	20-39	05-06	20.8	1.6	7
	40-59	05-06	36.3	4.3	7
	60+	05-06	77.4	12.5	5
Mexican American Male	12-19	05-06	15.3	1.1	11
	20-39	05-06	26.0	2.6	9
	40-59	05-06	34.5	3.6	4
	60+	05-06	40.3	5	4
Mexican American Female	12-19	05-06	13.2	0.7	16
	20-39	05-06	25.1	1.9	9
	40-59	05-06	41.8	4.3	6
	60+	05-06	59.1	6.9	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.6.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8-Heptachlorodibenzo-*p*-dioxin (HpCDD) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	354 (<LOD384)	564 (501-617)	749 (646-869)	1894
	01-02	252 (219-289)	267 (232-305)	440 (376-529)	779 (591-989)	1030 (840-1290)	1220
	03-04	155 (143-167)	150 (134-163)	288 (242-319)	465 (399-550)	618 (516-787)	1874
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	289 (<LOD340)	657
	01-02	†	†	†	†	†	†
	03-04	84.6 (77.4-92.5)	83.9 (76.1-92.0)	118 (106-131)	165 (139-185)	210 (176-319)	586
20 years and older	99-00	*	< LOD	385 (354-418)	610 (534-677)	802 (681-936)	1237
	01-02	252 (219-289)	267 (232-305)	440 (376-529)	779 (591-989)	1030 (840-1290)	1220
	03-04	169 (155-184)	170 (150-186)	314 (267-348)	506 (419-576)	628 (546-810)	1288
Gender							
Males	99-00	*	< LOD	< LOD	489 (436-543)	613 (535-681)	910
	01-02	243 (211-279)	245 (213-291)	422 (349-527)	766 (536-976)	983 (766-1260)	553
	03-04	150 (135-167)	144 (129-161)	273 (236-305)	424 (365-516)	554 (440-667)	920
Females	99-00	*	< LOD	373 (<LOD428)	637 (562-802)	907 (806-1040)	984
	01-02	260 (221-306)	281 (236-324)	466 (386-551)	795 (621-997)	1140 (849-1330)	667
	03-04	159 (147-172)	158 (138-170)	303 (249-345)	520 (429-591)	623 (545-848)	954
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	372 (<LOD404)	598 (546-657)	849 (657-1020)	621
	01-02	254 (232-278)	249 (229-286)	403 (360-521)	791 (643-929)	988 (817-1240)	262
	03-04	158 (139-180)	159 (138-193)	274 (241-299)	414 (349-469)	556 (451-592)	424
Non-Hispanic blacks	99-00	*	< LOD	323 (<LOD391)	567 (422-716)	778 (574-1130)	408
	01-02	258 (208-319)	262 (180-339)	478 (343-600)	852 (578-1170)	1170 (821-1660)	218
	03-04	148 (128-170)	137 (117-156)	247 (188-344)	434 (345-599)	715 (405-1090)	454
Non-Hispanic whites	99-00	*	< LOD	370 (<LOD401)	565 (491-634)	733 (637-816)	709
	01-02	255 (216-302)	277 (228-328)	442 (374-547)	782 (551-1020)	1020 (803-1290)	657
	03-04	155 (140-170)	152 (131-171)	292 (248-325)	492 (390-571)	621 (506-810)	875

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8-Heptachlorodibenzo-*p*-dioxin (HpCDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	106	11	9
	20-39	05-06	140	12	12
	40-59	05-06	274	39	12
	60+	05-06	325	28	15
Non-Hispanic White Female	12-19	05-06	79.4	5	10
	20-39	05-06	135	17	16
	40-59	05-06	236	23	13
	60+	05-06	365	24	17
Non-Hispanic Black Male	12-19	05-06	86.5	3.3	13
	20-39	05-06	95.6	7.3	6
	40-59	05-06	149	12	5
	60+	05-06	266	30	5
Non-Hispanic Black Female	12-19	05-06	78.3	7.8	14
	20-39	05-06	120	10	7
	40-59	05-06	234	31	7
	60+	05-06	522	90	5
Mexican American Male	12-19	05-06	82.6	5.3	11
	20-39	05-06	184	17	9
	40-59	05-06	284	41	4
	60+	05-06	286	26	4
Mexican American Female	12-19	05-06	72.1	4.2	16
	20-39	05-06	168	15	9
	40-59	05-06	311	37	6
	60+	05-06	458	88	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	10.7 (<LOD-13.9)	14.9 (11.7-20.0)	1239
	03-04	*	< LOD	< LOD	< LOD	< LOD	1861
Age group							
12-19 years	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	582
20 years and older	01-02	*	< LOD	< LOD	10.7 (<LOD-13.9)	14.9 (11.7-20.0)	1239
	03-04	*	< LOD	< LOD	< LOD	< LOD	1279
Gender							
Males	01-02	*	< LOD	< LOD	10.9 (<LOD-14.3)	14.7 (11.5-17.6)	566
	03-04	*	< LOD	< LOD	< LOD	< LOD	914
Females	01-02	*	< LOD	< LOD	10.7 (<LOD-14.1)	15.6 (11.1-23.0)	673
	03-04	*	< LOD	< LOD	< LOD	< LOD	947
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	9.20 (<LOD-11.8)	263
	03-04	*	< LOD	< LOD	< LOD	< LOD	421
Non-Hispanic blacks	01-02	*	< LOD	< LOD	13.9 (<LOD-17.6)	18.3 (13.9-23.0)	220
	03-04	*	< LOD	< LOD	< LOD	< LOD	450
Non-Hispanic whites	01-02	*	< LOD	< LOD	11.3 (<LOD-14.4)	15.1 (12.0-20.5)	672
	03-04	*	< LOD	< LOD	< LOD	< LOD	870

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 9.0 and 11.9.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	1.47	0.35	12
	40-59	05-06	4.47	0.42	12
	60+	05-06	7.15	0.76	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	3.58	0.21	13
	60+	05-06	6.40	0.33	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	1.54	0.32	6
	40-59	05-06	2.77	0.38	5
	60+	05-06	6.65	0.2	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	1.87	0.24	7
	40-59	05-06	3.82	0.43	7
	60+	05-06	12.7	1.7	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	4.98	0.17	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	3.64	0.8	6
	60+	05-06	7.26	0.68	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	74.7 (<LOD-90.7)	105 (81.2-139)	1239
	03-04	*	< LOD	< LOD	< LOD	< LOD	1861
Age group							
12-19 years	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	582
20 years and older	01-02	*	< LOD	< LOD	74.7 (<LOD-90.7)	105 (81.2-139)	1239
	03-04	*	< LOD	< LOD	< LOD	< LOD	1279
Gender							
Males	01-02	*	< LOD	< LOD	73.6 (<LOD-90.7)	105 (77.9-134)	566
	03-04	*	< LOD	< LOD	< LOD	< LOD	914
Females	01-02	*	< LOD	< LOD	74.7 (<LOD-90.7)	102 (78.0-152)	673
	03-04	*	< LOD	< LOD	< LOD	< LOD	947
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	71.2 (<LOD-118)	263
	03-04	*	< LOD	< LOD	< LOD	< LOD	421
Non-Hispanic blacks	01-02	*	< LOD	< LOD	82.1 (<LOD-130)	130 (75.7-184)	220
	03-04	*	< LOD	< LOD	< LOD	< LOD	450
Non-Hispanic whites	01-02	*	< LOD	< LOD	77.5 (<LOD-97.1)	105 (78.8-143)	672
	03-04	*	< LOD	< LOD	< LOD	< LOD	870

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	11.1	2.8	12
	40-59	05-06	33.7	3.5	12
	60+	05-06	48.4	4.7	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	26.6	2.2	13
	60+	05-06	46.7	2.5	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	9.39	1.97	6
	40-59	05-06	17.8	2.4	5
	60+	05-06	40.0	2.1	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	10.8	1.4	7
	40-59	05-06	24.6	3	7
	60+	05-06	84.0	9.5	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	36.0	2.7	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	27.2	5.6	6
	60+	05-06	55.9	8.6	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,6,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	32.6 (28.3-38.2)	56.9 (47.4-67.3)	74.0 (68.3-82.4)	1885
	01-02	34.6 (29.6-40.6)	39.2 (32.7-44.7)	60.8 (50.3-74.2)	95.2 (76.2-120)	128 (99.4-153)	1234
	03-04	17.2 (15.7-18.9)	20.0 (17.8-22.9)	36.5 (32.2-40.0)	53.0 (48.1-59.6)	68.5 (59.6-74.9)	1871
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	26.7 (20.2-29.6)	648
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	16.1 (14.3-18.1)	19.4 (16.4-27.7)	584
20 years and older	99-00	*	< LOD	36.2 (31.5-40.7)	62.8 (53.6-69.1)	75.6 (70.5-84.2)	1237
	01-02	34.6 (29.6-40.6)	39.2 (32.7-44.7)	60.8 (50.3-74.2)	95.2 (76.2-120)	128 (99.4-153)	1234
	03-04	19.7 (17.8-21.8)	23.8 (20.7-26.4)	39.3 (35.4-42.2)	56.6 (49.7-63.8)	70.8 (60.7-82.2)	1287
Gender							
Males	99-00	*	< LOD	31.5 (23.7-38.2)	55.0 (45.7-64.2)	71.3 (59.4-79.4)	908
	01-02	34.1 (28.3-41.1)	38.9 (32.1-44.7)	61.9 (50.0-79.5)	94.9 (70.8-131)	130 (88.5-181)	564
	03-04	17.5 (15.5-19.8)	19.8 (17.8-21.6)	35.5 (29.8-40.3)	52.9 (45.4-63.2)	70.2 (57.5-88.7)	920
Females	99-00	*	< LOD	34.9 (29.1-39.7)	61.2 (51.0-69.2)	74.9 (68.4-92.2)	977
	01-02	35.1 (29.9-41.2)	40.1 (32.4-46.3)	59.8 (49.8-72.3)	97.6 (77.1-114)	126 (108-142)	670
	03-04	16.9 (15.3-18.6)	20.5 (17.8-24.6)	36.9 (33.2-41.0)	53.6 (48.3-59.6)	65.6 (60.0-73.4)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	21.3 (<LOD-27.6)	43.3 (34.1-52.3)	58.0 (49.5-64.8)	624
	01-02	18.3 (15.6-21.4)	21.2 (19.4-25.0)	31.9 (27.5-40.3)	51.5 (40.3-69.9)	68.3 (48.0-111)	260
	03-04	*	< LOD	21.1 (16.3-26.5)	32.2 (24.5-47.4)	43.0 (31.5-65.3)	424
Non-Hispanic blacks	99-00	*	< LOD	31.9 (26.6-41.2)	56.7 (44.9-74.6)	81.6 (72.2-91.7)	402
	01-02	38.9 (33.6-45.0)	40.3 (33.5-47.3)	63.5 (54.6-76.9)	93.9 (78.7-133)	136 (92.6-185)	219
	03-04	16.2 (12.9-20.4)	18.1 (14.4-21.6)	34.9 (28.4-42.9)	54.5 (44.4-69.4)	74.0 (54.3-122)	454
Non-Hispanic whites	99-00	*	< LOD	35.5 (29.7-40.0)	60.9 (51.4-68.3)	74.3 (68.3-83.0)	703
	01-02	37.8 (31.5-45.4)	42.8 (33.9-51.2)	65.0 (52.3-82.9)	99.6 (78.4-130)	131 (103-165)	671
	03-04	18.7 (17.0-20.6)	22.9 (19.9-26.2)	38.0 (35.2-41.5)	56.6 (48.7-63.8)	69.0 (60.6-74.9)	872

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 20.1, 9.1, and 12.3, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,6,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	10.5	0.8	9
	20-39	05-06	14.7	1.4	12
	40-59	05-06	31.8	3.4	12
	60+	05-06	46.6	3.4	15
Non-Hispanic White Female	12-19	05-06	6.77	0.94	10
	20-39	05-06	13.1	1.4	16
	40-59	05-06	25.6	1.2	13
	60+	05-06	47.3	3.5	17
Non-Hispanic Black Male	12-19	05-06	9.16	0.4	13
	20-39	05-06	12.7	0.8	6
	40-59	05-06	20.4	1.3	5
	60+	05-06	45.5	6.1	5
Non-Hispanic Black Female	12-19	05-06	5.51	0.64	14
	20-39	05-06	12.3	0.8	7
	40-59	05-06	25.1	2.6	7
	60+	05-06	69.4	8	5
Mexican American Male	12-19	05-06	4.28	1	11
	20-39	05-06	9.77	0.78	9
	40-59	05-06	18.9	2.8	4
	60+	05-06	34.5	2.4	4
Mexican American Female	12-19	05-06	3.66	0.72	16
	20-39	05-06	8.51	0.78	9
	40-59	05-06	18.9	1.5	6
	60+	05-06	36.4	3.3	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.4.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,6,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	202 (168-239)	374 (314-426)	496 (429-571)	1885
	01-02	224 (192-261)	247 (219-287)	412 (347-498)	663 (549-785)	870 (696-1100)	1234
	03-04	105 (95.7-116)	123 (111-136)	240 (219-263)	357 (325-395)	443 (395-487)	1871
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	126 (93.2-155)	648
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	77.6 (67.4-84.8)	94.5 (77.6-122)	584
20 years and older	99-00	*	< LOD	232 (196-268)	403 (349-458)	521 (458-607)	1237
	01-02	224 (192-261)	247 (219-287)	412 (347-498)	663 (549-785)	870 (696-1100)	1234
	03-04	124 (112-138)	146 (125-172)	259 (235-285)	376 (334-422)	466 (403-528)	1287
Gender							
Males	99-00	*	< LOD	189 (134-239)	353 (278-417)	449 (378-526)	908
	01-02	226 (190-270)	263 (223-301)	425 (347-540)	678 (547-794)	883 (666-1180)	564
	03-04	108 (95.4-123)	123 (108-141)	231 (206-261)	351 (286-424)	450 (374-538)	920
Females	99-00	*	< LOD	213 (174-263)	400 (344-467)	553 (467-651)	977
	01-02	222 (190-260)	239 (207-281)	397 (338-474)	644 (526-771)	864 (720-1060)	670
	03-04	103 (92.8-113)	121 (101-142)	249 (225-277)	363 (336-395)	429 (395-512)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	128 (<LOD-167)	284 (242-334)	380 (328-419)	624
	01-02	117 (99.4-139)	130 (107-157)	233 (198-270)	409 (278-524)	524 (384-731)	260
	03-04	*	< LOD	141 (106-188)	221 (188-337)	327 (210-426)	424
Non-Hispanic blacks	99-00	*	< LOD	191 (150-234)	374 (267-424)	477 (389-568)	402
	01-02	229 (199-264)	229 (190-284)	371 (333-477)	643 (480-799)	821 (596-1260)	219
	03-04	92.9 (73.3-118)	106 (77.0-125)	219 (165-260)	362 (279-460)	461 (372-689)	454
Non-Hispanic whites	99-00	*	< LOD	222 (183-270)	403 (346-460)	520 (450-615)	703
	01-02	246 (206-293)	275 (233-325)	443 (365-553)	684 (549-870)	897 (718-1150)	671
	03-04	116 (105-127)	136 (120-165)	258 (231-286)	375 (333-418)	466 (402-519)	872

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,6,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	57.9	5.2	9
	20-39	05-06	109	14	12
	40-59	05-06	240	27	12
	60+	05-06	316	21	15
Non-Hispanic White Female	12-19	05-06	36.1	4.9	10
	20-39	05-06	86.0	10.4	16
	40-59	05-06	190	13	13
	60+	05-06	344	24	17
Non-Hispanic Black Male	12-19	05-06	46.0	1.5	13
	20-39	05-06	75.2	5.9	6
	40-59	05-06	130	8	5
	60+	05-06	270	32	5
Non-Hispanic Black Female	12-19	05-06	27.3	3.1	14
	20-39	05-06	70.2	4.3	7
	40-59	05-06	161	17	7
	60+	05-06	461	45	5
Mexican American Male	12-19	05-06	23.0	5.2	11
	20-39	05-06	68.7	4.7	9
	40-59	05-06	156	28	4
	60+	05-06	248	17	4
Mexican American Female	12-19	05-06	20.5	4	16
	20-39	05-06	56.7	5.7	9
	40-59	05-06	140	13	6
	60+	05-06	279	43	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8,9-Hexachlorodibenzo-*p*-dioxin (HxCDD) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1870
	01-02	*	< LOD	< LOD	12.5 (10.5-15.3)	17.0 (14.3-20.0)	1238
	03-04	*	< LOD	< LOD	< LOD	< LOD	1869
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	642
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	585
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1228
	01-02	*	< LOD	< LOD	12.5 (10.5-15.3)	17.0 (14.3-20.0)	1238
	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	895
	01-02	*	< LOD	< LOD	12.1 (<LOD-14.8)	15.1 (12.9-18.5)	567
	03-04	*	< LOD	< LOD	< LOD	< LOD	918
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	975
	01-02	*	< LOD	< LOD	13.0 (10.7-16.8)	18.3 (15.7-21.1)	671
	03-04	*	< LOD	< LOD	< LOD	< LOD	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	618
	01-02	*	< LOD	< LOD	9.60 (<LOD-11.6)	12.2 (<LOD-20.6)	262
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	396
	01-02	*	< LOD	< LOD	14.6 (11.2-20.0)	19.9 (14.6-23.9)	220
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	701
	01-02	*	< LOD	< LOD	12.9 (9.90-15.9)	17.3 (14.7-20.6)	672
	03-04	*	< LOD	< LOD	< LOD	< LOD	871

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 20.3, 9.3, and 12.3, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8,9-Hexachlorodibenzo-*p*-dioxin (HxCDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	1.72	0.42	9
	20-39	05-06	1.94	0.25	12
	40-59	05-06	3.99	0.38	12
	60+	05-06	5.48	0.47	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	2.46	0.2	16
	40-59	05-06	3.94	0.25	13
	60+	05-06	6.02	0.36	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	2.07	0.11	6
	40-59	05-06	2.57	0.18	5
	60+	05-06	5.15	0.91	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	2.42	0.23	7
	40-59	05-06	3.83	0.36	7
	60+	05-06	9.10	1.09	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	2.20	0.28	9
	40-59	05-06	3.39	0.54	4
	60+	05-06	4.49	0.22	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	1.84	0.32	9
	40-59	05-06	4.02	0.33	6
	60+	05-06	6.51	0.94	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.4.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8,9-Hexachlorodibenzo-*p*-dioxin (HxCDD) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1870
	01-02	*	< LOD	< LOD	86.5 (68.8-108)	121 (99.5-146)	1238
	03-04	*	< LOD	< LOD	< LOD	< LOD	1869
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	642
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	585
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1228
	01-02	*	< LOD	< LOD	86.5 (68.8-108)	121 (99.5-146)	1238
	03-04	*	< LOD	< LOD	< LOD	< LOD	1284
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	895
	01-02	*	< LOD	< LOD	84.1 (<LOD-104)	108 (90.6-142)	567
	03-04	*	< LOD	< LOD	< LOD	< LOD	918
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	975
	01-02	*	< LOD	< LOD	89.8 (67.9-121)	123 (102-157)	671
	03-04	*	< LOD	< LOD	< LOD	< LOD	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	618
	01-02	*	< LOD	< LOD	74.7 (<LOD-104)	107 (<LOD-167)	262
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	396
	01-02	*	< LOD	< LOD	92.6 (62.8-126)	123 (82.6-169)	220
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	701
	01-02	*	< LOD	< LOD	88.1 (67.9-119)	124 (96.6-152)	672
	03-04	*	< LOD	< LOD	< LOD	< LOD	871

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8,9-Hexachlorodibenzo-*p*-dioxin (HxCDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	9.70	2.31	9
	20-39	05-06	13.9	1.9	12
	40-59	05-06	30.2	3.2	12
	60+	05-06	37.3	3	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	16.1	1.5	16
	40-59	05-06	29.1	2.3	13
	60+	05-06	44.0	2.6	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	12.3	0.8	6
	40-59	05-06	16.4	1.2	5
	60+	05-06	30.7	5	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	14.0	1.3	7
	40-59	05-06	24.6	2.4	7
	60+	05-06	60.7	6.7	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	15.8	1.9	9
	40-59	05-06	27.8	5.5	4
	60+	05-06	32.5	1.2	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	12.5	2.1	9
	40-59	05-06	29.8	2.8	6
	60+	05-06	49.7	9.5	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8,9-Octachlorodibenzo-*p*-dioxin (OCDD) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	406 (359-453)	674 (597-767)	913 (787-1010)	1921
	01-02	346 (<LOD-394)	333 (<LOD-402)	573 (498-668)	944 (780-1090)	1260 (998-1610)	1171
	03-04	*	< LOD	336 (283-389)	582 (490-658)	767 (645-913)	1851
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	421 (363-517)	667
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	244 (<LOD-330)	352 (264-458)	581
20 years and older	99-00	*	< LOD	445 (389-496)	710 (624-802)	948 (822-1080)	1254
	01-02	346 (<LOD-394)	333 (<LOD-402)	573 (498-668)	944 (780-1090)	1260 (998-1610)	1171
	03-04	220 (<LOD-244)	223 (<LOD-243)	358 (297-421)	597 (502-719)	794 (665-978)	1270
Gender							
Males	99-00	*	< LOD	< LOD	517 (447-580)	704 (563-838)	919
	01-02	*	< LOD	442 (346-579)	767 (593-968)	1030 (837-1240)	517
	03-04	*	< LOD	270 (244-320)	457 (377-559)	668 (501-856)	910
Females	99-00	*	< LOD	504 (422-579)	802 (674-928)	1010 (928-1180)	1002
	01-02	410 (356-472)	405 (335-502)	647 (574-751)	1020 (858-1360)	1450 (1060-1780)	654
	03-04	235 (<LOD-256)	238 (225-248)	402 (321-486)	640 (551-749)	829 (675-1020)	941
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	418 (365-502)	703 (610-873)	940 (737-1230)	632
	01-02	*	< LOD	432 (394-545)	755 (578-1220)	1150 (696-1640)	250
	03-04	*	< LOD	296 (225-356)	452 (363-540)	588 (417-861)	419
Non-Hispanic blacks	99-00	*	< LOD	444 (371-519)	741 (566-983)	1120 (799-1560)	411
	01-02	421 (352-503)	420 (339-509)	682 (537-907)	1110 (956-1520)	1640 (1130-1900)	210
	03-04	*	< LOD	345 (276-455)	642 (513-883)	926 (636-1310)	448
Non-Hispanic whites	99-00	*	< LOD	391 (333-452)	625 (562-754)	861 (676-1010)	721
	01-02	349 (<LOD-409)	335 (<LOD-421)	574 (496-679)	945 (764-1170)	1290 (972-1660)	632
	03-04	*	< LOD	343 (282-403)	585 (464-674)	758 (635-922)	865

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 329.0, 319.0, and 218.0, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8,9-Octachlorodibenzo-*p*-dioxin (OCDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	125	13	9
	20-39	05-06	131	8	12
	40-59	05-06	240	25	12
	60+	05-06	382	38	15
Non-Hispanic White Female	12-19	05-06	109	9	10
	20-39	05-06	182	19	16
	40-59	05-06	306	22	13
	60+	05-06	548	66	17
Non-Hispanic Black Male	12-19	05-06	145	12	13
	20-39	05-06	130	9	6
	40-59	05-06	203	32	5
	60+	05-06	423	82	5
Non-Hispanic Black Female	12-19	05-06	156	22	14
	20-39	05-06	191	18	7
	40-59	05-06	384	43	7
	60+	05-06	929	134	5
Mexican American Male	12-19	05-06	114	9	11
	20-39	05-06	182	21	9
	40-59	05-06	*	*	4
	60+	05-06	353	46	4
Mexican American Female	12-19	05-06	114	5	16
	20-39	05-06	187	12	9
	40-59	05-06	412	60	6
	60+	05-06	542	64	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 7.7.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8,9-Octachlorodibenzo-*p*-dioxin (OCDD) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	2530 (2230-2880)	4260 (3770-4760)	5950 (5090-6790)	1921
	01-02	2230 (<LOD-2540)	2170 (<LOD-2550)	3860 (3180-4520)	6460 (5140-8290)	9110 (6940-11400)	1171
	03-04	*	< LOD	2180 (1830-2540)	3760 (3100-4540)	5020 (4190-6070)	1851
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	1910 (1600-2340)	667
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	1160 (<LOD1590)	1780 (1160-2850)	581
20 years and older	99-00	*	< LOD	2810 (2480-3110)	4570 (4100-5020)	6200 (5340-7300)	1254
	01-02	2230 (<LOD-2540)	2170 (<LOD-2550)	3860 (3180-4520)	6460 (5140-8290)	9110 (6940-11400)	1171
	03-04	1380 (<LOD-1540)	1360 (<LOD-1470)	2350 (1970-2900)	4030 (3370-4700)	5340 (4490-6370)	1270
Gender							
Males	99-00	*	< LOD	< LOD	3160 (2760-3760)	4160 (3320-5570)	919
	01-02	*	< LOD	3160 (2410-4070)	5270 (4070-7140)	7620 (6020-9760)	517
	03-04	*	< LOD	1780 (1540-2070)	3090 (2410-4020)	4190 (3410-5350)	910
Females	99-00	*	< LOD	3110 (2680-3560)	5090 (4610-5740)	6760 (5870-8710)	1002
	01-02	2590 (2250-2980)	2620 (2160-3000)	4340 (3880-4860)	6990 (5870-9040)	10000 (7070-12300)	654
	03-04	1420 (<LOD-1560)	1410 (1300-1510)	2700 (2210-3030)	4340 (3750-4830)	5770 (4590-6740)	941
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	2470 (2240-2990)	4350 (3660-5570)	6560 (5340-7590)	632
	01-02	*	< LOD	3410 (2760-3860)	5810 (4250-7670)	8050 (5760-11800)	250
	03-04	*	< LOD	1880 (1560-2220)	3240 (2450-3880)	3910 (3240-4630)	419
Non-Hispanic blacks	99-00	*	< LOD	2520 (2090-2950)	4770 (4040-5740)	7140 (4740-10700)	411
	01-02	2480 (2050-3000)	2460 (2060-2970)	4170 (3160-5560)	7250 (5470-9920)	9920 (7990-12000)	210
	03-04	*	< LOD	2150 (1780-2880)	3860 (3080-5640)	5720 (3750-7830)	448
Non-Hispanic whites	99-00	*	< LOD	2560 (2160-2910)	4120 (3580-4780)	5800 (4740-6790)	721
	01-02	2270 (<LOD-2660)	2200 (<LOD-2770)	3860 (3090-4720)	6530 (4890-8860)	9150 (6630-12300)	632
	03-04	*	< LOD	2220 (1830-2700)	3770 (3040-4680)	5050 (4190-6350)	865

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8,9-Octachlorodibenzo-*p*-dioxin (OCDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	690	73	9
	20-39	05-06	951	79	12
	40-59	05-06	1820	219	12
	60+	05-06	2600	245	15
Non-Hispanic White Female	12-19	05-06	577	44	10
	20-39	05-06	1200	153	16
	40-59	05-06	2260	191	13
	60+	05-06	3990	469	17
Non-Hispanic Black Male	12-19	05-06	727	59	13
	20-39	05-06	776	71	6
	40-59	05-06	1280	183	5
	60+	05-06	2480	424	5
Non-Hispanic Black Female	12-19	05-06	768	106	14
	20-39	05-06	1100	108	7
	40-59	05-06	2470	301	7
	60+	05-06	6240	925	5
Mexican American Male	12-19	05-06	621	47	11
	20-39	05-06	1300	156	9
	40-59	05-06	*	*	4
	60+	05-06	2520	251	4
Mexican American Female	12-19	05-06	621	23	16
	20-39	05-06	1250	104	9
	40-59	05-06	3070	469	6
	60+	05-06	4210	842	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8-Pentachlorodibenzo-*p*-dioxin (PeCDD) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1915
	01-02	*	< LOD	< LOD	11.3 (9.30-13.6)	15.8 (13.3-19.8)	1236
	03-04	*	< LOD	6.10 (5.50-6.80)	9.00 (8.30-9.70)	11.0 (9.90-12.2)	1878
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	659
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	4.80 (<LOD-5.90)	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1256
	01-02	*	< LOD	< LOD	11.3 (9.30-13.6)	15.8 (13.3-19.8)	1236
	03-04	*	< LOD	6.60 (5.90-7.20)	9.30 (8.60-10.1)	11.3 (10.1-12.7)	1290
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	920
	01-02	*	< LOD	< LOD	10.8 (9.10-13.3)	14.5 (11.7-19.4)	564
	03-04	*	< LOD	5.90 (5.30-6.40)	8.90 (7.90-9.60)	11.0 (9.60-12.7)	923
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	995
	01-02	*	< LOD	6.10 (<LOD-7.80)	11.8 (9.40-14.3)	16.6 (13.7-20.8)	672
	03-04	*	< LOD	6.50 (5.70-7.20)	9.10 (8.30-10.1)	11.0 (10.0-12.2)	955
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	632
	01-02	*	< LOD	< LOD	< LOD	8.70 (<LOD-12.7)	262
	03-04	*	< LOD	< LOD	6.50 (5.20-7.90)	7.80 (6.70-9.20)	425
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	408
	01-02	*	< LOD	7.70 (<LOD-9.30)	13.9 (9.60-18.4)	18.4 (14.2-24.0)	218
	03-04	*	< LOD	6.40 (5.30-8.20)	9.90 (8.50-13.4)	14.4 (9.60-20.1)	455
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	717
	01-02	*	< LOD	< LOD	11.7 (9.50-14.3)	16.7 (13.6-20.2)	672
	03-04	*	< LOD	6.50 (5.80-7.10)	9.30 (8.60-10.0)	11.1 (10.1-12.2)	877

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 14.2, 6.0, and 4.5, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8-Pentachlorodibenzo-*p*-dioxin (PeCDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	2.52	0.14	9
	20-39	05-06	2.47	0.24	12
	40-59	05-06	4.49	0.55	12
	60+	05-06	7.25	0.4	15
Non-Hispanic White Female	12-19	05-06	1.53	0.25	10
	20-39	05-06	2.55	0.23	16
	40-59	05-06	4.25	0.2	13
	60+	05-06	7.83	0.57	17
Non-Hispanic Black Male	12-19	05-06	2.35	0.14	13
	20-39	05-06	2.95	0.16	6
	40-59	05-06	3.99	0.23	5
	60+	05-06	8.17	1.25	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	2.53	0.1	7
	40-59	05-06	4.63	0.44	7
	60+	05-06	12.5	1.7	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	2.11	0.28	9
	40-59	05-06	3.59	0.62	4
	60+	05-06	5.58	0.14	4
Mexican American Female	12-19	05-06	1.07	0.19	16
	20-39	05-06	1.63	0.25	9
	40-59	05-06	3.21	0.64	6
	60+	05-06	6.49	0.81	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8-Pentachlorodibenzo-*p*-dioxin (PeCDD) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1915
	01-02	*	< LOD	< LOD	77.8 (62.8-96.3)	117 (90.3-133)	1236
	03-04	*	< LOD	40.7 (37.4-43.0)	60.3 (53.2-67.4)	76.1 (66.9-87.4)	1878
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	659
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	23.8 (<LOD-27.2)	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1256
	01-02	*	< LOD	< LOD	77.8 (62.8-96.3)	117 (90.3-133)	1236
	03-04	*	< LOD	43.0 (39.2-47.8)	63.5 (55.6-71.5)	80.7 (67.9-89.9)	1290
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	920
	01-02	*	< LOD	< LOD	76.1 (54.7-93.7)	107 (83.5-133)	564
	03-04	*	< LOD	39.3 (35.1-43.4)	57.3 (49.3-66.1)	67.8 (58.7-90.4)	923
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	995
	01-02	*	< LOD	37.8 (<LOD-50.2)	80.4 (62.9-107)	121 (85.7-167)	672
	03-04	*	< LOD	41.8 (37.7-45.2)	63.6 (55.9-72.9)	80.7 (70.5-88.3)	955
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	632
	01-02	*	< LOD	< LOD	< LOD	66.0 (<LOD-111)	262
	03-04	*	< LOD	< LOD	44.5 (34.2-52.0)	52.8 (44.5-76.6)	425
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	408
	01-02	*	< LOD	43.8 (<LOD-53.8)	81.9 (58.2-117)	123 (81.9-169)	218
	03-04	*	< LOD	39.7 (30.6-48.0)	67.3 (48.0-93.0)	93.0 (66.7-137)	455
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	717
	01-02	*	< LOD	< LOD	80.8 (61.6-107)	121 (88.6-142)	672
	03-04	*	< LOD	42.6 (38.6-45.1)	63.4 (55.4-67.9)	78.8 (67.9-84.7)	877

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8-Pentachlorodibenzo-*p*-dioxin (PeCDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	13.8	0.6	9
	20-39	05-06	17.9	2.1	12
	40-59	05-06	34.1	4.4	12
	60+	05-06	49.2	2.3	15
Non-Hispanic White Female	12-19	05-06	8.06	1.3	10
	20-39	05-06	16.7	1.7	16
	40-59	05-06	31.4	2.2	13
	60+	05-06	56.9	3.7	17
Non-Hispanic Black Male	12-19	05-06	11.7	0.6	13
	20-39	05-06	17.5	1	6
	40-59	05-06	25.5	1.5	5
	60+	05-06	48.9	7.8	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	14.5	0.7	7
	40-59	05-06	29.8	3.1	7
	60+	05-06	83.0	10	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	14.9	2	9
	40-59	05-06	29.7	6.3	4
	60+	05-06	40.5	2.9	4
Mexican American Female	12-19	05-06	5.94	1.02	16
	20-39	05-06	11.0	1.8	9
	40-59	05-06	24.0	4.7	6
	60+	05-06	49.3	8.2	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1898
	01-02	*	< LOD	< LOD	< LOD	< LOD	1228
	03-04	*	< LOD	< LOD	4.10 (<LOD-4.40)	5.20 (4.30-5.80)	1876
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	658
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1240
	01-02	*	< LOD	< LOD	< LOD	< LOD	1228
	03-04	*	< LOD	< LOD	4.30 (3.90-4.60)	5.30 (4.50-6.10)	1288
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	912
	01-02	*	< LOD	< LOD	< LOD	< LOD	559
	03-04	*	< LOD	< LOD	< LOD	4.60 (3.80-5.30)	921
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	986
	01-02	*	< LOD	< LOD	< LOD	6.40 (<LOD-9.20)	669
	03-04	*	< LOD	< LOD	4.40 (4.00-4.90)	5.50 (4.50-6.60)	955
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	630
	01-02	*	< LOD	< LOD	< LOD	< LOD	262
	03-04	*	< LOD	< LOD	< LOD	3.80 (<LOD-5.50)	424
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	404
	01-02	*	< LOD	< LOD	< LOD	7.50 (<LOD-10.0)	217
	03-04	*	< LOD	< LOD	4.50 (<LOD-6.10)	6.20 (4.40-10.3)	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	709
	01-02	*	< LOD	< LOD	< LOD	< LOD	665
	03-04	*	< LOD	< LOD	4.10 (<LOD-4.50)	5.20 (4.30-5.90)	877

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.1, 5.8, and 3.8, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	1.21	0.16	12
	60+	05-06	2.37	0.19	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	1.28	0.19	13
	60+	05-06	2.86	0.22	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	1.25	0.17	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	.691	0.131	7
	40-59	05-06	1.82	0.2	7
	60+	05-06	6.39	1.64	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	1.02	0.24	4
	60+	05-06	1.28	0.26	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	1.08	0.32	6
	60+	05-06	2.37	0.31	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1898
	01-02	*	< LOD	< LOD	< LOD	< LOD	1228
	03-04	*	< LOD	< LOD	27.5 (<LOD-30.1)	34.1 (30.2-39.6)	1876
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	658
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1240
	01-02	*	< LOD	< LOD	< LOD	< LOD	1228
	03-04	*	< LOD	< LOD	28.6 (26.4-31.2)	36.4 (31.5-42.7)	1288
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	912
	01-02	*	< LOD	< LOD	< LOD	< LOD	559
	03-04	*	< LOD	< LOD	< LOD	28.8 (25.4-34.5)	921
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	986
	01-02	*	< LOD	< LOD	< LOD	50.7 (<LOD-74.3)	669
	03-04	*	< LOD	< LOD	30.2 (27.4-33.1)	38.7 (32.3-44.5)	955
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	630
	01-02	*	< LOD	< LOD	< LOD	< LOD	262
	03-04	*	< LOD	< LOD	< LOD	23.6 (<LOD-32.5)	424
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	404
	01-02	*	< LOD	< LOD	< LOD	55.6 (<LOD-72.0)	217
	03-04	*	< LOD	< LOD	31.1 (<LOD-42.0)	42.4 (29.5-67.8)	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	709
	01-02	*	< LOD	< LOD	< LOD	< LOD	665
	03-04	*	< LOD	< LOD	28.0 (<LOD-30.0)	33.6 (31.1-38.8)	877

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	9.15	1.33	12
	60+	05-06	15.8	1.1	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	9.94	1.67	13
	60+	05-06	20.7	1.4	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	7.96	1.01	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	3.99	0.77	7
	40-59	05-06	11.7	1.5	7
	60+	05-06	42.6	10.8	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	8.37	2.32	4
	60+	05-06	9.13	1.9	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	7.71	2.17	6
	60+	05-06	17.9	3.2	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	14.7 (<LOD-18.0)	19.5 (17.4-23.0)	1709
	01-02	9.64 (8.53-10.9)	10.3 (8.80-11.7)	14.6 (12.7-16.7)	21.3 (18.0-25.5)	27.1 (22.5-32.0)	1219
	03-04	*	< LOD	10.6 (9.90-11.4)	14.6 (13.1-16.5)	18.7 (16.5-24.2)	1858
Age group							
12-19 years	99-00	*	< LOD	< LOD	18.4 (16.2-20.9)	24.7 (20.9-28.1)	600
	01-02	†	†	†	†	†	†
	03-04	9.36 (8.60-10.2)	8.80 (<LOD-9.80)	13.4 (12.3-14.6)	19.3 (16.2-29.6)	33.2 (21.2-54.2)	583
20 years and older	99-00	*	< LOD	< LOD	14.2 (<LOD-17.5)	18.4 (15.0-23.8)	1109
	01-02	9.64 (8.53-10.9)	10.3 (8.80-11.7)	14.6 (12.7-16.7)	21.3 (18.0-25.5)	27.1 (22.5-32.0)	1219
	03-04	*	< LOD	10.3 (9.60-11.1)	13.8 (12.5-16.0)	18.0 (16.0-20.9)	1275
Gender							
Males	99-00	*	< LOD	< LOD	16.4 (<LOD-20.0)	21.0 (18.1-26.8)	815
	01-02	10.1 (8.74-11.6)	11.0 (9.30-12.6)	15.2 (12.9-17.2)	20.8 (17.2-27.8)	28.9 (22.0-34.8)	557
	03-04	*	< LOD	11.3 (10.1-12.9)	17.1 (14.3-19.1)	23.9 (18.3-30.9)	913
Females	99-00	*	< LOD	< LOD	< LOD	17.5 (14.3-20.3)	894
	01-02	9.28 (8.20-10.5)	9.40 (8.20-11.2)	14.1 (12.4-16.6)	21.3 (17.4-25.5)	26.5 (22.3-31.9)	662
	03-04	*	< LOD	9.90 (9.30-10.5)	13.0 (12.2-14.3)	16.0 (14.3-18.4)	945
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	19.7 (<LOD-25.4)	570
	01-02	7.73 (7.19-8.31)	8.20 (7.80-8.70)	11.2 (10.6-12.5)	16.5 (14.5-19.6)	20.8 (17.2-26.8)	260
	03-04	*	< LOD	9.10 (<LOD-10.6)	12.0 (10.1-16.8)	15.1 (11.6-20.0)	420
Non-Hispanic blacks	99-00	*	< LOD	< LOD	22.4 (15.2-28.2)	28.2 (26.5-29.6)	359
	01-02	12.3 (10.6-14.4)	12.8 (10.9-15.2)	17.4 (15.4-21.0)	25.5 (22.4-31.2)	32.1 (25.7-37.6)	214
	03-04	*	< LOD	11.3 (10.7-12.2)	16.1 (13.4-19.0)	20.4 (16.7-26.8)	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	17.5 (14.9-18.9)	636
	01-02	9.50 (8.30-10.9)	10.1 (8.40-12.0)	14.4 (12.6-16.6)	20.5 (16.9-25.5)	25.8 (20.5-31.7)	665
	03-04	*	< LOD	10.5 (9.60-11.5)	14.6 (12.9-17.4)	19.0 (16.3-26.2)	864

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 13.5, 7.0, and 8.6, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	12.4	1.6	9
	20-39	05-06	7.09	0.32	12
	40-59	05-06	8.32	0.87	12
	60+	05-06	8.15	0.85	15
Non-Hispanic White Female	12-19	05-06	7.62	0.44	10
	20-39	05-06	6.94	0.58	16
	40-59	05-06	6.84	0.46	13
	60+	05-06	7.73	0.6	17
Non-Hispanic Black Male	12-19	05-06	10.8	1.2	13
	20-39	05-06	8.25	1.33	6
	40-59	05-06	7.87	1.02	5
	60+	05-06	10.6	2.5	5
Non-Hispanic Black Female	12-19	05-06	8.82	1.47	14
	20-39	05-06	6.20	0.57	7
	40-59	05-06	6.34	0.74	7
	60+	05-06	10.4	1.2	5
Mexican American Male	12-19	05-06	9.24	1.4	11
	20-39	05-06	6.55	0.48	9
	40-59	05-06	7.44	1.53	4
	60+	05-06	7.17	1	4
Mexican American Female	12-19	05-06	5.61	0.33	16
	20-39	05-06	5.99	0.41	9
	40-59	05-06	8.73	1.1	6
	60+	05-06	6.32	1.43	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	85.3 (<LOD-95.8)	108 (95.8-120)	1709
	01-02	62.2 (55.4-69.9)	64.5 (58.0-71.6)	94.1 (82.3-110)	134 (119-165)	181 (147-206)	1219
	03-04	*	< LOD	63.6 (59.9-67.7)	88.3 (81.0-100)	116 (101-137)	1858
Age group							
12-19 years	99-00	*	< LOD	< LOD	79.3 (69.5-84.2)	102 (83.4-120)	600
	01-02	†	†	†	†	†	†
	03-04	47.5 (44.0-51.3)	43.7 (<LOD-49.3)	63.7 (59.1-72.3)	102 (85.3-114)	175 (113-222)	583
20 years and older	99-00	*	< LOD	< LOD	85.7 (<LOD-97.7)	108 (93.6-127)	1109
	01-02	62.2 (55.4-69.9)	64.5 (58.0-71.6)	94.1 (82.3-110)	134 (119-165)	181 (147-206)	1219
	03-04	*	< LOD	63.5 (59.6-67.6)	86.5 (78.8-100)	112 (94.9-135)	1275
Gender							
Males	99-00	*	< LOD	< LOD	91.4 (<LOD-108)	112 (99.3-133)	815
	01-02	66.3 (58.2-75.6)	70.4 (59.8-81.8)	99.3 (85.5-120)	143 (120-181)	182 (146-237)	557
	03-04	*	< LOD	69.7 (62.9-76.0)	100 (85.9-116)	135 (113-193)	913
Females	99-00	*	< LOD	< LOD	< LOD	97.8 (86.2-130)	894
	01-02	58.8 (52.0-66.4)	60.2 (53.5-67.6)	89.5 (74.7-107)	131 (108-165)	169 (131-225)	662
	03-04	*	< LOD	58.5 (54.5-62.4)	83.5 (74.7-87.7)	102 (87.7-116)	945
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	113 (<LOD-133)	570
	01-02	49.5 (46.0-53.4)	50.4 (45.7-57.2)	76.9 (67.1-82.1)	99.7 (90.9-124)	133 (94.8-202)	260
	03-04	*	< LOD	56.0 (<LOD-64.6)	74.2 (62.6-91.4)	90.5 (76.4-107)	420
Non-Hispanic blacks	99-00	*	< LOD	< LOD	117 (89.0-143)	153 (109-197)	359
	01-02	72.6 (61.9-85.2)	70.9 (62.4-85.2)	107 (91.5-128)	147 (131-177)	193 (154-259)	214
	03-04	*	< LOD	66.2 (60.2-75.5)	91.3 (80.3-112)	119 (102-151)	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	97.7 (89.3-104)	636
	01-02	61.8 (54.3-70.4)	65.4 (57.4-72.2)	94.3 (78.1-112)	133 (114-166)	180 (137-222)	665
	03-04	*	< LOD	63.9 (59.6-69.2)	87.8 (79.8-100)	116 (102-139)	864

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	68.5	9.8	9
	20-39	05-06	51.2	3.2	12
	40-59	05-06	63.0	7.1	12
	60+	05-06	55.4	5.5	15
Non-Hispanic White Female	12-19	05-06	40.7	2.6	10
	20-39	05-06	45.0	3.8	16
	40-59	05-06	50.3	3.7	13
	60+	05-06	56.3	4.2	17
Non-Hispanic Black Male	12-19	05-06	54.8	6.7	13
	20-39	05-06	49.7	9.1	6
	40-59	05-06	50.1	6.7	5
	60+	05-06	62.2	13.5	5
Non-Hispanic Black Female	12-19	05-06	43.4	7.3	14
	20-39	05-06	35.6	3.3	7
	40-59	05-06	40.3	4.1	7
	60+	05-06	69.9	7.9	5
Mexican American Male	12-19	05-06	49.7	7.2	11
	20-39	05-06	46.5	3.4	9
	40-59	05-06	59.9	11.2	4
	60+	05-06	50.8	4.8	4
Mexican American Female	12-19	05-06	30.5	1.6	16
	20-39	05-06	39.8	2.8	9
	40-59	05-06	64.2	7.8	6
	60+	05-06	49.6	15.4	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	1224
	03-04	*	< LOD	< LOD	< LOD	< LOD	1852
Age group							
12-19 years	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	583
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1224
	03-04	*	< LOD	< LOD	< LOD	< LOD	1269
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	558
	03-04	*	< LOD	< LOD	< LOD	< LOD	908
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	666
	03-04	*	< LOD	< LOD	< LOD	< LOD	944
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	262
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	217
	03-04	*	< LOD	< LOD	< LOD	< LOD	452
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	661
	03-04	*	< LOD	< LOD	< LOD	< LOD	856

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 7.0 and 8.6.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	1224
	03-04	*	< LOD	< LOD	< LOD	< LOD	1852
Age group							
12-19 years	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	583
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1224
	03-04	*	< LOD	< LOD	< LOD	< LOD	1269
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	558
	03-04	*	< LOD	< LOD	< LOD	< LOD	908
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	666
	03-04	*	< LOD	< LOD	< LOD	< LOD	944
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	262
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	217
	03-04	*	< LOD	< LOD	< LOD	< LOD	452
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	661
	03-04	*	< LOD	< LOD	< LOD	< LOD	856

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1890
	01-02	*	< LOD	8.00 (6.90-9.60)	12.1 (9.40-14.9)	15.4 (12.9-18.6)	1223
	03-04	*	< LOD	< LOD	7.50 (<LOD-7.90)	8.90 (7.90-10.2)	1866
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	657
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	583
20 years and older	99-00	*	< LOD	< LOD	< LOD	12.8 (<LOD-14.5)	1233
	01-02	*	< LOD	8.00 (6.90-9.60)	12.1 (9.40-14.9)	15.4 (12.9-18.6)	1223
	03-04	*	< LOD	< LOD	7.60 (<LOD-8.20)	9.50 (8.00-10.5)	1283
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	908
	01-02	*	< LOD	8.20 (7.10-10.0)	12.8 (9.50-15.9)	15.9 (12.2-20.9)	562
	03-04	*	< LOD	< LOD	7.40 (<LOD-8.00)	8.50 (7.80-10.0)	916
Females	99-00	*	< LOD	< LOD	< LOD	12.9 (<LOD-16.4)	982
	01-02	*	< LOD	7.90 (6.70-9.00)	11.7 (9.40-13.8)	14.5 (12.2-18.6)	661
	03-04	*	< LOD	< LOD	7.50 (<LOD-8.40)	9.30 (7.80-10.4)	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	631
	01-02	*	< LOD	< LOD	7.30 (<LOD-8.00)	8.00 (7.00-10.9)	261
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	14.3 (<LOD-15.1)	399
	01-02	*	< LOD	8.90 (7.70-10.1)	13.6 (11.1-17.1)	18.6 (14.6-22.6)	214
	03-04	*	< LOD	< LOD	7.80 (<LOD-10.9)	10.9 (7.50-19.4)	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	703
	01-02	*	< LOD	8.30 (7.10-10.2)	12.5 (10.0-15.9)	15.9 (13.1-19.0)	664
	03-04	*	< LOD	< LOD	7.60 (<LOD-8.10)	9.40 (8.00-10.2)	869

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.7, 6.5, and 7.4, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	2.18	0.17	9
	20-39	05-06	2.51	0.29	12
	40-59	05-06	4.54	0.3	12
	60+	05-06	5.39	0.45	15
Non-Hispanic White Female	12-19	05-06	1.72	0.15	10
	20-39	05-06	2.37	0.1	16
	40-59	05-06	3.58	0.18	13
	60+	05-06	5.54	0.34	17
Non-Hispanic Black Male	12-19	05-06	1.78	0.18	13
	20-39	05-06	2.63	0.23	6
	40-59	05-06	3.32	0.33	5
	60+	05-06	5.34	0.88	5
Non-Hispanic Black Female	12-19	05-06	1.22	0.15	14
	20-39	05-06	2.13	0.13	7
	40-59	05-06	3.35	0.37	7
	60+	05-06	8.98	1.22	5
Mexican American Male	12-19	05-06	1.68	0.11	11
	20-39	05-06	2.30	0.22	9
	40-59	05-06	3.29	0.59	4
	60+	05-06	4.07	0.28	4
Mexican American Female	12-19	05-06	1.30	0.05	16
	20-39	05-06	1.84	0.17	9
	40-59	05-06	3.17	0.22	6
	60+	05-06	4.47	0.39	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1890
	01-02	*	< LOD	54.0 (46.5-63.6)	82.3 (65.5-104)	108 (88.0-138)	1223
	03-04	*	< LOD	< LOD	49.1 (<LOD-54.5)	60.1 (53.4-68.8)	1866
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	657
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	583
20 years and older	99-00	*	< LOD	< LOD	< LOD	84.2 (<LOD-97.2)	1233
	01-02	*	< LOD	54.0 (46.5-63.6)	82.3 (65.5-104)	108 (88.0-138)	1223
	03-04	*	< LOD	< LOD	51.0 (<LOD-57.2)	62.2 (56.2-72.7)	1283
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	908
	01-02	*	< LOD	56.9 (48.6-71.4)	90.1 (71.1-108)	108 (87.4-144)	562
	03-04	*	< LOD	< LOD	49.3 (<LOD-54.9)	59.3 (52.9-68.8)	916
Females	99-00	*	< LOD	< LOD	< LOD	82.8 (<LOD-102)	982
	01-02	*	< LOD	51.6 (43.7-59.6)	76.1 (62.6-97.0)	105 (81.6-140)	661
	03-04	*	< LOD	< LOD	48.9 (<LOD-57.2)	61.9 (53.3-72.7)	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	631
	01-02	*	< LOD	< LOD	48.4 (<LOD-56.3)	65.3 (48.6-79.4)	261
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	90.8 (<LOD-105)	399
	01-02	*	< LOD	52.7 (45.0-62.7)	87.1 (63.7-108)	122 (85.2-147)	214
	03-04	*	< LOD	< LOD	56.9 (<LOD-71.7)	71.7 (53.8-112)	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	703
	01-02	*	< LOD	56.0 (48.6-66.1)	84.2 (64.5-115)	112 (88.6-142)	664
	03-04	*	< LOD	< LOD	50.6 (<LOD-56.3)	61.4 (54.0-71.8)	869

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	12.1	1.1	9
	20-39	05-06	18.2	2.4	12
	40-59	05-06	34.3	2.6	12
	60+	05-06	36.8	2.8	15
Non-Hispanic White Female	12-19	05-06	9.06	0.73	10
	20-39	05-06	15.3	0.8	16
	40-59	05-06	26.5	1.9	13
	60+	05-06	40.4	2.4	17
Non-Hispanic Black Male	12-19	05-06	9.17	0.9	13
	20-39	05-06	15.5	1.5	6
	40-59	05-06	21.2	2.1	5
	60+	05-06	31.5	4.4	5
Non-Hispanic Black Female	12-19	05-06	6.17	0.74	14
	20-39	05-06	12.2	0.7	7
	40-59	05-06	21.5	2.4	7
	60+	05-06	60.0	7.8	5
Mexican American Male	12-19	05-06	9.02	0.49	11
	20-39	05-06	16.0	1.1	9
	40-59	05-06	27.2	5.9	4
	60+	05-06	29.3	0.9	4
Mexican American Female	12-19	05-06	7.08	0.26	16
	20-39	05-06	12.2	1.2	9
	40-59	05-06	23.3	2.1	6
	60+	05-06	34.2	5.3	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1898
	01-02	*	< LOD	7.10 (<LOD-8.20)	10.4 (9.00-13.1)	14.0 (11.0-17.1)	1236
	03-04	*	< LOD	< LOD	< LOD	8.80 (<LOD-9.80)	1868
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	656
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	585
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1242
	01-02	*	< LOD	7.10 (<LOD-8.20)	10.4 (9.00-13.1)	14.0 (11.0-17.1)	1236
	03-04	*	< LOD	< LOD	< LOD	9.00 (8.00-10.1)	1283
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	913
	01-02	*	< LOD	7.10 (<LOD-8.60)	11.3 (8.50-13.8)	15.1 (11.3-18.7)	566
	03-04	*	< LOD	< LOD	< LOD	9.10 (7.90-10.8)	918
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	985
	01-02	*	< LOD	7.00 (6.20-8.00)	10.0 (9.20-11.4)	13.1 (10.5-15.6)	670
	03-04	*	< LOD	< LOD	< LOD	8.50 (<LOD-9.20)	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	625
	01-02	*	< LOD	< LOD	< LOD	6.90 (<LOD-11.1)	262
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	408
	01-02	*	< LOD	7.60 (6.10-9.10)	12.1 (10.0-14.1)	16.0 (12.3-21.0)	219
	03-04	*	< LOD	< LOD	< LOD	8.50 (<LOD-12.1)	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	708
	01-02	*	< LOD	7.40 (6.20-9.00)	10.7 (9.20-14.3)	14.8 (11.6-17.2)	670
	03-04	*	< LOD	< LOD	< LOD	9.10 (8.00-10.6)	871

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.6, 6.1, and 7.9, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	2.45	0.17	9
	20-39	05-06	2.68	0.31	12
	40-59	05-06	4.91	0.42	12
	60+	05-06	5.43	0.36	15
Non-Hispanic White Female	12-19	05-06	1.91	0.12	10
	20-39	05-06	2.43	0.11	16
	40-59	05-06	3.69	0.2	13
	60+	05-06	5.51	0.3	17
Non-Hispanic Black Male	12-19	05-06	1.80	0.19	13
	20-39	05-06	2.31	0.19	6
	40-59	05-06	2.98	0.32	5
	60+	05-06	5.29	0.86	5
Non-Hispanic Black Female	12-19	05-06	1.27	0.18	14
	20-39	05-06	2.04	0.1	7
	40-59	05-06	2.97	0.34	7
	60+	05-06	7.44	0.96	5
Mexican American Male	12-19	05-06	1.69	0.25	11
	20-39	05-06	2.24	0.27	9
	40-59	05-06	3.18	0.43	4
	60+	05-06	4.34	0.37	4
Mexican American Female	12-19	05-06	1.41	0.07	16
	20-39	05-06	1.85	0.19	9
	40-59	05-06	2.85	0.13	6
	60+	05-06	4.11	0.34	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1898
	01-02	*	< LOD	46.2 (<LOD-56.5)	70.3 (58.4-90.7)	101 (77.6-120)	1236
	03-04	*	< LOD	< LOD	< LOD	59.8 (<LOD-65.4)	1868
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	656
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	585
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1242
	01-02	*	< LOD	46.2 (<LOD-56.5)	70.3 (58.4-90.7)	101 (77.6-120)	1236
	03-04	*	< LOD	< LOD	< LOD	61.0 (57.6-65.7)	1283
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	913
	01-02	*	< LOD	47.9 (<LOD-60.5)	72.3 (57.5-105)	104 (73.5-133)	566
	03-04	*	< LOD	< LOD	< LOD	61.0 (54.0-67.7)	918
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	985
	01-02	*	< LOD	45.0 (37.0-54.3)	66.2 (58.3-79.1)	90.7 (71.9-115)	670
	03-04	*	< LOD	< LOD	< LOD	58.1 (<LOD-65.0)	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	625
	01-02	*	< LOD	< LOD	< LOD	52.7 (<LOD-65.9)	262
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	408
	01-02	*	< LOD	44.4 (37.2-57.3)	71.5 (59.8-90.9)	107 (70.7-142)	219
	03-04	*	< LOD	< LOD	< LOD	58.4 (<LOD-73.0)	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	708
	01-02	*	< LOD	49.6 (41.9-57.9)	72.8 (58.4-102)	103 (82.6-121)	670
	03-04	*	< LOD	< LOD	< LOD	63.4 (57.3-67.7)	871

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	13.5	1.1	9
	20-39	05-06	19.4	2.6	12
	40-59	05-06	37.2	3.6	12
	60+	05-06	37.0	2.2	15
Non-Hispanic White Female	12-19	05-06	10.1	0.6	10
	20-39	05-06	15.8	0.8	16
	40-59	05-06	27.2	1.9	13
	60+	05-06	40.2	2.3	17
Non-Hispanic Black Male	12-19	05-06	9.35	0.92	13
	20-39	05-06	13.6	1.3	6
	40-59	05-06	19.1	2.1	5
	60+	05-06	31.3	5	5
Non-Hispanic Black Female	12-19	05-06	6.24	0.85	14
	20-39	05-06	11.6	0.4	7
	40-59	05-06	19.1	2.3	7
	60+	05-06	49.5	5.8	5
Mexican American Male	12-19	05-06	8.98	1.31	11
	20-39	05-06	15.7	1.5	9
	40-59	05-06	26.1	4.6	4
	60+	05-06	30.8	1.2	4
Mexican American Female	12-19	05-06	7.68	0.31	16
	20-39	05-06	12.5	1.4	9
	40-59	05-06	21.3	1.5	6
	60+	05-06	31.5	5	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1875
	01-02	*	< LOD	< LOD	< LOD	< LOD	1223
	03-04	*	< LOD	< LOD	< LOD	< LOD	1864
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	645
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	583
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1230
	01-02	*	< LOD	< LOD	< LOD	< LOD	1223
	03-04	*	< LOD	< LOD	< LOD	< LOD	1281
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	894
	01-02	*	< LOD	< LOD	< LOD	< LOD	559
	03-04	*	< LOD	< LOD	< LOD	< LOD	914
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	981
	01-02	*	< LOD	< LOD	< LOD	< LOD	664
	03-04	*	< LOD	< LOD	< LOD	< LOD	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	620
	01-02	*	< LOD	< LOD	< LOD	< LOD	261
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	400
	01-02	*	< LOD	< LOD	< LOD	< LOD	216
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	699
	01-02	*	< LOD	< LOD	< LOD	< LOD	665
	03-04	*	< LOD	< LOD	< LOD	< LOD	866

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.7, 6.0, and 8.3, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1875
	01-02	*	< LOD	< LOD	< LOD	< LOD	1223
	03-04	*	< LOD	< LOD	< LOD	< LOD	1864
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	645
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	583
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1230
	01-02	*	< LOD	< LOD	< LOD	< LOD	1223
	03-04	*	< LOD	< LOD	< LOD	< LOD	1281
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	894
	01-02	*	< LOD	< LOD	< LOD	< LOD	559
	03-04	*	< LOD	< LOD	< LOD	< LOD	914
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	981
	01-02	*	< LOD	< LOD	< LOD	< LOD	664
	03-04	*	< LOD	< LOD	< LOD	< LOD	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	620
	01-02	*	< LOD	< LOD	< LOD	< LOD	261
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	400
	01-02	*	< LOD	< LOD	< LOD	< LOD	216
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	699
	01-02	*	< LOD	< LOD	< LOD	< LOD	665
	03-04	*	< LOD	< LOD	< LOD	< LOD	866

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1884
	01-02	*	< LOD	< LOD	< LOD	< LOD	1230
	03-04	*	< LOD	< LOD	< LOD	< LOD	1866
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	652
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	584
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1232
	01-02	*	< LOD	< LOD	< LOD	< LOD	1230
	03-04	*	< LOD	< LOD	< LOD	< LOD	1282
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	900
	01-02	*	< LOD	< LOD	< LOD	< LOD	565
	03-04	*	< LOD	< LOD	< LOD	< LOD	916
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	984
	01-02	*	< LOD	< LOD	< LOD	< LOD	665
	03-04	*	< LOD	< LOD	< LOD	< LOD	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	614
	01-02	*	< LOD	< LOD	< LOD	< LOD	260
	03-04	*	< LOD	< LOD	< LOD	< LOD	422
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	408
	01-02	*	< LOD	< LOD	< LOD	< LOD	218
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	704
	01-02	*	< LOD	< LOD	< LOD	< LOD	671
	03-04	*	< LOD	< LOD	< LOD	< LOD	869

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.9, 5.8, and 8.2, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.573	0.133	9
	20-39	05-06	.876	0.13	12
	40-59	05-06	1.40	0.16	12
	60+	05-06	1.32	0.07	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	.706	0.08	16
	40-59	05-06	1.02	0.13	13
	60+	05-06	1.18	0.07	17
Non-Hispanic Black Male	12-19	05-06	.422	0.075	13
	20-39	05-06	*	*	6
	40-59	05-06	.799	0.117	5
	60+	05-06	.974	0.089	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	.543	0.101	7
	40-59	05-06	.779	0.099	7
	60+	05-06	1.08	0.18	5
Mexican American Male	12-19	05-06	.490	0.114	11
	20-39	05-06	.838	0.154	9
	40-59	05-06	1.16	0.16	4
	60+	05-06	1.18	0.16	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	.777	0.094	9
	40-59	05-06	1.10	0.08	6
	60+	05-06	.917	0.192	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1884
	01-02	*	< LOD	< LOD	< LOD	< LOD	1230
	03-04	*	< LOD	< LOD	< LOD	< LOD	1866
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	652
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	584
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1232
	01-02	*	< LOD	< LOD	< LOD	< LOD	1230
	03-04	*	< LOD	< LOD	< LOD	< LOD	1282
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	900
	01-02	*	< LOD	< LOD	< LOD	< LOD	565
	03-04	*	< LOD	< LOD	< LOD	< LOD	916
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	984
	01-02	*	< LOD	< LOD	< LOD	< LOD	665
	03-04	*	< LOD	< LOD	< LOD	< LOD	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	614
	01-02	*	< LOD	< LOD	< LOD	< LOD	260
	03-04	*	< LOD	< LOD	< LOD	< LOD	422
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	408
	01-02	*	< LOD	< LOD	< LOD	< LOD	218
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	704
	01-02	*	< LOD	< LOD	< LOD	< LOD	671
	03-04	*	< LOD	< LOD	< LOD	< LOD	869

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	3.14	0.74	9
	20-39	05-06	6.50	1.03	12
	40-59	05-06	10.6	1.4	12
	60+	05-06	8.97	0.53	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	4.66	0.54	16
	40-59	05-06	7.36	0.96	13
	60+	05-06	8.74	0.56	17
Non-Hispanic Black Male	12-19	05-06	2.29	0.39	13
	20-39	05-06	*	*	6
	40-59	05-06	5.10	0.8	5
	60+	05-06	5.78	0.53	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	3.14	0.57	7
	40-59	05-06	4.94	0.63	7
	60+	05-06	7.49	1.31	5
Mexican American Male	12-19	05-06	2.73	0.6	11
	20-39	05-06	5.77	0.92	9
	40-59	05-06	9.51	1.79	4
	60+	05-06	8.20	0.85	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	5.24	0.62	9
	40-59	05-06	8.07	0.72	6
	60+	05-06	7.33	2	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1884
	01-02	*	< LOD	< LOD	< LOD	< LOD	1202
	03-04	*	< LOD	< LOD	< LOD	< LOD	1849
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	652
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	581
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1232
	01-02	*	< LOD	< LOD	< LOD	< LOD	1202
	03-04	*	< LOD	< LOD	< LOD	< LOD	1268
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	904
	01-02	*	< LOD	< LOD	< LOD	< LOD	541
	03-04	*	< LOD	< LOD	< LOD	< LOD	905
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	980
	01-02	*	< LOD	< LOD	< LOD	< LOD	661
	03-04	*	< LOD	< LOD	< LOD	< LOD	944
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	623
	01-02	*	< LOD	< LOD	< LOD	< LOD	257
	03-04	*	< LOD	< LOD	< LOD	< LOD	420
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	404
	01-02	*	< LOD	< LOD	< LOD	< LOD	212
	03-04	*	< LOD	< LOD	< LOD	< LOD	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	705
	01-02	*	< LOD	< LOD	< LOD	< LOD	653
	03-04	*	< LOD	< LOD	< LOD	< LOD	857

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 35.6, 21.0, and 12.0, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	1.73	0.43	12
	40-59	05-06	1.39	0.35	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	1.68	0.49	10
	20-39	05-06	*	*	16
	40-59	05-06	1.42	0.4	13
	60+	05-06	1.86	0.4	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	2.74	0.63	14
	20-39	05-06	3.44	0.31	7
	40-59	05-06	3.08	0.53	7
	60+	05-06	3.28	0.37	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.5.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1884
	01-02	*	< LOD	< LOD	< LOD	< LOD	1202
	03-04	*	< LOD	< LOD	< LOD	< LOD	1849
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	652
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	581
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1232
	01-02	*	< LOD	< LOD	< LOD	< LOD	1202
	03-04	*	< LOD	< LOD	< LOD	< LOD	1268
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	904
	01-02	*	< LOD	< LOD	< LOD	< LOD	541
	03-04	*	< LOD	< LOD	< LOD	< LOD	905
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	980
	01-02	*	< LOD	< LOD	< LOD	< LOD	661
	03-04	*	< LOD	< LOD	< LOD	< LOD	944
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	623
	01-02	*	< LOD	< LOD	< LOD	< LOD	257
	03-04	*	< LOD	< LOD	< LOD	< LOD	420
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	404
	01-02	*	< LOD	< LOD	< LOD	< LOD	212
	03-04	*	< LOD	< LOD	< LOD	< LOD	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	705
	01-02	*	< LOD	< LOD	< LOD	< LOD	653
	03-04	*	< LOD	< LOD	< LOD	< LOD	857

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	11.6	2.8	12
	40-59	05-06	10.6	2.7	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	8.69	2.41	10
	20-39	05-06	*	*	16
	40-59	05-06	10.6	3.1	13
	60+	05-06	13.5	2.9	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	13.4	3	14
	20-39	05-06	19.7	1.7	7
	40-59	05-06	19.6	3.5	7
	60+	05-06	21.7	1.8	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8-Pentachlorodibenzofuran (PeCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1922
	01-02	*	< LOD	< LOD	< LOD	< LOD	1235
	03-04	*	< LOD	< LOD	< LOD	< LOD	1867
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	663
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	586
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1259
	01-02	*	< LOD	< LOD	< LOD	< LOD	1235
	03-04	*	< LOD	< LOD	< LOD	< LOD	1281
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	920
	01-02	*	< LOD	< LOD	< LOD	< LOD	565
	03-04	*	< LOD	< LOD	< LOD	< LOD	917
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1002
	01-02	*	< LOD	< LOD	< LOD	< LOD	670
	03-04	*	< LOD	< LOD	< LOD	< LOD	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	637
	01-02	*	< LOD	< LOD	< LOD	< LOD	263
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	217
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	717
	01-02	*	< LOD	< LOD	< LOD	< LOD	670
	03-04	*	< LOD	< LOD	< LOD	< LOD	869

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 13.2, 5.8, and 7.1, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8-Pentachlorodibenzofuran (PeCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	.331	0.082	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8-Pentachlorodibenzofuran (PeCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1922
	01-02	*	< LOD	< LOD	< LOD	< LOD	1235
	03-04	*	< LOD	< LOD	< LOD	< LOD	1867
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	663
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	586
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1259
	01-02	*	< LOD	< LOD	< LOD	< LOD	1235
	03-04	*	< LOD	< LOD	< LOD	< LOD	1281
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	920
	01-02	*	< LOD	< LOD	< LOD	< LOD	565
	03-04	*	< LOD	< LOD	< LOD	< LOD	917
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1002
	01-02	*	< LOD	< LOD	< LOD	< LOD	670
	03-04	*	< LOD	< LOD	< LOD	< LOD	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	637
	01-02	*	< LOD	< LOD	< LOD	< LOD	263
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	217
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	717
	01-02	*	< LOD	< LOD	< LOD	< LOD	670
	03-04	*	< LOD	< LOD	< LOD	< LOD	869

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 1,2,3,7,8-Pentachlorodibenzofuran (PeCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	2.32	0.61	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,4,7,8-Pentachlorodibenzofuran (PeCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	15.9 (13.7-17.1)	1895
	01-02	*	< LOD	9.20 (7.50-11.1)	14.3 (12.3-16.2)	18.1 (16.1-21.1)	1230
	03-04	*	< LOD	6.80 (<LOD-7.20)	9.90 (8.90-10.7)	12.3 (11.0-13.3)	1871
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	656
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	586
20 years and older	99-00	*	< LOD	< LOD	12.7 (<LOD-13.9)	16.1 (14.1-17.6)	1239
	01-02	*	< LOD	9.20 (7.50-11.1)	14.3 (12.3-16.2)	18.1 (16.1-21.1)	1230
	03-04	*	< LOD	7.20 (<LOD-7.70)	10.3 (9.40-11.2)	13.0 (11.4-14.7)	1285
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	13.9 (12.8-15.4)	906
	01-02	*	< LOD	9.40 (7.30-11.6)	14.1 (11.6-16.7)	16.7 (14.0-22.8)	560
	03-04	*	< LOD	7.00 (<LOD-7.30)	9.80 (8.70-10.6)	12.2 (10.4-14.8)	920
Females	99-00	*	< LOD	< LOD	13.1 (<LOD-16.1)	16.7 (14.3-19.2)	989
	01-02	*	< LOD	9.10 (7.50-10.7)	14.5 (13.0-16.1)	18.5 (16.4-21.2)	670
	03-04	*	< LOD	< LOD	10.3 (8.90-11.0)	12.6 (11.0-13.7)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	632
	01-02	*	< LOD	< LOD	7.80 (6.10-9.80)	9.80 (7.80-12.5)	264
	03-04	*	< LOD	< LOD	< LOD	7.90 (<LOD-10.5)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	16.3 (13.4-19.0)	400
	01-02	*	< LOD	8.40 (6.80-9.30)	14.1 (11.7-16.4)	19.1 (15.1-23.0)	216
	03-04	*	< LOD	< LOD	9.80 (8.00-13.9)	15.9 (9.70-23.9)	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	15.6 (13.8-17.1)	706
	01-02	*	5.90 (<LOD-6.90)	10.3 (8.10-12.0)	15.5 (13.0-17.6)	18.5 (16.5-22.2)	665
	03-04	*	< LOD	7.10 (<LOD-7.70)	10.3 (9.10-11.2)	12.4 (11.0-13.7)	873

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.7, 5.5, and 6.8, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,4,7,8-Pentachlorodibenzofuran (PeCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	2.25	0.14	9
	20-39	05-06	3.46	0.3	12
	40-59	05-06	5.82	0.26	12
	60+	05-06	8.03	0.43	15
Non-Hispanic White Female	12-19	05-06	1.78	0.12	10
	20-39	05-06	2.71	0.17	16
	40-59	05-06	4.56	0.17	13
	60+	05-06	8.49	0.5	17
Non-Hispanic Black Male	12-19	05-06	1.84	0.1	13
	20-39	05-06	3.49	1.01	6
	40-59	05-06	3.84	0.28	5
	60+	05-06	7.19	0.94	5
Non-Hispanic Black Female	12-19	05-06	1.17	0.12	14
	20-39	05-06	2.12	0.15	7
	40-59	05-06	3.98	0.4	7
	60+	05-06	12.7	2.5	5
Mexican American Male	12-19	05-06	1.64	0.25	11
	20-39	05-06	2.64	0.26	9
	40-59	05-06	3.86	0.39	4
	60+	05-06	5.88	0.45	4
Mexican American Female	12-19	05-06	.941	0.165	16
	20-39	05-06	2.06	0.23	9
	40-59	05-06	3.49	0.19	6
	60+	05-06	6.10	0.42	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,4,7,8-Pentachlorodibenzofuran (PeCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	103 (96.0-113)	1895
	01-02	*	< LOD	60.8 (51.3-70.6)	99.4 (83.4-114)	134 (113-152)	1230
	03-04	*	< LOD	44.2 (<LOD-46.9)	65.7 (60.5-71.1)	83.9 (74.9-92.9)	1871
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	656
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	586
20 years and older	99-00	*	< LOD	< LOD	85.2 (<LOD-96.0)	107 (98.3-125)	1239
	01-02	*	< LOD	60.8 (51.3-70.6)	99.4 (83.4-114)	134 (113-152)	1230
	03-04	*	< LOD	46.5 (<LOD-50.6)	69.3 (62.2-74.6)	86.8 (78.3-99.8)	1285
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	95.7 (80.1-101)	906
	01-02	*	< LOD	62.7 (49.3-76.6)	98.5 (76.6-134)	135 (105-160)	560
	03-04	*	< LOD	43.4 (<LOD-46.3)	62.0 (56.8-71.9)	83.2 (72.4-95.0)	920
Females	99-00	*	< LOD	< LOD	90.0 (<LOD-101)	111 (97.2-129)	989
	01-02	*	< LOD	59.8 (52.0-67.6)	100 (90.5-105)	126 (107-146)	670
	03-04	*	< LOD	< LOD	66.9 (59.7-74.3)	84.4 (74.7-95.7)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	632
	01-02	*	< LOD	< LOD	55.8 (48.7-69.7)	76.3 (61.9-92.4)	264
	03-04	*	< LOD	< LOD	< LOD	57.1 (<LOD-70.2)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	105 (82.7-116)	400
	01-02	*	< LOD	50.0 (41.3-60.4)	89.5 (72.3-107)	121 (92.0-154)	216
	03-04	*	< LOD	< LOD	66.8 (48.5-93.2)	97.4 (79.1-129)	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	103 (96.8-122)	706
	01-02	*	37.5 (<LOD-44.1)	65.7 (52.9-78.2)	104 (90.7-118)	136 (118-154)	665
	03-04	*	< LOD	45.9 (<LOD-50.6)	68.3 (61.3-73.6)	85.0 (74.6-102)	873

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,4,7,8-Pentachlorodibenzofuran (PeCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	12.4	0.9	9
	20-39	05-06	25.6	3.2	12
	40-59	05-06	43.9	2.4	12
	60+	05-06	54.6	2.6	15
Non-Hispanic White Female	12-19	05-06	9.51	0.56	10
	20-39	05-06	17.7	1.3	16
	40-59	05-06	33.7	2.2	13
	60+	05-06	62.0	3.8	17
Non-Hispanic Black Male	12-19	05-06	9.25	0.4	13
	20-39	05-06	20.7	6.2	6
	40-59	05-06	24.5	2.1	5
	60+	05-06	42.9	5.6	5
Non-Hispanic Black Female	12-19	05-06	5.82	0.59	14
	20-39	05-06	12.2	0.9	7
	40-59	05-06	25.4	2.5	7
	60+	05-06	84.2	16	5
Mexican American Male	12-19	05-06	8.87	1.26	11
	20-39	05-06	18.5	1.4	9
	40-59	05-06	31.2	4.2	4
	60+	05-06	42.4	2.1	4
Mexican American Female	12-19	05-06	5.26	0.9	16
	20-39	05-06	13.8	1.7	9
	40-59	05-06	26.0	2	6
	60+	05-06	46.7	6.2	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,7,8-Tetrachlorodibenzofuran (TCDF) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1903
	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	*	< LOD	< LOD	< LOD	< LOD	1868
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	660
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	586
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1243
	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	*	< LOD	< LOD	< LOD	< LOD	1282
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	912
	01-02	*	< LOD	< LOD	< LOD	< LOD	558
	03-04	*	< LOD	< LOD	< LOD	< LOD	917
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	991
	01-02	*	< LOD	< LOD	< LOD	< LOD	671
	03-04	*	< LOD	< LOD	< LOD	< LOD	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	628
	01-02	*	< LOD	< LOD	< LOD	< LOD	262
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	217
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	707
	01-02	*	< LOD	< LOD	< LOD	< LOD	667
	03-04	*	< LOD	< LOD	< LOD	< LOD	870

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 11.9, 5.2, and 6.0, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,7,8-Tetrachlorodibenzofuran (TCDF) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,7,8-Tetrachlorodibenzofuran (TCDF) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1903
	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	*	< LOD	< LOD	< LOD	< LOD	1868
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	660
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	586
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1243
	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	*	< LOD	< LOD	< LOD	< LOD	1282
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	912
	01-02	*	< LOD	< LOD	< LOD	< LOD	558
	03-04	*	< LOD	< LOD	< LOD	< LOD	917
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	991
	01-02	*	< LOD	< LOD	< LOD	< LOD	671
	03-04	*	< LOD	< LOD	< LOD	< LOD	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	628
	01-02	*	< LOD	< LOD	< LOD	< LOD	262
	03-04	*	< LOD	< LOD	< LOD	< LOD	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	217
	03-04	*	< LOD	< LOD	< LOD	< LOD	454
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	707
	01-02	*	< LOD	< LOD	< LOD	< LOD	667
	03-04	*	< LOD	< LOD	< LOD	< LOD	870

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,7,8-Tetrachlorodibenzofuran (TCDF) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	*	*	12
	40-59	05-06	*	*	12
	60+	05-06	*	*	15
Non-Hispanic White Female	12-19	05-06	*	*	10
	20-39	05-06	*	*	16
	40-59	05-06	*	*	13
	60+	05-06	*	*	17
Non-Hispanic Black Male	12-19	05-06	*	*	13
	20-39	05-06	*	*	6
	40-59	05-06	*	*	5
	60+	05-06	*	*	5
Non-Hispanic Black Female	12-19	05-06	*	*	14
	20-39	05-06	*	*	7
	40-59	05-06	*	*	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	*	*	11
	20-39	05-06	*	*	9
	40-59	05-06	*	*	4
	60+	05-06	*	*	4
Mexican American Female	12-19	05-06	*	*	16
	20-39	05-06	*	*	9
	40-59	05-06	*	*	6
	60+	05-06	*	*	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4'-Tetrachlorobiphenyl (PCB 77) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	18.2	2.6	12
	40-59	05-06	17.6	2.2	12
	60+	05-06	22.7	3.1	15
Non-Hispanic White Female	12-19	05-06	26.2	4.2	10
	20-39	05-06	29.3	8.4	16
	40-59	05-06	21.4	3.2	13
	60+	05-06	22.0	3	17
Non-Hispanic Black Male	12-19	05-06	26.8	2.8	13
	20-39	05-06	23.1	5	6
	40-59	05-06	15.0	2.6	5
	60+	05-06	21.5	3.9	5
Non-Hispanic Black Female	12-19	05-06	26.6	5.6	14
	20-39	05-06	14.9	1.9	7
	40-59	05-06	13.1	2.1	7
	60+	05-06	18.5	3.2	5
Mexican American Male	12-19	05-06	36.7	5.7	11
	20-39	05-06	18.1	3.1	9
	40-59	05-06	15.8	2.3	4
	60+	05-06	24.9	6.6	4
Mexican American Female	12-19	05-06	22.6	2.2	16
	20-39	05-06	21.2	3.3	9
	40-59	05-06	17.4	2.4	6
	60+	05-06	15.5	2.4	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.6.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4'-Tetrachlorobiphenyl (PCB 77) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
	20-39	05-06	131	19	12
	40-59	05-06	131	15	12
	60+	05-06	154	20	15
Non-Hispanic White Female	12-19	05-06	142	25	10
	20-39	05-06	183	48	16
	40-59	05-06	154	24	13
	60+	05-06	162	23	17
Non-Hispanic Black Male	12-19	05-06	136	14	13
	20-39	05-06	133	26	6
	40-59	05-06	95.4	16.8	5
	60+	05-06	128	23	5
Non-Hispanic Black Female	12-19	05-06	131	28	14
	20-39	05-06	86.0	11.8	7
	40-59	05-06	84.4	14.1	7
	60+	05-06	123	21	5
Mexican American Male	12-19	05-06	198	31	11
	20-39	05-06	124	17	9
	40-59	05-06	127	16	4
	60+	05-06	175	44	4
Mexican American Female	12-19	05-06	124	13	16
	20-39	05-06	139	21	9
	40-59	05-06	129	20	6
	60+	05-06	120	29	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,4,4',5-Tetrachlorobiphenyl (PCB 81) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1883
	01-02	*	< LOD	< LOD	< LOD	< LOD	1215
	03-04	*	< LOD	< LOD	< LOD	13.4 (<LOD-16.4)	1860
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	651
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	14.7 (<LOD-16.5)	579
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1232
	01-02	*	< LOD	< LOD	< LOD	< LOD	1215
	03-04	*	< LOD	< LOD	< LOD	13.1 (<LOD-16.5)	1281
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	900
	01-02	*	< LOD	< LOD	< LOD	< LOD	554
	03-04	*	< LOD	< LOD	< LOD	14.4 (<LOD-17.9)	913
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	983
	01-02	*	< LOD	< LOD	< LOD	< LOD	661
	03-04	*	< LOD	< LOD	< LOD	< LOD	947
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	621
	01-02	*	< LOD	< LOD	< LOD	< LOD	259
	03-04	*	< LOD	< LOD	< LOD	18.7 (<LOD-25.2)	420
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	405
	01-02	*	< LOD	< LOD	< LOD	< LOD	218
	03-04	*	< LOD	< LOD	< LOD	< LOD	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	699
	01-02	*	< LOD	< LOD	< LOD	< LOD	657
	03-04	*	< LOD	< LOD	< LOD	13.1 (<LOD-16.4)	867

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 68.4, 26.8, and 13.1, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,4,4',5-Tetrachlorobiphenyl (PCB 81) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	1.90	0.45	9
	20-39	05-06	1.29	0.13	12
	40-59	05-06	1.85	0.27	12
	60+	05-06	2.52	0.44	15
Non-Hispanic White Female	12-19	05-06	1.74	0.19	10
	20-39	05-06	1.90	0.3	16
	40-59	05-06	1.82	0.15	13
	60+	05-06	2.74	0.19	17
Non-Hispanic Black Male	12-19	05-06	1.75	0.14	13
	20-39	05-06	1.41	0.22	6
	40-59	05-06	1.35	0.14	5
	60+	05-06	2.22	0.2	5
Non-Hispanic Black Female	12-19	05-06	1.81	0.26	14
	20-39	05-06	1.36	0.18	7
	40-59	05-06	1.48	0.17	7
	60+	05-06	2.74	0.27	5
Mexican American Male	12-19	05-06	2.14	0.28	11
	20-39	05-06	1.56	0.21	9
	40-59	05-06	1.58	0.2	4
	60+	05-06	2.06	0.31	4
Mexican American Female	12-19	05-06	1.60	0.11	16
	20-39	05-06	1.63	0.16	9
	40-59	05-06	1.69	0.12	6
	60+	05-06	2.10	0.12	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.6.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,4,4',5-Tetrachlorobiphenyl (PCB 81) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1883
	01-02	*	< LOD	< LOD	< LOD	< LOD	1215
	03-04	*	< LOD	< LOD	< LOD	80.3 (<LOD-96.2)	1860
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	651
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	71.4 (<LOD-82.8)	579
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1232
	01-02	*	< LOD	< LOD	< LOD	< LOD	1215
	03-04	*	< LOD	< LOD	< LOD	80.7 (<LOD-99.3)	1281
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	900
	01-02	*	< LOD	< LOD	< LOD	< LOD	554
	03-04	*	< LOD	< LOD	< LOD	81.1 (<LOD-104)	913
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	983
	01-02	*	< LOD	< LOD	< LOD	< LOD	661
	03-04	*	< LOD	< LOD	< LOD	< LOD	947
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	621
	01-02	*	< LOD	< LOD	< LOD	< LOD	259
	03-04	*	< LOD	< LOD	< LOD	103 (<LOD-199)	420
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	405
	01-02	*	< LOD	< LOD	< LOD	< LOD	218
	03-04	*	< LOD	< LOD	< LOD	< LOD	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	699
	01-02	*	< LOD	< LOD	< LOD	< LOD	657
	03-04	*	< LOD	< LOD	< LOD	80.5 (<LOD-106)	867

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,4,4',5-Tetrachlorobiphenyl (PCB 81) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	10.2	2.3	9
	20-39	05-06	9.45	1.01	12
	40-59	05-06	14.0	2.2	12
	60+	05-06	17.0	3	15
Non-Hispanic White Female	12-19	05-06	9.34	1.08	10
	20-39	05-06	12.1	1.7	16
	40-59	05-06	13.4	1.3	13
	60+	05-06	20.0	1.4	17
Non-Hispanic Black Male	12-19	05-06	8.90	0.72	13
	20-39	05-06	8.17	1.21	6
	40-59	05-06	8.53	0.92	5
	60+	05-06	13.1	0.9	5
Non-Hispanic Black Female	12-19	05-06	8.95	1.27	14
	20-39	05-06	7.93	1.13	7
	40-59	05-06	9.50	0.99	7
	60+	05-06	18.3	1.6	5
Mexican American Male	12-19	05-06	11.7	1.5	11
	20-39	05-06	11.0	1.2	9
	40-59	05-06	13.0	1.8	4
	60+	05-06	14.6	1.9	4
Mexican American Female	12-19	05-06	8.79	0.66	16
	20-39	05-06	10.7	1	9
	40-59	05-06	12.5	1.1	6
	60+	05-06	16.1	2.3	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4',5-Pentachlorobiphenyl (PCB 126) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	28.5 (24.9-32.8)	53.2 (45.7-59.9)	80.5 (62.8-100)	1896
	01-02	22.7 (20.9-24.7)	24.5 (22.2-26.8)	40.8 (36.1-47.5)	69.3 (61.6-80.8)	108 (92.7-116)	1226
	03-04	16.3 (14.9-17.9)	14.7 (<LOD-16.5)	24.8 (22.4-27.4)	46.7 (41.6-51.9)	68.7 (58.1-84.4)	1860
Age group							
12-19 years	99-00	*	< LOD	< LOD	24.3 (<LOD-27.5)	31.1 (26.4-36.4)	658
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	17.0 (15.2-20.2)	21.2 (17.3-25.6)	577
20 years and older	99-00	*	< LOD	30.8 (27.2-36.3)	57.1 (50.5-65.8)	89.5 (66.1-110)	1238
	01-02	22.7 (20.9-24.7)	24.5 (22.2-26.8)	40.8 (36.1-47.5)	69.3 (61.6-80.8)	108 (92.7-116)	1226
	03-04	17.6 (16.0-19.3)	16.0 (14.2-18.6)	26.8 (24.2-30.3)	49.8 (43.5-59.1)	74.8 (60.2-94.4)	1283
Gender							
Males	99-00	*	< LOD	25.8 (<LOD-28.9)	41.6 (36.1-45.7)	62.0 (49.7-75.0)	911
	01-02	20.3 (18.5-22.3)	23.1 (20.8-25.4)	36.6 (32.8-39.4)	52.7 (46.4-62.3)	81.9 (61.0-101)	561
	03-04	14.9 (<LOD-16.4)	< LOD	21.8 (20.1-25.4)	38.5 (31.0-46.1)	51.5 (46.5-60.5)	912
Females	99-00	*	< LOD	33.6 (27.4-41.4)	59.4 (53.0-78.7)	98.1 (69.9-120)	985
	01-02	25.1 (22.9-27.4)	26.1 (23.0-28.8)	48.6 (41.4-54.4)	82.9 (71.1-96.8)	116 (98.6-128)	665
	03-04	17.8 (16.0-19.7)	15.7 (<LOD-18.4)	27.6 (24.3-33.4)	57.1 (46.0-67.1)	82.5 (63.0-109)	948
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	23.8 (<LOD-30.3)	43.2 (37.0-52.6)	66.1 (52.4-79.2)	631
	01-02	17.8 (15.5-20.4)	19.9 (18.3-21.2)	28.5 (26.5-34.2)	47.9 (39.7-54.9)	69.2 (49.3-103)	262
	03-04	15.5 (<LOD-17.9)	14.2 (<LOD-17.5)	22.5 (18.9-26.8)	31.2 (27.4-37.8)	42.0 (33.5-48.0)	420
Non-Hispanic blacks	99-00	*	< LOD	30.6 (25.2-43.1)	67.4 (45.0-126)	126 (67.4-224)	404
	01-02	22.2 (18.2-27.0)	22.1 (19.6-25.4)	44.3 (37.9-51.5)	88.4 (59.0-111)	115 (96.1-153)	217
	03-04	16.5 (13.9-19.5)	< LOD	23.5 (18.9-34.6)	63.4 (41.5-126)	142 (52.1-292)	452
Non-Hispanic whites	99-00	*	< LOD	28.3 (<LOD-34.0)	50.5 (41.6-57.0)	67.8 (57.0-94.1)	704
	01-02	23.1 (20.9-25.4)	24.7 (22.0-27.6)	42.0 (35.4-49.6)	72.3 (63.9-82.9)	114 (91.0-128)	663
	03-04	16.0 (14.3-17.8)	14.5 (<LOD-16.9)	24.3 (21.4-28.2)	46.5 (41.6-50.9)	64.0 (52.1-76.1)	869

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 23.2, 10.8, and 13.9, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4',5-Pentachlorobiphenyl (PCB 126) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	6.23	0.47	9
	20-39	05-06	8.13	0.78	12
	40-59	05-06	17.6	2.4	12
	60+	05-06	26.8	3.3	15
Non-Hispanic White Female	12-19	05-06	5.64	0.28	10
	20-39	05-06	9.56	1.07	16
	40-59	05-06	16.4	2	13
	60+	05-06	35.6	2.6	17
Non-Hispanic Black Male	12-19	05-06	5.77	0.42	13
	20-39	05-06	6.75	1	6
	40-59	05-06	15.0	3.4	5
	60+	05-06	30.9	5.4	5
Non-Hispanic Black Female	12-19	05-06	5.64	0.31	14
	20-39	05-06	9.71	1.08	7
	40-59	05-06	20.2	2.5	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	6.01	0.34	11
	20-39	05-06	8.96	0.46	9
	40-59	05-06	16.0	2.2	4
	60+	05-06	26.4	4.8	4
Mexican American Female	12-19	05-06	5.87	0.42	16
	20-39	05-06	8.21	0.62	9
	40-59	05-06	19.2	2.6	6
	60+	05-06	40.2	7.1	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.5.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4',5-Pentachlorobiphenyl (PCB 126) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	177 (151-212)	336 (294-404)	564 (455-663)	1896
	01-02	147 (135-160)	158 (143-175)	270 (249-298)	482 (436-541)	738 (625-840)	1226
	03-04	100 (91.0-110)	89.8 (<LOD-101)	159 (146-178)	308 (273-349)	475 (378-590)	1860
Age group							
12-19 years	99-00	*	< LOD	< LOD	108 (<LOD-126)	144 (116-196)	658
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	88.2 (78.8-106)	109 (89.2-135)	577
20 years and older	99-00	*	< LOD	204 (173-232)	381 (319-451)	596 (479-709)	1238
	01-02	147 (135-160)	158 (143-175)	270 (249-298)	482 (436-541)	738 (625-840)	1226
	03-04	111 (100-122)	100 (88.7-115)	176 (158-194)	326 (286-394)	534 (394-668)	1283
Gender							
Males	99-00	*	< LOD	154 (<LOD182)	260 (241-274)	382 (308-509)	911
	01-02	134 (121-149)	152 (130-169)	249 (233-267)	399 (346-467)	566 (467-693)	561
	03-04	92.3 (<LOD-102)	< LOD	148 (131-168)	245 (203-296)	360 (296-442)	912
Females	99-00	*	< LOD	212 (159-250)	429 (350-531)	648 (534-744)	985
	01-02	159 (146-172)	160 (149-186)	301 (270-348)	550 (483-637)	818 (658-958)	665
	03-04	108 (96.2-121)	94.5 (<LOD-110)	180 (153-212)	370 (303-455)	559 (438-843)	948
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	146 (<LOD213)	285 (229-364)	424 (361-534)	631
	01-02	114 (97.8-133)	125 (106-150)	217 (175-257)	335 (272-460)	593 (354-784)	262
	03-04	95.3 (<LOD-111)	91.4 (<LOD-99.4)	145 (116-180)	238 (172-288)	298 (238-461)	420
Non-Hispanic blacks	99-00	*	< LOD	192 (147-258)	457 (308-691)	786 (457-1520)	404
	01-02	131 (106-161)	128 (110-158)	266 (214-304)	532 (361-662)	779 (585-1060)	217
	03-04	94.3 (79.0-112)	< LOD	143 (106-254)	483 (254-847)	976 (483-1430)	452
Non-Hispanic whites	99-00	*	< LOD	175 (<LOD215)	311 (274-356)	508 (394-660)	704
	01-02	150 (136-164)	160 (146-179)	276 (249-301)	486 (431-553)	746 (600-937)	663
	03-04	98.4 (87.5-111)	89.4 (<LOD-104)	158 (142-182)	297 (260-346)	431 (365-541)	869

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4',5-Pentachlorobiphenyl (PCB 126) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	34.3	3.1	9
	20-39	05-06	59.8	7.5	12
	40-59	05-06	134	21	12
	60+	05-06	181	19	15
Non-Hispanic White Female	12-19	05-06	30.0	1.6	10
	20-39	05-06	62.7	8.1	16
	40-59	05-06	123	17	13
	60+	05-06	259	18	17
Non-Hispanic Black Male	12-19	05-06	29.2	2.3	13
	20-39	05-06	40.3	6.6	6
	40-59	05-06	96.2	22.7	5
	60+	05-06	183	31	5
Non-Hispanic Black Female	12-19	05-06	27.9	1.5	14
	20-39	05-06	56.3	6.6	7
	40-59	05-06	129	16	7
	60+	05-06	*	*	5
Mexican American Male	12-19	05-06	32.6	1.7	11
	20-39	05-06	63.0	2.2	9
	40-59	05-06	131	22	4
	60+	05-06	188	32	4
Mexican American Female	12-19	05-06	32.0	2.3	16
	20-39	05-06	54.9	4.8	9
	40-59	05-06	143	21	6
	60+	05-06	301	54	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4',5,5'-Hexachlorobiphenyl (PCB 169) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	34.4 (31.0-38.7)	44.9 (40.6-48.9)	1888
	01-02	17.9 (16.0-19.9)	19.0 (17.0-22.0)	33.1 (27.7-38.6)	50.0 (43.9-55.0)	60.9 (56.1-65.8)	1223
	03-04	*	< LOD	19.5 (16.8-22.7)	31.0 (27.9-36.0)	40.6 (36.5-47.3)	1866
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	648
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	581
20 years and older	99-00	*	< LOD	< LOD	36.4 (33.8-40.3)	47.8 (42.8-51.0)	1240
	01-02	17.9 (16.0-19.9)	19.0 (17.0-22.0)	33.1 (27.7-38.6)	50.0 (43.9-55.0)	60.9 (56.1-65.8)	1223
	03-04	*	< LOD	21.9 (19.0-24.2)	32.7 (28.6-37.3)	43.2 (37.3-49.5)	1285
Gender							
Males	99-00	*	< LOD	< LOD	36.2 (31.9-40.1)	44.3 (39.0-51.3)	908
	01-02	20.2 (17.8-22.8)	22.1 (18.5-25.3)	36.0 (29.5-43.2)	53.7 (46.5-57.3)	60.9 (55.7-69.0)	559
	03-04	*	< LOD	22.3 (17.7-24.8)	36.3 (29.3-41.1)	46.1 (38.8-52.0)	917
Females	99-00	*	< LOD	< LOD	34.0 (29.4-38.6)	46.5 (37.9-51.1)	980
	01-02	16.0 (14.2-18.1)	17.0 (14.7-19.2)	30.2 (24.5-36.5)	46.4 (40.7-51.9)	60.9 (52.2-70.0)	664
	03-04	*	< LOD	18.4 (<LOD-20.4)	28.4 (27.2-30.6)	35.8 (31.0-41.1)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	31.9 (28.8-35.2)	622
	01-02	*	< LOD	15.0 (11.3-18.3)	25.4 (18.3-32.3)	32.5 (26.7-41.9)	260
	03-04	*	< LOD	< LOD	< LOD	20.9 (16.5-30.2)	420
Non-Hispanic blacks	99-00	*	< LOD	< LOD	40.3 (28.7-48.6)	52.2 (44.3-63.6)	403
	01-02	17.2 (15.4-19.1)	18.5 (15.7-20.0)	31.7 (25.8-35.4)	47.5 (42.0-54.0)	57.3 (48.3-64.5)	217
	03-04	*	< LOD	17.3 (<LOD-22.6)	31.5 (26.8-36.9)	58.7 (31.5-97.9)	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	34.6 (31.7-40.1)	45.3 (40.1-50.9)	709
	01-02	19.5 (17.2-22.2)	21.4 (17.4-24.7)	36.0 (29.8-41.1)	53.5 (47.0-58.3)	64.3 (59.2-72.8)	662
	03-04	*	< LOD	21.2 (17.4-24.5)	32.6 (27.3-38.8)	41.1 (35.9-50.2)	873

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 27.0, 11.0, and 15.9, respectively.

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4',5,5'-Hexachlorobiphenyl (PCB 169) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in pg/g lipid or parts per trillion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	5.14	0.42	9
	20-39	05-06	8.17	0.77	12
	40-59	05-06	19.9	1.4	12
	60+	05-06	35.7	1.8	15
Non-Hispanic White Female	12-19	05-06	3.68	0.19	10
	20-39	05-06	6.72	0.58	16
	40-59	05-06	14.2	0.7	13
	60+	05-06	31.4	2.2	17
Non-Hispanic Black Male	12-19	05-06	5.19	0.31	13
	20-39	05-06	7.95	0.57	6
	40-59	05-06	17.5	0.7	5
	60+	05-06	39.1	3.8	5
Non-Hispanic Black Female	12-19	05-06	3.19	0.11	14
	20-39	05-06	6.19	0.45	7
	40-59	05-06	13.9	1.6	7
	60+	05-06	45.0	8.5	5
Mexican American Male	12-19	05-06	3.74	0.5	11
	20-39	05-06	6.10	0.14	9
	40-59	05-06	11.1	0.3	4
	60+	05-06	27.2	2.5	4
Mexican American Female	12-19	05-06	3.17	0.15	16
	20-39	05-06	4.40	0.3	9
	40-59	05-06	10.2	1.5	6
	60+	05-06	18.7	1.8	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey year 05-06 is 0.6.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4',5,5'-Hexachlorobiphenyl (PCB 169) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	230 (219-244)	287 (274-319)	1888
	01-02	115 (103-129)	125 (108-146)	217 (189-253)	344 (300-376)	416 (379-490)	1223
	03-04	*	< LOD	133 (114-149)	203 (191-226)	269 (243-305)	1866
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	648
	01-02	†	†	†	†	†	†
	03-04	*	< LOD	< LOD	< LOD	< LOD	581
20 years and older	99-00	*	< LOD	< LOD	241 (227-261)	303 (281-339)	1240
	01-02	115 (103-129)	125 (108-146)	217 (189-253)	344 (300-376)	416 (379-490)	1223
	03-04	*	< LOD	146 (130-160)	216 (198-242)	287 (245-325)	1285
Gender							
Males	99-00	*	< LOD	< LOD	229 (215-258)	286 (256-338)	908
	01-02	133 (117-152)	151 (130-168)	244 (198-293)	363 (309-409)	449 (372-529)	559
	03-04	*	< LOD	143 (122-159)	225 (190-259)	291 (245-334)	917
Females	99-00	*	< LOD	< LOD	234 (197-257)	293 (261-346)	980
	01-02	102 (90.0-115)	107 (91.0-123)	195 (170-234)	319 (282-368)	402 (373-462)	664
	03-04	*	< LOD	120 (<LOD140)	194 (182-203)	249 (222-284)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	215 (181-261)	622
	01-02	*	< LOD	98.1 (79.3-138)	184 (143-250)	266 (179-341)	260
	03-04	*	< LOD	< LOD	< LOD	168 (109-237)	420
Non-Hispanic blacks	99-00	*	< LOD	< LOD	251 (190-299)	320 (262-410)	403
	01-02	101 (91.2-113)	104 (88.0-118)	190 (147-234)	299 (267-335)	362 (309-423)	217
	03-04	*	< LOD	110 (<LOD-144)	216 (162-257)	338 (221-556)	453
Non-Hispanic whites	99-00	*	< LOD	< LOD	238 (222-257)	286 (267-340)	709
	01-02	127 (112-144)	145 (124-157)	229 (195-275)	368 (317-388)	438 (388-515)	662
	03-04	*	< LOD	141 (116-165)	210 (189-242)	277 (239-324)	873

† Data not collected for this age group for Survey year 01-02.

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 3,3',4,4',5,5'-Hexachlorobiphenyl (PCB 169) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in fg/g of serum or parts per quadrillion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	28.1	2.4	9
	20-39	05-06	60.0	7.2	12
	40-59	05-06	150	13	12
	60+	05-06	242	10	15
Non-Hispanic White Female	12-19	05-06	19.7	1.3	10
	20-39	05-06	44.1	4.5	16
	40-59	05-06	106	9	13
	60+	05-06	228	16	17
Non-Hispanic Black Male	12-19	05-06	26.0	1.3	13
	20-39	05-06	46.9	3.4	6
	40-59	05-06	112	5	5
	60+	05-06	235	26	5
Non-Hispanic Black Female	12-19	05-06	15.8	0.5	14
	20-39	05-06	35.8	2.8	7
	40-59	05-06	88.8	10.6	7
	60+	05-06	299	54	5
Mexican American Male	12-19	05-06	20.0	2.4	11
	20-39	05-06	43.3	1.7	9
	40-59	05-06	90.6	5.2	4
	60+	05-06	194	10	4
Mexican American Female	12-19	05-06	17.3	0.8	16
	20-39	05-06	29.6	2.4	9
	40-59	05-06	75.8	11.8	6
	60+	05-06	144	23	3

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 105) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1915
	01-02	*	< LOD	< LOD	< LOD	< LOD	2307
	03-04	1.20 (1.09-1.31)	1.09 (1.00-1.20)	1.90 (1.75-2.20)	4.04 (3.40-4.89)	6.24 (5.20-7.79)	1879
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	665
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.686 (.603-.781)	.700 (.600-.720)	1.00 (.900-1.11)	1.50 (1.20-2.00)	2.26 (1.50-3.50)	593
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1250
	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	1.30 (1.18-1.43)	1.15 (1.04-1.30)	2.10 (1.86-2.40)	4.44 (3.80-5.20)	6.82 (5.70-8.30)	1286
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	913
	01-02	*	< LOD	< LOD	< LOD	< LOD	1075
	03-04	1.02 (.923-1.12)	.980 (.850-1.08)	1.60 (1.40-1.80)	3.20 (2.76-3.42)	4.70 (3.83-5.50)	937
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1002
	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	1.40 (1.25-1.57)	1.20 (1.05-1.40)	2.30 (2.00-2.79)	5.17 (4.47-5.90)	7.70 (6.00-9.55)	942
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	635
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	.719 (.624-.829)	.700 (.600-.800)	1.10 (.900-1.36)	1.90 (1.80-2.20)	2.90 (1.98-3.52)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	12.8 (<LOD-18.0)	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	1.41 (1.15-1.74)	1.14 (.980-1.32)	2.47 (1.70-3.90)	6.70 (4.09-10.1)	11.9 (6.40-18.1)	456
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	714
	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	1.21 (1.08-1.36)	1.10 (1.00-1.20)	1.90 (1.74-2.28)	3.91 (3.22-4.85)	5.93 (4.98-7.79)	878

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.4, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 105) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.507	0.097	9
		07-08	*	*	6
	20-39	05-06	.671	0.134	12
		07-08	.808	0.058	15
	40-59	05-06	*	*	12
		07-08	1.02	0.07	16
	60+	05-06	2.15	0.26	15
		07-08	3.01	0.36	23
Non-Hispanic White Female	12-19	05-06	.482	0.041	10
		07-08	.555	0.065	7
	20-39	05-06	.598	0.044	16
		07-08	1.06	0.16	13
	40-59	05-06	1.29	0.15	13
		07-08	1.36	0.13	17
	60+	05-06	3.84	0.41	17
		07-08	4.09	0.7	21
Non-Hispanic Black Male	12-19	05-06	.524	0.041	13
		07-08	*	*	6
	20-39	05-06	.517	0.083	6
		07-08	.580	0.117	6
	40-59	05-06	1.31	0.33	5
		07-08	1.52	0.31	6
	60+	05-06	3.46	0.76	5
		07-08	3.00	0.32	8
Non-Hispanic Black Female	12-19	05-06	.431	0.026	14
		07-08	*	*	5
	20-39	05-06	.728	0.102	7
		07-08	.698	0.129	8
	40-59	05-06	2.19	0.26	7
		07-08	2.43	0.42	8
	60+	05-06	*	*	5
		07-08	7.46	1.29	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 105) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.341	0.032	11
		07-08	*	*	6
	20-39	05-06	.484	0.031	9
		07-08	.745	0.085	9
	40-59	05-06	.835	0.076	4
		07-08	1.45	0.34	6
60+	05-06	1.34	0.12	4	
	07-08	1.78	0.2	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.421	0.043	9
		07-08	.669	0.027	8
	40-59	05-06	1.32	0.14	6
		07-08	1.33	0.16	6
60+	05-06	2.81	0.51	3	
	07-08	2.89	0.44	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 105) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1915
	01-02	*	< LOD	< LOD	< LOD	< LOD	2307
	03-04	.007 (.007-.008)	.007 (.006-.007)	.012 (.011-.014)	.027 (.023-.031)	.043 (.037-.049)	1879
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	665
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.003 (.003-.004)	.003 (.003-.004)	.005 (.005-.006)	.008 (.006-.010)	.011 (.008-.020)	593
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1250
	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	.008 (.007-.009)	.007 (.006-.008)	.014 (.012-.015)	.030 (.026-.035)	.047 (.039-.053)	1286
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	913
	01-02	*	< LOD	< LOD	< LOD	< LOD	1075
	03-04	.006 (.006-.007)	.006 (.005-.007)	.010 (.009-.011)	.019 (.016-.023)	.030 (.027-.039)	937
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1002
	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	.008 (.007-.010)	.007 (.006-.009)	.015 (.013-.017)	.033 (.029-.040)	.049 (.043-.067)	942
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	635
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	.004 (.004-.005)	.004 (.004-.005)	.007 (.006-.009)	.014 (.011-.017)	.019 (.015-.023)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	.090 (<LOD-.110)	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.008 (.007-.010)	.007 (.005-.008)	.015 (.010-.020)	.041 (.027-.067)	.082 (.041-.116)	456
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	714
	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	.007 (.007-.008)	.007 (.006-.007)	.012 (.011-.014)	.027 (.022-.031)	.043 (.035-.049)	878

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 105) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.003	0.001	9
		07-08	*	*	6
	20-39	05-06	.005	0.001	12
		07-08	.005	0	15
	40-59	05-06	*	*	12
		07-08	.007	0.001	16
	60+	05-06	.014	0.001	15
		07-08	.019	0.002	23
Non-Hispanic White Female	12-19	05-06	.003	0	10
		07-08	.003	0	7
	20-39	05-06	.004	0	16
		07-08	.006	0.001	13
	40-59	05-06	.010	0.001	13
		07-08	.009	0.001	17
	60+	05-06	.028	0.003	17
		07-08	.027	0.005	21
Non-Hispanic Black Male	12-19	05-06	.003	0	13
		07-08	*	*	6
	20-39	05-06	.003	0.001	6
		07-08	.003	0.001	6
	40-59	05-06	.008	0.002	5
		07-08	.010	0.002	6
	60+	05-06	.020	0.005	5
		07-08	.018	0.002	8
Non-Hispanic Black Female	12-19	05-06	.002	0	14
		07-08	*	*	5
	20-39	05-06	.004	0.001	7
		07-08	.004	0.001	8
	40-59	05-06	.014	0.002	7
		07-08	.015	0.003	8
	60+	05-06	*	*	5
		07-08	.044	0.006	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 105) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.002	0	11
		07-08	*	*	6
	20-39	05-06	.003	0	9
		07-08	.005	0.001	9
	40-59	05-06	.007	0.001	4
		07-08	.010	0.003	6
60+	05-06	.010	0.001	4	
	07-08	.013	0.002	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.003	0	9
		07-08	.004	0	8
	40-59	05-06	.010	0.001	6
		07-08	.009	0.001	6
	60+	05-06	.021	0.003	3
		07-08	.020	0.003	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 114) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	.597	0.052	12
		07-08	.686	0.062	16
	60+	05-06	1.22	0.09	15
		07-08	1.60	0.16	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	.723	0.058	13
		07-08	.798	0.095	17
	60+	05-06	1.62	0.13	17
		07-08	1.99	0.18	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	.458	0.059	5
		07-08	.572	0.129	6
	60+	05-06	1.52	0.18	5
		07-08	1.48	0.17	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	.712	0.092	7
		07-08	*	*	8
	60+	05-06	2.83	0.57	5
		07-08	2.32	0.21	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 114) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	.689	0.113	4	
	07-08	.934	0.076	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.361	0.059	6
		07-08	.450	0.125	6
	60+	05-06	.869	0.068	3
		07-08	.992	0.121	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 114) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	.005	0	12
		07-08	.005	0	16
	60+	05-06	.008	0.001	15
		07-08	.010	0.001	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	.005	0.001	13
		07-08	.005	0.001	17
	60+	05-06	.012	0.001	17
		07-08	.013	0.001	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	.003	0	5
		07-08	.004	0.001	6
	60+	05-06	.009	0.001	5
		07-08	.009	0.001	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	.005	0.001	7
		07-08	*	*	8
	60+	05-06	.019	0.004	5
		07-08	.014	0.001	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4'-Pentachlorobiphenyl (PCB 114) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	.005	0.001	4	
	07-08	.007	0	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.003	0	6
		07-08	.003	0.001	6
	60+	05-06	.007	0	3
		07-08	.007	0.001	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5-Pentachlorobiphenyl (PCB 118) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	13.1 (<LOD-15.1)	27.0 (22.1-30.3)	40.8 (32.8-48.6)	1926
	01-02	*	< LOD	15.1 (13.0-17.5)	29.0 (26.1-33.7)	44.6 (39.6-48.9)	2307
	03-04	6.00 (5.54-6.50)	5.19 (4.80-5.61)	10.4 (9.70-11.6)	21.8 (19.3-23.8)	31.3 (28.2-36.6)	1887
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	3.06 (2.76-3.39)	2.83 (2.70-3.10)	4.15 (3.60-4.56)	6.80 (4.70-9.60)	9.60 (6.80-13.2)	596
20 years and older	99-00	*	< LOD	14.9 (12.9-17.4)	28.1 (24.8-33.8)	43.8 (35.6-52.3)	1259
	01-02	*	< LOD	16.9 (15.1-19.0)	32.4 (28.3-37.3)	46.5 (41.1-51.0)	1549
	03-04	6.62 (6.10-7.18)	5.65 (5.21-6.40)	11.5 (10.2-12.6)	22.8 (20.7-25.9)	34.3 (29.6-39.6)	1291
Gender							
Males	99-00	*	< LOD	< LOD	19.6 (16.6-24.4)	28.0 (24.4-38.1)	919
	01-02	*	< LOD	12.0 (<LOD-14.9)	22.0 (17.6-27.1)	32.7 (25.4-47.5)	1075
	03-04	5.11 (4.70-5.57)	4.71 (4.16-5.06)	8.72 (7.30-9.70)	16.4 (14.0-18.9)	23.2 (20.7-25.1)	940
Females	99-00	*	< LOD	16.7 (13.6-19.7)	32.0 (27.1-40.3)	46.8 (41.2-57.8)	1007
	01-02	*	< LOD	18.7 (16.5-21.0)	36.4 (31.6-40.9)	48.9 (44.5-54.3)	1232
	03-04	6.99 (6.32-7.73)	6.02 (5.31-6.84)	12.3 (11.1-14.4)	27.3 (22.7-30.0)	37.8 (31.3-45.0)	947
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	17.3 (14.5-19.6)	23.7 (20.9-27.1)	636
	01-02	*	< LOD	< LOD	16.8 (12.5-21.7)	26.4 (19.3-33.2)	567
	03-04	3.56 (3.04-4.18)	3.15 (2.60-3.82)	5.02 (4.26-6.66)	9.37 (7.45-11.7)	14.0 (10.0-17.6)	426
Non-Hispanic blacks	99-00	*	< LOD	19.0 (13.6-21.7)	38.7 (28.4-54.4)	59.7 (45.7-80.8)	413
	01-02	*	< LOD	15.6 (12.6-19.9)	37.3 (26.1-46.4)	54.9 (40.0-66.6)	515
	03-04	6.70 (5.52-8.14)	5.25 (4.40-6.10)	12.7 (8.80-18.0)	35.1 (21.1-45.0)	57.8 (33.5-110)	462
Non-Hispanic whites	99-00	*	< LOD	13.1 (<LOD-15.6)	25.6 (20.8-32.0)	40.3 (30.4-46.8)	720
	01-02	*	< LOD	15.9 (13.6-18.1)	31.1 (27.1-35.9)	45.3 (38.7-50.1)	1061
	03-04	6.19 (5.57-6.87)	5.46 (4.85-6.20)	10.9 (9.62-12.1)	22.0 (19.3-24.4)	30.7 (26.2-36.6)	878

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.5, 10.5, and 0.6, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5-Pentachlorobiphenyl (PCB 118) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	2.24	0.31	9
		07-08	2.29	0.32	6
	20-39	05-06	3.30	0.67	12
		07-08	3.30	0.22	15
	40-59	05-06	7.56	1.56	12
		07-08	5.16	0.23	16
60+	05-06	13.3	1.4	15	
	07-08	15.5	1.6	23	
Non-Hispanic White Female	12-19	05-06	2.07	0.15	10
		07-08	2.41	0.23	7
	20-39	05-06	2.99	0.19	16
		07-08	4.18	0.58	13
	40-59	05-06	7.83	0.8	13
		07-08	7.56	0.73	17
	60+	05-06	21.9	2	17
		07-08	21.5	2.8	21
Non-Hispanic Black Male	12-19	05-06	2.47	0.24	13
		07-08	2.04	0.2	6
	20-39	05-06	2.62	0.49	6
		07-08	2.85	0.35	6
	40-59	05-06	7.30	1.8	5
		07-08	7.31	1.3	6
	60+	05-06	19.9	3.6	5
		07-08	16.1	1.7	8
Non-Hispanic Black Female	12-19	05-06	2.02	0.11	14
		07-08	1.92	0.24	5
	20-39	05-06	3.70	0.44	7
		07-08	3.37	0.3	8
	40-59	05-06	11.7	1.3	7
		07-08	11.2	1.9	8
	60+	05-06	45.0	13.5	5
		07-08	35.1	5	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.72 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5-Pentachlorobiphenyl (PCB 118) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	1.50	0.11	11
		07-08	1.52	0.18	6
	20-39	05-06	2.17	0.14	9
		07-08	2.66	0.22	9
	40-59	05-06	4.06	0.35	4
		07-08	6.03	0.93	6
60+	05-06	8.12	0.99	4	
	07-08	8.69	0.8	5	
Mexican American Female	12-19	05-06	1.36	0.08	16
		07-08	*	*	5
	20-39	05-06	1.94	0.2	9
		07-08	2.50	0.17	8
	40-59	05-06	6.79	0.67	6
		07-08	6.11	0.85	6
	60+	05-06	15.3	2.7	3
		07-08	14.0	2.5	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.72 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5-Pentachlorobiphenyl (PCB 118) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	.090 (<LOD-.100)	.180 (.150-.200)	.260 (.220-.320)	1926
	01-02	*	< LOD	.100 (.090-.110)	.190 (.160-.220)	.300 (.270-.320)	2307
	03-04	.037 (.034-.040)	.032 (.029-.036)	.066 (.061-.073)	.143 (.127-.160)	.216 (.192-.233)	1887
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.015 (.014-.017)	.014 (.013-.016)	.021 (.018-.023)	.036 (.024-.046)	.047 (.035-.076)	596
20 years and older	99-00	*	< LOD	.100 (.090-.110)	.190 (.170-.220)	.280 (.230-.360)	1259
	01-02	*	< LOD	.110 (.100-.130)	.210 (.180-.250)	.310 (.280-.370)	1549
	03-04	.041 (.038-.045)	.036 (.032-.041)	.072 (.066-.081)	.159 (.137-.174)	.226 (.202-.249)	1291
Gender							
Males	99-00	*	< LOD	< LOD	.130 (.110-.140)	.190 (.160-.220)	919
	01-02	*	< LOD	.080 (<LOD-.100)	.140 (.120-.170)	.210 (.160-.300)	1075
	03-04	.032 (.029-.035)	.030 (.026-.033)	.054 (.046-.062)	.110 (.083-.125)	.160 (.136-.180)	940
Females	99-00	*	< LOD	.110 (.090-.130)	.210 (.180-.260)	.320 (.260-.410)	1007
	01-02	*	< LOD	.130 (.100-.140)	.240 (.210-.260)	.360 (.300-.380)	1232
	03-04	.042 (.038-.047)	.036 (.030-.043)	.081 (.068-.097)	.176 (.152-.206)	.242 (.217-.290)	947
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	.110 (.090-.130)	.140 (.130-.180)	636
	01-02	*	< LOD	< LOD	.120 (.080-.160)	.180 (.130-.220)	567
	03-04	.022 (.018-.026)	.019 (.016-.024)	.034 (.029-.039)	.069 (.046-.086)	.094 (.075-.117)	426
Non-Hispanic blacks	99-00	*	< LOD	.120 (.080-.150)	.220 (.180-.320)	.400 (.260-.500)	413
	01-02	*	< LOD	.100 (.070-.120)	.210 (.170-.270)	.320 (.230-.460)	515
	03-04	.038 (.031-.046)	.030 (.026-.034)	.074 (.054-.106)	.218 (.133-.277)	.370 (.218-.618)	462
Non-Hispanic whites	99-00	*	< LOD	.090 (<LOD-.100)	.180 (.150-.200)	.250 (.200-.330)	720
	01-02	*	< LOD	.100 (.090-.120)	.210 (.170-.240)	.300 (.270-.360)	1061
	03-04	.038 (.034-.043)	.034 (.029-.040)	.067 (.062-.076)	.143 (.127-.165)	.209 (.178-.234)	878

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5-Pentachlorobiphenyl (PCB 118) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.012	0.002	9
		07-08	.011	0.002	6
	20-39	05-06	.025	0.006	12
		07-08	.022	0.002	15
	40-59	05-06	.058	0.013	12
		07-08	.038	0.002	16
60+	05-06	.090	0.008	15	
	07-08	.099	0.009	23	
Non-Hispanic White Female	12-19	05-06	.011	0.001	10
		07-08	.011	0.001	7
	20-39	05-06	.019	0.001	16
		07-08	.025	0.004	13
	40-59	05-06	.058	0.007	13
		07-08	.050	0.005	17
60+	05-06	.158	0.013	17	
	07-08	.145	0.019	21	
Non-Hispanic Black Male	12-19	05-06	.012	0.001	13
		07-08	.010	0.001	6
	20-39	05-06	.016	0.003	6
		07-08	.016	0.002	6
	40-59	05-06	.047	0.012	5
		07-08	.047	0.008	6
60+	05-06	.117	0.02	5	
	07-08	.098	0.011	8	
Non-Hispanic Black Female	12-19	05-06	.010	0.001	14
		07-08	.009	0.001	5
	20-39	05-06	.021	0.003	7
		07-08	.017	0.002	8
	40-59	05-06	.075	0.01	7
		07-08	.068	0.012	8
60+	05-06	.300	0.091	5	
	07-08	.208	0.024	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5-Pentachlorobiphenyl (PCB 118) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.008	0.001	11
		07-08	.008	0.001	6
	20-39	05-06	.015	0.001	9
		07-08	.018	0.002	9
	40-59	05-06	.033	0.003	4
		07-08	.043	0.007	6
60+	05-06	.059	0.007	4	
	07-08	.063	0.007	5	
Mexican American Female	12-19	05-06	.007	0	16
		07-08	*	*	5
	20-39	05-06	.013	0.001	9
		07-08	.015	0.001	8
	40-59	05-06	.050	0.005	6
		07-08	.041	0.006	6
	60+	05-06	.114	0.016	3
		07-08	.096	0.018	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2',3,4,4',5-Pentachlorobiphenyl (PCB 123) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
60+	05-06	*	*	15	
	07-08	*	*	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
60+	05-06	*	*	17	
	07-08	*	*	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
60+	05-06	*	*	5	
	07-08	*	*	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
60+	05-06	*	*	5	
	07-08	*	*	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2',3,4,4',5-Pentachlorobiphenyl (PCB 123) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2',3,4,4',5-Pentachlorobiphenyl (PCB 123) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2',3,4,4',5-Pentachlorobiphenyl (PCB 123) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5-Hexachlorobiphenyl (PCB 156) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	12.6 (<LOD-13.8)	17.0 (15.5-18.0)	1907
	01-02	*	< LOD	< LOD	14.3 (12.1-16.0)	18.3 (15.6-21.1)	2296
	03-04	2.54 (2.36-2.74)	3.29 (2.90-3.80)	7.00 (6.20-8.07)	11.4 (10.4-12.6)	15.3 (13.8-17.5)	1880
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	665
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	.500 (<LOD-.660)	1.20 (1.00-1.43)	2.30 (2.06-2.45)	3.00 (2.37-3.60)	586
20 years and older	99-00	*	< LOD	< LOD	13.6 (<LOD-14.7)	17.5 (16.0-20.1)	1242
	01-02	*	< LOD	< LOD	15.0 (13.3-17.0)	19.7 (17.0-22.1)	1540
	03-04	3.31 (3.05-3.60)	4.10 (3.40-4.66)	7.81 (6.80-8.70)	12.0 (10.9-13.5)	16.8 (14.7-18.6)	1294
Gender							
Males	99-00	*	< LOD	< LOD	12.6 (<LOD-14.0)	17.0 (14.6-18.0)	912
	01-02	*	< LOD	< LOD	13.7 (11.3-16.4)	18.4 (14.3-22.1)	1069
	03-04	2.57 (2.34-2.82)	3.10 (2.80-3.80)	6.92 (5.82-8.00)	11.8 (9.79-14.2)	17.1 (14.7-19.3)	940
Females	99-00	*	< LOD	< LOD	12.8 (<LOD-14.7)	16.5 (15.4-20.7)	995
	01-02	*	< LOD	< LOD	14.8 (13.2-15.5)	18.2 (16.2-21.0)	1227
	03-04	2.51 (2.28-2.76)	3.42 (2.80-4.10)	7.09 (6.35-8.10)	11.1 (10.4-12.0)	14.1 (12.7-16.8)	940
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	631
	01-02	*	< LOD	< LOD	< LOD	< LOD	566
	03-04	.858 (.640-1.15)	1.00 (.700-1.40)	2.48 (1.50-3.60)	4.78 (4.00-5.68)	6.40 (5.40-7.70)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	14.6 (<LOD-20.5)	23.1 (17.2-32.1)	412
	01-02	*	< LOD	< LOD	15.5 (12.7-21.0)	23.5 (16.4-31.4)	511
	03-04	2.32 (1.87-2.89)	3.04 (2.40-3.60)	7.40 (5.48-9.80)	14.7 (11.5-18.6)	24.8 (15.0-33.0)	456
Non-Hispanic whites	99-00	*	< LOD	< LOD	13.6 (<LOD-15.2)	17.4 (16.0-18.9)	711
	01-02	*	< LOD	< LOD	15.0 (13.2-17.5)	19.4 (16.2-22.1)	1056
	03-04	3.03 (2.70-3.39)	4.03 (3.15-4.66)	7.76 (6.51-8.79)	11.7 (10.5-13.1)	15.7 (13.5-18.5)	880

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.5, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5-Hexachlorobiphenyl (PCB 156) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.822	0.123	9
		07-08	.985	0.283	6
	20-39	05-06	1.61	0.26	12
		07-08	1.73	0.21	15
	40-59	05-06	5.40	0.36	12
		07-08	5.41	0.32	16
60+	05-06	8.93	0.48	15	
	07-08	11.9	1	23	
Non-Hispanic White Female	12-19	05-06	.614	0.064	10
		07-08	.751	0.136	7
	20-39	05-06	1.08	0.09	16
		07-08	1.50	0.16	13
	40-59	05-06	4.72	0.39	13
		07-08	5.33	0.33	17
60+	05-06	9.10	0.84	17	
	07-08	10.5	0.7	21	
Non-Hispanic Black Male	12-19	05-06	.659	0.077	13
		07-08	*	*	6
	20-39	05-06	1.35	0.23	6
		07-08	1.35	0.19	6
	40-59	05-06	4.39	0.48	5
		07-08	5.13	0.48	6
60+	05-06	13.5	1.9	5	
	07-08	12.5	1.4	8	
Non-Hispanic Black Female	12-19	05-06	.383	0.036	14
		07-08	*	*	5
	20-39	05-06	1.17	0.17	7
		07-08	1.03	0.15	8
	40-59	05-06	4.73	0.62	7
		07-08	5.20	0.37	8
60+	05-06	17.1	3.9	5	
	07-08	13.0	0.7	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5-Hexachlorobiphenyl (PCB 156) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.435	0.073	11
		07-08	.436	0.091	6
	20-39	05-06	.809	0.117	9
		07-08	.914	0.147	9
	40-59	05-06	2.48	0.24	4
		07-08	4.17	0.83	6
60+	05-06	5.39	1.05	4	
	07-08	5.80	0.39	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.524	0.07	9
		07-08	.700	0.062	8
	40-59	05-06	2.45	0.26	6
		07-08	3.02	0.47	6
	60+	05-06	4.81	0.26	3
		07-08	5.07	0.65	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5-Hexachlorobiphenyl (PCB 156) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	.090 (<LOD-.090)	.110 (.100-.130)	1907
	01-02	*	< LOD	< LOD	.100 (.080-.110)	.130 (.110-.140)	2296
	03-04	.016 (.014-.017)	.021 (.018-.024)	.048 (.041-.052)	.075 (.069-.085)	.103 (.094-.112)	1880
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	665
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	.003 (<LOD-.003)	.006 (.005-.007)	.012 (.011-.012)	.015 (.012-.018)	586
20 years and older	99-00	*	< LOD	< LOD	.090 (<LOD-.100)	.120 (.100-.140)	1242
	01-02	*	< LOD	< LOD	.100 (.090-.120)	.130 (.120-.140)	1540
	03-04	.021 (.019-.023)	.026 (.021-.030)	.051 (.045-.057)	.082 (.072-.093)	.109 (.098-.121)	1294
Gender							
Males	99-00	*	< LOD	< LOD	.090 (<LOD-.100)	.110 (.090-.130)	912
	01-02	*	< LOD	< LOD	.090 (.080-.120)	.120 (.110-.150)	1069
	03-04	.016 (.014-.018)	.021 (.017-.025)	.045 (.037-.054)	.076 (.068-.094)	.108 (.094-.123)	940
Females	99-00	*	< LOD	< LOD	.090 (<LOD-.090)	.110 (.090-.140)	995
	01-02	*	< LOD	< LOD	.100 (.090-.100)	.130 (.110-.140)	1227
	03-04	.015 (.014-.017)	.020 (.017-.025)	.049 (.044-.052)	.073 (.064-.085)	.101 (.089-.112)	940
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	631
	01-02	*	< LOD	< LOD	< LOD	< LOD	566
	03-04	.005 (.004-.007)	.007 (.004-.009)	.015 (.011-.022)	.036 (.025-.041)	.046 (.037-.070)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	.090 (<LOD-.120)	.150 (.110-.190)	412
	01-02	*	< LOD	< LOD	.100 (.080-.130)	.150 (.100-.180)	511
	03-04	.013 (.010-.017)	.018 (.014-.022)	.044 (.032-.056)	.099 (.067-.127)	.166 (.108-.198)	456
Non-Hispanic whites	99-00	*	< LOD	< LOD	.090 (<LOD-.100)	.110 (.100-.130)	711
	01-02	*	< LOD	< LOD	.100 (.090-.120)	.130 (.120-.140)	1056
	03-04	.019 (.017-.021)	.025 (.020-.031)	.051 (.044-.057)	.078 (.069-.089)	.104 (.094-.119)	880

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5-Hexachlorobiphenyl (PCB 156) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.005	0.001	9
		07-08	.005	0.001	6
	20-39	05-06	.012	0.002	12
		07-08	.012	0.002	15
	40-59	05-06	.041	0.003	12
		07-08	.039	0.002	16
60+	05-06	.061	0.003	15	
	07-08	.075	0.006	23	
Non-Hispanic White Female	12-19	05-06	.003	0	10
		07-08	.004	0.001	7
	20-39	05-06	.007	0.001	16
		07-08	.009	0.001	13
	40-59	05-06	.035	0.004	13
		07-08	.035	0.002	17
60+	05-06	.066	0.005	17	
	07-08	.070	0.004	21	
Non-Hispanic Black Male	12-19	05-06	.003	0	13
		07-08	*	*	6
	20-39	05-06	.008	0.001	6
		07-08	.008	0.001	6
	40-59	05-06	.028	0.003	5
		07-08	.033	0.003	6
60+	05-06	.080	0.012	5	
	07-08	.076	0.009	8	
Non-Hispanic Black Female	12-19	05-06	.002	0	14
		07-08	*	*	5
	20-39	05-06	.007	0.001	7
		07-08	.005	0.001	8
	40-59	05-06	.030	0.004	7
		07-08	.031	0.003	8
60+	05-06	.114	0.026	5	
	07-08	.078	0.004	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5-Hexachlorobiphenyl (PCB 156) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.002	0	11
		07-08	.002	0.001	6
	20-39	05-06	.006	0.001	9
		07-08	.006	0.001	9
	40-59	05-06	.020	0.002	4
		07-08	.030	0.006	6
60+	05-06	.039	0.007	4	
	07-08	.042	0.003	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.004	0.001	9
		07-08	.004	0	8
	40-59	05-06	.018	0.002	6
		07-08	.020	0.003	6
	60+	05-06	.037	0.004	3
		07-08	.035	0.005	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1897
	01-02	*	< LOD	< LOD	< LOD	< LOD	2294
	03-04	.605 (.554-.661)	.800 (.700-.940)	1.73 (1.47-1.93)	2.80 (2.50-3.10)	3.80 (3.36-4.30)	1858
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	654
	01-02	*	< LOD	< LOD	< LOD	< LOD	755
	03-04	*	< LOD	< LOD	.690 (.540-.760)	.980 (.750-1.12)	580
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1243
	01-02	*	< LOD	< LOD	< LOD	< LOD	1539
	03-04	.743 (.675-.817)	.980 (.800-1.10)	1.88 (1.69-2.05)	2.98 (2.68-3.30)	3.97 (3.51-4.59)	1278
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	901
	01-02	*	< LOD	< LOD	< LOD	< LOD	1068
	03-04	.594 (.541-.653)	.750 (.700-.900)	1.65 (1.38-1.88)	2.90 (2.39-3.50)	4.00 (3.57-4.87)	928
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	996
	01-02	*	< LOD	< LOD	< LOD	< LOD	1226
	03-04	.615 (.552-.685)	.890 (.700-1.00)	1.79 (1.60-1.99)	2.70 (2.50-2.91)	3.40 (3.20-3.93)	930
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	622
	01-02	*	< LOD	< LOD	< LOD	< LOD	566
	03-04	*	< LOD	.600 (<LOD-.820)	1.12 (.960-1.32)	1.59 (1.20-1.80)	422
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	405
	01-02	*	< LOD	< LOD	< LOD	< LOD	510
	03-04	.568 (.488-.662)	.720 (.600-.930)	1.80 (1.39-2.26)	3.55 (3.10-4.40)	5.35 (3.60-8.80)	448
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	716
	01-02	*	< LOD	< LOD	< LOD	< LOD	1055
	03-04	.713 (.635-.801)	.980 (.760-1.12)	1.81 (1.58-2.06)	2.86 (2.54-3.27)	3.80 (3.27-4.73)	869

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.5, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.367	0.055	12
		07-08	*	*	15
	40-59	05-06	1.15	0.07	12
		07-08	1.31	0.09	16
	60+	05-06	2.03	0.11	15
		07-08	2.91	0.27	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	1.06	0.09	13
		07-08	1.38	0.09	17
	60+	05-06	2.15	0.21	17
		07-08	2.72	0.19	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	.988	0.12	5
		07-08	1.30	0.12	6
	60+	05-06	3.25	0.45	5
		07-08	3.26	0.39	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	1.13	0.16	7
		07-08	1.38	0.11	8
	60+	05-06	4.31	1.09	5
		07-08	3.41	0.19	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	.561	0.072	4
		07-08	.979	0.195	6
60+	05-06	1.13	0.2	4	
	07-08	1.30	0.07	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.567	0.057	6
		07-08	.756	0.101	6
	60+	05-06	1.08	0.08	3
		07-08	1.46	0.26	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1897
	01-02	*	< LOD	< LOD	< LOD	< LOD	2294
	03-04	.004 (.003-.004)	.005 (.004-.006)	.011 (.010-.013)	.018 (.016-.021)	.024 (.022-.027)	1858
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	654
	01-02	*	< LOD	< LOD	< LOD	< LOD	755
	03-04	*	< LOD	< LOD	.004 (.003-.004)	.005 (.004-.005)	580
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1243
	01-02	*	< LOD	< LOD	< LOD	< LOD	1539
	03-04	.005 (.004-.005)	.006 (.005-.007)	.012 (.011-.014)	.019 (.017-.022)	.026 (.023-.031)	1278
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	901
	01-02	*	< LOD	< LOD	< LOD	< LOD	1068
	03-04	.004 (.003-.004)	.005 (.004-.006)	.011 (.009-.012)	.018 (.016-.022)	.025 (.022-.027)	928
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	996
	01-02	*	< LOD	< LOD	< LOD	< LOD	1226
	03-04	.004 (.003-.004)	.005 (.004-.006)	.012 (.010-.013)	.017 (.016-.020)	.024 (.021-.027)	930
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	622
	01-02	*	< LOD	< LOD	< LOD	< LOD	566
	03-04	*	< LOD	.004 (<LOD-.006)	.008 (.006-.009)	.011 (.009-.014)	422
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	405
	01-02	*	< LOD	< LOD	< LOD	< LOD	510
	03-04	.003 (.003-.004)	.004 (.003-.006)	.011 (.008-.014)	.025 (.016-.031)	.040 (.026-.050)	448
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	716
	01-02	*	< LOD	< LOD	< LOD	< LOD	1055
	03-04	.004 (.004-.005)	.006 (.005-.008)	.012 (.010-.014)	.018 (.016-.021)	.025 (.022-.027)	869

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.003	0	12
		07-08	*	*	15
	40-59	05-06	.009	0.001	12
		07-08	.009	0.001	16
	60+	05-06	.014	0.001	15
		07-08	.018	0.002	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	.008	0.001	13
		07-08	.009	0.001	17
	60+	05-06	.016	0.001	17
		07-08	.018	0.001	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	.006	0.001	5
		07-08	.008	0.001	6
	60+	05-06	.019	0.003	5
		07-08	.020	0.003	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	.007	0.001	7
		07-08	.008	0.001	8
	60+	05-06	.029	0.007	5
		07-08	.020	0.001	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	.005	0	4
		07-08	.007	0.001	6
60+	05-06	.008	0.001	4	
	07-08	.009	0	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.004	0	6
		07-08	.005	0.001	6
	60+	05-06	.008	0	3
		07-08	.010	0.002	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1908
	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	.494 (.441-.553)	.700 (.560-.800)	1.60 (1.40-1.70)	2.99 (2.74-3.36)	4.10 (3.80-4.47)	1864
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	666
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	< LOD	.640 (.500-.770)	.870 (.700-1.10)	584
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1242
	01-02	*	< LOD	< LOD	< LOD	< LOD	1540
	03-04	.592 (.521-.673)	.860 (.700-1.00)	1.72 (1.55-2.03)	3.30 (2.90-3.60)	4.30 (3.90-4.81)	1280
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	908
	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	.423 (<LOD-.471)	.500 (.440-.600)	1.33 (1.16-1.60)	2.70 (2.25-3.50)	3.80 (3.50-4.70)	931
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1000
	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	.573 (.492-.669)	.880 (.660-1.06)	1.71 (1.51-2.10)	3.21 (2.82-3.62)	4.30 (3.80-4.70)	933
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	627
	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	*	< LOD	.600 (.400-.800)	1.20 (.960-1.44)	1.70 (1.30-1.94)	419
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	411
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.521 (.432-.628)	.560 (.460-.800)	1.95 (1.25-2.40)	4.30 (3.27-6.79)	7.90 (4.70-11.8)	451
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	715
	01-02	*	< LOD	< LOD	< LOD	< LOD	1056
	03-04	.538 (.458-.632)	.800 (.600-.990)	1.66 (1.42-1.90)	2.98 (2.70-3.45)	3.94 (3.62-4.47)	875

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.4, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.397	0.087	12
		07-08	*	*	15
	40-59	05-06	1.07	0.12	12
		07-08	.967	0.07	16
60+	05-06	2.30	0.16	15	
	07-08	2.73	0.23	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	.334	0.026	16
		07-08	*	*	13
	40-59	05-06	1.19	0.12	13
		07-08	1.30	0.1	17
60+	05-06	2.67	0.23	17	
	07-08	2.93	0.23	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.346	0.065	6
		07-08	*	*	6
	40-59	05-06	1.05	0.16	5
		07-08	1.28	0.2	6
60+	05-06	3.71	0.57	5	
	07-08	3.44	0.44	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.388	0.047	7
		07-08	*	*	8
	40-59	05-06	1.57	0.19	7
		07-08	1.60	0.21	8
60+	05-06	6.11	1.71	5	
	07-08	4.73	0.3	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	.589	0.034	4
		07-08	1.08	0.2	6
60+	05-06	1.35	0.26	4	
	07-08	1.55	0.18	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.893	0.088	6
		07-08	1.00	0.13	6
	60+	05-06	1.69	0.17	3
		07-08	1.92	0.25	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1908
	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	.003 (.003-.003)	.004 (.003-.005)	.010 (.009-.012)	.020 (.018-.022)	.026 (.024-.027)	1864
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	666
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	< LOD	.004 (.003-.004)	.005 (.004-.005)	584
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1242
	01-02	*	< LOD	< LOD	< LOD	< LOD	1540
	03-04	.004 (.003-.004)	.005 (.004-.006)	.012 (.010-.013)	.021 (.019-.023)	.027 (.025-.030)	1280
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	908
	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	.003 (<LOD-.003)	.003 (.003-.004)	.009 (.008-.010)	.018 (.015-.021)	.023 (.021-.026)	931
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1000
	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	.003 (.003-.004)	.005 (.004-.006)	.012 (.010-.013)	.022 (.019-.024)	.027 (.025-.031)	933
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	627
	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	*	< LOD	.004 (.003-.004)	.009 (.007-.011)	.013 (.010-.015)	419
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	411
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.003 (.002-.004)	.003 (.003-.005)	.012 (.008-.015)	.028 (.019-.042)	.051 (.030-.072)	451
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	715
	01-02	*	< LOD	< LOD	< LOD	< LOD	1056
	03-04	.003 (.003-.004)	.005 (.004-.006)	.011 (.009-.013)	.020 (.018-.022)	.025 (.023-.027)	875

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.003	0.001	12
		07-08	*	*	15
	40-59	05-06	.008	0.001	12
		07-08	.007	0	16
	60+	05-06	.016	0.001	15
		07-08	.017	0.001	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	.002	0	16
		07-08	*	*	13
	40-59	05-06	.009	0.001	13
		07-08	.009	0.001	17
	60+	05-06	.019	0.001	17
		07-08	.020	0.001	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.002	0	6
		07-08	*	*	6
	40-59	05-06	.007	0.001	5
		07-08	.008	0.001	6
	60+	05-06	.022	0.003	5
		07-08	.021	0.003	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.002	0	7
		07-08	*	*	8
	40-59	05-06	.010	0.001	7
		07-08	.010	0.001	8
	60+	05-06	.041	0.011	5
		07-08	.028	0.002	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	.005	0	4
		07-08	.008	0.001	6
60+	05-06	.010	0.002	4	
	07-08	.011	0.001	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.007	0.001	6
		07-08	.007	0.001	6
	60+	05-06	.013	0	3
		07-08	.013	0.002	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	*	< LOD	< LOD	.900 (.700-1.16)	1.47 (1.10-2.18)	1817
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	752
	03-04	*	< LOD	< LOD	< LOD	1.00 (<LOD-3.09)	570
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1546
	03-04	*	< LOD	< LOD	1.00 (.760-1.20)	1.50 (1.10-2.14)	1247
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1070
	03-04	*	< LOD	< LOD	1.00 (.800-1.20)	1.54 (1.20-2.23)	903
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1228
	03-04	*	< LOD	< LOD	.830 (.600-1.14)	1.39 (.890-2.18)	914
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	*	< LOD	< LOD	< LOD	.700 (<LOD-1.00)	406
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	514
	03-04	*	< LOD	< LOD	1.20 (.840-1.60)	1.85 (1.20-2.83)	444
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1057
	03-04	*	< LOD	< LOD	.900 (.700-1.26)	1.51 (1.04-2.19)	851

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 10.5 and 0.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	.474	0.075	12
		07-08	.525	0.041	16
60+	05-06	.799	0.047	15	
	07-08	1.04	0.06	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	.386	0.05	17
60+	05-06	.615	0.059	17	
	07-08	.856	0.056	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	.423	0.022	5
		07-08	.626	0.056	6
60+	05-06	1.11	0.05	5	
	07-08	1.42	0.18	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	.373	0.073	7
		07-08	.538	0.041	8
60+	05-06	1.24	0.24	5	
	07-08	1.21	0.1	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	.445	0.128	6
60+	05-06	.506	0.146	4	
	07-08	.729	0.068	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	.407	0.015	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	*	< LOD	< LOD	.006 (.005-.008)	.010 (.007-.014)	1817
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	752
	03-04	*	< LOD	< LOD	< LOD	.005 (<LOD-.017)	570
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1546
	03-04	*	< LOD	< LOD	.006 (.005-.008)	.010 (.007-.014)	1247
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1070
	03-04	*	< LOD	< LOD	.006 (.005-.009)	.011 (.008-.014)	903
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1228
	03-04	*	< LOD	< LOD	.005 (.004-.008)	.009 (.006-.014)	914
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	*	< LOD	< LOD	< LOD	.005 (<LOD-.007)	406
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	514
	03-04	*	< LOD	< LOD	.008 (.005-.010)	.012 (.007-.016)	444
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1057
	03-04	*	< LOD	< LOD	.006 (.005-.008)	.010 (.007-.014)	851

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	.004	0.001	12
		07-08	.004	0	16
	60+	05-06	.005	0	15
		07-08	.007	0	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	.003	0	17
	60+	05-06	.004	0	17
		07-08	.006	0	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	.003	0	5
		07-08	.004	0	6
	60+	05-06	.007	0	5
		07-08	.009	0.001	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	.002	0	7
		07-08	.003	0	8
	60+	05-06	.008	0.002	5
		07-08	.007	0.001	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	.003	0.001	6
60+	05-06	.004	0.001	4	
	07-08	.005	0	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	.003	0	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary:

http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_BiomonitoringSummary.html

Factsheet: http://www.cdc.gov/biomonitoring/DioxinLikeChemicals_FactSheet.html

Serum 2,4,4'-Trichlorobiphenyl (PCB 28) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1849
	03-04	4.90 (4.60-5.22)	4.96 (4.65-5.26)	6.79 (6.40-7.31)	9.39 (8.70-10.1)	11.3 (10.7-11.8)	1866
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	647
	03-04	5.02 (4.48-5.63)	4.88 (4.26-5.48)	7.20 (5.90-8.60)	10.2 (8.40-11.5)	11.7 (10.7-13.3)	590
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1202
	03-04	4.88 (4.61-5.17)	4.98 (4.67-5.26)	6.78 (6.40-7.25)	9.10 (8.70-9.87)	11.1 (10.6-11.8)	1276
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	886
	03-04	4.81 (4.47-5.16)	4.86 (4.45-5.20)	6.70 (6.07-7.50)	9.39 (8.34-10.4)	10.8 (10.5-11.4)	926
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	963
	03-04	4.99 (4.66-5.35)	5.07 (4.73-5.37)	6.90 (6.49-7.40)	9.37 (8.72-10.1)	11.6 (10.7-13.1)	940
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	618
	03-04	4.95 (4.55-5.38)	4.90 (4.40-5.70)	6.70 (6.07-7.90)	8.94 (8.17-9.70)	10.5 (9.02-14.7)	413
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	392
	03-04	4.95 (4.20-5.84)	5.03 (4.10-5.60)	7.20 (5.62-8.80)	10.2 (8.51-12.4)	13.1 (10.8-15.3)	459
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	687
	03-04	4.89 (4.58-5.22)	4.90 (4.61-5.26)	6.76 (6.33-7.32)	9.40 (8.70-10.3)	11.1 (10.6-11.8)	876

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00 and 03-04 are 32.4 and 1.7.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4'-Trichlorobiphenyl (PCB 28) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	.726	0.065	15
	40-59	05-06	*	*	12
		07-08	1.05	0.17	16
	60+	05-06	*	*	15
		07-08	1.73	0.15	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	1.36	0.38	7
	20-39	05-06	*	*	16
		07-08	.942	0.095	13
	40-59	05-06	*	*	13
		07-08	1.12	0.1	17
	60+	05-06	2.93	0.33	17
		07-08	2.64	0.32	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	.865	0.089	6
	60+	05-06	2.74	0.55	5
		07-08	1.40	0.19	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	.832	0.115	8
	40-59	05-06	*	*	7
		07-08	1.38	0.17	8
	60+	05-06	3.81	0.59	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 2.54 and 0.7.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4'-Trichlorobiphenyl (PCB 28) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	.715	0.091	9
	40-59	05-06	*	*	4
		07-08	.988	0.109	6
60+	05-06	*	*	4	
	07-08	1.26	0.15	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	1.13	0.24	8
	40-59	05-06	*	*	6
		07-08	1.04	0.06	6
60+	05-06	*	*	3	
	07-08	1.69	0.19	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 2.54 and 0.7.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4'-Trichlorobiphenyl (PCB 28) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1849
	03-04	.030 (.028-.032)	.030 (.028-.032)	.041 (.039-.043)	.057 (.055-.060)	.067 (.063-.071)	1866
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	647
	03-04	.025 (.023-.028)	.025 (.021-.028)	.035 (.030-.042)	.051 (.042-.061)	.061 (.051-.070)	590
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1202
	03-04	.031 (.029-.033)	.031 (.029-.032)	.042 (.040-.043)	.058 (.055-.061)	.067 (.063-.071)	1276
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	886
	03-04	.030 (.028-.032)	.030 (.027-.032)	.041 (.037-.044)	.056 (.051-.061)	.065 (.061-.070)	926
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	963
	03-04	.030 (.028-.032)	.030 (.028-.032)	.042 (.039-.044)	.058 (.055-.061)	.070 (.064-.075)	940
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	618
	03-04	.030 (.028-.033)	.031 (.027-.035)	.041 (.036-.044)	.053 (.048-.059)	.063 (.054-.070)	413
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	392
	03-04	.028 (.024-.033)	.029 (.024-.033)	.039 (.034-.047)	.059 (.047-.070)	.070 (.064-.082)	459
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	687
	03-04	.030 (.028-.032)	.030 (.028-.032)	.041 (.039-.043)	.058 (.055-.061)	.068 (.062-.075)	876

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4'-Trichlorobiphenyl (PCB 28) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	.005	0	15
	40-59	05-06	*	*	12
		07-08	.008	0.001	16
	60+	05-06	*	*	15
		07-08	.011	0.001	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	.006	0.002	7
	20-39	05-06	*	*	16
		07-08	.006	0.001	13
	40-59	05-06	*	*	13
		07-08	.007	0.001	17
	60+	05-06	.021	0.002	17
		07-08	.018	0.002	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	.005	0.001	6
	60+	05-06	.016	0.003	5
		07-08	.008	0.001	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	.004	0.001	8
	40-59	05-06	*	*	7
		07-08	.008	0.001	8
	60+	05-06	.025	0.004	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4'-Trichlorobiphenyl (PCB 28) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	.005	0.001	9
	40-59	05-06	*	*	4
		07-08	.007	0.001	6
60+	05-06	*	*	4	
	07-08	.009	0.001	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	.006	0.001	8
	40-59	05-06	*	*	6
		07-08	.007	0	6
	60+	05-06	*	*	3
		07-08	.012	0.001	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2'3,5'-Tetrachlorobiphenyl (PCB 44) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	2.06 (1.93-2.19)	2.05 (1.90-2.20)	3.03 (2.90-3.20)	4.40 (4.10-4.86)	5.70 (5.40-6.10)	1890
Age group							
12-19 years	03-04	2.45 (2.22-2.70)	2.44 (2.10-2.70)	3.68 (3.19-4.37)	5.78 (5.13-6.30)	7.99 (6.20-9.10)	597
20 years and older	03-04	2.00 (1.88-2.13)	2.00 (1.90-2.11)	2.97 (2.80-3.20)	4.27 (3.85-4.70)	5.44 (5.00-5.89)	1293
Gender							
Males	03-04	2.12 (2.01-2.25)	2.12 (2.00-2.30)	3.11 (2.97-3.31)	4.52 (4.10-5.20)	5.89 (5.38-6.80)	942
Females	03-04	1.99 (1.82-2.18)	1.98 (1.80-2.20)	2.94 (2.72-3.21)	4.30 (3.77-5.05)	5.60 (5.00-6.20)	948
Race/ethnicity							
Mexican Americans	03-04	2.09 (1.88-2.31)	2.10 (1.90-2.39)	3.10 (2.52-3.73)	4.20 (3.30-6.10)	5.69 (3.80-7.40)	427
Non-Hispanic blacks	03-04	2.21 (1.94-2.52)	2.12 (1.80-2.56)	3.18 (2.72-3.94)	4.90 (3.90-6.32)	6.32 (5.01-9.62)	464
Non-Hispanic whites	03-04	2.03 (1.87-2.19)	2.01 (1.90-2.20)	3.07 (2.90-3.29)	4.40 (4.01-4.95)	5.70 (5.25-6.35)	877

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 0.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2'3,5'-Tetrachlorobiphenyl (PCB 44) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
60+	05-06	*	*	15	
	07-08	*	*	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
60+	05-06	*	*	17	
	07-08	*	*	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
60+	05-06	*	*	5	
	07-08	*	*	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
60+	05-06	*	*	5	
	07-08	*	*	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.36 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,5'-Tetrachlorobiphenyl (PCB 44) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.36 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2'3,5'-Tetrachlorobiphenyl (PCB 44) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	.013 (.012-.013)	.013 (.012-.014)	.018 (.017-.020)	.026 (.024-.028)	.032 (.030-.034)	1890
Age group							
12-19 years	03-04	.012 (.011-.014)	.013 (.011-.014)	.019 (.016-.021)	.029 (.024-.031)	.037 (.030-.042)	597
20 years and older	03-04	.013 (.012-.013)	.013 (.012-.014)	.018 (.017-.020)	.025 (.023-.027)	.031 (.029-.034)	1293
Gender							
Males	03-04	.013 (.012-.014)	.013 (.012-.014)	.018 (.017-.020)	.026 (.024-.028)	.034 (.030-.038)	942
Females	03-04	.012 (.011-.013)	.012 (.011-.013)	.018 (.016-.020)	.025 (.023-.028)	.030 (.028-.032)	948
Race/ethnicity							
Mexican Americans	03-04	.013 (.011-.014)	.013 (.011-.014)	.017 (.015-.020)	.023 (.018-.031)	.027 (.021-.049)	427
Non-Hispanic blacks	03-04	.013 (.011-.014)	.012 (.010-.014)	.018 (.015-.022)	.026 (.021-.034)	.035 (.026-.040)	464
Non-Hispanic whites	03-04	.012 (.012-.014)	.012 (.011-.014)	.019 (.017-.020)	.026 (.024-.028)	.032 (.029-.035)	877

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2'3,5'-Tetrachlorobiphenyl (PCB 44) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2'3,5'-Tetrachlorobiphenyl (PCB 44) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5'-Tetrachlorobiphenyl (PCB 49) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	1.29 (1.20-1.39)	1.35 (1.24-1.45)	1.90 (1.80-2.10)	2.80 (2.60-3.08)	3.53 (3.33-3.80)	1876
Age group							
12-19 years	03-04	1.54 (1.37-1.72)	1.59 (1.33-1.80)	2.30 (2.00-2.84)	3.60 (3.10-4.13)	4.66 (3.80-5.73)	590
20 years and older	03-04	1.26 (1.17-1.35)	1.33 (1.22-1.40)	1.90 (1.78-2.03)	2.69 (2.50-2.80)	3.36 (3.15-3.63)	1286
Gender							
Males	03-04	1.36 (1.27-1.45)	1.40 (1.30-1.50)	2.00 (1.90-2.20)	3.00 (2.60-3.23)	3.79 (3.36-4.10)	932
Females	03-04	1.23 (1.12-1.35)	1.30 (1.13-1.40)	1.80 (1.70-2.04)	2.66 (2.38-3.01)	3.39 (3.02-3.73)	944
Race/ethnicity							
Mexican Americans	03-04	1.33 (1.20-1.48)	1.46 (1.20-1.57)	2.00 (1.67-2.30)	2.70 (2.17-3.80)	3.56 (2.63-4.20)	426
Non-Hispanic blacks	03-04	1.40 (1.23-1.59)	1.40 (1.20-1.64)	2.06 (1.70-2.80)	3.23 (2.80-3.70)	3.88 (3.33-5.53)	453
Non-Hispanic whites	03-04	1.26 (1.15-1.38)	1.32 (1.20-1.40)	1.90 (1.78-2.11)	2.77 (2.50-3.10)	3.45 (3.20-3.90)	876

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 0.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5'-Tetrachlorobiphenyl (PCB 49) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5'-Tetrachlorobiphenyl (PCB 49) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5'-Tetrachlorobiphenyl (PCB 49) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	.008 (.007-.008)	.008 (.007-.009)	.012 (.011-.012)	.016 (.015-.017)	.019 (.018-.021)	1876
Age group							
12-19 years	03-04	.008 (.007-.009)	.008 (.007-.009)	.012 (.010-.013)	.018 (.015-.019)	.022 (.018-.025)	590
20 years and older	03-04	.008 (.007-.008)	.008 (.007-.009)	.012 (.011-.013)	.015 (.015-.017)	.019 (.018-.021)	1286
Gender							
Males	03-04	.008 (.008-.009)	.008 (.008-.009)	.012 (.011-.013)	.017 (.015-.018)	.021 (.019-.022)	932
Females	03-04	.007 (.007-.008)	.008 (.007-.008)	.011 (.010-.012)	.015 (.014-.017)	.018 (.017-.019)	944
Race/ethnicity							
Mexican Americans	03-04	.008 (.007-.009)	.008 (.007-.009)	.011 (.010-.014)	.015 (.012-.018)	.018 (.015-.023)	426
Non-Hispanic blacks	03-04	.008 (.007-.009)	.008 (.007-.009)	.012 (.010-.014)	.017 (.015-.020)	.021 (.017-.026)	453
Non-Hispanic whites	03-04	.008 (.007-.009)	.008 (.007-.009)	.012 (.011-.013)	.016 (.015-.017)	.019 (.018-.021)	876

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5'-Tetrachlorobiphenyl (PCB 49) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5'-Tetrachlorobiphenyl (PCB 49) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',5,5'-Tetrachlorobiphenyl (PCB 52) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1912
	01-02	*	< LOD	< LOD	< LOD	16.5 (14.3-17.2)	1537
	03-04	2.66 (2.43-2.91)	2.74 (2.50-3.00)	4.17 (3.72-4.60)	5.91 (5.40-6.67)	7.60 (7.01-8.00)	1897
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	664
	01-02	*	< LOD	< LOD	16.2 (<LOD-23.8)	22.9 (16.9-32.3)	291
	03-04	3.16 (2.81-3.56)	3.22 (2.60-3.96)	5.15 (4.46-5.79)	7.60 (6.55-8.33)	9.20 (7.80-12.5)	597
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1248
	01-02	*	< LOD	< LOD	< LOD	16.0 (13.5-16.9)	1246
	03-04	2.59 (2.36-2.84)	2.70 (2.46-2.92)	4.06 (3.60-4.43)	5.70 (5.16-6.49)	7.15 (6.62-7.80)	1300
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	908
	01-02	*	< LOD	< LOD	< LOD	16.0 (<LOD-17.5)	716
	03-04	2.75 (2.54-2.98)	2.80 (2.55-3.20)	4.36 (3.90-4.88)	5.94 (5.51-6.80)	7.80 (7.29-8.49)	946
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1004
	01-02	*	< LOD	< LOD	< LOD	16.6 (14.0-18.2)	821
	03-04	2.57 (2.30-2.87)	2.70 (2.40-2.96)	3.96 (3.52-4.30)	5.86 (4.96-6.67)	7.15 (6.57-8.10)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	631
	01-02	*	< LOD	< LOD	< LOD	16.9 (<LOD-20.6)	366
	03-04	2.88 (2.56-3.24)	3.00 (2.70-3.30)	4.41 (3.60-5.30)	6.00 (4.71-8.11)	7.83 (5.70-12.0)	426
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	408
	01-02	*	< LOD	< LOD	< LOD	16.6 (<LOD-20.9)	282
	03-04	2.74 (2.35-3.19)	2.58 (2.08-3.40)	4.11 (3.30-5.60)	6.92 (5.10-8.70)	8.70 (7.10-10.2)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	716
	01-02	*	< LOD	< LOD	< LOD	16.6 (14.7-17.5)	773
	03-04	2.60 (2.35-2.88)	2.70 (2.40-2.98)	4.17 (3.70-4.60)	5.79 (5.20-6.75)	7.37 (6.80-7.87)	885

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.5, 12.4, and 0.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',5,5'-Tetrachlorobiphenyl (PCB 52) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	.348	0.071	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	.562	0.135	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	.807	0.149	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.33 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',5,5'-Tetrachlorobiphenyl (PCB 52) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.33 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',5,5'-Tetrachlorobiphenyl (PCB 52) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1912
	01-02	*	< LOD	< LOD	< LOD	.090 (.080-.100)	1537
	03-04	.016 (.015-.018)	.017 (.015-.019)	.024 (.022-.028)	.035 (.032-.037)	.043 (.039-.046)	1897
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	664
	01-02	*	< LOD	< LOD	.080 (<LOD-.100)	.100 (.080-.140)	291
	03-04	.016 (.014-.018)	.017 (.014-.019)	.026 (.022-.029)	.037 (.032-.041)	.042 (.037-.056)	597
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1248
	01-02	*	< LOD	< LOD	< LOD	.090 (.080-.090)	1246
	03-04	.016 (.015-.018)	.017 (.015-.019)	.024 (.022-.028)	.034 (.031-.036)	.043 (.038-.045)	1300
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	908
	01-02	*	< LOD	< LOD	< LOD	.090 (<LOD-.110)	716
	03-04	.017 (.016-.018)	.018 (.016-.020)	.025 (.023-.029)	.036 (.032-.039)	.044 (.040-.048)	946
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1004
	01-02	*	< LOD	< LOD	< LOD	.090 (.080-.100)	821
	03-04	.016 (.014-.017)	.016 (.014-.018)	.024 (.021-.028)	.033 (.030-.036)	.042 (.036-.046)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	631
	01-02	*	< LOD	< LOD	< LOD	.090 (<LOD-.110)	366
	03-04	.018 (.016-.020)	.018 (.016-.021)	.025 (.022-.030)	.035 (.028-.044)	.044 (.032-.064)	426
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	408
	01-02	*	< LOD	< LOD	< LOD	.090 (<LOD-.110)	282
	03-04	.016 (.014-.018)	.015 (.012-.019)	.022 (.019-.031)	.035 (.029-.043)	.043 (.035-.049)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	716
	01-02	*	< LOD	< LOD	< LOD	.090 (.080-.100)	773
	03-04	.016 (.014-.018)	.017 (.014-.019)	.024 (.022-.028)	.035 (.032-.037)	.043 (.038-.047)	885

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',5,5'-Tetrachlorobiphenyl (PCB 52) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	.003	0	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	.003	0.001	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	.005	0.001	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',5,5'-Tetrachlorobiphenyl (PCB 52) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†]
(in ng/g of serum or parts per billion) for the U.S. population from
the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3',4,4'-Tetrachlorobiphenyl (PCB 66) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1931
	01-02	*	< LOD	< LOD	< LOD	< LOD	2250
	03-04	1.39 (1.32-1.47)	1.37 (1.30-1.40)	1.97 (1.90-2.10)	3.10 (2.94-3.26)	4.10 (3.90-4.55)	1898
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	671
	01-02	*	< LOD	< LOD	< LOD	< LOD	724
	03-04	1.24 (1.14-1.34)	1.20 (1.10-1.26)	1.76 (1.62-2.00)	2.40 (2.30-2.70)	3.25 (2.70-4.10)	598
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1260
	01-02	*	< LOD	< LOD	< LOD	< LOD	1526
	03-04	1.42 (1.34-1.49)	1.40 (1.30-1.43)	2.00 (1.90-2.12)	3.10 (2.97-3.40)	4.20 (3.90-4.71)	1300
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	919
	01-02	*	< LOD	< LOD	< LOD	< LOD	1047
	03-04	1.29 (1.20-1.39)	1.30 (1.20-1.38)	1.87 (1.72-2.01)	2.70 (2.50-3.00)	3.30 (2.96-3.82)	947
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1012
	01-02	*	< LOD	< LOD	< LOD	< LOD	1203
	03-04	1.50 (1.42-1.58)	1.41 (1.38-1.50)	2.10 (1.92-2.30)	3.70 (3.20-4.00)	5.08 (4.10-5.46)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	636
	01-02	*	< LOD	< LOD	< LOD	< LOD	548
	03-04	1.19 (1.09-1.30)	1.14 (1.00-1.30)	1.60 (1.48-1.73)	2.23 (1.80-3.20)	3.20 (2.46-3.60)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	414
	01-02	*	< LOD	< LOD	< LOD	< LOD	495
	03-04	1.50 (1.34-1.68)	1.38 (1.21-1.50)	2.31 (1.80-2.80)	3.80 (3.10-4.80)	5.46 (4.30-8.60)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	723
	01-02	*	< LOD	< LOD	< LOD	< LOD	1047
	03-04	1.39 (1.31-1.47)	1.39 (1.30-1.45)	1.96 (1.90-2.10)	3.01 (2.71-3.28)	4.09 (3.70-4.33)	885

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.4, 12.4, and 0.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3',4,4'-Tetrachlorobiphenyl (PCB 66) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.415	0.061	12
		07-08	.389	0.063	15
	40-59	05-06	.827	0.138	12
		07-08	.657	0.066	16
	60+	05-06	*	*	15
		07-08	1.27	0.18	23
Non-Hispanic White Female	12-19	05-06	.421	0.04	10
		07-08	.605	0.146	7
	20-39	05-06	.471	0.054	16
		07-08	.629	0.157	13
	40-59	05-06	.671	0.091	13
		07-08	.750	0.066	17
	60+	05-06	1.91	0.21	17
		07-08	1.84	0.21	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	.784	0.154	5
		07-08	.645	0.155	6
	60+	05-06	1.49	0.23	5
		07-08	1.29	0.18	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.677	0.112	7
		07-08	*	*	8
	40-59	05-06	1.09	0.1	7
		07-08	1.13	0.21	8
	60+	05-06	3.40	0.95	5
		07-08	2.49	0.63	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.4 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3',4,4'-Tetrachlorobiphenyl (PCB 66) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.459	0.074	9
		07-08	.414	0.088	9
	40-59	05-06	.585	0.103	4
		07-08	*	*	6
	60+	05-06	.702	0.057	4
		07-08	.869	0.051	5
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	.597	0.07	8
	40-59	05-06	.751	0.095	6
		07-08	.684	0.073	6
	60+	05-06	1.32	0.36	3
		07-08	1.18	0.14	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.4 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3',4,4'-Tetrachlorobiphenyl (PCB 66) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1931
	01-02	*	< LOD	< LOD	< LOD	< LOD	2250
	03-04	.008 (.008-.009)	.008 (.008-.009)	.012 (.012-.013)	.019 (.018-.021)	.025 (.024-.030)	1898
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	671
	01-02	*	< LOD	< LOD	< LOD	< LOD	724
	03-04	.006 (.006-.007)	.006 (.005-.007)	.009 (.008-.010)	.013 (.012-.014)	.017 (.013-.019)	598
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1260
	01-02	*	< LOD	< LOD	< LOD	< LOD	1526
	03-04	.009 (.008-.009)	.008 (.008-.009)	.013 (.012-.013)	.020 (.019-.022)	.026 (.024-.032)	1300
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	919
	01-02	*	< LOD	< LOD	< LOD	< LOD	1047
	03-04	.008 (.007-.009)	.008 (.007-.008)	.011 (.010-.013)	.017 (.016-.019)	.020 (.019-.022)	947
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1012
	01-02	*	< LOD	< LOD	< LOD	< LOD	1203
	03-04	.009 (.009-.010)	.009 (.008-.009)	.013 (.012-.014)	.022 (.021-.024)	.032 (.026-.034)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	636
	01-02	*	< LOD	< LOD	< LOD	< LOD	548
	03-04	.007 (.007-.008)	.007 (.006-.008)	.010 (.009-.011)	.015 (.011-.018)	.018 (.014-.020)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	414
	01-02	*	< LOD	< LOD	< LOD	< LOD	495
	03-04	.009 (.008-.009)	.008 (.007-.009)	.013 (.011-.015)	.022 (.019-.030)	.036 (.024-.070)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	723
	01-02	*	< LOD	< LOD	< LOD	< LOD	1047
	03-04	.009 (.008-.009)	.008 (.008-.009)	.012 (.012-.013)	.019 (.018-.022)	.025 (.023-.030)	885

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3',4,4'-Tetrachlorobiphenyl (PCB 66) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.003	0.001	12
		07-08	.003	0	15
	40-59	05-06	.006	0.001	12
		07-08	.005	0.001	16
60+	05-06	*	*	15	
	07-08	.008	0.001	23	
Non-Hispanic White Female	12-19	05-06	.002	0	10
		07-08	.003	0.001	7
	20-39	05-06	.003	0	16
		07-08	.004	0.001	13
	40-59	05-06	.005	0.001	13
		07-08	.005	0	17
60+	05-06	.014	0.001	17	
	07-08	.012	0.001	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	.005	0.001	5
		07-08	.004	0.001	6
60+	05-06	.009	0.001	5	
	07-08	.008	0.001	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.004	0.001	7
		07-08	*	*	8
	40-59	05-06	.007	0.001	7
		07-08	.007	0.001	8
60+	05-06	.023	0.006	5	
	07-08	.015	0.004	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3',4,4'-Tetrachlorobiphenyl (PCB 66) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.003	0.001	9
		07-08	.003	0.001	9
	40-59	05-06	.005	0.001	4
		07-08	*	*	6
	60+	05-06	.005	0.001	4
		07-08	.006	0	5
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	.003	0	8
	40-59	05-06	.006	0.001	6
		07-08	.005	0	6
	60+	05-06	.010	0.002	3
		07-08	.008	0.001	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4',5-Tetrachlorobiphenyl (PCB 74) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	20.8 (17.9-23.4)	29.0 (24.5-32.3)	1924
	01-02	*	< LOD	13.2 (11.0-15.7)	23.5 (20.2-27.7)	33.0 (26.9-38.7)	2307
	03-04	4.81 (4.63-4.99)	4.36 (3.90-4.88)	8.72 (8.30-9.28)	15.8 (14.7-17.7)	22.3 (19.7-25.5)	1898
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	671
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	2.20 (2.02-2.39)	2.20 (2.03-2.33)	2.98 (2.80-3.30)	4.19 (3.60-4.73)	5.32 (4.20-6.90)	598
20 years and older	99-00	*	< LOD	12.6 (<LOD-14.4)	22.4 (19.5-25.6)	30.0 (26.2-35.8)	1253
	01-02	*	< LOD	14.7 (12.4-17.1)	25.2 (21.4-28.9)	34.8 (28.8-41.4)	1549
	03-04	5.38 (5.16-5.62)	5.00 (4.60-5.50)	9.60 (8.90-10.6)	17.1 (15.7-18.5)	24.1 (20.8-27.7)	1300
Gender							
Males	99-00	*	< LOD	< LOD	15.4 (13.0-17.8)	21.6 (18.3-24.4)	915
	01-02	*	< LOD	10.6 (<LOD-12.3)	20.2 (15.4-24.9)	28.8 (20.8-38.2)	1075
	03-04	4.06 (3.82-4.31)	3.62 (3.30-4.00)	6.71 (6.20-7.77)	12.4 (10.8-13.1)	15.8 (14.6-18.5)	947
Females	99-00	*	< LOD	13.9 (<LOD-16.4)	24.6 (21.9-28.5)	31.9 (28.8-40.3)	1009
	01-02	*	< LOD	15.8 (13.5-18.7)	26.1 (23.2-29.3)	35.9 (31.4-41.4)	1232
	03-04	5.65 (5.33-5.98)	5.38 (4.84-5.95)	10.9 (9.30-12.1)	19.1 (17.4-21.1)	27.4 (22.9-29.6)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	15.8 (12.8-18.2)	636
	01-02	*	< LOD	< LOD	13.3 (<LOD-18.1)	19.6 (16.2-23.0)	567
	03-04	2.43 (2.10-2.80)	2.12 (1.90-2.40)	3.34 (2.90-4.16)	6.10 (5.25-7.40)	10.4 (6.61-12.2)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	29.0 (18.6-38.4)	43.8 (35.4-64.0)	411
	01-02	*	< LOD	12.0 (<LOD-14.2)	21.7 (16.4-26.9)	31.9 (24.5-41.0)	515
	03-04	4.96 (4.21-5.83)	4.00 (3.45-4.65)	9.17 (6.80-12.4)	23.0 (14.6-35.1)	40.2 (23.0-68.2)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	21.6 (18.9-24.0)	29.0 (24.3-32.3)	719
	01-02	*	< LOD	14.7 (12.2-17.3)	25.8 (21.3-30.1)	35.9 (29.0-42.5)	1061
	03-04	5.23 (4.98-5.50)	4.91 (4.38-5.65)	9.30 (8.60-10.4)	16.1 (15.1-18.2)	21.9 (19.2-25.2)	885

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.4, 10.5, and 0.8, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4',5-Tetrachlorobiphenyl (PCB 74) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	1.84	0.15	9
		07-08	1.75	0.27	6
	20-39	05-06	2.63	0.45	12
		07-08	2.22	0.08	15
	40-59	05-06	5.30	0.32	12
		07-08	4.09	0.24	16
60+	05-06	11.7	1.1	15	
	07-08	10.5	0.9	23	
Non-Hispanic White Female	12-19	05-06	1.82	0.15	10
		07-08	1.65	0.11	7
	20-39	05-06	2.64	0.17	16
		07-08	2.45	0.3	13
	40-59	05-06	7.21	0.59	13
		07-08	6.25	0.52	17
60+	05-06	17.8	1.7	17	
	07-08	15.8	1.1	21	
Non-Hispanic Black Male	12-19	05-06	2.03	0.16	13
		07-08	1.37	0.2	6
	20-39	05-06	2.30	0.31	6
		07-08	1.82	0.17	6
	40-59	05-06	4.49	0.6	5
		07-08	4.41	0.74	6
60+	05-06	16.5	2.4	5	
	07-08	11.5	1.4	8	
Non-Hispanic Black Female	12-19	05-06	1.34	0.17	14
		07-08	1.03	0.15	5
	20-39	05-06	2.89	0.34	7
		07-08	1.83	0.16	8
	40-59	05-06	8.74	1.02	7
		07-08	8.01	1.27	8
60+	05-06	36.0	9.3	5	
	07-08	21.5	2.1	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.8.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4',5-Tetrachlorobiphenyl (PCB 74) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	1.22	0.17	11
		07-08	.992	0.124	6
	20-39	05-06	1.57	0.1	9
		07-08	1.55	0.14	9
	40-59	05-06	2.94	0.34	4
		07-08	3.13	0.29	6
60+	05-06	6.38	0.49	4	
	07-08	6.54	0.59	5	
Mexican American Female	12-19	05-06	1.08	0.12	16
		07-08	*	*	5
	20-39	05-06	1.46	0.17	9
		07-08	1.60	0.14	8
	40-59	05-06	4.28	0.43	6
		07-08	4.65	0.82	6
60+	05-06	11.9	2.1	3	
	07-08	9.71	2.27	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.8.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4',5-Tetrachlorobiphenyl (PCB 74) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	.140 (.120-.150)	.180 (.160-.220)	1924
	01-02	*	< LOD	.090 (.070-.100)	.150 (.140-.180)	.210 (.180-.270)	2307
	03-04	.029 (.028-.031)	.027 (.024-.029)	.058 (.053-.062)	.104 (.093-.121)	.153 (.135-.171)	1898
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	671
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.011 (.010-.012)	.011 (.010-.012)	.015 (.014-.018)	.021 (.019-.025)	.026 (.021-.033)	598
20 years and older	99-00	*	< LOD	.080 (<LOD-.100)	.150 (.140-.160)	.200 (.170-.230)	1253
	01-02	*	< LOD	.100 (.080-.110)	.170 (.140-.190)	.230 (.200-.280)	1549
	03-04	.034 (.032-.035)	.031 (.028-.033)	.064 (.059-.068)	.115 (.103-.128)	.167 (.143-.186)	1300
Gender							
Males	99-00	*	< LOD	< LOD	.100 (.090-.120)	.150 (.130-.170)	915
	01-02	*	< LOD	.070 (<LOD-.080)	.130 (.100-.170)	.190 (.140-.250)	1075
	03-04	.025 (.023-.027)	.023 (.021-.027)	.043 (.040-.050)	.077 (.067-.087)	.103 (.092-.118)	947
Females	99-00	*	< LOD	.100 (<LOD-.110)	.160 (.140-.180)	.220 (.190-.250)	1009
	01-02	*	< LOD	.100 (.090-.110)	.170 (.150-.190)	.240 (.210-.290)	1232
	03-04	.034 (.032-.037)	.031 (.027-.036)	.071 (.062-.085)	.130 (.120-.143)	.186 (.154-.216)	951
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	.110 (.080-.130)	636
	01-02	*	< LOD	< LOD	.090 (<LOD-.120)	.140 (.110-.160)	567
	03-04	.015 (.013-.017)	.013 (.011-.016)	.022 (.019-.026)	.042 (.032-.054)	.071 (.049-.097)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	.170 (.110-.250)	.280 (.220-.420)	411
	01-02	*	< LOD	.070 (<LOD-.080)	.130 (.100-.170)	.200 (.150-.250)	515
	03-04	.028 (.024-.033)	.022 (.019-.025)	.055 (.038-.077)	.153 (.090-.215)	.236 (.143-.337)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	.150 (.130-.160)	.190 (.160-.210)	719
	01-02	*	< LOD	.100 (.080-.110)	.170 (.140-.200)	.230 (.190-.280)	1061
	03-04	.032 (.030-.034)	.030 (.027-.033)	.064 (.058-.067)	.113 (.096-.123)	.153 (.131-.176)	885

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4',5-Tetrachlorobiphenyl (PCB 74) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.010	0.001	9
		07-08	.009	0.001	6
	20-39	05-06	.020	0.004	12
		07-08	.015	0.001	15
	40-59	05-06	.040	0.003	12
		07-08	.030	0.002	16
60+	05-06	.079	0.007	15	
	07-08	.066	0.005	23	
Non-Hispanic White Female	12-19	05-06	.010	0.001	10
		07-08	.008	0	7
	20-39	05-06	.017	0.001	16
		07-08	.015	0.002	13
	40-59	05-06	.054	0.006	13
		07-08	.041	0.004	17
60+	05-06	.129	0.011	17	
	07-08	.106	0.007	21	
Non-Hispanic Black Male	12-19	05-06	.010	0.001	13
		07-08	.006	0.001	6
	20-39	05-06	.014	0.002	6
		07-08	.010	0.001	6
	40-59	05-06	.029	0.004	5
		07-08	.028	0.005	6
60+	05-06	.098	0.013	5	
	07-08	.070	0.009	8	
Non-Hispanic Black Female	12-19	05-06	.007	0.001	14
		07-08	.005	0.001	5
	20-39	05-06	.017	0.002	7
		07-08	.009	0.001	8
	40-59	05-06	.057	0.008	7
		07-08	.049	0.008	8
60+	05-06	.239	0.062	5	
	07-08	.128	0.01	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,4,4',5-Tetrachlorobiphenyl (PCB 74) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.007	0.001	11
		07-08	.005	0.001	6
	20-39	05-06	.011	0.001	9
		07-08	.011	0.001	9
	40-59	05-06	.024	0.003	4
		07-08	.022	0.002	6
60+	05-06	.046	0.005	4	
	07-08	.046	0.002	5	
Mexican American Female	12-19	05-06	.006	0.001	16
		07-08	*	*	5
	20-39	05-06	.010	0.001	9
		07-08	.009	0.001	8
	40-59	05-06	.031	0.003	6
		07-08	.031	0.006	6
	60+	05-06	.088	0.008	3
		07-08	.067	0.016	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,5'-Pentachlorobiphenyl (PCB 87) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	.656 (.579-.744)	.900 (.800-.980)	1.32 (1.24-1.46)	2.02 (1.90-2.17)	2.70 (2.40-3.03)	1892
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.706 (.629-.792)	.860 (.800-1.00)	1.56 (1.32-1.69)	2.25 (2.05-2.60)	3.44 (2.27-3.98)	596
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1540
	03-04	.650 (.568-.742)	.900 (.800-.970)	1.30 (1.22-1.40)	2.00 (1.85-2.17)	2.60 (2.35-2.91)	1296
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	.665 (.606-.730)	.900 (.800-.970)	1.40 (1.30-1.50)	2.05 (1.90-2.27)	2.70 (2.33-3.08)	945
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	.648 (.545-.771)	.870 (.780-1.00)	1.30 (1.20-1.41)	2.00 (1.84-2.23)	2.60 (2.35-3.08)	947
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	.693 (.569-.843)	.840 (.700-1.00)	1.49 (1.14-1.70)	2.00 (1.64-2.40)	2.30 (2.00-3.10)	427
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.824 (.723-.939)	1.05 (.950-1.10)	1.70 (1.32-2.10)	2.81 (2.48-3.09)	3.42 (3.00-4.14)	462
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1056
	03-04	.625 (.531-.735)	.830 (.780-.940)	1.30 (1.20-1.40)	1.97 (1.85-2.17)	2.49 (2.30-2.80)	882

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 10.5 and 0.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,5'-Pentachlorobiphenyl (PCB 87) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
60+	05-06	*	*	15	
	07-08	*	*	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
60+	05-06	.416	0.073	17	
	07-08	*	*	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
60+	05-06	*	*	5	
	07-08	*	*	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
60+	05-06	*	*	5	
	07-08	*	*	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,5'-Pentachlorobiphenyl (PCB 87) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,5'-Pentachlorobiphenyl (PCB 87) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	.004 (.004-.005)	.005 (.005-.006)	.008 (.008-.009)	.012 (.012-.013)	.017 (.015-.017)	1892
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.004 (.003-.004)	.005 (.004-.005)	.008 (.007-.008)	.011 (.010-.013)	.016 (.011-.021)	596
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1540
	03-04	.004 (.004-.005)	.006 (.005-.006)	.008 (.008-.009)	.012 (.011-.014)	.017 (.014-.018)	1296
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	.004 (.004-.005)	.006 (.005-.006)	.009 (.008-.009)	.013 (.012-.014)	.017 (.015-.018)	945
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	.004 (.003-.005)	.005 (.005-.006)	.008 (.007-.009)	.012 (.011-.013)	.016 (.014-.017)	947
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	.004 (.003-.005)	.005 (.004-.006)	.008 (.007-.010)	.011 (.009-.014)	.014 (.011-.022)	427
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.005 (.004-.005)	.006 (.006-.006)	.009 (.008-.012)	.015 (.014-.017)	.020 (.017-.027)	462
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1056
	03-04	.004 (.003-.005)	.005 (.005-.006)	.008 (.007-.009)	.012 (.011-.013)	.016 (.014-.017)	882

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,5'-Pentachlorobiphenyl (PCB 87) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
60+	05-06	*	*	15	
	07-08	*	*	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	.003	0	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,5'-Pentachlorobiphenyl (PCB 87) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†]
(in ng/g of serum or parts per billion) for the U.S. population from
the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5-Pentachlorobiphenyl (PCB 99) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	13.1 (<LOD-14.7)	19.1 (16.2-20.6)	1897
	01-02	*	< LOD	< LOD	17.6 (15.3-21.0)	26.3 (22.1-30.5)	2281
	03-04	4.16 (3.82-4.54)	3.79 (3.43-4.10)	6.53 (5.76-7.56)	13.0 (10.5-16.1)	18.0 (16.7-19.4)	1877
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	654
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	2.34 (2.08-2.64)	2.30 (2.05-2.60)	3.20 (2.80-3.86)	5.16 (3.80-6.12)	6.00 (4.77-10.0)	587
20 years and older	99-00	*	< LOD	< LOD	13.9 (<LOD-16.0)	20.0 (17.2-21.6)	1243
	01-02	*	< LOD	10.8 (<LOD-12.3)	19.4 (16.7-22.3)	29.0 (23.5-32.1)	1523
	03-04	4.52 (4.14-4.94)	4.08 (3.75-4.45)	7.10 (6.08-8.53)	14.7 (10.9-17.0)	18.6 (17.1-21.3)	1290
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	16.7 (13.8-20.5)	905
	01-02	*	< LOD	< LOD	17.0 (13.8-21.0)	24.9 (19.8-30.1)	1061
	03-04	3.97 (3.64-4.33)	3.69 (3.31-4.00)	6.11 (5.44-7.00)	11.1 (9.40-14.3)	16.8 (12.9-19.0)	936
Females	99-00	*	< LOD	< LOD	13.9 (<LOD-16.2)	20.3 (17.3-23.5)	992
	01-02	*	< LOD	< LOD	18.0 (15.8-22.2)	28.5 (22.5-33.2)	1220
	03-04	4.35 (3.94-4.81)	3.90 (3.43-4.70)	7.05 (6.00-8.46)	15.3 (11.5-17.3)	18.9 (17.5-22.8)	941
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	624
	01-02	*	< LOD	< LOD	< LOD	13.8 (10.7-17.4)	562
	03-04	2.33 (2.08-2.62)	2.20 (1.90-2.47)	3.35 (2.80-4.02)	5.21 (4.50-6.10)	6.35 (5.53-7.55)	426
Non-Hispanic blacks	99-00	*	< LOD	< LOD	21.1 (17.0-31.1)	32.0 (22.9-57.4)	400
	01-02	*	< LOD	11.7 (10.6-13.1)	22.5 (18.8-25.8)	29.0 (23.0-37.2)	510
	03-04	5.54 (4.53-6.77)	4.81 (3.80-5.80)	10.4 (7.20-15.3)	23.1 (16.7-29.7)	31.7 (24.2-49.1)	447
Non-Hispanic whites	99-00	*	< LOD	< LOD	12.5 (<LOD-14.5)	18.2 (14.3-20.8)	715
	01-02	*	< LOD	< LOD	18.5 (15.3-22.3)	28.5 (22.1-32.4)	1046
	03-04	4.23 (3.80-4.72)	3.90 (3.40-4.44)	6.70 (5.70-8.00)	12.6 (9.73-15.8)	17.3 (15.5-18.1)	883

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.5, 10.5, and 0.6, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5-Pentachlorobiphenyl (PCB 99) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	1.51	0.17	9
		07-08	1.47	0.16	6
	20-39	05-06	2.60	0.41	12
		07-08	2.29	0.19	15
	40-59	05-06	4.29	0.4	12
		07-08	3.80	0.32	16
60+	05-06	8.00	0.53	15	
	07-08	8.46	0.78	23	
Non-Hispanic White Female	12-19	05-06	1.30	0.07	10
		07-08	.931	0.206	7
	20-39	05-06	1.86	0.1	16
		07-08	2.06	0.32	13
	40-59	05-06	4.50	0.3	13
		07-08	3.93	0.29	17
60+	05-06	10.0	1	17	
	07-08	8.84	0.87	21	
Non-Hispanic Black Male	12-19	05-06	2.09	0.22	13
		07-08	1.43	0.13	6
	20-39	05-06	2.46	0.25	6
		07-08	2.18	0.19	6
	40-59	05-06	5.47	0.79	5
		07-08	6.22	1.22	6
60+	05-06	14.2	1.8	5	
	07-08	11.5	1.2	8	
Non-Hispanic Black Female	12-19	05-06	1.37	0.09	14
		07-08	1.20	0.16	5
	20-39	05-06	2.39	0.25	7
		07-08	1.94	0.26	8
	40-59	05-06	7.60	0.79	7
		07-08	6.43	0.75	8
60+	05-06	21.5	5.7	5	
	07-08	17.9	2.9	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5-Pentachlorobiphenyl (PCB 99) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.912	0.089	11
		07-08	.810	0.143	6
	20-39	05-06	1.46	0.09	9
		07-08	1.52	0.13	9
	40-59	05-06	2.39	0.15	4
		07-08	3.19	0.52	6
60+	05-06	4.19	0.31	4	
	07-08	4.50	0.45	5	
Mexican American Female	12-19	05-06	.816	0.043	16
		07-08	.722	0.075	5
	20-39	05-06	1.13	0.12	9
		07-08	1.27	0.07	8
	40-59	05-06	3.32	0.33	6
		07-08	2.74	0.26	6
	60+	05-06	5.65	0.74	3
		07-08	5.21	1.01	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5-Pentachlorobiphenyl (PCB 99) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	.090 (<LOD-.100)	.120 (.100-.150)	1897
	01-02	*	< LOD	< LOD	.120 (.100-.140)	.180 (.150-.210)	2281
	03-04	.025 (.023-.028)	.024 (.021-.026)	.042 (.038-.049)	.082 (.067-.102)	.119 (.102-.140)	1877
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	654
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.012 (.011-.013)	.012 (.010-.013)	.017 (.015-.019)	.025 (.019-.032)	.032 (.025-.045)	587
20 years and older	99-00	*	< LOD	< LOD	.090 (<LOD-.100)	.130 (.110-.150)	1243
	01-02	*	< LOD	.070 (<LOD-.080)	.130 (.110-.160)	.190 (.160-.210)	1523
	03-04	.028 (.026-.031)	.026 (.024-.028)	.046 (.041-.054)	.093 (.073-.110)	.127 (.110-.153)	1290
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	.110 (.090-.130)	905
	01-02	*	< LOD	< LOD	.110 (.090-.140)	.160 (.120-.200)	1061
	03-04	.025 (.022-.027)	.023 (.021-.025)	.040 (.036-.045)	.070 (.059-.090)	.110 (.078-.132)	936
Females	99-00	*	< LOD	< LOD	.100 (<LOD-.110)	.130 (.100-.170)	992
	01-02	*	< LOD	< LOD	.130 (.110-.160)	.200 (.160-.210)	1220
	03-04	.026 (.024-.029)	.024 (.020-.028)	.045 (.040-.053)	.095 (.078-.111)	.130 (.110-.158)	941
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	624
	01-02	*	< LOD	< LOD	< LOD	.100 (.070-.120)	562
	03-04	.014 (.012-.016)	.013 (.011-.016)	.022 (.019-.026)	.036 (.030-.040)	.046 (.036-.058)	426
Non-Hispanic blacks	99-00	*	< LOD	< LOD	.140 (.110-.180)	.210 (.150-.400)	400
	01-02	*	< LOD	.070 (.060-.090)	.140 (.110-.150)	.190 (.140-.230)	510
	03-04	.032 (.026-.039)	.027 (.023-.033)	.060 (.042-.086)	.149 (.112-.172)	.211 (.150-.306)	447
Non-Hispanic whites	99-00	*	< LOD	< LOD	.090 (<LOD-.100)	.110 (.090-.150)	715
	01-02	*	< LOD	< LOD	.120 (.100-.160)	.190 (.160-.210)	1046
	03-04	.026 (.023-.029)	.025 (.022-.027)	.043 (.038-.051)	.081 (.065-.102)	.112 (.098-.127)	883

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5-Pentachlorobiphenyl (PCB 99) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.008	0.001	9
		07-08	.007	0.001	6
	20-39	05-06	.019	0.004	12
		07-08	.015	0.002	15
	40-59	05-06	.032	0.003	12
		07-08	.028	0.002	16
60+	05-06	.054	0.003	15	
	07-08	.054	0.005	23	
Non-Hispanic White Female	12-19	05-06	.007	0	10
		07-08	.004	0.001	7
	20-39	05-06	.012	0.001	16
		07-08	.012	0.002	13
	40-59	05-06	.033	0.003	13
		07-08	.026	0.002	17
60+	05-06	.072	0.007	17	
	07-08	.059	0.006	21	
Non-Hispanic Black Male	12-19	05-06	.010	0.001	13
		07-08	.007	0.001	6
	20-39	05-06	.015	0.002	6
		07-08	.012	0.001	6
	40-59	05-06	.035	0.005	5
		07-08	.040	0.008	6
60+	05-06	.084	0.009	5	
	07-08	.070	0.008	8	
Non-Hispanic Black Female	12-19	05-06	.007	0	14
		07-08	.006	0.001	5
	20-39	05-06	.014	0.002	7
		07-08	.010	0.001	8
	40-59	05-06	.049	0.006	7
		07-08	.039	0.005	8
60+	05-06	.144	0.039	5	
	07-08	.106	0.014	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5-Pentachlorobiphenyl (PCB 99) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†]
(in ng/g of serum or parts per billion) for the U.S. population from
the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.005	0	11
		07-08	.004	0.001	6
	20-39	05-06	.010	0.001	9
		07-08	.010	0.001	9
	40-59	05-06	.019	0.002	4
		07-08	.023	0.004	6
60+	05-06	.031	0.004	4	
	07-08	.032	0.003	5	
Mexican American Female	12-19	05-06	.004	0	16
		07-08	.003	0	5
	20-39	05-06	.008	0.001	9
		07-08	.007	0.001	8
	40-59	05-06	.024	0.002	6
		07-08	.019	0.002	6
	60+	05-06	.042	0.005	3
		07-08	.036	0.007	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5,5'-Pentachlorobiphenyl (PCB 101) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1929
	01-02	*	< LOD	< LOD	< LOD	< LOD	2307
	03-04	1.65 (1.51-1.81)	1.70 (1.50-1.80)	2.70 (2.50-2.94)	4.40 (3.97-4.82)	5.83 (5.29-6.66)	1897
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	669
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	1.93 (1.76-2.11)	1.73 (1.51-2.10)	3.20 (3.00-3.60)	5.05 (4.23-5.87)	7.25 (5.10-8.30)	598
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1260
	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	1.62 (1.46-1.78)	1.67 (1.50-1.80)	2.64 (2.40-2.90)	4.40 (3.88-4.75)	5.51 (5.00-6.60)	1299
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	918
	01-02	*	< LOD	< LOD	< LOD	< LOD	1075
	03-04	1.71 (1.59-1.84)	1.75 (1.60-1.90)	2.80 (2.60-3.10)	4.50 (3.97-5.06)	6.00 (5.35-6.75)	947
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1011
	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	1.60 (1.41-1.81)	1.61 (1.41-1.82)	2.57 (2.30-2.98)	4.40 (3.71-4.94)	5.60 (4.88-7.25)	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	634
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	1.72 (1.55-1.92)	1.70 (1.50-1.96)	3.00 (2.30-3.80)	4.40 (3.56-5.38)	5.38 (4.10-6.60)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	413
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	1.69 (1.38-2.07)	1.70 (1.50-2.16)	3.00 (2.57-3.80)	5.88 (3.88-7.39)	7.68 (6.60-11.9)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	724
	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	1.61 (1.46-1.79)	1.61 (1.41-1.80)	2.60 (2.37-2.90)	4.40 (3.75-4.61)	5.50 (5.00-6.15)	885

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 25.7, 10.5, and 0.6, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5,5'-Pentachlorobiphenyl (PCB 101) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	.632	0.15	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.59 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5,5'-Pentachlorobiphenyl (PCB 101) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.59 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5,5'-Pentachlorobiphenyl (PCB 101) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1929
	01-02	*	< LOD	< LOD	< LOD	< LOD	2307
	03-04	.010 (.009-.011)	.010 (.009-.012)	.016 (.015-.018)	.027 (.024-.028)	.033 (.031-.037)	1897
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	669
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.010 (.009-.011)	.009 (.008-.011)	.016 (.014-.018)	.024 (.021-.028)	.030 (.025-.044)	598
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1260
	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	.010 (.009-.011)	.011 (.009-.012)	.016 (.015-.018)	.027 (.024-.029)	.034 (.031-.037)	1299
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	918
	01-02	*	< LOD	< LOD	< LOD	< LOD	1075
	03-04	.011 (.010-.011)	.011 (.010-.012)	.017 (.016-.019)	.025 (.024-.029)	.037 (.031-.041)	947
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1011
	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	.010 (.009-.011)	.010 (.009-.011)	.015 (.014-.018)	.027 (.022-.029)	.032 (.030-.036)	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	634
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	.011 (.009-.012)	.011 (.009-.012)	.018 (.014-.021)	.024 (.020-.030)	.029 (.022-.052)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	413
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.010 (.008-.012)	.011 (.008-.013)	.016 (.013-.022)	.032 (.023-.041)	.045 (.032-.059)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	724
	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	.010 (.009-.011)	.010 (.009-.011)	.016 (.014-.018)	.025 (.024-.027)	.031 (.029-.037)	885

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5,5'-Pentachlorobiphenyl (PCB 101) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	.005	0.001	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,5,5'-Pentachlorobiphenyl (PCB 101) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3,3',4',6-Pentachlorobiphenyl (PCB 110) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	1.22 (1.11-1.33)	1.20 (1.10-1.36)	1.96 (1.80-2.20)	3.40 (3.10-3.57)	4.42 (3.88-4.95)	1882
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	1.44 (1.30-1.59)	1.30 (1.20-1.50)	2.50 (2.19-2.91)	4.13 (3.40-4.90)	5.40 (4.30-7.68)	593
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1540
	03-04	1.19 (1.08-1.30)	1.20 (1.09-1.33)	1.88 (1.71-2.11)	3.30 (2.95-3.50)	4.18 (3.66-4.94)	1289
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	1.26 (1.18-1.35)	1.30 (1.19-1.40)	2.05 (1.80-2.30)	3.42 (3.07-3.70)	4.61 (3.80-5.00)	939
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	1.17 (1.04-1.32)	1.20 (1.00-1.36)	1.82 (1.70-2.20)	3.40 (2.77-3.57)	4.40 (3.57-5.54)	943
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	1.33 (1.18-1.51)	1.29 (1.13-1.50)	2.30 (1.80-2.95)	3.40 (2.70-4.30)	4.10 (3.30-5.50)	420
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	1.36 (1.20-1.53)	1.30 (1.14-1.50)	2.10 (1.72-3.00)	4.19 (2.91-5.10)	5.32 (4.40-6.89)	464
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1056
	03-04	1.17 (1.05-1.31)	1.20 (1.00-1.33)	1.86 (1.70-2.16)	3.40 (2.93-3.57)	4.19 (3.69-4.95)	877

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 10.5 and 0.8.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3,3',4',6-Pentachlorobiphenyl (PCB 110) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	.374	0.077	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	.315	0.082	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3,3',4',6-Pentachlorobiphenyl (PCB 110) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3,3',4',6-Pentachlorobiphenyl (PCB 110) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	.007 (.007-.008)	.007 (.007-.008)	.012 (.011-.013)	.019 (.018-.021)	.026 (.023-.028)	1882
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.007 (.007-.008)	.007 (.006-.008)	.013 (.010-.015)	.020 (.017-.022)	.026 (.019-.037)	593
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1540
	03-04	.007 (.007-.008)	.007 (.007-.008)	.012 (.011-.013)	.019 (.017-.021)	.025 (.023-.028)	1289
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	.008 (.007-.008)	.008 (.007-.009)	.013 (.011-.014)	.019 (.018-.021)	.028 (.022-.031)	939
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	.007 (.006-.008)	.007 (.006-.008)	.011 (.010-.013)	.019 (.016-.022)	.024 (.021-.027)	943
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	.008 (.007-.009)	.008 (.007-.010)	.014 (.011-.016)	.019 (.016-.022)	.022 (.017-.043)	420
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.008 (.007-.009)	.008 (.006-.009)	.012 (.010-.016)	.023 (.017-.028)	.028 (.023-.032)	464
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1056
	03-04	.007 (.006-.008)	.007 (.006-.008)	.012 (.010-.013)	.019 (.017-.021)	.026 (.022-.029)	877

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3,3',4',6-Pentachlorobiphenyl (PCB 110) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	.002	0	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	.002	0	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,3,3',4',6-Pentachlorobiphenyl (PCB 110) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4'-Hexachlorobiphenyl (PCB 128) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1927
	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	*	< LOD	< LOD	< LOD	.600 (.500-.700)	1877
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	668
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	< LOD	< LOD	.510 (<LOD-.700)	589
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1259
	01-02	*	< LOD	< LOD	< LOD	< LOD	1540
	03-04	*	< LOD	< LOD	< LOD	.620 (.490-.800)	1288
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	917
	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	*	< LOD	< LOD	< LOD	.600 (.420-.770)	937
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1010
	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	*	< LOD	< LOD	< LOD	.630 (.500-.800)	940
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	636
	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	*	< LOD	< LOD	< LOD	< LOD	424
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	*	< LOD	< LOD	.670 (.500-.800)	1.00 (.770-2.10)	455
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	725
	01-02	*	< LOD	< LOD	< LOD	< LOD	1056
	03-04	*	< LOD	< LOD	< LOD	.500 (.420-.600)	878

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.4, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4'-Hexachlorobiphenyl (PCB 128) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	.410	0.103	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4'-Hexachlorobiphenyl (PCB 128) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4'-Hexachlorobiphenyl (PCB 128) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1927
	01-02	*	< LOD	< LOD	< LOD	< LOD	2298
	03-04	*	< LOD	< LOD	< LOD	.004 (.003-.004)	1877
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	668
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	< LOD	< LOD	.003 (<LOD-.004)	589
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1259
	01-02	*	< LOD	< LOD	< LOD	< LOD	1540
	03-04	*	< LOD	< LOD	< LOD	.004 (.004-.005)	1288
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	917
	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	*	< LOD	< LOD	< LOD	.004 (.003-.005)	937
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1010
	01-02	*	< LOD	< LOD	< LOD	< LOD	1229
	03-04	*	< LOD	< LOD	< LOD	.004 (.003-.005)	940
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	636
	01-02	*	< LOD	< LOD	< LOD	< LOD	564
	03-04	*	< LOD	< LOD	< LOD	< LOD	424
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	*	< LOD	< LOD	.004 (.003-.005)	.006 (.005-.013)	455
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	725
	01-02	*	< LOD	< LOD	< LOD	< LOD	1056
	03-04	*	< LOD	< LOD	< LOD	.003 (.002-.004)	878

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4'-Hexachlorobiphenyl (PCB 128) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	.003	0.001	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	*	*	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4'-Hexachlorobiphenyl (PCB 128) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5' and 2,3,3',4,4',6-Hexachlorobiphenyl (PCB 138 & 158) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	49.3 (42.9-55.8)	71.2 (59.8-82.7)	1930
	01-02	19.9 (18.0-22.0)	20.2 (18.2-23.1)	40.4 (35.3-45.1)	70.1 (61.8-78.8)	94.6 (82.5-107)	2293
	03-04	15.1 (14.1-16.1)	15.1 (13.6-16.6)	30.5 (28.1-34.0)	55.4 (47.3-63.3)	75.3 (69.0-81.8)	1896
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	669
	01-02	*	< LOD	< LOD	17.0 (13.7-20.2)	23.1 (17.7-41.2)	748
	03-04	4.97 (4.45-5.55)	4.57 (4.10-5.02)	7.39 (6.35-8.63)	12.7 (10.4-14.5)	15.2 (13.1-21.3)	598
20 years and older	99-00	*	< LOD	< LOD	54.7 (47.4-60.8)	72.8 (66.0-88.9)	1261
	01-02	23.3 (21.1-25.8)	24.0 (21.4-26.8)	44.6 (40.4-49.2)	73.9 (66.2-83.0)	100 (88.8-109)	1545
	03-04	17.7 (16.5-19.0)	17.6 (15.5-20.4)	34.3 (30.0-38.8)	59.6 (50.6-66.9)	77.4 (72.3-87.7)	1298
Gender							
Males	99-00	*	< LOD	< LOD	47.2 (<LOD-55.7)	68.2 (55.7-83.8)	918
	01-02	20.1 (17.9-22.6)	20.5 (17.6-23.5)	39.2 (32.2-46.0)	69.2 (58.1-82.5)	94.6 (77.8-109)	1066
	03-04	14.9 (13.8-15.9)	14.4 (12.9-15.8)	29.6 (26.6-33.8)	56.3 (45.3-62.3)	75.8 (63.3-87.7)	947
Females	99-00	*	< LOD	< LOD	53.7 (44.7-61.2)	72.1 (61.2-88.9)	1012
	01-02	19.7 (17.8-21.7)	20.2 (18.2-23.1)	41.8 (37.5-45.9)	70.1 (63.4-78.5)	93.9 (80.2-109)	1227
	03-04	15.3 (14.0-16.8)	15.8 (14.2-17.8)	31.8 (28.3-36.6)	54.6 (47.3-64.4)	72.5 (68.4-80.9)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	636
	01-02	*	< LOD	18.6 (13.6-23.4)	33.3 (26.5-44.2)	51.8 (42.3-57.6)	559
	03-04	7.06 (6.02-8.28)	6.52 (5.20-7.60)	12.3 (8.91-16.0)	23.6 (18.9-28.3)	29.2 (24.7-36.1)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	72.6 (61.7-89.4)	122 (86.5-185)	412
	01-02	22.3 (19.3-25.6)	22.0 (18.4-27.1)	46.5 (38.0-55.6)	91.5 (70.1-112)	122 (91.5-169)	513
	03-04	17.8 (14.9-21.3)	15.9 (12.5-21.0)	41.2 (30.6-57.3)	86.5 (69.4-110)	153 (94.6-191)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	49.3 (41.4-55.8)	70.1 (55.7-85.2)	727
	01-02	21.5 (19.2-24.0)	22.1 (19.6-24.9)	43.0 (37.7-48.0)	70.7 (62.3-79.7)	96.1 (79.1-110)	1057
	03-04	16.0 (14.5-17.7)	15.8 (14.6-18.4)	31.4 (28.2-36.6)	55.1 (47.2-62.3)	71.6 (64.0-75.8)	883

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 41.1, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5' and 2,3,3',4,4',6-Hexachlorobiphenyl (PCB 138 & 158) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†]
(in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from
the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	4.13	0.65	9
		07-08	4.81	0.89	6
	20-39	05-06	7.98	1.5	12
		07-08	8.74	0.84	15
	40-59	05-06	19.6	2.5	12
		07-08	20.7	1.6	16
60+	05-06	32.6	1.5	15	
	07-08	47.5	3.9	23	
Non-Hispanic White Female	12-19	05-06	3.24	0.3	10
		07-08	3.53	0.73	7
	20-39	05-06	5.23	0.28	16
		07-08	7.85	1.28	13
	40-59	05-06	16.5	1.2	13
		07-08	21.5	1.2	17
60+	05-06	33.5	3.1	17	
	07-08	45.0	3.3	21	
Non-Hispanic Black Male	12-19	05-06	4.83	0.62	13
		07-08	4.63	0.64	6
	20-39	05-06	7.69	1.17	6
		07-08	7.95	0.77	6
	40-59	05-06	20.9	2.4	5
		07-08	28.6	4.5	6
60+	05-06	56.5	7.5	5	
	07-08	68.2	8	8	
Non-Hispanic Black Female	12-19	05-06	2.97	0.19	14
		07-08	3.25	0.5	5
	20-39	05-06	6.94	0.87	7
		07-08	7.16	0.93	8
	40-59	05-06	25.5	2.5	7
		07-08	28.2	2.8	8
60+	05-06	81.1	23.7	5	
	07-08	79.9	7.1	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

**Serum 2,2',3,4,4',5' and 2,3,3',4,4',6-Hexachlorobiphenyl (PCB 138 & 158)
(lipid adjusted)**

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†]
(in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from
the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	2.68	0.35	11
		07-08	2.58	0.55	6
	20-39	05-06	4.46	0.27	9
		07-08	5.67	0.55	9
	40-59	05-06	9.75	0.58	4
		07-08	15.2	2	6
60+	05-06	19.6	3	4	
	07-08	23.9	3.3	5	
Mexican American Female	12-19	05-06	1.92	0.19	16
		07-08	1.84	0.53	5
	20-39	05-06	3.22	0.4	9
		07-08	4.41	0.22	8
	40-59	05-06	12.5	1	6
		07-08	14.6	1.9	6
60+	05-06	21.7	2.3	3	
	07-08	25.6	4	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5' and 2,3,3',4,4',6-Hexachlorobiphenyl (PCB 138 & 158) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	.340 (.300-.370)	.460 (.390-.530)	1930
	01-02	.122 (.110-.135)	.120 (.110-.140)	.270 (.230-.290)	.460 (.410-.510)	.650 (.560-.700)	2293
	03-04	.092 (.086-.099)	.095 (.082-.105)	.206 (.180-.231)	.359 (.326-.392)	.477 (.450-.528)	1896
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	669
	01-02	*	< LOD	< LOD	.090 (.060-.110)	.110 (.090-.210)	748
	03-04	.025 (.023-.028)	.023 (.020-.027)	.037 (.030-.047)	.062 (.049-.074)	.079 (.067-.103)	598
20 years and older	99-00	*	< LOD	< LOD	.360 (.320-.390)	.490 (.400-.560)	1261
	01-02	.148 (.133-.163)	.150 (.140-.170)	.290 (.270-.320)	.510 (.450-.550)	.680 (.610-.720)	1545
	03-04	.111 (.103-.120)	.114 (.099-.128)	.232 (.204-.256)	.383 (.354-.416)	.506 (.467-.594)	1298
Gender							
Males	99-00	*	< LOD	< LOD	.320 (<LOD-.370)	.420 (.370-.530)	918
	01-02	.125 (.112-.140)	.130 (.110-.140)	.260 (.230-.290)	.460 (.370-.540)	.630 (.510-.710)	1066
	03-04	.092 (.085-.099)	.091 (.079-.103)	.198 (.172-.229)	.354 (.316-.379)	.474 (.389-.562)	947
Females	99-00	*	< LOD	< LOD	.340 (.300-.370)	.490 (.390-.540)	1012
	01-02	.119 (.107-.132)	.120 (.110-.150)	.270 (.230-.300)	.460 (.420-.500)	.660 (.590-.700)	1227
	03-04	.093 (.084-.102)	.098 (.085-.115)	.212 (.178-.244)	.373 (.325-.417)	.482 (.450-.568)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	636
	01-02	*	< LOD	.110 (.090-.150)	.240 (.170-.330)	.350 (.270-.430)	559
	03-04	.043 (.036-.051)	.040 (.032-.049)	.080 (.068-.100)	.148 (.122-.203)	.239 (.147-.346)	427
Non-Hispanic blacks	99-00	*	< LOD	< LOD	.470 (.360-.670)	.830 (.530-1.22)	412
	01-02	.125 (.108-.146)	.130 (.090-.160)	.290 (.230-.350)	.560 (.420-.680)	.770 (.560-1.06)	513
	03-04	.101 (.084-.122)	.087 (.068-.113)	.239 (.186-.326)	.568 (.424-.811)	.965 (.698-1.09)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	.340 (.300-.370)	.440 (.370-.530)	727
	01-02	.133 (.119-.149)	.140 (.120-.160)	.280 (.250-.300)	.460 (.420-.530)	.650 (.550-.720)	1057
	03-04	.099 (.089-.109)	.104 (.091-.121)	.216 (.179-.252)	.355 (.320-.390)	.467 (.410-.520)	883

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

**Serum 2,2',3,4,4',5' and 2,3,3',4,4',6-Hexachlorobiphenyl (PCB 138 & 158)
(whole weight)**

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†]
(in ng/g of serum or parts per billion) for the U.S. population from the National Health and
Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]	
Non-Hispanic White Male	12-19	05-06	.023	0.003	9	
		07-08	.024	0.004	6	
	20-39	05-06	.060	0.013	12	
		07-08	.058	0.006	15	
	40-59	05-06	.150	0.022	12	
		07-08	.150	0.01	16	
	60+	05-06	.221	0.008	15	
		07-08	.303	0.023	23	
	Non-Hispanic White Female	12-19	05-06	.017	0.002	10
			07-08	.017	0.004	7
20-39		05-06	.034	0.002	16	
		07-08	.047	0.009	13	
40-59		05-06	.123	0.011	13	
		07-08	.143	0.009	17	
60+		05-06	.242	0.02	17	
		07-08	.302	0.022	21	
Non-Hispanic Black Male		12-19	05-06	.024	0.003	13
			07-08	.022	0.003	6
	20-39	05-06	.046	0.007	6	
		07-08	.045	0.005	6	
	40-59	05-06	.133	0.016	5	
		07-08	.183	0.028	6	
	60+	05-06	.335	0.042	5	
		07-08	.416	0.054	8	
	Non-Hispanic Black Female	12-19	05-06	.015	0.001	14
			07-08	.016	0.003	5
20-39		05-06	.040	0.005	7	
		07-08	.036	0.005	8	
40-59		05-06	.165	0.019	7	
		07-08	.171	0.02	8	
60+		05-06	.541	0.16	5	
		07-08	.477	0.035	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

**Serum 2,2',3,4,4',5' and 2,3,3',4,4',6-Hexachlorobiphenyl (PCB 138 & 158)
(whole weight)**

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†]
(in ng/g of serum or parts per billion) for the U.S. population from
the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]	
Mexican American Male	12-19	05-06	.014	0.002	11	
		07-08	.014	0.003	6	
	20-39	05-06	.032	0.003	9	
		07-08	.039	0.004	9	
	40-59	05-06	.079	0.004	4	
		07-08	.108	0.015	6	
	60+	05-06	.142	0.023	4	
		07-08	.174	0.025	5	
	Mexican American Female	12-19	05-06	.010	0.001	16
			07-08	.009	0.002	5
20-39		05-06	.022	0.003	9	
		07-08	.026	0.001	8	
40-59		05-06	.091	0.007	6	
		07-08	.099	0.014	6	
60+		05-06	.162	0.011	3	
		07-08	.177	0.029	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5'-Hexachlorobiphenyl (PCB 146) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	13.3 (<LOD-16.1)	1923
	01-02	*	< LOD	< LOD	11.0 (<LOD-12.5)	15.4 (13.6-16.9)	2299
	03-04	2.17 (2.02-2.32)	2.21 (2.03-2.54)	4.80 (4.38-5.21)	8.27 (7.10-9.50)	11.7 (10.2-13.3)	1894
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.620 (.562-.685)	.600 (.520-.690)	1.02 (.870-1.20)	1.67 (1.50-1.80)	2.30 (1.79-2.97)	594
20 years and older	99-00	*	< LOD	< LOD	< LOD	14.3 (<LOD-17.1)	1256
	01-02	*	< LOD	< LOD	11.8 (10.7-13.5)	16.5 (14.6-18.1)	1541
	03-04	2.60 (2.41-2.79)	2.60 (2.40-2.90)	5.27 (4.82-5.88)	8.81 (8.00-9.80)	12.7 (10.7-15.2)	1300
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	13.1 (<LOD-17.1)	915
	01-02	*	< LOD	< LOD	11.0 (<LOD-13.0)	15.4 (13.0-17.2)	1069
	03-04	2.16 (1.97-2.37)	2.10 (1.90-2.50)	4.80 (4.09-5.30)	8.30 (6.90-10.1)	12.4 (10.0-14.9)	945
Females	99-00	*	< LOD	< LOD	< LOD	13.9 (<LOD-16.3)	1008
	01-02	*	< LOD	< LOD	11.0 (<LOD-12.2)	15.2 (13.2-17.3)	1230
	03-04	2.17 (2.02-2.34)	2.35 (2.04-2.63)	4.83 (4.47-5.37)	8.00 (7.04-9.10)	11.3 (9.95-13.2)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	633
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	.940 (.826-1.07)	.820 (.780-1.00)	1.74 (1.21-2.60)	3.31 (2.80-4.00)	4.60 (3.47-6.00)	425
Non-Hispanic blacks	99-00	*	< LOD	< LOD	17.0 (13.4-22.9)	29.8 (20.2-42.2)	412
	01-02	*	< LOD	< LOD	16.6 (12.8-21.4)	23.8 (16.9-31.6)	515
	03-04	2.81 (2.37-3.34)	2.60 (2.05-3.50)	7.10 (4.97-8.90)	16.8 (11.5-23.2)	28.5 (17.8-41.6)	463
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	723
	01-02	*	< LOD	< LOD	10.8 (<LOD-12.0)	15.2 (12.8-16.8)	1054
	03-04	2.25 (2.06-2.45)	2.35 (2.10-2.60)	4.82 (4.37-5.41)	7.76 (6.94-8.70)	10.5 (9.51-11.3)	885

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.4, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5'-Hexachlorobiphenyl (PCB 146) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.633	0.079	9
		07-08	*	*	6
	20-39	05-06	1.41	0.24	12
		07-08	1.21	0.11	15
	40-59	05-06	3.69	0.22	12
		07-08	3.32	0.21	16
	60+	05-06	7.11	0.38	15
		07-08	7.67	0.73	23
Non-Hispanic White Female	12-19	05-06	.514	0.05	10
		07-08	*	*	7
	20-39	05-06	.908	0.06	16
		07-08	1.09	0.25	13
	40-59	05-06	3.13	0.19	13
		07-08	3.20	0.16	17
	60+	05-06	7.04	0.65	17
		07-08	7.10	0.53	21
Non-Hispanic Black Male	12-19	05-06	.890	0.147	13
		07-08	.694	0.182	6
	20-39	05-06	1.73	0.33	6
		07-08	1.26	0.15	6
	40-59	05-06	5.46	0.74	5
		07-08	5.79	0.97	6
	60+	05-06	14.3	1.5	5
		07-08	14.7	2	8
Non-Hispanic Black Female	12-19	05-06	.517	0.047	14
		07-08	*	*	5
	20-39	05-06	1.36	0.19	7
		07-08	1.17	0.13	8
	40-59	05-06	5.30	0.61	7
		07-08	4.78	0.53	8
	60+	05-06	21.0	5.7	5
		07-08	15.5	1.5	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5'-Hexachlorobiphenyl (PCB 146) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.358	0.042	11
		07-08	*	*	6
	20-39	05-06	.828	0.059	9
		07-08	.841	0.079	9
	40-59	05-06	1.94	0.1	4
		07-08	3.10	0.43	6
	60+	05-06	4.17	0.84	4
		07-08	4.91	0.43	5
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.538	0.075	9
		07-08	.610	0.092	8
	40-59	05-06	2.21	0.19	6
		07-08	2.49	0.31	6
	60+	05-06	4.29	0.14	3
		07-08	4.67	0.82	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5'-Hexachlorobiphenyl (PCB 146) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	.080 (<LOD-.110)	1923
	01-02	*	< LOD	< LOD	.070 (<LOD-.080)	.110 (.090-.120)	2299
	03-04	.013 (.012-.014)	.014 (.013-.016)	.032 (.028-.035)	.054 (.047-.062)	.077 (.069-.090)	1894
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.003 (.003-.003)	.003 (.003-.003)	.006 (.004-.007)	.008 (.007-.010)	.011 (.009-.014)	594
20 years and older	99-00	*	< LOD	< LOD	< LOD	.090 (<LOD-.120)	1256
	01-02	*	< LOD	< LOD	.080 (.070-.090)	.110 (.100-.120)	1541
	03-04	.016 (.015-.018)	.017 (.015-.020)	.035 (.032-.038)	.058 (.053-.064)	.084 (.073-.099)	1300
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	.080 (<LOD-.110)	915
	01-02	*	< LOD	< LOD	.070 (<LOD-.090)	.100 (.090-.120)	1069
	03-04	.013 (.012-.015)	.014 (.012-.016)	.032 (.027-.036)	.054 (.045-.063)	.074 (.064-.096)	945
Females	99-00	*	< LOD	< LOD	< LOD	.090 (<LOD-.110)	1008
	01-02	*	< LOD	< LOD	.070 (<LOD-.080)	.110 (.090-.120)	1230
	03-04	.013 (.012-.014)	.015 (.013-.016)	.032 (.028-.036)	.054 (.047-.065)	.079 (.069-.088)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	633
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	.006 (.005-.007)	.005 (.004-.007)	.012 (.008-.016)	.024 (.017-.033)	.034 (.024-.046)	425
Non-Hispanic blacks	99-00	*	< LOD	< LOD	.110 (.080-.130)	.180 (.120-.270)	412
	01-02	*	< LOD	< LOD	.100 (.080-.130)	.150 (.110-.200)	515
	03-04	.016 (.013-.019)	.016 (.012-.019)	.041 (.030-.061)	.100 (.069-.167)	.180 (.126-.224)	463
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	723
	01-02	*	< LOD	< LOD	.070 (<LOD-.080)	.100 (.090-.120)	1054
	03-04	.014 (.013-.015)	.015 (.013-.018)	.033 (.028-.036)	.053 (.047-.057)	.069 (.064-.075)	885

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5'-Hexachlorobiphenyl (PCB 146) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.004	0	9
		07-08	*	*	6
	20-39	05-06	.010	0.002	12
		07-08	.008	0.001	15
	40-59	05-06	.028	0.002	12
		07-08	.024	0.001	16
	60+	05-06	.048	0.002	15
		07-08	.049	0.004	23
Non-Hispanic White Female	12-19	05-06	.003	0	10
		07-08	*	*	7
	20-39	05-06	.006	0	16
		07-08	.007	0.002	13
	40-59	05-06	.023	0.002	13
		07-08	.021	0.001	17
	60+	05-06	.051	0.004	17
		07-08	.048	0.003	21
Non-Hispanic Black Male	12-19	05-06	.004	0.001	13
		07-08	.003	0.001	6
	20-39	05-06	.010	0.002	6
		07-08	.007	0.001	6
	40-59	05-06	.035	0.005	5
		07-08	.037	0.006	6
	60+	05-06	.086	0.009	5
		07-08	.090	0.013	8
Non-Hispanic Black Female	12-19	05-06	.003	0	14
		07-08	*	*	5
	20-39	05-06	.008	0.001	7
		07-08	.006	0.001	8
	40-59	05-06	.034	0.004	7
		07-08	.029	0.004	8
	60+	05-06	.140	0.038	5
		07-08	.093	0.009	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5'-Hexachlorobiphenyl (PCB 146) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.002	0	11
		07-08	*	*	6
	20-39	05-06	.006	0.001	9
		07-08	.006	0.001	9
	40-59	05-06	.016	0.001	4
		07-08	.022	0.003	6
60+	05-06	.030	0.006	4	
	07-08	.035	0.002	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.004	0.001	9
		07-08	.004	0.001	8
	40-59	05-06	.016	0.001	6
		07-08	.017	0.002	6
	60+	05-06	.033	0.003	3
		07-08	.032	0.006	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5',6-Hexachlorobiphenyl (PCB 149) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2307
	03-04	.598 (.556-.642)	.600 (.550-.680)	.900 (.830-1.00)	1.45 (1.28-1.60)	1.90 (1.68-2.20)	1873
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.650 (.605-.698)	.640 (.600-.700)	1.00 (.900-1.10)	1.70 (1.40-2.15)	2.40 (1.80-2.98)	590
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	.590 (.546-.638)	.600 (.540-.680)	.900 (.800-1.00)	1.40 (1.20-1.60)	1.89 (1.60-2.17)	1283
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1075
	03-04	.623 (.578-.671)	.650 (.600-.700)	.960 (.880-1.06)	1.50 (1.30-1.66)	1.90 (1.60-2.40)	930
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	.575 (.521-.634)	.600 (.500-.620)	.890 (.800-.950)	1.40 (1.20-1.67)	1.90 (1.69-2.20)	943
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	.549 (.496-.608)	.530 (.500-.600)	.900 (.700-1.04)	1.28 (1.03-1.60)	1.60 (1.26-2.30)	408
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.760 (.678-.853)	.800 (.700-.880)	1.20 (1.00-1.58)	1.98 (1.70-2.59)	2.90 (2.20-4.20)	463
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	.574 (.524-.628)	.600 (.510-.670)	.880 (.800-.950)	1.33 (1.19-1.50)	1.80 (1.50-2.02)	881

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 10.5 and 0.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5',6-Hexachlorobiphenyl (PCB 149) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	*	*	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	.373	0.106	5
		07-08	*	*	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5',6-Hexachlorobiphenyl (PCB 149) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5',6-Hexachlorobiphenyl (PCB 149) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2307
	03-04	.004 (.003-.004)	.004 (.004-.004)	.006 (.005-.006)	.009 (.008-.009)	.011 (.010-.013)	1873
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.003 (.003-.004)	.003 (.003-.003)	.005 (.004-.006)	.008 (.007-.009)	.012 (.008-.016)	590
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	.004 (.003-.004)	.004 (.004-.004)	.006 (.005-.006)	.009 (.007-.010)	.011 (.010-.013)	1283
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1075
	03-04	.004 (.004-.004)	.004 (.004-.004)	.006 (.005-.007)	.009 (.008-.010)	.011 (.010-.013)	930
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	.003 (.003-.004)	.004 (.003-.004)	.005 (.005-.006)	.008 (.007-.010)	.011 (.009-.013)	943
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	.003 (.003-.004)	.003 (.003-.004)	.005 (.004-.006)	.007 (.006-.008)	.009 (.007-.014)	408
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.004 (.004-.005)	.004 (.004-.005)	.007 (.006-.008)	.012 (.010-.013)	.019 (.013-.022)	463
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	.004 (.003-.004)	.004 (.003-.004)	.005 (.005-.006)	.008 (.007-.009)	.010 (.009-.012)	881

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5',6-Hexachlorobiphenyl (PCB 149) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
60+	05-06	*	*	15	
	07-08	*	*	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
60+	05-06	*	*	17	
	07-08	*	*	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
60+	05-06	*	*	5	
	07-08	*	*	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
60+	05-06	.002	0.001	5	
	07-08	*	*	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5',6-Hexachlorobiphenyl (PCB 149) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,5,5',6-Hexachlorobiphenyl (PCB 151) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2307
	03-04	*	< LOD	.420 (.400-.500)	.700 (.640-.820)	1.00 (.840-1.30)	1870
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	.400 (<LOD-.470)	.700 (.510-.820)	.880 (.700-1.10)	588
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	*	< LOD	.440 (.400-.500)	.720 (.650-.850)	1.02 (.850-1.40)	1282
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1075
	03-04	*	< LOD	.430 (.400-.500)	.720 (.620-.900)	1.10 (.830-1.50)	931
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	*	< LOD	.410 (.400-.500)	.700 (.600-.820)	1.00 (.830-1.20)	939
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	*	< LOD	< LOD	.500 (.400-.700)	.700 (.530-.700)	426
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	*	.400 (<LOD-.500)	.700 (.500-.900)	1.30 (.900-1.90)	2.20 (1.40-4.55)	461
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	*	< LOD	.400 (<LOD-.450)	.650 (.590-.700)	.850 (.750-1.00)	864

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 10.5 and 0.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,5,5',6-Hexachlorobiphenyl (PCB 151) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	.439	0.048	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	.468	0.07	5
		07-08	.580	0.148	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,5,5',6-Hexachlorobiphenyl (PCB 151) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,5,5',6-Hexachlorobiphenyl (PCB 151) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean (95% conf. interval)	Selected percentiles (95% confidence interval)				Sample size
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2307
	03-04	*	< LOD	.003 (.002-.003)	.004 (.004-.005)	.006 (.005-.008)	1870
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	.002 (<LOD-.002)	.003 (.003-.004)	.004 (.003-.006)	588
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	*	< LOD	.003 (.003-.003)	.004 (.004-.006)	.007 (.005-.010)	1282
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1075
	03-04	*	< LOD	.003 (.003-.003)	.005 (.004-.006)	.006 (.005-.010)	931
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	*	< LOD	.003 (.002-.003)	.004 (.004-.005)	.006 (.005-.008)	939
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	*	< LOD	< LOD	.003 (.003-.004)	.004 (.003-.005)	426
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	*	.002 (<LOD-.003)	.004 (.003-.006)	.008 (.006-.010)	.013 (.008-.028)	461
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	*	< LOD	.003 (<LOD-.003)	.004 (.004-.004)	.005 (.004-.006)	864

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,5,5',6-Hexachlorobiphenyl (PCB 151) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	*	*	12
		07-08	*	*	15
	40-59	05-06	*	*	12
		07-08	*	*	16
	60+	05-06	*	*	15
		07-08	*	*	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	*	*	13
		07-08	*	*	17
	60+	05-06	*	*	17
		07-08	*	*	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	*	*	6
		07-08	*	*	6
	40-59	05-06	*	*	5
		07-08	*	*	6
	60+	05-06	.003	0	5
		07-08	*	*	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	*	*	7
		07-08	*	*	8
	60+	05-06	.003	0	5
		07-08	.004	0.001	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,5,5',6-Hexachlorobiphenyl (PCB 151) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	*	*	4
		07-08	*	*	6
60+	05-06	*	*	4	
	07-08	*	*	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	*	*	6
		07-08	*	*	6
	60+	05-06	*	*	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	77.8 (70.2-87.3)	114 (93.0-133)	1926
	01-02	27.2 (24.7-30.1)	30.1 (26.1-34.3)	57.8 (52.1-63.2)	94.7 (86.5-104)	126 (109-142)	2306
	03-04	19.8 (18.8-20.9)	20.8 (18.4-22.2)	43.3 (39.1-46.9)	71.8 (64.4-82.8)	97.1 (88.8-111)	1896
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	668
	01-02	*	< LOD	12.5 (11.1-14.1)	21.2 (17.4-26.7)	31.9 (23.1-64.7)	757
	03-04	5.86 (5.25-6.55)	5.40 (4.70-6.21)	8.50 (7.80-9.85)	15.7 (12.9-18.4)	20.7 (16.9-28.3)	596
20 years and older	99-00	*	< LOD	< LOD	83.2 (75.9-91.8)	122 (100-139)	1258
	01-02	32.6 (29.5-36.1)	35.1 (31.1-39.0)	62.8 (57.6-68.0)	99.5 (90.7-110)	132 (116-146)	1549
	03-04	23.7 (22.3-25.1)	24.2 (21.8-27.4)	47.1 (43.3-50.5)	77.5 (68.0-87.9)	101 (92.9-119)	1300
Gender							
Males	99-00	*	< LOD	< LOD	75.0 (66.7-86.2)	111 (87.7-128)	917
	01-02	28.5 (25.5-32.0)	31.5 (26.7-35.2)	57.7 (48.3-66.2)	97.5 (82.1-110)	126 (104-150)	1074
	03-04	20.0 (18.7-21.3)	19.7 (17.7-21.2)	42.9 (37.4-47.6)	72.7 (60.4-88.8)	107 (86.8-122)	947
Females	99-00	*	< LOD	< LOD	79.0 (70.2-92.0)	119 (91.4-142)	1009
	01-02	26.1 (23.6-28.8)	29.0 (25.1-33.4)	57.9 (52.1-62.9)	94.3 (87.8-98.2)	128 (105-145)	1232
	03-04	19.7 (18.4-21.1)	21.9 (19.0-24.1)	43.8 (39.4-47.7)	70.9 (63.0-81.5)	93.3 (83.8-100)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	67.5 (59.5-71.8)	634
	01-02	12.5 (10.8-14.4)	11.1 (<LOD-13.3)	24.5 (18.2-33.9)	47.4 (36.2-60.3)	66.7 (55.2-72.3)	567
	03-04	8.75 (7.39-10.4)	7.86 (6.17-9.40)	15.6 (11.4-22.2)	30.3 (25.2-34.9)	37.8 (31.1-45.2)	425
Non-Hispanic blacks	99-00	*	< LOD	59.4 (<LOD-82.0)	121 (90.3-159)	176 (130-287)	412
	01-02	30.0 (26.2-34.4)	31.0 (25.8-36.4)	65.1 (54.2-82.7)	127 (97.1-152)	170 (126-246)	515
	03-04	22.8 (19.1-27.2)	20.9 (17.0-28.7)	54.1 (37.3-69.2)	126 (92.9-158)	194 (126-294)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	76.4 (69.3-83.9)	102 (87.8-127)	725
	01-02	29.9 (26.8-33.4)	33.0 (28.7-37.1)	61.2 (55.8-66.7)	96.3 (86.5-109)	126 (104-142)	1061
	03-04	21.3 (19.7-23.1)	22.2 (20.4-25.9)	44.9 (39.7-49.5)	70.9 (60.4-82.1)	91.3 (82.1-103)	885

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 55.6, 10.5, and 1.1, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	6.27	1.11	9
		07-08	5.80	1.04	6
	20-39	05-06	13.2	2.6	12
		07-08	11.0	1	15
	40-59	05-06	36.4	5.1	12
		07-08	28.2	1.6	16
	60+	05-06	60.4	2.7	15
		07-08	63.9	4.8	23
Non-Hispanic White Female	12-19	05-06	4.59	0.49	10
		07-08	4.34	0.99	7
	20-39	05-06	8.47	0.59	16
		07-08	9.36	1.4	13
	40-59	05-06	27.8	1.4	13
		07-08	27.7	1.3	17
	60+	05-06	58.0	5.1	17
		07-08	56.4	3.7	21
Non-Hispanic Black Male	12-19	05-06	7.42	1	13
		07-08	5.74	0.81	6
	20-39	05-06	13.8	2.1	6
		07-08	10.3	0.9	6
	40-59	05-06	38.8	4.2	5
		07-08	41.0	5.5	6
	60+	05-06	103	13	5
		07-08	94.5	11	8
Non-Hispanic Black Female	12-19	05-06	4.20	0.29	14
		07-08	3.76	0.52	5
	20-39	05-06	10.8	1.4	7
		07-08	8.99	1.17	8
	40-59	05-06	41.2	4	7
		07-08	35.7	3.2	8
	60+	05-06	130	33	5
		07-08	102	8	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	3.70	0.45	11
		07-08	3.32	0.69	6
	20-39	05-06	6.93	0.31	9
		07-08	6.85	0.64	9
	40-59	05-06	16.5	0.8	4
		07-08	22.4	2.5	6
60+	05-06	37.3	6.8	4	
	07-08	39.5	4.1	5	
Mexican American Female	12-19	05-06	2.83	0.2	16
		07-08	*	*	5
	20-39	05-06	4.75	0.67	9
		07-08	5.58	0.41	8
	40-59	05-06	19.1	1.5	6
		07-08	20.9	2.8	6
	60+	05-06	36.2	1.6	3
		07-08	40.9	8.5	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	.530 (.490-.560)	.750 (.610-.840)	1926
	01-02	.167 (.151-.185)	.190 (.170-.210)	.380 (.340-.410)	.620 (.560-.690)	.860 (.760-.950)	2306
	03-04	.121 (.114-.128)	.135 (.120-.144)	.283 (.258-.310)	.477 (.439-.518)	.624 (.575-.733)	1896
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	668
	01-02	*	< LOD	.060 (.050-.070)	.110 (.080-.140)	.150 (.110-.310)	757
	03-04	.030 (.027-.033)	.027 (.025-.031)	.044 (.039-.055)	.076 (.062-.098)	.101 (.079-.129)	596
20 years and older	99-00	*	< LOD	< LOD	.560 (.510-.610)	.790 (.670-.880)	1258
	01-02	.206 (.187-.228)	.220 (.200-.250)	.410 (.380-.450)	.670 (.600-.740)	.900 (.820-1.04)	1549
	03-04	.148 (.139-.158)	.156 (.141-.179)	.313 (.283-.339)	.512 (.452-.563)	.671 (.603-.756)	1300
Gender							
Males	99-00	*	< LOD	< LOD	.530 (.470-.560)	.690 (.580-.850)	917
	01-02	.177 (.159-.198)	.200 (.170-.220)	.380 (.340-.430)	.610 (.510-.730)	.850 (.700-1.04)	1074
	03-04	.123 (.115-.133)	.127 (.111-.146)	.277 (.253-.310)	.474 (.413-.522)	.608 (.533-.794)	947
Females	99-00	*	< LOD	< LOD	.530 (.480-.590)	.770 (.610-.880)	1009
	01-02	.158 (.142-.175)	.180 (.150-.210)	.380 (.340-.400)	.630 (.570-.710)	.860 (.760-.950)	1232
	03-04	.119 (.110-.128)	.138 (.113-.149)	.291 (.253-.319)	.492 (.439-.541)	.624 (.578-.689)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	.470 (.380-.540)	634
	01-02	.075 (.063-.089)	.060 (<LOD-.080)	.150 (.120-.210)	.330 (.270-.420)	.470 (.380-.550)	567
	03-04	.053 (.044-.064)	.047 (.038-.057)	.100 (.078-.144)	.205 (.164-.241)	.323 (.213-.435)	425
Non-Hispanic blacks	99-00	*	< LOD	.370 (<LOD-.510)	.750 (.620-.890)	1.27 (.820-1.64)	412
	01-02	.169 (.146-.195)	.180 (.140-.200)	.390 (.330-.490)	.780 (.580-.950)	1.05 (.830-1.43)	515
	03-04	.129 (.108-.156)	.126 (.098-.149)	.330 (.236-.425)	.734 (.562-1.06)	1.26 (.892-1.60)	464
Non-Hispanic whites	99-00	*	< LOD	< LOD	.520 (.480-.560)	.740 (.580-.850)	725
	01-02	.185 (.165-.207)	.210 (.180-.230)	.390 (.370-.430)	.640 (.570-.720)	.840 (.740-.990)	1061
	03-04	.131 (.121-.143)	.144 (.127-.170)	.295 (.268-.329)	.474 (.412-.518)	.600 (.519-.689)	885

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.034	0.006	9
		07-08	.028	0.005	6
	20-39	05-06	.099	0.023	12
		07-08	.074	0.008	15
	40-59	05-06	.279	0.045	12
		07-08	.205	0.011	16
60+	05-06	.409	0.015	15	
	07-08	.407	0.028	23	
Non-Hispanic White Female	12-19	05-06	.025	0.003	10
		07-08	.021	0.005	7
	20-39	05-06	.055	0.004	16
		07-08	.057	0.009	13
	40-59	05-06	.207	0.016	13
		07-08	.184	0.01	17
	60+	05-06	.420	0.033	17
		07-08	.378	0.024	21
Non-Hispanic Black Male	12-19	05-06	.037	0.005	13
		07-08	.027	0.003	6
	20-39	05-06	.082	0.013	6
		07-08	.058	0.006	6
	40-59	05-06	.247	0.027	5
		07-08	.263	0.037	6
	60+	05-06	.614	0.073	5
		07-08	.576	0.073	8
Non-Hispanic Black Female	12-19	05-06	.021	0.002	14
		07-08	.018	0.003	5
	20-39	05-06	.062	0.009	7
		07-08	.045	0.006	8
	40-59	05-06	.264	0.029	7
		07-08	.216	0.023	8
	60+	05-06	.865	0.218	5
		07-08	.614	0.045	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.020	0.002	11
		07-08	.018	0.004	6
	20-39	05-06	.049	0.003	9
		07-08	.047	0.005	9
	40-59	05-06	.134	0.01	4
		07-08	.160	0.02	6
60+	05-06	.268	0.045	4	
	07-08	.281	0.022	5	
Mexican American Female	12-19	05-06	.015	0.001	16
		07-08	*	*	5
	20-39	05-06	.032	0.005	9
		07-08	.033	0.003	8
	40-59	05-06	.140	0.01	6
		07-08	.141	0.018	6
	60+	05-06	.274	0.023	3
		07-08	.283	0.061	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5-Heptachlorobiphenyl (PCB 170) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	23.6 (22.0-25.4)	30.9 (28.1-35.1)	1798
	01-02	*	< LOD	17.5 (15.7-19.3)	26.7 (24.9-29.0)	35.0 (32.4-37.3)	2301
	03-04	5.46 (5.22-5.71)	6.30 (5.96-7.10)	12.9 (12.3-13.9)	21.7 (19.2-23.5)	28.2 (25.8-29.8)	1888
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	645
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	1.12 (.974-1.30)	1.12 (1.00-1.30)	1.90 (1.70-2.50)	4.00 (3.53-4.37)	5.60 (4.35-6.69)	592
20 years and older	99-00	*	< LOD	< LOD	24.9 (23.3-26.4)	33.9 (29.6-37.5)	1153
	01-02	10.5 (<LOD-11.3)	11.1 (<LOD-12.2)	19.1 (17.3-20.9)	28.4 (26.2-30.9)	36.8 (33.4-39.4)	1545
	03-04	6.86 (6.48-7.26)	7.83 (6.87-8.70)	14.2 (13.1-14.9)	22.7 (20.7-24.9)	29.5 (26.7-32.2)	1296
Gender							
Males	99-00	*	< LOD	< LOD	24.1 (22.0-27.0)	32.6 (28.1-37.0)	863
	01-02	*	10.9 (<LOD-11.9)	18.4 (16.3-20.9)	28.1 (24.7-31.9)	36.1 (32.1-38.3)	1073
	03-04	5.81 (5.43-6.22)	6.40 (5.61-7.68)	13.8 (12.4-14.4)	23.4 (20.4-25.6)	29.4 (26.0-32.7)	942
Females	99-00	*	< LOD	< LOD	22.4 (20.7-25.4)	29.7 (25.8-36.3)	935
	01-02	*	< LOD	16.4 (15.0-18.0)	25.9 (24.0-28.4)	34.0 (30.6-38.4)	1228
	03-04	5.14 (4.82-5.48)	6.28 (5.81-7.18)	12.5 (11.5-13.8)	19.2 (17.9-21.7)	26.1 (24.0-29.8)	946
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	23.2 (20.3-26.5)	606
	01-02	*	< LOD	< LOD	15.4 (11.4-18.7)	21.1 (17.1-25.1)	565
	03-04	2.34 (1.91-2.86)	2.20 (1.54-3.20)	4.91 (3.40-8.17)	10.4 (8.40-13.5)	13.9 (10.8-20.1)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	30.9 (23.1-36.3)	41.9 (33.7-59.1)	382
	01-02	*	< LOD	19.0 (14.9-23.2)	32.5 (25.5-43.9)	44.4 (32.5-61.7)	514
	03-04	5.42 (4.63-6.35)	6.28 (5.11-7.75)	14.3 (10.7-18.3)	31.7 (23.1-40.7)	48.5 (32.1-66.1)	461
Non-Hispanic whites	99-00	*	< LOD	< LOD	24.1 (22.1-25.9)	31.0 (26.8-37.3)	658
	01-02	*	10.9 (<LOD-12.0)	18.4 (16.8-20.4)	27.7 (25.4-30.2)	35.0 (32.3-38.1)	1059
	03-04	6.13 (5.74-6.56)	7.40 (6.40-8.18)	13.8 (12.6-14.6)	21.7 (19.2-24.2)	27.0 (24.9-29.5)	882

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 17.2, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5-Heptachlorobiphenyl (PCB 170) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	1.41	0.32	9
		07-08	1.52	0.29	6
	20-39	05-06	3.93	0.94	12
		07-08	3.36	0.28	15
	40-59	05-06	13.7	2.6	12
		07-08	10.3	0.4	16
60+	05-06	20.5	1.3	15	
	07-08	22.4	1.4	23	
Non-Hispanic White Female	12-19	05-06	.896	0.155	10
		07-08	1.14	0.32	7
	20-39	05-06	2.28	0.35	16
		07-08	2.75	0.41	13
	40-59	05-06	8.44	0.46	13
		07-08	9.09	0.45	17
60+	05-06	17.2	1.4	17	
	07-08	18.1	0.9	21	
Non-Hispanic Black Male	12-19	05-06	1.44	0.28	13
		07-08	1.12	0.28	6
	20-39	05-06	3.71	0.52	6
		07-08	2.82	0.3	6
	40-59	05-06	11.4	1	5
		07-08	13.1	1.5	6
60+	05-06	30.9	3.4	5	
	07-08	30.6	3.4	8	
Non-Hispanic Black Female	12-19	05-06	.589	0.099	14
		07-08	.828	0.105	5
	20-39	05-06	2.15	0.52	7
		07-08	2.29	0.35	8
	40-59	05-06	11.1	1.2	7
		07-08	10.8	1	8
60+	05-06	33.3	6.7	5	
	07-08	29.0	1.7	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5-Heptachlorobiphenyl (PCB 170) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.575	0.138	11
		07-08	.891	0.17	6
	20-39	05-06	2.01	0.11	9
		07-08	2.19	0.24	9
	40-59	05-06	6.41	0.43	4
		07-08	8.71	0.91	6
60+	05-06	14.0	3.2	4	
	07-08	16.1	1.9	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	1.39	0.21	9
		07-08	1.62	0.11	8
	40-59	05-06	5.93	0.46	6
		07-08	7.50	1.08	6
	60+	05-06	11.7	0.1	3
		07-08	13.5	2.5	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5-Heptachlorobiphenyl (PCB 170) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	.160 (.140-.180)	.210 (.200-.230)	1798
	01-02	*	< LOD	.110 (.110-.120)	.180 (.170-.200)	.250 (.210-.270)	2301
	03-04	.033 (.032-.035)	.041 (.037-.045)	.087 (.080-.092)	.144 (.134-.149)	.188 (.173-.202)	1888
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	645
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	.006 (.005-.006)	.006 (.005-.007)	.010 (.009-.012)	.020 (.017-.022)	.029 (.022-.033)	592
20 years and older	99-00	*	< LOD	< LOD	.170 (.150-.190)	.220 (.200-.250)	1153
	01-02	.067 (<LOD-.072)	.070 (<LOD-.080)	.120 (.110-.130)	.200 (.180-.210)	.250 (.230-.290)	1545
	03-04	.043 (.040-.046)	.051 (.045-.054)	.093 (.089-.099)	.149 (.141-.160)	.199 (.180-.223)	1296
Gender							
Males	99-00	*	< LOD	< LOD	.170 (.140-.200)	.220 (.200-.250)	863
	01-02	*	.070 (<LOD-.080)	.120 (.110-.130)	.190 (.170-.210)	.250 (.210-.280)	1073
	03-04	.036 (.033-.039)	.043 (.036-.051)	.089 (.080-.093)	.148 (.132-.164)	.200 (.173-.225)	942
Females	99-00	*	< LOD	< LOD	.150 (.130-.170)	.210 (.190-.230)	935
	01-02	*	< LOD	.110 (.100-.120)	.180 (.160-.200)	.250 (.210-.280)	1228
	03-04	.031 (.029-.033)	.040 (.034-.044)	.083 (.075-.092)	.140 (.121-.149)	.176 (.160-.194)	946
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	.170 (.130-.190)	606
	01-02	*	< LOD	< LOD	.110 (.080-.130)	.150 (.120-.190)	565
	03-04	.014 (.012-.018)	.015 (.010-.018)	.032 (.025-.047)	.073 (.054-.113)	.119 (.078-.134)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	.190 (.150-.210)	.260 (.190-.370)	382
	01-02	*	< LOD	.110 (.090-.140)	.210 (.150-.270)	.290 (.210-.390)	514
	03-04	.031 (.026-.037)	.038 (.027-.044)	.087 (.069-.117)	.210 (.136-.280)	.338 (.221-.380)	461
Non-Hispanic whites	99-00	*	< LOD	< LOD	.160 (.140-.200)	.210 (.200-.240)	658
	01-02	*	.070 (<LOD-.080)	.120 (.110-.130)	.190 (.170-.210)	.250 (.210-.280)	1059
	03-04	.038 (.035-.041)	.046 (.042-.052)	.091 (.083-.096)	.145 (.132-.151)	.180 (.164-.207)	882

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5-Heptachlorobiphenyl (PCB 170) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.008	0.002	9
		07-08	.007	0.001	6
	20-39	05-06	.030	0.008	12
		07-08	.022	0.002	15
	40-59	05-06	.105	0.022	12
		07-08	.075	0.003	16
60+	05-06	.139	0.008	15	
	07-08	.143	0.008	23	
Non-Hispanic White Female	12-19	05-06	.005	0.001	10
		07-08	.006	0.002	7
	20-39	05-06	.015	0.002	16
		07-08	.017	0.003	13
	40-59	05-06	.063	0.005	13
		07-08	.060	0.003	17
60+	05-06	.125	0.009	17	
	07-08	.122	0.006	21	
Non-Hispanic Black Male	12-19	05-06	.007	0.001	13
		07-08	.005	0.001	6
	20-39	05-06	.022	0.003	6
		07-08	.016	0.002	6
	40-59	05-06	.072	0.007	5
		07-08	.084	0.009	6
60+	05-06	.185	0.021	5	
	07-08	.186	0.022	8	
Non-Hispanic Black Female	12-19	05-06	.003	0	14
		07-08	.004	0.001	5
	20-39	05-06	.012	0.003	7
		07-08	.011	0.002	8
	40-59	05-06	.071	0.008	7
		07-08	.066	0.007	8
60+	05-06	.222	0.044	5	
	07-08	.175	0.011	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5-Heptachlorobiphenyl (PCB 170) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.003	0.001	11
		07-08	.005	0.001	6
	20-39	05-06	.014	0.001	9
		07-08	.015	0.002	9
	40-59	05-06	.052	0.005	4
		07-08	.062	0.007	6
60+	05-06	.100	0.021	4	
	07-08	.115	0.01	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.009	0.001	9
		07-08	.010	0.001	8
	40-59	05-06	.043	0.003	6
		07-08	.050	0.007	6
	60+	05-06	.089	0.009	3
		07-08	.094	0.018	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5'-Heptachlorobiphenyl (PCB 172) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1901
	01-02	*	< LOD	< LOD	< LOD	< LOD	2199
	03-04	.647 (.606-.691)	.900 (.800-1.00)	1.80 (1.69-1.90)	2.98 (2.70-3.30)	4.16 (3.65-4.55)	1878
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	660
	01-02	*	< LOD	< LOD	< LOD	< LOD	679
	03-04	*	< LOD	< LOD	.610 (.460-.840)	1.10 (.700-1.43)	585
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1241
	01-02	*	< LOD	< LOD	< LOD	< LOD	1520
	03-04	.805 (.746-.869)	1.08 (1.00-1.16)	1.95 (1.80-2.10)	3.20 (2.87-3.60)	4.38 (3.79-5.00)	1293
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	911
	01-02	*	< LOD	< LOD	< LOD	< LOD	1027
	03-04	.668 (.612-.730)	.900 (.800-1.01)	1.82 (1.70-2.00)	3.30 (2.80-3.80)	4.43 (3.76-5.50)	937
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	990
	01-02	*	< LOD	< LOD	< LOD	< LOD	1172
	03-04	.627 (.581-.678)	.900 (.800-1.03)	1.72 (1.51-1.90)	2.80 (2.50-3.11)	3.70 (3.37-4.26)	941
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	630
	01-02	*	< LOD	< LOD	< LOD	< LOD	519
	03-04	*	< LOD	.700 (.470-1.20)	1.40 (1.20-1.95)	2.00 (1.40-2.91)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	494
	03-04	.688 (.579-.818)	.900 (.700-1.10)	2.20 (1.70-2.60)	5.00 (3.38-6.53)	7.60 (5.00-11.3)	458
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	706
	01-02	*	< LOD	< LOD	< LOD	< LOD	1027
	03-04	.720 (.665-.781)	1.00 (.870-1.10)	1.80 (1.69-2.00)	2.94 (2.70-3.20)	3.90 (3.53-4.20)	877

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.5, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5'-Heptachlorobiphenyl (PCB 172) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.480	0.126	12
		07-08	*	*	15
	40-59	05-06	1.57	0.18	12
		07-08	1.29	0.07	16
60+	05-06	2.61	0.16	15	
	07-08	2.98	0.2	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	1.03	0.06	13
		07-08	1.13	0.06	17
60+	05-06	2.19	0.17	17	
	07-08	2.31	0.13	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.404	0.074	6
		07-08	*	*	6
	40-59	05-06	1.57	0.11	5
		07-08	1.85	0.24	6
60+	05-06	4.16	0.33	5	
	07-08	4.64	0.61	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	1.48	0.16	7
		07-08	1.48	0.15	8
60+	05-06	4.61	0.95	5	
	07-08	4.09	0.3	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5'-Heptachlorobiphenyl (PCB 172) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	.808	0.073	4
		07-08	1.32	0.18	6
60+	05-06	1.86	0.45	4	
	07-08	2.34	0.29	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.789	0.059	6
		07-08	.967	0.118	6
	60+	05-06	1.50	0.09	3
		07-08	1.90	0.37	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5'-Heptachlorobiphenyl (PCB 172) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1901
	01-02	*	< LOD	< LOD	< LOD	< LOD	2199
	03-04	.004 (.004-.004)	.006 (.005-.006)	.012 (.011-.012)	.021 (.019-.022)	.027 (.024-.031)	1878
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	660
	01-02	*	< LOD	< LOD	< LOD	< LOD	679
	03-04	*	< LOD	< LOD	.003 (.002-.004)	.005 (.004-.009)	585
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1241
	01-02	*	< LOD	< LOD	< LOD	< LOD	1520
	03-04	.005 (.005-.005)	.007 (.006-.007)	.013 (.012-.014)	.022 (.020-.024)	.028 (.025-.033)	1293
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	911
	01-02	*	< LOD	< LOD	< LOD	< LOD	1027
	03-04	.004 (.004-.005)	.006 (.005-.007)	.012 (.011-.013)	.021 (.018-.024)	.029 (.024-.036)	937
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	990
	01-02	*	< LOD	< LOD	< LOD	< LOD	1172
	03-04	.004 (.003-.004)	.006 (.005-.006)	.012 (.010-.012)	.019 (.017-.021)	.026 (.023-.028)	941
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	630
	01-02	*	< LOD	< LOD	< LOD	< LOD	519
	03-04	*	< LOD	.005 (.003-.007)	.010 (.007-.015)	.017 (.011-.022)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	409
	01-02	*	< LOD	< LOD	< LOD	< LOD	494
	03-04	.004 (.003-.005)	.006 (.004-.006)	.013 (.010-.017)	.030 (.020-.048)	.052 (.028-.067)	458
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	706
	01-02	*	< LOD	< LOD	< LOD	< LOD	1027
	03-04	.004 (.004-.005)	.006 (.005-.007)	.012 (.011-.013)	.020 (.018-.022)	.026 (.023-.029)	877

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5'-Heptachlorobiphenyl (PCB 172) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.004	0.001	12
		07-08	*	*	15
	40-59	05-06	.012	0.002	12
		07-08	.009	0.001	16
	60+	05-06	.018	0.001	15
		07-08	.019	0.001	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	.008	0.001	13
		07-08	.007	0	17
	60+	05-06	.016	0.001	17
		07-08	.015	0.001	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.002	0	6
		07-08	*	*	6
	40-59	05-06	.010	0.001	5
		07-08	.012	0.002	6
	60+	05-06	.025	0.002	5
		07-08	.028	0.004	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	*	*	7
		07-08	*	*	8
	40-59	05-06	.009	0.001	7
		07-08	.009	0.001	8
	60+	05-06	.031	0.006	5
		07-08	.025	0.002	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5'-Heptachlorobiphenyl (PCB 172) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	.007	0.001	4
		07-08	.009	0.001	6
60+	05-06	.013	0.003	4	
	07-08	.017	0.002	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.006	0	6
		07-08	.007	0.001	6
	60+	05-06	.011	0.002	3
		07-08	.013	0.003	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5',6'-Heptachlorobiphenyl (PCB 177) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1873
	01-02	*	< LOD	< LOD	< LOD	< LOD	2287
	03-04	1.13 (1.07-1.20)	1.30 (1.20-1.40)	2.77 (2.50-3.09)	5.30 (4.70-5.90)	7.20 (6.38-8.50)	1882
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	653
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	< LOD	.600 (.500-.720)	1.04 (.800-1.40)	1.70 (1.20-1.95)	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1220
	01-02	*	< LOD	< LOD	< LOD	< LOD	1531
	03-04	1.40 (1.30-1.50)	1.50 (1.40-1.71)	3.10 (2.80-3.44)	5.70 (4.99-6.22)	7.80 (6.50-9.50)	1294
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	887
	01-02	*	< LOD	< LOD	< LOD	< LOD	1065
	03-04	1.12 (1.03-1.21)	1.30 (1.17-1.41)	2.70 (2.40-3.00)	5.30 (4.72-5.80)	7.82 (6.22-9.50)	940
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	986
	01-02	*	< LOD	< LOD	< LOD	< LOD	1222
	03-04	1.15 (1.06-1.23)	1.30 (1.18-1.49)	2.80 (2.57-3.19)	5.30 (4.50-6.06)	6.97 (6.15-7.92)	942
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	622
	01-02	*	< LOD	< LOD	< LOD	< LOD	562
	03-04	.500 (<LOD-.649)	.550 (.400-.700)	1.36 (1.00-1.63)	2.40 (1.86-3.30)	3.40 (2.33-4.63)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	399
	01-02	*	< LOD	< LOD	< LOD	10.6 (<LOD-12.4)	515
	03-04	1.52 (1.25-1.84)	1.56 (1.29-2.05)	4.40 (3.20-5.70)	9.80 (6.30-13.3)	15.8 (10.0-25.7)	458
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	698
	01-02	*	< LOD	< LOD	< LOD	< LOD	1048
	03-04	1.16 (1.08-1.25)	1.34 (1.20-1.50)	2.74 (2.41-3.09)	5.00 (4.57-5.56)	6.50 (5.98-7.40)	880

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.5, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5',6'-Heptachlorobiphenyl (PCB 177) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.606	0.123	12
		07-08	.469	0.065	15
	40-59	05-06	1.59	0.2	12
		07-08	1.26	0.07	16
	60+	05-06	2.92	0.19	15
		07-08	3.26	0.37	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	.369	0.022	16
		07-08	*	*	13
	40-59	05-06	1.24	0.1	13
		07-08	1.16	0.08	17
	60+	05-06	2.65	0.21	17
		07-08	2.81	0.2	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.684	0.098	6
		07-08	.553	0.065	6
	40-59	05-06	2.09	0.29	5
		07-08	2.36	0.33	6
	60+	05-06	5.62	0.59	5
		07-08	5.17	0.63	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.556	0.07	7
		07-08	*	*	8
	40-59	05-06	2.36	0.22	7
		07-08	2.18	0.31	8
	60+	05-06	7.31	1.57	5
		07-08	6.47	0.57	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5',6'-Heptachlorobiphenyl (PCB 177) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.420	0.032	9
		07-08	.360	0.068	9
	40-59	05-06	1.02	0.13	4
		07-08	1.48	0.22	6
60+	05-06	2.10	0.4	4	
	07-08	2.34	0.27	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	1.20	0.08	6
		07-08	1.23	0.15	6
60+	05-06	2.48	0.21	3	
	07-08	2.64	0.53	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5',6'-Heptachlorobiphenyl (PCB 177) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1873
	01-02	*	< LOD	< LOD	< LOD	< LOD	2287
	03-04	.007 (.007-.007)	.008 (.008-.009)	.018 (.016-.021)	.034 (.031-.037)	.047 (.043-.056)	1882
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	653
	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	*	< LOD	.003 (.003-.004)	.006 (.005-.007)	.007 (.006-.011)	588
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1220
	01-02	*	< LOD	< LOD	< LOD	< LOD	1531
	03-04	.009 (.008-.009)	.010 (.009-.011)	.021 (.018-.023)	.036 (.033-.041)	.053 (.045-.058)	1294
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	887
	01-02	*	< LOD	< LOD	< LOD	< LOD	1065
	03-04	.007 (.006-.007)	.009 (.008-.009)	.018 (.016-.021)	.032 (.030-.036)	.045 (.039-.062)	940
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	986
	01-02	*	< LOD	< LOD	< LOD	< LOD	1222
	03-04	.007 (.006-.008)	.008 (.007-.009)	.018 (.017-.021)	.035 (.032-.042)	.051 (.043-.056)	942
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	622
	01-02	*	< LOD	< LOD	< LOD	< LOD	562
	03-04	.003 (<LOD-.004)	.003 (.002-.005)	.008 (.006-.011)	.018 (.012-.025)	.025 (.018-.032)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	399
	01-02	*	< LOD	< LOD	< LOD	.070 (<LOD-.100)	515
	03-04	.009 (.007-.011)	.009 (.007-.012)	.026 (.018-.036)	.057 (.044-.090)	.105 (.058-.143)	458
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	698
	01-02	*	< LOD	< LOD	< LOD	< LOD	1048
	03-04	.007 (.007-.008)	.009 (.008-.010)	.018 (.016-.021)	.032 (.029-.036)	.044 (.040-.052)	880

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5',6'-Heptachlorobiphenyl (PCB 177) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.005	0.001	12
		07-08	.003	0	15
	40-59	05-06	.012	0.002	12
		07-08	.009	0.001	16
	60+	05-06	.020	0.001	15
		07-08	.021	0.002	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	.002	0	16
		07-08	*	*	13
	40-59	05-06	.009	0.001	13
		07-08	.008	0.001	17
	60+	05-06	.019	0.001	17
		07-08	.019	0.001	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.004	0.001	6
		07-08	.003	0	6
	40-59	05-06	.013	0.002	5
		07-08	.015	0.002	6
	60+	05-06	.033	0.003	5
		07-08	.032	0.004	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.003	0	7
		07-08	*	*	8
	40-59	05-06	.015	0.001	7
		07-08	.013	0.002	8
	60+	05-06	.049	0.011	5
		07-08	.039	0.004	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5',6'-Heptachlorobiphenyl (PCB 177) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.003	0	9
		07-08	.002	0	9
	40-59	05-06	.008	0.001	4
		07-08	.010	0.002	6
60+	05-06	.015	0.003	4	
	07-08	.016	0.002	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.009	0	6
		07-08	.008	0.001	6
	60+	05-06	.019	0.003	3
		07-08	.018	0.004	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',5,5',6-Heptachlorobiphenyl (PCB 178) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1932
	01-02	*	< LOD	< LOD	< LOD	< LOD	2299
	03-04	.933 (.894-.974)	1.20 (1.10-1.30)	2.50 (2.30-2.67)	4.24 (3.90-4.47)	6.10 (5.13-7.10)	1887
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	669
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	.400 (<LOD-.480)	.850 (.620-1.04)	1.12 (1.00-1.32)	590
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1263
	01-02	*	< LOD	< LOD	< LOD	< LOD	1541
	03-04	1.18 (1.11-1.25)	1.46 (1.30-1.60)	2.74 (2.60-2.90)	4.41 (4.10-4.80)	6.50 (5.69-7.58)	1297
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	919
	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	.991 (.918-1.07)	1.20 (1.10-1.37)	2.66 (2.45-2.80)	4.50 (3.80-5.10)	6.40 (5.30-7.70)	940
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1013
	01-02	*	< LOD	< LOD	< LOD	< LOD	1230
	03-04	.881 (.835-.930)	1.20 (1.10-1.30)	2.30 (2.20-2.59)	4.01 (3.71-4.34)	5.40 (4.47-6.70)	947
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	635
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	*	< LOD	.900 (.600-1.34)	1.99 (1.37-2.60)	3.10 (2.00-3.60)	424
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	415
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	1.09 (.911-1.30)	1.30 (.910-1.65)	3.10 (2.30-4.21)	7.10 (5.13-10.1)	11.4 (6.80-16.5)	460
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	724
	01-02	*	< LOD	< LOD	< LOD	< LOD	1054
	03-04	1.03 (.963-1.10)	1.30 (1.19-1.50)	2.61 (2.40-2.80)	4.20 (3.80-4.40)	5.33 (4.58-6.30)	882

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.4, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',5,5',6-Heptachlorobiphenyl (PCB 178) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.735	0.151	12
		07-08	.591	0.067	15
	40-59	05-06	2.39	0.22	12
		07-08	1.81	0.09	16
60+	05-06	4.18	0.26	15	
	07-08	4.07	0.3	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	.460	0.036	16
		07-08	*	*	13
	40-59	05-06	1.59	0.12	13
		07-08	1.49	0.08	17
	60+	05-06	3.55	0.29	17
		07-08	3.44	0.17	21
Non-Hispanic Black Male	12-19	05-06	.351	0.049	13
		07-08	*	*	6
	20-39	05-06	.854	0.136	6
		07-08	.564	0.098	6
	40-59	05-06	2.78	0.15	5
		07-08	2.85	0.3	6
	60+	05-06	7.22	0.75	5
		07-08	6.82	0.77	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.545	0.092	7
		07-08	*	*	8
	40-59	05-06	2.39	0.27	7
		07-08	2.11	0.23	8
	60+	05-06	7.83	1.63	5
		07-08	6.33	0.56	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',5,5',6-Heptachlorobiphenyl (PCB 178) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.382	0.039	9
		07-08	*	*	9
	40-59	05-06	1.14	0.08	4
		07-08	1.58	0.21	6
60+	05-06	2.79	0.67	4	
	07-08	3.23	0.45	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	1.12	0.08	6
		07-08	1.37	0.23	6
60+	05-06	2.42	0.2	3	
	07-08	3.00	0.59	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',5,5',6-Heptachlorobiphenyl (PCB 178) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1932
	01-02	*	< LOD	< LOD	< LOD	< LOD	2299
	03-04	.006 (.005-.006)	.008 (.007-.008)	.016 (.015-.018)	.029 (.026-.032)	.041 (.035-.045)	1887
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	669
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	*	< LOD	.002 (<LOD-.002)	.004 (.003-.005)	.006 (.004-.007)	590
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1263
	01-02	*	< LOD	< LOD	< LOD	< LOD	1541
	03-04	.007 (.007-.008)	.009 (.008-.010)	.018 (.017-.019)	.031 (.028-.034)	.043 (.038-.048)	1297
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	919
	01-02	*	< LOD	< LOD	< LOD	< LOD	1069
	03-04	.006 (.006-.007)	.008 (.007-.009)	.017 (.016-.018)	.030 (.025-.034)	.042 (.034-.050)	940
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1013
	01-02	*	< LOD	< LOD	< LOD	< LOD	1230
	03-04	.005 (.005-.006)	.008 (.006-.008)	.016 (.014-.017)	.028 (.025-.031)	.040 (.034-.045)	947
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	635
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	*	< LOD	.005 (.004-.008)	.013 (.009-.020)	.022 (.014-.028)	424
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	< LOD	415
	01-02	*	< LOD	< LOD	< LOD	< LOD	515
	03-04	.006 (.005-.008)	.008 (.006-.008)	.019 (.016-.027)	.044 (.032-.061)	.077 (.045-.089)	460
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	724
	01-02	*	< LOD	< LOD	< LOD	< LOD	1054
	03-04	.006 (.006-.007)	.008 (.007-.010)	.017 (.015-.018)	.028 (.025-.031)	.037 (.033-.045)	882

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',5,5',6-Heptachlorobiphenyl (PCB 178) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.006	0.001	12
		07-08	.004	0	15
	40-59	05-06	.018	0.002	12
		07-08	.013	0.001	16
60+	05-06	.028	0.002	15	
	07-08	.026	0.002	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	.003	0	16
		07-08	*	*	13
	40-59	05-06	.012	0.001	13
		07-08	.010	0.001	17
	60+	05-06	.026	0.002	17
		07-08	.023	0.001	21
Non-Hispanic Black Male	12-19	05-06	.002	0	13
		07-08	*	*	6
	20-39	05-06	.005	0.001	6
		07-08	.003	0.001	6
	40-59	05-06	.018	0.001	5
		07-08	.018	0.002	6
	60+	05-06	.043	0.005	5
		07-08	.041	0.005	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.003	0.001	7
		07-08	*	*	8
	40-59	05-06	.015	0.002	7
		07-08	.013	0.002	8
	60+	05-06	.052	0.011	5
		07-08	.038	0.004	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',5,5',6-Heptachlorobiphenyl (PCB 178) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.003	0	9
		07-08	*	*	9
	40-59	05-06	.009	0.001	4
		07-08	.011	0.001	6
60+	05-06	.020	0.004	4	
	07-08	.023	0.002	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.008	0	6
		07-08	.009	0.002	6
	60+	05-06	.019	0.003	3
		07-08	.021	0.005	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	37.5 (32.7-41.8)	62.0 (56.6-66.6)	79.3 (72.1-89.2)	1924
	01-02	19.2 (17.4-21.1)	21.8 (19.0-24.6)	42.2 (38.3-47.5)	69.7 (63.7-75.9)	87.3 (83.3-93.0)	2302
	03-04	15.1 (14.5-15.7)	18.0 (16.4-18.9)	37.1 (34.9-39.5)	63.7 (56.2-68.2)	81.5 (75.8-89.9)	1896
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	12.2 (<LOD-14.8)	21.3 (14.2-26.0)	755
	03-04	3.06 (2.65-3.54)	2.97 (2.57-3.41)	5.00 (4.20-5.80)	11.1 (8.70-13.0)	14.7 (13.0-18.8)	598
20 years and older	99-00	*	< LOD	41.1 (37.6-45.2)	65.6 (60.6-69.4)	84.8 (75.6-96.1)	1257
	01-02	23.0 (20.8-25.5)	26.5 (22.7-28.6)	46.7 (41.4-51.3)	74.5 (66.7-79.8)	91.4 (85.0-99.5)	1547
	03-04	19.0 (17.9-20.1)	21.5 (18.9-24.3)	40.4 (37.9-42.8)	66.7 (61.0-70.1)	88.0 (77.8-96.7)	1298
Gender							
Males	99-00	*	< LOD	40.6 (34.5-45.0)	65.1 (58.6-71.4)	83.8 (75.8-96.3)	919
	01-02	21.1 (18.8-23.7)	25.1 (21.2-30.0)	46.7 (39.8-51.8)	73.8 (63.2-79.8)	86.9 (78.0-99.4)	1073
	03-04	16.1 (14.8-17.4)	18.5 (15.8-21.3)	39.6 (35.8-41.7)	68.6 (63.7-75.0)	88.3 (76.8-96.7)	947
Females	99-00	*	< LOD	34.4 (29.8-39.3)	56.7 (52.2-62.6)	74.6 (66.6-90.5)	1005
	01-02	17.5 (15.9-19.3)	18.5 (16.0-21.6)	39.7 (34.8-43.9)	64.3 (58.0-74.8)	87.9 (79.1-98.1)	1229
	03-04	14.2 (13.4-15.0)	17.8 (15.8-19.2)	35.4 (31.8-38.5)	55.7 (51.2-59.9)	74.3 (64.8-87.6)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	41.7 (33.2-50.5)	57.8 (49.7-63.8)	633
	01-02	*	< LOD	18.0 (11.5-22.7)	36.9 (28.2-45.7)	54.2 (42.2-60.0)	566
	03-04	6.14 (4.91-7.67)	5.80 (3.80-7.50)	12.6 (8.97-21.5)	27.2 (22.3-35.4)	38.8 (27.2-60.7)	427
Non-Hispanic blacks	99-00	*	< LOD	39.3 (32.2-48.4)	78.4 (64.3-93.7)	117 (89.6-144)	414
	01-02	19.5 (16.5-23.1)	21.0 (16.9-23.8)	48.4 (36.7-57.9)	90.5 (73.7-101)	116 (96.1-167)	514
	03-04	15.1 (12.8-17.7)	18.1 (12.6-20.8)	39.0 (29.7-51.1)	89.9 (64.4-122)	137 (91.4-206)	464
Non-Hispanic whites	99-00	*	< LOD	39.9 (34.5-45.3)	62.3 (56.5-68.6)	79.0 (71.3-91.4)	719
	01-02	21.4 (19.1-23.9)	24.8 (21.5-28.1)	45.6 (40.2-51.0)	72.4 (63.8-78.0)	87.9 (83.8-94.3)	1059
	03-04	16.9 (15.9-18.0)	19.8 (18.4-23.1)	39.3 (35.8-42.0)	63.7 (55.0-69.1)	79.8 (70.1-89.3)	883

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 28.2, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	4.20	0.71	9
		07-08	3.50	0.65	6
	20-39	05-06	10.7	2.6	12
		07-08	8.43	0.64	15
	40-59	05-06	37.8	6.8	12
		07-08	25.8	1	16
	60+	05-06	58.6	3.4	15
		07-08	57.6	3.4	23
Non-Hispanic White Female	12-19	05-06	2.70	0.27	10
		07-08	2.82	0.82	7
	20-39	05-06	6.45	0.74	16
		07-08	6.65	0.93	13
	40-59	05-06	22.4	1.2	13
		07-08	22.3	1.1	17
	60+	05-06	48.3	3.7	17
		07-08	45.7	2.2	21
Non-Hispanic Black Male	12-19	05-06	4.06	0.56	13
		07-08	3.54	0.68	6
	20-39	05-06	10.5	1.3	6
		07-08	7.36	0.82	6
	40-59	05-06	32.8	2.3	5
		07-08	34.2	3.9	6
	60+	05-06	88.0	8.6	5
		07-08	82.5	9.9	8
Non-Hispanic Black Female	12-19	05-06	2.10	0.16	14
		07-08	2.02	0.3	5
	20-39	05-06	6.93	1.06	7
		07-08	5.69	0.88	8
	40-59	05-06	30.5	3.1	7
		07-08	26.4	2.5	8
	60+	05-06	88.6	17.3	5
		07-08	73.1	4.2	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	2.04	0.29	11
		07-08	2.24	0.37	6
	20-39	05-06	5.12	0.28	9
		07-08	5.47	0.6	9
	40-59	05-06	15.9	1.3	4
		07-08	20.9	2.6	6
60+	05-06	39.3	10.1	4	
	07-08	42.0	5.5	5	
Mexican American Female	12-19	05-06	1.78	0.18	16
		07-08	*	*	5
	20-39	05-06	3.45	0.6	9
		07-08	3.88	0.21	8
	40-59	05-06	15.5	1.4	6
		07-08	18.2	2.6	6
	60+	05-06	30.9	0.3	3
		07-08	33.7	6.7	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	.250 (.220-.270)	.410 (.370-.450)	.540 (.490-.600)	1924
	01-02	.118 (.106-.130)	.140 (.120-.160)	.280 (.250-.300)	.460 (.400-.510)	.610 (.540-.690)	2302
	03-04	.092 (.088-.096)	.114 (.102-.127)	.246 (.230-.253)	.409 (.386-.438)	.534 (.495-.598)	1896
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	.060 (<LOD-.080)	.090 (.070-.140)	755
	03-04	.016 (.014-.018)	.015 (.013-.017)	.025 (.022-.030)	.050 (.046-.056)	.076 (.056-.096)	598
20 years and older	99-00	*	< LOD	.270 (.240-.300)	.440 (.410-.480)	.560 (.500-.620)	1257
	01-02	.146 (.131-.162)	.170 (.150-.190)	.300 (.280-.340)	.490 (.430-.560)	.640 (.570-.740)	1547
	03-04	.119 (.112-.126)	.138 (.125-.151)	.270 (.249-.284)	.433 (.402-.474)	.572 (.511-.635)	1298
Gender							
Males	99-00	*	< LOD	.260 (.220-.300)	.460 (.370-.510)	.570 (.500-.650)	919
	01-02	.131 (.117-.147)	.170 (.140-.200)	.300 (.260-.340)	.480 (.410-.550)	.620 (.530-.720)	1073
	03-04	.099 (.091-.108)	.117 (.098-.148)	.250 (.231-.274)	.422 (.388-.495)	.596 (.499-.654)	947
Females	99-00	*	< LOD	.230 (.200-.270)	.380 (.340-.410)	.510 (.430-.560)	1005
	01-02	.106 (.096-.117)	.120 (.100-.140)	.260 (.230-.280)	.420 (.380-.470)	.610 (.520-.700)	1229
	03-04	.085 (.080-.092)	.110 (.094-.123)	.230 (.209-.252)	.386 (.349-.421)	.494 (.459-.541)	949
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	.280 (.200-.320)	.390 (.320-.450)	633
	01-02	*	< LOD	.110 (.080-.160)	.260 (.200-.310)	.360 (.280-.420)	566
	03-04	.037 (.030-.047)	.035 (.024-.051)	.086 (.059-.118)	.191 (.138-.313)	.343 (.191-.405)	427
Non-Hispanic blacks	99-00	*	< LOD	.250 (.190-.300)	.500 (.350-.660)	.730 (.530-1.04)	414
	01-02	.110 (.092-.131)	.120 (.090-.140)	.290 (.220-.350)	.580 (.440-.650)	.720 (.520-1.01)	514
	03-04	.086 (.072-.102)	.098 (.070-.126)	.248 (.194-.327)	.580 (.373-.824)	.919 (.592-1.12)	464
Non-Hispanic whites	99-00	*	< LOD	.260 (.240-.300)	.430 (.380-.470)	.540 (.480-.600)	719
	01-02	.132 (.118-.148)	.160 (.140-.180)	.290 (.260-.330)	.470 (.410-.530)	.620 (.530-.730)	1059
	03-04	.104 (.097-.112)	.132 (.115-.151)	.253 (.235-.280)	.409 (.382-.449)	.512 (.478-.572)	883

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.023	0.004	9
		07-08	.017	0.003	6
	20-39	05-06	.081	0.023	12
		07-08	.056	0.005	15
	40-59	05-06	.288	0.058	12
		07-08	.188	0.008	16
60+	05-06	.397	0.02	15	
	07-08	.369	0.021	23	
Non-Hispanic White Female	12-19	05-06	.014	0.002	10
		07-08	.014	0.004	7
	20-39	05-06	.042	0.005	16
		07-08	.040	0.006	13
	40-59	05-06	.167	0.014	13
		07-08	.147	0.008	17
	60+	05-06	.350	0.025	17
		07-08	.306	0.015	21
Non-Hispanic Black Male	12-19	05-06	.020	0.003	13
		07-08	.017	0.003	6
	20-39	05-06	.062	0.008	6
		07-08	.041	0.004	6
	40-59	05-06	.209	0.015	5
		07-08	.219	0.025	6
	60+	05-06	.527	0.054	5
		07-08	.501	0.063	8
Non-Hispanic Black Female	12-19	05-06	.010	0.001	14
		07-08	.010	0.002	5
	20-39	05-06	.040	0.006	7
		07-08	.029	0.005	8
	40-59	05-06	.196	0.021	7
		07-08	.160	0.018	8
	60+	05-06	.589	0.113	5
		07-08	.439	0.028	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.011	0.001	11
		07-08	.012	0.002	6
	20-39	05-06	.036	0.002	9
		07-08	.037	0.004	9
	40-59	05-06	.130	0.015	4
		07-08	.148	0.018	6
60+	05-06	.281	0.066	4	
	07-08	.298	0.029	5	
Mexican American Female	12-19	05-06	.010	0.001	16
		07-08	*	*	5
	20-39	05-06	.023	0.004	9
		07-08	.023	0.002	8
	40-59	05-06	.113	0.009	6
		07-08	.122	0.017	6
	60+	05-06	.236	0.025	3
		07-08	.233	0.049	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5',6-Heptachlorobiphenyl (PCB 183) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1928
	01-02	*	< LOD	< LOD	< LOD	< LOD	2306
	03-04	1.45 (1.38-1.54)	1.60 (1.50-1.77)	3.29 (3.15-3.54)	5.86 (5.50-6.29)	7.90 (7.50-8.74)	1886
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	668
	01-02	*	< LOD	< LOD	< LOD	< LOD	757
	03-04	*	.440 (<LOD-.500)	.730 (.660-.840)	1.30 (.990-1.62)	1.91 (1.50-2.24)	590
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1260
	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	1.77 (1.67-1.88)	1.88 (1.73-2.09)	3.70 (3.34-4.00)	6.25 (5.71-6.83)	8.40 (7.70-9.55)	1296
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	919
	01-02	*	< LOD	< LOD	< LOD	< LOD	1074
	03-04	1.47 (1.34-1.61)	1.54 (1.40-1.77)	3.29 (3.10-3.54)	5.80 (5.30-6.50)	7.90 (7.20-9.20)	940
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1009
	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	1.44 (1.34-1.55)	1.69 (1.50-1.82)	3.30 (3.05-3.80)	5.88 (5.21-6.30)	7.90 (7.31-9.24)	946
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	635
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	.703 (.579-.854)	.700 (.600-.890)	1.40 (1.19-1.90)	2.62 (2.20-3.47)	3.73 (2.60-5.60)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	12.7 (<LOD-23.1)	413
	01-02	*	< LOD	< LOD	< LOD	13.0 (<LOD-14.9)	514
	03-04	1.64 (1.41-1.90)	1.60 (1.20-2.00)	4.31 (3.25-5.63)	11.0 (6.69-13.7)	14.8 (11.3-17.4)	459
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	722
	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	1.56 (1.43-1.69)	1.74 (1.55-1.90)	3.30 (3.14-3.73)	5.71 (5.20-6.29)	7.59 (6.70-8.04)	883

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.4, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5',6-Heptachlorobiphenyl (PCB 183) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.443	0.059	9
		07-08	*	*	6
	20-39	05-06	1.17	0.31	12
		07-08	.964	0.069	15
	40-59	05-06	3.14	0.76	12
		07-08	2.15	0.08	16
	60+	05-06	4.82	0.23	15
		07-08	5.17	0.41	23
Non-Hispanic White Female	12-19	05-06	.311	0.03	10
		07-08	*	*	7
	20-39	05-06	.694	0.043	16
		07-08	.839	0.194	13
	40-59	05-06	2.04	0.11	13
		07-08	2.08	0.12	17
	60+	05-06	4.13	0.34	17
		07-08	4.20	0.26	21
Non-Hispanic Black Male	12-19	05-06	.598	0.087	13
		07-08	*	*	6
	20-39	05-06	1.10	0.13	6
		07-08	.927	0.087	6
	40-59	05-06	3.05	0.33	5
		07-08	3.92	0.7	6
	60+	05-06	7.74	0.74	5
		07-08	8.16	1.05	8
Non-Hispanic Black Female	12-19	05-06	.335	0.035	14
		07-08	*	*	5
	20-39	05-06	.861	0.111	7
		07-08	.801	0.164	8
	40-59	05-06	3.48	0.24	7
		07-08	3.14	0.33	8
	60+	05-06	9.38	2.28	5
		07-08	9.22	0.69	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5',6-Heptachlorobiphenyl (PCB 183) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.686	0.053	9
		07-08	.781	0.056	9
	40-59	05-06	1.52	0.12	4
		07-08	2.22	0.34	6
60+	05-06	3.05	0.58	4	
	07-08	3.84	0.67	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.476	0.076	9
		07-08	.475	0.087	8
	40-59	05-06	1.86	0.13	6
		07-08	2.09	0.25	6
60+	05-06	3.36	0.07	3	
	07-08	4.27	1.05	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5',6-Heptachlorobiphenyl (PCB 183) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	< LOD	< LOD	1928
	01-02	*	< LOD	< LOD	< LOD	< LOD	2306
	03-04	.009 (.008-.009)	.010 (.009-.011)	.021 (.019-.024)	.039 (.036-.042)	.054 (.048-.059)	1886
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	668
	01-02	*	< LOD	< LOD	< LOD	< LOD	757
	03-04	*	.002 (<LOD-.003)	.004 (.004-.004)	.007 (.005-.009)	.009 (.007-.010)	590
20 years and older	99-00	*	< LOD	< LOD	< LOD	< LOD	1260
	01-02	*	< LOD	< LOD	< LOD	< LOD	1549
	03-04	.011 (.010-.012)	.012 (.011-.014)	.024 (.022-.027)	.042 (.038-.045)	.057 (.052-.064)	1296
Gender							
Males	99-00	*	< LOD	< LOD	< LOD	< LOD	919
	01-02	*	< LOD	< LOD	< LOD	< LOD	1074
	03-04	.009 (.008-.010)	.010 (.009-.011)	.022 (.019-.025)	.038 (.035-.042)	.052 (.043-.064)	940
Females	99-00	*	< LOD	< LOD	< LOD	< LOD	1009
	01-02	*	< LOD	< LOD	< LOD	< LOD	1232
	03-04	.009 (.008-.009)	.010 (.009-.012)	.021 (.019-.023)	.042 (.036-.044)	.055 (.049-.061)	946
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	< LOD	635
	01-02	*	< LOD	< LOD	< LOD	< LOD	567
	03-04	.004 (.003-.005)	.005 (.004-.006)	.009 (.007-.013)	.019 (.013-.029)	.029 (.017-.044)	423
Non-Hispanic blacks	99-00	*	< LOD	< LOD	< LOD	.070 (<LOD-.150)	413
	01-02	*	< LOD	< LOD	< LOD	.080 (<LOD-.110)	514
	03-04	.009 (.008-.011)	.009 (.008-.012)	.027 (.019-.037)	.065 (.043-.084)	.090 (.058-.120)	459
Non-Hispanic whites	99-00	*	< LOD	< LOD	< LOD	< LOD	722
	01-02	*	< LOD	< LOD	< LOD	< LOD	1061
	03-04	.010 (.009-.010)	.011 (.010-.012)	.022 (.019-.025)	.038 (.035-.042)	.049 (.046-.055)	883

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5',6-Heptachlorobiphenyl (PCB 183) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.002	0	9
		07-08	*	*	6
	20-39	05-06	.009	0.003	12
		07-08	.006	0.001	15
	40-59	05-06	.024	0.006	12
		07-08	.016	0.001	16
	60+	05-06	.033	0.001	15
		07-08	.033	0.003	23
Non-Hispanic White Female	12-19	05-06	.002	0	10
		07-08	*	*	7
	20-39	05-06	.005	0	16
		07-08	.005	0.001	13
	40-59	05-06	.015	0.001	13
		07-08	.014	0.001	17
	60+	05-06	.030	0.002	17
		07-08	.028	0.002	21
Non-Hispanic Black Male	12-19	05-06	.003	0	13
		07-08	*	*	6
	20-39	05-06	.007	0.001	6
		07-08	.005	0.001	6
	40-59	05-06	.019	0.002	5
		07-08	.025	0.005	6
	60+	05-06	.046	0.004	5
		07-08	.050	0.007	8
Non-Hispanic Black Female	12-19	05-06	.002	0	14
		07-08	*	*	5
	20-39	05-06	.005	0.001	7
		07-08	.004	0.001	8
	40-59	05-06	.022	0.002	7
		07-08	.019	0.002	8
	60+	05-06	.063	0.015	5
		07-08	.055	0.004	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4,4',5',6-Heptachlorobiphenyl (PCB 183) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.005	0	9
		07-08	.005	0	9
	40-59	05-06	.012	0.001	4
		07-08	.016	0.002	6
60+	05-06	.022	0.004	4	
	07-08	.027	0.004	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.003	0.001	9
		07-08	.003	0	8
	40-59	05-06	.014	0.001	6
		07-08	.014	0.002	6
	60+	05-06	.026	0.003	3
		07-08	.030	0.008	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5',6-Heptachlorobiphenyl (PCB 187) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	17.7 (15.6-20.4)	24.8 (22.0-27.7)	1930
	01-02	*	< LOD	12.2 (10.6-13.8)	21.7 (19.6-23.7)	28.1 (26.8-29.7)	2307
	03-04	4.23 (3.96-4.50)	4.60 (4.20-5.23)	10.1 (9.42-10.5)	17.2 (16.1-18.1)	24.3 (20.5-27.8)	1889
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	1.00 (.859-1.17)	1.00 (.880-1.10)	1.66 (1.50-1.99)	3.20 (2.50-4.34)	5.00 (3.70-6.07)	594
20 years and older	99-00	*	< LOD	< LOD	19.3 (16.8-21.8)	26.0 (24.1-29.0)	1263
	01-02	*	< LOD	13.3 (12.0-15.2)	23.0 (20.9-24.7)	29.2 (27.3-32.0)	1549
	03-04	5.20 (4.83-5.60)	5.71 (5.01-6.19)	10.9 (10.3-11.7)	18.1 (16.9-19.5)	25.9 (22.1-32.1)	1295
Gender							
Males	99-00	*	< LOD	< LOD	17.8 (15.8-19.6)	25.9 (21.6-29.8)	917
	01-02	*	< LOD	12.7 (<LOD-15.8)	22.5 (19.4-24.1)	27.7 (25.2-30.3)	1075
	03-04	4.34 (3.97-4.75)	4.50 (4.00-5.40)	10.4 (9.21-10.9)	17.9 (16.1-19.5)	25.1 (20.4-30.9)	939
Females	99-00	*	< LOD	< LOD	17.7 (15.1-21.7)	24.2 (21.6-27.0)	1013
	01-02	*	< LOD	11.9 (<LOD-13.0)	21.3 (18.5-23.5)	28.5 (26.2-31.3)	1232
	03-04	4.12 (3.84-4.41)	4.67 (4.10-5.30)	9.85 (8.83-10.5)	16.4 (15.3-17.9)	24.2 (19.1-27.8)	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	17.6 (15.1-19.1)	636
	01-02	*	< LOD	< LOD	10.8 (<LOD-13.4)	15.7 (12.7-18.2)	567
	03-04	1.79 (1.47-2.20)	1.58 (1.26-2.00)	3.96 (2.70-5.90)	7.86 (5.96-10.3)	11.6 (9.41-14.9)	426
Non-Hispanic blacks	99-00	*	< LOD	15.5 (<LOD-21.1)	31.6 (26.0-39.0)	47.1 (36.6-72.1)	412
	01-02	*	< LOD	16.5 (12.3-21.0)	30.3 (22.9-40.6)	44.8 (30.3-61.9)	515
	03-04	5.09 (4.39-5.90)	5.40 (4.20-6.21)	13.9 (10.6-17.7)	31.7 (22.7-46.4)	51.5 (31.7-76.5)	458
Non-Hispanic whites	99-00	*	< LOD	< LOD	16.8 (14.9-18.6)	22.0 (19.1-25.2)	727
	01-02	*	< LOD	12.5 (10.8-14.6)	21.7 (18.9-23.7)	27.4 (25.2-29.5)	1061
	03-04	4.47 (4.17-4.79)	5.00 (4.40-5.70)	10.1 (9.26-10.8)	16.5 (15.3-17.9)	21.1 (19.1-24.2)	884

Limit of detection (LOD, see Data Analysis section) for Survey years 99-00, 01-02, and 03-04 are 12.4, 10.5, and 0.4, respectively.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5',6-Heptachlorobiphenyl (PCB 187) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	1.15	0.17	9
		07-08	.986	0.243	6
	20-39	05-06	3.08	0.71	12
		07-08	2.65	0.22	15
	40-59	05-06	8.32	0.66	12
		07-08	6.81	0.33	16
	60+	05-06	16.6	1	15
		07-08	16.8	1.3	23
Non-Hispanic White Female	12-19	05-06	.832	0.069	10
		07-08	*	*	7
	20-39	05-06	1.77	0.13	16
		07-08	2.19	0.54	13
	40-59	05-06	6.07	0.42	13
		07-08	5.76	0.32	17
	60+	05-06	14.0	1.1	17
		07-08	13.8	0.8	21
Non-Hispanic Black Male	12-19	05-06	1.68	0.29	13
		07-08	1.56	0.3	6
	20-39	05-06	3.61	0.61	6
		07-08	2.73	0.31	6
	40-59	05-06	12.3	1	5
		07-08	12.8	1.9	6
	60+	05-06	32.1	3.5	5
		07-08	31.5	4.1	8
Non-Hispanic Black Female	12-19	05-06	.956	0.08	14
		07-08	.872	0.162	5
	20-39	05-06	2.57	0.42	7
		07-08	2.20	0.31	8
	40-59	05-06	11.3	1.1	7
		07-08	9.55	1.07	8
	60+	05-06	37.6	9.4	5
		07-08	32.7	3.2	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5',6-Heptachlorobiphenyl (PCB 187) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.633	0.083	11
		07-08	.656	0.133	6
	20-39	05-06	1.72	0.15	9
		07-08	1.92	0.22	9
	40-59	05-06	4.78	0.5	4
		07-08	6.80	0.92	6
60+	05-06	10.6	2.5	4	
	07-08	13.1	1.9	5	
Mexican American Female	12-19	05-06	.615	0.063	16
		07-08	*	*	5
	20-39	05-06	1.05	0.2	9
		07-08	1.40	0.13	8
	40-59	05-06	4.80	0.3	6
		07-08	5.66	0.8	6
	60+	05-06	10.4	0.6	3
		07-08	12.7	2.8	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5',6-Heptachlorobiphenyl (PCB 187) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	99-00	*	< LOD	< LOD	.120 (.110-.130)	.170 (.140-.180)	1930
	01-02	*	< LOD	.080 (.070-.090)	.140 (.130-.160)	.200 (.180-.220)	2307
	03-04	.026 (.024-.028)	.029 (.027-.032)	.065 (.061-.069)	.115 (.106-.125)	.167 (.139-.197)	1889
Age group							
12-19 years	99-00	*	< LOD	< LOD	< LOD	< LOD	667
	01-02	*	< LOD	< LOD	< LOD	< LOD	758
	03-04	.005 (.004-.006)	.005 (.004-.006)	.009 (.007-.010)	.016 (.013-.019)	.023 (.016-.029)	594
20 years and older	99-00	*	< LOD	< LOD	.130 (.110-.140)	.180 (.160-.190)	1263
	01-02	*	< LOD	.090 (.080-.100)	.150 (.130-.180)	.210 (.190-.230)	1549
	03-04	.033 (.030-.035)	.036 (.032-.040)	.071 (.067-.075)	.124 (.112-.139)	.172 (.143-.215)	1295
Gender							
Males	99-00	*	< LOD	< LOD	.120 (.100-.140)	.180 (.140-.200)	917
	01-02	*	< LOD	.080 (<LOD-.100)	.150 (.130-.160)	.190 (.170-.230)	1075
	03-04	.027 (.024-.030)	.030 (.026-.035)	.067 (.061-.072)	.116 (.102-.137)	.160 (.137-.215)	939
Females	99-00	*	< LOD	< LOD	.120 (.110-.130)	.160 (.140-.180)	1013
	01-02	*	< LOD	.080 (<LOD-.090)	.140 (.120-.150)	.200 (.180-.220)	1232
	03-04	.025 (.023-.027)	.029 (.026-.033)	.063 (.056-.070)	.115 (.104-.125)	.168 (.141-.191)	950
Race/ethnicity							
Mexican Americans	99-00	*	< LOD	< LOD	< LOD	.120 (.100-.130)	636
	01-02	*	< LOD	< LOD	.080 (<LOD-.100)	.100 (.090-.120)	567
	03-04	.011 (.009-.013)	.010 (.007-.015)	.025 (.018-.038)	.057 (.042-.079)	.094 (.059-.110)	426
Non-Hispanic blacks	99-00	*	< LOD	.090 (<LOD-.130)	.200 (.160-.230)	.290 (.210-.530)	412
	01-02	*	< LOD	.100 (.070-.120)	.190 (.150-.260)	.290 (.190-.380)	515
	03-04	.029 (.025-.034)	.032 (.024-.037)	.083 (.070-.109)	.191 (.140-.273)	.330 (.228-.468)	458
Non-Hispanic whites	99-00	*	< LOD	< LOD	.110 (.100-.130)	.150 (.130-.170)	727
	01-02	*	< LOD	.080 (.070-.090)	.140 (.120-.160)	.190 (.170-.220)	1061
	03-04	.028 (.026-.030)	.032 (.029-.036)	.065 (.061-.071)	.111 (.101-.124)	.153 (.132-.172)	884

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5',6-Heptachlorobiphenyl (PCB 187) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.006	0.001	9
		07-08	.005	0.001	6
	20-39	05-06	.023	0.006	12
		07-08	.018	0.002	15
	40-59	05-06	.063	0.005	12
		07-08	.049	0.002	16
60+	05-06	.112	0.006	15	
	07-08	.107	0.008	23	
Non-Hispanic White Female	12-19	05-06	.004	0	10
		07-08	*	*	7
	20-39	05-06	.011	0.001	16
		07-08	.013	0.004	13
	40-59	05-06	.045	0.004	13
		07-08	.038	0.002	17
60+	05-06	.102	0.008	17	
	07-08	.093	0.005	21	
Non-Hispanic Black Male	12-19	05-06	.008	0.001	13
		07-08	.007	0.001	6
	20-39	05-06	.021	0.004	6
		07-08	.015	0.002	6
	40-59	05-06	.078	0.006	5
		07-08	.082	0.012	6
60+	05-06	.191	0.02	5	
	07-08	.193	0.028	8	
Non-Hispanic Black Female	12-19	05-06	.005	0	14
		07-08	.004	0.001	5
	20-39	05-06	.015	0.003	7
		07-08	.011	0.002	8
	40-59	05-06	.072	0.007	7
		07-08	.058	0.008	8
60+	05-06	.251	0.063	5	
	07-08	.197	0.021	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,4',5,5',6-Heptachlorobiphenyl (PCB 187) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.003	0	11
		07-08	.004	0.001	6
	20-39	05-06	.012	0.001	9
		07-08	.013	0.002	9
	40-59	05-06	.039	0.006	4
		07-08	.048	0.007	6
60+	05-06	.076	0.017	4	
	07-08	.092	0.01	5	
Mexican American Female	12-19	05-06	.003	0	16
		07-08	*	*	5
	20-39	05-06	.007	0.001	9
		07-08	.008	0.001	8
	40-59	05-06	.035	0.001	6
		07-08	.038	0.005	6
	60+	05-06	.080	0.012	3
		07-08	.088	0.02	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5'-Octachlorobiphenyl (PCB 194) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	11.1 (<LOD-11.9)	18.2 (16.5-19.5)	23.7 (21.0-27.0)	2279
	03-04	2.69 (2.53-2.85)	4.19 (3.55-4.70)	8.47 (7.79-9.06)	14.3 (13.3-16.0)	19.1 (17.6-20.8)	1835
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	746
	03-04	*	< LOD	1.11 (.700-1.57)	2.40 (1.70-3.57)	5.18 (2.80-6.81)	572
20 years and older	01-02	*	< LOD	12.0 (11.0-13.3)	19.1 (17.7-20.5)	25.3 (22.7-28.0)	1533
	03-04	3.64 (3.40-3.89)	4.95 (4.56-5.42)	9.30 (8.62-9.90)	15.3 (13.9-17.1)	20.1 (18.1-21.9)	1263
Gender							
Males	01-02	*	< LOD	12.3 (10.6-14.1)	19.2 (17.2-21.2)	25.2 (21.7-29.0)	1059
	03-04	2.95 (2.61-3.35)	4.47 (3.55-5.20)	9.60 (8.30-10.9)	17.1 (14.2-18.8)	21.8 (19.8-25.6)	913
Females	01-02	*	< LOD	< LOD	17.0 (14.7-18.8)	21.9 (19.3-26.2)	1220
	03-04	2.45 (2.28-2.64)	4.00 (3.39-4.49)	7.50 (7.01-8.39)	12.7 (11.2-13.6)	16.6 (13.7-18.6)	922
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	14.3 (10.5-17.6)	561
	03-04	.845 (.582-1.23)	.900 (.640-1.49)	2.90 (1.79-4.76)	7.21 (4.80-8.90)	9.50 (7.40-15.0)	410
Non-Hispanic blacks	01-02	*	< LOD	12.3 (<LOD-15.3)	22.6 (18.0-27.5)	29.8 (23.7-41.1)	508
	03-04	2.45 (1.94-3.08)	3.62 (2.70-4.49)	8.83 (6.80-11.2)	18.1 (13.0-23.2)	28.3 (18.9-35.6)	450
Non-Hispanic whites	01-02	*	< LOD	11.9 (10.9-13.1)	18.8 (17.1-20.2)	24.1 (20.9-28.0)	1048
	03-04	3.25 (2.97-3.56)	4.90 (4.19-5.42)	9.08 (8.36-9.80)	14.7 (13.5-16.6)	19.1 (17.4-21.2)	861

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 10.5 and 0.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5'-Octachlorobiphenyl (PCB 194) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.599	0.081	9
		07-08	.622	0.099	6
	20-39	05-06	2.31	0.63	12
		07-08	1.91	0.2	15
	40-59	05-06	8.60	1.06	12
		07-08	7.17	0.32	16
60+	05-06	15.1	0.9	15	
	07-08	15.5	0.7	23	
Non-Hispanic White Female	12-19	05-06	.386	0.061	10
		07-08	.691	0.184	7
	20-39	05-06	1.30	0.16	16
		07-08	1.64	0.16	13
	40-59	05-06	4.30	0.54	13
		07-08	5.43	0.35	17
60+	05-06	11.3	0.8	17	
	07-08	11.7	0.6	21	
Non-Hispanic Black Male	12-19	05-06	.604	0.084	13
		07-08	*	*	6
	20-39	05-06	2.19	0.28	6
		07-08	1.75	0.2	6
	40-59	05-06	8.55	0.39	5
		07-08	8.95	0.86	6
60+	05-06	21.6	2.2	5	
	07-08	20.8	2.1	8	
Non-Hispanic Black Female	12-19	05-06	.362	0.042	14
		07-08	*	*	5
	20-39	05-06	1.30	0.2	7
		07-08	1.11	0.26	8
	40-59	05-06	6.26	0.84	7
		07-08	6.00	0.7	8
60+	05-06	18.1	3.4	5	
	07-08	16.9	1	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5'-Octachlorobiphenyl (PCB 194) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.501	0.138	11
		07-08	.465	0.077	6
	20-39	05-06	.988	0.082	9
		07-08	*	*	9
	40-59	05-06	3.83	0.39	4
		07-08	5.26	0.76	6
60+	05-06	9.53	2.54	4	
	07-08	12.2	1.7	5	
Mexican American Female	12-19	05-06	.331	0.038	16
		07-08	*	*	5
	20-39	05-06	.641	0.109	9
		07-08	.943	0.049	8
	40-59	05-06	3.25	0.45	6
		07-08	3.73	0.95	6
	60+	05-06	6.74	0.23	3
		07-08	7.77	1.42	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5'-Octachlorobiphenyl (PCB 194) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	.070 (<LOD-.080)	.120 (.110-.140)	.160 (.150-.190)	2279
	03-04	.016 (.015-.017)	.026 (.022-.028)	.056 (.052-.060)	.096 (.091-.103)	.129 (.116-.139)	1835
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	746
	03-04	*	< LOD	.006 (.004-.008)	.013 (.009-.020)	.023 (.013-.035)	572
20 years and older	01-02	*	< LOD	.080 (.070-.090)	.130 (.110-.150)	.170 (.150-.200)	1533
	03-04	.023 (.021-.024)	.031 (.027-.035)	.061 (.057-.065)	.103 (.094-.113)	.133 (.122-.145)	1263
Gender							
Males	01-02	*	< LOD	.080 (.070-.090)	.140 (.110-.160)	.170 (.150-.210)	1059
	03-04	.018 (.016-.021)	.027 (.023-.034)	.061 (.054-.069)	.106 (.095-.121)	.142 (.129-.166)	913
Females	01-02	*	< LOD	< LOD	.110 (.090-.130)	.150 (.130-.180)	1220
	03-04	.015 (.014-.016)	.024 (.020-.027)	.051 (.047-.056)	.088 (.080-.096)	.112 (.103-.117)	922
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	.100 (.080-.110)	561
	03-04	.005 (.004-.008)	.006 (.004-.010)	.019 (.012-.031)	.049 (.033-.064)	.081 (.051-.118)	410
Non-Hispanic blacks	01-02	*	< LOD	.080 (<LOD-.090)	.150 (.100-.180)	.200 (.140-.250)	508
	03-04	.014 (.011-.018)	.020 (.014-.026)	.054 (.045-.063)	.120 (.081-.155)	.172 (.123-.202)	450
Non-Hispanic whites	01-02	*	< LOD	.080 (.070-.090)	.130 (.110-.150)	.170 (.150-.190)	1048
	03-04	.020 (.018-.022)	.029 (.026-.034)	.061 (.056-.066)	.099 (.090-.106)	.132 (.116-.143)	861

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5'-Octachlorobiphenyl (PCB 194) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.003	0	9
		07-08	.003	0	6
	20-39	05-06	.018	0.005	12
		07-08	.013	0.001	15
	40-59	05-06	.065	0.008	12
		07-08	.052	0.003	16
60+	05-06	.102	0.005	15	
	07-08	.100	0.005	23	
Non-Hispanic White Female	12-19	05-06	.002	0	10
		07-08	.003	0.001	7
	20-39	05-06	.008	0.001	16
		07-08	.010	0.001	13
	40-59	05-06	.032	0.005	13
		07-08	.036	0.003	17
60+	05-06	.082	0.006	17	
	07-08	.079	0.004	21	
Non-Hispanic Black Male	12-19	05-06	.003	0	13
		07-08	*	*	6
	20-39	05-06	.013	0.002	6
		07-08	.010	0.001	6
	40-59	05-06	.054	0.003	5
		07-08	.057	0.006	6
60+	05-06	.131	0.017	5	
	07-08	.127	0.014	8	
Non-Hispanic Black Female	12-19	05-06	.002	0	14
		07-08	*	*	5
	20-39	05-06	.007	0.001	7
		07-08	.006	0.001	8
	40-59	05-06	.040	0.005	7
		07-08	.036	0.005	8
60+	05-06	.120	0.022	5	
	07-08	.102	0.007	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5'-Octachlorobiphenyl (PCB 194) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.003	0.001	11
		07-08	.002	0	6
	20-39	05-06	.007	0.001	9
		07-08	*	*	9
	40-59	05-06	.031	0.004	4
		07-08	.037	0.005	6
	60+	05-06	.069	0.018	4
		07-08	.086	0.009	5
Mexican American Female	12-19	05-06	.002	0	16
		07-08	*	*	5
	20-39	05-06	.004	0.001	9
		07-08	.005	0	8
	40-59	05-06	.024	0.003	6
		07-08	.025	0.006	6
	60+	05-06	.052	0.007	3
		07-08	.054	0.011	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6-Octachlorobiphenyl (PCB 195) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2230
	03-04	*	.900 (<LOD-1.01)	1.98 (1.80-2.10)	3.40 (3.10-3.60)	4.51 (3.90-5.18)	1820
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	716
	03-04	*	< LOD	< LOD	< LOD	1.74 (.720-3.82)	567
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1514
	03-04	.800 (.721-.888)	1.10 (.980-1.20)	2.10 (2.00-2.30)	3.53 (3.20-3.74)	4.68 (4.12-5.19)	1253
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1035
	03-04	*	.800 (<LOD-1.00)	2.08 (1.82-2.20)	3.50 (3.12-3.70)	4.60 (3.82-5.40)	904
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1195
	03-04	*	.970 (<LOD-1.10)	1.85 (1.64-2.09)	3.30 (2.86-3.60)	4.50 (3.57-5.25)	916
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	544
	03-04	*	< LOD	< LOD	1.80 (1.11-2.40)	2.40 (1.80-3.50)	407
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	490
	03-04	.709 (<LOD-.870)	.820 (<LOD-1.14)	2.00 (1.65-2.57)	4.18 (3.16-5.30)	6.90 (4.18-8.10)	444
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1037
	03-04	.749 (<LOD-.850)	1.00 (.800-1.18)	2.10 (1.90-2.30)	3.40 (3.10-3.61)	4.44 (3.80-5.19)	856

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 28.1 and 0.7.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6-Octachlorobiphenyl (PCB 195) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.487	0.129	12
		07-08	*	*	15
	40-59	05-06	1.50	0.29	12
		07-08	1.25	0.05	16
60+	05-06	2.52	0.16	15	
	07-08	2.74	0.16	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	.936	0.067	13
		07-08	1.14	0.08	17
60+	05-06	2.04	0.16	17	
	07-08	2.38	0.13	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.435	0.047	6
		07-08	*	*	6
	40-59	05-06	1.35	0.13	5
		07-08	1.80	0.23	6
60+	05-06	3.40	0.27	5	
	07-08	3.99	0.53	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.348	0.044	7
		07-08	*	*	8
	40-59	05-06	1.41	0.15	7
		07-08	1.35	0.17	8
60+	05-06	3.82	0.76	5	
	07-08	3.94	0.26	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6-Octachlorobiphenyl (PCB 195) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	.766	0.083	4
		07-08	1.02	0.11	6
60+	05-06	1.69	0.4	4	
	07-08	2.24	0.29	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.780	0.083	6
		07-08	.986	0.187	6
	60+	05-06	1.47	0.06	3
		07-08	1.98	0.39	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6-Octachlorobiphenyl (PCB 195) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2230
	03-04	*	.005 (<LOD-.006)	.012 (.012-.013)	.022 (.020-.025)	.031 (.027-.034)	1820
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	716
	03-04	*	< LOD	< LOD	< LOD	.007 (.004-.017)	567
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1514
	03-04	.005 (.004-.006)	.007 (.006-.007)	.014 (.012-.015)	.023 (.021-.026)	.032 (.028-.035)	1253
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1035
	03-04	*	.005 (<LOD-.006)	.013 (.012-.014)	.022 (.020-.025)	.031 (.025-.035)	904
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1195
	03-04	*	.006 (<LOD-.007)	.012 (.011-.013)	.021 (.019-.025)	.031 (.026-.035)	916
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	544
	03-04	*	< LOD	< LOD	.012 (.008-.018)	.018 (.013-.025)	407
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	490
	03-04	.004 (<LOD-.005)	.004 (<LOD-.006)	.012 (.010-.016)	.027 (.019-.034)	.041 (.029-.048)	444
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1037
	03-04	.005 (<LOD-.005)	.006 (.005-.007)	.013 (.012-.014)	.022 (.020-.025)	.030 (.027-.033)	856

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6-Octachlorobiphenyl (PCB 195) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.004	0.001	12
		07-08	*	*	15
	40-59	05-06	.011	0.002	12
		07-08	.009	0	16
	60+	05-06	.017	0.001	15
		07-08	.018	0.001	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	*	*	16
		07-08	*	*	13
	40-59	05-06	.007	0.001	13
		07-08	.008	0.001	17
	60+	05-06	.015	0.001	17
		07-08	.016	0.001	21
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.003	0	6
		07-08	*	*	6
	40-59	05-06	.009	0.001	5
		07-08	.012	0.002	6
	60+	05-06	.020	0.001	5
		07-08	.024	0.004	8
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.002	0	7
		07-08	*	*	8
	40-59	05-06	.009	0.001	7
		07-08	.008	0.001	8
	60+	05-06	.025	0.005	5
		07-08	.024	0.002	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6-Octachlorobiphenyl (PCB 195) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	*	*	9
		07-08	*	*	9
	40-59	05-06	.006	0.001	4
		07-08	.007	0.001	6
60+	05-06	.012	0.003	4	
	07-08	.016	0.001	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.006	0.001	6
		07-08	.006	0.001	6
	60+	05-06	.011	0.002	3
		07-08	.014	0.003	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6' and 2,2',3,4,4',5,5',6-Octachlorobiphenyl (PCB 196 & 203) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	14.2 (13.0-15.7)	19.2 (17.4-20.9)	2299
	01-02	*	< LOD	< LOD	14.2 (13.0-15.7)	19.2 (17.4-20.9)	2299
	03-04	2.61 (2.44-2.79)	3.40 (3.01-3.80)	6.70 (6.16-7.23)	11.8 (10.8-12.7)	15.0 (13.2-17.4)	1878
	03-04	2.61 (2.44-2.79)	3.40 (3.01-3.80)	6.70 (6.16-7.23)	11.8 (10.8-12.7)	15.0 (13.2-17.4)	1878
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	755
	01-02	*	< LOD	< LOD	< LOD	< LOD	755
	03-04	.437 (<LOD-.546)	.520 (.400-.700)	1.10 (.900-1.42)	2.41 (1.86-3.19)	3.59 (2.93-4.20)	587
	03-04	.437 (<LOD-.546)	.520 (.400-.700)	1.10 (.900-1.42)	2.41 (1.86-3.19)	3.59 (2.93-4.20)	587
20 years and older	01-02	*	< LOD	< LOD	15.0 (13.7-17.0)	19.9 (18.2-21.4)	1544
	01-02	*	< LOD	< LOD	15.0 (13.7-17.0)	19.9 (18.2-21.4)	1544
	03-04	3.37 (3.15-3.61)	4.07 (3.74-4.46)	7.30 (6.79-8.20)	12.4 (11.5-13.2)	15.9 (13.4-19.5)	1291
	03-04	3.37 (3.15-3.61)	4.07 (3.74-4.46)	7.30 (6.79-8.20)	12.4 (11.5-13.2)	15.9 (13.4-19.5)	1291
Gender							
Males	01-02	*	< LOD	< LOD	14.6 (13.3-16.5)	19.4 (16.3-21.2)	1071
	01-02	*	< LOD	< LOD	14.6 (13.3-16.5)	19.4 (16.3-21.2)	1071
	03-04	2.77 (2.49-3.08)	3.45 (3.07-3.98)	6.92 (6.12-7.71)	12.5 (11.2-13.7)	16.3 (13.7-23.1)	935
	03-04	2.77 (2.49-3.08)	3.45 (3.07-3.98)	6.92 (6.12-7.71)	12.5 (11.2-13.7)	16.3 (13.7-23.1)	935
Females	01-02	*	< LOD	< LOD	13.8 (11.9-15.9)	19.1 (17.0-21.3)	1228
	01-02	*	< LOD	< LOD	13.8 (11.9-15.9)	19.1 (17.0-21.3)	1228
	03-04	2.46 (2.27-2.66)	3.32 (2.83-3.80)	6.68 (6.12-7.20)	11.1 (9.74-12.3)	13.4 (12.8-14.9)	943
	03-04	2.46 (2.27-2.66)	3.32 (2.83-3.80)	6.68 (6.12-7.20)	11.1 (9.74-12.3)	13.4 (12.8-14.9)	943
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	11.1 (<LOD-12.5)	565
	01-02	*	< LOD	< LOD	< LOD	11.1 (<LOD-12.5)	565
	03-04	.947 (.689-1.30)	1.00 (.700-1.50)	2.80 (1.60-4.30)	5.90 (4.30-7.40)	8.35 (6.00-12.3)	424
	03-04	.947 (.689-1.30)	1.00 (.700-1.50)	2.80 (1.60-4.30)	5.90 (4.30-7.40)	8.35 (6.00-12.3)	424
Non-Hispanic blacks	01-02	*	< LOD	< LOD	17.6 (14.2-21.0)	23.1 (19.6-30.2)	513
	01-02	*	< LOD	< LOD	17.6 (14.2-21.0)	23.1 (19.6-30.2)	513
	03-04	2.45 (2.06-2.90)	2.90 (2.33-3.80)	7.18 (5.80-9.00)	13.8 (11.5-16.8)	21.6 (14.0-32.3)	453
	03-04	2.45 (2.06-2.90)	2.90 (2.33-3.80)	7.18 (5.80-9.00)	13.8 (11.5-16.8)	21.6 (14.0-32.3)	453
Non-Hispanic whites	01-02	*	< LOD	< LOD	14.7 (13.6-16.5)	19.6 (18.1-21.3)	1058
	01-02	*	< LOD	< LOD	14.7 (13.6-16.5)	19.6 (18.1-21.3)	1058
	03-04	3.01 (2.76-3.27)	3.88 (3.44-4.20)	7.10 (6.32-8.20)	11.8 (10.8-13.1)	15.4 (12.9-19.0)	882
	03-04	3.01 (2.76-3.27)	3.88 (3.44-4.20)	7.10 (6.32-8.20)	11.8 (10.8-13.1)	15.4 (12.9-19.0)	882

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 10.5 and 0.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6' and 2,2',3,4,4',5,5',6-Octachlorobiphenyl (PCB 196 & 203) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]	
Non-Hispanic White Male	12-19	05-06	.793	0.071	9	
		07-08	.667	0.151	6	
	20-39	05-06	2.63	0.69	12	
		07-08	2.15	0.14	15	
	40-59	05-06	8.33	0.94	12	
		07-08	6.45	0.25	16	
	60+	05-06	15.4	0.8	15	
		07-08	15.3	0.8	23	
Non-Hispanic White Female	12-19	05-06	.593	0.046	10	
		07-08	*	*	7	
	20-39	05-06	1.49	0.12	16	
		07-08	1.72	0.21	13	
	40-59	05-06	5.23	0.24	13	
		07-08	5.27	0.3	17	
	60+	05-06	12.6	1	17	
		07-08	12.2	0.7	21	
	Non-Hispanic Black Male	12-19	05-06	.937	0.103	13
			07-08	*	*	6
20-39		05-06	2.34	0.22	6	
		07-08	1.79	0.18	6	
40-59		05-06	8.31	0.66	5	
		07-08	8.17	0.91	6	
60+		05-06	20.9	2.2	5	
		07-08	19.8	2.3	8	
Non-Hispanic Black Female	12-19	05-06	.557	0.039	14	
		07-08	*	*	5	
	20-39	05-06	1.61	0.22	7	
		07-08	1.40	0.21	8	
	40-59	05-06	7.01	0.65	7	
		07-08	6.50	0.73	8	
	60+	05-06	21.2	4.7	5	
		07-08	18.1	1.4	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6' and 2,2',3,4,4',5,5',6-Octachlorobiphenyl (PCB 196 & 203) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.500	0.047	11
		07-08	.514	0.087	6
	20-39	05-06	1.16	0.1	9
		07-08	*	*	9
	40-59	05-06	3.85	0.42	4
		07-08	4.71	0.62	6
	60+	05-06	9.56	2.38	4
		07-08	10.6	1.8	5
Mexican American Female	12-19	05-06	.464	0.05	16
		07-08	*	*	5
	20-39	05-06	.770	0.121	9
		07-08	1.02	0.04	8
	40-59	05-06	3.87	0.54	6
		07-08	4.02	0.61	6
	60+	05-06	7.87	0.44	3
		07-08	7.87	1.56	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6' and 2,2',3,4,4',5,5',6-Octachlorobiphenyl (PCB 196 & 203) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	.100 (.090-.110)	.130 (.110-.140)	2299
	01-02	*	< LOD	< LOD	.100 (.090-.110)	.130 (.110-.140)	2299
	03-04	.016 (.015-.017)	.022 (.019-.025)	.044 (.042-.048)	.082 (.071-.087)	.101 (.091-.112)	1878
	03-04	.016 (.015-.017)	.022 (.019-.025)	.044 (.042-.048)	.082 (.071-.087)	.101 (.091-.112)	1878
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	755
	01-02	*	< LOD	< LOD	< LOD	< LOD	755
	03-04	.002 (<LOD-.003)	.003 (.002-.004)	.006 (.005-.007)	.013 (.009-.016)	.016 (.014-.025)	587
	03-04	.002 (<LOD-.003)	.003 (.002-.004)	.006 (.005-.007)	.013 (.009-.016)	.016 (.014-.025)	587
20 years and older	01-02	*	< LOD	< LOD	.100 (.090-.110)	.130 (.120-.150)	1544
	01-02	*	< LOD	< LOD	.100 (.090-.110)	.130 (.120-.150)	1544
	03-04	.021 (.020-.023)	.026 (.024-.028)	.049 (.045-.052)	.084 (.077-.090)	.104 (.092-.123)	1291
	03-04	.021 (.020-.023)	.026 (.024-.028)	.049 (.045-.052)	.084 (.077-.090)	.104 (.092-.123)	1291
Gender							
Males	01-02	*	< LOD	< LOD	.100 (.090-.120)	.130 (.110-.150)	1071
	01-02	*	< LOD	< LOD	.100 (.090-.120)	.130 (.110-.150)	1071
	03-04	.017 (.015-.019)	.023 (.020-.026)	.045 (.042-.050)	.084 (.077-.091)	.106 (.093-.127)	935
	03-04	.017 (.015-.019)	.023 (.020-.026)	.045 (.042-.050)	.084 (.077-.091)	.106 (.093-.127)	935
Females	01-02	*	< LOD	< LOD	.090 (.080-.100)	.130 (.110-.140)	1228
	01-02	*	< LOD	< LOD	.090 (.080-.100)	.130 (.110-.140)	1228
	03-04	.015 (.014-.016)	.020 (.017-.023)	.043 (.038-.047)	.076 (.066-.084)	.093 (.085-.111)	943
	03-04	.015 (.014-.016)	.020 (.017-.023)	.043 (.038-.047)	.076 (.066-.084)	.093 (.085-.111)	943
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	.070 (<LOD-.090)	565
	01-02	*	< LOD	< LOD	< LOD	.070 (<LOD-.090)	565
	03-04	.006 (.004-.008)	.007 (.004-.010)	.017 (.011-.026)	.038 (.028-.054)	.058 (.035-.087)	424
	03-04	.006 (.004-.008)	.007 (.004-.010)	.017 (.011-.026)	.038 (.028-.054)	.058 (.035-.087)	424
Non-Hispanic blacks	01-02	*	< LOD	< LOD	.110 (.090-.140)	.150 (.110-.210)	513
	01-02	*	< LOD	< LOD	.110 (.090-.140)	.150 (.110-.210)	513
	03-04	.014 (.012-.017)	.017 (.012-.022)	.046 (.038-.054)	.090 (.065-.111)	.141 (.094-.209)	453
	03-04	.014 (.012-.017)	.017 (.012-.022)	.046 (.038-.054)	.090 (.065-.111)	.141 (.094-.209)	453
Non-Hispanic whites	01-02	*	< LOD	< LOD	.100 (.090-.110)	.130 (.110-.140)	1058
	01-02	*	< LOD	< LOD	.100 (.090-.110)	.130 (.110-.140)	1058
	03-04	.019 (.017-.020)	.025 (.022-.028)	.048 (.043-.052)	.084 (.070-.090)	.100 (.090-.120)	882
	03-04	.019 (.017-.020)	.025 (.022-.028)	.048 (.043-.052)	.084 (.070-.090)	.100 (.090-.120)	882

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6' and 2,2',3,4,4',5,5',6-Octachlorobiphenyl (PCB 196 & 203) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]	
Non-Hispanic White Male	12-19	05-06	.004	0	9	
		07-08	.003	0.001	6	
	20-39	05-06	.020	0.006	12	
		07-08	.014	0.001	15	
	40-59	05-06	.063	0.008	12	
		07-08	.047	0.002	16	
	60+	05-06	.104	0.004	15	
		07-08	.098	0.005	23	
	Non-Hispanic White Female	12-19	05-06	.003	0	10
			07-08	*	*	7
20-39		05-06	.010	0.001	16	
		07-08	.010	0.001	13	
40-59		05-06	.039	0.003	13	
		07-08	.035	0.002	17	
60+		05-06	.091	0.007	17	
		07-08	.082	0.005	21	
Non-Hispanic Black Male		12-19	05-06	.005	0	13
			07-08	*	*	6
	20-39	05-06	.014	0.001	6	
		07-08	.010	0.001	6	
	40-59	05-06	.053	0.004	5	
		07-08	.053	0.006	6	
	60+	05-06	.125	0.013	5	
		07-08	.121	0.016	8	
	Non-Hispanic Black Female	12-19	05-06	.003	0	14
			07-08	*	*	5
20-39		05-06	.009	0.001	7	
		07-08	.007	0.001	8	
40-59		05-06	.045	0.004	7	
		07-08	.039	0.005	8	
60+		05-06	.140	0.031	5	
		07-08	.108	0.008	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,6' and 2,2',3,4,4',5,5',6-Octachlorobiphenyl (PCB 196 & 203) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.003	0	11
		07-08	.003	0	6
	20-39	05-06	.008	0.001	9
		07-08	*	*	9
	40-59	05-06	.032	0.004	4
		07-08	.033	0.004	6
	60+	05-06	.069	0.017	4
		07-08	.075	0.011	5
Mexican American Female	12-19	05-06	.003	0	16
		07-08	*	*	5
	20-39	05-06	.005	0.001	9
		07-08	.006	0	8
	40-59	05-06	.028	0.003	6
		07-08	.027	0.004	6
	60+	05-06	.060	0.008	3
		07-08	.054	0.011	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5',6-Octachlorobiphenyl (PCB 199) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	17.0 (15.4-18.1)	22.4 (19.9-25.9)	2292
	03-04	2.81 (2.64-3.00)	3.80 (3.26-4.26)	8.30 (7.70-9.08)	14.9 (13.8-15.7)	18.9 (17.9-21.3)	1861
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	.416 (<LOD-.522)	.500 (<LOD-.700)	1.17 (.890-1.59)	2.43 (2.02-2.73)	3.92 (2.90-4.10)	579
20 years and older	01-02	*	< LOD	10.6 (<LOD-11.8)	17.9 (16.2-19.2)	24.4 (20.7-27.4)	1536
	03-04	3.71 (3.41-4.02)	4.60 (4.10-5.08)	9.40 (8.73-10.0)	15.7 (14.7-16.9)	20.6 (18.3-23.6)	1282
Gender							
Males	01-02	*	< LOD	< LOD	17.3 (15.4-18.7)	21.3 (19.3-24.6)	1066
	03-04	3.01 (2.68-3.39)	3.95 (3.30-4.67)	9.10 (7.60-9.94)	16.6 (14.0-17.3)	22.1 (18.1-27.4)	925
Females	01-02	*	< LOD	< LOD	16.2 (14.5-17.9)	22.9 (19.4-27.3)	1226
	03-04	2.63 (2.49-2.79)	3.70 (3.20-4.00)	7.70 (6.92-8.81)	14.0 (11.6-15.3)	17.0 (15.7-18.9)	936
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	12.3 (<LOD-14.2)	565
	03-04	.943 (.664-1.34)	1.00 (.600-1.50)	3.35 (1.50-5.33)	7.37 (5.10-10.2)	10.2 (7.40-13.5)	418
Non-Hispanic blacks	01-02	*	< LOD	12.5 (<LOD-15.3)	21.9 (17.0-28.1)	30.7 (22.4-37.4)	513
	03-04	3.03 (2.46-3.74)	3.60 (2.70-4.70)	10.0 (7.70-13.7)	21.3 (15.9-30.4)	35.1 (21.3-43.3)	452
Non-Hispanic whites	01-02	*	< LOD	< LOD	17.3 (15.9-18.9)	22.7 (19.8-26.3)	1051
	03-04	3.24 (3.03-3.47)	4.19 (3.78-4.69)	9.08 (7.88-9.86)	14.8 (13.5-16.3)	18.6 (16.9-21.7)	871

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 10.5 and 0.4.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5',6-Octachlorobiphenyl (PCB 199) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.629	0.101	9
		07-08	*	*	6
	20-39	05-06	2.45	0.63	12
		07-08	2.12	0.18	15
	40-59	05-06	8.40	0.83	12
		07-08	6.73	0.4	16
	60+	05-06	17.5	1.1	15
		07-08	16.9	1	23
Non-Hispanic White Female	12-19	05-06	.540	0.058	10
		07-08	*	*	7
	20-39	05-06	1.37	0.12	16
		07-08	1.61	0.18	13
	40-59	05-06	5.45	0.34	13
		07-08	5.27	0.38	17
	60+	05-06	15.1	1.2	17
		07-08	14.7	0.9	21
Non-Hispanic Black Male	12-19	05-06	.851	0.108	13
		07-08	*	*	6
	20-39	05-06	2.43	0.32	6
		07-08	1.70	0.17	6
	40-59	05-06	10.7	0.9	5
		07-08	9.25	1.05	6
	60+	05-06	28.0	3.6	5
		07-08	24.6	2.9	8
Non-Hispanic Black Female	12-19	05-06	.490	0.051	14
		07-08	*	*	5
	20-39	05-06	1.59	0.24	7
		07-08	1.36	0.17	8
	40-59	05-06	8.16	1.04	7
		07-08	6.94	0.87	8
	60+	05-06	27.6	6	5
		07-08	22.0	1.8	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5',6-Octachlorobiphenyl (PCB 199) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.395	0.045	11
		07-08	*	*	6
	20-39	05-06	1.03	0.11	9
		07-08	*	*	9
	40-59	05-06	4.23	0.49	4
		07-08	5.47	0.92	6
	60+	05-06	10.1	2.6	4
		07-08	11.4	1.6	5
Mexican American Female	12-19	05-06	.386	0.052	16
		07-08	*	*	5
	20-39	05-06	.634	0.11	9
		07-08	.843	0.031	8
	40-59	05-06	3.67	0.52	6
		07-08	4.09	0.69	6
	60+	05-06	8.79	0.55	3
		07-08	8.71	1.93	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5',6-Octachlorobiphenyl (PCB 199) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	.110 (.100-.130)	.150 (.140-.170)	2292
	03-04	.017 (.016-.018)	.023 (.020-.027)	.056 (.053-.060)	.100 (.091-.106)	.127 (.119-.144)	1861
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	756
	03-04	.002 (<LOD-.003)	.002 (<LOD-.003)	.006 (.005-.008)	.012 (.009-.015)	.016 (.015-.019)	579
20 years and older	01-02	*	< LOD	.070 (<LOD-.080)	.120 (.110-.140)	.160 (.140-.180)	1536
	03-04	.023 (.021-.025)	.029 (.026-.033)	.062 (.058-.067)	.106 (.097-.113)	.135 (.120-.158)	1282
Gender							
Males	01-02	*	< LOD	< LOD	.110 (.100-.130)	.150 (.130-.180)	1066
	03-04	.019 (.016-.021)	.025 (.020-.030)	.059 (.053-.068)	.105 (.090-.113)	.138 (.120-.162)	925
Females	01-02	*	< LOD	< LOD	.110 (.100-.130)	.150 (.130-.170)	1226
	03-04	.016 (.015-.017)	.022 (.019-.025)	.054 (.047-.058)	.094 (.085-.105)	.120 (.108-.133)	936
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	.090 (<LOD-.110)	565
	03-04	.006 (.004-.008)	.007 (.004-.010)	.019 (.011-.031)	.051 (.029-.079)	.079 (.051-.093)	418
Non-Hispanic blacks	01-02	*	< LOD	.070 (<LOD-.090)	.140 (.100-.180)	.190 (.130-.300)	513
	03-04	.017 (.014-.021)	.021 (.015-.026)	.064 (.054-.073)	.135 (.102-.173)	.207 (.147-.297)	452
Non-Hispanic whites	01-02	*	< LOD	< LOD	.120 (.100-.130)	.150 (.130-.180)	1051
	03-04	.020 (.019-.022)	.027 (.023-.030)	.059 (.054-.065)	.101 (.087-.108)	.125 (.108-.144)	871

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5',6-Octachlorobiphenyl (PCB 199) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.003	0.001	9
		07-08	*	*	6
	20-39	05-06	.019	0.006	12
		07-08	.014	0.001	15
	40-59	05-06	.063	0.005	12
		07-08	.049	0.003	16
	60+	05-06	.118	0.006	15
		07-08	.109	0.006	23
Non-Hispanic White Female	12-19	05-06	.003	0	10
		07-08	*	*	7
	20-39	05-06	.009	0.001	16
		07-08	.010	0.001	13
	40-59	05-06	.041	0.004	13
		07-08	.035	0.003	17
	60+	05-06	.110	0.009	17
		07-08	.099	0.006	21
Non-Hispanic Black Male	12-19	05-06	.004	0.001	13
		07-08	*	*	6
	20-39	05-06	.014	0.002	6
		07-08	.010	0.001	6
	40-59	05-06	.069	0.006	5
		07-08	.059	0.007	6
	60+	05-06	.168	0.024	5
		07-08	.150	0.02	8
Non-Hispanic Black Female	12-19	05-06	.002	0	14
		07-08	*	*	5
	20-39	05-06	.009	0.001	7
		07-08	.007	0.001	8
	40-59	05-06	.052	0.006	7
		07-08	.042	0.006	8
	60+	05-06	.183	0.039	5
		07-08	.132	0.011	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,5,5',6-Octachlorobiphenyl (PCB 199) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	.002	0	11
		07-08	*	*	6
	20-39	05-06	.007	0.001	9
		07-08	*	*	9
	40-59	05-06	.035	0.005	4
		07-08	.038	0.006	6
	60+	05-06	.073	0.02	4
		07-08	.081	0.009	5
Mexican American Female	12-19	05-06	.002	0	16
		07-08	*	*	5
	20-39	05-06	.004	0.001	9
		07-08	.005	0	8
	40-59	05-06	.027	0.003	6
		07-08	.028	0.004	6
	60+	05-06	.068	0.01	3
		07-08	.060	0.014	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6-Nonachlorobiphenyl (PCB 206) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric	Selected percentiles				Sample size
		mean (95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2208
	03-04	2.13 (1.97-2.31)	2.34 (2.00-2.60)	5.00 (4.69-5.50)	9.20 (8.55-9.79)	13.7 (11.5-15.6)	1867
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	723
	03-04	*	< LOD	.900 (.750-1.20)	1.75 (1.23-2.92)	3.12 (1.60-6.91)	585
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1485
	03-04	2.59 (2.37-2.83)	2.80 (2.42-3.06)	5.53 (5.10-6.10)	9.61 (9.03-11.0)	14.2 (12.2-17.1)	1282
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1033
	03-04	2.23 (2.03-2.45)	2.30 (1.90-2.70)	5.10 (4.78-5.53)	9.31 (8.18-10.8)	13.8 (10.8-15.7)	933
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1175
	03-04	2.05 (1.86-2.25)	2.34 (1.95-2.78)	5.00 (4.30-5.70)	9.00 (8.00-9.79)	13.6 (11.2-16.5)	934
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	533
	03-04	.861 (<LOD-1.15)	.730 (<LOD-1.10)	1.70 (1.00-3.14)	4.40 (2.61-6.20)	6.20 (4.40-8.40)	421
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	483
	03-04	2.27 (1.98-2.60)	2.26 (1.85-2.51)	5.60 (4.30-7.40)	11.5 (9.49-16.5)	20.9 (12.1-31.7)	456
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1034
	03-04	2.40 (2.19-2.63)	2.70 (2.37-3.01)	5.37 (4.88-6.06)	9.20 (8.65-10.1)	13.8 (11.2-16.6)	872

Limit of detection (LOD, see Data Analysis section) for Survey years 01-02 and 03-04 are 28.1 and 0.7.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6-Nonachlorobiphenyl (PCB 206) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.392	0.038	9
		07-08	*	*	6
	20-39	05-06	1.24	0.26	12
		07-08	1.19	0.09	15
	40-59	05-06	3.97	0.23	12
		07-08	3.46	0.26	16
60+	05-06	10.3	0.7	15	
	07-08	8.72	0.48	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	.486	0.089	7
	20-39	05-06	.778	0.066	16
		07-08	.970	0.116	13
	40-59	05-06	2.94	0.22	13
		07-08	2.93	0.21	17
60+	05-06	11.2	1.3	17	
	07-08	9.65	0.75	21	
Non-Hispanic Black Male	12-19	05-06	.519	0.052	13
		07-08	.427	0.112	6
	20-39	05-06	1.27	0.13	6
		07-08	1.10	0.13	6
	40-59	05-06	6.34	0.29	5
		07-08	4.69	0.43	6
60+	05-06	17.8	3.2	5	
	07-08	13.7	1.7	8	
Non-Hispanic Black Female	12-19	05-06	.333	0.028	14
		07-08	*	*	5
	20-39	05-06	.953	0.119	7
		07-08	.861	0.122	8
	40-59	05-06	4.56	0.47	7
		07-08	3.75	0.52	8
60+	05-06	21.1	4.7	5	
	07-08	14.9	1.7	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6-Nonachlorobiphenyl (PCB 206) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.528	0.046	9
		07-08	*	*	9
	40-59	05-06	2.01	0.24	4
		07-08	2.47	0.34	6
60+	05-06	5.50	1.47	4	
	07-08	5.60	0.7	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.351	0.047	9
		07-08	.509	0.087	8
	40-59	05-06	1.91	0.28	6
		07-08	2.01	0.29	6
	60+	05-06	5.76	0.27	3
		07-08	5.10	1.24	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6-Nonachlorobiphenyl (PCB 206) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	01-02	*	< LOD	< LOD	< LOD	< LOD	2208
	03-04	.013 (.012-.014)	.015 (.012-.017)	.034 (.031-.037)	.060 (.054-.068)	.086 (.077-.104)	1867
Age group							
12-19 years	01-02	*	< LOD	< LOD	< LOD	< LOD	723
	03-04	*	< LOD	.005 (.004-.006)	.010 (.006-.013)	.014 (.008-.035)	585
20 years and older	01-02	*	< LOD	< LOD	< LOD	< LOD	1485
	03-04	.016 (.015-.018)	.017 (.016-.020)	.038 (.034-.042)	.066 (.059-.074)	.090 (.081-.112)	1282
Gender							
Males	01-02	*	< LOD	< LOD	< LOD	< LOD	1033
	03-04	.014 (.012-.015)	.015 (.012-.018)	.034 (.032-.038)	.060 (.051-.070)	.082 (.070-.097)	933
Females	01-02	*	< LOD	< LOD	< LOD	< LOD	1175
	03-04	.012 (.011-.014)	.014 (.012-.016)	.035 (.029-.038)	.061 (.054-.069)	.088 (.078-.118)	934
Race/ethnicity							
Mexican Americans	01-02	*	< LOD	< LOD	< LOD	< LOD	533
	03-04	.005 (<LOD-.007)	.005 (<LOD-.007)	.011 (.008-.018)	.031 (.019-.044)	.044 (.030-.062)	421
Non-Hispanic blacks	01-02	*	< LOD	< LOD	< LOD	< LOD	483
	03-04	.013 (.011-.015)	.012 (.010-.014)	.035 (.028-.045)	.078 (.052-.107)	.128 (.080-.213)	456
Non-Hispanic whites	01-02	*	< LOD	< LOD	< LOD	< LOD	1034
	03-04	.015 (.013-.016)	.017 (.015-.020)	.036 (.033-.040)	.063 (.058-.069)	.088 (.077-.107)	872

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6-Nonachlorobiphenyl (PCB 206) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	.002	0	9
		07-08	*	*	6
	20-39	05-06	.009	0.002	12
		07-08	.008	0.001	15
	40-59	05-06	.030	0.002	12
		07-08	.025	0.002	16
	60+	05-06	.070	0.004	15
		07-08	.056	0.003	23
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	.002	0	7
	20-39	05-06	.005	0	16
		07-08	.006	0.001	13
	40-59	05-06	.022	0.002	13
		07-08	.019	0.001	17
	60+	05-06	.081	0.009	17
		07-08	.065	0.005	21
Non-Hispanic Black Male	12-19	05-06	.003	0	13
		07-08	.002	0	6
	20-39	05-06	.007	0.001	6
		07-08	.006	0.001	6
	40-59	05-06	.040	0.003	5
		07-08	.030	0.003	6
	60+	05-06	.107	0.02	5
		07-08	.084	0.012	8
Non-Hispanic Black Female	12-19	05-06	.002	0	14
		07-08	*	*	5
	20-39	05-06	.005	0.001	7
		07-08	.004	0.001	8
	40-59	05-06	.029	0.003	7
		07-08	.023	0.004	8
	60+	05-06	.140	0.03	5
		07-08	.090	0.01	7

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6-Nonachlorobiphenyl (PCB 206) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.004	0	9
		07-08	*	*	9
	40-59	05-06	.016	0.002	4
		07-08	.017	0.002	6
	60+	05-06	.040	0.012	4
		07-08	.040	0.004	5
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	.002	0	9
		07-08	.003	0	8
	40-59	05-06	.014	0.002	6
		07-08	.014	0.002	6
	60+	05-06	.044	0.005	3
		07-08	.035	0.009	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6,6'-Decachlorobiphenyl (PCB 209) (lipid adjusted)

Geometric mean and selected percentiles of serum concentrations (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	1.40 (1.21-1.61)	1.18 (1.00-1.48)	3.20 (2.67-4.01)	7.58 (6.20-8.81)	11.1 (9.00-15.7)	1854
Age group							
12-19 years	03-04	*	< LOD	< LOD	1.12 (.800-4.60)	3.49 (1.12-6.64)	578
20 years and older	03-04	1.63 (1.42-1.88)	1.40 (1.18-1.70)	3.80 (3.04-4.44)	8.09 (6.60-9.30)	12.3 (9.05-18.0)	1276
Gender							
Males	03-04	1.37 (1.18-1.60)	1.15 (1.00-1.50)	3.00 (2.70-3.58)	6.22 (4.97-8.20)	9.10 (7.09-13.0)	925
Females	03-04	1.42 (1.22-1.66)	1.20 (1.00-1.50)	3.46 (2.45-4.61)	8.78 (7.45-9.47)	13.9 (10.0-18.7)	929
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	1.10 (.770-1.60)	2.40 (1.60-3.60)	3.71 (2.40-4.97)	416
Non-Hispanic blacks	03-04	1.56 (1.37-1.78)	1.35 (1.10-1.52)	3.40 (2.62-4.82)	9.83 (6.70-13.6)	18.0 (11.8-27.7)	455
Non-Hispanic whites	03-04	1.53 (1.31-1.79)	1.30 (1.09-1.70)	3.89 (3.04-4.50)	7.95 (6.49-9.09)	11.3 (9.07-16.3)	864

Limit of detection (LOD, see Data Analysis section) for Survey year 03-04 is 0.7.

< LOD means less than the limit of detection, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6,6'-Decachlorobiphenyl (PCB 209) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.664	0.144	12
		07-08	.654	0.097	15
	40-59	05-06	1.78	0.14	12
		07-08	1.86	0.18	16
60+	05-06	7.07	0.6	15	
	07-08	7.12	0.68	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	.374	0.033	16
		07-08	.420	0.073	13
	40-59	05-06	1.36	0.12	13
		07-08	1.64	0.16	17
60+	05-06	11.2	1.5	17	
	07-08	11.0	1.2	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.797	0.226	6
		07-08	.500	0.107	6
	40-59	05-06	4.04	0.4	5
		07-08	2.72	0.39	6
60+	05-06	14.7	2.6	5	
	07-08	13.4	1.9	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.570	0.093	7
		07-08	*	*	8
	40-59	05-06	2.80	0.3	7
		07-08	2.36	0.38	8
60+	05-06	21.6	4.9	5	
	07-08	16.0	2.5	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6,6'-Decachlorobiphenyl (PCB 209) (lipid adjusted)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of lipid or parts per billion on a lipid-weight basis) for the U.S. population from the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.317	0.029	9
		07-08	.330	0.075	9
	40-59	05-06	.846	0.084	4
		07-08	1.21	0.17	6
60+	05-06	*	*	4	
	07-08	4.41	0.65	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	1.01	0.15	6
		07-08	1.44	0.39	6
60+	05-06	4.98	0.24	3	
	07-08	*	*	5	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

Limit of detection (LOD, see Data Analysis section) for Survey years 05-06 and 07-08 are 0.31 and 0.3.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6,6'-Decachlorobiphenyl (PCB 209) (whole weight)

Geometric mean and selected percentiles of serum concentrations (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the following page for data from later NHANES Survey periods.

	Survey years	Geometric mean	Selected percentiles				Sample size
		(95% conf. interval)	(95% confidence interval)				
			50th	75th	90th	95th	
Total	03-04	.009 (.007-.010)	.008 (.006-.009)	.021 (.017-.026)	.049 (.040-.058)	.073 (.059-.097)	1854
Age group							
12-19 years	03-04	*	< LOD	< LOD	.006 (.004-.018)	.017 (.005-.033)	578
20 years and older	03-04	.010 (.009-.012)	.009 (.008-.011)	.024 (.020-.028)	.053 (.044-.063)	.077 (.065-.105)	1276
Gender							
Males	03-04	.008 (.007-.010)	.008 (.006-.010)	.019 (.017-.023)	.039 (.032-.051)	.063 (.042-.077)	925
Females	03-04	.009 (.007-.010)	.008 (.006-.009)	.024 (.016-.032)	.057 (.049-.064)	.092 (.068-.134)	929
Race/ethnicity							
Mexican Americans	03-04	*	< LOD	.007 (.005-.010)	.018 (.010-.023)	.024 (.018-.035)	416
Non-Hispanic blacks	03-04	.009 (.008-.010)	.008 (.006-.009)	.022 (.018-.027)	.065 (.039-.077)	.097 (.065-.144)	455
Non-Hispanic whites	03-04	.009 (.008-.011)	.009 (.007-.011)	.024 (.020-.029)	.053 (.044-.061)	.076 (.063-.099)	864

< LOD means less than the limit of detection for the lipid adjusted serum level, which may vary for some chemicals by year and by individual sample.

* Not calculated: proportion of results below limit of detection was too high to provide a valid result.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6,6'-Decachlorobiphenyl (PCB 209) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†] (in ng/g of serum or parts per billion) for the U.S. population from the National Health and Nutrition Examination Survey.

See the previous page for data from earlier NHANES Survey periods.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Non-Hispanic White Male	12-19	05-06	*	*	9
		07-08	*	*	6
	20-39	05-06	.005	0.001	12
		07-08	.004	0.001	15
	40-59	05-06	.013	0.001	12
		07-08	.014	0.001	16
60+	05-06	.048	0.004	15	
	07-08	.045	0.004	23	
Non-Hispanic White Female	12-19	05-06	*	*	10
		07-08	*	*	7
	20-39	05-06	.002	0	16
		07-08	.003	0	13
	40-59	05-06	.010	0.001	13
		07-08	.011	0.001	17
60+	05-06	.080	0.01	17	
	07-08	.074	0.008	21	
Non-Hispanic Black Male	12-19	05-06	*	*	13
		07-08	*	*	6
	20-39	05-06	.005	0.001	6
		07-08	.003	0	6
	40-59	05-06	.026	0.003	5
		07-08	.017	0.003	6
60+	05-06	.089	0.017	5	
	07-08	.082	0.013	8	
Non-Hispanic Black Female	12-19	05-06	*	*	14
		07-08	*	*	5
	20-39	05-06	.003	0.001	7
		07-08	*	*	8
	40-59	05-06	.018	0.002	7
		07-08	.014	0.003	8
60+	05-06	.142	0.03	5	
	07-08	.096	0.015	7	

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Serum 2,2',3,3',4,4',5,5',6,6'-Decachlorobiphenyl (PCB 209) (whole weight)

Weighted arithmetic mean and unadjusted standard error of pooled serum concentrations[†]
(in ng/g of serum or parts per billion) for the U.S. population from
the National Health and Nutrition Examination Survey.

Category	Age (years)	Survey years	Weighted arithmetic mean	Unadjusted standard error	Number of pools [‡]
Mexican American Male	12-19	05-06	*	*	11
		07-08	*	*	6
	20-39	05-06	.002	0	9
		07-08	.002	0.001	9
	40-59	05-06	.007	0.001	4
		07-08	.008	0.001	6
60+	05-06	*	*	4	
	07-08	.031	0.004	5	
Mexican American Female	12-19	05-06	*	*	16
		07-08	*	*	5
	20-39	05-06	*	*	9
		07-08	*	*	8
	40-59	05-06	.007	0.001	6
		07-08	.010	0.003	6
	60+	05-06	.038	0.002	3
		07-08	*	*	5

[†]See Special Analysis of Pooled Samples for Select Chemicals for additional explanation.

[‡] Each pool was composed of serum from 8 persons.

Weighted arithmetic means are not comparable to weighted geometric means.

Unadjusted standard errors do not incorporate survey design effects.

* Not reported: proportion of results below limit of detection was too high to provide a valid result or standard error >30% of mean.

Biomonitoring Summary: http://www.cdc.gov/biomonitoring/NDL-PCBs_BiomonitoringSummary.html

Appendix C. References for Biomonitoring Analytical Methods

Tobacco Smoke

Cotinine

Bernert JT, McGuffey JE, Morrison MA, Pirkle JL. Comparison of serum and salivary cotinine measurements by a sensitive high-performance liquid chromatography/tandem mass spectrometry method as an indicator of exposure to tobacco smoke among smokers and nonsmokers. *J Anal Toxicol* 2000;24:333-9.

Bernert JT, Turner WE, Pirkle JL, Sosnoff CS, Akins JR, Waldrep MK, et al. Development and validation of a sensitive measurement of serum cotinine in both smokers and nonsmokers by liquid chromatography/atmospheric pressure ionization tandem mass spectrometry. *Clin Chem* 1997;43:2281-91.

NNAL

Xia Y, Bernert JT. Stability of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) in urine samples stored at various temperatures. *J Anal Toxicol* 2010;34:411-5.

Xia Y, McGuffey JE, Bhattacharyya S, Sellergren B, Yilmaz E, Wang L, et al. Analysis of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol in urine by extraction on a molecularly imprinted polymer column and liquid chromatography/atmospheric pressure ionization tandem mass spectrometry. *Anal Chem*. 2005;77(23):7639-45.

Environmental Phenols

Ye X, Kuklennyik Z, Needham LL, Calafat AM. Automated on-line column-switching HPLC-MS/MS method with peak focusing for the determination of nine environmental phenols in urine. *Anal Chem* 2005;77:5407-13.

Fungicides and Metabolites

Bravo R, Caltabiano LM, Fernandez C, Smith KD, Gallegos M, Whitehead RD, et al. Quantification of phenolic metabolites of environmental chemicals in human urine using gas chromatography-tandem mass spectrometry and isotope dilution quantification. *J Chromatog B* 2005;820(2):229-36.

Hill RH Jr, Shealy DB, Head SL, Williams CC, Bailey SL, Gregg M, et al. Determination of pesticide metabolites in human urine using isotope dilution technique and tandem mass spectrometry. *J Anal Toxicol* 1995;19(5):323-9.

Montesano MA, Olsson AO, Kuklennyik P, Needham LL, Bradman AS, Barr DB. Method for determination of acephate, methamidophos, omethoate, dimethoate, ethylenethiourea and propylenethiourea in human urine using high-performance liquid chromatography-atmospheric pressure chemical ionization tandem mass spectrometry. *J Expo Sci Environ Epidemiol*. 2007;17(4):321-30.

Herbicides and Metabolites

2,4-D and 2,4,5-T

Olsson AO, Baker SE, Nguyen JV, Romanoff LC, Udunka SO, Walker RD, et al. A liquid chromatography-tandem mass spectrometry multiresidue method for quantification of specific metabolites of organophosphorus pesticides, synthetic pyrethroids, selected herbicides, and DEET in human urine. *Anal Chem* 2004;76(9):2453-61.

Sulfonylurea Herbicides

Nguyen JV, Olsson AO, Bravo R, Needham LL, Barr DB. Quantification of atrazine, phenylurea, and sulfonylurea herbicide metabolites in urine by high-performance liquid chromatography-tandem mass spectrometry. *J Anal Toxicol.* 2007;31(4):181-6.

Baker SE, Olsson AO, Needham LL, Barr DB. High-performance liquid chromatography-tandem mass spectrometry method for quantifying sulfonylurea herbicides in human urine: reconsidering the validation process. *Anal Bioanal Chem* 2005; 383:963-76.

Insecticides and Pesticides

Carbamate Pesticide Metabolites

Bravo R, Caltabiano LM, Fernandez C, Smith KD, Gallegos M, Whitehead RD, et al. Quantification of phenolic metabolites of environmental chemicals in human urine using gas chromatography-tandem mass spectrometry and isotope dilution quantification. *J Chromatog B* 2005;820(2):229-36.

Hill RH Jr, Shealy DB, Head SL, Williams CC, Bailey SL, Gregg M, et al. Determination of pesticide metabolites in human urine using isotope dilution technique and tandem mass spectrometry. *J Anal Toxicol* 1995;19(5):323-9.

Organochlorine Pesticides and Metabolites (Individual and Pooled Samples)

Barr JB, Maggio VL, Barr DB, Turner WE, Sjodin A, Sandau CD, et al. New high-resolution mass spectrometric approach for the measurement of polychlorinated biphenyls and organochlorine pesticides in human serum. *J Chromatog B* 2003;794:137-48.

Jones R, Edenfield E, Anderson S, Zhang Y and Sjodin A. Semi-automated Extraction and Cleanup Method for Measuring Persistent Organic Pollutants in Human Serum. *Organohalogen Compounds* 2012;74:97-8.

Sjodin A, Jones RS, Lapeza CR, Focant J-F, McGahee EE III, Patterson DG. Semiautomated high-throughput extraction and cleanup method for the measurement of polybrominated diphenyl ethers, polybrominated biphenyls, and polychlorinated biphenyls in human serum. *Anal Chem* 2004;76:1921-7.

2,4,5- and 2,4,6-Trichlorophenol

Ye X, Kuklennyik Z, Needham LL, Calafat AM. Automated on-line column-switching HPLC-MS/MS method with peak focusing for the determination of nine environmental phenols in urine. *Anal Chem* 2005;77:5407-13.

Other Pesticide Metabolites (2,4- and 2,5-Dichlorophenol)

Ye X, Kuklennyik Z, Needham LL, Calafat AM. Automated on-line column-switching HPLC-MS/MS method with peak focusing for the determination of nine environmental phenols in urine. *Anal Chem* 2005;77:5407-13.

Organophosphorus Insecticides: Specific Metabolites

Montesano MA, Olsson AO, Kuklennyik P, Needham LL, Bradman AS, Barr DB. Method for determination of acephate, methamidophos, omethoate, dimethoate, ethylenethiourea and propylenethiourea in human urine using high-performance liquid chromatography-atmospheric pressure chemical ionization tandem mass spectrometry. *J Expo Sci Environ Epidemiol* 2007; 17(4):321-30.

Organophosphorus Insecticides: Dialkyl Phosphate Metabolites

Bravo R, Caltabiano LM, Weerasekera G, Whitehead RD, Fernandez C, Needham LL, et al. Measurement of dialkyl phosphate metabolites of organophosphorus pesticides in human urine using lyophilization with gas chromatography-tandem mass spectrometry and isotope dilution quantification. *J Expo Anal Environ Epidemiol* 2004;14:249-59.

Metals and Metalloids

Caldwell KL, Hartel J, Jarrett J, Jones RL. Inductively coupled plasma mass spectrometry to measure multiple toxic elements in urine in NHANES 1999-2000. *Atomic Spectroscopy*. 2005;26(1):1-7.

Chen HP, Paschal DC, Miller DT, Morrow JC. Determination of total and inorganic mercury in whole blood by on-line digestion with flow injection. *Atomic Spectroscopy* 1998;19:176-9.

Jarrett JM, Jones RL, Caldwell KL, Verdon CP. Total urine arsenic measurements using inductively coupled plasma mass spectrometry with a dynamic reaction cell. *Atomic Spectroscopy* 2007;28(4):113-22.

Miller, DT, Paschal DC, Gunter EW, Stroud PE, D'Angelo J. Determination of lead in blood using electrothermal atomization atomic absorption spectrometry with a L'vov platform and matrix modifier. *Analyst* 1987;112:1701-4.

Paschal DC, Ting BG, Morrow JC, Pirkle JL, Jackson RJ, Sampson EJ, et al. Trace metals in urine of United States residents: reference range concentrations. *Environ Res* 1998;76:53-9.

Stoeppler M, Brandt K. Determination of cadmium in whole blood and urine by electrothermal atomic-absorption spectrophotometry. *Fresenius A Anal Chem* 1980;300:372-80.

Verdon CP, Caldwell KL, Fresques MR, Jones RL. Determination of seven arsenic compounds in urine by HPLC-ICP-DRC-MS: a CDC population biomonitoring method. *Anal Bioanal Chem* 2008;393(3):939-47.

Parabens

Ye X, Kuklennyik Z, Bishop AM, Needham LL, Calafat AM. Quantification of the urinary concentrations of parabens in humans by on-line solid phase extraction-high performance liquid chromatography–isotope dilution tandem mass spectrometry. *J Chromatogr B* 2006;844:53-9.

Perchlorate and Other Anions

Valentin-Blasini L, Blount BC, Delinsky A. Quantification of iodide and sodium-iodide symporter inhibitors in human urine using ion chromatography tandem mass spectrometry. *J Chrom A* 2007;1155(1):40-6.

Valentin-Blasini L, Mauldin JP, Maple D and Blount BC. Analysis of perchlorate in human urine using ion chromatography and electrospray tandem mass spectrometry. *Anal Chem* 2005;77(8):2475-81.

Perfluorinated Compounds: Surfactants

Kuklennyik Z, Needham LL, Calafat AM. Measurement of 18 Perfluorinated Organic Acids and Amides in Human Serum Using On-Line Solid-Phase Extraction. *Anal Chem* 2005;77:6085-91.

Kuklennyik Z, Reich JA, Tully JS, Needham LL, Calafat AM. Automated solid phase extraction and measurement of perfluorinated organic acids and amides in human serum and milk. *Environ Sci Technol* 2004;38:3698-704.

Phthalate Metabolites

Blount BC, Milgram KE, Silva M, Malek N, Reidy J, Needham LL, et al. Quantitative detection of eight phthalate metabolites in human urine using HPLC-APCI-MS/MS. *Anal Chem* 2000;72:4127-34.

Kato K, Silva MJ, Needham LL, Calafat AM. Determination of 16 Phthalate Metabolites in Urine Using Automated Sample Preparation and On-line Preconcentration/High-Performance Liquid Chromatography/Tandem Mass Spectrometry. *Anal Chem* 2005;77(9):2985-91.

Silva MJ, Malek NA, Hodge CC, Reidy JA, Kato K, Barr DB, et al. Improved quantitative detection of 11 urinary phthalate metabolites in humans using liquid chromatography-atmospheric pressure chemical ionization tandem mass spectrometry. *J Chromatogr B* 2003;789:393-404.

Silva MJ, Samandar E, Preau JL, Reidy JA, Needham LL, Calafat AM. Quantification of 22 phthalate metabolites in human urine. *J Chromatogr B* 2007;860:106-12.

Silva MJ, Slakman AR, Reidy JA, Preau JL, Herbert AR, Samandar E, et al. Analysis of human urine for 15 phthalate metabolites using automated solid-phase extraction. *J Chromatogr B* 2004;805:161-7.

Phytoestrogens and Metabolites

Kuklenyik Z, Ye X, Reich JA, Needham LL, Calafat AM. Automated on-line and off-line solid phase extraction methods for measuring isoflavones and lignans in urine. *J Chromatogr Sci* 2004; 42:495-500. (NHANES 1999-2000)

Parker DL, Rybak ME, Pfeiffer CM. Phytoestrogen biomonitoring: an extractionless LC-MS/MS method for measuring urinary isoflavones and lignans by use of atmospheric pressure photoionization (APPI). *Anal Bioanal Chem* 2012;402:1123-1136. (NHANES 2005-2006)

Rybak ME, Parker DL, Pfeiffer CM. Determination of urinary phytoestrogens by HPLC-MS/MS: A comparison of atmospheric pressure chemical ionization (APCI) and electrospray ionization (ESI). *J Chromatogr R* 2008;861:145-150. (NHANES 2003-2004)

Valentin-Blasini L, Blount BC, Rogers HS, Needham LL. HPLC-MS/MS method for the measurement of seven phytoestrogens in human serum and urine. *J Expo Anal Environ Epidemiol* 2000;10:799-807. (NHANES 2001-2002)

Polycyclic Aromatic Hydrocarbon Metabolites

Li Z, Romanoff LC, Trinidad DA, Hussain N, Jones RS, Porter EN, et al. Measurement of urinary Monohydroxy polycyclic aromatic hydrocarbons using automated liquid-liquid extraction and gas chromatography/isotope dilution high-resolution mass spectrometry. *Anal Chem* 2006;78:5744-51.

Volatile Organic Compounds, including Disinfection By-Products

Alwis KU, Blount BC, Silva LK, Smith MM, Loose KH. Method for quantifying nitromethane in blood as a potential biomarker of halonitromethane exposure. *Environ Sci Technol* 2008;42(7): 2522-7.

Bonin MA, Silva LK, Smith MM, Ashley DL and Blount BC. Measurement of trihalomethanes and methyl tert-butyl ether in whole blood using gas chromatography with high-resolution mass spectrometry. *J Anal Toxicol* 2005;29(2):81-9.

Chambers DM, Blount BC, McElprang DO, Waterhouse MG, Morrow J. Picogram measurement of volatile n-alkanes (n-hexane through n-dodecane) in blood using solid-phase microextraction to assess nonoccupational petroleum-based fuel exposure. *Anal Chem* 2008;80(12): 4666-74.

Chambers DM, McElprang DO, Mauldin JP, Hughes TM and Blount BC. Identification and elimination of polysiloxane curing agent interference encountered in the quantification of low-picogram per milliliter methyl tert-butyl ether in blood by solid-phase microextraction headspace analysis. *Anal Chem* 2005;77(9): 2912-9.

Chambers DM, McElprang DO, Waterhouse MG, and Blount BC. An improved approach for accurate quantitation of benzene, toluene, ethylbenzene, xylene, and styrene in blood. *Anal Chem* 2006;78(15): 5375-83.

Polybrominated Diphenyl Ethers and PBB 153 (Pooled and Individual Samples)

Jones R, Edenfield E, Anderson S, Zhang Y, Sjodin A. Semi-automated extraction and cleanup method for measuring persistent organic pollutants in human serum. *Organohalogen Compounds* 2012;74: 97-8.

Sjodin A, Jones RS, Lapeza CR, Focant J-F, McGahee EE III, Patterson DG. Semiautomated high-throughput extraction and cleanup method for the measurement of polybrominated diphenyl ethers, polybrominated biphenyls, and polychlorinated biphenyls in human serum. *Anal Chem* 2004;76:1921-7.

Polybrominated Dibenzo-*p*-dioxins and Dibenzofurans (Pooled and Individual Samples)

Turner WE, Whitfield WE, Cash TP, DiPietro ES, Anderson CJ, McClure PC, Sjodin A. Same sample cleanup for PCDDs/PCDFs/cPCBs, PCNs and PBDDs/PBDFs in human serum and quantification by GC/ID-HRMS. *Organohalogen Compounds* 2011;73: 661-4.

Polychlorinated Dibenzo-*p*-dioxins and Dibenzofurans (Pooled and Individual Samples)

Patterson DG Jr., Alexander LR, Turner WE, Isaacs SG, Needham LL. The development and application of a high resolution mass spectrometry method for measuring polychlorinated dibenzo-*p*-dioxins and dibenzofurans in serum. In: Clement RE, Sui KM, Hill HH Jr., eds. *Instrumentation for Trace Organic Monitoring*. Chelsea, MI: Lewis Publishers; 1990. pp. 119-53.

Patterson DG Jr., Hampton L, Lapeza CR Jr., Belser WT, Green V, Alexander L, et al. High resolution gas chromatographic/high resolution mass spectrometric analysis of human serum on a whole weight and lipid basis for 2,3,7,8 TCDD. *Anal Chem* 1987;59: 2000-5.

Patterson, DG Jr., Wong L-Y, Turner WE, Caudill SP, DiPietro ES, McClure PC, et al. Levels in the U.S. population of those persistent organic pollutants (2003-2004) included in the Stockholm Convention or in other Long-Range Tran boundary Air Pollution Agreements. *Environ Sci Technol* 2009;43(4):1211-8.

Turner WE, DiPietro E, Cash TP, McClure PC, Patterson DG Jr., Shirkhan H. An improved SPE extraction and automated sample cleanup method for serum PCDDs, PCDFs, and coplanar PCBs. *Organohalogen Compounds* 1994; 19: 31-5.

Polychlorinated Biphenyls (Pooled and Individual Samples)

Barr JB, Maggio VL, Barr DB, Turner WE, Sjodin A, Sandau CD, et al. New high-resolution mass spectrometric approach for the measurement of polychlorinated biphenyls and organochlorine pesticides in human serum. *J Chromatog B* 2003;794:137-48.

Jones R, Edenfield E, Anderson S, Zhang Y and Sjodin A. Semi-automated extraction and cleanup method for measuring persistent organic pollutants in human serum. *Organohalogen Compounds* 2012;74:97-8.

Sjodin A, Jones RS, Lapeza CR, Focant J-F, McGahee EE III, Patterson DG. Semiautomated high-throughput extraction and cleanup method for the measurement of polybrominated diphenyl ethers, polybrominated biphenyls, and polychlorinated biphenyls in human serum. *Anal Chem* 2004;76:1921-7.

Appendix D. Limit of Detection Table

The analytical limit of detection (LOD) for each of the different chemical measurements is presented in the table below. The LOD is the level at which the measurement has a 95% probability of being greater than zero (Taylor, 1987). As analytical methods improve, LODs will often change. For this reason, LOD results are reported by survey periods (e.g., 1999-2000, 2001-2002, 2003-2004, 2005-2006, 2007-2008, 2009-2010).

Reference: Taylor JK. Quality Assurance of Chemical Measurements. Chelsea (MI): Lewis Publishing, 1987

Chemical	Matrix	Units	1999-2000	2001-2002	2003-2004	2005-2006	2007-2008	2009-2010
Tobacco Smoke								
Cotinine	serum	ng/mL	0.05	0.05	0.015	0.015	0.015	0.015
NNAL	urine	pg/mL					0.6	0.6
Disinfection By-Products								
Bromodichloromethane	blood	pg/mL		0.233	0.62	0.62		
Dibromochloromethane	blood	pg/mL		0.271	0.62	0.62		
Tribromomethane (Bromoform)	blood	pg/mL		0.596	1.5	1.0		
Trichloromethane (Chloroform)	blood	pg/mL		2.37	2.11	2.1		
Environmental Phenols								
Benzophenone-3	urine	µg/L			0.3	0.4	0.4	0.4
Bisphenol A	urine	µg/L			0.4	0.4	0.4	0.4
4- <i>tert</i> -Octylphenol	urine	µg/L				0.2	0.2	0.2
Triclosan	urine	µg/L			2.3	2.3	2.3	2.3
Fungicides and Metabolites								
<i>ortho</i> -Phenylphenol	urine	µg/L			0.1	0.1	0.1	0.2
Ethylene thiourea	urine	µg/L			0.1	0.24		
Pentachlorophenol	urine	µg/L			0.5			
Propylene thiourea	urine	µg/L			0.1	0.37		
Herbicides and Metabolites								
2,4-Dichlorophenoxyacetic acid	urine	µg/L	0.89	0.2	0.1			
2,4,5-Trichlorophenoxyacetic acid	urine	µg/L	1.1	0.1	0.1			
Sulfonyl Urea Herbicides								
Bensulfuron-methyl	urine	µg/L			0.05	0.05	0.05	
Chlorsulfuron	urine	µg/L			0.06	0.06	0.06	
Ethametsulfuron-methyl	urine	µg/L			0.1	0.1	0.1	
Foramsulfuron	urine	µg/L			0.05	0.05	0.05	
Halosulfuron	urine	µg/L			0.1	0.1	0.1	
Mesosulfuron-methyl	urine	µg/L			0.06	0.06	0.06	
Metsulfuron-methyl	urine	µg/L			0.05	0.05	0.05	
Nicosulfuron	urine	µg/L			0.1	0.1	0.1	
Oxasulfuron	urine	µg/L			0.06	0.06	0.06	
Primisulfuron-methyl	urine	µg/L			0.07	0.07	0.07	
Prosulfuron	urine	µg/L			0.05	0.05	0.05	
Rimsulfuron	urine	µg/L			0.05	0.05	0.05	
Sulfometuron-methyl	urine	µg/L			0.05	0.05	0.05	
Sulfosulfuron	urine	µg/L			0.1	0.1	0.1	
Thifensulfuron-methyl	urine	µg/L			0.08	0.08	0.08	
Triasulfuron	urine	µg/L			0.07	0.07	0.07	
Triflusulfuron-methyl	urine	µg/L			0.05	0.05	0.05	
Insecticides and Metabolites								
Carbamate Pesticide Metabolites								
Carbofuranphenol	urine	µg/L	0.4	0.4	0.4			
2-Isopropoxyphenol	urine	µg/L	1.1	0.4	0.4			
Organochlorine Pesticides and Metabolites								
Aldrin	serum	ng/g lipid		5.94	7.8			
Dieldrin	serum	ng/g lipid		10.5	7.8			
Endrin	serum	ng/g lipid		5.09	7.8			
Heptachlor epoxide	serum	ng/g lipid	14.6	10.5	7.8			
<i>o,p'</i> -DDT	serum	ng/g lipid	20.7	17.4	7.8			
2,4,5-Trichlorophenol	urine	µg/L			0.1	0.1	0.1	0.1
2,4,6-Trichlorophenol	urine	µg/L			0.5	0.5	0.5	0.5
Other Pesticide Metabolites								
2,4-Dichlorophenol	urine	µg/L			0.17	0.2	0.2	0.2
2,5-Dichlorophenol	urine	µg/L			0.12	0.2	0.2	0.2
Organophosphorus Insecticides: Specific Metabolites								
Acephate	urine	µg/L			0.1	0.25		
Dimethoate	urine	µg/L			0.1	0.25		
Methamidophos	urine	µg/L			0.1	0.37		
Omethoate	urine	µg/L			0.1	0.14		

Appendix D. Limit of Detection Table

The analytical limit of detection (LOD) for each of the different chemical measurements is presented in the table below. The LOD is the level at which the measurement has a 95% probability of being greater than zero (Taylor, 1987). As analytical methods improve, LODs will often change. For this reason, LOD results are reported by survey periods (e.g., 1999-2000, 2001-2002, 2003-2004, 2005-2006, 2007-2008, 2009-2010).

Reference: Taylor JK. Quality Assurance of Chemical Measurements. Chelsea (MI): Lewis Publishing, 1987

Chemical	Matrix	Units	1999-2000	2001-2002	2003-2004	2005-2006	2007-2008	2009-2010
Organophosphorus Insecticides: Dialkyl Phosphate Metabolites								
Diethylphosphate (DEP)	urine	µg/L	0.2	0.2	0.1	0.37	0.37	
Dimethylphosphate (DMP)	urine	µg/L	0.58	0.5	0.5	0.47	0.47	
Diethylthiophosphate (DETP)	urine	µg/L	0.09	0.1	0.2	0.56	0.56	
Dimethylthiophosphate (DMTP)	urine	µg/L	0.18	0.4	0.5	0.55	0.55	
Diethyldithiophosphate (DEDTP)	urine	µg/L	0.05	0.1	0.1	0.39	0.39	
Dimethyldithiophosphate (DMDTP)	urine	µg/L	0.08	0.1	0.1	0.51	0.51	
Metals and Metalloids								
Antimony	urine	µg/L	0.04	0.04	0.07	0.032	0.032	0.032
Arsenic, Total	urine	µg/L			0.74	0.74	0.74	0.74
Arsenic (V) Acid	urine	µg As/L			1.0	1.0	1.0	1.0
Arsenobetaine	urine	µg As/L			0.4	0.4	0.4	0.4
Arsenocholine	urine	µg As/L			0.6	0.6	0.6	0.6
Arsenous (III) Acid	urine	µg As/L			1.2	1.2	1.2	1.2
Dimethylarsinic Acid	urine	µg As/L			1.7	1.7	1.7	1.7
Monomethylarsonic Acid	urine	µg As/L			0.9	0.9	0.9	0.9
Trimethylarsine oxide	urine	µg As/L			1.0	1.0	1.0	1.0
Barium	urine	µg/L	0.12	0.12	0.31	0.12	0.12	0.12
Beryllium	urine	µg/L	0.13	0.13	0.13	0.072	0.072	0.072
Cadmium	blood	µg/L	0.3	0.3	0.14	0.2	0.2	0.2
Cadmium	urine	µg/L	0.06	0.06	0.06	0.042	0.042	0.042
Cesium	urine	µg/L	0.14	0.14	0.2	0.066	0.066	0.066
Cobalt	urine	µg/L	0.07	0.07	0.08	0.041	0.041	0.041
Lead	blood	µg/dL	0.3	0.3	0.28	0.25	0.25	0.25
Lead	urine	µg/L	0.1	0.1	0.33	0.1	0.1	0.1
Mercury, Inorganic	blood	µg/L			0.42	0.4	0.35	0.35
Mercury, Total	blood	µg/L			0.2	0.33	0.33	0.33
Mercury	urine	µg/L			0.14	0.11	0.08	0.08
Molybdenum	urine	µg/L	0.8	0.8	1.5	0.92	0.92	0.92
Platinum	urine	µg/L	0.04	0.04	0.07	0.009	0.009	0.009
Thallium	urine	µg/L	0.02	0.02	0.02	0.015	0.015	0.015
Tungsten	urine	µg/L	0.04	0.04	0.04	0.021	0.021	0.021
Uranium	urine	µg/L	0.004	0.004	0.005	0.002	0.002	0.002
Parabens								
Butyl paraben	urine	µg/L				0.2	0.2	0.2
Ethyl paraben	urine	µg/L				1.0	1.0	1.0
Methyl paraben	urine	µg/L				1.0	1.0	1.0
n-Propyl paraben	urine	µg/L				0.2	0.2	0.2
Perchlorate and Other Anions								
Nitrate	urine	mg/L		0.7		0.7	0.7	
Perchlorate	urine	µg/L		0.05	0.05	0.05	0.05	
Thiocyanate	urine	mg/L		0.02		0.02	0.02	
Perfluorinated Compounds: Surfactants								
Perfluorobutane sulfonic acid (PFBuS)	serum	µg/L			0.4	0.1	0.1	0.1
Perfluorodecanoic acid (PFDeA)	serum	µg/L	0.2		0.3	0.2	0.2	0.1
Perfluorododecanoic acid (PFDoA)	serum	µg/L	0.2		1.0	0.2	0.2	0.1
Perfluoroheptanoic acid (PFHpA)	serum	µg/L	0.4		0.3	0.4	0.4	0.1
Perfluorohexane sulfonic acid (PFHxS)	serum	µg/L	0.1		0.3	0.1	0.1	0.1
Perfluorononanoic acid (PFNA)	serum	µg/L	0.1		0.1	0.1	0.082	0.082
Perfluorooctanoic acid (PFOA)	serum	µg/L	0.1		0.1	0.1	0.1	0.1
Perfluorooctane sulfonic acid (PFOS)	serum	µg/L	0.2		0.4	0.2	0.2	0.2
Perfluorooctane sulfonamide (PFOSA)	serum	µg/L	0.05		0.2	0.1	0.1	0.1
2-(N-Ethyl-perfluorooctane sulfonamido) acetic acid (Et-PFOSA-AcOH)	serum	µg/L	0.2		0.4	0.2	0.2	0.1
2-(N-Methyl-perfluorooctane sulfonamido) acetic acid (Me-PFOSA-AcOH)	serum	µg/L	0.2		0.5	0.2	0.2	0.1
Perfluoroundecanoic acid (PFUA)	serum	µg/L	0.2		0.3	0.2	0.2	0.1

Appendix D. Limit of Detection Table

The analytical limit of detection (LOD) for each of the different chemical measurements is presented in the table below. The LOD is the level at which the measurement has a 95% probability of being greater than zero (Taylor, 1987). As analytical methods improve, LODs will often change. For this reason, LOD results are reported by survey periods (e.g., 1999-2000, 2001-2002, 2003-2004, 2005-2006, 2007-2008, 2009-2010).

Reference: Taylor JK. Quality Assurance of Chemical Measurements. Chelsea (MI): Lewis Publishing, 1987

Chemical	Matrix	Units	1999-2000	2001-2002	2003-2004	2005-2006	2007-2008	2009-2010
Phthalate Metabolites								
Mono-benzyl phthalate (MBzP)	urine	µg/L	0.576	0.216	0.072	0.216	0.216	0.216
Mono-isobutyl phthalate (MiBP)	urine	µg/L		1.0	0.3	0.3	0.3	0.2
Mono- <i>n</i> -butyl phthalate (MnBP)	urine	µg/L	0.9	1.1	0.4	0.6	0.6	0.4
Mono-cyclohexyl phthalate (MCHP)	urine	µg/L	1.81	0.603	0.402	0.603	0.603	0.402
Mono-ethyl phthalate (MEP)	urine	µg/L	0.792	0.594	0.264	0.528	0.462	0.396
Mono-2-ethylhexyl phthalate (MEHP)	urine	µg/L	1.2	1.0	0.9	1.2	1.1	0.5
Mono-(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP)	urine	µg/L		1.0	0.3	0.7	0.7	0.2
Mono-(2-ethyl-5-oxohexyl) phthalate (MEOHP)	urine	µg/L		1.1	0.5	0.7	0.6	0.2
Mono-(2-ethyl-5-carboxypentyl) phthalate (MECPP)	urine	µg/L			0.3	0.6	0.5	0.2
Mono-(carboxynonyl) phthalate (MCNP)	urine	µg/L				0.6	0.5	0.2
Mono-isononyl phthalate (MiNP)	urine	µg/L	1.23	1.23	1.54	1.23	1.23	0.77
Mono-(carboxyoctyl) phthalate (MCOP)	urine	µg/L				0.7	0.7	0.2
Mono-methyl phthalate (MMP)	urine	µg/L		0.2	1.0	1.1	1.1	0.5
Mono-(3-carboxypropyl) phthalate (MCPP)	urine	µg/L		0.4	0.2	0.2	0.2	0.2
Mono- <i>n</i> -octyl phthalate (MOP)	urine	µg/L	1.51	1.68	1.68	1.85	1.85	0.84
Phytoestrogens and Metabolites								
Urinary Daidzein	urine	µg/L	0.5	1.6	0.3	0.4	0.4	0.4
Urinary Enterodiol	urine	µg/L	0.8	1.5	0.3	0.04	0.04	0.04
Urinary Enterolactone	urine	µg/L	0.6	1.9	0.3	0.1	0.1	0.1
Urinary Equol	urine	µg/L	3.0	3.3	0.3	0.06	0.06	0.06
Urinary Genistein	urine	µg/L	0.3	0.8	0.3	1.0	0.2	0.2
Urinary O-Desmethylangolensin	urine	µg/L	0.2	0.4	0.2	0.2	0.2	0.2
Polycyclic Aromatic Hydrocarbon Metabolites								
2-Hydroxyfluorene	urine	ng/L			5.0	5.0	5.0	
3-Hydroxyfluorene	urine	ng/L		2.0	5.0	5.0	5.0	
9-Hydroxyfluorene	urine	ng/L			5.0	5.0	5.0	
1-Hydroxyphenanthrene	urine	ng/L		3.5	5.0	5.0	5.0	
2-Hydroxyphenanthrene	urine	ng/L		3.2	5.0	5.0	5.0	
3-Hydroxyphenanthrene	urine	ng/L		3.6	5.0	5.0	5.0	
4-Hydroxyphenanthrene	urine	ng/L			5.0	5.0		
1-Hydroxypyrene	urine	ng/L			5.0	5.0	5.0	
1-Hydroxynaphthalene (1-Naphthol)	urine	ng/L			46.7	47.9	44.7	
2-Hydroxynaphthalene (2-Naphthol)	urine	ng/L		2.4	31.1	13.2	42.0	
Volatile Organic Compounds								
1,1,1-Trichloroethane (Methyl chloroform)	blood	ng/mL			0.048	0.01		
1,1,2,2-Tetrachloroethane	blood	ng/mL			0.05	0.01		
1,1,2-Trichloroethane	blood	ng/mL			0.01	0.01		
1,1-Dichloroethane	blood	ng/mL			0.01	0.01		
1,1-Dichloroethane (Vinylidene chloride)	blood	ng/mL			0.009	0.009		
1,2-Dibromo-3-chloropropane (DBCP)	blood	ng/mL			0.1	0.1		
1,2-Dichlorobenzene	blood	ng/mL			0.1	0.1		
1,2-Dichloroethane (Ethylene dichloride)	blood	ng/mL			0.01	0.01		
<i>cis</i> -1,2-Dichloroethene	blood	ng/mL			0.01	0.01		
<i>trans</i> -1,2-Dichloroethene	blood	ng/mL			0.01	0.01		
1,2-Dichloropropane	blood	ng/mL			0.008	0.008		
1,3-Dichlorobenzene	blood	ng/mL			0.05	0.05		
1,4-Dichlorobenzene (Paradichlorobenzene)	blood	ng/mL		0.12	0.12	0.011		
2,5-Dimethylfuran	blood	ng/mL			0.012	0.011		
Benzene	blood	ng/mL		0.024	0.024	0.024		
Chlorobenzene	blood	ng/mL			0.011	0.011		
Dibromomethane	blood	ng/mL			0.03	0.03		
Dichloromethane (Methylene chloride)	blood	ng/mL			0.07	0.25		
Ethylbenzene	blood	ng/mL		0.024	0.024	0.024		
Hexachloroethane	blood	ng/mL			0.011	0.011		
Methyl- <i>tert</i> -butyl ether (MTBE)	blood	pg/mL		0.232	2.0	1.4		
Nitrobenzene	blood	ng/mL			0.3	0.3		
Styrene	blood	ng/mL		0.03	0.03	0.03		
Tetrachloroethene (Perchloroethylene)	blood	ng/mL		0.048	0.048	0.048		
Tetrachloromethane (Carbon tetrachloride)	blood	ng/mL		0.01	0.005	0.005		
Toluene	blood	ng/mL		0.025	0.025	0.025		
Trichloroethene (Trichloroethylene)	blood	ng/mL		0.012	0.012	0.012		
<i>m</i> - <i>p</i> -Xylene	blood	ng/mL		0.340	0.034	0.034		
<i>o</i> -Xylene	blood	ng/mL		0.049	0.049	0.024		

Appendix D. Limit of Detection Table

The analytical limit of detection (LOD) for each of the different chemical measurements is presented in the table below. The LOD is the level at which the measurement has a 95% probability of being greater than zero (Taylor, 1987). As analytical methods improve, LODs will often change. For this reason, LOD results are reported by survey periods (e.g., 1999-2000, 2001-2002, 2003-2004, 2005-2006, 2007-2008, 2009-2010).

Reference: Taylor JK. Quality Assurance of Chemical Measurements. Chelsea (MI): Lewis Publishing, 1987

Chemical	Matrix	Units	1999-2000	2001-2002	2003-2004	2005-2006	2007-2008	2009-2010
Organochlorine Pesticides and Metabolites (Pooled Samples beginning in 2005-2006)								
Oxychlorodane	serum	ng/g lipid	14.5	10.5	7.8	1.46	1.4	
<i>trans</i> -Nonachlor	serum	ng/g lipid	14.5	10.5	7.8	0.77	1.4	
<i>p,p'</i> -DDT	serum	ng/g lipid	20.7	17.4	7.8	1.46	1.4	
<i>p,p'</i> -DDE	serum	ng/g lipid	18.6	8.3	7.8	4.02	1.4	
Hexachlorobenzene	serum	ng/g lipid	118.0	31.4	7.8	4.17	1.4	
<i>beta</i> -Hexachlorocyclohexane	serum	ng/g lipid	9.36	6.76	7.8	1.46	1.4	
<i>gamma</i> -Hexachlorocyclohexane	serum	ng/g lipid	14.5	10.5	7.8	1.46	1.4	
Mirex	serum	ng/g lipid	14.6	10.5	7.8	1.46	1.4	
Polybrominated Diphenyl Ethers and PBB 153 (Pooled Samples beginning in 2005-2006)								
2,2',4'-Tribromodiphenyl ether (BDE 17)	serum	ng/g lipid			1.0	0.6	0.6	
2,4,4'-Tribromodiphenyl ether (BDE 28)	serum	ng/g lipid			0.8	0.6	0.6	
2,2',4,4'-Tetrabromodiphenyl ether (BDE 47)	serum	ng/g lipid			4.2	1.6	0.6	
2,3',4,4'-Tetrabromodiphenyl ether (BDE 66)	serum	ng/g lipid			1.0	0.6	0.6	
2,2',3,4,4'-Pentabromodiphenyl ether (BDE 85)	serum	ng/g lipid			2.4	0.6	0.6	
2,2',4,4',5-Pentabromodiphenyl ether (BDE 99)	serum	ng/g lipid			5.0	1.2	0.6	
2,2',4,4',6-Pentabromodiphenyl ether (BDE 100)	serum	ng/g lipid			1.4	0.6	0.6	
2,2',4,4',5,5'-Hexabromodiphenyl ether (BDE 153)	serum	ng/g lipid			2.2	0.6	0.6	
2,2',4,4',5,6'-Hexabromodiphenyl ether (BDE 154)	serum	ng/g lipid			0.8	0.6	0.6	
2,2',3,4,4',5',6-Heptabromodiphenyl ether (BDE 183)	serum	ng/g lipid			1.7	0.6	0.6	
2,2',3,3',4,4',5,5',6,6'-Decabromodiphenyl ether (BDE 209)	serum	ng/g lipid				6.0	5.8	
2,2',4,4',5,5'-Hexabromobiphenyl (PBB 153)	serum	ng/g lipid			0.8	0.6	0.6	
Polybrominated Dibenzo-<i>p</i>-dioxins (Pooled Samples)								
1,2,3,4,6,7,8-Heptabromodibenzo- <i>p</i> -dioxin (HpBDD)	serum	pg/g lipid				11.1		
1,2,3,4,7,8 and 1,2,3,6,7,8-Hexabromodibenzo- <i>p</i> -dioxin (HxBDD)	serum	pg/g lipid				19.9		
1,2,3,7,8,9-Hexabromodibenzo- <i>p</i> -dioxin (HxBDD)	serum	pg/g lipid				11.1		
1,2,3,7,8-Pentabromodibenzo- <i>p</i> -dioxin (PeBDD)	serum	pg/g lipid				8.8		
2,3,7,8-Tetrabromodibenzo- <i>p</i> -dioxin (TBDD)	serum	pg/g lipid				6.6		
Polybrominated Dibenzofurans (Pooled Samples)								
1,2,3,4,6,7,8-Heptabromodibenzofuran (HpBDF)	serum	pg/g lipid				15.5		
1,2,3,4,7,8-Hexabromodibenzofuran (HxBDF)	serum	pg/g lipid				11.1		
1,2,3,7,8-Pentabromodibenzofuran (PeBDF)	serum	pg/g lipid				8.8		
2,3,4,7,8-Pentabromodibenzofuran (PeBDF)	serum	pg/g lipid				8.8		
2,3,7,8-Tetrabromodibenzofuran (TBDF)	serum	pg/g lipid				11.1		
Polychlorinated Dibenzo-<i>p</i>-dioxins (Pooled Samples beginning in 2005-2006)								
1,2,3,4,6,7,8-Heptachlorodibenzo- <i>p</i> -dioxin (HpCDD)	serum	pg/g lipid	55.9	10.3	13.0	0.6		
1,2,3,4,7,8-Hexachlorodibenzo- <i>p</i> -dioxin (HxCDD)	serum	pg/g lipid		9.0	11.9	0.4		
1,2,3,6,7,8-Hexachlorodibenzo- <i>p</i> -dioxin (HxCDD)	serum	pg/g lipid	20.1	9.1	12.3	0.4		
1,2,3,7,8,9-Hexachlorodibenzo- <i>p</i> -dioxin (HxCDD)	serum	pg/g lipid	20.3	9.3	12.3	0.4		
1,2,3,4,6,7,8,9-Octachlorodibenzo- <i>p</i> -dioxin (OCDD)	serum	pg/g lipid	329.0	319.0	218.0	7.7		
1,2,3,7,8-Pentachlorodibenzo- <i>p</i> -dioxin (PeCDD)	serum	pg/g lipid	14.2	6.0	4.5	0.3		
2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin (TCDD)	serum	pg/g lipid	12.1	5.8	3.8	0.3		

Appendix D. Limit of Detection Table

The analytical limit of detection (LOD) for each of the different chemical measurements is presented in the table below. The LOD is the level at which the measurement has a 95% probability of being greater than zero (Taylor, 1987). As analytical methods improve, LODs will often change. For this reason, LOD results are reported by survey periods (e.g., 1999-2000, 2001-2002, 2003-2004, 2005-2006, 2007-2008, 2009-2010).

Reference: Taylor JK. Quality Assurance of Chemical Measurements. Chelsea (MI): Lewis Publishing, 1987

Chemical	Matrix	Units	1999-2000	2001-2002	2003-2004	2005-2006	2007-2008	2009-2010
Polychlorinated Dibenzofurans (Pooled Samples beginning in 2005-2006)								
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	serum	pg/g lipid	13.5	7.0	8.6	0.3		
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	serum	pg/g lipid		7.0	8.6	0.3		
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	serum	pg/g lipid	12.7	6.5	7.4	0.3		
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	serum	pg/g lipid	12.6	6.1	7.9	0.3		
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	serum	pg/g lipid	12.7	6.0	8.3	0.3		
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	serum	pg/g lipid	12.9	5.8	8.2	0.3		
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	serum	pg/g lipid	35.6	21.0	12.0	0.5		
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	serum	pg/g lipid	13.2	5.8	7.1	0.3		
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	serum	pg/g lipid	12.7	5.5	6.8	0.3		
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	serum	pg/g lipid	11.9	5.2	6.0	0.3		
Dioxin-like Polychlorinated Biphenyls: Coplanar PCBs (Pooled Samples beginning in 2005-2006)								
3,3',4,4'-Tetrachlorobiphenyl (PCB 77)	serum	pg/g lipid				0.6		
3,4,4',5'-Tetrachlorobiphenyl (PCB 81)	serum	pg/g lipid	68.4	26.8	13.1	0.6		
3,3',4,4',5'-Pentachlorobiphenyl (PCB 126)	serum	pg/g lipid	23.2	10.8	13.9	0.5		
3,3',4,4',5,5'-Hexachlorobiphenyl (PCB 169)	serum	pg/g lipid	27.0	11.0	15.9	0.6		
Dioxin-like Polychlorinated Biphenyls: Mono-ortho-substituted PCBs (Pooled Samples beginning in 2005-2006)								
2,3,3',4,4'-Pentachlorobiphenyl (PCB 105)	serum	ng/g lipid	12.4	10.5	0.4	0.31	0.3	
2,3,4,4',5'-Pentachlorobiphenyl (PCB 114)	serum	ng/g lipid				0.31	0.3	
2,3',4,4',5'-Pentachlorobiphenyl (PCB 118)	serum	ng/g lipid	12.5	10.5	0.6	0.72	0.3	
2',3,4,4',5'-Pentachlorobiphenyl (PCB 123)	serum	ng/g lipid				0.31	0.3	
2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 156)	serum	ng/g lipid	12.5	10.5	0.4	0.31	0.3	
2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157)	serum	ng/g lipid	12.5	10.5	0.4	0.31	0.3	
2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167)	serum	ng/g lipid	12.4	10.5	0.4	0.31	0.3	
2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189)	serum	ng/g lipid		10.5	0.4	0.31	0.3	
Non-Dioxin-Like Polychlorinated Biphenyls (Pooled Samples beginning in 2005-2006)								
2,4,4'-Trichlorobiphenyl (PCB 28)	serum	ng/g lipid	32.4		1.7	2.54	0.7	
2,2',3,5'-Tetrachlorobiphenyl (PCB 44)	serum	ng/g lipid			0.4	0.36	0.3	
2,2',4,5'-Tetrachlorobiphenyl (PCB 49)	serum	ng/g lipid			0.4	0.31	0.3	
2,2',5,5'-Tetrachlorobiphenyl (PCB 52)	serum	ng/g lipid	12.5	12.4	0.8	0.33	0.3	
2,3',4,4'-Tetrachlorobiphenyl (PCB 66)	serum	ng/g lipid	12.4	12.4	0.8	0.4	0.3	
2,4,4',5'-Tetrachlorobiphenyl (PCB 74)	serum	ng/g lipid	12.4	10.5	0.8	0.31	0.8	
2,2',3,4,5'-Pentachlorobiphenyl (PCB 87)	serum	ng/g lipid		10.5	0.4	0.31	0.3	
2,2',4,4',5'-Pentachlorobiphenyl (PCB 99)	serum	ng/g lipid	12.5	10.5	0.6	0.31	0.3	
2,2',4,5,5'-Pentachlorobiphenyl (PCB 101)	serum	ng/g lipid	25.7	10.5	0.6	0.59	0.3	
2,3,3',4',6'-Pentachlorobiphenyl (PCB 110)	serum	ng/g lipid		10.5	0.8	0.31	0.3	
2,2',3,3',4,4'-Hexachlorobiphenyl (PCB 128)	serum	ng/g lipid	12.4	10.5	0.4	0.31	0.3	
2,2',3,4,4',5' and 2,3,3',4,4',6'-Hexachlorobiphenyl (PCB 138 & 158)	serum	ng/g lipid	41.1	10.5	0.4	0.31	0.3	
2,2',3,4',5,5'-Hexachlorobiphenyl (PCB 146)	serum	ng/g lipid	12.4	10.5	0.4	0.31	0.3	
2,2',3,4',5',6'-Hexachlorobiphenyl (PCB 149)	serum	ng/g lipid	10.5		0.4	0.31	0.3	
2,2',3,5,5',6'-Hexachlorobiphenyl (PCB 151)	serum	ng/g lipid		10.5	0.4	0.31	0.3	
2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153)	serum	ng/g lipid	55.6	10.5	1.1	0.31	0.3	
2,2',3,3',4,4',5'-Heptachlorobiphenyl (PCB 170)	serum	ng/g lipid	17.2	10.5	0.4	0.31	0.3	
2,2',3,3',4,5,5'-Heptachlorobiphenyl (PCB 172)	serum	ng/g lipid	12.5	10.5	0.4	0.31	0.3	
2,2',3,3',4,5',6'-Heptachlorobiphenyl (PCB 177)	serum	ng/g lipid	12.5	10.5	0.4	0.31	0.3	

Appendix D. Limit of Detection Table

The analytical limit of detection (LOD) for each of the different chemical measurements is presented in the table below. The LOD is the level at which the measurement has a 95% probability of being greater than zero (Taylor, 1987). As analytical methods improve, LODs will often change. For this reason, LOD results are reported by survey periods (e.g., 1999-2000, 2001-2002, 2003-2004, 2005-2006, 2007-2008, 2009-2010).

Reference: Taylor JK. Quality Assurance of Chemical Measurements. Chelsea (MI): Lewis Publishing, 1987

Chemical	Matrix	Units	1999-2000	2001-2002	2003-2004	2005-2006	2007-2008	2009-2010
Non-Dioxin-Like Polychlorinated Biphenyls (Pooled Samples beginning in 2005-2006)								
2,2',3,3',5,5',6-Heptachlorobiphenyl (PCB 178)	serum	ng/g lipid	12.4	10.5	0.4	0.31	0.3	
2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180)	serum	ng/g lipid	28.2	10.5	0.4	0.31	0.3	
2,2',3,4,4',5',6-Heptachlorobiphenyl (PCB 183)	serum	ng/g lipid	12.4	10.5	0.4	0.31	0.3	
2,2',3,4',5,5',6-Heptachlorobiphenyl (PCB 187)	serum	ng/g lipid	12.4	10.5	0.4	0.31	0.3	
2,2',3,3',4,4',5,5'-Octachlorobiphenyl (PCB 194)	serum	ng/g lipid		10.5	0.4	0.31	0.3	
2,2',3,3',4,4',5,6-Octachlorobiphenyl (PCB 195)	serum	ng/g lipid		28.1	0.7	0.31	0.3	
2,2',3,3',4,4',5,6' and 2,2',3,4,4',5,5',6-Octachlorobiphenyl (PCB 196 & 203)	serum	ng/g lipid		10.5	0.4	0.31	0.3	
2,2',3,3',4,5,5',6-Octachlorobiphenyl (PCB 199)	serum	ng/g lipid		10.5	0.4	0.31	0.3	
2,2',3,3',4,4',5,5',6-Nonachlorobiphenyl (PCB 206)	serum	ng/g lipid		28.1	0.7	0.31	0.3	
2,2',3,3',4,4',5,5',6,6'-Decachlorobiphenyl (PCB 209)	serum	ng/g lipid			0.7	0.31	0.3	



Centers for Disease Control and Prevention
National Center for Environmental Health
Division of Laboratory Sciences
Mail Stop F-20
4770 Buford Highway, NE
Atlanta, GA 30341-3724

Telephone: (toll free) 1-800-CDC-INFO (1-800-232-4636)
Email: CDCCINFO@cdc.gov
Website: <http://www.cdc.gov/exposurereport>