

HONEY MESQUITE

Prosopis glandulosa Torr. var.
torreyana (L. Benson) M.C.
Johnston
Plant Symbol = PRGLT

Contributed by: USDA NRCS National Plant Data Center & East Bay Regional Park Botanic Garden



Charles Webber
©California Academy of Sciences
@ CalPhotos

Uses

Ethnobotanic: Many parts of honey mesquite were useful to indigenous groups of California, the Southwest and northern Mexico, and some of these uses are still enacted today. The Cahuilla picked the flowers of honey mesquite in the spring and roasted them in a pit of heated stones, after which they were formed into balls and eaten. The blossoms were also utilized in making a tea. Many tribes in southern California and the Southwest including the Timbisha Shoshone, Nevada and Utah Southern Paiute, Kawaiisu, Cahuilla, Panamint, Luiseno, Pima, Yuma, and Commanche ate the edible pods. The seeds made into a fermented pinole was a favorite intoxicating drink. The Cahuilla gathered the pods in two stages, green or ripe. The green pods were pounded into a juicy pulp and consumed as a summer beverage. The ripe pods were eaten without any processing or were pounded into a flour in a mortar and then placed in a vessel, dampened with water, and left for 24 hours to harden. The meal was formed into cakes and eaten dry, made into a mush, or mixed with water for a beverage.

All other parts of honey mesquite were used. The leaves were used to neutralize stomach acidity by the Commanche. The Cahuilla, Kamia, and Yuma tribes used the inner bark of trunks or branches for cordage and rough wrapping material. The roots of young trees were harvested and made into cordage for harpoon lines, to bundle reeds, and make carrying nets by the Seri of northwestern Mexico. The trunk was made into wooden mortars and the smaller branches were used for bow making. Sometimes arrow foreshafts were of mesquite wood. Mesquite made excellent firewood for cooking, baking pottery, and warmth and mesquite bark was good for kindling. Large branches were used for construction such as posts for houses and rafters. The Seri use mesquite to make the ribs of wooden boats. Thorns were used as a needle in puncturing the skin for tattoos. Mesquite gum was used to glue foreshafts of arrows and adhere baskets to mortars. Tribes of the southwest used the gum secreted from the bark to mend pottery, make candy, and furnish a black dye.

Wildlife: Honey mesquite is an important tree to wildlife. The seeds are eaten by jack rabbits, Gambel quail, songbirds, various small mammal, and domestic livestock. Western chipmunks, ground squirrels, pocket mice, and various species of kangaroo and wood rats consume the foliage. Different birds also nest in the tree's canopy.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

Description

General: Pea Family (Fabaceae). The honey mesquite can penetrate downward into the water table, often at great depths. This shrub or small tree reaches less than 10 m in height and has a broad crown of arched branchlets. The ascending branches have straight, stipular spines. The jugate leaves have many long, narrow, leaflets that are more than five times longer than wide, glabrous, and widely spaced on the rachilla or stem. The greenish yellow flowers have fused petals and in dense racemes, 5-12 cm. and spikelike. The fruit is a pod with a prominent beak at the end.

Distribution

For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site. This plant is the common mesquite tree of the subtropical scrub south and southeast of the Sonoran desert in Sonora. In the desert it is found northward from Guaymas on the coast to the Colorado River and into the low elevation deserts of southeastern California and western Arizona. It occurs in plains, riverbanks, washes and playas in grasslands and deserts below 1800 m.

Establishment

Gather seed pods and remove the seeds from the pods. Scarify the seeds before planting. Plant scarified seeds in the fall in a well-drained soil in deep pots. Sprinkle soil on top of the seeds and cover soil with one-quarter inch of gravel. The seeds should be spaced one-half inch apart and the pots placed in partial shade. Water the pots right away. Keep pots moist if the rains are insufficient. As soon as the plants form one true leaf, transplant one plant per pot and water. Keep the pots damp. Out plant each seedling, in the ground, the following winter when dormant in full sun, and well-drained soil. Protect the plants from animals. Water the plants and keep them damp if annual rains are insufficient. Continue to water throughout the life of the plant once in awhile. Mesquite should be lightly damp all summer long.

Management

The Timbisha Shoshone pruned honey mesquite, keeping areas around the trees clear of undergrowth, and also of dead limbs and lower branches. The Cahuilla in southern California also pruned mesquite trees and broke and cut branches regularly to provide easier access to the seedpods.

Cultivars, Improved and Selected Materials (and area of origin)

This plant is available through most sources in the southwest specializing in native seed. Contact your local Natural Resources Conservation Service (formerly Soil Conservation Service) office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation Service will be listed under the subheading "Department of Agriculture."

References

Barney, R.C. 1989. *Prosopis*. Pages 4-6 IN: *Fabales, Vol. 3, Part B in Intermountain Flora: Vascular Plants of the Intermountain West, U.S.A.* Cronquist, A., A.H. Holmgren, N.H. Holmgren, J.L.

Reveal, & P.K. Holmgren (authors). New York Botanical Garden, Bronx, New York.

Bean L.J. & K.S. Saubel 1972. *Temalpakh: Cahuilla Indian knowledge and usage of plants*. Malki Museum Press, Morongo Indian Reservation, Banning, California.

Bell, W.H. & E. F. Castetter 1937. *The utilization of mesquite and screwbean by the aborigines in the American Southwest*. Ethnobiol. Studies in the American Southwest Bulletin 5. Biological Service. University of New Mexico 4(5):3-63.

Carlson, G.G. and V.H. Jones 1939. *Some notes on uses of plants by the Commanche Indians*. Michigan Academy of Science, Arts, and Letters Paper 25:517-542.

Coville, F.V. 1892. *The Panamint Indians of California*. The American Anthropologist 5(4):351-362.

Emery, D.E. 1988. *Seed propagation of native California plants*. Santa Barbara Botanic Garden, Santa Barbara, California.

Felger, R.S. and M.B. Moser 1985. *People of the desert and sea: Ethnobotany of the Seri Indians*. University of Arizona Press, Tucson, Arizona.

Fowler, C.S. 1986. *Subsistence*. Pages 64-97 IN: *Handbook of North American Indians Volume 11, Great Basin*. Smithsonian Institution, Washington, D.C.

Fowler, C.S. 1996. *Historical perspectives on Timbisha Shoshone land management practices, Death Valley, California*. IN: *Case Studies in Environmental Archaeology*. E.J. Reitz, L.A. Newsom, and S.J. Scudder (eds.). Plenum Press, New York, New York.

Krochmal, A., S. Paur, and P. Duisberg 1954. *Useful native plants in the American Southwestern deserts*. Economic Botany 8(2):3-20.

Martin, A.C., H.S. Zim, & A.L. Nelson 1951. *American wildlife and plants: A guide to wildlife food habits*. Dover Publications, New York, New York.

Shreve, F. & I.L. Wiggins 1964. *Prosopis*. Pages 602-605, IN: *Vegetation and Flora of the Sonoran Desert Vol. I*. Stanford University Press, Stanford, California.

Prepared By & Species Coordinators

M. Kat Anderson

USDA, NRCS, National Plant Data Center
c/o Plant Sciences Department University of
California, Davis, California

Wayne Roderick

Former Director of the East Bay Regional Parks
Botanic Garden, Berkeley, California

Edited: 30nov00 jsp; 29may03 ahv; 060808 jsp

For more information about this and other plants, please contact
your local NRCS field office or Conservation District, and visit the
PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials
Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

*The U.S. Department of Agriculture (USDA) prohibits
discrimination in all its programs and activities on the basis of
race, color, national origin, sex, religion, age, disability, political
beliefs, sexual orientation, and marital or family status. (Not all
prohibited bases apply to all programs.) Persons with disabilities
who require alternative means for communication of program
information (Braille, large print, audiotape, etc.) should contact
USDA's TARGET Center at 202-720-2600 (voice and TDD).*

*To file a complaint of discrimination write USDA, Director, Office
of Civil Rights, Room 326-W, Whitten Building, 14th and
Independence Avenue, SW, Washington, DC 20250-9410 or call
202-720-5964 (voice or TDD). USDA is an equal opportunity
provider and employer.*

*Read about [Civil Rights at the Natural Resources Conservation
Service](#).*