

# **Plant Guide**

## **SLIPPERY ELM**

### Ulmus rubra Muhl.

Plant Symbol = ULRU

Contributed By: USDA NRCS National Plant Data Center



Robert Mohlenbrock USDA, NRCS, Wetland Science Institute @ PLANTS

#### Alternate Names Red elm

#### Uses

Ethnobotanic: This tree was valued for its bark, which supplied material for the sides of winter houses and roofs of the Meskwaki. The inner bark was used for cordage by many tribes. The Menomini gathered the bark, boiled it, and used it for making fiber bags and large storage baskets. The Dakota, Omaha-Ponca, Winnebago, and Pawnee used the inner bark fiber for making ropes and cords. Slippery elm was also used extensively as a medicine. The Iroquois scraped the bark of the tree and used it in combination with other plants to treat infected and swollen glands. The inner bark was made into an eye wash for sore eyes. The Menomini used the inner bark in a tea and it was taken as a physic. The inner bark was used by the Menomini and the Meskwaki in a poultice to heal sores on the body. Meskwaki women drank a tea of the bark to make childbirth easier. The tree also was used by the Ojibwe to treat sore throats. The fresh inner bark was boiled and the Dakota, Omaha-Ponca, and other tribes drank the resulting decoction as a laxative. The indigenous people generously taught some of these uses to early non-Indian settlers. Today slippery elm is found in health food stores and is used to relieve sore throats,

coughs and other bronchial ailments, and used as a laxative. The wood is used commercially for making furniture, paneling, and containers.

*Wildlife:* Birds often nest in the thick elm foliage, and the seeds and buds are food to songbirds, game birds, and squirrels. Deer and rabbits browse on the twigs.

#### Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status, and wetland indicator values.

#### **Description**

General: Elm Family (Ulmaceae). This graceful, arching tree reaches 20 m, with twigs that are scabrous-pubescent. It can live to be 200 years old and is identified by its "slippery" inner bark. The winter-buds are densely covered with red-brown hairs. The leaves are oblong to obovate, thick and stiff and 10-20 cm. They are pinnately veined and not equilateral. The flowers are subsessile in dense fascicles with 5-9 stamens. They appear before the leaves in the spring. The fruit is a flat, 1-seeded samara. It is suborbicular, 1.5-2 cm and pubescent over the seed.

#### Distribution

For current distribution, please consult the Plant Profile page for this species on the PLANTS web site. This plant is found in moist woods, in southern Maine and southern Quebec to eastern North Dakota, and south to Florida and Texas.

#### **Establishment**

Propagation by seeds: Slippery elm samaras can be gathered when green and ripe from April to June, by sweeping them up from the ground soon after they fall or by knocking the branches with poles and collecting seeds that fall onto tarps. The seeds should then be air-dried for several days, but too much drying can reduce germination. Sow the seeds in flats as soon as they are mature. Sow them with their wings, as de-winging them damages them. There are approximately 90 seeds per gram. Seeds sometimes show dormancy and if so, need stratification. They should be stratified at 41° F for 60-90 days in a moist medium. If storing the seeds before planting, for best results, store them at low moisture content in sealed containers at cool temperatures. Seedlings are susceptible to damping off. In the seedling stage,

Plant Materials <a href="http://plant-materials.nrcs.usda.gov/">http://plant-materials.nrcs.usda.gov/</a> Plant Fact Sheet/Guide Coordination Page <a href="http://plant-materials.nrcs.usda.gov/">http://plant-materials.nrcs.usda.gov/</a> intranet/pfs.html> National Plant Data Center <a href="http://npdc.usda.gov/">http://npdc.usda.gov/</a>

transplant them into larger containers. The seeds can also be directly planted in the garden and the tree grows in a range of soil types, but prefers moist, rich, bottomland soils. This species can become a weed as it tends to inhabit unkempt shrub borders, hedges, fence-rows, and other open ground. It is susceptible to Dutch elm disease, but not to the degree of American elm.

*Propagation by cuttings*: take cuttings in early summer and root with IBA treatment.

## Cultivars, Improved and Selected Materials (and area of origin)

This species is somewhat available through native plant nurseries within its range.

#### References

Cooley, J.H., & J.W. Van Sambeek 1990. *Ulmus* rubra Muhl. Slippery elm. pp. 812-816 IN: Silvics of North America volume 2. hardwoods. USDA, Forest Service, Agriculture Handbook 654. Washington, D.C.

Densmore, F. 1974. *How Indians use wild plants for food, medicine and crafts.* Dover Publications, Inc., New York, New York.

Dirr, M.A. 1997. *Dirr's hardy trees and shrubs: an illustrated encyclopedia*. Timber Press, Portland, Oregon.

Dirr, M.A. 1975. Manual of woody landscape plants: their identification, ornamental characteristics, culture, propagation and uses. Fifth Edition. Stipes Publishing L.L.C. Champaign, Illinois.

Gilmore, M.R. 1977. *Uses of plants by the Indians of the Missouri River region*. University of Nebraska Press, Lincoln, Nebraska.

Gleason, H.A. & A. Cronquist 1991. *Manual of vascular plants of Northeastern United States and adjacent Canada*. Second Edition. The New York Botanical Garden. Bronx, New York.

Herrick, J.W. 1995. *Iroquois medical botany*. Syracuse University Press. Syracuse, New York.

Moerman, D.E. 1998. *Native American ethnobotany*. Timber Press. Portland, Oregon.

Smith, H.H. 1923. *Ethnobotany of the Menomini*. Bulletin of the Public Museum of the City of Milwaukee 4:1(8-175).

Smith, H.H. 1928. *Ethnobotany of the Meskwaki*. Bulletin of the Public Museum of the City of Milwaukee 4:2(175-326).

Young, J.A. & C.G. Young 1992. *Seeds of woody plants in North America*. Discorides Press. Portland, Oregon.

#### **Prepared By & Species Coordinator**

M. Kat Anderson

USDA NRCS National Plant Data Center, c/o Plant Science Horticulture, University of California, Davis, California

Edited 30nov00 jsp; 04jun03 ahv; 30may06jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site<a href="http://plants.usda.gov">http://plants.usda.gov</a> or the Plant Materials Program Web site <a href="http://Plant-Materials.nrcs.usda.gov">http://Plant-Materials.nrcs.usda.gov</a>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Read about <u>Civil Rights at the Natural Resources Convervation</u> Service.