

RED GRAMA

Bouteloua trifida Thurb.

Plant Symbol = BOTR2

Contributed By: USDA NRCS National Plant Data Center



From Hitchcock (1950)
@ plants.usda.gov

Alternate names

Sixweeks grama

Uses

All livestock graze red grama. Forage production is low, but quality is good when plants are green.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

Description

Grass Family (Poaceae). Red grama is a native, warm-season, weak perennial grass. The height is

between 5 to 10 inches. The leaf blade is short, narrow flat or rolling, and the surface slightly rough with scattered long hairs on the margin. The leaf sheath is shorter than the internodes and smooth or slightly rough. The ligule has a ring of short hairs. The stem is smooth. Young plants are erect and the old plants are bent at the base. The nodes are naked and darker than the rest of stem. The seedhead is 3 to 7 spikes per seedhead and about 12 spikelets per spike. The rachis is hairy and the fertile lemma of each spikelet is tipped with 3 short awns.

Distribution: For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

Management

After seedheads appear, nutritive value declines rapidly. This grass helps to protect soil from wind and water erosion if lightly grazed. Growing points and basal leaves are so close to the ground that most livestock cannot harvest more than 50 percent of current year's growth by weight. Deferred grazing during the growing season every 2 to 3 years maintains plant vigor and allows the plants to mature seed for maintaining a stand. This grass is seldom a key management species, except on ranges where it grows in almost pure stands.

Establishment

Growth starts in the spring when daytime temperature is about 70° F and moisture is adequate, and seeds mature 6 weeks after growth starts. Because of this characteristic, red grama is often referred to as 6 weeks' grama. It reproduces from seed, has a shallow root system, and becomes semi-dormant during the summer. It is adapted to shallow, gravelly, stony soils in southwest Texas that have a poor soil-moisture relationship. It invades deep, loamy soils denuded by overgrazing.

Cultivars, Improved and Selected Materials (and area of origin)

Please contact your local NRCS Field Office.

Reference

Leithead, H.L., L.L. Yarlett, & T.N. Shiflett. 1976. 100 native forage grasses in 11 southern states. USDA SCS *Agriculture Handbook No. 389*, Washington, DC.

Prepared By & Species Coordinator:

Percy Magee, USDA NRCS National Plant Data
Center, Baton Rouge, Louisiana

Edited: 13may02 ahv; jul03 ahv; 20sep05 jsp; 070116 jsp

For more information about this and other plants, please contact
your local NRCS field office or Conservation District, and visit the
PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials
Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

*The U.S. Department of Agriculture (USDA) prohibits
discrimination in all its programs and activities on the basis of
race, color, national origin, sex, religion, age, disability, political
beliefs, sexual orientation, and marital or family status. (Not all
prohibited bases apply to all programs.) Persons with disabilities
who require alternative means for communication of program
information (Braille, large print, audiotape, etc.) should contact
USDA's TARGET Center at 202-720-2600 (voice and TDD).*

*To file a complaint of discrimination write USDA, Director, Office
of Civil Rights, Room 326-W, Whitten Building, 14th and
Independence Avenue, SW, Washington, DC 20250-9410 or call
202-720-5964 (voice or TDD). USDA is an equal opportunity
provider and employer.*

*Read about [Civil Rights at the Natural Resources Conservation
Service](#).*