

Table 26. Natural gas home customer-weighted heating degree days

Month/Year/Type of data	New England	Middle Atlantic	East North Central	West North Central	South Atlantic
	CT, ME, MA, NH, RI, VT	NJ, NY, PA	IL, IN, MI, OH, WI	IA, KS, MN, MO, ND, NE, SD	DE, FL, GA, MD, DC, NC, SC, VA, WV
November					
Normal	703	665	757	841	442
2015	584	509	596	653	325
2016	639	576	575	578	384
% Diff (normal to 2016)	-9.1	-13.4	-24.0	-31.3	-13.1
% Diff (2015 to 2016)	9.4	13.2	-3.5	-11.5	18.2
November to November					
Normal	703	665	757	841	442
2015	584	509	596	653	325
2016	639	576	575	578	384
% Diff (normal to 2016)	-9.1	-13.4	-24.0	-31.3	-13.1
% Diff (2015 to 2016)	9.4	13.2	-3.5	-11.5	18.2

Month/Year/Type of data	East South Central	West South Central	Mountain	Pacific ^b	U.S. Average ^b
	AL, KY, MS, TN	AR, LA, OK, TX	AZ, CO, ID, MT, NV, NM, UT, WY	CA, OR, WA	
November					
Normal	455	305	740	366	589
2015	322	229	731	393	486
2016	351	172	550	263	452
% Diff (normal to 2016)	-22.9	-43.6	-25.7	-28.1	-23.3
% Diff (2015 to 2016)	9.0	-24.9	-24.8	-33.1	-7.0
November to November					
Normal	455	305	740	366	589
2015	322	229	731	393	486
2016	351	172	550	263	452
% Diff (normal to 2016)	-22.9	-43.6	-25.7	-28.1	-23.3
% Diff (2015 to 2016)	9.0	-24.9	-24.8	-33.1	-7.0

^a Normal is based on calculations of data from 1971 through 2000.

^b Excludes Alaska and Hawaii.

Note: See Appendix A, Explanatory Note 10, for discussion of Heating Degree-Days computations.

Source: National Oceanic and Atmospheric Administration.