

Table A-1a. Age-adjusted percentages (with standard errors) of selected circulatory diseases among adults aged 18 and over, by selected characteristics: United States, 2014

Selected characteristic	All types of heart disease ¹	Coronary heart disease ¹	Hypertension ¹	Stroke ¹
Total	10.9 (0.21)	5.5 (0.16)	24.5 (0.27)	2.4 (0.10)
Sex				
Male	12.2 (0.31)	7.3 (0.26)	25.3 (0.41)	2.5 (0.15)
Female	9.8 (0.27)	4.0 (0.17)	23.7 (0.35)	2.4 (0.13)
Age (years)				
18-44	4.3 (0.25)	1.1 (0.14)	8.7 (0.29)	0.5 (0.07)
45-64	12.0 (0.40)	6.0 (0.27)	34.0 (0.57)	3.0 (0.21)
65-74	24.6 (0.83)	15.4 (0.70)	54.6 (0.98)	5.4 (0.42)
75 and over	35.0 (1.04)	21.6 (0.95)	60.0 (1.10)	9.7 (0.62)
Race				
One race ²	10.8 (0.21)	5.5 (0.16)	24.5 (0.27)	2.4 (0.10)
White	11.1 (0.24)	5.6 (0.18)	23.5 (0.30)	2.3 (0.11)
Black or African American	10.3 (0.53)	5.5 (0.41)	33.1 (0.73)	4.0 (0.34)
American Indian or Alaska Native	13.7 (2.80)	6.0 (1.44)	26.4 (3.02)	3.0 (0.84)
Asian	6.0 (0.65)	3.3 (0.47)	19.5 (0.96)	1.5 (0.34)
Native Hawaiian or Other Pacific Islander	19.1 (5.51)	6.9 (2.06)	36.4 (6.50)	*8.6 (3.84)
Two or more races ³	16.7 (1.86)	5.8 (1.03)	28.6 (2.32)	3.0 (0.75)
Black or African American, white	12.4 (3.18)	*	26.7 (6.61)	*
American Indian or Alaska Native, white	19.1 (2.98)	6.6 (1.46)	33.8 (3.80)	*3.1 (1.07)
Hispanic or Latino origin ⁴ and race				
Hispanic or Latino	7.8 (0.47)	4.9 (0.40)	23.0 (0.64)	2.4 (0.27)
Mexican or Mexican American	7.6 (0.64)	5.0 (0.57)	22.8 (0.85)	2.6 (0.38)
Not Hispanic or Latino	11.3 (0.24)	5.6 (0.17)	24.9 (0.30)	2.4 (0.11)
White, single race	11.6 (0.28)	5.7 (0.20)	23.8 (0.34)	2.2 (0.12)
Black or African American, single race	10.4 (0.55)	5.6 (0.43)	33.5 (0.75)	4.1 (0.35)
Education ⁵				
Less than a high school diploma	13.7 (0.61)	8.5 (0.48)	31.6 (0.81)	3.8 (0.33)
High school diploma or GED ⁶	12.4 (0.52)	7.4 (0.45)	31.0 (0.64)	3.1 (0.22)
Some college	12.5 (0.43)	6.0 (0.31)	29.6 (0.61)	3.4 (0.25)
Bachelor's degree or higher	10.8 (0.41)	5.0 (0.27)	22.3 (0.52)	1.3 (0.14)
Current employment status ⁷				
Employed	9.3 (0.42)	4.3 (0.33)	21.5 (0.47)	1.3 (0.20)
Full-time	8.7 (0.59)	4.0 (0.49)	22.2 (0.67)	1.5 (0.43)
Part-time	11.1 (0.73)	5.4 (0.55)	20.4 (0.85)	1.4 (0.27)
Not employed but has worked previously	14.5 (0.47)	8.0 (0.40)	30.4 (0.59)	3.9 (0.20)
Not employed and has never worked	10.0 (0.99)	5.6 (0.71)	23.6 (1.20)	2.9 (0.53)

Table A-1a. Age-adjusted percentages (with standard errors) of selected circulatory diseases among adults aged 18 and over, by selected characteristics: United States, 2014

Selected characteristic	All types of heart disease ¹	Coronary heart disease ¹	Hypertension ¹	Stroke ¹
Family income⁸				
Less than \$35,000	13.3 (0.36)	7.5 (0.26)	29.3 (0.50)	3.9 (0.20)
\$35,000 or more	10.2 (0.30)	5.0 (0.22)	22.9 (0.36)	1.7 (0.13)
\$35,000–\$49,999	11.3 (0.71)	6.4 (0.58)	25.0 (0.74)	2.6 (0.31)
\$50,000–\$74,999	10.8 (0.66)	5.6 (0.44)	24.0 (0.70)	1.8 (0.22)
\$75,000–\$99,999	9.6 (0.60)	4.5 (0.46)	24.0 (0.90)	1.3 (0.25)
\$100,000 or more	10.1 (0.52)	4.3 (0.39)	20.9 (0.66)	1.5 (0.26)
Poverty status⁹				
Poor	13.8 (0.56)	8.3 (0.44)	30.3 (0.74)	3.9 (0.31)
Near poor	12.5 (0.51)	7.2 (0.43)	27.9 (0.60)	4.1 (0.29)
Not poor	10.2 (0.27)	4.8 (0.19)	23.0 (0.34)	1.8 (0.12)
Health insurance coverage¹⁰				
Under 65:				
Private	6.0 (0.25)	2.0 (0.13)	16.0 (0.33)	0.9 (0.10)
Medicaid	11.5 (0.69)	6.1 (0.49)	27.4 (0.89)	4.1 (0.38)
Other	12.6 (1.18)	6.8 (0.89)	28.1 (1.53)	3.7 (0.54)
Uninsured	6.1 (0.49)	2.9 (0.39)	15.1 (0.65)	0.9 (0.15)
65 and over:				
Private	30.4 (1.00)	18.0 (0.86)	55.8 (1.13)	6.3 (0.49)
Medicare and Medicaid	34.3 (2.27)	23.7 (2.07)	67.9 (2.34)	14.4 (1.90)
Medicare only	26.0 (1.08)	16.3 (0.92)	55.9 (1.22)	7.8 (0.65)
Other	37.9 (2.59)	26.8 (2.38)	65.0 (2.63)	7.2 (1.46)
Uninsured	*19.0 (8.35)	*16.5 (8.12)	45.0 (8.90)	*
Marital status				
Married	10.5 (0.32)	5.4 (0.25)	24.0 (0.39)	2.1 (0.14)
Widowed	16.5 (3.16)	7.2 (1.31)	38.7 (4.08)	2.6 (0.30)
Divorced or separated	13.8 (0.61)	7.1 (0.41)	30.5 (0.86)	3.7 (0.32)
Never married	10.3 (0.58)	5.1 (0.45)	24.1 (0.74)	2.3 (0.28)
Living with a partner	12.3 (1.24)	7.2 (1.13)	24.6 (1.42)	3.1 (0.78)
Place of residence¹¹				
Large MSA	9.7 (0.28)	5.1 (0.20)	23.2 (0.37)	2.3 (0.14)
Small MSA	12.0 (0.39)	5.6 (0.27)	24.8 (0.47)	2.4 (0.17)
Not in MSA	12.7 (0.59)	6.8 (0.47)	29.1 (0.74)	3.0 (0.22)
Region				
Northeast	10.0 (0.44)	5.1 (0.32)	22.0 (0.60)	1.9 (0.21)
Midwest	11.9 (0.48)	5.8 (0.40)	25.3 (0.56)	2.6 (0.23)
South	11.5 (0.35)	6.2 (0.26)	26.7 (0.45)	2.8 (0.17)
West	9.4 (0.40)	4.5 (0.25)	21.9 (0.56)	2.1 (0.19)

Table A-1a. Age-adjusted percentages (with standard errors) of selected circulatory diseases among adults aged 18 and over, by selected characteristics: United States, 2014

Selected characteristic	All types of heart disease ¹	Coronary heart disease ¹	Hypertension ¹	Stroke ¹
Hispanic or Latino origin ⁴ , race, and sex				
Hispanic or Latino, male	8.6 (0.67)	6.2 (0.61)	22.6 (0.96)	2.8 (0.45)
Hispanic or Latina, female	7.3 (0.63)	3.7 (0.47)	23.2 (0.84)	2.1 (0.34)
Not Hispanic or Latino:				
White, single race, male	13.2 (0.42)	7.8 (0.34)	25.5 (0.52)	2.3 (0.18)
White, single race, female	10.3 (0.36)	3.9 (0.19)	22.1 (0.44)	2.2 (0.15)
Black or African American, single race, male	10.1 (0.82)	5.6 (0.58)	30.0 (1.13)	3.3 (0.45)
Black or African American, single race, female	10.6 (0.75)	5.6 (0.62)	36.5 (1.04)	4.7 (0.51)

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) greater than 30% and less than or equal to 50% and should be used with caution. Data not shown have an RSE greater than 50%.

- Quantity zero.

¹In separate questions, respondents were asked if they had ever been told by a doctor or other health professional that they had: hypertension (or high blood pressure), coronary heart disease, angina (or angina pectoris), heart attack (or myocardial infarction), any other heart condition or disease not already mentioned, or a stroke. A person may be represented in more than one column.

²"All types of heart disease" includes coronary heart disease, angina, heart attack, or any other heart condition or disease.

³"Coronary heart disease" includes coronary heart disease, angina, or heart attack.

Persons had to have been told on two or more different visits that they had hypertension or high blood pressure to be classified as hypertensive.

²Refers to persons who indicated only a single race group, including those of Hispanic or Latino origin.

³Refers to persons who indicated more than one race group, including those of Hispanic or Latino origin. Only two combinations of multiple race groups are shown due to small sample sizes for other combinations.

⁴Refers to persons who are of Hispanic or Latino origin and may be of any race or combination of races. "Not Hispanic or Latino" refers to persons who are not of Hispanic or Latino origin, regardless of race.

⁵Shown only for adults aged 25 and over. Estimates are age adjusted to the projected 2000 U.S. population as the standard population using four age groups: 25-44, 45-64, 65-74, and 75 and over.

⁶GED is General Educational Development high school equivalency diploma.

⁷"Full-time" employment is 35 or more hours per week. "Part-time" employment is 34 or fewer hours per week.

⁸Includes persons who reported a dollar amount or who would not provide a dollar amount but provided an income interval.

⁹"Poor" persons are defined as having income below the poverty threshold. "Near poor" persons have incomes of 100% to less than 200% of the poverty threshold. "Not poor" persons have incomes that are 200% of the poverty threshold or greater.

¹⁰Based on a hierarchy of mutually exclusive categories. Adults with more than one type of health insurance were assigned to the first appropriate category in the hierarchy. "Uninsured" includes adults who had no coverage as well as those who had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care.

¹¹MSA is metropolitan statistical area. Large MSAs have a population size of 1 million or more; small MSAs have a population size of less than 1 million. "Not in MSA" consists of persons not living in a metropolitan statistical area.

NOTES: Estimates are based on household interviews of a sample of the civilian noninstitutionalized population. This table is based on data from the Sample Adult file and was weighted using the Sample Adult weight. Unknowns for the columns were not included in the denominators when calculating percentages. Percentages may not add to totals due to rounding. "Total" includes other races not shown separately and persons with unknown education, family income, poverty status, and health insurance characteristics. Unless otherwise specified, estimates are age-adjusted using the projected 2000 U.S. population as the standard population and using four age groups: 18-44, 45-64, 65-74, and 75 and over. Estimates for age groups are not age-adjusted. For more information on the data source, methods, and definitions used for this table, refer to Technical Notes for Summary Health Statistics Tables: National Health Interview Survey (available from <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>). Suggested citation: Blackwell DL, Lucas JW. Tables of Summary Health Statistics for U.S. Adults: 2014 National Health Interview Survey, 2015. Available from: <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, 2014.

Table A-1b. Frequencies (in thousands) of selected circulatory diseases among adults aged 18 and over, by selected characteristics: United States, 2014

Selected characteristic	All adults aged 18 and over	All types of heart disease ¹	Coronary heart disease ¹	Hypertension ¹	Stroke ¹
Total	239,688	27,613	14,299	63,208	6,286
Sex					
Male	115,541	14,422	8,703	30,707	2,931
Female	124,148	13,191	5,596	32,501	3,355
Age (years)					
18–44	112,149	4,798	1,283	9,721	598
45–64	82,605	9,858	4,964	28,022	2,463
65–74	26,362	6,470	4,059	14,366	1,435
75 and over	18,573	6,487	3,993	11,098	1,790
Race					
One race ²	235,831	27,057	14,104	62,267	6,190
White	190,462	23,176	12,078	49,703	4,825
Black or African American	29,355	2,857	1,513	9,512	1,120
American Indian or Alaska Native	1,948	268	125	504	*59
Asian	13,733	717	386	2,461	163
Native Hawaiian or Other Pacific Islander	333	*	*	*87	*
Two or more races ³	3,858	556	195	940	96
Black or African American, white	760	*70	*	*95	*
American Indian or Alaska Native, white	1,590	333	128	564	*57
Hispanic or Latino origin ⁴ and race					
Hispanic or Latino	36,571	2,265	1,301	6,725	649
Mexican or Mexican American	22,326	1,237	765	3,843	406
Not Hispanic or Latino	203,117	25,348	12,998	56,483	5,637
White, single race	157,364	21,108	10,867	43,648	4,202
Black or African American, single race	27,875	2,794	1,480	9,247	1,114
Education ⁵					
Less than a high school diploma	27,612	4,578	2,951	10,248	1,297
High school diploma or GED ⁶	52,697	7,469	4,469	18,397	1,941
Some college	59,919	7,539	3,576	18,313	2,068
Bachelor's degree or higher	67,940	6,856	3,117	14,951	822
Current employment status ⁷					
Employed	146,624	9,809	3,812	26,734	1,329
Full-time	117,112	7,147	2,627	21,540	985
Part-time	27,205	2,505	1,110	4,838	319
Not employed but has worked previously	79,316	16,514	9,801	33,578	4,639
Not employed and has never worked	13,623	1,289	687	2,870	313

Table A-1b. Frequencies (in thousands) of selected circulatory diseases among adults aged 18 and over, by selected characteristics: United States, 2014

Selected characteristic	All adults aged 18 and over	All types of heart disease ¹	Coronary heart disease ¹	Hypertension ¹	Stroke ¹
Family income ⁸					
Less than \$35,000	69,793	10,409	5,995	22,389	3,078
\$35,000 or more	144,503	14,253	6,853	33,702	2,379
\$35,000–\$49,999	28,044	3,301	1,875	7,402	751
\$50,000–\$74,999	35,733	3,914	2,050	9,115	671
\$75,000–\$99,999	27,053	2,504	1,137	6,563	363
\$100,000 or more	53,673	4,534	1,791	10,622	594
Poverty status ⁹					
Poor	31,383	3,909	2,251	8,510	1,058
Near poor	42,675	5,753	3,345	12,548	1,901
Not poor	152,199	16,432	7,912	38,413	2,926
Health insurance coverage ¹⁰					
Under 65:					
Private	129,666	8,518	3,056	23,336	1,319
Medicaid	23,239	2,618	1,373	6,216	914
Other	9,095	1,583	922	3,615	504
Uninsured	31,587	1,873	875	4,468	268
65 and over:					
Private	22,525	6,654	3,924	12,452	1,349
Medicare and Medicaid	2,951	1,009	694	1,999	421
Medicare only	15,536	3,965	2,491	8,616	1,173
Other	3,513	1,267	892	2,245	235
Uninsured	313	*56	*	137	*
Marital status					
Married	126,926	14,790	7,881	35,311	3,081
Widowed	14,312	4,026	2,385	8,398	1,136
Divorced or separated	26,802	4,310	2,311	9,692	1,193
Never married	53,788	3,059	1,069	6,714	608
Living with a partner	17,497	1,396	643	2,972	258
Place of residence ¹¹					
Large MSA	130,402	12,744	6,652	31,089	2,966
Small MSA	73,885	9,438	4,558	19,904	1,989
Not in MSA	35,402	5,430	3,089	12,215	1,331
Region					
Northeast	41,490	4,744	2,473	10,558	927
Midwest	55,095	6,839	3,392	14,896	1,521
South	89,270	10,975	6,043	25,742	2,688
West	53,834	5,055	2,391	12,012	1,150

Table A-1b. Frequencies (in thousands) of selected circulatory diseases among adults aged 18 and over, by selected characteristics: United States, 2014

Selected characteristic	All adults aged 18 and over	All types of heart disease ¹	Coronary heart disease ¹	Hypertension ¹	Stroke ¹
Hispanic or Latino origin ⁴ , race, and sex					
Hispanic or Latino, male	18,309	1,141	773	3,211	337
Hispanic or Latina, female	18,262	1,125	529	3,514	312
Not Hispanic or Latino:					
White, single race, male	76,277	11,361	6,853	22,108	2,045
White, single race, female	81,087	9,747	4,014	21,540	2,157
Black or African American, single race, male	12,626	1,218	643	3,629	400
Black or African American, single race, female	15,249	1,576	837	5,618	715

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) greater than 30% and less than or equal to 50% and should be used with caution. Data not shown have an RSE greater than 50%.

- Quantity zero.

¹In separate questions, respondents were asked if they had ever been told by a doctor or other health professional that they had: hypertension (or high blood pressure), coronary heart disease, angina (or angina pectoris), heart attack (or myocardial infarction), any other heart condition or disease not already mentioned, or a stroke. A person may be represented in more than one column.

"All types of heart disease" includes coronary heart disease, angina, heart attack, or any other heart condition or disease.

"Coronary heart disease" includes coronary heart disease, angina, or heart attack.

Persons had to have been told on two or more different visits that they had hypertension or high blood pressure to be classified as hypertensive.

²Refers to persons who indicated only a single race group, including those of Hispanic or Latino origin.

³Refers to persons who indicated more than one race group, including those of Hispanic or Latino origin. Only two combinations of multiple race groups are shown due to small sample sizes for other combinations.

⁴Refers to persons who are of Hispanic or Latino origin and may be of any race or combination of races. "Not Hispanic or Latino" refers to persons who are not of Hispanic or Latino origin, regardless of race.

⁵Shown only for adults aged 25 and over.

⁶GED is General Educational Development high school equivalency diploma.

⁷"Full-time" employment is 35 or more hours per week. "Part-time" employment is 34 or fewer hours per week.

⁸Includes persons in families that reported either a dollar amount or would not provide a dollar amount but provided an income interval.

⁹"Poor" persons live in families defined as below the poverty threshold. "Near poor" persons live in families with incomes of 100% to less than 200% of the poverty threshold. "Not poor" persons live in families with incomes that are 200% of the poverty threshold or greater.

¹⁰Based on a hierarchy of mutually exclusive categories. Adults with more than one type of health insurance were assigned to the first appropriate category in the hierarchy. "Uninsured" includes adults who had no coverage as well as those who had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care.

¹¹MSA is metropolitan statistical area. Large MSAs have a population size of 1 million or more; small MSAs have a population size of less than 1 million. "Not in MSA" consists of persons not living in a metropolitan statistical area.

NOTES: Estimates are based on household interviews of a sample of the civilian noninstitutionalized population. This table is based on data from the Sample Adult file and was weighted using the Sample Adult weight. Unknowns for the columns were not included in the frequencies, but they are included in the "All adults aged 18 and over" column. "Total" includes other races not shown separately and persons with unknown education, family income, poverty status, and health insurance characteristics. For more information on the data source, methods, and definitions used for this table, refer to Technical Notes for Summary Health Statistics Tables: National Health Interview Survey (available from <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>). Suggested citation: Blackwell DL, Lucas JW. Tables of Summary Health Statistics for U.S. Adults: 2014 National Health Interview Survey. 2015. Available from: <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, 2014.

Table A-1c. Crude percentages (with standard errors) of selected circulatory diseases among adults aged 18 and over, by selected characteristics: United States, 2014

Selected characteristic	All types of heart disease ¹	Coronary heart disease ¹	Hypertension ¹	Stroke ¹
Total	11.5 (0.23)	6.0 (0.17)	26.4 (0.33)	2.6 (0.11)
Sex				
Male	12.5 (0.35)	7.5 (0.28)	26.6 (0.48)	2.5 (0.16)
Female	10.6 (0.29)	4.5 (0.19)	26.2 (0.42)	2.7 (0.15)
Age (years)				
18–44	4.3 (0.25)	1.1 (0.14)	8.7 (0.29)	0.5 (0.07)
45–64	12.0 (0.40)	6.0 (0.27)	34.0 (0.57)	3.0 (0.21)
65–74	24.6 (0.83)	15.4 (0.70)	54.6 (0.98)	5.4 (0.42)
75 and over	35.0 (1.04)	21.6 (0.95)	60.0 (1.10)	9.7 (0.62)
Race				
One race ²	11.5 (0.23)	6.0 (0.17)	26.4 (0.33)	2.6 (0.11)
White	12.2 (0.27)	6.4 (0.20)	26.1 (0.37)	2.5 (0.12)
Black or African American	9.7 (0.53)	5.2 (0.40)	32.5 (0.84)	3.8 (0.34)
American Indian or Alaska Native	13.8 (2.78)	6.4 (1.57)	25.9 (3.61)	*3.1 (0.95)
Asian	5.2 (0.60)	2.8 (0.41)	17.9 (1.06)	1.2 (0.27)
Native Hawaiian or Other Pacific Islander	*	*	*26.1 (8.01)	*
Two or more races ³	14.4 (1.74)	5.1 (0.93)	24.7 (2.46)	2.5 (0.63)
Black or African American, white	*9.2 (3.08)	*	*12.5 (3.75)	*
American Indian or Alaska Native, white	20.9 (3.29)	8.1 (1.84)	35.8 (4.38)	*3.6 (1.29)
Hispanic or Latino origin ⁴ and race				
Hispanic or Latino	6.2 (0.40)	3.6 (0.32)	18.4 (0.63)	1.8 (0.20)
Mexican or Mexican American	5.5 (0.51)	3.4 (0.43)	17.2 (0.79)	1.8 (0.26)
Not Hispanic or Latino	12.5 (0.26)	6.4 (0.19)	27.9 (0.36)	2.8 (0.12)
White, single race	13.4 (0.32)	6.9 (0.23)	27.8 (0.43)	2.7 (0.15)
Black or African American, single race	10.0 (0.55)	5.3 (0.42)	33.2 (0.87)	4.0 (0.35)
Education ⁵				
Less than a high school diploma	16.6 (0.72)	10.7 (0.60)	37.2 (0.97)	4.7 (0.39)
High school diploma or GED ⁶	14.2 (0.55)	8.5 (0.46)	35.0 (0.73)	3.7 (0.27)
Some college	12.6 (0.46)	6.0 (0.32)	30.6 (0.67)	3.5 (0.26)
Bachelor's degree or higher	10.1 (0.41)	4.6 (0.26)	22.0 (0.56)	1.2 (0.12)
Current employment status ⁷				
Employed	6.7 (0.23)	2.6 (0.14)	18.3 (0.34)	0.9 (0.08)
Full-time	6.1 (0.25)	2.2 (0.15)	18.4 (0.40)	0.8 (0.09)
Part-time	9.2 (0.64)	4.1 (0.42)	17.8 (0.79)	1.2 (0.21)
Not employed but has worked previously	20.9 (0.48)	12.4 (0.41)	42.4 (0.64)	5.9 (0.26)
Not employed and has never worked	9.5 (0.89)	5.0 (0.59)	21.1 (1.14)	2.3 (0.38)

Table A-1c. Crude percentages (with standard errors) of selected circulatory diseases among adults aged 18 and over, by selected characteristics: United States, 2014

Selected characteristic	All types of heart disease ¹	Coronary heart disease ¹	Hypertension ¹	Stroke ¹
Family income⁸				
Less than \$35,000	14.9 (0.40)	8.6 (0.31)	32.1 (0.61)	4.4 (0.23)
\$35,000 or more	9.9 (0.30)	4.7 (0.21)	23.3 (0.41)	1.6 (0.13)
\$35,000–\$49,999	11.8 (0.72)	6.7 (0.58)	26.4 (0.90)	2.7 (0.33)
\$50,000–\$74,999	11.0 (0.64)	5.7 (0.46)	25.5 (0.81)	1.9 (0.23)
\$75,000–\$99,999	9.3 (0.61)	4.2 (0.44)	24.3 (1.01)	1.3 (0.25)
\$100,000 or more	8.5 (0.46)	3.3 (0.28)	19.8 (0.68)	1.1 (0.18)
Poverty status⁹				
Poor	12.5 (0.55)	7.2 (0.42)	27.2 (0.88)	3.4 (0.28)
Near poor	13.5 (0.54)	7.8 (0.44)	29.4 (0.71)	4.5 (0.32)
Not poor	10.8 (0.29)	5.2 (0.21)	25.3 (0.41)	1.9 (0.13)
Health insurance coverage¹⁰				
Under 65:				
Private	6.6 (0.25)	2.4 (0.15)	18.0 (0.38)	1.0 (0.11)
Medicaid	11.3 (0.71)	5.9 (0.50)	26.8 (1.00)	3.9 (0.38)
Other	17.4 (1.34)	10.1 (1.05)	39.9 (1.99)	5.5 (0.77)
Uninsured	5.9 (0.49)	2.8 (0.38)	14.2 (0.65)	0.8 (0.15)
65 and over:				
Private	29.6 (1.00)	17.5 (0.84)	55.4 (1.14)	6.0 (0.47)
Medicare and Medicaid	34.2 (2.27)	23.6 (2.06)	67.8 (2.32)	14.3 (1.88)
Medicare only	25.6 (1.05)	16.1 (0.89)	55.7 (1.22)	7.6 (0.63)
Other	36.2 (2.53)	25.6 (2.33)	64.0 (2.67)	6.7 (1.32)
Uninsured	*17.7 (6.86)	*14.4 (6.43)	43.9 (7.97)	*
Marital status				
Married	11.7 (0.33)	6.2 (0.26)	27.9 (0.46)	2.4 (0.16)
Widowed	28.3 (1.06)	16.8 (0.89)	59.0 (1.12)	8.0 (0.64)
Divorced or separated	16.1 (0.62)	8.7 (0.47)	36.3 (0.86)	4.5 (0.34)
Never married	5.7 (0.33)	2.0 (0.17)	12.5 (0.45)	1.1 (0.14)
Living with a partner	8.0 (0.78)	3.7 (0.57)	17.0 (1.06)	1.5 (0.29)
Place of residence¹¹				
Large MSA	9.8 (0.30)	5.1 (0.21)	23.9 (0.44)	2.3 (0.14)
Small MSA	12.8 (0.41)	6.2 (0.30)	27.0 (0.55)	2.7 (0.20)
Not in MSA	15.3 (0.68)	8.7 (0.55)	34.5 (0.94)	3.8 (0.27)
Region				
Northeast	11.4 (0.57)	6.0 (0.39)	25.5 (0.74)	2.2 (0.23)
Midwest	12.4 (0.45)	6.2 (0.40)	27.1 (0.68)	2.8 (0.26)
South	12.3 (0.42)	6.8 (0.31)	28.9 (0.56)	3.0 (0.18)
West	9.4 (0.43)	4.4 (0.27)	22.3 (0.66)	2.1 (0.19)

Table A-1c. Crude percentages (with standard errors) of selected circulatory diseases among adults aged 18 and over, by selected characteristics: United States, 2014

Selected characteristic	All types of heart disease ¹	Coronary heart disease ¹	Hypertension ¹	Stroke ¹
Hispanic or Latino origin ⁴ , race, and sex				
Hispanic or Latino, male	6.2 (0.53)	4.2 (0.42)	17.6 (0.91)	1.8 (0.30)
Hispanic or Latina, female	6.2 (0.58)	2.9 (0.41)	19.3 (0.85)	1.7 (0.27)
Not Hispanic or Latino:				
White, single race, male	14.9 (0.48)	9.0 (0.38)	29.0 (0.63)	2.7 (0.21)
White, single race, female	12.0 (0.39)	5.0 (0.25)	26.6 (0.53)	2.7 (0.18)
Black or African American, single race, male	9.7 (0.83)	5.1 (0.54)	28.8 (1.28)	3.2 (0.43)
Black or African American, single race, female	10.3 (0.75)	5.5 (0.61)	36.9 (1.18)	4.7 (0.53)

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) greater than 30% and less than or equal to 50% and should be used with caution. Data not shown have an RSE greater than 50%.

- Quantity zero.

¹In separate questions, respondents were asked if they had ever been told by a doctor or other health professional that they had: hypertension (or high blood pressure), coronary heart disease, angina (or angina pectoris), heart attack (or myocardial infarction), any other heart condition or disease not already mentioned, or a stroke. A person may be represented in more than one column.

"All types of heart disease" includes coronary heart disease, angina, heart attack, or any other heart condition or disease.

"Coronary heart disease" includes coronary heart disease, angina, or heart attack.

Persons had to have been told on two or more different visits that they had hypertension or high blood pressure to be classified as hypertensive.

²Refers to persons who indicated only a single race group, including those of Hispanic or Latino origin.

³Refers to persons who indicated more than one race group, including those of Hispanic or Latino origin. Only two combinations of multiple race groups are shown due to small sample sizes for other combinations.

⁴Refers to persons who are of Hispanic or Latino origin and may be of any race or combination of races. "Not Hispanic or Latino" refers to persons who are not of Hispanic or Latino origin, regardless of race.

⁵Shown only for adults aged 25 and over.

⁶GED is General Educational Development high school equivalency diploma.

⁷"Full-time" employment is 35 or more hours per week. "Part-time" employment is 34 or fewer hours per week.

⁸Includes persons in families that reported either a dollar amount or would not provide a dollar amount but provided an income interval.

⁹"Poor" persons live in families defined as below the poverty threshold. "Near poor" persons live in families with incomes of 100% to less than 200% of the poverty threshold. "Not poor" persons live in families with incomes that are 200% of the poverty threshold or greater.

¹⁰Based on a hierarchy of mutually exclusive categories. Adults with more than one type of health insurance were assigned to the first appropriate category in the hierarchy. "Uninsured" includes adults who had no coverage as well as those who had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care.

¹¹MSA is metropolitan statistical area. Large MSAs have a population size of 1 million or more; small MSAs have a population size of less than 1 million. "Not in MSA" consists of persons not living in a metropolitan statistical area.

NOTES: Estimates are based on household interviews of a sample of the civilian noninstitutionalized population. This table is based on data from the Sample Adult file and was weighted using the Sample Adult weight. Unknowns for the columns were not included in the denominators when calculating percentages. Percentages may not add to totals due to rounding.

"Total" includes other races not shown separately and persons with unknown education, family income, poverty status, and health insurance characteristics. For more information on the data source, methods, and definitions used for this table, refer to Technical Notes for Summary Health Statistics Tables: National Health Interview Survey (available from <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>). Suggested citation: Blackwell DL, Lucas JW. Tables of Summary Health Statistics for U.S. Adults: 2014 National Health Interview Survey. 2015. Available from: <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, 2014.