

Acknowledgments

We thank Jesse Gallagher, Melinda Vonkungthong, Anne Hurley-Bacon, Jasmina Luczo, James Doster, and Charles Foley for technical assistance with this work.

Severe acute respiratory syndrome coronavirus 2, isolate USA-WA1/2020, NR-52281 was deposited by the Centers for Disease Control and Prevention and obtained through BEI Resources, NIAID, NIH. Middle East respiratory syndrome coronavirus, Florida/USA-2_Saudi Arabia_2014, NR-50415 was obtained through BEI Resources, NIAID, NIH. Vero African green monkey kidney cells (ATCC CCL-81), FR-243, were obtained through the International Reagent Resource, Influenza Division, WHO Collaborating Center for Surveillance, Epidemiology and Control of Influenza, Centers for Disease Control and Prevention, Atlanta, GA, USA.

This work was supported by USDA-Agricultural Research Service (project no. 6040-32000-066-00-D).

About the Author

Dr. Suarez is the research leader for the Exotic and Emerging Avian Viral Disease Research Unit of the Agricultural Research Service, USDA. His primary research interests are in the understanding and control of avian influenza and Newcastle disease viruses in poultry and other emerging viral diseases that threaten the poultry industry.

References

1. Corman VM, Muth D, Niemeyer D, Drosten C. Hosts and sources of endemic human coronaviruses. *Adv Virus Res.* 2018;100:163–88. <https://doi.org/10.1016/bs.aivir.2018.01.001>
2. Chan JF, Kok KH, Zhu Z, Chu H, To KK, Yuan S, et al. Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect.* 2020;9:221–36. <https://doi.org/10.1080/22221751.2020.1719902>
3. Schlottau K, Rissmann M, Graaf A, Schön J, Sehl J, Wylezich C, et al. SARS-CoV-2 in fruit bats, ferrets, pigs, and chickens: an experimental transmission study. *Lancet Microbe.* 2020;1:e218–e225. [https://doi.org/10.1016/S2666-5247\(20\)30089-6](https://doi.org/10.1016/S2666-5247(20)30089-6)
4. Shi J, Wen Z, Zhong G, Yang H, Wang C, Huang B, et al. Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2. *Science.* 2020;368:1016–20. <https://doi.org/10.1126/science.abb7015>
5. The Lancet. MERS-CoV: a global challenge. *Lancet.* 2013; 381:1960. [https://doi.org/10.1016/S0140-6736\(13\)61184-8](https://doi.org/10.1016/S0140-6736(13)61184-8)
6. Hui DS, Azhar EI, Memish ZA, Zumla A. Human coronavirus infections—severe acute respiratory syndrome (SARS), Middle East respiratory syndrome (MERS), and SARS-CoV-2. Reference Module in Biomedical Sciences. 2020 [cited 2020 Sep 23]. <https://doi.org/10.1016/B978-0-12-801238-3.11634-4>
7. Ferguson NM, Van Kerkhove MD. Identification of MERS-CoV in dromedary camels. *Lancet Infect Dis.* 2014;14:93–4. [https://doi.org/10.1016/S1473-3099\(13\)70691-1](https://doi.org/10.1016/S1473-3099(13)70691-1)
8. Kandeil A, Gomaa M, Shehata M, El-Taweel A, Kayed AE, Abiadh A, et al. Middle East respiratory syndrome coronavirus infection in non-camelid domestic mammals. *Emerg Microbes Infect.* 2019;8:103–8. <https://doi.org/10.1080/22221751.2018.1560235>
9. Hemida MG, Perera RA, Wang P, Alhammadi MA, Siu LY, Li M, et al. Middle East respiratory syndrome (MERS) coronavirus seroprevalence in domestic livestock in Saudi Arabia, 2010 to 2013. *Euro Surveill.* 2013;18:20659. <https://doi.org/10.2807/1560-7917.ES2013.18.50.20659>
10. Barr IG, Rynehart C, Whitney P, Druce J. SARS-CoV-2 does not replicate in embryonated hen's eggs or in MDCK cell lines. *Euro Surveill.* 2020;25. <https://doi.org/10.2807/1560-7917.ES.2020.25.25.2001122>

Address for correspondence: Erica Spackman, US National Poultry Research Center, USDA Agricultural Research Service, 934 Station Rd, Athens, GA 30605, USA; email: erica.spackman@usda.gov

Serologic Responses in Healthy Adult with SARS-CoV-2 Reinfection, Hong Kong, August 2020

Paul K.S. Chan, Grace Lui, Asmaa Hachim, Ronald L.W. Ko, Siaw S. Boon, Timothy Li, Niloufar Kavian, Fion Luk, Zigui Chen, Emily M. Yau, Kin H. Chan, Chi-hang Tsang, Samuel M.S. Cheng, Daniel K.W. Chu, Ranawaka A.P.M. Perera, Wendy C.S. Ho, Apple C.M. Yeung, Chit Chow, Leo L.M. Poon, Sophie A. Valkenburg, David S.C. Hui, Malik Peiris

Author affiliations: The Chinese University of Hong Kong, Hong Kong, China (P.K.S. Chan, G. Lui, S.S. Boon, T. Li, F. Luk, Z. Chen, W.C.S. Ho, A.C.M. Yeung, C. Chow, D.S.C. Hui); The University of Hong Kong, Hong Kong (A. Hachim, R.L.W. Ko, N. Kavian, E.M. Yau, K.H. Chan, C. Tsang, S.M.S. Cheng, D.K.W. Chu, R.A.P.M. Perera, L.L.M. Poon, S.A. Valkenburg, M. Peiris)

DOI: <https://doi.org/10.3201/eid2612.203833>

In March 2020, mild signs and symptoms of coronavirus disease developed in a healthy 33-year-old man in Hong Kong. His first infection did not produce virus neutralizing antibodies. In August, he had asymptomatic reinfection, suggesting that persons without a robust neutralizing antibody response might be at risk for reinfection.

response suggests that antibody against SARS-CoV-2 developed on reinfection.

Levels of adaptive cytokine interleukin-2 were elevated on days 10 and 43 (Appendix Figure 3, panels A, B). Reinfection coincided with a stronger interleukin-21 memory type response on day 148 than on days 10 and 43.

Previous studies show that most patients with mild, severe, or asymptomatic SARS-CoV-2 infection produce neutralizing antibodies and antibodies against spike RBD and N proteins (3,4). This case was unusual because the patient had low or undetectable levels of neutralizing and binding antibodies against multiple viral proteins during his primary infection and acute stage of asymptomatic reinfection. He was not immunodeficient because he had IgG against measles and varicella zoster viruses and no history of recurrent infections. The virus from the first infection had a truncation in the 58AA open reading frame 8 gene, which mediates immune evasion through downregulation of major histocompatibility complex and interferon responses (Y. Zhang et al., unpub. data, <https://www.biorxiv.org/content/10.1101/2020.05.24.111823v1>) (8). However, it is unclear if this mutation contributed to the patient's lack of antibody production.

Reasons for this patient's unusual response need to be further investigated. He recovered from his primary infection within 3 weeks, and his secondary infection was asymptomatic. These findings indicate that, in the absence of primary neutralizing antibodies, T cells and mucosal immunity might have played a critical role in resolving the infection. Given the unusual antibody response in this patient to his first infection, researchers must be cautious about generalizing more widely from this patient's experience.

Acknowledgments

We thank Rity Wong, Vickie Li, Miu Ling Chin, Barry Wong, and Kitty Fung for their assistance in this study.

The study was supported by the Health and Medical Research Fund—Commissioned Research on the Novel Coronavirus Disease (COVID-19) (reference nos. COVID190107, COVID190126, COVID190205, and COVID190115) from the Food and Health Bureau, Hong Kong Special Administrative Region Government; and the US National Institutes of Health (contract no. HHSN272201400006C).

About the Author

Prof. Chan is a clinical virologist at the Chinese University of Hong Kong, Hong Kong. His research interests include diagnostics, epidemiology, and pathogenesis.

References

1. Kiyuka PK, Agoti CN, Munywoki PK, Njeru R, Bett A, Otiemo JR, et al. Human coronavirus NL63 molecular epidemiology and evolutionary patterns in rural coastal Kenya. *J Infect Dis*. 2018;217:1728–39. <https://doi.org/10.1093/infdis/jiy098>
2. Callow KA. Effect of specific humoral immunity and some non-specific factors on resistance of volunteers to respiratory coronavirus infection. *J Hyg (Lond)*. 1985;95:173–89. <https://doi.org/10.1017/S0022172400062410>
3. Perera RA, Mok CK, Tsang OT, Lv H, Ko RL, Wu NC, et al. Serological assays for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), March 2020. *Euro Surveill*. 2020;25:2000421. <https://doi.org/10.2807/1560-7917.ES.2020.25.16.2000421>
4. Hachim A, Kaviani N, Cohen CA, Chin AWH, Chu DKW, Mok CKP, et al. ORF8 and ORF3b antibodies are accurate serological markers of early and late SARS-CoV-2 infection. [Erratum in: *Nat Immunol*. 2020 Aug 27; Epub ahead of print]. *Nat Immunol*. 2020 Aug 17 [Epub ahead of print].
5. Addetia A, Crawford KHD, Dingens A, Zhu H, Roychoudhury P, Huang ML, et al. Neutralizing antibodies correlate with protection from SARS-CoV-2 in humans during a fishery vessel outbreak with high attack rate. *J Clin Microbiol*. 2020;Aug 21;JCM.02107-20. <https://doi.org/10.1128/JCM.02107-20>
6. To KK, Hung IF, Ip JD, Chu AW, Chan WM, Tam AR, et al. COVID-19 re-infection by a phylogenetically distinct SARS-coronavirus-2 strain confirmed by whole genome sequencing. *Clin Infect Dis*. 2020 Aug 25 [Epub ahead of print]. <https://doi.org/10.1093/cid/ciaa1275>
7. Tan CW, Chia WN, Qin X, Liu P, Chen MI, Tiu C, et al. A SARS-CoV-2 surrogate virus neutralization test based on antibody-mediated blockage of ACE2-spike protein-protein interaction. *Nat Biotechnol*. 2020;38:1073–8. <https://doi.org/10.1038/s41587-020-0631-z>
8. Li JY, Liao CH, Wang Q, Tan YJ, Luo R, Qiu Y, et al. The ORF6, ORF8 and nucleocapsid proteins of SARS-CoV-2 inhibit type I interferon signaling pathway. *Virus Res*. 2020;286:198074. <https://doi.org/10.1016/j.virusres.2020.198074>

Address for correspondence: Malik Peiris, School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, No. 7 Sassoon Rd., Pokfulam, Hong Kong, China; email: malik@hku.hk; Grace Lui, Department of Medicine and Therapeutics, Faculty of Medicine, The Chinese University of Hong Kong, Prince of Wales Hospital, 30-32 Ngan Shing St., Shatin, New Territories, Hong Kong, China; email: gracelui@cuhk.edu.hk