

**United States Department of Agriculture
Center for Veterinary Biologics
Testing Protocol**

SAM 624

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

Date: **August 5, 2008** Approved, pending Standard Requirement

Number: SAM 624.03

Supersedes: SAM 624.02, June 17, 2006

Standard Requirement:

Contact: Mary C. Rasmusson, (515) 663-7831

Approvals: /s/Geetha B. Srinivas Date: 16Sep08
Geetha B. Srinivas, Section Leader
Bacteriology

/s/Byron E. Rippke Date: 29Sep08
Byron E. Rippke, Director
Policy, Evaluation, and Licensing
Center for Veterinary Biologics

/s/Rebecca L.W. Hyde Date: 07Oct08
Rebecca L.W. Hyde, Section Leader
Quality Management
Center for Veterinary Biologics

United States Department of Agriculture
Animal and Plant Health Inspection Service
P. O. Box 844
Ames, IA 50010

Mention of trademark or proprietary product does not constitute a guarantee or warranty of the product by USDA and does not imply its approval to the exclusion of other products that may be suitable.

UNCONTROLLED COPY

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

Table of Contents

- 1. Introduction**
- 2. Materials**
 - 2.1 Equipment/instrumentation**
 - 2.2 Reagents/supplies**
- 3. Preparation for the Test**
 - 3.1 Personnel qualifications/training**
 - 3.2 Preparation of equipment/instrumentation**
 - 3.3 Preparation of reagents/control procedures**
 - 3.4 Preparation of the sample**
- 4. Performance of the Test**
- 5. Interpretation of the Test Results**
 - 5.1 Relative potency calculation method**
 - 5.2 Requirements for a valid assay**
 - 5.3 Requirements for a satisfactory test bacterin**
- 6. Report of Test Results**
- 7. Summary of Revisions**

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

1. Introduction

This is a Supplemental Assay Method (SAM) using a sandwich enzyme-linked immunosorbent assay (ELISA) system to measure the relative potency of bacterins containing *Leptospira interrogans* serovar *pomona* compared to a suitably qualified, nonexpired reference.

2. Materials

2.1 Equipment/instrumentation

Equivalent equipment or instrumentation may be substituted for any brand name listed below.

- 2.1.1 Micropipettor to cover the range of 5- μ L to 1000- μ L
- 2.1.2 8- or 12-channel micropipettor, to cover the range of 50- μ L to 200- μ L
- 2.1.3 Orbital shaker
- 2.1.4 Automatic microplate washer (optional)
- 2.1.5 Microplate reader with dual wavelengths (405 nm and 490 nm)
- 2.1.6 Balance, validated from 150 mg to 15 g
- 2.1.7 Relative Potency Calculation Software (United States Department of Agriculture [USDA], Veterinary Services, Center for Veterinary Biologics [CVB]), current version of **SAM 318**

2.2 Reagents/supplies

Equivalent reagents or supplies may be substituted for any brand name listed below.

- 2.2.1 96-well flat-bottom microtitration plates (Immulon 2, Dynatech Laboratories, Inc.)
- 2.2.2 96-well non-binding microtitration plates suitable for making serial dilutions (transfer plate)
- 2.2.3 Plate sealers
- 2.2.4 Carbonate coating buffer

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

- 2.2.5** Phosphate buffer for elution (optional)
- 2.2.6** Sodium citrate for elution (optional)
- 2.2.7** Sodium deoxycholate (desoxycholate) elution buffer (optional)
- 2.2.8** Phosphate-buffered saline with 0.05% Tween 20 (PBS-Tween 20)
- 2.2.9** Antibody diluent with Normal rabbit serum (negative for *Leptospira* spp.)
- 2.2.10** ABTS (2,2'-azino-di-3-ethylbenziazoline sulfonate) substrate, 1- or 2-component (Kirkegaard and Perry Laboratories, Inc.)
- 2.2.11** *L. pomona*-specific (clone 2D7F10) monoclonal antibody (MAb).
L. pomona 2D7F10 MAb is available from the CVB. Refer to the current reagent data sheet for additional information.
- 2.2.12** Polyclonal *L. pomona* antiserum (PAb), rabbit origin. *L. pomona* PAb is available from the CVB. Refer to the current reagent data sheet for additional information.
- 2.2.13** Goat anti-mouse IgA (H+L) horseradish peroxidase-labeled antibody (Kirkegaard and Perry Laboratories, Inc.)
- 2.2.14** Test bacterin(s) containing *L. pomona*
- 2.2.15** Reference bacterin (nonexpired) containing *L. pomona* (must be approved by the Animal and Plant Health Inspection Service [APHIS])

3. Preparation for the Test

3.1 Personnel qualifications/training

Technical personnel need a working knowledge of the use of general laboratory chemicals, equipment, and glassware; automated microplate washer and microplate reader; and data analysis software. They need specific training in the performance of this assay.

3.2 Preparation of equipment/instrumentation

Operate and maintain all equipment according to manufacturers' recommendations and applicable in-house standard operating procedures.

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

3.3 Preparation of reagents/control procedures

3.3.1 Carbonate coating buffer--National Veterinary Services Laboratories (NVSL) Media #20034

Na ₂ CO ₃	0.159 g
NaHCO ₃	0.293 g
Deionized water	q.s. to 100 mL

Adjust pH to 9.6 ± 0.1. Store at 2°- 7°C for up to 1 week.

3.3.2 Phosphate buffered saline (PBS)--NVSL Media #10559

NaCl	8.00 g
KCl	0.20 g
Na ₂ HPO ₄	1.15 g
KH ₂ PO ₄	0.20 g
Deionized water	q.s. to 1 L

Adjust pH to 7.2 ± 0.1. Store at 20°- 25°C for no longer than 6 months.

3.3.3 Polyvinyl alcohol, 1%, in PBS

PBS (see Section 3.3.2)	70 mL
Polyvinyl alcohol, 88% hydrolyzed, MW 13,000-23,000 (Aldrich Chemical, Cat. No. 36,317-0, or equivalent)	0.7 g

Stir to dissolve. Store at 20°- 25°C for no longer than 1 week.

3.3.4 Phosphate buffered saline with 0.05% Tween 20 (PBS-Tween 20)--NVSL Media #30179

PBS (see Section 3.3.2)	1000 mL
Tween 20	0.50 mL

Store at 20°- 25°C for no longer than 6 months.

3.3.5 Antibody diluent with normal rabbit serum (negative for *Leptospira* spp.)

Polyvinyl alcohol, 1%, in PBS (see Section 3.3.3)	43.6 mL
Normal rabbit serum	400 µL

Mix immediately prior to use. Store at 2°- 7°C for up to 1 week.

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

3.3.6 Phosphate buffer for antigen elution

KH ₂ PO ₄ (Mallinckrodt, Inc., 7100, or equivalent)	8.2 g
Deionized water	94 mL

Adjust pH to 9.3 ± 0.1, or other appropriate pH as optimized for use with a specific bacterin. Store at 20°- 25°C for no longer than 1 month.

3.3.7 Sodium deoxycholate (desoxycholate) elution buffer

Sodium deoxycholate (Sigma Chemical D6750, or equivalent)	0.50 g
PBS (see Section 3.3.2)	100 mL

Store at 2°- 7°C for up to 30 days. Warm to 20°- 25°C prior to use (buffer gels at 2°- 7°C).

3.3.8 MAb specific for *L. pomona*, clone 2D7F10

Obtain MAb 2D7F10 from the CVB. Store long-term at -70°C. The MAb may be stored at 2°- 7°C for several weeks. Refer to the current reagent data sheet for additional details.

3.3.9 Polyclonal *L. pomona* antiserum, rabbit origin

Obtain the rabbit antiserum from the CVB. Store long-term at -70°C. The antiserum may be stored at 2°- 7°C for several weeks. Refer to the current reagent data sheet for additional details.

3.3.10 Bacterins containing *L. pomona* antigen

1. Reference bacterin
2. Test bacterin(s)

CRITICAL CONTROL POINT: Ideally, the reference and test bacterins should be produced by the same Outline of Production. If reference formulation differs from that of the test bacterin, the assay must be validated to show that this does not adversely affect assay performance or accuracy of results.

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

3.4 Preparation of the sample

Many bacterins do not require antigen-elution treatment prior to being serially diluted in twofold increments with PBS-Tween 20. Test representative batches of each adjuvanted product with and without each antigen-elution treatment to determine if the treatment specifically enhances antigen capture. If no enhancement of antigen capture can be demonstrated, test the bacterins without antigen-elution treatment. Treat the reference bacterin and the test bacterins by the same elution procedure. Alternative elution procedures, other than those described here, may be more appropriate for some bacterins.

3.4.1 Aluminum hydroxide-adjuvanted bacterins

Bacterins adjuvanted with aluminum hydroxide may be treated with either sodium citrate or phosphate buffer prior to making serial twofold dilutions in PBS-Tween 20.

1. Sodium citrate elution

Mix 1 g sodium citrate with 10 mL of bacterin (10% w/v). Place on an orbital shaker (100-120 rpm) overnight at 36°- 38°C. Consider treated bacterin to be undiluted.

2. Phosphate buffer elution

Mix 1 mL of phosphate elution buffer with 1 mL of bacterin. Place on an orbital shaker (100-120 rpm) overnight at 36°- 38°C. Consider treated bacterin to be diluted 1:2.

3.4.2 Oil-adjuvanted bacterins

Mix 1 mL of sodium deoxycholate (desoxycholate) elution buffer with 1 mL of bacterin. Place on an orbital shaker (100-120 rpm) overnight at 20°- 25°C. Consider treated bacterin to be diluted 1:2.

4. Performance of the Test

4.1 Dilute *L. pomona* rabbit antiserum to the current use dilution (refer to current reagent data sheet) in cold carbonate coating buffer. Dispense 100 µL into each well of a 96-well microtitration plate (test plate). Seal plate and incubate at 2°- 7°C for 16 to 20 hours. Coated plates may be stored at 2°- 7°C for up to 5 days.

4.2 Make twofold dilutions of reference and test bacterins, using PBS-Tween 20 as a diluent. Add 125 µL PBS-Tween 20 to each well of a clean microtitration plate (transfer

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

plate). Place 125 μL of bacterin in the first well of each row. Test each bacterin in at least 2 replicate rows. Test the reference bacterin and the test bacterin on the same plate. Use a multichannel pipetting device to make serial twofold dilutions of each bacterin across the plate (125 μL transfer volume). Reserve at least 2 unused wells on each plate to serve as blanks.

The use of at least 7 serial twofold dilutions per bacterin is recommended. Ideally, the selected bacterin dilutions should delineate the sigmoid curve from antigen saturation to antigen extinction for each bacterin. The dilutions used for the reference bacterin and the test bacterin may differ.

4.3 Wash the test plate(s) once with PBS-Tween 20. An automatic plate washer (200-250 μL /well, 10- to 40-second soak cycle, 3 wash cycles) may be used, or the plates may be washed by hand. Tap the plates upside down on absorbent material to remove residual fluid.

4.4 Use a multichannel pipetting device to transfer the bacterin dilutions from the transfer plates to the coated ELISA plates (100 μL /well). Seal the ELISA plates and incubate the plates for 60-90 minutes at 36°- 38°C.

4.5 Wash the ELISA plates 3 times with PBS-Tween 20 as in **Section 4.3**.

4.6 Dilute MAb 2D7 to the current use dilution (refer to current reagent data sheet) in antibody diluent, and add 100 μL to each well of the test plate. Incubate the plates for 55 to 65 minutes at 36°- 38°C.

4.7 Wash the ELISA plates 3 times with PBS-Tween 20 as in **Section 4.3**.

4.8 Dilute the goat anti-mouse IgA horseradish peroxidase conjugate 1:1000 in antibody diluent; or to the appropriate use dilution as specified by manufacturer or determined by checkerboard titration for that lot. Add 100 μL to each well of the test plate. Incubate for 30 to 60 minutes at 36°- 38°C.

4.9 Wash the ELISA plates 3 times with PBS-Tween 20 as in **Section 4.3**.

4.10 Add ABTS substrate (100 μL) to all wells, and incubate plates for 15 to 30 minutes at 36°- 38°C.

4.11 Read plates at 405/490 nm. Calculate the mean absorbance for the blank wells. Subtract the mean absorbance of the blank wells from each bacterin test well absorbance value prior to data analysis.

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

5. Interpretation of the Test Results

5.1 Relative potency calculation method

5.1.1 Use the current version of the *Relative Potency Calculation Software* (RelPot) to calculate the relative potency of the test bacterin as compared to that of the reference bacterin.

5.1.2 Do not use bacterin dilutions with mean absorbance values <0.050 (after subtraction of the mean absorbance of the blank) in the relative potency calculations.

5.1.3 Do not use regression lines with slopes >-0.100 in relative potency calculations. Enter a minimum slope (Min Slope) assay parameter of 0.100 in the spreadsheet in place of the 0.000 default.

5.1.4 Enter the reference and test bacterin data, and execute the relative potency program as outlined in the current version of **SAM 318**.

5.1.5 Report the highest relative potency (RP) value included in the top scores from each test as the RP for the test bacterin.

5.2 Requirements for a valid assay

5.2.1 An assay must meet the validity requirements in the current version of **SAM 318** to be considered valid.

5.2.2 Lines determined by first-order linear regression of at least 3 contiguous points must have a correlation coefficient (r) of ≥ 0.95 .

5.2.3 The reference regression line and the test bacterin regression line must show parallelism (slope ratio ≥ 0.80).

5.2.4 Assays that are not valid may be repeated up to a maximum of 3 times. If a valid assay cannot be achieved with 3 independent assays, the test bacterin is unsatisfactory.

5.3 Requirements for a satisfactory test bacterin

To be considered satisfactory, a test bacterin must have an RP value of ≥ 1.0 . Test bacterins with RP values <1.0 on a valid assay may be retested by conducting 2 independent replicate tests in a manner identical to the initial test. If both retests are valid and the reported RP values of both of the retests are ≥ 1.0 , the test bacterin is satisfactory.

Supplemental Assay Method for *In vitro* Potency Testing of *Leptospira interrogans* Serovar *pomona* Bacterins

6. Report of Test Results

Report the results of the test(s) as described by standard operating procedures.

7. Summary of Revisions

Version .03

- Minor changes have been made to update the document to current formatting practices.

Version .02

This document was revised to clarify practices currently in use at the Center for Veterinary Biologics and to provide additional detail. While no significant changes were made that impact the outcome of the test, the following changes were made to the document:

- **2.2.7** Desoxycholate changed to deoxycholate to reflect Sigma catalog listing (the two chemicals are equivalent).
- **4.2** Test well replicates changed from 3 to 2.
- References to the current reagent data sheet were added throughout the document as appropriate.
- The contact person has been changed to Mary C. Rasmusson.