

Work Task C10: Razorback Sucker Growth Studies

FY07 Estimates	FY07 Actual	Cumulative Accomplishment Through FY07	FY08 Approved Estimate	FY09 Proposed Estimate	FY10 Proposed Estimate	FY11 Proposed Estimate
\$125,000	\$106,383	\$169,901	\$125,000	\$125,000	\$125,000	\$125,000

Contact: Ty Wolters, (702) 293-8463, twolters@lc.usbr.gov

Start Date: FY06

Expected Duration: FY11

Long-term Goal: Seek measures to improve quantity, quality, and cost effectiveness of RASU reared for the Fish Augmentation Program.

Conservation Measures: RASU3, RASU4, and RASU6.

Location: Various locations including hatcheries, rearing ponds, universities, and private research facilities.

Purpose: Evaluate factors affecting growth of subadult RASU to maximize total length at release and reduce rearing time in hatchery.

Connections with Other Work Tasks (past and future): This work is similar to actions in C11 and shares some activities (concurrent studies at same locations). Also, a workshop for fish culturists was held in FY07, which focused on culture needs for both RASU (C10) and BONY (C11).

Project Description: Provides funding over a 5-year period for investigations into rearing and culture of RASU. The goal is to investigate ways to accelerate growth of RASU through manipulation of physical, chemical, and biological attributes of the rearing environment (e.g., manipulate feed, fish density, water temperature, water hardness, turbidity, lighting, presence/absence of cover). Current hatchery practices rear fish that are 250-300 mm TL in roughly 3 years. However, numerous observations during recent rearing and culture of RASU show a wide range in growth rates for this species, and it is possible to have 100, 200, and 300 mm TL fish from the same year class on station at the same time. In general, 25% of a RASU year class exhibit accelerated growth, 50% show moderate growth, and 25% demonstrate slow growth.

The species is a rare fish for which only limited life-history data exist, and data that exist are mostly for adults, not young life stages such as those being reared in hatcheries. As more fish are reared, released, and followed, more life-history data are being collected. Much of this information may be important to fish culturists. For example, the fact that young RASU were nocturnal was determined in 1992 by observations of biologists from the Lake Mohave NFWG.

Even so, hatchery managers are just now testing night-time feeding regimes. Active culture of RASU is a young science; many of the techniques initially used for rearing this species originated in the culture of rainbow trout, a species actively cultured for more than 50 years. Only during the past decade was it conclusively determined that a high-protein trout diet results in spinal deformities in fingerling RASU. As a final example, it was not recognized until the 1980s that adult RASU can feed successfully in open water areas on zooplankton. Much of the existing literature up to that time was for riverine populations, and assumed that the adult RASU were only bottom feeders. This information may be vital in determining where feed should be introduced within the water column during the culturing process (sinking, floating, or suspension). These types of observations need to be recongnized, then hypotheses developed, and finally tests of the hypotheses designed and conducted.

Previous Activities: Literature reviews were conducted to compile information on rearing these fish. These included site visits to facilities acitvely culturing RASU to document successes and failures. Inquiries were made to field biologists and technicians to document behavior of fish in the wild (i.e., daily activities such as feeding, resting, and use of cover in wild habitat). Ideas and hypotheses are being formulated into numerous small experiments, testing one variable at a time.

FY07 Accomplishments: Reclamation is working cooperatively with AGFD to study factors affecting growth of RASU. A report was submitted by AGFD titled, *Factors Affecting Growth of RASU in Captivity: Literature Review and Knowledge Assessment*. This report is a compilation of information regarding RASU rearing techniques in both hatcheries and natural rearing areas. The intent is to incorporate learned information and the best technologies into Bubbling Ponds SFH for facility improvements to accelerate RASU growth and improve survival. A Native Fish Culture Workshop was held in August 2007, which focused on RASU and BONY culturing practices and shared results of summary studies on existing culture techniques for RASU and BONY. Testing-apparatus designs were reviewed for installation at Bubbling Ponds SFH.

FY08 Activities: A new agreement was signed early in FY08 between Reclamation and USFWS testing the feasibility of polyculture for RASU and BONY in the same ponds. RASU and BONY that had been polycultured were harvested in December 2007, and BONY reach 300+ mm TL and RASU grew to be 400+ mm TL. The USFWS is also conducting RASU growth studies at Willow Beach NFH to determine density levels and feeding rates for rearing RASU from 300 mm up to 500 mm TL to accelerate brood stock development in Lake Mohave. Preliminary results from these studies will be written into reports in FY08.

Proposed FY09 Activities: Research investigations from a priority list of research needs developed at the fish culture workshop will be designed and conducted.

Pertinent Reports: The scope of work is available upon request from the LCR MSCP.