




HIV Surveillance in Women

2018

(preliminary)

Diagnoses of HIV Infection among Female Adults and Adolescents, by Race/Ethnicity, 2010–2017—United States and 6 Dependent Areas



^a Hispanics/Latinos can be of any race.



Deaths of Female Adults and Adolescents with Diagnosed HIV Infection by Race/Ethnicity, 2017—United States

| Race/ethnicity | No. | Rate | % |
|--|--------------|------------|------------|
| American Indian/Alaska Native | 10 | 1.0 | 0.3 |
| Asian ^a | 14 | 0.2 | 0.4 |
| Black/African American | 2,242 | 12.7 | 57.5 |
| Hispanic/Latino ^b | 564 | 2.5 | 14.5 |
| Native Hawaiian/other Pacific Islander | 4 | 1.7 | 0.1 |
| White | 801 | 0.9 | 20.5 |
| Multiple races | 267 | 11.4 | 6.8 |
| Total | 3,902 | 2.8 | 100 |

Note. Deaths of persons with diagnosed HIV infection may be due to any cause. Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.



^b Hispanics/Latinos can be of any race.



Rates of Female Adults and Adolescents Living with Diagnosed HIV Infection Year-end 2017—United States and 6 Dependent Areas

N = 241,585


Total Rate = 171.4




Note. Data are based on address of residence as of December 31, 2017 (i.e., most recent known address).



Diagnosed HIV Infection Classified as Stage 3 (AIDS) among Female Adults and Adolescents, 1985–2017—United States and 6 Dependent Areas



Stage 3 (AIDS) Classifications among Female Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity and Year of Classification 1985–2017—United States and 6 Dependent Areas




^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Percentages of Stage 3 (AIDS) Classifications among Female Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity and Year of Classification 1985–2017—United States and 6 Dependent Areas



^a Hispanics/Latinos can be of any race.


^b Includes Asian/Pacific Islander legacy cases.



Rates of Female Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2017—United States and 6 Dependent Areas

N = 127,177


Total Rate = 90.2



Note. Data are based on address of residence as of December 31, 2017 (i.e., most recent known address).



Diagnoses of HIV Infection and Population among Female Adults and Adolescents, by Race/Ethnicity, 2018—United States



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Female Adults and Adolescents by Race/Ethnicity, 2018—United States


| Race/ethnicity | No. | Rate | % |
|--|--------------|------------|------------|
| American Indian/Alaska Native | 32 | 3.2 | 0.4 |
| Asian | 103 | 1.2 | 1.4 |
| Black/African American | 4,114 | 23.1 | 57.6 |
| Hispanic/Latino ^a | 1,183 | 5.2 | 16.6 |
| Native Hawaiian/other Pacific Islander | 5 | 2.1 | 0.1 |
| White | 1,526 | 1.7 | 21.4 |
| Multiple races | 176 | 7.3 | 2.5 |
| Total | 7,139 | 5.1 | 100 |

Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.




Diagnoses of HIV Infection among Female Adults and Adolescents by Region and Race/Ethnicity, 2018—United States



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay.
^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Female Adults and Adolescents, by Race/Ethnicity and Transmission Category, 2018—United States and 6 Dependent Areas



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Female Adults and Adolescents by Transmission Category and Age at Diagnosis 2018—United States and 6 Dependent Areas

| Transmission category | Age group (years), % | | | | |
|-----------------------------------|----------------------|----------------|------------------|------------------|----------------|
| | 13–19 N=263 | 20–24 N=715 | 25–34 N=1,937 | 35–44 N=1,730 | ≥45 N=2,575 |
| Injection drug use | 8.7 | 13.8 | 16.1 | 15.7 | 13.4 |
| Heterosexual contact ^a | 86.4 | 85.1 | 83.3 | 84.1 | 86.4 |
| Other ^b | 4.9 | 1.0 | 0.6 | 0.2 | 0.2 |
| Total | 100 | 100 | 100 | 100 | 100 |

Note. Data for the year 2017 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Rates of Diagnoses of HIV Infection among Female Adults and Adolescents 2018—United States and 6 Dependent Areas

N = 7,220

Total Rate = 5.1



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay.



Stage 3 (AIDS) Classifications among Female Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity, 2018—United States

| Race/ethnicity | No. | Rate | % |
|--|--------------|------------|------------|
| American Indian/Alaska Native | 15 | 1.5 | 0.4 |
| Asian ^a | 57 | 0.7 | 1.4 |
| Black/African American | 2,492 | 14.0 | 60.7 |
| Hispanic/Latino ^b | 625 | 2.7 | 15.2 |
| Native Hawaiian/other Pacific Islander | 0 | 0.0 | 0.0 |
| White | 743 | 0.8 | 18.1 |
| Multiple races | 174 | 7.2 | 4.2 |
| Total | 4,106 | 2.9 | 100 |


Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.



Stage 3 (AIDS) Classifications among Female Adults and Adolescents with Diagnosed HIV Infection, by Region and Race/Ethnicity, 2018—United States



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay.

^a Hispanics/Latinos can be of any race.



Stage 3 (AIDS) Classifications among Female Adults and Adolescents with Diagnosed HIV Infection, by Transmission Category and Age at Diagnosis 2018—United States and 6 Dependent Areas

| Transmission category | Age group (years), % | | | | |
|-----------------------------------|----------------------|----------------|----------------|------------------|----------------|
| | 13–19 N=44 | 20–24 N=162 | 25–34 N=812 | 35–44 N=1,109 | ≥45 N=2,022 |
| Injection drug use | 6.4 | 8.9 | 13.5 | 16.4 | 21.1 |
| Heterosexual contact ^a | 50.7 | 73.1 | 83.8 | 83.4 | 78.4 |
| Other ^b | 43.0 | 18.0 | 2.7 | 0.2 | 0.5 |
| Total | 100 | 100 | 100 | 100 | 100 |

Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Rates of Stage 3 (AIDS) Classifications among Female Adults and Adolescents with Diagnosed HIV Infection, 2018—United States and 6 Dependent Areas

N = 4,149

Total Rate = 2.9



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay.

