



Observed and forecasted cumulative COVID-19 deaths in the US




Forecasts shown here fall into one of three categories. The LANL model is explicitly 'unconditional' on any particular interventions being in place. The IHME and MOBS_NEU models are conditional on existing social distancing measures continuing through the projected time-period. The CU models make different assumptions about the effectiveness of current interventions. Intervals shown are at the 95% uncertainty level.

Observed and forecasted cumulative COVID-19 deaths in the US




Forecasts shown here fall into one of three categories. The LANL model is explicitly 'unconditional' on any particular interventions being in place. The IHME and MOBS_NEU models are conditional on existing social distancing measures continuing through the projected time-period. The CU models make different assumptions about the effectiveness of current interventions. Intervals shown are at the 95% uncertainty level.

Observed and forecasted cumulative COVID-19 deaths in the US




Forecasts shown here fall into one of three categories. The LANL model is explicitly 'unconditional' on any particular interventions being in place. The IHME and MOBS_NEU models are conditional on existing social distancing measures continuing through the projected time-period. The CU models make different assumptions about the effectiveness of current interventions. Intervals shown are at the 95% uncertainty level.

Observed and forecasted cumulative COVID-19 deaths in the US




Forecasts shown here fall into one of three categories. The LANL model is explicitly 'unconditional' on any particular interventions being in place. The IHME and MOBS_NEU models are conditional on existing social distancing measures continuing through the projected time-period. The CU models make different assumptions about the effectiveness of current interventions. Intervals shown are at the 95% uncertainty level.

Observed and forecasted cumulative COVID-19 deaths in the US




Forecasts shown here fall into one of three categories. The LANL model is explicitly 'unconditional' on any particular interventions being in place. The IHME and MOBS_NEU models are conditional on existing social distancing measures continuing through the projected time-period. The CU models make different assumptions about the effectiveness of current interventions. Intervals shown are at the 95% uncertainty level.

Observed and forecasted cumulative COVID-19 deaths in the US



Forecasts shown here fall into one of three categories. The LANL model is explicitly 'unconditional' on any particular interventions being in place. The IHME and MOBS_NEU models are conditional on existing social distancing measures continuing through the projected time-period. The CU models make different assumptions about the effectiveness of current interventions. Intervals shown are at the 95% uncertainty level.

Observed and forecasted cumulative COVID-19 deaths in the US



- CU 20% contact reduction
- CU 40% contact reduction
- LANL
- observed data
- CU 30% contact reduction
- IHME
- MOBS

Forecasts shown here fall into one of three categories. The LANL model is explicitly 'unconditional' on any particular interventions being in place. The IHME and MOBS_NEU models are conditional on existing social distancing measures continuing through the projected time-period. The CU models make different assumptions about the effectiveness of current interventions. Intervals shown are at the 95% uncertainty level.